PHASE TRANSITIONS OF COMPOSITION SCHEMES: MITTAG-LEFFLER AND MIXED POISSON DISTRIBUTIONS

BY CYRIL BANDERIER ¹, MARKUS KUBA ² AND MICHAEL WALLNER ³

¹CNRS & Université Sorbonne Paris Nord, https://lipn.fr/~banderier/

²University of Applied Sciences Technikum Wien, https://dmg.tuwien.ac.at/kuba/

³Technische Universität Wien, https://dmg.tuwien.ac.at/mwallner/

Multitudinous probabilistic and combinatorial objects are associated with generating functions satisfying a composition scheme F(z) = G(H(z)). The analysis becomes challenging when this scheme is critical (i.e., G and H are simultaneously singular). Motivated by many examples (random mappings, planar maps, directed lattice paths), we consider a natural extension of this scheme, namely F(z,u) = G(uH(z))M(z). We also consider a variant of this scheme, which allows us to analyse the number of H-components of a given size in F.

We prove that these two models lead to a rich world of limit laws, where we identify the key rôle played by a new universal law introduced in this article: the three-parameter Mittag-Leffler distribution, which is essentially the product of a beta and a Mittag-Leffler distribution. We also prove (double) phase transitions, additionally involving Boltzmann and mixed Poisson distributions, bringing a unified explanation of the associated thresholds. In all cases we obtain moment convergence and local limit theorems. We end with extensions of the critical composition scheme to a cycle scheme and to the multivariate case, leading to product distributions. Applications are presented for random walks, trees (supertrees of trees, increasingly labelled trees, preferential attachment trees), triangular Pólya urns, and the Chinese restaurant process.

** This article is kindly devoted to Alois Panholzer, on the occasion of his 50th birthday. **

CONTENTS

1	murc	duction	4
2	New	main results	4
	2.1	Composition schemes analysed in this article	4
	2.2	Phase transitions	6
	2.3	Plan of the paper	7
3	Sing	ularity analysis, stable laws, and mixed Poisson distributions	7
	3.1	Singularity analysis and asymptotic expansions	8
	3.2	Probability distribution melting pot	. 1
4	Exte	nded composition scheme	8
5	Size-	-refined composition scheme	1/2
6	Appl	lications and examples	3
	6.1	Core size of supertrees and a confluent example	3
	6.2	Root degree and branching structure in bilabelled increasing trees	1
	6.3	Returns to zero: walks and bridges with drift zero	13
	6.4	Initial returns in coloured bridges	5
	6.5	Sign changes in walks	,

MSC2020 subject classifications: Primary 60C05; secondary 60E07, 60G50, 05A15.

Keywords and phrases: Mixed Poisson distributions, Mittag-Leffler distributions, stable laws, Boltzmann distributions, analytic combinatorics, generating functions, singularity analysis, critical composition schemes.

	6.6	Tables in the Chinese restaurant process	39
	6.7	Triangular urn models and the three-parameter Mittag-Leffler distribution	42
7	Furt	her extensions	44
	7.1	Critical cycle scheme	44
	7.2	Multivariate critical composition schemes	45
8	Con	clusion	52
Αc	know	eledgements	54
Re	feren	ces	54

1. Introduction. Many combinatorial structures are an assemblage of more basic building blocks, and this situation is ubiquitous in many different fields, such as combinatorics, probability theory, and statistical mechanics. It appears for example in permutations, random walks, random mappings, random forests, parking functions, Pólya–Eggenberger urn models, (Bienaymé)–Galton–Watson processes, destruction procedures in simply generated trees, inversions in labelled tree families, generalized plane-oriented recursive trees (scale-free trees), set or integer partitions with some constraints, sequences of words, tilings, different families of graphs or maps, etc.; see, e.g., [6–8, 13–15, 30, 31, 39, 42–44, 53, 90, 109, 110, 112, 123, 131]. In the language of generating functions, one then has a functional composition scheme such as

$$F(z) = G(H(z)).$$

Let us illustrate this composition scheme with some examples (each of them being in fact the starting point of many theorems in the literature): a random forest is a set of random trees, a permutation is a set of cycles, a bridge (a random walk on \mathbb{Z}) is a sequence of arches, functional mappings are sets of cycles of Cayley trees, supertrees are trees in which each leaf is replaced by another family of trees, an integer partition is a sequence of parts, the factorization of a polynomial in a finite field is a multiset of irreducible factors, and, following the work of Tutte, several important families of planar maps can be seen as a "simple core" in which each node is replaced by some "simple map", etc. The reader can find many other examples illustrating the universality of the scheme F(z) = G(H(z)) in the wonderful book by Flajolet and Sedgewick on analytic combinatorics [42, Sec. VI.9]. Structurally, this composition scheme is at the heart of many fascinating phase transition phenomena (analytically corresponding, e.g., to coalescing saddle points or to confluence of singularities). More precisely, let $G(z) = \sum_{n \geq 0} g_n z^n$ and $H(z) = \sum_{n \geq 0} h_n z^n$ be analytic functions at the origin with nonnegative coefficients and H(0) = 0. Let ρ_G and ρ_H be the radii of convergence of G(z) and H(z), respectively. Then, following [7, 42], we focus on critical composition schemes.

DEFINITION 1.1 (Critical composition scheme). The composition scheme F(z) = G(H(z)) is *critical* if it satisfies $H(\rho_H) = \rho_G$.

In other words, G and H are here concomitantly singular. We will assume throughout this work that we are always in the critical case (the asymptotic analysis is straightforward otherwise). Note that this terminology is a generalization of the notion of critical/supercritical/subcritical Galton–Watson processes, initially popularized by Harris for neutron branching processes [60].

Often, G(z) and H(z) are the counting series of certain combinatorial families $\mathcal G$ and $\mathcal H$ such that $\mathcal F=\mathcal G(\mathcal H)$. We refer to the first part of [42] for a more detailed presentation of this combinatorial approach, starting from the so-called atoms, and then assembling them into more elaborate structured blocks via combinatorial constructors. Some important subclasses of such structures were also subject of more probabilistic approaches; see, e.g., [1,4,53,80].

Now, our goal is to analyse probabilistic properties of critical compositions like

$$F(z, u) = G(uH(z)),$$

where u marks each occurrence of objects of \mathcal{H} . From a combinatorial perspective, F(z,u) enumerates \mathcal{F} -structures of size n made of k "building blocks from \mathcal{H} " (also simply called \mathcal{H} -components), i.e., \mathcal{G} -structures made of k atoms, where each atom is then replaced by an \mathcal{H} -block (which is itself a structured set of atoms). For any combinatorial structure in \mathcal{F} , its corresponding \mathcal{G} -structure is sometimes called its $core^1$ (or "skeleton", or "backbone").

A first natural question is what is the typical size of this core, i.e., what is the *typical* number of \mathcal{H} -components? Such insight helps, for example, to make many algorithms on combinatorial structures more efficient, as Knuth shows in [79] (see also [7], where this insight is used to design faster random generation algorithms). To answer this question, one considers the discrete random variable X_n associated with this core size in a uniformly chosen object of size n. Its probability mass function is obtained by extraction of coefficients:

$$\mathbb{P}\{X_n = k\} = \frac{[z^n u^k] F(z, u)}{[z^n] F(z, 1)} = g_k \frac{[z^n] H(z)^k}{f_n},$$

where $[z^n]$ denotes the extraction of coefficient operator $[z^n] \sum_n f_n z^n = f_n$. As H(z) has typically a singular expansion of the type

$$H(z) = \tau_H + c_H \left(1 - \frac{z}{\rho_H}\right)^{\lambda_H} + \dots,$$

this implies that the asymptotic behaviour of $\mathbb{P}\{X_n = k\}$ depends on the exponent λ_H (which is called the *singular exponent* of H). Actually, this exponent even plays a key rôle, as it entails four types of asymptotic behaviour for X_n :

- For $\lambda_H > 2$ the limit law is related to a Gaussian law.
- For 1 < λ_H < 2 the limit law is related to a stable law of parameter λ_H (this distribution is supported on ℝ and possesses moments up to order λ_H; e.g., for λ_H = 3/2 this gives the map-Airy distribution).
- For $0 < \lambda_H < 1$ we will show that the limit law is related to a stable law of parameter λ_H , or more precisely to a generalized Mittag-Leffler distribution (this distribution is supported on \mathbb{R}^+ and has moments of any order; e.g., for $\lambda = 1/2$, this gives the Rayleigh distribution).
- For $\lambda_H < 0$ the scheme is not critical because the function H(z) diverges at $z = \rho_H$, and thus leads to a singularity of G(H(z)) at some $z < \rho_H$. Such a scheme is called supercritical and typically leads to a Gaussian limit law.

The case $\lambda_H < 0$ is analysed by Flajolet and Sedgewick $[42]^2$, building on the seminal work of Bender [17]. The three cases $0 < \lambda_H < 1$, $1 < \lambda_H < 2$, and $\lambda_H > 2$ were partially analysed by Banderier, Flajolet, Soria, and Schaeffer [7, Theorem 12], but without a precise statement for the right renormalizations in the limit laws. It is partly due to the fact that the initial motivation of the authors of [7] was to analyse the core of planar maps, so they focused on the subcase $\lambda_H = \frac{3}{2}$, which corresponds to the map-Airy distribution. Thus, a more complete analysis of the composition scheme in these three regions of λ_H remained to be done.

¹The word "core" comes from the theory of graphs and maps, where this composition scheme is natural and was, e.g., analysed by Janson, Knuth, Łuczak, and Pittel [75] for graphs and by Banderier, Flajolet, Schaeffer, and Soria [7] for maps.

²We refer to *The Hitchhiker's Guide to the Galaxy*, by Douglas Adams, for an interesting epistemology of [42].

For sure, we expected that the different possible analytic behaviours of G introduce further subcases, but we were surprised that the detailed analyses were much more challenging than expected: As we shall see, they require several new ingredients. Our identification of the limit laws involves moment-tilted distributions, product distributions, and Boltzmann distributions (see Section 3.2 for a formal presentation of these three types of distributions and their key properties). Therefore, the first main objective of our work is to give a complete landscape of the limit laws associated with critical composition schemes. We analyse the case $0 < \lambda_H < 1$ in this article, and the cases $1 < \lambda_H < 2$ and $\lambda_H > 2$ in our companion article [11].

Our second main objective is to explain the phase transitions observed for the *number of* \mathcal{H} -components of a given size. This builds on the work of Panholzer and the second author [90], in which they started to unify the diversity of limit laws encountered in these phase transitions under the umbrella of *mixed Poisson distributions*, and relies on the study of a size-refined composition scheme that we detail in the next section.

2. New main results.

2.1. Composition schemes analysed in this article. In this work we complete the analysis of critical compositions with exponent $0 < \lambda_H < 1$. Motivated by models associated with directed lattice paths [6, 13–15, 131] and triangular Pólya urns [39, 72, 74], we consider two natural schemes (Equations (2) and (4) hereafter). They contain a multiplication by a factor M(z) which plays a non-trivial rôle in our identification of the corresponding limit laws as a three-parameter generalization of Mittag-Leffler distributions. In Section 6, we illustrate via various examples how these two schemes unify and refine many previous studies.

Firstly, we consider the following extended composition scheme

(1)
$$F(z) = G(H(z)) \cdot M(z),$$


for some functions F/G/H/M analytic at the origin, with nonnegative coefficients. Such schemes are *critical* if $\rho_G = H(\rho_H)$ (like in Definition 1.1) and additionally satisfy $\rho_M \ge \rho_H$, where ρ_M is the radius of convergence of M(z) (the analysis is straightforward if the extended composition scheme is not critical). In order to enumerate the family $\mathcal F$ according to the occurrences of $\mathcal H$ -components, we consider

(2)
$$F(z,u) = G(uH(z)) \cdot M(z),$$

which from now on we will refer to as the extended composition scheme. Equivalently, $[z^n u^k] F(z,u)$ is the number of \mathcal{F} -structures of size n having k \mathcal{H} -components. This generalizes the schemes G(uH(z)) and A(uz)B(z) analysed in [7] and [42,58], respectively. The corresponding random variable X_n has a probability mass function given by

(3)
$$\mathbb{P}\{X_n = k\} = \frac{[z^n u^k] F(z, u)}{[z^n] F(z, 1)} = g_k \frac{[z^n] H(z)^k \cdot M(z)}{f_n}.$$

Our first main result is Theorem 4.1, in which we give explicit expressions for the asymptotics of the factorial moments of X_n , the limit distribution of X_n (suitably normalized), and its density function. It appears that this limit distribution differs depending on some relationship between the singular exponents of G(z), H(z), and M(z), as summarized in Table 1 (where we write $X_n \sim c_n \cdot \mathcal{D}$ when $c_n^{-1} \cdot X_n \to \mathcal{D}$ in distribution for $n \to \infty$). In addition to these convergences in distribution, we prove moment convergence for the continuous limit laws.



Three asymptotic behaviours according to the singular exponents $\lambda_G/\lambda_H/\lambda_M$ of G/H/M: the number X_n of \mathcal{H} -components in \mathcal{F} -structures of size n in the extended scheme (1) is given by three completely different types of limit laws, depending on whether the scheme is analytically pure/confluent/degenerate (Theorems 4.1, 4.3, and 4.4).

Secondly, we consider a *size-refined composition scheme* which allows us to capture some threshold phenomenon via a bivariate generating function $F(z, v)^3$:

(4)
$$F(z,v) = G(H(z) - (1-v)h_j z^j) \cdot M(z), \quad j \in \mathbb{N}.$$

In this scheme, $[z^n v^k] F(z, v)$ is therefore the number of \mathcal{F} -structures of size n having k \mathcal{H} -components each of size j; this is combinatorially summarized by:

$$\mathcal{F} = \mathcal{G}(\mathcal{H}_{\neq i} + v\mathcal{H}_{=i}) \times \mathcal{M}.$$

Assuming the uniform distribution amongst structures of size n, let $X_{n,j}$ be the number of \mathcal{H} -components of size j inside \mathcal{F} -structures⁴ of size n. Thus, the random variables $X_{n,j}$ naturally refine the distribution of the core size X_n given in (3), since

$$\sum_{j \in \mathbb{N}} X_{n,j} = X_n.$$

Formula (4) implies that one has

(5)
$$\mathbb{P}\{X_{n,j} = k\} = \frac{[z^n v^k] F(z, v)}{[z^n] F(z, 1)} = \frac{h_j^k}{k!} \frac{[z^{n-jk}] G^{(k)} (H(z) - h_j z^j) M(z)}{f_n}.$$

Our second main result is Theorem 5.1, in which we prove that the factorial moments of $X_{n,j}$ are asymptotically of mixed Poisson type, and establish a convergence in distribution, with convergence of all moments, towards explicit limit distributions.

We extend these two main results to the composition schemes involving a logarithmic singularity in Theorem 7.1 and 7.3, leading to Mittag-Leffler distributions.

Then, our third main result is Theorem 7.4 in which we give a multivariate generalization with arbitrary many variables, leading to Dirichlet product distributions.


³Note that we still use the letter F to denote the main function. The auxiliary variable is now v and no more u as we are marking a different parameter. This choice avoids more cumbersome notation and is also motivated by the fact that we always have F(z,1) = F(z).

⁴Flajolet and Sedgewick call the corresponding distribution the *profile* of the combinatorial object, by analogy with the profile of integer compositions; see [42, p. 169, p. 451, p. 632].

2.2. *Phase transitions*. Analogously to what can happen in physics or chemistry for some small change of temperature, pressure, or concentration, a phase transition in mathematics corresponds to a sudden non-smooth change of properties under smooth variation of the parameters. Such non-smooth changes are thus analytically reflected by a singularity of some function associated with these properties. For combinatorial structures, generating function methods were successfully used to analyse such phase transitions, e.g., for random graphs or planar maps [7, 50, 51, 75], for satisfiability problems [105, 123], and for many other problems [42, 46, 47, 63–65].

Usually, the main problems are to locate the phase transition, to properly describe the phase transition via special functions in terms of the involved parameters, and to give intuitive explanations of the observed phenomena. Some cases even exhibit two successive phase transitions. In probability theory, such a double phase transition occurs for example with the binomial distribution $\mathbb{P}\{X_n=k\}=\binom{n}{k}p^k(1-p)^{n-k}$ when p depends on n, with p bounded away from one. It indeed leads to the following trichotomy involving a continuous to discrete to degenerate phase transition: Firstly, when both $\mathbb{E}(X_n)=p(n)\cdot n$ and the variance $\mathbb{V}(X_n)$ tend to infinity, then a central limit theorem for X_n follows. Secondly, if $p(n)\cdot n\to \lambda>0$, then X_n degenerates to a Dirac distribution with all mass at 0.

The analysis of the size-refined composition scheme unravels, though more subtly, such a double phase transition: the random variable $X_{n,j}$ given by Equation (5) has three phases, each leading to its own limit law, visualized in Table 2.



Three successive régimes for the number $X_{n,j}$ of \mathcal{H} -components of size j in \mathcal{F} -structures of size n in the critical size-refined scheme (4), depending on the relation between j and n; this double phase transition is proven in Theorem 5.1.

Table 2 also motivates the following important remark.

REMARK 2.1 (Ubiquity of the exponent 1/3). Generating functions often have a dominant singularity of the square-root type (i.e., $\lambda_H=1/2$). This phenomenon is explained by the Drmota–Lalley–Woods theorem: Whenever H(z) can be defined by a strongly connected set of polynomial equations with nonnegative coefficients, it has a singular exponent 1/2 (see, e.g., [5,29,42]). Accordingly, in conjunction with Table 2, this explains why one observes a threshold at $j=n^{1/3}$ in many phase transitions; see Section 6.

One pleasant consequence of our work is that it gives a unified explanation of phase transitions from continuous to discrete observed in many examples: descendants in increasing trees [82], node degrees in increasing trees [83], block sizes in k-Stirling permutations [85], stopping times in urn models [87], death processes [88], inversions in labelled tree families [112], ancestors and descendants in evolving k-tree models [113].

These case by case studies lacked a proper comprehensive and uniform description of the arising phase transitions. So, instead of treating these combinatorial structures individually, we directly study the size-refined composition scheme (4). As summarized in Table 2, we show how the phase transitions for the random variable $X_{n,j}$ depend on the growth of j=j(n) with respect to the size n. We prove that the distribution of $X_{n,j(n)}$ is continuous for small values of j (a three-parameter generalization of the Mittag-Leffler distribution), or discrete for some threshold values of j (a Poisson distribution mixed with the previous Mittag-Leffler distribution), or a Dirac distribution for large values of j. We further exemplify these results on different processes, like the Chinese restaurant process, sign changes and returns to zero in random walks, and the branching structure of random trees.

- 2.3. Plan of the paper. In Section 3 we collect results from analytic combinatorics. We also present our basic assumptions on the generalized composition scheme and collect properties of various distributions that appear later in our main results. Section 4 is devoted to our results on the random variable X_n corresponding to the extended composition scheme (2) involving F(z,u). Section 5 contains the results for the random variable $X_{n,j}$ corresponding to the size-refined composition scheme (4) involving F(z,v) and exhibiting phase transitions. We also give the covariance and the correlation coefficient of X_{n,j_1}, X_{n,j_2} , observing again some phase transitions. In Section 6 we discuss various examples to which we apply our results. Finally, in Section 7, we analyse further extensions for a cycle scheme and for a multivariate critical composition scheme. We also present new examples for these two extensions.
- 3. Singularity analysis, stable laws, and mixed Poisson distributions. In this section, we first present a few important notions from analytic combinatorics [42] which we use to identify the radius of convergence and the singular exponents in our composition schemes. Then, we present a few results on the family of moment-tilted stable laws, based on James [67–69] and Janson [74]. We also collect properties of mixed Poisson distributions and their factorial moments. All of this allows us to identify in Sections 4 and 5 the distribution of the \mathcal{H} -components in our composition schemes.




FIG 1. As visually summarized by Flajolet in [38], three fundamental methods of asymptotic analysis require information on the function in different parts (shown here in red) of the complex plane. Flajolet and Odlyzko's singularity analysis [41] offers more powerful results, but requires analyticity in a Δ -domain (tastefully also sometimes called "Camembert domain" or "Pac-Man" by Flajolet himself!). This is the domain inside the blue curve, defined by $\Delta = \{z \in \mathbb{C} \text{ such that } |z| < \rho + \varepsilon \text{ and } \arg(z - \rho) > \theta \}$, for some $\varepsilon > 0$ and $0 < \theta < \pi/2$. This analyticity is agreeably typically granted for most combinatorial constructions (e.g., for the ones leading to meromorphic, algebraic-logarithmic, hypergeometric, or D-finite functions).

3.1. Singularity analysis and asymptotic expansions. Let

$$F(z) = \sum_{n \ge 0} f_n z^n$$

be a function with nonnegative coefficients f_n that is analytic in a Δ -domain (see Figure 1 for this notion) with a finite radius of convergence ρ and singular expansion

(6)
$$F(z) = P\left(1 - \frac{z}{\rho}\right) + c_F \cdot \left(1 - \frac{z}{\rho}\right)^{\lambda_F} (1 + o(1)),$$

where $\lambda_F \in \mathbb{R} \setminus \{0, 1, 2, ...\}$ is called the *singular exponent* (of F(z) at $z = \rho$), and where $P(x) \in \mathbb{C}[x]$ is a polynomial (of degree ≥ 1 for $\lambda_F > 1$, of degree 0 for $0 < \lambda_F < 1$, and P = 0 for $\lambda_F < 0$). Then, by standard singularity analysis using the transfer theorem [41, Sec. 2], if ρ is the unique singularity of F(z) in $|z| \leq \rho$, the Taylor series coefficients of F(z) satisfy

(7)
$$[z^n]F(z) = f_n = \frac{c_F}{\rho^n} \cdot \frac{n^{-\lambda_F - 1}}{\Gamma(-\lambda_F)} \cdot (1 + o(1)).$$

As the f_n 's are nonnegative, this implies the sign property

(8)
$$\operatorname{sgn}(c_F) = \operatorname{sgn}(\Gamma(-\lambda_F)),$$

i.e., due to the sign change of the gamma function at each negative integer we have $c_F > 0$ for $\lambda_F < 0$, $c_F < 0$ for $0 < \lambda_F < 1$, and $\operatorname{sgn}(c_F) = (-1)^{\lceil \lambda_F \rceil}$ for $\lambda_F > 1$.

Note that it is in general easy to get more asymptotic terms in (6), and that singularity analysis directly translates them into more asymptotic terms in (7). Moreover, if one has several dominant singularities, one has to sum the contributions of the local expansions at each singularity to get the asymptotics of the coefficients (for more details see [42, Chapter IV.6]; see also the rotation law in [16] for walks or trees with periodic offspring).

Note that algebraic functions constitute one of the main sources of functions satisfying the conditions of the expansion (6); indeed, they admit a Puiseux expansion

$$F(z) = \sum_{k>k_0} c_k \cdot (1 - z/\rho)^{k/r}$$
,

for $k_0, r \in \mathbb{Z}$ with $r \ge 1$. For example, in Section 6.1 we will encounter the Catalan generating function $1/2 - \sqrt{1 - 4z}/2$, for which one has $\rho = 1/4$, P(x) = 1/2, $k_0 = 0$, and r = 2.

This Catalan example is pleasantly simple, and thus obviously not generic, as its Puiseux expansion contains only two terms. In full generality such an expansion can involve an infinite number of terms whose sum is converging. Let us now analyse these singular expansions.

LEMMA 3.1 (Singular expansion). Let F(z) be a power series with nonnegative coefficients satisfying (6). Then F(z) has the following singular expansion

$$F(z) = \begin{cases} c_F \left(1 - \frac{z}{\rho}\right)^{\lambda_F} \left(1 + o(1)\right) & \text{if } \lambda_F < 0, \\ \tau_F + c_F \left(1 - \frac{z}{\rho}\right)^{\lambda_F} \left(1 + o(1)\right) & \text{if } 0 < \lambda_F < 1, \\ \tau_F + \sum_{i=1}^{\lfloor \lambda_F \rfloor} p_i \left(1 - \frac{z}{\rho}\right)^i + c_F \left(1 - \frac{z}{\rho}\right)^{\lambda_F} \left(1 + o(1)\right) & \text{if } \lambda_F > 1, \end{cases}$$

where $\tau_F = F(\rho) > 0$ for $\lambda_F > 0$ and $p_1 = -\rho F'(\rho) < 0$ for $\lambda_F > 1$.

PROOF. Firstly, if $\lambda_F < 0$, then the lowest order of the Puiseux expansion is λ_F . Secondly, for $\lambda_F > 0$, we rewrite (6) into

$$F(z) = \sum_{i=0}^{k} p_i (1 - z/\rho)^i + c_F (1 - z/\rho)^{\lambda_F} + \dots$$

Then, we have $p_0 = P(0) = F(\rho)$ which we define to be τ_F . We get $\tau_F > 0$ as it is an infinite convergent sum of nonnegative not-all-zero terms. Next, observe that the nonnegative coefficients of F(z) imply that $F'(\rho) > 0$. Thus, taking the derivative in the expansion of F(z) we get $\lim_{z\to\rho} F'(z) = +\infty$ for $0 < \lambda_F < 1$ and $p_1 = -\rho F'(\rho) \neq 0$ for $\lambda_F > 1$.

We now consider the critical scheme F(z) = G(H(z))M(z), assuming that G(z), H(z), and M(z) have a finite radius of convergence and a unique dominant singularity (i.e., the one of smallest modulus). By Pringsheim's theorem [42, p. 240] applied to each of these functions, the nonnegativity of its coefficients implies that this dominant singularity lies on the positive real axis and corresponds therefore to its radius of convergence denoted by ρ_G , ρ_H , and ρ_M .

Note that if M(z) has an infinite radius of convergence (denoted by $\rho_M = +\infty$) or if $\rho_M \neq \rho_H$, then the asymptotics are easily obtained via

$$[z^n]F(z) \sim \begin{cases} G(H(\rho_M))[z^n]M(z) & \text{if } \rho_M < \rho_H, \\ M(\rho_H)[z^n]G(H(z)) & \text{if } \rho_M > \rho_H. \end{cases}$$

Now, as in Lemma 3.1, we define for each function:

- the singular exponents λ_G , λ_H , and λ_M ,
- the constant terms τ_G , τ_H , and τ_M ,
- and the singular coefficients c_G , c_H , and c_M .

Thus, thanks to Equations (9a) and (9b), we can now focus (without loss of generality, or rather "without loss of difficulty!") on the case $\rho_M = \rho_H$ which is more involved as here G(z), H(z), and M(z) are all contributing to the asymptotics in a nontrivial way. Then, one gets different régimes (depending on λ_H) for the asymptotics of the coefficient of F(z). In this article we focus on the range $0 < \lambda_H < 1$, while we treat the other range $\lambda_H > 1$ in a companion article [11].

Note that, as our work extends to some interesting combinatorial cases where M(z) has a radius of convergence $\rho_M > \rho_H$, we will also encompass this case, for which it is then convenient to set $\lambda_M = +\infty$ (the singular exponent of M(z) at $z = \rho_H$ is infinite whenever M is analytical there). The case $\lambda_M = +\infty$ is thus archetypal of cases where M(z) only affects the asymptotics of f_n by a multiplicative constant like in Equation (9b).

We can now express the singular exponent of F in terms of those of G/H/M.

LEMMA 3.2. Let F(z) = G(H(z))M(z) be a critical composition scheme that is singular at ρ_H . Then, the singular exponent λ_H of H(z) satisfies $\lambda_H > 0$.

Moreover, for the range $0 < \lambda_H < 1$, the singular exponent λ_F of F(z) satisfies

$$\lambda_F = \min(\lambda_G \lambda_H, \lambda_H, \lambda_M, \lambda_G \lambda_H + \lambda_M).$$

For $\lambda_H > 1$, the singular exponent λ_F of F(z) satisfies

$$\lambda_F = \min(\lambda_G, \lambda_H, \lambda_M, \lambda_G + \lambda_M).$$

PROOF. The claim $\lambda_H > 0$ follows from $H(\rho_H) = \rho_G \in (0, \infty)$ as one would have $H(\rho_H) = +\infty$ if H(z) had a singularity of negative singular exponent at $z = \rho_H$.

Now, we plug the singular expansions from Lemma 3.1 at $z = \rho_H$ for G(z), H(z), and M(z) into F(z) = G(H(z))M(z). When $0 < \lambda_H < 1$ we get the following expansions (in which we omit the terms not contributing to the first-order asymptotics):

This lemma motivates the following definition.

DEFINITION 3.3 (Pure/confluent/degenerate composition schemes). Consider an extended or size-refined composition scheme (1) or (4) with a unique dominant singularity $\rho_F = \rho_H$, and with $0 < \lambda_H < 1$. It is either analytically

$$\begin{array}{ll} \bullet & \textit{pure} & \text{ if } \begin{cases} \lambda_G < 0 \text{ or } \\ 0 < \lambda_G < 1 \text{ and } \lambda_M > \min(\lambda_G \lambda_H, \lambda_H); \end{cases} \\ \bullet & \textit{confluent} & \text{ if } 0 < \lambda_G < 1 \text{ and } \lambda_M = \min(\lambda_G \lambda_H, \lambda_H); \end{cases} \\ \bullet & \textit{degenerate} & \text{ if } \begin{cases} \lambda_G > 1 \text{ or } \\ 0 < \lambda_G < 1 \text{ and } \lambda_M < \min(\lambda_G \lambda_H, \lambda_H). \end{cases} \end{array}$$

This is pictorially summarized by Figure 2.




FIG 2. The three different régimes (pure, confluent, degenerate) for extended or size-refined composition schemes: The Puiseux expansions of G/H/M go into resonance (or not), thus leading to these three cases.

We characterize the distributions associated with critical composition schemes in the analytically pure/confluent/degenerate cases in Theorems 4.1, 4.3, and 4.4. We now present some probabilistic results on the distributions which will appear in these theorems.

3.2. Probability distribution melting pot. First, we discuss properties of tilted distributions, their link with positive stable distributions, and we introduce the three-parameter Mittag-Leffler distribution. Then, we collect properties of a family of discrete distributions called mixed Poisson distributions, and we end this section with a brief introduction to Boltzmann distributions.

For a random variable X of density f(x), the tilt of f(x) by a nonnegative integrable function g(x) is the density $\frac{g(x)}{\mathbb{E}(g(X))} \cdot f(x)$. An important class of tilted densities are the polynomially tilted densities, where one tilts by a polynomial $g(x) = x^c$ (with c being any real value such that $\mathbb{E}(X^c)$ is well defined). We then use the notation

$$\operatorname{tilt}_c(f(x)) = \frac{x^c}{\mathbb{E}(X^c)} \cdot f(x).$$

Such tilted densities occur in many places: in the degree distribution in preferential attachment trees [68, 69], in Lamberti-type laws [67], in triangular urn schemes [72, 74], in node-degrees in plane-oriented recursive trees [83], and in table sizes in the Chinese restaurant process [3, 90, 118, 119]. Note that many classes of distributions like the beta distribution, generalized gamma distribution [12], the F-distribution, the beta-prime distribution, and distributions with gamma-type moments [74] are closed under the tilting operation.

The following lemma shows that the operator tilt, admits in fact several equivalent definitions using the density, the moments, or the Laplace transform (see also [74, Remark 2.11]).

LEMMA 3.4 (Polynomially tilted density functions and moment shifts). Consider a random variable X with moment sequence $(\mu_s)_{s>0}$ and density f(x) with support $[0,\infty)$. Now consider a random variable X_c with $c \in \mathbb{N}$, having a distribution uniquely determined by its moments⁵. Then the following properties are equivalent:

- 1. Tilted density: X_c is a random variable with density $f_c(x) = \frac{x^c}{\mu_c} \cdot f(x)$.
- 2. Shifted moments: X_c is a random variable with moments $\mathbb{E}(X_c^s) = \frac{\mu_{s+c}}{\mu_c}$.
- 3. Differentiated moment generating function: X_c is such that $\mathbb{E}(e^{tX_c}) = \frac{1}{u_c} \frac{d^c}{dt^c} \mathbb{E}(e^{tX})$.

REMARK 3.5 (Tilt with $c \in \mathbb{R}$). For the properties (1) and (2) of Lemma 3.4, it is possible to extend their equivalence to $c \in \mathbb{R}$, assuming that the corresponding moments exist. More generally, the equivalence between properties (1) and (2) stays valid for any random variable with density f(x) such that only moments μ_1, \ldots, μ_n exist up to a certain value $n \ge 1$.

REMARK 3.6 (The tilt operator for densities/moments/random variables). This lemma justifies a slight abuse of notation: Starting with the densities of X and X_c linked by $\operatorname{tilt}_c(f(x)) = f_c(x)$, the operator tilt_c is also used to denote the corresponding tilted random variable tilt_c(X) := \hat{X}_c and the corresponding tilted moments tilt_c($\hat{\mu}_s$) := $\frac{\hat{\mu}_{s+c}}{\mu_s}$.

PROOF OF LEMMA 3.4. For (1) \Rightarrow (2), first observe that $f_c(x)$ is indeed a density: One has $f_c(x) \ge 0$ on $[0,\infty)$ and $\int_0^\infty f_c(x) \, dx = \frac{\mu_c}{\mu_c} = 1$. Then, one checks that

$$\mathbb{E}(X_c^s) = \int_0^\infty x^s f_c(x) \, dx = \int_0^\infty x^{s+c} \frac{f(x)}{\mu_c} \, dx = \frac{\mu_{s+c}}{\mu_c}.$$

The fact that X_c is uniquely determined by its moments then implies (2) \Rightarrow (1). For (2) \Leftrightarrow (3), observe that $\frac{d^c}{dt^c}\mathbb{E}(e^{tX}) = \frac{d^c}{dt^c}\sum_{s\geq 0}\frac{\mu_st^s}{s!} = \sum_{s\geq c}\frac{\mu_st^{s-c}}{(s-c)!} = \sum_{s\geq 0}\frac{\mu_{s+c}t^s}{s!}$, and, on the other hand, (1) and (2) give

$$\mathbb{E}(e^{tX_c}) = \int_0^\infty f_c(x)e^{tx} \, dx = \frac{1}{\mu_c} \sum_{s>0} \frac{t^s}{s!} \int_0^\infty x^{s+c} f(x) \, dx = \frac{1}{\mu_c} \sum_{s>0} \frac{t^s}{s!} \mu_{s+c}. \qquad \Box$$

⁵See e.g. Feller [34, Chapter VII.3] for conditions implying that the moments define uniquely the distribution.

Let us now introduce *positive stable laws* (also called *one-sided stable laws*, as their density has support $(0, +\infty)$).

DEFINITION 3.7 (Positive stable laws and their negative powers). We say that a positive random variable S_{α} follows a stable law of parameter $\alpha \in (0,1)$ if its Laplace transform is $\mathbb{E}(e^{-tS_{\alpha}}) = e^{-t^{\alpha}}$ (see [130] for a general presentation involving skewness, scale, and location parameters; they are respectively always 0, 1, and 0 in our work). The density of S_{α} is 0

(11)
$$f_{S_{\alpha}}(x) = \sum_{n=1}^{\infty} \frac{(-1)^n}{n!\Gamma(-n\alpha)} x^{-n\alpha-1}$$

$$= \frac{1}{\pi} \sum_{n=1}^{\infty} (-1)^{n+1} \frac{\Gamma(n\alpha+1)\sin(\pi n\alpha)}{n!} x^{-n\alpha-1}$$

$$= \frac{1}{\pi} \frac{\alpha}{1-\alpha} \int_0^{\pi} \frac{K(\phi)}{x^{1/(1-\alpha)}} \exp\left(-\frac{K(\phi)}{x^{\alpha/(1-\alpha)}}\right) d\phi,$$
(12)

where

$$K(\phi) = \left(\frac{\sin(\alpha\phi)}{\sin(\phi)}\right)^{1/(1-\alpha)} \frac{\sin((1-\alpha)\phi)}{\sin(\alpha\phi)}.$$

Formula (11) was first obtained by Humbert [61], and then rigorously proven by Pollard [121] (see also Feller [35, Chapter XVII.6, Lemma 1], with the parameter $\gamma = -\alpha$ therein). Formula (12) is due, up to a typo that we corrected here, to Ibragimov and Chernin [66].

Now, let r > 0 be some real number. Since $\mathbb{P}\{S_{\alpha}^{-r} \leq x\} = 1 - \mathbb{P}\{S_{\alpha} < x^{-1/r}\}$, we directly obtain from (11) the density of S_{α}^{-r} on its support $(0, +\infty)$:

(13)
$$f_{S_{\alpha}^{-r}}(x) = \frac{1}{r} \sum_{n=1}^{\infty} \frac{(-1)^n}{n! \Gamma(-n\alpha)} x^{n\alpha/r-1} = \frac{x^{-1/r-1}}{r} f_{S_{\alpha}}(x^{-1/r}).$$

Its moments are given by (see, e.g., Janson's survey on moments of Gamma type [74]):

$$\mathbb{E}(S_{\alpha}^{-rs}) = \frac{\Gamma(\frac{sr}{\alpha} + 1)}{\Gamma(sr + 1)}, \quad s > -\frac{\alpha}{r}.$$

We will encounter composition schemes leading to powers of stable laws, an important subcase of it being the Mittag-Leffler distribution.⁷

DEFINITION 3.8 (Mittag-Leffler distribution). We say that a random variable M_{α} follows a Mittag-Leffler distribution $\mathrm{ML}(\alpha)$ if $M_{\alpha} \stackrel{d}{=} S_{\alpha}^{-\alpha}$. Its moment generating function $\mathbb{E}(e^{tM_{\alpha}})$ is the Mittag-Leffler function $E_{\alpha}(t) = \sum_{k \geq 0} \frac{t^k}{\Gamma(1+\alpha k)}$. An important special case is $M_{\frac{1}{2}}$, the half-normal distribution $|\mathcal{N}(0,\sigma^2)|$ with $\sigma = \sqrt{2}$; see Example 3.17 hereafter.

⁶Throughout this article, we use that, by analytical continuation, $1/\Gamma(m) = 0$ whenever m is an integer ≤ 0 .

 $^{^{7}}$ In the literature, there are unfortunately two distinct distributions which are called *Mittag-Leffler distribution*. Both of them are defined in terms of the function $E_{\alpha}(x)$ introduced in 1903 by Mittag-Leffler [102, 103]. The first distribution (which we use in this article) was popularized by Feller [34] (with a slight change of variable) and by Darling and Kac [25] for the study of the local time of Markov processes. It has an exponentially bounded tail. The second one, which has a heavy tail, was introduced by Pillai [117], and should rather be called the Pillai–Mittag-Leffler distribution.

More generally, we will encounter computations leading to the following moments of random variables of shape $X_c = \operatorname{tilt}_c(S_{\alpha}^{-r})$ (which are well defined for $c > -\frac{\alpha}{r}$):

$$\mathbb{E}(X_c^s) = \frac{\Gamma(\frac{(s+c)r}{\alpha}+1)}{\Gamma((s+c)r+1)} \frac{\Gamma(rc+1)}{\Gamma(\frac{cr}{\alpha}+1)} = \frac{\Gamma(\frac{(s+c)r}{\alpha})}{\Gamma((s+c)r)} \frac{\Gamma(rc)}{\Gamma(\frac{cr}{\alpha})}.$$

Note that, by (13) and Lemma 3.4, the density of X_c is given by

$$f_{X_c}(x) = \frac{\Gamma(rc+1)}{r\Gamma(\frac{cr}{\alpha}+1)} \sum_{n=1}^{\infty} \frac{(-1)^n}{n!\Gamma(-n\alpha)} x^{n\alpha/r+c-1}.$$

E.g., $\mathrm{tilt}_1(M_{\frac{1}{2}})$ is the *Rayleigh distribution* of parameter $\sqrt{2}$; see Example 3.18 hereafter.

Another fundamental instance of such a tilted power of stable distribution is a two-parameter generalization of the Mittag-Leffler distribution, which was considered in the literature in link with different probabilistic processes, for example by Pitman [119], James [69], Goldschmidt, Haas, and Sénizergues [52, 53]. We will establish in the next section that the analytic phenomenon hiding behind this distribution is a critical composition scheme (and we will explain in subsequent sections how it is related to these probabilistic processes). For now, let us give a formal definition of this distribution.

DEFINITION 3.9 (Two-parameter Mittag-Leffler distribution). For $\alpha \in (0,1)$ and $\beta > -\alpha$, we say that a random variable X follows a two-parameter Mittag-Leffler distribution $\mathrm{ML}(\alpha,\beta)$ if $X \stackrel{d}{=} (\mathrm{tilt}_{-\beta}(S_\alpha))^{-\alpha}$. Note that one has $\mathrm{ML}(\alpha,0) = \mathrm{ML}(\alpha)$ and that $\mathrm{ML}(\alpha,\beta)$ is uniquely defined by its moments

(15)
$$\mathbb{E}(X^s) = \frac{\Gamma\left(s + \frac{\beta}{\alpha} + 1\right)\Gamma(\beta + 1)}{\Gamma(\alpha s + \beta + 1)\Gamma\left(\frac{\beta}{\alpha} + 1\right)} = \frac{\Gamma\left(s + \frac{\beta}{\alpha}\right)\Gamma(\beta)}{\Gamma(\alpha s + \beta)\Gamma\left(\frac{\beta}{\alpha}\right)}.$$

Comparing these moments with (14) we directly get that for $\beta = \alpha$ and $c = \beta/\alpha$ we have

(16)
$$\operatorname{tilt}_{\beta/\alpha}(\operatorname{ML}(\alpha)) \stackrel{d}{=} \operatorname{ML}(\alpha, \beta), \quad \text{i.e.,} \quad \operatorname{tilt}_{\beta/\alpha}(S_{\alpha}^{-\alpha}) = (\operatorname{tilt}_{-\beta}(S_{\alpha}))^{-\alpha}.$$

In other words, the permutation of the tilt and the power creates a change of the tilt parameter.

Next, we discuss product distributions. First, we recall properties of the beta distribution.

DEFINITION 3.10 (Beta distribution). A *beta-distributed* random variable $B \stackrel{d}{=} \text{Beta}(a,b)$ with parameters a,b>0 has a probability density function defined on (0,1) by

$$f(x) = \frac{\Gamma(a+b)}{\Gamma(a)\Gamma(b)} x^{a-1} (1-x)^{b-1}.$$

The moments of B are given by

(17)
$$\mathbb{E}(B^s) = \frac{\Gamma(s+a)\Gamma(a+b)}{\Gamma(s+a+b)\Gamma(a)}, \quad s > 0,$$

and the beta distribution is uniquely determined by the sequence of its moments. Furthermore, let the reader be convinced of the convenient convention $\operatorname{Beta}(a,0) \stackrel{d}{=} 1$.

We now have all the ingredients to present the main properties of the distribution which will play a key rôle in the next sections, namely, the three-parameter Mittag-Leffler distribution.

DEFINITION 3.11 (Three-parameter Mittag-Leffler distribution). We define the *three-parameter Mittag-Leffler distribution* $\mathrm{ML}(\alpha,\beta,\gamma)$ as the distribution of the product of independent random variables

$$Z \stackrel{d}{=} Y \cdot B^{\alpha}$$

where $Y \stackrel{d}{=} {
m tilt}_{\beta/\alpha}(S_{\alpha}^{-\alpha}) \stackrel{d}{=} {
m ML}(\alpha,\beta)$ and $B \stackrel{d}{=} {
m Beta}(\beta,\gamma)$ are respectively distributed like a Mittag-Leffler and a beta distribution, with $0 < \alpha < 1, \, \beta > 0$, and $\gamma \geq 0$.

LEMMA 3.12. The three-parameter Mittag-Leffler distribution $\mathrm{ML}(\alpha,\beta,\gamma)$ is uniquely determined by its moments

(18)
$$\mathbb{E}(Z^{s}) = \frac{\Gamma\left(s + \frac{\beta}{\alpha}\right)\Gamma\left(\beta + \gamma\right)}{\Gamma\left(\alpha s + \beta + \gamma\right)\Gamma\left(\frac{\beta}{\alpha}\right)}, \quad s > 0.$$

What is more, one has the following identity

(19)
$$\mathrm{ML}(\alpha, \beta, \gamma) \stackrel{d}{=} \mathrm{ML}(\alpha, \beta) \operatorname{Beta}(\beta, \gamma)^{\alpha} \stackrel{d}{=} \mathrm{ML}(\alpha, \beta + \gamma) \operatorname{Beta}(\frac{\beta}{\alpha}, \frac{\gamma}{\alpha}).$$

In particular, one has $ML(\alpha, \beta, 0) = ML(\alpha, \beta)$. Furthermore, one also has the simplification

$$ML(\alpha, \alpha, 1 - \alpha) = ML(\alpha),$$

and, more generally, for any real $\kappa \geq 1$, $\mathrm{ML}(\alpha, \alpha\kappa, 1 - \alpha) = \mathrm{ML}(\alpha, \alpha(\kappa - 1))$.

PROOF. The moments characterize a unique distribution via Carleman's criterion [24, pp. 189–220]. Now, due to the independence of the random variables we have $\mathbb{E}(Z^s) = \mathbb{E}(Y^s)\mathbb{E}(B^{\alpha s})$. Then, using (14) and (17), one gets (18). With the relation (16), this gives (19). The last simplifications of the lemma follow from the identity (15).

Note that $ML(\alpha, \beta, \gamma)$ is one important instance of a "distribution with moments of Gamma type", a class of distributions popularized by Janson in his nice thorough survey [74], where he obtained the following asymptotics of the tail of their density f(x).

$$(20) \qquad f(x) \sim C x^{d-1} \exp(-c x^{\frac{1}{1-\alpha}}) \qquad \text{for } x \sim +\infty,$$
 with $c = (1-\alpha)\alpha^{\frac{\alpha}{1-\alpha}}, \quad d = \frac{\beta/\alpha - \beta - \gamma + 1/2}{1-\alpha}, \quad C = \frac{\Gamma(\beta+\gamma)}{\Gamma(\beta/\alpha)} \frac{\alpha^{\frac{1-2\gamma}{2(1-\alpha)}}}{\sqrt{2\pi(1-\alpha)}}.$

The behaviour at 0 of the density of $ML(\alpha, \beta, \gamma)$ follows from Eq. (22) in the next section and is summarized in Table 3.




FIG 3. Several instances of Mittag-Leffler distributions. All of them have an exponential tail while the behaviour at 0 is more diverse: One has $\mathrm{ML}(\alpha) \sim \frac{1}{\alpha|\Gamma(-\alpha)|}$, $\mathrm{ML}(\alpha,\beta) \sim Cst \cdot x^{\beta/\alpha}$, and $\mathrm{ML}(\alpha,\beta,\gamma) \sim Cst \cdot x^{\beta/\alpha-1}$.

REMARK 3.13 (A link with electromagnetism). The moment generating function of $Z = \mathrm{ML}(\alpha, \beta, \gamma)$ can be expressed via the function

$$E_{\alpha,\beta'}^{\gamma'}(t) := \sum_{k=0}^{\infty} \frac{\Gamma(k+\gamma')}{\Gamma(\alpha k+\beta')\Gamma(\gamma')} \frac{t^k}{k!};$$

one has indeed

$$\mathbb{E}(e^{tZ}) = \Gamma(\beta') E_{\alpha,\beta'}^{\gamma'}(t) \qquad \text{(with } \beta' = \gamma + \beta \text{ and } \gamma' = \beta/\alpha\text{)}.$$

Our article thus provides a unified probabilistic interpretation of the special function $E_{\alpha,\beta'}^{\gamma'}(t)$ which also appears in physics, where it is sometimes called the *Prabhakar function*, or the *three-parameter Mittag-Leffler function* [54, 122]. This special function is also the inverse Laplace transform of the right-hand side of the Havriliak–Negami generalization of the Debye and Cole–Cole equations, which are classical models of dielectric relaxation in electromagnetism [23,49,55]. What is more, it is proven by Górska et al. in [56] that $g(t) := E_{\alpha,\beta'}^{\gamma'}(-t)$ (with $0 < \alpha \le 1$, $\gamma' > 0$, and $\beta' > \alpha \gamma'$, conditions which match exactly our probabilistic setting!) is *totally monotone*: $(-1)^n \partial_t^n g(t) > 0$ for $t \in \mathbb{R}^+$. Bernstein's theorem on totally monotone functions then implies that this function is associated to a density. Our article unravels this mysterious density, and explains its universality.

REMARK 3.14 (Other appearances of $\mathrm{ML}(\alpha,\beta,\gamma)$). Since the initial publication of our work on arXiv, the three-parameter Mittag-Leffler distribution also appeared in two other articles. In the first one, Möhle [104], independently of our results, established that it is the limit law appearing in a generalization of the Chinese restaurant process (see our Section 6.6). In the second one, Sibisi [125], with a motivation coming from special functions, considered a four-parameter generalization $\mathrm{ML}_4(a,b,c,d)$. Comparing moments, we then see that

$$\mathrm{ML}(\alpha,\beta,\gamma) = \mathrm{ML}_4(\alpha,\gamma,0,\beta) = \mathrm{ML}_4(\alpha,\beta+\gamma,\frac{\beta}{\alpha},0).$$

Another key ingredient that we shall need is the mixed Poisson distributions. These distributions were first introduced by Dubourdieu in 1938 for actuarial mathematics/insurance modelling [32], and then also studied by Lundberg and others (sometimes under the name "compound Poisson processes", a term that has a different meaning nowadays); they were also used for applications in bacteriology by Neyman [107], in combinatorics by Kuba and Panholzer [90], for expectation-maximization algorithms by Karlis [78], or for the analysis of some point processes by Grandell [59]. Their unimodality properties are studied by Masse and Theodorescu [99], and their tail asymptotics are analysed by Wilmot and Lin in [133, 134].

DEFINITION 3.15 (Mixed Poisson distributions). Let X denote a nonnegative random variable with cumulative distribution function U. We say that the discrete random variable Y has a mixed Poisson distribution with mixing distribution U and scale parameter $\xi \geq 0$, if its probability mass function is given for $\ell \geq 0$ by

$$\mathbb{P}\{Y=\ell\} = \frac{\xi^\ell}{\ell!} \int_{\mathbb{R}^+} X^\ell e^{-\xi X} dU = \frac{\xi^\ell}{\ell!} \mathbb{E}(X^\ell e^{-\xi X}).$$

This is summarized by the notation $Y \stackrel{d}{=} MPo(\xi U)$, or, indifferently, $Y \stackrel{d}{=} MPo(\xi X)$.

Note that mixed Poisson distributions provide a common generalization of three major discrete laws ubiquitous in combinatorics (see [42, Figure IX.5]), namely the Poisson, the geometric, and the negative binomial distributions, making them of great importance per se.

We emphasize that the *factorial moments*⁸ of a mixed Poisson distribution are closely related to the classical *raw moments* of its mixing distribution: $\mathbb{E}(Y^{\underline{s}}) = \xi^s \mathbb{E}(X^s)$, $s \ge 1$. Additionally, like for any distribution, the factorial and raw moments of Y are related via the Stirling set partition numbers $\binom{s}{k}$ (also called Stirling numbers of the second kind):

$$\mathbb{E}(Y^s) = \sum_{k=0}^r \binom{s}{k} \mathbb{E}(Y^{\underline{k}}).$$

Such relations are called Stirling transforms [19]. We refer to [90] for more properties of the Stirling transform and mixed Poisson distributions, like the following useful expression for the probability mass function of MPo(ξX) in terms of its *factorial* moments.

PROPOSITION 3.16. Let X denote a random variable with moment sequence given by $(\mu_s)_{s\in\mathbb{N}}$. If a random variable Y has factorial moments given by $\mathbb{E}(Y^{\underline{s}}) = \xi^s \mu_s$, then $Y \stackrel{d}{=} MPo(\xi X)$. What is more, the sequence of moments of Y is the Stirling transform of the moment sequence $(\mu_s)_{s\in\mathbb{N}}$, and the probability mass function of Y is given by

$$\mathbb{P}\{Y=\ell\} = \sum_{s>\ell} (-1)^{s-\ell} \binom{s}{\ell} \mu_s \frac{\xi^s}{s!}, \quad \ell \ge 0.$$

Let us give two short examples of mixed Poisson distributions, which, as we shall later see, correspond to ubiquitous cases in combinatorics.

EXAMPLE 3.17 (Mixed Poisson half-normal distribution). A half-normally distributed random variable $X \stackrel{d}{=} HN(\sigma)$ with parameter σ is the absolute value of a normally distributed random variable, i.e. $X \stackrel{d}{=} |\mathcal{N}(0, \sigma^2)|$. Consequently, X has the probability density function

$$f(x;\sigma) = \frac{\sqrt{2}}{\sigma\sqrt{\pi}}e^{-\frac{x^2}{2\sigma^2}}, \quad x \ge 0;$$

alternatively, it is fully characterized by its moment sequence $\mathbb{E}(X^s) = \sigma^s 2^{s/2} \Gamma(\frac{s+1}{2}) / \Gamma(\frac{1}{2})$. Thus, a discrete random variable Y with probability mass function

$$\mathbb{P}\{Y=\ell\} = \frac{\xi^{\ell}}{\ell!} \cdot \frac{\sqrt{2}}{\sigma\sqrt{\pi}} \int_0^\infty x^{\ell} e^{-\xi x - \frac{x^2}{2\sigma^2}} dx, \quad \ell \ge 0,$$

has a mixed Poisson distribution: $Y \stackrel{d}{=} \mathrm{MPo}(\xi X)$ with $X \stackrel{d}{=} \mathrm{HN}(\sigma)$. Note that we can readily expand the exponential function and obtain various series representations of $\mathbb{P}\{Y=\ell\}$.

EXAMPLE 3.18 (Mixed Poisson Rayleigh distribution). A Rayleigh distributed random variable $X \stackrel{d}{=} \text{Rayleigh}(\sigma)$ with parameter σ has the probability density function

$$f(x;\sigma) = \frac{x}{\sigma^2} e^{-\frac{x^2}{2\sigma^2}}, \quad x \ge 0;$$

alternatively, it is fully characterized by its moment sequence $\mathbb{E}(X^s) = \sigma^s \, 2^{s/2} \, \Gamma\left(\frac{s}{2} + 1\right)$. Thus, a discrete random variable Y with probability mass function

$$\mathbb{P}\{Y = \ell\} = \frac{\xi^{\ell}}{\ell!\sigma^2} \int_0^\infty x^{\ell+1} e^{-\xi x - \frac{x^2}{2\sigma^2}} dx, \quad \ell \ge 0,$$

has a mixed Poisson distribution: $Y \stackrel{d}{=} MPo(\xi X)$ with $X \stackrel{d}{=} Rayleigh(\sigma)$.

⁸Throughout this work we denote by $x^{\underline{n}}$ the n^{th} falling factorial, $x^{\underline{n}} = x(x-1)\cdots(x-n+1)$, $n \geq 0$, with $x^{\underline{0}} = 1$. It will be used for $\mathbb{E}(X^{\underline{n}})$, the factorial moment of order n of a random variable X.

Last but not least, in our results, we shall also encounter another important family of discrete distributions: the *Boltzmann distributions*. They were introduced in combinatorics by Duchon, Flajolet, Louchard, and Schaeffer in order to perform sampling of combinatorial structures [33]. The starting point of these authors was the idea to give, like in statistical mechanics, a Gibbs measure/Boltzmann weight x^n (for some fixed real number x) to each combinatorial object of size n. Note that the objects of same size then follow a uniform distribution. It was then a nice surprise that, if one deals with an assemblage of combinatorial objects, the corresponding Boltzmann weights are given by very simple probabilistic laws (similarly to the symbolic method [42] which directly gives the generating functions of unions/products/cycles of objects). This led to an outstanding generic linear time sampling algorithm: Its astonishing efficiency is partially due to the fact that, thanks to these Boltzmann weights, the sampling of a product of two combinatorial structures is simply obtained by two independent recursive subsamplings. The sampling algorithm is thus essentially based on the following definition:

DEFINITION 3.19 (Boltzmann distribution). For any generating function $G(z) = \sum_{n\geq 0} g_n z^n$, and for any parameter x>0 smaller than the radius of convergence of G, a random variable X follows a *Boltzmann distribution (associated with G) of parameter x*, denoted by $\mathcal{B}_G(x)$, if

$$\mathbb{P}\{X=n\} = \frac{g_n x^n}{G(x)}, \quad n \ge 0.$$

Then, the key idea behind Boltzmann sampling (of objects of size n) is to choose x adequately to maximize $\mathbb{P}\{X=n\}$. If the object generated is not of size n, one rejects it and restarts the sampling. This leads to a uniform sampling algorithm of optimal efficiency when x is the unique real root of the equation xG'(x)=nG(x). This equation is reminiscent of many probabilistic results with mean $\mu=xG'(x)/G(x)$ (e.g., when G encodes the offspring of a Galton–Watson process). This is no coincidence: By design, Boltzmann sampling "reverse-engineers" these results (see e.g. [10], or the recent works of Sportiello [127] or Panagiotou, Ramzews, and Stufler [108, 128] for more on these aspects).

As we shall see, these Boltzmann distributions also occur in our *critical* composition schemes. Retrospectively, it explains and puts in a unified framework earlier sporadic occurrences of such distributions for the limit law of the degree of a random node in simply-generated trees, the root degree in simply-generated trees, as well as in *subcritical* composition schemes; see [42, pages 460, 629–633].




FIG 4. Boltzmann distributions $\mathcal{B}_G(x)$ have different shapes 10 , depending on the singular exponent λ_G of the function G: It goes from an asymptotic Gaussian shape if $\lambda_G \ll -1$ (and for entire functions) to a spread shape if $\lambda_G \gg -1$. The above distributions are drawn with a value of the parameter x such that $\mathbb{E}(\mathcal{B}_G(x)) = xG'(x)/G(x) = n$ (with n = 200). It is interesting that they appear in the context of critical schemes, independently of uniform random generation motivations, but, as explained above, both aspects are in fact intimately related.

4. Extended composition scheme. In the following we state and prove our main theorem on the extended critical composition scheme (2) for pure schemes. For the terms critical and pure, we refer to Definitions 1.1 and 3.3, respectively. This theorem shows the universality of the three-parameter Mittag-Leffler distribution (introduced in Definition 3.11).

THEOREM 4.1 (Extended composition scheme: pure case). In a pure extended critical composition scheme F(z,u) = G(uH(z))M(z), the core size X_n , rescaled, converges in distribution and in moments¹¹ to a random variable X distributed like the three-parameter Mittag-Leffler distribution:

(21)
$$\frac{X_n}{\kappa \cdot n^{\lambda_H}} \xrightarrow{d} X, \quad \text{with} \quad X \stackrel{d}{=} \mathrm{ML}(\alpha, \beta, \gamma),$$

where

$$\alpha = \lambda_H, \quad \beta = -\lambda_G \lambda_H, \quad \gamma = -\lambda_M^- = -\min(0, \lambda_M), \quad and \quad \kappa = \frac{\tau_H}{-c_H}.$$

What is more, one has a local limit theorem

$$\mathbb{P}\{X_n = x \cdot \kappa n^{\lambda_H}\} \sim \frac{1}{\kappa n^{\lambda_H}} \cdot f_X(x),$$

with uniform convergence over any interval in $(0, +\infty)$, and where $f_X(x)$ is the density of X:

(22)
$$f_X(x) = \frac{\Gamma(\beta + \gamma)}{\Gamma(\beta/\alpha)} \sum_{j \ge 0} \frac{(-1)^j}{j! \Gamma(\gamma - j\alpha)} x^{\beta/\alpha + j - 1}.$$

REMARK 4.2 (Two simplifications of the three-parameter Mittag-Leffler distribution). In the above theorem, if $\lambda_M \geq 0$ (which includes the critical scheme F(z,u) = G(uH(z)) as $\lambda_M = +\infty$), then $\gamma = 0$ and the three-parameter Mittag-Leffler distribution (21) simplifies into a two-parameter Mittag-Leffler distribution:

$$X \stackrel{d}{=} \operatorname{tilt}_{-\lambda_G}(\operatorname{ML}(\lambda_H)) \stackrel{d}{=} \operatorname{ML}(\lambda_H, -\lambda_G \lambda_H).$$

In particular, for $\lambda_H = \frac{1}{2}$ and $\lambda_G = -1$ the random variable X follows a Rayleigh distribution of parameter $\sigma = \sqrt{2}$; see Example 3.8.

Another noteworthy simplification occurs for $\lambda_M < 0$, in the special case $\lambda_G = -1$ and $\lambda_H - \lambda_M = 1$: We then obtain a Mittag-Leffler distribution of parameter λ_H :

$$X \stackrel{d}{=} \mathrm{ML}(\lambda_H).$$

In particular, for $\lambda_H = \frac{1}{2}$ the random variable X follows a half-normal distribution of parameter $\sigma = \sqrt{2}$; see again Example 3.8.

Note that the cases with $\lambda_G = -1$ occur in a great many places in applied probability theory; they indeed correspond to a natural combinatorial framework where \mathcal{F} -objects are essentially *sequences* of \mathcal{H} -components (see [42] and examples in our Section 6).

¹⁰See https://lipn.fr/~cb/Papers/CriticalSchemes/ for several animations of the different limit laws occurring in this article

 $^{^{11}}$ We write $X_n \xrightarrow{d} X$ to denote that X_n converges in distribution to X, with convergence of all moments, i.e., $\mathbb{E}(X_n^s) \to \mathbb{E}(X^s)$ for all s. This notion of convergence in moments was, e.g., used in [48,77]. It is also indirectly used in [124], which deals with more constrained models that offer a convergence in L^p , for all p > 1. Note that in all our results involving convergence in distribution or in moments, we omit the speed of convergence, which is in fact easily obtained by considering the Puiseux expansions of order 2 of G/H/M.

PROOF OF THEOREM 4.1. The factorial moments satisfy 12:

$$\mathbb{E}(X_n^{\underline{s}}) = \frac{[z^n] \partial_u^s(F)(z,1)}{[z^n] F(z,1)} = \frac{[z^n] H(z)^s G^{(s)} \big(H(z)\big) M(z)}{[z^n] G(H(z)) M(z)}.$$

In the following we use the notation ρ_F , τ_F , λ_F , and c_F from Section 3.1 for the singular expansions of F = G/H/M. As we are in a pure critical scheme (Definition 3.3), the unique singularity of F(z) = G(H(z))M(z) is at $z = \rho_H$. Then, we unify the three cases (10a), (10b), and (10c) by using $\lambda_M^- = \min(0, \lambda_M)$ and choosing C_M according to the specific case $(C_M$ is for $\lambda_M \neq \lambda_G \lambda_H$ either τ_M or c_M ; this quantity C_M will anyway cancel in the end). Note that the case (10d) does not hold in a pure scheme. This gives

$$F(z) \sim C_M c_G \left(\frac{-c_H}{\rho_G}\right)^{\lambda_G} \left(1 - \frac{z}{\rho_H}\right)^{\lambda_G \lambda_H + \lambda_M^-}$$
.

Therefore, using the transfer theorem from singularity analysis we get

(23)
$$f_n = [z^n] F(z, 1) \sim \frac{C_M c_G (-c_H/\rho_G)^{\lambda_G}}{\rho_H^n} \cdot \frac{n^{-\lambda_G \lambda_H - \lambda_M^- - 1}}{\Gamma(-\lambda_G \lambda_H - \lambda_M^-)}.$$

Using singular differentiation (see [42, Theorem VI.8] or [37]) for G(z), we get the following singular expansion of the higher-order derivatives $G^{(s)}(z)$, for integer $s \ge 1$:

$$G^{(s)}(z) \sim (-1)^s \frac{c_G}{\rho_G^s} \lambda_{\overline{G}}^{\underline{s}} \left(1 - \frac{z}{\rho_G}\right)^{\lambda_G - s}.$$

Next, from the singular expansion of H(z) we directly get

$$H(z)^s \sim \tau_H^s - s\tau_H^{s-1} \cdot (-c_H) \left(1 - \frac{z}{\rho_H}\right)^{\lambda_H}.$$

Therefore, we get (recall that our scheme is critical, i.e., $H(\rho_H) = \tau_H = \rho_G$)

$$G^{(s)}(H(z)) \sim (-1)^s c_G \rho_G^{-\lambda_G} \lambda_G^{\underline{s}} (-c_H)^{\lambda_G - s} \left(1 - \frac{z}{\rho_H}\right)^{\lambda_G \lambda_H - s \lambda_H}.$$

Combining these expansions with the one of M(z) gives the required expansion

$$H(z)^s G^{(s)}(H(z)) M(z) \sim (-1)^s \tau_H^s C_M c_G \rho_G^{-\lambda_G} \lambda_G^{\underline{s}} (-c_H)^{\lambda_G - s} \left(1 - \frac{z}{\rho_H}\right)^{\lambda_M^- + \lambda_H \lambda_G - s\lambda_H}.$$

Next we rewrite $(-1)^s \lambda_G^s$ using the gamma function, that is, we use $(-1)^s \lambda_G^s = \frac{\Gamma(s - \lambda_G)}{\Gamma(-\lambda_G)}$. Hence, we obtain by extraction of coefficients and singularity analysis

(24)
$$[z^{n}]H(z)^{s}G^{(s)}(H(z))M(z) \sim \left(\frac{\tau_{H}}{-c_{H}}\right)^{s} \frac{C_{M}c_{G}(-c_{H}/\rho_{G})^{\lambda_{G}}}{\rho_{H}^{n}} \times \frac{\Gamma(s-\lambda_{G})}{\Gamma(-\lambda_{G})} \cdot \frac{n^{-\lambda_{G}\lambda_{H}-\lambda_{M}^{-}-1+s\lambda_{H}}}{\Gamma(s\lambda_{H}-\lambda_{G}\lambda_{H}-\lambda_{M}^{-})}.$$

Combining this expression with (23) gives $\mathbb{E}(X_n^{\underline{s}}) \sim n^{s\lambda_H} \kappa^s \cdot \mu_s$, where $\kappa := \frac{\tau_H}{-c_H}$ and $\Gamma(s - \lambda_C)\Gamma(-\lambda_C\lambda_H - \lambda_M)$

$$\mu_s := \frac{\Gamma(s-\lambda_G)\Gamma(-\lambda_G\lambda_H - \lambda_M^-)}{\Gamma(-\lambda_G)\Gamma(s\lambda_H - \lambda_G\lambda_H - \lambda_M^-)}$$

¹²Throughout this work, we denote by ∂_u the differentiation operator with respect to the variable u. Accordingly, we use the shorthand notation $\partial_u(F)(z,1) = (\partial_u F(z,u))\big|_{u=1}$.

What is more, one has $\mathbb{E}(X_n^s) \sim \mathbb{E}(X_n^s)$ since we can express the raw moments by using the factorial moments and the Stirling numbers of the second kind:

(25)
$$\mathbb{E}(X_n^s) = \mathbb{E}\left(\sum_{j=0}^s {s \brace j} X_n^{\underline{j}}\right) = \sum_{j=0}^s {s \brace j} \mathbb{E}(X_n^{\underline{j}}).$$

Consequently, we obtain the moment convergence to the stated moment sequence:

$$\frac{\mathbb{E}(X_n^s)}{n^{s\lambda_H}\kappa^s} \to \mu_s.$$

By Stirling's formula for the gamma function, one has $\Gamma(z) = \left(\frac{z}{e}\right)^z \frac{\sqrt{2\pi}}{\sqrt{z}} \left(1 + \mathcal{O}\left(\frac{1}{z}\right)\right)$; this entails that for $s \to \infty$ the moments satisfy

$$(\mu_s)^{-1/(2s)} \sim \sqrt{e^{1-\lambda_H} \lambda_H^{\lambda_H}} s^{\frac{H-1}{2}} \left(1 + \mathcal{O}\left(\frac{1}{s}\right)\right).$$

As $0 < \lambda_H < 1$, the divergence in Carleman's criterion [24, pp. 189–220] is satisfied:

(26)
$$\sum_{s=0}^{\infty} \mu_s^{-1/(2s)} = +\infty;$$

consequently, the moment sequence $(\mu_s)_{s\in\mathbb{N}}$ characterizes a unique distribution. Now, the Fréchet–Shohat theorem (see their original article [45] or the book by Loève [93, Sec. 11.4]) states that if $\mathbb{E}[Y_n^r] \to \mathbb{E}[Y^r]$ for all $r \in \mathbb{N}$, where Y is uniquely characterized by its moments, then Y_n converges to Y in distribution. In our case, this implies the convergence in distribution of the random variable $\frac{X_n}{\kappa \cdot n^{\lambda_H}}$ to a random variable X with moment sequence $(\mu_s)_{s\in\mathbb{N}}$.

Concerning the local limit theorem, we have to analyse $\mathbb{P}\{X_n=k\}=\frac{g_k[z^n]H(z)^kM(z)}{f_n}$, for $k=x\cdot\kappa\cdot n^{\lambda_H}$, with x in a compact subinterval of $(0,\infty)$. First, we note that by (7) applied to G(z) we directly obtain

(27)
$$g_k \sim \frac{c_G}{\rho_G^k} \cdot \frac{(\kappa x)^{-\lambda_G - 1} n^{-\lambda_H \lambda_G - \lambda_H}}{\Gamma(-\lambda_G)}.$$

It remains to determine the asymptotics of $[z^n]H(z)^kM(z)$. We have

$$[z^n]H(z)^kM(z) = \frac{1}{2\pi i} \oint \frac{H(z)^k M(z)}{z^{n+1}} dz.$$

Introducing the point A of coordinates $(\frac{1}{n}, \rho_H(1+\frac{\log^2 n}{n}))$, this Cauchy integral can be transformed into an integral over a larger contour in the Delta-domain (in blue in Figures 1 and 5). Then, setting $z=\rho_H(1+t/n)$ leads to an integral asymptotically concentrated on the Hankel contour $\mathcal C$ (which starts and ends at $+\infty$):

$$[z^n]H(z)^kM(z) \sim \frac{\tau_H^k C_M}{\rho_H^n n^{1+\lambda_M^-}} \cdot \frac{1}{2\pi i} \int_{\mathcal{C}} (-t)^{\lambda_M^-} e^{-t-x(-t)^{\lambda_H}} dt.$$

Then, expanding $e^{-x(-t)^{\lambda_H}}$ leads to

(28)
$$[z^n]H(z)^k M(z) \sim \frac{\tau_H^k C_M}{\rho_H^n n^{1+\lambda_M^-}} \cdot \frac{1}{2\pi i} \int_{\mathcal{C}} e^{-t} \sum_{i>0} \frac{(-x)^j}{j!} (-t)^{j\lambda_H + \lambda_M^-} dt,$$

in which we recognize the following Hankel contour representation [132, Section 12.22]:

$$\frac{1}{\Gamma(-z)} = \frac{1}{2i\pi} \int_{\mathcal{C}} (-t)^z e^{-t} dt.$$

Recall that we use, by analytic continuation, $1/\Gamma(m)=0$ for any integer m<0; this avoids heavier expressions relying on Euler's reflection formula $\frac{1}{\Gamma(-z)}=-\frac{1}{\pi}\sin(\pi z)\Gamma(1+z)$.




FIG 5. The integration contour of the Cauchy integral (left) is transformed into a larger contour (in red in the middle image) inside a Delta-domain (in blue), becoming in the limit the Hankel contour \mathcal{C} (right), allowing the identification of the gamma function.

Now, combining the expansions of f_n from (23), g_k from (27), and term-wise integration in (28), we get the desired local limit theorem:

(29)
$$\mathbb{P}\{X_n = k\} = \frac{g_k}{f_n} [z^n] H(z)^k M(z)$$

$$\sim \frac{\Gamma(-\lambda_G \lambda_H - \lambda_M^-)}{\kappa n^{\lambda_H} \Gamma(-\lambda_G)} \sum_{j>0} \frac{(-1)^j}{j! \Gamma(-j\lambda_H - \lambda_M^-)} x^{j-\lambda_G - 1}.$$

It remains to verify that $f_X(x)$ is the density function of X with moments

$$\mu_s = \frac{\Gamma(s - \lambda_G)\Gamma(-\lambda_G\lambda_H - \lambda_M^-)}{\Gamma(-\lambda_G)\Gamma(s\lambda_H - \lambda_G\lambda_H - \lambda_M^-)}.$$

Note that the gamma function is never zero and that its only singularities are simple poles at the negative integers. Therefore μ_s (considered as a complex function of s) has simple poles at $\rho_0 = \lambda_G$ (if $\lambda_M^- \neq 0$, which entails $\lambda_G < 0$ as we have a pure scheme) and at $\rho_j = \lambda_G - j$ (for $j \in \{1, 2, \dots\}$). Then, as μ_{s-1} is the Mellin transform of the density $f_X(x)$ of X, an inverse Mellin computation implies that $f_X(x)$ is expressible in terms of the residues of μ_s :

$$f_X(x) = \frac{1}{2\pi i} \int_{\sigma-i\infty}^{\sigma+i\infty} \mu_{s-1} x^{-s} ds = \sum_{\substack{\sigma_j \, = \, 1 \, + \, \rho_j \text{ pole of } \mu_{s-1} \\ \sigma_j \, < \, \sigma}} \operatorname{Res}_{s=\sigma_j}(\mu_{s-1}) x^{-\sigma_j},$$

which is valid for x > 0, and where $\sigma = 1 + \rho_0 + \epsilon$ is in the fundamental strip of μ_{s-1} , and

$$\operatorname{Res}_{s=\sigma_j}(\mu_{s-1}) = \operatorname{Res}_{s=\rho_j}(\mu_s) = \frac{(-1)^j}{j!} \cdot \frac{\Gamma(-\lambda_G \lambda_H - \lambda_M^-)}{\Gamma(-\lambda_G)\Gamma(-j\lambda_H - \lambda_M^-)}.$$

Summing for $j \ge 0$ (we can include ρ_0 , even if it is not a pole, as the corresponding residue is then 0) gives the same density function $f_X(x)$ as in the local limit theorem (29).

Now, thanks to our probability distribution melting pot section 3.2, we identify the corresponding limit law by matching the parameters in the moments of the three-parameter Mittag-Leffler product given in Lemma 3.12.

Above, we have established the limit laws occurring for analytically *pure* composition schemes. Next we deal with the analytically *confluent* and *degenerate* cases of Definition 3.3. They lead either to discrete distributions, or, more interestingly, to a mixture of discrete and continuous distributions. This generalizes the phenomenon observed in [7], where for $\lambda_H = \frac{3}{2}$ the limit law also consists of a discrete part *plus* a continuous part, a map-Airy distribution (this phenomenon also occurs for variants of 3-connected graphs; see [51]). The following two theorems explain the connection between this discrete part and Boltzmann distributions (usually used for random sampling; see, e.g., [21,33]).

¹³See [42, Appendix B.7] for more on this method; see also [74, Theorem 5.4] and [74, Equation (6.12)] for similar results on the class of functions with moments of Gamma type.

THEOREM 4.3 (Extended composition scheme: degenerate case). Let a degenerate extended critical composition scheme F(z,u) = G(uH(z))M(z) with $0 < \lambda_H < 1$ be given.

For $0 < \lambda_G < 1$ and $\lambda_M < \lambda_G \lambda_H$, the core size X_n converges for $k \ge 0$ and $n \to \infty$ to a Boltzmann distribution $\mathcal{B}_G(\rho_G)$ (see Definition 3.19):

$$\mathbb{P}\{X_n = k\} \to \mathbb{P}\{\mathcal{B}_G(\rho_G) = k\} = \frac{g_k \rho_G^k}{G(\rho_G)}.$$

For $\lambda_G > 1$, the core size X_n has for $k \ge 0$ and $n \to \infty$ the following behaviour:

• For $\lambda_M < \lambda_H$, the random variable X_n converges to a Boltzmann distribution $\mathcal{B}_G(\rho_G)$:

$$\mathbb{P}\{X_n = k\} \to \frac{g_k \rho_G^k}{G(\rho_G)}.$$

• For $\lambda_M = \lambda_H$, the random variable X_n converges to a convex combination of two discrete independent Boltzmann distributions:

$$Be(p) \cdot \mathcal{B}_G(\rho_G) + (1 - Be(p)) \cdot \mathcal{B}_{G'}(\rho_G),$$

with $p = \frac{c_M G(\rho_G)}{c_M G(\rho_G) + c_H \tau_M G'(\rho_G)}$ and where the Bernoulli distribution Be(p) is independent of $\mathcal{B}_G(\rho_G)$ and $\mathcal{B}_{G'}(\rho_G)$. We have

$$\mathbb{P}\{X_n = k\} \to p \cdot \frac{g_k \rho_G^k}{G(\rho_G)} + (1-p) \cdot \frac{kg_k \rho_G^{k-1}}{G'(\rho_G)}.$$

• For $\lambda_M > \lambda_H$, the random variable X_n converges to a Boltzmann distribution $\mathcal{B}_{G'}(\rho_G)$:

$$\mathbb{P}\{X_n = k\} \to \frac{kg_k \rho_G^{k-1}}{G'(\rho_G)}.$$

PROOF. Let us start with the case $0 < \lambda_G < 1$. We study for arbitrary but fixed $k \in \mathbb{N}$ the probability $\mathbb{P}\{X_n = k\}$, as n tends to infinity. Using (3) and applying the singular expansion of Lemma 3.1 to H(z) and M(z), one gets

(30)
$$\mathbb{P}\{X_{n} = k\} = g_{k} \frac{[z^{n}]H(z)^{k} \cdot M(z)}{f_{n}} \\
\sim \frac{g_{k}}{f_{n}}[z^{n}] \left(\left(\tau_{H}^{k} + k\tau_{H}^{k-1}c_{H} \left(1 - \frac{z}{\rho_{H}}\right)^{\lambda_{H}} \right) \left(\tau_{M} + c_{M} \left(1 - \frac{z}{\rho_{H}}\right)^{\lambda_{M}} \right) \right).$$

As $\lambda_M < \lambda_G \lambda_H < \lambda_H$, the singular exponent λ_M dominates the asymptotics and applying singularity analysis (7) leads to

$$\mathbb{P}\{X_n = k\} \sim \frac{g_k}{f_n} [z^n] \tau_H^k c_M (1 - \frac{z}{\rho_H})^{\lambda_M} \sim \frac{g_k \tau_H^k}{\tau_G} = \frac{g_k \rho_G^k}{G(\rho_G)},$$

where we simplified with the asymptotics $f_n \sim \tau_G c_M \frac{1}{\rho_H^n} \frac{n^{-\lambda_M-1}}{\Gamma(-\lambda_M)}$ coming from the expansion (10c), and where we used $\tau_H = \rho_G$ and $\tau_G = G(\rho_G)$ which follows by our assumptions on criticality.

Let us continue with the case $\lambda_G > 1$. From expansion (10d) we see that we now have to distinguish three subcases: $\lambda_M < \lambda_H$, $\lambda_M = \lambda_H$, or $\lambda_H < \lambda_M$. The case $\lambda_M < \lambda_H$ is exactly the same as before. The case $\lambda_M > \lambda_H$ is different, yet the results are derived analogously: the asymptotics of f_n depend now on λ_H and are given by

$$f_n \sim G'(\rho_G) \tau_M c_H \frac{1}{\rho_H^n} \frac{n^{-\lambda_H - 1}}{\Gamma(-\lambda_H)}.$$

Then, we obtain from (30) where now again the contribution from H(z) dominates

$$\mathbb{P}\{X_n = k\} \sim \frac{kg_k \tau_H^{k-1}}{G'(\rho_G)} = \frac{kg_k \rho_G^{k-1}}{G'(\rho_G)}.$$

Finally, in the case $\lambda_M = \lambda_H$ the previous two contributions are mixed as the first two terms in (10d) contribute: The asymptotics of f_n are given by

$$f_n \sim (G'(\rho_G)\tau_M c_H + c_M \tau_G) \frac{1}{\rho_H^n} \frac{n^{-\lambda_H - 1}}{\Gamma(-\lambda_H)}.$$

Note that the coefficient is not zero, as both terms are negative: $G'(\rho_g), \tau_M, \tau_M > 0$ and $c_H, c_M < 0$ due to the sign property (8) for $0 < \lambda_H < 1$. Then, we obtain from (30) where again both contributions have to be taken into account

$$\mathbb{P}\{X_n = k\} \sim \frac{c_M g_k \rho_G^k + \tau_M c_H k g_k \rho_G^{k-1}}{G'(\rho_G) \tau_M c_H + c_M \tau_G} = p \cdot \frac{g_k \rho_G^k}{G(\rho_G)} + (1 - p) \cdot \frac{k g_k \rho_G^{k-1}}{G'(\rho_G)},$$

where $p = \frac{c_M G(\rho_G)}{c_M G(\rho_G) + c_H \tau_M G'(\rho_G)} \in (0,1)$ by the sign property (8). We thus get a linear combination of two Boltzmann distributions, weighted by a Bernoulli random variable Be(p).

THEOREM 4.4 (Extended composition scheme: confluent case). Let a confluent extended critical composition scheme F(z,u) = G(uH(z))M(z) with $0 < \lambda_H < 1$ be given (i.e., $0 < \lambda_G < 1$ and $\lambda_M = \lambda_G \lambda_H$). Then the core size X_n is a convex combination of a Boltzmann distribution $\mathcal{B}_G(\rho_G)$ and an asymptotically continuous random variable Z_n :

$$X_n \stackrel{d}{=} Be(p) \cdot \mathcal{B}_G(\rho_G) + (1 - Be(p)) \cdot Z_n, \qquad \frac{Z_n}{\kappa \cdot n^{\lambda_H}} \stackrel{d}{\longrightarrow} X,$$

where κ and the limit law $X \stackrel{d}{=} \mathrm{ML}(\lambda_H, -\lambda_G \lambda_H)$ are the same as in Remark 4.2, and where $\mathrm{Be}(p)$ (with $p = \frac{c_M G(\rho_G)}{c_M G(\rho_G) + \tau_M c_G (-c_H/\rho_G)^{\lambda_G}}$) is independent of $\mathcal{B}_G(\rho_G)$, Z_n , and X.

PROOF. The proof follows the same lines as the one of Theorem 4.3. We start from expansion (10c). Due to $\lambda_M = \lambda_G \lambda_H$ both terms contribute, and we get

(31)
$$f_n \sim \frac{\tau_M c_G (-c_H/\rho_G)^{\lambda_G} + \tau_G c_M}{\rho_H^n} \frac{n^{-\lambda_M - 1}}{\Gamma(-\lambda_M)}.$$

Then we extract the asymptotics from (30), where now the contributions from M(z) dominate, as $\lambda_M < \lambda_G$:

(32)
$$\mathbb{P}\{X_n = k\} = \frac{c_M g_k \tau_H^k}{\tau_M c_G(-c_H/\rho_G)^{\lambda_G} + \tau_G c_M} + o(1) = p \cdot \frac{g_k \rho_G^k}{G(\rho_G)} + o(1),$$

Using the sign property (8) we get $p \in (0,1)$. So, for large n, X_n behaves with probability p like $\mathcal{B}_G(\rho_G)$, but what happens with probability 1-p? Where is this missing mass in (32)? For sure, it is sneakily spread in $\sum_k o(1)$: It turns out that more and more mass is distributed in the range $k \sim \Theta(\kappa n^{\lambda_H})$, leading to an asymptotic continuous distribution therein.

In order to identify this distribution, we compute the factorial moments of X_n like in the proof of Theorem 4.1. We again use the singular expansions of $G^{(s)}(H(z))$, $H(z)^s$, and $H(z)^sG^{(s)}(H(z))M(z)$. Formula (24) holds verbatim with $C_M=\tau_M$. The big difference lies now in the asymptotics of f_n : It is given by (23) in the pure case and by (31) in the confluent case. Thus, after rescaling (24) by f_n , the factorial moments have the same shape, but with an additional prefactor 1-p: $\mathbb{E}(X_n^{\underline{s}}) \sim (1-p) \cdot \mathbb{E}(Z_n^{\underline{s}})$, which proves the claim. \square

REMARK 4.5 (Physical interpretation of the bimodal behaviour). As illustrated by Table 1 on page 5, the confluent case gives a bimodal distribution. The first mode is dictated by small values of k, where a Boltzmann distribution is dominant, while for larger values of $k \approx n^{\lambda_H}$, a second mode appears, which is associated with a continuous Mittag-Leffler distribution. This phenomenon explains the following behaviour: If one picks an atom at random in a structure of size n, then with probability p it lies in a large \mathcal{H} -block of size $\Theta(n)$ and with probability 1-p in a smaller \mathcal{H} -block of size $\Theta(n^{1-\lambda_H})$.

In the next section, we refine these considerations by having a closer look at the distribution of the \mathcal{H} -blocks of any given size.

5. Size-refined composition scheme. In this section, we give the limit laws for the profile of combinatorial structures given by a critical composition scheme. We focus here on schemes which are analytically pure (see Definition 3.3), while we handle the confluent and degenerate cases in our companion article [11] (as they require additional technical details and different families of limit laws, which also pop up for $\lambda_H > 1$). The profile is captured by the size-refined composition scheme (4). As we see in the theorem below, we get three successive asymptotic régimes, each leading to its own limit law. Two of these limit laws are expressible in terms of the three-parameter Mittag-Leffler distribution of Theorem 4.1 (see also Definition 3.9 and Definition 3.11).

THEOREM 5.1 (Mixed Poisson limit behaviour for the size-refined scheme). *Consider a size-refined pure critical composition scheme*

$$F(z,v) = G(H(z) - (1-v)h_j z^j)M(z)$$
, with $j \in \mathbb{N}$.

Let $\xi_{n,j} = \frac{\rho_H^j}{-c_H} h_j n^{\lambda_H}$, and X be the three-parameter Mittag-Leffler distribution of Theorem 4.1:

$$X \stackrel{d}{=} \mathrm{ML}(\alpha, \beta, \gamma),$$

where $\alpha = \lambda_H$, $\beta = -\lambda_G \lambda_H$, and $\gamma = -\min(0, \lambda_M)$.

Then, the random variable $X_{n,j}$, which counts the number of \mathcal{H} -components of size j possesses three successive asymptotic régimes, with a phase transition at $j = \Theta(n^{\frac{\lambda_H}{1+\lambda_H}})$:

(i) For $j \ll n^{\frac{\lambda_H}{1+\lambda_H}}$, we have $\xi_{n,j} \to +\infty$ and convergence in distribution and in moments:

$$\frac{X_{n,j}}{\xi_{n,j}} \xrightarrow{d} X.$$

(ii) For $j \sim r \cdot n^{\frac{\lambda_H}{1+\lambda_H}}$, $r \in (0,\infty)$, we have $\xi_{n,j} \to \xi$ with $\xi = r^{-\frac{\lambda_H}{1+\lambda_H}} \cdot \frac{1}{-\Gamma(-\lambda_H)}$ and convergence in distribution and in moments towards a mixed Poisson distribution:

$$X_{n,j} \xrightarrow{d} MPo(\xi X).$$

(iii) For $j \gg n^{\frac{\lambda_H}{1+\lambda_H}}$, we have $\xi_{n,j} \to 0$, and $X_{n,j}$ converges to a Dirac distribution at 0.

REMARK 5.2 (Phase transition I). The intuition behind the phase transition is as follows: In the limit $n\to\infty$, there are many small $(j\ll n^{\frac{\lambda_H}{1+\lambda_H}})$, some giant $(j\sim rn^{\frac{\lambda_H}{1+\lambda_H}})$, and no super-giant $(j\gg n^{\frac{\lambda_H}{1+\lambda_H}})$ \mathcal{H} -components of size j. It is interesting to compare this situation with the birth of the giant component in Erdős–Rényi random graphs [75]. Note that the case $j\in\mathbb{N}$ fixed (i.e., independent of n as n tends to infinity) falls into the case $j\ll n^{\frac{\lambda_H}{1+\lambda_H}}$.

REMARK 5.3 (Phase transition II). For the often observed case of a square-root singularity of H(z) (i.e., $\lambda_H = \frac{1}{2}$), we reobtain the critical range $j = \Theta(n^{1/3})$, which was already observed in the mixed Poisson Rayleigh distributions in [90]. Furthermore, as one has

$$\mathbb{E}(X_{n,j}^{\underline{s}}) = \xi_{n,j}^{s} \cdot \mathbb{E}(X^{s}) \cdot (1 + o(1)),$$

this offers en passant a link between the $\xi_{n,j}$'s and the rescaling factor κn^{λ_H} in Theorem 4.1:

$$\sum_{j>1} \xi_{n,j} = \sum_{j>1} \frac{\rho_H^j}{-c_H} h_j n^{\lambda_H} = \frac{H(\rho_H)}{-c_H} n^{\lambda_H} = \frac{\tau_H}{-c_H} n^{\lambda_H} = \kappa n^{\lambda_H}.$$

This link can be seen as an asymptotic avatar of the combinatorial relation $\sum_{j\geq 1} X_{n,j} = X_n$, implying

$$\sum_{j\geq 1} \mathbb{E}(X_{n,j}) = \mathbb{E}(X_n).$$

In order to prove Theorem 5.1, we first need the following lemma concerning the convergence of mixed Poisson distributions.

LEMMA 5.4 (Factorial moments and limit laws of mixed Poisson type [90]). Let $(X_n)_{n\in\mathbb{N}}$ denote a sequence of random variables, whose factorial moments are asymptotically of mixed Poisson type, i.e., they satisfy for $n \to \infty$ the asymptotic expansion

$$\mathbb{E}(X_{\underline{s}}^{\underline{s}}) = \xi_n^s \cdot \mu_s \cdot (1 + o(1)), \quad s \ge 1,$$

with $\mu_s \ge 0$ and $\xi_n > 0$. Furthermore, assume that the moment sequence $(\mu_s)_{s \in \mathbb{N}}$ determines a unique distribution X satisfying Carleman's condition. Then, the following limit distribution results hold:

- (i) If $\xi_n \to \infty$, the random variable $\frac{X_n}{\xi_n}$ converges in distribution, with convergence of all moments, to X.
- (ii) If $\xi_n \to \xi \in (0, \infty)$, the random variable X_n converges in distribution, with convergence of all moments, to a mixed Poisson distributed random variable $Y \stackrel{d}{=} MPo(\xi X)$.
- (iii) If $\xi_n \to 0$, the random variable X_n converges to a Dirac distribution: $X_n \stackrel{d}{\to} 0$.

PROOF OF THEOREM 5.1. The factorial moments $\mathbb{E}(X_{n,j}^{\underline{s}}) = \sum_{k\geq 0} \mathbb{P}\{X_{n,j} = k\}k^{\underline{s}}$ of $X_{n,j}$ are obtained from F(z,v) by repeated differentiation and evaluation at s=1:

$$\mathbb{E}(X_{n,j}^{\underline{s}}) = \frac{[z^n]\partial_v^s(F)(z,1)}{[z^n]F(z,1)} = h_j^s \frac{[z^{n-j \cdot s}]G^{(s)}(H(z))M(z)}{f_n}.$$

We already know the asymptotics of f_n from (23). For fixed j we can proceed by extraction of coefficients, while for j = j(n) tending to infinity, the asymptotic expansion of h_j follows by singularity analysis applied to H(z) (see Equation (7)):

(33)
$$h_j = \frac{c_H}{\rho_H^j} \cdot \frac{j^{-\lambda_H - 1}}{\Gamma(-\lambda_H)} \cdot (1 + o(1)).$$

What is more, one has

$$(34) \qquad G^{(s)}(H(z))M(z) \sim (-1)^s c_M c_G \rho_G^{-\lambda_G} \lambda_G^{\underline{s}} (-c_H)^{\lambda_G - s} \left(1 - \frac{z}{\rho_H}\right)^{\lambda_H \lambda_G - s\lambda_H + \lambda_M^-}.$$

This implies that $X_{n,j}$ has factorial moments of mixed Poisson type:

$$\mathbb{E}(X_{n,j}^{\underline{s}}) \sim h_j^s(-c_H)^{-s} \rho_H^{js} \cdot \mu_s \cdot n^{s\lambda_H} \quad \text{with} \quad \mu_s := \frac{\Gamma(s - \lambda_G)\Gamma(-\lambda_G\lambda_H - \lambda_M^{\underline{-}})}{\Gamma(-\lambda_G)\Gamma(-\lambda_H\lambda_G + s\lambda_H - \lambda_M^{\underline{-}})}.$$

We already observed that the moment sequence $(\mu_s)_{s\in\mathbb{N}}$ determines a unique distribution, by Carleman's criterion (26). Thus, the limit laws follow by using Lemma 5.4.

Finally, the critical growth range is obtained via the closed-form expression for $\xi_{n,j}$ (in which one inserts the expansion (33)): Indeed, $\xi_{n,j}$ is of growth order $\frac{n^{\lambda_H}}{j^{1+\lambda_H}\Gamma(-\lambda_H)}$ and converges to a nonzero constant if and only if $j(n) \sim r \cdot n^{\frac{\lambda_H}{1+\lambda_H}}$, therefore the critical growth range is $\Theta(n^{\frac{\lambda_H}{1+\lambda_H}})$. Collecting all contributions from $\xi_{n,j}$ (for j=o(n) tending to infinity) gives the constant ξ . The Dirac case is now finally obtained by an additional analysis of the expected value in the remaining range $j\gg n^{\frac{\lambda_H}{1+\lambda_H}}$. There, we directly obtain $\mathbb{E}(X_{n,j})\to 0$, which proves the stated result.

REMARK 5.5 (A probabilistic approach). Note that our Theorem 5.1 was recently revisited by Stufler in [128, Section 3.4] with a probabilistic point of view. He also describes a nice link with a point process giving access to the distribution of the largest components; see [128, Proposition 3.16]. Compare also with the results of Gourdon [58].

Next we turn to the dependence between the number of \mathcal{H} -components of size j_1 and j_2 , determining the covariance and the correlation coefficient.

THEOREM 5.6. In a size-refined pure critical composition scheme, the covariance of the random variables X_{n,j_1} and X_{n,j_2} , counting the number of \mathcal{H} -components of size j_1 and j_2 (with $j_1, j_2 = o(n)$), satisfies

(35)
$$\operatorname{Cov}(X_{n,j_1}, X_{n,j_2}) \sim \xi_{n,j_1} \xi_{n,j_2} \cdot \mathbb{V}(X),$$

where $\xi_{n,j} = \frac{\rho_H^j}{-c_H} h_j n^{\lambda_H} > 0$ and $X \stackrel{d}{=} \mathrm{ML}(\alpha, \beta, \gamma)$ denotes the three-parameter Mittag-Leffler distribution from Theorem 4.1. Furthermore, the correlation coefficient between X_{n,j_1} and X_{n,j_2} satisfies

$$\rho(X_{n,j_1}, X_{n,j_2}) \sim \frac{1}{\sqrt{1 + \frac{\mathbb{E}(X)}{\xi_1}}} \frac{1}{\sqrt{1 + \frac{\mathbb{E}(X)}{\xi_2}}},$$

where, for k=1,2, one has $\xi_k:=\lim_n \xi_{n,j_k}$ and $\sqrt{1+\frac{\mathbb{E}(X)}{\xi_k}}\sim 1$ if $j_k\ll n^{\frac{\lambda_H}{1+\lambda_H}}$.

REMARK 5.7. We observe that for small j_1, j_2 (e.g., if $j_1, j_2 = O(1)$), the random variables are asymptotically highly correlated: $\rho(X_{n,j_1}, X_{n,j_2}) \sim 1$.

PROOF OF THEOREM 5.6. The combinatorial scheme

$$\mathcal{F} = \mathcal{G}(\mathcal{H}_{\neq j_1, j_2} + v_1 \mathcal{H}_{=j_1} + v_2 \mathcal{H}_{=j_2}) \times \mathcal{M}$$

(where one takes distinct sizes $1 \le j_1 < j_2$) directly translates into

$$F(z; v_1, v_2) = G(H(z) - (1 - v_1)h_{j_1}z^{j_1} - (1 - v_2)h_{j_2}z^{j_2})M(z).$$

Accordingly, the random variables X_{n,j_1} and X_{n,j_2} have the joint distribution

$$\mathbb{P}\{X_{n,j_1} = k_1, X_{n,j_2} = k_2\} = \frac{[z^n v_1^{k_1} v_2^{k_2}] F(z; v_1, v_2)}{[z^n] F(z; 1, 1)}.$$

We already know the asymptotics of $f_n = [z^n]F(z;1,1)$, given in (23). We get by differentiation and evaluation at $v_1 = v_2 = 1$

$$\mathbb{E}(X_{n,j_1}X_{n,j_2}) = \frac{[z^n]\partial_{v_1}\partial_{v_2}(F)(z;1,1)}{[z^n]F(z;1,1)} = h_{j_1}h_{j_2}\frac{[z^{n-j_1-j_2}]G''(H(z))M(z)}{f_n}.$$

The asymptotics of h_{j_1} and h_{j_2} are given in (33). The singular expansion of G''(H(z))M(z) is a special case of (34), so we obtain for $j_1, j_2 = o(n)$:

$$\mathbb{E}(X_{n,j_1}X_{n,j_2}) \sim h_{j_1}h_{j_2}c_H^{-2}\rho_H^{j_1+j_2} \cdot \mathbb{E}(X^2) \cdot n^{2\lambda_H}.$$

Hence, using the explicit form of $\mathbb{E}(X)$ in (18), we obtain for the covariance:

$$Cov(X_{n,j_1}, X_{n,j_2}) = \mathbb{E}(X_{n,j_1} X_{n,j_2}) - \mathbb{E}(X_{n,j_1}) \mathbb{E}(X_{n,j_2})$$
$$\sim \frac{h_{j_1} h_{j_2} \rho_H^{j_1 + j_2}}{c_H^2} (\mathbb{E}(X^2) - \mathbb{E}(X)^2) n^{2\lambda_H}.$$

By $\mathbb{V}(X)=\mathbb{E}(X^2)-\mathbb{E}(X)^2$ and the definition of $\xi_{n,j}$, this implies (35). For the correlation coefficient, we observe that Theorem 5.1 together with Lemma 5.4 implies $\mathbb{V}(X_{n,j})\sim \xi_{n,j}^2\mathbb{V}(X)+\xi_{n,j}\mathbb{E}(X)$. Collecting all contributions, this implies that

$$\rho(X_{n,j_1}, X_{n,j_2}) = \frac{\text{Cov}(X_{n,j_1}, X_{n,j_2})}{\sqrt{\mathbb{V}(X_{n,j_1})} \sqrt{\mathbb{V}(X_{n,j_2})}} \sim \frac{1}{\sqrt{1 + \frac{\mathbb{E}(X)}{\xi_{n,j_1}}} \sqrt{1 + \frac{\mathbb{E}(X)}{\xi_{n,j_2}}}}.$$

Taking the limit of this expression for n going to infinity concludes the proof.

Before to present in Section 7 a multivariate generalization of these results, we now list several applications to a variety of combinatorial structures.

6. Applications and examples. Our main results, Theorems 4.1 and 5.1, can be readily applied to the problems considered by Drmota and Soria [31], Flajolet and Sedgewick [42], Dumas, Flajolet and Puyhaubert [39], Janson [73], Meir and Moon [100], Panholzer and Seitz [112] (see also [90] for many additional pointers to the literature) once the required singular expansions of the involved generating functions are established. This includes returns to record-subtrees in Cayley trees, edge-cutting in Cayley trees, returns to zero in Dyck paths, cyclic points and trees in graphs of random mappings, all leading to (mixed Poisson) Rayleigh laws, as well as block sizes in *k*-Stirling permutations.

In the following we discuss several new results for the distribution of different statistics such as returns to zero and sign changes in walks with arbitrary steps, the number of subtrees satisfying some constraint in different fundamental families of trees, as well as the table sizes in the Chinese restaurant process, and the evolution of the number of balls in triangular urn models. These examples illustrate that composition schemes $\mathcal{F} = \mathcal{G}(\mathcal{H}) \times \mathcal{M}$ lead universally to three-parameter Mittag-Leffler distributions (Definition 3.11), which simplify into two-parameter Mittag-Leffler distributions if M(z) has a nonnegative singular exponent.

6.1. Core size of supertrees and a confluent example. Let C denote the family of plane trees (i.e., trees with all arities allowed and embedded into the plane) defined by

$$\mathcal{C} = \mathcal{Z} \times \mathrm{SEQ}(\mathcal{C}), \quad \text{ which translates to } \quad C(z) = \frac{1 - \sqrt{1 - 4z}}{2}.$$

Then, following [42, pp. 412–414, 714], we consider supertrees, or "trees of trees", defined by

$$\mathcal{K} = \mathcal{C}\Big((\mathcal{Z}_{\text{red}} + \mathcal{Z}_{\text{blue}}) \times \mathcal{C}\Big), \quad \text{ which translates to } \quad K(z) = C\Big(2zC(z)\Big).$$

Note that this is a critical scheme as one has $\rho_C = 1/4$ and $\tau_C = C(\rho_C) = 1/2$.

The tree family \mathcal{K} corresponds to trees where onto each node we graft a red or blue tree; one can also draw them like done in Figure 6. By Lagrange inversion, these supertrees \mathcal{K} are thus enumerated by a neat combinatorial sum:

$$K_n = \sum_{k=1}^{\lfloor n/2 \rfloor} \frac{2^k}{n-k} {2k-2 \choose k-1} {2n-3k-1 \choose n-k-1},$$

thus the sequence K_n for $n \ge 2$ starts like 2, 2, 8, 18, 64, 188, 656, 2154, ..., constituting sequence A168506 in the OEIS (On-Line Encyclopedia of Integer Sequence: https://oeis.org).




FIG 6. A bicoloured supertree is a "tree of trees": Each node (here drawn in a potato shape) of some initial plane tree is replaced by a red or a blue node to which one attaches another plane tree.

By the Laplace method or by singularity analysis, this directly leads to the asymptotic expansion $K_n \sim \frac{4^n}{8\Gamma(3/4)n^{5/4}}$. (See also DeVries [26, 116] for an alternative approach to this expansion via multivariate analysis.) This asymptotic behaviour is noteworthy, because one sees here an unusual occurrence of the exponent $-\frac{5}{4}$, while most tree-like structures in combinatorics usually involve the exponent $-\frac{3}{2}$. In fact, one could similarly define supersupertrees, super-super-supertrees, and so on, by further iterations of the critical scheme: $\mathcal{C}_{k+1} = \mathcal{C}_k \left(2\mathcal{Z}\mathcal{C}_k \right)$ with $\mathcal{C}_0 = \mathcal{C}$. This gives an interesting family of combinatorial structures whose asymptotic enumeration involves a dyadic exponent $-1-1/2^{k+1}$; see [5] for a complete characterization of the possible singular exponents for \mathbb{N} -algebraic functions (i.e., generating functions of any structure which can be generated by a context-free language).

With respect to supertrees, the critical scheme is

$$K(z) = G(H(z)), \quad G(z) = C(z), \quad H(z) = 2zC(z),$$

where H(z) has the following Puiseux expansion at $z \sim \frac{1}{4}$:

$$H(z) \sim \frac{1}{4} - \frac{1}{4}\sqrt{1 - 4z}.$$

Now, we can study the core size X_n via the bivariate generating function

$$K(z, u) = C(u \cdot 2zC(z)),$$

as well as the number of \mathcal{H} -components of size j, as captured by

$$K(z,v) = C(2zC(z) + (v-1)2c_{j-1}z^{j}).$$

We can then apply our main Theorems 4.1 and 5.1 (with $\lambda_G = \lambda_H = 1/2$, $\tau_H = \frac{1}{4}$, $c_H = -\frac{1}{4}$, and $\kappa = 1$). This directly gives the following corollaries.

COROLLARY 6.1. The core size X_n in supertrees of size n has factorial moments

$$\mathbb{E}(X_n^{\underline{s}}) \sim n^{s/2} \cdot \mu_s, \qquad \mu_s = \frac{\Gamma(s - \frac{1}{2})\Gamma(-\frac{1}{4})}{\Gamma(-\frac{1}{2})\Gamma(\frac{s}{2} - \frac{1}{4})}.$$

The scaled random variable $X_n/n^{1/2}$ converges in distribution with convergence of all moments to a c=-1/2 moment-tilted stable distribution (a tilted Mittag-Leffler distribution of index 1/2):

$$\frac{X_n}{n^{1/2}} \xrightarrow{d} X, \qquad \textit{with} \qquad X \stackrel{d}{=} \operatorname{tilt}_{-\frac{1}{2}} \left(\operatorname{ML} \left(\frac{1}{2} \right) \right) \stackrel{d}{=} \operatorname{ML} \left(\frac{1}{2}, -\frac{1}{4} \right).$$

Moreover, we have the local limit theorem

$$\mathbb{P}{X_n = x \cdot n^{1/2}} \sim \frac{1}{n^{1/2}} \cdot f_X(x),$$

with $f_X(x)$ denoting the density of the random variable X.

Note that by Legendre's duplication formula one has $\mu_s = 2^s \Gamma(\frac{s}{2} + \frac{1}{4}) / \Gamma(\frac{1}{4})$, so the random variable X can also be seen as equal in law to the chi distribution $\chi(\frac{1}{2})$ of parameter $\frac{1}{2}$, which is itself a generalized gamma distribution [12].

COROLLARY 6.2. The number of coloured trees of size j-1 in supertrees of size n has factorial moments of mixed Poisson type given by

$$\mathbb{E}(X_{n,j}^{\underline{s}}) = \xi_{n,j}^{s} \cdot \mu_{s}(1 + o(1)),$$

with

$$\xi_{n,j} = 2 \cdot (\frac{1}{4})^{j-1} c_{j-1} \cdot n^{1/2}$$

and mixing distribution $X = \chi(\frac{1}{2})$ with $\mathbb{E}(X^s) = \mu_s$.

Furthermore, the random variable $X_{n,j}$ possesses the three successive asymptotic régimes of Theorem 5.1, with a phase transition at $j = \Theta(n^{1/3})$.

We note in passing that a very similar result holds for the family of binary supertrees S, occurring in Bousquet-Mélou's study [22] of the integrated super-Brownian excursion. This family is defined in terms of the family of complete binary trees B:

$$S = \mathcal{B}(\mathcal{Z} \times \mathcal{B}), \qquad \mathcal{B} = \mathcal{Z} + \mathcal{Z} \times \mathcal{B} \times \mathcal{B},$$

with initial values of S_n given by $1,1,3,8,25,80,267,911,\ldots$, constituting sequence A101490 in the OEIs. These functional equations lead to $B(z)=\frac{1-\sqrt{1-4z^2}}{2z}$, and to a critical scheme leading to limit laws similar to the ones of supertrees in Corollaries 6.1 and 6.2.

A confluent example. Next, we consider pairs of supertrees in which we mark the number of \mathcal{H} -trees in the first part of the pair. This translates to the scheme F(z,u)=G(uH(z))K(z) (with G,H, and K defined as in (6.1)), which is confluent because we have $\lambda_H\lambda_G=\lambda_M=\frac{1}{4}$. This case is interesting as $\mathbb{P}\{X_n=k\}$ (the probability that a pair of supertrees has k \mathcal{H} -trees in its first part) converges to the sum of a discrete law and a continuous law. More precisely, as given by Theorem 4.4, and illustrated in Figure 7, the limit is a linear combination of the Boltzmann distribution $\mathcal{B}_C(\frac{1}{4})^{14}$ and the two-parameter Mittag-Leffler distribution $\mathrm{ML}\left(\frac{1}{2},-\frac{1}{4}\right)$.




FIG 7. The empirical distribution $\mathbb{P}\{X_n=k\}$ (drawn with red dots), and its theoretical limiting distribution (in blue). This blue curve is a linear combination of a discrete and a continuous distribution (the middle and right curves drawn in black): $\frac{1}{2}\mathcal{B}_C\left(\frac{1}{4}\right) + \frac{1}{2}\sqrt{n}\operatorname{ML}\left(\frac{1}{2}, -\frac{1}{4}\right)$. The blue theoretical limit curve agrees almost perfectly with the red empirical distribution even for small values of n (here, n=500).

 $^{^{14}}C$ denoting the Catalan generating function, it could be natural to call this Boltzmann distribution $\mathcal{B}_C(\frac{1}{4})$ the *Catalan distribution*, but this name is already used for some other distributions popping up in random matrix theory, and having moments related to the Catalan numbers, like the Marchenko–Pastur distribution.




FIG 8. An increasing diamond and the bijectively equivalent bilabelled increasing plane-oriented recursive tree.

6.2. Root degree and branching structure in bilabelled increasing trees. Bilabelled increasing trees are a natural generalization of increasing trees [84] where every node is assigned two labels instead of just one. General families of bilabelled trees are in bijection with increasing diamonds, which are a natural type of directed acyclic graphs; see [91] for the general statement and Figure 8 for a concrete example. Increasing diamonds model partial orders and their linear extensions, as well as computational processes and their executions in parallel computing [20]. They possess nice combinatorial properties and are enumerated by variants of hook-length formulas [86, 89, 91].

Given a degree-weight sequence $(\varphi_j)_{j\geq 0}$, the corresponding degree-weight generating function is defined as $\varphi(t) = \sum_{j\geq 0} \varphi_j t^j$. The family $\mathcal T$ of bilabelled increasing trees can be described by the following symbolic equation $\mathcal T = \mathcal Z^\square * (\mathcal Z^\square * \varphi(\mathcal T))$, where $\mathcal Z$ denotes single unilabelled nodes, $\mathcal A^\square * \mathcal B$ denotes the boxed product (i.e., the smallest label is constrained to lie in the $\mathcal A$ component) of the combinatorial classes $\mathcal A$ and $\mathcal B$, and $\varphi(\mathcal A) = \varphi_0 \cdot \{\epsilon\} + \varphi_1 \cdot \mathcal A + \varphi_2 \cdot \mathcal A^2 + \ldots$ denotes the class containing all weighted finite labelled sequences of objects of $\mathcal A$ (i.e., each sequence of length k is weighted by φ_k ; ϵ denotes the neutral object of size 0); see [42]. Note that increasing diamonds are associated with $\varphi(t) = \frac{1}{1-t}$. For the exponential generating function $T(z) = \sum_{n\geq 1} T_n \frac{z^n}{n!}$ where n counts the number of labels, the above construction translates into

(36)
$$T''(z) = \varphi(T(z)), \quad T(0) = 0, \quad T'(0) = 0.$$

We now focus on the case of 3-bundled bilabelled increasing trees; see Figure 9.




FIG 9. An increasing plane-oriented recursive tree and the bijectively equivalent 3-bundled bilabelled increasing plane-oriented recursive tree. (The grey ellipses are just here to sketch the bijective correspondences.)

The family of 3-bundled trees is defined by Equation (36) with the following degree-weight generating function

(37)
$$\varphi(t) = \frac{1}{(1-t)^3} = \sum_{k \ge 0} {k+2 \choose 2} t^k.$$

In other words, each node may have any number k of children, and the binomial indicates two bars between these children, thus creating 3 (possibly empty) sequences (or *bundles*) of children. From [89] we get the remarkably simple closed form

(38)
$$T(z) = 1 - \sqrt{1 - z^2} = \sum_{n \ge 1} (2n - 1)!! (2n - 3)!! \frac{z^{2n}}{(2n)!},$$

where the double factorials are defined as

$$(2n-1)!! = \prod_{k=1}^{n} (2k-1) = \frac{(2n)!}{n! \cdot 2^{n}},$$

with initial values of T_{2n} given by 1, 3, 45, 1575, 99225, 9823275, 1404728325, ..., constituting sequence A079484 in the OEIS.

We are interested in the random variable X_n counting the root degree of these 3-bundled bilabelled increasing trees of size n, under the uniform random tree model. Note that by definition there are no bilabelled trees with an odd number of labels, so $T_{2n+1}=0$ and, consequently, $X_{2n+1}=0$. In the following we use the notation $R_n=X_{2n}$ for the root degree.

By (36) and (37), the generating function $T(z,u) = \sum_{n\geq 1} T_n \mathbb{E}(u^{X_n}) \frac{z^n}{n!}$ satisfies

$$\frac{\partial^2}{\partial z^2} T(z, u) = \varphi \left(u T(z) \right) = \frac{1}{\left(1 - u \left(1 - \sqrt{1 - z^2} \right) \right)^3}.$$

Since a derivative with respect to z is simply a shift in the coefficient sequence of exponential generating functions, we obtain

$$\mathbb{E}(u^{R_{n+1}}) = \frac{(2n)!}{T_{2n+2}} [z^n] \frac{1}{(1 - u(1 - \sqrt{1 - z}))^3}.$$

Now we observe, by Stirling's formula for the gamma function and singularity analysis (7) applied to (38), that

(39)
$$\frac{T_{2n+2}}{(2n)!} \sim \frac{2\sqrt{n}}{\sqrt{\pi}} \sim [z^n] \frac{1}{(1-z)^{3/2}}.$$

This implies that, except for the non-standard shift in the random variable, the problem is equivalent (for first-order asymptotics) to the composition scheme $(1-u(1-\sqrt{1-z}))^{-3}$, i.e., $\rho_H=1, \lambda_H=\frac{1}{2}, \lambda_G=-3$, and $\lambda_M=+\infty$ (as M(z)=1 is entire). We note in passing that in this special case, it is also possible to obtain a quite simple closed-form expression for the probability mass function, as well as the (factorial) moments, due to the explicit expressions for the involved generating functions:

$$\mathbb{E}(X_{2n+2}^{\underline{s}}) = \frac{(2n)!}{T_{2n+2}} [z^n] \partial_u^s \frac{1}{(1 - u(1 - \sqrt{1 - z}))^3} \bigg|_{u=1}.$$

However, here we use our general scheme and Theorem 4.1. We apply Legendre's duplication formula and obtain

$$\frac{\Gamma(s+3)\Gamma(\frac{3}{2})}{\Gamma(3)\Gamma(\frac{s+3}{2})} = 2^s \cdot \Gamma\left(\frac{s+4}{2}\right).$$

This leads to the following result.

COROLLARY 6.3. The random variable R_n , counting the root degree in a random strict bilabelled increasing 3-bundled tree with 2n labels, with tree generating function given by $\varphi(t) = (1-t)^{-3}$, satisfies

$$\mathbb{E}(R_{\overline{n}}^s) \sim n^{s/2} \cdot 2^s \cdot \Gamma\left(\frac{s+4}{2}\right).$$

The random variable $R_n/n^{1/2}$ converges (in distribution and in moments) to a multiple of a chi-distributed random variable $X \stackrel{d}{=} \chi(4)$, with four degrees of freedom:

$$\frac{R_n}{n^{1/2}} \xrightarrow{d} \sqrt{2} \cdot X, \quad X \stackrel{d}{=} \chi(4).$$

We can refine the root degree by looking at the branching structure. We denote by $R_{n,j} = X_{2n,j}$ the random variable counting the number of branches (i.e., subtrees) with 2j labels, $1 \le j \le n$, attached to the root:

$$R_n = \sum_{j>1} R_{n,j}.$$

Such random variables naturally arise in the context of the Chinese restaurant process [90, 118, 119] and generalized plane-oriented recursive trees [81]. See also Feng et al. [129] for the analysis of the branching structure of recursive trees. The generating function $T(z,v) = \sum_{n\geq 1} T_n \mathbb{E}(v^{X_{n,j}}) \frac{z^n}{n!}$ satisfies

$$\frac{\partial^2}{\partial z^2} T(z, v) = \varphi \left(T(z) - (1 - v) \frac{T_{2j}}{(2j)!} z^{2j} \right).$$

Consequently,

$$\mathbb{E}(u^{R_{n+1,j}}) = \frac{(2n)!}{T_{2n+2}}[z^{2n}] \left(1 - \left((1 - \sqrt{1 - z^2}) - (1 - v)\frac{T_{2j}}{(2j)!}z^{2j}\right)\right)^{-3}.$$

We can use the asymptotics (39) and apply Theorem 5.1 to obtain the following result.

COROLLARY 6.4. The random variable $R_{n,j}$ counting the number of size 2j branches attached to the root in a random strict bilabelled increasing 3-bundled tree with 2n labels, associated with $\varphi(t) = (1-t)^{-3}$, has factorial moments of mixed Poisson type,

$$\mathbb{E}(R_{n,j}^{\underline{s}}) = \xi_{n,j}^{s} \cdot \mathbb{E}(X^{s})(1 + o(1)),$$

with $\xi_{n,j} = \sqrt{2} \cdot \frac{T_{2j}}{(2j)!} \cdot n^{1/2}$ and mixing distribution $X = \chi(4)$.

Furthermore, the random variable $R_{n,j}$ possesses the three successive asymptotic régimes of Theorem 5.1, with a phase transition at $j = \Theta(n^{1/3})$.

6.3. Returns to zero: walks and bridges with drift zero. A lattice path of length n is a sequence (s_1, \ldots, s_n) of steps $s_i \in \mathcal{S}$ for a fixed finite subset $\mathcal{S} \subseteq \mathbb{Z}$ called step set. Geometrically, we fix the starting point 0 and consider the partial sums $\sum_{i=1}^k s_i$ which can be interpreted as appending the steps one after another. Each step s_i gets a weight $p_i > 0$ and the weight of a path is the product of the weights of its steps. The step polynomial

$$P(u) = \sum_{i} p_i u^i$$

connects the weights and the steps. We call a step set *periodic* if there exist integers $b, p \in \mathbb{Z}$, $p \ge 2$ such that $P(u) = u^b P(u^p)$; otherwise we call it *aperiodic*.




FIG 10. A walk consists of an initial bridge which contains all returns to zero (red dots) and a final walk never returning to zero. The bridge is further decomposed into minimal bridges touching zero only twice.

Here and in the next section, we assume that the step set is aperiodic. Note that this is not a major constraint as the asymptotics of walks with periodic steps can be deduced from the ones with aperiodic ones; see [16]. We call a lattice path a walk if it is unconstrained, and a bridge if it ends at zero, i.e., $\sum_{i=1}^{n} s_i = 0$. A *return to zero* is a point in the path such that $\sum_{i=1}^{k} s_i = 0$; see Figure 10.

Generalizing results from Feller [35, Problems 9–10] and Barton [126, Discussion p. 115], it was shown in [131, Section 3.2] that for drift P'(1) = 0 the law of the number of returns to zero follows a Rayleigh distribution for bridges, while it follows a half-normal distribution for walks. This result follows easily from our Theorem 4.1.

Let $w_{n,k}$ be the number of walks of length n with k returns to zero. The bivariate generating function of walks $W(z,u)=\sum_{n,k\geq 0}w_{n,k}z^nu^k$ is given by

(40)
$$W(z,u) = \frac{1}{1 - u(1 - 1/B(z))} \frac{W(z)}{B(z)},$$

where B(z) and W(z) = 1/(1-zP(1)) are the generating functions of bridges and walks, respectively; see [131, Equation (3.3)]. To explain (40), observe that every bridge is a sequence of *minimal bridges*, which are bridges that never return to the x-axis between the start- and endpoint; see Figure 10. Therefore, minimal bridges are enumerated by 1-1/B(z). Hence, this is exactly the situation of the extended composition scheme (2) with G(z) = 1/(1-z), H(z) = 1-1/B(z), and M(z) = W(z)/B(z). Now, for zero drift, one has

(41)
$$B(z) = \frac{c_B}{\sqrt{1 - zP(1)}} + \mathcal{O}(1) \quad \text{with} \quad c_B = \sqrt{\frac{P(1)}{2P''(1)}}.$$

Hence, one has $\lambda_G = -1$, $\lambda_H = \frac{1}{2}$, and $\lambda_M^- = -\frac{1}{2}$, therefore we are in the pure régime; see Definition 3.3. This is exactly the situation in Remark 4.2 and the number of returns to zero in walks thus follows a half-normal distribution with parameter $\sigma = \sqrt{2}c_B = \sqrt{P(1)/P''(1)}$.

Now, the generating function for bridges is nearly the same as W(z,u) from (40) except that the last factor W(z)/B(z) is omitted. So, by Corollary 4.2, the number of returns to zero here follows a Rayleigh distribution with the same parameter $\sigma = \sqrt{P(1)/P''(1)}$.

We can refine this result for the random variable $X_{n,j}$ counting the number of distance-j-zeroes (which were introduced in [90]). These are the number of returns to zero which have a distance of exactly j steps to the *previous* zero contact. The union over j of distance-j-zeroes gives all returns to zero, and they therefore clearly represent a partition of all returns to zero. Using Theorem 5.1, we then get the following limit theorem.

COROLLARY 6.5. Let $X_{n,j}$ be the number of distance-j-zeroes in lattice paths of length n. For walks (resp. bridges) with zero drift (i.e., P'(1) = 0), $X_{n,j}$ has factorial moments of mixed Poisson half-normal type (resp. mixed Poisson Rayleigh type)

(42)
$$\mathbb{E}(X_{n,j}^{\underline{s}}) = \xi_{n,j}^{s} \cdot \mathbb{E}(X^{s}) \left(1 + o(1)\right),$$

with $\xi_{n,j} = \sqrt{\frac{P(1)}{2P''(1)}} \frac{h_j}{P(1)^j} \cdot n^{1/2}$, where X is given by

$$X = \begin{cases} \mathit{HN}(\sigma) & \mathit{for walks}, \\ \mathit{Rayleigh}(\sigma) & \mathit{for bridges}, \end{cases} \qquad \sigma = \sqrt{\frac{P(1)}{P''(1)}}.$$

Furthermore, the random variable $X_{n,j}$ possesses the three successive asymptotic régimes of Theorem 5.1, with a phase transition at $j = \Theta(n^{1/3})$.

REMARK 6.6 (Universality of the rescaling factor). Note that the rescaling factor $\xi_{n,j}$ in (42) is the same for walks and bridges, while the moment sequence μ_s changes. This independence of $\xi_{n,j}$ can be explained: In Figure 10, the last factor of the walk is a walk not touching zero and is encoded by M(z) = W(z)/B(z). Then by Theorem 5.1 we know that $\xi_{n,j}$ is independent of this factor, and thus has the same value for walks and bridges.

Furthermore, this factor M(z) is also responsible for the often observed dichotomy between half-normal and Rayleigh distributions in the extended composition scheme which we will also observe in the next examples of initial returns and sign changes.

Note that Formula (42) offers a neat factorization for the moments: One can regroup in one factor the quantities with a probabilistic flavour (involving the variance P''(1) of the allowed steps, and μ_s), while the remaining factor $(h_j$, the number of minimal bridges of length j) corresponds to a quantity with a combinatorial flavour. This could also be explained using renewal theory.

6.4. Initial returns in coloured bridges. We generalize the previous model by introducing m-coloured bridges B_m (see Figure 11): We append m non-empty bridges one after the other (and each one with a different colour): $B_m = (B-1)^m$. Then, we are interested in the number of returns to zero in the first bridge, i.e., the initial one that we uniformly coloured. We call such returns the *initial returns*.

Reusing the combinatorial constructions of the previous section, this gives for the bivariate generating function $B_m(z, u)$ the following decomposition

(43)
$$B_m(z,u) = \left(\frac{1}{1 - u(1 - 1/B(z))} - 1\right) (B(z) - 1)^{m-1}.$$




FIG 11. A 4-coloured bridge, with all its initial returns to zero marked by red dots.

For m=1 this is (40) except the factor W(z)/B(z) and the constraint to be non-empty. Asymptotically, and therefore for the law, the non-emptiness is negligible. The generating function $W_m(z,u)$ of m-coloured walks (m-tuples of bridges with a few more steps coloured in the same colour as the final bridge) is given by

$$W_m(z,u) = (1 + B_m(z,u)) \frac{W(z)}{B(z)}.$$

Now, we can directly apply Theorem 4.1. From the reasoning above we see that $\lambda_G = -1$, $\lambda_H = \frac{1}{2}$, and $\lambda_M = -\frac{m-1}{2}$ for bridges and $\lambda_M = -\frac{m}{2}$ for walks.

COROLLARY 6.7. The random variable X_n counting the number of initial returns in a m-coloured walk (resp. bridge) of length n satisfies

$$\mathbb{E}(X_{n}^{\underline{s}}) \sim n^{s/2} \left(\frac{\sigma}{\sqrt{2}}\right)^{s} \mu_{s}, \quad \sigma = \sqrt{\frac{P(1)}{P''(1)}}, \quad \mu_{s} = \begin{cases} \frac{\Gamma(s+1)\Gamma((m+1)/2)}{\Gamma((m+s+1)/2)}, & \textit{for walks,} \\ \frac{\Gamma(s+1)\Gamma(m/2)}{\Gamma((m+s)/2)}, & \textit{for bridges.} \end{cases}$$

The random variable $X_n/n^{1/2}$ converges in distribution with convergence of all moments

$$\frac{X_n}{n^{1/2}} \xrightarrow{d} X, \qquad X \stackrel{d}{=} \begin{cases} \frac{\sigma}{\sqrt{2}} \operatorname{ML}\left(\frac{1}{2}, \frac{1}{2}, \frac{m}{2}\right), & \text{for walks,} \\ \frac{\sigma}{\sqrt{2}} \operatorname{ML}\left(\frac{1}{2}, \frac{1}{2}, \frac{m-1}{2}\right), & \text{for bridges.} \end{cases}$$

These two limit laws can also be seen as the product of independent random variables, namely a Rayleigh and a scaled beta distribution (see Definition 3.10 and Example 3.18):

$$X \stackrel{d}{=} \textit{Rayleigh}(\sigma) \cdot B^{1/2}, \quad \textit{with} \quad B = \begin{cases} \text{Beta}\left(\frac{1}{2}, \frac{m}{2}\right), & \textit{for walks,} \\ \text{Beta}\left(\frac{1}{2}, \frac{m-1}{2}\right), & \textit{for bridges,} \end{cases}$$

where $Beta(\alpha, 0) = 1$. Moreover, we have the local limit theorem

$$\mathbb{P}{X_n = x \cdot n^{1/2}} \sim n^{-1/2} \cdot f_X(x),$$

where, for bridges, the density $f_X(x)$ is given by

$$f_X(x) = \sqrt{\frac{2}{\pi\sigma^2}} \Gamma\left(\frac{m}{2}\right) e^{-\frac{x^2}{2\sigma^2}} U\left(\frac{m}{2} - 1, \frac{1}{2}, \frac{x^2}{2\sigma^2}\right),$$

where U(a,b,x) is the confluent hypergeometric function of the second kind which is the solution of zy''+(b-z)y'-ay=0 such that $U(a,b,x)\sim z^{-a}$ for $z\to\infty$ and $|\arg(z)|<3\pi/2$; see [27, Section 13.2]. For walks, one replaces m by m+1.

Observe the special cases $U(-1/2, 1/2, x) = \sqrt{x}$ and U(0, 1/2, x) = 1 which nicely give the density functions of a Rayleigh (see Example 3.18) and a half-normal distribution (see Example 3.17). Hence, for m = 1 we recover the results of the previous section and uncover a large family of connected probability distributions. It is interesting that this family also appears in the context of preferential attachments in graphs [114, Formula (1.1)].

It is also interesting to consider multicoloured bridges, where we allow any number of colours. We still mark by u the initial returns. The corresponding generating function is

$$B(z,u) = \sum_{m>1} B_m(z,u) = \left(\frac{1}{1 - u(1 - 1/B(z))} - 1\right) \frac{1}{2 - B(z)}.$$

The generating function for the number of multicoloured bridges is thus $B(z,1) = \frac{1}{2-B(z)} - 1$. From (41) we see that B(z) possesses a singularity of order -1/2 at z = 1/P(1), and hence B(z,1) becomes singular at some $z_0 > 0$ which is the unique solution of $B(z_0) = 2$. Hence, the probability generating function reveals a geometric distribution of parameter 1/2:

$$\frac{[z^n]B(z,u)}{[z^n]B(z,1)} \sim \frac{1}{1-u\left(1-1/B(z_0)\right)} - 1 = \frac{u/2}{1-u/2}.$$

As the truncated sum $\sum_{m=1}^{m_0} B_m(z,u)$ behaves asymptotically like $B_{m_0}(z,u)$, we see here a phase transition from a continuous law (for any finite m_0) to a discrete law (when m_0 goes to infinity). Note that this phenomenon holds verbatim for walks.

Finally, let us apply the size-refined scheme Theorem 5.1, counting initial returns in *m*-coloured bridges (or walks) which are a certain distance apart:

COROLLARY 6.8. Let $X_{n,j}$ be the number of initial returns at distance j from the previous zero in m-coloured walks or bridges of length n. Then, $X_{n,j}$ has mixed Poisson type moments

$$\mathbb{E}(X_{n,j}^{\underline{s}}) = \xi_{n,j}^{s} \cdot \mathbb{E}(X^{s}) \left(1 + o(1)\right),\,$$

with
$$\xi_{n,j} = \sqrt{\frac{P(1)}{2P''(1)}} \frac{h_j}{P(1)^j} \cdot n^{1/2}$$
, $h_j = [z^j](1 - 1/B(z))$, and where $X \stackrel{d}{=} \frac{\sigma}{\sqrt{2}} \operatorname{ML}\left(\frac{1}{2}, \frac{1}{2}, \frac{m}{2}\right)$ for walks and $X \stackrel{d}{=} \frac{\sigma}{\sqrt{2}} \operatorname{ML}\left(\frac{1}{2}, \frac{1}{2}, \frac{m-1}{2}\right)$ for bridges, as given in Corollary 6.7. Furthermore, $X_{n,j}$ possesses the three successive asymptotic régimes of Theorem 5.1, with a phase transition at $j = \Theta(n^{1/3})$.

These results also hold for other variants of paths that are in bijection with sequences that already appeared in the literature; see Table 3.

Steps	GF	Sequence	OEIS		
{U, D}	$\frac{8z^2 - 2 - \sqrt{1 - 4z^2}}{16z^2 - 3}$	$1, 0, 2, 0, 10, 0, 52, 0, 274, 0, 1452, 0, 7716, \dots$	A075436		
$\{U,D,H_1\}$	$\frac{\frac{5z-2}{16z^2-3}}{\frac{z+\sqrt{1-4z^2}}{1-5z^2}}$	$1, 1, 3, 5, 13, 25, 61, 125, 295, 625, 1447, \dots$	A098615		
$\{U,D,H_2\}$	$\frac{z^2 + \sqrt{1 - 4z^2}}{1 - 4z^2 - z^4}$	$1, 0, 3, 0, 11, 0, 43, 0, 173, 0, 707, 0, 2917, \dots$	A026671		
TABLE 3					

Multicoloured bridge models: they end at 0 and use up steps U=(1,1), down steps D=(1,-1), and horizontal steps $H_i=(i,0)$ allowed only at altitude 0. The limit laws of initial returns to zero in these models are all the same and special cases of Corollaries 6.7 and 6.8.

6.5. Sign changes in walks. Using the same notation as in Example 6.3, we now define the sign of the path (s_1,\ldots,s_n) after k steps as $\mathrm{sgn}(\sum_{i=1}^k s_i) \in \{-1,0,1\}$. Thereby every lattice path is associated with a sequence of signs. A sign change is therein any subsequence $(-1,0^*,1)$ or $(1,0^*,-1)$ where 0^* denotes a (possibly empty) sequence of 0s; see Figure 12. In this section we consider Motzkin paths. They are composed of up steps +1, down steps -1, and horizontal steps 0; see again Figure 12. Their step polynomial is therefore given by $P(u) = \frac{p-1}{u} + p_0 + p_1 u$ (with $p_{-1}p_0p_1 \neq 0$). In the case of zero drift, let us show how to apply our results to get that the number of sign changes follows asymptotically a Rayleigh distribution for bridges and a half-normal distribution for walks, while for nonzero drift it

follows a geometric distribution; see [131].




FIG 12. A Motzkin walk (i.e., step set $S = \{-1, 0, 1\}$) with 4 sign changes marked in red.

Combinatorially, we see that the bivariate generating function of bridges is

$$B(z,u) = S(z)\left(1 + \frac{2H(z)}{1 - uH(z)}\right), \quad \text{ where } S(z) = \frac{1}{1 - p_0 z} \quad \text{ and } \quad H(z) = \frac{E(z)}{S(z)} - 1.$$

Here, S(z) counts sequences of horizontal steps, E(z) counts excursions (bridges constrained to be nonnegative; see [6]), and H(z) counts excursions which start with an up or a down step (and not with a horizontal step).

We now give the main corresponding Puiseux expansions. First one has

$$H(z) = 1 - 2\sqrt{\frac{2P(1)}{P''(1)}}\sqrt{1 - zP(1)} + \mathcal{O}(1 - zP(1)).$$

Then, as the radius of convergence $1/p_0$ of S(z) is strictly larger than 1/P(1), which is the one of H(z), we see that the additive term S(z) is negligible for the limit law. Thus, we have a composition scheme (2) where M(z) = 2S(z)H(z) has the asymptotic expansion

$$M(z) = 2E\left(\frac{1}{P(1)}\right) + \mathcal{O}\left(\sqrt{1 - zP(1)}\right).$$

Hence, we have $\lambda_M^-=0$, which means that the factor M(z) is asymptotically negligible for the law. The asymptotic dominant part arises from $\frac{1}{1-uH(z)}$ and we get from Corollary 4.2 the

expected convergence to a Rayleigh distribution with parameter
$$\sigma = -\sqrt{2} \frac{\tau_H}{c_H} = \frac{1}{2} \sqrt{\frac{P''(1)}{P(1)}}$$
.

A similar reasoning (and Remark 4.2) allows us to prove that the number of sign changes in walks asymptotically follows a half-normal distribution with the same parameter σ . We now refine the analysis by counting sign changes which are j steps apart. Then we can apply Theorem 5.1 to get the following refined result which strongly depends on $H(z) = \sum_{j \geq 0} h_j z^j$. Note that a statement analogous to Remark 6.6 also applies here.

COROLLARY 6.9. For walks of length n of Motzkin paths, let the random variable $X_{n,j}$ be the number of sign changes at distance j from the previous sign change or the origin. For walks (resp. bridges) with zero drift (i.e., P'(1) = 0), $X_{n,j}$ has factorial moments of mixed Poisson half-normal type (resp. mixed Poisson Rayleigh type)

$$\mathbb{E}(X^{\underline{s}}_{n,j}) = \xi^s_{n,j} \cdot \mathbb{E}(X^s) \left(1 + o(1)\right),$$

with $\xi_{n,j}=\frac{1}{2}\sqrt{\frac{P''(1)}{2P(1)}}\frac{h_j}{P(1)^j}\cdot n^{1/2}$ and mixing distributions

$$X \stackrel{d}{=} \begin{cases} \mathit{HN}(\sigma) & \mathit{for walks}, \\ \mathit{Rayleigh}(\sigma) & \mathit{for bridges}, \end{cases} \qquad \sigma = \frac{1}{2} \sqrt{\frac{P''(1)}{P(1)}}.$$

Furthermore, the random variable $X_{n,j}$ (for walks and for bridges) possesses the three successive asymptotic régimes of Theorem 5.1, with a phase transition at $j = \Theta(n^{1/3})$.

6.6. Tables in the Chinese restaurant process. Following Aldous, Pitman, and Dubins (see [3,118,119]), we now consider the Chinese restaurant process. This a discrete-time stochastic process having as value at time n one of the B_n partitions of the set $[n] = \{1, 2, \ldots, n\}$ (where B_n denotes the Bell numbers found as sequence A000110 in the OEIS). One fancifully imagines a Chinese restaurant with an infinite number of tables, where each table has a possibly infinite number of seats. In the beginning the first customer takes a seat at the first table. At each discrete time step a new customer arrives and either joins one of the existing tables, or takes a seat at the next empty table. Each table corresponds to a block of a random partition. The process thus starts at time n=1 with the partition $\{\{1\}\}$. Now, given a partition $T=\{t_1,\ldots,t_k\}$ of [n] with |T|=k parts t_i , at time n+1 the element n+1 is either added to one of the existing parts $t_i \in T$ with probability

$$\mathbb{P}\{n+1 \prec t_i\} = \frac{|t_i| - \alpha}{n+\beta}, \quad 1 \le i \le k,$$

where $n+1 \prec t_i$ denotes that n+1 is a costumer sitting at table t_i , or as a new singleton block with probability

$$\mathbb{P}\{n+1 \prec t_{|T|+1}\} = \frac{\beta + k \cdot \alpha}{n+\beta}.$$

This model (parametrized by the two parameters $0 < \alpha < 1$ and $\beta > -\alpha$) thus assigns a probability to any particular partition T of [n]. We are interested in the random variable C_n , counting the total number of tables in the Chinese restaurant process, as well as the random variable $C_{n,j}$, counting the number of parts of size j in a partition of [n]. As pointed out in [90], this process can be embedded into a variant of the growth process of generalized plane-oriented recursive trees with two different connectivity parameters a > 0 and b > -1. This allows us to study properties of the Chinese restaurant process using analytic combinatorial tools. For the reader's convenience, we restate this embedding below.

We collect the results of [90] and complement them by extending the constraint b > 0 to the full range b > -1 as well as by providing the missing identification of the limit law as a (moment-shifted) stable law.

Combinatorially, we consider a family $\mathcal{T}_{a,b}$ of generalized plane-oriented recursive trees, where the degree-weight generating function $\psi(t) = \frac{1}{(1-t)^b}$, b>0, associated with the root of the tree, is different to the one for non-root nodes in the tree, $\varphi(t) = \frac{1}{(1-t)^a}$, a>0. Then, the family $\mathcal{T}_{a,b}$ is closely related to the corresponding family \mathcal{T} of generalized plane-oriented recursive trees with degree-weight generating $\varphi(t) = \frac{1}{(1-t)^a}$, a>0, via the following formal recursive equations (see Section 6.2 for the definition of the boxed product):

$$\mathcal{T}_{a,b} = \mathcal{Z}^{\square} * \psi(\mathcal{T}), \qquad \mathcal{T} = \mathcal{Z}^{\square} * \varphi(\mathcal{T}).$$




FIG 13. A plane-oriented recursive tree of size 15 and the corresponding tables in the Chinese restaurant model.

The weight w(T) of a tree $T \in \mathcal{T}_{a,b}$ is then defined by

$$w(T) = \psi_{d(\text{root})} \prod_{v \in T \setminus \{\text{root}\}} \varphi_{d(v)},$$

where d(v) denotes the outdegree of node v. Thus, the generating functions

$$T_{a,b}(z) = \sum_{n\geq 1} T_{a,b;n} \frac{z^n}{n!}$$
 and $T(z) = \sum_{n\geq 1} T_n \frac{z^n}{n!}$

of the total weight of size-n trees in $\mathcal{T}_{a,b}$ and \mathcal{T} , respectively, satisfy the differential equations

$$T'_{a,b}(z) = \psi(T(z))$$
 and $T'(z) = \varphi(T(z))$.

The ordinary tree evolution process to generate a random tree of arbitrary given size in the family \mathcal{T} (see [111] for a detailed discussion) can be extended in the following way to generate a random tree in the family $\mathcal{T}_{a,b}$. The process, evolving in discrete time, starts with the root labelled zero. At step n+1, with $n \ge 0$, the node with label n+1 is attached as a new child to any previous node v (this is denoted by $n+1 \prec v$) with probabilities

$$\mathbb{P}\{n+1 \prec v\} = \begin{cases} \frac{d(v)+b}{b+(a+1)n} & \text{if } v \text{ is the root,} \\ \frac{d(v)+a}{b+(a+1)n} & \text{if } v \text{ is not the root.} \end{cases}$$

We recall the following result from [90].

PROPOSITION 6.10 (Chinese restaurant process and generalized plane-oriented recursive trees). A random partition of $\{1, \dots, n\}$ generated by the Chinese restaurant process with parameters $0 < \alpha < 1$ and $\beta > 0$ can be generated equivalently by the growth process of the family of generalized plane-oriented recursive trees $\mathcal{T}_{a,b}$ when generating such a tree of size n+1. The parameters α, β and a,b>0, respectively, are related via

$$\alpha = \frac{1}{1+a}, \qquad \beta = \frac{b}{1+a}.$$

The random variable C_n is distributed as the outdegree X_{n+1} of the root of a random generalized plane-oriented recursive trees of size n+1 from the family $\mathcal{T}_{a,b}$: $C_n \stackrel{d}{=} X_{n+1}$. The random variable $C_{n,j}$ is distributed as the number $X_{n+1,j}$ of branches of size j attached

to the root of a random tree of size n+1 from the family $\mathcal{T}_{a,b}$: $C_{n,i} \stackrel{d}{=} X_{n+1,i}$.

Note that in the above relation, β cannot be negative, since b is assumed to be positive. As already observed in [90], the correspondence can be extended to the full range $0 < \alpha < 1$ and $\beta > -a$, where one has $a = \frac{1}{\alpha} - 1 > 0$ and $b = \frac{\beta}{\alpha} > -1$. For $-1 < b \le 0$, we cannot directly use the degree-weight generating function $\psi(t) = (1-t)^{-b}$. Indeed, for -1 < b < 0 we would have $\psi(t) = 1 + bt + \dots$, involving a negative weight; while for b = 0 we would have $\psi(t) = 1$, a degenerate case. However, we can use a modified generating function, leading to a correct model of the Chinese restaurant process in the range $-1 < b \le 0$ (see [90, 111] for more details on the growth process):

$$\psi(t) = 1 + \int_0^t \frac{1}{(1-x)^{1+b}} dx = 1 + \frac{1}{b} \left(\frac{1}{(1-t)^b} - 1 \right) = 1 + \sum_{k \ge 1} \binom{b+k}{k-1} \frac{t^k}{k},$$

for -1 < b < 0, while for b = 0 one uses

$$\psi(t) = 1 - \log(1 - t) = 1 + \sum_{k>1} \frac{t^k}{k}.$$

Thus, we have some generalized plane-oriented recursive trees attached to a root with a different tree-weight generating function $\psi(t)$. Summarizing, we have

$$\psi(t) = \begin{cases} \frac{1}{(1-t)^b} & \text{if } b > 0, \\ 1 - \log(1-t) & \text{if } b = 0, \\ 1 + \frac{1}{b} \left(\frac{1}{(1-t)^b} - 1 \right) & \text{if } -1 < b < 0. \end{cases}$$

Here (except for the special case b = 0, which is handled by a slightly different approach, detailed later in Theorem 7.1), we can directly apply our results from Theorem 4.1 to

$$T'_{a,b}(z,u) = \sum_{n>1} T_n \mathbb{E}(u^{X_n}) \frac{z^{n-1}}{(n-1)!} = \psi(u \cdot T(z)), \qquad T'(z) = \varphi(T(z)),$$

for the total number of tables, and from Theorem 5.1 to

$$R'_{a,b}(z,v) = \sum_{n>1} T_n \mathbb{E}(v^{X_{n,j}}) \frac{z^{n-1}}{(n-1)!} = \psi(T(z) - (1-v)z^j \frac{T_j}{j!}),$$

for the number of tables of size j. This allows us to extend the corresponding result of [90] to the full range of b > -1, also providing the missing identification of the limit law as a (moment-tilted) stable law:

THEOREM 6.11. Let a > 0, b > -1. The random variable $X_{n,j}$ counting the number of branches of size j in a random $\mathcal{T}_{a,b}$ tree of size n (or, equivalently, the number of tables with j seated customers in a Chinese restaurant process of parameter $\alpha = 1/(1+a)$ and $\beta = b/(1+a)$, with a total of n-1 customers) possesses the three successive asymptotic régimes of Theorem 5.1, with a phase transition at $j = \Theta(n^{1/(a+2)})$:

(i) For $j \ll n^{\frac{1}{a+2}}$ we have $\xi_{n,j} = \frac{\alpha n^{\alpha}}{j} {j-1-\alpha \choose j-1} \to \infty$ and $\frac{X_{n,j}}{\xi_{n,j}}$ converges in distribution, with convergence of all moments, to a two-parameter Mittag-Leffler distribution:

$$\frac{X_{n,j}}{\xi_{n,j}} \xrightarrow{d} X \qquad with \qquad X \stackrel{d}{=} \mathrm{ML}(\alpha,\beta).$$

(ii) For $j \sim r \cdot n^{\frac{1}{a+2}}$, $r \in (0, \infty)$, we have $\xi_{n,j} \to \xi$, and the random variable $X_{n,j}$ converges in distribution, with convergence of all moments, to a mixed Poisson distribution:

$$X_{n,j} \xrightarrow{d} MPo(\xi X).$$

(iii) For $j \gg n^{\frac{1}{a+2}}$ we have $\xi_{n,j} \to 0$, so $X_{n,j}$ converges to a Dirac distribution at 0.

REMARK 6.12. Our result above implies that there are only a few giant tables in the Chinese restaurant process (a mixed-Poisson number of tables with a number of customers proportional to $n^{\frac{1}{a+2}}$). In contrast, there are much more tables with a smaller number of customers, and an asymptotically negligible number of tables of size $\gg n^{\frac{1}{a+2}}$.

REMARK 6.13. Our results also allow recovering the limit theorem in [119] for the total number of tables C_n in the Chinese restaurant process (via X_n), albeit with a totally different proof, as the normalized random variable X_n/n^{α} converges in distribution with convergence of all moments to a random variable X, with X given in the theorem before. For the reader's convenience, we state the moments in terms of β and α , compare with [119, Theorem 3.8]:

$$\mathbb{E}(X^s) = \frac{\Gamma(s + \frac{\beta}{\alpha})\Gamma(\beta)}{\Gamma(\beta + s \cdot \alpha)\Gamma(\frac{\beta}{\alpha})}.$$

PROOF OF THEOREM 6.11. We follow very closely [90] and sketch the remaining steps. We solve the differential equation $T'(z) = \varphi(T(z))$ and get $T(z) = 1 - (1 - (a+1)z)^{\frac{1}{a+1}}$. Thus, the probability generating function is given by

$$\mathbb{E}(v^{X_{n+1,j}}) = \frac{n!}{T_{n+1}} [z^n] \psi \left(T(z) - (1-v)z^j \frac{T_j}{j!} \right),$$

where the coefficient $\frac{T_{n+1}}{n!} = [z^n]\psi(T(z))$ is computed by standard singularity analysis. Therefore, except for the non-standard shift, we can readily apply our scheme to

$$\psi\left(T(z)-(1-v)z^j\frac{T_j}{j!}\right).$$

Here, T=H and $\psi=G$, whose Puiseux exponents are $\lambda_H=\frac{1}{a+1}$ and $\lambda_G=-b$ for $b\neq 0$, where a>0 and b>-1. Hence, the critical range is given by $j(n)=\Theta(n^{\frac{\lambda}{1+\lambda H}})=\Theta(n^{\frac{1}{a+2}})$. In this model, no additional factor M(z) is present, so $\lambda_M^-=0$, and if b=0 we apply the cycle scheme of Theorem 7.3. This gives

$$\mathbb{E}(X^s) = \mu_s = \begin{cases} \frac{\Gamma(s+b)\Gamma(\frac{b}{a+1})}{\Gamma(b)\Gamma(\frac{b+s}{a+1})} & \text{if } b \neq 0, \\ \frac{\Gamma(s+1)}{\Gamma(\frac{s}{a+1}+1)} & \text{if } b = 0. \end{cases}$$

Finally, we unify both expressions by simply using $\Gamma(x+1) = x\Gamma(x)$.

REMARK 6.14 (Chinese restaurant with a bar). Recently, Möhle introduced in [104] a generalization of the classical Chinese restaurant process, in which in addition to the tables, the customers can sit at an infinite bar. After proper rescaling, the number of customers at the bar converges to a beta-distributed random variable, and the number of occupied tables to the three-parameter Mittag-Leffler distribution.

6.7. Triangular urn models and the three-parameter Mittag-Leffler distribution. Two-colour triangular urns are instances of generalized Pólya urn models [39, 70, 94]. At each time step $n \geq 1$, a ball is drawn uniformly at random, reinserted, and depending on the observed colour, balls of both colours are added to the urn: If a white ball was drawn, we add a white and b black balls, whereas, if a black ball was drawn, we add c white and d black balls. The addition/replacement of balls can be described by the so-called ball replacement matrix $M = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$, where for balanced urn models it holds that a + b = c + d, such that the total number $\sigma = a + b$ of added balls in each step is independent of the observed colour. The initial configuration of the urn consists of w_0 white balls and b_0 black balls, and the random variable \mathcal{W}_n counts the number of white balls in the urn after n draws. For balanced urns with replacement matrix

$$M = \begin{pmatrix} a & b \\ 0 & d \end{pmatrix}, \quad a, b > 0, \quad d = \sigma = a + b,$$

it was shown by Flajolet, Dumas, and Puyhaubert [39] (and also by Janson [72,74] via different analytic methods) that $\frac{W_n}{an^{\alpha/\sigma}} \stackrel{d}{\to} W$, for a random variable with moments

$$\mathbb{E}(\mathcal{W}^s) = \frac{\Gamma(\frac{b_0 + w_0}{\sigma})}{\Gamma(\frac{w_0}{a})} \cdot \frac{\Gamma(s + \frac{w_0}{a})}{\Gamma(s \cdot \frac{a}{\sigma} + \frac{b_0 + w_0}{\sigma})}.$$

The limit law thus depends on the initial composition of the urn, as it is often the case for urn models. In the special case $(w_0, b_0) = (a, b)$, and thus $w_0 + b_0 = \sigma$, \mathcal{W} follows a Mittag-Leffler distribution $\mathrm{ML}(a/\sigma)$. For $b_0 > 0$ and either $w_0 = 0$ or $w_0 = b$, Janson observed a moment-tilted stable law, leaving the other cases open; see [72, Theorem 1.8 and Problem 1.15].

Let us now show how the general case can be solved using our extended scheme. The key tool is the history generating function $F(z,u) = \sum_{n,k\geq 0} f_{n,k} u^k \frac{z^n}{n!}$ where $f_{n,k}$ is equal to the number of transitions (or histories) leading to a configuration with k white balls after n steps.

The closed form of this history generating function was derived in [39, Proposition 14]:

$$F(z,u) = u^{w_0} (1 - \sigma z)^{-b_0/\sigma} \left(1 - u^a \left(1 - (1 - \sigma z)^{a/\sigma} \right) \right)^{-w_0/a}.$$

Putting aside the prefactor u^{w_0} , and after a change of variable $u^a \mapsto u$, this equation can be interpreted as an extended critical composition scheme

(44)
$$F(z,u) = M(z) \cdot G(uH(z)),$$

involving the exponential generating functions with nonnegative integer coefficients

$$M(z) = (1 - \sigma z)^{-b_0/\sigma}, \ G(z) = (1 - az)^{-w_0/a}, \ \text{and} \ H(z) = (1 - (1 - \sigma z)^{a/\sigma})/a.$$

The fact that the singular exponents depend on b_0 and w_0 explains en passant why the limit distribution of W_n differs according to the initial composition of the urn. Indeed, as the number of white balls at time n satisfies

$$\mathbb{P}\{\mathcal{W}_n = ak + w_0\} = \frac{n![z^n u^k] F(z, u)}{n![z^n] F(z, 1)} = \frac{g_k}{k!} \frac{[z^n] H(z)^k M(z)}{[z^n] F(z, 1)},$$

we can apply Theorem 4.1 and we then get the following limit distributions of W_n for balanced triangular urn models, completing and extending earlier results [72, Theorem 1.8]:

COROLLARY 6.15. Let W_n be the random variable for the number of white balls in a balanced triangular Pólya urn with initially $w_0 > 0$ white and $b_0 \ge 0$ black balls. Then, we have a convergence in distribution, with convergence of all moments, towards a three-parameter Mittag-Leffler distribution (see Definition 3.11)

$$\frac{\mathcal{W}_n}{an^{a/\sigma}} \xrightarrow{d} \mathrm{ML}\left(\frac{a}{\sigma}, \frac{w_0}{a}, \frac{b_0}{a}\right).$$

REMARK 6.16 (Almost sure convergence and beyond). This limit was also recently identified by Goldschmidt, Haas, and Sénizergues [53] for urns with non-integer weights: A link with the Chinese restaurant model (for $b_0 = 0$) leads to a Mittag-Leffler distribution, then they show that the impact of $b_0 > 0$ on the process leads to a distribution with an additional beta law factor. It is interesting to stress that their approach implies an almost sure convergence. Note that the fluctuations around this almost sure limit are known: A second-order central limit theorem (that is, the random variable minus its almost sure limit converges, rescaled, to a Gaussian distribution), as well as a law of the iterated logarithm was obtained using a discrete martingale [92]. What is more, following Gouet [57], a continuous-time reparametrization leads to a functional second-order limit theorem for balanced urn models. There is currently no systematic way to obtain an almost sure convergence for all combinatorial models covered by our composition schemes; however, in a few cases (e.g., for walks, trees, and maps), some ad-hoc clever constructions entail this almost sure convergence [96–98, 101, 124].

Note that applying Theorem 5.1 to the size-refined version of the composition scheme (44), we get factorial moments of mixed Poisson type for a size-refined random variable $X_{n,j}$ and the corresponding limit laws. However, the combinatorial interpretation of the random variable(s) $X_{n,j}$ is more involved and will be given elsewhere.

We stress the fact that the methods and results presented in this article are thus holding both for ordinary generating functions (typically used for unlabelled structures) and for exponential generating functions (typically used for labelled structures); see [42].

This concludes the list of applications for our results on the extended and size-refined composition schemes. We now give some extensions of our work to other schemes.

7. Further extensions.

7.1. *Critical cycle scheme*. Many combinatorial structures are cycles of more basic building blocks (e.g., cyclic permutations or functional applications are cycles of Cayley trees). If one marks the number of such basic building blocks, this corresponds to

$$\mathcal{F} = \mathcal{G}(\mathcal{H}) = \text{CYC}(\mathcal{H}) \implies F(z, u) = -\log(1 - uH(z)),$$

where $\mathcal{G}=\mathrm{CYC}$ denotes the cycle operator. This scheme is analysed in Flajolet and Sedgewick's magnum opus [42, page 414] in the supercritical case, and we now extend this analysis to the critical case (i.e., by Definition 1.1 for $H(\rho_H)=1$). Note that the previous sections were assuming Puiseux-like expansions for the generating function F(z,1) at its dominant singularity $z=\rho=\rho_H$. Now, for critical cycle schemes, F does not have a Puiseux expansion, so the previous results need to be adapted.

Let us begin with an example: For $H(z) = 1 - \sqrt{1 - 2z}$ we get the sequence

$$n![z^n]F(z) = n![z^n]\frac{1}{2}\log\left(\frac{1}{1-2z}\right) = 2^{n-1}(n-1)! = (2n-2)!!, \quad n \ge 1,$$

which starts with $1, 2, 8, 48, 384, 384, \dots$, and constitutes the entry A000165 in the OEIS. Here, the moments $\mathbb{E}(X_n^s)$ are of order $n^{s/2}$, so the scaling with $1/\sqrt{n}$ leads directly to moment convergence. This is just one instance of the following more general result.

THEOREM 7.1 (Critical schemes with a log). In a critical cycle composition scheme

(45)
$$F(z,u) = -\log(1 - uH(z)),$$

if H(z) has a singular exponent $0 < \lambda_H < 1$, the core size X_n (i.e., the number of H-components in structures of size n) has factorial moments given by

$$\mathbb{E}(X_n^{\underline{s}}) \sim \kappa n^{s\lambda_H} \mu_s, \quad \text{with} \qquad \kappa = \frac{1}{-c_H} \quad \text{and} \quad \mu_s = \frac{\Gamma(s+1)}{\Gamma(s\lambda_H + 1)}.$$

The scaled random variable $X_n/(\kappa n^{\lambda_H})$ converges in distribution with convergence of all moments to a Mittag-Leffler distributed random variable $X \stackrel{d}{=} M_{\lambda_H}$.

REMARK 7.2. Observe that this scheme leads to a distribution similar (except for a shift in the moments) to the one obtained for the scheme involving the sequence operator $\mathcal{G}=\operatorname{SEQ}$, i.e., $G(z)=\frac{1}{1-z}$, for which one has $c_G=1$, $\lambda_G=-1$, and $\rho_G=1$. Alternatively, we may think of this cycle scheme as the limit case of Theorem 4.1 when $\lambda_G\to 0$. Indeed, for $\lambda_M^-=0$ the moments (18) of the extended composition scheme can be rewritten into

$$\mathbb{E}(X^s) = \frac{\Gamma(s - \lambda_G + 1)\Gamma(-\lambda_G \lambda_H + 1)}{\Gamma(s\lambda_H - \lambda_G \lambda_H + 1)\Gamma(-\lambda_G + 1)}.$$

Thus, for $\lambda_G \to 0$ the moments of the random variable X converge to the moments of an ordinary Mittag-Leffler distribution; see Definition 3.7. Similarly, taking the limit $\lambda_G \to 0$ in Remark 4.2 gives $\lambda_G = 0$ as the tilting parameter, resulting again in the ordinary Mittag-Leffler distribution.

Now, for $j \in \mathbb{N}$, we can also look at the size-refined scheme

$$\mathcal{F} = \operatorname{CYC}(v\mathcal{H}_{=j} + \mathcal{H}_{\neq j}),$$

for which we get the following theorem.

THEOREM 7.3 (Size-refined critical schemes with a log). *In the size-refined critical cycle composition scheme*

(46)
$$F(z,v) = -\log\left(1 - \left(H(z) - (1-v)h_{j}z^{j}\right)\right), \quad j \in \mathbb{N},$$

if H(z) has a singular exponent $0 < \lambda_H < 1$, then the number $X_{n,j}$ of \mathcal{H} -components of size j in structures of size n has factorial moments of mixed Poisson type,

$$\mathbb{E}(X_{n,j}^{\underline{s}}) = \xi_{n,j}^{s} \cdot \mu_{s} \cdot (1 + o(1)),$$

with $\xi_{n,j} = \frac{\rho_H^j}{-c_H} h_j n^{\lambda_H}$ and Mittag-Leffler mixing distribution $X \stackrel{d}{=} M_{\lambda_H}$. The random variable $X_{n,j}$ converges to one of the three limit laws given in Theorem 5.1, depending on whether j=j(n) is smaller, equal, or larger than the critical growth range $j=\Theta(n^{\frac{\lambda_H}{1+\lambda_H}})$.

PROOFS OF THEOREM 7.1 AND 7.3. The proofs are analogous to those of Theorems 4.1 and 5.1, and we only point out the differences next. Let us start with the proof of Theorem 7.1. The factorial moments of order s satisfy

$$\mathbb{E}(X_{\underline{n}}^{\underline{s}}) = \frac{[z^n]\partial_u^s(F)(z,1)}{[z^n]F(z,1)},$$

where F is defined by Equation (45). As the scheme is critical (i.e. $H(\rho_H) = 1$), one has

$$F(z,1) = -\log(1 - H(z)) = -\log\left((1 - z/\rho_H)^{\lambda_H}(1 + o(1))\right) \sim -\lambda_H \cdot \log\left(1 - z/\rho_H\right).$$

Using the transfer theorems of [42] we directly obtain $[z^n]F(z,1) \sim \lambda_H \frac{\rho_h^{-n}}{n}$. It remains to compute $\partial_u^s F$. Note that the log function can be replaced by a quasi-inverse using

$$\partial_u^s \log \left(\frac{1}{1-u} \right) = \partial_u^{s-1} \frac{1}{1-u}.$$

Thus, the sth factorial moment is obtained from the asymptotics in (24) computed for s-1 and with $G(z) = \frac{1}{1-z}$. Then, we obtain the final result: the normalized moments converge to the moments of a Mittag-Leffler distribution.

For the proof of Theorem 7.3 one replaces (45) by (46) and $\partial_u^s F$ by $\partial_v^s F$.

7.2. *Multivariate critical composition schemes*. It is possible to generalize the critical composition scheme by looking at combinatorial constructions of the form

$$\mathcal{F} = \mathcal{M} imes \mathcal{G}_1ig(\mathcal{H}_1ig) imes \mathcal{G}_2ig(\mathcal{H}_2ig) imes \cdots imes \mathcal{G}_mig(\mathcal{H}_mig) = \mathcal{M} imes \prod_{\ell=1}^m \mathcal{G}_\ellig(\mathcal{H}_\ellig).$$

We measure the size of the \mathcal{G}_{ℓ} component by the variable u_{ℓ} ; accordingly this gives

(47)
$$F(z, u_1, \dots, u_m) = M(z) \cdot \prod_{\ell=1}^m G_\ell(u_\ell H_\ell(z)).$$

The random vector $\mathbf{X}_n = (X_{n,1}, \dots, X_{n,m})$ measures the sizes of the \mathcal{G}_ℓ -components,

$$\mathbb{P}\{X_{n,1} = k_1, \dots, X_{n,m} = k_m\} = \frac{[z^n u_1^{k_1} \dots u_m^{k_m}] F(z, u_1, \dots, u_m)}{[z^n] F(z, 1, \dots, 1)}.$$

We now introduce a suitable extension of the terms critical (Definition 1.1) and pure (Definition 3.3) for multivariate schemes. We call a multivariate scheme critical if all functions $H_{\ell}(z)$ have the identical radius of convergence $\rho_{H_{\ell}} = \rho_H$ such that $\tau_{\ell} := H_{\ell}(\rho_H) = \rho_{G_{\ell}}$ and M(z) has radius of convergence $\rho_M \ge \rho_H$. We call a multivariate scheme pure if

- $H_{\ell}(z)$ has a singular exponent $0 < \lambda_{H_{\ell}} < 1$ for $1 \le \ell \le m$;
- $G_{\ell}(z)$ has a singular exponent $\lambda_{G_{\ell}} < 0$ for $1 \le \ell \le m$;
- M(z) has a singular exponent $\lambda_M \leq 0$ or M(z) is analytic at ρ_H .

We can now state our multivariate result.

THEOREM 7.4. In a multivariate pure extended critical composition scheme (47), the joint moments of the random vector $\mathbf{X}_n = (X_{n,1}, \dots, X_{n,m})$ satisfy

$$\mathbb{E}(X_{n,1}^{s_1} \dots X_{n,m}^{s_m}) \sim \mu_{s_1,\dots,s_m} \prod_{\ell=1}^m n^{s_\ell \lambda_{H_\ell}} \kappa_\ell^{s_\ell},$$

with $\kappa_{\ell} = \frac{\tau_{H_{\ell}}}{-c_{H_{\ell}}}$ and $\mu_{s_1,...,s_m}$ given by

(48)
$$\mu_{s_1,\dots,s_m} = \frac{\Gamma\left(-\sum_{\ell=1}^m \lambda_{G_\ell} \lambda_{H_\ell} - \lambda_M\right)}{\Gamma\left(\sum_{\ell=1}^m s_\ell \lambda_{H_\ell} - \sum_{\ell=1}^m \lambda_{G_\ell} \lambda_{H_\ell} - \lambda_M\right)} \prod_{\ell=1}^m \frac{\Gamma(s_\ell - \lambda_{G_\ell})}{\Gamma(-\lambda_{G_\ell})}.$$

Consequently, one gets a convergence in distribution and in moments

$$\left(\frac{X_{n,1}}{\kappa_1 n^{\lambda_1}}, \dots, \frac{X_{n,m}}{\kappa_m n^{\lambda_m}}\right) \xrightarrow{d} \mathbf{X},$$

where **X** is determined by its joint moment sequence $\mu_s = \mu_{s_1,...,s_m}$. Moreover, the random vector **X** has a scaled Dirichlet-stable product distribution,

(49)
$$\mathbf{X} = (X_1, \dots, X_m) \stackrel{d}{=} (V_1 \cdot W_1^{\lambda_{H_1}}, \dots, V_m \cdot W_m^{\lambda_{H_m}}),$$

where $\mathbf{W} = (W_1, \dots, W_m, W_{m+1})$ follows a Dirichlet distribution

$$\mathbf{W} \stackrel{d}{=} \operatorname{Dir}(-\lambda_{G_1}\lambda_{H_1}, \dots, -\lambda_{G_m}\lambda_{H_m}, -\lambda_{M}),$$

and where the V_{ℓ} 's, for $1 \leq \ell \leq m$, are m independent two-parameter Mittag-Leffler distributions $V_{\ell} \stackrel{d}{=} \mathrm{ML}(\lambda_{H_{\ell}}, -\lambda_{G_{\ell}}\lambda_{H_{\ell}})$, also independent of \mathbf{W} .

PROOF. We proceed similarly to the proof of the first part of Theorem 4.1. First, the mixed factorial moments of X_n , which are obtained by differentiation and extraction of coefficients:

$$\mathbb{E}(X_{n,1}^{\underline{s_1}} \cdots X_{n,m}^{\underline{s_m}}) = \frac{[z^n] \partial_{u_1}^{s_1} \dots \partial_{u_m}^{s_m}(F)(z,1,\dots,1)}{[z^n] F(z,1,\dots,1)}.$$

The differentiation with respect to u_{ℓ} only affects the factor $G_{\ell}(u_{\ell}H_{\ell}(z))$, leading to a singular expansion covered in Section 3.1. Extraction of coefficients then gives an asymptotic expansion of the factorial moments. Converting all the factorial moments into moments using (25) gives the desired asymptotics and moments in (48).

It remains to identify the distribution. To this aim, note that a Dirichlet distributed random vector $(W_1,\ldots,W_{m+1})\stackrel{d}{=} \operatorname{Dir}(a_1,\ldots,a_{m+1})$ with positive parameters a_1,\ldots,a_{m+1} has a density function supported on the m simplex $\{(x_1,\ldots,x_{m+1})\in\mathbb{R}^{m+1}_{\geq 0}\mid\sum_{j=1}^{m+1}x_j=1\}$:

$$f(x_1,\ldots,x_{m+1}) = \frac{\Gamma(\sum_{j=1}^{m+1} a_j)}{\prod_{j=1}^{m+1} \Gamma(a_j)} \prod_{j=1}^{m+1} x_j^{a_j-1}.$$

П

Accordingly, the joint moments are given by

$$\mathbb{E}(W_1^{s_1}\cdots W_{m+1}^{s_{m+1}}) = \frac{\Gamma(\sum_{j=1}^{m+1} a_j)}{\Gamma(\sum_{j=1}^{m+1} (s_j + a_j))} \frac{\prod_{j=1}^{m+1} \Gamma(s_j + a_j)}{\prod_{j=1}^{m+1} \Gamma(a_j)}.$$

Now, consider a random vector (Z_1, \ldots, Z_m) satisfying

$$(Z_1,\ldots,Z_m) \stackrel{d}{=} (V_1 \cdot W_1^{\alpha_1},\ldots,V_m \cdot W_m^{\alpha_m})$$

with two-parameter Mittag-Leffler distributions $V_{\ell} \stackrel{d}{=} \mathrm{ML}(\alpha_{\ell}, a_{\ell})$ for $\ell = 1, \dots, m$, such that all random variables are mutually independent. Using the closed form (15) we get

$$\mathbb{E}\left(Z_1^{s_1}\cdots Z_m^{s_m}\right) = \frac{\Gamma\left(\sum_{\ell=1}^{m+1} a_\ell\right)}{\Gamma\left(a_{m+1} + \sum_{\ell=1}^{m} (\alpha_\ell s_\ell + a_\ell)\right)} \prod_{\ell=1}^m \frac{\Gamma\left(s_\ell + a_\ell/\alpha_\ell\right)}{\Gamma\left(a_\ell/\alpha_\ell\right)}.$$

Comparing this expression with the moments (48), the claim follows.

REMARK 7.5. The marginals X_{ℓ} of the random vector \mathbf{X} are also covered by Theorem 4.1. The random vector \mathbf{X} is closely related to Poisson–Dirichlet distributions PD (α, β) , [68, 69, 120] and the joint limit law of node degrees in preferential attachment trees or generalized plane-oriented recursive trees [106]; see also the subsequent example.

Now, if one considers the multivariate size-refined scheme

$$\mathcal{F} = \mathcal{M} \times \prod_{\ell=1}^{m} \mathcal{G}_{\ell} \big(\mathcal{H}_{\ell, \neq j_{\ell}} + v_{\ell} \mathcal{H}_{\ell, = j_{\ell}} \big),$$

one gets the following multivariate version of Theorem 5.1.

THEOREM 7.6 (Multivariate pure size-refined critical scheme). *In a multivariate pure size-refined critical composition scheme*

$$F(z, v_1, \dots, v_m) = M(z) \cdot \prod_{\ell=1}^m G_{\ell} (H_{\ell}(z) - (1 - v_{\ell}) h_{\ell, j_{\ell}} z^{j_{\ell}})$$

the random variables X_{n,ℓ,j_ℓ} , which count the number of H_ℓ -components of size j_ℓ , have joint factorial moments of mixed Poisson type:

$$\mathbb{E}(X_{n,1,j_1}^{\underline{s_1}} \dots X_{n,m,j_m}^{\underline{s_m}}) = \mu_{s_1,\dots,s_m} \cdot \prod_{\ell=1}^m \xi_{n,\ell,j_\ell}^{s_\ell} \cdot (1 + o(1)),$$

with $\xi_{n,\ell,j_\ell} = \frac{\rho_{H_\ell}^{j_\ell}}{-c_{H_\ell}} h_{\ell,j_\ell} n^{\lambda_{H_\ell}}$ and joint mixing distribution $\mathbf{X} = (X_1, \dots, X_m)$ as in Equation (49).

Let X_{ℓ} , for $1 \leq \ell \leq m$, denote the marginal distribution of the ℓ th coordinate of $\mathbf{X} = (X_1, \dots, X_m)$. For $n \to \infty$, the limiting distributions of $X_{n,\ell,j_{\ell}}$ jointly undergo mixed Poisson type phase transitions with mixing distributions X_{ℓ} . The phase transitions depend on the growth of $j_{\ell} = j_{\ell}(n)$, with critical growth ranges given by $j_{\ell} = j_{\ell}(n) = \Theta(n^{\frac{\lambda_{H_{\ell}}}{1 + \lambda_{H_{\ell}}}})$.

In particular, for $j_{\ell}(n) \sim \xi_{\ell} \cdot n^{\frac{\lambda_{H_{\ell}}}{1+\lambda_{H_{\ell}}}}$, the random vector $\mathbf{X}_{n,\mathbf{j}} = (X_{n,1,j_1}, \dots, X_{n,m,j_m})$ converges in distribution with convergence of all (factorial) moments to a multivariate distribution $MPo(\boldsymbol{\xi}\mathbf{X})$.

For properties of multivariate mixed Poisson distributions we refer to [36] or [90]. The key tool for proving Theorem 7.6 is the following multivariate extension of Lemma 5.4.

LEMMA 7.7 (Joint factorial moments and limit laws of mixed Poisson type). Let $(\mathbf{X}_n)_{n \in \mathbb{N}}$ denote a sequence of m-dimensional random vectors, whose factorial moments are asymptotically of mixed Poisson type, i.e., they satisfy for $n \to \infty$ the asymptotic expansion

$$\mathbb{E}(\mathbf{X}_{\overline{n}}^{\underline{s}}) = \mathbb{E}\left(X_{\overline{n},1}^{\underline{s_1}} \dots X_{\overline{n},m}^{\underline{s_m}}\right) = \mu_{s_1,\dots,s_m} \cdot \prod_{\ell=1}^m \xi_{n,\ell}^{s_\ell} \cdot (1 + o(1)), \quad s \ge 1,$$

with $\mu_{s_1,...,s_m} \ge 0$, and $\xi_{n,\ell} > 0$ for $1 \le \ell \le m$. Furthermore, assume that the sequence of joint moments $(\mu_{\mathbf{s}})_{\mathbf{s} \in \mathbb{N}^m}$ determines a unique distribution $\mathbf{L} = (L_1, \ldots, L_m)$. Then, for $n \to \infty$, one has the following joint limit distributions:

- (i) If $\xi_{n,\ell} \to \infty$, the random variable $\frac{X_{n,\ell}}{\xi_{n,\ell}}$ converges in distribution, with convergence of all moments, to L_{ℓ} .
- (ii) If $\xi_{n,\ell} \to \xi \in (0,\infty)$, the random variable $X_{n,\ell}$ converges in distribution, with convergence of all moments, to $Y \stackrel{d}{=} MPo(\xi L_{\ell})$.
- (iii) If $\xi_{n,\ell} \to 0$, $X_{n,\ell}$ converges to a Dirac distribution: $X_{n,\ell} \stackrel{d}{\to} 0$.

PROOF. We follow the proof of the univariate case [90, Lemma 2]. Assume that the set $\{1,\ldots,m\}$ decomposes into two disjoint sets C and D such that for indices $k \in C$ we have $\lambda_{n,k} \to \infty$, whereas for an index $k \in D$ it holds $\lambda_{n,k} \to \rho_k$ for $n \to \infty$. We observe that the mixture of joint raw moments and joint factorial moments converge:

$$\mathbb{E}\left(\left(\prod_{k\in C}\frac{X_{n,k}^{s_k}}{\lambda_{n,k}^{s_k}}\right)\cdot\left(\prod_{k\in D}X_{n,k}^{\underline{s_k}}\right)\right)\to\mu_{s_1,\dots,s_m}\cdot\prod_{k\in D}\rho_k^{s_k}.$$

The latter joint moment sequence, both raw and factorial moments, is exactly the joint moment sequence of a random vector $\mathbf{Z} = (Z_1, \dots, Z_m)$, with

$$\forall k \in C \colon Z_k \stackrel{d}{=} L_k, \quad \forall k \in D \colon Z_k \stackrel{d}{=} \mathsf{MPo}(\rho_k L_k),$$

such that

$$\mathbb{E}\left(\left(\prod_{k\in C} Z_k^{s_k}\right) \cdot \left(\prod_{k\in D} Z_k^{s_k}\right)\right) = \mu_{s_1,\dots,s_m} \cdot \prod_{k\in D} \rho_k^{s_k}.$$

PROOF OF THEOREM 7.6. The proof is very similar to the proofs of Theorems 5.1 and 7.4, so we will be brief again. The mixed factorial moments of $\mathbf{X}_{n,j_{\ell}}$ are obtained by differentiation and extraction of coefficients:

$$\mathbb{E}\left(X_{n,1,j_1}^{\underline{s_1}}\cdots X_{n,m,j_m}^{\underline{s_m}}\right) = \frac{[z^n]\partial_{v_1}^{s_1}\dots\partial_{v_m}^{s_m}(F)(z,1,\dots,1)}{[z^n]F(z,1,\dots,1)}.$$

The differentiation with respect to v_{ℓ} only affects the factor

$$G_{\ell}(H_{\ell}(z)-(1-v_{\ell})h_{\ell,j_{\ell}}z^{j_{\ell}}),$$

leading to singular expansions covered in Section 3.1 and additional factors $h_{\ell,j_\ell}^{s_\ell}$, $1 \leq \ell \leq m$. The asymptotics of these factors for $j_\ell \to \infty$ are all governed by (33). The individual singular expansions are similar to (34). Thus, extracting the coefficient $z^{n-s_1j_1-\cdots-s_mj_m}$ from $M(z) \cdot \prod_{\ell=1}^m G_\ell^{(s_\ell)} \big(H_\ell(z)\big)$ leads to the asymptotic expansion of the joint factorial moments. Lemma 7.7 then yields the stated limit law.

We end this section with four examples covered by the multivariate scheme (47).

EXAMPLE 7.8 (m-bundled plane-oriented recursive trees). We have previously encountered 3-bundled trees in Section 6.2 in the framework of bilabelled trees. In the following we study ordinary increasing trees [18,29,76,83,95,111], where each node has only one label. As before, given a degree-weight sequence $(\varphi_j)_{j\geq 0}$, the corresponding degree-weight generating function is defined by

$$\varphi(t) = \sum_{j \ge 0} \varphi_j t^j.$$

The associated family \mathcal{T} of increasing trees can be described by the following symbolic equation using the boxed product (see Section 6.2):

$$\mathcal{T} = \mathcal{Z}^{\square} * \varphi(\mathcal{T}).$$

For the exponential generating function

$$T(z) = \sum_{n>1} T_n \frac{z^n}{n!},$$

we thus have the following equation

$$T'(z) = \varphi(T(z)), \quad T(0) = 0.$$

We are interested in families of generalized plane-oriented recursive trees with degree-weight generating function $\varphi(t) = 1/(1-t)^m$, with $m \in \mathbb{N}$.

For m=1 we get the ordinary plane-oriented recursive tree, whereas for m>1 we get the so-called m-bundled trees, with generating function

$$T(z) = 1 - (1 - (m+1)z)^{1/(m+1)}$$
.

One may think of each node holding m-1 additional separation bars [76], which can be regarded as a special edge type. Naturally, this refines the root degree X_n in a random tree of size n, since we may look at the number of nodes attached to the root in a specific cluster, induced by the m-1 bars:

$$X_n = \sum_{\ell=1}^m X_{n,\ell}.$$

By construction, the random variables $X_{n,\ell}$ are exchangeable, but not independent. Writing $\mathbf{u}^{\mathbf{X_n}} = u_1^{X_{n,1}} \cdots u_m^{X_{n,m}}$, the generating function

$$T(z, \mathbf{u}) = \sum_{n \ge 1} T_n \mathbb{E}(\mathbf{u}^{\mathbf{X}_n}) \frac{z^n}{n!}$$

of the random vector $\mathbf{X}_n = (X_{n,1}, \dots, X_{n,m})$ is given by

$$\frac{\partial}{\partial z}T(z,\mathbf{u}) = \prod_{\ell=1}^m \frac{1}{1 - u_\ell T(z)}.$$

This refinement is covered by the multivariate scheme (47) (with a shift due to the derivative: Theorem 7.4 thus gives here the asymptotic behaviour of $X_{n+1,\ell}$, which, of course, then gives the asymptotic behaviour of $X_{n,\ell}$). Similarly, one may study the outdegree of a node labelled j, as well as multiple nodes, leading to closely related generating functions [69, 83, 106].

EXAMPLE 7.9 (Bilabelled increasing trees and refined root degree). Continuing from Section 6.2, we can refine the root-degree X_n in 3-bundled bilabelled increasing trees of size 2n, such that $X_n = X_{n,1} + X_{n,2} + X_{n,3}$, where $X_{n,\ell}$ is the root-degree in the ℓ th bundle. Note that the random variables $X_{n,\ell}$ are exchangeable. The corresponding generating function

$$T(z, u_1, u_2, u_3) = \sum_{n \ge 1} T_n \mathbb{E}(u_1^{X_{n,1}} u_2^{X_{n,2}} u_3^{X_{n,3}}) \frac{z^n}{n!}$$

then satisfies

$$\frac{\partial^2}{\partial z^2} T(z, u_1, u_2, u_3) = \prod_{\ell=1}^3 \frac{1}{1 - u_\ell (1 - \sqrt{1 - z^2})}.$$

Except for the non-standard shift in the random variable, the problem corresponds directly to a multivariate pure critical composition scheme.

EXAMPLE 7.10 (Returns in coloured bridges and walks). Consider k-coloured bridges B_k as defined in Section 6.4: A bridge is coloured in exactly k colours, where each colour consists of a non-empty bridge. Combinatorially, we append k non-empty bridges one after the other. We are interested in the individual number of returns to zero in each of the first k_1 bridges, followed by k_2 additional bridges, such that $k_1 + k_2 = k$. Reusing the combinatorial construction (43), the multivariate generating function $B_k(z, \mathbf{u})$ satisfies

$$B_k(z, u_1, \dots, u_{k_1}) = \left(\prod_{j=1}^{k_1} \left(\frac{1}{1 - u_j \left(1 - \frac{1}{B(z)}\right)} - 1\right)\right) (B(z) - 1)^{k_2}.$$

By our previous reasoning we see that the corresponding random variables are exchangeable and the multivariate scheme directly applies. Moreover, we can also consider walks and a refined generating function $W_k(z, \mathbf{u})$ of k-coloured walks with the tail coloured in the same colour as the final bridge, keeping track of the individual returns to zero of the first k_1 bridges,

$$W_k(z, \mathbf{u}) = (1 + B_k(z, \mathbf{u})) \frac{W(z)}{B(z)}.$$

Again, the corresponding random variables are exchangeable and the multivariate scheme directly applies again.

EXAMPLE 7.11 (Triangular urn models and node degrees in generalized recursive trees). We discuss a specific balanced triangular urn model with m colours, which generalizes the case with 2 colours from Section 6.7. Our urn model is specified by the following $m \times m$ balanced ball replacement matrix M with $a_k, b_k \in \mathbb{N}$ and $a_k + b_k = \sigma$, for $1 \le k \le m$:

(50)
$$M = \begin{pmatrix} a_1 & 0 & \dots & 0 & b_1 \\ 0 & a_2 & \dots & 0 & b_2 \\ \vdots & \ddots & \ddots & & \vdots \\ 0 & \dots & 0 & a_{m-1} & b_{m-1} \\ 0 & 0 & \dots & 0 & a_m + b_m \end{pmatrix}.$$

We consider the random variables $A_{n,k}$ which count the number of balls of colour k after n draws. To analyse this urn model, it is natural to introduce the history generating function

$$F(z, x_1, \dots, x_m) = \sum_{n, k_1, \dots, k_m \ge 0} f_{n, k_1, \dots, k_m} x_1^{k_1} \dots x_m^{k_m} \frac{z^n}{n!},$$

where $f_{n,k_1,...,k_m}$ is equal to the number of transitions leading after n steps to a configuration with $k_1,...,k_m$ balls of colours 1,...,m, respectively. Thus, the joint distribution of the number of balls of each colour is given by

$$\mathbb{P}\{A_{n,1} = k_1, \dots, A_{n,m} = k_m\} = \frac{[z^n \, x_1^{k_1} \dots x_m^{k_m}] F(z, x_1, \dots, x_m)}{[z^n] F(z, 1, \dots, 1)}.$$

where F has, surprisingly, the following simple algebraic closed form.

PROPOSITION 7.12. The history generating function $F(z, x_1, ..., x_m)$, associated with the balanced triangular urn model with ball replacement matrix M in (50), is given by

(51)
$$F(z, x_1, \dots, x_m) = \prod_{k=1}^m X_k(z)^{A_{0,k}}$$

where

(52)
$$X_m(z) = x_m \cdot \left(1 - \sigma x_m^{\sigma} z\right)^{-1/\sigma}$$

and, for $1 \le k < m$,

(53)
$$X_k(z) = x_k \cdot \left(1 - x_k^{a_k} x_m^{-a_k} \left(1 - (1 - \sigma x_m^{\sigma} z)^{a_k/\sigma}\right)\right)^{-1/a_k}.$$

PROOF. We follow the history-counting approach introduced by Flajolet et al. in [39,40], which shows that the evolution of the urn is captured by the matricial equation $\partial_z \mathbf{X} = \mathbf{X}^{1+M}$, that is, one has the differential system

$$\dot{X}_k(z) = X_k(z)^{1+a_k} X_m^{b_k}(z) \ \ (\text{for } 1 \leq k < m) \qquad \text{ and }$$

$$\dot{X}_m(z) = X_m^{1+\sigma}(z),$$

with initial conditions $X_k(0) = x_k$. Now, by separation of variables, we obtain Formula (52). ¹⁵ Then, by integration, we get Formula (53). Finally, using the basic isomorphism between differential systems and history generating functions [39, Theorem 6], we obtain the desired closed form (51) for the generating function of urn histories.

As the urn is balanced, we add the same number of balls at each step. Therefore, knowing the number of balls of colour 1 to m-1 immediately gives the number of balls of colour m, so, without loss of information, we can stop tracking this parameter (i.e., we set $x_m=1$). One then recognizes that this urn model has a multivariate pure critical composition scheme (with $M(z)=(1-\sigma z)^{-1/\sigma}$, and for k< m, $G_k(z)=(1-z)^{-1/a_k}$, $H_k(z)=1-(1-\sigma z)^{a_k/\sigma}$, and $u_k=x_k^{a_k}$); thus Theorem 7.4 applies: The corresponding joint distribution (i.e., the *number of drawings* of balls of each colour) converges to a Dirichlet-stable product distribution.

Note that the node degrees in generalized plane-oriented recursive trees can be modelled by such urns [71, 106]. So, as an application, let us set $a_k = 1$ and $b_k = m-1$ for $1 \le k \le m$ and $A_{0,k} = 1$ (for $1 \le k < m$) and $A_{0,m} = 0$ as initial values. Then, the random variables $A_{n,1}$ up to $A_{n,m-1}$ are exchangeable and count the refined root degree of (m-1)-bundled plane-oriented recursive trees, i.e., the number of children of the root in each bundle.

Moreover, the urn model M can be used to study the joint degree distribution of the nodes labelled $1, 2, \ldots, m-1$ in generalized plane-oriented recursive trees, as well as to obtain second order asymptotics and central limit theorem analogues.

¹⁵There is a small sign typo in [39, page 94] where, in the corresponding result for the 2×2 model, $y_0^{-\sigma}$ should be replaced by y_0^{σ} .

8. Conclusion. In this article, we introduced the three-parameter Mittag-Leffler distribution and showed its universality for capturing transition phases related to critical compositions. It opens a new chapter in the long history of the Mittag-Leffler function. This special function, $E_{\alpha}(t) := \sum_{k \geq 0} \frac{1}{\Gamma(\alpha k + 1)} t^k$, was introduced by Mittag-Leffler in 1903 [102, 103]. It was generalized to a two-parameter function $E_{\alpha,\beta}(t) := \sum_{k \geq 0} \frac{1}{\Gamma(\alpha k + \beta)} t^k$ by Wiman in 1905 [135] and to a three-parameter function $E_{\alpha,\beta}^{\gamma}(t) := \sum_{k=0}^{\infty} \frac{\Gamma(k+\gamma)}{\Gamma(\alpha k + \beta)\Gamma(\gamma)} \frac{t^k}{k!}$ by Prabhakar in 1971 [122]. While these functions were used for decades by mathematicians and physicists working on differential equations (see Remark 3.13 on a link with electromagnetism), special functions, or fractional calculus, it is a nice surprise that this three-parameter Mittag-Leffler function is the key to unify different fundamental distributions in probability theory, as summarized in Table 4.

This work makes explicit the limit laws of structures associated with schemes like G(H(z))M(z), where we additionally mark either the number of \mathcal{H} -components, or the number of \mathcal{H} -components of a given size. We focused on the technically more delicate and mathematically richer case where G,H, and M are simultaneously singular. We proved that when these functions have algebraic dominant singularities with exponents between 0 and 1, the limit laws of the schemes are moment-tilted stable distributions and distributions involving Mittag-Leffler laws. In Table 5 we give an overview of the covered schemes and their limit laws, where we have convergence in distribution and convergence of all moments.

In the extended scheme, in which one follows the total number of \mathcal{H} -components, we proved the appearance of three régimes for the corresponding limit law (continuous, Boltzmann, or a linear combination of these two), depending on the relation of the singular exponents of G, H, and M; see Table 1 on page 4. In the size-refined scheme, in which one follows the number of \mathcal{H} -components of a given size, we proved the appearance of mixed Poisson distributions (see Definition 3.15) and a double phase transition for the limit law from continuous to discrete to degenerate, with explicit threshold sizes depending on the exponents; see Table 2 on page 6.

We also presented several extensions (logarithmic singularities, multivariate cases) and a variety of applications to important probabilistic and combinatorial objects. This allowed us to obtain new results for the core size of supertrees, the number of subtrees in different increasing trees, the returns to zero and sign changes in walks and bridges, the table sizes in the Chinese restaurant process, and the number of balls in some urn models.

$\mathbb{E}(e^{tX}) = E_{\alpha}(t)$	MGF appeared as Laplace transform of stable distributions and for the local time of Markov processes [25, 34]. MGF introduced by Pitman in 2002 [119] for the Chinese rectangent process and
	,
e^{tX}) = $\Gamma(\beta')E_{\alpha,\beta'}^{\gamma'}(t)$ $(\beta',\gamma') = (\beta,\frac{\beta}{\alpha})$	for the Chinese restaurant process, and used for a line-breaking construction of stable trees by Goldschmidt and Haas [52]; it also occurs for triangular Pólya urns (see Flajolet, Dumas, Puyhaubert [39] and Janson [72,74]).
e^{tX}) = $\Gamma(\beta')E_{\alpha,\beta'}^{\gamma'}(t)$ β',γ') = $(\beta+\gamma,\frac{\beta}{\alpha})$	MGF introduced in 2021 (in this article) for critical composition schemes.
	$(\beta', \gamma') = (\beta, \frac{\beta}{\alpha})$ $e^{tX}) = \Gamma(\beta') E_{\alpha, \beta'}^{\gamma'}(t)$

The successive generalizations of the Mittag-Leffler distribution, its moment generating function (MGF), its connection to the corresponding generalized Mittag-Leffler function, and some occurrences in probability theory.

The results of Table 5 hold for functions H(z) having a dominant singularity of algebraic type. Our methods can also deal with schemes having other types of singularities. For example, one could allow algebraic-logarithmic behaviours of the form

$$H(z) \sim \left(1 - \frac{z}{\rho_H}\right)^{\lambda_H} \left(\frac{1}{z} \log \left(1 - \frac{z}{\rho_H}\right)\right)^{\psi_H}.$$

Such algebraic-logarithmic schemes cover some instances of 3-colour balanced triangular urn models [39, 123], whose complete analysis, however, remains a challenge as other types of singularities appear, thus leading to new families of limit laws whose nature is unclear. Compare also with the related open problem by Janson [74], asking for a more detailed description of the limit law of unbalanced 2-colour triangular urn models.

To keep this article readable, we eluded the question of the speed of convergence of X_n , properly normalized, to its limit law X. Recently, a few articles addressed this for some preferential attachment rules [115] and for the Chinese restaurant process [28] (see Remark 6.16). In a future work we plan to give uniform bounds on the moments, leading to Berry-Esseenlike inequalities for the speed of convergence of all the limit laws associated with critical composition schemes; see also [2,62].

Building on the current work, we consider in [10] an extension of our results on critical schemes by imposing a Gibbs measure on the parameter of the combinatorial structure: that is, instead of the uniform distribution model imposed by Formula (3), we consider the model $\mathbb{P}\{X_n=k\}=\frac{f_{n,k}q^k}{f_n(q)}$. This leads to universal phase transitions where the three-parameter Mittag-Leffler distribution strikes again. (For q=1, this gives back the uniform distribution model.)

Last but not least, in our companion article [11], we further extend our analysis to schemes with algebraic singularities for $\lambda_H > 1$: We analyse a generalization of the composition scheme from [7], leading to stable laws (e.g., the map-Airy distribution), Gaussian laws, as well as bimodal distributions. For extended composition schemes we observe a new behaviour, obtaining for example beta limit laws. In size-refined schemes we anticipate further continuous-to-discrete phase transitions. This will achieve the exhaustive exploration of the landscape of critical composition schemes with algebraic singularities and their phase transitions.

Composition scheme	Symbolic form	Limit law	Thm.
Ordinary	F(z,u) = G(uH(z))	generalized Mittag-Leffler	4.2
Extended	$F(z,u)=M(z)G\bigl(uH(z)\bigr)$	three-parameter Mittag-Leffler and Boltzmann distribution	4.1 4.3 4.4
Cyclic	$F(z, u) = -\log(1 - uH(z))$	Mittag-Leffler	7.1
Multivariate extended	$F(z, \mathbf{u}) = M(z) \prod_{\ell=1}^{m} G_{\ell} \left(u_{\ell} H_{\ell}(z) \right)$	multivariate product distribution	7.4
Size-refined	$F(z,v) = M(z)G(H(z) - z^{j}h_{j}(1-v))$	mixed Poisson type phase transition	5.1
Size-refined cyclic	$F(z,v) = -\log\left(1 - (H(z) - (1-v)h_jz^j/j!)\right)$	mixed Poisson type phase transition	7.3
Multivariate size-refined	$F(z, \mathbf{v}) = M(z) \prod_{\ell=1}^{m} G_{\ell} (H_{\ell}(z) - z^{j_{\ell}} h_{\ell, j_{\ell}} (1 - v_{\ell}))$	mv. mixed Poisson type phase transition	7.6

TABLE 5

Overview of our results on critical composition schemes, where $\mathbf{u} = (u_1, \dots, u_m)$ and $\mathbf{v} = (v_1, \dots, v_m)$.

Acknowledgements. We thank the two referees for their careful reading and their feedback. We also thank Katarzyna Górska for pointing out that the moment generating function of our three-parameter Mittag-Leffler distribution is called the Prabhakar function in physics. ¹⁶

Last but not least, we are pleased to dedicate this article to our colleague and mentor Alois Panholzer, on the occasion of his 50th birthday. In fact, Alois played a central rôle in the birth of this article in several ways:

- As a byproduct of Alois' analyses of combinatorial structures and stochastic processes (such as permutations, parking functions, lattice paths, urns, and trees), several phase transitions involving the mixed Poisson distribution were uncovered [90,112]. This guided the second author to study a general framework for these phase transitions, and to join forces with the other authors to extend the results of [7,131] involving Rayleigh, half-normal, and stable distributions. This led us to the composition schemes analysed in this work.
- Alois also connected the first and second authors via a French-Austrian international project (PHC Amadeus) in 2005/06, perpetuating a long tradition of exchanges in combinatorics between Austria and France. A part of this project led to the article [9], which established Gaussian limit laws for some composition schemes related to trees.
- Alois was also a constant source of inspiration and guidance for the second author. Without him, we would have never started this work.

Funding. Michael Wallner was funded by the FWF Erwin Schrödinger Fellowship J 4162 and the FWF Stand-alone Project P 34142. The authors also thank the French-Austrian PHC Amadeus project "Asymptotic behaviour of combinatorial structures" (OeAD WTZ project FR 01/2023), which funded mutual visits.

REFERENCES

- [1] Louigi Addario-Berry. A probabilistic approach to block sizes in random maps. ALEA, Lat. Am. J. Probab. Math. Stat., 16(1):1–13, 2019.
- [2] José A. Adell and Jesús de la Cal. On the uniform convergence of normalized Poisson mixtures to their mixing distribution. Statistics & Probability Letters, 18:227–232, 1993.
- [3] David J. Aldous. *Exchangeability and Related Topics*, volume 1117 of *Lecture Notes in Math.* Springer, 1985.
- [4] Richard Arratia, Andrew D. Barbour, and Simon Tavaré. *Logarithmic Combinatorial Structures: a Probabilistic Approach*. EMS Monographs in Mathematics. European Mathematical Society, 2003.
- [5] Cyril Banderier and Michael Drmota. Formulae and asymptotics for coefficients of algebraic functions. *Combin. Probab. Comput.*, 24(1):1–53, 2015.
- [6] Cyril Banderier and Philippe Flajolet. Basic analytic combinatorics of directed lattice paths. *Theoret. Comput. Sci.*, 281(1-2):37–80, 2002.
- [7] Cyril Banderier, Philippe Flajolet, Gilles Schaeffer, and Michèle Soria. Random maps, coalescing saddles, singularity analysis, and Airy phenomena. *Random Struct. Algorithms*, 19(3-4):194–246, 2001.
- [8] Cyril Banderier and Paweł Hitczenko. Enumeration and asymptotics of restricted compositions having the same number of parts. *Discrete Appl. Math.*, 160(18):2542–2554, 2012.
- [9] Cyril Banderier, Markus Kuba, and Alois Panholzer. Analysis of three graph parameters for random trees. *Random Struct. Algorithms*, 35(1):42–69, 2009.
- [10] Cyril Banderier, Markus Kuba, Stephan Wagner, and Michael Wallner. Composition schemes: q-enumerations and phase transitions in Gibbs models. In 35rd International Conference on Probabilistic, Combinatorial and Asymptotic Methods for the Analysis of Algorithms (AofA 2024), volume 302 of Leibniz International Proceedings in Informatics (LIPIcs), pages 23:1–23:17, 2024.
- [11] Cyril Banderier, Markus Kuba, and Michael Wallner. Phase transitions of composition schemes II: stable laws and Boltzmann distributions. *In preparation*, 2024.

¹⁶A recording of this surprising feedback is even available in our IHÉS talk, at the conference *Combinatorics* and *Arithmetic for Physics: Special Days 2022*. We thank Gérard Duchamp, Maxim Kontsevich, Gleb Koshevoy, Sergei Nechaev, and Karol Penson, who had the great idea to organize this interdisciplinary event!

- [12] Cyril Banderier, Philippe Marchal, and Michael Wallner. Periodic Pólya urns, the density method and asymptotics of Young tableaux. Ann. Probab., 48(4):1921–1965, 2020.
- [13] Cyril Banderier and Michael Wallner. Some reflections on directed lattice paths. In *Proceedings of the 25th International Conference on Probabilistic, Combinatorial and Asymptotic Methods for the Analysis of Algorithms (AofA'14)*, pages 25–36. Discrete Math. Theor. Comput. Sci., 2014.
- [14] Cyril Banderier and Michael Wallner. Lattice paths with catastrophes. Discrete Math. Theor. Comput. Sci., Vol. 19 no. 1, 2017.
- [15] Cyril Banderier and Michael Wallner. Local time for lattice paths and the associated limit laws. In Proceedings of GASCom 2018, volume 2113, pages 69–78. CEUR Workshop Proceedings, 2018.
- [16] Cyril Banderier and Michael Wallner. The kernel method for lattice paths below a rational slope. In *Lattice Path Combinatorics and Applications*, volume 58 of *Developments in Mathematics Series*, pages 119–154. Springer, 2019.
- [17] Edward A. Bender. Central and local limit theorems applied to asymptotic enumeration. *J. Comb. Theory*, *Ser. A*, 15:91–111, 1973.
- [18] François Bergeron, Philippe Flajolet, and Bruno Salvy. Varieties of increasing trees. In *Colloque sur les arbres en algèbre et programmation (CAAP'92)*, volume 581 of *Lect. Notes Comput. Science*, pages 24–48. Springer, 1992.
- [19] Mira Bernstein and Neil J. A. Sloane. Some canonical sequences of integers. Linear Algebra Appl., 226-228:57–72, 1995.
- [20] Olivier Bodini, Matthieu Dien, Xavier Fontaine, Antoine Genitrini, and Hsien-Kuei Hwang. Increasing diamonds. In LATIN 2016, volume 9644 of Lect. Notes Comput. Science, pages 207–219. Springer, 2016.
- [21] Olivier Bodini, Olivier Roussel, and Michèle Soria. Boltzmann samplers for first-order differential specifications. *Discrete Appl. Math.*, 160(18):2563–2572, 2012.
- [22] Mireille Bousquet-Mélou. Limit laws for embedded trees: applications to the integrated superBrownian excursion. *Random Struct. Algorithms*, 29(4):475–523, 2006.
- [23] Edmundo Capelas de Oliveira, Francesco Mainardi, and Jayme Jr. Vaz. Models based on Mittag-Leffler functions for anomalous relaxation in dielectrics. Eur. Phys. J., Special Topics, 193:161–171, 2011.
- [24] Torsten Carleman. Sur les équations intégrales singulières à noyau réel et symétrique. Uppsala Universitets Årsskrift, 1923.
- [25] Donald A. Darling and Mark Kac. On occupation times for Markoff processes. Trans. Amer. Math. Soc., 84:444–458, 1957.
- [26] Timothy DeVries. Algorithms for Bivariate Singularity Analysis. PhD thesis, Univ. of Pennsylvania, 2011.
- [27] NIST Digital Library of Mathematical Functions. http://dlmf.nist.gov/, release 1.1.3 of 2021-09-15. F. W. J. Olver, A. B. Olde Daalhuis, D. W. Lozier, B. I. Schneider, R. F. Boisvert, C. W. Clark, B. R. Miller, B. V. Saunders, H. S. Cohl, and M. A. McClain, eds.
- [28] Emanuele Dolera and Stefano Favaro. A Berry–Esseen theorem for Pitman's α -diversity. Ann. Appl. Probab., 30(2):847–869, 2020.
- [29] Michael Drmota. Random Trees. Springer, 2009.
- [30] Michael Drmota and Michèle Soria. Marking in combinatorial constructions: generating functions and limiting distributions. *Theoret. Comput. Sci.*, 144(1-2):67–99, 1995.
- [31] Michael Drmota and Michèle Soria. Images and preimages in random mappings. SIAM J. Discrete Math., 10(2):246–269, 1997.
- [32] Jules Dubourdieu. Les principes fondamentaux du calcul des probabilités et la théorie de l'assurance-maladie. Gauthier-Villars, 1939.
- [33] Philippe Duchon, Philippe Flajolet, Guy Louchard, and Gilles Schaeffer. Boltzmann samplers for the random generation of combinatorial structures. *Combin. Probab. Comput.*, 13(4-5):577–625, 2004.
- [34] William Feller. Fluctuation theory of recurrent events. Trans. Amer. Math. Soc., 67:98–119, 1949.
- [35] William Feller. An Introduction to Probability Theory and its Applications. Vol. II. Second edition (1st ed.: 1966). John Wiley & Sons, 1971.
- [36] Anthony Ferrari, Gérard Letac, and Jean-Yves Tourneret. Multivariate mixed Poisson distributions. In *12th European Signal Processing Conference, IEEE*, pages 1067–1070, 2004.
- [37] James Allen Fill, Philippe Flajolet, and Nevin Kapur. Singularity analysis, Hadamard products, and tree recurrences. J. Comput. Appl. Math., 174(2):271–313, 2005.
- [38] Philippe Flajolet. Some exactly solvable models of urn process theory. Slides for the conference Mathinfo in Nancy, 2006.

- [39] Philippe Flajolet, Philippe Dumas, and Vincent Puyhaubert. Some exactly solvable models of urn process theory. In Fourth Colloquium on Mathematics and Computer Science Algorithms, Trees, Combinatorics and Probabilities, pages 59–118. Discrete Math. Theor. Comput. Sci., 2006.
- [40] Philippe Flajolet, Joaquim Gabarró, and Helmut Pekari. Analytic urns. Ann. Probab., 33(3):1200–1233, 2005.
- [41] Philippe Flajolet and Andrew Odlyzko. Singularity analysis of generating functions. SIAM J. Discrete Math., 3(2):216–240, 1990.
- [42] Philippe Flajolet and Robert Sedgewick. Analytic Combinatorics. Cambridge University Press, 2009.
- [43] Philippe Flajolet and Michèle Soria. Gaussian limiting distributions for the number of components in combinatorial structures. *J. Combin. Theory Ser. A*, 53(2):165–182, 1990.
- [44] Philippe Flajolet and Michèle Soria. General combinatorial schemas: Gaussian limit distributions and exponential tails. *Discrete Math.*, 114(1-3):159–180, 1993.
- [45] Maurice Fréchet and James Shohat. A proof of the generalized second limit theorem in the theory of probability. *Trans. Amer. Math. Soc.*, 33(2):533–543, 1931.
- [46] Michael Fuchs. Subtree sizes in recursive trees and binary search trees: Berry–Esseen bounds and Poisson approximations. Combin. Probab. Comput., 17(5):661–680, 2008.
- [47] Michael Fuchs. Limit theorems for subtree size profiles of increasing trees. *Combin. Probab. Comput.*, 21(3):412–441, 2012.
- [48] Michael Fuchs, Hsien-Kuei Hwang, and Ralph Neininger. Profiles of random trees: Limit theorems for random recursive trees and binary search trees. *Algorithmica*, 46(3-4):367–407, 2006.
- [49] Roberto Garra and Roberto Garrappa. The Prabhakar or three parameter Mittag-Leffler function: theory and application. Commun. Nonlinear Sci. Numer. Simul., 56:314–329, 2018.
- [50] Omer Giménez and Marc Noy. Asymptotic enumeration and limit laws of planar graphs. *J. Am. Math. Soc.*, 22(2):309–329, 2009.
- [51] Omer Giménez, Marc Noy, and Juanjo Rué. Graph classes with given 3-connected components: asymptotic enumeration and random graphs. *Random Struct. Algorithms*, 42(4):438–479, 2013.
- [52] Christina Goldschmidt and Bénédicte Haas. A line-breaking construction of the stable trees. Electron. J. Probab., 20:24, 2015. Id/No 16.
- [53] Christina Goldschmidt, Bénédicte Haas, and Delphin Sénizergues. Stable graphs: distributions and line-breaking construction. Ann. H. Lebesgue, 5:841–904, 2022.
- [54] Rudolf Gorenflo, Anatoly A. Kilbas, Francesco Mainardi, and Sergei V. Rogosin. *Mittag-Leffler functions*, related topics and applications. Springer Monogr. Math. Berlin: Springer, 2014.
- [55] Katarzyna Górska, Andrzej Horzela, Łukasz Bratek, Giuseppe Dattoli, and Karol Penson. The Havriliak—Negami relaxation and its relatives: the response, relaxation and probability density functions. *J. Phys. A, Math. Theor.*, 51(13):15, 2018. Id/No 135202.
- [56] Katarzyna Górska, Andrzej Horzela, Ambra Lattanzi, and Tibor K. Pogány. On complete monotonicity of three parameter Mittag-Leffler function. Appl. Anal. Discrete Math., 15(1):118–128, 2021.
- [57] Raúl Gouet. Martingale Functional Central Limit Theorems for a Generalized Pólya Urn. Ann. Probab., 21:1624–1639, 1993.
- [58] Xavier Gourdon. Largest component in random combinatorial structures. Discrete Math., 180(1-3):185–209, 1998
- [59] Jan Grandell. Mixed Poisson Processes. Chapman & Hall, 1997.
- [60] Theodore Edward Harris. *The Theory of Branching Processes*. Springer, 1963.
- [61] Pierre Humbert. Nouvelles correspondances symboliques. Bull. Sci. Math., II. Sér., 69:121–129, 1945.
- [62] Hsien-Kuei Hwang. Second phase changes in random *m*-ary search trees and generalized quicksort: convergence rates. *Ann. Probab.*, 31(2):609–629, 1998.
- [63] Hsien-Kuei Hwang. Asymptotics of Poisson approximation to random discrete distributions: an analytic approach. Adv. in Appl. Probab., 31(2):448–491, 1999.
- [64] Hsien-Kuei Hwang. Phase changes in random recursive structures and algorithms. In Proceedings of the Workshop on Probability with Applications to Finance and Insurance, pages 82–97. World Scientific, 2004.
- [65] Hsien-Kuei Hwang and Vytas Zacharovas. Uniform asymptotics of Poisson approximation to the Poissonbinomial distribution. Theory Probab. Appl., 55(2):198–224, 2011.
- [66] Ildar Abdulovich Ibragimov and Kalman Eremeevich Chernin. On the unimodality of stable laws. Theory Probab. Appl., 4:417–419, 1959.
- [67] Lancelot Fitzgerald James. Lamperti-type laws. Ann. Appl. Probab., 20(4):1303–1340, 2010.
- [68] Lancelot Fitzgerald James. Stick-breaking $PG(\alpha, \zeta)$ -generalized gamma processes. arXiv, 2013.

- [69] Lancelot Fitzgerald James. Generalized Mittag Leffler distributions arising as limits in preferential attachment models. arXiv, 2015.
- [70] Svante Janson. Functional limit theorems for multitype branching processes and generalized Pólya urns. *Stochastic Process. Appl.*, 110(2):177–245, 2004.
- [71] Svante Janson. Asymptotic degree distribution in random recursive trees. *Random Struct. Algorithms*, 26(1-2):69–83, 2005.
- [72] Svante Janson. Limit theorems for triangular urn schemes. Probab. Theory Related Fields, 134(3):417–452, 2006.
- [73] Svante Janson. Random cutting and records in deterministic and random trees. *Random Struct. Algorithms*, 29:139–179, 2006.
- [74] Svante Janson. Moments of Gamma type and the Brownian supremum process area. *Probab. Surv.*, 7:1–52, 2010. With addendum on pages 207–208.
- [75] Svante Janson, Donald E. Knuth, Tomasz Łuczak, and Boris Pittel. The birth of the giant component. *Random Struct. Algorithms*, 4(3):233–358, 1993.
- [76] Svante Janson, Markus Kuba, and Alois Panholzer. Generalized Stirling permutations, families of increasing trees and urn models. J. Combin. Theory Ser. A, 118(1):94–114, 2011.
- [77] Svante Janson and Nicolas Pouyanne. Moment convergence of balanced Pólya processes. *Electron. J. Probab.*, 23:13, 2018. Id/No 34.
- [78] Dimitris Karlis. EM algorithm for mixed Poisson and other discrete distributions. Astin Bull., 35(1):3–24, 2005.
- [79] Donald E. Knuth. The Art of Computer Programming. Vol. 3: Sorting and searching. Addison-Wesley, 1998. Second edition (1st ed: 1973).
- [80] Valentin F. Kolchin. *Random Graphs*, volume 53. Cambridge University Press, 1999.
- [81] Markus Kuba and Alois Panholzer. Analysis of label-based parameters in increasing trees. In Fourth Colloquium on Mathematics and Computer Science Algorithms, Trees, Combinatorics and Probabilities, pages 321–330. Discrete Math. Theor. Comput. Sci., 2006.
- [82] Markus Kuba and Alois Panholzer. Descendants in increasing trees. Electron. J. Combin., 13 (1)(Paper 8), 2006
- [83] Markus Kuba and Alois Panholzer. On the degree distribution of the nodes in increasing trees. J. Combin. Theory Ser. A, 114(4):597–618, 2007.
- [84] Markus Kuba and Alois Panholzer. A combinatorial approach to the analysis of bucket recursive trees. *Theoret. Comput. Sci.*, 411(34-36):3255–3273, 2010.
- [85] Markus Kuba and Alois Panholzer. Analysis of statistics for generalized Stirling permutations. Discrete Math., 20:875–910, 2011.
- [86] Markus Kuba and Alois Panholzer. Bilabelled increasing trees and hook-length formulae. *European J. Combin.*, 33(2):248–258, 2012.
- [87] Markus Kuba and Alois Panholzer. Limiting distributions for a class of diminishing urn models. *Adv. in Appl. Probab.*, 44(4):1–31, 2012.
- [88] Markus Kuba and Alois Panholzer. On death processes and urn models. In *Proceedings of the 23rd International Conference on Probabilistic, Combinatorial, and Asymptotic Methods in the Analysis of Algorithms (AofA'12)*, pages 29–42. Discrete Math. Theor. Comput. Sci., 2012.
- [89] Markus Kuba and Alois Panholzer. Combinatorial families of multilabelled increasing trees and hook-length formulas. Discrete Math., 339(1):227–254, 2016.
- [90] Markus Kuba and Alois Panholzer. On moment sequences and mixed Poisson distributions. Probab. Surv., 13:89–155, 2016.
- [91] Markus Kuba and Alois Panholzer. On bucket increasing trees, clustered increasing trees and increasing diamonds. *Combin. Probab. Comput.*, 31(4):629–661, 2022.
- [92] Markus Kuba and Henning Sulzbach. On martingale tail sums in affine two-color urn models with multiple drawings. *J. Appl. Probab.*, 54:1–21, 2017.
- [93] Michel Loève. Probability Theory. Vol. I. Fourth edition (1st ed.: 1955). Springer, 1977.
- [94] Hosam M. Mahmoud. *Pólya Urn Models*. CRC Press, 2009.
- [95] Hosam M. Mahmoud, Robert Thomas Smythe, and Jerzy Szymański. On the structure of random plane-oriented recursive trees and their branches. *Random Struct. Algorithms*, 4(2):151–176, 1993.
- [96] Philippe Marchal. Constructing a sequence of random walks strongly converging to Brownian motion. In *Proceedings of Discrete Random Walks (DRW'03)*, pages 181–190. Discr. Math. Theor. Comput. Sci., 2003.

- [97] Philippe Marchal. A note on the fragmentation of a stable tree. In *Proceedings of the Fifth Colloquium on Mathematics and Computer Science*, pages 489–500. Discrete Math. Theor. Comput. Sci., 2008.
- [98] Jean-François Marckert. Growing random uniform d-ary trees. Ann. Comb., 27:51–66, 2023.
- [99] Jean-Claude Massé and Radu Theodorescu. Neyman type A distribution revisited. *Statist. Neerlandica*, 59:206–213, 2005.
- [100] Amram Meir and John W. Moon. Cutting down random trees. J. Austral. Math. Soc., 11:313–324, 1970.
- [101] Grégory Miermont. Random maps and their scaling limits. In Fractal Geometry and Stochastics IV. 4th conference, Greifswald, Germany, September 8–12, 2008, pages 197–224. Birkhäuser, 2009.
- [102] Gösta Mittag-Leffler. Sur la nouvelle fonction $E_{\alpha}(x)$. C. R. Acad. Sci., Paris, 137:554–558, 1903.
- [103] Gösta Mittag-Leffler. Une généralisation de l'intégrale de Laplace-Abel. C. R. Acad. Sci., Paris, 136:537–539, 1903.
- [104] Martin Möhle. A restaurant process with cocktail bar and relations to the three-parameter Mittag-Leffler distribution. J. Appl. Probab., 58(4):978–1006, 2021.
- [105] Rémi Monasson. A generating function method for the average-case analysis of DPLL. In Approximation, randomization and combinatorial optimization, volume 3624 of Lecture Notes in Comput. Sci., pages 402–413. Springer, 2005.
- [106] Tamás F. Móri. The maximum degree of the Barabási–Albert random tree. *Combin. Probab. Comput.*, 14(3):339–348, 2005.
- [107] Jerzy Neyman. On a new class of "contagious" distributions applicable in entomology and bacteriology. *Ann. Math. Stat.*, 10:35–57, 1939.
- [108] Konstantinos Panagiotou, Leon Ramzews, and Benedikt Stufler. Exact-size sampling of enriched trees in linear time. SIAM J. Comput., 52(5):1097–1131, 2023.
- [109] Alois Panholzer. Non-crossing trees revisited: cutting down and spanning subtrees. In *Proceedings of Discrete Random Walks (DRW'03)*, pages 265–276. Discrete Math. Theor. Comput. Sci., 2003.
- [110] Alois Panholzer. Cutting down very simple trees. Quaest. Math., 29(2):211–227, 2006.
- [111] Alois Panholzer and Helmut Prodinger. Level of nodes in increasing trees revisited. *Random Struct. Algorithms*, 31(2):203–226, 2007.
- [112] Alois Panholzer and Georg Seitz. Limiting distributions for the number of inversions in labelled tree families. *Ann. Comb.*, 16(4):847–870, 2012.
- [113] Alois Panholzer and Georg Seitz. Ancestors and descendants in evolving *k*-tree models. *Random Struct. Algorithms*, 44:465–489, 2014.
- [114] Erol A. Peköz, Adrian Röllin, and Nathan Ross. Degree asymptotics with rates for preferential attachment random graphs. Ann. Appl. Probab., 23(3):1188–1218, 2013.
- [115] Erol A. Peköz, Adrian Röllin, and Nathan Ross. Generalized gamma approximation with rates for urns, walks and trees. *Ann. Probab.*, 44(3):1776–1816, 2016.
- [116] Robin Pemantle and Mark C. Wilson. Analytic Combinatorics in Several Variables. Cambridge University Press, 2013.
- [117] Raman N. Pillai. On Mittag-Leffler functions and related distributions. Ann. Inst. Statist. Math., 42(1):157–161, 1990.
- [118] Jim Pitman. Exchangeable and partially exchangeable random partitions. Probab. Theory Relat. Fields, 102(2):145–158, 1995.
- [119] Jim Pitman. Combinatorial Stochastic Processes, volume 1875 of Lecture Notes in Math. Springer, 2006. Lectures from the 32nd Summer School on Probability Theory held in Saint-Flour, July 2002.
- [120] Jim Pitman and Marc Yor. The two-parameter Poisson-Dirichlet distribution derived from a stable subordinator. Ann. Probab., 25(2):855–900, 1997.
- [121] Harry Pollard. The representation of $e^{-x^{\lambda}}$ as a Laplace integral. Bull. Amer. Math. Soc., 52:908–910, 1946.
- [122] Tilak Raj Prabhakar. A singular integral equation with a generalized Mittag Leffler function in the kernel. Yokohama Math. J., 19:7–15, 1971.
- [123] Vincent Puyhaubert. *Modèles d'urnes et phénomènes de seuils en combinatoire analytique*. PhD thesis, École polytechnique, Palaiseau, 2005.
- [124] Uwe Rösler and Ludger Rüschendorf. The contraction method for recursive algorithms. *Algorithmica*, 29(1-2):3–33, 2001.
- [125] Nomvelo Karabo Sibisi. Convolutions and mixtures of gamma, stable and mittag-leffler distributions. *arXiv*, 2024.
- [126] John G. Skellam and Leonard R. Shenton. Distributions associated with random walk and recurrent events. *J. Roy. Statist. Soc. Ser. B.*, 19:64–111 (and discussion pp. 111–118), 1957.

- [127] Andrea Sportiello. Boltzmann sampling of irreducible context-free structures in linear time. arXiv, 2021.
- [128] Benedikt Stufler. Gibbs partitions: a comprehensive phase diagram. *Ann. Inst. Henri Poincaré, Probab. Stat.*, 2024 (to appear).
- [129] Chun Su, Qunqiang Feng, and Zhishui Hu. Uniform recursive trees: branching structure and simple random downward walk. J. Math. Anal. Appl., 315(1):225–243, 2006.
- [130] Vladimir Vasilevich Uchaikin and Vladimir Mikhailovich Zolotarev. *Chance and Stability. Stable Distributions and their Applications*. De Gruyter, 1999.
- [131] Michael Wallner. A half-normal distribution scheme for generating functions. *European J. Combin.*, 87:103138, 21, 2020.
- [132] Edmund Taylor Whittaker and George Neville Watson. A Course of Modern Analysis. Fifth edition (1st ed.: 1902). Cambridge University Press, 2021.
- [133] Gordon E. Willmot. Mixed compound Poisson distributions. ASTIN Bulletin, 16(S1):S59-S79, 1986.
- [134] Gordon E. Willmot and X. Sheldon Lin. *Lundberg Approximations for Compound Distributions with Insurance Applications*. Springer, 2001.
- [135] Anders Wiman. Über den Fundamentalsatz in der Theorie der Funktionen $E_{\alpha}(x)$. Acta Math., 29:191–201, 1905.