

CONTACT POINTS WITH INTEGER FREQUENCIES IN THE THIN OBSTACLE PROBLEM

OVIDIU SAVIN AND HUI YU

ABSTRACT. For the thin obstacle problem, we develop a unified approach that leads to rates of convergence to blow-up profiles at contact points with integer frequencies. For these points, we also obtain a stratification result.

1. INTRODUCTION

The *thin obstacle problem* studies the following system

$$(1.1) \quad \begin{cases} \Delta u \leq 0 & \text{in } B_1, \\ u \geq 0 & \text{on } B_1 \cap \{x_d = 0\}, \\ \Delta u = 0 & \text{in } B_1 \cap (\{u > 0\} \cup \{x_d \neq 0\}). \end{cases}$$

Here we denote by B_1 the unit ball in the Euclidean space \mathbb{R}^d . For a point $x \in \mathbb{R}^d$, we decompose its coordinate as $x = (x', x_d)$ with $x' \in \mathbb{R}^{d-1}$ and $x_d \in \mathbb{R}$. Since the odd part of the solution, $(u(x', x_d) - u(x', -x_d))/2$, is harmonic, it is customary to remove it and assume that *the solution u is even in the x_d -direction*.

After earlier results by Richardson [R] and Uraltseva [U], Athanasopoulos and Caffarelli showed in [AC] that the solution is locally Lipschitz in B_1 , and is locally $C^{1,1/2}$ when restricted to either $B_1 \cap \{x_d \geq 0\}$ or $B_1 \cap \{x_d \leq 0\}$. This optimal regularity of the solution opened the door to the study of the *contact set*

$$\Lambda(u) := \{u = 0\} \cap \{x_d = 0\}$$

and the *free boundary*

$$\Gamma(u) := \partial\{u > 0\} \cap \{x_d = 0\}.$$

In Athanasopoulos-Caffarelli-Salsa [ACS], the authors made a breakthrough by applying Almgren's monotonicity formula to show that for each point $q \in \Lambda(u)$, there is a constant λ_q such that

$$\|u\|_{\mathcal{L}^2(\partial B_r(q))} \sim r^{\frac{d-1}{2} + \lambda_q}$$

as $r \rightarrow 0$. This constant λ_q is called the *frequency of the solution at q* . They also showed that the normalized rescalings converge to a *blow-up profile*, that is,

$$(1.2) \quad u_{q,r}(\cdot) := r^{\frac{d-1}{2}} \frac{u(r \cdot + q)}{\|u\|_{\mathcal{L}^2(\partial B_r(q))}} \rightarrow u_0$$

along a subsequence of $r \rightarrow 0$. The limit u_0 is a λ_q -homogeneous solution to (1.1) in \mathbb{R}^d .

It is interesting to study admissible values of frequencies, to classify homogeneous solutions, and to establish regularity of the contact set/ the free boundary.

O. S. is supported by NSF grant DMS-1800645.

H. Y. is supported by NSF grant DMS-1954363.

So far this program is completed only when $d = 2$. See, for instance, Petrosyan-Shahgholian-Uraltseva [PSU].

Let \mathcal{A} denote the set of admissible frequencies, that is,

$$\mathcal{A} = \{\lambda \in \mathbb{R} : \text{there is a non-trivial } \lambda\text{-homogeneous solution to (1.1)}\}.$$

For a solution u to (1.1) and $\lambda \in \mathcal{A}$, let $\Lambda_\lambda(u)$ denote the set of contact points with frequency λ , that is,

$$\Lambda_\lambda(u) = \{q \in \Lambda(u) : \lambda_q = \lambda\}.$$

In general dimensions, Athanasopoulos-Caffarelli-Salsa [ACS] showed that $\mathcal{A} \subset \{1, \frac{3}{2}\} \cup [2, +\infty)$. Explicit examples give that $\mathbb{N} \cup \{2k - \frac{1}{2} : k \in \mathbb{N}\} \subset \mathcal{A}$. See, for instance, [PSU]. Around each $m \in \mathbb{N}$, there is a frequency gap [CSV, SY2], in the sense that we can find $\alpha_m > 0$, depending on m and d , such that

$$\mathcal{A} \cap (m - \alpha_m, m + \alpha_m) = \{m\} \text{ for each } m \in \mathbb{N}.$$

As for the classification of homogeneous solutions and for the regularity of the contact set/ the free boundary, most results center on points with frequencies in $\{\frac{3}{2}\} \cup \mathbb{N}$.

Already in [ACS], it was known that the only $\frac{3}{2}$ -homogeneous solutions are rotations and multiples of

$$u_{\frac{3}{2}}(x', x_d) = \operatorname{Re}(x_{d-1} + i|x_d|)^{3/2}.$$

For a solution to (1.1), the set $\Lambda_{\frac{3}{2}}(u)$ is relatively open in $\Gamma(u)$. The free boundary is an analytic manifold of dimension $(d - 2)$ near $\Lambda_{\frac{3}{2}}(u)$ [DS, KPS].

For an even integer $2k$, Garofalo-Petrosyan [GP] classified all $2k$ -homogeneous solutions to (1.1) as

$$(1.3) \quad \mathcal{P}_{2k}^+ = \{p : \Delta p = 0 \text{ and } x \cdot \nabla p = 2kp \text{ in } \mathbb{R}^d, \\ p(\cdot, 0) \geq 0 \text{ and } p(\cdot, x_d) = p(\cdot, -x_d)\}.$$

Let u be a solution to (1.1) and $q \in \Lambda(u)$. Garofalo-Petrosyan gave a geometric characterization of contact points with even frequencies

$$q \in \Lambda_{2k}(u) \text{ for some } k \in \mathbb{N} \iff \mathcal{H}^{d-1}(\Lambda(u) \cap B_r(q)) = o(r^{d-1}) \text{ as } r \rightarrow 0.$$

Here \mathcal{H}^{d-1} is the $(d - 1)$ -dimensional Hausdorff measure. In particular, we have $\Lambda_{2k}(u) \subset \Gamma(u)$. They also showed that $\Lambda_{2k}(u)$ is locally covered by C^1 -manifolds. By quantifying the rate of convergence in (1.2) at points in $\Lambda_{2k}(u)$, the regularity of the covering manifolds was improved to C^{1, \log^c} in Colombo-Spolaor-Velichkov [CSV].

More recently, there has been some interest in the study of contact points with odd frequencies, mainly motivated by the connection to the singular set in the obstacle problem, see [FRS]. While points in $\Lambda_1(u)$ lie in the interior of $\Lambda(u)$, it remains an open question whether $\Lambda_{2k+1}(u)$ can contain points in the free boundary for $k \geq 1$. On the other hand, it is known that around $\Lambda_{2k+1}(u)$, the contact set has full density, that is,

$$q \in \Lambda_{2k+1}(u) \text{ for some } k \in \mathbb{N} \iff \mathcal{H}^{d-1}(\{u > 0, x_d = 0\} \cap B_r(q)) = o(r^{d-1}) \text{ as } r \rightarrow 0.$$

See, for instance, Proposition 7.1 in Fernández-Real [Fe].

The family of $(2k + 1)$ -homogeneous solutions to (1.1) was recently classified by Figalli, Ros-Oton and Serra in [FRS] as

$$(1.4) \quad \mathcal{P}_{2k+1}^+ = \{p : \Delta p = 0 \text{ in } \{x_d \neq 0\} \text{ and } \Delta p \leq 0 \text{ in } \mathbb{R}^d, \\ x \cdot \nabla p = (2k + 1)p \text{ in } \mathbb{R}^d, \\ p(\cdot, 0) = 0 \text{ and } p(\cdot, x_d) = p(\cdot, -x_d)\}.$$

They also proved uniqueness of the blow-up profile u_0 in (1.2) at $q \in \Lambda_{2k+1}(u)$.

Along a different direction, Focardi-Spadaro proved that the free boundary $\Gamma(u)$ is countably $(d - 2)$ -rectifiable in [FoS1, FoS2]. They also showed that outside a set of dimension at most $(d - 3)$, all free boundary points have frequencies in $\{2k, 2k + 1, 2k - \frac{1}{2} : k \in \mathbb{N} \setminus \{0\}\}$. For generic boundary data, Fernández-Real and Ros-Oton showed that the free boundary is smooth outside a set of dimension at most $(d - 3)$ in [FeR].

In this paper, we focus on contact points with integer frequencies, that is, points in $\cup_{k \in \mathbb{N}} \Lambda_k(u)$. Around these points, we develop a unified approach that gives a uniform rate for the convergence in (1.2).

Our main result is:

Theorem 1.1. *Suppose that u is a solution to the thin obstacle problem (1.1), and that $0 \in \Lambda_m(u)$ for some $m \in \mathbb{N}$.*

If $m = 2k + 1$ is odd, then there is a constant $\alpha \in (0, 1)$, depending only on k and the dimension d , such that

$$(1.5) \quad u(x) = p(x) + O(|x|^{2k+1+\alpha}) \text{ as } x \rightarrow 0$$

for some $p \in \mathcal{P}_{2k+1}^+$.

If $m = 2k$ is even, then there is a constant $c > 0$, depending only on k and d , such that

$$(1.6) \quad u(x) = p(x) + O(|x|^{2k} (-\log|x|)^{-c}) \text{ as } x \rightarrow 0$$

for some $p \in \mathcal{P}_{2k}^+$.

Remark 1.1. While it is known in [FRS] that rescaled solutions converge to some $p \in \mathcal{P}_{2k+1}^+$ at points with odd frequencies, this is the first time a quantified rate of convergence has been obtained. Corresponding results at even frequency points were known in Colombo-Spolaor-Velichkov [CSV]. Our method is different and applies to all points with integer frequencies. It also leads to an improved exponent c in (1.6), and in the corresponding log-epiperimetric inequality from [CSV] at points with even frequencies, see Remark 4.1.

With a standard application of Whitney's extension theorem and the implicit function theorem, Theorem 1.1 leads to the following stratification result of $\Lambda_m(u)$:

Theorem 1.2. *Suppose that u is a solution to the thin obstacle problem (1.1).*

For each $m \in \mathbb{N}$, we have the following decomposition

$$\Lambda_m(u) = \cup_{j=0,1,\dots,d-2} \Lambda_m^j(u).$$

The lowest stratum $\Lambda_m^0(u)$ is locally isolated.

If m is odd, then $\Lambda_m^j(u)$ is locally covered by a j -dimensional $C^{1,\alpha}$ manifold for each $j = 1, \dots, d - 2$.

If m is even, then $\Lambda_m^j(u)$ is locally covered by a j -dimensional $C^{1,\log}$ manifold for each $j = 1, \dots, d - 2$.

Remark 1.2. Points in $\Lambda_1(u)$ and $\Lambda_3^0(u)$ lie in the interior of the contact set $\Lambda(u)$. It remains open whether other strata of $\Lambda_{2k+1}(u)$ can contain points on the free boundary, see Remark 5.1.

To obtain the results at even-frequency points, the approach taken by Colombo-Spolaor-Velichkov [CSV] is based on the decomposition of the energy in terms of Fourier modes. This leads to a log-epiperimetric inequality for the $2k$ -Weiss energy functional (see (2.3)). On the other hand, our method is based on the classic technique of linearization as in De Silva [D]. By working directly in the physical space instead of the Fourier space, it seems that we are able to get more detailed information.

The main challenge is that solutions to the linearized problem do not have to satisfy the constraints in (1.1) (they might fail $\Delta u \leq 0$ or $u(x', 0) \geq 0$). In our approach, this issue is fixed by solving a ‘boundary layer problem’ near the hyperplane $\{x_d = 0\}$. For each unconstrained m -homogenous harmonic polynomial p , we associate its approximation \bar{p} that satisfies the constraints on $\{x_d = 0\}$ and is harmonic up to an error κ_p away from this hyperplane. We use the class of functions \bar{p} to approximate the solution u inductively in dyadic balls B_r , while keeping track of the rescaled error $\varepsilon \geq \kappa_p$. We introduce the notation $u \in \mathcal{S}_m(p, \varepsilon, r)$ when a solution u is ε -approximated by \bar{p} at scale r , see Definition 2.2.

With this notation, the main lemma is the following:

Lemma 1.1. *Given $m \in \mathbb{N}$, there are constants, $\tilde{\varepsilon}$, r_0 , c small, C big, such that if $u \in \mathcal{S}_m(p, \varepsilon, 1)$ with $\varepsilon < \tilde{\varepsilon}$ and $1 \leq \|p\|_{\mathcal{L}^2(\mathbb{S}^{d-1})} \leq 2$, then we have the following dichotomy:*

a) *Either*

$$W_m(u; 1) - W_m(u; r_0) \geq c\varepsilon^2,$$

and

$$u \in \mathcal{S}_m(p, C\varepsilon, r_0);$$

b) *or*

$$u \in \mathcal{S}_m(p', \frac{1}{2}\varepsilon, r_0)$$

for some p' with

$$\|p' - p\|_{\mathcal{L}^2(\mathbb{S}^{d-1})} \leq C\varepsilon.$$

Here the integer m denotes the frequency of the contact point, and W_m is the Weiss energy functional, see (2.3).

Lemma 1.1 states that when moving to a smaller scale, we can improve, by a definite amount, either the Weiss energy or the error in approximation. On the other hand, the Weiss energy is controlled by the error ε in the approximation, see Lemma 2.8. It follows that both quantities decay in a quantified fashion. Similar ideas have been applied to the nonlinear obstacle problem in [SY1] as well as the triple-membrane problem in [SY3].

This paper is organized as follows: In Section 2, we collect some preliminary results and introduce the boundary layer problem. In Sections 3 and 4, we prove Lemma 1.1 at points with the odd and even frequencies. These two sections contain the heart of this work. In Section 5, we conclude with the proof of our main results.

2. PRELIMINARIES

In this section, we collect some useful results and introduce the boundary layer problem.

Throughout this paper, we denote by u a solution to the thin obstacle problem (1.1) on some domain inside \mathbb{R}^d with $d \geq 3$. This space is decomposed as

$$\mathbb{R}^d = \{(x', x_d) : x' \in \mathbb{R}^{d-1}, x_d \in \mathbb{R}\}.$$

For a set $E \subset \mathbb{R}^d$, we define the following subsets relative to $\{x_d = 0\}$:

$$E' = E \cap \{x_d = 0\}, E^+ = E \cap \{x_d > 0\} \text{ and } E^- = E \cap \{x_d < 0\}.$$

In particular, the contact set is

$$\Lambda(u) = \{u = 0\}'.$$

By applying Almgren's monotonicity formula to the thin obstacle problem, Athanassopoulos, Caffarelli and Salsa showed in [ACS] that the contact set $\Lambda(u)$ can be decomposed according to the frequencies of the contact points. In this work, we focus on points with integer frequencies. Thanks to [GP] and [FRS], these points can be characterized as

$$(2.1) \quad \Lambda_{2k}(u) = \{q \in \Lambda(u) : u_{q,r} \rightarrow u_0 \in \mathcal{P}_{2k}^+ \text{ as } r \rightarrow 0\}$$

and

$$(2.2) \quad \Lambda_{2k+1}(u) = \{q \in \Lambda(u) : u_{q,r} \rightarrow u_0 \in \mathcal{P}_{2k+1}^+ \text{ as } r \rightarrow 0\}.$$

Recall the definition of normalized rescalings $u_{q,r}$ from (1.2). The spaces of homogeneous solutions, \mathcal{P}_{2k}^+ and \mathcal{P}_{2k+1}^+ , were introduced in (1.3) and (1.4).

When focusing on a particular frequency, constants depending only on that frequency and the dimension d are called *universal constants*.

2.1. Weiss monotonicity formula and consequences. First used by Weiss for the obstacle problem in [W], the Weiss monotonicity formula has been indispensable in the study of free boundary problems. Garofalo-Petrosyan [GP] introduced its analogue to the thin obstacle problem.

For each $\lambda \in \mathbb{R}$, the λ -Weiss energy functional is

$$(2.3) \quad W_\lambda(u; r) = \frac{1}{r^{d-2+2\lambda}} \int_{B_r} |\nabla u|^2 - \frac{\lambda}{r^{d-1+2\lambda}} \int_{\partial B_r} u^2.$$

We collect some of its properties in the following lemma. For its proof, see Theorem 1.4.1 and Theorem 1.5.4 in [GP].

Lemma 2.1. *Suppose that u solves the thin obstacle problem in B_1 . Then for $r \in (0, 1)$, we have*

$$(2.4) \quad \frac{d}{dr} W_\lambda(u; r) = \frac{2}{r} \int_{\partial B_1} (\nabla u_r \cdot \nu - \lambda u_r)^2,$$

where $u_r(x) = u(rx)/r^\lambda$. In particular, $r \mapsto W_\lambda(u; r)$ is non-decreasing.

If we further assume that $0 \in \Lambda_\lambda(u)$, then $\lim_{r \rightarrow 0} W_\lambda(u; r) = 0$.

Under the same assumptions as in Lemma 2.1, we can integrate (2.4) and apply Hölder's inequality to get

$$(2.5) \quad \int_{\partial B_1} |u_r - u_s| \leq (\log(r/s))^{\frac{1}{2}} [W_\lambda(u; r) - W_\lambda(u; s)]^{\frac{1}{2}}$$

for $0 < s < r < 1$.

2.2. The boundary layer problem. When dealing with the linearized problem, we need to work with polynomials that may fail the constraints in (1.1). These polynomials form the following spaces:

$$(2.6) \quad \mathcal{P}_{2k} = \{p : \Delta p = 0 \text{ and } x \cdot \nabla p = 2kp \text{ in } \mathbb{R}^d, \text{ and } p(\cdot, x_d) = p(\cdot, -x_d)\}$$

and

$$(2.7) \quad \mathcal{P}_{2k+1} = \{p : \Delta p = 0 \text{ in } \{x_d \neq 0\}, x \cdot \nabla p = (2k+1)p \text{ in } \mathbb{R}^d, \\ p(\cdot, 0) = 0 \text{ and } p(\cdot, x_d) = p(\cdot, -x_d)\}.$$

Compared with (1.3) and (1.4), polynomials in \mathcal{P}_{2k} may fail to be non-negative along $\{x_d = 0\}$, and polynomials in \mathcal{P}_{2k+1} may fail to be superharmonic. We ‘correct’ such error by solving a thin obstacle problem in a boundary layer on the sphere \mathbb{S}^{d-1} .

To be precise, for small $\eta > 0$, the *boundary layer of width η* is defined as

$$L_\eta = \{(x', x_d) : |x_d| < \eta|x|\}.$$

This is the region trapped by the following surfaces

$$S_\eta^+ = \{(x', x_d) : x_d = \eta|x|\} \text{ and } S_\eta^- = \{(x', x_d) : x_d = -\eta|x|\}.$$

When there is no ambiguity, we denote their intersections with \mathbb{S}^{d-1} by the same expressions.

Remark 2.1. Given $m \in \mathbb{N}$, we fix η small enough, depending only on m and d , so that the first Dirichlet eigenvalue of the operator $\Delta_{\mathbb{S}^{d-1}} + \lambda(m)$ in L_η is negative, where $\Delta_{\mathbb{S}^{d-1}}$ is the spherical Laplacian, and

$$\lambda(m) = m(m + d - 2).$$

In particular, the following is well-defined:

Definition 2.1. Given $m \in \mathbb{N}$ and $p \in \mathcal{P}_m$, the *replacement of p* , denoted by \bar{p} , is the minimizer of the following energy

$$w \mapsto \int_{\mathbb{S}^{d-1}} |\nabla_{\mathbb{S}^{d-1}} w|^2 - \lambda(m)w^2$$

over functions satisfying $w \geq 0$ on $\{x_d = 0\}$ and $w = p$ outside L_η .

Here $\nabla_{\mathbb{S}^{d-1}}$ denotes the tangential gradient on \mathbb{S}^{d-1} .

Denote the difference of p and its replacement \bar{p} by v_p , that is,

$$v_p = \bar{p} - p.$$

Remark 2.2. The replacement solves the thin obstacle problem for the operator $\Delta_{\mathbb{S}^{d-1}} + \lambda(m)$ in the boundary layer, namely,

$$\begin{cases} (\Delta_{\mathbb{S}^{d-1}} + \lambda(m))\bar{p} \leq 0 & \text{in } L_\eta, \\ \bar{p} \geq 0 & \text{on } (\mathbb{S}^{d-1})', \\ (\Delta_{\mathbb{S}^{d-1}} + \lambda(m))\bar{p} = 0 & \text{in } L_\eta \cap (\{x_d \neq 0\} \cup \{\bar{p} > 0\}). \end{cases}$$

As a result, we can view $(\Delta_{\mathbb{S}^{d-1}} + \lambda(m))\bar{p}$ as a signed measure, supported along S_η^\pm and $\mathbb{S}^{d-1} \cap \{\bar{p} = 0\}'$, of the following form

$$(\Delta_{\mathbb{S}^{d-1}} + \lambda(m))\bar{p} = f_p d\mathcal{H}^{d-2}|_{S_\eta^\pm} + g_p d\mathcal{H}^{d-2}|_{(\mathbb{S}^{d-1})'}.$$

With an abuse of notation, we denote the m -homogeneous extension of \bar{p} and the corresponding extensions of f_p and g_p by the same notations. This way, we have

$$\Delta \bar{p} = f_p d\mathcal{H}^{d-1}|_{S_\eta^\pm} + g_p d\mathcal{H}^{d-1}|_{\{x_d=0\}}.$$

For each $p \in \mathcal{P}_m$, the following constant, κ_p , measures the extent to which p fails to be a solution to the thin obstacle problem:

$$\kappa_p := \int_{S_\eta^+ \cap \mathbb{S}^{d-1}} f_p d\mathcal{H}^{d-2}.$$

For these functions and constants, we often omit the subscript p when there is no ambiguity.

Lemma 2.2. *Using the notations in Definition 2.1 and Remark 2.2, we have*

- 1) $v_p \geq 0$ on \mathbb{S}^{d-1} , $\kappa_p \geq 0$.
- 2) *There are universal constants, c and C , such that*

$$c\kappa_p \leq f_p \leq C\kappa_p \text{ on } S_\eta^\pm \cap \mathbb{S}^{d-1}.$$

Proof. The first statement follows directly from the maximum principle.

To see the second statement, we first note that v is a non-negative harmonic function in L_η^+ . By the strong maximum principle, it suffices to consider the case when $v > 0$ in L_η^+ .

In this case, we can apply the Harnack principle to v inside $(B_2 \setminus B_{1/2}) \cap L_\eta^+$ to get $c \leq \frac{\sup_K v}{\inf_K v} \leq C$, where $K = \mathbb{S}^{d-1} \cap \{x_d = \frac{1}{2}\eta\}$.

If we denote by ν the unit normal along S_η^+ that is exterior to L_η , by the boundary Harnack principle, we have

$$(2.8) \quad c \inf_K v \leq \partial_\nu v(x) \leq C \sup_K v \quad \forall x \in \mathbb{S}^{d-1} \cap S_\eta^+.$$

Note that $f = \partial_\nu v$ along S_η^+ , the conclusion follows. \square

We also have the following bounds for κ_p :

Lemma 2.3. *Using the notations in Definition 2.1 and Remark 2.2, and further assume $\|p\|_{\mathcal{L}^2(\mathbb{S}^{d-1})} \leq 1$, we can find a universal constant C such that*

$$\kappa_p \leq C \sup_{\mathbb{S}^{d-1}} v_p.$$

Moreover, when m is even, we have

$$\left(\sup_{\mathbb{S}^{d-1}} v_p \right)^{\frac{d-1}{2}} \leq C\kappa_p.$$

Proof. Directly from (2.8), we have the bound $\kappa_p \leq C \sup_{\mathbb{S}^{d-1}} v_p$.

For the second comparison, we note that when m is even, $v = \bar{p} - p$ solves the thin obstacle problem in L_η with $-p$ as the obstacle and 0 as boundary data. Suppose

$$\varepsilon = -p(e_1) = \sup_{(\mathbb{S}^{d-1})'} (-p),$$

where e_1 is the unit vector in the x_1 -direction, then $\sup_{\mathbb{S}^{d-1}} v \leq \varepsilon$. Regularity of p gives $v \geq -p \geq \frac{7}{8}\varepsilon$ in $B'_{c\varepsilon^{1/2}}(e_1) \cap \mathbb{S}^{d-1}$, which leads to $v \geq \frac{7}{8}\varepsilon$ in $B_{c\varepsilon^{1/2}}(e_1) \cap \mathbb{S}^{d-1}$ by a scaling argument.

From here we have $v(e_1, \frac{1}{2}\eta) \geq c\varepsilon^{\frac{d-1}{2}}$ by comparing with a truncation and rescaling of the Green's function for $\Delta_{\mathbb{S}^{d-1}}$ with a pole at e_1 and vanishes outside $B_\eta(e_1) \cap \mathbb{S}^{d-1}$. Combining this with (2.8) gives the desired result. \square

With this, we have the following control over the size of $\|v\|_{H^1}$ in terms of κ :

Lemma 2.4. *Using the notations in Definition 2.1 and Remark 2.2, and further assume $\|p\|_{\mathcal{L}^2(\mathbb{S}^{d-1})} \leq 1$, we have a universal constant C such that*

$$\|v_p\|_{H^1(B_1)} \leq C\kappa_p^{\frac{1}{2} + \frac{1}{d-1}} \text{ if } p \in \mathcal{P}_{2k}$$

and

$$\|v_p\|_{H^1(B_1)} \leq C\kappa_p^{1/2} \text{ if } p \in \mathcal{P}_{2k+1}.$$

Proof. We first deal with the case when $p \in \mathcal{P}_{2k}$.

With the homogeneity and harmonicity of p , we have

$$\begin{aligned} \int_{\mathbb{S}^{d-1}} |\nabla_{\mathbb{S}^{d-1}} \bar{p}|^2 - \lambda(2k) \bar{p}^2 &= - \int_{\mathbb{S}^{d-1}} v(\Delta_{\mathbb{S}^{d-1}} + \lambda(2k)) \bar{p} \\ &\leq - \sup_{\mathbb{S}^{d-1}} v \int_{(\mathbb{S}^{d-1})'} g. \end{aligned}$$

Now let P denote the $2k$ -homogeneous harmonic polynomial with $P = 1$ on $(\mathbb{S}^{d-1})'$. Then $\int_{\mathbb{S}^{d-1}} P(\Delta_{\mathbb{S}^{d-1}} + \lambda(2k)) \bar{p} = \int_{\mathbb{S}^{d-1}} \bar{p}(\Delta_{\mathbb{S}^{d-1}} + \lambda(2k)) P = 0$ gives

$$- \int_{(\mathbb{S}^{d-1})'} g \sim \int_{S_\eta^\pm} f \sim \kappa.$$

Thus

$$(2.9) \quad \int_{\mathbb{S}^{d-1}} |\nabla_{\mathbb{S}^{d-1}} \bar{p}|^2 - \lambda \bar{p}^2 \leq C\kappa^{1 + \frac{2}{d-1}}$$

by Lemma 2.3.

Using again the homogeneity and harmonicity of $p \in \mathcal{P}_{2k}$, this implies

$$\int_{\mathbb{S}^{d-1}} |\nabla_{\mathbb{S}^{d-1}} v|^2 - \lambda v^2 = \int_{\mathbb{S}^{d-1}} |\nabla_{\mathbb{S}^{d-1}} \bar{p}|^2 - \lambda \bar{p}^2 \leq C\kappa^{1 + \frac{2}{d-1}}.$$

To conclude the estimate for $p \in \mathcal{P}_{2k}$, we simply note that the left-hand side is comparable to $\|v\|_{H^1(B_1)}^2$ when η is chosen small.

Now we deal with the case when $p \in \mathcal{P}_{2k+1}$.

In this case, note that p is admissible in the minimization problem in Definition 2.1, we have

$$(2.10) \quad \int_{\mathbb{S}^{d-1}} |\nabla_{\mathbb{S}^{d-1}} \bar{p}|^2 - \lambda(2k+1) \bar{p}^2 \leq \int_{\mathbb{S}^{d-1}} |\nabla_{\mathbb{S}^{d-1}} p|^2 - \lambda(2k+1) p^2 = 0,$$

which implies

$$\int_{\mathbb{S}^{d-1}} |\nabla_{\mathbb{S}^{d-1}} (\bar{p} - p)|^2 - \lambda(\bar{p} - p)^2 \leq 2 \int_{\mathbb{S}^{d-1}} p(\Delta_{\mathbb{S}^{d-1}} + \lambda)(\bar{p} - p) = 2 \int_{\mathbb{S}^{d-1} \cap S_\eta^\pm} p f d\mathcal{H}^{d-2}.$$

For the last equality, we used $p = 0$ along $\{x_d = 0\}$, and $\Delta p = 0$ away from $\{x_d = 0\}$. This gives

$$\int_{\mathbb{S}^{d-1}} |\nabla_{\mathbb{S}^{d-1}} v|^2 - \lambda v^2 \leq C\kappa.$$

We conclude by noting the left-hand side is comparable to $\|v\|_{H^1(B_1)}^2$ when η is small. \square

For $p \in \mathcal{P}_m$ with $\bar{p} \neq p$, we sometimes need to absorb the right-hand side f_p from Remark 2.2. To do so, we need some auxiliary functions.

Firstly, let $\varphi_p : \mathbb{S}^{d-1} \rightarrow \mathbb{R}$ denote the projection of the normalized f_p onto \mathcal{P}_m , that is,

$$(2.11) \quad \varphi_p = \sum \left\langle \frac{f_p}{\kappa_p}, p_j \right\rangle p_j,$$

where $\{p_j\}$ is an orthonormal basis for \mathcal{P}_m in $\mathcal{L}^2(\mathbb{S}^{d-1})$ and

$$\left\langle \frac{f_p}{\kappa_p}, p_j \right\rangle = \frac{1}{\kappa_p} \int_{\mathbb{S}^{d-1} \cap S_\eta^\pm} p_j f_p d\mathcal{H}^{d-2}.$$

In particular, the difference $\frac{f_p}{\kappa_p} - \varphi_p$ is perpendicular to \mathcal{P}_m . Consequently, the theory of Fredholm implies that we can find a unique function H_p on \mathbb{S}^{d-1} , that is even with respect to x_d and satisfies:

If $m = 2k$, then

$$(2.12) \quad (\Delta_{\mathbb{S}^{d-1}} + \lambda(2k))H_p = \frac{f_p}{\kappa_p} - \varphi_p \text{ on } \mathbb{S}^{d-1};$$

If $m = 2k + 1$, then

$$(2.13) \quad (\Delta_{\mathbb{S}^{d-1}} + \lambda(2k + 1))H_p = \frac{f_p}{\kappa_p} - \varphi_p \text{ on } (\mathbb{S}^{d-1})^\pm, \text{ and } H_p(\cdot, 0) = 0.$$

If we denote its m -homogeneous extension also by H_p , then

$$(2.14) \quad \Phi_p := H_p + \frac{1}{d + 2m - 2} \varphi_p \left(\frac{x}{|x|} \right) |x|^m \log |x|$$

satisfies

$$\Delta \Phi_p = \frac{f_p}{\kappa_p} d\mathcal{H}^{d-1} \Big|_{S_\eta^\pm} \text{ in } \mathbb{R}^d \text{ if } m = 2k;$$

and

$$\Delta \Phi_p = \frac{f_p}{\kappa_p} d\mathcal{H}^{d-1} \Big|_{S_\eta^\pm} \text{ in } (\mathbb{R}^d)^\pm, \text{ and } \Phi_p(\cdot, 0) = 0 \text{ if } m = 2k + 1.$$

We often omit the subscript when there is no ambiguity.

For our argument, it is crucial that f_p has a non-trivial projection into \mathcal{P}_m :

Lemma 2.5. *If $\kappa_p \neq 0$, then there are universal positive constants c and C such that*

$$c \leq \|\varphi_p\|_{\mathcal{L}^2(\mathbb{S}^{d-1})} \leq C,$$

and

$$\|\Phi_p\|_{C^{0,1}(B_1)} \leq C.$$

Proof. Both upper bounds follow from the definitions of φ , Φ and Lemma 2.2.

For the lower bound for φ , it suffices to note that we can find $q \in \mathcal{P}_m$ with $\|q\|_{\mathcal{L}^2(\mathbb{S}^{d-1})} = 1$ and $q \geq c > 0$ along S_η^\pm if η is small. \square

Lemma 2.6. *If u solves (1.1) in B_1 , and $p \in \mathcal{P}_{2k} \cup \mathcal{P}_{2k+1}$, then*

$$\|u - \bar{p}\|_{\mathcal{L}^\infty(B_{1/2})} + \|u - \bar{p}\|_{H^1(B_{1/2})} \leq C(\|u - \bar{p}\|_{\mathcal{L}^2(B_1)} + \kappa_p)$$

for a universal constant C .

Proof. By Definition 2.1, we have $\bar{p} \geq 0$ along $\{x_d = 0\}$. Thus $\Delta u = 0$ inside $\{u - \bar{p} > 0\}$. As a result,

$$\Delta(u - \bar{p}) = -\Delta\bar{p} \geq -fd\mathcal{H}^{d-1}|_{S_\eta^\pm} \text{ in } \{u - \bar{p} > 0\}.$$

Similarly, inside $\{\bar{p} - u > 0\}$, we have $\Delta\bar{p} \geq 0$. Thus

$$\Delta(u - \bar{p}) \leq \Delta u \leq 0 \text{ in } \{\bar{p} - u > 0\}.$$

Combining these, we have

$$\Delta|u - \bar{p}| \geq -fd\mathcal{H}^{d-1}|_{S_\eta^\pm} \text{ in } B_1.$$

Meanwhile, let ζ be a smooth non-negative function on \mathbb{S}^{d-1} satisfying

$$\zeta = 0 \text{ on } (\mathbb{S}^{d-1})' \text{ and } \zeta = 1 \text{ on } \{|x_d| \geq \frac{\eta}{2}|x'|\}.$$

Recall the auxiliary function from (2.14), we have

$$\Delta(\zeta\Phi) = \frac{1}{\kappa}fd\mathcal{H}^{d-1}|_{S_\eta^\pm} + R,$$

where the remainder R is universally bounded in B_1 . Consequently, we have

$$\Delta(|u - \bar{p}| + \kappa\zeta\Phi) \geq -C\kappa \text{ in } B_1.$$

Together with Lemma 2.5, this gives the estimates on $\|u - \bar{p}\|_{\mathcal{L}^\infty(B_{1/2})}$ and $\|u - \bar{p}\|_{H^1(B_{1/2})}$. \square

Remark 2.3. Lemma 2.6 is the reason why it is preferable to work with the replacement \bar{p} rather than the original p .

2.3. Well-approximated solutions. The heart of this paper is Lemma 1.1, where we improve the approximation of a solution u by replacements of polynomials from \mathcal{P}_{2k} or \mathcal{P}_{2k+1} , defined in (2.6) and (2.7).

Let u be a solution to (1.1) in B_1 , and let $p \in \mathcal{P}_m$ for $m = 2k$ or $2k + 1$. The distance between them is denoted by

$$(2.15) \quad \delta(u, p) := \max\{\|u - \bar{p}\|_{H^1(B_1)}, \kappa_p\}.$$

Here we use notations from Definition 2.1 and Remark 2.2.

Definition 2.2. Given $\varepsilon > 0$, we say that u is ε -approximated by $p \in \mathcal{P}_m$ at scale $r > 0$, and write

$$u \in \mathcal{S}_m(p, \varepsilon, r)$$

if

$$\delta(u_r, p) < \varepsilon,$$

where $u_r(x) = \frac{1}{r^m}u(rx)$.

We collect some immediate consequences.

Lemma 2.7. *If $u \in \mathcal{S}_m(p, \varepsilon, 1)$, then*

$$W_m(u; 3/4) \leq W_m(\bar{p}; 3/4) + C\varepsilon^2$$

for a universal C .

Proof. With $\|u - \bar{p}\|_{H^1(B_1)} \leq \varepsilon$, we can find $\rho \in [\frac{3}{4}, \frac{7}{8}]$ such that

$$\int_{\partial B_\rho} (u_\nu - \bar{p}_\nu)^2 + (u - \bar{p})^2 d\mathcal{H}^{d-1} \leq C\varepsilon^2.$$

A direct computation gives

$$\begin{aligned} W_m(\bar{p}; \rho) - W_m(u; \rho) &= \frac{1}{\rho^{d+2m-2}} \int_{B_\rho} |\nabla(\bar{p} - u)|^2 - 2\Delta u(\bar{p} - u) \\ &\quad + \frac{1}{\rho^{d+2m-2}} \int_{\partial B_\rho} 2u_\nu(\bar{p} - u) - \frac{m}{\rho^{d+2m-1}} \int_{\partial B_\rho} (\bar{p} - u)^2 + 2u(\bar{p} - u). \end{aligned}$$

With $u\Delta u = 0$ and $\bar{p}\Delta u \leq 0$, this implies

$$\begin{aligned} W_m(\bar{p}; \rho) - W_m(u; \rho) &\geq \frac{1}{\rho^{d+2m-2}} \int_{\partial B_\rho} 2(\rho u_\nu - mu)(\bar{p} - u) - m(\bar{p} - u)^2 \\ &\geq -C\varepsilon^2. \end{aligned}$$

where we have used $\rho u_\nu - mu = \rho(u - \bar{p})_\nu - m(u - \bar{p})$.

With monotonicity of W_m and homogeneity of \bar{p} , this implies

$$W_m(u; 3/4) \leq W_m(\bar{p}; 3/4) + C\varepsilon^2.$$

□

A consequence of Lemma 2.7 is the following relation between W_m and ε .

Lemma 2.8. *If $u \in \mathcal{S}_m(p, \varepsilon, 1)$, then*

$$W_m(u; 3/4) \leq C\varepsilon^2, \quad \text{if } m \text{ is odd,}$$

and

$$W_m(u; 3/4) \leq C\varepsilon^{1+\frac{2}{d-1}}, \quad \text{if } m \text{ is even,}$$

with C universal.

Proof. By Lemma 2.7 we only need to bound $W_m(\bar{p}; 1)$.

When m is odd, $W_{2k+1}(\bar{p}; 1) \leq 0$ by (2.10).

When m is even, note that

$$(2.16) \quad W(\bar{p}; 1) = C \int_{\mathbb{S}^{d-1}} |\nabla_{\mathbb{S}^{d-1}} \bar{p}|^2 - \lambda(2k)\bar{p}^2 \leq C\varepsilon^{1+\frac{2}{d-1}}$$

by (2.9). □

Remark 2.4. The difference between the exponents in Lemma 2.8 leads to the different rates of convergence in Theorem 1.1.

The following is a version of Lemma B.2 from [FRS]. It follows by quantifying the proof in [FRS] and it is left to the reader:

Lemma 2.9. *Suppose that u is a solution to (1.1) in B_1 .*

If $u \leq p + \varepsilon$ in B_1 for some $p \in \mathcal{P}_{2k+1}$ with $\|p\|_{\mathcal{L}^2(\mathbb{S}^{d-1})} \leq 1$, then

$$u = 0 \quad \text{in } B'_{1-\varepsilon^{1/2}} \cap \left\{ \frac{\partial}{\partial x_d} p \leq -M\varepsilon^{1/2} \right\},$$

where M is a universal constant.

Remark 2.5. For a function $w \in C^1(B_1 \cap \{x_d \geq 0\})$, $\frac{\partial}{\partial x_d} w(x', 0)$ denotes the one-sided derivative in the x_d -direction taken in $B_1 \cap \{x_d \geq 0\}$, that is,

$$\frac{\partial}{\partial x_d} w(x', 0) = \lim_{t \rightarrow 0^+} \frac{w(x', t) - w(x', 0)}{t}.$$

3. THE DICHOTOMY AT A POINT WITH ODD FREQUENCY

Suppose that u is a solution to the thin obstacle problem (1.1), and that 0 is a point with integer frequency. By results in [GP, FRS], up to an initial scaling, the solution u is well-approximated in B_1 by some homogeneous solution from either \mathcal{P}_{2k+1}^+ or \mathcal{P}_{2k}^+ as in (1.4) and (1.3). To get a rate of convergence as in Theorem 1.1, we need to improve this approximation at smaller scales.

This is achieved through the dichotomy as in Lemma 1.1, which states that at a smaller scale, either the approximation can be improved in a quantified fashion, or the Weiss energy drops in a quantified fashion. In some sense, this method combines the strengths of the epiperimetric inequality approach as in [CSV] and the approach by linearization as in [D].

In this section and the next, we prove this dichotomy for points with odd and even frequencies, respectively. In the final section of this paper, we show how to deduce the main result from them.

We state the main lemma for this section:

Lemma 3.1 (Dichotomy at a point with odd frequency). *Given $k \in \mathbb{N}$, there are universal constants, $\tilde{\varepsilon}$, r_0 , c small and C big, such that*

If $u \in \mathcal{S}_{2k+1}(p, \varepsilon, 1)$ with $\varepsilon < \tilde{\varepsilon}$ and $1 \leq \|p\|_{\mathcal{L}^2(\mathbb{S}^{d-1})} \leq 2$, then we have the following dichotomy:

a) *Either*

$$W_{2k+1}(u; 1) - W_{2k+1}(u; r_0) \geq c\varepsilon^2,$$

and

$$u \in \mathcal{S}_{2k+1}(p, C\varepsilon, r_0);$$

b) *or*

$$u \in \mathcal{S}_{2k+1}(p', \frac{1}{2}\varepsilon, r_0)$$

for some p' with

$$\|p' - p\|_{\mathcal{L}^2(\mathbb{S}^{d-1})} \leq C\varepsilon.$$

Recall the space of well-approximated solutions \mathcal{S}_{2k+1} from Definition 2.2, and the Weiss energy from (2.3).

The remaining part of this section is devoted to the proof of Lemma 3.1. We argue by contradiction.

Suppose, on the contrary, the lemma is not true. Then we find a sequence (u_n, p_n) satisfying

$$1 \leq \|p_n\|_{\mathcal{L}^2(\mathbb{S}^{d-1})} \leq 2, \text{ and } u_n \in \mathcal{S}_{2k+1}(p_n, \varepsilon_n, 1) \text{ with } \varepsilon_n \rightarrow 0.$$

However, neither a) nor b) holds, that is

$$(3.1) \quad W_{2k+1}(u_n; 1) - W_{2k+1}(u_n; r_0) < \frac{1}{n^2} \varepsilon_n^2,$$

and

$$(3.2) \quad u_n \notin \mathcal{S}_{2k+1}(p', \frac{1}{2}\varepsilon_n, r_0) \quad \forall p' \in \mathcal{P}_{2k+1} \text{ with } \|p' - p_n\|_{\mathcal{L}^2(\mathbb{S}^{d-1})} \leq C\varepsilon_n.$$

The constants r_0 and C_0 will be chosen depending on universal constants.

We will choose $r_0 \leq 1/2$. Hence by monotonicity of the Weiss energy and (3.1), we have

$$(3.3) \quad W_{2k+1}(u_n; 1) - W_{2k+1}(u_n; 1/2) < \frac{1}{n^2} \varepsilon_n^2.$$

Corresponding to this sequence p_n , we have auxiliary functions $f_n, g_n, v_n, \varphi_n, H_n$ and Φ_n , and constants κ_n as in Definition 2.1, Remark 2.2, (2.11) and (2.14).

Now with p_n uniformly bounded in the finite dimensional space \mathcal{P}_{2k+1} , we have, up to a subsequence,

$$p_n \rightarrow p_\infty \text{ uniformly in } C^1(B_1 \cap \{x_d \geq 0\}) \text{ and } C^1(B_1 \cap \{x_d \leq 0\}).$$

As a result, we have $p_\infty \in \mathcal{P}_{2k+1}$. Actually, it is in the more restrictive space \mathcal{P}_{2k+1}^+ (see (1.4)):

Lemma 3.2. $\frac{\partial}{\partial x_d} p_\infty \leq 0$ in B_1' .

Proof. By Lemma 2.4, we have

$$(3.4) \quad \|u_n - p_n\|_{H^1(B_1)} \leq \|u_n - \bar{p}_n\|_{H^1(B_1)} + \|v_n\|_{H^1(B_1)} \leq C\varepsilon_n^{1/2}.$$

Suppose $\frac{\partial}{\partial x_d} p_\infty(x', 0) = \beta > 0$ at some $(x', 0) \in B_1'$.

Since $\frac{\partial}{\partial x_d} u_n(\cdot, 0) \leq 0$, we have $\frac{\partial}{\partial x_d} (p_n - u_n) \geq \frac{1}{2}\beta$ in a neighborhood of $(x', 0)$ for large n . This contradicts (3.4) eventually. \square

Since $p_\infty \in \mathcal{P}_{2k+1}$, the following set

$$\mathcal{N} := \left\{ \frac{\partial}{\partial x_d} p_\infty = 0 \right\}'$$

is of dimension at most $(d-2)$.

Lemma 3.3. *Given a compact set $K \subset B_1' \setminus \mathcal{N}$, we can find $N \in \mathbb{N}$ such that*

$$u_n = 0 \text{ on } K$$

for all $n \geq N$.

Proof. By compactness of K , we can find $\beta > 0$ such that

$$\frac{\partial}{\partial x_d} p_\infty \leq -\beta \text{ on } K.$$

With Lemma 2.6 and (3.4), we have

$$u_n \leq p_n + C_K \varepsilon_n^{1/2} \text{ in a neighborhood of } K.$$

Together with Lemma 2.9, this gives the desired result. \square

Now define the *normalized solutions*

$$(3.5) \quad \hat{u}_n = \frac{u_n - \bar{p}_n}{\varepsilon_n}.$$

With Lemma 2.6 and (2.14), we have that

$$\Delta(\hat{u}_n + \frac{\kappa_n}{\varepsilon_n} \Phi_n) = 0 \text{ in } B_1^+$$

and

$$\|\hat{u}_n + \frac{\kappa_n}{\varepsilon_n} \Phi_n\|_{H^1(B_\rho)} \leq C(\rho) \text{ for any } \rho < 1.$$

Thus, up to a subsequence, $\hat{u}_n + \frac{\kappa_n}{\varepsilon_n} \Phi_n$ converges in $L^2_{loc}(B_1)$ to a limit function $h \in H^1_{loc}(B_1)$, which is harmonic in B_1^+ and B_1^- , and

$$(3.6) \quad \|\hat{u}_n + \frac{\kappa_n}{\varepsilon_n} \Phi_n - h\|_{\mathcal{L}^2(B_{7/8})} = o(1) \text{ as } n \rightarrow \infty.$$

Moreover, Lemma 3.3 implies that $\hat{u}_n + \frac{\kappa_n}{\varepsilon_n} \Phi_n$ vanishes eventually on any compact subsets of $B_1' \setminus \mathcal{N}$, where \mathcal{N} is a subset of $\{x_d = 0\}$ of dimension at most $(d-2)$. Consequently, $\hat{u}_n + \frac{\kappa_n}{\varepsilon_n} \Phi_n$ converges uniformly to h on compact sets in $B_1 \setminus \mathcal{N}$, which implies that $h = 0$ on $B_1' \setminus \mathcal{N}$. With \mathcal{N} having 0 capacity, this gives

$$\Delta h = 0 \text{ in } B_1^+, \text{ and } h = 0 \text{ on } B_1'.$$

Denote the $(2k+1)$ -order Taylor expansion of h at the origin (in B_1^+ , then evenly reflected to B_1^-) by $\sum_{\ell=0}^{2k+1} h_\ell$, with each h_ℓ being the ℓ -homogeneous part. Then we have

Lemma 3.4. *There is a universal constant C , such that for $r \in (0, 1/4)$ we have*

$$\|(\hat{u}_n)_r - h_{2k+1}\|_{\mathcal{L}^2(B_2)} \leq Cr(1 + |\log r|) + o(1)$$

and

$$\frac{\kappa_n}{\varepsilon_n} \leq Cr + o(1) \text{ as } n \rightarrow \infty,$$

where $(\hat{u}_n)_r(x) = \frac{1}{r^{2k+1}} \hat{u}_n(rx)$.

Proof. Throughout this proof, for a function w , we use w_r to denote its rescaling

$$w_r(x) = \frac{1}{r^{2k+1}} w(rx).$$

Firstly, with (2.5) and (3.3), we have

$$\int_{\partial B_1} |u - u_{\frac{1}{2}}| \leq \varepsilon o(1),$$

which implies, by maximum principle and the homogeneity of \bar{p} , that

$$(3.7) \quad |\hat{u} - \hat{u}_{\frac{1}{2}}| = o(1) \text{ in } B_{7/8}.$$

With (3.6) and regularity of the harmonic function h , we have

$$(3.8) \quad \|\hat{u} + \frac{\kappa}{\varepsilon} \Phi - \sum_{\ell=0}^{2k+1} h_\ell\|_{\mathcal{L}^2(B_{2r})} \leq Cr^{2k+2+\frac{d}{2}} + o(1).$$

A rescaling gives

$$\|\hat{u}_{\frac{1}{2}} + \frac{\kappa}{\varepsilon} \Phi_{\frac{1}{2}} - \sum_{\ell=0}^{2k+1} (h_\ell)_{\frac{1}{2}}\|_{\mathcal{L}^2(B_{4r})} \leq Cr^{2k+2+\frac{d}{2}} + o(1).$$

Combining these with (3.7), we get

$$\|\frac{\kappa}{\varepsilon} (\Phi - \Phi_{\frac{1}{2}}) + \sum [(h_\ell)_{\frac{1}{2}} - h_\ell]\|_{\mathcal{L}^2(B_{2r})} \leq Cr^{2k+2+\frac{d}{2}} + o(1).$$

That is,

$$\frac{\kappa}{\varepsilon} \frac{\log(2)}{d+4k} \|\varphi|x|^{2k+1}\|_{\mathcal{L}^2(B_{2r})} + \sum_{\ell=0}^{2k} (2^{2k+1-\ell} - 1) \|h_\ell\|_{\mathcal{L}^2(B_{2r})} \leq Cr^{2k+2+\frac{d}{2}} + o(1),$$

where we used the definition of Φ from (2.14), and the orthogonality of φ and h_ℓ in $\mathcal{L}^2(\mathbb{S}^{d-1})$ for $\ell \leq 2k$.

With Lemma 2.5, we can use the bound on the first term to get

$$\frac{\kappa}{\varepsilon} \leq Cr + o(1).$$

Similarly, the bound on each of the remaining terms gives

$$\|h_\ell\|_{\mathcal{L}^2(B_{2r})} \leq Cr^{2k+2+\frac{d}{2}} + o(1) \text{ for each } 0 \leq \ell \leq 2k.$$

Putting these into (3.8) gives

$$(3.9) \quad \|\hat{u}_r - h_{2k+1}\|_{\mathcal{L}^2(B_2)} \leq Cr(1 + |\log r|) + o(1).$$

This is the desired estimate. \square

As an immediate consequence of Lemma 3.4, we have

$$(3.10) \quad \|(u_n)_r - (\bar{p}_n + \varepsilon_n h_{2k+1})\|_{\mathcal{L}^2(B_2)} \leq \varepsilon_n [Cr(1 + |\log r|) + o(1)].$$

Lemma 3.5. *As $n \rightarrow \infty$, we have*

$$\|\overline{p_n + \varepsilon_n h_{2k+1}} - \bar{p}_n - \varepsilon_n h_{2k+1}\|_{\mathcal{L}^2(B_2)} = \varepsilon_n o(1).$$

Proof. In this proof, define $w = \frac{1}{\varepsilon}(\overline{p + \varepsilon h_{2k+1}} - \bar{p} - \varepsilon h_{2k+1})$.

Firstly, note that by the maximum principle for $\Delta_{\mathbb{S}^{d-1}} + \lambda(2k+1)$ in L_η^+ , we have $|\overline{p + \varepsilon h_{2k+1}} - \bar{p}| \leq C\varepsilon$ in L^+ . Thus $|w| \leq C$ in L^+ .

Meanwhile, in any compact subset of $B_1' \setminus \mathcal{N}$, the same argument as in Lemma 3.3 implies $w = 0$ for large n . We also have $w = 0$ along S_η^+ .

To summarize, w is a bounded solution to $(\Delta_{\mathbb{S}^{d-1}} + \lambda)w = 0$ in L_η^+ with $w = 0$ on S_η^+ and eventually vanishing in any compact subset of $B_1' \setminus \mathcal{N}$, where \mathcal{N} is of dimension at most $(d-2)$.

Note that w is even in the x_d -direction and vanishes outside L_η , this implies that $\|w\|_{\mathcal{L}^2(\mathbb{S}^{d-1})} = o(1)$, which gives the desired estimate. \square

If we define $p'_n \in \mathcal{P}_{2k+1}$ as

$$p'_n = p_n + \varepsilon_n h_{2k+1},$$

then $|p'_n - p_n| \leq C\varepsilon_n$.

With Lemma 3.4 and Lemma 3.5, we have

$$\kappa_{p'_n} \leq \kappa_{p_n} + \varepsilon_n o(1) \leq \varepsilon_n (Cr + o(1)).$$

By choosing r small, we have $\kappa_{p'_n} < \iota\varepsilon_n$ for all large n , where $\iota < \frac{1}{2}$ is a small universal constant to be chosen.

Combining (3.10) and Lemma 3.5, we have

$$\|(u_n)_r - \overline{p'_n}\|_{\mathcal{L}^2(B_2)} \leq \varepsilon_n [Cr(1 + |\log r|) + o(1)].$$

By choosing r small, depending on universal constants, such that $Cr(1 + |\log r|) < \iota/2$, we have

$$\|(u_n)_r - \overline{p'_n}\|_{\mathcal{L}^2(B_2)} < \iota\varepsilon_n$$

for all large n . Together with Lemma 2.6, this implies

$$\|(u_n)_r - \overline{p'_n}\|_{H^1(B_1)} < C\iota\varepsilon_n < \frac{1}{2}\varepsilon_n$$

if ι is small.

Consequently, we have

$$u_n \in \mathcal{S}_{2k+1}(p'_n, \frac{1}{2}\varepsilon_n, r),$$

contradicting (3.2).

This concludes the proof of Lemma 3.1.

4. THE DICHOTOMY AT A POINT WITH EVEN FREQUENCY

In this section, we establish a dichotomy similar to Lemma 3.1 but at a contact point with even frequency. We also explain how to get a log-epiperimetric inequality with a slightly improved exponent than the one in Colombo-Spolaor-Velichkov [CSV].

The ideas are similar to those in the previous section. We only sketch the proof.

This main lemma for this section is:

Lemma 4.1 (Dichotomy at a point with even frequency). *Given $k \in \mathbb{N}$, there are universal constants, $\tilde{\varepsilon}$, r_0 , c small and C big, such that*

If $u \in \mathcal{S}_{2k}(p, \varepsilon, 1)$ with $\varepsilon < \tilde{\varepsilon}$ and $1 \leq \|p\|_{\mathcal{L}^2(\mathbb{S}^{d-1})} \leq 2$, then we have the following dichotomy:

a) Either

$$W_{2k}(u; 1) - W_{2k}(u; r_0) \geq c\varepsilon^2$$

and

$$u \in \mathcal{S}_{2k}(p, C\varepsilon, r_0);$$

b) or

$$u \in \mathcal{S}_{2k}(p', \frac{1}{2}\varepsilon, r_0)$$

for some p' with

$$\|p' - p\|_{\mathcal{L}^2(\mathbb{S}^{d-1})} \leq C\varepsilon.$$

We prove this lemma by contradiction.

Suppose the lemma is not true, then we find a sequence (u_n, p_n) satisfying

$$1 \leq \|p_n\|_{\mathcal{L}^2(\mathbb{S}^{d-1})} \leq 2, \text{ and } u_n \in \mathcal{S}_{2k}(p_n, \varepsilon_n, 1) \text{ with } \varepsilon_n \rightarrow 0.$$

However,

$$(4.1) \quad W_{2k}(u_n; 1) - W_{2k}(u_n; \frac{1}{2}) < \frac{1}{n^2}\varepsilon_n^2,$$

and

$$(4.2) \quad u_n \notin \mathcal{S}_{2k}(p', \frac{1}{2}\varepsilon_n, r_0) \quad \forall p' \in \mathcal{P}_{2k} \text{ with } \|p' - p_n\|_{\mathcal{L}^2(\mathbb{S}^{d-1})} \leq C\varepsilon_n.$$

The constants r_0 and C will be chosen depending on universal constants.

Similar to the previous case, up to a subsequence, we have

$$p_n \rightarrow p_\infty \in \mathcal{P}_{2k} \text{ in } C^\infty(B_1).$$

With Lemma 2.3, we have $p_n \geq -v_n \geq -C\varepsilon_n^{\frac{2}{d-1}}$ on B'_1 . Thus

$$p_\infty \geq 0 \text{ on } B'_1.$$

The set where p_∞ vanishes on B'_1 , $\mathcal{N} = \{p_\infty = 0\}'$, has dimension at most $(d-2)$.

Define normalized solutions \hat{u}_n as in (3.5), we have, up to a subsequence,

$$\|\hat{u}_n + \frac{\kappa_n}{\varepsilon_n}\Phi_n - h\|_{\mathcal{L}^2(B_{7/8})} = o(1) \text{ as } n \rightarrow \infty$$

for some h satisfying

$$\Delta h = 0 \text{ in } B_1.$$

With similar ideas as in Lemma 3.4, we can rule out lower order terms in the Taylor polynomial of h at 0 and obtain for $r \in (0, 1/4)$,

$$\|(u_n)_r - (\bar{p}_n + \varepsilon_n h_{2k})\|_{\mathcal{L}^2(B_2)} \leq \varepsilon_n [Cr(1 + |\log r|) + o(1)].$$

If we choose r small, then $p'_n = p_n + \varepsilon_n h_{2k} \in \mathcal{P}_{2k}$ satisfies $|p'_n - p_n| \leq C\varepsilon_n$, and

$$\|(u_n)_r - \bar{p}'_n\|_{\mathcal{L}^2(B_2)} < \iota\varepsilon_n \text{ and } \kappa_{p'_n} \leq \iota\varepsilon_n$$

for large n , where ι is a universally small constant.

An application of Lemma 2.6 again gives

$$u_n \in \mathcal{S}_{2k}(p'_n, \frac{1}{2}\varepsilon_n, r)$$

if ι is small, which contradicts (4.2).

This completes the proof for Lemma 4.1.

Remark 4.1. We sketch how similar ideas lead to a log-epiperimetric inequality for the $2k$ -Weiss energy functional. We get an improved exponent than the one currently known in the literature.

To be precise, let w be a $2k$ -homogeneous function satisfying

$$w \geq 0 \text{ on } \{x_d = 0\},$$

and

$$\|w\|_{\mathcal{L}^2(\mathbb{S}^{d-1})} \leq 1, \quad |W_{2k}(w; 1)| \leq 1,$$

then we will show

$$(4.3) \quad W_{2k}(w; 1) - W_{2k}(u; 1) \geq cW_{2k}(w; 1)^{1 + \frac{d-3}{d+1}},$$

where u is the solution to (1.1) with $u|_{\mathbb{S}^{d-1}} = w$, and c is a universal constant.

A similar result is known in [CSV] with the exponent on the right-hand side as $1 + \frac{d-2}{d}$.

It suffices to prove (4.3) under the assumption $\|w\|_{\mathcal{L}^2(\mathbb{S}^{d-1})} = 1$. For such w , choose $p \in \mathcal{P}_{2k}$ that minimizes $\delta(w, p)$ from (2.15).

With $W(p; 1) = 0$, $\Delta p = 0$ and the homogeneity of p , we have

$$W(w; 1) \leq \int_{B_1} |\nabla w - \nabla p|^2 \leq C\delta(w, p)^2 + C\|\bar{p} - p\|_{H^1(B_1)}^2.$$

With homogeneity and harmonicity of p , we also have

$$W(\bar{p}; 1) = W(\bar{p} - p; 1) = C\left[\int_{L_\eta} |\nabla_{\mathbb{S}^{d-1}}(\bar{p} - p)|^2 - \lambda(2k)(\bar{p} - p)^2\right],$$

where the definitions of L_η and $\lambda(2k)$ are given at the beginning of Subsection 2.2. By making η smaller, if necessary, we have $\int_{L_\eta} |\nabla_{\mathbb{S}^{d-1}}(\bar{p} - p)|^2 - \lambda(2k)(\bar{p} - p)^2 \sim \|\bar{p} - p\|_{H^1(B_1)}^2$. Therefore,

$$W(w; 1) \leq C\delta(w, p)^{1 + \frac{2}{d-1}},$$

where we used (2.16).

With $u = w$ on \mathbb{S}^{d-1} and $w \geq 0$ on $\{x_d = 0\}$, we have

$$W(w; 1) - W(u; 1) \geq \int_{B_1} |\nabla(w - u)|^2.$$

Therefore, it suffices to show that for $\delta(w, p)$ small, we have

$$(4.4) \quad \int_{B_1} |\nabla(w - u)|^2 \geq c\delta(w, p)^2$$

for some universal constant c .

Suppose, on the contrary, this fails. Then we find a sequence (w_n, u_n, p_n) as described above with $\delta_n = \delta(w_n, p_n) \rightarrow 0$ but

$$(4.5) \quad \int_{B_1} |\nabla(w_n - u_n)|^2 \leq \frac{1}{n^2} \delta_n^2.$$

Similar ideas as in the proof for Lemma 4.1 gives, for large n ,

$$\delta(u_n, p'_n) < \frac{1}{2} \delta_n$$

for some $p'_n \in \mathcal{P}_{2k}$, where (4.5) can be used in place of (4.1) to control terms with lower homogeneities. With (4.5), this gives $\delta(w_n, p'_n) < \delta_n(w_n, p_n)$, contradicting the minimizing property of p_n .

5. CONVERGENCE RATE TO THE BLOW-UP PROFILE

In this final section, we prove our main result Theorem 1.1. Our result on stratification of contact points with integer frequencies, Theorem 1.2, follows with Whitney's extension theorem and the implicit function theorem. See, for instance, [GP].

We first give a technical lemma about sequences. We will apply this to the sequences of Weiss energy and errors in approximations at different scales.

Lemma 5.1. *Let (w_n) and (e_n) be two sequences of nonnegative real numbers with $e_0 \leq 1$. Suppose that for some constants, A big, a small and $\gamma \in (0, 1]$, we have*

$$w_{n+1} \leq A e_n^{1+\gamma},$$

and the following dichotomy:

- either $w_{n+1} \leq w_n - a e_n^2$ and $e_{n+1} = A e_n$;
- or $w_{n+1} \leq w_n$ and $e_{n+1} = \frac{1}{2} e_n$.

Then

$$\sum e_n < \sigma(e_0) \rightarrow 0 \quad \text{as } e_0 \rightarrow 0,$$

and

$$(5.1) \quad \sum_{n \geq N} e_n \leq C(1 - c)^N \quad \text{if } \gamma = 1;$$

and

$$(5.2) \quad \sum_{n \geq N} e_n \leq C N^{\frac{-\gamma}{1-\gamma}} \quad \text{if } \gamma \in (0, 1).$$

Here $c \in (0, 1)$ and C are constants depending only on A , a and γ .

Proof. Define a new sequence

$$\alpha_n := w_n + \mu e_n^2.$$

If $\mu > 0$ is small enough, then we have

$$(5.3) \quad \alpha_{n+1} \leq \alpha_n - c \alpha_n^{\frac{2}{1+\gamma}},$$

and $\alpha_1 \leq C e_0^{1+\gamma}$.

For $\gamma = 1$, we have $\alpha_n \leq (1-c)^{n-1} \alpha_1$, which gives the desired estimate for this case.

For $\gamma \in (0, 1)$, from (5.3) we have that for all $n \geq 1$,

$$\alpha_n \leq C(n+M)^{-\frac{1+\gamma}{1-\gamma}},$$

with $M \rightarrow \infty$ as $e_0 \rightarrow 0$, and the first estimate follows. Meanwhile, by our definition of α_n , we have $e_n^2 \leq C(\alpha_n - \alpha_{n+1})$. Thus

$$\begin{aligned} \sum_{n=N}^{2N} e_n &\leq C \sum_N^{2N} (\alpha_n - \alpha_{n+1})^{1/2} \\ &\leq CN^{1/2} \left[\sum_N^{2N} (\alpha_n - \alpha_{n+1}) \right]^{1/2} \\ &\leq CN^{1/2} \alpha_N^{1/2} \\ &\leq CN^{-\frac{\gamma}{1-\gamma}}. \end{aligned}$$

This gives the desired estimate for $\gamma \in (0, 1)$. \square

Now we give the proof of our main result.

Proof of Theorem 1.1. Suppose that $0 \in \Lambda_m(u)$ for some $m \in \mathbb{N}$, then up to an initial rescaling, we have

$$u \in \mathcal{S}_m(p, \varepsilon, 1), \quad \|p\|_{L^2(B_1)} = 3/2,$$

for some $\varepsilon < \tilde{\varepsilon}$. Here $\tilde{\varepsilon}$ is the constant from Lemma 1.1, and the solution class \mathcal{S}_m is from Definition 2.2.

As the initial set up, let $p_0 = p$, $\rho_0 = 1$, $e_0 = \varepsilon$, and $w_0 = W_m(u; 1)$.

Suppose that we have found p_n , e_n , w_n small, and $\rho_n \in (0, 1)$ such that $u \in \mathcal{S}_m(p_n, e_n, \rho_n)$ with $e_n < \tilde{\varepsilon}$. Then we apply Lemma 1.1. If possibility a) happens in the dichotomy, we let $p_{n+1} = p_n$, $e_{n+1} = C e_n$. If possibility b) happens, we let $p_{n+1} = p'$ and $e_{n+1} = \frac{1}{2} e_n$. In both cases, we let $\rho_{n+1} = r_0 \rho_n$ and $w_{n+1} = W_m(u; \rho_{n+1})$.

By Lemma 1.1 and Lemma 2.8, the sequences (w_n) and (e_n) satisfy the assumptions in Lemma 5.1, with $\gamma = 1$ if m is odd, and $\gamma = \frac{2}{d-1}$ if m is even. In particular, we have $\sum e_n < \tilde{\varepsilon}$ along the sequence if e_0 is chosen small enough. Consequently, Lemma 1.1 can be applied indefinitely.

Now note that $\|p_{n+1} - p_n\|_{L^2(B_1)} \leq C e_n$. The summability of (e_n) implies the convergence of p_n to some limit p_∞ .

If we denote by u_r the rescaled solution $u_r(x) = \frac{1}{r^m} u(rx)$. When m is odd, we use (5.1) to get $\|u_{r_0^n} - p_\infty\|_{H^1(B_1)} \leq C(1-c)^n$. This gives the estimate in (1.5). When m is even, we use (5.2) to get $\|u_{r_0^n} - p_\infty\|_{H^1(B_1)} \leq C n^{-\frac{2}{d-3}}$. This gives the estimate in (1.6). \square

The stratification in Theorem 1.2 follows by the same strategy as in Garofalo-Petrosyan [GP] or Colombo-Spolaor-Velichkov [CSV]. In the following remark, we point out that $\Lambda_1(u)$ and $\Lambda_3^0(u)$ always lie in the interior of the contact set.

Remark 5.1. Suppose $0 \in \Lambda_{2k+1}(u)$, then there is $p \in \mathcal{P}_{2k+1}^+$ such that

$$u(x) = p(x) + O(|x|^{2k+1+\alpha})$$

as $x \rightarrow 0$.

If $0 \in \Lambda_1(u)$, then p is a positive multiple of $-|x_d|$.

If $0 \in \Lambda_3(u)$, we have

$$p(x', x_d) = -|x_d|(p_1(x') + x_d^2 p_2(x', x_d)),$$

where p_1 is a 2-homogeneous polynomial with $p_1 \geq 0$ on $\{x_d = 0\}$. The zero stratum of $\Lambda_3(u)$ is defined as those points where p_1 depends on all $(d-1)$ -variables. This implies $p_1 > 0$ on $(\mathbb{S}^{d-1})'$.

Consequently, if $0 \in \Lambda_1(u)$ or $\Lambda_3^0(u)$, then $\frac{\partial}{\partial x_d} p < 0$ on $(\mathbb{S}^{d-1})'$. With Lemma 2.9, we have $u = 0$ in B'_r for some $r > 0$.

REFERENCES

- [AC] Athanasopoulos, I.; Caffrelli, L.A. Optimal regularity of lower dimensional obstacle problems. *Zap. Nauchn. Sem. S.-Petersburg. Otdel. Mat. Inst. Steklov.* 310 (2004), Kraev. Zadachi Mat. Fiz. i Smezh. Vopr. Teor. Funkts. 35, 49-66.
- [ACS] Athanasopoulos, I.; Caffrelli, L.A.; Salsa, S. The structure of the free boundary for lower dimensional obstacle problems. *Amer. J. Math.* 130 (2008), no. 2, 485-498.
- [CSV] Colombo, M.; Spolaor, L.; Velichkov, B. Direct epiperimetric inequalities for the thin obstacle problem and applications. *Comm. Pure Appl. Math.* 73 (2020), no. 2, 384-420.
- [D] De Silva, D. Free boundary regularity for a problem with right hand side. *Interfaces Free Bound.* 13 (2011), no. 2, 223-238.
- [DS] De Silva, D.; Savin, O. Boundary Harnack estimates in slip domains and applications to thin free boundary problems. *Rev. Mat. Iberoam.* 32 (2016), no. 3, 891-912.
- [Fe] Fernández-Real, X. The thin obstacle problem: a survey. *Publicacions Matemàtiques*, to appear.
- [FeR] Fernández-Real, X.; Ros-Oton, X. Free boundary regularity for almost every solution to the Signorini problem. *Arch. Ration. Mech. Anal.* to appear.
- [FRS] Figalli, A.; Ros-Oton, X.; Serra, J. Generic regularity of free boundaries for the obstacle problem. *Publ. Math. Inst. Hautes Études Sci.* 132 (2020), 181-292.
- [FoS1] Focardi, M.; Spadaro, E. On the measure and structure of the free boundary of the lower dimensional obstacle problem. *Arch. Ration. Mech. Anal.* 230 (2018), no. 1, 125-184.
- [FoS2] Focardi, M.; Spadaro, E. Correction to : On the measure and structure of the free boundary of the lower dimensional obstacle problem. *Arch. Ration. Mech. Anal.* 230 (2018), no. 2, 783-784.
- [GP] Garofalo, N.; Petrosyan, A. Some new monotonicity formulas and the singular set in the lower dimensional obstacle problem. *Invent. Math.* 177 (2009), no. 2, 415-461.
- [KPS] Koch, H.; Petrosyan, A.; Shi, W. Higher regularity of the free boundary in the elliptic Signorini problem. *Nonlinear Anal.* 126 (2015), 3-44.
- [PSU] Petrosyan, A.; Shahgholian, H.; Uraltseva, N. Regularity of free boundaries in obstacle-type problems. Graduate Studies in Mathematics, 136. *American Mathematical Society, Providence, RI*, 2012.
- [R] Richardson, D. Variational problems with thin obstacles. *Thesis-The University of British Columbia* 1978.
- [SY1] Savin, O.; Yu, H. Regularity of the singular set in the fully nonlinear obstacle problem. *J. Euro. Math. Soc.* to appear.
- [SY2] Savin, O.; Yu, H. On the fine regularity of the singular set in the nonlinear obstacle problem. *Preprint*: arXiv:2101.11759.
- [SY3] Savin, O.; Yu, H. Free boundary regularity in the triple membrane problem. *Preprint*: arXiv:2002.10628.
- [U] Uraltseva, N. On the regularity of solutions of variational inequalities. *Uspekhi Mat. Nauk* 42 (1987), no. 6, 151-174.

- [W] Weiss, G.S. A homogeneity improvement approach to the obstacle problem. *Invent. Math.* 138 (1999), no. 1, 23-50.

DEPARTMENT OF MATHEMATICS, COLUMBIA UNIVERSITY, NEW YORK, USA
Email address: `savin@math.columbia.edu`

DEPARTMENT OF MATHEMATICS, COLUMBIA UNIVERSITY, NEW YORK, USA
Email address: `huiyu@math.columbia.edu`