

On a Conjecture Concerning the Approximates of Complete Elliptic Integral of the First Kind by Inverse Hyperbolic Tangent

Song-Liang Qiu*, Qi Bao, Xiao-Yan Ma, Hong-Biao Jiang

Abstract: Let \mathcal{K} be the complete elliptic integral of the first kind. In this paper, the authors prove that the function $r \mapsto r^{-2}\{[\log(2\mathcal{K}(r)/\pi)]/\log(\operatorname{arth}r/r) - 3/4\}$ is strictly increasing from $(0, 1)$ onto $(1/320, 1/4)$, so that $[(\operatorname{arth}r)/r]^{3/4+r^2/320} < 2\mathcal{K}(r)/\pi < [(\operatorname{arth}r)/r]^{3/4+r^2/4}$ for $r \in (0, 1)$, in which all the coefficients of the exponents of the two bounds are best possible, thus proving a conjecture raised by Alzer and Qiu to be true, and giving better bounds of $\mathcal{K}(r)$ than those they conjectured and put in an open problem. Some other analytic properties of the complete elliptic integrals, including other kind of approximates for $\mathcal{K}(r)$, are obtained, too.

Keywords: The complete elliptic integrals; inverse hyperbolic tangent; monotonicity; convexity; inequality

MSC (2010): 33C75, 33E05, 33F05

1 Introduction

Throughout this paper, we let \mathbb{N} (arth) denote the set of positive integers (the inverse hyperbolic tangent, respectively), $\mathbb{N}_0 = \mathbb{N} \cup \{0\}$, and let $r' = \sqrt{1-r^2}$ for each $r \in [0, 1]$. For $r \in (0, 1)$, the complete elliptic integrals of the first and second kinds are defined as

$$\begin{cases} \mathcal{K} = \mathcal{K}(r) = \int_0^{\pi/2} (1 - r^2 \sin^2 t)^{-1/2} dt, \\ \mathcal{K}' = \mathcal{K}'(r) = \mathcal{K}(r'), \\ \mathcal{K}(0) = \pi/2, \mathcal{K}(1) = \infty, \end{cases} \quad (1.1)$$

and

$$\begin{cases} \mathcal{E} = \mathcal{E}(r) = \int_0^{\pi/2} (1 - r^2 \sin^2 t)^{1/2} dt, \\ \mathcal{E}' = \mathcal{E}'(r) = \mathcal{E}(r'), \\ \mathcal{E}(0) = \pi/2, \mathcal{E}(1) = 1, \end{cases} \quad (1.2)$$

respectively, which have wide applications and are the particular cases of the Gaussian hypergeometric function

$$F(a, b; c; x) = {}_2F_1(a, b; c; x) = \sum_{n=0}^{\infty} \frac{(a)_n (b)_n}{(c)_n n!} x^n \quad (|x| < 1) \quad (1.3)$$

for complex numbers a, b and c with $c \neq 0, -1, -2, \dots$, where $(a)_n = a(a+1)(a+2) \cdots (a+n-1) = \Gamma(a+n)/\Gamma(a)$ for $n \in \mathbb{N}$, and $(a)_0 = 1$, while $\Gamma(x) = \int_0^{\infty} t^{x-1} e^{-t} dt$ is the Euler gamma function. As a matter of fact, it is well known that

$$\mathcal{K}(r) = \frac{\pi}{2} F\left(\frac{1}{2}, \frac{1}{2}; 1; r^2\right) \text{ and } \mathcal{E}(r) = \frac{\pi}{2} F\left(-\frac{1}{2}, \frac{1}{2}; 1; r^2\right) \quad (1.4)$$

(cf. [4, 17.3.9–17.3.10]). Moreover, by [4, 15.1.4],

$$\frac{\operatorname{arth}r}{r} = F\left(\frac{1}{2}, 1; \frac{3}{2}; r^2\right) = \sum_{n=0}^{\infty} \frac{1}{2n+1} r^{2n}. \quad (1.5)$$

*Corresponding author.

As we known, numerous properties of the complete elliptic integrals have been obtained (cf. [1–13, 15, 16, 18–21, 23, 25] and bibliographies therein).

One of the topics of studying the properties of the complete elliptic integrals is to approximate $\mathcal{K}(r)$ by means of the function $r \mapsto (\operatorname{arth}r)/r$. Following this topic, G.D. Anderson et al. proved in [8, Theorem 3.10] that the function $r \mapsto r\mathcal{K}(r)^2/\operatorname{arth}r$ ($r \mapsto r\mathcal{K}(r)/\operatorname{arth}r$) is strictly increasing (decreasing) from $(0, 1)$ onto $(\pi^2/4, \infty)$ ($(1, \pi/2)$, respectively), so that

$$\max \left\{ \left(\frac{\operatorname{arth}r}{r} \right)^{1/2}, \frac{2}{\pi} \frac{\operatorname{arth}r}{r} \right\} < \frac{2}{\pi} \mathcal{K}(r) < \frac{\operatorname{arth}r}{r} \quad (1.6)$$

for $r \in (0, 1)$. In [3, Theorem 18], Alzer and Qiu improved (1.6) to the following double inequality

$$\frac{\pi}{2} \left(\frac{\operatorname{arth}r}{r} \right)^{\alpha_1} < \mathcal{K}(r) < \frac{\pi}{2} \left(\frac{\operatorname{arth}r}{r} \right)^{\beta_1} \quad (1.7)$$

with best possible exponents $\alpha_1 = 3/4$ and $\beta_1 = 1$, and in [3, Section 4], they posed the following

Open Problem 1.1. *What is the largest (smallest) constant α^* (β^* , respectively) such that the double inequality*

$$\frac{\pi}{2} \left(\frac{\operatorname{arth}r}{r} \right)^{3/4+\alpha^*r} < \mathcal{K}(r) < \frac{\pi}{2} \left(\frac{\operatorname{arth}r}{r} \right)^{3/4+\beta^*r} \quad (1.8)$$

holds for all $r \in (0, 1)$?

The double inequality (1.8) is obviously equivalent to

$$\alpha^* < \frac{1}{r} \left[G(r) - \frac{3}{4} \right] < \beta^*, \quad \text{where } G(r) = \frac{\log(2\mathcal{K}(r)/\pi)}{\log(\operatorname{arth}r/r)} \quad (1.9)$$

for $r \in (0, 1)$. In [3, Section 4], it was shown that the best possible constants $\alpha^* = 0$ and $\beta^* \geq 1/4$, and the following conjecture was put forward.

Conjecture 1.2. *The function G in (1.9) is strictly increasing and convex from $(0, 1)$ onto $(3/4, 1)$.*

If Conjecture 1.2 is true, then the best possible constants α^* and β^* in (1.8) are 0 and 1/4, respectively.

In 2011, Y.-M. Chu et al. proved in [14] that (1.8) holds for all $r \in (0, 1)$ with the best possible constant $\beta^* = 1/4$. In [17], Pál A. Kupán and Róbert Szász studied Open Problem 1.1 and Conjecture 1.2, and proved that for $r \in (0, 1)$, the following double inequality

$$\frac{\pi}{2} \left(\frac{\operatorname{arth}r}{r} \right)^{3/4+r^4/200} < \mathcal{K}(r) < \frac{\pi}{2} \left(\frac{\operatorname{arth}r}{r} \right)^{3/4+r^2/4} \quad (1.10)$$

holds, which improves the result obtained in [14]. In [17, Section 1], it was stated that “It seems very difficult to prove the conjecture regarding the monotonicity of G ”. Here “the conjecture” means Conjecture 1.2.

Based on the results above-mentioned and our computation, the following question is natural and more reasonable than Open Problem 1.1.

Question 1.3. *What are the best possible constants α and β such that for $r \in (0, 1)$,*

$$\frac{\pi}{2} \left(\frac{\operatorname{arth}r}{r} \right)^{3/4+\alpha r^2} < \mathcal{K}(r) < \frac{\pi}{2} \left(\frac{\operatorname{arth}r}{r} \right)^{3/4+\beta r^2} ? \quad (1.11)$$

Let α and β be the best possible constants such that (1.11) holds for $r \in (0, 1)$, G as in (1.9), and let

$$f(r) = \frac{1}{r^2} \left[G(r) - \frac{3}{4} \right] \quad \text{for } r \in (0, 1). \quad (1.12)$$

Then it is easy to obtain the limiting values $f(0^+) = 1/320$ and $f(1^-) = 1/4$ (see Theorem 1.7(3)). It is clear that (1.11) is equivalent to

$$\alpha = \inf_{0 < r < 1} f(r) < f(r) < \sup_{0 < r < 1} f(r) = \beta \quad (1.13)$$

for $r \in (0, 1)$. Observe that (1.8), with the best possible constants $\alpha^* = 0$ and $\beta^* = 1/4$, is weaker than (1.11) if $0 < \alpha < \beta \leq 1/4$ in (1.13). From this point of view, the convexity of G in Conjecture 1.2 is no longer so important, if our following conjecture is true.

Conjecture 1.4. *The function f defined by (1.12) is strictly increasing from $(0, 1)$ onto $(1/320, 1/4)$.*

Clearly, the truth of Conjecture 1.4 implies that the function G in Conjecture 1.2 is strictly increasing from $(0, 1)$ onto $(3/4, 1)$, and gives the answer to Question 1.3 (see Theorem 1.7(3)). Naturally, the proof of Conjecture 1.4 is much more difficult than that of the monotonicity of G .

In addition, based on [8, Theorem 3.10], the following question is natural.

Question 1.5. *Is the function $r \mapsto r\mathcal{K}(r)/\operatorname{arth}r$ concave on $(0, 1)$?*

The main purpose of this paper is to prove that Conjecture 1.4 is true, to give the complete answers to Questions 1.3 and 1.5 (see Theorems 1.6–1.8 stated at the end of this section), and to present some other related properties of the complete elliptic integrals of the first and second kinds, including the sharp approximates of $\mathcal{K}(r)$ by certain combinations in terms of $(\operatorname{arth}r)/r$ and polynomials (see Theorems 3.1–3.2 proved in Section 3), by showing the monotonicity and convexity properties of certain combinations defined in terms of $\mathcal{K}(r)$, $\mathcal{E}(r)$ and $\operatorname{arth}r$. In order to prove these results, we shall establish several lemmas in Section 2. We shall put forward three conjectures and an open problem in Section 5, for further related studies.

In the sequel, we always let $a_n = [(1/2)_n/n!]^2$ and $\tilde{a}_n = 3/[4(2n+1)] - a_n$ for $n \in \mathbb{N}_0$, and let

$$F_0(x) = F\left(\frac{1}{2}, \frac{1}{2}; 1; x\right), \quad F_1(x) = F\left(\frac{1}{2}, 1; \frac{3}{2}; x\right), \quad (1.14)$$

$$G_0(x) = F\left(\frac{1}{2}, \frac{1}{2}; 2; x\right) \quad \text{and} \quad G_1(x) = F\left(\frac{1}{2}, 1; \frac{5}{2}; x\right) \quad (1.15)$$

for $x \in (0, 1)$. By [4, 15.2.1&15.3.3], we have the following relations

$$F'_0(x) = \frac{G_0(x)}{4(1-x)} \quad \text{and} \quad F'_1(x) = \frac{G_1(x)}{3(1-x)}. \quad (1.16)$$

Now we state some of our main results, which will be proved in Section 4.

Theorem 1.6. *For each $c \in (0, \infty)$, define the function g on $(0, 1)$ by*

$$g(r) = \left(\frac{3}{4} + cr^2\right) \log \frac{\operatorname{arth}r}{r} - \log \frac{2\mathcal{K}(r)}{\pi}.$$

Then g is strictly increasing (decreasing) on $(0, 1)$ if and only if $c \geq 1/4$ ($c \leq 1/320$, respectively), with

$$g(0^+) = 0 \quad \text{and} \quad g(1^-) = \begin{cases} \log(\pi/2), & \text{if } c = 1/4, \\ \infty, & \text{if } c > 1/4, \\ -\infty, & \text{if } c < 1/4. \end{cases}$$

Moreover, if $c \geq 1/4$ ($c \leq 1/320$), then g is convex (concave, respectively) on $(0, 1)$.

Theorem 1.7. (1) *The function*

$$g_1(r) \equiv \frac{1}{r^2} \left[\frac{3+r^2}{4} \log \frac{\operatorname{arth}r}{r} - \log \frac{2\mathcal{K}(r)}{\pi} \right]$$

is strictly increasing from $(0, 1)$ onto $(0, \log(\pi/2))$, while the function

$$g_2(r) \equiv \frac{g_1(r)}{\log((\operatorname{arth}r)/r)} = \frac{1}{r^2 \log((\operatorname{arth}r)/r)} \left[\frac{3+r^2}{4} \log \frac{\operatorname{arth}r}{r} - \log \frac{2\mathcal{K}(r)}{\pi} \right]$$

is strictly decreasing from $(0, 1)$ onto $(0, 79/320)$.

(2) Define the functions g_3 and g_4 on $(0, 1)$ by

$$g_3(r) = \frac{1}{r^2 \log((\operatorname{arth}r)/r)} \left[\log \frac{2\mathcal{K}(r)}{\pi} - \left(\frac{3}{4} + \frac{1}{320}r^2 \right) \log \frac{\operatorname{arth}r}{r} \right],$$

$$g_4(r) = \frac{1}{r^4} \left[\log \frac{2\mathcal{K}(r)}{\pi} - \left(\frac{3}{4} + \frac{1}{320}r^2 \right) \log \frac{\operatorname{arth}r}{r} \right].$$

Then g_3 and g_4 are both strictly increasing on $(0, 1)$, with ranges $(0, 79/320)$ and $(0, \infty)$, respectively.

(3) The function f defined by (1.12) is strictly increasing from $(0, 1)$ onto $(1/320, 1/4)$, namely, Conjecture 1.4 is true. In particular, the function

$$G(r) \equiv \frac{\log(2\mathcal{K}(r)/\pi)}{\log((\operatorname{arth}r)/r)}$$

defined as in (1.9) is strictly increasing from $(0, 1)$ onto $(3/4, 1)$, and the double inequality

$$\frac{\pi}{2} \left(\frac{\operatorname{arth}r}{r} \right)^{3/4+\alpha r^2} < \mathcal{K}(r) < \frac{\pi}{2} \left(\frac{\operatorname{arth}r}{r} \right)^{3/4+\beta r^2} \quad (1.17)$$

holds for $r \in (0, 1)$ if and only if $\alpha \leq 1/320$ and $\beta \geq 1/4$. Moreover, for $r \in (0, 1)$,

$$\left(\frac{\pi}{2} \right)^{1-r^2} \left(\frac{\operatorname{arth}r}{r} \right)^{3/4+r^2/4} < \mathcal{K}(r) < \frac{\pi}{2} \left(\frac{\operatorname{arth}r}{r} \right)^{3/4+r^2/4}. \quad (1.18)$$

Theorem 1.8. (1) The function $h_1(r) \equiv r\mathcal{K}(r)/\operatorname{arth}r$ is strictly decreasing and concave from $(0, 1)$ onto $(1, \pi/2)$.

(2) For each $n \in \mathbb{N}$, define the function h_2 on $(0, 1)$ by

$$h_2(r) = \frac{1}{r^{2(n+1)}} \left[\frac{3\operatorname{arth}r}{4r} - \frac{2}{\pi} \mathcal{K}(r) - \sum_{k=0}^n \tilde{a}_k r^{2k} \right].$$

Then the coefficients of the Maclaurin series of h_2 are all positive. In particular, h_2 is strictly absolutely monotone on $(0, 1)$, namely, $h_2^{(k)}(r) > 0$ for all $k \in \mathbb{N}_0$ and $r \in (0, 1)$, with $h_2(0^+) = \tilde{a}_{n+1} > 0$ and $h_2(1^-) = \infty$.

(3) The function $h_3(r) \equiv rh_2(r)/\operatorname{arth}r$ is strictly increasing from $(0, 1)$ onto $(\tilde{a}_{n+1}, (3/4) - 2/\pi)$.

(4) The function

$$h_4(r) \equiv \frac{1}{r^6} \left[\frac{\pi}{2} \left(1 - \frac{1}{12}r^2 - \frac{91}{2880}r^4 \right) - \frac{r\mathcal{K}(r)}{\operatorname{arth}r} \right]$$

is strictly increasing from $(0, 1)$ onto $(871\pi/96768, 2549\pi/5760 - 1)$.

(5) For each $n \in \mathbb{N}$, and for all $r \in (0, 1)$,

$$\frac{1}{4} + \frac{3\operatorname{arth}r}{4r} - \sum_{k=2}^{n+1} \tilde{a}_k r^{2k} - P_1(r)r^{2n+2} < \frac{2}{\pi} \mathcal{K}(r) < \frac{1}{4} + \frac{3\operatorname{arth}r}{4r} - \sum_{k=2}^{n+1} \tilde{a}_k r^{2k} - P_2(r)r^{2n+2}, \quad (1.19)$$

$$\left[P_3(r) - \alpha r^6 \right] \frac{\operatorname{arth}r}{r} < \frac{2}{\pi} \mathcal{K}(r) < \left[P_3(r) - \beta r^6 \right] \frac{\operatorname{arth}r}{r} \quad (1.20)$$

and

$$\frac{1}{4} + \frac{3}{4} (1 - \delta r^4) \frac{\operatorname{arth}r}{r} < \frac{2}{\pi} \mathcal{K}(r) < \frac{1}{4} + \frac{3}{4} (1 - \eta r^4) \frac{\operatorname{arth}r}{r}, \quad (1.21)$$

where

$$P_1(r) = \frac{3\pi - 8 \operatorname{arth}r}{4\pi} \frac{1}{r} - \tilde{a}_{n+1}, \quad P_2(r) = \tilde{a}_{n+1} \left(\frac{\operatorname{arth}r}{r} - 1 \right),$$

$$P_3(r) = 1 - \frac{1}{12}r^2 - \frac{91}{2880}r^4, \quad \alpha = \frac{2549}{2880} - \frac{2}{\pi} = 0.2484496 \dots,$$

$\beta = 871/48384 = 0.0180018 \dots$, $\delta = 1 - 8/(3\pi) = 0.151173 \dots$ and $\eta = 1/80 = 0.0125$. Moreover, the constants α , β , δ and η in (1.20) and (1.21) are all best possible.

2 Preliminaries

In order to prove our theorems stated in Section 1, we shall prove four lemmas in this section. Firstly, let us recall the following well-known formulas: For $r \in (0, 1)$,

$$\frac{d\mathcal{K}}{dr} = \frac{\mathcal{E}(r) - r'^2 \mathcal{K}(r)}{rr'^2}, \quad \frac{d\mathcal{E}}{dr} = \frac{\mathcal{E}(r) - \mathcal{K}(r)}{r}, \quad (2.1)$$

$$\mathcal{K}(r) = \log \frac{4}{r'} + O(r'^2 \log r') \quad (r \rightarrow 1) \quad \text{and} \quad (2.2)$$

$$\frac{\mathcal{E}(r) - r'^2 \mathcal{K}(r)}{r^2} = \frac{\pi}{4} \sum_{n=0}^{\infty} \frac{a_n}{n+1} r^{2n} \quad (2.3)$$

(see [4, 15.3.10], [6, Theorem 4.1] and [6, (5.3)], respectively), which will be frequently applied later.

Our first lemma presents several properties of the elementary function arth and the special function \mathcal{K} , most of which will be applied in the proofs of our main results.

- Lemma 2.1.** (1) The function $f_1(r) \equiv (r - r'^2 \text{arth}r)/r^3$ is strictly increasing and convex from $(0, 1)$ onto $(2/3, 1)$.
(2) The function $f_2(r) \equiv (r - r'^2 \text{arth}r)/(r^2 r'^2 \text{arth}r)$ is strictly increasing from $(0, 1)$ onto $(2/3, \infty)$.
(3) The function $f_3(r) \equiv (r - r'^2 \text{arth}r)/[(1 + r^2) \text{arth}r - r]$ is strictly decreasing from $(0, 1)$ onto $(0, 1/2)$.
(4) The function $f_4(r) \equiv [(1 + r^2) \text{arth}r - r]/[r^3 \mathcal{K}(r)]$ is strictly increasing from $(0, 1)$ onto $(8/(3\pi), 2)$.
(5) The function $f_5(r) \equiv [1 + (r'^2 \text{arth}r)/r]/r'$ is strictly increasing and convex from $(0, 1)$ onto $(2, \infty)$.
(6) The function $f_6(r) \equiv (r/r' - r' \text{arth}r)/(\text{arth}r - r)$ is strictly increasing from $(0, 1)$ onto $(2, \infty)$.
(7) The function $f_7(r) \equiv [\log((\text{arth}r)/r)]/r^2$ is strictly increasing from $(0, 1)$ onto $(1/3, \infty)$.
(8) The functions

$$f_8(r) \equiv \frac{r'^2 \text{arth}r}{r - r'^2 \text{arth}r} \log \frac{\text{arth}r}{r} \quad \text{and} \quad f_9(r) \equiv \frac{r'^2 \mathcal{K}(r)}{\mathcal{E}(r) - r'^2 \mathcal{K}(r)} \log \frac{2\mathcal{K}(r)}{\pi}$$

are both strictly decreasing from $(0, 1)$ onto $(0, 1/2)$.

(9) For all $r \in (0, 1)$,

$$f_{10}(r) \equiv \frac{1}{r^4} \left[72 - 126r^2 + 19r^4 - (72 - 150r^2 - 253r^4 + 167r^6) \frac{\text{arth}r}{r} \right] > \frac{25688}{105}, \quad (2.4)$$

$$f_{11}(r) \equiv \frac{1}{r^2} \left[(13 - 9r^2) \frac{\text{arth}r}{r} - 13 + 6r^2 \right] > \frac{14}{15} \quad (2.5)$$

and

$$f_{12}(r) \equiv \frac{1}{r^2} \left[(1 - 5r^2) \frac{\text{arth}r}{r} - 1 + 2r^2 \right] < -\frac{8}{3}. \quad (2.6)$$

(10) The functions

$$f_{13}(r) \equiv \frac{1}{r^6} \left[(15 - 12r^2 + r^4) \frac{\text{arth}r}{r} - 15 + 7r^2 \right] \text{ and}$$

$$f_{14}(r) \equiv \frac{1}{r^4} \left[3r'^2 - (3 - 4r^2 - r^4) \frac{\text{arth}r}{r} \right]$$

are both absolutely monotone on $(0, 1)$, with ranges $(8/105, \infty)$ and $(26/15, \infty)$, respectively.

Proof. (1) By (1.5), $f_1(r)$ can be written as

$$f_1(r) = \frac{1 - (r'^2 \text{arth}r)/r}{r^2} = 2 \sum_{n=0}^{\infty} \frac{r^{2n}}{(2n+1)(2n+3)}, \quad (2.7)$$

yielding the result for f_1 .

- (2) Since $f_2(r) = f_1(r)[r/(r'^2 \text{arth}r)]$, part (2) follows from part (1) and [22, Lemma 3(1)].
(3) Clearly, $f_3(1^-) = 0$. By (1.5),

$$\begin{aligned} f_3(r) &= \frac{1 - (r'^2 \text{arth}r)/r}{(1 + r^2)(\text{arth}r)/r - 1} \\ &= \frac{1}{2} \left[\sum_{n=0}^{\infty} \frac{r^{2n}}{(2n+1)(2n+3)} \right] \left[\sum_{n=0}^{\infty} \frac{(n+1)r^{2n}}{(2n+1)(2n+3)} \right]^{-1}. \end{aligned} \quad (2.8)$$

Applying [19, Lemma 2.1], one can easily obtain the monotonicity of f_3 . By (2.8), $f_3(0^+) = 1/2$.

- (4) By (1.4) and (1.5), we can write $f_4(r)$ as

$$\begin{aligned} f_4(r) &= \frac{r^{-3}[(1+r^2)\text{arth}r - r]}{\mathcal{K}(r)} \\ &= \frac{8}{\pi} \left[\sum_{n=0}^{\infty} \frac{n+1}{(2n+1)(2n+3)} r^{2n} \right] \left(\sum_{n=0}^{\infty} a_n r^{2n} \right)^{-1}. \end{aligned} \quad (2.9)$$

Let $\lambda_n = (n+1)/[(2n+1)(2n+3)a_n]$ for $n \in \mathbb{N}_0$. Then

$$\frac{\lambda_{n+1}}{\lambda_n} = \frac{(n+2)(2n+1)a_n}{(n+1)(2n+5)a_{n+1}} = \frac{4n^2 + 12n + 8}{4n^2 + 12n + 5} > 1,$$

which shows that the sequence $\{\lambda_n\}$ is strictly increasing, and hence by [19, Lemma 2.1], f_4 is strictly increasing on $(0, 1)$. By (2.9), $f_4(0^+) = 8/(3\pi)$. Since

$$\lim_{r \rightarrow 1} r\mathcal{K}(r)/\text{arth}r = 1 \quad (2.10)$$

by [8, Theorem 3.10] or (2.2), we obtain the limiting value $f_4(1^-) = 2$.

(5) The limiting values of f_5 are clear. By differentiation, $f_5'(r) = rf_1(r)/r'^3$, and hence the monotonicity and convexity properties of f_5 follow from part (1).

(6) Let $G_2(r) = (r/r') - r'\text{arth}r$ and $G_3(r) = \text{arth}r - r$. Then $G_2(0) = G_3(0) = 0$, $G_2(1^-) = G_3(1^-) = \infty$, $f_6(r) = G_2(r)/G_3(r)$, and

$$G_2'(r)/G_3'(r) = f_5(r). \quad (2.11)$$

Hence the monotonicity of f_6 follows from [9, Theorem 1.25] and part (5).

By l'Hôpital's rule and (2.11), $f_6(0^+) = 2$ and $f_6(1^-) = \infty$.

- (7) Let $G_4(r) = \log((\text{arth}r)/r)$ and $G_5(r) = r^2$. Then $G_4(0^+) = G_5(0) = 0$, $f_7(r) = G_4(r)/G_5(r)$ and

$$G_4'(r)/G_5'(r) = f_2(r)/2.$$

Hence the monotonicity of f_7 follows from [9, Theorem 1.25] and part (2). By l'Hôpital's rule, $f_7(0^+) = f_2(0^+)/2 = 1/3$. Clearly, $f_7(1^-) = \infty$.

(8) Let $G_6(r) = r/(r'^2 \text{arth}r) - 1$. Then $f_8(r) = G_4(r)/G_6(r)$, $G_4(0^+) = G_6(0^+) = 0$, $G_4(1^-) = G_6(1^-) = \infty$, and by differentiation,

$$\frac{G_4'(r)}{G_6'(r)} = f_3(r) \frac{r'^2 \text{arth}r}{r}$$

which is strictly decreasing from $(0, 1)$ onto $(0, 1/2)$ by part (3) and [22, Lemma 3(1)]. Hence the result for f_8 follows from [9, Theorem 1.25], part (3) and l'Hôpital's rule.

Similarly, since

$$\lim_{r \rightarrow 0} \log \frac{2\mathcal{K}(r)}{\pi} = \lim_{r \rightarrow 0} \frac{\mathcal{E}(r) - r'^2 \mathcal{K}(r)}{r'^2 \mathcal{K}(r)} = 0,$$

and since the ratio

$$\begin{aligned} & \left[\frac{d}{dr} \left(\log \frac{2\mathcal{K}(r)}{\pi} \right) \right] \left[\frac{d}{dr} \left(\frac{\mathcal{E}(r) - r'^2 \mathcal{K}(r)}{r'^2 \mathcal{K}(r)} \right) \right]^{-1} \\ &= \frac{r'^2 \mathcal{K}(r) [\mathcal{E}(r) - r'^2 \mathcal{K}(r)]}{\mathcal{E}(r)[\mathcal{K}(r) - \mathcal{E}(r)] + \mathcal{K}(r)[\mathcal{E}(r) - r'^2 \mathcal{K}(r)]} \\ &= \left[\frac{1}{r'^2} + \frac{\mathcal{E}(r)}{r'} \cdot \frac{1}{r' \mathcal{K}(r)} \cdot \frac{\mathcal{K}(r) - \mathcal{E}(r)}{\mathcal{E}(r) - r'^2 \mathcal{K}(r)} \right]^{-1} \end{aligned}$$

is strictly decreasing on $(0, 1)$ by [9, Theorem 3.21(6)&(8)], the result for f_9 follows from [9, Theorem 1.25] and l'Hôpital's rule.

(9) It follows from (1.5) that

$$\begin{aligned} f_{10}(r) &= \frac{1}{r^4} \left[19r^4 - 126r^2 - 72 \sum_{n=1}^{\infty} \frac{r^{2n}}{2n+1} + 150r^2 \sum_{n=0}^{\infty} \frac{r^{2n}}{2n+1} \right. \\ &\quad \left. + 253r^4 \sum_{n=0}^{\infty} \frac{r^{2n}}{2n+1} - 167r^6 \sum_{n=0}^{\infty} \frac{r^{2n}}{2n+1} \right] \\ &= \frac{1}{r^4} \left[19r^4 - 72 \sum_{n=2}^{\infty} \frac{r^{2n}}{2n+1} + 150r^2 \sum_{n=1}^{\infty} \frac{r^{2n}}{2n+1} \right. \\ &\quad \left. + 253r^4 \sum_{n=0}^{\infty} \frac{r^{2n}}{2n+1} - 167r^6 \sum_{n=0}^{\infty} \frac{r^{2n}}{2n+1} \right] \\ &= 19 - \sum_{n=0}^{\infty} \frac{72r^{2n}}{2n+5} + \sum_{n=0}^{\infty} \left(\frac{150}{2n+3} + \frac{253}{2n+1} \right) r^{2n} - r^2 \sum_{n=0}^{\infty} \frac{167r^{2n}}{2n+1} \\ &= \frac{1538}{5} - \frac{1322}{21} r^2 + 2r^4 \sum_{n=0}^{\infty} \frac{656n^3 + 5540n^2 + 12140n + 3951}{(2n+9)(2n+7)(2n+5)(2n+3)} r^{2n} \\ &> \frac{1538}{5} - \frac{1322}{21} + 2r^4 \sum_{n=0}^{\infty} \frac{656n^3 + 5540n^2 + 12140n + 3951}{(2n+9)(2n+7)(2n+5)(2n+3)} r^{2n} \\ &> \frac{1538}{5} - \frac{1322}{21} = \frac{25688}{105}, \\ f_{11}(r) &= 13 \sum_{n=0}^{\infty} \frac{r^{2n}}{2n+3} - 9 \sum_{n=0}^{\infty} \frac{r^{2n}}{2n+1} + 6 \\ &= \frac{4}{3} - \frac{2}{5} r^2 + 2 \sum_{n=2}^{\infty} \frac{4n-7}{(2n+1)(2n+3)} r^{2n} > \frac{4}{3} - \frac{2}{5} r^2 > \frac{14}{15} \end{aligned}$$

and

$$f_{12}(r) = 2 - 2 \sum_{n=0}^{\infty} \frac{4n+7}{(2n+1)(2n+3)} r^{2n} < 2 - \frac{14}{3} = -\frac{8}{3}.$$

(10) Clearly, $f_{13}(1^-) = f_{14}(1^-) = \infty$. Applying (1.5), we obtain the following series expansions

$$f_{13}(r) = 8 \sum_{n=0}^{\infty} \frac{(n+1)(2n+1)}{(2n+3)(2n+5)(2n+7)} r^{2n}$$

and

$$f_{14}(r) = 2 \sum_{n=0}^{\infty} \frac{4n^2 + 20n + 13}{(2n+1)(2n+3)(2n+5)} r^{2n},$$

from which the absolute monotonicity and the limiting values of f_{13} and f_{14} as $r \rightarrow 0$ follow. \square

Our next lemma presents some relations of $F_0(r^2) = 2\mathcal{K}(r)/\pi$ and $F_1(r^2) = (\operatorname{arth}r)/r$.

Lemma 2.2. (1) The function $f_{15}(r) \equiv F_1(r)/F_0(r)$ is strictly increasing from $(0, 1)$ onto $(1, \pi/2)$.

(2) The function $f_{16}(r) \equiv G_1(r)/G_0(r) = 3F_1'(r)/[4F_0'(r)]$ is strictly increasing from $(0, 1)$ onto $(1, 3\pi/8)$.

(3) The function

$$f_{17}(r) \equiv F_0(r) - F_1(r)F_0'(r)/F_1'(r)$$

is strictly increasing from $(0, 1)$ onto $(1/4, (\log 4)/\pi)$.

Proof. Part (1) is implied by [8, Theorem 3.10], and part (2) follows from [20, Lemmas 3.2(2)].

By (1.16), $f_{17}(r) = F_0(r) - 3F_1(r)G_0(r)/[4G_1(r)]$. Since

$$f_{17}'(r) = -F_1(r) \frac{d}{dr} \left[\frac{F_0'(r)}{F_1'(r)} \right] = -\frac{3}{4} F_1(r) \frac{d}{dr} \left[\frac{1}{f_{16}(r)} \right] > 0$$

by part (2), the monotonicity of f_{17} follows.

Clearly, $f_{17}(0) = 1/4$. By (1.16),

$$\begin{aligned} f_{17}(r) &= -\frac{F_0'(r)}{F_1'(r)} \left[F_1(r) - F_0(r) \frac{F_1'(r)}{F_0'(r)} \right] \\ &= -\frac{3}{4f_{16}(r)} \left[F_1(r) - F_0(r) \frac{F_1'(r)}{F_0'(r)} \right]. \end{aligned}$$

Hence the limiting value $f_{17}(1^-) = (\log 4)/\pi$ follows from part (2) and [20, Lemma 3.7(2)]. \square

Lemma 2.3. For $r \in (0, 1)$, the coefficients of the Maclaurin series of the function

$$f_{18}(r) \equiv \frac{1}{r^8} \left[\left(1 - \frac{1}{2}r^2 - \frac{1}{16}r^4 - \frac{1}{32}r^6 \right) \mathcal{K}(r) - \mathcal{E}(r) \right]$$

are all positive, with $f_{18}(0^+) = 41\pi/4096$ and $f_{18}(1^-) = \infty$.

Proof. By (1.3) and (1.4), we have

$$\begin{aligned} f_{18}(r) &= \frac{\pi}{2r^8} \left[\left(1 - \frac{1}{2}r^2 - \frac{1}{16}r^4 - \frac{1}{32}r^6 \right) \sum_{n=0}^{\infty} a_n r^{2n} + \sum_{n=0}^{\infty} \frac{a_n}{2n-1} r^{2n} \right] \\ &= \frac{\pi}{2r^8} \left[\sum_{n=1}^{\infty} \left(\frac{2n}{2n-1} a_n - \frac{1}{2} a_{n-1} \right) r^{2n} - \frac{1}{16} \sum_{n=2}^{\infty} a_{n-2} r^{2n} - \frac{1}{32} \sum_{n=3}^{\infty} a_{n-3} r^{2n} \right] \\ &= \frac{\pi}{2r^8} \left[\sum_{n=2}^{\infty} \left(\frac{2n}{2n-1} a_n - \frac{1}{2} a_{n-1} \right) r^{2n} - \frac{1}{16} r^4 - \sum_{n=3}^{\infty} \left(\frac{1}{16} a_{n-2} + \frac{1}{32} a_{n-3} \right) r^{2n} \right] \\ &= \frac{\pi}{2r^8} \sum_{n=4}^{\infty} \left(\frac{2n}{2n-1} a_n - \frac{1}{2} a_{n-1} - \frac{1}{16} a_{n-2} - \frac{1}{32} a_{n-3} \right) r^{2n} \\ &= \frac{\pi}{2} \sum_{n=0}^{\infty} \left(\frac{2n+8}{2n+7} a_{n+4} - \frac{1}{2} a_{n+3} - \frac{1}{16} a_{n+2} - \frac{1}{32} a_{n+1} \right) r^{2n} \\ &= \frac{\pi}{2} \sum_{n=0}^{\infty} b_n r^{2n}, \end{aligned} \tag{2.12}$$

where

$$b_n = \frac{2n+8}{2n+7} a_{n+4} - \frac{1}{2} a_{n+3} - \frac{1}{16} a_{n+2} - \frac{1}{32} a_{n+1}.$$

Since $4a_{n+1} = [(2n+1)/(n+1)]^2 a_n$, we have

$$\begin{aligned}
b_n &= \left[\frac{n+4}{2(2n+7)} \left(\frac{2n+7}{n+4} \right)^2 - \frac{1}{2} \right] a_{n+3} - \frac{1}{32} \left(2 + \frac{a_{n+1}}{a_{n+2}} \right) a_{n+2} \\
&= \frac{n+3}{2(n+4)} a_{n+3} - \frac{6n^2+20n+17}{16(2n+3)^2} a_{n+2} \\
&= \left[\frac{n+3}{8(n+4)} \left(\frac{2n+5}{n+3} \right)^2 - \frac{6n^2+20n+17}{16(2n+3)^2} \right] a_{n+2} \\
&= \frac{26n^4+194n^3+523n^2+601n+246}{16(n+3)(n+4)(2n+3)^2} a_{n+2} \\
&= \frac{(n+2)(n+1)(26n^2+116n+123)}{16(n+3)(n+4)(2n+3)^2} a_{n+2} > 0
\end{aligned} \tag{2.13}$$

for all $n \in \mathbb{N}_0$. Hence it follows from (2.12) and (2.13) that all the coefficients of the Maclaurin series of f_{18} are positive. By (2.12), $f_{18}(0^+) = \pi b_0/2 = 41\pi/4096$. Clearly, $f_{18}(1^-) = \infty$. \square

The following lemma plays an important role in the proof of Theorem 1.8.

Lemma 2.4. For $n \in \mathbb{N}_0$, let $c_n = (2n+3)a_{n+1}$,

$$\begin{aligned}
\tilde{c}_n &= \frac{(22n+83)(2n+9)(2n+7)(2n+5)(2n+3)}{(n+4)^2(32276n^2+123808n+110907)} a_{n+3}, \\
d_n &= (2n+1) \left[\frac{10196n^2+39096n+34975}{(2n+7)(2n+5)(2n+3)} - 2880a_{n+3} \right].
\end{aligned}$$

(1) The sequence $\{c_n\}$ is strictly decreasing with $c_0 = 3/4$ and $c_\infty = \lim_{n \rightarrow \infty} c_n = 2/\pi$.

(2) The sequence $\{\tilde{c}_n\}$ is strictly decreasing with $\tilde{c}_0 = 217875/50475008 = 0.0043164 \dots$ and $\tilde{c}_\infty \equiv \lim_{n \rightarrow \infty} \tilde{c}_n = 88/(8069\pi) = 0.0034714 \dots$.

(3) The sequence $\{d_n\}$ is strictly increasing with $d_0 = 4355/84 = 51.845238 \dots$ and $d_\infty \equiv \lim_{n \rightarrow \infty} d_n = 2549 - 5760/\pi = 715.535055 \dots$.

Proof. (1) Clearly, $c_0 = 3/4$. By applying [4, 6.1.8 & 6.1.47], we obtain the value $c_\infty = \lim_{n \rightarrow \infty} c_n = 2/\pi$. Computation gives

$$\frac{c_{n+1}}{c_n} = \frac{(2n+5)a_{n+2}}{(2n+3)a_{n+1}} = \frac{2n+5}{2n+3} \left(\frac{n+3/2}{n+2} \right)^2 = \frac{4n^2+16n+15}{4n^2+16n+16} < 1,$$

yielding the monotonicity of the sequence $\{c_n\}$.

(2) It is easy to see that $\tilde{c}_0 = 217875/50475008$. By [4, 6.1.8 & 6.1.47], $\lim_{n \rightarrow \infty} \tilde{c}_n = 88/(8069\pi)$.

By computation, we obtain

$$\begin{aligned}
\frac{\tilde{c}_{n+1}}{\tilde{c}_n} &= \frac{(n+4)^2(22n+105)(2n+11)(32276n^2+123808n+110907)}{(n+5)^2(22n+83)(2n+3)(32276n^2+188360n+266991)} \frac{a_{n+4}}{a_{n+3}} \\
&= \frac{(2n+7)^2(22n+105)(2n+11)(32276n^2+123808n+110907)}{4(n+5)^2(22n+83)(2n+3)(32276n^2+188360n+266991)} = \frac{\Delta_1(n)}{\Delta_2(n)}
\end{aligned}$$

for $n \in \mathbb{N}_0$, where

$$\begin{aligned}
\Delta_1(n) &= 5680576n^6 + 119909248n^5 + 1023083184n^4 + 4501649024n^3 \\
&\quad + 10727855348n^2 + 13050014376n + 6276781665 \quad \text{and} \\
\Delta_2(n) &= 5680576n^6 + 119909248n^5 + 1026984672n^4 + 4552315328n^3 \\
&\quad + 10968049916n^2 + 13543585560n + 6648075900.
\end{aligned}$$

Since

$$\begin{aligned}\Delta_2(n) - \Delta_1(n) &= 3901488n^4 + 50666304n^3 + 240194568n^2 \\ &\quad + 493571184n + 371294235 > 371294235\end{aligned}$$

for $n \in \mathbb{N}_0$, $\tilde{c}_{n+1} < \tilde{c}_n$ for $n \in \mathbb{N}_0$, namely, the sequence $\{\tilde{c}_n\}$ is strictly decreasing.

(3) Put $\Delta = d_{n+1} - d_n$. Then by computation, we obtain

$$\begin{aligned}\Delta &= (2n+3) \frac{10196n^2 + 59488n + 84267}{(2n+9)(2n+7)(2n+5)} - (2n+1) \frac{10196n^2 + 39096n + 34975}{(2n+7)(2n+5)(2n+3)} \\ &\quad - 2880 \left[(2n+3) \frac{a_{n+4}}{a_{n+3}} - (2n+1) \right] a_{n+3} \\ &= 4 \left[\frac{32276n^2 + 123808n + 110907}{(2n+9)(2n+7)(2n+5)(2n+3)} - 180 \frac{22n+83}{(n+4)^2} a_{n+3} \right],\end{aligned}$$

and hence

$$\begin{aligned}d_{n+1} > d_n &\iff \frac{32276n^2 + 123808n + 110907}{(2n+9)(2n+7)(2n+5)(2n+3)} > 180 \frac{22n+83}{(n+4)^2} a_{n+3} \\ &\iff \frac{(22n+83)(2n+9)(2n+7)(2n+5)(2n+3)}{(n+4)^2 (32276n^2 + 123808n + 110907)} a_{n+3} < \frac{1}{180} \\ &\iff \tilde{c}_n < \frac{1}{180} = 0.0055555 \dots\end{aligned}$$

which is true by part (2). Hence the monotonicity of the sequence $\{d_n\}$ follows. Clearly, $d_0 = 34975/105 - 2880a_3 = 4355/84$. By [4, 6.1.8 & 6.1.47], we obtain the value $d_\infty = 2549 - 5760/\pi$. \square

3 Some Properties of the Complete Elliptic Integrals

In this section, we present several properties of the complete elliptic integrals $\mathcal{K}(r)$ and $\mathcal{E}(r)$, which are also needed in the proofs of Theorems 1.6 and 1.7.

Theorem 3.1. *Let f_{18} be as in Lemma 2.3. Then the function $f_{19}(r) \equiv f_{18}(r)/\mathcal{K}(r)$ is strictly increasing from $(0, 1)$ onto $(41/2048, 13/32)$. In particular, for $r \in (0, 1)$,*

$$P(r) < \mathcal{E}(r)/\mathcal{K}(r) < Q(r), \quad (3.1)$$

where $Q(r) = 1 - r^2/2 - r^4/16 - r^6/32 - 41r^8/2048$ and

$$P(r) = 1 - \frac{1}{2}r^2 - \frac{1}{16}r^4 - \frac{1}{32}r^6 - \frac{13}{32}r^8 = r'^2 \left(1 + r^2 \frac{16 + 14r^2 + 13r^4}{32} \right).$$

Proof. For $n \in \mathbb{N}_0$, let b_n be as in the proof of Lemma 2.3, and $C_n = b_n/a_n$. Then by (2.12), we have

$$f_{19}(r) = \left(\sum_{n=0}^{\infty} b_n r^{2n} \right) \left(\sum_{n=0}^{\infty} a_n r^{2n} \right)^{-1}. \quad (3.2)$$

It follows from (2.13) that

$$\begin{aligned}C_n &= \frac{(n+2)(n+1)(26n^2 + 116n + 123)}{256(n+3)(n+4)(2n+3)^2} \left(\frac{2n+3}{n+2} \right)^2 \left(\frac{2n+1}{n+1} \right)^2 \\ &= \frac{(2n+1)^2 (26n^2 + 116n + 123)}{256(n+4)(n+3)(n+2)(n+1)}\end{aligned}$$

and

$$\begin{aligned}\frac{C_{n+1}}{C_n} &= \frac{(n+1)(2n+3)^2(26n^2+168n+265)}{(n+5)(2n+1)^2(26n^2+116n+123)} \\ &= \frac{104n^5+1088n^4+4294n^3+8002n^2+7077n+2385}{104n^5+1088n^4+3822n^3+5518n^2+3163n+615} > 1\end{aligned}$$

which shows that the sequence $\{C_n\}$ is strictly increasing. Consequently, the monotonicity of f_{19} follows from [19, Lemma 2.1] and (3.2).

Clearly, $f_{19}(0^+) = C_0 = 41/2048$ and $f_{19}(1^-) = 13/32$. The double inequality (3.1) is clear. \square

Theorem 3.2. (1) Define the functions f_{20} and f_{21} on $(0, 1)$ by

$$\begin{aligned}f_{20}(r) &= \left(\frac{3}{4} + \frac{1}{4}r^2\right)(r - r'^2 \operatorname{arth} r) \mathcal{K}(r) - [\mathcal{E}(r) - r'^2 \mathcal{K}(r)] \operatorname{arth} r, \\ f_{21}(r) &= [\mathcal{E}(r) - r'^2 \mathcal{K}(r)] \operatorname{arth} r - \left(\frac{3}{4} + \frac{1}{160}r^2\right)(r - r'^2 \operatorname{arth} r) \mathcal{K}(r).\end{aligned}$$

Then f_{20} and f_{21} are both strictly increasing and convex on $(0, 1)$, with ranges $(0, \log 2)$ and $(0, \infty)$, respectively.

(2) The function

$$f_{22}(r) \equiv \frac{(r - r'^2 \operatorname{arth} r) \mathcal{E}(r) - 2r'^2 (\operatorname{arth} r - r) \mathcal{K}(r)}{r^5}$$

is strictly increasing from $(0, 1)$ onto $(\pi/30, 1)$.

(3) The function

$$f_{23}(r) \equiv 2 \frac{r'^2 \operatorname{arth} r}{r} \mathcal{K}(r) + \frac{r - r'^2 \operatorname{arth} r}{r^3} \mathcal{E}(r)$$

is strictly decreasing from $(0, 1)$ onto $(1, 4\pi/3)$.

(4) The function

$$f_{24}(r) \equiv \frac{1}{r^2} \left\{ 4 \frac{[\mathcal{E}(r) - r'^2 \mathcal{K}(r)] \operatorname{arth} r}{(r - r'^2 \operatorname{arth} r) \mathcal{K}(r)} - 3 \right\}$$

is strictly increasing from $(0, 1)$ onto $(1/40, 1)$. In particular, for $r \in (0, 1)$,

$$\frac{3}{4} + \frac{r^2}{4} P_4(r) < \frac{[\mathcal{E}(r) - r'^2 \mathcal{K}(r)] \operatorname{arth} r}{(r - r'^2 \operatorname{arth} r) \mathcal{K}(r)} < \frac{3}{4} + \frac{r^2}{4}, \quad (3.3)$$

where

$$P_4(r) = \max \left\{ \frac{1}{40}, 1 - \frac{\log 16}{r(r - r'^2 \operatorname{arth} r) \mathcal{K}(r)} \right\}.$$

Proof. (1) By (2.1) and by differentiation, we obtain

$$\begin{aligned}f'_{20}(r) &= \frac{1}{2} \left[2r(1+r^2) \operatorname{arth} r + r^2 \right] \mathcal{K} + \frac{1}{4} (3+r^2) (r - r'^2 \operatorname{arth} r) \frac{\mathcal{E} - r'^2 \mathcal{K}}{r r'^2} \\ &\quad - r \mathcal{K} \operatorname{arth} r - \frac{\mathcal{E} - r'^2 \mathcal{K}}{r'^2} \\ &= \frac{1}{4} \left\{ \left[(3r^4 - 2r^2 + 3) \frac{\operatorname{arth} r}{r} + r^2 + 1 \right] \mathcal{K} - \left[(3+r^2) \frac{\operatorname{arth} r}{r} + 1 \right] \mathcal{E} \right\} \\ &= \frac{1}{4} \left\{ \left[1 + (3+r^2) \frac{\operatorname{arth} r}{r} \right] (\mathcal{K} - \mathcal{E}) + r^2 \left(1 - 3 \frac{r'^2 \operatorname{arth} r}{r} \right) \mathcal{K} \right\} \\ &= \frac{1}{4} r^2 \mathcal{K} \left\{ \left[1 + (3+r^2) \frac{\operatorname{arth} r}{r} \right] \frac{\mathcal{K} - \mathcal{E}}{r^2 \mathcal{K}} + 1 - 3 \frac{r'^2 \operatorname{arth} r}{r} \right\} \\ &= \frac{1}{4} G_7(r) G_8(r),\end{aligned} \quad (3.4)$$

where $G_7(r) = r^2 \mathcal{K}$ and

$$G_8(r) = \left[1 + (3 + r^2) \frac{\operatorname{arth} r}{r} \right] \frac{\mathcal{K} - \mathcal{E}}{r^2 \mathcal{K}} + 1 - 3 \frac{r'^2 \operatorname{arth} r}{r}.$$

Clearly, G_7 is strictly increasing from $(0, 1)$ onto $(0, \infty)$. By [6, Lemma 5.2(3)] and [22, Lemma 3(1)], we see that G_8 is strictly increasing from $(0, 1)$ onto $(0, \infty)$. Hence it follows from (3.4) that f_{20}' is strictly increasing from $(0, 1)$ onto $(0, \infty)$. This yields the monotonicity and convexity of f_{20} .

Clearly, $f_{20}(0^+) = 0$. By l'Hôpital's rule,

$$\lim_{r \rightarrow 1} \frac{4\mathcal{E} - r(3 + r^2)}{r'} = \lim_{r \rightarrow 1} \frac{1}{r} \left[4r' \frac{\mathcal{K} - \mathcal{E}}{r} + 3r'(1 + r^2) \right] = 0,$$

and hence by (2.2), we obtain

$$\begin{aligned} f_{20}(1^-) &= \lim_{r \rightarrow 1} \left[\frac{3 + r^2}{4} (r - r'^2 \operatorname{arth} r) \mathcal{K} - (\mathcal{E} - r'^2 \mathcal{K}) \operatorname{arth} r \right] \\ &= \lim_{r \rightarrow 1} \left[\frac{r(3 + r^2)}{4} \log \frac{4}{r'} - \frac{\mathcal{E}}{2} \log \frac{1+r}{1-r} \right] \\ &= \log 2 + \frac{1}{8} \lim_{r \rightarrow 1} \frac{4\mathcal{E} - r(3 + r^2)}{r'} \cdot r' \log(1 - r) = \log 2. \end{aligned}$$

Next, for $r \in (0, 1)$, let

$$\begin{aligned} G_9(r) &= \left[40r - r^3 + (120 + r^2) r'^2 \operatorname{arth} r \right] \mathcal{E} \\ &\quad - r'^2 \left[40r + r^3 + (120 - 41r^2 + 3r^4) \operatorname{arth} r \right] \mathcal{K}, \\ G_{10}(r) &= 4 \left[40 - (2r^3 + 79r) \operatorname{arth} r - r^2 \right] \mathcal{E} \\ &\quad - \left[160 - (15r^5 - 131r^3 + 280r) \operatorname{arth} r - r^4 - 120r^2 \right] \mathcal{K}, \\ G_{11}(r) &= 2G_{12}(r) \mathcal{K} + \left[(47r^4 + 469r^2 - 352) \operatorname{arth} r + 5r^3 - 368r \right] \mathcal{E} \text{ and} \\ G_{12}(r) &= r'^2 (30r^4 - 127r^2 + 158) \operatorname{arth} r + 6r^5 - 126r^3 + 202r. \end{aligned}$$

Then by differentiation,

$$\begin{aligned} f_{21}'(r) &= r \mathcal{K} \operatorname{arth} r + \frac{\mathcal{E} - r'^2 \mathcal{K}}{r'^2} - \frac{r}{80} \left[(2r^2 + 119) \operatorname{arth} r + r \right] \mathcal{K} \\ &\quad - \frac{120 + r^2}{160} (r - r'^2 \operatorname{arth} r) \frac{\mathcal{E} - r'^2 \mathcal{K}}{r r'^2} = \frac{G_9(r)}{160 r r'^2}, \end{aligned} \quad (3.5)$$

$$\begin{aligned} G_9'(r) &= \left[(18r^5 - 176r^3 + 322r) \operatorname{arth} r + 2r^4 + 158r^2 - 160 \right] \mathcal{K} \\ &\quad + \left[(3r^6 - 44r^4 + 161r^2 - 120) \operatorname{arth} r + r^5 + 39r^3 - 40r \right] \frac{\mathcal{E} - r'^2 \mathcal{K}}{r r'^2} \\ &\quad - \left[(4r^3 + 238r) \operatorname{arth} r + 2r^2 - 160 \right] \mathcal{E} \\ &\quad + \left[(r^4 + 119r^2 - 120) \operatorname{arth} r + r^3 - 40r \right] \frac{\mathcal{K} - \mathcal{E}}{r} = G_{10}(r), \end{aligned} \quad (3.6)$$

$$\begin{aligned} G_{10}'(r) &= \frac{1}{r'^2} \left[520r - 367r^3 + 11r^5 - (75r^6 - 468r^4 + 673r^2 - 280) \operatorname{arth} r \right] \mathcal{K} \\ &\quad + \left[(15r^5 - 131r^3 + 280r) \operatorname{arth} r + r^4 + 120r^2 - 160 \right] \frac{\mathcal{E} - r'^2 \mathcal{K}}{r r'^2} \\ &\quad + \frac{4}{r'^2} \left[(6r^4 + 73r^2 - 79) \operatorname{arth} r - 81r \right] \mathcal{E} \\ &\quad + 4 \left[(2r^3 + 79r) \operatorname{arth} r + r^2 - 40 \right] \frac{\mathcal{K} - \mathcal{E}}{r} = \frac{G_{11}(r)}{r'^2}. \end{aligned} \quad (3.7)$$

It is clear that for all $r \in (0, 1)$,

$$G_{12}(r) = r'^2 (158r'^2 + 31r^2 + 30r^4) \operatorname{arth} r + r (202r'^2 + 76r^2 + 6r^4) > 0.$$

Let f_{10} be as in Lemma 2.1(9). Then by (2.4) and (3.1), we have

$$\begin{aligned} \frac{G_{11}(r)}{\mathcal{E}} &> \frac{4G_{12}(r)}{2-r^2} + (47r^4 + 469r^2 - 352) \operatorname{arth} r \\ &+ 5r^3 - 368r = \frac{r^5}{2-r^2} f_{10}(r) > 0. \end{aligned} \quad (3.8)$$

It follows from (2.4) and (3.7)–(3.8) that

$$G_{10}(r) = G'_9(r) \left[\frac{d}{dr}(r) \right]^{-1}$$

is strictly increasing on $(0, 1)$ with $G_{10}(0^+) = 0$, and hence by [9, Theorem 1.25], the function $r \mapsto G_9(r)/r$ is strictly increasing on $(0, 1)$ with $G_9(0^+) = 0$. Consequently, by (3.5), f'_{21} is positive and strictly increasing on $(0, 1)$. This yields the monotonicity and convexity of f_{21} .

Clearly, $f_{21}(0^+) = 0$. By (2.10),

$$\lim_{r \rightarrow 1} \left[\mathcal{E}(r) - r \left(\frac{3}{4} + \frac{1}{160} r^2 \right) \frac{\mathcal{K}(r)}{\operatorname{arth} r} \right] = \frac{39}{160},$$

so that we obtain the limiting value

$$f_{21}(1^-) = \lim_{r \rightarrow 1} \left[\mathcal{E}(r) - r \left(\frac{3}{4} + \frac{1}{160} r^2 \right) \frac{\mathcal{K}(r)}{\operatorname{arth} r} \right] \operatorname{arth} r = \infty.$$

(2) Let f_{11} and f_{13} be as in Lemma 2.1. Then by differentiation,

$$\begin{aligned} r^6 f'_{22}(r) &= r \left[2r\mathcal{E} \operatorname{arth} r + (r - r'^2 \operatorname{arth} r) \frac{\mathcal{E} - \mathcal{K}}{r} - 2r(3r - 2\operatorname{arth} r)\mathcal{K} \right. \\ &\quad \left. - 2(\operatorname{arth} r - r) \frac{\mathcal{E} - r'^2 \mathcal{K}}{r} \right] - 5 \left[(r - r'^2 \operatorname{arth} r) \mathcal{E} - 2r'^2 (\operatorname{arth} r - r) \mathcal{K} \right] \\ &= \left[(13 - 9r^2) \operatorname{arth} r - 13r + 6r^3 \right] \mathcal{K} - 2(r - r'^2 \operatorname{arth} r) \mathcal{E} \\ &= r^3 f_{11}(r) \mathcal{K} - 2(r - r'^2 \operatorname{arth} r) \mathcal{E}. \end{aligned} \quad (3.9)$$

By (3.1) and Lemma 2.1(10), and by (3.9),

$$\begin{aligned} r^6 f'_{22}(r) &> r^3 f_{11}(r) \frac{2\mathcal{E}}{2-r^2} - 2(r - r'^2 \operatorname{arth} r) \mathcal{E} \\ &= \frac{2\mathcal{E}}{2-r^2} \left[r^3 f_{11}(r) - (2-r^2)(r - r'^2 \operatorname{arth} r) \right] \\ &= \frac{2\mathcal{E}}{2-r^2} \left[(15 - 12r^2 + r^4) \operatorname{arth} r - 15r + 7r^3 \right] \\ &= \frac{2r^7 \mathcal{E}}{2-r^2} f_{13}(r) > \frac{16r^7 \mathcal{E}}{105(2-r^2)} > 0 \end{aligned}$$

for all $r \in (0, 1)$, and hence the monotonicity of f_{22} follows.

Clearly, $f_{22}(1^-) = 1$. By Lemma 2.1(1), (1.5) and (2.3), we obtain

$$\begin{aligned} f_{22}(0^+) &= \lim_{r \rightarrow 0} \frac{1}{r^5} \left\{ (r - r'^2 \operatorname{arth} r) (\mathcal{E} - r'^2 \mathcal{K}) - r'^2 \left[(3 - r^2) \operatorname{arth} r - 3r \right] \mathcal{K} \right\} \\ &= \lim_{r \rightarrow 0} \left[\frac{(r - r'^2 \operatorname{arth} r) \mathcal{E} - r'^2 \mathcal{K}}{r^3} - 4r'^2 \mathcal{K} \sum_{n=0}^{\infty} \frac{n+1}{(2n+3)(2n+5)} r^{2n} \right] \\ &= \frac{\pi}{6} - \frac{2\pi}{15} = \frac{\pi}{30}. \end{aligned}$$

(3) Let f_{12} and f_{14} be as in Lemma 2.1. Differentiation gives

$$\begin{aligned}
r^4 f'_{23}(r) &= 2r^2 \left\{ \left[r - (1+r^2) \operatorname{arth} r \right] \mathcal{K} + (\mathcal{E} - r'^2 \mathcal{K}) \operatorname{arth} r \right\} \\
&\quad + \left[(3-r^2) \operatorname{arth} r - 3r \right] \mathcal{E} + (r - r'^2 \operatorname{arth} r) (\mathcal{E} - \mathcal{K}) \\
&= 2 \left[(1+r^2) \operatorname{arth} r - r \right] \mathcal{E} + \left[(1-5r^2) \operatorname{arth} r - r + 2r^3 \right] \mathcal{K} \\
&= 2 \left[(1+r^2) \operatorname{arth} r - r \right] \mathcal{E} + r^3 f_{12}(r) \mathcal{K}.
\end{aligned} \tag{3.10}$$

It follows from (2.6), (3.1), (3.10) and Lemma 2.1(10) that

$$\begin{aligned}
r^4 f'_{23}(r) &< 2 \left[(1+r^2) \operatorname{arth} r - r \right] \mathcal{E} + 2r^3 f_{12}(r) \frac{\mathcal{E}}{2-r^2} \\
&= -\frac{2r\mathcal{E}}{2-r^2} \left[3r'^2 - (3-4r^2-r^4) \frac{\operatorname{arth} r}{r} \right] \\
&= -\frac{2r^5 \mathcal{E}}{2-r^2} f_{14}(r) < -\frac{52r^5 \mathcal{E}}{15(2-r^2)} < 0,
\end{aligned}$$

yielding the monotonicity of f_{23} .

By Lemma 2.1(1), we obtain the limiting values $f_{23}(0^+) = 4\pi/3$ and $f_{23}(1^-) = 1$.

(4) For $r \in (0, 1)$, let

$$\begin{aligned}
G_{13}(r) &= 4 \left(\mathcal{E} - r'^2 \mathcal{K} \right) \frac{\operatorname{arth} r}{r} - 3 \left(1 - \frac{r'^2 \operatorname{arth} r}{r} \right) \mathcal{K} \text{ and} \\
G_{14}(r) &= r^2 \left(1 - \frac{r'^2 \operatorname{arth} r}{r} \right) \mathcal{K}.
\end{aligned}$$

Then $f_{24}(r) = G_{13}(r)/G_{14}(r)$, $G_{13}(0^+) = G_{14}(0^+) = 0$, $G_{14}(1^-) = \infty$, and by (2.10),

$$G_{13}(1^-) = \lim_{r \rightarrow 1} \mathcal{K} \left[4 \left(\mathcal{E} - r'^2 \mathcal{K} \right) \frac{\operatorname{arth} r}{r \mathcal{K}} + 3 \frac{r'^2 \operatorname{arth} r}{r} - 3 \right] = \infty.$$

Differentiation gives

$$\begin{aligned}
G'_{13}(r) &= \frac{1}{r^2 r'^2} \left[(r - r'^2 \operatorname{arth} r) \mathcal{E} - 2r'^2 (\operatorname{arth} r - r) \mathcal{K} \right], \\
G'_{14}(r) &= \frac{1}{r'^2} \left[(r - r'^2 \operatorname{arth} r) \mathcal{E} + 2r^2 r'^2 \mathcal{K} \operatorname{arth} r \right], \text{ and} \\
\frac{G'_{13}(r)}{G'_{14}(r)} &= \frac{(r - r'^2 \operatorname{arth} r) \mathcal{E} - 2r'^2 (\operatorname{arth} r - r) \mathcal{K}}{r^2 [(r - r'^2 \operatorname{arth} r) \mathcal{E} + 2r^2 r'^2 \mathcal{K} \operatorname{arth} r]} = \frac{f_{22}(r)}{f_{23}(r)}
\end{aligned}$$

which is strictly increasing on $(0, 1)$ by parts (2) and (3). Hence the monotonicity of f_{24} follows from [9, Theorem 1.25].

By l'Hôpital's rule, we obtain the limiting values $f_{24}(0^+) = f_{22}(0^+)/f_{23}(0^+) = 1/40$ and $f_{24}(1^-) = f_{22}(1^-)/f_{23}(1^-) = 1$.

The first lower bound and the upper bound in (3.3) are clear. The second lower bound in (3.3) follows from the convexity of f_{20} . \square

Corollary 3.3. *The function*

$$f_{25}(r) \equiv \frac{[\mathcal{E}(r) - r'^2 \mathcal{K}(r)] \operatorname{arth} r}{(r - r'^2 \operatorname{arth} r) \mathcal{K}(r)}$$

is strictly increasing from $(0, 1)$ onto $(3/4, 1)$. In particular, for $r \in (0, 1)$,

$$1 - \frac{r - r'^2 \operatorname{arth} r}{r^2 \operatorname{arth} r} < \frac{\mathcal{K}(r) - \mathcal{E}(r)}{r^2 \mathcal{K}(r)} < 1 - \frac{3r - r'^2 \operatorname{arth} r}{4r^2 \operatorname{arth} r}. \tag{3.11}$$

The coefficients 1 and 3/4 of $(r - r'^2 \operatorname{arth} r)/(r^2 \operatorname{arth} r)$ in (3.11) are both best possible.

Proof. Let f_{24} be as in Theorem 3.2(4). Then $f_{25}(r)$ can be rewritten as

$$f_{25}(r) = \frac{1}{4}r^2 f_{24}(r) + \frac{3}{4},$$

and hence the result for f_{25} follows from Theorem 3.2(4). The remaining conclusions are clear. \square

4 Proofs of Theorems 1.6–1.8

In this section, we prove Theorems 1.6–1.8 stated in Section 1.

3.1 Proof of Theorem 1.6

Let f_2 , f_7 and f_8 be as in Lemma 2.1, f_{24} as in Theorem 3.2(4), and let

$$H_1(r) = \frac{f_{24}(r)}{2f_8(r) + 1} \quad \text{and} \quad H_2(r) = c - \frac{1}{4}H_1(r).$$

It follows from Lemma 2.1(8) and Theorem 3.2(4) that H_1 is strictly increasing from $(0, 1)$ onto $(1/80, 1)$. Differentiation gives

$$\begin{aligned} g'(r) &= 2cr \log \frac{\operatorname{arth} r}{r} + \left(\frac{3}{4} + cr^2 \right) \frac{r - r'^2 \operatorname{arth} r}{rr'^2 \operatorname{arth} r} - \frac{\mathcal{E} - r'^2 \mathcal{K}}{rr'^2 \mathcal{K}} \\ &= cr \left(2 \log \frac{\operatorname{arth} r}{r} + \frac{r - r'^2 \operatorname{arth} r}{r'^2 \operatorname{arth} r} \right) \\ &\quad - \frac{4(\mathcal{E} - r'^2 \mathcal{K}) \operatorname{arth} r - 3(r - r'^2 \operatorname{arth} r) \mathcal{K}}{4rr'^2 \mathcal{K} \operatorname{arth} r} \\ &= cr \left(2 \log \frac{\operatorname{arth} r}{r} + \frac{r - r'^2 \operatorname{arth} r}{r'^2 \operatorname{arth} r} \right) - r \frac{r - r'^2 \operatorname{arth} r}{4r'^2 \operatorname{arth} r} f_{24}(r) \\ &= r \left(2 \log \frac{\operatorname{arth} r}{r} + \frac{r - r'^2 \operatorname{arth} r}{r'^2 \operatorname{arth} r} \right) H_2(r) \\ &= r^3 [f_2(r) + 2f_7(r)] H_2(r). \end{aligned} \tag{4.1}$$

By Lemma 2.1(8) and Theorem 3.2(4), H_2 is strictly decreasing from $(0, 1)$ onto $(c-1/4, c-1/320)$. Hence it follows from and (4.1) that g is strictly increasing (decreasing) on $(0, 1)$ if and only if $c \geq 1/4$ ($c \leq 1/320$, respectively).

Clearly, $g(0^+) = 0$. By (2.10),

$$\begin{aligned} g(1^-) &= \lim_{r \rightarrow 1} \log \left(\frac{\pi}{2\mathcal{K}} \left(\frac{\operatorname{arth} r}{r} \right)^{3/4 + cr^2} \right) \\ &= \log \frac{\pi}{2} + \lim_{r \rightarrow 1} \log \left(\frac{\operatorname{arth} r}{r\mathcal{K}} \left(\frac{\operatorname{arth} r}{r} \right)^{cr^2 - 1/4} \right) \\ &= \log \frac{\pi}{2} + \lim_{r \rightarrow 1} \left(cr^2 - \frac{1}{4} \right) \log \frac{\operatorname{arth} r}{r} \\ &= \begin{cases} \log(\pi/2), & \text{if } c = 1/4, \\ \infty, & \text{if } c > 1/4, \\ -\infty, & \text{if } c < 1/4. \end{cases} \end{aligned}$$

Next, let f_{17} be as in Lemma 2.2(3), and for $r \in (0, 1)$, let

$$\begin{aligned} x &= r^2, \quad H_3(r) = g(r)|_{c=1/4}, \\ H_4(x) &= H_3(\sqrt{x}) = \frac{3+x}{4} \log F_1(x) - \log F_0(x), \\ H_5(x) &= (1-x)^{1/4} F_0(x), \quad H_6(x) = (1-x)^{1/3} F_1(x) \end{aligned}$$

and $H_7(x) = (1-x)F_0(x)$. Then by differentiation and (1.16),

$$\begin{aligned}
H_3'(r) &= 2r \left[\frac{1}{4} \log F_1(x) + \frac{(3+x)F_1'(x)}{4F_1(x)} - \frac{F_0'(x)}{F_0(x)} \right] \\
&= 2r \left\{ \frac{1}{4} \log F_1(x) + \frac{F_1'(x)}{F_0(x)F_1(x)} \left[\frac{3+x}{4} F_0(x) - F_1(x) \frac{F_0'(x)}{F_1(x)} \right] \right\} \\
&= \frac{r}{2} \left\{ \log F_1(x) + \frac{G_1(x)}{3(1-x)F_0(x)F_1(x)} [4f_{17}(x) - (1-x)F_0(x)] \right\} \\
&= \frac{r}{2} \left\{ \log F_1(x) + \frac{G_1(x)}{3(1-x)^{5/12}H_5(x)H_6(x)} [4f_{17}(x) - H_7(x)] \right\},
\end{aligned}$$

which gives

$$\frac{2}{r} H_3'(r) = H_8(x) \equiv \log F_1(x) + \frac{G_1(x) [4f_{17}(x) - H_7(x)]}{3(1-x)^{5/12}H_5(x)H_6(x)}. \quad (4.2)$$

Clearly, $H_8(0^+) = 0$. By [24, Lemma 2.15(1)], the functions H_5 , H_6 and H_7 are all strictly decreasing from $(0, 1)$ onto itself, and by Lemma 2.2(3), the function $x \mapsto 4f_{17}(x) - H_7(x)$ is strictly increasing from $(0, 1)$ onto $(0, 4(\log 4)/\pi)$. Hence H_8 is positive and strictly increasing on $(0, 1)$, and so is H_3' by (4.2). This shows that g is convex on $(0, 1)$ in the case when $c = 1/4$.

Let $H_9(r) = r^2 \log((\operatorname{arth} r)/r)$. Then

$$H_9'(r) = r \frac{r - r'^2 \operatorname{arth} r}{r'^2 \operatorname{arth} r} [1 + 2f_8(r)] = r^3 [f_2(r) + 2f_7(r)], \quad (4.3)$$

which is strictly increasing from $(0, 1)$ onto $(0, \infty)$ by Lemma 2.1(2) and (7), so that H_9 is strictly increasing and convex on $(0, 1)$. If $c > 1/4$, then $g(r)$ can be written as

$$g(r) = H_3(r) + \left(c - \frac{1}{4} \right) H_9(r),$$

which is a sum of two strictly increasing and convex functions. Hence, if $c \geq 1/4$, then g is convex on $(0, 1)$.

Finally, if $c \leq 1/320$, then by (4.1), we have

$$\begin{aligned}
-g'(r) &= r^3 \left(\frac{2}{r^2} \log \frac{\operatorname{arth} r}{r} + \frac{r - r' \operatorname{arth} r}{r^2 r' \operatorname{arth} r} \right) \left[\frac{1}{4} H_1(r) - c \right] \\
&= r^3 \cdot [f_2(r) + 2f_7(r)] \cdot \left[\frac{1}{4} H_1(r) - c \right],
\end{aligned} \quad (4.4)$$

which is a product of three positive and strictly increasing functions on $(0, 1)$ by Lemma 2.1(2) and (7), and by the monotonicity of H_1 . Hence g is concave on $(0, 1)$ provided that $c \leq 1/320$. \square

3.2 Proof of Theorem 1.7

Let H_1, H_3, H_8 and H_9 be as in the Proof of Theorem 1.6.

(1) Clearly, $g_1(r) = H_3(r)/r^2$. Since

$$H_3'(r) \left[\frac{d}{dr} (r^2) \right]^{-1} = \frac{H_3'(r)}{2r} = \frac{1}{4} H_8(x) \quad (4.5)$$

by (4.2), the monotonicity of g_1 follows from that of H_8 and [9, Theorem 1.25].

Clearly, $g_1(1^-) = H_3(1^-) = \log(\pi/2)$. By l'Hôpital's rule and (4.5), we obtain

$$g_1(0^+) = \lim_{r \rightarrow 0} \frac{H_3(r)}{r^2} = \frac{1}{4} H_8(0^+) = 0.$$

Observe that $g_2(r) = H_3(r)/H_9(r)$ with $H_3(0^+) = H_9(0^+) = 0$. It follows from (4.1) and (4.3) that

$$H_3'(r)/H_9'(r) = [1 - H_1(r)]/4, \quad (4.6)$$

which is strictly decreasing from $(0, 1)$ onto $(0, 79/320)$ by the monotonicity property of H_1 . Hence the monotonicity of g_2 follows from [9, Theorem 1.25].

By l'Hôpital's rule and (4.6), $g_2(0^+) = 79/320$. Since $H_3(1^-) = \log(\pi/4)$ by part (1), and since $H_9(1^-) = \infty$, we obtain the limiting value $g_2(1^-) = 0$.

(2) For $r \in (0, 1)$, let

$$H_{10}(r) = -g(r)|_{c=1/320} = \log \frac{2\mathcal{K}(r)}{\pi} - \left(\frac{3}{4} + \frac{1}{320}r^2 \right) \log \frac{\operatorname{arth} r}{r}.$$

Then $H_9(0^+) = H_{10}(0^+) = 0$, $g_3(r) = H_{10}(r)/H_9(r)$, and it follows from (4.3) and (4.4) that

$$\frac{H_{10}'(r)}{H_9'(r)} = \frac{1}{4} \left[H_1(r) - \frac{1}{80} \right], \quad (4.7)$$

which is strictly increasing on $(0, 1)$ by the monotonicity property of H_1 . Hence the monotonicity of g_3 follows from [9, Theorem 1.25].

By l'Hôpital's rule, (4.7), and Corollary 3.3, we obtain the limiting values $g_3(0^+) = 0$ and

$$\begin{aligned} g_3(1^-) &= \lim_{r \rightarrow 1} \frac{\log(2\mathcal{K}/\pi)}{\log((\operatorname{arth} r)/r)} - \frac{3}{4} - \frac{1}{320} \\ &= \lim_{r \rightarrow 1} \frac{[\mathcal{E}(r) - r'^2 \mathcal{K}(r)] \operatorname{arth} r}{(r - r'^2 \operatorname{arth} r) \mathcal{K}(r)} - \frac{241}{320} = \frac{79}{320}. \end{aligned}$$

Next, let f_7 be as in Lemma 2.1. Then

$$g_4(r) = \left(\frac{1}{r^2} \log \frac{\operatorname{arth} r}{r} \right) g_3(r) = f_7(r) g_3(r),$$

and hence the result for g_4 follows from Lemma 2.1(7) and the property of g_3 .

(3) It is clear that

$$f(r) = \frac{1}{4} - g_2(r) \quad \text{and} \quad G(r) = r^2 f(r) + \frac{3}{4}.$$

Hence the results for f and G follow from part (1).

The assertion on (1.17) is obvious. The double inequality (1.18) follows from the property of g_1 . \square

3.3 Proof of Theorem 1.8

(1) The monotonicity and limiting values of h_1 were obtained in [8, Theorem 3.10]. By differentiation,

$$h_1'(r) = \frac{\mathcal{E}(r) \operatorname{arth} r - r \mathcal{K}(r)}{r'^2 \operatorname{arth}^2 r} = -H_{11}(r) H_{12}(r), \quad (4.8)$$

where

$$H_{11}(r) = r \left(\frac{r}{r' \operatorname{arth} r} \right)^2 \quad \text{and} \quad H_{12}(r) = \frac{r \mathcal{K}(r) - \mathcal{E}(r) \operatorname{arth} r}{r^3}.$$

By [22, Lemma 3(1)], H_{11} is strictly increasing from $(0, 1)$ onto $(0, \infty)$. Let $H_{13}(r) = r \mathcal{K}(r) - \mathcal{E}(r) \operatorname{arth} r$ and $H_{14}(r)$. Then $H_{13}(0) = H_{14}(0) = 0$, $H_{12}(r) = H_{13}(r)/H_{14}(r)$, and

$$\frac{H_{13}'(r)}{H_{14}'(r)} = \frac{\mathcal{K}(r) - \mathcal{E}(r)}{3r^2} \cdot \frac{\operatorname{arth} r}{r}$$

which is strictly increasing from $(0, 1)$ onto $(\pi/12, \infty)$ by [6, Lemma 5.2 (3)], so that H_{12} is positive and strictly increasing on $(0, 1)$ by [9, Theorem 1.25]. Consequently, it follows from (4.8) that h_1' is strictly decreasing on $(0, 1)$. This yields the concavity of h_1 .

(2) Let c_n be as in Lemma 2.4. Then by Lemma 2.4(1),

$$\tilde{a}_{k+1} = \frac{1}{2k+3} \left[\frac{3}{4} - (2k+3)a_{k+1} \right] = \frac{1}{2k+3} \left(\frac{3}{4} - c_k \right) > 0 \quad (4.9)$$

for $k \in \mathbb{N}_0$. It follows from (1.3)–(1.5) that

$$\begin{aligned} h_2(r) &= \frac{1}{r^{2(n+1)}} \left(\frac{3}{4} \sum_{n=0}^{\infty} \frac{1}{2n+1} r^{2n} - \sum_{n=0}^{\infty} a_n r^{2n} - \sum_{k=0}^n \tilde{a}_k r^{2k} \right) \\ &= \frac{1}{r^{2(n+1)}} \left(\sum_{n=0}^{\infty} \tilde{a}_n r^{2n} - \sum_{k=0}^n \tilde{a}_k r^{2k} \right) = \sum_{k=0}^{\infty} \tilde{a}_{k+n+1} r^{2k}. \end{aligned} \quad (4.10)$$

Hence by (4.9), all the coefficients of the Maclaurin series of h_2 are positive, so that h_2 is strictly absolutely monotone on $(0, 1)$.

By (4.9) and (4.10), $h_2(0^+) = \tilde{a}_{n+1} > 0$, and by (2.2), we obtain

$$\begin{aligned} h_2(1^-) &= \lim_{r \rightarrow 1} \left(\frac{3}{8r} \log \frac{1+r}{1-r} - \frac{2}{\pi} \log \frac{4}{r'} \right) - \sum_{k=0}^n \tilde{a}_k \\ &= \lim_{r \rightarrow 1} \left(\frac{3}{8r} - \frac{1}{\pi} \right) \log \frac{1}{1-r} - \sum_{k=0}^n \tilde{a}_k - \frac{8-\pi}{8\pi} \log 8 = \infty. \end{aligned}$$

(3) It follows from (1.5) and (4.10)

$$h_3(r) = \left(\sum_{k=0}^{\infty} \tilde{a}_{k+n+1} r^{2k} \right) \left(\sum_{k=0}^{\infty} \frac{1}{2k+1} r^{2k} \right)^{-1}. \quad (4.11)$$

The ratio of the coefficients of the two series expansions in (4.11) is equal to

$$\frac{2k+1}{2(k+n)+3} \left(\frac{3}{4} - c_{k+n} \right) = \left[1 - 2 \frac{n+1}{2(k+n)+3} \right] \left(\frac{3}{4} - c_{k+n} \right). \quad (4.12)$$

By Lemma 2.4(1), the left side of (4.12) is a product of two factors which are both positive and strictly increasing in $k \in \mathbb{N}_0$. Hence the monotonicity of h_3 follows from [19, Lemma 2.1].

Clearly, $h_3(0^+) = h_2(0^+) = \tilde{a}_{n+1}$. By part (1), we obtain the limiting value

$$h_3(1^-) = \lim_{r \rightarrow 1} \left[\frac{3}{4} - \frac{2}{\pi} h_1(r) - \frac{r}{\text{arth}r} \sum_{k=0}^n \tilde{a}_k \right] = \frac{3}{4} - \frac{2}{\pi}.$$

(4) Let d_n be as in Lemma 2.4. Then by (1.3)–(1.5), we have

$$\begin{aligned} & \frac{\pi}{2} \left(1 - \frac{1}{12} r^2 - \frac{91}{2880} r^4 \right) \frac{\text{arth}r}{r} - \mathcal{H}(r) \\ &= \frac{\pi}{2} \left(\sum_{n=1}^{\infty} \frac{r^{2n}}{2n+1} - \frac{1}{12} r^2 \sum_{n=0}^{\infty} \frac{r^{2n}}{2n+1} - \frac{91}{2880} r^4 \sum_{n=0}^{\infty} \frac{r^{2n}}{2n+1} - \sum_{n=1}^{\infty} a_n r^{2n} \right) \\ &= \frac{\pi}{2} \left(\sum_{n=2}^{\infty} \frac{r^{2n}}{2n+1} - \frac{1}{12} r^2 \sum_{n=1}^{\infty} \frac{r^{2n}}{2n+1} - \frac{91}{2880} r^4 \sum_{n=0}^{\infty} \frac{r^{2n}}{2n+1} - \sum_{n=2}^{\infty} a_n r^{2n} \right) \\ &= \frac{\pi}{2} r^4 \sum_{n=0}^{\infty} \left[\frac{1}{2n+5} - \frac{1}{12(2n+3)} - \frac{91}{2880(2n+1)} - a_{n+2} \right] r^{2n} \\ &= \frac{\pi}{5760} r^4 \sum_{n=0}^{\infty} \left[\frac{10196n^2 + 18704n + 6075}{(2n+5)(2n+3)(2n+1)} - 2880a_{n+2} \right] r^{2n} \\ &= \frac{\pi}{5760} r^6 \sum_{n=0}^{\infty} \left[\frac{10196n^2 + 39096n + 34975}{(2n+7)(2n+5)(2n+3)} - 2880a_{n+3} \right] r^{2n} \\ &= \frac{\pi}{5760} r^6 \sum_{n=0}^{\infty} \frac{d_n}{2n+1} r^{2n}, \end{aligned}$$

and hence

$$\begin{aligned} h_4(r) &= \frac{1}{(\operatorname{arth} r)/r} \cdot \frac{1}{r^6} \left[\frac{\pi}{2} \left(1 - \frac{1}{12} r^2 - \frac{91}{2880} r^4 \right) \frac{\operatorname{arth} r}{r} - \mathcal{K}(r) \right] \\ &= \frac{\pi}{5760} \left(\sum_{n=0}^{\infty} \frac{d_n}{2n+1} r^{2n} \right) \cdot \left(\sum_{n=0}^{\infty} \frac{1}{2n+1} r^{2n} \right)^{-1}. \end{aligned} \quad (4.13)$$

Clearly, the ratio of the coefficients of the two series expansions in (4.13) is d_n . Hence the monotonicity of h_4 follows from (4.13), Lemma 2.4(3) and [19, Lemma 2.1].

By (4.13), $h_4(0^+) = \pi d_0/5760 = 871\pi/96768$, and by part (1),

$$h_4(1^-) = \frac{\pi}{2} \left(1 - \frac{1}{12} - \frac{91}{2880} \right) - h_1(1^-) = \frac{2549\pi}{5760} - 1.$$

(5) The double inequality (1.19) ((1.20)) follows from part (3) (part (4), respectively). Taking $n = 1$ in (1.19), we obtain the double inequality (1.21).

By parts (3)–(4), the remaining conclusion is clear. \square

5 Concluding Remark

(i) The double inequality (1.17) can also be obtained from Theorem 1.6, or Theorem 1.7(1), or Theorem 1.7(2).

(ii) The upper bound of $\mathcal{K}(r)$ given in (1.17) coincides with that given in (1.10) obtained by Kupán and Szász. However, our result is implied by a stronger conclusion, and our proof is more direct and simpler than theirs. In addition, it was not indicated in [17] that the constant $\beta = 1/4$ in (1.17) is best possible.

On the other hand, since the function

$$h_5(r) \equiv \left\{ \left(\frac{\pi}{2} \right)^{1-r^2} \left(\frac{\operatorname{arth} r}{r} \right)^{3/4+r^2/4} \left[\frac{\pi}{2} \left(\frac{\operatorname{arth} r}{r} \right)^{3/4+r^2/320} \right]^{-1} \right\}^{1/r^2} = \frac{2}{\pi} \left(\frac{\operatorname{arth} r}{r} \right)^{79/320}$$

is strictly increasing from $(0, 1)$ onto $(2/\pi, \infty)$, there exists a number $r_0 \in (0, 1)$ satisfying the condition $(\operatorname{arth} r_0)/r_0 = (\pi/2)^{320/79}$ such that the lower bound given in (1.17) ((1.18)) is better than that given in (1.18) ((1.17)) for $r \in (0, r_0)$ ($r \in (r_0, 1)$, respectively).

(iii) Let h_4 be as in Theorem 1.8. Then by Theorem 1.8(4), the function

$$h_6(r) \equiv \frac{r\mathcal{K}(r)}{\operatorname{arth} r} + \frac{\pi}{24} r^2 + \frac{91\pi}{5760} r^4 = \frac{\pi}{2} - r^6 h_4(r)$$

is strictly decreasing from $(0, 1)$ onto $(1 + 331\pi/5760, \pi/2)$. Therefore, Theorem 1.8(4) remarkably improves the monotonically decreasing property of $h_1(r) = r\mathcal{K}(r)/\operatorname{arth} r$ obtained in [8, Theorem 3.10].

(iv) It follows from Theorem 1.8(1) that for all $r \in (0, 1)$,

$$\left[1 - \left(1 - \frac{2}{\pi} \right) r \right] \frac{\operatorname{arth} r}{r} < \frac{2}{\pi} \mathcal{K}(r) < \frac{\operatorname{arth} r}{r}. \quad (5.1)$$

Clearly, each of the upper bounds of $2\mathcal{K}(r)/\pi$ given in (1.17)–(1.18) and in (1.20)–(1.21) is better than the upper bound given in (5.1).

Let P_3 , α and δ be as in Theorem 1.8(5), and for $r \in (0, 1)$, let

$$\begin{aligned} h_7(r) &= \frac{1}{r} \left\{ [P_3(r) - \alpha r^6] - \left[1 - \left(1 - \frac{2}{\pi} \right) r \right] \right\} \quad \text{and} \\ h_8(r) &= \frac{1}{r} \left\{ \left[\frac{1}{4} + \frac{3}{4} (1 - \delta r^4) \frac{\operatorname{arth} r}{r} \right] - \left[1 - \left(1 - \frac{2}{\pi} \right) r \right] \frac{\operatorname{arth} r}{r} \right\}. \end{aligned}$$

Then

$$h_7(r) = 1 - \frac{2}{\pi} - \frac{1}{12}r - \frac{91}{2880}r^3 - \alpha r^5 \quad \text{and} \quad (5.2)$$

$$\begin{aligned} h_8(r) &= \frac{1}{r} \left\{ \frac{1}{4} + \left[\left(1 - \frac{2}{\pi}\right)r - \left(\frac{3}{4} - \frac{2}{\pi}\right)r^4 - \frac{1}{4} \right] \frac{\operatorname{arth}r}{r} \right\} \\ &= \frac{1}{4r} \left[1 - \frac{(1-r)\operatorname{arth}r}{r} \right] + \left(\frac{3}{4} - \frac{2}{\pi} \right) (1-r^3) \frac{\operatorname{arth}r}{r}, \end{aligned} \quad (5.3)$$

from which we can easily see that h_7 is strictly decreasing from $(0, 1)$ onto $(0, 1 - 2/\pi)$, and $h_8(r) > 0$ for $r \in (0, 1)$ since the function

$$r \mapsto \frac{(1-r)\operatorname{arth}r}{r} = \frac{1}{1+r} \frac{r'^2 \operatorname{arth}r}{r}$$

is strictly decreasing from $(0, 1)$ onto itself by [22, Lemma 3(1)]. Consequently, each of the lower bounds of $2\mathcal{K}(r)/\pi$ given in (1.20)–(1.21) is better than the lower bound of this function given in (5.1).

(v) The pair of lower bounds, and the pair of upper bounds of $\mathcal{K}(r)$ given in (1.20)–(1.21) are both not directly comparable on $(0, 1)$. In order to explain this, we let

$$\begin{aligned} h_9(r) &= \left[1 - \frac{1}{12}r^2 - \frac{91}{2880}r^4 - \left(\frac{2549}{2880} - \frac{2}{\pi}\right)r^6 \right] \frac{\operatorname{arth}r}{r} - \left\{ \frac{1}{4} + \frac{3}{4} \left[1 - \left(1 - \frac{8}{3\pi}\right)r^4 \right] \frac{\operatorname{arth}r}{r} \right\} \\ &= \left[\frac{1}{4} - \frac{1}{12}r^2 + \left(\frac{2069}{2880} - \frac{2}{\pi}\right)r^4 - \left(\frac{2549}{2880} - \frac{2}{\pi}\right)r^6 \right] \frac{\operatorname{arth}r}{r} - \frac{1}{4}, \\ h_{10}(r) &= \left(1 - \frac{1}{12}r^2 - \frac{91}{2880}r^4 - \frac{871}{48384}r^6 \right) \frac{\operatorname{arth}r}{r} - \left[\frac{1}{4} + \left(\frac{3}{4} - \frac{3}{320}r^4\right) \frac{\operatorname{arth}r}{r} \right] \\ &= \frac{1}{4} \left(\frac{\operatorname{arth}r}{r} - 1 \right) - r^2 \left(\frac{1}{12} + \frac{1}{45}r^2 + \frac{871}{48384}r^4 \right) \frac{\operatorname{arth}r}{r}. \end{aligned}$$

Applying l'Hôpital's, one can easily obtain the limiting value

$$\lim_{r \rightarrow 1} \frac{1}{r'^2} \left[\frac{1}{4} - \frac{1}{12}r^2 + \left(\frac{2069}{2880} - \frac{2}{\pi}\right)r^4 - \left(\frac{2549}{2880} - \frac{2}{\pi}\right)r^6 \right] = \frac{1}{12},$$

and hence $h_9(1^-) = -1/4$. On the other hand, by (1.5), we obtain

$$\frac{3}{r^4} h_9(r) = \sum_{n=0}^{\infty} \left[\frac{n+1}{(2n+3)(2n+5)} + \left(\frac{2069}{960} - \frac{6}{\pi}\right) \frac{1}{2n+1} \right] r^{2n} - \left(\frac{2549}{960} - \frac{6}{\pi}\right) \sum_{n=0}^{\infty} \frac{r^{2(n+1)}}{2n+1},$$

so that

$$\lim_{r \rightarrow 0} \frac{3}{r^4} h_9(r) = \frac{1}{15} + \frac{2069}{960} - \frac{6}{\pi} = \frac{2133}{960} - \frac{6}{\pi} = 0.3120156 \dots > 0.$$

Consequently, there exist two numbers $r_1, r_2 \in (0, 1)$ such that $h_9(r)$ is positive (negative) for $r \in (0, r_1)$ ($r \in (r_2, 1)$, respectively). This shows that the lower bounds of $\mathcal{K}(r)$ given in (1.20)–(1.21) are not directly comparable on $(0, 1)$.

Similarly, we have

$$\frac{241920}{r^6} h_{10}(r) = \sum_{n=0}^{\infty} \frac{244712n^3 + 749388n^2 + 408406n - 161595}{(2n+1)(2n+3)(2n+5)(2n+7)} r^{2n},$$

by which we obtain $\lim_{r \rightarrow 0} [241920 h_{10}(r)/r^6] = -161595/105 = -1539$. On the other hand, we have

$$\frac{h_{10}(r)}{r \operatorname{arth}r} = \frac{\operatorname{arth}r - r}{4r^2 \operatorname{arth}r} - \left(\frac{1}{12} + \frac{1}{45}r^2 + \frac{871}{48384}r^4 \right),$$

from which we see that

$$\lim_{r \rightarrow 1} \frac{h_{10}(r)}{r \operatorname{arth} r} = \frac{1}{4} - \frac{1}{12} - \frac{1}{45} - \frac{871}{48384} = \frac{30589}{241920} > 0.$$

Therefore, there exist two numbers $r_3, r_4 \in (0, 1)$ such that $h_{10}(r)$ is negative (positive) for $r \in (0, r_3)$ ($r \in (r_4, 1)$, respectively), showing that the upper bounds of $\mathcal{K}(r)$ given in (1.20)–(1.21) are also not directly comparable on $(0, 1)$.

(vi) Let f, G, g_1 and g_4 be as in Theorem 1.7. By computation, we have

$$\begin{aligned} f(r) = & \frac{1}{320} + \frac{517}{201600}r^2 + \frac{767341}{387072000}r^4 + \frac{4277471797}{2682408960000}r^6 \\ & + \frac{1851483120061}{1394852659200000}r^8 + \frac{2989339649544551}{2636271525888000000}r^{10} + \dots, \end{aligned} \quad (5.4)$$

$$\begin{aligned} G(r) = & \frac{3}{4} + \frac{1}{320}r^2 + \frac{517}{201600}r^4 + \frac{767341}{387072000}r^6 + \frac{4277471797}{2682408960000}r^8 \\ & + \frac{1851483120061}{1394852659200000}r^{10} + \frac{2989339649544551}{2636271525888000000}r^{12} + \dots, \end{aligned} \quad (5.5)$$

$$\begin{aligned} h_{11}(r) \equiv & \frac{g_1(r)}{r^2} = \frac{1}{r^4} \left[\frac{3+r^2}{4} \log \frac{\operatorname{arth} r}{r} - \log \frac{2\mathcal{K}(r)}{\pi} \right] \\ = & \frac{79}{960} + \frac{421}{12096}r^2 + \frac{690961}{33177600}r^4 + \frac{18414493}{1277337600}r^6 \\ & + \frac{164673431213}{15216574464000}r^8 + \dots \end{aligned} \quad (5.6)$$

and

$$\begin{aligned} h_{12}(r) \equiv & \frac{g_4(r)}{r^2} = \frac{1}{r^6} \left[\log \frac{2\mathcal{K}(r)}{\pi} - \left(\frac{3}{4} + \frac{1}{320}r^2 \right) \log \frac{\operatorname{arth} r}{r} \right] \\ = & \frac{517}{604800} + \frac{239497}{232243200}r^2 + \frac{741527}{709632000}r^4 \\ & + \frac{168874886801}{167382319104000}r^6 + \frac{2405137262477}{2510734786560000}r^8 + \dots \end{aligned} \quad (5.7)$$

Our computation seems to show that the following conjectures are true.

Conjecture 5.1. (1) The function f in Theorem 1.7(3) is convex on $(0, 1)$.

(2) The functions h_{11} and h_{12} are both strictly increasing and convex on $(0, 1)$, with ranges $(79/960, \log(\pi/2))$ and $(517/604800, \infty)$, respectively.

(3) Let f_7 be as in Lemma 2.1(7), and let $h_{13}(r) = r^{-2} \log(2\mathcal{K}(r)/\pi)$ for $r \in (0, 1)$. Then the coefficients of the Maclaurin series expansions of the functions $f, G, f_7, h_{11}, h_{12}$ and h_{13} are all positive, and in particular, these functions are all absolutely monotone on $(0, 1)$.

If these conjectures are true, then the inequalities in Theorem 1.7(3) can be improved. For example, the following double inequalities

$$\left(\frac{\operatorname{arth} r}{r} \right)^{\frac{3}{4} + \frac{1}{320}r^2 + \frac{517}{201600}r^4} < \frac{2\mathcal{K}(r)}{\pi} < \left(\frac{\operatorname{arth} r}{r} \right)^{\frac{3}{4} + \frac{1}{320}r^2 + \frac{79}{320}r^4} \quad (5.8)$$

and

$$\tau^{r^4} \left(\frac{\operatorname{arth} r}{r} \right)^{\frac{3}{4} + \frac{1}{4}r^2} < \frac{2\mathcal{K}(r)}{\pi} < \mu^{r^4} \left(\frac{\operatorname{arth} r}{r} \right)^{\frac{3}{4} + \frac{1}{4}r^2} \quad (5.9)$$

will be valid for all $r \in (0, 1)$, where $\tau = 2/\pi = 0.6366197 \dots$ and $\mu = e^{-79/960} = 0.9210032 \dots$.

(vii) We also would like to put forward the following

Open Problem 5.2. *What are the analogues of Theorems 1.6–1.8 for the generalized complete elliptic integrals (or even for the zero-balanced hypergeometric functions)?*

Acknowledgments

- (1) This work is supported by the National Natural Science Foundation of China (Grant No.11771400).
- (2) The two universities at which the corresponding author works are listed in no particular order.

Competing Interests

The authors declare that they have no competing interests.

References

- [1] R. Askey, in: P. Duren (Ed.), *Handbooks of Special Functions, A century of Mathematics in America, Part III*, Amer. Math. Soc. (1989)
- [2] G. Andrews, R. Askey and R. Roy, *Special Functions, Encyclopedia of Mathematics and Its Applications*, vol. 71, Cambridge Univ Press (1999)
- [3] H. Alzer and S.L. Qiu, Monotonicity theorems and inequalities for the complete elliptic integrals, *J. Comput. Appl. Math.* 172(2), 289–312 (2004)
- [4] M. Abramowitz and I.A. Stegun, *Handbook of Mathematical Functions With Formulas, Graphs and Mathematical Tables*. New York: Dover, 1965. Tenth Printing, with corrections (1972)
- [5] G.D. Anderson, S.L. Qiu, and M.K. Vamanamurthy, Elliptic integrals inequalities, with applications, *Constr. Approx.* 14, 195–207 (1998)
- [6] G.D. Anderson, S.L. Qiu, M.K. Vamanamurthy, and M. Vuorinen, Generalized elliptic integrals and modular equations, *Pacific J. Math.* 192(1), 1–37 (2000)
- [7] G.D. Anderson, M.K. Vamanamurthy, and M. Vuorinen, Functional inequalities for complete elliptic integrals and their ratios, *SIAM J. Math. Anal.* 21(2), 536–549 (1990)
- [8] G.D. Anderson, M.K. Vamanamurthy, and M. Vuorinen, Functional inequalities for hypergeometric functions and complete elliptic integrals, *SIAM J. Math. Anal.* 21(2), 512–524 (1992)
- [9] G.D. Anderson, M.K. Vamanamurthy, M. Vuorinen, *Conformal Invariants, Inequalities, and Quasiconformal Maps*, John Wiley and Sons, New York (1997)
- [10] A. Baricz, Turán type inequalities for generalized complete elliptic integrals, *Math. Z.* 256(4), 895–911 (2007)
- [11] B.C. Berndt, *Ramanujan’s Notebooks, Part II*, Springer-Verlag, New York (1989)
- [12] J.M. Borwein and P.B. Borwein, *Pi and the AGM*, John Wiley & Sons, New York (1987)
- [13] B.C. Carlson, *Special Functions of Applied Mathematics*, Academic Press, New York (1977)
- [14] Y.-M. Chu, M.-K. Wang and Y.-F. Qiu, On Alzer and Qiu’s conjecture for complete elliptic integral and inverse hyperbolic tangent function, *Abstract and Applied Analysis* 2011, Article ID 697547, 7 pages (<https://doi.org/10.1155/2011/697547>)
- [15] Y.M. Chu, M.K. Wang, S.L. Qiu, and Y.P. Jiang, Bounds for complete elliptic integrals of the second kind with applications, *Comput. Math. Appl.* 63(7), 117–1184 (2012)

- [16] T.R. Huang, S.L. Qiu and X.Y. Ma, Monotonicity properties and inequalities for the generalized elliptic integral of the first kind, *J. Math. Anal. Appl.* 469(1), 95–116 (2019)
- [17] Pál A. Kupán and Róbert Szász, About bounds for the elliptic integral of the first kind, *Rev. Anal. Numér. Théor. Approx.* 41(2), 149–156 (2012)
- [18] F.W.J. Olver, D.W. Lozier, R.F. Boisvert, et al., *NIST Handbook of Mathematical Functions*, Cambridge University Press, Cambridge (2010)
- [19] S. Ponnusamy, M. Vuorinen, Asymptotic expansions and inequalities for hypergeometric functions, *Mathematika* 44(2), 278–301 (1997)
- [20] S.L. Qiu, X.Y. Ma and Y.M. Chu, Sharp Landen transformation inequalities for hypergeometric functions, with applications, *J. Math. Anal. Appl.* 474(2), 1306–1337 (2019)
- [21] S.L. Qiu and M.K. Vamanamurthy, Sharp estimates for complete elliptic integrals, *SIAM J. Math. Anal.* 27(3), 823–834 (1996)
- [22] S.L. Qiu, M.K. Vamanamurthy and M. Vuorinen, Some inequalities for the Hersch-Pfluger distortion function. *J. of Inequal. & Appl.* 4, 115–139 (1999)
- [23] S.L. Qiu and M. Vuorinen, *Handbook of Complex Analysis: Special Function in Geometric Function Theory*, pp.621–659. Elsevier B.V., Amsterdam (2005)
- [24] S.L. Qiu and M. Vuorinen, Duplication inequalities for the ratios of hypergeometric functions, *Forum Math.* 12(1), 109–133 (2000)
- [25] G.D. Wang, X.H. Zhang, Y.M. Chu and S.L. Qiu, Complete elliptic integrals and the Hersch-Pfluger distortion function (in Chinese), *Acta Mathematica Scientia* 4, 731–734 (2008)

Authors' addresses:

Song-Liang Qiu:

Department of Mathematics, Lishui University, Lishui 323000, Zhejiang, China;

Department of Mathematics, Zhejiang Sci-Tech University, Hangzhou 310018, Zhejiang, China

E-mail address: sl_qiu@zstu.edu.cn

Qi Bao:

School of Mathematical Sciences, East China Normal University, Shanghai 200241, China

E-mail address: 52205500010@stu.ecnu.edu.cn

Xiao-Yan Ma:

Department of Mathematics, Zhejiang Sci-Tech University, Hangzhou 310018, Zhejiang, China

E-mail address: mxy@zstu.edu.cn

Hong-Biao Jiang:

Department of Mathematics, Lishui University, Lishui 323000, Zhejiang, China

E-mail address: lsxybj@126.com