

SIMPLICIAL RESOLUTIONS FOR THE SECOND POWER OF SQUARE-FREE MONOMIAL IDEALS

SUSAN M. COOPER, SABINE EL KHOURY, SARA FARIDI, SARAH MAYES-TANG, SUSAN MOREY,
LIANA M. ŞEGA, AND SANDRA SPIROFF

ABSTRACT. Given a square-free monomial ideal I , we define a simplicial complex labeled by the generators of I^2 which supports a free resolution of I^2 . As a consequence, we obtain (sharp) upper bounds on the Betti numbers of the second power of any square-free monomial ideal.

1. INTRODUCTION

The question of finding, or even effectively bounding, the Betti numbers of an ideal in a commutative ring is a difficult one. Even more complicated is using the structure of an ideal I to find information about the Betti numbers of its powers I^r : predicting something as basic as the minimal number of generators of I^r is a difficult problem.

Taylor's thesis [10] described a free resolution of any ideal minimally generated by q monomials using the simplicial chain complex of a simplex with q vertices. Taylor's construction, though often far from minimal, produces a resolution of *every* monomial ideal I . It gives upper bounds $\binom{q}{i+1} \geq \beta_i(I)$ for the Betti numbers of I where $\binom{q}{i+1}$ is the number of i -faces of a q -simplex. If I is generated by q monomials and r is a positive integer, then the number of generators of I^r generally grows exponentially and as a result, so do the bounds on the Betti numbers of I^r given by Taylor's resolution.

In this paper, we focus on the case where $r = 2$ and I is a square-free monomial ideal with q generators. In this case, we know that I^2 can be generated by at most $\binom{q+1}{2}$ monomials, and hence has a Taylor resolution supported on a simplex with at most $\binom{q+1}{2}$ vertices. The question that we address in this paper is: can we find a subcomplex of this simplex whose simplicial chain complex yields a free resolution of I^2 ? Such a resolution would be closer to minimal than the Taylor resolution.

We answer this question by constructing a simplicial complex on $\binom{q+1}{2}$ vertices which we call \mathbb{L}_q^2 in honor of the Lyubeznik resolution [9] which was our inspiration. While \mathbb{L}_q^2 has the same number of vertices as the $\binom{q+1}{2}$ -simplex, it is significantly smaller because it has far fewer faces. For a given square-free monomial ideal I , we can use further deletions of \mathbb{L}_q^2 specific to the generators of I to show that \mathbb{L}_q^2 has an induced subcomplex $\mathbb{L}^2(I)$ which supports a free resolution of I^2 . As a result, we find (sharp) upper bounds on the Betti numbers of the second power of any square-free monomial ideal. These bounds are often significantly smaller than the bounds provided by the Taylor resolution (see Section 4).

2010 *Mathematics Subject Classification.* 13D02; 13F55.

Key words and phrases. powers of ideals; simplicial complex; Betti numbers; free resolutions; monomial ideals.

Section 2 lays out the notation and terminology used in the paper including the construction of simplicial resolutions. In Section 3, we describe the complexes \mathbb{L}_q^2 (Definition 3.1) and $\mathbb{L}^2(I)$ (Definition 3.4) and prove that $\mathbb{L}^2(I)$ supports a free resolution of I^2 when I is a square-free monomial ideal (Theorem 3.9). Section 4 provides results on the bounds on the Betti numbers that follow from the main results.

This paper is part of a larger project [3] to study resolutions of powers of monomial ideals, which the authors started during the 2019 Banff workshop “Women in Commutative Algebra”.

2. BACKGROUND

Throughout this paper we let $S = k[x_1, \dots, x_n]$ be a polynomial ring over a field k . In this section we briefly recall some necessary background about simplicial complexes.

A **simplicial complex** Δ over a **vertex set** V is a set of subsets of V such that if $F \in \Delta$ and $G \subseteq F$ then $G \in \Delta$. An element σ of Δ is called a **face** and the maximal faces under inclusion are called **facets**. A simplicial complex can be uniquely determined by its facets, and we use the notation

$$\Delta = \langle F_0, \dots, F_q \rangle$$

to describe a simplicial complex whose facets are F_0, \dots, F_q .

The **dimension** of a face F in Δ is $\dim(F) = |F| - 1$, and the dimension of Δ is the maximum of the dimensions of its faces.

A simplicial complex with one facet is called a **simplex**.

If $W \subseteq V$, the subcomplex

$$\Delta_W = \{\sigma \in \Delta \mid \sigma \subseteq W\}$$

is called the **induced subcomplex** of Δ on W .

If Δ is a simplicial complex with vertex v , then when we **delete** v from Δ we obtain the simplicial complex


$$\Delta \setminus \{v\} = \{\sigma \in \Delta \mid v \notin \sigma\}.$$

A facet F of Δ is said to be a **leaf** if it is the only facet of Δ , or there is a different facet G of Δ , called a **joint**, such that

$$F \cap H \subseteq G$$

for all facets $H \neq F$. The joint G in this definition is not unique ([4]). A simplicial complex Δ is a **quasi-forest** if the facets of Δ can be ordered as F_0, \dots, F_q such that for $i = 0, \dots, q$, the facet F_i is a leaf of the simplicial complex $\langle F_0, \dots, F_i \rangle$. A connected quasi-forest is called a **quasi-tree** ([11]).

Example 2.1. The simplicial complex below is a quasi-tree, with leaf order: F_0, F_1, F_2, F_3 , meaning that each F_i is a leaf of $\langle F_0, \dots, F_i \rangle$. Note that in this case, the joint of F_i is F_0 for every $i \geq 1$.



This complex is in fact \mathbb{L}_3^2 as we will see later in Example 3.2.

If I is minimally generated by monomials m_1, \dots, m_q in S , a **minimal free resolution** of I is a (unique up to isomorphism) exact sequence of free S -modules


$$0 \rightarrow S^{\beta_p} \rightarrow S^{\beta_{p-1}} \rightarrow \dots \rightarrow S^{\beta_1} \rightarrow S^{\beta_0} \rightarrow I \rightarrow 0$$

where $p \in \mathbb{N}$, $\beta_0 = q$ and for each $i \in \{1, \dots, p\}$, β_i is the smallest possible rank of a free module in the i -th spot of any free resolution of I . The β_i , called the **Betti numbers** of I , are invariants of the ideal I .

Finding ways to describe a free resolution of a given ideal is an open and active area of research. For monomial ideals, combinatorics plays a big role. In her thesis in the 1960's, Diana Taylor introduced a method of labeling the faces of a simplex Δ with monomials, and then used this labeling to turn the simplicial chain complex of Δ into a free resolution of a monomial ideal. This technique has been generalized to other simplicial complexes by Bayer and Sturmfels [2], among others.

More precisely, if I is minimally generated by monomials m_1, \dots, m_q and Δ is a simplicial complex on q vertices v_1, \dots, v_q , we label each vertex v_i with the monomial m_i , and we label each face of Δ with the least common multiple of the labels of its vertices. Then, if the labeling of Δ satisfies certain properties, the simplicial chain complex of Δ can be "homogenized" using the monomial labels on the faces to give a free resolution of I . In this case, we say that Δ **supports a free resolution** of I and the resulting free resolution is called a **simplicial resolution** of I . Peeva's book [8] details this method for simplicial as well as other topological resolutions.

Example 2.2. Let $I = (x^2, y^2, z^2, xy, xz, yz)$. In the picture below, we label the simplicial complex Δ in Example 2.1 using the generators of I . To make the picture less busy, we have included the labels of the vertices and the facets only.



Our main result Theorem 3.9 will prove that, indeed, Δ does support a free resolution of I . This in particular implies that $\beta_i(I)$ is bounded above by the number of i -faces of Δ , which is the rank of the i -th chain group of Δ . That is,

$$\beta_0(I) \leq 6, \quad \beta_1(I) \leq 9, \quad \beta_2(I) \leq 4, \quad \beta_i(I) = 0 \text{ if } i > 2.$$

We calculate, using Macaulay2 [7], that the actual Betti numbers of I are:

$$\beta_0(I) = 6, \quad \beta_1(I) = 8, \quad \beta_2(I) = 3, \quad \beta_i(I) = 0 \text{ if } i > 2.$$

A major question in the theory of combinatorial resolutions is to determine whether a given simplicial complex supports a free resolution of a given monomial ideal. Taylor proved that a simplex with q vertices *always* supports a free resolution of an ideal with q generators, or in other words, *every* monomial ideal has a Taylor resolution. As a result $\binom{q}{i+1}$ (the number of i -faces of a simplex with q vertices) is an upper bound for $\beta_i(I)$ if I is *any* monomial ideal with q generators. We denote the q -simplex labeled with the q generators of I by $\text{Taylor}(I)$.

Taylor's resolution is usually far from minimal. However, if I is a monomial ideal with a free resolution supported on a (labeled) simplicial complex Δ , then Δ has to be a subcomplex of $\text{Taylor}(I)$. As a result, the question of finding smaller simplicial resolutions of I turns into a question of finding smaller subcomplexes of $\text{Taylor}(I)$ which support a resolution of I .

One of the best known tools to identify such subcomplexes of the Taylor complex is due to Bayer, Peeva, and Sturmfels [1], and reduces the problem to checking acyclicity of induced subcomplexes. This criterion was adapted in [5] to the class of simplicial trees, and then in [3] to quasi-trees. Theorem 2.3 is this latter adaptation, and will be used in the rest of the paper.

If Δ is a subcomplex of $\text{Taylor}(I)$ and \mathbf{m} is a monomial in S , let $\Delta_{\mathbf{m}}$ be the subcomplex of Δ induced on the vertices of Δ whose labels divide \mathbf{m} , and let $\text{LCM}(I)$ denote the set of monomials that are least common multiples of arbitrary subsets of the minimal monomial generating set of I .

Theorem 2.3 ([3] Criterion for quasi-trees supporting resolutions). *Let Δ be a quasi-tree whose vertices are labeled with the monomial generating set of a monomial ideal I in the polynomial ring S over a field k . Then Δ supports a resolution of I if and only if for every monomial \mathbf{m} in $\text{LCM}(I)$, $\Delta_{\mathbf{m}}$ is empty or connected.*

If I is minimally generated by q monomials, then I^2 is minimally generated by at most $\binom{q+1}{2}$ monomials. Our goal in this paper is to find a (smaller) subcomplex of the $\binom{q+1}{2}$ -simplex which produces a free resolution of I^2 , and only depends on q . The quasi-tree \mathbb{L}_q^2 , introduced in the next section, is such a candidate: it has exactly $\binom{q+1}{2}$ vertices, and for any given ideal I with q generators, it has an induced subcomplex $\mathbb{L}^2(I)$ contained in $\text{Taylor}(I^2)$ which supports a free resolution of I^2 .

3. THE QUASI-TREES \mathbb{L}_q^2 AND $\mathbb{L}^2(I)$

For an integer $q \geq 1$ we now give a description of a simplicial complex \mathbb{L}_q^2 , a subcomplex of the $\binom{q+1}{2}$ -simplex. We will show that if I is a monomial ideal generated by q square-free monomials, an induced subcomplex of \mathbb{L}_q^2 , which we denote by $\mathbb{L}^2(I)$, always supports a free resolution of I^2 . The complex \mathbb{L}_q^2 is a far smaller subcomplex of the $\binom{q+1}{2}$ -simplex, and its construction is motivated by the monomial orderings used to build the Lyubeznik complex [9].


Definition 3.1. For an integer $q \geq 3$, the simplicial complex \mathbb{L}_q^2 over the vertex set $\{\ell_{i,j} : 1 \leq i \leq j \leq q\}$ is defined by its facets as:

$$\mathbb{L}_q^2 = \langle \{\ell_{i,j} : 1 \leq j \leq q\}_{1 \leq i \leq q}, \{\ell_{i,j} : 1 \leq i < j \leq q\} \rangle,$$

where we define $\ell_{j,i}$ for $j > i$ by the equality $\ell_{j,i} = \ell_{i,j}$. For $q = 1$ and $q = 2$ we use the same construction but note that $\{\ell_{i,j} : 1 \leq i < j \leq q\}$ is empty for $q = 1$ and is a face but not a facet for $q = 2$.

When $q = 1$, the ideals I and I^r for all $r \geq 2$ are principal and \mathbb{L}_q^2 is a point. When $q = 2$, the complex \mathbb{L}_2^2 has only 2 facets, see Example 3.2. Note that \mathbb{L}_q^2 has $\binom{q+1}{2}$ vertices, which is the number of vertices of the $\binom{q+1}{2}$ -simplex, and, when $q > 2$, it has $q + 1$ facets, where one facet has dimension $\binom{q}{2} - 1$ and the remaining q facets have dimension $q - 1$.

Example 3.2. The complexes \mathbb{L}_3^2 and \mathbb{L}_2^2 are shown on the left and right, respectively.



Proposition 3.3. For $q \geq 1$, \mathbb{L}_q^2 is a quasi-tree.

Proof. If $q = 1$, then \mathbb{L}_q^2 is a simplex of dimension 0, and so is a quasi-tree. If $q = 2$, there are only two facets, namely F_1 and F_2 (as depicted above), and F_2 is a leaf of $\langle F_1, F_2 \rangle$ with joint F_1 , so \mathbb{L}_2^2 is a quasi-tree. For $q \geq 3$, order the facets of \mathbb{L}_q^2 by $F_0 = \{\ell_{i,j} : 1 \leq i < j \leq q\}$, and $F_i = \{\ell_{i,j} : 1 \leq j \leq q\}$ for $1 \leq i \leq q$. By definition, if $i \neq k$ are nonzero, then $F_i \cap F_k = \{\ell_{i,k}\} \subseteq F_0$. Thus each F_i is a leaf of $\langle F_0, \dots, F_i \rangle$ with joint F_0 , and we are done. \square

Given a square-free monomial ideal I , we now define a labeled induced subcomplex of \mathbb{L}_q^2 , denoted $\mathbb{L}^2(I)$, which is obtained by deleting vertices from \mathbb{L}_q^2 .

Definition 3.4 ($\mathbb{L}^2(I)$). For an ideal I minimally generated by the square-free monomials m_1, \dots, m_q , we define $\mathbb{L}^2(I)$ to be a labeled induced subcomplex of \mathbb{L}_q^2 formed by the following rules:

- (1) Label each vertex of $\ell_{i,j}$ of \mathbb{L}_q^2 with the monomial $m_i m_j$.
- (2) If for any indices $i, j, u, v \in [q]$ where $[q] = \{1, \dots, q\}$ with $\{i, j\} \neq \{u, v\}$ we have $m_i m_j \mid m_u m_v$, then
 - If $m_i m_j = m_u m_v$ and $i = \min\{i, j, u, v\}$, then delete the vertex $\ell_{i,j}$.
 - If $m_i m_j \neq m_u m_v$, then delete the vertex $\ell_{u,v}$.
- (3) Label each of the remaining faces with the least common multiple of the labels of its vertices.

The remaining labeled subcomplex of \mathbb{L}_q^2 is called $\mathbb{L}^2(I)$, and is a subcomplex of $\text{Taylor}(I^2)$.


Remark 3.5.

It follows from Proposition 3.7 below that if m_i^2 divides $m_u m_v$ then $u = v = i$, hence the vertices $\ell_{i,i}$ are not deleted in the construction of $\mathbb{L}^2(I)$.

In Step 2 above, when there is equality, the choice was made to eliminate the vertex $\ell_{i,j}$ with minimum index i so that one has a well-defined definition for $\mathbb{L}^2(I)$; in fact, one could show that a different choice of elimination would also serve our purposes.

Example 3.6. Let $I = (abe, bc, cdf, ad)$. Setting $m_1 = abe$, $m_2 = bc$ and $m_3 = cdf$, $m_4 = ad$, we first label all vertices of \mathbb{L}_4^2 with the products $m_i m_j$, but then note that

$$m_2m_4 \mid m_1m_3.$$



So the (labeled) facets of $\mathbb{L}^2(I)$ are the following five:

Facet	Dimension
$\{m_1^2, m_1m_2, m_1m_4\}$	2
$\{m_2^2, m_1m_2, m_2m_3, m_2m_4\}$	3
$\{m_3^2, m_2m_3, m_3m_4\}$	2
$\{m_4^2, m_1m_4, m_2m_4, m_3m_4\}$	3
$\{m_1m_2, m_1m_4, m_2m_3, m_2m_4, m_3m_4\}$	4

In particular, $\mathbb{L}^2(I)$ is a 4-dimensional complex labeled with the generators of I^2 .

We now present two preliminary results needed for the proof that when the ideal I is square-free, $\mathbb{L}^2(I)$ supports a free resolution of I^2 .

Proposition 3.7. *Let m_1, \dots, m_q be a minimal square-free monomial generating set for an ideal I , let r be a positive integer, and suppose that for some $i \in [q]$ and $1 \leq u_1 \leq \dots \leq u_r \leq q$,*

$$m_i^r \mid m_{u_1} \cdots m_{u_r} \quad \text{or} \quad m_{u_1} \cdots m_{u_r} \mid m_i^r.$$

Then $u_1 = \dots = u_r = i$.

Proof. If for all, or some, of $j \in [r]$ we have $u_j = i$, then those copies of m_i can be deleted from each side of the division, so one can assume, without loss of generality that $i = 1 < u_1 \leq \dots \leq u_r \leq q$. Suppose that

$$m_1 = x_1^{a_1} \cdots x_n^{a_n} \quad \text{and} \quad m_{u_j} = x_1^{b_1^j} \cdots x_n^{b_n^j},$$

where $a_v, b_v^j \in \{0, 1\}$ for $j \in [r]$ and $v \in [n]$. It follows that:

- if $m_i^r \mid m_{u_1} \cdots m_{u_r}$, then for every index $v \in [n]$ where $a_v \neq 0$, we have $ra_v = r$ and so $b_v^1 = \dots = b_v^r = 1$. Therefore, we have $m_1 \mid m_{u_j}$ for $j \in [r]$. This is a contradiction since these monomials are minimal generators of I .
- if $m_{u_1} \cdots m_{u_r} \mid m_i^r$, then for each nonzero exponent b_v^1 of m_{u_1} we must have $a_v \neq 0$, and so $m_{u_1} \mid m_1$, again a contradiction.

□

Proposition 3.8. *Let I be an ideal minimally generated by square-free monomials m_1, \dots, m_q with $q \geq 2$. Then for every $i \in [q]$ there is a $j \in [q] \setminus \{i\}$ such that*

$$m_u m_v \nmid m_i m_j \text{ for any choice of } u, v \in [q] \setminus \{i, j\}.$$

In particular, $m_i m_j$ is a minimal generator of I^2 .

Proof. Suppose, by way of contradiction, that there exists $i \in [q]$ such that for every $j \in [q] \setminus \{i\}$ there exist $u, v \in [q] \setminus \{i, j\}$ such that $m_u m_v \mid m_i m_j$.

With i as above, there exist functions $\varphi, \psi: [q] \setminus \{i\} \rightarrow [q] \setminus \{i\}$ such that

$$m_{\varphi(j)} m_{\psi(j)} \mid m_i m_j \quad \text{for all } j \in [q] \setminus \{i\}. \quad (1)$$

For each $k \geq 0$, let φ^k denote the composition $\varphi \circ \varphi \circ \dots \circ \varphi$ (k times). (When $k = 0$, φ^0 is the identity function.) Let $a \in [q] \setminus \{i\}$. For each $w \geq 1$, set $b_w = \psi(\varphi^{w-1}(a))$. Apply (1) with $j = \varphi^{k-1}(a)$ to get:

$$m_{\varphi^k(a)} m_{b_k} \mid m_i m_{\varphi^{k-1}(a)} \quad \text{for all } k \geq 1.$$

From this, it is easy to see that

$$\left(m_{\varphi^k(a)} \cdot \prod_{w=1}^k m_{b_w} \right) \mid \left(m_i m_{\varphi^{k-1}(a)} \cdot \prod_{w=1}^{k-1} m_{b_w} \right) \quad \text{for all } k \geq 2.$$

Inductively, we thus obtain

$$\left(m_{\varphi^k(a)} \cdot \prod_{w=1}^k m_{b_w} \right) \mid \left(m_i^s m_{\varphi^{k-s}(a)} \cdot \prod_{w=1}^{k-s} m_{b_w} \right) \quad \text{for all } k \geq 2 \text{ and } k > s \geq 1. \quad (2)$$

Assume $\varphi^k(a) = \varphi^{k-s}(a)$ for some $k \geq 2$ and some s with $k > s \geq 1$. After simplifying in (2) we obtain

$$\left(\prod_{w=k-s+1}^k m_{b_w} \right) \mid m_i^s.$$

For $s = 1$, this implies $m_{b_k} \mid m_i$, but since $b_k \neq i$, this contradicts the minimality of the generating set. If $s > 1$ this is a contradiction according to Proposition 3.7. Therefore, we have shown that the integers $\varphi(a), \varphi^2(a), \dots$ are distinct. This is a contradiction, since $\varphi^k(a) \in [q] \setminus \{i\}$ for all k , and $[q] \setminus \{i\}$ is a finite set. \square

We are now ready to prove the main result of the paper.

Theorem 3.9 (Main Result). *Let I be a square-free monomial ideal. Then $\mathbb{L}^2(I)$ supports a free resolution of I^2 .*

Proof. Suppose I is minimally generated by the square-free monomials m_1, \dots, m_q .

The simplicial complex $\mathbb{L}^2(I)$ is an induced subcomplex of the quasi-tree \mathbb{L}_q^2 (Proposition 3.3), and is therefore a quasi-forest itself (see [3, 6]). Let V denote the set of vertices of $\mathbb{L}^2(I)$. In view of Theorem 2.3, to show that $\mathbb{L}^2(I)$ supports a resolution of I^2 , we need to show that, for every $\mathbf{m} \in \text{LCM}(I^2)$, $\mathbb{L}^2(I)_{\mathbf{m}}$ is connected, where $\mathbb{L}^2(I)_{\mathbf{m}}$ is the induced subcomplex of the complex $\mathbb{L}^2(I)$ on the set $V_{\mathbf{m}} = \{\ell_{i,j} \in V : m_i m_j \mid \mathbf{m}\}$.

Suppose $\mathbf{m} \in \text{LCM}(I^2)$. If $q = 1$, then $\mathbb{L}^2(I)_{\mathbf{m}}$ is either empty or a point. If $q = 2$, then $I^2 = (m_1^2, m_1 m_2, m_2^2)$ and $\mathbb{L}^2(I)$, as pictured in Example 3.2, has two facets connected by the vertex $\ell_{1,2}$. If $\mathbf{m} \in \{m_1^2, m_2^2\}$, then $\mathbb{L}^2(I)_{\mathbf{m}}$ is a point, and hence

connected. Otherwise, $m_1 m_2 \mid \mathbf{m}$, so the vertex $\ell_{1,2}$ will be in $\mathbb{L}^2(I)_{\mathbf{m}}$. If either $\ell_{1,1}$ or $\ell_{2,2}$ are in $\mathbb{L}^2(I)_{\mathbf{m}}$, they will be connected to $\ell_{1,2}$. Therefore $\mathbb{L}^2(I)_{\mathbf{m}}$ is connected.

Now assuming $q \geq 3$, we use the notation introduced in the proof of Proposition 3.3 for the facets of \mathbb{L}_q^2 , namely F_0, \dots, F_q . The facets of $\mathbb{L}^2(I)_{\mathbf{m}}$ are the maximal sets among the sets $F_0 \cap V_{\mathbf{m}}, \dots, F_q \cap V_{\mathbf{m}}$.

If $\mathbf{m} = m_i^2$ for some $i \in [q]$, then Proposition 3.7 shows that $\mathbb{L}^2(I)_{\mathbf{m}}$ is one point, and hence is connected. Assume now that $\mathbf{m} \neq m_i^2$ for all $i \in [q]$, and hence $F_0 \cap V_{\mathbf{m}} \neq \emptyset$. To show that $\mathbb{L}^2(I)_{\mathbf{m}}$ is connected, it suffices to show that, for each $i \in [q]$ such that $F_i \cap V_{\mathbf{m}} \neq \emptyset$, the intersection between $F_i \cap V_{\mathbf{m}}$ and $F_0 \cap V_{\mathbf{m}}$ is nonempty. Note that any vertex in $F_i \cap V_{\mathbf{m}}$ other than $\ell_{i,i}$ is also in $F_0 \cap V_{\mathbf{m}}$. We thus need to show that if $\ell_{i,i} \in V_{\mathbf{m}}$ for some $i \in [q]$, then there exists $b \in [q]$ with $b \neq i$ such that $\ell_{i,b} \in V_{\mathbf{m}}$.

Assume $\ell_{i,i} \in V_{\mathbf{m}}$, hence $m_i^2 \mid \mathbf{m}$. Set

$$A = \{j \in [q] : m_j \mid \mathbf{m}\}.$$

Note that $i \in A$. Since $\mathbf{m} \neq m_i^2$, we see that $|A| \geq 2$. By Proposition 3.8 applied to the ideal generated by the monomials m_j with $j \in A$, there exists $b \in A \setminus \{i\}$ such that

$$m_u m_v \text{ does not divide } m_i m_b \text{ for all } u, v \in A \setminus \{i, b\}. \quad (3)$$

Since $b \in A$, we have $m_b \mid \mathbf{m}$. We claim that $m_i m_b \mid \mathbf{m}$ as well. Indeed, since m_b is a square-free monomial, setting $\mathbf{m} = m_i^2 \mathbf{n}$, one has

$$m_b \mid \mathbf{m} \Rightarrow m_b \mid m_i^2 \mathbf{n} \Rightarrow m_b \mid m_i \mathbf{n} \Rightarrow m_i m_b \mid m_i^2 \mathbf{n} \Rightarrow m_i m_b \mid \mathbf{m}. \quad (4)$$

In order to conclude $\ell_{i,b} \in V_{\mathbf{m}}$, we need to show that $\ell_{i,b} \in V$, that is, $\ell_{i,b}$ is a vertex of $\mathbb{L}^2(I)$. If $\ell_{i,b} \notin V$, then we must have $m_u m_v \mid m_i m_b$ for some $u, v \in [q] \setminus \{i, b\}$. Since $m_i m_b \mid \mathbf{m}$, we further have $m_u \mid \mathbf{m}$ and $m_v \mid \mathbf{m}$, hence $u, v \in A$. This contradicts (3) above. \square

Remark 3.10. Given any $q \geq 2$, there are square free monomial ideals I with q generators such that $\mathbb{L}_q^2(I) = \mathbb{L}_q^2$ and the resolution supported on $\mathbb{L}^2(I)$ is minimal. The ideal $I = (xabc, yade, zbd, wcef)$ is such an example when $q = 4$, (see [3]).

4. A BOUND ON THE BETTI NUMBERS OF I^2

We now consider bounds on the Betti numbers of the second power of a square-free monomial ideal I , as provided by the simplicial complex $\mathbb{L}^2(I)$. Since I^2 has a free resolution supported on $\mathbb{L}^2(I)$, $\beta_d(I^2)$ is bounded above by the number of d -faces of $\mathbb{L}^2(I)$, which itself is bounded above by the number of d -faces of \mathbb{L}_q^2 .

It can be seen from the proof of Theorem 4.1 below that the right-hand term of the inequality (a) below is precisely the number of d -faces of \mathbb{L}_q^2 . Note that the bound in (a) depends only on the number of generators q , and not on I itself. The right-hand term of the inequality (b) below is equal to the number of d -dimensional faces of $\mathbb{L}^2(I)$, which provides a more precise bound that is dependent on the ideal I .

Theorem 4.1. *Let I be a square-free monomial ideal minimally generated by $q \geq 2$ monomials. Then for each $d \geq 0$ the d^{th} Betti number $\beta_d(I^2)$ satisfies*

$$(a) \quad \beta_d(I^2) \leq \binom{\frac{1}{2}(q^2 - q)}{d+1} + q \binom{q-1}{d}.$$

Furthermore, setting s to be the minimal number of generators of I^2 and t_i to be the number of vertices of the form $\ell_{i,j}$ that were deleted from \mathbb{L}_q^2 when forming $\mathbb{L}^2(I)$, then

$$(b) \quad \beta_d(I^2) \leq \binom{s-q}{d+1} + \sum_{i=1}^q \binom{q-1-t_i}{d}.$$

By Remark 3.10, the bound in (a) is sharp.

Proof. We begin by proving inequality (b). Theorem 3.9 gives that for each $d \geq 0$, $\beta_d(I^2)$ is bounded above by the number of d -dimensional faces of $\mathbb{L}^2(I)$. We compute this number next.

The faces of $\mathbb{L}^2(I)$ are of two types:

- (1) Faces that do not contain any vertex of the form $\ell_{i,i}$ for $i \in [q]$.
- (2) Faces that contain a vertex $\ell_{i,i}$ for some $i \in [q]$, and, as a consequence, all the other vertices have the form $\ell_{i,j}$ with $j \in [q] \setminus \{i\}$.

Let s denote the minimal number of generators of I^2 and set $t = \binom{q+1}{2} - s$. Since $\binom{q+1}{2}$ is the number of vertices of \mathbb{L}_q^2 , the integer t is precisely the number of vertices that are deleted in the construction of $\mathbb{L}^2(I)$, as described in Definition 3.4. As noted in Remark 3.5, all the deleted vertices $\ell_{i,j}$ must satisfy $i \neq j$, hence the number of vertices $\ell_{i,j}$ of $\mathbb{L}^2(I)$ with $i, j \in [q]$ and $i \neq j$ is $\binom{q}{2} - t$, which is equal to $s - q$.

To construct a d -dimensional face of type (1), we need to choose $d+1$ vertices among the vertices $\ell_{i,j}$ of $\mathbb{L}^2(I)$ with $i, j \in [q]$ and $i \neq j$. As noted above, there are $s - q$ such vertices. Thus, the number of d -dimensional faces of type (1) is $\binom{s-q}{d+1}$.

Fix $i \in [q]$. To construct a d -dimensional face of type (2) that contains $\ell_{i,i}$, we need to choose d vertices among the vertices $\ell_{i,j}$ of $\mathbb{L}^2(I)$ that satisfy $j \neq i$. There are $q - 1 - t_i$ such vertices, where t_i denotes the number of vertices $\ell_{i,j}$ of \mathbb{L}_q^2 that are deleted in $\mathbb{L}^2(I)$. Thus the number of d -dimensional faces of type (2) is $\sum_{i=1}^q \binom{q-1-t_i}{d}$.

Putting the two computations above together, we have that the number of d -dimensional faces of $\mathbb{L}^2(I)$ is equal to $\binom{s-q}{d+1} + \sum_{i=1}^q \binom{q-1-t_i}{d}$, yielding the inequality (b).

Note that inequality (a) follows from (b) by setting $t_i = 0$ for all i and $s = \binom{q+1}{2}$. In view of our computation above, the right-hand side of inequality (a) is precisely the number of d -dimensional faces of \mathbb{L}_q^2 . \square

For comparison, the fact that $\text{Taylor}(I^2)$ supports a free resolution of I^2 gives an inequality

$$\beta_d(I^2) \leq \binom{\frac{1}{2}(q^2 + q)}{d+1},$$

where the binomial on the right side denotes the number of d -faces of a $\frac{1}{2}(q^2 + q)$ -simplex, which is the largest possible size for $\text{Taylor}(I^2)$.

To get an idea how much Theorem 4.1 improves on this bound, we present the following table, for $q = 4$:

d	0	1	2	3	4	5	6
d -faces of largest possible $\text{Taylor}(I^2)$ $\binom{10}{d+1}$	10	45	120	210	252	210	120
d -faces of \mathbb{L}_q^2 $\binom{6}{d+1} + 4\binom{3}{d}$	10	27	32	19	6	1	0

To put this in context, we examine two specific ideals with 4 generators, and use Macaulay2 to find the Betti numbers of these ideals.

Example 4.2. For the ideal $J = (x, y, z, w)$ Macaulay2 gives the following Betti table for J^2 :

d	0	1	2	3
$\beta_d(J^2)$	10	20	15	4

These Betti numbers should be compared with the bounds in the table above.

Now let $I = (abe, bc, cdf, ad)$ be the ideal Example 3.6. The Betti numbers of I^2 as calculated by Macaulay2 are the following.

d	0	1	2	3
$\beta_d(I^2)$	9	14	6	0

In this case we should compare these Betti numbers with the bounds given by the Taylor complex with 9 vertices and the bounds given by the Theorem 4.1(b). For the given ideal, we saw that $\mathbb{L}^2(I)$ has 9 vertices, and m_2m_4 is an eliminated vertex, hence $s = 9$, $t_2 = t_4 = 1$ and $t_1 = t_3 = 0$ in Theorem 4.1(b). We have:

d	0	1	2	3	4	5	6
d -faces of $\text{Taylor}(I^2)$ $\binom{9}{d+1}$	9	36	84	126	126	84	36
d -faces of $\mathbb{L}^2(I)$ $\binom{5}{d+1} + 2\binom{3}{d} + 2\binom{2}{d}$	9	20	18	7	1	0	0

Acknowledgements. The bulk of this work was done during the 2019 Banff workshop “Women in Commutative Algebra”. We are grateful to the organizers, the funding agencies (NSF DMS-1934391), and to the Banff International Research Station for their hospitality.

Author Şega and Spiroff were partially supported by grants from the Simons Foundation (#354594, #584932, respectively), and authors Cooper and Faridi were supported by the Natural Sciences and Engineering Research Council of Canada (NSERC).

REFERENCES

1. D. Bayer, I. Peeva, B. Sturmfels, *Monomial resolutions*, Math. Res. Lett. 5, no. 1-2 (1998) 31–46.
2. D. Bayer, B. Sturmfels, *Cellular resolutions of monomial modules*, J. Reine Angew. Math. 503 (1998) 123–140.
3. S. M. Cooper, S. El Khoury, S. Faridi, S. Mayes-Tang, S. Morey, L. M. Şega and S. Spiroff, *Simplicial resolutions of powers of square free monomial ideals*, in preparation.
4. S. Faridi, *The facet ideal of a simplicial complex*, Manuscripta Math. 109, no. 2 (2014) 159-174.
5. S. Faridi, *Monomial resolutions supported by simplicial trees*, J. Commut. Algebra 6, no. 3 (2014).
6. S. Faridi, B. Hersey, *Resolutions of monomial ideals of projective dimension 1*, Comm. Algebra 45, no. 12 (2017) 5453–5464.
7. D.R. Grayson, M.E. Stillman, *Macaulay2, a software system for research in algebraic geometry*, Available at <http://www.math.uiuc.edu/Macaulay2/>.
8. I. Peeva, *Graded syzygies*, Algebra and Applications, 14. Springer-Verlag London, Ltd., London, 2011.
9. G. Lyubeznik, *A new explicit finite free resolution of ideals generate by monomials in an R-sequence*, J. Pure Appl. Algebra 51 (1998) 193-195.
10. D. Taylor, *Ideals generated by monomials in an R-sequence*, Thesis, University of Chicago (1966).
11. X. Zheng, *Resolutions of facet ideals*, Comm. Algebra 32 no. 6 (2004) 2301-2324.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MANITOBA, 520 MACHRAY HALL, 186 DYSART ROAD, WINNIPEG, MB, CANADA R3T 2N2

Email address: susan.cooper@umanitoba.ca

DEPARTMENT OF MATHEMATICS, AMERICAN UNIVERSITY OF BEIRUT, BLISS HALL 315, P.O. BOX 11-0236, BEIRUT 1107-2020, LEBANON

Email address: se24@aub.edu.lb

DEPARTMENT OF MATHEMATICS & STATISTICS, DALHOUSIE UNIVERSITY, 6316 COBURG RD., PO BOX 15000, HALIFAX, NS, CANADA B3H 4R2

Email address: faridi@dal.ca

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF TORONTO, 40 ST. GEORGE STREET, ROOM 6290, TORONTO, ON, CANADA M5S 2E4

Email address: smt@math.toronto.edu

DEPARTMENT OF MATHEMATICS, TEXAS STATE UNIVERSITY, 601 UNIVERSITY DR., SAN MARCOS, TX 78666, U.S.A.

Email address: morey@txstate.edu

LIANA M. ŞEGA, DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF MISSOURI, KANSAS CITY, MO 64110, U.S.A.

Email address: segal@umkc.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MISSISSIPPI, HUME HALL 335, P.O. BOX 1848, UNIVERSITY, MS 38677 USA

Email address: spiroff@olemiss.edu