

# On coloring digraphs with forbidden induced subgraphs

Raphael Steiner <sup>\*†</sup>

March 9, 2021

## Abstract

We prove a conjecture by Aboulker, Charbit and Naserasr [1] by showing that every oriented graph in which the out-neighborhood of every vertex induces a transitive tournament can be partitioned into two acyclic induced subdigraphs. We prove multiple extensions of this result to larger classes of digraphs defined by a finite list of forbidden induced subdigraphs. We thereby resolve several special cases of an extension of the famous Gyárfás-Sumner conjecture to directed graphs stated in [1].

## 1 Introduction

**Notation.** All graphs and digraphs considered in this paper are simple, that is, they are loopless, between two vertices in a graph there is at most one connecting edge, and between two vertices in a digraph there is at most one arc in each direction. We say that a digraph is an *oriented graph* if it does not contain directed cycles of length two (digons). Given a digraph  $D$ , we denote by  $V(D)$  its vertex-set and by  $A(D) \subseteq V(D) \times V(D)$  its set of arcs. We put  $v(D) := |V(D)|$ ,  $a(D) := |A(D)|$ . Arcs are denoted as  $(u, v)$ , where  $u$  is the tail of the arc and  $v$  is its head. For  $v \in V(D)$  we denote by  $N_D^+(v)$ ,  $N_D^-(v)$  the sets of out- and in-neighbors of  $v$  in  $D$ , respectively. We drop the subscript  $D$  if it is clear from context. We generalize this notation to vertex subsets by putting  $N_D^+(X) := \bigcup_{x \in X} N_D^+(x) \setminus X$  and  $N_D^-(X) := \bigcup_{x \in X} N_D^-(x) \setminus X$  for all  $X \subseteq V(D)$ . We further denote by  $D[X]$  the subdigraph with vertex set  $X$  and arc-set  $(X \times X) \cap A(D)$ . Any digraph of the form  $D[X]$  with  $\emptyset \neq X \subseteq V(D)$  is called an *induced subdigraph* of  $D$ . Given a set  $X$  of vertices or arcs, we denote by  $D - X$  the digraph obtained from  $D$  by deleting the elements in  $X$ .

For a graph  $G$ , we denote by  $\vec{G}$  the directed graph with  $V(\vec{G}) := V(G)$  and  $A(\vec{G}) := \{(x, y), (y, x) \mid xy \in E(G)\}$  and call it the *biorientation* of  $G$ . For an integer  $k \geq 1$  we denote by  $\vec{K}_k$  the biorientation of  $K_k$  and call it the *complete digraph* of order  $k$ , by  $\vec{K}_k$  we denote the *transitive tournament* of order  $k$ , and by  $\overline{K}_k$  we denote the digraph consisting of  $k$  isolated vertices. By  $S_k^+, S_k^-$  we denote the orientation of the star with  $k$  leaves in which all arcs are oriented outwards (inwards). By  $W_k^+$  and  $W_k^-$  we denote

<sup>\*</sup>Institute of Mathematics, Technische Universität Berlin, Germany

<sup>†</sup>Email: [steiner@math.tu-berlin.de](mailto:steiner@math.tu-berlin.de). Funded by DFG-GRK 2434 Facets of Complexity.

the oriented wheel graphs obtained by connecting the leaves of  $S_k^+$  and  $S_k^-$ , respectively, by a directed cycle.

A classical research topic in the theory of graph coloring is to study the chromatic number of graph classes defined by forbidden *induced* subgraphs. Maybe the most famous open problem in this area is the *Gyárfás-Sumner Conjecture*, which states the following.

**Conjecture 1.1** (Gyárfás [13] 1975, Sumner [24] 1981). *If  $F$  is a forest and  $k \in \mathbb{N}$ , then there exists an integer  $c(F, k) \geq 1$  such that every graph  $G$  excluding  $F$  and  $K_k$  as induced subgraphs satisfies  $\chi(G) \leq c(F, k)$ .*

Note that the result claimed by the Gyárfás-Sumner-conjecture would be best-possible in the following strong sense: Let  $\mathcal{F}$  be finite set of graphs, such that the set of graphs  $\text{Forb}_{\text{ind}}(F)$  excluding each member of  $\mathcal{F}$  as an induced subgraph has bounded chromatic number. Then at least one member of  $\mathcal{F}$  must be a complete graph (for otherwise all complete graphs would be contained in  $\text{Forb}_{\text{ind}}(F)$ , and hence the chromatic number of graphs in this class would be unbounded). Similarly, at least one member of  $\mathcal{F}$  must be a forest: If every graph in  $\mathcal{F}$  would include a cycle, then let  $g$  be the maximum length of a cycle appearing in a member of  $\mathcal{F}$ . It is now clear that every graph whose girth exceeds  $g$  is contained in  $\text{Forb}_{\text{ind}}(F)$ , but such graphs may have arbitrarily large chromatic number due to a classical result of Erdős [11]. Hence, the Gyárfás-Sumner conjecture may be restated as follows: If  $F$  is a finite list of graphs, then  $\text{Forb}_{\text{ind}}(F)$  has bounded chromatic number if and only if  $F$  contains a clique and a forest.

Despite being quite popular, the Gyárfás-Sumner conjecture has not yet been resolved in full generality. Some special cases for which the conjecture has been proved are when  $F$  is a subdivision of a star [22], a tree of radius at most two [18], a certain kind of caterpillar [10] or a certain type of a so-called (multi-)broom [23].

A widely known generalization of the chromatic number to directed graphs is the *dichromatic number*, which was introduced around 1980 by Erdős [12] and Neumann-Lara [21]. Given a digraph  $D$ , an *acyclic  $k$ -coloring* of  $D$  is an assignment  $c : V(D) \rightarrow S$  of colors from a finite color set  $S$  of size  $k$  to the vertices such that every *color class*  $c^{-1}(i), i \in S$  induces an *acyclic* subdigraph of  $D$ . In other words, an acyclic coloring is a vertex-coloring avoiding monochromatic directed cycles. The *dichromatic number*  $\vec{\chi}(D)$  is the smallest integer  $k$  for which an acyclic  $k$ -coloring of  $D$  exists. Since around 2000, the dichromatic number has been quite popular in graph theoretical research, and many results have established that it shares interesting structural properties with the chromatic number, see [2, 3, 4, 6, 7, 8, 14, 15, 16, 17, 19, 20] for only a small fraction of recent results on the topic.

In the same spirit, Aboulker, Charbit and Naserasr [1] recently initiated the systematic study of the relation between excluded induced subdigraphs and the dichromatic number and asked the following intriguing question.

**Problem 1.2.** *Let  $\mathcal{F}$  be a finite set of digraphs. Under which circumstances does there exist  $C \in \mathbb{N}$  such that every digraph  $D$  without an induced subdigraph isomorphic to a member of  $\mathcal{F}$  satisfies  $\vec{\chi}(D) \leq C$ ?*

Following the terminology introduced by Aboulker et al. [1], we denote by  $\text{Forb}_{\text{ind}}(\mathcal{F})$  the set of digraphs containing no induced subdigraph isomorphic to a member of  $\mathcal{F}$ . We say that  $\mathcal{F}$  is *heroic* if the digraphs in  $\text{Forb}_{\text{ind}}(\mathcal{F})$  have bounded dichromatic number and in this case we denote  $\bar{\chi}(\text{Forb}_{\text{ind}}(\mathcal{F})) := \max\{\bar{\chi}(D) \mid D \in \text{Forb}_{\text{ind}}(\mathcal{F})\}$ .

Just as in the undirected case, Aboulker et al. [1] observed several necessary conditions for a finite set  $\mathcal{F}$  of digraphs to be heroic, which we summarize in the following.

**Proposition 1.3** (cf. [1]). *Let  $\mathcal{F}$  be a finite heroic set of digraphs. Then  $\mathcal{F}$  must contain*

- a complete digraph  $\vec{K}_k$  for some  $k \in \mathbb{N}$ ,
- a biorientation of a forest,
- an orientation of a forest,
- a tournament, i.e., an orientation of a complete graph.

Inspired by yet another important conjecture in graph theory, the *Erdős-Hajnal Conjecture*, in [7] Berger, Choromanski, Chudnovsky, Fox, Loeb, Scott, Seymour and Thomassé studied the dichromatic number of tournaments which exclude a *single* fixed tournament  $H$  as a(n induced) subdigraph. In this paper, the authors defined a *hero* as a tournament  $H$  such that the tournaments excluding isomorphic copies of  $H$  have bounded dichromatic number. In other words, a digraph  $H$  is a hero if the set  $\{\vec{K}_2, \overleftarrow{K}_2, H\}$  is heroic. The main result of Berger et al. in [7] is a recursive characterization of heroes. It follows directly from this characterization that every transitive tournament and all tournaments on at most four vertices are heroes.

It is a natural aim to characterize the finite heroic sets  $\mathcal{F}$  of digraphs similar to what is claimed by the Gyárfás-Sumner-Conjecture for undirected graphs. In contrast to undirected graphs, only heroic sets of size at least 3 are interesting to consider, as the necessary conditions from Proposition 1.3 directly imply that  $\{\vec{K}_2, \overleftarrow{K}_2\}$  is the only heroic set of size two (and is so trivially). Aboulker et al. in [1] proved that every heroic set of size three must be of one of the following types:

- $\{\vec{K}_2, \overleftrightarrow{F}, \vec{K}_k\}$  for a forest  $F$  and a number  $k \in \mathbb{N}$ ,
- $\{\vec{K}_k, \overleftarrow{K}_\alpha, H\}$  for  $k, \alpha \in \mathbb{N}$  and a hero  $H$  such that  $k = 2$  or  $H$  is transitive, or
- $\{\vec{K}_2, F, H\}$  for some oriented star forest<sup>1</sup>  $F$  and a hero  $H$ , or
- $\{\vec{K}_2, F, \vec{K}_k\}$  for some oriented forest  $F$  and some  $k \in \mathbb{N}$ .

They then ventured to propose the conjecture that every one of the above triples is indeed heroic, thus claiming a complete description of the heroic sets of size 3.

Note that since  $\vec{K}_2$ -free digraphs amount exactly to the biorientations of undirected graphs, and since dichromatic number and chromatic number coincide on these, the

---

<sup>1</sup>An oriented star forest is a disjoint union of orientations of stars.

conjecture of Aboulker et al. corresponds exactly to the undirected Gyárfás-Sumner-Conjecture when restricting to the triples of the first type above. Triples of the second type as above were shown to be indeed heroic by Aboulker et al. (cf. [1], Theorem 4.1), their proof is based on the results from [15]. Finally, for the third and fourth types of triples we deal with oriented graphs. Let us restate the conjectures for these cases.

**Conjecture 1.4.** *For every orientation of a star forest  $F$  and every hero  $H$  the oriented graphs excluding  $F$  and  $H$  as induced subdigraphs have bounded dichromatic number.*

**Conjecture 1.5.** *For every orientation of a forest  $F$  and every  $k \in \mathbb{N}$  the oriented graphs excluding  $F$  and  $\vec{K}_k$  as induced subdigraphs have bounded dichromatic number.*

Conjecture 1.4 and Conjecture 1.5 can be regarded as oriented variants of the Gyárfás-Sumner conjecture, and in the main results of this paper we solve several special cases of these conjectures. Let us remark that Conjecture 1.5 can be reduced to the case in which  $F$  is an oriented tree via the following observation.

**Proposition 1.6.** *Let  $F_1$  and  $F_2$  be oriented forests and let  $F$  be their disjoint union. If Conjecture 1.5 holds for  $F_1$  and  $F_2$ , then it also holds for  $F$ .*

*Proof.* Suppose that  $\{\vec{K}_2, F_i, \vec{K}_k\}$  is heroic for all  $i \in \{1, 2\}$  and  $k \in \mathbb{N}$ .

We prove by induction on  $k \geq 2$  that digraphs in  $\text{Forb}_{\text{ind}}(\vec{K}_2, F, \vec{K}_k)$  have bounded dichromatic number. This is obvious for  $k = 2$ , so let  $k \geq 3$  and suppose there is  $C \in \mathbb{N}$  such that every digraph in  $\text{Forb}_{\text{ind}}(\vec{K}_2, F, \vec{K}_{k-1})$  admits an acyclic  $C$ -coloring. Let

$$C' := \max\{\bar{\chi}(\text{Forb}_{\text{ind}}(\vec{K}_2, F_1, \vec{K}_k)), v(F_1)(1 + 2C) + \bar{\chi}(\text{Forb}_{\text{ind}}(\vec{K}_2, F_2, \vec{K}_k))\}.$$

We claim that every digraph in  $\text{Forb}_{\text{ind}}(\vec{K}_2, F, \vec{K}_k)$  is acyclically  $C'$ -colorable. Suppose towards a contradiction that there exists  $D \in \text{Forb}_{\text{ind}}(\vec{K}_2, F, \vec{K}_k)$  such that  $\bar{\chi}(D) > C'$ . Since  $C' \geq \bar{\chi}(\text{Forb}_{\text{ind}}(\vec{K}_2, F_1, \vec{K}_k))$ , we find that  $D \notin \text{Forb}_{\text{ind}}(\vec{K}_2, F_1, \vec{K}_k)$ , and hence  $D$  contains an induced copy of  $F_1$ . Let  $X \subseteq V(D)$  be the vertex-set of this copy, let  $Y$  be the set of vertices outside  $X$  having at least one neighbor in  $X$ , and let us put  $Z := V(D) \setminus (X \cup Y)$ . Then we have  $\bar{\chi}(D[X]) \leq |X| = v(F_1)$ . By definition,  $Y$  is contained in the union of the  $2v(F_1)$  sets  $N^+(x), x \in X$  and  $N^-(x), x \in X$ . It is now clear that since  $D \in \text{Forb}_{\text{ind}}(\vec{K}_2, F, \vec{K}_k)$ , we must have  $D[N^+(x)], D[N^-(x)] \in \text{Forb}_{\text{ind}}(\vec{K}_2, F, \vec{K}_{k-1})$  for every  $x \in X$ , and it follows that

$$\bar{\chi}(D[Y]) \leq \sum_{x \in X} (\bar{\chi}(D[N^+(x)]) + \bar{\chi}(D[N^-(x)])) \leq 2v(F_1)C$$

by definition of  $C$ . Finally, since there are no arcs in  $D$  connecting vertices of  $X$  and  $Z$ , it follows that  $F$  cannot contain an induced copy of  $F_2$ , as the vertex-set of this copy joined with  $X$  would induce a copy of  $F$  in  $D$ . Hence, we have  $\bar{\chi}(D[Z]) \leq \bar{\chi}(\text{Forb}_{\text{ind}}(\vec{K}_2, F_2, \vec{K}_k))$ . We conclude

$$\bar{\chi}(D) \leq \bar{\chi}(D[X]) + \bar{\chi}(D[Y]) + \bar{\chi}(D[Z]) \leq v(F_1) + 2v(F_1)C + \bar{\chi}(\text{Forb}_{\text{ind}}(\vec{K}_2, F_2, \vec{K}_k)) \leq C',$$

contradicting our assumptions on  $D$ . This shows that these assumptions were wrong. It follows that  $\{\vec{K}_2, F, \vec{K}_k\}$  is heroic for all  $k \geq 1$ , as required.  $\blacksquare$

Aboulker et al. [1] noted that in case that  $H$  is a transitive tournament, Conjecture 1.4 is implied by a result of Chudnovsky, Scott and Seymour [9]. They further observed that Conjecture 1.4 is true in the case that  $F$  has at most two vertices. Finally they focused on the case when  $H = \vec{C}_3$  is the smallest non-trivial hero and  $F$  has 3 vertices. Then  $F$  must be one of the following:

- $\overline{K}_3$ , the forest consisting of three isolated vertices,
- $\vec{P}_3$ , the directed path on three vertices,
- $\vec{K}_2 + K_1$ , the oriented star forest consisting of an arc plus an isolated vertex,
- $S_2^+$ , the 2-out-star, or
- $S_2^-$ , the 2-in-star.

They proved that  $\{\vec{K}_2, F, \vec{C}_3\}$  is indeed heroic if  $F$  is one of the first three star forests. Already in the case  $F \in \{S_2^+, S_2^-\}$  however, they could not prove heroicity and made the following explicit conjecture.

**Conjecture 1.7** (cf. [1], Conjecture 6.2).

$$\vec{\chi}(\text{Forb}_{\text{ind}}(\vec{K}_2, S_2^+, \vec{C}_3)) = \vec{\chi}(\text{Forb}_{\text{ind}}(\vec{K}_2, S_2^-, \vec{C}_3)) = 2.$$

Note that by symmetry of reversing all arcs, it suffices to prove Conjecture 1.7 for the out-star  $S_2^+$ . The digraphs in  $\text{Forb}_{\text{ind}}(\vec{K}_2, S_2^+, \vec{C}_3)$  are exactly the directed triangle-free oriented graphs such that the out-neighborhood of every vertex induces a tournament. As the first main result of this paper, we prove Conjecture 1.7.

**Theorem 1.**

$$\vec{\chi}(\text{Forb}_{\text{ind}}(\vec{K}_2, S_2^+, \vec{C}_3)) = \vec{\chi}(\text{Forb}_{\text{ind}}(\vec{K}_2, S_2^-, \vec{C}_3)) = 2.$$

In fact, we deduce Theorem 1 as an immediate Corollary of the following stronger result involving the hero  $W_3^+$ .

**Theorem 2.**  $\vec{\chi}(\text{Forb}_{\text{ind}}(\vec{K}_2, S_2^+, W_3^+)) = 2.$

In order to prove Theorem 2, we need to establish several auxiliary results which deal with the structure of digraphs in the class  $\text{Forb}_{\text{ind}}(\vec{K}_2, S_2^+, W_3^+)$ , which is surprisingly complicated (these are exactly the oriented graphs in which the out-neighborhood of every vertex induces a transitive tournament).

As a next step we verify Conjecture 1.4 for more triples of the form  $\{\vec{K}_2, S_2^+, H\}$ , where  $H$  is some hero. We start with  $W_3^-$ .

**Theorem 3.**  $\vec{\chi}(\text{Forb}_{\text{ind}}(\vec{K}_2, S_2^+, W_3^-)) \leq 4.$

Our last result concerning Conjecture 1.4 generalizes Theorem 3 qualitatively and proves that for every  $k \in \mathbb{N}$ , the triple  $\{\vec{K}_2, S_2^+, H_k\}$  is heroic, where  $H_k$  is the hero on  $k$  vertices obtained from the disjoint union of  $W_3^+$  and  $\vec{K}_{k-4}$  by adding all possible arcs from  $W_3^+$  towards  $\vec{K}_{k-4}$ . More generally, we show the following.

**Theorem 4.** *Let  $H$  be a hero and let  $H^-$  be the hero obtained from  $H$  by adding a dominating sink. If  $\{\vec{K}_2, S_2^+, H\}$  is heroic, then so is  $\{\vec{K}_2, S_2^+, H^-\}$ .*

Our last new result in this paper concerns Conjecture 1.5. As mentioned above, Conjecture 1.5 holds true whenever  $F$  is an oriented star forest, and therefore particularly for forests on at most 3 vertices. The first open cases therefore appear when  $F$  is an orientation of the  $P_4$ . Aboulker et al. considered the directed path  $\vec{P}_4$  and showed in one of their main results that the set  $\{\vec{K}_2, \vec{P}_4, \vec{K}_3\}$  is heroic. There are three other oriented paths on four vertices. Two of them, which are called  $P^+(2, 1)$  and  $P^-(2, 1)$  in [1], consist of two oppositely oriented dipaths of length two and one, respectively. Chudnovsky, Scott and Seymour proved in [9] that for every  $k \in \mathbb{N}$ , digraphs in the set  $\text{Forb}_{\text{ind}}(\vec{K}_2, P, \vec{K}_k)$  have underlying graphs with bounded chromatic number (and thus bounded dichromatic number) for  $P \in \{P^+(2, 1), P^-(2, 1)\}$ . Hence, Conjecture 1.5 holds for these two orientations of  $P_4$ . The same result however is wrong for the remaining orientation of  $P_4$ , denoted  $P^+(1, 1, 1)$  in [1], as it consists of 3 alternately oriented arcs. Here we complement the result of Aboulker et al. [1] concerning the directed path  $\vec{P}_4$  and  $k = 3$  by showing that also the set  $\{\vec{K}_2, P^+(1, 1, 1), \vec{K}_3\}$  is heroic.

**Theorem 5.**  $\vec{\chi}(\text{Forb}_{\text{ind}}(\vec{K}_2, P^+(1, 1, 1), \vec{K}_3)) = 2$ .

We remark that the class  $\text{Forb}_{\text{ind}}(\vec{K}_2, P^+(1, 1, 1), \vec{K}_3)$  is quite rich, as it (among others) contains all oriented *line digraphs*.

**Structure of the paper.** In Section 2 we investigate the structure of digraphs in the class  $\text{Forb}_{\text{ind}}(\vec{K}_2, S_2^+, W_3^+)$  and use these insights to prove Theorem 2. In Section 3 we give the proof of Theorem 3. In Section 4 we prove Theorem 4. Finally, in Section 5 we prove Theorem 5 and we conclude with final comments in Section 6.

## 2 $\{S_2^+, W_3^+\}$ -Free Oriented Graphs

In this section, we will prove Theorem 2 and thereby show that  $\{\vec{K}_2, S_2^+, W_3^+\}$  is a heroic set. Note that  $\text{Forb}_{\text{ind}}(\vec{K}_2, S_2^+, W_3^+)$  is the class of oriented graphs  $D$  with the property that the out-neighbourhood of every vertex in  $D$  induces a transitive tournament. Given  $D \in \text{Forb}_{\text{ind}}(\vec{K}_2, S_2^+, W_3^+)$ , we define  $F = F(D)$  to be the spanning subdigraph of  $D$  consisting of the arcs  $(x, y) \in A(D)$  such that  $y$  is the source in the transitive tournament induced by the out-neighbourhood of  $x$  in  $D$ . Observe that for every  $x \in V(D)$ , if  $d_D^+(x) \geq 1$  then  $d_F^+(x) = 1$ , and otherwise  $d_F^+(x) = 0$ . From the definition of  $F(D)$  we immediately obtain the following property:

**Claim 2.1.** Let  $D \in \text{Forb}_{\text{ind}}(\overleftrightarrow{K}_2, S_2^+, W_3^+)$  and  $(x, y) \in A(F(D))$ . Then we have

$$N_D^+(x) \subseteq N_D^+(y) \cup \{y\}.$$

The next claim follows immediately from Claim 2.1 via induction.

**Claim 2.2.** Let  $D \in \text{Forb}_{\text{ind}}(\overleftrightarrow{K}_2, S_2^+, W_3^+)$  and let  $(x_1, \dots, x_k)$  be a dipath in  $F(D)$ . Then

$$N_D^+(x_1) \setminus \{x_2, \dots, x_k\} \subseteq N_D^+(x_k).$$

From Claim 2.2 we can derive that the vertex-sets of directed cycles in  $F(D)$  form so-called *out-modules* in  $D$ .

**Definition 2.3.** Let  $D$  be a digraph, and  $\emptyset \neq M \subseteq V(D)$ . We say that  $M$  is an out-module in  $D$  if it holds that  $(x, z) \in A(D) \Rightarrow (y, z) \in V(D) \setminus M$  for every  $x, y \in M$  and  $z \in V(D) \setminus M$ . Equivalently,  $N_D^+(x) \setminus M = N_D^+(y) \setminus M$  for all  $x, y \in M$ .

**Claim 2.4.** Let  $D \in \text{Forb}_{\text{ind}}(\overleftrightarrow{K}_2, S_2^+, W_3^+)$ , and let  $C$  be a directed cycle in  $F(D)$ . Then  $V(C)$  is an out-module in  $D$ .

*Proof.* Let  $x_1, x_2, \dots, x_k, x_1$  be the vertex-trace of  $C$ . Let  $y \in V(D) \setminus \{x_1, \dots, x_k\}$  and  $1 \leq i \leq k$  be arbitrary such that  $(x_i, y) \in A(D)$ . Let  $j \in [k] \setminus \{i\}$ . By Claim 2.2, applied to the directed subpath of  $C$  starting in  $x_i$  and ending in  $x_j$ , we know that  $N_D^+(x_i) \setminus \{x_1, \dots, x_k\} \subseteq N_D^+(x_j) \setminus \{x_1, \dots, x_k\}$ . Hence  $(x_j, y) \in A(D)$ . This shows that  $\{x_1, \dots, x_k\}$  is indeed an out-module.  $\blacksquare$

For a non-empty vertex-set  $U$  in a digraph  $D$ , we denote by  $D/U$  the digraph obtained by *identifying*  $U$ , that is, the digraph with vertex set  $(V(D) \setminus U) \cup \{x_U\}$  where  $x_U \notin V(D)$  is some newly added vertex representing  $U$ , and the following arcs: the arcs of  $D$  inside  $V(D) \setminus U$ , the arc  $(x_U, v)$  for every  $v \in N_D^+(U)$ , and the arc  $(v, x_U)$  for every  $v \in N_D^-(U)$ .

In the following we prepare the proof of Theorem 2 with a set of useful Lemmas. We start with two lemmas yielding modifications of digraphs which preserve the containment in the class  $\text{Forb}_{\text{ind}}(\overleftrightarrow{K}_2, S_2^+, W_3^+)$ .

**Lemma 2.5.** For every  $D \in \text{Forb}_{\text{ind}}(\overleftrightarrow{K}_2, S_2^+, W_3^+)$  and for every out-module  $U \subseteq V(D)$  it holds that  $D/U \in \text{Forb}_{\text{ind}}(\overleftrightarrow{K}_2, S_2^+, W_3^+)$ .

*Proof.* We need to show that  $D/U$  is induced  $\{\overleftrightarrow{K}_2, S_2^+, W_3^+\}$ -free. We argue by contradiction. Suppose first that  $D/U$  contains a  $\overleftrightarrow{K}_2$ , namely a pair of vertices  $x, y$  with  $(x, y), (y, x) \in A(D/U)$ . If  $x, y \neq x_U$  then  $x, y$  also span a copy of  $\overleftrightarrow{K}_2$  in  $D$ , a contradiction. Suppose then that  $x = x_U$  or  $y = x_U$ ; say  $x = x_U$ . By the definition of  $D/U$ , there are (not necessarily distinct)  $u_1, u_2 \in U$  such that  $(u_1, y), (y, u_2) \in A(D)$ . Since  $U$  is an out-module,  $(u_2, y) \in A(D)$ . Hence,  $u_2, y$  span a copy of  $\overleftrightarrow{K}_2$  in  $D$ , a contradiction.

Suppose next that  $D/U$  contains an induced copy of  $S_2^+$ , namely, distinct vertices  $x, y, z \in V(D/U)$  with  $(x, y), (x, z) \in A(D/U)$  and with no arc in  $D/U$  between  $y$  and  $z$ . If  $x, y, z \neq x_U$  then  $x, y, z$  also span an induced  $S_2^+$  in  $D$ , a contradiction.

Suppose now that  $y = x_U$ , and let  $u \in U$  be such that  $(x, u) \in A(D)$ . We have  $(u, z), (z, u) \notin A(D)$  because  $(x_U, z), (z, x_U) \notin A(D/U)$ . Hence,  $x, u, z$  span an induced  $S_2^+$  in  $D$ , a contradiction. The case  $z = x_U$  is analogous. Suppose now that  $x = x_U$ . Since  $(x_U, y), (x_U, z) \in A(D/U)$  and  $U$  is an out-module, we must have  $(u, y), (u, z) \in A(D)$  for every  $u \in U$ , implying that  $u, y, z$  span an induced  $S_2^+$  in  $D$  for every such  $u$ , again yielding the desired contradiction.

Suppose now that  $D/U$  contains a copy of  $W_3^+$  with vertices  $x, y, z, w$  and arcs  $(x, y), (x, z), (x, w), (y, z), (z, w), (w, y)$ . Again, if  $x, y, z, w \neq x_U$  then  $D$  also has a copy of  $W_3^+$ , a contradiction. Suppose now that  $x = x_U$ , and fix any  $u \in U$ . Since  $U$  is an out-module, we have  $(u, y), (u, z), (u, w) \in A(D)$ , implying that  $u, y, z, w$  span a copy of  $W_3^+$  in  $D$ , a contradiction. Suppose finally that one of  $y, z, w$  equals  $x_U$ , say  $y = x_U$  (without loss of generality). Since  $(x, x_U), (x_U, z), (w, x_U) \in A(D/U)$ , there are  $u_1, u_2, u_3 \in U$  (not necessarily distinct) such that  $(x, u_1), (u_2, z), (w, u_3) \in A(D)$ . Since  $U$  is an out-module, we have  $(u_1, z), (u_3, z) \in A(D)$ . Since  $(x, w), (x, u_1) \in A(D)$  and  $D$  is induced  $S_2^+$ -free, we must have either  $(w, u_1) \in A(D)$  or  $(u_1, w) \in A(D)$ . If  $(u_1, w) \in A(D)$  then also  $(u_3, w) \in A(D)$  because  $U$  is an out-module, but this is impossible as then  $u_3, w$  would induce a digon in  $D$ . Finally, if  $(w, u_1) \in A(D)$  then  $x, u_1, z, w$  span a copy of  $W_3$  in  $D$ , again yielding a contradiction. This concludes the proof of the lemma.  $\blacksquare$

**Lemma 2.6.** *Let  $D \in \text{Forb}_{\text{ind}}(\vec{K}_2, S_2^+, W_3^+)$ , and let  $(x, y) \in A(F(D))$ . Let  $z \in N_D^+(y)$  such that  $(x, z), (z, x) \notin A(D)$ . Then the digraph  $D + (x, z)$  obtained from  $D$  by adding the arc  $(x, z)$  is contained in  $\text{Forb}_{\text{ind}}(\vec{K}_2, S_2^+, W_3^+)$ .*

*Proof.* We need to show that  $D + (x, z)$  is induced  $\{\vec{K}_2, S_2^+, W_3^+\}$ -free. Again, we argue by contradiction. Clearly  $D + (x, z)$  does not contain a  $\vec{K}_2$ , since  $(z, x) \notin A(D)$  by assumption. Suppose next that  $D + (x, z)$  contains an induced copy of  $S_2^+$ , i.e. distinct vertices  $a, b, c$  such that  $(a, b), (a, c) \in A(D + (x, z))$ , and  $(b, c), (c, b) \notin A(D + (x, z))$ . If  $(x, z) \notin \{(a, b), (a, c)\}$ , then  $a, b, c$  induce a copy of  $S_2^+$  also in  $D$ , a contradiction. We may therefore assume w.l.o.g. that  $(x, z) = (a, b)$ . Then we have  $c \neq y$ , since  $(c, b) \notin A(D)$ , but  $(y, b) = (y, z) \in A(D)$  by assumption. Since  $(x, c) = (a, c) \in A(D)$ , we have  $c \in N_D^+(x)$ . But  $(x, y) \in A(F(D))$ , and hence Claim 2.1 implies that  $c \in N_D^+(x) \subseteq \{y\} \cup N_D^+(y)$ . It follows that  $(y, c) \in A(D)$ . We further have  $(y, b) = (y, z) \in A(D)$  and  $(b, c), (c, b) \notin A(D)$ . Hence,  $y, b, c$  induce an  $S_2^+$  in  $D$ , a contradiction.

Moving on, suppose that  $D + (x, z)$  contains an induced copy of  $W_3^+$ , i.e., distinct vertices  $a, b, c, d$  such that  $(a, b), (a, c), (a, d), (b, c), (c, d), (d, b) \in A(D) \cup \{(x, z)\}$ .

Suppose first that  $(x, z) \notin \{(a, b), (a, c), (a, d)\}$ . Then  $(a, b), (a, c), (a, d) \in A(D)$  and since  $D$  does not contain an induced copy of  $S_2^+$ , the vertices  $b, c, d$  are pairwise adjacent. Since  $x$  and  $z$  are non-adjacent in  $D$ , it follows that  $\{x, z\} \not\subseteq \{b, c, d\}$ . Hence we have  $(b, c), (c, d), (d, b) \in A(D)$ , yielding that  $a, b, c, d$  induce a copy of  $W_3^+$  in  $D$ , a contradiction. Hence we may suppose that  $(x, z) \in \{(a, b), (a, c), (a, d)\}$ . By symmetry we may assume that  $(x, z) = (a, b)$  w.l.o.g. Again using Claim 2.1 we then have  $c, d \in N_D^+(x) \subseteq N_D^+(y) \cup \{y\}$ .

Let us first consider the case that  $c, d \neq y$ . Then  $(y, c), (y, d) \in A(D)$ , and since  $(y, b) = (y, z) \in A(D)$  by assumption, it follows that the vertices  $y, b, c, d$  induce a copy



of  $W_3^+$  in  $D$ , a contradiction. For the next case suppose that  $y \in \{c, d\}$ . The first option, namely that  $y = c$ , is impossible, since then we would have  $(y, z) \in A(D)$  (by assumption) and  $(z, y) = (b, c) \in A(D)$ , a contradiction since  $D$  is  $\overleftrightarrow{K}_2$ -free. Therefore, we must have  $y = d$ . Then  $c \in N_D^+(y)$  as well as  $(c, y) = (c, d) \in A(D)$ . It follows that  $y, c$  induce a  $\overleftrightarrow{K}_2$  in  $D$ , so again, we conclude with a contradiction.

Having reached a contradiction in all cases, it follows that our initial assumption was wrong, indeed,  $D + (x, z) \in \text{Forb}_{\text{ind}}(\overleftrightarrow{K}_2, S_2^+, W_3^+)$ . This concludes the proof.  $\blacksquare$

The next lemma shows the existence of out-modules with special properties.

**Lemma 2.7.** *Let  $D \in \text{Forb}_{\text{ind}}(\overleftrightarrow{K}_2, S_2^+, W_3^+)$ , and let  $v \in V(D)$ . If  $N_D^-(v) \neq \emptyset$ , then there exists an out-module  $M$  in  $D$  such that  $M \subseteq N_D^-(v)$  and  $N_D^+(M) \subseteq N_D^+(v) \cup \{v\}$ .*

*Proof.* We prove by induction on  $n \geq 1$  the statement of the lemma for all digraphs  $D \in \text{Forb}_{\text{ind}}(\overleftrightarrow{K}_2, S_2^+, W_3^+)$  and vertices  $v \in V(D)$  such that  $d_D^-(v) \leq n$ .

If  $n = 1$ , then  $N_D^-(v) = \{w\}$  for a vertex  $w \in V(D)$ . Then  $M := \{w\}$  is an out-module of  $D$ . Hence, it suffices to verify that  $N_D^+(w) \subseteq N_D^+(v) \cup \{v\}$ . Suppose towards a contradiction that  $(w, v') \in A(D)$  for  $v' \in N_D^+(w) \setminus (N_D^+(v) \cup \{v\})$ . Since  $N_D^-(v) = \{w\}$ , we have  $v' \notin N_D^-(v)$ , and hence  $v, v'$  are non-adjacent in  $D$ , while  $(w, v), (w, v') \in A(D)$ . Hence,  $w, v, v'$  induce an  $S_2^+$  in  $D$ , a contradiction.

Now let  $D \in \text{Forb}_{\text{ind}}(\overleftrightarrow{K}_2, S_2^+, W_3^+)$  and  $v \in V(D)$  such that  $d_D^-(v) = n \geq 2$ , and assume that the claim holds for all pairs of digraphs in  $\text{Forb}_{\text{ind}}(\overleftrightarrow{K}_2, S_2^+, W_3^+)$  and vertices whose in-degree is less than  $n$ .

First let us assume that there exists a vertex  $w \in N_D^-(v)$  such that  $(w, v) \in A(F(D))$ . Then  $M := \{w\}$  is an out-module of  $D$ , and by Claim 2.1 we have  $N_D^+(w) \subseteq N_D^+(v) \cup \{v\}$ . This proves the assertion in this case.

Hence, for the rest of this proof we may suppose that  $(w, v) \notin A(F(D))$  for every  $w \in N_D^-(v)$ . For any  $w \in N_D^-(v)$  we clearly have  $d_D^+(w) \geq 1$  and hence it follows that  $d_{F(D)}^+(w) = 1$ . Furthermore, for every arc  $(w, w') \in A(F(D))$  such that  $w \in N_D^-(v)$  by Claim 2.1 we must have  $v \in N_D^+(w) \subseteq N_D^+(w') \cup \{w'\}$ . Since  $(w, v) \notin A(F(D))$ , we have  $v \neq w'$  and hence  $(w', v) \in A(D)$ . This shows that the out-neighbor in  $F(D)$  of any vertex in  $N_D^-(v)$  is again contained in  $N_D^-(v)$ . It follows that  $F(D)$  restricted to  $N_D^-(v)$  has minimum out-degree 1 and therefore contains a directed cycle  $C$  such that  $V(C) \subseteq N_D^-(v)$ .

By Claim 2.4,  $N := V(C)$  is an out-module in  $D$ . Consider the digraph  $D' := D/N$ , which by Lemma 2.5 is a member of  $\text{Forb}_{\text{ind}}(\overleftrightarrow{K}_2, S_2^+, W_3^+)$ . Then by definition of  $D/N$  and since  $N \subseteq N_D^-(v)$ , we have  $v \in V(D')$  and  $N_{D'}^-(v) = (N_D^-(v) \setminus N) \cup \{x_N\} \neq \emptyset$ . Since  $|N| = |V(C)| \geq 3$ , this implies that  $d_{D'}^-(v) = 1 + d_D^-(v) - |N| \leq d_D^-(v) - 2 < n$ . We may therefore apply the induction hypothesis to the digraph  $D' \in \text{Forb}_{\text{ind}}(\overleftrightarrow{K}_2, S_2^+, W_3^+)$  and the vertex  $v \in V(D')$ . We thus find an out-module  $M'$  in  $D'$  with the properties  $M' \subseteq N_{D'}^-(v) = (N_D^-(v) \setminus N) \cup \{x_N\}$  and  $N_{D'}^+(M') \subseteq N_{D'}^+(v) \cup \{v\} = N_D^+(v) \cup \{v\}$ . Let us define the set  $M \subseteq N_D^-(v)$  as  $M := M'$ , if  $x_N \notin M'$ , and  $M := (M' \setminus \{x_N\}) \cup N$  if  $x_N \in M'$ . We claim that  $M$  satisfies the assertions of the Lemma with respect to  $D$  and  $v$ . In the following, we verify both parts of the inductive claim separately.

**Claim 1.**  $N_D^+(M) \subseteq N_D^+(v) \cup \{v\}$ .

*Proof.* In the proof we will use the fact that

$$N_{D'}^+(M') \subseteq N_{D'}^+(v) \cup \{v\} = N_D^+(v) \cup \{v\},$$

which holds by the induction hypothesis.

Let  $x \in N_D^+(M)$  be given arbitrarily. Let  $m \in M$  such that  $(m, x) \in A(D)$ . Our goal is to show that  $x \in N_D^+(v) \cup \{v\}$ .

Let us first consider the case that  $x_N \notin M'$  and hence  $M = M'$ . By definition of  $D' = D/N$  we either have  $x \notin N$  and  $(m, x) \in A(D')$ , or  $x \in N$  and  $(m, x_N) \in A(D')$ . Then since  $m \in M = M'$  and  $x, x_N \notin M = M'$ , we obtain that either  $x \notin N$  and  $x \in N_{D'}^+(M')$ , or  $x \in N$  and  $x_N \in N_{D'}^+(M')$ . As  $N_{D'}^+(M') \subseteq N_D^+(v) \cup \{v\}$ , in the first case we have  $x \in N_D^+(v) \cup \{v\}$ , as desired. The second case does not occur, since it yields  $x_N \in N_D^+(v) \cup \{v\}$ , which is impossible as  $x_N$  is not a vertex of  $D$ . Hence, we have shown the claim that  $x \in N_D^+(v) \cup \{v\}$ .

For the second case, suppose that  $x_N \in M'$  and hence  $M = (M' \setminus \{x_N\}) \cup N$ . Note that  $x \notin N$ , since  $x \notin M \supseteq N$ . Hence, the existence of the arc  $(m, x) \in A(D)$  yields that either  $m \in N$  and  $(x_N, x) \in A(D')$ , or  $m \notin N$  and  $(m, x) \in A(D')$ . In both cases, this implies that  $x \in N_{D'}^+(M') \subseteq N_D^+(v) \cup \{v\}$ , proving the assertion. ■

**Claim 2.**  $M$  is an out-module in  $D$ .

*Proof.* Let  $x \neq y \in M$  and  $z \in V(D) \setminus M$  arbitrary, and assume that  $(x, z) \in A(D)$ . We need to show that also  $(y, z) \in A(D)$ . Note that by Claim 1 we have  $z \notin N_D^-(v)$ , as otherwise  $z \in N_D^+(M) \cap N_D^-(v) = \emptyset$ . In particular,  $z \notin N$ .

Observe that  $z \in N_{D'}^+(M')$ . Indeed, if  $x \notin N$  then  $x \in M'$  and  $(x, z) \in A(D')$ , and if  $x \in N$  then  $x_N \in M'$  and  $(x_N, z) \in A(D')$ ; in any case,  $z \in N_{D'}^+(M')$ .

Since  $y \in M$ , we have either  $y \in M'$  or  $y \in N$  and  $x_N \in M'$ . Suppose first that  $y \in M'$ . Then  $(y, z) \in A(D')$  because  $z \in N_{D'}^+(M')$  and  $M'$  is an out-module. Hence, in this case  $(y, z) \in A(D)$ , as required. Now suppose that  $y \in N$  and  $x_N \in M'$ . Since  $z \in N_{D'}^+(M')$  and  $M'$  is an out-module, we have  $(x_N, z) \in A(D')$ . This means that there is  $w \in N$  such that  $(w, z) \in A(D)$ . Now, as  $N$  is itself an out-module in  $D$  and  $y \in N, z \notin N$ , we have  $(y, z) \in A(D)$ , as required. ■

By Claim 1 and 2 the out-module  $M$  certifies that the pair  $(D, v)$  satisfies the inductive claim. This concludes the proof of the Lemma by induction. ■

**Lemma 2.8.** Let  $D \in \text{Forb}_{\text{ind}}(\overleftrightarrow{K}_2, S_2^+, W_3^+)$ , let  $M \subseteq V(D)$  be an out-module in  $D$  and let  $v \in V(D) \setminus M$ . Let  $T$  be the set of vertices defined by

$$T := \{t \in M \mid \exists u \in V(D) \setminus M : (v, u), (u, t) \in A(D)\}.$$

Then  $D[T]$  is a (possibly empty) transitive tournament.

*Proof.* The assertion will follow directly from the following two claims.

**Claim 1.** If  $t_1 \neq t_2 \in T$ , then  $t_1$  and  $t_2$  are adjacent in  $D$ .

*Proof.* By definition of  $T$  there exist vertices  $u_1, u_2 \in V(D) \setminus M$  (not necessarily distinct) such that  $(v, u_i), (u_i, t_i) \in A(T)$  for  $i = 1, 2$ . If  $u_1 = u_2$ , then  $t_1, t_2$  must be adjacent, for otherwise the vertices  $u_1, t_1, t_2$  would induce an  $S_2^+$  in  $D$ , a contradiction. Suppose now that  $u_1 \neq u_2$ . Since  $u_1, u_2 \in N^+(v)$ , they must be adjacent, w.l.o.g. let  $(u_1, u_2) \in A(D)$ . Then  $t_1 \in M$  and  $u_2 \in V(D) \setminus M$  are distinct out-neighbors of  $u_1$ , and hence they must be adjacent in  $D$ . If  $(t_1, u_2) \in A(D)$ , then  $M$  being an out-module implies that also  $(t_2, u_2) \in A(D)$ , yielding a  $\overleftrightarrow{K}_2$  in  $D$  induced by  $t_2$  and  $u_2$ , a contradiction. Therefore we have  $(u_2, t_1) \in A(D)$ . Then  $t_1$  and  $t_2$  are distinct out-neighbors of  $u_2$  in  $D$ , which implies that they must be adjacent. This concludes the proof.  $\blacksquare$

**Claim 2.**  $D[T]$  contains no directed triangle.

*Proof.* Suppose towards a contradiction that there are three distinct  $t_1, t_2, t_3 \in T$  inducing a directed triangle in  $D$ . Let  $u_1, u_2, u_3 \in V(D) \setminus M$  be (not necessarily distinct) such that  $(v, u_i), (u_i, t_i) \in A(D), i = 1, 2, 3$ . We distinguish three different cases depending on the size of the set  $\{u_1, u_2, u_3\}$ .

For the first case, suppose that  $u_1 = u_2 = u_3$ . Then  $t_1, t_2, t_3$  are three distinct out-neighbors of  $u_1$  spanning a directed triangle. Hence,  $u_1, t_1, t_2, t_3$  induce a  $W_3^+$  in  $D$ , a contradiction to our assumption that  $D \in \text{Forb}_{\text{ind}}(\overleftrightarrow{K}_2, S_2^+, W_3^+)$ .

For the second case, suppose that  $\{u_1, u_2, u_3\}$  contains exactly two distinct vertices, w.l.o.g.  $u_1 \neq u_2 = u_3$ . Since  $u_1$  and  $u_2$  are two distinct out-neighbors of  $v$  in  $D$ , they must be adjacent. Suppose first that  $(u_1, u_2) \in A(D)$ . Then  $t_1$  and  $u_2$  are two distinct out-neighbors of  $u_1$  in  $D$ , and hence they must be adjacent. If  $(t_1, u_2) \in A(D)$ , then by the module-property of  $M$ , also  $(t_2, u_2) \in A(D)$ , and hence  $t_2, u_2$  induce a  $\overleftrightarrow{K}_2$  in  $D$ , a contradiction. If  $(u_2, t_1) \in A(D)$ , then  $t_1, t_2, t_3 \in N_D^+(u_2)$  and hence  $u_2, t_1, t_2, t_3$  induce a  $W_3^+$  in  $D$ , a contradiction. Next suppose that  $(u_2, u_1) \in A(D)$ . Then  $u_1, t_2, t_3$  are three distinct out-neighbors of  $u_2$  in  $D$ , and hence  $u_1$  must be adjacent to both  $t_2$  and  $t_3$ . If  $(t_2, u_1) \in A(D)$  or  $(t_3, u_1) \in A(D)$ , then  $(t_1, u_1) \in A(D)$  since  $M$  is an out-module, and hence  $u_1, t_1$  induce a  $\overleftrightarrow{K}_2$  in  $D$ , a contradiction. Otherwise, we have  $(u_1, t_2), (u_1, t_3) \in A(D)$  and hence  $u_1, t_1, t_2, t_3$  induce a  $W_3^+$  in  $D$ , again yielding the desired contradiction.

For the third case, suppose that  $u_1, u_2, u_3$  are pairwise distinct. Since  $u_1, u_2, u_3$  are three distinct vertices in the transitive tournament  $D[N^+(v)]$ , they form a transitive triangle, and we may assume w.l.o.g. that  $(u_1, u_2), (u_1, u_3), (u_2, u_3) \in A(D)$ . Then  $t_1$  and  $u_3$  are distinct out-neighbors of  $u_1$  in  $D$ , while  $t_2$  and  $u_3$  are distinct out-neighbors of  $u_2$  in  $D$ . Hence,  $u_3$  must be adjacent to both  $t_1$  and  $t_2$ . If  $(t_i, u_3) \in A(D)$  for some  $i = 1, 2$ , then we also have  $(t_3, u_3) \in A(D)$  since  $M$  is an out-module, and hence  $u_3, t_3$  induce a  $\overleftrightarrow{K}_2$  in  $D$ , contradiction. Finally, if  $(u_3, t_1), (u_3, t_2) \in A(D)$ , then  $u_3, t_1, t_2, t_3$  induce a  $W_3^+$  in  $D$ , yielding again a contradiction to the containment of  $D$  in  $\text{Forb}_{\text{ind}}(\overleftrightarrow{K}_2, S_2^+, W_3^+)$ .

Since we arrived at a contradiction in each case, we conclude that the initial assumption concerning the existence of  $t_1, t_2, t_3$  was wrong. This concludes the proof.  $\blacksquare$

We are now sufficiently prepared to give the proof of Theorem 2. In fact, we will prove the following slightly stronger version of the result, which allows to enforce a monochromatic coloring on the closed out-neighborhood of an arbitrarily chosen vertex.

**Theorem 6.** *Let  $D \in \text{Forb}_{\text{ind}}(\overleftrightarrow{K}_2, S_2^+, W_3^+)$ , and  $v \in V(D)$ . Then there exists an acyclic coloring  $c : V(D) \rightarrow \{1, 2\}$  of  $D$  such that  $c(u) = c(v)$  for every  $u \in N_D^+(v)$ .*

*Proof.* Suppose towards a contradiction that the claim is wrong, and let  $D$  be a counterexample to the claim minimizing  $v(D)$ . Let  $v \in V(D)$  be a vertex such that  $D$  does not admit an acyclic 2-coloring  $c$  with the property that  $c(u) = c(v)$  for every  $u \in N_D^+(v)$ .

**Claim 1.**  $N_D^-(v) \neq \emptyset$ .

*Proof.* Suppose towards a contradiction that  $N_D^-(v) = \emptyset$ . If also  $N_D^+(v) = \emptyset$ , then  $v$  is an isolated vertex of  $D$ . Then any acyclic 2-coloring of  $D - v$  could be extended to an acyclic 2-coloring of  $D$  by coloring  $v$  with color 1, and the statement that  $v$  has the same color as its out-neighbors would hold vacuously. Since this is impossible, we must have  $\bar{\chi}(D - v) \geq 3$ , which however contradicts the minimality of  $D$  as a counterexample. This shows that  $N_D^+(v) \neq \emptyset$ . Let  $u \in N_D^+(v)$  be the unique out-neighbor of  $v$  in  $F(D)$ . Then  $N_D^+(v) \subseteq N_D^+(u) \cup \{u\}$  by Claim 2.1. The minimality of  $D$  as a counterexample now implies that the digraph  $D - v \in \text{Forb}_{\text{ind}}(\overleftrightarrow{K}_2, S_2^+, W_3^+)$  admits an acyclic 2-coloring  $c^- : V(D) \rightarrow \{1, 2\}$  satisfying  $c^-(x) = c^-(u)$  for every  $x \in N_{D-v}^+(u) = N_D^+(u)$ . Let  $c : V(D) \rightarrow \{1, 2\}$  be defined as  $c(x) := c^-(x)$  for every  $x \in V(D) \setminus \{v\}$  and  $c(v) := c^-(u)$ . Since  $c$  restricted to  $V(D) \setminus \{v\}$  is an acyclic coloring, and no directed cycle in  $D$  contains  $v$  (recall  $N_D^-(v) = \emptyset$ ), it follows that  $c$  is an acyclic coloring of  $D$ . Moreover, for every  $x \in N_D^+(v) \subseteq N_D^+(u) \cup \{u\}$  we have  $c(x) = c^-(x) = c^-(u) = c(v)$ . This is a contradiction to our initial assumption that  $D$  does not admit an acyclic 2-coloring with this property. This shows that our assumption  $N_D^-(v) = \emptyset$  was wrong, concluding the proof.  $\blacksquare$

By Lemma 2.7 applied to the vertex  $v$  of  $D$ , there exists an out-module  $M$  in  $D$  such that  $M \subseteq N_D^-(v)$  and  $N_D^+(M) \subseteq N_D^+(v) \cup \{v\}$ . Let  $T \subseteq M$  be the set of vertices  $t \in M$  for which there exists  $u \in N_D^+(v)$  such that  $(u, t) \in A(D)$ . Since  $N_D^+(v) \cap M = \emptyset$ , the definition of  $T$  here coincides with the one in Lemma 2.8. Now, Lemma 2.8 implies that  $D[T]$  is a (possibly empty) transitive tournament.

**Claim 2.** The digraph  $D[M]$  admits an acyclic 2-coloring  $c_M : M \rightarrow \{1, 2\}$  satisfying  $c_M(t) = 2$  for all  $t \in T$ .

*Proof.* Since  $v(D[M]) \leq v(D - v) < v(D)$ , the minimality of the counterexample  $D$  implies that  $D[M] \in \text{Forb}_{\text{ind}}(\overleftrightarrow{K}_2, S_2^+, W_3^+)$  satisfies the assertion of Theorem 6. If  $T = \emptyset$ , Claim 2 is satisfied by an arbitrary choice of an acyclic 2-coloring for  $D[M]$ . If  $T \neq \emptyset$ , let  $t_0 \in T$  be the source of the transitive tournament  $D[T]$ . Applying the assertion of the theorem to  $D[M]$  and the vertex  $t_0$ , we find that there exists an acyclic 2-coloring

of  $D[M]$  in which  $t_0$  has the same color as all its out-neighbors. W.l.o.g. we may choose this color to be 2, and since  $\{t_0\} \cup N_{D[M]}^+(t_0) \supseteq T$ , the claim follows. ■

**Claim 3.**  $D[M]$  contains a directed cycle.

*Proof.* Suppose towards a contradiction that  $D[M]$  is acyclic. Let  $D' := D - M$ . Clearly,  $D' \in \text{Forb}_{\text{ind}}(\overleftrightarrow{K}_2, S_2^+, W_3^+)$ . Since  $v(D') \leq v(D) - 1$  and by the minimality of  $D$  as a counterexample, we know that  $D'$  admits an acyclic 2-coloring  $c' : V(D) \setminus M \rightarrow \{1, 2\}$  in which  $c'(v) = c'(u) = 1$  for every  $u \in N_D^+(v)$ . Let  $c : V(D) \rightarrow \{1, 2\}$  be defined by  $c(x) := c'(x)$  for all  $x \in V(D) \setminus M$  and  $c(x) := 2$  for all  $x \in M$ . We claim that  $c$  is an acyclic coloring of  $D$ . Suppose towards a contradiction that  $C$  is a directed cycle in  $D$  which is monochromatic in the coloring  $c$ . We must have  $V(C) \cap M \neq \emptyset$ , for otherwise  $C$  would form a monochromatic directed cycle in the coloring  $c'$  of  $D'$ . Since  $D[M]$  is acyclic, we must also have  $V(C) \setminus M \neq \emptyset$ . It follows that there exists an arc  $(x, y) \in A(C)$  such that  $x \in M$  and  $y \in V(D) \setminus M$ . Then  $y \in N_D^+(M) \subseteq N_D^+(v) \cup \{v\}$ , and therefore  $c(y) = c'(y) = 1$ , while  $c(x) = 2$  by definition. This contradicts the fact that  $C$  is monochromatic, and hence we have shown that indeed  $c$  is an acyclic coloring of  $D$ . Moreover,  $c(v) = c(u) = 1$  for every  $u \in N_D^+(v)$ . This contradicts our initial assumptions on  $D$  that such a coloring does not exist. Hence,  $D[M]$  cannot be acyclic, proving the claim. ■

Claim 3 in particular implies that  $|M| \geq 3$  and  $M \setminus T \neq \emptyset$ .

Let us further note that since  $M$  forms an out-module in  $D$ ,  $M \setminus T \neq \emptyset$  is an out-module in the digraph  $D - T \in \text{Forb}_{\text{ind}}(\overleftrightarrow{K}_2, S_2^+, W_3^+)$ , and hence by Lemma 2.5 we also have  $D_0 := (D - T)/(M \setminus T) \in \text{Forb}_{\text{ind}}(\overleftrightarrow{K}_2, S_2^+, W_3^+)$ . Also note that since  $T \subseteq M \subseteq N_D^-(v)$ , we still have  $N_D^+(v) \cup \{v\} \subseteq \{x_{M \setminus T}\} \cup (V(D) \setminus M) = V(D_0)$ , where we denote by  $x_{M \setminus T}$  the vertex in  $D_0$  obtained by identifying  $M \setminus T$ .

**Claim 4.** We have  $N_{D_0}^+(v) = N_D^+(v)$ ,  $(x_{M \setminus T}, v) \in A(F(D_0))$ , and for every  $u \in N_{D_0}^+(v)$ , we have  $(u, x_{M \setminus T}) \notin A(D_0)$ .

*Proof.* The very first claim follows directly from the definition of  $D_0$ .

We have  $M \setminus T \subseteq N_D^-(v)$  and  $N_D^+(M) \subseteq \{v\} \cup N_D^+(v)$ . This directly implies that  $(x_{M \setminus T}, v) \in A(D_0)$  and that  $N_{D_0}^+(x_{M \setminus T}) \subseteq N_D^+(M) \subseteq N^+(v) \cup \{v\} = N_{D_0}^+(v) \cup \{v\}$ . Hence,  $v \in N_{D_0}^+(x_{M \setminus T})$  has an out-arc to every other out-neighbor of  $x_{M \setminus T}$  in  $D_0$ , and this shows (by definition) that  $(x_{M \setminus T}, v) \in A(F(D_0))$ .

For the second claim, suppose towards a contradiction that there exists  $u \in N_{D_0}^+(v)$  such that  $(u, x_{M \setminus T}) \in A(D_0)$ . By definition of  $D_0$ , this means that  $u \in N_D^+(v)$  and that there exists a vertex  $m \in M \setminus T$  such that  $(u, m) \in A(D)$ . By definition of  $T$ , this however shows that  $m \in T$ , a contradiction. ■

In the following, let  $D^*$  be the digraph defined by

$$V(D^*) := V(D_0), A(D^*) := A(D_0) \cup \{(x_{M \setminus T}, u) \mid u \in N_{D_0}^+(v)\}.$$

**Claim 5.**  $D^* \in \text{Forb}_{\text{ind}}(\overleftrightarrow{K}_2, S_2^+, W_3^+)$ .

*Proof.* Let  $e_i = (x_{M \setminus T}, u_i), i = 1, \dots, k$  be a list of the arcs contained in  $A(D^*) \setminus A(D_0)$  for some  $k \geq 0$ . For  $0 \leq i \leq k$  let  $D_i$  denote the digraph defined by  $V(D_i) := V(D_0)$  and  $A(D_i) := A(D_0) \cup \{e_1, \dots, e_i\}$ . Note that  $D_k = D^*$ .

We now claim that  $D_i \in \text{Forb}_{\text{ind}}(\overleftrightarrow{K}_2, S_2^+, W_3^+)$  and  $(x_{M \setminus T}, v) \in A(F(D_i))$  for every  $i \in \{0, 1, \dots, k\}$  and prove this claim by induction on  $i$ .

For  $i = 0$  the claim holds true by the previous discussions and Claim 4. Now let  $1 \leq i \leq k$  and suppose we know that the claim holds for  $D_{i-1}$ .

Note that  $D_i$  is the digraph obtained from  $D_{i-1}$  by adding the arc  $e_i = (x_{M \setminus T}, u_i)$ , where  $u_i \in N_{D_0}^+(v) = N_{D_{i-1}}^+(v)$ ,  $(x_{M \setminus T}, v) \in A(F(D_{i-1}))$ . Note that  $e_i \notin A(D_{i-1})$ , as well as  $(u_i, x_{M \setminus T}) \notin A(D_{i-1})$  by Claim 4. Hence, we may apply Lemma 2.6 to the digraph  $D_{i-1}$  with  $x = x_{M \setminus T}, y = v, z = u_i$  to find that indeed  $D_i \in \text{Forb}_{\text{ind}}(\overleftrightarrow{K}_2, S_2^+, W_3^+)$ . It remains to show that  $(x_{M \setminus T}, v) \in A(F(D_i))$ . However, the only new out-neighbor of  $x_{M \setminus T}$  in  $D_i$  compared to  $D_{i-1}$  is the vertex  $u_i$ , which is still dominated by the vertex  $v \in N_{D_i}^+(x_{M \setminus T})$  via the arc  $(v, u_i) \in A(D_i)$ , and hence  $v$  still dominates all other out-neighbors of  $x_{M \setminus T}$  in  $D_i$ . This shows that  $D_i$  satisfies the induction claim.

We have shown  $D^* = D_k \in \text{Forb}_{\text{ind}}(\overleftrightarrow{K}_2, S_2^+, W_3^+)$ , concluding the proof of Claim 5. ■

The number of vertices of  $D^*$  satisfies

$$v(D^*) = v(D_0) = v(D) - |T| - (|M \setminus T| - 1) \leq v(D) - (|M| - 1) \leq v(D) - 2 < v(D)$$

since  $|M| \geq 3$  by Claim 3. Hence, the minimality of  $D$  implies that the assertion of the theorem holds for  $D^*$ . Applying this assertion to the vertex  $x_{M \setminus T}$  in  $D^*$ , we find that there exists an acyclic 2-coloring  $c^* : V(D^*) \rightarrow \{1, 2\}$  of  $D^*$  such that  $c^*(x_{M \setminus T}) = 1 = c^*(u)$  for every  $u \in N_{D^*}^+(x_{M \setminus T})$ . Using the facts  $N_{D_0}^+(x_{M \setminus T}) \subseteq N_D^+(v) \cup \{v\}$ ,  $N_{D_0}^+(v) = N_D^+(v)$  and  $(x_{M \setminus T}, v) \in A(D_0)$ , the definition of  $D^*$  yields that  $N_{D^*}^+(x_{M \setminus T}) = N_D^+(v) \cup \{v\}$ . Hence, we have  $c^*(x_{M \setminus T}) = c^*(v) = c^*(u) = 1$  for every  $u \in N_D^+(v)$ .

Let  $c : V(D) \rightarrow \{1, 2\}$  be the coloring of  $D$  defined by  $c(x) := c_M(x)$  for every  $x \in M$ , and  $c(x) := c^*(x)$  for every  $x \in V(D) \setminus M$ . We note that  $c(v) = c(u)$  for all  $u \in N_D^+(v)$ . Hence, by the initial assumption on  $D$ , the coloring  $c$  cannot be acyclic, i.e., there is a directed cycle  $C$  in  $D$  which is monochromatic in the coloring  $c$ . Then we must have  $V(C) \setminus M \neq \emptyset$ , for otherwise  $C$  would be a monochromatic directed cycle in the acyclic coloring  $c_M$  of  $D[M]$ . Analogously, if  $V(C) \cap M = \emptyset$ , then  $C$  would be a directed cycle in  $D - M \subseteq (D - T)/(M \setminus T) = D_0 \subseteq D^*$ , a contradiction. Therefore we also have  $V(C) \cap M \neq \emptyset$ , and hence there must be an arc  $(x, y) \in A(C)$  such that  $x \in M$  and  $y \notin M$ . However, this means that  $y \in N_D^+(M) \subseteq \{v\} \cup N_D^+(v)$ , and hence  $c(y) = 1$  by the above. Since  $C$  is monochromatic, it follows that  $V(C) \subseteq c^{-1}(1)$ . In particular, since  $c(t) = c_M(t) = 2$  for every  $t \in T$ , it follows that  $C$  is a directed cycle in  $D - T$ . Let  $z$  be the first vertex of  $M$  we meet when traversing the directed cycle  $C$  in forward-direction, starting at  $y$ . Then  $z \in M \setminus T$ . Let  $P$  be the unique directed  $x, z$ -path contained in  $C$ . Then  $P$  has length at least two and satisfies  $V(P) \cap M = \{x, z\}$  and

$V(P) \subseteq c^{-1}(1)$ . Now  $(V(P) \setminus \{x, z\}) \cup \{x_{M \setminus T}\}$  forms the vertex-set of a directed cycle  $C^*$  in  $(D - T)/(M \setminus T) = D_0 \subseteq D^*$  containing  $x_{M \setminus T}$ , and we have  $c^*(x) = c(x) = 1$  for every vertex  $x \in V(C^*) \setminus \{x_{M \setminus T}\} = V(P) \setminus \{x, z\} \subseteq V(D) \setminus M$ . We have  $c^*(x_{M \setminus T}) = 1$  by definition of  $c^*$ , and hence  $C^*$  forms a monochromatic directed cycle of color 1 in the acyclic coloring  $c^*$  of  $D^*$ . This contradiction finally shows that our very first assumption, namely that a (smallest) counterexample  $D$  to the claim of the theorem exists, was wrong. This concludes the proof of the theorem.  $\blacksquare$

### 3 $\{S_2^+, W_3^-\}$ -Free Oriented Graphs

In this section we prove Theorem 3, showing that every digraph in  $\text{Forb}_{\text{ind}}(\vec{K}_2, S_2^+, W_3^-)$  is acyclically 4-colorable. Note that  $\text{Forb}_{\text{ind}}(\vec{K}_2, S_2^+, W_3^-)$  is the class of oriented graphs  $D$  such that the out-neighborhood of any vertex in  $D$  spans a tournament, and the in-neighborhood of any vertex spans a directed triangle-free graph. In fact, we show the following strengthened statement.

**Theorem 7.** *Let  $D \in \text{Forb}_{\text{ind}}(\vec{K}_2, S_2^+, W_3^-)$  and let  $(u, v) \in A(D)$ . Then  $D$  admits an acyclic coloring  $c : V(D) \rightarrow \{1, 2, 3, 4\}$  satisfying the additional conditions  $c(u) = 1$ ,  $c(x) = 1$  for all  $x \in N_D^+(u) \setminus N_D^+(v)$  (so  $c(v) = 1$ ) and  $c(x) \in \{1, 2\}$  for all  $x \in N_D^+(v)$ .*

*Proof.* Suppose towards a contradiction the claim of the theorem was wrong, and let  $D$  be a counterexample minimizing  $v(D)$ . Then there exists an arc  $(u, v) \in A(D)$  such that  $D$  does not admit an acyclic coloring  $c : V(D) \rightarrow \{1, 2, 3, 4\}$  satisfying the additional conditions  $c(u) = 1$ ,  $c(x) = 1$  for all  $x \in N_D^+(u) \setminus N_D^+(v)$  and  $c(x) \in \{1, 2\}$  for all  $x \in N_D^+(v)$ . Let us define  $A := N_D^+(u) \setminus (N_D^+(v) \cup \{v\})$  and  $B := N_D^-(u) \cap N_D^+(v)$ . We start with some useful observations concerning these sets.

**Claim 1.**  $A \subseteq N_D^-(v)$ , and  $D[A]$  and  $D[B]$  are transitive tournaments.

*Proof.* To show  $A \subseteq N_D^-(v)$ , let  $x \in A = N_D^+(u) \setminus (N_D^+(v) \cup \{v\})$  be arbitrary. Since  $(u, x), (u, v) \in A(D)$  and  $x, u, v$  cannot induce an  $S_2^+$  in  $D$ , the vertices  $x$  and  $v$  must be equal or adjacent in  $D$ . Since  $x \notin N_D^+(v) \cup \{v\}$ , it follows that  $x \in N_D^-(v)$ , as claimed.

Since  $D[N_D^+(u)]$  is a tournament and  $A \subseteq N_D^+(u)$ , also  $D[A]$  is a tournament. Furthermore  $D[N_D^-(v)]$  is directed triangle-free, and with  $A \subseteq N_D^-(v)$  also  $D[A]$  is directed triangle-free, i.e., a transitive tournament, as claimed.

Similarly, since  $D[N_D^-(u)]$  is directed triangle-free, and since  $D[N_D^+(v)]$  is a tournament,  $B = N_D^-(u) \cap N_D^+(v)$  implies that  $D[B]$  must be both directed triangle-free and a tournament, i.e., a transitive tournament.  $\blacksquare$

In the following, let us denote by  $D' := D - (N_D^-(u) \cup \{u\})$  the induced subdigraph of  $D$  obtained by deleting the closed in-neighborhood of  $u$ . We clearly have  $D' \in \text{Forb}_{\text{ind}}(\vec{K}_2, S_2^+, W_3^-)$  and  $v(D') < v(D)$ , and hence by minimality of  $D$  the theorem statement holds for  $D'$ .

**Claim 2.** There exists an acyclic coloring  $c' : V(D') \rightarrow \{1, 2, 3, 4\}$  of  $D'$  such that  $c'(v) = 1$ ,  $c'(x) = 1$  for all  $x \in A$  and  $c'(x) \in \{1, 2\}$  for all  $x \in N_{D'}^+(v)$ .

*Proof.* We distinguish the two cases  $A = \emptyset$  and  $A \neq \emptyset$ .

Suppose first that  $A = \emptyset$ . If  $N_{D'}^+(v) = \emptyset$ , then applying the theorem statement to  $D'$  (for an arbitrarily chosen arc) yields that  $\vec{\chi}(D') \leq 4$ , and hence there exists an acyclic coloring  $c' : V(D') \rightarrow \{1, 2, 3, 4\}$ . Since  $v$  is a sink in  $D'$ , no directed cycle in  $D'$  contains  $v$ . Consequently, we may assume w.l.o.g. (possibly by recoloring) that  $c'(v) = 1$ . In particular, since  $A = \emptyset$ ,  $N_{D'}^+(v) = \emptyset$ , the remaining two statements of Claim 2 are satisfied vacuously for  $c'$ , concluding the proof in this case.

On the other hand, if  $N_{D'}^+(v) \neq \emptyset$ , then there exists an arc in  $D'$  leaving  $v$ . Fix an arbitrary such arc  $(v, y)$ . Applying the Theorem statement to this arc in  $D'$ , we find that there is an acyclic coloring  $c' : V(D') \rightarrow \{1, 2, 3, 4\}$  such that  $c'(v) = 1$ ,  $c'(x) = 1$  for all  $x \in N_{D'}^+(v) \setminus N_{D'}^+(y)$  and  $c'(x) \in \{1, 2\}$  for all  $x \in N_{D'}^+(y)$ ; in particular, and  $c'(x) \in \{1, 2\}$  for all  $x \in N_{D'}^+(v)$ . Again, this shows that the claim holds true.

Next suppose that  $A \neq \emptyset$ . By Claim 1,  $D[A]$  is a transitive tournament. Let  $a$  be the unique source-vertex of this tournament. Since  $a \in A \subseteq N_D^-(v)$ , it follows that  $(a, v) \in A(D')$ . Hence, we may apply the Theorem statement to the arc  $(a, v)$  in  $D'$  and find that there exists an acyclic coloring  $c' : V(D') \rightarrow \{1, 2, 3, 4\}$  such that  $c'(a) = 1$ ,  $c'(x) = 1$  for all  $x \in N_{D'}^+(a) \setminus N_{D'}^+(v)$  (in particular  $c'(v) = 1$ ), and  $c'(x) \in \{1, 2\}$  for all  $x \in N_{D'}^+(v)$ . Since  $a$  is the source of  $D[A] = D'[A]$  and since  $A \cap N_D^+(v) = \emptyset$ , we have  $\{a\} \cup (N_{D'}^+(a) \setminus N_{D'}^+(v)) \supseteq A$  and thus  $c'(x) = 1$  for all  $x \in A$ , as required. This shows the assertion of Claim 2 and concludes the proof.  $\blacksquare$

**Claim 3.** There exists an acyclic coloring  $c^- : N_D^-(u) \rightarrow \{2, 3, 4\}$  of  $D[N_D^-(u)]$  such that  $c^-(x) = 2$  for all  $x \in B$  and  $c^-(x) \in \{3, 4\}$  for all  $x \in N_D^-(u) \setminus B$ .

*Proof.* Since  $D \in \text{Forb}_{\text{ind}}(\vec{K}_2^+, S_2^+, W_3^-)$ , we have  $D[N_D^-(u)] \in \text{Forb}_{\text{ind}}(\vec{K}_2^+, S_2^+, \vec{C}_3)$ . By Theorem 1 there exists an acyclic coloring of  $D[N_D^-(u)] - B$  using only colors 3 and 4. Clearly,  $D[B]$  as a transitive tournament (see Claim 1) admits an acyclic coloring only with color 2. Putting these colorings together yields an acyclic coloring  $c'$  of  $D[N_D^-(u)]$  with the required properties.  $\blacksquare$

Let  $c : V(D) \rightarrow \{1, 2, 3, 4\}$  be the coloring defined by  $c(u) := 1$ ,  $c(x) := c^-(x)$  for all  $x \in N_D^-(u)$  and  $c(x) := c'(x)$  for all  $x \in V(D) \setminus (N_D^-(u) \cup \{u\})$ .

Note that from the properties of  $c'$  given by Claim 2 we have  $c(u) = c(v) = 1$  and  $c(x) = 1$  for all  $x \in A = N_D^+(u) \setminus (N_D^+(v) \cup \{v\})$ . Furthermore, since  $N_D^+(v) = N_{D'}^+(v) \cup B$ , the properties of  $c'$  and  $c^-$  imply that  $c(x) \in \{1, 2\}$  for all  $x \in N_D^+(v)$ .

Given these properties, our initial assumption concerning  $D$  implies that  $c$  cannot be an acyclic coloring of  $D$ , that is, there is a directed cycle  $C$  in  $D$  which is monochromatic under  $c$ . Since  $c' = c|_{V(D')}$  and  $c^- = c|_{N_D^-(u)}$ , we must have  $V(C) \cap (N_D^-(u) \cup \{u\}) \neq \emptyset$  and  $V(C) \setminus N_D^-(u) \neq \emptyset$ , for otherwise  $c'$  resp.  $c^-$  would not be acyclic. Further note that  $u \notin V(C)$ , for every arc in  $D$  entering  $u$  has its tail colored with either 2, 3 or 4, while its head,  $u$ , receives color 1 under  $c$  (so a directed cycle containing  $u$  cannot be



monochromatic). Hence, there must exist an arc  $(x, y) \in A(C)$  such that  $x \in N_D^-(u)$  and  $y \in V(D) \setminus (N_D^-(u) \cup \{u\})$ . Since  $(x, u), (x, y) \in A(D)$  and  $D$  is induced  $S_2^+$ -free,  $u$  and  $y$  must be equal or adjacent, and since  $y \notin N_D^-(u) \cup \{u\}$ , we have  $y \in N_D^+(u)$ . By the properties of  $c'$  and  $c^-$ , we have  $N_D^+(u) \setminus N_D^+(v) = A \cup \{v\} \subseteq c^{-1}(\{1\})$ ,  $B \subseteq c^{-1}(\{2\})$  and  $N_D^-(u) \setminus B \subseteq c^{-1}(\{3, 4\})$ . The cycle  $C$  is monochromatic, therefore  $c(x) = c(y)$ . From this we conclude that  $y \in N_D^+(v)$ , and hence  $c(y) \in \{1, 2\}$ . This is only possible if  $c(x) = c(y) = 2$ , and hence  $x \in B$ . It follows that  $x, y, u, v \in V(D)$  are distinct vertices satisfying  $(x, y), (u, y), (v, y) \in A(D)$ , as well as  $(x, u), (u, v), (v, x) \in A(D)$  (here we used that  $x \in B = N_D^-(u) \cap N_D^+(v)$ ). This however means that  $x, y, u, v$  induce a copy of  $W_3^-$  in  $D$ , which is absurd considering that  $D \in \text{Forb}_{\text{ind}}(\vec{K}_2, S_2^+, W_3^-)$ . This shows that our very first assumption concerning the existence of a smallest counterexample  $D$  was wrong. This concludes the proof of the theorem.  $\blacksquare$

## 4 Adding a Dominating Sink to a Hero

In this section our goal is to prove Theorem 4. Let us first prove the following lemma.

**Lemma 4.1.** *Let  $D \in \text{Forb}_{\text{ind}}(\vec{K}_2, S_2^+)$  and let  $C \in \mathbb{N}$  be such that  $\vec{\chi}(D[N_D^-(x)]) \leq C$  for every  $x \in V(D)$ . Let  $u, v \in V(D)$  and let  $P$  be a shortest  $u$ - $v$ -dipath in  $D$ . Let  $X := V(P) \cup N_D^-(V(P))$ . Then  $\vec{\chi}(D[X]) \leq 3C + 2$ .*

*Proof.* Let  $u = x_0, x_1, \dots, x_{\ell-1}, x_\ell = v$  be the vertex-trace of  $P$  and consider the partition  $(A_i)_{i=1}^\ell$  of  $N_D^-(V(P))$  where  $A_i := N_D^-(x_i) \setminus (V(P) \cup \bigcup_{1 \leq j < i} A_j)$ ,  $i = 0, \dots, \ell$ .

**Claim.** Let  $0 \leq i < j \leq \ell$  with  $j - i \geq 3$ . Then there exists no arc in  $D$  starting in  $A_i$  and ending in  $A_j$ .

*Proof.* Suppose towards a contradiction that there are vertices  $x \in A_i, y \in A_j$  with  $(x, y) \in A(D)$ . Since  $(x, x_i) \in A(D), (x, y) \in A(D)$  and  $x_i \neq y$  (since  $x_i \in V(P)$  and  $y \notin V(P)$ ),  $x_i$  and  $y$  must be adjacent in  $D$ . By definition of  $A_j$  we have  $A_j \cap N_D^-(x_i) = \emptyset$  and hence  $(x_i, y) \in A(D)$ . However, now the directed path described by the vertices  $u = x_0, x_1, \dots, x_i, y, x_j, \dots, x_\ell = v$  is a  $u$ - $v$ -dipath in  $D$  shorter than  $P$ , a contradiction. This proves the claim.  $\blacksquare$

For every  $0 \leq i \leq \ell$  we have  $\vec{\chi}(D[A_i]) \leq \vec{\chi}(D[N_D^-(x_i)]) \leq C$ . Let us define the set  $B_r := \bigcup \{A_i \mid i \equiv r \pmod{3}\}$  for every  $r \in \{0, 1, 2\}$ . From the above claim it follows that no directed cycle in  $D[B_r]$  intersects two different sets  $A_i, A_j$ . Hence, we have

$$\vec{\chi}(D[B_r]) \leq \max\{\vec{\chi}(D[A_i]) \mid i \equiv r \pmod{3}\} \leq C$$

for  $r = 0, 1, 2$ . Further note that the two sets

$$V_0 := \{x_i \mid i \in \{0, \dots, \ell\} \text{ even}\}, V_1 := \{x_i \mid i \in \{0, \dots, \ell\} \text{ odd}\}$$

both induce acyclic subdigraphs of  $D$ , for otherwise  $D$  would not be a shortest  $u$ - $v$ -dipath in  $D$ . Since  $X$  is the disjoint union of  $B_0, B_1, B_2, V_0, V_1$ , we conclude

$$\bar{\chi}(D[X]) \leq \bar{\chi}(D[B_0]) + \bar{\chi}(D[B_1]) + \bar{\chi}(D[B_2]) + \bar{\chi}(D[V_0]) + \bar{\chi}(D[V_1]) \leq 3C + 2,$$

as required. ■

*Proof of Theorem 4.* Let  $\{\vec{K}_2, S_2^+, H\}$  be heroic and  $C := \bar{\chi}(\text{Forb}_{\text{ind}}(\vec{K}_2, S_2^+, H))$ .

We claim that every digraph  $D \in \text{Forb}_{\text{ind}}(\vec{K}_2, S_2^+, H^-)$  admits an acyclic coloring with  $C^- := v(H)(C + 1) + 3C + 2$  colors.

Suppose towards a contradiction that there exists some  $D \in \text{Forb}_{\text{ind}}(\vec{K}_2, S_2^+, H^-)$  with  $\bar{\chi}(D) > C'$ , and choose such a  $D$  minimizing  $v(D)$ . Then we have  $\bar{\chi}(D) > C' \geq C$  and hence there is  $Y \subseteq V(D)$  such that  $D[Y]$  is isomorphic to  $H$ . Furthermore, since the dichromatic number of  $D$  is the maximum of the dichromatic numbers of its strong components, the minimality of  $v(D)$  implies that  $D$  is strongly connected.

Let  $S \supseteq Y$  denote a set of vertices in  $D$  defined as follows:

If  $D[Y]$  (resp.  $H$ ) is strongly connected, put  $S := Y$ . Otherwise, let  $Y_1, \dots, Y_t$  be a partition of  $Y$  into the  $t \geq 2$  strong components of  $D[Y]$  such that all arcs between  $Y_i$  and  $Y_j$  start in  $Y_i$  and end in  $Y_j$ , for any  $1 \leq i < j \leq t$  (note that since  $D[Y]$  is a tournament all elements of  $Y_i \times Y_j$  are arcs of  $D[Y]$  for  $1 \leq i < j \leq t$ ). Now pick  $u \in Y_t$ ,  $v \in Y_1$  arbitrarily, let  $P$  be a shortest  $u$ - $v$ -dipath in  $D$  and put  $S := V(P) \cup Y$ . Let us note that in any case,  $D[S]$  is strongly connected.

Let  $Z := S \cup N_D^-(S)$ . Then we have  $Z = X \cup Y \cup N_D^-(Y)$ , where  $X$  is defined as  $X := \emptyset$  if  $S = Y$  and as  $X := V(P) \cup N_D^-(V(P))$  otherwise. For every  $x \in V(D)$  we know that since  $D$  is  $H^-$ -free, the digraph  $D[N_D^-(x)]$  is contained in  $\text{Forb}_{\text{ind}}(\vec{K}_2, S_2^+, H)$ , and hence  $\bar{\chi}(D[N_D^-(x)]) \leq C$ . Using Lemma 4.1 we obtain that  $\bar{\chi}(D[X]) \leq 3C + 2$ . Putting it all together, we find that

$$\bar{\chi}(D[Z]) \leq \sum_{y \in Y} \underbrace{\bar{\chi}(D[\{y\} \cup N_D^-(y)])}_{\leq C+1} + \bar{\chi}(D[X]) \leq v(H)(C + 1) + 3C + 2 = C'.$$

**Claim.** No arc in  $D$  leaves  $Z$ .

*Proof.* We first show that there do not exist  $z \in S$ ,  $w \in V(D) \setminus Z$  such that  $(z, w) \in A(D)$ . Suppose towards a contradiction that such an arc  $(z, w)$  exists. We claim that then  $(s, w) \in A(D)$  for every  $s \in S$ . Consider  $s \in S$  arbitrarily. Since  $D[S]$  is strongly connected, there exist vertices  $z = s_0, s_1, \dots, s_k = s$  in  $S$  such that  $(s_{i-1}, s_i) \in A(D)$ ,  $i = 1, \dots, k$ . We show  $(s_i, w) \in A(D)$  for all  $i = 0, \dots, k$  by induction on  $i$ . Clearly it is true for  $i = 0$ , so suppose that  $1 \leq i \leq k$  and we have established that  $(s_{i-1}, w) \in A(D)$ . Since  $w \notin Z$ ,  $s_i \in Z$ , we have  $w \neq s_i$  and  $(s_{i-1}, w), (s_{i-1}, s_i) \in A(D)$ . Since  $D$  is  $S_2^+$ -free, it follows that  $s_i$  and  $w$  are adjacent. However, since  $w \notin Z = S \cup N_D^-(S) \supseteq N_D^-(s_i)$ , we must have  $(s_i, w) \in A(D)$ , as claimed.

This shows that indeed  $(s, w) \in A(D)$  for all  $s \in S$ . However, since  $S \supseteq Y$  and since  $D[Y]$  is isomorphic to  $H$ , it follows that  $D[Y \cup \{w\}]$  is an induced subdigraph of  $D$

isomorphic to  $H^-$ , a contradiction to  $D \in \text{Forb}_{\text{ind}}(\vec{K}_2, S_2^+, H^-)$ . This shows that there are no arcs from  $S$  to  $V(D) \setminus Z$ .

To complete the proof, let us show that there are no arcs starting in  $Z \setminus S = N_D^-(S)$  that end in  $V(D) \setminus Z$ . Suppose towards a contradiction that there exist  $z \in N_D^-(S)$  and  $w \in V(D) \setminus Z$  with  $(z, w) \in A(D)$ . Then there is a vertex  $s \in S$  such that  $(z, s) \in A(D)$ . Since  $s \neq w$ ,  $(z, s), (z, w) \in A(D)$  and  $D$  is  $S_2^+$ -free we find that  $s$  and  $w$  are adjacent. Since  $w \notin Z \supseteq N_D^-(s)$ , it follows that  $(s, w) \in A(D)$ . However, this yields a contradiction, since we showed above that no arc in  $D$  starts in  $S$  and ends in  $V(D) \setminus Z$ . All in all, the claim follows.  $\blacksquare$

Since  $D$  is strongly connected and  $Z \neq \emptyset$  (since  $Z \supseteq Y$ ), it follows that  $Z = V(D)$ , and hence that  $\vec{\chi}(D) = \vec{\chi}(D[Z]) \leq C'$ , a contradiction to our initial assumption. This concludes the proof of the theorem.  $\blacksquare$

## 5 Oriented 4-Vertex-Paths

In this section we establish that  $\{\vec{K}_2, \vec{K}_3, P^+(1, 1, 1)\}$  is heroic proving Theorem 5.

*Proof of Theorem 5.* We prove by induction on  $n$  that every directed graph on  $n$  vertices  $D \in \text{Forb}_{\text{ind}}(\vec{K}_2, \vec{K}_3, P^+(1, 1, 1))$  admits an acyclic 2-coloring. The claim trivially holds for  $n = 1$ , so let  $n \geq 2$  and suppose that every digraph in  $\text{Forb}_{\text{ind}}(\vec{K}_2, \vec{K}_3, P^+(1, 1, 1))$  having less than  $n$  vertices is 2-colorable. Pick some  $x \in V(D)$  arbitrarily. Let us define a sequence  $X_0, X_1, X_2, \dots$  of subsets of  $V(D)$  as follows:

$$X_i := \begin{cases} \{x\}, & \text{if } i = 0, \\ N^+(X_{i-1}) \setminus \bigcup_{j=0}^{i-1} X_j, & \text{if } i \text{ odd,} \\ N^-(X_{i-1}) \setminus \bigcup_{j=0}^{i-1} X_j, & \text{if } i \geq 2 \text{ even.} \end{cases}$$

The sets  $(X_i)_{i \geq 0}$  are by definition pairwise disjoint, and so there exists  $k \geq 1$  such that  $X_1, \dots, X_k \neq \emptyset$  and  $X_i = \emptyset$  for all  $i > k$ .

**Claim.**  $X_i$  is an independent set of  $D$  for every  $i \geq 0$ .

*Proof.* We prove the claim by induction on  $i$ . The claim trivially holds for  $i = 0$  since  $X_0 = \{x\}$ , and since  $D$  does not contain a transitive triangle, also  $X_1 = N^+(x)$  must be an independent set in  $D$ . Now let  $i \geq 2$  and suppose that we already established that  $X_0, \dots, X_{i-1}$  are independent. To show that  $X_i$  is independent, let us suppose towards a contradiction that there are  $x, y \in X_i$  such that  $(x, y) \in A(D)$ . By definition of the sets  $X_i$  there are vertices  $x_1, y_1 \in X_{i-1}$  and  $x_2, y_2 \in X_{i-2}$  such that the following holds:  $(x_1, x_2), (x_1, x), (y_1, y_2), (y_1, y) \in A(D)$  if  $i$  is odd, respectively  $(x_2, x_1), (x, x_1), (y_2, y_1), (y, y_1) \in A(D)$  if  $i$  is even. We must have  $x_1 \neq y_1$  in any case, since otherwise the vertices  $x_1 = y_1, x, y$  would induce a  $\vec{K}_3$  in  $D$ . Let us now consider the oriented 4-vertex-path  $P$  in  $D$  defined as  $P = x, (x, y), y, (y_1, y), y_1, (y_1, y_2), y_2$  if  $i$  is odd, respectively as  $P = x_2, (x_2, x_1), x_1, (x, x_1), x, (x, y), y$  if  $i$  is even. In order for

this path not to be an induced copy of  $P^+(1, 1, 1)$ , two non-consecutive vertices of the path must be adjacent. However, since  $D$  does not contain transitive triangles, this is only possible if  $x$  and  $y_2$  ( $i$  odd) respectively  $x_2$  and  $y$  ( $i$  even) are adjacent. Since  $x \notin X_{i-1}$ , we have  $x \notin N^-(X_{i-2})$  if  $i$  is odd and  $y \notin N^+(X_{i-2})$  if  $i$  is even. Since  $x_2, y_2 \in X_{i-2}$  we conclude that  $(y_2, x) \in A(D)$  if  $i$  is odd and  $(y, x_2) \in A(D)$  if  $i$  is even. In both cases we conclude that  $x_2 \neq y_2$ , since otherwise the vertices  $x_2 = y_2, x_1, x$  respectively  $x_2 = y_2, y_1, y$  would induce a transitive triangle in  $D$ . Now consider the oriented path  $Q$  in  $D$  defined as  $Q = y_2, (y_2, x), x, (x_1, x), x_1, (x_1, x_2), x_2$  if  $i$  is odd and as  $Q = y_2, (y_2, y_1), y_1, (y, y_1), y, (y, x_2), x_2$  if  $i$  is even. In order for  $Q$  not to be an induced copy of  $P^+(1, 1, 1)$  in  $D$  and since  $D$  does not contain transitive triangles, this implies in both cases that the endpoints  $x_2$  and  $y_2$  of  $Q$  must be adjacent. This contradicts the induction hypothesis that  $X_{i-2}$  is an independent set. Hence, our assumption was wrong,  $X_i$  is indeed independent. This concludes the proof of the claim. ■

Let  $X := X_0 \cup \dots \cup X_k$  and  $D' := D - X$ . By the induction hypothesis  $D'$  admits an acyclic 2-coloring  $c' : V(D') \rightarrow \{1, 2\}$ . Let us now define  $c : V(D) \rightarrow \{1, 2\}$  by  $c(x) := c'(x)$  for every  $x \in V(D) \setminus X$ ,  $c(x) := 1$  for every  $x \in X_i$  such that  $i$  is even, and  $c(x) := 2$  for every  $x \in X_i$  such that  $i$  is odd. We claim that  $D$  defines an acyclic coloring of  $D$ . Suppose towards a contradiction that there exists a monochromatic directed cycle  $C$  in  $(D, c)$ . Since  $c'$  is an acyclic coloring of  $D'$ , we must have  $V(C) \cap (X_0 \cup \dots \cup X_k) \neq \emptyset$ . Note that by definition of the sets  $(X_i)_{i \geq 0}$  we have  $N^+(\bigcup_{i \text{ even}} X_i), N^-(\bigcup_{i \text{ odd}} X_i) \subseteq X$ . Hence, no arc of  $D$  starts in  $c^{-1}(\{1\}) \cap X$  and ends in  $V(D) \setminus X$ , and no arc of  $D$  starts in  $c^{-1}(\{2\}) \cap X$  and ends in  $V(D) \setminus X$ . Since  $V(C) \subseteq c^{-1}(t)$  for some  $t \in \{1, 2\}$ , the strong connectivity of  $C$  shows that in fact  $V(C) \subseteq c^{-1}(t) \cap X$  for some  $t \in \{1, 2\}$ . Let  $i_0 \geq 0$  be smallest such that  $X_{i_0} \cap V(C) \neq \emptyset$ . Let  $u \in X_{i_0} \cap V(C) \neq \emptyset$ , and let  $u^-, u^+ \in V(C)$  be such that  $(u^-, u), (u, u^+) \in A(C)$ . Since  $X_{i_0}$  is an independent set, and by definition of the coloring  $c$ , we must have  $u^- \in X_{i_0+2s^-}, u^+ \in X_{i_0+2s^+}$  for integers  $s^-, s^+ \geq 1$ . On the other hand, we have  $u^+ \in N^+(X_{i_0}) \setminus \bigcup_{j=0}^{i_0-1} X_j = X_{i_0+1}$  if  $i_0$  is even and  $u^- \in N^-(X_{i_0}) \setminus \bigcup_{j=0}^{i_0-1} X_j = X_{i_0+1}$  if  $i_0$  is odd, in both cases yielding a contradiction since  $X_{i_0+2s^+}, X_{i_0+2s^-}$  are disjoint from  $X_{i_0+1}$ . This shows that our assumption was wrong, indeed,  $c$  is an acyclic coloring of  $D$ . Hence,  $\bar{\chi}(D) \leq 2$ , concluding the proof. ■

## 6 Conclusion

In the first three sections of this paper we have proved that set  $\{\vec{K}_2, S_2^+, H\}$  is heroic for several small heroes  $H$ , and in particular we resolved Conjecture 1.7. It would be interesting to prove that in fact, for any hero  $H$ ,  $\{\vec{K}_2, S_2^+, H\}$  is heroic, as this would be a broad generalization of the main result of Berger et al. [7] from tournaments to *locally out-complete* oriented graphs, i.e., oriented graphs in which the out-neighborhood of every vertex induces a tournament. This class of digraphs has been thoroughly studied in the past, see for instance [5] for a survey of results on locally complete digraphs.

The smallest open case of this problem would be to show that  $\{\vec{K}_2, S_2^+, \vec{K}_4^s\}$  is heroic, where  $\vec{K}_4^s$  denotes the unique strong tournament on four vertices. It seems that already for this case a new method is required. We do however believe that the following is true.

**Conjecture 6.1.**  $\vec{\chi}(\text{Forb}_{\text{ind}}(\vec{K}_2, S_2^+, \vec{K}_4^s)) = 3$ .

Here, a tight lower bound would be provided by the following construction: Take a 3-fold blow-up of a directed 4-cycle (every arc being replaced by an oriented  $K_{3,3}$ ) and connect each of the three blow-up triples by a directed triangle. This oriented graph is contained in  $\text{Forb}_{\text{ind}}(\vec{K}_2, S_2^+, \vec{K}_4^s)$  and has dichromatic number 3.

Let us further remark at this point that there exists a very simple proof that if we exclude *both*  $S_2^+$  and  $S_2^-$ , i.e., we consider *locally complete* oriented graphs (where the in- and out-neighborhood of every vertex induces a tournament), then we can show that the exclusion of any hero indeed bounds the dichromatic number as follows.

**Remark 6.2.** *For any hero  $H$ , we have*

$$\vec{\chi}(\text{Forb}_{\text{ind}}(\vec{K}_2, S_2^+, S_2^-, H)) \leq 2\vec{\chi}(\text{Forb}_{\text{ind}}(\vec{K}_2, \overline{K}_2, H)) < \infty.$$

*Proof.* By the result of Berger et al. [7] we have  $C := \vec{\chi}(\text{Forb}_{\text{ind}}(\vec{K}_2, \overline{K}_2, H)) < \infty$ . Let us now prove that  $\vec{\chi}(\text{Forb}_{\text{ind}}(\vec{K}_2, S_2^+, S_2^-, H)) \leq 2C$ . Towards a contradiction suppose that  $\vec{\chi}(D) > 2C$  for some  $D \in \text{Forb}_{\text{ind}}(\vec{K}_2, S_2^+, S_2^-, H)$ , and pick  $D$  such that  $v(D)$  is minimum. Pick  $v \in V(D)$  arbitrarily and define  $D' := D - (\{v\} \cup N_D(v))$ .

Since  $v(D') < v(D)$ , there exists an acyclic  $2C$ -coloring  $c' : V(D') \rightarrow \{1, \dots, 2C\}$  of  $D'$ . Since  $D$  is induced  $S_2^+, S_2^-$ -free, we further know that  $D^+ := D[\{v\} \cup N_D^+(v)]$  and  $D^- := D[N_D^-(v)]$  are tournaments excluding  $H$ . It follows from the definition of  $C$  that there exists an acyclic  $C$ -coloring  $c^+ : V(D^+) \rightarrow \{1, \dots, C\}$  of  $D^+$  as well as an acyclic coloring  $c^- : V(D^-) \rightarrow \{C+1, \dots, 2C\}$  of  $D^-$ . Let  $c$  be the  $2C$ -coloring of  $D$  defined as the common extension of  $c', c^+, c^-$  to  $V(D)$ . Since  $\vec{\chi}(D) > 2C$  there exists a directed cycle  $C$  which is monochromatic under  $c$ . Since  $c', c^+, c^-$  are acyclic colorings and since the color sets used by  $c^+$  and  $c^-$  are disjoint, we must have  $V(C) \cap (\{v\} \cup N_D(v)) \neq \emptyset$ ,  $V(C) \setminus (\{v\} \cup N_D(v)) \neq \emptyset$ . Since all in-neighbors of  $v$  have a distinct color from  $v$ , we further have  $v \notin V(C)$ . We conclude that there are vertices  $x_1, x_2 \in V(C) \cap N_D(v)$ ,  $y_1, y_2 \in V(C) \setminus (\{v\} \cup N_D(v))$  such that  $(x_1, y_1), (y_2, x_2) \in A(C)$ . We claim that we must have  $x_1 \in N_D^+(v)$  and  $x_2 \in N_D^-(v)$ . Indeed, otherwise we would have  $(x_1, v) \in A(D)$  or  $(v, x_2) \in A(D)$ , and then either the vertices  $x_1, v, y_1$  induce an  $S_2^+$  in  $D$ , or  $x_2, v, y_2$  induce an  $S_2^-$  in  $D$ , in each case yielding a contradiction to  $D \in \text{Forb}_{\text{ind}}(\vec{K}_2, S_2^+, S_2^-, H)$ . Finally, we conclude that  $c(x_1) = c^+(x_1) \leq C < c^-(x_2) = c(x_2)$ , a contradiction to the facts that  $C$  is monochromatic and  $x_1, x_2 \in V(C)$ . This shows that our initial assumption concerning the existence of  $D$  was wrong, concluding the proof of the remark.  $\blacksquare$

In the last section of this paper we investigated oriented graphs excluding the anti-directed 4-vertex-path  $P^+(1, 1, 1)$ . It would certainly be very interesting and insightful to generalize both Theorem 5 as well as the result of Aboulker et al. concerning  $\vec{P}_4$  by proving that  $\{\vec{K}_2, \vec{P}_4, \vec{K}_k\}$  and  $\{\vec{K}_2, P^+(1, 1, 1), \vec{K}_k\}$  are heroic for all  $k \geq 4$ .

**Acknowledgements.** My sincerest thanks go to Lior Gishboliner and Tibor Szabó for stimulating and fruitful discussions on the topic which contributed to the results presented in this paper. I would also like to thank Lior Gishboliner for improving the presentation of some parts of the write-up.

## References

- [1] P. Aboulker, P. Charbit and R. Naserasr. Extension of the Gyárfás-Sumner conjecture to digraphs. *arXiv preprint*, arXiv:2009.13319, 2020.
- [2] P. Aboulker, N. Cohen, F. Havet, W. Lochet, P. F. S. Moura and S. Thomassé. Subdivisions in digraphs of large outdegree or large dichromatic number. *Electronic Journal of Combinatorics*, 26(3), P3.19, 2019.
- [3] P. Aboulker, F. Havet, K. Knauer and C. Rambaud. On the dichromatic number of surfaces. *arXiv preprint*, arXiv:2102.01034, 2021.
- [4] S. D. Andres and W. Hochstättler. Perfect digraphs. *Journal of Graph Theory*, 79(1), 21–29, 2015.
- [5] J. Bang-Jensen. Locally Semicomplete Digraphs and Generalizations. *Chapter 6 of Book “Classes of Directed Graphs” (J. Bang-Jensen and G. Gutin ed.)*, Springer Monographs in Mathematics, ISBN: 978-3-319-71839-2, 2018.
- [6] J. Bang-Jensen, T. Bellitto, T. Schweser and M. Stiebitz. On DP-colorings of digraphs. *Journal of Graph Theory*, 95(1), 76–98, 2020.
- [7] E. Berger, K. Choromanski, M. Chudnovsky, J. Fox, M. Loeb, A. Scott, P. Seymour and S. Thomassé. Tournaments and colouring. *Journal of Combinatorial Theory, Series B*, 103(1), 1–20, 2013.
- [8] D. Bokal, G. Fijavz, M. Juvan, P. M. Kayll and B. Mohar. The circular chromatic number of a digraph. *Journal of Graph Theory*, 46(3), 227–240, 2004.
- [9] M. Chudnovsky, A. Scott and P. Seymour. Induced subgraphs of graphs with large chromatic number. XI. Orientations. *European Journal of Combinatorics*, 76, 53–61, 2019.
- [10] M. Chudnovsky, A. Scott and P. Seymour. Induced subgraphs of graphs with large chromatic number. XII. Distant stars. *Journal of Graph Theory*, 92(3), 237–254, 2019.
- [11] P. Erdős. Graph theory and probability. *Canadian Journal of Mathematics*, 11, 34–38, 1959.
- [12] P. Erdős. Problems and results in number theory and graph theory. *Congress. Numer. XXVII, (Proc. 9th Manitoba Conf. Num. Math. Comput. 1979) Utilitas Math., Winnipeg*, 3–21, 1980.

- [13] A. Gyárfás. On Ramsey covering-numbers. *Infinite and finite sets (Colloq., Keszthely, 1973, dedicated to P. Erdős on his 60th birthday), Vol. II.* Colloq. Math. Soc. János Bolyai, 10, Amsterdam: North-Holland, 801–816, 1975.
- [14] A. Harutyunyan and B. Mohar. Two results on the digraph chromatic number. *Discrete Mathematics*, 312(10), 1823–1826, 2012.
- [15] A. Harutyunyan, T.-N. Le, A. Newman and S. Thomassé. Coloring dense digraphs. *Combinatorica*, 39, 1021–1053, 2019.
- [16] A. Harutyunyan, T.-N. Le, S. Thomassé and H. Wu. Coloring tournaments: From local to global. *Journal of Combinatorial Theory, Series B*, 138, 166–171, 2019.
- [17] W. Hochstättler. A flow theory for the dichromatic number. *European Journal of Combinatorics*, 66, 160–167, 2017.
- [18] H. A. Kierstead and S. G. Penrice. Radius two trees specify  $\chi$ -bounded classes. *Journal of Graph Theory*, 18(2), 119–129, 1994.
- [19] Eigenvalues and colorings of digraphs. *Linear Algebra and Its Applications*, 432(9), 2273–2277, 2010.
- [20] B. Mohar and H. Wu. Dichromatic number and fractional chromatic number. *Forum of Mathematics, Sigma*, 4, e32, 2016.
- [21] V. Neumann-Lara. The dichromatic number of a digraph. *Journal of Combinatorial Theory, Series B*, 33(3), 265–270, 1982.
- [22] A. Scott. Induced trees in graphs of large chromatic number. *Journal of Graph Theory*, 24, 297–311, 1997.
- [23] A. Scott and P. Seymour. Induced subgraphs of graphs with large chromatic number XIII. New brooms. *European Journal of Combinatorics*, 84, 103024, 2020.
- [24] D. P. Sumner. Subtrees of a graph and the chromatic number. *The theory and applications of graphs (Kalamazoo, Mich., 1980)*. Wiley, New York, 557–576, 1981.