

A UNIFYING VIEW TOWARD POLYHEDRAL PRODUCTS THROUGH PANEL STRUCTURES

LI YU

ABSTRACT. A panel structure on a topological space is just a locally finite family of closed subspaces. A space together with a panel structure is called a space with faces. In this paper, we introduce a notion of polyhedral product over a space with faces. This notion provides a unifying viewpoint on the constructions of polyhedral products and generalized moment-angle complexes in various settings. We compute the stable decomposition of these spaces and use it to study their cohomology ring structures. Moreover, we can compute the equivariant cohomology ring of the moment-angle complex over a space with faces with respect to the canonical torus action. The calculation leads to the notion of topological face ring of a space with faces, which generalizes the classical notion of face ring (Stanley-Reisner ring) of a simplicial complex. We will see that many known results in the study of polyhedral products and moment-angle complexes can be reinterpreted from our general theorems on the polyhedral product over a space with faces. Moreover, we can derive some new results via our approach in some settings.

1. Introduction

1.1. Moment-angle complex and polyhedral product.

The construction of moment-angle complex \mathcal{Z}_K over a simplicial complex K first appeared in Davis and Januszkiewicz’s pioneering work [12]. The name “moment-angle complex” was given by Buchstaber and Panov [7] afterwards. The more general definition of polyhedral product determined by a simplicial complex and a sequence of space pairs was introduced by Bahri, Bendersky, Cohen and Gitler [2], which has become the major subject in the homotopy theoretic study in toric topology. The reader is also referred to [2] for more information on the earlier works related to polyhedral products by different authors.

Let K denote an (abstract) *simplicial complex* with m vertices labeled by the set $[m] = \{1, 2, \dots, m\}$. We think of each $(k - 1)$ -simplex σ of K as a subset

Key words and phrases. Polyhedral product, moment-angle complex, panel structure, torus action, topological face ring.

2020 *Mathematics Subject Classification.* 57S12 (Primary), 57S25, 55N91, 13F55.

$\{i_1, \dots, i_k\} \subseteq [m]$ such that if $\tau \subseteq \sigma$, then τ is a simplex of K . In particular, the empty set \emptyset is a simplex of K .

Let $(\mathbb{X}, \mathbb{A}) = \{(X_j, A_j, a_j)\}_{j=1}^m$ be a sequence of based CW-complexes where $a_j \in A_j \subseteq X_j$, $1 \leq j \leq m$. The *polyhedral product* (or *generalized moment-angle complex*) determined by (\mathbb{X}, \mathbb{A}) and K , denoted by $(\mathbb{X}, \mathbb{A})^K$ or $\mathcal{Z}(K; (\mathbb{X}, \mathbb{A}))$, is defined to be

$$(1) \quad (\mathbb{X}, \mathbb{A})^K = \bigcup_{\sigma \in K} (\mathbb{X}, \mathbb{A})^\sigma \subseteq \prod_{j \in [m]} X_j, \text{ where } (\mathbb{X}, \mathbb{A})^\sigma = \prod_{j \in \sigma} X_j \times \prod_{j \in [m] \setminus \sigma} A_j.$$

If $(\mathbb{X}, \mathbb{A}) = \{(X_j, A_j, a_j) = (X, A, a_0)\}_{j=1}^m$, we also denote $(\mathbb{X}, \mathbb{A})^K$ by $(X, A)^K$. In particular, $(D^2, S^1)^K$ and $(D^1, S^0)^K$ are called the *moment-angle complex* and *real moment-angle complex* over K , respectively.

We can extend the construction of $(\mathbb{X}, \mathbb{A})^K$ to polyhedral product over any simplicial poset (or simplicial cell complex) via the colimit construction. Such a generalization has been done by Lü and Panov [17] for moment-angle complexes.

Another generalization is the *moment-angle space* of a convex polytope (not necessarily simple) introduced by Ayzenberg and Buchstaber [1].

More recently, a notion of polyhedral product over a nice manifold with corners was introduced by Yu [24]. The main idea of the construction in [24] suggests that we can define a more general notion of polyhedral product which unifies the constructions of various polyhedral products (up to homotopy equivalences). This is main purpose of this paper.

1.2. Cohomology ring.

The cohomology ring of the moment-angle complex over a simplicial complex K was computed by Franz [13] and Baskakov, Buchstaber and Panov [6]. It is shown in [6] that the cohomology ring of \mathcal{Z}_K with coefficients \mathbf{k} (a commutative ring with unit) is isomorphic to the Tor-algebra of the *face ring* (or *Stanley-Reisner ring*) $\mathbf{k}[K]$ of K using the Koszul resolution of $\mathbf{k}[K]$. Recall $\mathbf{k}[K]$ is the quotient

$$(2) \quad \mathbf{k}[K] = \mathbf{k}[v_1, \dots, v_m] / \mathcal{I}_K$$

where \mathcal{I}_K is the ideal generated by the monomials $v_{i_1} \cdots v_{i_s}$ for which $\{i_1, \dots, i_s\}$ does not span a simplex of K . A linear basis of $\mathbf{k}[K]$ over \mathbf{k} is given by

$$(3) \quad \{1\} \cup \{v_{i_1}^{n_1} \cdots v_{i_s}^{n_s} \mid \emptyset \neq \{i_1, \dots, i_s\} \in K, n_1 > 0, \dots, n_s > 0\}.$$

The cohomology ring of \mathcal{Z}_K was also computed by Wang and Zheng [21] using the simplicial complement of K and the Taylor resolution of $\mathbf{k}[K]$.

For a general polyhedral product $(\mathbb{X}, \mathbb{A})^K$, its cohomology ring structure was computed by Bahri, Bendersky, Cohen and Gitler [3] via partial diagonal maps and by Bahri, Bendersky, Cohen and Gitler [4] by a spectral sequence under

certain freeness conditions (coefficients in a field for example). The study in this direction is further extended by Bahri, Bendersky, Cohen and Gitler [5]. A computation using a different method was carried out by Zheng [25]. In this paper, we will mainly use the strategy in [3] to study the cohomology ring of our polyhedral products.

1.3. Equivariant cohomology ring.

For a topological group G , denote by $G \rightarrow EG \rightarrow BG$ the universal principal G -bundle, with EG a contractible space endowed with a free G -action, and BG the orbit space. We may view G as a subspace of EG (the orbit of a basepoint).

The *Borel construction* of a space X with a G -action is the quotient space

$$EG \times_G X = EG \times X / \sim$$

where $(e, x) \sim (eg, g^{-1}x)$ for any $e \in EG$, $x \in X$ and $g \in G$.

Associated to the Borel construction, there is a canonical fiber bundle

$$X \rightarrow EG \times_G X \rightarrow BG.$$

The equivariant cohomology ring of X , denoted by $H_G^*(X)$ is the (ordinary) cohomology ring of the Borel construction of X .

In addition, if a subspace A of X is invariant under the G -action on X , we call (X, A) a *G -space pair*. Then if in $(\mathbb{X}, \mathbb{A}) = \{(X_j, A_j, a_j)\}_{j=1}^m$, each (X_j, A_j) is a G_j -space pair, the G_j -actions on the pairs (X_j, A_j) extend canonically to an action of the product group $\prod_{j \in [m]} G_j$ on the pair $(\prod_{j \in [m]} X_j, \prod_{j \in [m]} A_j)$. It is easy to check that this action of $\prod_{j \in [m]} G_j$ preserves the subspace $(\mathbb{X}, \mathbb{A})^K \subseteq \prod_{j \in [m]} X_j$ for any abstract simplicial complex K on $[m]$. Then it is meaningful to study the equivariant cohomology ring of $(\mathbb{X}, \mathbb{A})^K$ with respect to the $\prod_{j \in [m]} G_j$ -action.

When $G_1 = \cdots = G_m = S^1$, the equivariant cohomology ring of $(D^2, S^1)^K$ with respect to the canonical $(S^1)^m$ -action is shown to be isomorphic to the face ring $\mathbb{Z}[K]$ of K (see [12] and Buchstaber and Panov [9]). This result was generalized to moment-angle complexes over simplicial posets in [17].

1.4. Panel structure.

A *panel structure* on a topological space Y is just a locally finite family of closed subspaces $\mathcal{P} = \{P_j\}_{j \in \Lambda}$ (see Davis [11, Section 6]). Each subspace P_j is called a *panel*. The space Y together with a panel structure \mathcal{P} is called a *space with faces*, denoted by (Y, \mathcal{P}) .

Suppose $\Lambda = [m] = \{1, \dots, m\}$ is a finite set. For any subset $J \subseteq [m]$, let

$$P_J = \bigcup_{j \in J} P_j, \quad P_\emptyset = \emptyset; \quad P_{\cap J} = \bigcap_{j \in J} P_j, \quad P_{\cap \emptyset} = Y.$$

It is clear that

$$P_{\cap J} \subseteq P_J, \quad P_J \subseteq P_{J'}, \quad P_{\cap J'} \subseteq P_{\cap J}, \quad \forall J \subseteq J' \subseteq [m].$$

Let

$$Y_{c(\mathcal{P})} = Y \setminus P_{[m]} = Y \setminus (P_1 \cup \cdots \cup P_m)$$

We called $Y_{c(\mathcal{P})}$ the *core* of (Y, \mathcal{P}) .

Each path-connected component of $P_{\cap J}$ is called a *face* of (Y, \mathcal{P}) . In particular, each path-connected component of $Y = P_{\cap \emptyset}$ is a face of (Y, \mathcal{P}) .

In addition, let $\mathcal{S}_{(Y, \mathcal{P})}$ denote the *face poset* of (Y, \mathcal{P}) , which is the set of all faces of (Y, \mathcal{P}) ordered by inclusion.

1.5. Polyhedral product over a space with faces.

Let $\mathcal{P} = \{P_j\}_{j \in [m]}$ be a panel structure on a topological space Y . For any face f of Y , the following subset of $[m]$ is called the *panel index* of f .

$$(4) \quad I_f = \{j \in [m] \mid f \subseteq P_j\} \subseteq [m].$$

Let $(\mathbb{X}, \mathbb{A}) = \{(X_j, A_j, a_j)\}_{j=1}^m$ be a sequence of based CW-complexes where $a_j \in A_j \subseteq X_j$, $1 \leq j \leq m$. The *polyhedral product* of (\mathbb{X}, \mathbb{A}) over (Y, \mathcal{P}) is

$$\begin{aligned} (\mathbb{X}, \mathbb{A})^{(Y, \mathcal{P})} &= \bigcup_{f \in \mathcal{S}_{(Y, \mathcal{P})}} (\mathbb{X}, \mathbb{A})^f \subseteq Y \times \prod_{j \in [m]} X_j, \text{ where} \\ (\mathbb{X}, \mathbb{A})^f &= f \times \prod_{j \in I_f} X_j \times \prod_{j \in [m] \setminus I_f} A_j, \quad \forall f \in \mathcal{S}_{(Y, \mathcal{P})}. \end{aligned}$$

If $(\mathbb{X}, \mathbb{A}) = \{(X_j, A_j, a_j) = (X, A, a_0)\}_{j=1}^m$, we also denote $(\mathbb{X}, \mathbb{A})^{(Y, \mathcal{P})}$ by $(X, A)^{(Y, \mathcal{P})}$. In particular, we call $(D^2, S^1)^{(Y, \mathcal{P})}$ and $(D^1, S^0)^{(Y, \mathcal{P})}$ the *moment-angle complex* and *real moment-angle complex* over (Y, \mathcal{P}) , respectively.

Example 1.1. Let $\mathcal{P} = \{P_j\}_{j \in [m]}$ be a panel structure on Y .

- If $P_1 = \cdots = P_m = Y$, then $(\mathbb{X}, \mathbb{A})^{(Y, \mathcal{P})} = Y \times \prod_{j \in [m]} X_j$.
- If $P_1 = \cdots = P_m = \emptyset$, then $(\mathbb{X}, \mathbb{A})^{(Y, \mathcal{P})} = Y \times \prod_{j \in [m]} A_j$.
- Let $\mathcal{P} = \{P_j\}_{j \in [m]}$ and $\mathcal{P}' = \{P'_j\}_{j \in [m]}$ be two panel structures on Y . If $P_j \subset P'_j$ for all $j \in [m]$, we call \mathcal{P} a *refinement* of \mathcal{P}' and write $\mathcal{P} \preccurlyeq \mathcal{P}'$. It is clear that if $\mathcal{P} \preccurlyeq \mathcal{P}'$, then $(\mathbb{X}, \mathbb{A})^{(Y, \mathcal{P})} \subseteq (\mathbb{X}, \mathbb{A})^{(Y, \mathcal{P}'})$.

So we obtain a hierarchy of spaces between $Y \times \prod_{j \in [m]} A_j$ and $Y \times \prod_{j \in [m]} X_j$ given by the polyhedral products of (\mathbb{X}, \mathbb{A}) over all the panel structures on Y .

In categorical languages, let $\mathcal{L}_m(Y)$ be the category of all the panel structures on Y with m panels where the morphisms between two panel structures $\mathcal{P} \preccurlyeq \mathcal{P}'$ are the inclusions of corresponding panels. Then every $(\mathbb{X}, \mathbb{A}) = \{(X_j, A_j, a_j)\}_{j=1}^m$

determines a functor from $\mathcal{L}_m(Y)$ to the category of topological spaces which maps any $(Y, \mathcal{P}) \in \mathcal{L}_m(Y)$ to the polyhedral product $(\mathbb{X}, \mathbb{A})^{(Y, \mathcal{P})}$.

Similarly to the study of moment-angle complexes and polyhedral products over simplicial complexes, we consider the following problems in this paper.

- Compute the stable decomposition of $(\mathbb{X}, \mathbb{A})^{(Y, \mathcal{P})}$.
- Describe the cohomology ring structure of $(\mathbb{X}, \mathbb{A})^{(Y, \mathcal{P})}$.
- Compute the equivariant cohomology of $(D^2, S^1)^{(Y, \mathcal{P})}$ and $(D^1, S^0)^{(Y, \mathcal{P})}$.

We will study these problems in Section 2.1, 2.2 and 2.3, respectively. The philosophy in our study is to treat a CW-complex with a panel structure as a polyhedron where each panel plays the role of a facet (codimension-one face). Therefore, many theorems and their proofs in Section 2.1, 2.2 and 2.3 are parallel to the theorems in [24, Section 4, Section 5]. Because of this, we will only write the statements of these theorems and omit their proofs.

In Section 3, Section 4 and Section 5, we will reinterpret the constructions of polyhedral products over simplicial complexes, simplicial posets and manifold with corners, respectively, from the viewpoint of panel structures on a space; and recover many known results on these spaces from our general theorems of polyhedral products over a space with faces. Moreover, we can derive some new results via our approach in these settings (e.g. Proposition 3.2, Proposition 3.4, Proposition 4.2 and Proposition 4.3). Especially, we will give a new way to write the cohomology ring structure of the ordinary (real) moment-angle complex of a simplicial complex.

Convention: In the rest of the paper, we assume that Y is a CW-complex and every panel of a panel structure \mathcal{P} on Y is a CW-subcomplex of Y . In addition, we assume that the number of faces in (Y, \mathcal{P}) is finite, i.e. $\mathcal{S}_{(Y, \mathcal{P})}$ is a finite poset.

In this paper, the term ‘‘cohomology’’ of a space X , denoted by $H^*(X)$, always mean singular cohomology with integral coefficients if not specified otherwise.

2. Topology of $(\mathbb{X}, \mathbb{A})^{(Y, \mathcal{P})}$

2.1. Stable decomposition of $(\mathbb{X}, \mathbb{A})^{(Y, \mathcal{P})}$.

To do the stable decomposition of $(\mathbb{X}, \mathbb{A})^{(Y, \mathcal{P})}$ as the stable decomposition of a polyhedral product over a simplicial complex in [2], we want to first think of $(\mathbb{X}, \mathbb{A})^{(Y, \mathcal{P})}$ as the colimit of a diagram of CW-complexes over a poset. The following are some basic definitions (see [26]).

- Let CW be the category of CW-complexes and continuous maps.
- Let CW_* be the category of based CW-complexes and based continuous maps.

- A *diagram* \mathcal{D} of CW-complexes or based CW-complexes over a finite poset \mathcal{P} is a functor

$$\mathcal{D} : \mathcal{P} \rightarrow CW \text{ or } CW_*$$

so that for every $p \leq p'$ in \mathcal{P} , there is a map $d_{pp'} : \mathcal{D}(p') \rightarrow \mathcal{D}(p)$ with

$$d_{pp} = id_{\mathcal{D}(p)}, \quad d_{pp'}d_{p'p''} = d_{pp''}, \quad \forall p \leq p' \leq p''.$$

- The *colimit* of \mathcal{D} is the space

$$\text{colim}(\mathcal{D}) = \text{colim}_{p \in \mathcal{P}} \mathcal{D}(p) := \left(\prod_{p \in \mathcal{P}} \mathcal{D}(p) \right) / \sim$$

where \sim denotes the equivalence relation generated by requiring that for each $x \in \mathcal{D}(p')$, $x \sim d_{pp'}(x)$ for every $p < p'$.

Let (Y, \mathcal{P}) be a space with faces where $\mathcal{P} = \{P_j\}_{j \in [m]}$ is a panel structure on Y . We can think of $(\mathbb{X}, \mathbb{A})^{(Y, \mathcal{P})}$ as a colimit of CW-complexes as follows. Let

$$(5) \quad \mathcal{P}_{(Y, \mathcal{P})} := \{(f, L) \mid f \in \mathcal{S}_{(Y, \mathcal{P})}, L \subseteq I_f\}$$

be a poset where the partial order \leq on $\mathcal{P}_{(Y, \mathcal{P})}$ is defined by

$$(f, L) \leq (f', L') \text{ if and only if } f \supseteq f' \text{ and } I_f \setminus L \supseteq I_{f'} \setminus L'.$$

Note that $\mathcal{P}_{(Y, \mathcal{P})}$ is a finite poset since by our convention (Y, \mathcal{P}) only has finitely many faces.

For any $(f, L) \in \mathcal{P}_{(Y, \mathcal{P})}$, let

$$(6) \quad (\mathbb{X}, \mathbb{A})^{(f, L)} = f \times \prod_{j \in I_f \setminus L} X_j \times \prod_{j \in [m] \setminus (I_f \setminus L)} A_j.$$

Clearly, $(\mathbb{X}, \mathbb{A})^{(f, L)} \subseteq (\mathbb{X}, \mathbb{A})^{(f, \emptyset)} = (\mathbb{X}, \mathbb{A})^f$. So

$$(\mathbb{X}, \mathbb{A})^{(Y, \mathcal{P})} = \bigcup_{f \in \mathcal{S}_{(Y, \mathcal{P})}} (\mathbb{X}, \mathbb{A})^f = \bigcup_{(f, L) \in \mathcal{P}_{(Y, \mathcal{P})}} (\mathbb{X}, \mathbb{A})^{(f, L)}.$$

Next, define a diagram of CW-complexes

$$\mathbf{D}_{(\mathbb{X}, \mathbb{A})} : \mathcal{P}_{(Y, \mathcal{P})} \rightarrow CW, \quad \mathbf{D}_{(\mathbb{X}, \mathbb{A})}((f, L)) = (\mathbb{X}, \mathbb{A})^{(f, L)}$$

where $(d_{(\mathbb{X}, \mathbb{A})})_{(f, L), (f', L')} : \mathbf{D}_{(\mathbb{X}, \mathbb{A})}((f', L')) \rightarrow \mathbf{D}_{(\mathbb{X}, \mathbb{A})}((f, L))$ is the natural inclusion for any $(f, L) \leq (f', L') \in \mathcal{P}_{(Y, \mathcal{P})}$.

It is easy to see that $(\mathbb{X}, \mathbb{A})^{(Y, \mathcal{P})}$ is the colimit of $\mathbf{D}_{(\mathbb{X}, \mathbb{A})}$, that is

$$(7) \quad (\mathbb{X}, \mathbb{A})^{(Y, \mathcal{P})} = \text{colim}(\mathbf{D}_{(\mathbb{X}, \mathbb{A})}).$$

Next, we compute the homotopy type of the *suspension* $\Sigma((\mathbb{X}, \mathbb{A})^{(Y, \mathcal{P})})$ of $(\mathbb{X}, \mathbb{A})^{(Y, \mathcal{P})}$ under some special conditions on (\mathbb{X}, \mathbb{A}) . The following two theorems

are generalizations of [24, Theorem 4.4] and [24, Theorem 4.12], respectively. The “ \vee ” and “ \wedge ” are wedge sum and smash product of spaces, respectively.

Theorem 2.1. *Let $(\mathbb{X}, \mathbb{A}) = \{(X_j, A_j, a_j)\}_{j=1}^m$ where each X_j is contractible and each A_j is either connected or is a disjoint union of a connected CW-complex with its basepoint. For any panel structure $\mathcal{P} = \{P_j\}_{j \in [m]}$ on a CW-complex Y , there is a homotopy equivalence*

$$\Sigma((\mathbb{X}, \mathbb{A})^{(Y, \mathcal{P})}) \simeq \bigvee_{J \subseteq [m]} \Sigma\left(Y/P_J \wedge \bigwedge_{j \in J} A_j\right).$$

So the reduced homology group

$$\tilde{H}_*((\mathbb{X}, \mathbb{A})^{(Y, \mathcal{P})}) \cong \bigoplus_{J \subseteq [m]} \tilde{H}_*(Y/P_J \wedge \bigwedge_{j \in J} A_j).$$

Proof. The proof is completely parallel to the proof of [24, Theorem 4.4]. \square

Theorem 2.2. *Let $(\mathbb{X}, \mathbb{A}) = \{(X_j, A_j, a_j)\}_{j=1}^m$ where each A_j is contractible and each X_j is either connected or is a disjoint union of a connected CW-complex with its basepoint. For any panel structure $\mathcal{P} = \{P_j\}_{j \in [m]}$ on a CW-complex Y , there is a homotopy equivalence*

$$S^1 \vee \Sigma((\mathbb{X}, \mathbb{A})^{(Y, \mathcal{P})}) \simeq \bigvee_{J \subseteq [m]} \Sigma\left((P_{\cap J} \cup y_0) \wedge \bigwedge_{j \in J} X_j\right).$$

where y_0 is a point disjoint from Y . This implies

$$H_*((\mathbb{X}, \mathbb{A})^{(Y, \mathcal{P})}) \cong \bigoplus_{J \subseteq [m]} \tilde{H}_*\left((P_{\cap J} \cup y_0) \wedge \bigwedge_{j \in J} X_j\right).$$

Proof. The proof is parallel to the proof of [24, Theorem 4.12]. \square

2.2. Cohomology ring of $(\mathbb{X}, \mathbb{A})^{(Y, \mathcal{P})}$.

It was shown in [3] that the cohomology ring structure of a polyhedral product over a simplicial complex can be computed through the stable decomposition and partial diagonal maps of the polyhedral product. For a panel structure \mathcal{P} on a finite CW-complex Y , since we also have the stable decomposition of $(\mathbb{X}, \mathbb{A})^{(Y, \mathcal{P})}$, we can describe the cohomology ring of $(\mathbb{X}, \mathbb{A})^{(Y, \mathcal{P})}$ in a similar way.

Under some special conditions on (\mathbb{X}, \mathbb{A}) (e.g. the conditions in Theorem 2.1 and Theorem 2.2), we can write the cohomology ring structure of $(\mathbb{X}, \mathbb{A})^{(Y, \mathcal{P})}$ explicitly. For example, let

$$(\mathbb{D}, \mathbb{S}) = \{(D^{n_j+1}, S^{n_j}, a_j)\}_{j=1}^m$$

where D^{n+1} is the unit ball in \mathbb{R}^{n+1} and $S^n = \partial D^{n+1}$. We can describe the cohomology ring structure of $(\mathbb{D}, \mathbb{S})^{(Y, \mathcal{P})}$ by the following ring. Let

$$(8) \quad \mathcal{R}_{Y, \mathcal{P}}^* := \bigoplus_{J \subseteq [m]} H^*(Y, P_J)$$

Define a graded ring structure $\mathfrak{U}^{(\mathbb{D}, \mathbb{S})}$ on $\mathcal{R}_{Y, \mathcal{P}}^*$ as follows.

- If $J \cap J' = \emptyset$ or $J \cap J' \neq \emptyset$ but $n_j = 0$ for all $j \in J \cap J'$,
 $H^*(Y, P_J) \otimes H^*(Y, P_{J'}) \xrightarrow{\mathfrak{U}^{(\mathbb{D}, \mathbb{S})}} H^*(Y, P_{J \cup J'})$ is the relative cup product \cup .
- If $J \cap J' \neq \emptyset$ and there exists $n_j \geq 1$ for some $j \in J \cap J'$,
 $H^*(Y, P_J) \otimes H^*(Y, P_{J'}) \xrightarrow{\mathfrak{U}^{(\mathbb{D}, \mathbb{S})}} H^*(Y, P_{J \cup J'})$ is trivial.

Let

$$\tilde{\mathcal{R}}_{Y, \mathcal{P}}^* := \bigoplus_{J \subseteq [m]} \tilde{H}^*(Y/P_J), \text{ then } \mathcal{R}_{Y, \mathcal{P}}^* = \tilde{\mathcal{R}}_{Y, \mathcal{P}}^* \oplus \mathbb{Z}.$$

By [24, Lemma 3.2], there is a natural product on $\tilde{\mathcal{R}}_{Y, \mathcal{P}}^*$, denoted by $\tilde{\mathfrak{U}}^{(\mathbb{D}, \mathbb{S})}$, that is induced from the product $\mathfrak{U}^{(\mathbb{D}, \mathbb{S})}$ on $\mathcal{R}_{Y, \mathcal{P}}^*$. So we have a commutative diagram

$$(9) \quad \begin{array}{ccc} H^*(Y, P_J) \otimes H^*(Y, P_{J'}) & \xrightarrow{\mathfrak{U}^{(\mathbb{D}, \mathbb{S})}} & H^*(Y, P_{J \cup J'}) \\ \downarrow & & \downarrow \\ \tilde{H}^*(Y/P_J) \otimes \tilde{H}^*(Y/P_{J'}) & \xrightarrow{\tilde{\mathfrak{U}}^{(\mathbb{D}, \mathbb{S})}} & \tilde{H}^*(Y/P_{J \cup J'}) \end{array}$$

- If $J \cap J' = \emptyset$ or $J \cap J' \neq \emptyset$ but $n_j = 0$ for all $j \in J \cap J'$, then
 $\tilde{H}^*(Y/P_J) \otimes \tilde{H}^*(Y/P_{J'}) \xrightarrow{\tilde{\mathfrak{U}}^{(\mathbb{D}, \mathbb{S})}} \tilde{H}^*(Y/P_{J \cup J'})$ is the product $\tilde{\cup}$ induced from the relative cup product $H^*(Y, P_J) \otimes H^*(Y, P_{J'}) \xrightarrow{\cup} H^*(Y, P_{J \cup J'})$.
- If $J \cap J' \neq \emptyset$ and there exists $n_j \geq 1$ for some $j \in J \cap J'$, then
 $\tilde{H}^*(Y/P_J) \otimes \tilde{H}^*(Y/P_{J'}) \xrightarrow{\tilde{\mathfrak{U}}^{(\mathbb{D}, \mathbb{S})}} \tilde{H}^*(Y/P_{J \cup J'})$ is trivial.

It is clear that $(\mathcal{R}_{Y, \mathcal{P}}^*, \mathfrak{U}^{(\mathbb{D}, \mathbb{S})})$ and $(\tilde{\mathcal{R}}_{Y, \mathcal{P}}^*, \tilde{\mathfrak{U}}^{(\mathbb{D}, \mathbb{S})})$ determine each other.

Theorem 2.3. *For any panel structure $\mathcal{P} = \{P_j\}_{j \in [m]}$ on a CW-complex Y , there is a ring isomorphism (up to a sign) from $(\mathcal{R}_{Y, \mathcal{P}}^*, \mathfrak{U}^{(\mathbb{D}, \mathbb{S})})$ to the integral cohomology ring of $(\mathbb{D}, \mathbb{S})^{(Y, \mathcal{P})}$. Moreover, we can make the ring isomorphism degree-preserving by shifting the degrees of all the elements in $H^*(Y, P_J)$ up by $\sum_{j \in J} n_j$ for every $J \subseteq [m]$.*

Proof. The proof is parallel to the proof of [24, Theorem 4.8 (b)]. \square

By Theorem 2.3, we have the following results for $(D^2, S^1)^{(Y, \mathcal{P})}$ and $(D^1, S^0)^{(Y, \mathcal{P})}$.

Corollary 2.4. *For any panel structure $\mathcal{P} = \{P_j\}_{j \in [m]}$ on a finite CW-complex Y , there is a ring isomorphism (up to a sign) from $(\mathcal{R}_{Y, \mathcal{P}}^*, \mathbb{U})$ to the integral cohomology ring of $(D^2, S^1)^{(Y, \mathcal{P})}$, where the \mathbb{U} on $\mathcal{R}_{Y, \mathcal{P}}^*$ is defined by:*

- *If $J \cap J' \neq \emptyset$, $H^*(Y, P_J) \otimes H^*(Y, P_{J'}) \xrightarrow{\mathbb{U}} H^*(Y, P_{J \cup J'})$ is trivial.*
- *If $J \cap J' = \emptyset$, $H^*(Y, P_J) \otimes H^*(Y, P_{J'}) \xrightarrow{\mathbb{U}} H^*(Y, P_{J \cup J'})$ is the relative cup product \cup .*

Moreover, we can make this ring isomorphism degree-preserving by shifting the degrees of all the elements in $H^*(Y, P_J)$ up by $|J|$ for every $J \subseteq [m]$.

Corollary 2.5. *For any panel structure $\mathcal{P} = \{P_j\}_{j \in [m]}$ on a finite CW-complex Y , there is a graded ring isomorphism from $(\mathcal{R}_{Y, \mathcal{P}}^*, \cup)$ to the integral cohomology ring of $(D^1, S^0)^{(Y, \mathcal{P})}$, where the \cup on $\mathcal{R}_{Y, \mathcal{P}}^*$ is the relative cup product*

$$H^*(Y, P_J) \otimes H^*(Y, P_{J'}) \xrightarrow{\cup} H^*(Y, P_{J \cup J'}), \quad \forall J, J' \subseteq [m].$$

Note that here we have an authentic graded ring isomorphism (without the sign factors) from $(\mathcal{R}_{Y, \mathcal{P}}^*, \cup)$ to the integral cohomology ring of $(D^1, S^0)^{(Y, \mathcal{P})}$ (see the remark of [24, Corollary 4.10]).

2.3. Equivariant Cohomology of $(D^2, S^1)^{(Y, \mathcal{P})}$ and $(D^1, S^0)^{(Y, \mathcal{P})}$.

Let Y be a finite CW-complex with a panel structure $\mathcal{P} = \{P_j\}_{j \in [m]}$. Let

$$D^2 = \{z \in \mathbb{C} \mid \|z\| \leq 1\}, \quad S^1 = \partial D^2.$$

The natural action of S^1 on (D^2, S^1) induces a canonical action of $T^m = (S^1)^m$ on the polyhedral product $(D^2, S^1)^{(Y, \mathcal{P})}$.

The equivariant cohomology of $(D^2, S^1)^{(Y, \mathcal{P})}$, denoted by $H_{T^m}^*((D^2, S^1)^{(Y, \mathcal{P})})$, is the cohomology of the *Borel construction*

$$ET^m \times_{T^m} (D^2, S^1)^{(Y, \mathcal{P})} = ET^m \times (D^2, S^1)^{(Y, \mathcal{P})} / \sim$$

where $(e, x) \sim (eg, g^{-1}x)$ for any $e \in ET^m$, $x \in (D^2, S^1)^{(Y, \mathcal{P})}$ and $g \in T^m$. Here

$$ET^m = (ES^1)^m = (S^\infty)^m.$$

Associated to the Borel construction, there is a canonical fiber bundle

$$(10) \quad (D^2, S^1)^{(Y, \mathcal{P})} \rightarrow ET^m \times_{T^m} (D^2, S^1)^{(Y, \mathcal{P})} \rightarrow BT^m$$

where $BT^m = (BS^1)^m = (S^\infty/S^1)^m = (\mathbb{C}P^\infty)^m$ is the *classifying space* of T^m .

By the colimit construction of $(D^2, S^1)^{(Y, \mathcal{P})}$ in (7), the Borel construction

$$\begin{aligned} ET^m \times_{T^m} (D^2, S^1)^{(Y, \mathcal{P})} &= \bigcup_{(f, L) \in \mathcal{P}(Y, \mathcal{P})} ET^m \times_{T^m} (D^2, S^1)^{(f, L)} \\ &= \bigcup_{(f, L) \in \mathcal{P}(Y, \mathcal{P})} (S^\infty \times_{S^1} D^2, S^\infty \times_{S^1} S^1)^{(f, L)} \\ &= (S^\infty \times_{S^1} D^2, S^\infty \times_{S^1} S^1)^{(Y, \mathcal{P})}. \end{aligned}$$

Then by the homotopy equivalence of the pairs

$$(S^\infty \times_{S^1} D^2, S^\infty \times_{S^1} S^1) \rightarrow (\mathbb{C}P^\infty, *),$$

we can derive from [2, Corollary 4.5] that there exists a homotopy equivalence

$$(S^\infty \times_{S^1} D^2, S^\infty \times_{S^1} S^1)^{(Y, \mathcal{P})} \simeq (\mathbb{C}P^\infty, *)^{(Y, \mathcal{P})}.$$

We call $(\mathbb{C}P^\infty, *)^{(Y, \mathcal{P})}$ the *Davis-Januszkiewicz space* of (Y, \mathcal{P}) , denoted by $\mathcal{DJ}(Y, \mathcal{P})$. So the equivariant cohomology ring of $(D^2, S^1)^{(Y, \mathcal{P})}$ with respect to the canonical T^m -action is isomorphic to the ordinary cohomology ring of $\mathcal{DJ}(Y, \mathcal{P})$. Similarly, we can prove that the equivariant cohomology ring of $(D^1, S^0)^{(Y, \mathcal{P})}$ with respect to the canonical $(\mathbb{Z}_2)^m$ -action is isomorphic to the ordinary cohomology ring of $(\mathbb{R}P^\infty, *)^{(Y, \mathcal{P})}$.

To describe the cohomology ring structures of $(\mathbb{C}P^\infty, *)^{(Y, \mathcal{P})}$ and $(\mathbb{R}P^\infty, *)^{(Y, \mathcal{P})}$, we introduce the following definition which generalizes the classical notion of face ring (Stanley-Reisner ring) of a simplicial complex; see Lemma 3.4 for an explanation.

Let \mathbf{k} denote a commutative ring with unit. For any $J \subseteq [m]$, let $R_{\mathbf{k}}^J$ be the subring of the polynomial ring $\mathbf{k}[x_1, \dots, x_m]$ defined by

$$R_{\mathbf{k}}^J := \begin{cases} \text{span}_{\mathbf{k}}\{x_{j_1}^{n_1} \cdots x_{j_s}^{n_s} \mid n_1 > 0, \dots, n_s > 0\}, & \text{if } J = \{j_1, \dots, j_s\} \neq \emptyset; \\ \mathbf{k}, & \text{if } J = \emptyset. \end{cases}$$

We can multiply $f(x) \in R_{\mathbf{k}}^J$ and $f'(x) \in R_{\mathbf{k}}^{J'}$ in $\mathbf{k}[x_1, \dots, x_m]$ and obtain an element $f(x)f'(x) \in R_{\mathbf{k}}^{J \cup J'}$.

Definition 2.6 (Topological Face Ring of a Space with Faces). Suppose Y is a CW-complex with a panel structure $\mathcal{P} = \{P_j\}_{j \in [m]}$. For any coefficients ring \mathbf{k} , the *topological face ring of (Y, \mathcal{P})* over \mathbf{k} is defined to be

$$\mathbf{k}(Y, \mathcal{P}) := \bigoplus_{J \subseteq [m]} H^*(P_{\cap J}; \mathbf{k}) \otimes R_{\mathbf{k}}^J.$$

The product \star on $\mathbf{k}(Y, \mathcal{P})$ is defined by: for any $J, J' \subseteq [m]$,

$$\left(H^*(P_{\cap J}; \mathbf{k}) \otimes R_{\mathbf{k}}^J \right) \otimes \left(H^*(P_{\cap J'}; \mathbf{k}) \otimes R_{\mathbf{k}}^{J'} \right) \xrightarrow{\star} \left(H^*(P_{\cap (J \cup J')}; \mathbf{k}) \otimes R_{\mathbf{k}}^{J \cup J'} \right)$$

for any $\phi \in H^*(P_{\cap J}; \mathbf{k})$, $\phi' \in H^*(P_{\cap J'}; \mathbf{k})$ and $f(x) \in R_{\mathbf{k}}^J$, $f'(x) \in R_{\mathbf{k}}^{J'}$,

$$(11) \quad (\phi \otimes f(x)) \star (\phi' \otimes f'(x)) := (\kappa_{J \cup J', J}^*(\phi) \cup \kappa_{J \cup J', J'}^*(\phi')) \otimes f(x)f'(x)$$

where $\kappa_{I', I} : P_{\cap I'} \rightarrow P_{\cap I}$ is the inclusion map for any subsets $I \subseteq I' \subseteq [m]$ and $\kappa_{I', I}^* : H^*(P_{\cap I}; \mathbf{k}) \rightarrow H^*(P_{\cap I'}; \mathbf{k})$ is the induced homomorphism on cohomology.

In addition, we can consider $\mathbf{k}(Y, \mathcal{P})$ as a graded ring if we choose a degree for every indeterminate x_j in $\mathbf{k}[x_1, \dots, x_m]$ and define

$$\deg(\phi \otimes (x_{j_1}^{n_1} \cdots x_{j_s}^{n_s})) = \deg(\phi) + n_1 \deg(x_{j_1}) + \cdots + n_s \deg(x_{j_s}).$$

We have the following theorem which generalizes [24, Theorem 1.7].

Theorem 2.7. *Suppose Y is a finite CW-complex and $\mathcal{P} = \{P_j\}_{j \in [m]}$ is a panel structure on Y . Then the equivariant cohomology ring of $(D^2, S^1)^{(Y, \mathcal{P})}$ (or $(D^1, S^0)^{(Y, \mathcal{P})}$) with \mathbb{Z} -coefficients (or \mathbb{Z}_2 -coefficients) with respect to the canonical $(S^1)^m$ -action (or $(\mathbb{Z}_2)^m$ -action) is isomorphic as a graded ring to the topological face ring $\mathbb{Z}(Y, \mathcal{P})$ (or $\mathbb{Z}_2(Y, \mathcal{P})$) of (Y, \mathcal{P}) by choosing $\deg(x_j) = 2$ (or $\deg(x_j) = 1$) for all $1 \leq j \leq m$.*

Proof. The proof is parallel to the proof of [24, Theorem 1.7]. \square

From the canonical fiber bundle associated to the Borel construction in (10), we have a natural $H^*(BT^m)$ -module structure on $H_{T^m}^*((D^2, S^1)^{(Y, \mathcal{P})})$. By the identification $H^*(BT^m) = \mathbb{Z}[x_1, \dots, x_m]$, we can write the $H^*(BT^m)$ -module structure on $H_{T^m}^*((D^2, S^1)^{(Y, \mathcal{P})})$ as: for each $1 \leq i \leq m$,

$$(12) \quad x_i \cdot (\phi \otimes f(x)) = (1 \otimes x_i) \star (\phi \otimes f(x)) \stackrel{(11)}{=} \kappa_{J \cup \{i\}, J}^*(\phi) \otimes x_i f(x)$$

where $\phi \in H^*(P_{\cap J})$ and $f(x) \in R_{\mathbb{Z}}^J$, $J \subseteq [m]$.

3. Reexamining moment-angle complexes and polyhedral products over simplicial complexes

In this section, we reexamine the constructions of (generalized) moment-angle complexes and polyhedral products over simplicial complexes from the viewpoint of panel structures. We will reinterpret some known results using our theorems in the last three sections and obtain a few new results.

Let K be a simplicial complex on $[m] = \{1, 2, \dots, m\}$. We assume that the set $[m]$ is *minimal* for K in the sense that every $j \in [m]$ belongs to at least one simplex of K . We also use v_1, \dots, v_m to refer to the vertices (0-dimensional simplex) of K . We do not distinguish K and its geometrical realization below.

- For any $J \subseteq [m]$, let K_J denote the *full subcomplex* of K consisting of all simplices of K which have all of their vertices in J , i.e.

$$K_J = \{\sigma \cap J \mid \sigma \in K\}.$$




FIGURE 1. The cone of the barycentric subdivision of K

- Let K' denote the barycentric subdivision of K . We can consider K' as the set of chains of simplices in K ordered by inclusions. For each nonempty simplex $\sigma \in K$, let F_σ denote the geometrical realization of the poset

$$K_{\geq \sigma} = \{\tau \in K \mid \sigma \subseteq \tau\}.$$

Thus, F_σ is the subcomplex of K' consisting of all simplices of the form

$$\sigma = \sigma_0 \subsetneq \sigma_1 \subsetneq \cdots \subsetneq \sigma_l.$$

It is clear that F_σ is contractible.

- Let $Y^K = u_0 K'$ denote the cone on K' over a point u_0 (see Figure 1). Notice that $Y_K = F_\emptyset$.
- There is a natural panel structure \mathcal{P} on Y^K defined by:

$$\mathcal{P} = \{P_j = F_{v_j}\}_{j \in [m]}.$$

This panel structure \mathcal{P} on Y^K has been studied in [11, p.314] and [12, p.428]. Then by definition, for any $J \subseteq [m]$,

- If $J \neq \emptyset$, we have

$$P_{\cap J} = \bigcap_{j \in J} F_{v_j} = \begin{cases} F_\sigma, & \text{if } J \text{ is the vertex set of a simplex } \sigma \in K; \\ \emptyset, & \text{otherwise.} \end{cases}$$

If $J = \emptyset$, $P_{\cap \emptyset} = Y^K = F_\emptyset$. So all the faces of (Y^K, \mathcal{P}) are $\{F_\sigma \mid \sigma \in K\}$. The space Y^K together with its faces is called a *simple polyhedral complex* in [12, p.428].

- For any $\sigma \in K$, the panel index of the face F_σ in (Y^K, \mathcal{P}) is exactly $\sigma \subseteq [m]$, i.e. $I_{F_\sigma} = \sigma$.
- P_j is homotopy equivalent to the full subcomplex K_j of K . This is because every P_j and any of their intersections $P_{\cap J}$ are contractible (if not empty),

so K_J can be thought of as the nerve complex of P_J covered by $\{P_j\}_{j \in J}$ (see [14, Corollary 4G.3]). Recall that the *nerve complex* of a cover of a set X by a family of subsets $\{U_i\}_{1 \leq i \leq n}$ is a simplicial complex with n vertices whose $(k-1)$ -dimensional simplices correspond to the nonempty intersections of k different U_i in the cover.

Lemma 3.1. *For any $(\mathbb{X}, \mathbb{A}) = \{(X_j, A_j, a_j)\}_{j=1}^m$, there is a homotopy equivalence*

$$(\mathbb{X}, \mathbb{A})^{(Y^K, \mathcal{P})} \simeq (\mathbb{X}, \mathbb{A})^K.$$

Proof. Since the panel index of a face F_σ in (Y^K, \mathcal{P}) is exactly $\sigma \subseteq [m]$, the poset $\mathcal{P}_{(Y^K, \mathcal{P})}$ (see (5)) determined by (Y^K, \mathcal{P}) is isomorphic to the following poset

$$\mathcal{P}_K := \{(\sigma, \tau) \mid \tau \subseteq \sigma \in K\}$$

where $(\sigma, \tau) \leq (\sigma', \tau')$ if and only if $\sigma \subseteq \sigma'$ and $\sigma \setminus \tau \supseteq \sigma' \setminus \tau'$. Let

$$(\mathbb{X}, \mathbb{A})^{(\sigma, \tau)} := \prod_{j \in \sigma \setminus \tau} X_j \times \prod_{j \in [m] \setminus (\sigma \setminus \tau)} A_j, \quad (\mathbb{X}, \mathbb{A})^{(F_\sigma, \tau)} := F_\sigma \times \prod_{j \in \sigma \setminus \tau} X_j \times \prod_{j \in [m] \setminus (\sigma \setminus \tau)} A_j.$$

Then we can think of both $(\mathbb{X}, \mathbb{A})^K$ and $(\mathbb{X}, \mathbb{A})^{(Y^K, \mathcal{P})}$ as colimits of diagrams over \mathcal{P}_K :

$$\begin{aligned} (\mathbb{X}, \mathbb{A})^K &= \bigcup_{\sigma \in K} \prod_{j \in \sigma} X_j \times \prod_{j \in [m] \setminus \sigma} A_j = \operatorname{colim}_{(\sigma, \tau) \in \mathcal{P}_K} (\mathbb{X}, \mathbb{A})^{(\sigma, \tau)}, \\ (\mathbb{X}, \mathbb{A})^{(Y^K, \mathcal{P})} &= \bigcup_{\sigma \in K} F_\sigma \times \prod_{j \in \sigma} X_j \times \prod_{j \in [m] \setminus \sigma} A_j = \operatorname{colim}_{(\sigma, \tau) \in \mathcal{P}_K} (\mathbb{X}, \mathbb{A})^{(F_\sigma, \tau)}. \end{aligned}$$

Moreover, there is an obvious map of diagrams over \mathcal{P}_K defined by:

$$(\mathbb{X}, \mathbb{A})^{(F_\sigma, \tau)} \longrightarrow (\mathbb{X}, \mathbb{A})^{(\sigma, \tau)}, \quad (\sigma, \tau) \in \mathcal{P}_K.$$

Since F_σ is contractible for all $\sigma \in K$ (including $\sigma = \emptyset$), the space $(\mathbb{X}, \mathbb{A})^{(Y^K, \mathcal{P})}$ is homotopy equivalent to $(\mathbb{X}, \mathbb{A})^K$ by [2, Corollary 4.5]. \square

For (D^2, S^1) (or (D^1, S^0)), we can see that the homotopy equivalence in the above proof from $(D^2, S^1)^{(Y^K, \mathcal{P})}$ (or $(D^1, S^0)^{(Y^K, \mathcal{P})}$) to $(D^2, S^1)^K$ (or $(D^1, S^0)^K$) is equivariant with respect to the canonical $(S^1)^m$ -actions (or $(\mathbb{Z}_2)^m$ -actions).

So by Lemma 3.1, we can compute the stable decomposition, cohomology ring and equivariant cohomology ring of $(\mathbb{X}, \mathbb{A})^K$ in terms of $(\mathbb{X}, \mathbb{A})^{(Y^K, \mathcal{P})}$. Next, we explain how to derive some known results of $(\mathbb{X}, \mathbb{A})^K$ by applying our theorems to $(\mathbb{X}, \mathbb{A})^{(Y^K, \mathcal{P})}$.

3.1. $(\mathbb{X}, \mathbb{A})^K$ where every X_j in $(\mathbb{X}, \mathbb{A}) = \{(X_j, A_j, a_j)\}_{j=1}^m$ is contractible.

We can derive a homotopy equivalence from Theorem 2.1 for $(\mathbb{X}, \mathbb{A})^{(Y^K, \mathcal{P})}$:

$$\Sigma((\mathbb{X}, \mathbb{A})^{(Y^K, \mathcal{P})}) \simeq \bigvee_{J \subseteq [m]} \Sigma\left((Y^K/P_J) \wedge \bigwedge_{j \in J} A_j\right).$$

Since P_J is homotopy equivalent to K_J ,

$$(13) \quad Y^K/P_J = u_0 K/P_J \simeq u_0 P_J/P_J \simeq \Sigma(P_J) \simeq \Sigma(K_J).$$

In the following if $J \subseteq [m]$ is the vertex set of a simplex in K , we simply write $J \in K$ by abuse of notation; otherwise we write $J \notin K$.

If J is the vertex set of a simplex $\sigma \in K$, then $K_J = \sigma$, which implies that Y^K/P_J is contractible by (13). Then since the smash product of a contractible space with any space is contractible, we obtain

$$(14) \quad \begin{aligned} \Sigma((\mathbb{X}, \mathbb{A})^{(Y^K, \mathcal{P})}) &\simeq \bigvee_{J \notin K, J \subseteq [m]} \Sigma\left(\Sigma(K_J) \wedge \bigwedge_{j \in J} A_j\right) \\ &\simeq \bigvee_{J \notin K, J \subseteq [m]} \Sigma(K_J) * \bigwedge_{j \in J} A_j \simeq \bigvee_{J \notin K, J \subseteq [m]} \Sigma\left(K_J * \bigwedge_{j \in J} A_j\right) \end{aligned}$$

where $*$ denotes the join of two spaces. The homotopy equivalences in (14) are due to the standard fact that for any based spaces X and Y ,

$$\Sigma(X) \wedge Y \simeq \Sigma(X \wedge Y) \simeq X * Y.$$

Note that the stable decomposition in (14) recovers [2, Theorem 2.21]. In particular, for $(D^2, S^1)^K$ we recover the following result in [2, Corollary 2.23]:

$$(15) \quad \Sigma((D^2, S^1)^K) \simeq \Sigma((D^2, S^1)^{(Y^K, \mathcal{P})}) \simeq \bigvee_{J \notin K, J \subseteq [m]} \Sigma^{|J|+2}(K_J).$$

Moreover, for $(D^1, S^0)^K$, we obtain the following new result.

Proposition 3.2. $\Sigma((D^1, S^0)^K) \simeq \Sigma((D^1, S^0)^{(Y^K, \mathcal{P})}) \simeq \bigvee_{J \notin K, J \subseteq [m]} \Sigma^2(K_J).$

3.2. $(\mathbb{X}, \mathbb{A})^K$ where every A_j in $(\mathbb{X}, \mathbb{A}) = \{(X_j, A_j, a_j)\}_{j=1}^m$ is contractible.

We can derive a homotopy equivalence from Theorem 2.2 for $(\mathbb{X}, \mathbb{A})^{(Y^K, \mathcal{P})}$:

$$S^1 \vee \Sigma((\mathbb{X}, \mathbb{A})^{(Y^K, \mathcal{P})}) \simeq \bigvee_{J \subseteq [m]} \Sigma\left((P_{\cap J} \cup y_0) \wedge \bigwedge_{j \in J} X_j\right).$$

- If $J \neq \emptyset$ and $J \notin K$, then $P_{\cap J}$ is empty, which implies that the space $(P_{\cap J} \cup y_0) \wedge \bigwedge_{j \in J} X_j$ is contractible.
- If $J \neq \emptyset$ and $J \in K$ (suppose J is the vertex set of σ), then $P_{\cap J} = F_\sigma$ is contractible, which implies that $(P_{\cap J} \cup y_0) \wedge \bigwedge_{j \in J} X_j \simeq \bigwedge_{j \in J} X_j$.

- If $J = \emptyset$, then $(P_{\cap J} \cup y_0) \wedge \bigwedge_{j \in J} X_j = Y^K \cup y_0$.

From the above discussion, we obtain

$$S^1 \vee \Sigma((\mathbb{X}, \mathbb{A})^{(Y^K, \mathcal{P})}) \simeq \Sigma(Y^K \cup y_0) \vee \bigvee_{\emptyset \neq J \in K} \Sigma\left(\bigwedge_{j \in J} X_j\right).$$

Then since $Y^K = u_0 K'$ is contractible, $\Sigma(Y^K \cup y_0) \simeq S^1$ implies

$$(16) \quad \Sigma((\mathbb{X}, \mathbb{A})^{(Y^K, \mathcal{P})}) \simeq \bigvee_{\emptyset \neq J \in K} \Sigma\left(\bigwedge_{j \in J} X_j\right).$$

Note that the stable decomposition in (16) recovers [2, Theorem 2.15].

3.3. Cohomology rings of $(D^2, S^1)^K$ and $(D^1, S^0)^K$.

The cohomology rings of $(D^2, S^1)^K$ and $(D^2, S^1)^{(Y^K, \mathcal{P})}$ are isomorphic since they are homotopy equivalent. So by Corollary 2.4, we obtain the following proposition.

Proposition 3.3. *There is a ring isomorphism (up to a sign) from $(\mathcal{R}_{Y^K, \mathcal{P}}^*, \mathbb{U})$ to the integral cohomology ring of $(D^2, S^1)^K$. Moreover, we can make this isomorphism degree-preserving by shifting the degrees of all the elements in $H^*(Y^K, P_J)$ up by $|J|$ for every $J \subseteq [m]$.*

Since Y^K is contractible, for any subset $J \subseteq [m]$, we have an isomorphism

$$\Xi_J : H^q(Y^K, P_J) \xrightarrow{\cong} \tilde{H}^{q-1}(P_J) \xrightarrow{\cong} \tilde{H}^{q-1}(K_J), \quad \forall q \in \mathbb{Z}.$$

So for any subsets $J, J' \subseteq [m]$, we have a commutative diagram

$$(17) \quad \begin{array}{ccc} H^q(Y^K, P_J) \otimes H^{q'}(Y^K, P_{J'}) & \xrightarrow{\mathbb{U}} & H^{q+q'}(Y^K, P_{J \cup J'}) \\ \cong \downarrow & & \downarrow \cong \\ \tilde{H}^{q-1}(P_J) \otimes \tilde{H}^{q'-1}(P_{J'}) & \longrightarrow & \tilde{H}^{q+q'-1}(P_{J \cup J'}) \\ \cong \downarrow & & \downarrow \cong \\ \tilde{H}^{q-1}(K_J) \otimes \tilde{H}^{q'-1}(K_{J'}) & \xrightarrow{\sqcup} & \tilde{H}^{q+q'-1}(K_{J \cup J'}) \end{array}$$

where \sqcup is induced by \mathbb{U} through the above isomorphisms Ξ_J . More precisely, for any $\phi \in \tilde{H}^*(K_J)$, $\phi' \in \tilde{H}^*(K_{J'})$, we have (see Corollary 2.4)

$$\phi \sqcup \phi' = \begin{cases} \Xi_{J \cup J'}(\Xi_J^{-1}(\phi) \cup \Xi_{J'}^{-1}(\phi')), & \text{if } J \cap J' = \emptyset; \\ 0, & \text{if } J \cap J' \neq \emptyset. \end{cases}$$

From the definitions of the isomorphism Ξ_J and the relative cup product \cup , we can check that \sqcup is equivalent to the cohomology product of $(D^2, S^1)^K$ given in [9, Proposition 3.2.10].

Similarly, by Corollary 2.5 we obtain a parallel result for $(D^1, S^0)^K$ which gives a new description of integral cohomology ring of $(D^1, S^0)^K$.

Proposition 3.4. *The integral cohomology ring of the real moment-angle complex $(D^1, S^0)^K$ is isomorphic to $(\mathcal{R}_{Y^K, \mathcal{P}}^*, \cup)$ where $\mathcal{R}_{Y^K, \mathcal{P}}^* = \bigoplus_{J \subseteq [m]} H^*(Y^K, P_J)$ and $H^*(Y^K, P_J) \otimes H^*(Y^K, P_{J'}) \xrightarrow{\cup} H^*(Y^K, P_{J \cup J'})$ is the relative cup product for all $J, J' \subseteq [m]$.*

Remark 3.5. An earlier description of the integral cohomology ring of $(D^1, S^0)^K$ was given by Cai [10] in a very different way.

3.4. The equivariant cohomology rings of $(D^2, S^1)^K$ and $(D^1, S^0)^K$.

Since $(D^2, S^1)^K$ and $(D^2, S^1)^{(Y^K, \mathcal{P})}$ are equivariantly homotopy equivalent, their equivariant cohomology rings are isomorphic. According to Theorem 2.7, the equivariant cohomology ring of $(D^2, S^1)^K$ is isomorphic to the topological face ring of (Y^K, \mathcal{P}) :

$$\mathbb{Z}(Y^K, \mathcal{P}) = \bigoplus_{J \subseteq [m]} H^*(P_{\cap J}) \otimes R_{\mathbb{Z}}^J,$$

where the product \star on $\mathbb{Z}(Y^K, \mathcal{P})$ is defined by (see (11)): for any $J, J' \subseteq [m]$,

$$\begin{aligned} & \left(H^*(P_{\cap J}) \otimes R_{\mathbb{Z}}^J \right) \otimes \left(H^*(P_{\cap J'}) \otimes R_{\mathbb{Z}}^{J'} \right) \xrightarrow{\star} \left(H^*(P_{\cap (J \cup J')}) \otimes R_{\mathbb{Z}}^{J \cup J'} \right) \\ & (\phi \otimes f(x)) \star (\phi' \otimes f'(x)) := (\kappa_{J \cup J', J}^*(\phi) \cup \kappa_{J \cup J', J'}^*(\phi')) \otimes f(x)f'(x) \end{aligned}$$

where $\phi \in H^*(P_{\cap J})$, $\phi' \in H^*(P_{\cap J'})$ and $f(x) \in R_{\mathbb{Z}}^J$, $f'(x) \in R_{\mathbb{Z}}^{J'}$.

Lemma 3.6. $\mathbb{Z}(Y^K, \mathcal{P})$ is isomorphic to the face ring $\mathbb{Z}[K]$.

Proof. For $J \subseteq [m]$, if J is the vertex set of a simplex σ in K , then $P_{\cap J} = F_{\sigma}$ is contractible which implies $H^*(P_{\cap J}) \otimes R_{\mathbb{Z}}^J \cong R_{\mathbb{Z}}^J$. Otherwise $P_{\cap J}$ is empty and $H^*(P_{\cap J}) \otimes R_{\mathbb{Z}}^J$ vanishes. So we can write

$$(18) \quad \mathbb{Z}(Y^K, \mathcal{P}) \cong \left(\bigoplus_{\sigma \subseteq [m], \sigma \in K} R_{\mathbb{Z}}^{\sigma}, \star \right),$$

where for any $f(x) \in R_{\mathbb{Z}}^{\sigma}$, $f'(x) \in R_{\mathbb{Z}}^{\sigma'}$ with $\sigma, \sigma' \in K$,

$$f(x) \star f'(x) = \begin{cases} f(x)f'(x), & \text{if } \sigma \cup \sigma' \in K; \\ 0, & \text{otherwise.} \end{cases}$$

By comparing (18) with the linear basis of $\mathbb{Z}[K]$ given in (3), we can easily check that $\mathbb{Z}(Y^K, \mathcal{P})$ is isomorphic to $\mathbb{Z}[K]$. The lemma is proved. \square

The above lemma implies that the integral equivariant cohomology ring of $(D^2, S^1)^K$ is isomorphic to the face ring $\mathbb{Z}[K]$ of K . Similarly, we can derive from Theorem 2.7 that the equivariant \mathbb{Z}_2 -cohomology ring of $(D^1, S^0)^K$ is isomorphic to $\mathbb{Z}_2[K]$. So we recover the statement of [12, Theorem 4.8].

From the above discussion, we see that the topological face ring of a panel structure is a generalization of the face ring of a simplicial complex from the equivariant cohomology viewpoint.

3.5. Polyhedral product over the panel structure on Y^K determined by a partition of the vertex set of K .

Let $\mathcal{J} = \{J_1, \dots, J_k\}$ be a partition of the vertex set $[m]$ of K . Then we can define a panel structure on Y^K associated to \mathcal{J} by:

$$\mathcal{P}^{\mathcal{J}} := \{P_i^{\mathcal{J}} = \bigcup_{j \in J_i} F_{v_j}\}_{i=1}^k.$$

For any simplex $\sigma \in K$, let $V(\sigma) \subseteq [m]$ be the vertex set of σ . By a similar argument as for (Y^K, \mathcal{P}) , we can deduce that for any $L \subseteq [k] = \{1, \dots, k\}$,

- If $L \neq \emptyset$, $P_{\cap L}^{\mathcal{J}}$ is the union of all the F_σ where $\sigma \in K$ and $V(\sigma) \cap J_i \neq \emptyset$ for all $i \in L$. If $L = \emptyset$, $P_{\cap \emptyset}^{\mathcal{J}} = Y^K$.
- $P_L^{\mathcal{J}}$ is homotopy equivalent to the full subcomplex K_{J_L} of K where

$$J_L = \bigcup_{i \in L} J_i.$$

For any $(\mathbb{X}, \mathbb{A}) = \{(X_i, A_i, a_i)\}_{i=1}^k$, we can apply our theorems to compute the stable decomposition, cohomology ring and equivariant cohomology ring of $(\mathbb{X}, \mathbb{A})^{(Y^K, \mathcal{P}^{\mathcal{J}})}$ under the same conditions on (\mathbb{X}, \mathbb{A}) in the previous subsections. The precise statements are left to the reader.

For the pair (D^2, S^1) , we can think of $(D^2, S^1)^{(Y^K, \mathcal{P}^{\mathcal{J}})}$ as the quotient space of $(D^2, S^1)^{(Y^K, \mathcal{P})}$ by the canonical action of an $(m - k)$ -dimensional subtorus of $(S^1)^m = \mathbb{R}^m / \mathbb{Z}^m$ determined by

$$\{\varepsilon_j - \varepsilon_{j'} \mid j, j' \text{ belong to the same } J_i \text{ for some } 1 \leq i \leq k\} \subseteq \mathbb{Z}^m.$$

where $\varepsilon_j = \{0, \dots, \overset{j}{1}, \dots, 0\} \in \mathbb{Z}^m$.

Similarly, for the pair (D^1, S^0) , we can think of $(D^1, S^0)^{(Y^K, \mathcal{P}^{\mathcal{J}})}$ as the quotient space of $(D^1, S^0)^{(Y^K, \mathcal{P})}$ by the canonical action of a subgroup of rank $m - k$ in $(\mathbb{Z}_2)^m$.

Using Theorem 2.1, we obtain the stable decomposition of $(D^2, S^1)^{(Y^K, \mathcal{P}^{\mathcal{J}})}$:

$$(D^2, S^1)^{(Y^K, \mathcal{P}^{\mathcal{J}})} \simeq \bigvee_{L \subseteq [k]} \Sigma^{|L|+2}(K_{J_L}).$$

Note that this formula recovers [23, Theorem 1.3].

In addition, for any partition $\mathcal{J} = \{J_1, \dots, J_k\}$ of $[m]$ with $k = \dim(K) + 1$, $(D^2, S^1)^{(Y^K, \mathcal{P}^{\mathcal{J}})}$ and $(D^1, S^0)^{(Y^K, \mathcal{P}^{\mathcal{J}})}$ can be considered as the generalization of the *pull-back from the linear model* (see [12, Example 1.15]) in the study of quasitoric manifolds and small covers.

4. Reexamining moment-angle complexes and polyhedral products over simplicial posets

The notion of moment-angle complex $\mathcal{Z}_{\mathcal{S}}$ associated to a simplicial poset \mathcal{S} is introduced by L\"u-Panov [17] as a generalization of the moment-angle complex of a simplicial complex. A poset (partially ordered set) \mathcal{S} with the order relation \leq is called *simplicial* if it has an initial element $\hat{0}$ and for each $\sigma \in \mathcal{S}$ the lower segment $[\hat{0}, \sigma] = \{\tau \in \mathcal{S} : \hat{0} \leq \tau \leq \sigma\}$ is the face poset of a simplex.

By the definition in [17], $\mathcal{Z}_{\mathcal{S}}$ is glued from the “moment-angle blocks” $(D^2, S^1)^\sigma$ according to the poset relation in \mathcal{S} . We can write $\mathcal{Z}_{\mathcal{S}}$ as the colimit of a diagram on \mathcal{S} defined by $\sigma \rightarrow (D^2, S^1)^\sigma$, that is

$$\mathcal{Z}_{\mathcal{S}} = \operatorname{colim}_{\sigma \in \mathcal{S}} (D^2, S^1)^\sigma.$$

For each $\sigma \in \mathcal{S}$ we assign a geometrical simplex Δ^σ whose face poset is $[\hat{0}, \sigma]$. By gluing these geometrical simplices together according to the order relation in \mathcal{S} , we get a cell complex $\Delta^{\mathcal{S}}$ in which the closure of each cell is identified with a simplex preserving the face structure, and all attaching maps are inclusions. We call $\Delta^{\mathcal{S}}$ the *geometrical realization* of \mathcal{S} . For convenience, we still use Δ^σ to denote the image of each geometrical simplex Δ^σ in $\Delta^{\mathcal{S}}$.

In the rest of this section, we assume that \mathcal{S} is a finite simplicial poset. Let the *vertex set* of $\Delta^{\mathcal{S}}$ be

$$V(\Delta^{\mathcal{S}}) = [m] = \{1, \dots, m\}.$$

Besides, we also use $v_1, \dots, v_m \in \mathcal{S}$ to denote the elements corresponding to the vertices of $\Delta^{\mathcal{S}}$.

For any $\sigma \in \mathcal{S}$, let $V(\Delta^\sigma) \subseteq [m]$ be the *vertex set* of the simplex Δ^σ . Note that here two different simplices of $\Delta^{\mathcal{S}}$ may have the same vertex set.

- Let $\operatorname{Sd}(\Delta^{\mathcal{S}})$ denote the barycentric subdivision of $\Delta^{\mathcal{S}}$. For each nonempty simplex $\Delta^\sigma \in \Delta^{\mathcal{S}}$, let F_σ denote the geometrical realization of the poset $\mathcal{S}_{\geq \sigma} = \{\tau \in \mathcal{S} \mid \sigma \leq \tau\}$. So F_σ is the subcomplex of $\operatorname{Sd}(\Delta^{\mathcal{S}})$ consisting of all simplices of the form $\sigma = \sigma_0 \lessdot \sigma_1 \lessdot \dots \lessdot \sigma_l$.




FIGURE 2.

- For any $J \subseteq [m]$, let \mathcal{S}_J denote the *full subposet* of \mathcal{S} consisting of all $\sigma \in \mathcal{S}$ with $V(\Delta^\sigma) \subseteq J$. Then the geometrical realization $\Delta^{\mathcal{S}_J} = \emptyset$ if and only if $\mathcal{S}_J = \hat{0}$.
- Let $Y^{\mathcal{S}} = u_0 \text{Sd}(\Delta^{\mathcal{S}})$ denote the cone of $\text{Sd}(\Delta^{\mathcal{S}})$ over a point u_0 .
- There is a natural panel structure \mathcal{P} on $Y^{\mathcal{S}}$ defined by:

$$\mathcal{P} = \{P_j = F_{v_j}\}_{j=1}^m.$$

For a subset $J \subseteq [m]$,

- If $J \neq \emptyset$, $P_{\cap J}$ is the disjoint union of all F_σ with $V(\Delta^\sigma) = J$. Let c_J = the number of connected components of $P_{\cap J}$.

Note that there is a one-to-one correspondence between the components F_σ in $P_{\cap J}$ and the simplices $\Delta^\sigma \subseteq \Delta^{\mathcal{S}}$ with $V(\Delta^\sigma) = J$ (see Figure 2 for an example). If $J = \emptyset$, $P_{\cap \emptyset} = Y^{\mathcal{S}}$ is contractible.

Lemma 4.1. *For any $J \subseteq [m]$, P_J is homotopy equivalent to $\Delta^{\mathcal{S}_J}$.*

Proof. The proof is analogous to the proof of [14, Corollary 4G.3]. Let

$$\mathcal{U}_J = \text{the cover of } P_J \text{ by } \{P_j = F_{v_j} \mid j \in J\}.$$

Then \mathcal{U}_J determines a *diagram of spaces* $X_{\mathcal{U}_J}$ (see [14, Section 4G] for the definition) where

- the vertices of $X_{\mathcal{U}_J}$ are the finite intersections of P_j 's in \mathcal{U}_J , and the edges of $X_{\mathcal{U}_J}$ are inclusions;
- the base of $X_{\mathcal{U}_J}$, denoted by Γ_J , is the barycentric subdivision of the nerve complex $N(\mathcal{U}_J)$ of \mathcal{U}_J .

From the diagram $X_{\mathcal{U}_J}$, one can construct a space $\Delta X_{\mathcal{U}_J}$ called the *realization* of $X_{\mathcal{U}_J}$ (see [14, Section 4G]). Moreover, by [14, Proposition 4G.2], the space $\Delta X_{\mathcal{U}_J}$ is homotopy equivalent to P_J .

For each $I \subseteq J$, the intersection $\bigcap_{j \in I} P_j = P_{\cap I}$ is the disjoint union of all F_σ 's with $V(\Delta^\sigma) = I$, which has c_I components. So we define another diagram of spaces over Γ_J , denoted by Z_J , where

- the vertices of Z_J are disjoint union of c_I points;

- the edges of Z_J consist of maps sending a vertex corresponding to F_σ to a vertex corresponding to F_τ with $\tau \subsetneq \sigma$.

Then by [14, Proposition 4G.1], the map $\Delta X_{\mathcal{U}_J} \rightarrow \Delta Z_J$ induced by sending each F_σ to a point is a homotopy equivalence. On the other hand, since Γ_J is the barycentric subdivision of the nerve complex of \mathcal{U}_J , it is routine to check that ΔZ_J is homeomorphic to Δ^{S_J} . This proves the lemma. \square

For any $(\mathbb{X}, \mathbb{A}) = \{(X_j, A_j, a_j)\}_{j=1}^m$, we can construct the polyhedral product $(\mathbb{X}, \mathbb{A})^{(Y^S, \mathcal{P})}$. From our general theorems of polyhedral products over a manifold with faces, we can derive the following results.

- If every X_j in $(\mathbb{X}, \mathbb{A}) = \{(X_j, A_j, a_j)\}_{j=1}^m$ is contractible, we obtain from Theorem 2.1 that

$$\Sigma((\mathbb{X}, \mathbb{A})^{(Y^S, \mathcal{P})}) \simeq \bigvee_{J \subseteq [m]} \Sigma\left((Y^S/P_J) \wedge \bigwedge_{j \in J} A_j\right).$$

Then since $Y^S/P_J = u_0 \text{Sd}(\Delta^S)/P_J \simeq u_0 P_J/P_J \simeq \Sigma(P_J) \simeq \Sigma(\Delta^{S_J})$,

$$(19) \quad \Sigma((\mathbb{X}, \mathbb{A})^{(Y^S, \mathcal{P})}) \simeq \bigvee_{J \subseteq [m]} \Sigma\left(\Sigma(\Delta^{S_J}) \wedge \bigwedge_{j \in J} A_j\right) \simeq \bigvee_{J \subseteq [m]} \Sigma\left(\Delta^{S_J} * \bigwedge_{j \in J} A_j\right).$$

- If every A_j in $(\mathbb{X}, \mathbb{A}) = \{(X_j, A_j, a_j)\}_{j=1}^m$ is contractible, Theorem 2.2 gives

$$S^1 \vee \Sigma((\mathbb{X}, \mathbb{A})^{(Y^S, \mathcal{P})}) \simeq \bigvee_{J \subseteq [m]} \Sigma\left((P_{\cap J} \cup y_0) \wedge \bigwedge_{j \in J} X_j\right).$$

– If $J \neq \emptyset$, $P_{\cap J}$ is the disjoint union of all F_σ with $V(\Delta^\sigma) = J$. Then since each F_σ is contractible, $(P_{\cap J} \cup y_0) \wedge \bigwedge_{j \in J} X_j$ is homotopy equivalent to a wedge of c_J copies of $\bigwedge_{j \in J} X_j$, denoted by $\bigvee_{c_J} \bigwedge_{j \in J} X_j$.

– If $J = \emptyset$, $(P_{\cap J} \cup y_0) \wedge \bigwedge_{j \in J} X_j = Y^S \cup y_0$.

From the above discussion, we obtain

$$S^1 \vee \Sigma((\mathbb{X}, \mathbb{A})^{(Y^S, \mathcal{P})}) \simeq \Sigma(Y^S \cup y_0) \vee \bigvee_{\emptyset \neq J \subseteq [m]} \Sigma\left(\bigvee_{c_J} \bigwedge_{j \in J} X_j\right).$$

Since $Y^S = u_0 \text{Sd}(\Delta^S)$ is contractible, $\Sigma(Y^S \cup y_0) \simeq S^1$. So we have

$$(20) \quad \Sigma((\mathbb{X}, \mathbb{A})^{(Y^S, \mathcal{P})}) \simeq \bigvee_{\emptyset \neq J \subseteq [m]} \Sigma\left(\bigvee_{c_J} \bigwedge_{j \in J} X_j\right).$$

In the rest of this section, we focus on the study of $(D^2, S^1)^{(Y^S, \mathcal{P})}$. There is a canonical $(S^1)^m$ -action on $(D^2, S^1)^{(Y^S, \mathcal{P})}$. By an argument completely parallel to the proof of Lemma 3.1, we can prove that $(D^2, S^1)^{(Y^S, \mathcal{P})}$ is equivariantly

homotopy equivalent to the moment-angle complex $\mathcal{Z}_{\mathcal{S}}$ over \mathcal{S} . Then by our Theorem 2.1 (also see (19)), we obtain the following proposition.

Proposition 4.2. *For any finite simplicial poset \mathcal{S} on $[m]$,*

$$(21) \quad \Sigma(\mathcal{Z}_{\mathcal{S}}) \simeq \Sigma((D^2, S^1)^{(Y^{\mathcal{S}}, \mathcal{P}})}) \simeq \bigvee_{J \subseteq [m]} \Sigma^{|J|+2}(\Delta^{\mathcal{S}_J}).$$

It was shown in [17, Theorem 3.5] that the integral cohomology ring of $\mathcal{Z}_{\mathcal{S}}$ is isomorphic as a graded algebra to the Tor algebra of the face ring $\mathbb{Z}[\mathcal{S}]$:

$$H^*(\mathcal{Z}_{\mathcal{S}}; \mathbb{Z}) \cong \mathrm{Tor}_{\mathbb{Z}[v_1, \dots, v_m]}(\mathbb{Z}[\mathcal{S}], \mathbb{Z}).$$

On the other hand, by our Corollary 2.4, there is a ring isomorphism (up to a sign) between the integral cohomology ring of $(D^2, S^1)^{(Y^{\mathcal{S}}, \mathcal{P}})}$ and $(\mathcal{R}_{Y^{\mathcal{S}}, \mathcal{P}}^*, \mathfrak{U})$, where

$$\mathcal{R}_{Y^{\mathcal{S}}, \mathcal{P}}^* = \bigoplus_{J \subseteq [m]} H^*(Y^{\mathcal{S}}, P_J).$$

Moreover, we can make this isomorphism degree-preserving by shifting the degrees of all the elements in $H^*(Y^{\mathcal{S}}, P_J)$ up by $|J|$ for every $J \subseteq [m]$. Then since $\mathcal{Z}_{\mathcal{S}}$ is homotopy equivalent to $(D^2, S^1)^{(Y^{\mathcal{S}}, \mathcal{P}})}$, we obtain the following proposition which gives a new topological description of $\mathrm{Tor}_{\mathbb{Z}[v_1, \dots, v_m]}(\mathbb{Z}[\mathcal{S}], \mathbb{Z})$.

Proposition 4.3. *For a finite simplicial poset \mathcal{S} on $[m]$, there is a graded algebra isomorphism (up to a sign) between $\mathrm{Tor}_{\mathbb{Z}[v_1, \dots, v_m]}(\mathbb{Z}[\mathcal{S}], \mathbb{Z})$ and $(\mathcal{R}_{Y^{\mathcal{S}}, \mathcal{P}}^*, \mathfrak{U})$ by shifting the degrees of all the elements in $H^*(Y^{\mathcal{S}}, P_J)$ up by $|J|$ for every $J \subseteq [m]$.*

In addition, it was shown in [17, Corollary 3.10] that $\mathrm{Tor}_{\mathbb{Z}[v_1, \dots, v_m]}(\mathbb{Z}[\mathcal{S}], \mathbb{Z})$ can be computed by a generalized Hochster's formula for the poset \mathcal{S} , from which one obtains the following formula:

$$(22) \quad H^p(\mathcal{Z}_{\mathcal{S}}) \cong \bigoplus_{J \subseteq [m]} \tilde{H}^{p-|J|-1}(\Delta^{\mathcal{S}_J}), \quad \forall p \in \mathbb{Z}.$$

Clearly, this formula also follows from our stable decomposition of $\mathcal{Z}_{\mathcal{S}}$ in (21).

According to [17, Remark 3.2], the equivariant cohomology ring of $\mathcal{Z}_{\mathcal{S}}$ with integral coefficients is isomorphic to the face ring $\mathbb{Z}[\mathcal{S}]$ of \mathcal{S} , that is:

$$(23) \quad H_{T^m}^*(\mathcal{Z}_{\mathcal{S}}) = H_{T^m}^*(\mathrm{colim}_{\sigma \in \mathcal{S}} (D^2, S^1)^{\sigma}) \cong H^*(\mathrm{colim}_{\sigma \in \mathcal{S}} (\mathbb{C}P^{\infty}, pt)^{\sigma}) \cong \mathbb{Z}[\mathcal{S}].$$

Recall that the *face ring* of a simplicial poset \mathcal{S} with \mathbf{k} -coefficients is the quotient ring (see Stanley [19])

$$(24) \quad \mathbf{k}[\mathcal{S}] := \mathbf{k}[\mathbf{v}_{\sigma} : \sigma \in \mathcal{S}] / \mathcal{I}_{\mathcal{S}}$$

where $\mathcal{I}_{\mathcal{S}}$ is the ideal generated by all the elements of the form

$$\mathbf{v}_{\hat{0}} - 1, \quad \mathbf{v}_{\sigma} \mathbf{v}_{\tau} - \mathbf{v}_{\sigma \wedge \tau} \cdot \sum_{\eta \in \sigma \vee \tau} \mathbf{v}_{\eta},$$

where $\sigma \wedge \tau$ denotes the greatest common lower bound of σ and τ , and $\sigma \vee \tau$ denotes the set of least common upper bounds of σ and τ . The sum over the empty set is assumed to be zero. So we have $\mathbf{v}_{\sigma} \mathbf{v}_{\tau} = 0$ in $\mathbf{k}[\mathcal{S}]$ if $\sigma \vee \tau = \emptyset$. If \mathcal{S} is a simplicial complex K , the definition (24) of $\mathbf{k}[K]$ is related to the definition (2) of $\mathbf{k}[K]$ by mapping \mathbf{v}_{σ} to $\prod_{j \in \sigma} v_j$.

On the other hand, by our Theorem 2.7, the equivariant cohomology ring of $(D^2, S^1)^{(Y^{\mathcal{S}}, \mathcal{P})}$ with integral coefficients is isomorphic to the topological face ring

$$\mathbb{Z}(Y^{\mathcal{S}}, \mathcal{P}) := \bigoplus_{J \subseteq [m]} H^*(P_{\cap J}) \otimes R_{\mathbb{Z}}^J$$

where the product is defined in (11). The following proposition implies that our result agrees with the conclusion (23) from [17].

Proposition 4.4. *There is a ring isomorphism from $\mathbb{Z}[\mathcal{S}]$ to $\mathbb{Z}(Y^{\mathcal{S}}, \mathcal{P})$.*

Proof. By our preceding discussion,

$$H^n(P_{\cap J}) \cong \begin{cases} \mathbb{Z}^{c_J}, & \text{if } n = 0; \\ 0, & \text{otherwise;} \end{cases}$$

where c_J is the number of connected components of $P_{\cap J}$ which correspond to all the simplices $\Delta^{\sigma} \subseteq \Delta^{\mathcal{S}}$ with $V(\Delta^{\sigma}) = J$. For convenience, denote the vertex set of Δ^{σ} by J_{σ} , that is

$$J_{\sigma} = V(\Delta^{\sigma}) \subseteq [m].$$

So we have

$$\mathbb{Z}(Y^{\mathcal{S}}, \mathcal{P}) \cong \bigoplus_{J \subseteq [m]} \bigoplus_{\sigma \in \mathcal{S}, J_{\sigma} = J} R_{\mathbb{Z}}^{J_{\sigma}} = \bigoplus_{\sigma \in \mathcal{S}} R_{\mathbb{Z}}^{J_{\sigma}, \sigma}.$$

where $R_{\mathbb{Z}}^{J_{\sigma}, \sigma} = R_{\mathbb{Z}}^{J_{\sigma}}$ for any $\sigma \in \mathcal{S}$.

On the other hand, by [18, Corollary 5.7], every element of $\mathbb{Z}[\mathcal{S}]$ can be written uniquely as a linear combination

$$\sum_{\sigma_1 \preceq \cdots \preceq \sigma_q} A(\sigma_1 \preceq \cdots \preceq \sigma_q; l_1, \cdots, l_q) \cdot \mathbf{v}_{\sigma_1}^{l_1} \cdots \mathbf{v}_{\sigma_q}^{l_q}$$

where $A(\sigma_1 \preceq \cdots \preceq \sigma_q; l_1, \cdots, l_q) \in \mathbb{Z}$ and the sum is taken over all chains of elements $\sigma_1 \preceq \cdots \preceq \sigma_q$ in \mathcal{S} with positive integers l_1, \cdots, l_q .

Then according to the maximal element in the chain $\sigma_1 \preceq \cdots \preceq \sigma_q$ in \mathcal{S} , we can write $\mathbb{Z}[\mathcal{S}]$ as a direct sum of subrings

$$\mathbb{Z}[\mathcal{S}] = \mathbb{Z} \oplus \bigoplus_{\hat{0} \neq \omega \in \mathcal{S}} \mathbb{Z}[\mathcal{S}]^\omega$$

where $\mathbb{Z}[\mathcal{S}]^\omega$ consists of elements of the form

$$\sum_{\sigma_1 \preceq \cdots \preceq \sigma_{q-1} \preceq \omega} A(\sigma_1 \preceq \cdots \preceq \sigma_{q-1} \preceq \omega; l_1, \dots, l_{q-1}, l_q) \cdot \mathbf{v}_{\sigma_1}^{l_1} \cdots \mathbf{v}_{\sigma_{q-1}}^{l_{q-1}} \mathbf{v}_\omega^{l_q}.$$

We can construct a ring homomorphism $\rho : \mathbb{Z}[\mathcal{S}] \rightarrow \mathbb{Z}(Y^\mathcal{S}, \mathcal{P})$ as follows. Define $\rho(\hat{0}) = 1$, and for any chain $\sigma_1 \preceq \cdots \preceq \sigma_q$ in \mathcal{S} , let

$$\rho(\mathbf{v}_{\sigma_1}^{l_1} \cdots \mathbf{v}_{\sigma_q}^{l_q}) = f_{\sigma_1}(x)^{l_1} \cdots f_{\sigma_q}(x)^{l_q} \in R_{\mathbb{Z}}^{J_{\sigma_q}, \sigma_q}$$

where for any $\hat{0} \neq \sigma \in \mathcal{S}$, $f_\sigma(x) = x_{i_1} \cdots x_{i_s}$ if $J_\sigma = \{v_{i_1}, \dots, v_{i_s}\}$.

We claim that ρ is a ring isomorphism. Indeed, for each $\hat{0} \neq \sigma \in \mathcal{S}$, we can think of $R_{\mathbb{Z}}^{J_\sigma, \sigma}$ as a subring of the face ring $\mathbb{Z}(\Delta^\sigma)$ of Δ^σ . So by the equivalence of the two definitions of $\mathbb{Z}(\Delta^\sigma)$ given by (2) and (24), the restriction of ρ to $\mathbb{Z}[\mathcal{S}]^\sigma$ is a ring isomorphism from $\mathbb{Z}[\mathcal{S}]^\sigma$ onto $R_{\mathbb{Z}}^{J_\sigma, \sigma}$. Then the proposition follows. \square

By the above proposition, we can consider the topological face ring of a panel structure as a generalization of the face ring of a simplicial poset as well.

5. Moment-angle complexes and polyhedral products over manifolds with corners

An n -dimensional manifold with corners Q is a Hausdorff space together with a maximal atlas of local charts onto open subsets of $\mathbb{R}_{\geq 0}^n$ such that the transitional functions are homeomorphisms which preserve the codimension of each point. Here the *codimension* $c(x)$ of a point $x = (x_1, \dots, x_n)$ in $\mathbb{R}_{\geq 0}^n$ is the number of x_i which are 0. So we have a well defined map $c : Q \rightarrow \mathbb{Z}_{\geq 0}$ where $c(q)$ is the codimension of a point $q \in Q$. In particular, the interior Q° of Q consists of points of codimension 0, i.e. $Q^\circ = c^{-1}(0)$.

Suppose Q is an n -dimensional manifold with corners with $\partial Q \neq \emptyset$. An *open face* of Q of codimension k is a connected component of $c^{-1}(k)$. A *closed face* is the closure of an open face. A face of codimension 1 is called a *facet* of Q . Let F_1, \dots, F_m be all the facets of Q .

A cell decomposition of Q is called *compatible* with its manifold with corners structure if all facets of Q are subcomplexes in the cell decomposition. In the following, we assume that Q is equipped with a compatible cell decomposition.

Then there is a natural panel structure on Q defined by

$$(25) \quad \mathcal{P} = \{P_j = F_j\}_{j=1}^m.$$

Then we can apply all our theorems proved in Section 2.1, 2.2 and 2.3 to obtain the corresponding results for any polyhedral product $(\mathbb{X}, \mathbb{A})^{(Q, \mathcal{P})}$.

- If Q is a convex polytope in an Euclidean space, then Q is a manifold with corners with an obvious compatible cell decomposition. The faces of (Q, \mathcal{P}) are exactly the faces of Q . Let K_Q be the nerve complex of Q covered by its facets. Since each face of Q is also a convex polytope hence contractible, we can use a similar argument as the proof of Lemma 3.1 to show that $(\mathbb{X}, \mathbb{A})^{(Q, \mathcal{P})}$ is homotopy equivalent to $(\mathbb{X}, \mathbb{A})^{K_Q}$. In particular, we have $(D^2, S^1)^{(Q, \mathcal{P})} \simeq (D^2, S^1)^{K_Q}$. In addition, there is another notion called *moment-angle space* of Q (denoted by \mathcal{Z}_Q) defined in [1]. It was proved in [1] that \mathcal{Z}_Q is also homotopy equivalent to $(D^2, S^1)^{K_Q}$.
- The polyhedral product $(\mathbb{X}, \mathbb{A})^{(Q, \mathcal{P})}$ generalizes the polyhedral products over a nice manifold with corners defined in [24, Section 4]. Recall that a manifold with corners Q is said to be *nice* if either its boundary ∂Q is empty or ∂Q is non-empty and any codimension- k face of Q is a connected component of the intersection of k different facets in Q . Moreover, the definition of $(\mathbb{X}, \mathbb{A})^{(Q, \mathcal{P})}$ clearly makes sense for an arbitrary manifold with corners (not necessarily nice).

Acknowledgment

This work is partially supported by National Natural Science Foundation of China (grant no.11871266) and the PAPD (priority academic program development) of Jiangsu higher education institutions.

Competing Interests. The author declares no competing interests pertaining to the undertaken research.

References

- [1] A. A. Ayzenberg and V. M. Buchstaber, *Nerve complexes and moment-angle spaces of convex polytopes*. Proc. Steklov Inst. Math. 275 (2011), no. 1, 15–46.
- [2] A. Bahri, M. Bendersky, F. Cohen and S. Gitler, *The Polyhedral Product Functor: a method of computation for moment-angle complexes, arrangements and related spaces*. Adv. Math. **225** (2010) no. 3, 1634–1668.
- [3] A. Bahri, M. Bendersky, F. Cohen and S. Gitler, *Cup products in generalized moment-angle complexes*. Math. Proc. Cambridge Philos. Soc. 153 (2012), no. 3, 457–469.
- [4] A. Bahri, M. Bendersky, F. Cohen and S. Gitler, *A spectral sequence for polyhedral products*. Adv. Math. 308 (2017), 767–814.

- [5] A. Bahri, M. Bendersky, F. Cohen and S. Gitler, *A Cartan formula for the cohomology of polyhedral products and its application to the ring structure*. [arXiv:2009.06818](https://arxiv.org/abs/2009.06818).
- [6] I. V. Baskakov, V. M. Buchstaber and T. E. Panov, *Algebras of cellular cochains, and torus actions*. (Russian) Uspekhi Mat. Nauk 59 (2004), no. **3** (357), 159–160; translation in Russian Math. Surveys 59 (2004), no. **3**, 562–563.
- [7] V. M. Buchstaber and T. E. Panov, *Actions of tori, combinatorial topology and homological algebra*. Russian Math. Surveys, 55 (2000), 825–921.
- [8] V. M. Buchstaber and T. E. Panov, *Torus actions and their applications in topology and combinatorics*. University Lecture Series, **24**. American Mathematical Society, Providence, RI, 2002.
- [9] V. M. Buchstaber and T. E. Panov, *Toric Topology*. Mathematical Surveys and Monographs, Vol. 204, American Mathematical Society, Providence, RI, 2015.
- [10] L. Cai, *On products in a real moment-angle manifold*. J. Math. Soc. Japan 69 (2017), no. **2**, 503–528.
- [11] M. W. Davis, *Groups generated by reflections and aspherical manifolds not covered by Euclidean space*. Ann. of Math. 117 (1983), 293–324.
- [12] M. W. Davis and T. Januszkiewicz, *Convex polytopes, Coxeter orbifolds and torus actions*. Duke Math. J. **62** (1991), no. **2**, 417–451.
- [13] M. Franz, *On the integral cohomology of smooth toric varieties*. Proc. Steklov Inst. Math. 252 (2006), 53–62.
- [14] A. Hatcher, *Algebraic topology*. Cambridge University Press, Cambridge, 2002.
- [15] L. Hoa, *On Segre products of affine semigroup rings*. Nagoya Math. J. 110 (1988), 113–128.
- [16] F. E. A. Johnson, *On the triangulation of stratified sets and singular varieties*. Trans. Amer. Math. Soc. 275 (1983), no. **1**, 333–343.
- [17] Z. Lü and T. E. Panov, *Moment-angle complexes from simplicial posets*. Cent. Eur. J. Math. 9 (2011), no. **4**, 715–730.
- [18] M. Masuda and T. Panov, *On the cohomology of torus manifolds*. Osaka J. Math. 43 (2006), no. **3**, 711–746.
- [19] R. P. Stanley, *f-vectors and h-vectors of simplicial posets*, J. Pure Appl. Algebra. 71 (1991), 319–331.
- [20] R. Vogt, *Homotopy limits and colimits*. Math. Z. 134 (1973) 11–52.
- [21] X. J. Wang, Q. B. Zheng, *The homology of simplicial complements and the cohomology of polyhedral products*. Forum Math. 27 (2015), no. **4**, 2267–2299.
- [22] V. Welker, G. Ziegler, R. Živaljević, *Homotopy colimits-comparison lemmas for combinatorial applications*, J. Reine Angew. Math. 509 (1999) 117–149.
- [23] L. Yu, *On Hochster’s formula for a class of quotient spaces of moment-angle complexes*. Osaka Journal of Mathematics 56 (2019), no. **1**, 33–50.
- [24] L. Yu, *A generalization of moment-angle manifolds with non-contractible orbit spaces*, Algebr. Geom. Topol. 24 (2024) 449–492.
- [25] Q. B. Zheng, *The cohomology algebra of polyhedral product spaces*. J. Pure Appl. Algebra 220 (2016), no. **11**, 3752–3776.
- [26] G. Ziegler, R. Živaljević, *Homotopy types of sub-space arrangements via diagrams of spaces*. Math. Ann. 295 (1993), 527–548.

SCHOOL OF MATHEMATICS, NANJING UNIVERSITY, NANJING, 210093, P.R.CHINA.

Email address: yuli@nju.edu.cn