

MATRIX FORMS OF IMMANANT INEQUALITIES

FELIX HUBER AND HANS MAASSEN

Abstract: We translate inequalities and conjectures for immanants and generalized matrix functions into inequalities in the Löwner order. These have the form of trace polynomials and generalize the inequalities from [FH, J. Math. Phys. 62 (2021), 2, 022203].

1. IMMANANTS

Let $A = (a_{ij})$ be a complex $n \times n$ matrix. The determinant and permanent of A are given by

$$\det(A) = \sum_{\sigma \in S_n} \text{sign}(\sigma) \prod_{t=1}^n a_{t\sigma(t)}$$

$$\text{per}(A) = \sum_{\sigma \in S_n} \prod_{t=1}^n a_{t\sigma(t)}.$$

A generalization of these functions are *immanants*. Given an irreducible character χ of the symmetric group S_n , the corresponding immanant is defined as

$$(1) \quad \text{imm}_\chi(A) = \sum_{\sigma \in S_n} \chi(\sigma) \prod_{t=1}^n a_{t\sigma(t)}.$$

Now one sees that the determinant and the permanent arise when $\chi(\sigma) = \text{sign}(\sigma)$ and $\chi(\sigma) = 1$ are the alternating and trivial representations respectively.

This can be generalized further: given a subgroup H of S_n and χ a character on H , one can define a *generalized matrix function* [7, 8]

$$(2) \quad d_\chi^H(A) = \sum_{\sigma \in H} \chi(\sigma) \prod_{t=1}^n a_{t\sigma(t)},$$

where the sum is now over the subgroup H instead of S_n . Note that \det , per , and imm_χ are special cases of the function d_χ^H .

It is well known that, when A is positive semidefinite, all the above quantities are non-negative

$$(3) \quad d_\chi^H(A) \geq 0.$$

Many more inequalities are known. A short selection appears in Appendix A, a larger overview can be found in Chapter 7 of Ref. [8].

We write $A \geq B$ whenever $A - B$ is positive semidefinite; this is the Löwner order. The aim of this note is to turn *scalar* inequalities for positive semidefinite operators, such as e.g. appearing in Formula (3), into *matrix* inequalities in the Löwner order.

Date: April 13, 2021.

FH acknowledges support by the Government of Spain (FIS2020-TRANQI and Severo Ochoa CEX2019-000910-S), Fundació Mir-Puig, Generalitat de Catalunya (AGAUR SGR 1381 and CERCA), the European Union under Horizon2020 (PROBIST 754510), and the Foundation for Polish Science through TEAM-NET (POIR.04.04.00-00-17C1/18-00).

2. MAPPING TO TENSOR PRODUCTS

To approach this subject, it is convenient to consider any given positive semidefinite matrix A as the Gram matrix $A = (a_{ij})$ of some set of vectors $\{|v_1\rangle, \dots, |v_n\rangle\}$. So we write

$$(4) \quad A = \begin{pmatrix} \langle v_1|v_1\rangle & \langle v_1|v_2\rangle & \dots & \langle v_1|v_n\rangle \\ \langle v_2|v_1\rangle & \langle v_2|v_2\rangle & & \vdots \\ \vdots & & \ddots & \\ \langle v_n|v_1\rangle & \dots & & \langle v_n|v_n\rangle \end{pmatrix}$$

where $|v_1\rangle, \dots, |v_n\rangle$ are complex vectors in a sufficiently large Hilbert space \mathbb{C}^m . Every Gram matrix is positive semidefinite, and in turn, every positive semidefinite matrix A can be written as the Gram matrix of some set of vectors, which is unique up to unitary equivalence. From now we call them *Gram vectors* for A . The canonical choice will be through the matrix square root: for every matrix $A \geq 0$ there is a unique matrix $B \geq 0$ such that $A = B^2$. Then the columns of B can be used as Gram vectors for A .

Now when $A \geq 0$ is thus understood as a Gram matrix one can relate immanants and generalised matrix functions of A with an expression on a tensor product space involving its Gram vectors.

For this we consider the action of the symmetric group on a tensor product space. Namely, let $\sigma \in S_n$ act on $(\mathbb{C}^m)^{\otimes n} = \mathbb{C}^m \otimes \dots \otimes \mathbb{C}^m$ (n times) by permuting its tensor factors,

$$\sigma |v_1\rangle \otimes \dots \otimes |v_n\rangle = |v_{\sigma^{-1}(1)}\rangle \otimes \dots \otimes |v_{\sigma^{-1}(n)}\rangle .$$

For example, (143)(2) $|v_1\rangle \otimes |v_2\rangle \otimes |v_3\rangle \otimes |v_4\rangle = |v_3\rangle \otimes |v_2\rangle \otimes |v_4\rangle \otimes |v_1\rangle$.

Our aim is to express the generalized matrix function d_χ^H as a multilinear function on a tensor product space; see also Theorem 7.26 in Ref. [8] and Ref. [6] which take such an approach.

Lemma 1. *Let $A \geq 0$ with Gram vectors $|v_1\rangle, \dots, |v_n\rangle$ and define $X_i = |v_i\rangle\langle v_i|$, $1 \leq i \leq n$. Then*

$$d_\chi^H(A) = \sum_{\sigma \in H} \chi(\sigma) \operatorname{tr} [\sigma^{-1} X_1 \otimes \dots \otimes X_n]$$

Proof. Put $v := v_1 \otimes \dots \otimes v_n$. Then

$$\begin{aligned} \prod_{t=1}^n a_{t\sigma(t)} &= \prod_{t=1}^n \langle v_t | v_{\sigma(t)} \rangle = \langle v | \sigma^{-1} v \rangle \\ &= \operatorname{tr}(\sigma^{-1} |v\rangle\langle v|) = \operatorname{tr}(\sigma^{-1} X_1 \otimes \dots \otimes X_n) . \end{aligned}$$

The statement follows from the definition (2) of d_χ^H . \square

Now it is interesting to understand the immanant of positive semidefinite matrices as arising from projectors: let H be subgroup of S_n . The centrally primitive idempotent in the group ring $\mathbb{C}[H]$ associated with a character χ is [9, Theorem 3.6.2]

$$p_\chi^H = \frac{\chi(e)}{|H|} \sum_{\sigma \in H} \chi(\sigma) \sigma^{-1} .$$

One has that $(p_\chi^H)^2 = p_\chi^H$. Under a representation, p_χ^H is projects onto an isotypic component and thus $p_\chi^H \geq 0$. In terms of the above proof it is then immediate that

Eq. (3) holds:

$$(5) \quad d_\chi^H(A) = \frac{|H|}{\chi(e)} \langle v | p_\chi^H v \rangle = \frac{|H|}{\chi(e)} \|p_\chi^H v\|^2 \geq 0.$$

3. TRACE POLYNOMIAL INEQUALITIES

Each summand in Lemma 1 is linear in the X_i 's. This allows to turn scalar inequalities for immanants into inequalities in terms of the Löwner order.

For this we require a canonical ordering when writing permutations. Given a permutation decomposed into cycles as $\sigma = \sigma_1 \dots \sigma_l \in S_n$, its canonical order is such that its largest element appears at the end of the last cycle, $\sigma_l = (\dots n)$. Now given a set of complex matrices X_1, \dots, X_n of equal size, we define for every permutation $(\alpha_1 \dots \alpha_r) \dots (\zeta_1 \dots \zeta_t)(\xi_1 \dots \xi_w n)$ the scalar

$$T_\sigma = \text{tr}(X_{\alpha_1} \dots X_{\alpha_r}) \cdots \text{tr}(X_{\zeta_1} \dots X_{\zeta_t}) \text{tr}(X_{\xi_1} \cdots X_{\xi_w} X_n).$$

and the matrix

$$\tilde{T}_\sigma = \text{tr}(X_{\alpha_1} \dots X_{\alpha_r}) \cdots \text{tr}(X_{\zeta_1} \dots X_{\zeta_t}) X_{\xi_1} \cdots X_{\xi_w} \mathbb{1},$$

where the last matrix X_n is replaced by $\mathbb{1}$ and its trace opened. It is clear that $T_\sigma = \text{tr}(\tilde{T}_\sigma X_n)$ holds. Furthermore, it is well-known that [4]

$$T_\sigma = \text{tr}(\sigma^{-1} X_1 \otimes \dots \otimes X_n)$$

Now suppose one is given a function $f : S_n \rightarrow \mathbb{C}$. To it we associate, on the one hand, the generalized matrix function $d_f : M_n \rightarrow \mathbb{C}$,

$$d_f(A) = \sum_{\sigma \in S_n} f(\sigma) \prod_{t=1}^n a_{t\sigma(t)}.$$

On the other hand, we define for complex $m \times m$ matrices X_1, \dots, X_{n-1} the matrix-valued map $\tilde{d}_f : M_m^{n-1} \rightarrow M_m$,

$$\tilde{d}_f(X_1, \dots, X_{n-1}) = \sum_{\sigma \in S_n} f(\sigma) \tilde{T}_\sigma.$$

Starting from an inequality for some d_f a corresponding matrix inequality for \tilde{d}_f can be made.

Theorem 2. *Suppose for all positive semidefinite $n \times n$ matrices A it holds that*

$$d_f(A) \geq 0.$$

Then for all positive semidefinite matrices X_1, \dots, X_{n-1} of equal size it holds that

$$\tilde{d}_f(X_1, \dots, X_{n-1}) \geq 0.$$

Proof. Suppose the X_i 's have size $m \times m$. Then $\tilde{d}_f \geq 0$ if and only if $\langle w | \tilde{d}_f | w \rangle \geq 0$ holds for all $|w\rangle \in \mathbb{C}^m$. We rewrite this as

$$(6) \quad \begin{aligned} \langle w | \tilde{d}_f | w \rangle &= \langle w | \sum_{\sigma \in S_n} f(\sigma) \tilde{T}_\sigma | w \rangle \\ &= \sum_{\sigma \in S_n} f(\sigma) T_\sigma \\ &= \sum_{\sigma \in S_n} f(\sigma) \text{tr}(\sigma^{-1} X_1 \otimes \dots \otimes X_n), \end{aligned}$$

where we have set $X_n = |w\rangle\langle w|$.

Now every such tensor product $X_1 \otimes \dots \otimes X_n$ where all X_i are positive semidefinite can be decomposed as a convex combination of a finite set of rank one matrices of

the form $|v_1\rangle\langle v_1| \otimes \dots \otimes |v_n\rangle\langle v_n|$. It thus suffices to show that Eq. (6) holds for each term in the decomposition. With Lemma 1 and Eq. (5) we see that

$$\sum_{\sigma \in S_n} f(\sigma) \operatorname{tr}(\sigma^{-1} |v_1\rangle\langle v_1| \otimes \dots \otimes |v_n\rangle\langle v_n|) = d_f(A) \geq 0,$$

where A is the Gram matrix of the vectors $|v_1\rangle, \dots, |v_n\rangle$.

This ends the proof. \square

In particular, if for some $\alpha, \beta \in \mathbb{R}$ an inequality $\alpha d_\chi^H(A) \geq \beta d_\psi^K(A)$ holds for all complex positive semidefinite $n \times n$ matrices A , then $\alpha \widetilde{d}_\chi^H(X_1, \dots, X_{n-1}) \geq \beta \widetilde{d}_\psi^K(X_1, \dots, X_{n-1})$ for all $X_1, \dots, X_{n-1} \geq 0$ of equal size.

Remark. In Ref. [3] the author has derived a special case of this inequality, namely when χ is a character of the symmetric group, in the framework of the Choi-Jamiołkowski isomorphism.

Example 3. Let $H = A_4$ be the alternating group of degree four. The conjugacy class of (123) in S_4 splits in A_4 into two: the conjugacy classes

$$\begin{aligned} C_{A_4}((123)) &= \{(123), (142), (243), (134)\}, \\ C_{A_4}((132)) &= \{(132), (124), (234), (143)\}. \end{aligned}$$

The conjugacy class of (12)(34) does not split. So one has the representatives $(e), (12)(34), (123), (132)$. Define $\omega = e^{2\pi i/3}$. The non-trivial characters of A_4 are

	(e)	(12)(34)	(123)	(132)
χ_1	3	-1	0	0
χ_2	1	1	ω	$\bar{\omega}$
χ_3	1	1	$\bar{\omega}$	ω

The element $\frac{\chi_i(e)}{|A_4|} \sum_{\sigma \in A_4} \chi_i(\sigma^{-1}) \sigma$ is idempotent and thus $d_{\chi_i}^{A_4}(A) \geq 0$ for all $A \geq 0$ as reasoned in Eq. (5). The expressions are multilinear and it suffices to consider matrices of trace one. Then for square matrices X, Y, Z define

$$\begin{aligned} L &= \operatorname{tr}(XY)Z + \operatorname{tr}(XZ)Y + \operatorname{tr}(YZ)X, \\ M &= \operatorname{tr}(ZYX)\mathbb{1} + XY + YZ + ZX. \end{aligned}$$

The matrix functions $\widetilde{d}_{\chi_i}^{A_4} = \widetilde{d}_{\chi_i}^{A_4}(X, Y, Z)$ are then

$$\begin{aligned} \widetilde{d}_{\chi_1}^{A_4} &= 3\mathbb{1} - L \\ \widetilde{d}_{\chi_2}^{A_4} &= \mathbb{1} + L + \omega M + \bar{\omega} M^* \\ \widetilde{d}_{\chi_3}^{A_4} &= \mathbb{1} + L + \bar{\omega} M + \omega M^*. \end{aligned}$$

where M^* is the hermitian conjugate of M . Theorem 2 implies that for all $X, Y, Z \geq 0$ with $\operatorname{tr}(X) = \operatorname{tr}(Y) = \operatorname{tr}(Z) = 1$ the above three expressions are positive semidefinite.

Example 4. Set $f = \chi_2^{A_4}$. Again it suffices to consider matrices of trace one. It follows from Theorem 2 and Watkins's theorem (see Appendix A) that

$$(\omega - 1)M + (\bar{\omega} - 1)M^* + \operatorname{tr}(L)\mathbb{1} + X + Y + Z + X\{Y, Z\} + Y\{Z, X\} + Z\{X, Y\} \geq 0$$

holds for all $X, Y, Z \geq 0$ with $\operatorname{tr}(X) = \operatorname{tr}(Y) = \operatorname{tr}(Z) = 1$.

Example 5. In $\mathbb{C}S_3$ the centrally primitive idempotent associated to the partition $(2, 1)$ reads $p_{(2,1)} = \frac{1}{3}[2(e) - (123) - (132)]$. Watkin's theorem applied to $p_{(1,2)}$ and the non-negativity of the determinant yields the following upper and lower bounds on the anti-commutator of positive semidefinite matrices X, Y of trace 1:

$$X + Y + [\text{tr}(XY) - 1]\mathbf{1} \leq XY + YX \leq \frac{2}{3}[X + Y + \text{tr}(XY)]\mathbf{1}.$$

Remark. We note when $\widetilde{\text{imm}}_\chi := \widetilde{d}_\chi^{S_n}(X_1, \dots, X_{n-1})$ becomes a matrix identity: one has that $\widetilde{\text{imm}}_\chi(X_1, \dots, X_{n-1}) = 0$ for all complex $m \times m$ matrices X_1, \dots, X_{n-1} when χ corresponds to a Young diagram with more than m rows. This result is also known as Lew's polarized (or generalized) Cayley-Hamilton theorem [5] and it can be understood as a consequence of

$$p_\chi |v_1\rangle \otimes \dots \otimes |v_n\rangle = 0.$$

whenever χ corresponds to a Young diagram with r rows but $|v_1\rangle, \dots, |v_n\rangle \in \mathbb{C}^m$ with $m < r$ [3].

APPENDIX A. SOME IMMANANT INEQUALITIES

The following inequalities for the immanants of positive semidefinite matrices are well-known [1, 2, 8, 10].

Hadamard inequality (1893): For all $A \geq 0$

$$\det(A) \leq \prod_i a_{ii}.$$

Schur inequality (1918): For all $A \geq 0$

$$\det(A) \leq \frac{\text{imm}_\lambda(A)}{\chi_\lambda(e)}.$$

Marcus inequality (1963): For all $A \geq 0$

$$\text{per}(A) \geq \prod_i a_{ii}.$$

Permanent dominance conjecture (open): For all $A \geq 0$

$$\text{per}(A) \geq \frac{\text{imm}_\lambda(A)}{\chi_\lambda(e)}.$$

Heyfron's theorem (1988): Define the normalized immanant $\overline{\text{imm}}_\chi(A) = \text{imm}_\chi(A)/\chi(e)$. For all $A \geq 0$

$$\begin{aligned} \det(A) = \overline{\text{imm}}_{[1^n]}(A) &\leq \overline{\text{imm}}_{[2,1^{n-2}]}(A) \leq \overline{\text{imm}}_{[3,1^{n-3}]}(A) \leq \dots \\ &\dots \leq \overline{\text{imm}}_{[n-3,1^3]}(A) \leq \overline{\text{imm}}_{[n-2,1^2]}(A) \leq \overline{\text{imm}}_{[n]}(A) = \text{per}(A) \end{aligned}$$

where the dots consist of a chain of immanants corresponding to single-hook Young Tableaux with increasing row and decreasing column lengths.

Watkins's theorem (1988): Suppose $f : S_n \rightarrow \mathbb{C}$ is such that $d_f(A) \geq 0$ for all complex semidefinite $m \times m$ matrices A . Then it holds that $d_f(A) \geq f(e) \det(A)$ for all complex semidefinite $m \times m$ matrices A .

REFERENCES

1. Gi-Sang Cheon and Ian M. Wanless, *An update on Minc's survey of open problems involving permanents*, Linear Algebra Appl. **403** (2005), 314–342.
2. Robert Grone, Russel Merris, and William Watkins, *Cones in the group algebra related to Schur's determinantal inequality*, Rocky Mountain J. Math. **18** (1988), no. 1, 137–146.
3. Felix Huber, *Positive maps and trace polynomials from the symmetric group*, J. Math. Phys. **62** (2021), no. 2, 022203.
4. Bertram Kostant, *A Theorem of Frobenius, a Theorem of Amitsur-Levitski and Cohomology Theory*, pp. 64–91, Springer New York, New York, NY, 2009.
5. John S. Lew, *The generalized Cayley-Hamilton theorem in n dimensions*, J. App. Math. Phys. **17** (1966), 650.
6. Hans Maassen and Burkhard Kümmerer, *Entanglement of symmetric Werner states*, available online at <http://www.bjadres.nl/MathQuantWorkshop/Slides/SymmWernerHandout.pdf>, May 2019.
7. Marvin Marcus and Henryk Minc, *Generalized matrix functions*, Trans. Amer. Math. Soc. **116** (1965), 316–329.
8. R. Merris, *Multilinear Algebra*, CRC Press, 1997.
9. Peter Webb, *A course in finite group representation theory*, Cambridge Studies in Advanced Mathematics, Cambridge University Press, 2016.
10. Fuzhen Zhang, *An update on a few permanent conjectures*, Spec. Matrices **4** (2016), 305–316.

FELIX HUBER, ATOMIC OPTICS DEPARTMENT, JAGIELLONIAN UNIVERSITY, 30-348 KRAKÓW, POLAND, E-MAIL: FELIX.HUBER@UJ.EDU.PL

HANS MAASSEN, DEPARTMENT OF MATHEMATICS, RADBOUD UNIVERSITY, 6525 AJ NIJMEGEN, THE NETHERLANDS