

DIFFERENTIAL SUBORDINATION IMPLICATIONS FOR CERTAIN CARATHÉODORY FUNCTIONS

MEGHNA SHARMA, SUSHIL KUMAR, AND NAVEEN KUMAR JAIN

ABSTRACT. In this article, we wish to establish some first order differential subordination relations for certain Carathéodory functions with nice geometrical properties. Moreover, several implications are determined so that the normalized analytic function belongs to various subclasses of starlike functions.

1. INTRODUCTION

Denote the collection of all functions f which are analytic on the open unit disc by \mathcal{H} . Let $\mathcal{A} \subset \mathcal{H}$ be the subclass consisting of analytic functions given by $f(z) = z + \sum_{n=2}^{\infty} a_n z^n$ and normalised by the conditions $f(0) = 0$ and $f'(0) - 1 = 0$. Further, let \mathcal{S}^* and \mathcal{C} denote the subclasses of univalent function consisting of starlike and convex functions, characterised by the quantities $zf'(z)/f(z)$ and $1 + zf''(z)/f'(z)$ lying in the interior of the right half plane respectively. Let f and g be members of \mathcal{H} . We say f is subordinate to g (written as $f \prec g$) if there exists a function $w \in \mathcal{H}$ with $w(0) = 0$ and $|w(z)| < 1$ such that $f(z) = g(w(z))$. Equivalently, if g is univalent in \mathbb{D} , then the conditions $f(0) = g(0)$ and $f(\mathbb{D}) \subset g(\mathbb{D})$ together gives $f \prec g$. For more details, see [16]. The unified class of starlike functions $\mathcal{S}_{\varphi}^* := \{f \in \mathcal{A} : zf'(z)/f(z) \prec \varphi(z); \text{ for all } z \in \mathbb{D}\}$ where φ is analytic, univalent, $\varphi(\mathbb{D})$ is starlike with respect to $\varphi(0) = 1$ and $\text{Re}(\varphi) > 0$ was introduced and studied by Ma and Minda [14]. Various subclasses of starlike functions have been studied by considering different choices of φ in recent years. For $\varphi(z) := (1 + Az)/(1 + Bz)$, $(-1 \leq B < A \leq 1)$, the class \mathcal{S}_{φ}^* reduces to the class $\mathcal{S}^*[A, B]$, introduced by Janowski [9]. A function $f \in \mathcal{H}$ is said to be a Carathéodory function if $f(0) = 1$ and $\text{Re}(f(z)) > 0$. The class of such functions is denoted by \mathcal{P} . On taking some Carathéodory functions $\varphi(z) := e^z, \phi_q(z), \phi_0(z), \phi_c(z), \phi_{\text{lim}}(z), \mathcal{Q}(z), \phi_{SG}, \phi_s(z)$, the class \mathcal{S}_{φ}^* reduce to subclasses \mathcal{S}_e^* [15], \mathcal{S}_q^* [20], \mathcal{S}_R^* [12], \mathcal{S}_c^* [21], \mathcal{S}_{LC}^* [23], \mathcal{S}_B^* [5], \mathcal{S}_{SG}^* [8], \mathcal{S}_s^* [4] respectively, where

$$\begin{aligned} \phi_q(z) &= z + \sqrt{1+z^2}, \phi_0(z) := 1 + \frac{z}{k} \cdot \frac{k+z}{k-z}; \quad k = 1 + \sqrt{2}, \\ \phi_c(z) &:= 1 + \frac{4z}{3} + \frac{2z^2}{3}, \phi_{\text{lim}}(z) := 1 + \sqrt{2}z + \frac{z^2}{2}, \phi_s(z) := 1 + \sin z. \end{aligned}$$

Recently, Kumar *et al.* [5] introduced and studied differential subordination relations and radius estimates for the class $\mathcal{S}_B^* := \mathcal{S}^*(\mathcal{Q}(z))$, where

$$\mathcal{Q}(z) := e^{e^z - 1} \tag{1.1}$$

2010 *Mathematics Subject Classification.* 30C45.

Key words and phrases. Differential Subordination; Carathéodory function; Starlike Function; sufficient conditions.

The first author is supported by Junior Research Fellowship from Council of Scientific and Industrial Research, New Delhi, Ref. No.:1753/(CSIR-UGC NET JUNE, 2018).

In 2020, Goel and Kumar [8] studied the subclass $\mathcal{S}_{SG}^* := \mathcal{S}^*(\phi_{SG})$, where

$$\phi_{SG}(z) = 2/(1 + e^{-z}) \text{ for all } z \in \mathbb{D}. \quad (1.2)$$

Several first order differential subordination results, radius estimates and coefficient estimates were investigated. These subclasses of starlike functions are well associated with the right half plane of the complex plane and introduced and discussed by various authors.

In 1989, for $p \in \mathcal{P}$, Nunokawa *et al.* [18] proved that the differential subordination $1 + zp'(z) \prec 1 + z$ implies $p(z) \prec 1 + z$. Further, authors [19] established sufficient conditions for starlike functions discussed by Silverman [22] to be strongly convex and strongly starlike in \mathbb{D} . In 2006, Kanas [10] determined the conditions for the functions to map \mathbb{D} onto hyperbolic and parabolic regions using the concept of differential subordination. In 2007, Ali *et al.* [2] obtained conditions on $\beta \in \mathbb{R}$ for which $1 + \beta zp'(z)/p^j(z) \prec (1 + Dz)/(1 + Ez)$, $j = 0, 1, 2$ implies $p(z) \prec (1 + Az)/(1 + Bz)$, where $A, B, D, E \in [-1, 1]$. Later, Kumar and Ravichandran [13] determined sharp upper bounds on β such that $1 + \beta zp'(z)/p^j(z)$, $j = 0, 1, 2$ is subordinate to some Carathéodory functions like $e^z, \phi_0(z)$ etc. implies $p(z) \prec e^z$ and $(1 + Az)/(1 + Bz)$. For more such results, we refer [1, 3, 7, 6].

In the present paper, we determine sharp estimate on β so that $p(z) \prec \phi_q(z), \mathcal{Q}(z), \phi_c(z), \phi_0(z), \phi_{\text{lim}}(z), \phi_s(z), \phi_{SG}(z)$ whenever $1 + \beta zp'(z)/p^j(z) \prec \mathcal{Q}(z)$ and $\phi_{SG}(z)$; ($j = 0, 1, 2$). Further the best possible bound on β is computed such that $p(z) \prec \mathcal{Q}(z)$ whenever $1 + \beta zp'(z)/p^j(z) \prec \phi_c(z)$; ($j = 0, 1, 2$). At last, the upper bound on β is estimated so that the subordination $1 + \beta zp'(z)/p^j(z) \prec \phi_0(z)$ and $\phi_c(z)$ implies $p(z) \prec \phi_{SG}(z)$. Moreover, sufficient conditions are obtained for an analytic function f to be a member of a certain subclass of starlike function.

2. MAIN RESULTS

Before we prove our main results, we recall following lemma which plays a vital role in our proofs.

Lemma 2.1. [17, Theorem 3.4h, p.132] *Let $q : \mathbb{D} \rightarrow \mathbb{C}$ be analytic, and ψ and v be analytic in a domain $U \supseteq q(\mathbb{D})$ with $\psi(w) \neq 0$ whenever $w \in q(\mathbb{D})$. Set*

$$Q(z) := zq'(z)\psi(q(z)) \quad \text{and} \quad h(z) := v(q(z)) + Q(z), \quad z \in \mathbb{D}.$$

Suppose that

- (i) either $h(z)$ is convex, or $Q(z)$ is starlike univalent in \mathbb{D} and
- (ii) $\text{Re} \left(\frac{zh'(z)}{Q(z)} \right) > 0, z \in \mathbb{D}$.

If p is analytic in \mathbb{D} , with $p(0) = q(0), p(\mathbb{D}) \subset U$ and

$$v(p(z)) + zp'(z)\psi(p(z)) \prec v(q(z)) + zq'(z)\psi(q(z))$$

then $p \prec q$, and q is the best dominant.

Throughout this paper, the following notations will be used:

$$\Psi_\beta(z, p(z)) = 1 + \beta zp'(z), \quad \Lambda_\beta(z, p(z)) = 1 + \beta \frac{zp'(z)}{p(z)}, \quad \text{and} \quad \Theta_\beta(z, p(z)) = 1 + \beta \frac{zp'(z)}{p^2(z)}.$$

Theorem 2.2. Let $\mathcal{Q}(z) \in \mathcal{P}$ be defined by (1.1) and further

$$\mathcal{L} = \int_{-1}^0 \frac{e^{e^t-1} - 1}{t} dt \quad \text{and} \quad \mathfrak{U} = \int_0^1 \frac{e^{e^t-1} - 1}{t} dt. \tag{2.1}$$

Assume p to be an analytic function in \mathbb{D} with $p(0) = 1$. If $\Psi_\beta(z, p(z)) \prec \mathcal{Q}(z)$, then

- (a) $p(z) \prec \phi_q(z)$ for $\beta \geq \frac{1}{\sqrt{2}}\mathfrak{U} \approx 1.49762$.
- (b) $p(z) \prec \mathcal{Q}(z)$ for $\beta \geq \frac{1}{1-e^{(e^{-1}-1)}}\mathcal{L} \approx 1.446103$.
- (c) $p(z) \prec \phi_c(z)$ for $\beta \geq \frac{1}{2}\mathfrak{U} \approx 1.05898$.
- (d) $p(z) \prec \phi_0(z)$ for $\beta \geq (3 + 2\sqrt{2})\mathcal{L} \approx 3.94906$.
- (e) $p(z) \prec \phi_{\lim}(z)$ for $\beta \geq \frac{2}{2\sqrt{2}+1}\mathfrak{U} \approx 1.10643$.
- (f) $p(z) \prec \phi_s(z)$ for $\beta \geq \frac{1}{\sin 1}\mathfrak{U} \approx 2.51696$.
- (g) $p(z) \prec \phi_{SG}(z)$ for $\beta \geq \frac{e+1}{e-1}\mathfrak{U} \approx 4.583145$.

The bounds in each case are sharp.

Proof. The analytic function $q_\beta : \overline{\mathbb{D}} \rightarrow \mathbb{C}$ defined by

$$q_\beta(z) = 1 + \frac{1}{\beta} \int_0^z \frac{e^{e^t-1} - 1}{t} dt$$

is a solution of the first order linear differential equation $1 + \beta z q'_\beta(z) = e^{e^z-1}$. For $w \in \mathbb{C}$, define the functions $v(w) = 1$ and $\psi(w) = \beta$. Now, the function $Q : \overline{\mathbb{D}} \rightarrow \mathbb{C}$ defined by

$$Q(z) = z q'_\beta(z) \psi(q_\beta(z)) = \beta z q'_\beta(z) = e^{e^z-1} - 1$$

is starlike in \mathbb{D} . Also, note that by analytic characterization of starlike functions, the function $h : \overline{\mathbb{D}} \rightarrow \mathbb{C}$ defined by $h(z) := v(q_\beta(z)) + Q(z)$ satisfies the inequality

$$\operatorname{Re} \left(\frac{z h'(z)}{Q(z)} \right) = \operatorname{Re} \left(\frac{z Q'(z)}{Q(z)} \right) > 0.$$

Therefore, the subordination $1 + \beta z p'(z) \prec 1 + \beta z q'_\beta(z)$ implies $p \prec q_\beta$ by Lemma 2.1. For suitable $\mathcal{P}(z)$, as $r \rightarrow 1$, $q_\beta(z) \prec \mathcal{P}(z)$ holds if the following inequalities holds:

$$\mathcal{P}(-1) < q_\beta(-1) < q_\beta(1) < \mathcal{P}(1). \tag{2.2}$$

By the transitivity property, the required subordination $p(z) \prec \mathcal{P}(z)$ holds if $q_\beta(z) \prec \mathcal{P}(z)$. The condition (2.2) turns out to be both necessary and sufficient for the subordination $p \prec \mathcal{P}$ to hold.

- (a) Consider $\mathcal{P}(z) = \phi_q(z)$. Then the inequalities $q_\beta(-1) > -1 + \sqrt{2}$ and $q_\beta(1) < 1 + \sqrt{2}$ reduce to $\beta \geq \beta_1$ and $\beta \geq \beta_2$, where

$$\beta_1 = \frac{1}{2 - \sqrt{2}}\mathcal{L} \quad \text{and} \quad \beta_2 = \frac{1}{\sqrt{2}}\mathfrak{U}$$

respectively. Thus, the subordination $q_\beta \prec \phi_q$ holds whenever $\beta \geq \max\{\beta_1, \beta_2\} = \beta_2$.

- (b) For $\mathcal{P}(z) = \mathcal{Q}(z)$, the inequalities $q_\beta(-1) > \mathcal{Q}(-1)$ and $q_\beta(1) < \mathcal{Q}(1)$ give $\beta \geq \beta_1$ and $\beta \geq \beta_2$, where

$$\beta_1 = \frac{1}{1 - e^{e^{-1}-1}}\mathcal{L} \quad \text{and} \quad \beta_2 = \frac{1}{e^{e^{-1}-1} - 1}\mathfrak{U}$$

respectively. Therefore, $q_\beta \prec \mathcal{Q}$ whenever $\beta \geq \max\{\beta_1, \beta_2\} = \beta_1$.

- (c) On taking $\mathcal{P}(z) = \phi_c(z)$, a simple calculation shows that the inequalities $q_\beta(-1) > \phi_c(-1)$ and $q_\beta(1) < \phi_c(1)$ give $\beta \geq \beta_1$ and $\beta \geq \beta_2$, where $\beta_1 = (3/2)\mathcal{L}$ and $\beta_2 = (1/2)\mathfrak{U}$ respectively. Therefore, $q_\beta \prec \phi_c$ holds whenever $\beta \geq \max\{\beta_1, \beta_2\} = \beta_2$.
- (d) On substituting $\mathcal{P}(z) = \phi_0(z)$, the inequalities $q_\beta(-1) > \phi_0(-1)$ and $q_\beta(1) < \phi_0(1)$ give $\beta \geq \beta_1$ and $\beta \geq \beta_2$, where $\beta_1 = (1/(3 - 2\sqrt{2}))\mathcal{L}$ and $\beta_2 = \mathfrak{U}$ respectively. Therefore, the subordination $q_\beta \prec \phi_0$ holds if $\beta \geq \max\{\beta_1, \beta_2\} = \beta_1$.
- (e) Take $\mathcal{P}(z) = \phi_{\lim}(z)$. Then the inequalities $q_\beta(-1) > \frac{3}{2} - \sqrt{2}$ and $q_\beta(1) < \frac{3}{2} + \sqrt{2}$ reduce to $\beta \geq \beta_1$ and $\beta \geq \beta_2$, where $\beta_1 = (2/(2\sqrt{2} - 1))\mathcal{L}$ and $\beta_2 = 2/(2\sqrt{2} + 1)\mathfrak{U}$ respectively. Thus, the required subordination $q_\beta \prec \phi_{\lim}$ holds if $\beta \geq \max\{\beta_1, \beta_2\} = \beta_2$.
- (f) Take $\mathcal{P}(z) = \phi_s(z)$. Then the inequalities $q_\beta(-1) > 1 + \sin(-1)$ and $q_\beta(1) < 1 + \sin(1)$ give $\beta \geq \beta_1$ and $\beta \geq \beta_2$, where $\beta_1 = \mathcal{L}/\sin 1$ and $\beta_2 = \mathfrak{U}/\sin 1$ respectively. This shows that the subordination $q_\beta \prec \phi_s$ holds if $\beta \geq \max\{\beta_1, \beta_2\} = \beta_2$.
- (g) Set $\mathcal{P}(z) = 2/(1 + e^{-z})$. Then $q_\beta(-1) > 2/(e + 1)$ and $q_\beta(1) < 2e/(e + 1)$ gives

$$\beta_1 = \frac{e + 1}{e - 1}\mathcal{L} \quad \text{and} \quad \beta_2 = \frac{e + 1}{e - 1}\mathfrak{U}.$$

Hence, the subordination holds true for $\beta \geq \beta_2$ since $\max\{\beta_1, \beta_2\} = \beta_2$.




FIGURE 1. Sharpness for the case (a) and (b).

□

As an application of Theorem 2.2, we have the following sufficient conditions for starlikeness:

Corollary 2.3. Set $\mathfrak{M}(z) := 1 - zf'(z)/f(z) + zf''(z)/f'(z)$. If the function $f \in \mathcal{A}$ satisfies $1 + \beta \frac{zf'(z)}{f(z)}\mathfrak{M}(z) \prec \mathcal{Q}(z)$, then

- (a) $f \in \mathcal{S}_q^*$ if $\beta \geq (1/\sqrt{2})\mathfrak{U}$,
- (b) $f \in \mathcal{S}_B^*$ if $\beta \geq \left(1/(1 - e^{(e^{-1}-1)})\right)\mathcal{L}$,
- (c) $f \in \mathcal{S}_c^*$ if $\beta \geq (1/2)\mathfrak{U}$,
- (d) $f \in \mathcal{S}_R^*$ if $\beta \geq (3 + 2\sqrt{2})\mathcal{L}$,
- (e) $f \in \mathcal{S}_{LC}^*$ if $\beta \geq (2/(2\sqrt{2} + 1))\mathfrak{U}$,
- (f) $f \in \mathcal{S}_s^*$ if $\beta \geq (1/(\sin 1))\mathfrak{U}$
- (g) $f \in \mathcal{S}_{SG}^*$ if $\beta \geq ((e + 1)/(e - 1))\mathfrak{U}$,

where \mathfrak{U} and \mathcal{L} are given by (2.1).

Theorem 2.4. Let \mathfrak{U} and \mathcal{L} be given by (2.1) and $\mathcal{Q}(z)$ be given by (1.1). Let p be an analytic function in \mathbb{D} with $p(0) = 1$. If $\Lambda_\beta(z, p(z)) \prec \mathcal{Q}(z)$, then

- (a) $p(z) \prec \phi_q(z)$ for $\beta \geq \frac{1}{\log(1+\sqrt{2})}\mathfrak{U} \approx 2.40301$.
- (b) $p(z) \prec \mathcal{Q}(z)$ for $\beta \geq \frac{1}{e-1}\mathfrak{U} \approx 1.23260$.
- (c) $p(z) \prec \phi_c(z)$ for $\beta \geq \frac{1}{\log 3}\mathfrak{U} \approx 1.92784$.
- (d) $p(z) \prec \phi_0(z)$ for $\beta \geq \frac{1}{\log\left(\frac{1+\sqrt{2}}{2}\right)}\mathcal{L} \approx 3.59966$.
- (e) $p(z) \prec \phi_{\lim}(z)$ for $\beta \geq \frac{1}{\log(\sqrt{2}+3/2)}\mathfrak{U} \approx 1.98013$.
- (f) $p(z) \prec \phi_s(z)$ for $\beta \geq \frac{1}{\log(1+\sin 1)}\mathfrak{U} \approx 3.4688$.
- (g) $p(z) \prec \phi_{SG}(z)$ for $\beta \geq \frac{1}{1+\log 2-\log(1+e)}\mathfrak{U} \approx 5.57523$.

The bounds on β are best possible.

Proof. Consider the first order differential equation given by

$$1 + \beta \frac{z\check{q}'_\beta(z)}{\check{q}_\beta(z)} = e^{e^z-1}. \tag{2.3}$$

It is easy to verify that the analytic function $\check{q}_\beta : \mathbb{D} \rightarrow \mathbb{C}$ defined by

$$\check{q}_\beta(z) = \exp\left(\frac{1}{\beta} \int_0^z \frac{e^{e^t-1} - 1}{t} dt\right)$$

is a solution of differential equation (2.3). On taking $v(w) = 1$ and $\psi(w) = \beta/w$, the functions $Q, h : \mathbb{D} \rightarrow \mathbb{C}$ reduces to

$$Q(z) = z\check{q}'_\beta(z)\psi(\check{q}_\beta(z)) = \beta z\check{q}'_\beta(z)/\check{q}_\beta(z) = e^{e^z-1} - 1$$

and

$$h(z) = v(\check{q}_\beta(z)) + Q(z) = 1 + Q(z) = e^{e^z-1}.$$

It is seen that the function Q is starlike and $\text{Re}(zh'(z)/Q(z)) > 0$, for $z \in \mathbb{D}$. Hence,

$$1 + \beta \frac{zp'(z)}{p(z)} \prec 1 + \beta \frac{z\check{q}'_\beta(z)}{\check{q}_\beta(z)} \quad \text{implies} \quad p(z) \prec \check{q}_\beta(z)$$

which follows from Lemma 2.1. Proceeding as Theorem in 2.2, proof is completed. □

Theorem 2.5. Let \mathfrak{U} and \mathcal{L} be given by (2.1) and $\mathcal{Q}(z)$ be given by (1.1). Assume p to be an analytic function in \mathbb{D} with $p(0) = 1$. If $\Theta_\beta(z, p(z)) \prec \mathcal{Q}(z)$, then each of the following subordination holds:

- (a) $p(z) \prec \phi_q(z)$ for $\beta \geq \frac{1}{2-\sqrt{2}}\mathfrak{U} \approx 3.61556$.
- (b) $p(z) \prec \mathcal{Q}(z)$ for $\beta \geq \frac{e^e-1}{e^{e-1}-1}\mathfrak{U} \approx 2.58089$.
- (c) $p(z) \prec \phi_c(z)$ for $\beta \geq \frac{3}{2}\mathfrak{U} \approx 3.17692$.
- (d) $p(z) \prec \phi_0(z)$ for $\beta \geq 2\mathfrak{U} \approx 4.2359$.
- (e) $p(z) \prec \phi_{\lim}(z)$ for $\beta \geq \frac{5+4\sqrt{2}}{7}\mathfrak{U} \approx 3.22438$.
- (f) $p(z) \prec \phi_s(z)$ for $\beta \geq \frac{1+\sin 1}{\sin 1}\mathfrak{U} \approx 4.63491$.
- (g) $p(z) \prec \phi_{SG}(z)$ for $\beta \geq \frac{2e}{e-1}\mathfrak{U} \approx 6.7011$.

The estimates on β cannot be improved further.

Proof. The function

$$\hat{q}_\beta(z) = \left(1 - \frac{1}{\beta} \int_0^z \frac{e^{e^t-1} - 1}{t} dt\right)^{-1}$$

is the analytic solution of the differential equation

$$\beta \frac{z\hat{q}'_\beta(z)}{\hat{q}_\beta^2(z)} = e^{e^z-1} - 1.$$

Consider the functions $v(w) = 1$ and $\psi(w) = \beta/w^2$. Moreover, the function $Q(z) = z\hat{q}'_\beta(z)\psi(\hat{q}_\beta(z)) = e^{e^z-1} - 1$ is starlike in \mathbb{D} . Simple computation shows that the function $h(z) := 1 + Q(z)$ satisfies the inequality $\operatorname{Re}(zh'(z)/Q(z)) > 0$, ($z \in \mathbb{D}$). Now, by Lemma 2.1, we see that the subordination

$$1 + \beta \frac{zp'(z)}{p^2(z)} \prec 1 + \beta \frac{z\hat{q}'_\beta(z)}{\hat{q}_\beta^2(z)}$$

implies $p(z) \prec \hat{q}_\beta(z)$. Proceeding as in Theorem 2.2, we conclude the proof. \square

Theorem 2.6. Let ϕ_{SG} be given by (1.2) and further

$$I_- = \int_{-1}^0 \frac{e^t - 1}{t(e^t + 1)} dt \quad \text{and} \quad I_+ = \int_0^1 \frac{e^t - 1}{t(e^t + 1)} dt. \quad (2.4)$$

Assume p to be an analytic function in \mathbb{D} with $p(0) = 1$. If the subordination

$$\Psi_\beta(z, p(z)) \prec \phi_{SG}(z)$$

holds, then each of the following subordination inclusion hold:

- (a) $p(z) \prec \phi_q(z)$ for $\beta \geq \frac{1}{2-\sqrt{2}}I_- \approx 0.83117$.
- (b) $p(z) \prec \phi_c(z)$ for $\beta \geq \frac{3}{2}I_- \approx 0.730335$.
- (c) $p(z) \prec \phi_0(z)$ for $\beta \geq (3 + 2\sqrt{2})I_- \approx 2.837797$.
- (d) $p(z) \prec \mathcal{Q}(z)$ for $\beta \geq \frac{1}{1-e^{e^{-1}-1}}I_- \approx 1.039170$.
- (e) $p(z) \prec \phi_{\lim}(z)$ for $\beta \geq \frac{2}{2\sqrt{2}-1}I_- \approx 0.53257$.
- (f) $p(z) \prec \phi_s(z)$ for $\beta \geq \frac{1}{\sin 1}I_- \approx 0.578616$.
- (g) $p(z) \prec \phi_{SG}(z)$ for $\beta \geq \frac{e+1}{e-1}I_- \approx 1.05361$.

The bounds in each of the above case are sharp.

Proof. Consider the functions v and ψ defined as in Theorem 2.2. Define the function $q_\beta : \overline{\mathbb{D}} \rightarrow \mathbb{C}$ by

$$q_\beta(z) = 1 + \frac{1}{\beta} \int_0^z \frac{e^t - 1}{t(e^t + 1)} dt$$

Note that the function $q_\beta(z)$ is analytic solution of the differential equation $1 + \beta zq'_\beta(z) = 2/(1 + e^{-z})$. The function $Q(z) = zq'_\beta(z)\psi(q_\beta(z)) = (e^z - 1)/(e^z + 1)$ is starlike in \mathbb{D} and $h(z) = 1 + Q(z)$ satisfies the inequality $\operatorname{Re}(zh'(z)/Q(z)) > 0$, $z \in \mathbb{D}$. Thus, applying Lemma 2.1, it follows that the subordination $1 + \beta zp'(z) \prec 1 + \beta zq'_\beta(z)$ implies $p(z) \prec$

$q_\beta(z)$. Each of the subordination $p(z) \prec \mathcal{P}(z)$, for appropriate \mathcal{P} , from (a) to (g) holds if $q_\beta(z) \prec \mathcal{P}(z)$ holds. This subordination holds provided

$$\mathcal{P}(-1) < q_\beta(-1) < q_\beta(1) < \mathcal{P}(1).$$

These inequalities yield necessary and sufficient condition for the required subordination.

- (a) Take $\mathcal{P}(z) = \phi_q(z)$. Then, the inequalities $q_\beta(-1) > -1 + \sqrt{2}$ and $q_\beta(1) < 1 + \sqrt{2}$ reduce to $\beta \geq \beta_1$ and $\beta \geq \beta_2$, where

$$\beta_1 = \frac{1}{2 - \sqrt{2}}I_- \quad \text{and} \quad \beta_2 = \frac{1}{\sqrt{2}}I_+$$

respectively. Therefore, $q_\beta \prec \phi_q$ whenever $\beta \geq \max\{\beta_1, \beta_2\} = \beta_1$.

- (b) Consider $\mathcal{P}(z) = \phi_c(z)$. A simple calculation shows that the inequalities $q_\beta(-1) > \phi_c(-1)$ and $q_\beta(1) < \phi_c(1)$ gives $\beta \geq \beta_1$ and $\beta \geq \beta_2$, where

$$\beta_1 = \frac{3}{2}I_- \quad \text{and} \quad \beta_2 = \frac{1}{2}I_+$$

respectively. Therefore, the subordination $q_\beta \prec \phi_c$ holds if $\beta \geq \max\{\beta_1, \beta_2\} = \beta_1$.

- (c) On taking $\mathcal{P}(z) = \phi_0(z)$, the inequalities $q_\beta(-1) > \phi_0(-1)$ and $q_\beta(1) < \phi_0(1)$ give $\beta \geq \beta_1$ and $\beta \geq \beta_2$, where

$$\beta_1 = \frac{1}{3 - 2\sqrt{2}}I_- \quad \text{and} \quad \beta_2 = I_+$$

respectively. Therefore, $q_\beta \prec \phi_0$ if $\beta \geq \max\{\beta_1, \beta_2\} = \beta_1$.

- (d) Consider $\mathcal{P}(z) = \mathcal{Q}(z)$. From the inequalities $q_\beta(-1) > \mathcal{Q}(-1)$ and $q_\beta(1) < \mathcal{Q}(1)$, we note that $\beta \geq \beta_1$ and $\beta \geq \beta_2$, where

$$\beta_1 = \frac{1}{1 - e^{e^{-1}-1}}I_- \quad \text{and} \quad \beta_2 = \frac{1}{e^{e^{-1}-1} - 1}I_+$$

respectively. Thus, $q_\beta \prec \mathcal{Q}$ if $\beta \geq \max\{\beta_1, \beta_2\}$.

- (e) Take $\mathcal{P}(z) = \phi_{\lim}(z)$. Then, the inequalities $q_\beta(-1) > \frac{3}{2} - \sqrt{2}$ and $q_\beta(1) < \frac{3}{2} + \sqrt{2}$ reduce to $\beta \geq \beta_1$ and $\beta \geq \beta_2$, where

$$\beta_1 = \frac{2}{2\sqrt{2} - 1}I_- \quad \text{and} \quad \beta_2 = \frac{2}{2\sqrt{2} + 1}I_+$$

respectively. Thus, $q_\beta \prec \phi_{\lim}$ whenever $\beta \geq \max\{\beta_1, \beta_2\} = \beta_1$.

- (f) Take $\mathcal{P}(z) = \phi_s(z)$. Then, the inequalities $q_\beta(-1) > 1 + \sin(-1)$ and $q_\beta(1) < 1 + \sin(1)$ give $\beta \geq \beta_1$ and $\beta \geq \beta_2$, where

$$\beta_1 = \frac{1}{\sin 1}I_- \quad \text{and} \quad \beta_2 = \frac{1}{\sin 1}I_+$$

respectively. Therefore, the subordination $q_\beta \prec \phi_s$ holds if $\beta \geq \max\{\beta_1, \beta_2\} = \beta_1$.

- (g) Let $\mathcal{P}(z) = \phi_{SG}(z)$. On simplifying the inequalities $q_\beta(-1) > 2/(e + 1)$ and $q_\beta(1) < 2e/(e + 1)$, we get β_1 and β_2 , where

$$\beta_1 = \frac{e + 1}{e - 1}I_- \quad \text{and} \quad \beta_2 = \frac{e + 1}{e - 1}I_+$$

respectively and thus, $q_\beta \prec \phi_{SG}$ whenever $\beta \geq \max\{\beta_1, \beta_2\} = \beta_1$.




FIGURE 2. Sharpness for the case (b) and (f).

□

As an application of Theorem 2.6, we have the following sufficient conditions for starlikeness:

Corollary 2.7. *Let $f \in \mathcal{A}$ be analytic function which satisfies*

$$1 + \beta \frac{zf'(z)}{f(z)} \mathfrak{M}(z) \prec \phi_{SG}(z).$$

Then,

- (a) $f \in \mathcal{S}_q^*$ if $\beta \geq (1/(2 - \sqrt{2})) I_-$,
- (b) $f \in \mathcal{S}_c^*$ if $\beta \geq (3/2) I_-$,
- (c) $f \in \mathcal{S}_R^*$ if $\beta \geq (3 + 2\sqrt{2}) I_-$,
- (d) $f \in \mathcal{S}_B^*$ if $\beta \geq (1/(1 - e^{e^{-1}-1})) I_-$,
- (e) $f \in \mathcal{S}_{LC}^*$ if $\beta \geq (2/(2\sqrt{2} - 1)) I_-$,
- (f) $f \in \mathcal{S}_s^*$ if $\beta \geq (1/(\sin 1)) I_-$,

where $\mathfrak{M}(z)$ is defined in Corollary 2.3.

Theorem 2.8. *Let I_+ and I_- be given by 2.4 and ϕ_{SG} be given by (1.2). Assume p to be an analytic function in \mathbb{D} with $p(0) = 1$. If $\Lambda_\beta(z, p(z)) \prec \phi_{SG}(z)$, then each of the following holds.*

- (a) $p(z) \prec \phi_q(z)$ for $\beta \geq \frac{1}{\log(1+\sqrt{2})} I_- \approx 0.55242$.
- (b) $p(z) \prec \phi_c(z)$ for $\beta \geq \frac{1}{\log 3} I_- \approx 0.443185$.
- (c) $p(z) \prec \phi_0(z)$ for $\beta \geq \frac{1}{\log\left(\frac{1+\sqrt{2}}{2}\right)} I_- \approx 2.58671$.
- (d) $p(z) \prec \mathcal{Q}(z)$ for $\beta \geq \frac{1}{1-e^{-1}} I_- \approx 0.77024$.
- (e) $p(z) \prec \phi_{\lim}(z)$ for $\beta \geq \frac{1}{\log(\sqrt{2}+3/2)} I_+ \approx 0.455206$.
- (f) $p(z) \prec \phi_s(z)$ for $\beta \geq \frac{1}{\log(1+\sin 1)} I_+ \approx 0.79744$.
- (g) $p(z) \prec \phi_{SG}(z)$ for $\beta \geq \frac{1}{1+\log 2 - \log(1+e)} I_+ \approx 1.28167$.

The estimates on β are best possible.

Proof. Let the functions v and ψ be defined as in Theorem 2.4. Define the analytic function $\check{q}_\beta : \mathbb{D} \rightarrow \mathbb{C}$ by

$$\check{q}_\beta(z) = \exp \left(\frac{1}{\beta} \int_0^z \frac{e^t - 1}{t(e^t + 1)} dt \right),$$

which satisfies the differential equation

$$\frac{d\check{q}'_\beta(z)}{dz} = \frac{1}{\beta z} \left(\frac{1 - e^{-z}}{1 + e^{-z}} \right) \check{q}_\beta(z).$$

Now, observe that the function $Q(z) = z\check{q}'_\beta(z)\psi(\check{q}_\beta(z)) = \frac{1-e^{-z}}{1+e^{-z}}$ is starlike in \mathbb{D} . Also, it can be easily seen that the function h defined by $h(z) := v(\check{q}_\beta(z)) + Q(z) = 1 + Q(z)$ satisfies the inequality $\operatorname{Re}(zh'(z)/Q(z)) > 0$, $z \in \mathbb{D}$. Therefore, the Lemma 2.1 states that the subordination $1 + \beta \frac{zp'(z)}{p(z)} \prec 1 + \beta \frac{z\check{q}'_\beta(z)}{\check{q}_\beta(z)}$ implies $p(z) \prec \check{q}_\beta(z)$. As in the proof of Theorem 2.6, we conclude the result. \square

Theorem 2.9. *Let I_+ and I_- be given by (2.4). Assume p to be an analytic function in \mathbb{D} with $p(0) = 1$. If $\Theta_\beta(z, p(z)) \prec \phi_{SG}(z)$, then*

- (a) $p(z) \prec \phi_q(z)$ for $\beta \geq \frac{1}{2-\sqrt{2}}I_+ \approx 0.83117$.
- (b) $p(z) \prec \phi_c(z)$ for $\beta \geq \frac{3}{2}I_+ \approx 0.73033$.
- (c) $p(z) \prec \phi_0(z)$ for $\beta \geq (2 + 2\sqrt{2})I_- \approx 2.35090$.
- (d) $p(z) \prec \mathcal{Q}(z)$ for $\beta \geq \frac{e^e-1}{e^{e-1}-1}I_+ \approx 0.59331$.
- (e) $p(z) \prec \phi_{\lim}(z)$ for $\beta \geq \frac{5+4\sqrt{2}}{7}I_+ \approx 0.74124$.
- (f) $p(z) \prec \phi_s(z)$ for $\beta \geq \frac{1+\sin 1}{\sin 1}I_+ \approx 1.06550$.
- (g) $p(z) \prec \phi_{SG}(z)$ for $\beta \geq \frac{2e}{e-1}I_+ \approx 1.54049$.

All these estimates are sharp.

Proof. The function $\hat{q}_\beta : \mathbb{D} \rightarrow \mathbb{C}$ defined by

$$\hat{q}_\beta(z) = \left(1 - \frac{1}{\beta} \int_0^z \frac{e^t - 1}{t(e^t + 1)} dt \right)^{-1}$$

is clearly analytic in \mathbb{D} . It is noted that the function $\hat{q}_\beta(z)$ is a solution of the differential equation

$$1 + \beta \frac{z\hat{q}'_\beta(z)}{\hat{q}_\beta^2(z)} = \frac{2}{1 + e^{-z}}.$$

We take the functions v and ψ as in Theorem 2.5. Note that the function Q defined by $Q(z) = z\hat{q}'_\beta(z)\psi(\hat{q}_\beta(z)) = (1 - e^{-z})/(1 + e^{-z})$ is starlike in \mathbb{D} and the function h defined as $h(z) := v(\hat{q}_\beta(z)) + Q(z) = 1 + Q(z)$ follows the inequality $\operatorname{Re}(zh'(z)/Q(z)) = \operatorname{Re}(zQ'(z)/Q(z)) > 0$. Therefore, as in view of Lemma 2.1, the subordination

$$1 + \beta \frac{zp'(z)}{p^2(z)} \prec 1 + \beta \frac{z\hat{q}'_\beta(z)}{\hat{q}_\beta^2(z)}$$

implies $p(z) \prec \hat{q}_\beta(z)$. Proceeding as in Theorem 2.6, proof is completed. \square

Theorem 2.10. *Let p be an analytic function in \mathbb{D} with $p(0) = 1$. Then each of the following subordination implies $p(z) \prec \mathcal{Q}(z) := e^{e^z-1}$:*

- (a) $\Psi_\beta(z, p(z)) \prec \phi_c(z)$ if $\beta \geq \frac{1}{1-e^{(e^{-1}-1)}} \approx 2.13430$.
- (b) $\Lambda_\beta(z, p(z)) \prec \phi_c(z)$ if $\beta \geq \frac{e}{e-1} \approx 1.581976$.
- (c) $\Theta_\beta(z, p(z)) \prec \phi_c(z)$ if $\beta \geq \frac{5e^{e-1}}{3(e^{e-1}-1)} \approx 2.030970$.

The bounds in each case are sharp.

Proof. (a) Define the analytic function $q_\beta : \mathbb{D} \rightarrow \mathbb{C}$ by

$$q_\beta(z) = 1 + \frac{1}{\beta} \left(\frac{4z}{3} + \frac{z^2}{3} \right)$$

It is easy to see that the function q_β satisfies the differential equation $\beta z q'(z) = \phi_c(z) - 1$. Proceeding as similar lines in Theorem 2.2, the required subordination holds if and only if,

$$e^{e^{-1}-1} < q_\beta(-1) < q_\beta(1) < e^{e-1}. \quad (2.5)$$

Simplifying the condition (2.5), we obtain the inequalities

$$\beta \geq \frac{1}{1-e^{(e^{-1}-1)}} = \beta_1 \quad \text{and} \quad \beta \geq \frac{5e}{3(e^e-e)} = \beta_2.$$

Thus, the required subordination holds if $\beta \geq \max\{\beta_1, \beta_2\} = \beta_1$.

(b) Define the analytic function $\check{q}_\beta(z)$ by,

$$\check{q}_\beta(z) = \exp \left(\frac{1}{\beta} \left(\frac{4z}{3} + \frac{z^2}{3} \right) \right)$$

which is a solution of the equation

$$\frac{d\check{q}'_\beta(z)}{dz} = \frac{2(2+z)}{3\beta} \check{q}_\beta(z).$$

Proceeding as similar lines in Theorem 2.4, the subordination $p(z) \prec e^{e^z-1}$ holds if $\beta \geq \max\{\check{\beta}_1, \check{\beta}_2\}$, where

$$\check{\beta}_1 = \frac{e}{e-1} \quad \text{and} \quad \check{\beta}_2 = \frac{5}{3(e-1)}$$

are obtained from the inequalities $\check{q}_\beta(-1) > e^{e^{-1}-1}$ and $\check{q}_\beta(1) < e^{e-1}$ respectively.

(c) The differential equation

$$\frac{d\hat{q}'_\beta(z)}{dz} = \frac{2(2+z)}{3\beta} \hat{q}_\beta^2(z)$$

has an analytic solution

$$\hat{q}_\beta(z) = \left(1 - \frac{1}{\beta} \left(\frac{4z}{3} + \frac{z^2}{3} \right) \right)^{-1}$$

in \mathbb{D} . Therefore, proceeding as in Theorem 2.5, the required subordination $p(z) \prec e^{e^z-1}$ holds if $\beta \geq \max\{\hat{\beta}_1, \hat{\beta}_2\} = \hat{\beta}_2$, where

$$\hat{\beta}_1 = \frac{e^{\frac{1}{e}-1}}{1-e^{\frac{1}{e}-1}} \quad \text{and} \quad \hat{\beta}_2 = \frac{5e^{e-1}}{3(e^{e-1}-1)}.$$

□

Corollary 2.11. Let $f \in \mathcal{A}$ be given by $f(z) = z + \sum_{n=2}^{\infty} a_n z^n$. If one of the following subordinations holds

(a) $1 + \beta \frac{zf'(z)}{f(z)} \mathfrak{M}(z) \prec \phi_c(z)$ for $\beta \geq \frac{1}{1-e^{(e^{-1}-1)}}$,

(b) $1 + \beta \mathfrak{M}(z) \prec \phi_c(z)$ for $\beta \geq \frac{e}{e-1}$,

(c) $1 + \beta \left(\frac{zf'(z)}{f(z)} \right)^{-1} \mathfrak{M}(z) \prec \phi_c(z)$ for $\beta \geq \frac{5e^{e-1}}{(3(e^{e-1}-1))}$,

then $f \in \mathcal{S}_B^*$, where $\mathfrak{M}(z)$ is defined in Corollary 2.3.

The next results provide best possible bound on β so that the subordination $1 + \beta zp'(z)/p^j(z) \prec \phi_c(z)$, $\phi_0(z)$ ($j = 0, 1, 2$) implies the subordination $p(z) \prec \phi_{SG}(z)$. Proofs of the following results are omitted as similar to the previous Theorem 2.10.

Theorem 2.12. Let p be an analytic function in \mathbb{D} with $p(0) = 1$. Then the following subordinations hold for $p(z) \prec \phi_{SG}(z) := 2/(1 + e^{-z})$.

(a) $\Psi_{\beta}(z, p(z)) \prec \phi_0(z)$ if $\beta \geq \frac{(e+1)(1-\sqrt{2}-2\log(2-\sqrt{2}))}{e-1} \approx 1.418226$.

(b) $\Lambda_{\beta}(z, p(z)) \prec \phi_0(z)$ if $\beta \geq \frac{1-\sqrt{2}-2\log(2-\sqrt{2})}{1+\log 2-\log(1+e)} \approx 1.725221$.

(c) $\Theta_{\beta}(z, p(z)) \prec \phi_0(z)$ if $\beta \geq \frac{2e(1-\sqrt{2}-2\log(2-\sqrt{2}))}{e-1} \approx 2.073612$.

The bounds on β in each case are sharp.

Theorem 2.13. Let p be an analytic function in \mathbb{D} which satisfies $p(0) = 1$. Then each of the following subordination is sufficient for $p(z) \prec \phi_{SG}(z)$.

(a) $\Psi_{\beta}(z, p(z)) \prec \phi_c(z)$ if $\beta \geq \frac{5(e+1)}{3(e-1)} \approx 3.60659$.

(b) $\Lambda_{\beta}(z, p(z)) \prec \phi_c(z)$ if $\beta \geq \frac{5}{3(1+\log 2-\log(1+e))} \approx 4.387286$.

(c) $\Theta_{\beta}(z, p(z)) \prec \phi_c(z)$ if $\beta \geq \frac{10e}{3(e-1)} \approx 5.27326$.

The bounds on β in each case are sharp.

REFERENCES

- [1] O. P. Ahuja, S. Kumar and V. Ravichandran, Applications of first order differential subordination for functions with positive real part, Stud. Univ. Babeş-Bolyai Math. **63** (2018), no. 3, 303–311.
- [2] R. M. Ali, V. Ravichandran and N. Seenivasagan, Sufficient conditions for Janowski starlikeness, Int. J. Math. Math. Sci. **2007**, Art. ID 62925, 7 pp.
- [3] N. Bohra, S. Kumar and V. Ravichandran, Some special differential subordinations, Hacet. J. Math. Stat. **48** (2019), no. 4, 1017–1034.
- [4] N. E. Cho, V. Kumar, S. S. Kumar and V. Ravichandran, Radius problems for starlike functions associated with the sine function, Bull. Iranian Math. Soc. **45** (2019), no. 1, 213–232.
- [5] N. E. Cho, S. Kumar, V. Kumar, V. Ravichandran, and H. M. Srivastava, Starlike functions related to the Bell numbers, Symmetry **11** (2019), Article ID 219, 1–17.
- [6] A. Ebadian, N. H. Mohammed, E. A. Adegani, and T. Bulboacă, New results for some generalizations of starlike and convex functions, J. Funct. Spaces **2020**, Art. ID 7428648, 12 pp.
- [7] S. Gandhi, S. Kumar and V. Ravichandran, First order differential subordinations for Carathéodory functions, Kyungpook Math. J. **58** (2018), no. 2, 257–270.
- [8] P. Goel and S. S. Kumar, Certain class of starlike functions associated with modified sigmoid function, Bull. Malays. Math. Sci. Soc. **43** (2020), no. 1, 957–991.
- [9] W. Janowski, Extremal problems for a family of functions with positive real part and for some related families, Ann. Polon. Math. **23** (1970/71), 159–177.
- [10] S. Kanas, Differential subordination related to conic sections, J. Math. Anal. Appl. **317** (2006), no. 2, 650–658.

- [11] R. Kargar, A. Ebadian and J. Sokół, Radius problems for some subclasses of analytic functions, *Complex Anal. Oper. Theory* **11** (2017), no. 7, 1639–1649.
- [12] S. Kumar and V. Ravichandran, A subclass of starlike functions associated with a rational function, *Southeast Asian Bull. Math.* **40** (2016), no. 2, 199–212.
- [13] S. Kumar and V. Ravichandran, Subordinations for functions with positive real part, *Complex Anal. Oper. Theory* **12** (2018), no. 5, 1179–1191.
- [14] W. C. Ma and D. Minda, A unified treatment of some special classes of univalent functions, in *Proceedings of the Conference on Complex Analysis (Tianjin, 1992)*, 157–169, Conf. Proc. Lecture Notes Anal., I, Int. Press, Cambridge, MA.
- [15] R. Mendiratta, S. Nagpal and V. Ravichandran, On a subclass of strongly starlike functions associated with exponential function, *Bull. Malays. Math. Sci. Soc.* **38** (2015), no. 1, 365–386.
- [16] S. S. Miller and P. T. Mocanu, On some classes of first-order differential subordinations, *Michigan Math. J.* **32** (1985), no. 2, 185–195.
- [17] S. S. Miller and P. T. Mocanu, *Differential subordinations*, Monographs and Textbooks in Pure and Applied Mathematics, 225, Marcel Dekker, Inc., New York, 2000.
- [18] M. Nunokawa, M. Obradović and S. Owa, One criterion for univalence, *Proc. Amer. Math. Soc.* **106** (1989), no. 4, 1035–1037.
- [19] M. Nunokawa, S. Owa, S. Saitoh and O. P. Ahuja, On the quotient of the analytic representations of convex and starlike functions, *Research Institute of Mathematical Analysis*, No. 1112 (1999), 63–69.
- [20] R. K. Raina and J. Sokół, On coefficient estimates for a certain class of starlike functions, *Hacet. J. Math. Stat.* **44** (2015), no. 6, 1427–1433.
- [21] K. Sharma, N. K. Jain and V. Ravichandran, Starlike functions associated with a cardioid, *Afr. Mat.* **27** (2016), no. 5-6, 923–939.
- [22] H. Silverman, Convex and starlike criteria, *Int. J. Math. Math. Sci.* **22** (1999), no. 1, 75–79.
- [23] Y. Yunus, S.A. Halim, A.B. Akbarally, Subclass of starlike functions associated with a limaçon. *AIP Conf. Proc.* 1974(1), 030023 (2018)

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF DELHI, DELHI-110 007, INDIA

Email address: meghnasharma203@gmail.com

BHARATI VIDYAPEETH'S COLLEGE OF ENGINEERING, DELHI-110063, INDIA

Email address: sushilkumar16n@gmail.com

DEPARTMENT OF MATHEMATICS, ARYABHATTA COLLEGE, DELHI-110021, INDIA

Email address: naveenjain05@gmail.com