

# A NOTE ON THE HIT PROBLEM FOR THE STEENROD ALGEBRA AND ITS APPLICATIONS

NGUYEN KHAC TIN

ABSTRACT. Let  $P_k = H^*((\mathbb{R}P^\infty)^k)$  be the modulo-2 cohomology algebra of the direct product of  $k$  copies of infinite dimensional real projective spaces  $\mathbb{R}P^\infty$ . Then,  $P_k$  is isomorphic to the graded polynomial algebra  $\mathbb{F}_2[x_1, \dots, x_k]$  of  $k$  variables, in which each  $x_j$  is of degree 1, and let  $GL_k$  be the general linear group over the prime field  $\mathbb{F}_2$  which acts naturally on  $P_k$ . Here the cohomology is taken with coefficients in the prime field  $\mathbb{F}_2$  of two elements. We study the *hit problem*, set up by Frank Peterson, of finding a minimal set of generators for the polynomial algebra  $P_k$  as a module over the mod-2 Steenrod algebra,  $\mathcal{A}$ .

In this Note, we explicitly compute the hit problem for  $k = 5$  and the degree  $5(2^s - 1) + 24.2^s$  with  $s$  an arbitrary non-negative integer. These results are used to study the Singer algebraic transfer which is a homomorphism from the homology of the mod-2 Steenrod algebra,  $\text{Tor}_{k, k+n}^{\mathcal{A}}(\mathbb{F}_2, \mathbb{F}_2)$ , to the subspace of  $\mathbb{F}_2 \otimes_{\mathcal{A}} P_k$  consisting of all the  $GL_k$ -invariant classes of degree  $n$ . We show that Singer's conjecture for the algebraic transfer is true in the case  $k = 5$  and the above degrees. This method is different from that of Singer in studying the image of the algebraic transfer. Moreover, as a consequence, we get the dimension results for polynomial algebra in some generic degrees in the case  $k = 6$ .

## 1. Introduction

Denote by  $P_k = H^*((\mathbb{R}P^\infty)^k)$  the modulo-2 cohomology algebra of the direct product of  $k$  copies of infinite dimensional real projective spaces  $\mathbb{R}P^\infty$ . Then,  $P_k$  is isomorphic to the graded polynomial algebra  $\mathbb{F}_2[x_1, x_2, \dots, x_k]$  of  $k$  variables, in which each  $x_j$  is of degree 1. Here the cohomology is taken with coefficients in the prime field  $\mathbb{F}_2$  of two elements.

Being the cohomology of a group,  $P_k$  is a module over the mod-2 Steenrod algebra,  $\mathcal{A}$ . The action of  $\mathcal{A}$  on  $P_k$  is determined by the elementary properties of the Steenrod squares  $Sq^i$  and the Cartan formula (see Steenrod and Epstein [18]).

A polynomial  $f$  in  $P_k$  is called *hit* if it can be written as a finite sum  $f = \sum_{u \geq 0} Sq^{2^u}(h_u)$  for suitable polynomials  $h_u$ . That means  $f$  belongs to  $\mathcal{A}^+ P_k$ , where  $\mathcal{A}^+$  denotes the augmentation ideal in  $\mathcal{A}$ .

Let  $GL_k$  be the general linear group over the field  $\mathbb{F}_2$ . This group acts naturally on  $P_k$  by matrix substitution. Since the two actions of  $\mathcal{A}$  and  $GL_k$  upon  $P_k$  commute with each other, there is an action of  $GL_k$  on  $\mathbb{F}_2 \otimes_{\mathcal{A}} P_k$ .

Many authors study the *hit problem* of determination of a minimal set of generators for  $P_k$  as a module over the Steenrod algebra, or equivalently, a basis of  $\mathbb{F}_2 \otimes_{\mathcal{A}} P_k$ . This problem has first been studied by Peterson [11], Wood [32], Singer

<sup>1</sup>2000 *Mathematics Subject Classification*. Primary 55S10; 55S05, 55T15.

<sup>2</sup>*Keywords and phrases*: Polynomial algebra, Steenrod squares, hit problem, Algebraic transfer.

[16], Priddy [15], who show its relationship to several classical problems in homotopy theory. Then, this problem was investigated by Nam [10], Silverman [17], Wood [32], Sum [19, 21, 22], Tin-Sum [24], Tin [26, 27, 29] and others.

For a positive integer  $n$ , by  $\mu(n)$  one means the smallest number  $r$  for which it is possible to write  $n = \sum_{1 \leq i \leq r} (2^{u_i} - 1)$ , where  $u_i > 0$ . This result implies a result of Wood, which originally is a conjecture of Peterson [11].

**Theorem 1.1** (Wood [32]). *If  $\mu(n) > k$ , then  $(\mathbb{F}_2 \otimes_{\mathcal{A}} P_k)_n = 0$ .*

From the above result of Wood, the hit problem is reduced to the case of degree  $n$  with  $\mu(n) \leq k$ .

Let  $GL_k$  be the general linear group over the field  $\mathbb{F}_2$ . This group acts naturally on  $P_k$  by matrix substitution. Since the two actions of  $\mathcal{A}$  and  $GL_k$  upon  $P_k$  commute with each other, there is an action of  $GL_k$  on  $\mathbb{F}_2 \otimes_{\mathcal{A}} P_k$ .

For a nonnegative integer  $d$ , denote by  $(P_k)_d$  the subspace of  $P_k$  consisting of all the homogeneous polynomials of degree  $d$  in  $P_k$  and by  $(\mathbb{F}_2 \otimes_{\mathcal{A}} P_k)_d$  the subspace of  $QP_k$  consisting of all the classes represented by the elements in  $(P_k)_d$ . One of the major applications of hit problem is in surveying a homomorphism introduced by Singer. In [16], Singer defined the algebraic transfer, which is a homomorphism

$$\varphi_k : \text{Tor}_{k, k+d}^{\mathcal{A}}(\mathbb{F}_2, \mathbb{F}_2) \longrightarrow (\mathbb{F}_2 \otimes_{\mathcal{A}} P_k)_d^{GL_k}$$

from the homology of the Steenrod algebra to the subspace of  $(\mathbb{F}_2 \otimes_{\mathcal{A}} P_k)_d$  consisting of all the  $GL_k$ -invariant classes. It is a useful tool in describing the homology groups of the Steenrod algebra,  $\text{Tor}_{k, k+d}^{\mathcal{A}}(\mathbb{F}_2, \mathbb{F}_2)$ . The hit problem and the algebraic transfer were studied by many authors (see Boardman [1], Hưng [5], Chơn and Hà [2], Nam [10], Phuc [14], Phuc and Sum [12], Sum [19, 20, 21], Sum and Tín [23] and others).

Singer showed in [16] that  $\varphi_k$  is an isomorphism for  $k = 1, 2$ . Boardman showed in [1] that  $\varphi_3$  is also an isomorphism. However, for any  $k \geq 4$ ,  $\varphi_k$  is not a monomorphism in infinitely many degrees (see Singer [16], Hưng [5].) Singer made the following conjecture.

**Conjecture 1.2** (Singer [16]). *The algebraic transfer  $\varphi_k$  is an epimorphism for any  $k \geq 0$ .*

The conjecture is true for  $k \leq 3$ . However, for  $k > 3$ , it is still open. Recently, the hit problem and its applications to representations of general linear groups have been presented in the books of Walker and Wood [30, 31].

One of the extremely useful tools for computing the hit problem and studying Singer's transfer is the Kameko squaring operation

$$\widetilde{Sq}_*^0 := (\widetilde{Sq}_*^0)_{(k, d)} : (\mathbb{F}_2 \otimes_{\mathcal{A}} P_k)_{2k+d} \rightarrow (\mathbb{F}_2 \otimes_{\mathcal{A}} P_k)_d,$$

which is induced by an  $\mathbb{F}_2$ -linear map  $\varphi_k : P_k \rightarrow P_k$ , given by

$$\varphi_k(x) = \begin{cases} y, & \text{if } x = x_1 x_2 \dots x_k y^2, \\ 0, & \text{otherwise,} \end{cases}$$

for any monomial  $x \in P_k$ . The map  $\varphi_k$  is not an  $\mathcal{A}$ -homomorphism. However,  $\varphi_k Sq^{2i} = Sq^i \varphi_k$  and  $\varphi_k Sq^{2i+1} = 0$  for any non-negative integer  $i$ .

**Theorem 1.3** (Kameko [6]). *Let  $d$  be a non-negative integer. If  $\mu(2d + k) = k$ , then*

$$(\widetilde{Sq}_*)^0_{(k,d)} : (\mathbb{F}_2 \otimes_{\mathcal{A}} P_k)_{2d+k} \longrightarrow (\mathbb{F}_2 \otimes_{\mathcal{A}} P_k)_d$$

*is an isomorphism of  $GL_k$ -modules.*

Thus, the hit problem is reduced to the case of degree  $n$  of the form

$$n = a(2^s - 1) + 2^s b,$$

where  $a, b, m$  are non-negative integers such that  $0 \leq \mu(b) < a \leq k$ .

Now, the  $\mathbb{F}_2$ -vector space  $\mathbb{F}_2 \otimes_{\mathcal{A}} P_k$  was explicitly calculated by Peterson [11] for  $k = 1, 2$ , by Kameko [6] for  $k = 3$  and by Sum [21] for  $k = 4$ . However, for  $k > 4$ , it is still unsolved, even in the case of  $k = 5$  with the help of computers.

For  $a = k - 1 = 4$  and  $b = 0$ , the vector space  $(\mathbb{F}_2 \otimes_{\mathcal{A}} P_5)_n$  is explicitly computed by Phuc and Sum [12], and in the case  $a = k - 2 = 3, b = 1$  by Phuc [13] and by the present author [28] in the case  $a = k - 1 = 4, b \in \{8; 10; 11\}$  with  $s$  an arbitrary non-negative integer.

In this Note, we explicitly compute the hit problem for  $k = 5$  and the degree  $5(2^s - 1) + 24 \cdot 2^s$  with  $s$  an arbitrary non-negative integer. Using this result, we show that Singer's conjecture for the algebraic transfer is true in the case  $k = 5$  and the above degrees. Moreover, as a consequence, we get the dimension results for polynomial algebra in some generic degrees in the case  $k = 6$ .

## 2. Preliminaries

In this section, we recall some needed information from Kameko [6], Singer [16] and Sum [19], which will be used in the next section.

**Notation 2.1.** We denote  $\mathbb{N}_k = \{1, 2, \dots, k\}$  and

$$X_{\mathbb{J}} = X_{\{j_1, j_2, \dots, j_s\}} = \prod_{j \in \mathbb{N}_k \setminus \mathbb{J}} x_j, \quad \mathbb{J} = \{j_1, j_2, \dots, j_s\} \subset \mathbb{N}_k,$$

Let  $\alpha_i(a)$  denote the  $i$ -th coefficient in dyadic expansion of a non-negative integer  $a$ . That means  $a = \alpha_0(a)2^0 + \alpha_1(a)2^1 + \alpha_2(a)2^2 + \dots$ , for  $\alpha_i(a) = 0$  or  $1$  with  $i \geq 0$ .

Let  $x = x_1^{\alpha_1} x_2^{\alpha_2} \dots x_k^{\alpha_k} \in P_k$ . Denote  $\nu_j(x) = \alpha_j, 1 \leq j \leq k$ . Set

$$\mathbb{J}_t(x) = \{j \in \mathbb{N}_k : \alpha_t(\nu_j(x)) = 0\},$$

for  $t \geq 0$ . Then, we have  $x = \prod_{t \geq 0} X_{\mathbb{J}_t(x)}^{2^t}$ .

**Definition 2.2.** For a monomial  $x$  in  $P_k$ , define two sequences associated with  $x$  by

$$\omega(x) = (\omega_1(x), \omega_2(x), \dots, \omega_i(x), \dots), \quad \sigma(x) = (\nu_1(x), \nu_2(x), \dots, \nu_k(x)),$$

where  $\omega_i(x) = \sum_{1 \leq j \leq k} \alpha_{i-1}(\nu_j(x)) = \deg X_{\mathbb{J}_{i-1}(x)}$ ,  $i \geq 1$ . The sequence  $\omega(x)$  is called the weight vector of  $x$ .

Let  $\omega = (\omega_1, \omega_2, \dots, \omega_i, \dots)$  be a sequence of non-negative integers. The sequence  $\omega$  is called the weight vector if  $\omega_i = 0$  for  $i \gg 0$ .

The sets of all the weight vectors and the exponent vectors are given the left lexicographical order.

For a weight vector  $\omega$ , we define  $\deg \omega = \sum_{i > 0} 2^{i-1} \omega_i$ . Denote by  $P_k(\omega)$  the subspace of  $P_k$  spanned by all monomials  $y$  such that  $\deg y = \deg \omega$ ,  $\omega(y) \leq \omega$ ,

and by  $P_k^-(\omega)$  the subspace of  $P_k$  spanned by all monomials  $y \in P_k(\omega)$  such that  $\omega(y) < \omega$ .

**Definition 2.3.** Let  $\omega$  be a weight vector and  $f, g$  two polynomials of the same degree in  $P_k$ .

- i)  $f \equiv g$  if and only if  $f - g \in \mathcal{A}^+ P_k$ . If  $f \equiv 0$  then  $f$  is called *hit*.
- ii)  $f \equiv_\omega g$  if and only if  $f - g \in \mathcal{A}^+ P_k + P_k^-(\omega)$ .

Obviously, the relations  $\equiv$  and  $\equiv_\omega$  are equivalence ones. Denote by  $QP_k(\omega)$  the quotient of  $P_k(\omega)$  by the equivalence relation  $\equiv_\omega$ . Then, we have

$$QP_k(\omega) = P_k(\omega) / ((\mathcal{A}^+ P_k \cap P_k(\omega)) + P_k^-(\omega)).$$

For a polynomial  $f \in P_k$ , we denote by  $[f]$  the class in  $QP_k$  represented by  $f$ . If  $\omega$  is a weight vector, then denote by  $[f]_\omega$  the class represented by  $f$ . Denote by  $|S|$  the cardinal of a set  $S$ .

**Definition 2.4.** Let  $x, y$  be monomials of the same degree in  $P_k$ . We say that  $x < y$  if and only if one of the following holds:

- i)  $\omega(x) < \omega(y)$ ;
- ii)  $\omega(x) = \omega(y)$  and  $\sigma(x) < \sigma(y)$ .

**Definition 2.5.** A monomial  $x$  is said to be inadmissible if there exist monomials  $y_1, y_2, \dots, y_m$  such that  $y_t < x$  for  $t = 1, 2, \dots, m$  and  $x - \sum_{t=1}^m y_t \in \mathcal{A}^+ P_k$ .

A monomial  $x$  is said to be admissible if it is not inadmissible.

Obviously, the set of all the admissible monomials of degree  $n$  in  $P_k$  is a minimal set of  $\mathcal{A}$ -generators for  $P_k$  in degree  $n$ .

Now, we recall some notations and definitions in [21], which will be used in the next sections. We set

$$\begin{aligned} P_k^0 &= \langle \{x = x_1^{a_1} x_2^{a_2} \dots x_k^{a_k} : a_1 a_2 \dots a_k = 0\} \rangle, \\ P_k^+ &= \langle \{x = x_1^{a_1} x_2^{a_2} \dots x_k^{a_k} : a_1 a_2 \dots a_k > 0\} \rangle. \end{aligned}$$

It is easy to see that  $P_k^0$  and  $P_k^+$  are the  $\mathcal{A}$ -submodules of  $P_k$ . Furthermore, we have the following.

**Proposition 2.6.** *We have a direct summand decomposition of the  $\mathbb{F}_2$ -vector spaces  $\mathbb{F}_2 \otimes_{\mathcal{A}} P_k = QP_k^0 \oplus QP_k^+$ . Here  $QP_k^0 = \mathbb{F}_2 \otimes_{\mathcal{A}} P_k^0$  and  $QP_k^+ = \mathbb{F}_2 \otimes_{\mathcal{A}} P_k^+$ .*

**Definition 2.7.** For any  $1 \leq i \leq k$ , define the homomorphism  $f_i : P_{k-1} \rightarrow P_k$  of algebras by substituting

$$f_i(x_j) = \begin{cases} x_j, & \text{if } 1 \leq j < i, \\ x_{j+1}, & \text{if } i \leq j < k. \end{cases}$$

Then,  $f_i$  is a homomorphism of  $\mathcal{A}$ -modules.

For a subset  $B \subset P_k$ , we denote  $[B] = \{[f] : f \in B\}$ . Obviously, we have

**Proposition 2.8.** *It is easy to see that if  $B$  is a minimal set of generators for  $\mathcal{A}$ -module  $P_{k-1}$  in degree  $n$ , then  $f(B) = \bigcup_{i=1}^k f_i(B)$  is a minimal set of generators for  $\mathcal{A}$ -module  $P_k^0$  in degree  $n$ .*

From now on, we denote by  $B_k(n)$  the set of all admissible monomials of degree  $n$  in  $P_k$ ,  $B_k^0(n) = B_k(n) \cap P_k^0$ ,  $B_k^+(n) = B_k(n) \cap P_k^+$ . For a weight vector  $\omega$  of degree  $n$ , we set  $B_k(\omega) = B_k(n) \cap P_k(\omega)$ ,  $B_k^+(\omega) = B_k^+(n) \cap P_k(\omega)$ .

Then,  $[B_k(\omega)]_\omega$  and  $[B_k^+(\omega)]_\omega$ , are respectively the basses of the  $\mathbb{F}_2$ -vector spaces  $QP_k(\omega)$  and  $QP_k^+(\omega) := QP_k(\omega) \cap QP_k^+$ .

### 3. The Main Results

We first recall a result in [24] the following: Let  $d$  be an arbitrary non-negative integer. Set  $t(k, d) = \max\{0, k - \alpha(d + k) - \zeta(d + k)\}$ , where  $\zeta(n)$  the greatest integer  $u$  such that  $n$  is divisible by  $2^u$ , that means  $n = 2^{\zeta(n)}m$ , with  $m$  an odd integer.

**Theorem 3.1** (Tin-Sum [24]). *Let  $d$  be an arbitrary non-negative integer. Then*

$$(\widetilde{S}q_*^0)^{s-t} : (\mathbb{F}_2 \otimes_{\mathcal{A}} P_k)_{k(2^s-1)+2^s d} \longrightarrow (\mathbb{F}_2 \otimes_{\mathcal{A}} P_k)_{k(2^t-1)+2^t d}$$

*is an isomorphism of  $GL_k$ -modules for every  $s \geq t$  if and only if  $t \geq t(k, d)$ .*

It is easy to check that for  $k = 5$  and  $d = 53$  then

$$t(k, d) = \max\{0, k - \alpha(d + k) - \zeta(d + k)\} = 0.$$

Using the above theorem, we get  $(\mathbb{F}_2 \otimes_{\mathcal{A}} P_5)_{5(2^s-1)+2^s 53} \cong (\mathbb{F}_2 \otimes_{\mathcal{A}} P_5)_{5(2^0-1)+2^0 53}$  for all  $s \geq 0$ .

Therefore, we need only to study  $(\mathbb{F}_2 \otimes_{\mathcal{A}} P_5)_{5(2^s-1)+24 \cdot 2^s}$  for  $s = 0$  and  $s = 1$ .

#### Case $s=0$ .

Denote  $\omega_{(1)} = (4, 4, 3)$ ,  $\omega_{(2)} = (4, 4, 1, 1)$ ,  $\omega_{(3)} = (4, 2, 2, 1)$ ,  $\omega_{(4)} = (4, 2, 4)$ . We give a direct summand decomposition of the  $\mathbb{F}_2$ -vector spaces  $(\mathbb{F}_2 \otimes_{\mathcal{A}} P_5)_{5(2^0-1)+24 \cdot 2^0}$  as follows:

**Theorem 3.2.** *We have a direct summand decomposition of the  $\mathbb{F}_2$ -vector spaces*

$$(\mathbb{F}_2 \otimes_{\mathcal{A}} P_5)_{24} = (QP_5^0)_{24} \oplus QP_5^+(\omega_{(1)}) \oplus QP_5^+(\omega_{(2)}) \oplus QP_5^+(\omega_{(3)}) \oplus QP_5^+(\omega_{(4)}).$$

Recall that  $(\mathbb{F}_2 \otimes_{\mathcal{A}} P_4)_{10}$  is an  $\mathbb{F}_2$ -vector space of dimension 70 with a basis consisting of all the classes represented by the monomials  $w_j$ ,  $1 \leq j \leq 70$ . Consequently,  $|B_4(10)| = 70$ , (see Sum [21]).

Since  $\mu(24) = 4$ , Theorem 1.3 implies that the squaring operation

$$\widetilde{S}q_*^0 : (\mathbb{F}_2 \otimes_{\mathcal{A}} P_4)_{24} \longrightarrow (\mathbb{F}_2 \otimes_{\mathcal{A}} P_4)_{10}$$

is an isomorphism of  $GL_4$ -module. Hence,  $|B_4(24)| = |B_4(10)| = 70$ , and therefore the set  $[B_4(24)] = \{[\phi_4(w_j)] : w_j \in B_4(10)\}$  is a basis of the  $\mathbb{F}_2$ -vector space  $(\mathbb{F}_2 \otimes_{\mathcal{A}} P_4)_{24}$ , where  $\phi_4(u) = x_1 x_2 x_3 x_4 u^2$ .

Using Proposition 2.8, we obtain  $[B_5^0(24)] = \{[a_t] : a_t \in \bigcup_{i=1}^5 f_i(B_4(24)), 1 \leq t \leq 350\}$  is a basis of the  $\mathbb{F}_2$ -vector space  $(QP_5^0)_{24}$ . Consequently,  $\dim(QP_5^0)_{24} = |\bigcup_{i=1}^5 f_i(B_4(24))| = 350$ .

**Remark 3.3.** We recall a result in Mothebe-Kaelo-Ramatebele [9] as follows.

Set  $\mathcal{M}_{(k,r)} = \{J = (j_1, j_2, \dots, j_r) : 1 \leq j_1 < \dots < j_r \leq k\}$ ,  $1 \leq r < k$ . For  $J \in \mathcal{M}_{(k,r)}$ , define the homomorphism  $f_J : P_r \rightarrow P_k$  of algebras by substituting  $f_J(x_t) = x_{j_t}$  with  $1 \leq t \leq r$ . Then,  $f_J$  is a monomorphism of  $\mathcal{A}$ -modules. We have a direct summand decomposition of the  $\mathbb{F}_2$ -vector subspaces:

$$QP_k^0 = \bigoplus_{1 \leq r \leq k-1} \bigoplus_{J \in \mathcal{M}_{(k,r)}} (Qf_J(P_r^+)),$$

where,  $Qf_J(P_r^+) = \mathbb{F}_2 \otimes_{\mathcal{A}} f_J(P_r^+)$ .

In degree  $n$ , we have  $\dim(Qf_J(P_r^+))_n = \dim(QP_r^+)_n$  and  $|\mathcal{M}_{(k,r)}| = \binom{k}{r}$ . Hence, combining with Theorem 1.1 we get

$$\dim(QP_k^0)_n = \sum_{\mu(n) \leq r \leq k-1} \binom{k}{r} \dim(QP_r^+)_n.$$

And therefore, using the results in Sum [21], one gets  $\dim(QP_5^0)_{24} = 350$ .

**Theorem 3.4.**

$$\dim(QP_5(\omega_{(d)})) = \begin{cases} 75, & \text{if } d = 1, \\ 145, & \text{if } d = 2, \\ 390, & \text{if } d = 3, \\ 1, & \text{if } d = 4. \end{cases}$$

The proof of the above theorem is too long and very technical. It is proved by explicitly determining all admissible monomials of  $P_5(\omega_{(d)})$  for  $d \in \{1, 2, 3, 4\}$ .

From the above results, we get the corollary following.

**Corollary 3.5.** *There exist exactly 961 admissible monomials of degree twenty-four in  $P_5$ . Consequently,  $\dim(\mathbb{F}_2 \otimes_{\mathcal{A}} P_5)_{24} = 961$ .*

**Case s=1.**

Since Kameko's homomorphism  $(\widetilde{Sq}_*)_{(5,24)}^0$  is an epimorphism, we have

$$(\mathbb{F}_2 \otimes_{\mathcal{A}} P_5)_{53} \cong \text{Ker}(\widetilde{Sq}_*)_{(5,24)}^0 \oplus \text{Im}(\widetilde{Sq}_*)_{(5,24)}^0.$$

We have the following

**Theorem 3.6.**  *$\text{Im}(\widetilde{Sq}_*)_{(5,24)}^0$  is isomorphic to a subspace of  $(\mathbb{F}_2 \otimes_{\mathcal{A}} P_5)_{53}$  generated by all the classes represented by the admissible monomials of the form  $x_1 x_2 \dots x_5 u^2$ , for every  $u \in B_5(24)$ . Consequently,  $\dim(\text{Im}(\widetilde{Sq}_*)_{(5,24)}^0) = 961$ .*

Next, we explicitly determine  $\text{Ker}(\widetilde{Sq}_*)_{(5,24)}^0$  by giving a direct summand decomposition of the  $\mathbb{F}_2$ -vector spaces  $\text{Ker}(\widetilde{Sq}_*)_{(5,24)}^0$  as follows.

**Theorem 3.7.** *We have a direct summand decomposition of the  $\mathbb{F}_2$ -vector spaces*

$$\text{Ker}(\widetilde{Sq}_*)_{(5,24)}^0 = (QP_5^0)_{53} \oplus QP_5^+(3, 3, 3, 2, 1).$$

**Remark 3.8.** From the result in [9] we have

$$\dim(QP_k^0)_{53} = \sum_{\mu(53) \leq r \leq 4} \binom{5}{r} \dim(QP_r^+)_{53}.$$

Since  $\mu(53) = 3$ ,  $\dim(QP_3^+)_{53} = 8$  (see Kameko [6]) and  $\dim(QP_4^+)_{53} = 88$  (see Sum [21]), one gets

$$\dim(QP_5^0)_{53} = \binom{5}{3} \cdot \dim(QP_3^+)_{53} + \binom{5}{4} \cdot \dim(QP_4^+)_{53} = 520.$$

And therefore the set  $\{[b_t] : b_t \in \bigcup_{i=1}^5 f_i(B_4(53)), 1 \leq t \leq 520\}$  is a basis of the  $\mathbb{F}_2$ -vector space  $(QP_5^0)_{53}$ .

We denote by  $\mathbb{M}^d(n) = \{[\bigcup_{i=1}^5 x_i^{2^d-1} f_i(x)] : x \in B_{k-1}(n - 2^d + 1)\}$  and set  $\mathbb{M} = \text{Span}\{u : u \in \bigcup_{d=1}^5 \mathbb{M}^d(53) \text{ and } \omega(u) = (3, 3, 3, 2, 1)\}$ . Hence, one gets the theorem following.

**Theorem 3.9.** *The following statements are true:*

- i)  $\mathbb{M}$  is the  $\mathbb{F}_2$ -vector subspaces of  $QP_5^+(\omega)$ , where  $\omega = (3, 3, 3, 2, 1)$ .
- ii) Assume  $QP_5^+(\omega) = \mathbb{M} \oplus \mathbb{N}$ . Then,  $\dim(\mathbb{M}) = 389$  and  $\dim(\mathbb{N})$ .

The above theorem is proved by explicitly determining all admissible monomials in  $P_5^+(\omega)$ .

On the other hand, we see that  $|B_5(25)| = 1240$  (see Sum [22]) and  $|B_5(19)| = 905$  (see Tin [25]). Since  $\mu(25) = 3 = \alpha(25 + \mu(25))$ , using the result in Sum [21], one gets

$$|B_6(5(2^v - 1) + 2^v 25)| = (2^6 - 1)|B_5(25)|, \text{ for any integer } v \geq k - 1 = 5.$$

So, we obtain the corollary following.

**Corollary 3.10.** *There exist exactly 78120 admissible monomials of degree  $m_1 = 5(2^v - 1) + 25 \cdot 2^v$  in  $P_6$ , for any  $v > 4$ . Consequently,  $\dim(\mathbb{F}_2 \otimes_{\mathcal{A}} P_6)_{m_1} = 78120$ .*

Similarly, it is easy to check that  $\mu(19) = 3 = \alpha(\mu(19) + 19)$ , using the result in Sum [21] we get the following.

**Corollary 3.11.** *There exist exactly 57015 admissible monomials of degree  $m_2 = 5(2^r - 1) + 19 \cdot 2^r$  in  $P_6$ , for any  $r \geq 5$ . Consequently,  $\dim(\mathbb{F}_2 \otimes_{\mathcal{A}} P_6)_{m_2} = 57015$ .*

Next, the aim of our paper is to verify this conjecture for the case  $k = 5$  and the internal degree  $5(2^s - 1) + 24 \cdot 2^s$ , with  $s$  an arbitrary non-negative integer. We get the following theorem.

**Theorem 3.12.** *Singer's conjecture is true for  $k = 5$  and the degree  $5(2^s - 1) + 24 \cdot 2^s$ , with  $s$  an arbitrary non-negative integer.*

**Remark 3.13.** We prove the above theorem by using the admissible monomial basis of degree  $5(2^s - 1) + 24 \cdot 2^s$  in  $P_5$  to explicitly compute the vector space  $(\mathbb{F}_2 \otimes_{\mathcal{A}} P_5)_{5(2^s-1)+24 \cdot 2^s}^{GL_5}$  and combining the computation of  $\text{Ext}_{\mathcal{A}}^{5,5+*}(\mathbb{F}_2, \mathbb{F}_2)$  by Lin [7] and Chen [4]. Our method is different from Boardman's method in [1] for  $k = 3$ . Recall that Boardman [1] computed the space  $(\mathbb{F}_2 \otimes_{\mathcal{A}} P_3)_{GL_3}$  by using a basis consisting of the all the classes represented by certain polynomials in  $P_3$ . It is difficult to use his method for  $k \geq 4$ . Moreover, our approach can be applied for  $k = 4$  by using the admissible monomial basis for  $P_4$  (see Sum [21]). We hope that the conjecture is also true in this case.

The proofs of the results of this Note will be pulished in detail elsewhere.

**Acknowledgment.** I would like to thank Ho Chi Minh City University of Technology and Education for supporting this work.



## REFERENCES

- [1] J. M. Boardman, *Modular representations on the homology of power of real projective space*, in: M. C. Tangora (Ed.), *Algebraic Topology*, Oaxtepec, 1991, in: *Contemp. Math.*, vol. 146, 1993, pp. 49-70, MR1224907.
- [2] P. H. Chon and L. M. Ha, *On May spectral sequence and the algebraic transfer*, *Proc. Japan Acad. Ser. A Math. Sci.* 86 (2010), no. 9, 159-164.
- [3] P. H. Chon and L. M. Hà, *On the May spectral sequence and the algebraic transfer II*, *Topology Appl.* 178 (2014), 372-383.
- [4] T. W. Chen, *Determination of  $\text{Ext}_{\mathcal{A}}^{5,*}(\mathbb{Z}/2, \mathbb{Z}/2)$* , *Topology Appl.*, 158 (2011), 660-689, MR2774051.
- [5] N. H. V. Hung, *The cohomology of the Steenrod algebra and representations of the general linear groups*, *Trans. Amer. Math. Soc.* 357 (2005), 4065-4089.
- [6] M. Kameko, *Products of projective spaces as Steenrod modules*, Ph.D. Thesis, The Johns Hopkins University, ProQuest LLC, Ann Arbor, MI, 1990. 29 pp.
- [7] W. H. Lin,  *$\text{Ext}_{\mathcal{A}}^{4,*}(\mathbb{Z}/2, \mathbb{Z}/2)$  and  $\text{Ext}_{\mathcal{A}}^{5,*}(\mathbb{Z}/2, \mathbb{Z}/2)$* , *Topology Appl.*, 155 (2008), 459-496, MR2380930.
- [8] M. Mothebe, *Dimensions of the polynomial algebra  $\mathbf{F}_2[x_1, \dots, x_n]$  as a module over the Steenrod algebra*, *JP J. Algebra Number Theory Appl.* 13 (2009), No. 2, 161-170.
- [9] M. F. Mothebe, P. Kaelo and O. Ramatebele, *Dimension formulae for the polynomial algebra as a module over the Steenrod algebra in degrees less than or equal to 12*, *J. Math. Research*, 8, (2016), pp. 92-100.
- [10] T. N. Nam,  *$\mathcal{A}$ -générateurs génériques pour l'algèbre polynomiale*, *Adv. Math.*, 186, (2004), pp. 334-362.
- [11] F. P. Peterson, *Generators of  $H^*(\mathbb{R}P^\infty \times \mathbb{R}P^\infty)$  as a module over the Steenrod algebra*, *Abstracts Amer. Math. Soc. No. 833*, (1987), pp. 55-89.
- [12] D. V. Phuc and N. Sum, *On a minimal set of generators for the polynomial algebra of five variables as a module over the Steenrod algebra*, *Acta Math. Vietnam.* 42 (2017), no. 1, 149-162.
- [13] D. V. Phuc,  *$\mathcal{A}$ -generators for the polynomial algebra of five variables and the Singer transfer*, Preprint, available online at <http://arxiv.org/abs/1810.06061v10>.
- [14] D. V. Phuc, *A note on the epimorphism of the fifth algebraic transfer*, *JP J. Algebra, Number Theory Appl.* 48, No.2, (2020) pp. 193-201.
- [15] S. Priddy, *On characterizing summands in the classifying space of a group, I*, *Amer. Jour. Math.* 112 (1990), 737-748.
- [16] W. M. Singer, *The transfer in homological algebra*, *Math. Zeit.* 202 (1989) 493-523.
- [17] J. H. Silverman (1995), "Hit polynomials and the canonical antiautomorphism of the Steenrod algebra", *Proc. Amer. Math. Soc.*, 123, pp. 627-637.
- [18] N. E. Steenrod and D. B. A. Epstein, *Cohomology operations*, *Annals of Mathematics Studies* 50, Princeton University Press, Princeton N.J (1962).
- [19] N. Sum, *The negative answer to Kameko's conjecture on the hit problem*, *Adv. Math.* 225, (2010), 2365-2390.
- [20] N. Sum, *On the hit problem for the polynomial algebra*, *C. R. Acad. Sci. Paris, Ser. I*, 351 (2013) 565-568.
- [21] N. Sum, *On the Peterson hit problem*, *Adv. Math.* 274 (2015) 432-489.
- [22] N. Sum, *On a construction for the generators of the polynomial algebra as a module over the Steenrod algebra*, *Algebraic topology and related topics*, 265-286, *Trends Math.*, Birkh./Springer, Singapore, 2019.
- [23] N. Sum and N. K. Tin, *The hit problem for the polynomial algebra in some weight vectors*, *Topology Appl.* 290 (2021), 107579, 17pp.
- [24] N. K. Tin and N. Sum, *Kameko's homomorphism and the algebraic transfer*, *C. R. Acad. Sci. Paris, Ser. I*, 354, (2016), 940-943.
- [25] N. K. Tin, *The hit problem for the polynomial algebra in five variables and applications*, Ph.D. thesis, Quy Nhon University 2017.
- [26] N. K. Tin, *The admissible monomial basis for the polynomial algebra as a module over Steenrod algebra in some degrees*, *JP J. Algebra, Number Theory Appl.* 46, No.1, (2020) pp. 55-68.



- [27] N. K. Tin, *The hit problem for the polynomial algebra as a module over Steenrod algebra, I*, JP J. Algebra, Number Theory Appl. 47, No.1, (2020) pp. 67-86.
- [28] N. K. Tin, *A note on the Peterson hit problem for the Steenrod algebra*, Preprint, 2021.
- [29] N. K. Tin, *The hit problem for the polynomial algebra as a module over Steenrod algebra in some degrees*, Asian-European J. Math. 18 pp. (2021). (accepted)
- [30] G. Walker and R. M. W. Wood, *Polynomials and the mod 2 Steenrod algebra. Vol. 1. The Peterson hit problem*, London Mathematical Society Lecture Note Series, **441**. Cambridge University Press, 2018.
- [31] G. Walker and R. M. W. Wood, *Polynomials and the mod 2 Steenrod algebra. Vol. 2. Representations of  $GL(n, \mathbb{F}_2)$* , London Mathematical Society Lecture Note Series, **442**. Cambridge University Press, 2018.
- [32] R. M. W. Wood, *Steenrod squares of polynomials and the Peterson conjecture*, Math. Proc. Cambridge Phil. Soc. 105 (1989) 307-309.

FACULTY OF APPLIED SCIENCES, HO CHI MINH CITY UNIVERSITY OF TECHNOLOGY AND EDUCATION, 01 VÕ VĂN NGÂN, THỦ ĐỨC, HỒ CHÍ MINH CITY, VIET NAM.

*Email address:* `tinnk@hcmute.edu.vn`