

## New Flat surfaces in $S^3$

**Armando M. V. Corro**

IME, Universidade Federal de Goiás  
Caixa Postal 131, 74001-970, Goiânia, GO, Brazil  
e-mail:corro@ufg.br

**Marcelo Lopes Ferro**

IME, Universidade Federal de Goiás  
Caixa Postal 131, 74001-970, Goiânia, GO, Brazil  
e-mail: marceloferro@ufg.br

### Abstract

In this paper, we consider a method of constructing flat surfaces based on Ribaucour transformations in the sphere 3-space. By applying the theory to the flat torus, we obtain a families of complete flat surfaces in  $S^3$  which are determined by several parameters. we provide explicit examples.

### Introduction

Ribaucour transformations for hypersurfaces, parametrized by lines of curvature, were classically studied by Bianchi [2]. They can be applied to obtain surfaces of constant Gaussian curvature and surfaces of constant mean curvature, from a given such surface, respectively, with constant Gaussian curvature and constant mean curvature. The first application of this method to minimal and cmc surfaces in  $R^3$  was obtained by Corro, Ferreira, and Tenenblat in [4]-[6]. In [26], Tenenblat and Wang extended such transformations to surfaces of space forms. For more application this method, see [7], [11], [13], [24], [25] and [27].

Using Ribaucour transformations and applying the theory to the rotational flat surfaces in  $H^3$ , in [10] the authours obtained families of new such surfaces.

The study of flat surfaces in  $S^3$  traces back to Bianchi's works in the 19th century, and it has a very rich global theory, as evidenced by the existence of a large class of flat tori in  $S^3$ , see [14], [21], [22] and [28]. Indeed, these flat tori constitute the only examples of compact surfaces of constant curvature in space forms that are not totally umbilical round spheres. Flat surfaces in  $S^3$  admit a more explicit treatment than other constant curvature surfaces. Moreover, there are still important open problems regarding flat surfaces in  $S^3$ , some of them unanswered for more than 40 years. For instance, it remains unknown if there exists an isometric embedding of  $R^2$  into  $S^3$ . These facts show that the geometry of flat surfaces in  $S^3$  is a worth studying topic, although the number of contributions to the theory is not too large. Some important references of the theory are [12], [14]-[18], [21], [22] and [28].

In [20], the authors give a complete classification of helicoidal flat surfaces in  $S^3$  by means of asymptotic lines coordinates.

In [1], the authors characterized the flat surfaces in the unit 3-sphere that pass through a given regular curve of  $S^3$  with a prescribed tangent plane distribution along this curve.

In this paper, motivated by [10] we use the Ribaucour transformations to get a family of complete flat surfaces in  $S^3$  from a given such surface in  $S^3$ . As an application of the theory,

we obtain a families of complete flat surfaces in  $S^3$  associated to the flat torus. The families we obtain depend four parameters. One of these parameters is given by parameterizing of the flat torus. The other parameters, appear from integrating the Ribaucour transformation. We show explicit examples of these surfaces. This work is organized as follows. In Section 1, we give a brief description of Ribaucour transformations in space forms. In Section 2, we give an additional condition for the transformed surface to be flat. In Section 3, we describe all flat surfaces of the sphere 3-space obtained by applying the Ribaucour transformation to the flat torus. We prove that such surfaces are complete and provide explicit examples.

## 1. Preliminary

This section contains the definitions and the basic theory of Ribaucour transformations for surfaces in  $S^3$ . ( For more Details see [26] )

Let  $M$  be an orientable surface in  $S^3$  without umbilic points, with Gauss map we denote by  $N$ . Suppose that there exist 2 orthonormal principal vector fields  $e_1$  and  $e_2$  defined on  $M$ . We say that  $\widetilde{M} \subset S^3$  is associated to  $M$  by a Ribaucour transformation with respect to  $e_1$  and  $e_2$ , if there exist a differentiable function  $h$  defined on  $M$  and a diffeomorphism  $\psi : M \rightarrow \widetilde{M}$  such that

- (a) for all  $p \in M$ ,  $exp_p h(p)N(p) = exp_{\psi(p)} h(p)\widetilde{N}(\psi(p))$ , where  $\widetilde{N}$  is the Gauss map of  $\widetilde{M}$  and  $exp$  is the exponential map of  $M^3(\bar{k})$ .
- (b) The subset  $\{exp_p h(p)N(p), p \in M\}$ , is a two-dimensional submanifold of  $S^3$ .
- (c)  $d\psi(e_i) \ 1 \leq i \leq 2$  are orthogonal principal directions of  $\widetilde{M}$ .

The following result gives a characterization of Ribaucour transformations. ( see [26] for a proof and more details)

**Theorem 1.1** *Let  $M$  be an orientable surface of  $S^3$  parametrized by  $X : U \subseteq R^2 \rightarrow M$ , without umbilic points. Assume  $e_i = \frac{X_{,i}}{a_i}$ ,  $1 \leq i \leq 2$  where  $a_i = \sqrt{g_{ii}}$  are orthogonal principal directions,  $-\lambda_i$  the corresponding principal curvatures, and  $N$  is a unit vector field normal to  $M$ . A surface  $\widetilde{M}$  is locally associated to  $M$  by a Ribaucour transformation if and only if there is differentiable functions  $W, \Omega, \Omega_i : V \subseteq U \rightarrow R$  which satisfy*

$$\begin{aligned} \Omega_{i,j} &= \Omega_j \frac{a_{j,i}}{a_i}, \quad \text{for } i \neq j, \\ \Omega_{,i} &= a_i \Omega_i, \\ W_{,i} &= -a_i \Omega_i \lambda_i. \end{aligned} \tag{1}$$

$W(W + \lambda_i \Omega) \neq 0$  and  $\widetilde{X} : V \subseteq U \rightarrow \widetilde{M}$ , is a parametrization of  $\widetilde{M}$  given by

$$\widetilde{X} = \left(1 - \frac{2\Omega^2}{S}\right)X - \frac{2\Omega}{S} \left(\sum_{i=1}^2 \Omega_i e_i - WN\right), \tag{2}$$

where

$$S = \sum_{i=1}^2 (\Omega_i)^2 + W^2 + \Omega^2. \tag{3}$$

Moreover, the normal map of  $\widetilde{X}$  is given by

$$\widetilde{N} = N + \frac{2W}{S} \left(\sum_{i=1}^2 \Omega_i e_i - WN + \Omega X\right), \tag{4}$$

and the principal curvatures and coefficients of the first fundamental form of  $\tilde{X}$ , are given by

$$\tilde{\lambda}_i = \frac{WT_i + \lambda_i S}{S - \Omega T_i}, \quad \tilde{g}_{ii} = \left( \frac{S - \Omega T_i}{S} \right)^2 g_{ii} \quad (5)$$

where  $\Omega_i$ ,  $\Omega$  and  $W$  satisfy (1),  $S$  is given by (3),  $g_{ii}$ ,  $1 \leq i \leq 2$  are coefficients of the first fundamental form of  $X$ , and

$$T_1 = 2 \left( \frac{\Omega_{1,1}}{a_1} + \frac{a_{1,2}}{a_1 a_2} \Omega_2 - W \lambda_1 + \Omega \right), \quad T_2 = 2 \left( \frac{\Omega_{2,2}}{a_2} + \frac{a_{2,1}}{a_1 a_2} \Omega_1 - W \lambda_2 + \Omega \right) \quad (6)$$

## 2. Ribaucour transformation for flat surface in $S^3$

In this section we provides a sufficient condition for a Ribaucour transformation to transform a flat surface into another such surface.

**Theorem 2.1** *Let  $M$  be a surfaces of  $S^3$  parametrized by  $X : U \subseteq R^2 \rightarrow M$ , without umbilic points and let  $\tilde{M}$  parametrized by (2) be associated to  $M$  by a Ribaucour transformation, such that the normal lines intersect at a distance function  $h$ . Assume that  $h = \frac{\Omega}{W}$  is not constant along the lines of curvature and the function  $\Omega$ ,  $\Omega_i$  and  $W$  satisfy one of the additional relation*

$$\Omega_1^2 + \Omega_2^2 = c(\Omega^2 + W^2) \quad (7)$$

where  $c > 0$ ,  $S$  is given by (3) and  $W$ ,  $\Omega_i$ ,  $1 \leq i \leq 2$  satisfies (1). Then  $\tilde{M}$  parameterized by (2) is a flat surface, if and only if  $M$  is a flat surface.

**Proof:** See [26] Theorem 2.1, with  $\alpha = \gamma = 1$  and  $\beta = 0$ .

**Remark 2.2** Let  $X$  as in the previous theorem. Then the parameterization  $\tilde{X}$  of  $\tilde{M}$ , locally associated to  $X$  by a Ribaucour transformation, given by (2), is defined on

$$V = \{(u_1, u_2) \in U; (\Omega T_1 - S)(\Omega T_2 - S) \neq 0\}.$$

## 3. Families of flat surfaces associated to the Flat Torus in $S^3$ .

In this section, by applying Theorem 2.1 to the Flat Torus in  $S^3$ , we obtain a three parameter family of complete flat surfaces in  $S^3$ .

**Theorem 3.1** *Consider the Flat Torus in  $S^3$  parametrized by*

$$X(u_1, u_2) = (r_1 \cos(r_2 u_1), r_1 \sin(r_2 u_1), r_2 \cos(r_1 u_2), r_2 \sin(r_1 u_2)), \quad (u_1, u_2) \in R^2 \quad (8)$$

$r_i$ ,  $1 \leq i \leq 2$  are positive constants satisfying  $r_1^2 + r_2^2 = 1$ , as flat torus in  $S^3$  where the first fundamental form is  $I = r_1^2 r_2^2 (du_1^2 + du_2^2)$ . A parametrized surface  $\tilde{X}(u_1, u_2)$  is flat surface locally associated to  $X$  by a Ribaucour transformation as in Theorem 2.1 with additional relation given by (7), if and only if, up to an isometries of  $S^3$ , it is given by

$$\tilde{X} = \frac{1}{S} \begin{bmatrix} (r_1 S - 2r_1 \Omega^2 - 2r_2 \Omega W) \cos(r_2 u_1) + 2f' \Omega \sin(r_2 u_1) \\ (r_1 S - 2r_1 \Omega^2 - 2r_2 \Omega W) \sin(r_2 u_1) - 2f' \Omega \cos(r_2 u_1) \\ (r_2 S - 2r_2 \Omega^2 + 2r_1 \Omega W) \cos(r_1 u_2) + 2g' \Omega \sin(r_1 u_2) \\ (r_2 S - 2r_2 \Omega^2 + 2r_1 \Omega W) \sin(r_1 u_2) - 2g' \Omega \cos(r_1 u_2) \end{bmatrix} \quad (9)$$

defined on  $V = \{(u_1, u_2) \in \mathbb{R}^2; (r_1^2 g^2 - r_2^2 f^2 - 2r_2^2 fg)(r_2^2 f^2 - r_1^2 g^2 + 2r_1^2 fg) \neq 0\}$ , where

$$\Omega = r_1 r_2 (f(u_1) + g(u_2)), \quad W = r_2^2 f(u_1) - r_1^2 g(u_2), \quad S = (1 + c)(\Omega^2 + W^2) \quad (10)$$

$c > 0$ , and the functions  $f$  and  $g$  are given by

$$i) f(u_1) = \cosh(r_2 \sqrt{c} u_1), \quad g(u_2) = \frac{r_2}{r_1} \sinh(r_1 \sqrt{c} u_2), \quad (11)$$

or

$$ii) f(u_1) = \sinh(r_2 \sqrt{c} u_1), \quad g(u_2) = \frac{r_2}{r_1} \cosh(r_1 \sqrt{c} u_2), \quad (12)$$

or

$$iii) f(u_1) = a_1 e^{\epsilon_1 r_2 \sqrt{c} u_1}, \quad g(u_2) = b_1 e^{\epsilon_2 r_1 \sqrt{c} u_2}, \quad \epsilon_i^2 = 1, \quad 1 \leq i \leq 2. \quad (13)$$

Moreover, the normal map of  $\tilde{X}$  is given by

$$\tilde{N} = \frac{1}{S} \begin{bmatrix} \left( -r_2 S - 2r_2 W^2 + 2r_1 \Omega W \right) \cos(r_2 u_1) + 2f' \Omega \sin(r_2 u_1) \\ \left( -r_2 S - 2r_2 W^2 + 2r_1 \Omega W \right) \sin(r_2 u_1) - 2f' \Omega \cos(r_2 u_1) \\ \left( r_1 S + 2r_1 W^2 - 2r_2 \Omega W \right) \cos(r_1 u_2) + 2g' \Omega \sin(r_1 u_2) \\ \left( r_1 S + 2r_1 W^2 - 2r_2 \Omega W \right) \sin(r_1 u_2) - 2g' \Omega \cos(r_1 u_2) \end{bmatrix} \quad (14)$$

**Proof:** Consider the first fundamental form of the Flat Torus  $ds^2 = r_1^2 r_2^2 (du_1^2 + du_2^2)$  and the principal curvatures  $-\lambda_i$   $1 \leq i \leq 2$  given by  $\lambda_1 = \frac{-r_2}{r_1}$ ,  $\lambda_2 = \frac{r_1}{r_2}$ . Using (1), to obtain the Ribaucour transformations, we need to solve the following of equations

$$\Omega_{i,j} = 0, \quad \Omega_{,i} = r_1 r_2 \Omega_i, \quad W_{,i} = -r_1 r_2 \Omega_i \lambda_i, \quad 1 \leq i \neq j \leq 2. \quad (15)$$

Therefore we obtain

$$\Omega = r_1 r_2 (f_1(u_1) + f_2(u_2)), \quad W = -r_1 r_2 (\lambda_1 f_1 + \lambda_2 f_2) + \bar{c}, \quad \Omega_i = f'_i, \quad 1 \leq i \neq j \leq 2, \quad (16)$$

where  $\bar{c}$  is a real constant. Using (7) the associated surface will be flat when  $\Omega_1^2 + \Omega_2^2 = c(\Omega^2 + W^2)$ . Therefore, we obtain that  $c > 0$  and the functions  $f_1$  and  $f_2$  satisfy

$$(f'_1)^2 + (f'_2)^2 = c(\Omega^2 + W^2). \quad (17)$$

Differentiate this last equation with respect  $x_1$  and  $x_2$ , using (15) and (16) we get

$$\begin{aligned} f''_1 &= cr_2^2 f_1 + cr_2^2 \bar{c}, \\ f''_2 &= cr_1^2 f_2 - cr_1^2 \bar{c}. \end{aligned}$$

Defining  $f(u_1) = f_1(u_1) + \bar{c}$  and  $g(u_2) = f_2(u_2) - \bar{c}$ , we have

$$f'' - cr_2^2 f = 0, \quad g'' - cr_1^2 g = 0, \quad (18)$$

$$\Omega = r_1 r_2 (f(u_1) + g(u_2)), \quad W = r_2^2 f(u_1) - r_1^2 g(u_2). \quad (19)$$

By Theorem 1.1, we have that  $\tilde{X}$  and  $\tilde{N}$  are given by (9) and (14). Using (6) and (19), we get

$$T_1 = \frac{2r_2(1+c)f}{r_1} \quad T_2 = \frac{2r_1(1+c)g}{r_2}.$$

Thus, from Remark 2.2  $\tilde{X}$  is defined in  $V = \{(u_1, u_2) \in \mathbb{R}^2; (r_1^2 g^2 - r_2^2 f^2 - 2r_2^2 fg)(r_2^2 f^2 - r_1^2 g^2 + 2r_1^2 fg) \neq 0\}$ .

From (18), we get

$$f(u_1) = a_1 \cosh(r_2 \sqrt{c} u_1) + a_2 \sinh(r_2 \sqrt{c} u_1), \quad g(u_2) = b_1 \cosh(r_1 \sqrt{c} u_2) + b_2 \sinh(r_1 \sqrt{c} u_2). \quad (20)$$

Substituting (20) and (19) in (17), we have

$$(a_1^2 - a_2^2)r_2^2 = (b_2^2 - b_1^2)r_1^2. \quad (21)$$

Let  $A_1 = a_1^2 - a_2^2$ . If  $A_1 > 0$ , then from (21)  $b_2^2 > b_1^2$ . Hence (20) can be rewritten as

$$f(u_1) = \sqrt{A_1} \cosh(r_2 \sqrt{c} u_1 + A_2), \quad (22)$$

$$g(u_2) = \sqrt{A_1} \frac{r_2}{r_1} \sinh(r_1 \sqrt{c} u_2 + B_2), \quad (23)$$

where  $\cosh(A_2) = \frac{a_1}{\sqrt{a_1^2 - a_2^2}}$ ,  $\sinh(A_2) = \frac{a_2}{\sqrt{a_1^2 - a_2^2}}$ ,  $\sinh(B_2) = \frac{b_2}{\sqrt{b_2^2 - b_1^2}}$  and  $\cosh(B_2) = \frac{b_1}{\sqrt{b_2^2 - b_1^2}}$ .

The constants  $A_2$  and  $B_2$ , without loss of generality, may be considered to be zero. One can verify that the surfaces with different values of  $A_2$  and  $B_2$  are congruent. In fact, using the notation  $\tilde{X}_{A_2 B_2}$  for the surface  $\tilde{X}$  with fixed constants  $A_2$  and  $B_2$ , we have

$$\tilde{X}_{A_2 B_2} = R_{\left(\frac{-A_2}{r_2 \sqrt{c}}, \frac{-B_2}{r_1 \sqrt{c}}\right)} \tilde{X}_{00} \circ h$$

where  $h(u_1, u_2) = \left(u_1 + \frac{A_2}{r_2 \sqrt{c}}, u_1 + \frac{B_2}{r_1 \sqrt{c}}\right)$  with

$$R_{(\theta, \phi)}(x_1, x_2, x_3, x_4) = (x_1 \cos \theta - x_2 \sin \theta, x_1 \sin \theta + x_2 \cos \theta, x_3 \cos \phi - x_4 \sin \phi, x_3 \sin \phi + x_4 \cos \phi).$$

Now substituting (22) with  $A_2 = 0$ , (23) with  $B_2 = 0$  and (19) in (9) we obtain that  $\tilde{X}$  does not depend on  $A_1$ . Thus without loss of generality, we can consider  $A_1 = 1$ . Therefore we conclude that  $f$  and  $g$  are given by (11).

On the other hand, if  $A_1 < 0$ , then from (21)  $b_2^2 < b_1^2$ . Hence (20) can be rewritten as

$$f(u_1) = \sqrt{-A_1} \sinh(r_2 \sqrt{c} u_1 + A_2), \quad (24)$$

$$g(u_2) = \sqrt{-A_1} \frac{r_2}{r_1} \cosh(r_1 \sqrt{c} u_2 + B_2). \quad (25)$$

Proceeding in a similar way to the previous case, we obtain that  $f$  and  $g$  are given by (12).

If  $A_1 = 0$ , then  $a_2 = \epsilon_1 a_1$ , and from (21)  $b_2 = \epsilon_2 b_1$ , with  $\epsilon_i^2 = 1$ ,  $1 \leq i \leq 2$ . Thus, substituting this in (20), we obtain (13).

**Remark 3.2** Each flat surfaces associated to the flat torus as in Theorem 3.1, is parametrized by lines of curvature and from (5), the metric is given by  $ds^2 = \psi_1^2 du_1^2 + \psi_2^2 du_2^2$ , where

$$\psi_1 = \frac{(-r_2^2 f^2 + r_1^2 g^2 - 2r_2^2 f g)r_1 r_2}{r_1^2 g^2 + r_2^2 f^2}, \quad \psi_2 = \frac{(r_2^2 f^2 - r_1^2 g^2 - 2r_1^2 f g)r_1 r_2}{r_1^2 g^2 + r_2^2 f^2}. \quad (26)$$

Moreover, from (5), the principal curvatures of the  $\tilde{X}$  are given by

$$\tilde{\lambda}_1 = \frac{-r_2 \psi_2}{r_1 \psi_1}, \quad \tilde{\lambda}_2 = \frac{r_1 \psi_1}{r_2 \psi_2}. \quad (27)$$

**Proposition 3.3** Any flat surfaces associated to the flat torus  $\tilde{X}$ , given by Theorem 3.1 is complete.

**Proof:** For divergent curves  $\gamma(t) = (u_1(t), u_2(t))$ , such that  $\lim_{t \rightarrow \infty} (u_1^2 + u_2^2) = \infty$ , we have  $l(\tilde{X} \circ \gamma) = \infty$ .

In fact, the functions  $f$  and  $g$  are given by (11) or (12) or (13) and the coefficients of the first fundamental form  $\psi_i$ ,  $1 \leq i \leq 2$  of  $\tilde{X}$  given by (26). Therefore  $\lim_{|u_1| \rightarrow \infty} |\psi_i| = r_1 r_2$ ,  $1 \leq i \leq 2$ , uniformly in  $u_2$  and  $\lim_{|u_2| \rightarrow \infty} |\psi_i| = r_1 r_2$ ,  $1 \leq i \leq 2$ , uniformly in  $u_1$ . Hence, there are  $k_1 > 0$  and  $k_2 > 0$  such that  $|\psi_i(u_1, u_2)| > \frac{r_1 r_2}{2}$ ,  $1 \leq i \leq 2$  for all  $(u_1, u_2) \in \mathbb{R}^2$  with  $|u_1| > k_1$  and  $|u_2| > k_2$ . Let

$$m_i = \min\{|\psi_i(u_1, u_2)|; (u_1, u_2) \in [-k_1, k_1] \times [-k_2, k_2]\}.$$

Therefore  $|\psi_i(u_1, u_2)| \geq m_i$  in  $[-k_1, k_1] \times [-k_2, k_2]$ . Now consider  $m_0 = \min\{m_1, m_2, \frac{r_1 r_2}{2}\}$ , then  $|\psi_i(u_1, u_2)| \geq m_0$  in  $\mathbb{R}^2$ . We conclude that  $\tilde{X}$  is a complete surface.

In the following, we provide some examples

**Example 3.3** Consider stereographic projection  $\pi : S^3 \rightarrow \mathbb{R}^3$

$$\pi(x_1, x_2, x_3, x_4) = \frac{1}{1 - x_4} (x_1, x_2, x_3)$$

where  $x_1^2 + x_2^2 + x_3^2 + x_4^2 = 1$ .

In Figure 1, we provide the surface parametrized by  $\pi \circ \tilde{X}$  where  $\tilde{X}$  is given by (9) locally associated to flat torus in  $S^3$  by a Ribaucour transformation. In this surface, we have

$$f(u_1) = \cosh\left(\frac{8u_1}{5}\right) \text{ and } g(u_2) = \frac{4}{3} \sinh\left(\frac{6u_2}{5}\right)$$




Figure 1: In the figure above we have  $r_1 = \frac{3}{5}$ ,  $r_2 = \frac{4}{5}$ ,  $c = 4$ .

In Figure 2, we provide the surface parametrized by  $\pi \circ \tilde{X}$  where  $\tilde{X}$  is given by (9) locally associated to flat torus in  $S^3$  by a Ribaucour transformation. In this surface, we have

$$f(u_1) = \sinh\left(\frac{8u_1}{5}\right) \text{ and } g(u_2) = \frac{4}{3} \cosh\left(\frac{6u_2}{5}\right).$$

In Figure 3, we provide two surfaces parametrized by  $\pi \circ \tilde{X}$  where  $\tilde{X}$  is given by (9) locally associated to flat torus in  $S^3$  by a Ribaucour transformation. In this surfaces, we have  $r_1 = \frac{3}{5}$ ,  $r_2 = \frac{4}{5}$ ,  $c = \frac{1}{1000}$  with  $f$  and  $g$  given by (11) in the first surface and  $f$  and  $g$  given by (12) and the second.

In Figure 4, we provide two surfaces parametrized by  $\pi \circ \tilde{X}$  where  $\tilde{X}$  is given by (9) locally associated to flat torus in  $S^3$  by a Ribaucour transformation. In this surfaces, we have  $r_1 = \frac{3}{5}$ ,  $r_2 = \frac{4}{5}$ ,  $c = \frac{1}{1000}$ ,  $a_1 = b_1 = 1$  with  $f$  and  $g$  given by (13).




Figure 2: In the figure above we have  $r_1 = \frac{3}{5}$ ,  $r_2 = \frac{4}{5}$ ,  $c = 4$ .




Figure 3:




Figure 4:

## References

- [1] Aledo, J.A.; Gálvez, J.A.; Mira, P.; *A D'Alembert Formula for Flat Surfaces in the 3-Sphere*. Journal of Geometric Analysis, 19, 211-232, (2009).
- [2] Bianchi, L.: *Lezioni di geometria Differenziale*. Terza Edizione, Nicola Zanichelli Editore (1927).
- [3] Bobenko, A. I.: *All constant mean curvature tori in  $R^3$ ,  $S^3$ ,  $H^3$  in terms of theta functions*. Math. Ann. 290, 209–245, (1991).
- [4] Corro, A. M. V.; Ferreira, W. P.; Tenenblat, K.: *On Ribaucour transformations for hypersurfaces*, Mat. Contemp. 17, 137-160, (1999).
- [5] Corro, A. M. V.; Ferreira, W. P.; Tenenblat, K.: *Ribaucour transformations for Constant mean curvature and linear weingarten surfaces*. Pacific Journal of Mathematics. 212, 2, 265-296, (2003).
- [6] Corro, A. M. V.; Ferreira, W. P.; Tenenblat, K.: *Minimal surfaces obtained by Ribaucour transformations*. Geometriae Dedicata, Netherlands, 96, 1, 117-150, (2003).
- [7] Corro, A. M. V.; Tenenblat, K.: *Ribaucour transformation revisited*, Comum. Geom. 12, 5, 1055-1082, (2004).

- [8] Corro, A. M. V.: *Generalized Weingarten surfaces of Bryant type en Hiperbolic 3-space*, Mat. Contemp. 30, 71-89, (2006).
- [9] Corro, A. M. V.; Martinez A., Milán F.. *Complete flat surfaces with two isolated singularities in hyperbolic 3-space*, J. Math. Anal. Appl. 366, 582-592 (2010).
- [10] Corro, A. M. V.; Martinez A., Tenenblat, K.: *Ribaucour transformations for flat surfaces in the hyperbolic 3-space*. J. Math. Anal. Appl. 412, 720-743 (2014).
- [11] Cui N. W.; Tenenblat, K.: *New minimal surfaces in the hyperbolic space*. Science China Mathematics, 60, 1679-1704 (2017).
- [12] Dadok, J.; Sha, J.: *On embedded flat surfaces in  $S^3$* . J. Geom. Anal. 7, 47-55 (1997).
- [13] Ferreira, W. P.; Tenenblat, K.: *Hypersurfaces with flat  $r$ -mean curvature and Ribaucour transformations*. International Journal of Applied Mathematics and Statistics, 11, 38-51, (2007).
- [14] Kitagawa, Y.: *Periodicity of the asymptotic curves on flat tori in  $S^3$* . J. Math. Soc. Jpn. 40, 457-476, (1988).
- [15] Kitagawa, Y.: *Ridigity of the Clifford tori in  $S^3$* . Math. Z. 198, 591-599 (1988).
- [16] Kitagawa, Y.: *Embedded flat tori in the unit 3-sphere*. J. Math. Soc. Jpn. 47, 275-296 (1995).
- [17] Kitagawa, Y.: *Isometric deformations of flat tori in the 3-sphere with nonconstant mean curvature*. Tohoku Math. J. 52, 283-298 (2000).
- [18] Kitagawa, Y.: *Deformable flat tori in  $S^3$  with constant mean curvature*. Osaka J. Math. 40, 103-119 (2003).
- [19] Martinez A.; Santos, J.; Tenenblat, K.: *Helicoidal flat surfaces in hyperbolic 3-space*, Pacific Journal of Mathematics. 264, 1, 195-211 (2013).
- [20] Manfio, F.; Santos, J.: *Helicoidal flat surfaces in the 3-sphere*. Mathematische Nachrichten, 292, 127-136, (2019).
- [21] Pinkall, U.: *Hopf tori in  $S^3$* . Invent. Math. 81, 379-386 (1985).
- [22] Spivak, M.; Wang: *A Comprehensive Introduction to Differential Geometry, vol. 4*. Publish or Perish, Inc., Berkeley, (1979).
- [23] Tenenblat, K.: *Transformations of manifolds and applications to differential equations*, Longman Sci. and Tech. (1998).
- [24] Tenenblat, K.: *On Ribaucour transformations and applications to linear Weingarten surfaces*. Anais da Academia Brasileira de Ciências (Impresso), ABC, 74, 559-575, (2002).
- [25] Tenenblat, K.; Lemes, M. V.: *On Ribaucour transformations and minimal surfaces*. Matemática Contemporânea, 29, 13-40, (2005).
- [26] Tenenblat, K.; Wang, Q.: *Ribaucour Transformations for Hypersurfaces in Space Forms*. Annals of Global Analysis and Geometry, 29, 135-162, (2009).
- [27] Tenenblat, K.; Wang, Q.: *New surfaces of constant mean curvature in the hyperbolic space*. Illinois Journal of Mathematics, 53, 157-185, (2006).
- [28] Weiner, J.L.: *Flat tori in  $S^3$  and their Gauss maps*. Proc. Lond. Math. Soc. 62, 54-76 (1991).