

# ON LIPSCHITZ APPROXIMATIONS IN SECOND ORDER SOBOLEV SPACES AND THE CHANGE OF VARIABLES FORMULA

PAZ HASHASH AND ALEXANDER UKHLOV

**ABSTRACT.** In this paper we study approximations of functions of Sobolev spaces  $W_{p,\text{loc}}^2(\Omega)$ ,  $\Omega \subset \mathbb{R}^n$ , by Lipschitz continuous functions. We prove that if  $f \in W_{p,\text{loc}}^2(\Omega)$ ,  $1 \leq p < \infty$ , then there exists a sequence of closed sets  $\{A_k\}_{k=1}^\infty$ ,  $A_k \subset A_{k+1} \subset \Omega$ , such that the restrictions  $f|_{A_k}$  are Lipschitz continuous functions and  $\text{cap}_p(S) = 0$ ,  $S = \Omega \setminus \bigcup_{k=1}^\infty A_k$ . Using these approximations we prove the change of variables formula in the Lebesgue integral for mappings of Sobolev spaces  $W_{p,\text{loc}}^2(\Omega; \mathbb{R}^n)$  with the Luzin capacity-measure  $N$ -property.

## 1. INTRODUCTION

In this paper we study approximations of the second order Sobolev spaces  $W_{p,\text{loc}}^2(\Omega)$ , where  $\Omega \subset \mathbb{R}^n$  is an open set, by Lipschitz continuous functions. The Lipschitz approximations in Sobolev spaces  $W_{p,\text{loc}}^1(\Omega)$  have significant applications in geometric measure theory [9, 25] and in the analysis on metric measure spaces [10, 12, 15].

In [9] it was proved that if  $f \in W_{1,\text{loc}}^1(\Omega)$ , then there exists a sequence of closed sets  $\{A_k\}_{k=1}^\infty$ ,  $A_k \subset A_{k+1} \subset \Omega$ , for which the restrictions  $f^*|_{A_k}$  are Lipschitz continuous functions on the sets  $A_k$  and  $|S| = 0$ ,  $S = \Omega \setminus \bigcup_{k=1}^\infty A_k$ , where we denote by  $|S|$  the  $n$ -dimensional Lebesgue measure of the set  $S \subset \mathbb{R}^n$ . Here  $f^*$  stands for the precise representative of the function  $f$  (see (2.4) below).

In the case of the second order Sobolev spaces  $W_{1,\text{loc}}^2(\Omega)$  we prove the following refined version of this Lipschitz approximation: *Let  $f \in W_{1,\text{loc}}^2(\Omega)$ . Then there exists a sequence of closed sets  $\{A_k\}_{k=1}^\infty$ ,  $A_k \subset A_{k+1} \subset \Omega$ , such that the restrictions  $f^*|_{A_k}$  are Lipschitz continuous functions  $\mathcal{H}^{n-1}$ -a.e. in  $A_k$  and  $\mathcal{H}^{n-1}(S) = 0$ ,  $S = \Omega \setminus \bigcup_{k=1}^\infty A_k$ , where  $\mathcal{H}^{n-1}$  is the  $(n-1)$ -Hausdorff measure.*

Using this Lipschitz approximation we obtain the following change of variables formula in the Lebesgue integral for twice weakly differentiable mappings: Let  $\varphi : \Omega \rightarrow \mathbb{R}^n$  be a mapping which belongs to the Sobolev space  $W_{1,\text{loc}}^2(\Omega; \mathbb{R}^n)$ . Then there exists a Borel set  $S \subset \Omega$ ,  $\mathcal{H}^{n-1}(S) = 0$ , such that the mapping  $\varphi : \Omega \setminus S \rightarrow \mathbb{R}^n$  has the Luzin  $N$ -property (an image of a set of Lebesgue measure zero has Lebesgue measure zero) and the change of variables formula

$$(1.1) \quad \int_A u \circ \varphi(x) |J(x, \varphi)| \, dx = \int_{\mathbb{R}^n \setminus \varphi(S)} u(y) N_\varphi(A, y) \, dy,$$

where  $N_\varphi(A, y)$  is the multiplicity function which is defined as the number of preimages of  $y$  in  $A$  under  $\varphi$  and  $J(x, \varphi) = \det(D\varphi(x))$  is the Jacobian of  $\varphi$  at the point  $x$ , holds for every measurable set  $A \subset \Omega$  and every nonnegative measurable function  $u : \mathbb{R}^n \rightarrow \mathbb{R}$ .

---

<sup>0</sup>**Key words and phrases:** Sobolev spaces, Geometric measure theory, Potential theory.

<sup>0</sup>**2010 Mathematics Subject Classification:** 28A75, 31B15, 46E35.

If the mapping  $\varphi$  possesses the Luzin Hausdorff-Lebesgue measure  $N$ -property (an image of a set of  $\mathcal{H}^{n-1}$ -Hausdorff measure zero has  $n$ -dimensional Lebesgue measure zero), then  $|\varphi(S)| = 0$  and the integral on the right hand side of (1.1) can be rewritten as an integral on  $\mathbb{R}^n$ .

In the case of the second order Sobolev spaces  $W_{p,\text{loc}}^2(\Omega)$ ,  $1 < p < \infty$ , we prove the refined version of the Lipschitz approximation in capacity terms: *Let  $f \in W_{p,\text{loc}}^2(\Omega)$ ,  $1 < p < \infty$ . Then there exists a sequence of closed sets  $\{A_k\}_{k=1}^\infty$ ,  $A_k \subset A_{k+1} \subset \Omega$ , such that the restrictions  $f^*|_{A_k}$  are Lipschitz continuous functions  $p$ -quasieverywhere in  $A_k$  (means that outside a set of  $\text{cap}_p$  zero) and  $\text{cap}_p(S) = 0$ ,  $S = \Omega \setminus \bigcup_{k=1}^\infty A_k$ , where  $\text{cap}_p$  is the  $p$ -capacity.*

Using this Lipschitz approximation we obtain the following capacity version of the change of variables formula in the Lebesgue integral for twice weakly differentiable mappings: Let  $\varphi : \Omega \rightarrow \mathbb{R}^n$  be a mapping which belongs to the Sobolev space  $W_{p,\text{loc}}^2(\Omega; \mathbb{R}^n)$ ,  $1 < p < \infty$ . Then there exists a Borel set  $S \subset \Omega$ ,  $\text{cap}_p(S) = 0$ , such that the mapping  $\varphi : \Omega \setminus S \rightarrow \mathbb{R}^n$  has the Luzin  $N$ -property and the change of variables formula

$$(1.2) \quad \int_A u \circ \varphi(x) |J(x, \varphi)| dx = \int_{\mathbb{R}^n \setminus \varphi(S)} u(y) N_\varphi(A, y) dy,$$

holds for every measurable set  $A \subset \Omega$  and every nonnegative measurable function  $u : \mathbb{R}^n \rightarrow \mathbb{R}$ .

If the mapping  $\varphi$  possesses the Luzin capacity-measure  $N$ -property (an image of a set of capacity zero has Lebesgue measure zero), then  $|\varphi(S)| = 0$  and the integral on the right hand side of (1.2) can be rewritten as an integral on  $\mathbb{R}^n$ . Since for any  $E \subset \mathbb{R}^n$  we have the inequality  $|E| \leq \text{cap}_p(E)$ ,  $1 < p < \infty$  [14], then mappings which possess the Luzin measure  $N$ -property have the Luzin capacity-measure  $N$ -property.

Note, mappings which possess the Luzin capacity-capacity  $N$ -property (an image of a set of capacity zero has capacity zero) and so possess the Luzin capacity-measure  $N$ -property arise, in particular, in the geometric theory of composition operators on Sobolev spaces, see, for example, [8, 21, 26]. The geometric theory of composition operators on Sobolev spaces have applications in the spectral theory of elliptic operators, see, for example, [7]. We note also the generalized quasiconformal mappings, so-called  $Q$ -mappings, which possess the Luzin capacity-capacity  $N$ -property. The theory of  $Q$ -mappings is intensively developed in the last decades. See, for example, [19].

On the base of the refined Lipschitz approximation we obtain the Luzin type theorem for capacity for second order Sobolev spaces. It is known [1, 4, 13, 20] that if  $f \in W_{p,\text{loc}}^1(\Omega)$ , then for each  $\varepsilon > 0$  there exists an open set  $U_\varepsilon$  of  $p$ -capacity less than  $\varepsilon$  such that  $f^*$  is continuous on the set  $\Omega \setminus U_\varepsilon$ . We prove the following refined version: If  $f \in W_{p,\text{loc}}^2(\Omega)$ , then for each  $\varepsilon > 0$  there exists an open set  $U_\varepsilon$  of  $p$ -capacity less than  $\varepsilon$  such that  $f^*$  is Lipschitz continuous on the set  $\Omega \setminus U_\varepsilon$ .

The refined Luzin theorem was obtained in [16]. It was proved that Sobolev functions coincide with Hölder continuous functions on the complement of a set of arbitrary small capacity. This result plays a crucial role in the refined versions of the change of variables formula for the Sobolev mappings [17, 18]. In the case of higher order Sobolev spaces the Luzin type theorem considered in [27]. The refined Luzin theorem on the Hölder continuity of higher order Sobolev spaces was proved in [2]. In the recent work [3] the Luzin type theorem for functions of  $W_1^n(\mathbb{R}^n)$  in the terms of the Hausdorff content was proved.

We prove the following refined version of the Luzin type theorem: *If  $f \in W_{1,\text{loc}}^2(\Omega)$  and  $B \subset \Omega$  is a Borel set such that  $\mathcal{H}^{n-1}(B) < \infty$ , then for every  $\varepsilon > 0$  there exists a Borel set  $B_\varepsilon \subset B$  such that  $f^*|_{B_\varepsilon}$  is Lipschitz continuous  $\mathcal{H}^{n-1}$ -a.e in  $B_\varepsilon$  and  $\mathcal{H}^{n-1}(B \setminus B_\varepsilon) < \varepsilon$ .*

In the case of the second order Sobolev spaces  $W_{p,\text{loc}}^2(\Omega)$ ,  $1 < p < \infty$ , we prove: *If  $f \in W_{p,\text{loc}}^2(\Omega)$ ,  $1 < p < \infty$ , and  $A \subset \Omega$  is a compact set with an additional assumption (see Corollary 2.10), then for every  $\varepsilon > 0$  there exists a Borel set  $A_\varepsilon \subset A$  such that  $f^*|_{A_\varepsilon}$  is Lipschitz continuous  $p$ -quasieverywhere in  $A_\varepsilon$  and  $\text{cap}_p(A \setminus A_\varepsilon) < \varepsilon$ .*

The suggested methods are based in the non-linear potential theory [13, 20] and the Chebyshev type inequality for capacity [4, 22, 23]. This paper is organized as follows: Section 2 contains definitions and we consider Lipschitz approximations in Sobolev spaces  $W_{p,\text{loc}}^2(\Omega)$  and prove Luzin type theorems for Lipschitz  $p$ -quasicontinuity (see Definition 2.5). In Section 3 we prove the change of variables formula for twice weakly differentiable mappings which have the Luzin capacity-measure  $N$ -property.

## 2. LIPSCHITZ APPROXIMATIONS OF SOBOLEV FUNCTIONS

Let  $E \subset \mathbb{R}^n$  be a measurable set. Recall that the Lebesgue space  $L_p(E)$ ,  $1 \leq p < \infty$ , is defined as the space of  $p$ -integrable functions with the norm

$$\|f\|_{L_p(E)} = \left( \int_E |f(x)|^p dx \right)^{1/p}, \quad 1 \leq p < \infty.$$

By  $L_{p,\text{loc}}(E)$  we denote the space of locally  $p$ -integrable functions, means that,  $f \in L_{p,\text{loc}}(E)$  if and only if  $f \in L_p(F)$  for every compact subset  $F \subset E$ .

Let  $\Omega \subset \mathbb{R}^n$  be an open set. The Sobolev space  $W_p^m(\Omega)$ ,  $m \in \mathbb{N}$ ,  $1 \leq p < \infty$ , is defined as the normed space of functions  $f \in L_p(\Omega)$  such that the partial derivatives of order less than or equal to  $m$  exist in the weak sense and belong to  $L_p(\Omega)$ . The space is equipped with the norm

$$\|f\|_{W_p^m(\Omega)} = \sum_{|\alpha| \leq m} \left( \int_\Omega |D^\alpha f(x)|^p dx \right)^{\frac{1}{p}} < \infty,$$

where  $D^\alpha f$  is the weak derivative of order  $\alpha$  of the function  $f$  which is defined by the following formula:

$$\int_\Omega f D^\alpha \eta dx = (-1)^{|\alpha|} \int_\Omega (D^\alpha f) \eta dx, \quad \forall \eta \in C_c^\infty(\Omega),$$

where  $\alpha := (\alpha_1, \alpha_2, \dots, \alpha_n)$  is a multiindex,  $\alpha_i = 0, 1, \dots$ ,  $|\alpha| = \alpha_1 + \alpha_2 + \dots + \alpha_n$ . The space  $C_c^\infty(\Omega)$  is the space of smooth functions with compact support in  $\Omega$ . The Sobolev space  $W_{p,\text{loc}}^m(\Omega)$  is defined as follows:  $f \in W_{p,\text{loc}}^m(\Omega)$  if and only if  $f \in W_p^m(U)$  for every open and bounded set  $U \subset \Omega$  such that  $\bar{U} \subset \Omega$ , where  $\bar{U}$  stands for the topological closure of the set  $U$ .

Let us recall the definition of the capacity [4, 13, 20]. Suppose  $\Omega$  is an open set in  $\mathbb{R}^n$  and  $F \subset \Omega$  is a compact set. The  $p$ -capacity of  $F$  with respect to  $\Omega$  is defined by

$$(2.1) \quad \text{cap}_p(F; \Omega) = \inf \{ \|\nabla f\|_{L_p(\Omega)}^p \},$$

where the inferior is taken over all  $f \in C_c^\infty(\Omega)$  such that  $f \geq 1$  on  $F$ . If  $U \subset \Omega$  is an open set, we define

$$(2.2) \quad \text{cap}_p(U; \Omega) = \sup \{ \text{cap}_p(F; \Omega) : F \subset U, F \text{ is compact} \}.$$

In the case of an arbitrary set  $E \subset \Omega$  we define

$$(2.3) \quad \text{cap}_p(E; \Omega) = \inf \{ \text{cap}_p(U; \Omega) : E \subset U \subset \Omega, U \text{ is open} \}.$$

The  $p$ -capacity is an outer measure on  $\Omega$ , i.e it is a set function which is defined on every subset of  $\Omega$  such that  $\text{cap}_p(\emptyset; \Omega) = 0$  and it is  $\sigma$ -subadditive, means that, if  $E \subset \bigcup_{l \in \mathbb{N}} E_l$ , then  $\text{cap}_p(E; \Omega) \leq \sum_{l \in \mathbb{N}} \text{cap}_p(E_l; \Omega)$ . We write for short  $\text{cap}_p(E) = \text{cap}_p(E; \mathbb{R}^n)$ ,  $E \subset \mathbb{R}^n$ .

The notion of the  $p$ -capacity permits us to refine the notion of Sobolev functions [11, 20]. Let  $f \in L_{1,\text{loc}}(\Omega)$ . The **precise representative** of  $f$  is defined by

$$(2.4) \quad f^* : \Omega \rightarrow \mathbb{R}, \quad f^*(x) := \begin{cases} \lim_{\epsilon \downarrow 0} f_{B(x,\epsilon)}, & \text{if the limit exists;} \\ 0, & \text{otherwise;} \end{cases}$$

where

$$(2.5) \quad f_{B(x,\epsilon)} := \int_{B(x,\epsilon)} f(y) dy = \frac{1}{|B(x,\epsilon)|} \int_{B(x,\epsilon)} f(y) dy.$$

Recall that a function  $f$  is termed **p-quasicontinuous** in an open set  $\Omega$  if and only if for every  $\varepsilon > 0$  there exists an open set  $U_\varepsilon \subset \Omega$  such that the  $p$ -capacity of  $U_\varepsilon$  is less than  $\varepsilon$  and on the set  $\Omega \setminus U_\varepsilon$  the function  $f$  is continuous. If  $f \in W_{p,\text{loc}}^1(\Omega)$ , then  $f^*$  is  $p$ -quasicontinuous and there exists a Borel set  $E$  such that  $\text{cap}_p(E) = 0$  and  $f^*(x) = \lim_{\epsilon \rightarrow 0^+} f_{B(x,\epsilon)}$ ,  $\forall x \in \Omega \setminus E$  [4, 11, 13, 20]. In case  $f \in W_{p,\text{loc}}^1(\Omega)$  the refined function  $f^*$  is called **the unique quasicontinuous representation** (or **the canonical representation**) of  $f$ .

**2.1. Lipschitz  $p$ -quasicontinuous functions.** Let us recall the following two theorems from the theory of Sobolev spaces, see for example [4, 14, 20]:

**Theorem 2.1.** (The local Poincaré inequality for functions of  $W_{1,\text{loc}}^1(\mathbb{R}^n)$ ).  
There exists a constant  $C = C(n)$  such that

$$(2.6) \quad \int_{B(x,r)} |f(y) - f_{B(x,r)}| dy \leq Cr \int_{B(x,r)} |\nabla f(y)| dy,$$

for every ball  $B(x,r) \subset \mathbb{R}^n$  and every  $f \in W_1^1(B(x,r))$ .

**Theorem 2.2.** (The Chebyshev type inequality for capacity).  
Assume  $p \geq 1$ ,  $f \in W_p^1(\mathbb{R}^n)$  and let  $\alpha > 0$ . Define

$$R_\alpha := \left\{ x \in \mathbb{R}^n \mid \sup_{\epsilon > 0} f_{B(x,\epsilon)} \leq \alpha \right\}.$$

Then

$$\text{cap}_p(\mathbb{R}^n \setminus R_\alpha) \leq \frac{C(n,p)}{\alpha^p} \|\nabla f\|_{L_p(\mathbb{R}^n)}^p.$$

The following theorem gives us an approximation of twice weakly differentiable functions by Lipschitz continuous functions.

**Theorem 2.3.** Let  $\Omega \subset \mathbb{R}^n$  be an open set and  $f \in W_{p,\text{loc}}^2(\Omega)$ ,  $1 \leq p < \infty$ . Then there exists a sequence of closed sets  $\{C_k\}_{k=1}^\infty$  such that for every  $k = 1, 2, \dots$ ,  $C_k \subset C_{k+1} \subset \Omega$ , the restriction  $f^*|_{C_k}$  is a Lipschitz continuous function defined  $p$ -quasieverywhere in  $C_k$  and

$$(2.7) \quad \text{cap}_p \left( \Omega \setminus \bigcup_{k=1}^\infty C_k \right) = 0.$$

*Proof.* Assume first  $f \in W_p^2(\mathbb{R}^n)$ . Then for any  $\alpha > 0$  we define a set

$$(2.8) \quad R_\alpha = \left\{ x \in \mathbb{R}^n \mid \sup_{r>0} |\nabla f|_{B(x,r)} \leq \alpha \right\}.$$

The set  $R_\alpha$  is closed: fix an arbitrary number  $\varepsilon > 0$  and let  $\{x_i\}_{i=1}^\infty \subset R_\alpha$  be a sequence of points such that  $x_i \rightarrow x$  as  $i \rightarrow \infty$ . Since for every  $r > 0$  the function  $z \mapsto |\nabla f|_{B(z,r)}$  is continuous, there exists a number  $r_0 > 0$  such that

$$||\nabla f|_{B(z,r)} - |\nabla f|_{B(x,r)}| \leq \varepsilon, \quad \forall z \in B(x, r_0).$$

Thus, for sufficiency large numbers  $i$  such that  $x_i \in B(x, r_0)$  we have

$$|\nabla f|_{B(x,r)} \leq \varepsilon + |\nabla f|_{B(x_i,r)} \leq \varepsilon + \alpha.$$

Hence  $\sup_{r>0} |\nabla f|_{B(x,r)} \leq \varepsilon + \alpha$  and since  $\varepsilon > 0$  is arbitrary we get  $x \in R_\alpha$  and so  $R_\alpha$  is a closed set. Note, in addition, that by the definition of the sets  $R_\alpha$ ,  $\alpha > 0$ , we have  $R_{\alpha_1} \subset R_{\alpha_2}$  for numbers  $\alpha_2 \geq \alpha_1 > 0$ . Now, let  $x \in R_\alpha$ . Since  $|\nabla f| \in L_p(\mathbb{R}^n)$ , then by the local Poincaré inequality we have

$$(2.9) \quad \int_{B(x,r)} |f(y) - f_{B(x,r)}| dy \leq C(n,p)r \int_{B(x,r)} |\nabla f(y)| dy \leq Cr\alpha.$$

Hence

$$(2.10) \quad |f_{B(x,r/2^{k+1})} - f_{B(x,r/2^k)}| \leq \int_{B(x,r/2^{k+1})} |f(y) - f_{B(x,r/2^k)}| dy \\ \leq 2^n \int_{B(x,r/2^k)} |f(y) - f_{B(x,r/2^k)}| dy \leq \frac{Cr\alpha}{2^k},$$

where  $C$  is a constant which is dependent on  $n$  and  $p$  only. Since  $f \in W_p^2(\mathbb{R}^n)$ , then  $f \in W_p^1(\mathbb{R}^n)$  and [11, 20] there exists a Borel set  $E \subset \mathbb{R}^n$  such that  $\text{cap}_p(E) = 0$  and

$$(2.11) \quad \lim_{r \rightarrow 0^+} f_{B(x,r)} = f^*(x) \quad \text{for all } x \in \mathbb{R}^n \setminus E.$$

Therefore, for every  $x \in R_\alpha \setminus E$  it follows that

$$(2.12) \quad |f^*(x) - f_{B(x,r)}| = \left| \sum_{k=0}^\infty [f_{B(x,r/2^{k+1})} - f_{B(x,r/2^k)}] \right| \leq Cr\alpha.$$

Now, we take arbitrary points  $x, y \in R_\alpha$  such that  $x \neq y$  and we set  $r = |x - y|$ . Then

$$(2.13) \quad |f_{B(x,r)} - f_{B(y,r)}| \leq \int_{B(x,r) \cap B(y,r)} (|f_{B(x,r)} - f(z)| + |f(z) - f_{B(y,r)}|) dz \\ \leq 2^n \left( \int_{B(x,r)} |f_{B(x,r)} - f(z)| dz + \int_{B(y,r)} |f(z) - f_{B(y,r)}| dz \right) \leq Cr\alpha.$$

By the results (2.12) and (2.13) and the triangle inequality we obtain

$$(2.14) \quad |f^*(x) - f^*(y)| \leq C\alpha|x - y|, \quad \text{for any } x, y \in R_\alpha \setminus E.$$

In particular, for every integer  $k \geq 1$  the function  $f^*$  is Lipschitz continuous on  $R_k \setminus E$ ,  $\text{cap}_p(E) = 0$ . Now, since  $f \in W_p^2(\mathbb{R}^n)$ , then  $\nabla f \in W_p^1(\mathbb{R}^n; \mathbb{R}^n)$  and by the Chebyshev type inequality for capacity [4, 14] we have the following estimate

$$(2.15) \quad \text{cap}_p(\mathbb{R}^n \setminus R_\alpha) \leq \frac{C(n, p)}{\alpha^p} \|\nabla^2 f\|_{L_p(\mathbb{R}^n)}^p, \quad \text{for any number } \alpha > 0.$$

Therefore, by (2.15)

$$(2.16) \quad \text{cap}_p(\mathbb{R}^n \setminus \cup_{k=1}^\infty R_k) = 0.$$

It completes the proof in the case  $f \in W_p^2(\mathbb{R}^n)$ .

Now, assume  $f \in W_{p, \text{loc}}^2(\Omega)$ . Let  $\Omega_k \subset \Omega_{k+1} \subset \Omega$  be a nested sequence for  $\Omega$ , i.e.,  $\forall k$ ,  $\Omega_k$  is an open set such that  $\overline{\Omega}_k \subset \Omega_{k+1}$ ,  $\overline{\Omega}_k$  is compact and the union of the sets  $\Omega_k$  equals to  $\Omega$ .

For each  $k$  we choose a function  $\zeta_k \in C_c^\infty(\Omega_{k+1})$  such that  $\zeta_k \equiv 1$  on  $\overline{\Omega}_k$ .

It follows that  $f_k := f\zeta_k \in W_p^2(\Omega_{k+1})$ ,  $\text{Supp}(f_k) \subset \Omega_{k+1}$ , where  $\text{Supp}(f_k)$  denotes the support of  $f_k$ . We extend  $f_k$  from  $\Omega_{k+1}$  to  $\mathbb{R}^n$  by zero and denote the extension again by  $f_k$ . Then  $f_k \in W_p^2(\mathbb{R}^n)$  and we get by the previous case a non decreasing sequence of closed sets  $\{A_l^k\}_{l=1}^\infty$  such that  $f_k^*|_{A_l^k}$  is Lipschitz continuous in  $A_l^k$  up to a set of  $p$ -capacity zero and  $\text{cap}_p(\mathbb{R}^n \setminus \cup_{l=1}^\infty A_l^k) = 0$ . Let us choose a sequence of numbers  $\alpha_k \in \mathbb{N}$  such that

$$(2.17) \quad \alpha_k^p \geq 2^k \|\nabla^2 f_k\|_{L_p(\mathbb{R}^n)}^p.$$

Using the Chebyshev type inequality for capacity we have

$$(2.18) \quad \text{cap}_p(\mathbb{R}^n \setminus A_{\alpha_k}^k) \leq \frac{C(n, p)}{\alpha_k^p} \|\nabla^2 f_k\|_{L_p(\mathbb{R}^n)}^p \leq \frac{C(n, p)}{2^k}.$$

For a sequence  $\{\alpha_k\}_{k=1}^\infty$  as above let us define a sequence of sets

$$(2.19) \quad \{B_l\}_{l=1}^\infty, \quad B_l = \bigcap_{k=l}^\infty A_{\alpha_k}^k.$$

It follows that

$$(2.20) \quad \mathbb{R}^n \setminus \bigcup_{l=1}^\infty B_l = \mathbb{R}^n \setminus \liminf_{l \rightarrow \infty} A_{\alpha_l}^l = \limsup_{l \rightarrow \infty} (\mathbb{R}^n \setminus A_{\alpha_l}^l) = \bigcap_{j=1}^\infty \bigcup_{k=j}^\infty (\mathbb{R}^n \setminus A_{\alpha_k}^k).$$

Therefore, for any  $j \geq 1$  we obtain

$$(2.21) \quad \text{cap}_p\left(\mathbb{R}^n \setminus \bigcup_{l=1}^\infty B_l\right) \leq \sum_{k=j}^\infty \text{cap}_p(\mathbb{R}^n \setminus A_{\alpha_k}^k) \leq C(n, p) \sum_{k=j}^\infty \frac{1}{2^k}.$$

Thus,

$$(2.22) \quad \text{cap}_p\left(\mathbb{R}^n \setminus \bigcup_{l=1}^\infty B_l\right) = 0.$$

Now, define a sequence of sets

$$(2.23) \quad \{C_l\}_{l=1}^\infty, \quad C_l = B_l \cap \overline{\Omega}_l.$$

The sequence  $\{C_l\}_{l=1}^\infty$  is monotone increasing since  $\{B_l\}_{l=1}^\infty, \{\overline{\Omega}_l\}_{l=1}^\infty$  are monotone increasing.  $C_l$  is closed as an intersection of closed sets  $B_l, \overline{\Omega}_l$ . Since

$$(2.24) \quad \Omega \setminus \bigcup_{l=1}^\infty C_l = \Omega \setminus \bigcup_{l=1}^\infty B_l \cap \overline{\Omega}_l = \Omega \setminus \bigcup_{l=1}^\infty B_l \cap \Omega = \Omega \cap \left( \mathbb{R}^n \setminus \bigcup_{l=1}^\infty B_l \right) \subset \mathbb{R}^n \setminus \bigcup_{l=1}^\infty B_l,$$

then

$$(2.25) \quad \text{cap}_p \left( \Omega \setminus \bigcup_{l=1}^\infty C_l \right) = 0.$$

At last, since  $C_l \subset A_{\alpha_l}^l \cap \overline{\Omega}_l$  and  $f_l^*|_{A_{\alpha_l}^l \cap \overline{\Omega}_l} = f^*|_{A_{\alpha_l}^l \cap \overline{\Omega}_l}$  is Lipschitz continuous in  $A_{\alpha_l}^l \cap \overline{\Omega}_l$  up to a set of  $p$ -capacity zero, then  $f^*|_{C_l}$  is a Lipschitz continuous function in  $C_l$  up to a set of  $p$ -capacity zero. It completes the proof of the theorem.  $\square$

We denote by  $\mathcal{H}^s, s \geq 0$ , the  $s$ -dimensional Hausdorff measure. For Borel sets  $B \subset \mathbb{R}^n$  such that  $\mathcal{H}^{n-1}(B) < \infty$  it follows that

$$(2.26) \quad \mathcal{H}^{n-1}(B) = 0 \iff \text{cap}_1(B) = 0.$$

For a proof of this relation between the (outer) measures  $\mathcal{H}^{n-1}, \text{cap}_1$  see for example Theorem 3 in Section 5.6 in the book [4].

The following corollary is an immediate consequence of Theorem 2.3 and (2.26).

**Corollary 2.4.** *Let  $\Omega \subset \mathbb{R}^n$  be an open set and  $f \in W_{1,\text{loc}}^2(\Omega)$ . Then there exists a sequence of closed sets  $\{C_k\}_{k=1}^\infty$  such that for every  $k = 1, 2, \dots$ ,  $C_k \subset C_{k+1} \subset \Omega$ , the restriction  $f^*|_{C_k}$  is a Lipschitz continuous function defined 1-quasieverywhere in  $C_k$  and*

$$(2.27) \quad \mathcal{H}^{n-1} \left( \Omega \setminus \bigcup_{k=1}^\infty C_k \right) = 0.$$

Now we define the notion of Lipschitz  $p$ -quasicontinuous functions.

**Definition 2.5.** Let  $\Omega \subset \mathbb{R}^n$  be an open set and let  $f : \Omega \rightarrow \mathbb{R}$  be a function. The function  $f$  is called  **$p$ -Lipschitz quasicontinuous** if for each  $\varepsilon > 0$  there exists an open set  $V \subset \Omega$  such that

$$(2.28) \quad \text{cap}_p(V) \leq \varepsilon$$

and

$$(2.29) \quad f|_{\Omega \setminus V} \text{ is Lipschitz continuous.}$$

The following corollary is an immediate consequence of the previous theorem.

**Corollary 2.6.** *Let  $f \in W_p^2(\mathbb{R}^n), 1 \leq p < \infty$ . Then  $f^*$  is  $p$ -Lipschitz quasicontinuous.*

*Proof.* Fix  $\varepsilon > 0$ . By (2.15) there exists a big enough  $\alpha > 0$  such that  $\text{cap}_p(\mathbb{R}^n \setminus (R_\alpha \setminus E)) \leq \frac{\varepsilon}{2}$  and  $f^*$  is Lipschitz continuous in  $R_\alpha \setminus E$ , where  $R_\alpha$  and  $E$  are the same as in the proof of Theorem 2.3. By the outer regularity of  $p$ -capacity (2.3) there exists an open set  $V$  such that  $\mathbb{R}^n \setminus (R_\alpha \setminus E) \subset V$ ,  $\text{cap}_p(V) \leq \varepsilon$  and  $f^*$  is Lipschitz continuous in  $\mathbb{R}^n \setminus V$ .  $\square$

## 2.2. The refined Luzin type theorem for functions of second order Sobolev spaces.

**Theorem 2.7.** *Let  $\Omega \subset \mathbb{R}^n$  be an open set and  $f \in W_{1,\text{loc}}^2(\Omega)$ . Let  $B \subset \Omega$  be a Borel set such that  $\mathcal{H}^{n-1}(B) < \infty$ . Then for any real number  $\varepsilon > 0$  there exists a Borel set  $B_\varepsilon \subset B$  such that  $f^*|_{B_\varepsilon}$  is Lipschitz continuous in  $B_\varepsilon$  up to a set of 1-capacity zero and  $\mathcal{H}^{n-1}(B \setminus B_\varepsilon) < \varepsilon$ .*

*Proof.* By Theorem 2.3, since  $f \in W_{1,\text{loc}}^2(\Omega)$ , there exists a sequence of closed sets  $B_k \subset B_{k+1} \subset \Omega$  such that  $f^*|_{B_k}$  is Lipschitz continuous in  $B_k$  up to a set of 1-capacity zero and  $\text{cap}_1(\Omega \setminus \bigcup_{k=1}^\infty B_k) = 0$ . Since

$$(2.30) \quad \text{cap}_1 \left( B \setminus \bigcup_{k=1}^\infty B_k \right) = 0,$$

then

$$(2.31) \quad \mathcal{H}^{n-1} \left( B \setminus \bigcup_{k=1}^\infty B_k \right) = 0.$$

By the assumption that  $\mathcal{H}^{n-1}(B) < \infty$ , we have

$$(2.32) \quad \lim_{l \rightarrow \infty} \mathcal{H}^{n-1} \left( B \setminus \bigcup_{k=1}^l B_k \right) = 0.$$

Choose a big enough natural number  $l$  such that  $\mathcal{H}^{n-1}(B \setminus \bigcup_{k=1}^l B_k) < \varepsilon$  and set  $B_\varepsilon := \bigcup_{k=1}^l B_k \cap B$ . It follows that  $B_\varepsilon \subset B$  is a Borel set for which  $f^*|_{B_\varepsilon}$  is Lipschitz continuous in  $B_\varepsilon$  up to a set of 1-capacity zero and  $\mathcal{H}^{n-1}(B \setminus B_\varepsilon) < \varepsilon$ .  $\square$

**Remark 2.8.** *If we assume in Theorem 2.7 that the set  $B$  is closed, then we see from the proof of this theorem that we can also choose the set  $B_\varepsilon$  to be closed.*

Before we introduce another capacity version of Luzin's theorem we recall that for a monotone decreasing sequence of compact sets  $C_{k+1} \subset C_k \subset \mathbb{R}^n$  we have

$$\lim_{k \rightarrow \infty} \text{cap}_p(C_k) = \text{cap}_p \left( \bigcap_{k=1}^\infty C_k \right).$$

The proof of this formula can be found, for example, in [4]. We denote by  $\text{Int}(B)$  the topological interior of a set  $B$ .

**Theorem 2.9.** *Let  $\Omega \subset \mathbb{R}^n$  be an open set and  $f : \Omega \rightarrow \mathbb{R}$  be a function. Assume the existence of a sequence of closed sets  $\{B_k\}_{k=1}^\infty$ ,  $B_k \subset B_{k+1} \subset \Omega$  for which the restrictions  $f|_{B_k}$  are Lipschitz continuous functions on the sets  $B_k$  and*

$$\text{cap}_p \left( \Omega \setminus \bigcup_{k=1}^\infty B_k \right) = 0.$$

*Let  $A$  be a compact set such that  $A \subset \bigcup_{k=1}^\infty \text{Int}(B_k)$ . Then for every  $\varepsilon > 0$  there exists a Borel set  $A_\varepsilon \subset A$  such that  $f|_{A_\varepsilon}$  is Lipschitz continuous and  $\text{cap}_p(A \setminus A_\varepsilon) \leq \varepsilon$ .*

*Proof.* By assumption there exists a monotone increasing sequence of closed sets  $B_k$  such that  $f|_{B_k}$  is Lipschitz continuous for any integer  $k \geq 1$  and  $\text{cap}_p(\Omega \setminus \bigcup_{k=1}^{\infty} B_k) = 0$ . Thus,

$$(2.33) \quad 0 = \text{cap}_p \left( A \setminus \bigcup_{k=1}^{\infty} \text{Int}(B_k) \right) = \lim_{l \rightarrow \infty} \text{cap}_p \left( A \setminus \bigcup_{k=1}^l \text{Int}(B_k) \right).$$

Fix  $\varepsilon > 0$ . There exists a big enough integer  $l_0$  such that

$$(2.34) \quad \text{cap}_p \left( A \setminus \bigcup_{k=1}^l \text{Int}(B_k) \right) \leq \varepsilon, \quad \forall l \geq l_0.$$

Set  $A_\varepsilon := \bigcup_{k=1}^{l_0} \text{Int}(B_k) \cap A$ . It follows that  $A_\varepsilon \subset A$  is a Borel set such that  $f|_{A_\varepsilon}$  is Lipschitz continuous and  $\text{cap}_p(A \setminus A_\varepsilon) \leq \varepsilon$ .  $\square$

The following Corollary is an immediate consequence of Theorem 2.3 and Theorem 2.9.

**Corollary 2.10.** *Let  $\Omega \subset \mathbb{R}^n$  be an open set and  $f \in W_{p,\text{loc}}^2(\Omega)$ ,  $1 \leq p < \infty$ . There exists a sequence of closed sets  $\{B_k\}_{k=1}^{\infty}$ ,  $B_k \subset B_{k+1} \subset \Omega$  for which the restrictions  $f^*|_{B_k}$  are Lipschitz continuous in  $B_k$  up to a set of  $p$ -capacity zero and*

$$\text{cap}_p \left( \Omega \setminus \bigcup_{k=1}^{\infty} B_k \right) = 0.$$

Moreover, if  $A$  is a compact set such that  $A \subset \bigcup_{k=1}^{\infty} \text{Int}(B_k)$ , then for every  $\varepsilon > 0$  there exists a Borel set  $A_\varepsilon \subset A$  such that  $f^*|_{A_\varepsilon}$  is Lipschitz continuous in  $A_\varepsilon$  up to a set of  $p$ -capacity zero and  $\text{cap}_p(A \setminus A_\varepsilon) \leq \varepsilon$ .

### 3. THE CHANGE OF VARIABLES FORMULA

In this section we will derive from the results of the previous section the change of variables formula for mappings of the class  $W_{p,\text{loc}}^2(\Omega; \mathbb{R}^n)$ .

**Remark 3.1.** *Because differential and geometric properties of mappings  $\varphi : \Omega \rightarrow \mathbb{R}^n$  are defined by their coordinate functions, the previous results on Lipschitz approximations of Sobolev functions are valid for mappings  $\varphi : \Omega \rightarrow \mathbb{R}^n$ . More precisely, one can generalize the results of Section 2 from the case of functions  $f : \Omega \rightarrow \mathbb{R}$  to the case of mappings  $\varphi : \Omega \rightarrow \mathbb{R}^n$  using that a mapping  $\varphi : \Omega \rightarrow \mathbb{R}^n$  is a Sobolev mapping if and only if its coordinate functions are Sobolev functions and it is a Lipschitz mapping if and only if its coordinate functions are Lipschitz functions.*

The following theorem refines the formula of change of variables in the Lebesgue integral in the terms of the non-linear potential theory. The change of variables formula for Lipschitz and Sobolev mappings can be found, for example, in [5, 9, 24].

**Theorem 3.2.** *Let  $\Omega \subset \mathbb{R}^n$  be an open set and let  $\varphi : \Omega \rightarrow \mathbb{R}^n$  be a measurable mapping such that there exists a collection of closed sets  $\{A_k\}_{k=1}^{\infty}$ ,  $A_k \subset A_{k+1} \subset \Omega$  for which the restrictions  $\varphi|_{A_k}$  are Lipschitz continuous mappings on the sets  $A_k$  and*

$$\text{cap}_p \left( \Omega \setminus \bigcup_{k=1}^{\infty} A_k \right) = 0.$$

Then there exists a Borel set  $S \subset \Omega$ ,  $\text{cap}_p(S) = 0$ , such that the mapping  $\varphi : \Omega \setminus S \rightarrow \mathbb{R}^n$  has the Luzin  $N$ -property and the change of variables formula

$$(3.1) \quad \int_A u \circ \varphi(x) |J(x, \varphi)| dx = \int_{\mathbb{R}^n \setminus \varphi(S)} u(y) N_\varphi(A, y) dy,$$

where  $N_\varphi(A, y)$  is the multiplicity function defined as the number of preimages of  $y$  in  $A$  under  $\varphi$ , holds for every measurable set  $A \subset \Omega$  and every nonnegative Lebesgue measurable function  $u : \mathbb{R}^n \rightarrow \mathbb{R}$ .

*Proof.* Denote  $S := \Omega \setminus \bigcup_{k=1}^\infty A_k$ . By assumption  $\text{cap}_p(S) = 0$ . If  $N \subset \Omega$ ,  $|N| = 0$ , then  $\varphi(N \setminus S) = \bigcup_{k=1}^\infty \varphi(A_k \cap N)$  and  $|\bigcup_{k=1}^\infty \varphi(A_k \cap N)| = 0$  since  $\varphi$  is Lipschitz continuous on  $A_k$ .

For each  $k \in \mathbb{N}$  let us extend the Lipschitz continuous mapping  $\varphi|_{A_k}$  to a Lipschitz continuous mapping defined on  $\mathbb{R}^n$ . By the change of variables formula for Lipschitz continuous mappings we have

$$(3.2) \quad \int_{A \cap A_k} u \circ \varphi(x) |J(x, \varphi)| dx = \int_{\varphi(A \cap A_k)} u(y) N_\varphi(A \cap A_k, y) dy,$$

for every measurable set  $A \subset \Omega$  and every nonnegative measurable function  $u : \mathbb{R}^n \rightarrow \mathbb{R}$ . The following two nonnegative sequences of functions are monotone increasing a.e as  $k \rightarrow \infty$ :

$$(3.3) \quad \chi_{A \cap A_k} u \circ \varphi |J(\cdot, \varphi)| \nearrow \chi_{A \setminus S} u \circ \varphi |J(\cdot, \varphi)|,$$

$$(3.4) \quad \chi_{\varphi(A \cap A_k)} u N_\varphi(A \cap A_k, \cdot) \nearrow \chi_{\varphi(A \setminus S)} u N_\varphi(A \setminus S, \cdot).$$

Thus, by the Monotone Convergence Theorem we get

$$(3.5) \quad \int_{A \setminus S} u \circ \varphi(x) |J(x, \varphi)| dx = \int_{\varphi(A \setminus S)} u(y) N_\varphi(A \setminus S, y) dy.$$

Since  $\text{cap}_p(S) = 0$  we have  $|S| = 0$ , and one can replace  $A \setminus S$  on the left hand side of (3.5) by  $A$ . Since the function  $N_\varphi(A \setminus S, \cdot)$  vanishes outside  $\varphi(A)$ , then one can replace  $\varphi(A \setminus S)$  by  $\mathbb{R}^n \setminus \varphi(S)$ . It completes the proof.  $\square$

By Theorem 2.3 Sobolev mappings of the class  $W_{p,\text{loc}}^2(\Omega; \mathbb{R}^n)$  satisfy the conditions of the change of variables formula and so for Sobolev mappings  $\varphi : \Omega \rightarrow \mathbb{R}^n$  of the class  $W_{p,\text{loc}}^2(\Omega; \mathbb{R}^n)$  the change of variables formula (3.1) holds.

In the case of Sobolev mappings of the class  $W_{1,\text{loc}}^2(\Omega; \mathbb{R}^n)$ , by Theorem 2.3 with  $p = 1$ , Theorem 3.2 and (2.26) we obtain the following corollary:

**Corollary 3.3.** *Let  $\Omega \subset \mathbb{R}^n$  be an open set and  $\varphi \in W_{1,\text{loc}}^2(\Omega; \mathbb{R}^n)$ . Then there exists a Borel set  $S \subset \Omega$ ,  $\mathcal{H}^{n-1}(S) = 0$ , such that the mapping  $\varphi : \Omega \setminus S \rightarrow \mathbb{R}^n$  has the Luzin  $N$ -property and the change of variables formula*

$$(3.6) \quad \int_A u \circ \varphi(x) |J(x, \varphi)| dx = \int_{\mathbb{R}^n \setminus \varphi(S)} u(y) N_\varphi(A, y) dy,$$

holds for every measurable set  $A \subset \Omega$  and every nonnegative Lebesgue measurable function  $u : \mathbb{R}^n \rightarrow \mathbb{R}$ .

If the mapping  $\varphi$  possesses the Luzin capacity-measure  $N$ -property (the image of a set of capacity zero has Lebesgue measure zero), then  $|\varphi(S)| = 0$  and the integral on the right hand side of (3.1) can be rewritten as an integral on  $\mathbb{R}^n$ . We summarize it in the following corollary:

**Corollary 3.4.** *Let  $\Omega \subset \mathbb{R}^n$  be an open set and let  $\varphi \in W_{p,\text{loc}}^2(\Omega; \mathbb{R}^n)$  which has the Luzin capacity-measure  $N$ -property. Then the change of variables formula*

$$(3.7) \quad \int_A u \circ \varphi(x) |J(x, \varphi)| dx = \int_{\mathbb{R}^n} u(y) N_\varphi(A, y) dy,$$

holds for every measurable set  $A \subset \Omega$  and every nonnegative Lebesgue measurable function  $u : \mathbb{R}^n \rightarrow \mathbb{R}$ .

### Acknowledgments:

The authors thank Vladimir Gol'dshtein for his useful remarks.

### REFERENCES

- [1] B. Bojarski, P. Hajlasz, Pointwise inequalities for Sobolev functions and some applications, *Studia Math.*, 106 (1993), 77–92.
- [2] B. Bojarski, P. Hajlasz, P. Strzelecki, Improved  $C^{k,\lambda}$  approximation of higher order Sobolev functions in norm and capacity, *Indiana Univ. Mat. J.*, 51 (2002), 507–540.
- [3] J. Bourgain, M. V. Korobkov and J. Kristensen, On the Morse-Sard property and level sets of  $W^{n,1}$  Sobolev functions on  $\mathbb{R}^n$ . *J. Reine Angew. Math.* 700 (2015), 93–112.
- [4] L. C. Evans and R. F. Gariepy, *Measure theory and fine properties of functions*, CRC Press, 2015.
- [5] H. Federer, *Geometric measure theory*, Springer Verlag, Berlin, 1969.
- [6] V. M. Gol'dshtein, Yu. G. Reshetnyak, *Quasiconformal mappings and Sobolev spaces*, Dordrecht, Boston, London: Kluwer Academic Publishers, 1990.
- [7] V. Gol'dshtein, A. Ukhlov, The spectral estimates for the Neumann-Laplace operator in space domains, *Adv. in Math.*, 315 (2017), 166–193.
- [8] V. Gol'dshtein, A. Ukhlov, On the functional properties of weak  $(p, q)$ -quasiconformal homeomorphisms, *J. Math. Sci. (NY)*, 246 (2020), 18–29.
- [9] P. Hajlasz, Change of variables formula under minimal assumptions, *Colloq. Math.*, 64 (1993), 93–101.
- [10] P. Hajlasz, J. Kinnunen, Hölder quasicontinuity of Sobolev functions on metric spaces, *Revista Math. Iberoamericana*, 14 (1998), 601–622
- [11] V. Havin, V. Maz'ya, Nonlinear potential theory, *Russian Math. Surveys*, 27 (1972), 71–148
- [12] J. Heinonen, *Lectures on Analysis on Metric Spaces*, Springer, New York, NY, 2001.
- [13] J. Heinonen, T. Kilpelinen, O. Martio, *Nonlinear Potential Theory of Degenerate Elliptic Equations*, Clarendon Press. Oxford, New York, Tokio. 1993.
- [14] J. Kinnunen, *Sobolev spaces*, Aalto University, Espoo, 2021.
- [15] J. Heinonen, P. Koskela, N. Shanmugalingam, J. T. Tyson, *Sobolev spaces on metric measure spaces: An approach based on upper gradients*, Cambridge University Press. 2015.
- [16] J. Maly, Hölder type quasicontinuity, *Potential Analysis*, 2 (1993), 249–254.
- [17] J. Maly, The area formula for  $W^{1,n}$ -mappings, *Comment. Math. Univ. Carolinae*, 35 (1994), 291–298.
- [18] J. Maly, O. Martio, Luzin's condition (N) and mappings of the class  $W^{1,n}$ , *J. Reine Angew. Math.*, 458 (1995), 19–36.
- [19] O. Martio, V. Ryazanov, U. Srebro and E. Yakubov, *Moduli in modern mapping theory*. Springer Monographs in Mathematics. Springer, New York, (2009).
- [20] V. Maz'ya, *Sobolev spaces: with applications to elliptic partial differential equations*, Springer, Berlin/Heidelberg, 2010.
- [21] A. Ukhlov, On mappings, which induce embeddings of Sobolev spaces, *Siberian Math. J.*, 34 (1993), 185–192.
- [22] S. K. Vodop'yanov, V. M. Chernikov, Sobolev spaces and hypoelliptic equations. I, *Siberian Advances in Mathematics*, 6 (1996), 27–67.

- [23] S. K. Vodop'yanov, V. M. Chernikov, Sobolev spaces and hypoelliptic equations. II, *Siberian Advances in Mathematics*, 6 (1996), 64–96.
- [24] S. K. Vodop'yanov, V. M. Gol'dshtein, Yu. G. Reshetnyak, On geometric properties of functions with generalized first derivatives, *Uspekhi Mat. Nauk* 34 (1979), 17–65.
- [25] S. K. Vodop'yanov, A. D. Ukhlov, Approximately differentiable transformations and change of variables on nilpotent groups, *Siberian Math. J.*, 37 (1996), 79–80.
- [26] S. K. Vodop'yanov, A. D. Ukhlov, Set functions and their applications in the theory of Lebesgue and Sobolev spaces, *Siberian Adv. Math.*, 14 (2004), (2005) 78–125.
- [27] W. P. Ziemer, *Weakly Differentiable Functions*, Springer-Verlag, New York, 1989.

Paz Hashash; Department of Mathematics, Ben-Gurion University of the Negev, P.O.Box 653, Beer Sheva, 8410501, Israel

*E-mail address:* pazhash@post.bgu.ac.il

Alexander Ukhlov; Department of Mathematics, Ben-Gurion University of the Negev, P.O.Box 653, Beer Sheva, 8410501, Israel

*E-mail address:* ukhlov@math.bgu.ac.il