

# On faithful quasi-permutation representations of VZ-groups and Camina $p$ -groups

Sunil Kumar Prajapati\* and Ayush Udeep

ABSTRACT. For a finite group  $G$ , we denote by  $\mu(G)$  and  $c(G)$ , the minimal degree of faithful permutation representation of  $G$  and the minimal degree of faithful representation of  $G$  by quasi-permutation matrices over the complex field  $\mathbb{C}$ . In this paper, we examine  $c(G)$  and  $\mu(G)$  for VZ-groups and Camina  $p$ -groups.

## 1. Introduction

Throughout this paper,  $G$  is a finite group and  $\text{Irr}(G)$  is the set of irreducible complex characters of  $G$ . By a classical Cayley's theorem,  $G$  can be embedded into the Symmetric group  $S_{|G|}$ . It is interesting to search for smallest  $n$  such that  $G$  is contained in  $S_n$ . The minimal faithful permutation degree  $\mu(G)$  of  $G$  is the least positive integer  $n$  such that  $G$  embeds inside  $S_n$ . As we know, every permutation representation can be expressed as  $\mathcal{P} = \{H_1, \dots, H_n\}$  and  $\deg(\mathcal{P}) = \sum_{i=1}^n |G/H_i|$ , where  $H_i$ 's are subgroups of  $G$ . So we may restate the minimal faithful degree of a group  $G$  as

$$\mu(G) = \min \left\{ \sum_{i=1}^n |G/H_i| \mid H_i \leq G \text{ for } i = 1, 2, \dots, n \text{ and } \bigcap_{i=1}^n \bigcap_{x \in G} H_i^x = 1 \right\}$$

(see [5]). In a parallel direction to the definition of permutation group, in 1963, Wong [24] defined a quasi-permutation group in the following way: If  $G$  is a finite linear group of degree  $n$  (i.e. a finite group of automorphisms of an  $n$ -dimensional complex vector space) such that the trace of every element of  $G$  is a non-negative integer, then  $G$  is called a quasi-permutation group. This terminology allows us to have another degree  $c(G)$ , denoting the minimal degree of a faithful representation of  $G$  by complex quasi-permutation matrices. Since every permutation matrix is a quasi-permutation matrix, it is easy to see that  $c(G) \leq \mu(G)$ . Another interesting quantity is  $q(G)$ , which denotes the minimal degree of a faithful representation of  $G$  by quasi-permutation matrices over the rational field  $\mathbb{Q}$ . In past, several researchers have carried out the study of  $c(G)$ ,  $q(G)$  and  $\mu(G)$ . The interested reader may refer to [1, 2, 3, 4, 5, 7, 10, 11, 24, 25]. It is easy to see that, for a finite group  $G$ , we have the following relation between  $\mu(G)$ ,  $c(G)$  and  $q(G)$  :

$$c(G) \leq q(G) \leq \mu(G).$$

In this article, we study  $c(G)$  and  $\mu(G)$  for two types of finite non-abelian groups, type 1: group  $G$  for which the set  $\text{cd}(G) = \{\chi(1) \mid \chi \in \text{Irr}(G)\} = \{1, |G/Z(G)|^{1/2}\}$  and type 2: Camina  $p$ -groups. Groups of type 1 are also known as VZ-group and their nilpotency class is 2. A group is called a VZ-group if all its nonlinear irreducible complex characters vanish off the center. VZ-group was studied in [9, 17, 18]. A  $p$ -group  $G$  is called a Camina  $p$ -group if the conjugacy class of  $g$  is  $gG'$  for all  $g \in G \setminus G'$ . Study of Camina groups was influenced by investigating common generalization of Frobenius groups and extra-special groups, and it has been studied by several researchers ([8, 13, 16]).

In [3], authors gave an algorithm for the computation of  $c(G)$ . They proved that if  $X$  is a subset of  $\text{Irr}(G)$  such that  $\bigcap_{\chi \in X} \ker(\chi) = 1$ , and for any proper subset  $Y$  of  $X$ ,  $\bigcap_{\chi \in Y} \ker(\chi) \neq 1$ , then  $c(G) \leq \xi(1) + m(\xi)$ ,

2010 *Mathematics Subject Classification.* primary 20D15; secondary 20C15, 20B05.

*Key words and phrases.* Camina groups,  $p$ -groups, Quasi-permutation representations.

\* Corresponding author.

where  $\xi = \sum_{\chi \in X} \left[ \sum_{\sigma \in \Gamma(\chi)} \chi^\sigma \right]$ , where  $\Gamma(\chi)$  is the Galois group of  $\mathbb{Q}(\chi)$  over  $\mathbb{Q}$  and  $m(\xi)$  is the absolute value of the least value of  $\sum_{\chi \in X} \sum_{\sigma \in \Gamma(\chi)} \chi^\sigma$  over  $G$  (see [3, Lemma 2.2]). We call a set  $X_G \subset \text{Irr}(G)$ , a minimal faithful quasi-permutation representation of  $G$  if

$$(1) \quad \bigcap_{\chi \in X_G} \ker(\chi) = 1,$$

such that equation (1) does not hold for any proper subset of  $X_G$  and  $c(G) = \xi(1) + m(\xi)$ .

In [3], authors proved that for a finite  $p$ -group  $G$  with  $p$  odd prime,  $q(G) = \mu(G)$ . Moreover, for odd prime  $p$ ,  $c(G) = q(G) = \mu(G)$ . In [5, Theorem 4.12], author proved that if  $G$  is a finite  $p$ -group of class 2 and  $Z(G)$  is cyclic, then  $c(G) = |G/Z(G)|^{1/2} |Z(G)|$ . Note that a finite  $p$ -group of class 2 with cyclic center may or may not be a VZ  $p$ -group. Suppose  $\text{nl}(G)$  and  $\text{lin}(G)$  denote the set of all non-linear irreducible complex characters of  $G$  and the set of all linear characters of  $G$ , respectively. In this article, to prove our results, we use the following set up.

SET UP 1. Suppose  $d(G)$  denotes the minimal number of generators of  $G$ . Let  $G$  be a VZ  $p$ -group,  $d(Z(G)) = r$  and  $d(G') = k$ . Here  $G' \leq Z(G)$  and  $G'$  is elementary abelian  $p$ -group (by Lemma 14). Then we can choose  $a_1, a_2, \dots, a_r \in G$  such that

$$(2) \quad Z(G) = \langle a_1, a_2, \dots, a_r \rangle = H_1 \times \dots \times H_r \cong C_{p^{l_1}} \times C_{p^{l_2}} \times \dots \times C_{p^{l_r}}, \quad G' = \langle a_1^{p^{l_1-1}}, \dots, a_r^{p^{l_r-1}} \rangle,$$

where  $H_i = \langle a_i \rangle$  and

$$(3) \quad \frac{Z(G)}{G'} = \langle a_1 G', \dots, a_k G', a_{k+1} G', \dots, a_r G' \rangle \cong C_{p^{l_1-1}} \times \dots \times C_{p^{l_k-1}} \times C_{p^{l_{k+1}}} \dots \times C_{p^{l_r}}.$$

Set, for each  $i$  ( $1 \leq i \leq r$ )  $\widehat{H}_i = H_1 \times \dots \times H_{i-1} \times H_{i+1} \times \dots \times H_r$ .

Under the above set up, we have the following theorem.

THEOREM 2. *Let  $G$  be a VZ  $p$ -group, for any prime  $p$ . Suppose  $d(Z(G)) = r$  and  $d(G') = k$ . For each  $i$  ( $1 \leq i \leq k$ ), let  $\chi_i \in \text{nl}(G)$  be such that  $\ker(\chi_i) = \widehat{H}_i$  and for  $k+1 \leq i \leq r$ , let  $\psi_i \in \text{lin}(G)$  be such that  $\ker(\psi_i) \cap Z(G) = \widehat{H}_i$ . Then  $X_G = \{\chi_1, \dots, \chi_k, \psi_{k+1}, \dots, \psi_r\}$  is a minimal faithful quasi-permutation representation of  $G$ . Further, suppose  $|G/\ker(\psi_i)| = p^{m_i}$  for  $k+1 \leq i \leq r$ . Then*

$$(4) \quad c(G) = |G/Z(G)|^{1/2} \sum_{i=1}^k |H_i| + \sum_{i=k+1}^r p^{m_i}.$$

If  $p$  is an odd prime, then  $c(G) = \mu(G) = q(G) = |G/Z(G)|^{1/2} \sum_{i=1}^k |H_i| + \sum_{i=k+1}^r p^{m_i}$ .

For VZ-groups, in [1] authors proved the following result.

LEMMA 3. [1, Lemma 2.2] *If  $G$  is a finite group with  $G' = Z(G)$ , and  $\text{cd}(G) = \{1, |G/Z(G)|^{1/2}\}$ , then  $c(G) = |G/Z(G)|^{1/2} c(Z(G))$ .*

Note that a finite VZ-group is a nilpotent group of class 2 and thus can be written as a direct product of its Sylow subgroups. Since  $\text{cd}(G) = \{1, |G/Z(G)|^{1/2}\}$ , all the Sylow subgroups of  $G$ , except one, are abelian. Now by this observation, group mentioned in Lemma 3 must be a finite  $p$ -group. From Theorem 2, we deduce the following result, which is a generalization of Lemma 3.

COROLLARY 4. *Let  $G$  be a VZ  $p$ -group (prime  $p$ ) such that  $d(G') = d(Z(G))$ . Then*

$$c(G) = |G/Z(G)|^{1/2} c(Z(G)).$$

Since VZ-group is a nilpotent group of class 2, we deduce the following result for VZ-group of odd order.

COROLLARY 5. *Let  $G$  be a VZ group of odd order. Then  $c(G) = \mu(G) = q(G)$ .*

Further, we examine  $c(G)$  for the family of Camina  $p$ -groups. In [8], authors proved that if  $G$  is a finite Camina  $p$ -group, then the nilpotency class of  $G$  is at most 3. It is easy to see that if  $G$  is Camina  $p$ -group of nilpotency class 2, then  $G$  is indeed a VZ-group with  $Z(G) = G'$  and hence by Corollary 4, we get  $c(G) = |G/Z(G)|^{1/2}c(Z(G))$ . For Camina  $p$ -group of nilpotency class 3, we have proved the following result.

**THEOREM 6.** *Let  $G$  be a Camina  $p$ -group of nilpotency class 3 (prime  $p$ ). Then*

$$(5) \quad c(G) = |G/Z(G)|^{1/2}c(Z(G)).$$

Moreover, when  $p$  is an odd prime, we get  $c(G) = \mu(G) = q(G) = |G/Z(G)|^{1/2}c(Z(G))$ .

In Section 2, we briefly summarize the background results that will be used in the course of this paper. In Section 3, we first try to find out  $c(G)$  for VZ  $p$ -group by explicit description of the set  $X_G \subseteq \text{Irr}(G)$ . Since VZ-groups are nilpotent groups, we get  $c(G) = \mu(G)$ , for VZ-groups of odd order. In Section 4, by using the concept of isoclinism, we have shown that: If  $G$  and  $H$  are two isoclinic VZ  $p$ -groups of same order such that  $d(Z(G)) = d(Z(H))$ , then  $c(G) = c(H)$  if and only if  $Z(G)$  is isomorphic to  $Z(H)$  (Corollary 20). By using Corollary 20 and Theorem 2, we have computed  $c(G)$  for all the  $p$ -groups of size  $p^6$  (for an odd prime  $p$ ) (see Table 1, Table 2 and Corollary 25). In the last section, we have calculated  $c(G)$  for Camina  $p$ -group and provided  $X_G$ .

## 2. Basic results on $c(G)$ , $q(G)$ and $\mu(G)$

Suppose  $F$  is a subfield of  $E$ , where  $E$  is a splitting field for  $G$ . Let  $\chi, \psi \in \text{Irr}_E(G) = \text{Irr}(G)$ . We say that  $\chi$  and  $\psi$  are Galois conjugate over  $F$  if  $F(\chi) = F(\psi)$  and there exists  $\sigma \in \text{Gal}(F(\chi)/F)$  such that  $\chi^\sigma = \psi$ , where  $F(\chi)$  denotes the field obtained by adjoining the values  $\chi(g)$ , for all  $g \in G$ , to  $F$ . Indeed, this defines an equivalence relation on  $\text{Irr}_E(G)$ . Moreover, if  $\mathcal{C}$  denotes the equivalence class of  $\chi$  with respect to Galois conjugacy over  $F$ , then  $|\mathcal{C}| = |F(\chi) : F|$  (see [14, Lemma 9.17]).

Suppose  $\chi$  is an irreducible character of a group  $G$  and  $\Gamma(\chi)$  denotes the Galois group of  $\mathbb{Q}(\chi)$  over  $\mathbb{Q}$ . Let  $\mathcal{C}_i$ , for  $0 \leq i \leq r$ , be the Galois conjugacy classes of irreducible characters of the group  $G$ . For  $0 \leq i \leq r$ , let  $\psi_i$  be a representative of the class  $\mathcal{C}_i$  with  $\psi_0 = 1_G$  (trivial character). Let us write  $\Theta_i = \sum_{\chi \in \mathcal{C}_i} \chi$ , and  $K_i = \ker(\psi_i)$ . For  $I \subseteq \{0, 1, \dots, r\}$ , set  $K_I = \bigcap_{i \in I} K_i$ . Under the above set up, we have following definition and results which we use to calculate  $c(G)$ .

**DEFINITION 2.0.1.** *Let  $G$  be a finite group. Let  $\chi$  be an irreducible complex character of  $G$ . Then define*

$$(i) \quad d(\chi) = |\Gamma(\chi)|\chi(1).$$

$$(ii) \quad m(\chi) = \begin{cases} 0 & \text{if } \chi = 1_G, \\ \left| \min \left\{ \sum_{\sigma \in \Gamma(\chi)} \chi^\sigma(g) : g \in G \right\} \right| & \text{otherwise.} \end{cases}$$

**LEMMA 7.** [3, Lemma 2.2] *Let  $G$  be a finite group. Then*

$$c(G) = \min \left\{ \xi(1) + m(\xi) : \xi = \sum_{i \in I} \Theta_i, K_I = 1, \text{ for } I \subseteq \{1, \dots, r\} \text{ and } K_J \neq 1, \text{ if } J \subset I \right\}$$

and

$$q(G) = \min \left\{ \xi(1) + m(\xi) : \xi = \sum_{i \in I} m_{\mathbb{Q}}(\psi_i)\Theta_i, K_I = 1, \text{ for } I \subseteq \{1, \dots, r\} \text{ and } K_J \neq 1, \text{ if } J \subset I \right\},$$

where  $m_{\mathbb{Q}}(\psi_i)$  denotes the Schur index of  $\psi_i$  over  $\mathbb{Q}$ .

If  $G$  is a  $p$ -group ( $p$  odd prime), then by Lemma 7,  $c(G) = q(G)$  because  $m_{\mathbb{Q}}(\psi) = 1$  for every  $\psi \in \text{Irr}(G)$ .

**LEMMA 8.** [2, Corollary 2.6] *Let  $G$  be a  $p$ -group whose center  $Z(G)$  is minimally generated by  $d$  elements. Let  $c(G) = \xi(1) + m(\xi)$  and  $\xi = \sum_{i \in I} \Psi_i$ . Let  $\Psi'_i$ 's satisfy the conditions of the algorithm for  $c(G)$ . Then*

- (i)  $m(\xi) = \frac{1}{p-1} \sum_{i \in I} \Psi_i$ ,
- (ii)  $|I| = d$ .

Suppose  $G$  is an abelian group isomorphic to  $\prod_{i=1}^k C_{p_i^{r_i}}$ , where  $C_{p_i^{r_i}}$ 's are cyclic groups of prime power order  $p_i^{r_i}$ . Define  $T(G) = \sum_{i=1}^k p_i^{r_i}$ . In [4], author proved that  $c(G) = T(G) - n$ , where  $n$  is the largest integer such that,  $C_6^n$  is a direct summand of an abelian group  $G$ . Further, if  $G$  is an abelian  $p$ -group, then  $c(G) = q(G) = \mu(G) = T(G)$ .

**THEOREM 9.** [25, Corollary 2.2] *If  $H, K$  are non-trivial nilpotent groups, then  $\mu(H \times K) = \mu(H) + \mu(K)$ .*

**THEOREM 10.** [3, Theorem 3.2] *Let  $G$  be a finite  $p$ -group. Then  $q(G) = \mu(G)$ . Moreover, if  $p \neq 2$ , then  $c(G) = q(G) = \mu(G)$ .*

**THEOREM 11.** [5, Theorem 4.12] *Let  $G$  be a finite  $p$ -group of class 2 and let  $Z(G)$  be cyclic. Then  $c(G) = c(Z(G))|G/Z(G)|^{1/2}$ .*

### 3. VZ group

Suppose  $G$  is a VZ-group. Then  $\chi(g) = 0$ , for all  $g \in G \setminus Z(G)$  and for all  $\chi \in \text{nl}(G)$ , if and only if  $\text{cd}(G) = \{1, |G/Z(G)|^{1/2}\}$ . Moreover,  $|\text{nl}(G)| = |Z(G)| - \frac{|Z(G)|}{|G'|}$  and  $\text{nl}(G) = \{ |G/Z(G)|^{1/2} \lambda \mid \lambda \in \text{Irr}(Z(G)) \text{ such that } G' \not\subseteq \ker(\lambda) \}$ , where members of  $\text{nl}(G)$  are defined as follows:

$$|G/Z(G)|^{1/2} \lambda(g) := \begin{cases} |G/Z(G)|^{1/2} \lambda(g) & \text{if } g \in Z(G), \\ 0 & \text{otherwise} \end{cases}$$

(see [23, Subsection 3.1]). Obviously,  $\ker(\chi) \leq Z(G)$  for all  $\chi \in \text{nl}(G)$ .

**LEMMA 12.** *Suppose  $G$  is a VZ group. Then every  $\chi \in \text{nl}(G)$  is fully ramified with respect to  $G/Z(G)$ .*

**PROOF.** Follows from Problem 6.3 of [14]. □

**LEMMA 13.** *Suppose  $G$  is a VZ group. Let  $\chi, \psi \in \text{nl}(G)$ . Then  $\chi$  and  $\psi$  are Galois conjugate if and only if  $\ker(\chi) = \ker(\psi)$ .*

**PROOF.** It follows from the fact that two linear characters of an abelian group are Galois conjugate if and only if their kernels are same. □

Now we quote the group-theoretic information of  $G/Z(G)$  and  $G'$  for a VZ  $p$ -group.

**LEMMA 14.** [19, Lemma 2.4] *Let  $G$  be a VZ  $p$ -group. Then  $G/Z(G)$  and  $G'$  are elementary abelian  $p$ -groups.*

**LEMMA 15.** *Suppose  $G$  is a VZ  $p$ -group and  $d(G') \geq 2$ . Then for all  $\chi \in \text{nl}(G)$ ,  $\ker(\chi)$  intersects with  $G'$  non-trivially.*

**PROOF.** By the above observation, if  $\chi \in \text{nl}(G)$ , then  $\chi = |G/Z(G)|^{1/2} \lambda$  for some  $\lambda \in \text{Irr}(Z(G))$  such that  $G' \not\subseteq \ker(\lambda)$  and  $\ker(\chi) = \ker(\lambda) \leq Z(G)$ . On the contrary, suppose  $\ker(\chi) \cap G' = 1$ . Consider the natural group homomorphism  $\eta : Z(G) \rightarrow Z(G)/\ker(\chi)$  such that  $g \mapsto g \ker(\chi)$ . Take the restriction  $\eta \downarrow_{G'} : G' \rightarrow Z(G)/\ker(\chi)$ . Under the assumption  $\ker(\chi) \cap G' = 1$ ,  $\eta \downarrow_{G'}$  is injective group homomorphism and thus  $G'$  is isomorphic to a subgroup of the cyclic group  $Z(G)/\ker(\chi)$ . This is a contradiction because  $d(G') \geq 2$ . □

**LEMMA 16.** *Suppose  $G$  is a VZ  $p$ -group. Let  $G' = \langle g_1, g_2, \dots, g_k \rangle$ . Then for any  $\chi \in \text{nl}(G)$ , there exists  $g_i$  such that  $\langle g_i \ker(\chi) \rangle$  is a subgroup of cyclic group  $Z(G)/\ker(\chi)$ .*

**PROOF.** Suppose  $G' = \langle g_1, g_2, \dots, g_k \rangle \cong C_p \times C_p \times \dots \times C_p$  ( $k$  copy of  $C_p$ ). Let  $\chi \in \text{nl}(G)$ . Then there exists  $g \in G'$  such that  $g = g_1^{m_1} g_2^{m_2} \dots g_k^{m_k}$  for some  $m_i$  ( $1 \leq m_i \leq p$  and  $1 \leq i \leq k$ ) and

$$\chi(g) = \chi(1) \lambda(g_1^{m_1} g_2^{m_2} \dots g_k^{m_k}) = \chi(1) \lambda(g_1^{m_1}) \lambda(g_2^{m_2}) \dots \lambda(g_k^{m_k}) \neq \chi(1),$$

where  $\lambda \in \text{Irr}(Z(G))$  such that  $G' \not\leq \ker(\lambda)$ . This implies that there exists  $g_i^{m_i}$  such that  $\lambda(g_i^{m_i}) \neq 1$ , i.e.  $\chi(g_i^{m_i}) \neq \chi(1)$  for some  $i$ . Since order of  $g_i$  is  $p$ , we get  $\chi(g_i) \neq \chi(1)$ . Since  $Z(G)/\ker(\chi)$  is cyclic group,  $\langle g_i \ker(\chi) \rangle \leq Z(G)/\ker(\chi)$ .  $\square$

LEMMA 17. *Let  $G$  be a VZ  $p$ -group with  $d(Z(G)) = r$  and  $d(G') = k$ . Let  $X$  be a set of  $r$  irreducible complex characters of  $G$  such that*

$$\bigcap_{\chi \in X} \ker(\chi) = 1.$$

*Then the number of linear characters in  $X$  must be  $r - k$ . In fact, the set  $X_G$ , which represents a minimal faithful quasi-permutation representation of  $G$ , must contain  $k$  many non-linear irreducible characters and  $r - k$  many linear characters of  $G$ .*

PROOF. We will prove this lemma in two steps. In the first step, we will show that there are at least  $k$  many non-linear irreducible characters in  $X$ . In the second step, we will show that the number of non-linear irreducible characters in the set  $X$  is equal to  $k$ .

**Step 1:** We will prove this part in cases.

**Case 1** ( $d(G') = 1$ , i.e.  $G' \cong C_p$ ). Since all the irreducible characters in the set  $X$  can not be linear, we can take at most  $r - k = r - 1$  linear characters in  $X$ .

**Case 2** ( $d(G') > 1$ ). Let  $d(Z(G)) = r$  and let  $d(G') = k$ . On the contrary, suppose in  $X$ , we have only  $k - 1$  many non-linear irreducible characters of  $G$ , say  $\chi_1, \chi_2, \dots, \chi_{k-1}$ . Since  $d(G') = k$  and  $G'$  is an elementary abelian  $p$ -group,  $G'$  is isomorphic to direct product of  $k$  many copies of cyclic group of order  $p$ . For each  $i$ , since  $Z(G)/\ker(\chi_i)$  is cyclic,  $\ker(\chi_i)$  contains all cyclic components of  $G'$  except one. Hence  $G' \cap (\bigcap_{i=1}^{k-1} \ker(\chi_i)) \neq 1$ . Further,  $G'$  is contained in each  $\ker(\eta)$ , where  $\eta \in \text{lin}(G)$ . This shows that  $\bigcap_{\chi \in X} \ker(\chi) \neq 1$ , which is a contradiction. This proves that in the set  $X$ , we require at least  $k$  many non-linear irreducible characters.

**Step 2:** Now we will show that the number of non-linear irreducible characters in the set  $X$  must be  $k$ . In the view of Set Up 1, we have  $a_1, a_2, \dots, a_r \in G$  such that

$$Z(G) = \langle a_1, a_2, \dots, a_r \rangle = H_1 \times \dots \times H_r \cong C_{p^{t_1}} \times C_{p^{t_2}} \times \dots \times C_{p^{t_r}}, \quad G' = \langle a_1^{p^{t_1-1}}, \dots, a_k^{p^{t_k-1}} \rangle,$$

where  $H_i = \langle a_i \rangle$  and

$$\frac{Z(G)}{G'} = \langle a_1 G', \dots, a_k G', a_{k+1} G', \dots, a_r G' \rangle \cong C_{p^{t_1-1}} \times \dots \times C_{p^{t_k-1}} \times C_{p^{t_{k+1}}} \dots \times C_{p^{t_r}}.$$

By Lemma 16, without loss of generality, we can choose  $k$  many non-linear irreducible characters  $\chi_1, \dots, \chi_k$  such that  $\langle a_i^{p^{t_i-1}} \ker(\chi_i) \rangle \leq Z(G)/\ker(\chi_i)$ , for  $1 \leq i \leq k$ . This shows that, for each  $i$ , we have  $\langle a_i \ker(\chi_i) \rangle \leq Z(G)/\ker(\chi_i)$  and no power of  $a_i$  belongs to  $\ker(\chi_i)$ . Since  $|\langle a_i \ker(\chi_i) \rangle| = |\langle a_i \rangle|$ , we have  $Z(G)/\ker(\chi_i) \cong \langle a_i \rangle$  for  $1 \leq i \leq k$ . Now suppose if we add one more non-linear irreducible character, say  $\chi_{k+1}$ , in the set  $X$ , then by Lemma 16, we get  $\langle a_i^{p^{t_i-1}} \ker(\chi_{k+1}) \rangle \leq Z(G)/\ker(\chi_{k+1})$ , for some fixed  $i$ ,  $1 \leq i \leq k$ . This shows that  $\langle a_i \rangle \cong Z(G)/\ker(\chi_i) \cong Z(G)/\ker(\chi_{k+1})$  and hence  $\chi_i$  and  $\chi_{k+1}$  are Galois conjugate. Thus it is meaningless to keep  $\chi_{k+1}$  in the set  $X$ . Therefore, in the set  $X$ , the number of non-linear characters is equal to  $k$  and the rest  $r - k$  characters in  $X$  are linear characters.

Now, from Lemma 8, we get  $|X_G| = r$ , and from equation (1),  $\bigcap_{\chi \in X_G} \ker(\chi) = 1$ . Therefore,  $X_G$  contains  $k$  many non-linear irreducible characters and  $r - k$  many linear characters of  $G$ .  $\square$

**3.1. Proof of Theorem 2.** Let  $G$  be a VZ  $p$ -group,  $d(Z(G)) = r$  and  $d(G') = k$ . Here  $G' \leq Z(G)$  and by Lemma 14,  $G'$  is elementary abelian  $p$ -group. In the view of Set up 1, we have

$$(6) \quad Z(G) = \langle a_1, a_2, \dots, a_r \rangle \cong C_{p^{t_1}} \times C_{p^{t_2}} \times \dots \times C_{p^{t_r}}, \quad G' = \langle a_1^{p^{t_1-1}}, \dots, a_k^{p^{t_k-1}} \rangle$$

$$\text{and } \frac{Z(G)}{G'} = \langle a_1 G', \dots, a_k G', a_{k+1} G', \dots, a_r G' \rangle \cong C_{p^{t_1-1}} \times \dots \times C_{p^{t_k-1}} \times C_{p^{t_{k+1}}} \dots \times C_{p^{t_r}}.$$

**Step I: To get a set  $X$  of  $r$  irreducible characters which satisfies equation (1).**

For all  $i$  ( $1 \leq i \leq r$ ), let  $K_i = \langle a_i G' \rangle \leq Z(G)/G'$  and let  $\widehat{K}_i = K_1 \times \dots \times K_{i-1} \times K_{i+1} \times \dots \times K_r$ . Choose  $\psi_{k+1}, \dots, \psi_r \in \text{lin}(G/G')$  such that  $\ker(\psi_i \downarrow_{Z(G)/G'}) = \widehat{K}_i$  for all  $i$  ( $k+1 \leq i \leq r$ ). Set  $\eta_i = \psi_i \downarrow_{Z(G)/G'}$  for  $k+1 \leq i \leq r$ . One can view  $\eta_i$  as a faithful linear character of cyclic group  $\frac{Z(G)/G'}{\widehat{K}_i}$ . Observe that

$\ker(\eta_i) = \ker(\psi_i \downarrow_{Z(G)/G'}) = (\ker \psi_i \cap Z(G))/G'$ . On the other hand, let  $H_i = \langle a_i \rangle$  for all  $i$  ( $1 \leq i \leq r$ ) and let  $\widehat{H}_i = H_1 \times \cdots \times H_{i-1} \times H_{i+1} \times \cdots \times H_r$ . Now, for each  $i$  ( $1 \leq i \leq k$ ), let  $\chi_i \in \text{nl}(G)$  such that  $\chi_i \downarrow_{Z(G)} = \chi_i(1)\lambda_i$ , where  $\lambda_i \in \text{lin}(Z(G))$  and  $\ker(\chi_i) = \ker(\lambda_i) = \widehat{H}_i$ . One can view  $\lambda_i$  as a faithful linear character of cyclic group  $Z(G)/\widehat{H}_i$ . Now,

$$\bigcap_{i=k+1}^r \ker(\psi_i \downarrow_{Z(G)/G'}) = \bigcap_{i=k+1}^r \widehat{K}_i = K_1 \times \cdots \times K_k = \langle a_1 G', \dots, a_k G' \rangle = H_1 \times \cdots \times H_k,$$

and

$$\bigcap_{j=1}^k \ker(\chi_j) = \bigcap_{j=1}^k \widehat{H}_j = H_{k+1} \times \cdots \times H_r \leq Z(G).$$

This shows that  $(\bigcap_{i=1}^k \ker(\chi_i)) \cap (\bigcap_{i=k+1}^r \ker(\psi_i)) = 1$ . Therefore,

$$(7) \quad X = \{\chi_i \mid 1 \leq i \leq k\} \cup \{\psi_i \mid k+1 \leq i \leq r\}$$

is a set of  $r$  irreducible characters which satisfies equation (1). Now we will prove that set  $X$  is a minimal faithful quasi-permutation representation of  $G$ .

For the non-linear irreducible characters  $\chi_i$ ,  $1 \leq i \leq k$ , it is easy to observe the following.

- (1) For each  $i$ ,  $\Gamma(\chi_i) = \Gamma(\lambda_i)$  and  $|\Gamma(\lambda_i)| = \phi(p^{l_i}) = p^{l_i} - p^{l_i-1}$ , where  $\phi$  is Euler phi function.
- (2) For each  $i$ ,  $m(\chi_i) = \chi_i(1)m(\lambda_i)$  and  $\chi_i(1) = |G/Z(G)|^{1/2}$ .
- (3) For each  $i$ ,  $m(\lambda_i) = |\min \{ \sum_{\sigma \in \Gamma(\lambda_i)} \lambda_i^\sigma(g) \mid g \in G \}| = |-p^{l_i-1}| = p^{l_i-1}$ .

In the case of the linear characters  $\psi_i$  ( $k+1 \leq i \leq r$ ), we first note that  $\frac{G/G'}{\ker(\psi_i)}$  is cyclic. For each  $i$ , let

$$(8) \quad \frac{G/G'}{\ker(\psi_i)} \cong C_{p^{m_i}}, \quad k+1 \leq i \leq r.$$

Without loss of generality, we can assume that  $p^{m_{k+1}} \leq p^{m_{k+2}} \leq \cdots \leq p^{m_r}$ . Again, one can view  $\psi_i$  as a faithful linear character of cyclic group  $G/\ker(\psi_i)$ , for  $k+1 \leq i \leq r$ . Now  $|\Gamma(\psi_i)|\psi_i(1) = \phi(p^{m_i}) \cdot 1 = p^{m_i} - p^{m_i-1}$ , for each  $i$  ( $k+1 \leq i \leq r$ ). Also, it is easy to observe that  $m(\psi_i) = |\min \{ \sum_{\sigma \in \Gamma(\psi_i)} \psi_i^\sigma(g) \mid g \in G \}| = |-p^{m_i-1}| = p^{m_i-1}$ . Now consider the character

$$\xi = \sum_{i=1}^r \theta_i,$$

where  $\theta_i = \sum_{\sigma \in \Gamma(\chi_i)} \chi_i^\sigma = \chi_i(1) \left( \sum_{\sigma \in \Gamma(\lambda_i)} \lambda_i^\sigma \right) = |G/Z(G)|^{1/2} \left( \sum_{\sigma \in \Gamma(\lambda_i)} \lambda_i^\sigma \right)$ , for  $1 \leq i \leq k$ , and  $\theta_i = \sum_{\sigma \in \Gamma(\psi_i)} \psi_i^\sigma$ , for  $k+1 \leq i \leq r$ . Then by the above observations, we have

$$m(\xi) = |G/Z(G)|^{1/2} \sum_{i=1}^k p^{l_i-1} + \sum_{k+1}^r p^{m_i-1}$$

and

$$\xi(1) = |G/Z(G)|^{1/2} \sum_{i=1}^k p^{l_i-1}(p-1) + \sum_{k+1}^r p^{m_i-1}(p-1).$$

Therefore,

$$c(G) \leq \xi(1) + m(\xi) = |G/Z(G)|^{1/2} \sum_{i=1}^k p^{l_i} + \sum_{k+1}^r p^{m_i} = |G/Z(G)|^{1/2} \sum_{i=1}^k |H_i| + \sum_{i=k+1}^r p^{m_i}.$$

Thus we have got a collection  $X$ , from equation (7), which satisfies equation (1) and we get  $c(G) \leq |G/Z(G)|^{1/2} \sum_{i=1}^k |H_i| + \sum_{i=k+1}^r p^{m_i}$ .

**Step II:**  $\mathbf{c}(G) = \xi(1) + \mathbf{m}(\xi)$ .

In the set  $X$  given by (7), we have exactly  $k$  many non-linear irreducible characters and  $r-k$  many linear characters and also no proper subset of  $X$  satisfies (1). Thus the set  $X$  satisfies Lemma 17. In addition to this, set  $X$  gives a set  $Y = \{\lambda_i \mid 1 \leq i \leq k\} \cup \{\eta_i \mid k+1 \leq i \leq r\}$ , where  $\lambda_i$  for  $1 \leq i \leq k$  and  $\eta_i$  for

$k + 1 \leq i \leq r$  are defined above. It is easy to see that the set  $Y$ , which contains  $r$  irreducible characters of  $Z(G)$  gives  $c(Z(G)) = \widehat{\xi}(1) + m(\widehat{\xi})$ , where  $\widehat{\xi} = \sum_{i=1}^r \widehat{\theta}_i$ , where  $\widehat{\theta}_i = \sum_{\sigma \in \Gamma(\lambda_i)} \lambda_i^\sigma$  for  $1 \leq i \leq k$ , and  $\widehat{\theta}_i = \sum_{\sigma \in \Gamma(\eta_i)} \eta_i^\sigma$ , for  $k + 1 \leq i \leq r$ . This shows that the set  $X$  gives  $c(G)$ , i.e.,

$$c(G) = |G/Z(G)|^{1/2} \sum_{i=1}^k |H_i| + \sum_{i=k+1}^r p^{m_i}.$$

Also, from Theorem 10,  $c(G) = \mu(G) = q(G)$  for  $p \neq 2$ , and thus for VZ  $p$ -group, for odd prime  $p$ ,  $c(G) = \mu(G) = q(G) = |G/Z(G)|^{1/2} \sum_{i=1}^k |H_i| + \sum_{i=k+1}^r p^{m_i}$ . This proves the result.

**Proof of Corollary 4:** From Lemma 17, we know that a minimal faithful quasi-permutation representation  $X_G$  has exactly  $r - k$  linear characters. As in our case,  $r = k$ ,  $X_G$  does not contain any linear characters. Thus, we consider all the  $r$  characters to be non-linear irreducible. Let  $H_i$ ,  $1 \leq i \leq r$ , denote the  $r$  cyclic components of  $Z(G)$  such that  $|H_i| = p^{l_i}$ , and  $\widehat{H}_i = H_1 \times \cdots \times H_{i-1} \times H_{i+1} \times \cdots \times H_r$ . Let  $\chi_i \in \text{Irr}(G/Z(G))$ , for all  $i$ ,  $1 \leq i \leq r$ , be such that  $\chi_i \downarrow_{Z(G)} = \chi_i(1)\lambda_i$ , where  $\lambda_i \in \text{lin}(Z(G))$  is a faithful linear character of  $Z(G)/\widehat{H}_i$ . Then  $|\Gamma(\chi_i)|\chi_i(1) = \phi(|H_i|)|G/Z(G)|^{1/2} = (p^{l_i} - p^{l_i-1})|G/Z(G)|^{1/2}$  and  $m(\chi_i) = |G/Z(G)|^{1/2} | -p^{l_i-1} | = |G/Z(G)|^{1/2} p^{l_i-1}$ , for each  $i$ . Now consider the character  $\xi = \sum_{i=1}^r \theta_i$ , where  $\theta_i = \sum_{\sigma \in \Gamma(\chi_i)} \chi_i^\sigma = \chi_i(1) \left( \sum_{\sigma \in \Gamma(\lambda_i)} \lambda_i^\sigma \right) = |G/Z(G)|^{1/2} \left( \sum_{\sigma \in \Gamma(\lambda_i)} \lambda_i^\sigma \right)$ , for  $1 \leq i \leq r$ . Then by the above observations, we have

$$m(\xi) = |G/Z(G)|^{1/2} \sum_{i=1}^r p^{l_i-1}$$

and

$$\xi(1) = |G/Z(G)|^{1/2} \sum_{i=1}^r p^{l_i-1} (p - 1).$$

Then  $\xi(1) + m(\xi) = |G/Z(G)|^{1/2} \sum_{i=1}^r |H_i| = |G/Z(G)|^{1/2} c(Z(G))$ .

Note that the set  $X = \{\chi_i\}_{i=1}^r$  gives a set  $Y = \{\lambda_i\}_{i=1}^r$ , which is a minimal faithful quasi-permutation representation of  $Z(G)$ . This shows that

$$c(G) = \xi(1) + m(\xi) = |G/Z(G)|^{1/2} c(Z(G)).$$

REMARK 18.

- (1) *The converse of the Corollary 4 is not true in general, i.e., there exists VZ  $p$ -group for which  $c(G) = |G/Z(G)|^{1/2} c(Z(G))$  but  $d(Z(G)) > d(G')$ . For example, consider the group*

$$G = \phi_2(22) = \langle \alpha, \alpha_1, \alpha_2 \mid [\alpha_1, \alpha] = \alpha^p = \alpha_2, \alpha_1^{p^2} = \alpha_2^p = 1 \rangle,$$

*which is a VZ  $p$ -group of order  $p^4$  ( $p$  an odd prime) (see [15, Section 4.4]). Here  $Z(G) = \langle \alpha^p, \alpha_1^p \rangle \cong C_p \times C_p$ ,  $G' = \langle \alpha^p \rangle \cong C_p$  and  $|G/Z(G)|^{1/2} = p$ . Thus,  $d(Z(G)) > d(G')$ . However,  $c(G) = 2p^2 = |G/Z(G)|^{1/2} c(Z(G))$  (we have provided the details of the computation of  $c(\phi_2(22))$  in subsection 4.1).*

- (2) *There exists VZ  $p$ -group such that  $d(Z(G)) > d(G')$  but  $c(G) < |G/Z(G)|^{1/2} c(Z(G))$ . For example, consider the group*

$$G = \phi_2(321)e = \phi_2(21) \times C_{p^3} = \langle \alpha, \alpha_1, \alpha_2 \mid [\alpha_1, \alpha] = \alpha_2, \alpha^p = \alpha_2, \alpha_1^p = \alpha_2^p = 1 \rangle \times C_{p^3},$$

*which is a VZ  $p$ -group of order  $p^6$  ( $p$  an odd prime) (see [15, Section 4.6]). Now, let  $G_1 = \phi_2(21)$ . Then  $Z(G_1) = G'_1 = \langle \alpha^p \rangle \cong C_p$ . As  $G = G_1 \times C_{p^3}$ ,  $c(Z(G)) = c(Z(G_1)) + c(C_{p^3}) = p + p^3$ . Now, from Corollary 4,  $c(G_1) = |G_1/Z(G_1)|^{1/2} c(Z(G_1)) = p^2$ . Then from Theorem 9 and 10, we get  $c(G) = p^2 + p^3 < |G/Z(G)|^{1/2} c(Z(G)) = p(p + p^3) = p^2 + p^4$ .*

**Proof of Corollary 5.** Let  $G$  be a VZ-group of odd order. As  $G$  is a nilpotent group, it is isomorphic to the direct product of its Sylow subgroups. Since  $G$  has exactly two character degrees, all the Sylow subgroups of  $G$ , except one, are abelian. Let  $G \cong P_1 \times P_2 \times \cdots \times P_n$ , where  $P_i$  is a sylow  $p_i$ -subgroup

of  $G$ , for each  $i$ ,  $1 \leq i \leq n$ , where  $P_1$  is the only non-abelian VZ  $p_1$ -group and the rest are abelian Sylow subgroups. Then

$$\begin{aligned}\mu(G) &= \mu(P_1) + \mu(P_2) + \cdots + \mu(P_n) \quad (\text{by Theorem 9}) \\ &= c(P_1) + c(P_2) + \cdots + c(P_n) \quad (\text{by Theorem 10}) \\ &= c(G) \quad (\text{see [11]}).\end{aligned}$$

Thus, we finally get the values of  $\mu(G)$ ,  $q(G)$  and  $c(G)$  by using Theorem 2.

#### 4. Isoclinism and VZ $p$ -group of order $p^6$

In this section, we use the notation of the paper [15]. We will use not only the results of the previous sections but, more crucially, also use the classification of  $p$ -groups provided by James ([15]). The aim of this section is to calculate  $c(G)$  for all the VZ  $p$ -groups of order  $p^6$  ( $p$  an odd prime). We begin with the definition.

DEFINITION 4.0.1. *Two finite groups  $G$  and  $H$  are said to be isoclinic if there exist isomorphisms  $\theta : G/Z(G) \rightarrow H/Z(H)$  and  $\phi : G' \rightarrow H'$  such that the following diagram is commutative:*

$$\begin{array}{ccc} G/Z(G) \times G/Z(G) & \xrightarrow{a_G} & G' \\ \downarrow \theta \times \theta & & \downarrow \phi \\ H/Z(H) \times H/Z(H) & \xrightarrow{a_H} & H', \end{array}$$

where  $a_G(g_1Z(G), g_2Z(G)) = [g_1, g_2]$ , for  $g_1, g_2 \in G$ , and  $a_H(h_1Z(H), h_2Z(H)) = [h_1, h_2]$  for  $h_1, h_2 \in H$ .

The resulting pair  $(\theta, \phi)$  is called an *isoclinism* of  $G$  onto  $H$ . Notice that isoclinism is an equivalence relation on groups, weaker than isomorphism and it was first introduced by P. Hall [12] for a classification of  $p$ -groups. It is well-known that two isoclinic nilpotent groups have the same nilpotency class. Let the number of the irreducible characters of  $G$  of degree  $k$  be denoted by  $r_k(G)$ . Then we have the following theorem.

THEOREM 19. [6, Theorem 2.2] *Let  $G$  and  $H$  be isoclinic groups. Then*

- (1)  $|H|r_k(G) = |G|r_k(H)$ .
- (2) *The matrices of the irreducible complex representations of  $G$  and  $H$  only differ by scalar factors.*

COROLLARY 20. *Let  $G$  and  $H$  be two isoclinic VZ  $p$ -groups of same order such that  $d(Z(G)) = d(G')$ . Then  $c(G) = c(H)$  if and only if  $Z(G)$  is isomorphic to  $Z(H)$ .*

PROOF. Result follows from Corollary 4. □

**Note.** If we relax the condition  $d(Z(G)) = d(G')$  in Corollary 20, then Corollary 20 is no longer true. To produce counter examples, we have taken groups of order  $p^6$ , mentioned in the classification of  $p$ -groups, which is based on the concept of isoclinism (see [15]). Groups of order  $p^6$  (odd prime  $p$ ) have been divided into 43 isoclinic families denoted by  $\Phi_k$ ,  $1 \leq k \leq 43$  ([15, Section 4.6]). The groups are given by polycyclic presentation, in which all the relations of the form  $[x, y] = x^{-1}y^{-1}xy = 1$  between the generators have been omitted from the list. Since in the calculation of  $c(G)$ , we heavily use the presentations of the groups of order  $p^6$  ( $p$  odd) from the paper of James ([15]), the reader is advised to keep this paper handy. VZ  $p$ -groups of order  $p^6$  belong to isoclinic families  $\Phi_k$ , for  $k = 2, 5, 15$ .

EXAMPLE 21. **Isoclinic VZ  $p$ -groups  $G$  and  $H$  such that  $d(Z(G)) \neq d(G')$  and  $Z(G) \cong Z(H)$  but  $c(G) \neq c(H)$ .**

Consider two groups,  $G$  and  $H$ , of order  $p^6$  ( $p$  an odd prime) given by

$$G = \phi_2(3111)a = \phi_2(31) \times C_p \times C_p = \langle \alpha, \alpha_1, \alpha_2 \mid [\alpha_1, \alpha] = \alpha^{p^2} = \alpha_2, \alpha_1^p = \alpha_2^p = 1 \rangle \times C_p \times C_p$$

and

$$H = \phi_2(222)a = \phi_2(22) \times C_{p^2} = \langle \alpha, \alpha_1, \alpha_2 \mid [\alpha_1, \alpha] = \alpha^p = \alpha_2, \alpha_1^{p^2} = \alpha_2^p = 1 \rangle \times C_{p^2},$$



which belong to the isoclinic family  $\Phi_2$  ([15]). From Theorem 9 and 10,  $c(G) = c(\phi_2(31)) + c(C_p) + c(C_p)$ , and  $c(H) = c(\phi_2(22)) + c(C_{p^2})$ , and thus we need to calculate  $c(\phi_2(31))$  and  $c(\phi_2(22))$ . Let  $\phi_2(31) = G_1$ . Then  $Z(G_1) = \langle \alpha^p \rangle \cong C_{p^2}$  and thus from Theorem 11,  $c(G_1) = p^3$ , which finally gives  $c(G) = p^3 + 2p$ . Now, let  $\phi_2(22) = H_1$ . Then  $Z(H_1) = \langle \alpha^p, \alpha_1^p \rangle \cong C_p \times C_p$ ,  $H'_1 = \langle \alpha^p \rangle$  and  $Z(H_1)/H'_1 = \langle \alpha_1^p H'_1 \rangle$ . We have computed  $c(H_1)$  in subsection 4.1, and we get  $c(H_1) = 2p^2$ , and finally  $c(H) = 2p^2 + p^2 = 3p^2$ .

**EXAMPLE 22. Isoclinic VZ  $p$ -groups  $G$  and  $H$  such that  $d(Z(G)) \neq d(G')$  and  $Z(G) \not\cong Z(H)$  with  $c(G) \neq c(H)$ .**

Let us consider two groups  $G = \phi_2(3111)a = \phi_2(31) \times C_p \times C_p$ , and  $H = \phi_2(321)b = \phi_2(31) \times C_{p^2}$  of order  $p^6$  ( $p$  an odd prime) which belong to the isoclinic family  $\Phi_2$  ([15]). From Example 21, we get  $c(\phi_2(31)) = p^3$ . Here  $Z(G) = \langle \alpha^p \rangle \times C_p \times C_p \cong C_{p^2} \times C_p \times C_p$ ,  $Z(H) = \langle \alpha^p \rangle \times C_{p^2} \cong C_{p^2} \times C_{p^2}$  and  $d(Z(G)) \neq d(G')$ . Again by Theorem 9 and 10, we get  $c(G) = p^3 + 2p$  and  $c(H) = p^3 + p^2$ .

**EXAMPLE 23. Isoclinic VZ  $p$ -groups  $G$  and  $H$ , such that  $d(Z(G)) \neq d(G')$  and  $Z(G) \cong Z(H)$  with  $c(G) = c(H)$ .**

Consider two groups of order  $p^5$  ( $p$  an odd prime) given by

$$G = \phi_2(32)a_1 = \langle \alpha, \alpha_1, \alpha_2 \mid [\alpha_1, \alpha] = \alpha^{p^2} = \alpha_2, \alpha_1^{p^2} = \alpha_2^p = 1 \rangle$$

and

$$H = \phi_2(32)a_2 = \langle \alpha, \alpha_1, \alpha_2 \mid [\alpha_1, \alpha] = \alpha_1^p = \alpha_2, \alpha^{p^3} = \alpha_2^p = 1 \rangle,$$

which belong to the isoclinic family  $\Phi_2$  ([15]). Here  $Z(G) = Z(H) = \langle \alpha^p, \alpha_1^p \rangle \cong C_{p^2} \times C_p$ ,  $G' = \langle \alpha^{p^2} \rangle \cong C_p$ ,  $H' = \langle \alpha_1^p \rangle \cong C_p$ ,  $Z(G)/G' = \langle \alpha^p G', \alpha_1^p G' \rangle \cong C_p \times C_p$  and  $Z(H)/H' = \langle \alpha^p H' \rangle \cong C_{p^2}$ . Further,  $d(Z(G)) = d(Z(H)) = 2$  and  $d(G') = d(H') = 1$ . We first compute  $c(G)$  and then  $c(H)$ . In the view of the proof of Theorem 2, we consider a set  $X = \{\chi_1, \psi_2\}$ , where  $\chi_1 \in \text{nl}(G)$  and  $\psi_2 \in \text{lin}(G/G')$ . Here  $\chi_1 \downarrow_{Z(G)} = \chi_1(1)\lambda_1$ , where  $\lambda_1 \in \text{lin}(Z(G))$  is given by  $\lambda_1 = \lambda'_{\langle \alpha^p \rangle} \cdot 1_{\langle \alpha_1^p \rangle}$ , where  $\lambda'_{\langle \alpha^p \rangle}$  is a faithful linear character of  $\langle \alpha^p \rangle$ . On the other hand,  $\psi_2 = 1_{\langle \alpha G' \rangle} \cdot \psi_{\langle \alpha_1 G' \rangle}$ , where  $\psi_{\langle \alpha_1 G' \rangle}$  is a faithful linear character of  $\langle \alpha_1 G' \rangle$ , so that  $\psi_2 \downarrow_{Z(G)/G'} = 1_{\langle \alpha^p G' \rangle} \cdot \psi'_{\langle \alpha_1^p G' \rangle}$ , where  $\psi'_{\langle \alpha_1^p G' \rangle}$  is a faithful linear character of  $\langle \alpha_1^p G' \rangle$ . Then  $\ker \chi_1 = \langle \alpha_1^p \rangle$ , and  $\ker(\psi_2 \downarrow_{Z(G)/G'}) = \langle \alpha^p \rangle$ . Consider  $\xi = \sum_{i=1}^2 \theta_i$ , where  $\theta_1 = \sum_{\sigma \in \Gamma(\chi_1)} \chi_1^\sigma = p \left( \sum_{\sigma \in \Gamma(\lambda_1)} \lambda_1^\sigma \right)$ , and  $\theta_2 = \sum_{\sigma \in \Gamma(\psi_2)} \psi_2^\sigma$ . Then,  $m(\xi) = p^2 + p$  and  $\xi(1) = p^2(p-1) + p(p-1)$ . Therefore,  $c(G) = \xi(1) + m(\xi) = p^3 + p^2$ . In the similar way, we get  $c(H) = p^3 + p^2$ .

**EXAMPLE 24. Isoclinic VZ  $p$ -groups  $G$  and  $H$  such that  $d(Z(G)) \neq d(G')$  and  $Z(G) \cong Z(H)$  but  $c(G) = c(H)$ .**

Consider two groups,  $G$  and  $H$ , of order  $p^6$  ( $p$  an odd prime) given by

$$G = \phi_2(321)e = \phi_2(21) \times C_{p^3} = \langle \alpha, \alpha_1, \alpha_2 \mid [\alpha_1, \alpha] = \alpha_2, \alpha^p = \alpha_2, \alpha_1^p = \alpha_2^p = 1 \rangle \times C_{p^3},$$

and  $H = \phi_2(321)b = \phi_2(31) \times C_{p^2}$ , which belong to the isoclinic family  $\Phi_2$  ([15]). Here  $Z(G) = \langle \alpha^p \rangle \times C_{p^3} \cong C_p \times C_{p^3}$ ,  $Z(H) = \langle \alpha^p \rangle \times C_{p^2} \cong C_{p^2} \times C_{p^2}$  and  $d(Z(G)) \neq d(G')$ . From Example 22,  $c(H) = p^3 + p^2$ . On the other hand, from Remark 18, we get  $c(G) = p^3 + p^2$  and hence  $c(G) = c(H)$ .

In the next subsection, we give a complete list of  $c(G)$  for a VZ  $p$ -groups of order  $p^6$ .

#### 4.1. VZ $p$ -groups of order $p^6$ ( $p$ an odd prime).

**COROLLARY 25.** *Let  $G$  be a group of order  $p^6$  belonging to  $\Phi_{15}$  (odd prime  $p$ ). Then  $c(G) = 2p^3$ .*

**PROOF.** Consider the group

$$G = \phi_{15}(1^6) = \langle \alpha_1, \alpha_2, \alpha_3, \alpha_4, \beta_1, \beta_2 \mid [\alpha_1, \alpha_2] = \beta_1, [\alpha_1, \alpha_3] = \beta_2, \\ [\alpha_3, \alpha_4] = \beta_1, [\alpha_2, \alpha_4] = \beta_2^q, \alpha_1^p = \alpha_2^p = \alpha_3^p = \alpha_4^p = \beta_1^p = \beta_2^p = 1 \rangle,$$

where  $q$  denotes the smallest positive integer which is a primitive root (mod  $p$ ) ([15]). The group  $G$  is a VZ  $p$ -group. Here  $Z(G) = G' = \langle \beta_1, \beta_2 \rangle \cong C_p \times C_p$  and  $\text{cd}(G) = \{1, p^2\}$ . Now by Theorem 19, for all the groups  $H \in \Phi_{15}$ ,  $\text{cd}(H) = \{1, p^2\}$  and hence all the groups belonging to family  $\Phi_{15}$  are VZ groups.

Observe that, for each  $G \in \Phi_{15}$ ,  $Z(G) = G' \cong C_p \times C_p$ . Thus from Corollary 4 and Corollary 20, we have  $c(G) = 2p^3$  for every  $G \in \Phi_{15}$ .  $\square$

In Table 1 and 2, we provide a complete list of  $c(G)$ , where  $G$  is a  $p$ -group of order  $p^6$  lying in  $\Phi_2$  and  $\Phi_5$ . Calculation of  $c(G)$  is based upon the construction of a set  $X_G \subset \text{Irr}(G)$  (see in the proof of Theorem 2). Here we illustrate how to read Table 1 and 2.

Table 1 contains the groups belonging to family  $\Phi_2$  and Table 2 contains the groups belonging to family  $\Phi_5$ . For  $G \in \Phi_2$ ,  $|G/Z(G)| = p^2$  and thus  $\text{cd}(G) = \{1, p\}$ . For  $G \in \Phi_5$ ,  $|G/Z(G)| = p^4$  and hence  $\text{cd}(G) = \{1, p^2\}$ . There are two types of groups:

- (1) Type 1: groups which are not a direct product.
- (2) Type 2: groups which are expressed as a direct product of two or more groups.

The column with the heading ‘‘Group  $G$ ’’ contains the name of the group of order  $p^6$ . For groups of Type 2, the column with the heading ‘‘Group  $N$ ’’ contains the description of a non-abelian component  $N$  of  $G$ . For groups of Type 2, computation has been given for  $N$  and finally  $c(G)$  is computed from Theorems 9 and 10. In the column with the heading ‘‘Set  $X$ ’’, we have mentioned the members of the set  $X_G$  for the groups of Type 1, and the members of the set  $X_N$  for the groups of Type 2. Note that set  $X_G$  is in the form  $\{\chi_1, \dots, \chi_k, \psi_{k+1}, \dots, \psi_r\}$ , where  $\chi_i \in \text{nl}(G)$  (for  $1 \leq i \leq k$ ), and  $\psi_j \in \text{lin}(G/G')$  (for  $k+1 \leq j \leq r$ ) with  $d(Z(G)) = r$  and  $d(G') = k$ . Here  $\chi_i \downarrow_{Z(G)} = \chi_i(1)\lambda_i$ , where for each  $i$ ,  $\lambda_i \in \text{lin}(Z(G))$  such that  $G' \not\subseteq \ker(\lambda_i) = \ker(\chi_i)$ . But in Table 1 and 2, in place of  $\chi_i$  for  $(1 \leq i \leq k)$ , we have just mentioned  $\lambda_i$ . The notation  $\lambda_H$  stands for a faithful character of a subgroup  $H$ , and  $1_H$  stands for the trivial character of  $H$ . For better understanding on the working of table, we present an example. Let

$$G = \phi_2(222)a = \phi_2(22) \times C_{p^2} = \langle \alpha, \alpha_1, \alpha_2 \mid [\alpha_1, \alpha] = \alpha^p = \alpha_2, \alpha_1^{p^2} = \alpha_2^p = 1 \rangle \times C_{p^2}.$$

We take  $N = \phi_2(22)$ , which is a  $p$ -group of order  $p^4$  with nilpotency class 2. Then  $Z(N) = \langle \alpha^p, \alpha_1^p \rangle \cong C_p \times C_p$ ,  $N' = \langle \alpha^p \rangle$ ,  $Z(N)/N' = \langle \alpha_1^p N' \rangle$ ,  $N/N' = \langle \alpha N', \alpha_1 N' \rangle \cong C_p \times C_{p^2}$  and  $(\alpha N')^p = (\alpha_1 N')^{p^2} = N'$ . Here  $d(Z(N)) = 2$  and  $d(N') = 1$ , and so we have to use two irreducible characters of  $N$  to obtain a minimal faithful quasi-permutation representation (see Theorem 2). Consider a set  $X_N = \{\chi_1, \psi_2\} \subseteq \text{Irr}(N)$ , where  $\chi_1 \in \text{nl}(N)$  and  $\psi_2 \in \text{lin}(N/N')$ . Here  $\chi_1 \downarrow_{Z(N)} = \chi_1(1)\lambda_1$ , where  $\lambda_1 \in \text{lin}(Z(N))$  is given by  $\lambda_1 = \lambda'_{\langle \alpha^p \rangle} \cdot 1_{\langle \alpha_1^p \rangle}$ , where  $\lambda'_{\langle \alpha^p \rangle}$  is a faithful linear character of  $\langle \alpha^p \rangle$ . On the other hand,  $\psi_2 = 1_{\langle \alpha N' \rangle} \cdot \psi_{\langle \alpha_1 N' \rangle}$ , where  $\psi_{\langle \alpha_1 N' \rangle}$  is a faithful linear character of  $\langle \alpha_1 N' \rangle$ . Now,  $\psi_2 \downarrow_{Z(N)/N'} = \left( 1_{\langle \alpha N' \rangle} \cdot \psi_{\langle \alpha_1 N' \rangle} \right) \downarrow_{Z(N)/N'}$  is a faithful irreducible character of  $Z(N)/N' = \langle \alpha_1^p N' \rangle$ . Then  $\ker(\chi_1) = \langle \alpha_1^p \rangle$ ,  $\ker(\psi_2) = \langle \alpha N' \rangle$  and  $\ker(\psi_2 \downarrow_{Z(N)/N'}) = N' = \langle \alpha^p \rangle$ . Let  $\eta_{N'} : N \rightarrow N/N'$  be a natural group homomorphism. Then  $\widetilde{\psi}_2 := \psi_2 \circ \eta_{N'}$  is a linear character of  $N$  with  $\ker(\widetilde{\psi}_2) = \langle \alpha \rangle$ . So we can think  $\widetilde{\psi}_2$  as a faithful irreducible character of  $N/\ker(\widetilde{\psi}_2) = \langle \alpha_1 \rangle$ . Observe that

- (1)  $|\Gamma(\widetilde{\psi}_2)| = |\Gamma(\psi_2)| = \phi(p^2) = p(p-1)$ .
- (2)  $m\left(\sum_{\sigma \in \Gamma(\widetilde{\psi}_2)} \widetilde{\psi}_2^\sigma\right) = m\left(\sum_{\sigma \in \Gamma(\psi_2)} \psi_2^\sigma\right) = p$ .

Consider  $\xi = \sum_{i=1}^2 \theta_i$ , where  $\theta_1 = \sum_{\sigma \in \Gamma(\chi_1)} \chi_1^\sigma = p\left(\sum_{\sigma \in \Gamma(\lambda_1)} \lambda_1^\sigma\right)$ , and  $\theta_2 = \sum_{\sigma \in \Gamma(\psi_2)} \psi_2^\sigma$ . Then,  $m(\xi) = p + p = 2p$  and  $\xi(1) = p(p-1) + p(p-1) = 2p(p-1)$ . Therefore,  $c(N) = \xi(1) + m(\xi) = 2p^2$ . From Theorems 9 and 10,  $c(G) = c(\phi_2(22)) + c(C_{p^2})$ , and thus,  $c(G) = 2p^2 + p^2 = 3p^2$ . This group has been mentioned at Sl. no. 10 in Table 1.

Table 1:  $c(G)$  of VZ  $p$ -groups of order  $p^6$  belonging to  $\Phi_2$  family

Sl. no.	Group $G$	Group $N$	Set $X$	$c(G)$
1	$\phi_2(411)a$ $= \phi_2(41) \times C_p$	$\phi_2(41)$	$\lambda_1 = \lambda_{\langle \alpha^p \rangle}$	$c(G) = c(N) + c(C_p)$ $= p^4 + p$
2	$\phi_2(321)a_1$ $= \phi_2(32)a_1 \times C_p$	$\phi_2(32)a_1$	$\lambda_1 = \lambda_{\langle \alpha^p \rangle} \cdot 1_{\langle \alpha_1^p \rangle}$ , $\psi_2 = 1_{\langle \alpha N' \rangle} \cdot \psi_{\langle \alpha_1 N' \rangle}$	$c(G) = c(N) + c(C_p)$ $= p^3 + p^2 + p$
3	$\phi_2(321)a_2$ $= \phi_2(32)a_2 \times C_p$	$\phi_2(32)a_2$	$\lambda_1 = 1_{\langle \alpha^p \rangle} \cdot \lambda_{\langle \alpha_1^p \rangle}$ , $\psi_2 = \psi_{\langle \alpha N' \rangle} \cdot 1_{\langle \alpha_1 N' \rangle}$	$c(G) = c(N) + c(C_p)$ $= p^3 + p^2 + p$

Continuation of Table 1				
Sl. no.	Group $G$	Group $N$	Set $X$	$c(G)$
4	$\phi_2(321)b$ $= \phi_2(31) \times C_{p^2}$	$\phi_2(31)$	$\lambda_1 = \lambda_{\langle \alpha p \rangle}$	$c(G) = c(N) + c(C_{p^2})$ $= p^3 + p^2$
5	$\phi_2(321)e$ $= \phi_2(21) \times C_{p^3}$	$\phi_2(21)$	$\lambda_1 = \lambda_{\langle \alpha p \rangle}$	$c(G) = c(N) + c(C_{p^3})$ $= p^2 + p^3$
6	$\phi_2(3111)a$ $= \phi_2(31) \times C_p \times C_p$	$\phi_2(31)$	$\lambda_1 = \lambda_{\langle \alpha p \rangle}$	$c(G) = c(N) + 2c(C_p)$ $= p^3 + 2p$
7	$\phi_2(3111)b$ $= \phi_2(311)b \times C_p$	$\phi_2(311)b$	$\lambda_1 = \lambda_{\langle \gamma \rangle}$	$c(G) = c(N) + c(C_p)$ $= p^4 + p$
8	$\phi_2(3111)c$ $= \phi_2(311)c \times C_p$	$\phi_2(311)c$	$\lambda_1 = 1_{\langle \alpha p \rangle} \cdot \lambda_{\langle \alpha_2 \rangle}$ , $\psi_2 = \psi_{\langle \alpha N' \rangle} \cdot 1_{\langle \alpha_1 N' \rangle}$	$c(G) = c(N) + c(C_p)$ $= p^3 + p^2 + p$
9	$\phi_2(3111)d$ $= \phi_2(111) \times C_{p^3}$	$\phi_2(111)$	$\lambda_1 = \lambda_{\langle \alpha_2 \rangle}$	$c(G) = c(N) + c(C_{p^3})$ $= p^2 + p^3$
10	$\phi_2(222)a$ $= \phi_2(22) \times C_{p^2}$	$\phi_2(22)$	$\lambda_1 = \lambda_{\langle \alpha p \rangle} \cdot 1_{\langle \alpha_1 p \rangle}$ , $\psi_2 = 1_{\langle \alpha N' \rangle} \cdot \psi_{\langle \alpha_1 N' \rangle}$	$c(G) = c(N) + c(C_{p^2})$ $= 2p^2 + p^2 = 3p^2$
11	$\phi_2(2211)a$ $= \phi_2(22) \times C_p \times C_p$	$\phi_2(22)$	$\lambda_1 = \lambda_{\langle \alpha p \rangle} \cdot 1_{\langle \alpha_1 p \rangle}$ , $\psi_2 = 1_{\langle \alpha N' \rangle} \cdot \psi_{\langle \alpha_1 N' \rangle}$	$c(G) = c(N) + 2c(C_p)$ $= 2p^2 + 2p$
12	$\phi_2(2211)b$ $= \phi_2(21) \times C_{p^2} \times C_p$	$\phi_2(21)$	See Column 4 at Sl. no. 5 in this table	$c(G) = c(N) + c(C_{p^2}) + c(C_p)$ $= p^2 + p^2 + p = 2p^2 + p$
13	$\phi_2(2211)c$ $= \phi_2(221)c \times C_p$	$\phi_2(221)c$	$\lambda_1 = 1_{\langle \alpha p \rangle} \cdot \lambda_{\langle \gamma \rangle}$ , $\psi_2 = \psi_{\langle \alpha N' \rangle} \cdot 1_{\langle \alpha_1 N' \rangle} \cdot 1_{\langle \gamma N' \rangle}$	$c(G) = c(N) + c(C_p)$ $= p^3 + p^2 + p$
14	$\phi_2(2211)d$ $= \phi_2(221)d \times C_p$	$\phi_2(221)d$	$\lambda_1 = 1_{\langle \alpha p \rangle} \cdot 1_{\langle \alpha_1 p \rangle} \cdot \lambda_{\langle \alpha_2 \rangle}$ , $\psi_2 = \psi_{\langle \alpha N' \rangle} \cdot 1_{\langle \alpha_1 N' \rangle} \cdot 1_{\langle \alpha_2 N' \rangle}$ , $\psi_3 = 1_{\langle \alpha N' \rangle} \cdot \psi_{\langle \alpha_1 N' \rangle} \cdot 1_{\langle \alpha_2 N' \rangle}$	$c(G) = c(N) + c(C_p)$ $= 3p^2 + p$
15	$\phi_2(2211)e$ $= \phi_2(211)b \times C_{p^2}$	$\phi_2(211)b$	$\lambda_1 = \lambda_{\langle \gamma \rangle}$	$c(G) = c(N) + c(C_{p^2})$ $= p^3 + p^2$
16	$\phi_2(2211)f$ $= \phi_2(211)c \times C_{p^2}$	$\phi_2(211)c$	$\lambda_1 = 1_{\langle \alpha p \rangle} \cdot \lambda_{\langle \alpha_2 \rangle}$ , $\psi_2 = \psi_{\langle \alpha N' \rangle} \cdot 1_{\langle \alpha_1 N' \rangle} \cdot 1_{\langle \alpha_2 N' \rangle}$	$c(G) = c(N) + c(C_{p^2})$ $= 2p^2 + p^2 = 3p^2$
17	$\phi_2(21^4)a$ $= \phi_2(21) \times C_p \times C_p \times C_p$	$\phi_2(21)$	See Column 4 at Sl. no. 5 in this table	$c(G) = c(N) + 3c(C_p)$ $= p^2 + 3p$
18	$\phi_2(21^4)b$ $= \phi_2(21) \times C_{p^2} \times C_p$	$\phi_2(21)$	See Column 4 at Sl. no. 5 in this table	$c(G) = c(N) + c(C_{p^2}) + c(C_p)$ $= p^2 + p^2 + p = 2p^2 + p$
19	$\phi_2(21^4)c$ $= \phi_2(211)c \times C_p \times C_p$	$\phi_2(211)c$	See Column 4 at Sl. no. 16 in this table	$c(G) = c(N) + 2c(C_p)$ $= 2p^2 + 2p = 2p^2 + 2p$
20	$\phi_2(21^4)d$ $= \phi_2(111) \times C_{p^2} \times C_p$	$\phi_2(111)$	See Column 4 at Sl. no. 9 in this table	$c(G) = c(N) + c(C_{p^2}) + c(C_p)$ $= p^2 + p^2 + p = 2p^2 + p$
21	$\phi_2(1^6)$ $= \phi_2(111) \times C_p \times C_p \times C_p$	$\phi_2(111)$	See Column 4 at Sl. no. 9 in this table	$c(G) = c(N) + 3c(C_p)$ $= p^2 + 3p$
22	$\phi_2(51)$	-	$\lambda_1 = \lambda_{\langle \alpha p \rangle}$	$c(G) = p^5$
23	$\phi_2(42)a_1$	-	$\lambda_1 = \lambda_{\langle \alpha p \rangle} \cdot 1_{\langle \alpha_1 p \rangle}$ , $\psi_2 = 1_{\langle \alpha G' \rangle} \cdot \psi_{\langle \alpha_1 G' \rangle}$	$c(G) = p^4 + p^2$
24	$\phi_2(42)a_2$	-	$\lambda_1 = 1_{\langle \alpha p \rangle} \cdot \lambda_{\langle \alpha_1 p \rangle}$ , $\psi_2 = \psi_{\langle \alpha G' \rangle} \cdot 1_{\langle \alpha_1 G' \rangle}$	$c(G) = p^4 + p^2$
25	$\phi_2(411)b$	-	$\lambda_1 = \lambda_{\langle \gamma \rangle}$	$c(G) = p^5$
26	$\phi_2(411)c$	-	$\lambda_1 = 1_{\langle \alpha p \rangle} \cdot \lambda_{\langle \alpha_2 \rangle}$ , $\psi_2 = \psi_{\langle \alpha G' \rangle} \cdot 1_{\langle \alpha_1 G' \rangle}$	$c(G) = p^4 + p^2$
27	$\phi_2(33)$	-	$\lambda_1 = \lambda_{\langle \alpha p \rangle} \cdot 1_{\langle \alpha_1 p \rangle}$ , $\psi_2 = 1_{\langle \alpha G' \rangle} \cdot \psi_{\langle \alpha_1 G' \rangle}$	$c(G) = 2p^3$
28	$\phi_2(321)c$	-	$\lambda_1 = 1_{\langle \alpha p \rangle} \cdot \lambda_{\langle \gamma \rangle}$ , $\psi_2 = \psi_{\langle \alpha G' \rangle} \cdot 1_{\langle \alpha_1 G' \rangle} \cdot 1_{\langle \gamma G' \rangle}$	$c(G) = 2p^3$
29	$\phi_2(321)d$	-	$\lambda_1 = 1_{\langle \alpha p \rangle} \cdot \lambda_{\langle \gamma \rangle}$ , $\psi_2 = \psi_{\langle \alpha G' \rangle} \cdot 1_{\langle \alpha_1 G' \rangle} \cdot 1_{\langle \gamma G' \rangle}$	$c(G) = p^4 + p^2$
30	$\phi_2(321)f$	-	$\lambda_1 = 1_{\langle \alpha p \rangle} \cdot \lambda_{\langle \alpha_1 p \rangle} \cdot \lambda_{\langle \alpha_2 \rangle}$ , $\psi_2 = \psi_{\langle \alpha G' \rangle} \cdot 1_{\langle \alpha_1 G' \rangle} \cdot 1_{\langle \alpha_2 G' \rangle}$ , $\psi_3 = 1_{\langle \alpha G' \rangle} \cdot \psi_{\langle \alpha_1 G' \rangle} \cdot 1_{\langle \alpha_2 G' \rangle}$	$c(G) = p^3 + 2p^2$
31	$\phi_2(222)b$	-	$\lambda_1 = 1_{\langle \alpha p \rangle} \cdot 1_{\langle \alpha_1 p \rangle} \cdot \lambda_{\langle \gamma \rangle}$ , $\psi_2 = \psi_{\langle \alpha G' \rangle} \cdot 1_{\langle \alpha_1 G' \rangle} \cdot 1_{\langle \gamma G' \rangle}$ , $\psi_3 = 1_{\langle \alpha G' \rangle} \cdot \psi_{\langle \alpha_1 G' \rangle} \cdot 1_{\langle \gamma G' \rangle}$	$c(G) = p^3 + 2p^2$

 Table 2:  $c(G)$  of VZ  $p$ -groups of order  $p^6$  belonging to  $\Phi_5$  family

Sl. no.	Group $G$	Group $N$	Set $X$	$c(G)$
1	$\phi_5(21^4)a$ $= \phi_5(2111) \times C_p$	$\phi_5(2111)$	$\lambda_1 = \lambda_{\langle \alpha_1 p \rangle}$	$c(G) = c(N) + c(C_p)$ $= p^3 + p$
2	$\phi_5(1^6)$ $= \phi_5(1^5) \times C_p$	$\phi_5(1^5)$	$\lambda_1 = \lambda_{\langle \beta \rangle}$	$c(G) = c(N) + c(C_p)$ $= p^3 + p$
3	$\phi_5(3111)$	-	$\lambda_1 = \lambda_{\langle \alpha_1 p \rangle}$	$c(G) = p^4$
4	$\phi_5(2211)a$	-	$\lambda_1 = 1_{\langle \alpha_1 p \rangle} \cdot \lambda_{\langle \alpha_2 p \rangle}$ , $\psi_2 = \psi_{\langle \alpha_1 G' \rangle} \cdot 1_{\langle \alpha_2 G' \rangle} \cdot 1_{\langle \alpha_3 G' \rangle} \cdot 1_{\langle \alpha_4 G' \rangle}$	$c(G) = p^3 + p^2$
5	$\phi_5(2211)b$	-	$\lambda_1 = 1_{\langle \alpha_1 p \rangle} \cdot \lambda_{\langle \alpha_3 p \rangle}$ , $\psi_2 = \psi_{\langle \alpha_1 G' \rangle} \cdot 1_{\langle \alpha_2 G' \rangle} \cdot 1_{\langle \alpha_3 G' \rangle} \cdot 1_{\langle \alpha_4 G' \rangle}$	$c(G) = p^3 + p^2$
6	$\phi_5(21^4)b$	-	$\lambda_1 = \lambda_{\langle \gamma \rangle}$	$c(G) = p^4$

Continuation of Table 2				
Sl. no.	Group $G$	Group $N$	$X$	$c(G)$
7	$\phi_5(21^4)c$	-	$\lambda_1 = 1_{(\alpha_1^p)} \cdot \lambda_{(\beta)}$ , $\psi_2 = \psi_{(\alpha_1 G')} \cdot 1_{(\alpha_2 G')} \cdot 1_{(\alpha_3 G')} \cdot 1_{(\alpha_4 G')}$	$c(G) = p^3 + p^2$

THEOREM 26. *Suppose  $G$  is VZ  $p$ -group of order  $p^6$  (odd prime  $p$ ). Then  $c(G) = q(G) = p(G)$  and we have one of the following possibilities:*

- (1) *If  $|Z(G)| = p^4$ , then  $c(G) = p^4 + p, p^3 + p^2 + p, p^3 + p^2, p^3 + 2p, 3p^2, 2p^2 + 2p, 2p^2 + p, 3p^2 + p, p^2 + 3p, p^5, p^4 + p^2, 2p^3$  or  $p^3 + 2p^2$ .*
- (2) *If  $|Z(G)| = p^2$ , then  $c(G) = 2p^3, p^3 + p, p^4$  or  $p^3 + p^2$ .*

PROOF. The result follows from Corollary 25, Table 1 and 2.  $\square$

### 5. Camina $p$ -group of class 3

**Notation.** Throughout this section,  $G_2 = G'$  denotes the commutator subgroup of  $G$ . The nilpotency class of a nilpotent group  $G$  is the number  $n$  such that  $G_n \neq 1$  and  $G_{n+1} = 1$ , where  $G_2 = [G, G] = G'$  and  $G_{i+1} = [G_i, G]$ , for  $i > 2$ . Let  $N$  be a normal subgroup of  $G$ . Then we denote the set  $\text{Irr}(G) \setminus \text{Irr}(G/N)$  by  $\text{Irr}(G|N)$ .

We start by recalling some basic results that we will need later. The pair  $(G, N)$  is said to be a Camina pair if for every  $g \in G \setminus N$ ,  $gN \subseteq Cl_G(g)$ , the conjugacy class of  $g$ . When  $N = G'$ , the group  $G$  is called a Camina group. The following lemma gives us equivalent conditions for a pair  $(G, N)$  to be a Camina pair.

LEMMA 27 ([21], Lemma 3). *Let  $N$  be a normal subgroup of  $G$  and let  $g \in G \setminus N$ . Then the following statements are equivalent:*

- (i)  $\chi(g) = 0$  for all  $\chi \in \text{Irr}(G|N)$ .
- (ii)  $gN \subseteq Cl_G(g)$ , the conjugacy class of  $g$ .

It is easy to see that if  $(G, N)$  is a Camina pair, then  $Z(G) \leq N \leq G'$ . By Lemma 27, it is clear that if  $G$  is a Camina group, then for all  $g \in G \setminus G'$ ,  $\chi(g) = 0$  for all  $\chi \in \text{nl}(G)$ . Next theorem gives us bound on nilpotency class of Camina  $p$ -group.

THEOREM 28. [8] *If  $G$  is a finite Camina  $p$ -group, then the nilpotency class of  $G$  is at most 3, i.e.,  $G_4 = \{1\}$ .*

Now, we describe group theoretical property of Camina  $p$ -group and give a brief information about all non-linear irreducible characters of  $G$ .

LEMMA 29 ([20], Corollary 2.3). *Let  $G$  be a  $p$ -group of nilpotency class  $r$ . If  $(G, G_k)$  is a Camina pair, then  $G_i/G_{i+1}$  has exponent  $p$  for  $k - 1 \leq i \leq r$ .*

LEMMA 30 ([20], Theorem 5.2). *Let  $G$  be a Camina  $p$ -group of nilpotency class 3 and let  $|G/G_2| = p^m, |G_2/G_3| = p^n$ . Then*

- (i)  $(G, G_3)$  is a Camina pair,
- (ii)  $m = 2n$  and  $n$  is even.

LEMMA 31 ([20], Corollary 5.3). *If  $G$  is a Camina  $p$ -group of nilpotency class 3, then  $Z_2(G) = G_2$  and  $Z(G) = G_3$ , where  $Z_2(G)/Z(G) = Z(G/Z(G))$ .*

REMARK 32. *Suppose that nilpotency class of Camina  $p$ -group  $G$  is 2, i.e.,  $1 < G' \leq Z(G)$ . Since  $G$  is a Camina group, each nonlinear irreducible character of  $G$  vanishes outside  $G'$ . Therefore,  $G' = Z(G)$  and hence  $G$  becomes VZ  $p$ -group. So by Corollary 4, we get  $c(G)$ . Now Suppose  $G$  be a Camina  $p$ -group of nilpotency class 3. Then from Lemma 29 and Lemma 31,  $G/G', G'/Z(G)$  and  $Z(G)$  are elementary abelian  $p$ -groups and  $Z(G) \leq G'$ . Further, from Lemma 30,  $(G, Z(G))$  is a Camina pair,  $|G/G'| = p^{2n}, |G'/Z(G)| = p^n$  and  $|G/Z(G)| = p^{3n}$ , where  $n$  is even. Then, it is proved in ([22, Section 3, pp. 51-52]), that  $\text{nl}(G) = \text{Irr}(G|Z(G)) \sqcup \text{nl}(G/Z(G))$  as a disjoint union and  $\text{cd}(G) = \{1, p^n, p^{3n/2}\}$ .*

**5.1. Proof of the Theorem 6.** To prove Theorem 6, first we will prove the following lemma.

LEMMA 33. *Let  $G$  be a Camina  $p$ -group of nilpotency class 3. Let  $X$  be a subset of  $\text{Irr}(G)$  such that*

$$(9) \quad \bigcap_{\chi \in X} \ker(\chi) = 1$$

and equation (9) does not hold for any proper subset of  $X$ . Then  $X \subseteq \text{Irr}(G|Z(G))$ .

PROOF. Since  $(G, Z(G))$  is a Camina pair, from Lemma 27, for all  $\chi \in \text{Irr}(G|Z(G))$ ,  $\chi(g) = 0$ ,  $\forall g \in G \setminus Z(G)$ . Let  $\chi$  be an arbitrary irreducible character in  $\text{Irr}(G|Z(G))$ . Then  $\chi \downarrow_{Z(G)} = \chi(1)\lambda$ , where  $\lambda \in \text{lin}(Z(G))$ , and

$$(10) \quad \ker(\chi) = \ker(\lambda) < Z(G) < G', \quad \forall \chi \in \text{Irr}(G|Z(G)).$$

Secondly, if  $\psi \in \text{nl}(G/Z(G))$ , then  $Z(G) \leq \ker(\psi)$ . Suppose  $X$  is a collection of some irreducible characters of  $G$  satisfying equation (9) such that equation (9) does not hold for any proper subset of  $X$ . We know that all characters in  $X$  can not be linear. So, at least one irreducible character in  $X$ , say  $\chi$ , must be non-linear which either belongs to  $\text{Irr}(G|Z(G))$  or to  $\text{nl}(G/Z(G))$  (see Remark 32). If  $\chi \in \text{Irr}(G|Z(G))$ , then  $\ker(\chi) < Z(G) < G' \leq \ker(\eta)$ , where  $\eta \in \text{lin}(G)$ . Then  $\ker(\chi) \cap \ker(\eta) = \ker(\chi) \neq 1$ . Now, if  $\chi \in \text{nl}(G/Z(G))$ , then  $Z(G) \leq (\ker(\chi) \cap \ker(\eta))$ , where  $\eta \in \text{lin}(G)$ . Therefore, we can not put any linear characters in  $X$ . Hence  $X \subseteq \text{nl}(G) = \text{Irr}(G|Z(G)) \sqcup \text{nl}(G/Z(G))$  (disjoint union). Further, since  $X$  must satisfy equation (10),  $X$  can not be a subset of  $\text{nl}(G/Z(G))$ . Now, suppose  $X$  intersects with  $\text{Irr}(G|Z(G))$  and  $\text{nl}(G/Z(G))$ . Let  $\chi, \psi \in X$  such that  $\chi \in \text{Irr}(G|Z(G))$  and  $\psi \in \text{nl}(G/Z(G))$ . Then  $\ker \chi < Z(G) \leq \ker \psi \Rightarrow \ker \chi \cap \ker \psi = \ker \chi$  and this contradicts equation (10). Therefore,  $X$  must be a subset of  $\text{Irr}(G|Z(G))$ .  $\square$

*Proof of Theorem 6.* Suppose  $d(Z(G)) = r$ . Let  $H_i$ ,  $1 \leq i \leq r$ , denote the  $r$  cyclic components of  $Z(G)$ , i.e.  $Z(G) \cong \prod_{i=1}^r H_i$ , and let  $\widehat{H}_i = H_1 \times \cdots \times H_{i-1} \times H_{i+1} \times \cdots \times H_r$  for each  $i$  ( $1 \leq i \leq r$ ). By Remark 32,  $Z(G)$  is elementary abelian  $p$ -group, i.e.  $|H_i| = p$  for each  $i$ . From Lemma 33, it is clear that, to obtain  $c(G)$ , we have to consider irreducible characters from the set  $\text{Irr}(G|Z(G))$ . Now consider the set  $X = \{\chi_1, \dots, \chi_r\} \subseteq \text{Irr}(G|Z(G))$ , where  $\chi_i \downarrow_{Z(G)} = \chi_i(1)\lambda_i$ . Here  $\lambda_i \in \text{lin}(Z(G))$  is a linear character of  $Z(G)$  such that  $\ker(\lambda_i) = \widehat{H}_i$ . Then  $\bigcap_{i=1}^r \ker(\lambda_i) = \bigcap_{i=1}^r \ker(\chi_i) = 1$ . Moreover, no proper subset of  $X$  satisfies equation (9). It is easy to observe the following points.

- (1) For each  $i$ ,  $\Gamma(\chi_i) = \Gamma(\lambda_i)$  and  $|\Gamma(\chi_i)| = |\Gamma(\lambda_i)| = p - 1$ .
- (2) For each  $i$ ,  $m(\chi_i) = \chi_i(1)m(\lambda_i)$  and  $\chi_i(1) = p^{3n/2}$ .
- (3) For each  $i$ ,  $m(\lambda_i) = |\min \{ \sum_{\sigma \in \Gamma(\lambda_i)} \lambda_i^\sigma(g) \mid g \in G \}| = |-1| = 1$ .

In fact, for any  $\chi \in \text{Irr}(G|Z(G))$ ,  $m(\chi) = \chi(1)$  and all the above three observations holds. Now consider the character

$$\xi = \sum_{i=1}^r \theta_i,$$

where  $\theta_i = \sum_{\sigma \in \Gamma(\chi_i)} \chi_i^\sigma = \chi_i(1) \left( \sum_{\sigma \in \Gamma(\chi_i)} \lambda_i^\sigma \right)$ . Then by the above observations, we have  $m(\xi) = r\chi_i(1) = rp^{3n/2}$  and  $\xi(1) = \sum_{i=1}^r |\Gamma(\chi_i)|\chi_i(1) = r(p-1)p^{3n/2}$ . Thus,  $\xi(1) + m(\xi) = r(p-1)p^{3n/2} + rp^{3n/2} = rp^{\frac{3n}{2}+1}$ . Note that the set  $X = \{\chi_i\}_{i=1}^r$  gives a set  $Y = \{\lambda_i\}_{i=1}^r$ , which is a minimal faithful quasi-permutation representation of  $Z(G)$ . Therefore,

$$c(G) = \xi(1) + m(\xi) = rp^{\frac{3n}{2}+1}.$$

Moreover, we know that  $c(Z(G)) = rp$ . Hence we have

$$c(G) = p^{\frac{3n}{2}} c(Z(G)) = |G/Z(G)|^{1/2} c(Z(G)).$$

Further, from Theorem 10, for Camina  $p$ -group (odd prime  $p$ ) we get

$$c(G) = \mu(G) = q(G) = |G/Z(G)|^{1/2} c(Z(G)).$$

This proves the result.

COROLLARY 34. *Let  $G$  and  $H$  be two Camina  $p$ -groups of same order, and of same nilpotency class. Then  $c(G) = c(H)$  if and only if  $Z(G)$  is isomorphic to  $Z(H)$ .*

## 6. Acknowledgements

Ayush Udeep acknowledges University Grants Commission, Government of India. The corresponding author acknowledges SERB, Government of India for financial support through grant (MTR/2019/000118).

## References

- [1] M. H. Abbaspour, H. Behraves, A note on quasi-permutation representations of finite groups, *Int. J. Contemp. Math. Sci.*, **4**(27) (2009), 1315-1320.
- [2] H. Behraves and M. Delfani, On faithful quasi-permutation representations of groups of order  $p^5$ , *J. Algebra Appl.*, **17**(7) (2018), 1850127 (12 pages).
- [3] H. Behraves and G. Ghaffarzadeh, Minimal degree of faithful quasi-permutation representations of  $p$ -groups, *Algebra Colloq.*, **18**(Special Issue no. 1) (2011), 843-846.
- [4] H. Behraves, The minimal degree of a faithful quasi-permutation representation of an abelian group, *Glasgow Math. J.*, **39**(1) (1997), 51-57.
- [5] H. Behraves, Quasi-permutation Representations of  $p$ -groups of Class 2, *J. London Math. Soc. (2)*, **55**(2) (1997), 251-260.
- [6] J. C. Bioch, On  $n$ -isoclinic Groups. *Indag. Math. Proc.*, **79**(5) (1976), 400-407.
- [7] J. M. Burns, B. Goldsmith, B. Hartley and R. Sandling, On Quasi-permutation Representations of Finite Groups, *Glasgow Math. J.*, **36**(3) (1994), 301-308.
- [8] R. Dark and C. M. Scoppola, On Camina group of prime power order, *J. Algebra*, **181**(3) (1996), 787-802.
- [9] G. A. Fernández-Alcober and A. Moretó, Groups with two extreme character degrees and their normal subgroups, *Trans. Amer. Math. Soc.* **353**(6) (2001), 2171-2192.
- [10] G. Ghaffarzadeh and M. H. Abbaspour, Minimal degrees of faithful quasi-permutation representations for direct products of  $p$ -groups, *Proc. Indian Acad. Sci. Math. Sci.*, **122**(3) (2012), 329-334.
- [11] M. Ghaffarzadeh, On minimal degrees of faithful quasi-permutation representations of nilpotent groups, *Journal of New Researches in Mathematics*, **3**(12) (2018), 87-98.
- [12] P. Hall, The classification of prime-power groups, *J. Reine Angew. Math.*, **182** (1940), 130-141.
- [13] I. M. Isaacs, M. L. Lewis, Camina  $p$ -groups that are generalized Frobenius complements, *Arch. Math. (Basel)*, **104**(5) (2015), 401-405.
- [14] I. M. Isaacs, *Character theory of finite groups*, Academic, New York, 1976.
- [15] R. James, The groups of order  $p^6$  ( $p$  an odd prime), *Math. Comp.*, **34**(150) (1980), 613-637.
- [16] M. L. Lewis, Classifying Camina groups: a theorem of Dark and Scoppola. *Rocky Mountain J. Math.*, **44**(2) (2014), 591-597.
- [17] M. L. Lewis, The vanishing-off subgroup, *J. Algebra*, **321**(4) (2009), 1313-1325.
- [18] M.L. Lewis, Character tables of groups where all nonlinear irreducible characters vanish off the center, *Ischia group theory 2008*, 174-182, World Sci. Publ., Hackensack, NJ, 2009.
- [19] M. L. Lewis, Generalizing Camina groups and their character tables, *J. Group Theory*, **12**(2) (2009), 209-218.
- [20] I. D. Macdonald, Some  $p$ -groups of Frobenius and extra-special type, *Isr. J. Math.*, **40** (1981), 350-364.
- [21] S. Mattarei, Character Tables and Metabelian groups, *J. London Math. Soc.*, **46**(2) (1992), 92-100.
- [22] S. K. Prajapati and B. Sury, On the Total Character of Finite Groups, *Int. J. Group Theory*, **3**(3) (2014), 47-67.
- [23] S. K. Prajapati and R. Sarma, Total Character of a Group  $G$  with  $(G, Z(G))$  as a Generalized Camina Pair, *Canad. Math. Bull.*, **59**(2) (2016), 392-402.
- [24] W. J. Wong, Linear Groups Analogous to Permutation Groups, *J. Austral. Math. Soc., (Sec A)*, **3** (1963), 180-184.
- [25] D. Wright, Degrees of Minimal Embeddings for Some Direct Products, *Amer. J. Math.*, **97**(4) (1975), 897-903.

INDIAN INSTITUTE OF TECHNOLOGY, BHUBANESWAR, ARUGUL CAMPUS, JATNI, KHURDA-752050, INDIA.

*Email address:* [skprajapati@iitbbs.ac.in](mailto:skprajapati@iitbbs.ac.in)

INDIAN INSTITUTE OF TECHNOLOGY, BHUBANESWAR, ARUGUL CAMPUS, JATNI, KHURDA-752050, INDIA.

*Email address:* [udeepayush@gmail.com](mailto:udeepayush@gmail.com)