

AN EFFICIENT ALGORITHM FOR DECIDING VANISHING OF SCHUBERT POLYNOMIAL COEFFICIENTS

ANSHUL ADVE, COLLEEN ROBICHAUX, AND ALEXANDER YONG

ABSTRACT. *Schubert polynomials* form a basis of all polynomials and appear in the study of cohomology rings of flag manifolds. The vanishing problem for Schubert polynomials asks if a coefficient of a Schubert polynomial is zero. We give a tableau criterion to solve this problem, from which we deduce the first polynomial time algorithm. These results are obtained from new characterizations of the *Schubertope*, a generalization of the permutahedron defined for any subset of the $n \times n$ grid. In contrast, we show that computing these coefficients explicitly is #P-complete.

1. INTRODUCTION

Schubert polynomials form a linear basis of all polynomials $\mathbb{Z}[x_1, x_2, x_3, \dots]$. They were introduced by A. Lascoux–M.-P. Schützenberger [9] to study the cohomology ring of the flag manifold. These polynomials represent the Schubert classes under the Borel isomorphism. A reference is the textbook [6].

If $w_0 = n \ n - 1 \ \dots \ 2 \ 1$ is the longest length permutation in S_n , then

$$\mathfrak{S}_{w_0}(x_1, \dots, x_n) := x_1^{n-1} x_2^{n-2} \dots x_{n-1}.$$

Otherwise, $w \neq w_0$ and there exists i such that $w(i) < w(i+1)$. Then one sets

$$\mathfrak{S}_w(x_1, \dots, x_n) = \partial_i \mathfrak{S}_{ws_i}(x_1, \dots, x_n),$$

where s_i is the transposition swapping i and $i+1$ and

$$\partial_i f := \frac{f(\dots, x_i, x_{i+1}, \dots) - f(\dots, x_{i+1}, x_i, \dots)}{x_i - x_{i+1}}.$$

Since ∂_i satisfies

$$\partial_i \partial_j = \partial_j \partial_i \text{ for } |i - j| > 1, \text{ and } \partial_i \partial_{i+1} \partial_i = \partial_{i+1} \partial_i \partial_{i+1},$$

the above description of \mathfrak{S}_w is well-defined. In addition, under the inclusion $\iota : S_n \hookrightarrow S_{n+1}$ defined by $w(1) \dots w(n) \mapsto w(1) \dots w(n) \ n+1$, $\mathfrak{S}_w = \mathfrak{S}_{\iota(w)}$. Thus one unambiguously refers to \mathfrak{S}_w for each $w \in S_\infty = \bigcup_{n \geq 1} S_n$.

The *graph* $G(w)$ of a permutation $w \in S_n$ is the $n \times n$ grid, with a \bullet placed in position $(i, w(i))$ (in matrix coordinates). The *Rothe diagram* of w is given by


$$D(w) = \{(i, j) : 1 \leq i, j \leq n, j < w(i), i < w^{-1}(j)\}.$$

This is pictorially described with rays that strike out boxes south and east of each \bullet in $G(w)$. $D(w)$ are the remaining boxes.

The *code* of w , denoted $\text{code}(w)$ is the vector (c_1, c_2, \dots, c_L) where c_i is the number of boxes in the i -th row of $D(w)$ and L indexes the southmost row with a positive number of boxes. To each $w \in S_\infty$ there is a unique associated code; see [10, Proposition 2.1.2].

Date: March 3, 2021.

Example 1.1. If $w = 53841267 \in S_8$ (in one line notation) then $D(w)$ is depicted by:



Here, $\text{code}(w) = (4, 2, 5, 2)$.

Consider the monomial expansion

$$\mathfrak{S}_w = \sum_{\alpha \in \mathbb{Z}_{\geq 0}^n} c_{\alpha, w} x^\alpha.$$

Now, $c_{\alpha, w} = 0$ unless $\alpha_i = 0$ for $i > L$, and moreover, $c_{\alpha, w} \in \mathbb{Z}_{\geq 0}$. Let Schubert be the problem of deciding $c_{\alpha, w} \neq 0$, as measured in the input size of α and w (under the assumption that arithmetic operations take constant time). The INPUT is $\text{code} = (c_1, \dots, c_L) \in \mathbb{Z}_{\geq 0}^L$ with $c_L > 0$ and $\alpha \in \mathbb{Z}_{\geq 0}^L$. Schubert returns YES if $c_{\alpha, w} > 0$ and NO otherwise.


Theorem 1.2. Schubert \in P.

We prove Theorem 1.2 using another result. Fix $n \in \mathbb{Z}_{>0}$ and let $D \subseteq [n]^2$. We call D a *diagram* and visualize D as a subset of an $n \times n$ grid of boxes, oriented so that $(r, c) \in [n]^2$ represents the box in the r th row from the top and the c th column from the left. Let $\text{PerfectTab}(D, \alpha)$ be the fillings of D with α_k many k 's, where entries in each column are distinct, any entry in row i is $\leq i$, and each box contains exactly one entry. Let $\text{PerfectTab}_{\downarrow}(D, \alpha) \subseteq \text{PerfectTab}(D, \alpha)$ be fillings where entries in each column increase from top to bottom.

Theorem 1.3. $c_{\alpha, w} > 0 \iff \text{PerfectTab}(D(w), \alpha) \neq \emptyset \iff \text{PerfectTab}_{\downarrow}(D(w), \alpha) \neq \emptyset$

In general $\#\text{PerfectTab}(D(w), \alpha) \neq c_{\alpha, w}$ but rather $\#\text{PerfectTab}(D(w), \alpha) \geq c_{\alpha, w}$ (cf. [5]).

Example 1.4. Here are the tableaux in $\bigcup_{\alpha} \text{PerfectTab}_{\downarrow}(D(31524), \alpha)$:



Hence, for instance, $c_{(2,1,1), 31524} > 0$ but $c_{(4), 31524} = 0$.

To prove Theorems 1.2 and 1.3 we establish results about the *Schubitope* [11]. This polytope \mathcal{S}_D is defined with a halfspace description for any $D \subseteq [n]^2$. We prove (Theorem 2.13) that a lattice point α is in \mathcal{S}_D if and only if $\text{PerfectTab}(D, \alpha) \neq \emptyset$ where D is any diagram.

We then introduce the *indicator polytope* $\mathcal{P}(D, \alpha)$ whose lattice points $\mathcal{P}(D, \alpha)_{\mathbb{Z}}$ are in bijection with $\text{PerfectTab}(D, \alpha)$. We prove that $\mathcal{P}(D, \alpha) \neq \emptyset \iff \mathcal{P}(D, \alpha)_{\mathbb{Z}} \neq \emptyset$ (Theorem 2.27). Thus determining $\mathcal{P}(D, \alpha)_{\mathbb{Z}} \neq \emptyset$ (and equivalently $\alpha \in \mathcal{S}_D$) is in P using L. Khachiyan’s ellipsoid method for linear programming, see [14]. We give two proofs of Theorem 2.27. The first shows $\mathcal{P}(D, \alpha)$ is totally unimodular. Hence $\mathcal{P}(D, \alpha) \neq \emptyset$ implies $\mathcal{P}(D, \alpha)$ has integral vertices. Our second proof obviates total unimodularity and is potentially adaptable to problems lacking that property. However, only the high-level structure of the second proof is easily generalizable — the rest is necessarily *ad hoc*.

For the case of Rothe diagrams $D = D(w)$, using A. Fink-K. Mészáros-A. St. Dizier [4, Corollary 12 and Theorem 14] (conjectured in [11, Conjectures 5.1 and 5.13]),

$$(1) \quad \alpha \in \mathcal{S}_{D(w)} \iff c_{\alpha, w} > 0.$$

This, combined with our results on the Schubitope, proves Theorems 1.2 and 1.3.

The class #P in L. Valiant’s complexity theory of counting problems are those that count the number of accepting paths of a nondeterministic Turing machine running in polynomial time. A problem $\mathcal{P} \in \#P$ is *complete* if for any problem $\mathcal{Q} \in \#P$ there exists a polynomial-time counting reduction from \mathcal{Q} to \mathcal{P} . These are the hardest of the problems in #P. There does not exist a polynomial time algorithm for such problems unless $P = NP$.

In contrast with Theorem 1.2, we prove:

Theorem 1.5. *Counting $c_{\alpha, w}$ is #P-complete.*

Given $\{c_{\alpha, w} \in \mathbb{Z}_{\geq 0}\}$ it is standard to ask for a counting rule for $c_{\alpha, w}$. A complexity motivation is an *appropriate* rule that establishes a counting problem is in #P with respect to given input (length). The rule of [3] establishes that counting $c_{\alpha, w}$ is in #P if the input is (w, α) but not if the input is $(\text{code}(w), \alpha)$. For the latter input assumption, we use the transition algorithm of [8] and its *graphical* reformulation from [7]. This allows us to give a polynomial time counting reduction to the #P-complete problem of counting Kostka coefficients [12], (see Section 5).

Related discussion of complexity and the “nonvanishing problem” in algebraic combinatorics appears in the conference proceedings [1] and preprint [2] versions of this paper.

2. THE SCHUBITOPE


Consider a diagram $D \subseteq [n]^2$. Given $S \subseteq [n]$ and a column $c \in [n]$, construct a string denoted $\text{word}_{c, S}(D)$ by reading column c from top to bottom and recording

- (if $(r, c) \notin D$ and $r \in S$,
-) if $(r, c) \in D$ and $r \notin S$, and
- \star if $(r, c) \in D$ and $r \in S$.

Let $\theta_D^c(S) = \#\{\star\text{'s in } \text{word}_{c, S}(D)\} + \#\{\text{paired } ()\text{'s in } \text{word}_{c, S}(D)\}$ and

$$\theta_D(S) = \sum_{c=1}^n \theta_D^c(S).$$

Example 2.1. In the diagram D below, we labelled the corresponding strings for $\text{word}_{c,S}(D)$ for $S = \{1, 3\}$. For instance, we see $\text{word}_{5,\{1,3\}}(D) = (\star)$.



The *Schubitope* \mathcal{S}_D , as defined in [11], is the polytope

$$(2) \quad \left\{ (\alpha_1, \dots, \alpha_n) \in \mathbb{R}_{\geq 0}^n : \alpha_1 + \dots + \alpha_n = \#D \text{ and } \sum_{i \in S} \alpha_i \leq \theta_D(S) \text{ for all } S \subseteq [n] \right\}.$$

2.1. Characterizations via tableaux. A *tableau* of shape D is a map

$$\tau : D \rightarrow [n] \cup \{\circ\},$$


where $\tau(r, c) = \circ$ indicates that the box (r, c) is unlabelled. Let $\text{Tab}(D)$ denote the set of such tableaux.

It will be useful to reformulate the original definition of $\theta_D(S)$ into the language of tableaux. Given $S \subseteq [n]$, define $\pi_{D,S} \in \text{Tab}(D)$ by

$$(3) \quad \pi_{D,S}(r, c) = \begin{cases} r & \text{if } (r, c) \text{ contributes a ``}\star\text{'' to } \text{word}_{c,S}(D), \\ s & \text{if } (r, c) \text{ contributes a ``)'' to } \text{word}_{c,S}(D) \text{ which is} \\ & \text{paired with an ``('' from } (s, c), \\ \circ & \text{otherwise.} \end{cases}$$

In (3) and throughout, we pair by the standard “inside-out” convention.

Example 2.2. Continuing Example 2.1, below is $\pi_{D,\{1,3\}}(D)$




Proposition 2.3. For all $D \subseteq [n]^2$ and $S \subseteq [n]$, we have $\theta_D(S) = \#\pi_{D,S}^{-1}(S)$.

Proof. $\pi_{D,S}(r, c) \in S$ if and only if (r, c) falls into one of the first two cases in (3). □

Say $\tau \in \text{Tab}(D)$ is *flagged* if $\tau(r, c) \leq r$ whenever $\tau(r, c) \neq \circ$. It is *column-injective* if $\tau(r, c) \neq \tau(r', c)$ whenever $r \neq r'$ and $\tau(r, c) \neq \circ$. Let $\text{FCITab}(D) \subseteq \text{Tab}(D)$ be the set of tableaux of shape D which are flagged and column-injective.

Example 2.4. Of the tableaux of shape D below, only the second and fourth are flagged, and only the third and fourth are column-injective.



Proposition 2.5. $\pi_{D,S} \in \text{FCITab}(D)$ for all $D \subseteq [n]^2$ and $S \subseteq [n]$.

Proof. This is immediate from (3). □

A simple consequence of being flagged and column-injective is the following.

Proposition 2.6. Let $\tau \in \text{FCITab}(D)$. Then for all $(r, c) \in [n]^2$ and $S \subseteq [n]$, we have

$$(4) \quad \#\{(i, c) \in \tau^{-1}(S) : i < r\} \leq \#\{i \in S : i \leq r\},$$

with strict inequality whenever $(r, c) \in \tau^{-1}(S)$.

Proof. The map $(i, c) \mapsto \tau(i, c)$ from $\{(i, c) \in \tau^{-1}(S) : i \leq r\}$ to $\{i \in S : i \leq r\}$ is well-defined since τ is flagged. It is injective since τ is column-injective. Thus (4) holds, and

$$\#\{(i, c) \in \tau^{-1}(S) : i < r\} < \#\{(i, c) \in \tau^{-1}(S) : i \leq r\} \leq \#\{i \in S : i \leq r\}$$

whenever $(r, c) \in \tau^{-1}(S)$, establishing the strict inequality assertion. □

In fact, a stronger assertion holds when $\tau = \pi_{D,S}$.

Proposition 2.7. If $(r, c) \in D \subseteq [n]^2$ and $S \subseteq [n]$, then

$$(r, c) \in \pi_{D,S}^{-1}(S) \iff \#\{(i, c) \in \pi_{D,S}^{-1}(S) : i < r\} < \#\{i \in S : i \leq r\}.$$

Proof. (\Rightarrow) This direction follows from Propositions 2.5 and 2.6.

(\Leftarrow) If $r \in S$, then (r, c) contributes a “ \star ” to $\text{word}_{c,S}(D)$, so $\pi_{D,S}(r, c) = r \in S$, as desired. Thus we assume $r \notin S$. The hypothesis combined with this assumption says

$$\#\{(i, c) \in \pi_{D,S}^{-1}(S) : i < r\} < \#\{i \in S : i \leq r\} = \#\{i \in S : i < r\}.$$

Thus, there is a maximal $s \in S$ with $s < r$ such that $\pi_{D,S}(r', c) \neq s$ whenever $r' < r$. If $(s, c) \in D$, then (s, c) contributes a “ \star ” to $\text{word}_{c,S}(D)$, so $\pi_{D,S}(s, c) = s$, contradicting our choice of s . Therefore, (s, c) contributes an “(” to $\text{word}_{c,S}(D)$. If this “(” is paired by a “)” contributed by $(r', c) \in D$ with $r' < r$, then $\pi_{D,S}(r', c) = s$, again a contradiction. Thus, this “(” pairs the “)” from (r, c) , so $\pi_{D,S}(r, c) = s \in S$. Hence, $(r, c) \in \pi_{D,S}^{-1}(S)$ as desired. □

The previous two propositions combined assert that $\{(r, c) \in \pi_{D,S}^{-1}(S)\}$ is characterized by greedy selection as one moves down each column c . The next proposition shows that this greedy algorithm maximizes $\#\tau^{-1}(S)$ among all $\tau \in \text{FCITab}(D)$.

Proposition 2.8. Let $D \subseteq [n]^2$ and $S \subseteq [n]$. Then $\#\pi_{D,S}^{-1}(S) \geq \#\tau^{-1}(S)$ for all $\tau \in \text{FCITab}(D)$.

Proof. If not, then there exist $\tau \in \text{FCITab}(D)$ and $(r, c) \in [n]^2$ satisfying

$$\#\{(i, c) \in \pi_{D,S}^{-1}(S) : i \leq r\} < \#\{(i, c) \in \tau^{-1}(S) : i \leq r\}$$

and we can choose these such that r is minimized. Then because r is minimal,

$$\#\{(i, c) \in \pi_{D,S}^{-1}(S) : i < r\} = \#\{(i, c) \in \tau^{-1}(S) : i < r\}$$

and $(r, c) \in \tau^{-1}(S) \setminus \pi_{D,S}^{-1}(S)$, so in particular $(r, c) \in D$. Thus Proposition 2.6 implies

$$\#\{(i, c) \in \pi_{D,S}^{-1}(S) : i < r\} = \#\{(i, c) \in \tau^{-1}(S) : i < r\} < \#\{i \in S : i \leq r\}.$$

But then we must have $(r, c) \in \pi_{D,S}^{-1}(S)$ by Proposition 2.7, a contradiction. □

If τ has shape a subset of $[n]^2$ and $\alpha = (\alpha_1, \dots, \alpha_n) \in \mathbb{R}_{\geq 0}^n$, say τ exhausts α over S if

$$\sum_{i \in S} \alpha_i \leq \#\tau^{-1}(S).$$

Example 2.9. Only the left tableau below exhausts $\alpha = (3, 2, 2, 4)$ over $S = \{1, 3\}$.

1				
	1	1		
		3		
		4	4	
4	o	4	3	

1				
	1	2		
		3		
		4	4	
4	o	4	2	

Theorem 2.10. Let $D \subseteq [n]^2$ and $\alpha = (\alpha_1, \dots, \alpha_n) \in \mathbb{Z}_{\geq 0}^n$ with $\alpha_1 + \dots + \alpha_n = \#D$. Then $\alpha \in \mathcal{S}_D$ if and only if for each $S \subseteq [n]$, there exists $\tau_{D,S} \in \text{FCITab}(D)$ which exhausts α over S .

Proof of Theorem 2.10. (\Rightarrow) The inequalities in (2) combined with Proposition 2.3 imply

$$\sum_{i \in S} \alpha_i \leq \theta_D(S) = \#\pi_{D,S}^{-1}(S).$$

Thus, $\tau_{D,S} := \pi_{D,S}$ exhausts α over S .

(\Leftarrow) By Propositions 2.8 and 2.3,

$$\sum_{i \in S} \alpha_i \leq \#\tau_{D,S}^{-1}(S) \leq \#\pi_{D,S}^{-1}(S) = \theta_D(S),$$

so the inequalities in (2) hold. □

Remark 2.11. The proof of (\Rightarrow) shows that we can take $\tau_{D,S} = \pi_{D,S}$ in Theorem 2.10.

It would be nice if $\tau_{D,S}$ did not depend on S , i.e., if some τ_D exhausted α over all $S \subseteq [n]$, so we could take $\tau_{D,S} = \tau_D$ in Theorem 2.10. Indeed, this is shown in Theorem 2.13.

Say $\tau \in \text{Tab}(D)$ has content α if $\#\tau^{-1}(\{i\}) = \alpha_i$ for each $i \in [n]$. Let $\text{Tab}(D, \alpha)$ and $\text{FCITab}(D, \alpha)$ be the subsets of $\text{Tab}(D)$ and $\text{FCITab}(D)$, respectively, of those tableaux which have content α . In addition, call a tableau $\tau \in \text{Tab}(D)$ perfect if $\tau \in \text{FCITab}(D)$, and if no boxes are left unlabelled, i.e., $\tau^{-1}(\{o\}) = \emptyset$. Thus, the set of perfect tableaux of content α is precisely $\text{PerfectTab}(D, \alpha) \subseteq \text{FCITab}(D, \alpha)$ introduced in Section 1.

Proposition 2.12. Let $D \subseteq [n]^2$ and $\alpha = (\alpha_1, \dots, \alpha_n) \in \mathbb{Z}_{\geq 0}^n$. Then $\text{PerfectTab}(D, \alpha) \neq \emptyset$ if and only if $\alpha_1 + \dots + \alpha_n = \#D$ and $\text{FCITab}(D, \alpha) \neq \emptyset$.

Proof. (\Rightarrow) Let $\tau \in \text{PerfectTab}(D, \alpha)$. Then $\tau \in \text{FCITab}(D, \alpha)$, and since τ has content α and satisfies $\tau^{-1}(\{o\}) = \emptyset$,

$$\alpha_1 + \dots + \alpha_n = \#\tau^{-1}(\{1\}) + \dots + \#\tau^{-1}(\{n\}) = \#D.$$

(\Leftarrow) Let $\tau \in \text{FCITab}(D, \alpha)$. Then since τ has content α ,

$$\#\tau^{-1}(\{o\}) = \#D - \#\tau^{-1}(\{1\}) - \dots - \#\tau^{-1}(\{n\}) = \#D - \alpha_1 - \dots - \alpha_n = 0.$$

Thus, $\tau \in \text{PerfectTab}(D, \alpha)$. □

Theorem 2.13. Let $D \subseteq [n]^2$ and $\alpha = (\alpha_1, \dots, \alpha_n) \in \mathbb{Z}_{\geq 0}^n$. Then $\alpha \in \mathcal{S}_D$ if and only if $\text{PerfectTab}(D, \alpha) \neq \emptyset$.

The proof will require a lemma regarding tableaux of the form $\tau = \pi_{D,S}$.

Lemma 2.14. Let $D \subseteq [n]^2$, and $S, T \subseteq [n]$ be disjoint. Set

$$\tilde{D} = D \setminus \pi_{D,S}^{-1}(S) \text{ and } U = S \cup T.$$

Then

$$\pi_{D,U}^{-1}(U) = \pi_{D,S}^{-1}(S) \cup \pi_{\tilde{D},T}^{-1}(T).$$

Proof. Let $(r, c) \in D$, and assume by induction on r that

$$(5) \quad (i, c) \in \pi_{D,U}^{-1}(U) \iff (i, c) \in \pi_{D,S}^{-1}(S) \cup \pi_{\tilde{D},T}^{-1}(T)$$

whenever $i < r$. This clearly holds in the base case $r = 1$. By Proposition 2.7, $(r, c) \in \pi_{D,U}^{-1}(U)$ if and only if

$$(6) \quad \#\{(i, c) \in \pi_{D,U}^{-1}(U) : i < r\} < \#\{i \in U : i \leq r\}.$$

By (5) and the fact that

$$\pi_{D,S}^{-1}(S) \cap \tilde{D} = \emptyset = S \cap T,$$

(6) is equivalent to

$$\#\{(i, c) \in \pi_{D,S}^{-1}(S) : i < r\} + \#\{(i, c) \in \pi_{\tilde{D},T}^{-1}(T) : i < r\} < \#\{i \in S : i \leq r\} + \#\{i \in T : i \leq r\}.$$

By applying Proposition 2.6 twice, we see that this holds if and only if at least one of (i) and (ii) below hold.

- (i) $\#\{(i, c) \in \pi_{D,S}^{-1}(S) : i < r\} < \#\{i \in S : i \leq r\}$
- (ii) $\#\{(i, c) \in \pi_{\tilde{D},T}^{-1}(T) : i < r\} < \#\{i \in T : i \leq r\}$

By Proposition 2.7, (i) is equivalent to $(r, c) \in \pi_{D,S}^{-1}(S)$. If indeed $(r, c) \in \pi_{D,S}^{-1}(S)$ holds, then our induction step is complete. Otherwise, $(r, c) \notin \pi_{D,S}^{-1}(S)$, so by definition, $(r, c) \in \tilde{D}$. Thus, applying Proposition 2.7 to \tilde{D} , $T \subseteq [n]$ and $(r, c) \in \tilde{D}$, (ii) is equivalent to $(r, c) \in \pi_{\tilde{D},T}^{-1}(T)$. Hence, (5) holds for all $i \leq r$. \square

Corollary 2.15. Let $D \subseteq [n]^2$ and $S \subseteq U \subseteq [n]$. Then $\pi_{D,S}^{-1}(S) \subseteq \pi_{D,U}^{-1}(U)$.

Proof. Take $T = U \setminus S$ in Lemma 2.14. \square

Finally, we are ready to prove Theorem 2.13.

Proof of Theorem 2.13. (\Leftarrow) Let $\tau_D \in \text{PerfectTab}(D, \alpha)$. Then $\alpha_1 + \cdots + \alpha_n = \#D$ by Proposition 2.12. Also, for each $S \subseteq [n]$,

$$\sum_{i \in S} \alpha_i = \sum_{i \in S} \#\tau_D^{-1}(\{i\}) = \#\tau_D^{-1}(S),$$

so τ_D exhausts α over S . Thus, $\alpha \in \mathcal{S}_D$ by Theorem 2.10.

(\Rightarrow) We induct on the sum of the row indices of each box in D , i.e., $\sum_{(i,j) \in D} i$. The base case of an empty diagram is trivial, so we may assume $D \neq \emptyset$. Then since $\alpha \in \mathcal{S}_D$, (2) implies $\alpha_1 + \cdots + \alpha_n = \#D > 0$, so we can choose m maximal such that $\alpha_m > 0$.

Case 1: (D contains boxes below row m). Pick $(r, c) \in D$ below row m (so $r > m$).

Claim 2.16. There exists $r_1 < r$ such that $(r_1, c) \notin D$.

Proof of Claim 2.16. By Theorem 2.10, there exists $\tau_{D,[m]} \in \text{FCITab}(D)$ such that

$$(7) \quad \#\tau_{D,[m]}^{-1}([m]) \geq \alpha_1 + \cdots + \alpha_m = \alpha_1 + \cdots + \alpha_n = \#D.$$

Thus, $\tau_{D,[m]}(D) \subseteq [m]$. Consequently, by column-injectivity of $\tau_{D,[m]}$, there can be at most m boxes in each column of D . Since $(r, c) \in D$ with $r > m$, there are more than m boxes in column c if $(r_1, c) \in D$ for all $r_1 < r$. Hence there must be some $r_1 < r$ for which $(r_1, c) \notin D$, as asserted. \square

By Claim 2.16, we can choose $r_1 < r$ maximal such that $(r_1, c) \notin D$. Let

$$\tilde{D} = (D \setminus \{(r, c)\}) \cup \{(r_1, c)\}.$$

Claim 2.17. $\alpha \in \mathcal{S}_{\tilde{D}}$.

Proof of Claim 2.17. Since $\alpha \in \mathcal{S}_D$, $(r, c) \in D$, and $(r_1, c) \notin D$, we have

$$\alpha_1 + \cdots + \alpha_n = \#D = \#\tilde{D}.$$

Let $S \subseteq [n]$ and $T = S \cap [m]$. Then define $\tau_{\tilde{D},S} \in \text{Tab}(\tilde{D})$ by

$$\tau_{\tilde{D},S}(i, j) = \begin{cases} \pi_{D,T}(r, c) & \text{if } (i, j) = (r_1, c), \\ \pi_{D,T}(i, j) & \text{otherwise.} \end{cases}$$

If $\pi_{D,T}(r, c) = \circ$, then certainly $\tau_{\tilde{D},S} \in \text{FCITab}(\tilde{D})$. Otherwise, let $s = \pi_{D,T}(r, c)$. Since $(r, c) \in D$ but $r \notin T$, (r, c) contributes a “)” to $\text{word}_{c,S}(D)$. Thus, by (3), (s, c) contributes an “(”, so in particular $(s, c) \notin D$. From our choice of r_1 , we must therefore have $s \leq r_1$, so $\tau_{\tilde{D},S}$ is flagged. Hence, $\tau_{\tilde{D},S} \in \text{FCITab}(\tilde{D})$.

By construction,

$$\#\tau_{\tilde{D},S}^{-1}(\{i\}) = \#\pi_{D,T}^{-1}(\{i\})$$

for each $i \in [n]$, so $\tau_{\tilde{D},S}$ exhausts α over T by Theorem 2.10 and in particular Remark 2.11. Since $\alpha_i = 0$ for all $i > m$, we can write

$$\sum_{i \in S} \alpha_i = \sum_{i \in T} \alpha_i \leq \#\tau_{\tilde{D},S}^{-1}(T) \leq \#\tau_{\tilde{D},S}^{-1}(S).$$

Therefore, $\tau_{\tilde{D},S} \in \text{FCITab}(\tilde{D})$ exhausts α over S , so $\alpha \in \mathcal{S}_{\tilde{D}}$ by Theorem 2.10. \square

Since $r_1 < r$,

$$\sum_{(i,j) \in \tilde{D}} i < \sum_{(i,j) \in D} i.$$

Thus, Claim 2.17 and induction yields $\tau_{\tilde{D}} \in \text{PerfectTab}(\tilde{D}, \alpha)$. Define $\tau_D \in \text{Tab}(D)$ by

$$\tau_D(i, j) = \begin{cases} \tau_{\tilde{D}}(r_1, c) & \text{if } (i, j) = (r, c), \\ \tau_{\tilde{D}}(i, j) & \text{otherwise.} \end{cases}$$

Then it is easy to check that $\tau_D \in \text{PerfectTab}(D, \alpha)$, so Case 1 is complete.

Case 2: (D does not contain boxes below row m). We say an inequality $\sum_{i \in S} \alpha_i \leq \theta_D(S)$ from (2) is *nontrivial* if

$$(8) \quad \sum_{i \in S} \alpha_i > 0 \quad \text{and} \quad \theta_D(S) < \#D.$$

Case 2a: (All nontrivial inequalities from (2) are strict). Thus if (8) holds, then

$$(9) \quad \sum_{i \in S} \alpha_i < \theta_D(S).$$

Claim 2.18. *There exists $c \in [n]$ such that $(m, c) \in D$.*

Proof of Claim 2.18. By Theorem 2.10, there exists some $\tau_{D, \{m\}} \in \text{FCITab}(D)$ which exhausts α over $\{m\}$. Then

$$\#\tau_{D, \{m\}}^{-1}(\{m\}) \geq \alpha_m > 0,$$

so $\tau_{D, \{m\}}(r, c) = m$ for some $(r, c) \in D$. Since $\tau_{D, \{m\}}$ is flagged, we must have $r \geq m$. But by the assumption of Case 2, there are no boxes below row m , so $r = m$. \square

Pick $c \in [n]$ as in Claim 2.18. Then let $\tilde{D} = D \setminus \{(m, c)\}$ and $\tilde{\alpha} = (\tilde{\alpha}_1, \dots, \tilde{\alpha}_n) := (\alpha_1, \dots, \alpha_{m-1}, \alpha_m - 1, 0, \dots, 0)$.

Claim 2.19. $\tilde{\alpha} \in \mathcal{S}_{\tilde{D}}$.

Proof of Claim 2.19. Since $\alpha_i = 0$ for all $i > m$, and $(m, c) \in D$, we have

$$(10) \quad \tilde{\alpha}_1 + \dots + \tilde{\alpha}_n = \alpha_1 + \dots + \alpha_n - 1 = \#D - 1 = \#\tilde{D}.$$

For each $S \subseteq [n]$, let

$$\tau_{\tilde{D}, S} = \pi_{D, S}|_{\tilde{D}} \in \text{FCITab}(\tilde{D})$$

be the restriction of $\pi_{D, S}$ to \tilde{D} . Then by Proposition 2.3,

$$(11) \quad \#\tau_{\tilde{D}, S}^{-1}(S) \geq \#\pi_{D, S}^{-1}(S) - 1 = \theta_D(S) - 1.$$

If $\sum_{i \in S} \alpha_i = 0$, then

$$\sum_{i \in S} \tilde{\alpha}_i = 0 \leq \#\tau_{\tilde{D}, S}^{-1}(S).$$

If $\theta_D(S) = \#D$, then by (10) and (11),

$$\sum_{i \in S} \tilde{\alpha}_i \leq \tilde{\alpha}_1 + \dots + \tilde{\alpha}_n = \#D - 1 = \theta_D(S) - 1 \leq \#\tau_{\tilde{D}, S}^{-1}(S).$$

Finally, if $\sum_{i \in S} \alpha_i > 0$ and $\theta_D(S) < \#D$, then (9) must hold, so by (9) and (11),

$$\sum_{i \in S} \tilde{\alpha}_i \leq \sum_{i \in S} \alpha_i \leq \theta_D(S) - 1 \leq \#\tau_{\tilde{D}, S}^{-1}(S).$$

In all three cases, $\tau_{\tilde{D}, S}$ exhausts $\tilde{\alpha}$ over S , so $\tilde{\alpha} \in \mathcal{S}_{\tilde{D}}$ by Theorem 2.10. \square

By construction,

$$\sum_{(i, j) \in \tilde{D}} i < \sum_{(i, j) \in D} i.$$

Thus, Claim 2.19 and induction yield $\tau_{\tilde{D}} \in \text{PerfectTab}(\tilde{D}, \tilde{\alpha})$. Define $\tau_D \in \text{Tab}(D)$ by

$$\tau_D(i, j) = \begin{cases} m & \text{if } (i, j) = (m, c), \\ \tilde{\tau}(i, j) & \text{otherwise.} \end{cases}$$

Clearly, τ_D is flagged, has content α , and satisfies $\tau_D^{-1}(\{\circ\}) = \emptyset$. The only potential obstruction to column-injectivity is that there could be some $r \neq m$ for which $\tau_D(r, c) = m$.

This is impossible, since τ_D is flagged, so such an r must be greater than m , but by the assumption of Case 2 there are no boxes below row m . Thus, $\tau_D \in \text{PerfectTab}(D, \alpha)$, so Case 2a is complete.

Case 2b: (There exists a tight, nontrivial inequality in (2)). Thus, there exists $A \subseteq [n]$ satisfying

$$(12) \quad 0 < \sum_{i \in A} \alpha_i = \theta_D(A) < \#D.$$

Let $D^{(1)} = \pi_{D,A}^{-1}(A)$ and $D^{(2)} = D \setminus D^{(1)}$. Then for each $i \in [n]$, set

$$\alpha_i^{(1)} = \begin{cases} \alpha_i & \text{if } i \in A, \\ 0 & \text{if } i \notin A \end{cases} \quad \text{and} \quad \alpha_i^{(2)} = \begin{cases} \alpha_i & \text{if } i \notin A, \\ 0 & \text{if } i \in A. \end{cases}$$

Claim 2.20. $\alpha^{(1)} := (\alpha_1^{(1)}, \dots, \alpha_n^{(1)}) \in \mathcal{S}_{D^{(1)}}$.

Proof of Claim 2.20. By (12) and Proposition 2.3, we have

$$\alpha_1^{(1)} + \dots + \alpha_n^{(1)} = \sum_{i \in A} \alpha_i = \theta_D(A) = \#\pi_{D,A}^{-1}(A) = \#D^{(1)}.$$

Let $S \subseteq [n]$ and $T = S \cap A$. Then set

$$\tau_{D^{(1)},S} = \pi_{D,T}|_{D^{(1)}} \in \text{FCITab}(D^{(1)}).$$

By Corollary 2.15, $\pi_{D,T}^{-1}(T) \subseteq D^{(1)}$, so $\tau_{D^{(1)},S}^{-1}(T) = \pi_{D,T}^{-1}(T)$. Thus, by Remark 2.11, $\tau_{D^{(1)},S}$ exhausts α over T . Hence,

$$\sum_{i \in S} \alpha_i^{(1)} = \sum_{i \in T} \alpha_i \leq \#\tau_{D^{(1)},S}^{-1}(T) \leq \#\tau_{D^{(1)},S}^{-1}(S),$$

so $\tau_{D^{(1)},S}$ exhausts $\alpha^{(1)}$ over S , and consequently $\alpha^{(1)} \in \mathcal{S}_{D^{(1)}}$ by Theorem 2.10. \square

Claim 2.21. $\alpha^{(2)} := (\alpha_1^{(2)}, \dots, \alpha_n^{(2)}) \in \mathcal{S}_{D^{(2)}}$.

Proof of Claim 2.21. By (12) and Proposition 2.3,

$$\alpha_1^{(2)} + \dots + \alpha_n^{(2)} = \alpha_1 + \dots + \alpha_n - \sum_{i \in A} \alpha_i = \#D - \theta_D(A) = \#D - \#\pi_{D,A}^{-1}(A) = \#D^{(2)}.$$

Let $S \subseteq [n]$, $T = S \setminus A$, and $U = A \cup T$. Then by Theorem 2.10, Remark 2.11, (12), Proposition 2.3, and Lemma 2.14, we can write

$$\begin{aligned} \sum_{i \in S} \alpha_i^{(2)} &= \sum_{i \in U} \alpha_i - \sum_{i \in A} \alpha_i \leq \#\pi_{D,U}^{-1}(U) - \theta_D(A) \\ &= \#\pi_{D,U}^{-1}(U) - \#\pi_{D,A}^{-1}(A) = \#\pi_{D^{(2)},T}^{-1}(T) \leq \#\pi_{D^{(2)},T}^{-1}(S). \end{aligned}$$

Thus, $\tau_{D^{(2)},S} := \pi_{D^{(2)},T}$ exhausts $\alpha^{(2)}$ over S , so $\alpha^{(2)} \in \mathcal{S}_{D^{(2)}}$ by Theorem 2.10. \square

By (12) and Proposition 2.3, we have

$$0 < \#\pi_{D,A}^{-1}(A) < \#D,$$

so $D^{(1)}, D^{(2)} \subsetneq D$. Thus, by Claims 2.20 and 2.21 and induction, there exist

$$\tau_{D^{(1)}} \in \text{PerfectTab}(D^{(1)}, \alpha^{(1)}) \quad \text{and} \quad \tau_{D^{(2)}} \in \text{PerfectTab}(D^{(2)}, \alpha^{(2)}).$$

Define $\tau_D = \tau_{D^{(1)}} \cup \tau_{D^{(2)}} \in \text{Tab}(D)$ by

$$\tau_D(i, j) = \begin{cases} \tau_{D^{(1)}}(i, j) & \text{if } (i, j) \in D^{(1)}, \\ \tau_{D^{(2)}}(i, j) & \text{if } (i, j) \in D^{(2)}. \end{cases}$$

Clearly τ_D is flagged and satisfies $\tau_D^{-1}(\{\circ\}) = \emptyset$. It has content α because $\alpha = \alpha^{(1)} + \alpha^{(2)}$, and it is column-injective because the images of $\tau_{D^{(1)}}$ and $\tau_{D^{(2)}}$ are disjoint. Therefore, $\tau_D \in \text{PerfectTab}(D, \alpha)$ and Case 2b is complete.

This completes the proof of Theorem 2.13. \square

2.2. Polytopal descriptions of perfect tableaux. Given $D \subseteq [n]^2$ and $\alpha = (\alpha_1, \dots, \alpha_n) \in \mathbb{Z}_{\geq 0}^n$, define the indicator polytope

$$\mathcal{P}(D, \alpha) \subseteq \mathbb{R}^{n^2}$$

to be the polytope with points of the form $(\alpha_{ij})_{i,j \in [n]} = (\alpha_{11}, \dots, \alpha_{n1}, \dots, \alpha_{1n}, \dots, \alpha_{nn})$ governed by the inequalities (A)-(C) below.

(A) Column-Injectivity Conditions: For all $i, j \in [n]$,

$$0 \leq \alpha_{ij} \leq 1.$$

(B) Content Conditions: For all $i \in [n]$,

$$\sum_{j=1}^n \alpha_{ij} = \alpha_i.$$

(C) Flag Conditions: For all $s, j \in [n]$,

$$\sum_{i=1}^s \alpha_{ij} \geq \#\{(i, j) \in D : i \leq s\}.$$

Proposition 2.22. *Let $D \subseteq [n]^2$ and $\alpha = (\alpha_1, \dots, \alpha_n) \in \mathbb{Z}_{\geq 0}^n$ with $\alpha_1 + \dots + \alpha_n = \#D$. If $(\alpha_{ij}) \in \mathcal{P}(D, \alpha)$, then for each $j \in [n]$, we have*

$$\sum_{i=1}^n \alpha_{ij} = \#\{(i, j) \in D : i \in [n]\}.$$

Proof. From the flag conditions (C) where $s = n$, we have that

$$\sum_{i=1}^n \alpha_{ij} \geq \#\{(i, j) \in D : i \in [n]\}.$$

If this inequality is strict for any j , then using the content conditions (B), we can write

$$\#D = \alpha_1 + \dots + \alpha_n = \sum_{i=1}^n \sum_{j=1}^n \alpha_{ij} = \sum_{j=1}^n \sum_{i=1}^n \alpha_{ij} > \sum_{j=1}^n \#\{(i, j) \in D : i \in [n]\} = \#D,$$

a contradiction. \square

Theorem 2.23. *Let $D \subseteq [n]^2$ and $\alpha = (\alpha_1, \dots, \alpha_n) \in \mathbb{Z}_{\geq 0}^n$. Then $\text{PerfectTab}(D, \alpha) \neq \emptyset$ if and only if $\alpha_1 + \dots + \alpha_n = \#D$ and $\mathcal{P}(D, \alpha) \cap \mathbb{Z}^{n^2} \neq \emptyset$.*

Proof. (\Rightarrow) By Proposition 2.12, we have $\alpha_1 + \cdots + \alpha_n = \#D$. Let $\tau \in \text{PerfectTab}(D, \alpha)$. Then for each $i, j \in [n]$, set

$$\alpha_{ij} = \#\{r \in [n] : \tau(r, j) = i\} = \begin{cases} 1 & \text{if } \tau(r, j) = i \text{ for some } r \in [n], \\ 0 & \text{otherwise,} \end{cases}$$

where the second equality follows from the fact that τ is column-injective.

Claim 2.24. $(\alpha_{ij}) \in \mathcal{P}(D, \alpha) \cap \mathbb{Z}^{n^2}$.

Proof of Claim 2.24. Clearly $(\alpha_{ij}) \in \mathbb{Z}^{n^2}$ and the column-injectivity conditions (A) hold. Since τ has content α ,

$$\sum_{j=1}^n \alpha_{ij} = \sum_{j=1}^n \#\{r \in [n] : \tau(r, j) = i\} = \#\tau^{-1}(\{i\}) = \alpha_i$$

for each $i \in [n]$, so the content conditions (B) hold. Finally, for each $s, j \in [n]$, we have

$$\sum_{i=1}^s \alpha_{ij} = \#\{r \in [n] : \tau(r, j) \leq s\} \geq \#\{(r, j) \in D : r \leq s\}$$

since τ is flagged. Thus, the flag conditions (C) also hold. \square

(\Leftarrow) Let $(\alpha_{ij}) \in \mathcal{P}(D, \alpha) \cap \mathbb{Z}^{n^2}$. By the column-injectivity conditions (A), $\alpha_{ij} \in \{0, 1\}$. Thus, by Proposition 2.22, there exists for each $j \in [n]$ a bijection

$$\varphi_j : \{i \in [n] : (i, j) \in D\} \rightarrow \{i \in [n] : \alpha_{ij} = 1\}$$

that is order-preserving, i.e., φ_j satisfies $\varphi_j(i) < \varphi_j(i')$ whenever $i < i'$. Define $\tau \in \text{Tab}(D)$ by $\tau(i, j) = \varphi_j(i)$.

Claim 2.25. $\tau \in \text{PerfectTab}(D, \alpha)$.

Proof of Claim 2.25. By construction, $\tau^{-1}(\{\circ\}) = \emptyset$. Since φ_j is injective and order-preserving, τ is strictly increasing along columns, hence column-injective. For each $i \in [n]$, the content conditions (B) imply

$$\tau^{-1}(\{i\}) = \sum_{j=1}^n \#\varphi_j^{-1}(\{i\}) = \sum_{j=1}^n \alpha_{ij} = \alpha_i,$$

so τ has content α . Finally, the flag conditions (C) show that for each $s, j \in [n]$,

$$\#\{i \leq s : (i, j) \in D\} \leq \sum_{i=1}^s \alpha_{ij} = \#\{i \leq s : \alpha_{ij} = 1\},$$

so $\varphi_j(i) \leq i$ for each $(i, j) \in D$ since φ_j is order-preserving. Thus, $\tau(i, j) = \varphi_j(i) \leq i$ and τ is flagged. Hence, $\tau \in \text{PerfectTab}(D, \alpha)$. \square

This shows that $\text{PerfectTab}(D, \alpha) \neq \emptyset$ and completes the proof of the theorem. \square

Remark 2.26. The proof of Claim 2.25 shows that if $\text{PerfectTab}(D, \alpha) \neq \emptyset$, then we can find $\tau \in \text{PerfectTab}(D, \alpha)$ which is not only column-injective, but also strictly increasing along columns, so $\tau(i, j) < \tau(i', j)$ whenever $i < i'$. Thus $\text{PerfectTab}(D, \alpha) \neq \emptyset$ if and only if $\text{PerfectTab}(D, \alpha)_\downarrow \neq \emptyset$.

Theorem 2.23 formulates the problem of determining if $\text{PerfectTab}(D, \alpha) \neq \emptyset$ in terms of feasibility of an integer linear programming problem. In general, integral feasibility is NP-complete. We now show that in our case, feasibility of the problem is equivalent to feasibility of its LP-relaxation:

Theorem 2.27. *Let $D \subseteq [n]^2$ and $\alpha = (\alpha_1, \dots, \alpha_n) \in \mathbb{Z}^n$ with $\alpha_1 + \dots + \alpha_n = \#D$. Then $\mathcal{P}(D, \alpha) \cap \mathbb{Z}^{n^2} \neq \emptyset$ if and only if $\mathcal{P}(D, \alpha) \neq \emptyset$.*

For reasons given in the Introduction, we provide two proofs of this fact.

Proof 1 of Theorem 2.27. We write the constraints (A)-(C) in the form $M\vec{x} \leq \vec{b}$ where M is a $(3n^2 + n) \times n^2$ block matrix and \vec{b} is a vector of length $3n^2 + n$ of the form

$$M = \begin{pmatrix} M_{A_1} \\ M_{A_2} \\ M_B \\ M_C \end{pmatrix} \text{ and } \vec{b} = (b_i)_{i=1}^{3n^2+n}.$$

Let \vec{b}_I denote the subvector of \vec{b} containing those b_i with $i \in I \subseteq [3n^2 + n]$. Also, we use the following coordinatization:

$$\vec{x} = (\alpha_{11}, \dots, \alpha_{n1}, \alpha_{12}, \dots, \alpha_{n2}, \dots, \alpha_{nn})^T.$$

- M_{A_1} is the $n^2 \times n^2$ block corresponding to the condition $0 \leq \alpha_{ij}$ from (A). Thus, $M_{A_1} = -I_{n^2}$ and $b_r = 0$ for $r \in [1, n^2]$.
- M_{A_2} is the $n^2 \times n^2$ block corresponding to $\alpha_{ij} \leq 1$ from (A). Hence, $M_{A_2} = I_{n^2}$ and $b_r = 1$ for $r \in [n^2 + 1, 2n^2]$.
- M_C is the $n^2 \times n^2$ matrix for (C). Thus,

$$M_C = \begin{pmatrix} M_{C_T} & 0 & \dots & 0 \\ 0 & M_{C_T} & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & M_{C_T} \end{pmatrix}$$

where $M_{C_T} = (c_{ij})_{1 \leq i, j \leq n}$ is lower triangular such that $c_{ij} = -1$ for $i \geq j$. Also,

$$b_{(2n^2+n)+n(j-1)+s} = -\#\{(i, j) \in D : i \leq s\}, \text{ for } s, j \in [n].$$

- M_B is the $n \times n^2$ block encoding (B). Take $M_B = (I_n \ I_n \ \dots \ I_n)$ and $\vec{b}_{[2n^2+1, 2n^2+n]} = (\alpha_i)_{i \in [n]}$. Clearly $M_B \vec{x} \leq (\alpha_i)_{i \in [n]}$ encodes the inequalities $\sum_{j=1}^n \alpha_{ij} \leq \alpha_i$. Now, (B) requires $\sum_{j=1}^n \alpha_{ij} = \alpha_i$. However, $\alpha_1 + \dots + \alpha_n = \#D$ ensures that

$$\begin{pmatrix} M_B \\ M_C \end{pmatrix} \vec{x} \leq \vec{b}_{[2n^2+1, 3n^2+n]} \text{ only if } M_B \vec{x} = (\alpha_i)_{i \in [n]}.$$

Summarizing, $M\vec{x} \leq \vec{b}$ indeed encodes (A)-(C).

Example 2.28. For $n = 2$ consider $\vec{x} = (\alpha_{11}, \alpha_{21}, \alpha_{12}, \alpha_{22})^T$ with $D = \{(1, 1), (1, 2), (2, 2)\} \subset [2] \times [2]$ and $\alpha = (2, 1)$.

We have

$$M_{A_1} \vec{x} = \begin{pmatrix} -1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix} \begin{pmatrix} \alpha_{11} \\ \alpha_{21} \\ \alpha_{12} \\ \alpha_{22} \end{pmatrix} \leq \begin{pmatrix} 0 \\ 0 \\ 0 \\ 0 \end{pmatrix}$$

$$\begin{aligned}
M_{A_2}\vec{x} &= \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} \alpha_{11} \\ \alpha_{21} \\ \alpha_{12} \\ \alpha_{22} \end{pmatrix} \leq \begin{pmatrix} 1 \\ 1 \\ 1 \\ 1 \end{pmatrix} \\
M_B\vec{x} &= \begin{pmatrix} 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \end{pmatrix} \begin{pmatrix} \alpha_{11} \\ \alpha_{21} \\ \alpha_{12} \\ \alpha_{22} \end{pmatrix} \leq \begin{pmatrix} \alpha_1 \\ \alpha_2 \end{pmatrix} = \begin{pmatrix} 2 \\ 1 \end{pmatrix} \\
M_C\vec{x} &= \begin{pmatrix} -1 & 0 & 0 & 0 \\ -1 & -1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & -1 & -1 \end{pmatrix} \begin{pmatrix} \alpha_{11} \\ \alpha_{21} \\ \alpha_{12} \\ \alpha_{22} \end{pmatrix} \leq \begin{pmatrix} -\#\{(i, 1) \in D : i \leq 1\} \\ -\#\{(i, 1) \in D : i \leq 2\} \\ -\#\{(i, 2) \in D : i \leq 1\} \\ -\#\{(i, 2) \in D : i \leq 2\} \end{pmatrix} = \begin{pmatrix} -1 \\ -1 \\ -1 \\ -2 \end{pmatrix}
\end{aligned}$$

Theorem 2.29. M is a totally unimodular matrix; that is, every minor of M equals 0, 1, or -1 .

Proof. Suppose M' is a square submatrix of M with k rows from M_{A_1} or M_{A_2} . We show by induction on k that $\det(M') \in \{0, \pm 1\}$.

For the base case $k = 0$, consider M' an $\ell \times \ell$ submatrix of M with only rows from M_B and M_C . Let M'_B, M'_C be the corresponding blocks of M' , i.e. $M' = \begin{pmatrix} M'_B \\ M'_C \end{pmatrix}$ where M'_B , or M'_C , is the submatrix of M_B , or M_C respectively, using the rows and columns of M' . Since M_B has one 1 per column, M'_B has at most one 1 per column. By the form of M_C , it is straightforward to row reduce M'_C to obtain a $(0, -1)$ -matrix M''_C with at most one -1 in each column. Let $M'' = \begin{pmatrix} M'_B \\ M''_C \end{pmatrix}$, an $\ell \times \ell$ matrix. It is textbook (see [13, Theorem 13.3]) that if a $(0, \pm 1)$ -matrix N has at most one 1 and at most one -1 in each column, N is totally unimodular; hence $\det(M') = \pm \det(M'') \in \{0, -1, 1\}$ as desired. Thus the base case holds.

Now suppose M' is a square submatrix of M that contains $k \geq 1$ rows from M_{A_1} or M_{A_2} . Let R be such a row from M_{A_1} or M_{A_2} . If R contains all 0's, $\det(M') = 0$, and we are done. Otherwise R contains a single ± 1 . Hence the cofactor expansion for $\det(M')$ along R gives $\det(M') = \pm \det(M'')$ where M'' is a submatrix of M with $k - 1$ rows from M_{A_1} or M_{A_2} . So by induction, $\det(M') \in \{0, \pm 1\}$, as required. \square

Since M is totally unimodular then any vertices of $M\vec{x} \leq \vec{b}$ are integral [13, Theorem 13.2]. Thus, if $\mathcal{P}(D, \alpha) \neq \emptyset$ then its vertices are integral, i.e., $\mathcal{P}(D, \alpha) \cap \mathbb{Z}^{n^2} \neq \emptyset$. \square

Proof 2 of Theorem 2.27. Given a point $(\alpha_{ij}) \in \mathcal{P}(D, \alpha)$, we say a pair of sequences

$$(r_1, \dots, r_{m+1}; c_1, \dots, c_m) \in [n]^{m+1} \times [n]^m,$$

for some $m \in \mathbb{Z}_{>0}$, is *stable* at (α_{ij}) if the properties (i)-(iv) below hold. The purpose of each property will become clear later.

- (i) $r_{m+1} = r_1$.
- (ii) For all $k \in [m]$, $\alpha_{r_k c_k}, \alpha_{r_{k+1} c_k} \notin \mathbb{Z}$.
- (iii) For all $k \in [m]$, if $i > r_{k+1}$ and $\alpha_{i c_k} \notin \mathbb{Z}$, then $i = r_k$.

(iv) There exists $(r, c) \in [n]^2$ such that

$$\#\{k \in [m] : (r, c) = (r_k, c_k)\} \neq \#\{k \in [m] : (r, c) = (r_{k+1}, c_k)\}.$$

Claim 2.30. For any $(\alpha_{ij}) \in \mathcal{P}(D, \alpha) \setminus \mathbb{Z}^{n^2}$, there exists $(r_1, \dots, r_{m+1}; c_1, \dots, c_m)$ stable at (α_{ij}) .

Proof of Claim 2.30. Choose r_1, c_1 such that $\alpha_{r_1 c_1} \notin \mathbb{Z}$, and assume that we have fixed r_k, c_k such that $\alpha_{r_k c_k} \notin \mathbb{Z}$. By Proposition 2.22, we have

$$\sum_{i=1}^n \alpha_{i c_k} = \#\{(i, c_k) \in D : i \in [n]\} \in \mathbb{Z}.$$

Thus, as $\alpha_{r_k c_k} \notin \mathbb{Z}$, it makes sense to set

$$(13) \quad r_{k+1} = \max\{i \neq r_k : \alpha_{i c_k} \notin \mathbb{Z}\}.$$

If $r_{k+1} = r_\ell$ for some $\ell \in [k]$, then end the construction of these sequences. Otherwise, the content conditions (B) say that

$$\sum_{j=1}^n \alpha_{r_{k+1} j} = \alpha_{r_{k+1}} \in \mathbb{Z},$$

and since $\alpha_{r_{k+1} c_k} \notin \mathbb{Z}$, we can choose $c_{k+1} \neq c_k$ such that $\alpha_{r_{k+1} c_{k+1}} \notin \mathbb{Z}$, completing the recursive definition. By the pigeonhole principle, this process must halt, yielding sequences $r_1, \dots, r_\ell, \dots, r_{m+1}$ and $c_1, \dots, c_\ell, \dots, c_m$ with $r_{m+1} = r_\ell$.

By disregarding the first $\ell - 1$ terms of each sequence, we may assume $\ell = 1$ without loss of generality. Then we assert that $(r_1, \dots, r_{m+1}; c_1, \dots, c_m)$ is stable at (α_{ij}) . Indeed, (i) and (ii) are immediate from the construction, (iii) follows from (13), and (iv) holds because $(r, c) := (r_2, c_2)$ exists and satisfies

$$\#\{k \in [m] : (r, c) = (r_k, c_k)\} = 1 \quad \text{and} \quad \#\{k \in [m] : (r, c) = (r_{k+1}, c_k)\} = 0,$$

since $c_2 \neq c_1$ and $r_2 \neq r_k$ for all $k \neq 2$. □

We now fix a pair of sequences $(r_1, \dots, r_{m+1}; c_1, \dots, c_m)$. Given (α_{ij}) and $\delta > 0$, set

$$(14) \quad \alpha_{ij}^\delta = \alpha_{ij} + \delta[\#\{k \in [m] : (i, j) = (r_k, c_k)\} - \#\{k \in [m] : (i, j) = (r_{k+1}, c_k)\}].$$

Claim 2.31. If $(r_1, \dots, r_{m+1}; c_1, \dots, c_m)$ is stable at $(\alpha_{ij}) \in \mathcal{P}(D, \alpha)$, then $(\alpha_{ij}^\delta) \in \mathcal{P}(D, \alpha)$ for some $\delta > 0$.

Proof of Claim 2.31. First, note that the content conditions (B) are preserved regardless of our choice of δ . Indeed, for each $i \in [n]$,

$$\begin{aligned} \sum_{j=1}^n \alpha_{ij}^\delta &= \sum_{j=1}^n [\alpha_{ij} + \delta[\#\{k \in [m] : (i, j) = (r_k, c_k)\} - \#\{k \in [m] : (i, j) = (r_{k+1}, c_k)\}]] \\ &= \alpha_i + \delta[\#\{k \in [m] : i = r_k\} - \#\{k \in [m] : i = r_{k+1}\}], \end{aligned}$$

and the term in brackets vanishes by (i).

We next check the flag conditions (C). For each $s, j \in [n]$, we can write

$$\begin{aligned}
\sum_{i=1}^s \alpha_{ij}^\delta &= \sum_{i=1}^s [\alpha_{ij} + \delta[\#\{k \in [m] : (i, j) = (r_k, c_k)\} - \#\{k \in [m] : (i, j) = (r_{k+1}, c_k)\}]] \\
&= \sum_{i=1}^s \alpha_{ij} + \delta[\#\{k \in [m] : s \geq r_k \text{ and } j = c_k\} - \#\{k \in [m] : s \geq r_{k+1} \text{ and } j = c_k\}] \\
(15) \quad &\geq \sum_{i=1}^s \alpha_{ij} - \delta[\#\{k \in [m] : r_{k+1} \leq s < r_k \text{ and } j = c_k\}].
\end{aligned}$$

Thus, if $\#\{k \in [m] : r_{k+1} \leq s < r_k \text{ and } j = c_k\} = 0$, then the flag condition (C) for these s, j is preserved.

Otherwise, $r_{k+1} \leq s < r_k$ and $j = c_k$ for some $k \in [m]$, so (ii) and (iii) tell us that there is exactly one $i > s$ for which $\alpha_{ij} \notin \mathbb{Z}$, namely $i = r_k$. This, combined with Proposition 2.22, shows that

$$(16) \quad \sum_{i=1}^s \alpha_{ij} = \sum_{i=1}^n \alpha_{ij} - \sum_{i=s+1}^n \alpha_{ij} = \#\{(i, j) \in D : i \in [n]\} - \sum_{i=s+1}^n \alpha_{ij} \notin \mathbb{Z}.$$

By the nonintegrality from (16), the flag inequalities (C) for $(\alpha_{ij}) \in \mathcal{P}(D, \alpha)$ are strict:

$$(17) \quad \sum_{i=1}^s \alpha_{ij} > \#\{(i, j) \in D : i \leq s\}.$$

Hence, by taking δ sufficiently small and applying (15) and (17), we can ensure

$$\sum_{i=1}^s \alpha_{ij}^\delta \geq \sum_{i=1}^s \alpha_{ij} - \delta[\#\{k \in [m] : r_{k+1} \leq s < r_k \text{ and } j = c_k\}] \geq \#\{(i, j) \in D : i \leq s\}$$

for all $s, j \in [n]$, so the flag conditions (C) will be preserved. If $\alpha_{ij} \neq \alpha_{ij}^\delta$ then by (14) we must have $(i, j) = (r_k, c_k)$ or $(i, j) = (r_{k+1}, c_k)$ for some k , which by (ii) implies $0 < \alpha_{ij} < 1$. So we can require in addition that δ be small enough that $0 \leq \alpha_{ij}^\delta \leq 1$ for all $i, j \in [n]$. For such δ , the conditions (A)-(C) all hold, so $(\alpha_{ij}^\delta) \in \mathcal{P}(D, \alpha)$. \square

Finally, choose a point $(\alpha_{ij}) \in \mathcal{P}(D, \alpha)$ with the maximum number of integer coordinates. If $(\alpha_{ij}) \in \mathbb{Z}^{n^2}$, then we are done. Otherwise, there exists $(r_1, \dots, r_{m+1}; c_1, \dots, c_m)$ that is stable at (α_{ij}) by Claim 2.30. By (iv), there exists $(r, c) \in [n]^2$ such that $|\alpha_{rc}^\delta| \rightarrow \infty$ as $\delta \rightarrow \infty$, so α_{rc}^δ violates the column-injectivity conditions (A) for large δ . This, combined with Claim 2.31, shows that the set $S = \{\delta > 0 : (\alpha_{ij}^\delta) \in \mathcal{P}(D, \alpha)\}$ is nonempty and bounded above. Thus, we can define $\eta = \sup S$ and set $(\tilde{\alpha}_{ij}) = (\alpha_{ij}^\eta)$. Since $\mathcal{P}(D, \alpha)$ is closed and the map $\delta \mapsto (\alpha_{ij}^\delta)$ from S to $\mathcal{P}(D, \alpha)$ is continuous, this supremum is in fact a maximum, and $(\tilde{\alpha}_{ij}) \in \mathcal{P}(D, \alpha)$. By our choice of (α_{ij}) , we cannot have $\tilde{\alpha}_{r_k c_k} \in \mathbb{Z}$ or $\tilde{\alpha}_{r_{k+1} c_k} \in \mathbb{Z}$ for any $k \in [m]$, since then $(\tilde{\alpha}_{ij})$ has more integer coordinates than (α_{ij}) . Thus, $(r_1, \dots, r_{m+1}; c_1, \dots, c_m)$ is stable at $(\tilde{\alpha}_{ij})$, so by Claim 2.31, there exists $\delta > 0$ for which $(\tilde{\alpha}_{ij}^\delta) \in \mathcal{P}(D, \alpha)$. But then $(\alpha_{ij}^{\eta+\delta}) = (\tilde{\alpha}_{ij}^\delta) \in \mathcal{P}(D, \alpha)$, contradicting the maximality of η . \square

In summary, applying the results of this section to $D = D(w)$,

$$(18) \quad c_{\alpha, w} > 0 \stackrel{[4]}{\iff} \alpha \in \mathcal{S}_D \stackrel{\text{Thm 2.13}}{\iff} \text{PerfectTab}(D, \alpha) \neq \emptyset \stackrel{\text{Thm 2.23}}{\iff} \mathcal{P}(D, \alpha) \cap \mathbb{Z}^{n^2} \neq \emptyset \stackrel{\text{Thm 2.27}}{\iff} \mathcal{P}(D, \alpha) \neq \emptyset.$$

If $D \subseteq [n]^2$ has many identical columns, then many of the flag conditions (C) will look essentially the same. Thus, for efficiency of computation, we construct a “compressed” version of $\mathcal{P}(D, \alpha)$ that removes some of the repetitive inequalities.

A tuple $\mathcal{C} = (m, \{P_k\}_{k=1}^\ell, \{p_k\}_{k=1}^\ell, \{\lambda_k\}_{k=1}^\ell)$ is a *compression* of $D \subseteq [n]^2$ if:

- $m \leq n$ is a nonnegative integer such that $(r, p) \notin D$ whenever $r > m$ and $p \in [n]$,
- $P = P_1 \dot{\cup} \dots \dot{\cup} P_\ell \subseteq [n]$ such that if $p, p' \in P_k$ then

$$\{r \in [n] : (r, p) \in D\} = \{r \in [n] : (r, p') \in D\},$$

and moreover if D is nonempty in column p then $p \in P_k$ for some $k \in [\ell]$.

- $p_k \in P_k$ a representative for each $k \in [\ell]$, and
- $\lambda_k = \#P_k$ for each $k \in \ell$.

For $D \subseteq [n]^2$, a compression \mathcal{C} of D , and $\tilde{\alpha} = (\tilde{\alpha}_1, \dots, \tilde{\alpha}_m) \in \mathbb{Z}_{\geq 0}^m$ define

$$(19) \quad \mathcal{Q}(D, \mathcal{C}, \tilde{\alpha}) \subseteq \mathbb{R}^{m\ell}$$

to be the polytope with points of the form $(\tilde{\alpha}_{ik})_{i \in [m], k \in [\ell]}$ satisfying (A')-(C') below.

(A') Column-Injectivity Conditions: For all $i \in [m], k \in [\ell]$,

$$0 \leq \tilde{\alpha}_{ik} \leq 1.$$

(B') Content Conditions: For all $i \in [m]$,

$$\sum_{k=1}^{\ell} \lambda_k \tilde{\alpha}_{ik} = \alpha_i.$$

(C') Flag Conditions: For all $s \in [m], k \in [\ell]$,

$$\sum_{i=1}^s \tilde{\alpha}_{ik} \geq \#\{(i, p_k) \in D : i \leq s\}.$$

Remark 2.32. We can always take $m = \ell = n$ and $P_k = \{k\}$ for each $k \in [\ell]$, in which case $\mathcal{Q}(D, \mathcal{C}, \tilde{\alpha}) = \mathcal{P}(D, \alpha) \subseteq \mathbb{R}^{n^2}$.

Theorem 2.33. Let $\alpha = (\alpha_1, \dots, \alpha_n) \in \mathbb{Z}_{\geq 0}^n$ and $\tilde{\alpha} = (\tilde{\alpha}_1, \dots, \tilde{\alpha}_m) := (\alpha_1, \dots, \alpha_m)$. Then $\alpha_1 + \dots + \alpha_n = \#D$ and $\mathcal{P}(D, \alpha) \neq \emptyset$ if and only if $\alpha_1 + \dots + \alpha_m = \#D$, $\alpha_{m+1} = \dots = \alpha_n = 0$, and $\mathcal{Q}(D, \mathcal{C}, \tilde{\alpha}) \neq \emptyset$.

Proof. (\Rightarrow) Let $(\alpha_{ij}) \in \mathcal{P}(D, \alpha)$. Then by the content and flag conditions (B) and (C),

$$\begin{aligned} \#D = \alpha_1 + \dots + \alpha_n &\geq \alpha_1 + \dots + \alpha_m = \sum_{i=1}^m \sum_{j=1}^n \alpha_{ij} \\ &= \sum_{j=1}^n \sum_{i=1}^m \alpha_{ij} \geq \sum_{j=1}^n \#\{(i, j) \in D : i \leq m\} = \#D. \end{aligned}$$

Thus, $\alpha_1 + \dots + \alpha_m = \#D$ and $\alpha_{m+1} = \dots = \alpha_n = 0$. Now, for each $i \in [m]$ and $k \in [\ell]$, set

$$\tilde{\alpha}_{ik} = \frac{1}{\lambda_k} \sum_{j \in P_k} \alpha_{ij}.$$

We claim that $(\tilde{\alpha}_{ik}) \in \mathcal{Q}(D, \mathcal{C}, \alpha)$. First, for each $i \in [m]$ and $k \in [\ell]$, we have

$$0 \leq \tilde{\alpha}_{ik} = \frac{1}{\lambda_k} \sum_{j \in P_k} \alpha_{ij} \leq \frac{1}{\lambda_k} \sum_{j \in P_k} 1 = 1,$$

so the column-injectivity conditions (A') are satisfied. Next, for each $i \in [m]$, (B) implies

$$\sum_{k=1}^{\ell} \lambda_k \tilde{\alpha}_{ik} = \sum_{k=1}^{\ell} \sum_{j \in P_k} \alpha_{ij} = \sum_{j=1}^n \alpha_{ij} = \alpha_i,$$

so the content conditions (B') are satisfied. Finally, for each $s \in [m]$ and $k \in [\ell]$, (C) implies

$$\sum_{i=1}^s \tilde{\alpha}_{ik} = \frac{1}{\lambda_k} \sum_{j \in P_k} \sum_{i=1}^s \alpha_{ij} \geq \frac{1}{\lambda_k} \sum_{j \in P_k} \#\{(i, j) \in D : i \leq s\} = \#\{(i, p_k) \in D : i \leq s\},$$

so the flag conditions (C') are satisfied.

(\Leftarrow) Clearly $\alpha_1 + \dots + \alpha_n = \#D$. Let $(\tilde{\alpha}_{ik}) \in \mathcal{Q}(D, \mathcal{C}, \tilde{\alpha})$. For each $i, j \in [n]$, set

$$\alpha_{ij} = \begin{cases} 0 & \text{if } i > m, \\ \tilde{\alpha}_{ik} & \text{if } i \leq m \text{ and } j \in P_k. \end{cases}$$

We claim that $(\alpha_{ij}) \in \mathcal{P}(D, \alpha)$. The column-injectivity conditions (A) are clear. If $i > m$,

$$\sum_{j=1}^n \alpha_{ij} = 0 = \alpha_i.$$

Otherwise $i \leq m$, and (B') implies

$$\sum_{j=1}^n \alpha_{ij} = \sum_{k=1}^{\ell} \sum_{j \in P_k} \tilde{\alpha}_{ik} = \sum_{k=1}^{\ell} \lambda_k \tilde{\alpha}_{ik} = \alpha_i.$$

Thus, the content conditions (B) hold. Finally, if $s \in [n]$ and $j \in P_k$, then (C') implies

$$\sum_{i=1}^s \alpha_{ij} = \sum_{i=1}^{\min\{s, m\}} \tilde{\alpha}_{ik} \geq \#\{(i, p_k) \in D : i \leq \min\{s, m\}\} = \#\{(i, j) \in D : i \leq s\}.$$

Hence, the flag conditions (C) hold as well. \square

2.3. Deciding membership in the Schubitope. We use the above results of this section to give a polynomial time algorithm to check if a lattice point is in the Schubitope.

Let $D \subseteq [n]^2$, and fix a compression $\mathcal{C} = (m, \{P_k\}_{k=1}^{\ell}, \{p_k\}_{k=1}^{\ell}, \{\lambda_k\}_{k=1}^{\ell})$ of D (as in Section 2.2).

Theorem 2.34. *Let $\alpha = (\alpha_1, \dots, \alpha_n) \in \mathbb{Z}_{\geq 0}^n$. Then $\alpha \in \mathcal{S}_D$ if and only if $\alpha_1 + \dots + \alpha_m = \#D$, $\alpha_{m+1} = \dots = \alpha_n = 0$, and $\mathcal{Q}(D, \mathcal{C}, \tilde{\alpha}) \neq \emptyset$, where $\tilde{\alpha} = (\tilde{\alpha}_1, \dots, \tilde{\alpha}_m) := (\alpha_1, \dots, \alpha_m)$.*

Proof. This follows from Theorems 2.13, 2.23, 2.27, and 2.33. \square

For each $k \in [\ell]$, let $R_k(\mathcal{C}) = \{r \in [n] : (r, p_k) \in D\} \subseteq [m]$.

Theorem 2.35. *Given as input $\{R_k(\mathcal{C})\}_{k=1}^{\ell}$, $\{\lambda_k\}_{k=1}^{\ell}$, and $\tilde{\alpha} = (\tilde{\alpha}_1, \dots, \tilde{\alpha}_m) \in \mathbb{Z}_{\geq 0}^m$ satisfying $\tilde{\alpha}_1 + \dots + \tilde{\alpha}_m = \#D$, one can decide if $\alpha := (\tilde{\alpha}_1, \dots, \tilde{\alpha}_m, 0, \dots, 0) \in \mathbb{Z}_{\geq 0}^n$ lies in \mathcal{S}_D in polynomial time in m and ℓ .*

Remark 2.36. In view of Theorem 2.34, this input is most natural, because the conditions $\alpha_1 + \dots + \alpha_m = \#D$ and $\alpha_{m+1} = \dots = \alpha_n = 0$ are clearly necessary, and it contains the minimum amount of information we need to compute $\mathcal{Q}(D, \mathcal{C}, \tilde{\alpha})$.

Remark 2.37. As in Remark 2.32, we can take $m = \ell = n$ and $P_k = \{k\}$ for each $k \in [\ell]$, so we can check if α is in \mathcal{S}_D in polynomial time in n regardless of the structure of D .

Proof of Theorem 2.35. Since $R_k(\mathcal{C})$ takes m bits to encode for each $k \in [\ell]$, and $\mathcal{Q}(D, \mathcal{C}, \tilde{\alpha}) \subseteq \mathbb{R}^{m\ell}$ is governed by $O(m\ell)$ constraints, $\mathcal{Q}(D, \mathcal{C}, \tilde{\alpha})$ can be constructed in polynomial time in m and ℓ . By Theorem 2.34, we are done using L. Khachiyan's ellipsoid method [14]. \square

3. COMPUTING ROTHE DIAGRAMS

We will repeatedly use the following to establish the complexity of computing preliminary data of $D(w)$ given $\text{code}(w)$.

Proposition 3.1. *There exists an $O(L^2)$ -time algorithm to compute $(w(1), \dots, w(L))$ from the input $\text{code}(w) = (c_1, \dots, c_L)$.*

Proof. Clearly $w(1) = c_1 + 1$. After determining $w(1), \dots, w(i-1)$, we determine (in $O(L)$ -time) $\pi := \pi^{(i)} \in S_{i-1}$ such that $(w(\pi(1)) < w(\pi(2)) < \dots < w(\pi(i-1)))$. Next, set

$$B := (w(\pi(1)), w(\pi(2)) - w(\pi(1)), w(\pi(3)) - w(\pi(2)), \dots, w(\pi(i-1)) - w(\pi(i-2))).$$

Let

$$V_t := \sum_{j=1}^t (B_j - 1), \quad \text{for } 0 \leq t \leq i-1.$$

Set $w(i) := c_i + T + 1$ where $T := \max_{t \in [0, i-1]} \{t : c_i \geq V_t\}$. By construction, $w(1), \dots, w(i)$ is a partial permutation with code $(c_1, \dots, c_{i-1}, c_i)$. Each stage $1 \leq i \leq L$ takes $O(i)$ -time. \square

The *essential set* of w consists of the maximally southeast boxes of each connected component of $D(w)$, i.e.,

$$(20) \quad \text{Ess}(w) = \{(i, j) \in D(w) : (i+1, j), (i, j+1) \notin D(w)\}.$$

If it exists, we call the connected component of $D(w)$ involving $(1, 1)$ the *dominant component* and denote it by $\text{Dom}(w)$. For instance, in Example 1.1, $\text{Dom}(w)$ has shape $(4, 2, 2, 2)$. Further, if it exists, the *accessible box* \mathbf{z}_w is the southmost then eastmost box in $\text{Ess}(w) \setminus \text{Dom}(w)$. In Example 1.1,

$$\text{Ess}(w) = \{(1, 4), (3, 4), (3, 7), (4, 2)\} \text{ and } \mathbf{z}_w = (3, 7).$$

(Although $(4, 2)$ is the southmost box of $\text{Ess}(w)$, it is in $\text{Dom}(w)$, and hence not the accessible.)

We will need the following in Section 5:

Proposition 3.2. *Given $\text{code}(w)$, there exists an $O(L^2)$ -time algorithm to compute $\mathbf{z}_w = (r, c)$ or determine it does not exist.*

Proof. Use Proposition 3.1 to find $(w(1), \dots, w(L))$ in $O(L^2)$ -time. Next, compute

$$w_{NW}(i) := \{w(j) : w(j) \leq w(i), j \leq i\}.$$

Then take

$$Y(i) := \{q - 1 : q \in w_{NW}(i)\} \setminus w_{NW}(i), \quad \text{for } i \in [L].$$

Compute $k_i := \max Y(i)$ for $i \in [L]$ in $O(L^2)$ -time (if $k_i \geq 1$, then k_i is the column index of the eastmost box of $D(w)$ in row i). In $O(L^2)$ -time, calculate

$$I := \{i \in [2, \dots, L] : k_i > \min_{j < i} w(j)\}.$$

Let $Y := \{(i, k_i) : i \in I\}$. Hence, $Y \cap \text{Dom}(w) = \emptyset$. Thus, if $Y = \emptyset$, \mathbf{z}_w does not exist. Otherwise, $\mathbf{z}_w \in Y$. Thus, in $O(L)$ -time, determine $r := \max\{i : (i, k_i) \in Y\}$. Output $\mathbf{z}_w = (r, k_r)$. \square

The *pivots* of \mathbf{z}_w denoted $\text{Piv}(\mathbf{z}_w)$ are the \bullet 's of $D(w)$ that are maximally southeast, among those northwest of \mathbf{z}_w . In Example 1.1, $\text{Piv}((3, 7)) = \{(2, 3), (1, 5)\}$.

4. PROOFS OF THEOREMS 1.2 AND 1.3

4.1. Proof of Theorem 1.2. By (1) combined with Theorem 2.35, it remains to establish the complexity of computing a compression of $D(w)$. For this, we need the following lemmas and propositions. Fix $w \in S_\infty$ with $\text{code}(w) = (c_1, \dots, c_L)$. Let $\sigma \in S_L$ be such that $\{w(\sigma(1)) < w(\sigma(2)) < \dots < w(\sigma(L))\}$. For convenience, set $w(\sigma(0)) := 0$.

Lemma 4.1. *For $1 \leq h \leq L$, and for all*

$$j_1, j_2 \in \{w(\sigma(h-1)) + 1, w(\sigma(h-1)) + 2, \dots, w(\sigma(h)) - 1\},$$

we have $(i, j_1) \in D(w)$ if and only if $(i, j_2) \in D(w)$.

Proof. For each k , let $u_1^{(k)} < \dots < u_k^{(k)}$ be $w(1), w(2), \dots, w(k)$ sorted in increasing order. Set $u_0^{(k)} := 0$. The lemma follows from the inductive claim that in the first k rows of $D(w)$, the columns $u_{h-1}^{(k)} + 1, u_{h-1}^{(k)} + 2, \dots, u_h^{(k)} - 1$ are the same. The base case $k = 1$ is clear. The inductive step is straightforward by considering how, in row $k + 1$ of $D(w)$, the \bullet and its ray emanating east affects the columns. \square

Define a collection of intervals in $[n]$ by

$$P'_{2k-1} := [w(\sigma(k-1)) + 1, w(\sigma(k)) - 1] \text{ and } P'_{2k} := \{w(\sigma(k))\}, \text{ for } 1 \leq k \leq L.$$

Let $1 \leq h_1 < h_2 < \dots < h_\ell \leq 2L$ be indices of the intervals P'_h that are nonempty. Set $P_i := P'_{h_i}$.

Lemma 4.2. *If $j_1, j_2 \in P_k$ for some k , then $(i, j_1) \in D(w) \iff (i, j_2) \in D(w)$.*

Proof. This follows by the definition of $\{P_k\}_{k=1}^\ell$ together with Lemma 4.1. \square

Let $p_k := \min\{p \in P_k\}$ for each $k \in [\ell]$.

Proposition 4.3. *There exists an $O(L^2)$ -time algorithm to compute $\{P_k\}_{k=1}^\ell$, $\{p_k\}_{k=1}^\ell$, and $\{\#P_k\}_{k=1}^\ell$ from the input $\text{code}(w) = (c_1, \dots, c_L)$.*

Proof. Proposition 3.1 computes $(w(1), \dots, w(L))$ in $O(L^2)$ -time. It takes $O(L \log(L))$ -time to sort $(w(1), \dots, w(L))$, i.e., to compute $\sigma \in S_L$. Computing the endpoints, and thus cardinalities, of the P'_k takes $O(L)$ -time as there are at most $2L$ of them. Then we reindex $\{\#P'_k\}_{k=1}^{2L}$ to obtain $\{\#P_k\}_{k=1}^\ell$ in $O(L)$ -time. \square

For each $k \in [\ell]$, let

$$R_k := \{r \in [L] : (r, p_k) \in D(w)\}.$$

Proposition 4.4. *Computing $\{R_k\}_{k=1}^\ell$ from $\text{code}(w)$ takes $O(L^2)$ -time.*

Proof. By $D(w)$'s definition, $r \in R_k$ if and only if $w(r) > p_k$ and $p_k \notin \{w(i) : i < r\}$. Propositions 4.3 and 3.1 give $\{P_k\}_{k=1}^\ell$, $\{p_k\}_{k=1}^\ell$ and $\{w(1), \dots, w(L)\}$ in $O(L^2)$ -time. \square

Conclusion of proof of Theorem 1.2: Proposition 4.3 computes $\{P_k\}_{k=1}^\ell$, $\{p_k\}_{k=1}^\ell$, and $\{\#P_k\}_{k=1}^\ell$ in $O(L^2)$ -time. Proposition 4.4 finds $\{R_k\}_{k=1}^\ell$ in $O(L^2)$ -time. One checks, using Lemma 4.2, that $\mathcal{C} = (L, \{P_k\}_{k=1}^\ell, \{p_k\}_{k=1}^\ell, \{\#P_k\}_{k=1}^\ell)$ is a compression of $D(w)$. Hence we may apply Theorem 2.35 by taking $D := D(w)$, $R_k(\mathcal{C}) := R_k$, $\lambda_k := \#P_k$ for $k \in [\ell]$ and $m := L$. Thus the result follows by (1). \square

4.2. Proof of Theorem 1.3; an application. Remark 2.26 combined with (18) proves the theorem. \square

Let $n_{132}(w)$ be the number of 132-patterns in $w \in S_n$, that is,

$$n_{132}(w) = \#\{(i, j, k) : 1 \leq i < j < k \leq n, w(i) < w(k) < w(j)\}.$$

Corollary 4.5. *There are at least $n_{132}(w) + 1$ distinct vectors α such that $c_{\alpha, w} > 0$.*

Proof. Suppose $i < j < k$ index a 132 pattern in w . There is a box b of $D(w)$ in row j and column $w(k)$. There are $N := n_{132}(w)$ many such boxes, b_1, \dots, b_N (all distinct), listed in English language reading order. Let M_i be boxes in the same column and connected component as b_i that are weakly north of b_i and strictly south of any b_j , where $j < i$. Iteratively define fillings $F_0, F_1, F_2, \dots, F_N$ of $D(w)$:

(F_0) Fill each box c of $D(w)$ with the row number of c .

(F_i) For $1 \leq i \leq N$, F_i is the same as F_{i-1} except that $F_i(c) := F_{i-1}(c) - 1$ if $c \in M_i$.

Clearly, $F_0 \in \text{PerfectTab}_\downarrow(D(w)) := \bigcup_\alpha \text{PerfectTab}_\downarrow(D(w), \alpha)$. Inductively assume $F_{i-1} \in \text{PerfectTab}_\downarrow(D(w))$. Since labels only decrease, F_i satisfies the row bound condition. Next we check that each column is strictly increasing. Let m_i be the northmost box of M_i . If m_i is adjacent and directly below some b_j (for a $j < i$) then

$$F_i(b_j) = F_0(b_j) - 1 < F_0(m_i) - 1 = F_i(m_i),$$

as needed. Otherwise suppose m_i is adjacent and south of a non-diagram position. Let d_i (if it exists) be the first diagram box directly north of m_i . Then $F_0(d_i) < F_0(m_i) - 1$. Hence

$$F_i(d_i) \leq F_0(d_i) < F_0(m_i) - 1 = F_i(m_i),$$

verifying column increasingness here as well. That F_i is column increasing elsewhere is clear since F_{i-1} is column increasing (by induction) and only labels of M_i are changed.

It remains to check that every label of F_i is in $\mathbb{Z}_{>0}$. Since each box of $D(w)$ is decremented at most once, the only concern is there is a box x in the first row that appears in some M_i , since then $F_0(x) = 1$ and $F_i(x) = 0$. However, in this case b_i must be in $\text{Dom}(w)$, which implies that the "1" in the 132-pattern associated to b_i could not exist, a contradiction. Thus $F_i \in \text{PerfectTab}_\downarrow(D(w))$, completing the induction.

Finally, under Theorem 1.3, each F_i corresponds to a distinct exponent vector since the sum of the labels is strictly decreasing at each step $F_{i-1} \mapsto F_i$. \square

From Corollary 4.5, this result of A. Weigandt [17] is immediate:

Corollary 4.6 (A. Weigandt's 132-bound). $\mathfrak{S}_w(1, 1, 1, \dots, 1) \geq n_{132}(w) + 1$.

As shown in [17], Corollary 4.6 in turn implies $\mathfrak{S}_w(1, 1, \dots, 1) \geq 3$ if $n_{132}(w) \geq 2$, a recent conjecture of R. P. Stanley [15].

5. COUNTING $c_{\alpha, w}$ IS IN #P

5.1. Vexillary permutations. A permutation $w \in S_n$ is *vexillary* if there does not exist a 2143 *pattern*, i.e., indices $i < j < k < l$ such that w has the pattern $w(j) < w(i) < w(l) < w(k)$. For example, $w = \underline{53841267}$ is not vexillary; we underlined the positions of a 2143 pattern. *Fulton's criterion* states that w is vexillary if and only if there do not exist $(a, b), (c, d) \in \text{Ess}(w)$ such that $a < c$ and $b < d$. In Example 1.1, w is not vexillary due to $(1, 4)$ and $(3, 7)$. Our main reference for this subsection is [10, Chapter 2].

We will also use this characterization of vexillary permutations:

Theorem 5.1. [8] *Given $\text{code}(w) = (c_1, \dots, c_L) \in \mathbb{Z}_{\geq 0}^n$, w vexillary if and only if*

- (i) *if i is such that $c_i > c_{i+1}$, then $c_i > c_j$ for any $j > i$, and*
- (ii) *if i, h are such that $c_i \geq c_h$, then $\#\{j : i < j < h, c_j < c_h\} \leq c_i - c_h$.*




The *shape* of a vexillary permutation v is the partition $\lambda(v)$ formed by sorting $\text{code}(v) = (c_1, c_2, \dots)$ into decreasing order. Now, if $c_i \neq 0$, let e_i be the greatest integer $j \geq i$ such that $c_j \geq c_i$. The *flag*

$$\phi(v) = (\phi_1 \leq \phi_2 \leq \dots \leq \phi_m)$$

for v is the sequence of e_i 's sorted into increasing order; see, e.g., [10, Definition 2.2.9].

Example 5.2. Consider $\text{code}(v) = (5, 1, 3, 1, 2)$ for the vexillary $v = 6253714$. Here

$$e = (1, 5, 3, 5, 5), \phi(v) = (1, 3, 5, 5, 5) \text{ and } \lambda(v) = (5, 3, 2, 1, 1).$$



For a partition $\lambda = (\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_m \geq 0)$ and a flag $\phi = (\phi_1 \leq \phi_2 \leq \dots \leq \phi_m)$ of positive integers, define the *flagged Schur function*

$$S_\lambda(\phi) = \det |h_{\lambda_i - i + j}(\phi_i)|_{i,j=1, \dots, m},$$

where

$$h_k(n) = \sum_{1 \leq i_1 \leq \dots \leq i_k \leq n} x_{i_1} \cdots x_{i_k}$$

is the complete homogeneous symmetric polynomial of degree k . Furthermore,

$$(21) \quad \mathfrak{S}_v = S_{\lambda(v)}(\phi(v)), \text{ for } v \text{ vexillary.}$$

A semistandard Young tableau of shape λ is *flagged* by ϕ if its entries in row i are $\leq \phi_i$; see Example 5.2. Denote the set of such tableaux by $\text{SSYT}(\lambda, \phi)$. Then

$$(22) \quad S_\lambda(\phi) = \sum_{T \in \text{SSYT}(\lambda, \phi)} x^{\text{content}(T)},$$

where $\text{content}(T) = (\mu_1, \dots, \mu_{\ell(\lambda)})$ such that μ_i is the number of i 's in T .

5.2. Graphical transition. The transition recurrence for \mathfrak{S}_w was found by A. Lascoux and M.-P. Schützenberger [8]. This is transition for the case discussed in [7]:

Theorem 5.3 ([8], cf. [7]). *Let $\mathbf{z}_w = (r, c)$ and $w' = w \cdot (r \ k)$ where $k = w^{-1}(c)$. Then*

$$(23) \quad \mathfrak{S}_w = x_r \mathfrak{S}_{w'} + \sum_{w'' = w' \cdot (i \ k)} \mathfrak{S}_{w''},$$

where the summation is over $\{i : (i, w(i)) \in \text{Piv}(\mathbf{z}_w)\}$.


We will use the *graphical* transition tree $\mathcal{T}(w)$ of [7]. This reformulates (23) in terms of Rothe diagrams and certain moves on these diagrams. By definition, $D(w)$ (equivalently w) will label the root of $\mathcal{T}(w)$. If w is vexillary, stop. Otherwise, there exists an accessible box $\mathbf{z}_w = (r, c) \in D(w)$ (if not, $D(w) = \text{Dom}(w)$, contradicting w is not vexillary).

The children of $D(w)$ are Rothe diagrams resulting from two types of moves:

- (T.1) *Deletion moves:* remove \mathbf{z}_w from $D(w)$. The resulting diagram is $D(w')$. Add an edge $D(w) \xrightarrow{x_r} D(w')$.
- (T.2) *March moves:* There is a move for each $\mathbf{x}^{(i)} = (i, w(i)) \in \text{Piv}(\mathbf{z}_w)$. Let \mathcal{R} be the rectangle with corners \mathbf{z}_w and $\mathbf{x}^{(i)}$. Remove $\mathbf{x}^{(i)}$ and its rays from $G(w)$ to form $G^{(i)}(w)$. Order the boxes $\{b_i\}_{i=1}^r$ in \mathcal{R} in English reading order. Move b_1 strictly north and strictly west to the closest position not occupied by other boxes of $D(w)$ or rays from $G^{(i)}(w)$. Repeat with b_2, b_3, \dots where b_j may move into a square left unoccupied by earlier moves. The resulting diagram will be $D(w'')$ where $w'' = w' \cdot (i \ k)$. Add an edge $D(w) \xrightarrow{i} D(w'')$.

Repeat for each child $D(u)$. Stop when u vexillary; these permutations are the leaves $\mathcal{L}(w)$ of $\mathcal{T}(w)$. (Multiple leaves may be labelled by the same permutation.)

Example 5.4. Let $w = 53841267$. We compute the march move 2 for the pivot $(2, 3)$:



The moved boxes during $D(w) \mapsto D(w'')$ are shaded gray.

Example 5.5. Let $w = 53861247$. Using $\mathcal{T}(w)$ from Figure 1, we compute

$$\begin{aligned} \mathfrak{S}_w = & x_4 \cdot \mathfrak{S}_{73541268} + x_4 \cdot \mathfrak{S}_{57341268} + x_3^2 x_4 \cdot \mathfrak{S}_{53641278} + x_3 x_4 \cdot \mathfrak{S}_{63541278} + x_3 x_4 \cdot \mathfrak{S}_{56341278} \\ & + \mathfrak{S}_{74531268} + \mathfrak{S}_{57431268} + x_3^2 \cdot \mathfrak{S}_{54631278} + x_3 \cdot \mathfrak{S}_{64531278} + x_3 \cdot \mathfrak{S}_{56431278}. \end{aligned}$$

For instance, $c_{(4,2,5,3),w} := [x_1^4 x_2^2 x_3^5 x_4^3] \mathfrak{S}_w = 1$ is witnessed by

- the path $w \xrightarrow{x_4} \bullet \xrightarrow{x_3} \bullet \xrightarrow{x_3} u = 53641278$, and
- the semistandard tableau

$$T = \begin{array}{|c|c|c|c|} \hline 1 & 1 & 1 & 1 \\ \hline 2 & 2 & 3 & \\ \hline 3 & 3 & & \\ \hline 4 & 4 & & \\ \hline \end{array} \text{ of shape } \lambda(u), \text{ flagged by } \phi(u) = (1, 3, 4, 4).$$

Proposition 5.9 below formalizes a rule for $c_{\alpha,w}$ in terms of such pairs.




FIGURE 1. $\mathcal{T}(w)$ for $w = 53861247$ where the accessible boxes are marked with z and those boxes of the parent which moved are shaded gray.

5.3. **Proof of #P-ness.** The technical core of our proof of Theorem 1.5 is to show:

Theorem 5.6. *The problem of computing $c_{\alpha,w}$, given input α and $\text{code}(w)$, is in #P.*

Define X to be the set consisting of pairs (S, R) where:

- (X.1) $S = (s_1, \dots, s_h)$, $s_t \in [L] \cup \{(x_k, m_t) : k \in [L], m_t \in \mathbb{Z}_{>0}\}$ such that if $s_t = (x_k, m_t)$ then $s_{t+1} \neq (x_k, m_{t+1})$ for $t < h$, and
- (X.2) $R = (r_{ij})_{1 \leq i, j \leq L}$, where $r_{ij} \in \mathbb{Z}_{\geq 0}$.

Fix $w \in S_\infty$ and a vexillary permutation $v \in S_\infty$. A (w, v) -transition string is a sequence $S = (s_1, \dots, s_h)$ satisfying (X.1) such that if we interpret i as $\bullet \xrightarrow{i} \bullet$ and (x_k, m_t) as $\bullet \xrightarrow{x_k} \bullet \dots \bullet \xrightarrow{x_k} \bullet$ (m_t -times) then S describes a path from w to (a leaf labelled by) v in $\mathcal{T}(w)$. Let $\text{Trans}(w, v)$ be the set of such sequences.

The *deletion weight* of $S \in \text{Trans}(w, v)$ is

$$\text{delwt}(S) = \sum m_t \cdot \vec{e}_r,$$

where the summation is over $1 \leq t \leq h$ such that $s_t = (x_r, m_t) \in S$ for some $r \in [L]$ (depending on t). Here $\vec{e}_r \in \mathbb{Z}_{\geq 0}^L$ is the r -th standard basis vector and L is the length of $\text{code}(w) = (c_1, c_2, \dots, c_L)$.

Example 5.7. In Figure 1 we read the $(w = 53861247, v = 54631278)$ -transition string $S = (2, (x_3, 2))$ as the path $w \xrightarrow{2} \bullet \xrightarrow{x_3} \bullet \xrightarrow{x_3} v$. Here, $\text{delwt}(S) = (0, 0, 2, 0)$.

Suppose T is a tableau of shape $\lambda = (\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_L \geq 0)$, with entries in $[L]$ and weakly increasing along rows. Define

$$R(T) = (r_{ij})_{1 \leq i, j \leq L}$$

to be the $L \times L$ matrix where r_{ij} is the number of j 's in row i of T . $R(T)$ encodes T . As pointed out in (a preprint version of) [12], T might have exponentially many (in L) boxes, whereas $R(T)$ is a $O(L^2)$ description of T .

Example 5.8. If $\lambda = (4, 3, 1, 0, 0)$ and

$$T = \begin{array}{|c|c|c|c|} \hline 1 & 1 & 2 & 3 \\ \hline 2 & 4 & 5 & \\ \hline 4 & & & \\ \hline \end{array} \longleftrightarrow R(T) = \begin{pmatrix} 2 & 1 & 1 & 0 & 0 \\ 0 & 1 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix}$$

Let $X_{\alpha,w} = \{(S, R(T))\} \subseteq X$ such that the following hold:

- (X.1') $S \in \text{Trans}(w, v)$,
- (X.2') $T \in \text{SSYT}(\lambda(v), \phi(v))$, and
- (X.3') $\text{delwt}(S) + \text{content}(T) = \alpha$.

Proposition 5.9. $c_{\alpha,w} = \#X_{\alpha,w}$.

Proof. Iterating (23),

$$\mathfrak{S}_w = \sum_{\text{vexillary } v \in S_\infty} \sum_{S \in \text{Trans}(w,v)} x^{\text{delwt}(S)} \mathfrak{S}_v.$$

Hence

$$(24) \quad c_{\alpha, w} = \sum_{\text{vexillary } v \in S_\infty} \sum_{S \in \text{Trans}(w, v)} [x^\alpha] x^{\text{delwt}(S)} \mathfrak{S}_v.$$

The result then follows from by (21), (22), and (24) combined. \square

Proposition 5.10 (cf. [8]). *Let $\text{code}(w) = (c_1, \dots, c_L)$. Suppose $D(w')$ is obtained from $D(w)$ using move (T.1) and $D(w'')$ is obtained from $D(w)$ with move (T.2) for a pivot in row i . There is an $O(L^2)$ -time algorithm to compute*

- (I) $\text{code}(w') = (c_1, \dots, c_{r-1}, c_r - 1, c_{r+1}, \dots, c_L)$ and
- (II) $\text{code}(w'') = (c_1, \dots, c_{i-1}, c_i + b, c_{i+1}, \dots, c_{r-1}, c_r - b, c_{r+1}, \dots, c_L)$, for some $b \in \mathbb{Z}_{>0}$.

Proof. By Proposition 3.2, determine $\mathbf{z}_w := (r, c)$ in $O(L^2)$ -time.

For (I), $D(w')$ is obtained from $D(w)$ by deleting \mathbf{z}_w ; so the expression in (I) is clear.

For (II), using Proposition 3.1, we can find $\mathbf{x} = (i, w(i))$, in $O(L^2)$ -time; this is our (T.2) pivot. Notice that row r of $D(w) \cap \mathcal{R}$ is nonempty (it contains $\mathbf{z}_w = (r, c)$); let b be the number of boxes in this row. It is straightforward from the graphical description of \mathcal{R} in terms of Rothe diagrams that each row of $D(w) \cap \mathcal{R}$ either has zero boxes or $b > 0$ boxes. Moreover, the d -th box (say, from the left) of each row are in the same column.

Suppose $j_1, \dots, j_m \in [i + 1, r]$ index the rows where $D(w) \cap \mathcal{R} \neq \emptyset$ (and thus has b boxes). (T.2) moves the b boxes of j_1 to row i and moves the b boxes of j_q to row j_{q-1} for $q = 2, \dots, m$. As explained above $j_m = r$, so (T.2) moves no boxes into row r . Thus row r of $D(w'') \cap \mathcal{R}$ has zero boxes.

It remains to compute b in $O(L^2)$ -time. Using Proposition 3.1 compute, in $O(L^2)$ -time,

$$m := \#\{h < r : w(h) < w(i)\}.$$

Clearly $b = c_r - [(w(i) - 1) - m]$. \square

Let $s_t = (x_r, m_t)$, as in (X.1), be a valid (multi)-deletion move on $u \in \mathcal{T}(w)$. Let $u^{(m_t)} \in \mathcal{T}(w)$ be defined by $u \xrightarrow{x_k} \bullet \dots \bullet \xrightarrow{x_k} u^{(m_t)}$ (m_t -times).

Proposition 5.11. *Suppose $u \in \mathcal{T}(w)$ where $\text{code}(u) = (\tilde{c}_1, \dots, \tilde{c}_{L'})$. Let $s_t = (x_k, m_t)$ or $s_t = i$ be as in (X.1). Given input $\text{code}(u)$ and s_t , there is an $O(L^2)$ algorithm to respectively determine if $u \xrightarrow{x_k} \bullet \dots \bullet \xrightarrow{x_k} u^{(m_t)}$ (m_t -times) or $u \xrightarrow{i} u''$ occurs in $\mathcal{T}(w)$ and (if yes) to compute*

- $\text{code}(u^{(m_t)})$ in the case $s_t = (x_k, m_t)$ (a multi-deletion move (T.1)), or
- $\text{code}(u'')$ in the case $s_t = i$ (a march move (T.2)).

Proof. By Proposition 5.10, $L' \leq L$. Thus in our run-time analysis, we replace L' by L .

Proposition 3.2 finds $\mathbf{z}_u := (r, c)$ (or determines it does not exist) in $O(L^2)$ -time. If \mathbf{z}_u does not exist then u is dominant and thus vexillary; output s_t is invalid. Thus we assume henceforth that \mathbf{z}_u exists.

Case 1: ($s_t = (x_k, m_t)$.) Proposition 3.1 finds $u(1), \dots, u(L')$ in $O(L^2)$ -time. Determine (taking $O(L^2)$ time) if

$$(25) \quad c_r - \left(\left(\min_{i \in [r]} u(i) \right) - 1 \right) \geq m_t,$$

holds. We claim that s_t is valid if and only if (25) holds and $k = r$. Indeed, observe

$$(26) \quad \#\{\text{boxes in row } r \text{ of } \text{Dom}(u)\} = \left(\min_{i \in [r]} u(i) \right) - 1.$$

Thus, (25) is equivalent to the existence of m_t boxes in row r of $D(u) \setminus \text{Dom}(u)$. By (T.1), if $k = r$ this is equivalent to being able to apply $\bullet \xrightarrow{x_r} \bullet$ successively m_t -times.

Finally, if s_t is valid, by m_t applications of Proposition 5.10 (I),

$$(27) \quad \text{code}(u^{(m_t)}) = (\tilde{c}_1, \dots, \tilde{c}_{r-1}, \tilde{c}_r - m_t, \tilde{c}_{r+1}, \dots, \tilde{c}_{L'}).$$

Hence we can output (27) in $O(L^2)$ -time.

Case 2: ($s_t = i$.) By Proposition 3.1, determine $u(1), \dots, u(L')$ from $\text{code}(u)$ in $O(L^2)$ -time. In particular this computes $\mathbf{x} := (i, u(i))$ in $O(L^2)$ -time. To decide if s_t is valid we must determine if $\mathbf{x} \in \text{Piv}(\mathbf{z}_u)$. To do this, first calculate (in $O(L)$ -time)

$$u_{NW}(\mathbf{z}_u) := \{(j, u(j)) : j < r, u(j) < c\}.$$

By definition,

$$\text{Piv}(\mathbf{z}_u) = \{(j, u(j)) \in u_{NW}(\mathbf{z}_u) : \nexists (h, u(h)) \in u_{NW}(\mathbf{z}_u) \text{ with } h > j, u(h) > u(j)\}.$$

$\text{Piv}(\mathbf{z}_u)$ takes $O(L)$ -time to compute since $\#u_{NW}(\mathbf{z}_u) \leq r - 1 \leq L - 1$. Hence we check if $\mathbf{x} \in \text{Piv}(\mathbf{z}_u)$ in $O(L)$ -time. If this is false, we output a rejection. Otherwise, Proposition 5.10 outputs $\text{code}(u'')$ in $O(L^2)$ -time. \square

Proposition 5.12. *If $S = (s_1, \dots, s_h) \in \text{Trans}(w, v)$ then $h \leq L^2$.*

Proof. Let $w := w_0 \xrightarrow{s_1} w_1 \xrightarrow{s_2} \dots \xrightarrow{s_h} w_h = v$ be the path in $\mathcal{T}(w)$ associated to S . By (T.1) and (T.2), $\mathbf{z}_{w_{t+1}}$ is weakly northwest of \mathbf{z}_{w_t} . Hence, for any fixed r , those $t \in [0, h - 1]$ with \mathbf{z}_{w_t} in row r form an interval $I^{(r)} \subseteq [0, h - 1]$. Since $1 \leq r \leq L$, it suffices to prove

$$(28) \quad \#I^{(r)} \leq 2(r - 1).$$

By (X.1) the transition moves acting on row r alternate between multi-(T.1) moves (x_r, m_t) and (T.2) moves. Thus to show (28), it is enough to prove

$$(29) \quad \#\{t \in I^{(r)} : w_{t-1} \rightarrow w_t \text{ is a (T.2) move}\} \leq r - 1.$$

Consider a march move i with $\mathbf{z}_{w_{t-1}} = (r, c)$ and $\mathbf{x} = (i, w_{t-1}(i)) \in \text{Piv}(\mathbf{z}_{w_{t-1}})$. By (T.2), if $(r, c') \in D(w_{t-1})$ is in the same connected component as $\mathbf{z}_{w_{t-1}}$, the move i takes (r, c') strictly north of row r . Thus, each march move strictly reduces the number of components in row r . Let $t_0 = \min\{t \in I^{(r)}\}$. Since there are at most r \bullet 's weakly above row r , $D(w_{t_0})$ has at most $r - 1$ (non-dominant) components in row r . Hence (29) holds, as desired. \square

Proposition 5.13. *Let v be vexillary with $\text{code}(v) = (c_1, \dots, c_{L'})$ and $L' \leq L$. There exists an $O(L^2)$ -time algorithm to check if $R = (r_{ij})_{1 \leq i, j \leq L'}$ is $R = R(T)$ for some $T \in \text{SSYT}(\lambda(v), \phi(v))$.*

Proof. Since $L' \leq L$, it is $O(L^2)$ -time to calculate $\phi(v), \lambda(v)$. Let

$$\lambda_i := \sum_{j=1}^{L'} r_{ij}, \text{ for } 1 \leq i \leq L'.$$

First verify (in $O(L)$ -time) that $\lambda_i \geq \lambda_{i+1}$ for $1 \leq i \leq L' - 1$. Then $R = R(T)$ where T is the (unique) row weakly increasing tableau of shape λ with r_{ij} many j 's in row i .

To verify $T \in \text{SSYT}(\lambda(v), \phi(v))$ we must check that it is (i) is flagged by $\phi(v)$, (ii) has shape $\lambda(v)$, and (iii) is semistandard. For (i), we need

$$(30) \quad r_{ij} = 0 \text{ if } j > \phi(v)_i, \text{ for all } i, j \in [L'].$$

For (ii), we need

$$(31) \quad \lambda_i = \lambda(v)_i \text{ for each } i \in [L'].$$

For (iii), it remains to ensure that T is column strict, i.e.,

$$(32) \quad \sum_{j' \leq j} r_{i+1, j'} \leq \sum_{j' < j} r_{i, j'} \text{ for each } i \in [L' - 1], j \in [L'].$$

We found the inequalities (31) and (32) from a (preprint) version of [12]. The inequalities (30), (31), and (32) can be checked in $O(L^2)$ -time since $i, j \in [L'] \subseteq [L]$. \square

The following completes our proof that we can check that $(S, R) \in X_{\alpha, w}$ in $L^{O(1)}$ -time.

Proposition 5.14. *Given $(S, R) \in X$ and $(\text{code}(w), \alpha)$, one can determine if $(S, R) \in X_{\alpha, w}$ in $L^{O(1)}$ -time.*

Proof. By Propositions 5.11 and 5.12 combined, one determines in $O(L^4)$ -time if S encodes a path $w := w_0 \xrightarrow{s_1} w_1 \xrightarrow{s_2} \dots \xrightarrow{s_h} w_h = v$ in $\mathcal{T}(w)$. If so, the length of $\text{code}(v)$ is at most L . Thus, using Theorem 5.1, one checks v is vexillary in $O(L^3)$ -time. This decides if S satisfies (X.1'). Proposition 5.13 checks R satisfies (X.2') in $O(L^2)$ -time. Finally since $h \leq L^2$, computing $\text{delwt}(S)$ takes $O(L^2)$ -time. Hence (X.3') is checkable in $O(L^2)$ time. \square

Proof of Theorem 5.6: By Proposition 5.9, $\#X_{\alpha, w} = c_{\alpha, w}$. By Proposition 5.12, $(S, R) \in \#X_{\alpha, w}$ only if the list S has at most L^2 elements. Assuming this, we check (S, R) satisfies (X.1) and (X.2) in $O(L^2)$ -time. Using Proposition 5.14, we can verify $(S, R) \in X_{\alpha, w}$ in $L^{O(1)}$ -time. Thus, given input α and $\text{code}(w)$, computing $c_{\alpha, w}$ is in $\#\text{P}$. \square

5.4. Hardness, and the conclusion of the proof of Theorem 1.5. *Schur polynomials* are an important basis of the vector space of symmetric polynomials. The Schur polynomial $s_\lambda = a_{\lambda+\delta}/a_\delta$ where $\lambda = (\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_n \geq 0)$, $a_\gamma := \det(x_i^{\gamma_j})_{i,j=1}^n$, and $\delta = (n-1, n-2, \dots, 2, 1, 0)$. The flagged Schur function of Section 5.1 is a generalization of the Schur polynomial.

A permutation w is *grassmannian* if it has at most one *descent* i , i.e., where $w(i) > w(i+1)$. Given a partition $(\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_L \geq 0)$ define a grassmannian permutation w_λ by setting

$$w_\lambda(i) = i + \lambda_{L-i+1} \text{ for } 1 \leq i \leq L.$$

For w_λ grassmannian, it is well-known (see, e.g., [10]) that

$$(33) \quad \text{code}(w_\lambda) = (\lambda_L, \lambda_{L-1}, \dots, \lambda_1).$$

Moreover,

$$(34) \quad \mathfrak{S}_{w_\lambda} = s_\lambda(x_1, \dots, x_L) = \sum_{\alpha \in \mathbb{Z}_{\geq 0}^L} K_{\lambda, \alpha} x^\alpha,$$

where $K_{\lambda, \alpha}$ is the *Kostka coefficient*. This number counts semistandard tableaux of shape λ with content α .

By (34),

$$(35) \quad c_{\alpha, w_\lambda} = K_{\lambda, \alpha}.$$

By Theorem 5.6, counting $c_{\alpha, w}$ is in #P. Suppose there is an oracle to compute $c_{\alpha, w}$ in polynomial time in the input length of $(\text{code}(w), \alpha)$. This input length is the same as for the input λ, α for $K_{\lambda, \alpha}$. Hence (33) and (35) combined imply a polynomial-time counting reduction from $\{c_{\alpha, w}\}$ to Kostka coefficients. Now H. Narayanan [12] proved that counting $K_{\lambda, \alpha}$ is a #P-complete problem. Thus counting $c_{\alpha, w}$ is a #P-complete problem. \square

Remark 5.15. Suppose the input for counting $c_{\alpha, w}$ is (α, w) where $w \in S_n$ (in one-line notation). Then the above counting reduction is not polynomial time in the input length of the Kostka problem. For example, suppose $\lambda = \alpha = (2^L, 2^L, \dots, 2^L)$ (L -many). Then the input length of this instance of the Kostka problem is $2L^2 \in O(L^2)$. On the other hand, $w_\lambda \in S_{L+2^L}$. Therefore, a polynomial time algorithm for the Schubert coefficient problem in n would have $\Omega(2^L)$ run time for the Kostka problem.

It seems unlikely that there is a polynomial-time reduction under this input assumption. This is our justification to encode w via $\text{code}(w)$ rather than one line notation. \square

ACKNOWLEDGMENTS

AY was supported by an NSF grant, a Simons Collaboration grant and a UIUC Campus Research Board grant. CR was supported by the National Science Foundation Graduate Research Fellowship Program under Grant No. DGE – 1746047. We acknowledge Mathoverflow, S. Kintali’s blog “My brain is open” and R. O’Donnell’s Youtube videos (from his class at Carnegie Mellon) for background. We thank Philipp Hieronymi, Alexandr Kostochka, Cara Monical, Erik Walsberg, Douglas West, Alexander Woo, for helpful comments and conversations. We also thank the anonymous referee for their insightful suggestions that improved the clarity of this paper.

REFERENCES

- [1] A. Adve, C. Robichaux, and A. Yong, *Computational complexity, Newton polytopes, and Schubert polynomials*, Proceedings of the 31st Conference on Formal Power Series and Algebraic Combinatorics (Ljubljana), Sémin. Lothar. Combin. 82B (2020), Art. 52, 12 pp.
- [2] ———, *Complexity, combinatorial positivity, and Newton polytopes*, preprint, 2018. arXiv:1810.10361v1
- [3] S. Billey, W. Jockusch and R. P. Stanley, *Some combinatorial properties of Schubert polynomials*, J. Algebraic Combin. 2(1993), no. 4, 345–374.
- [4] A. Fink, K. Mészáros, and A. St. Dizier, *Schubert polynomials as integer point transforms of generalized permutahedra*, Adv. Math. 332 (2018), 465–475.
- [5] S. Fomin, C. Greene, V. Reiner, and M. Shimozono, *Balanced labellings and Schubert polynomials*, European J. Combin. 18 (1997), no. 4, 373–389.
- [6] W. Fulton, *Young tableaux. With applications to representation theory and geometry*. London Mathematical Society Student Texts, 35. Cambridge University Press, Cambridge, 1997.
- [7] A. Knutson and A. Yong, *A formula for K -theory truncation Schubert calculus*, Int. Math. Res. Not. 2004, no. 70, 3741–3756.
- [8] A. Lascoux and M. -P. Schützenberger, *Schubert polynomials and the Littlewood-Richardson rule*, Letters in Math. Physics 10 (1985), 111–124.
- [9] ———, *Polynômes de Schubert*, C. R. Acad. Sci. Paris Sér. I Math. 294 (1982), 447–450.
- [10] L. Manivel, *Symmetric functions, Schubert polynomials and degeneracy loci*. Translated from the 1998 French original by John R. Swallow. SMF/AMS Texts and Monographs, American Mathematical Society, Providence, 2001.

- [11] C. Monical, N. Tokcan and A. Yong, *Newton polytopes in algebraic combinatorics*, Sel. Math. 25(5) (2019), no. 66, 37 pp.
- [12] H. Narayanan, *On the complexity of computing Kostka numbers and Littlewood-Richardson coefficients*, J. Alg. Comb., Vol. 24, N. 3, 2006, 347–354.
- [13] C. H. Papadimitriou and K. Steiglitz, Kenneth, *Combinatorial optimization: algorithms and complexity*. Corrected reprint of the 1982 original. Dover Publications, Inc., Mineola, NY, 1998. xvi+496 pp.
- [14] A. Schrijver, *Theory of Linear and Integer Programming*, John Wiley & sons, 1998.
- [15] R. P. Stanley, *Some Schubert shenanigans*, preprint, 2017. arXiv:1704.00851
- [16] L. G. Valiant, *The complexity of computing the permanent*, Theoret. Comput. Sci., 8(2):189–201, 1979.
- [17] A. Weigandt, *Schubert polynomials, 132-patterns, and Stanley’s conjecture*, Algebr. Comb. 1 (2018), no. 4, 415–423.

UCLA, LOS ANGELES, CA 90095

Email address: aadve@g.ucla.edu

DEPT. OF MATHEMATICS, U. ILLINOIS AT URBANA-CHAMPAIGN, URBANA, IL 61801, USA

Email address: cer2@illinois.edu, ayong@illinois.edu