

A PROBLEM ON CONCATENATED INTEGERS

Josep M. Brunat

Retired from the Departament de Matemàtiques, Universitat Politècnica de Catalunya, Barcelona.

Josep.M.Brunat@upc.edu

Joan-C. Lario

Department de Matemàtiques, Universitat Politècnica de Catalunya, Barcelona.

Joan.Carles.Lario@upc.edu

Abstract

Motivated by a WhatsApp message, we find out the integers $x > y \geq 1$ such that $(x+1)/(y+1) = (x \circ (y+1))/(y \circ (x+1))$, where \circ means the concatenation of the strings of two natural numbers (for instance $783 \circ 56 = 78356$). The discussion involves the equation $x(x+1) = 10y(y+1)$, a slight variation of Pell's equation related to the arithmetic of the Dedekind ring $\mathbb{Z}[\sqrt{10}]$. We obtain the infinite sequence $\mathcal{S} = \{(x_n, y_n)\}_{n \geq 1}$ of all the solutions of the equation $x(x+1) = 10y(y+1)$, which turn out to have limit $1/\sqrt{10}$. The solutions of the initial problem on concatenated integers form the infinite subsequence of \mathcal{S} formed by the pairs (x_n, y_n) such that x_n has one more digit than y_n .

1. Introduction

A friend sent to us a WhatsApp message with the content of Figure 1; no words at all, only a sequence of numbers and mathematical symbols.

Letting \circ denote the action of concatenate the strings of two natural numbers, for instance $783 \circ 56 = 78356$, the first equalities in every row are of the form

$$\frac{x!(y+1)!}{y!(x+1)!} = \frac{x \circ (y+1)}{y \circ (x+1)}$$

with $x \geq y$. Simplifying the factorials, the above equality is equivalent to

$$\frac{y+1}{x+1} = \frac{x \circ (y+1)}{y \circ (x+1)}. \quad (1)$$

The real numbers at the end of each row of the message along with the last row suggest that these fractions are decreasing and tend to the limit $1/\sqrt{10}$ as x, y grow.

Our aim here is to solve the following two questions:




Figure 1: The motivating WhatsApp message.

- 1) Find out all pairs of integers (x, y) with $x > y \geq 1$ verifying (1).
- 2) Show that there exist infinitely many solutions and that the fractions $(y + 1)/(x + 1)$, with $x > y \geq 1$ solutions of (1), are decreasing as x, y grow and that its limit exists and equals $1/\sqrt{10}$.

Observe that the case $y = 0$ is not interesting, since then $x \circ (y + 1) = 10x + 1$ and the equality (1) reads

$$\frac{1}{x + 1} = \frac{10x + 1}{x + 1},$$

that is fulfilled only when $x = 0$. The case $x = y$ does not have any interest either, since the equality (1) is a tautology and the quotients $(y + 1)/(x + 1) = (x + 1)/(x + 1) = 1$ do not have limit $1/\sqrt{10}$. Therefore, we shall restrict ourselves to the cases $x > y \geq 1$.

Formally, the concatenation of the strings of two positive integers runs as follows.

Let a and b be two positive integers, written with base-10 number system:

$$\begin{aligned} a &= a_r 10^r + a_{r-1} 10^{r-1} + \cdots + a_1 10 + a_0 \quad (a_r \neq 0), \\ b &= b_s 10^s + b_{s-1} 10^{s-1} + \cdots + b_1 10 + b_0, \quad (b_s \neq 0). \end{aligned}$$

Then, one has

$$\begin{aligned} a \circ b &= 10^{s+1} a + b \\ &= a_r 10^{r+s+1} + \cdots + a_0 10^{s+1} + b_s 10^s + b_{s-1} 10^{s-1} + \cdots + b_1 10 + b_0. \end{aligned}$$

Hereafter, as usual, for a positive real number x , the functions $\lfloor x \rfloor$, $\lceil x \rceil$, $\lceil x \rceil$ denote the floor, ceiling, and round part of x . For future reference, we introduce the function $\delta(x) = \lfloor \log x \rfloor$. Note that $\delta(x) + 1$ counts the number of digits of the integer $\lfloor x \rfloor$ with the 10-base system.

The next lemma transforms our WhatsApp question it into a diophantine equation plus a condition on the digits function δ .

Lemma 1. *Let $x > y \geq 1$ be integers. Then, one has*

$$\frac{y+1}{x+1} = \frac{x \circ (y+1)}{y \circ (x+1)} \quad (2)$$

if, and only if, it holds $x(x+1) = 10y(y+1)$ and $\delta(x+1) = \delta(y+1) + 1$.

Proof. Let denote $r = \delta(x+1)$, and $s = \delta(y+1)$. Assume that

$$\frac{y+1}{x+1} = \frac{x \circ (y+1)}{y \circ (x+1)} = \frac{10^{s+1}x + y + 1}{10^{r+1}y + x + 1}.$$

By cross-multiplying and simplifying, one gets

$$x(x+1) = 10^{r-s}y(y+1).$$

Since $x > y$, one has $r - s \geq 1$. Let $v = \lfloor \log(x(x+1)) \rfloor = \lfloor \log(10^{r-s}y(y+1)) \rfloor$. We have $v \in \{2r-1, 2r, 2r+1\}$. The value $v = 2r-1$ occurs only when $x+1 = 10^r$. In this case, we have $x(x+1) = (10^r-1)10^r = 10^{r-s}y(y+1)$ so that

$$(10^r-1)10^s = y(y+1).$$

Since $10^s - 1 \leq y < 10^{s+1} - 1$, we have

$$(10^r-1)10^s = y(y+1) < (10^{s+1}-1)10^{s+1}$$

and $10^r - 1 < (10^{s+1} - 1)10$. Since $r - s \geq 1$, we must have $r = s + 1$. Hence, we have

$$\begin{aligned} v &= \lfloor \log(x(x+1)) \rfloor \in \{2r, 2r+1\}, \\ v &= \lfloor \log(10^{r-s}y(y+1)) \rfloor \in \{r+s-1, r+s, r+s+1\}, \end{aligned}$$

and $2r \leq v \leq r + s + 1$. It follows $r - s \leq 1$. Summarizing, we must have $10y(y + 1) = x(x + 1)$ and $r = s + 1$ as wanted.

Reciprocally, it is easy to check that if $10y(y + 1) = x(x + 1)$ and $r = s + 1$, then equation (2) holds. \square

Thus, our first task is to determine the set of solutions:

$$\mathcal{S} = \{(x, y) \in \mathbb{Z}^2 : x > y \geq 1, x(x + 1) = 10y(y + 1)\},$$

and then select the subset of those that concatenate well with regard to our What-App challenge:

$$\mathcal{C} = \{(x, y) \in \mathcal{S} : \delta(x + 1) = \delta(y + 1) + 1\}.$$

2. The associate Pell's equation

In this section, we deal with the set \mathcal{S} of positive solutions of the diophantine equation

$$x(x + 1) = 10y(y + 1).$$

Multiplying by 4 and completing the square, the equation is equivalent to

$$(2x + 1)^2 - 10(2y + 1)^2 = -9.$$

Letting $a = 2x + 1$ and $b = 2y + 1$, the above equation reads

$$a^2 - 10b^2 = -9. \tag{3}$$

Therefore, we need to find all odd positive integer solutions of (3).

To this end, we shall use the arithmetic properties of the ring $\mathbb{Z}[\sqrt{10}]$. It is a Dedekind domain; that is, every non-zero proper ideal factors into a product of prime ideals, and such a factorization is unique up to the order of the factors. We shall use some standard terminology and basic properties: the conjugate of $\alpha = a + b\sqrt{10} \in \mathbb{Z}[\sqrt{10}]$ is $\bar{\alpha} = a - b\sqrt{10}$; the norm of α is $N(\alpha) = \alpha\bar{\alpha} = a^2 - 10b^2$. The conjugation $\alpha \mapsto \bar{\alpha}$ is an automorphism of the ring $\mathbb{Z}[\sqrt{10}]$ and the norm map is multiplicative; that is, $N(\alpha\beta) = N(\alpha)N(\beta)$ for all $\alpha, \beta \in \mathbb{Z}[\sqrt{10}]$. An element $u \in \mathbb{Z}[\sqrt{10}]$ is a unit if, and only if, $|N(u)| = 1$. Note that the units of $\mathbb{Z}[\sqrt{10}]$ are just the numbers $a + b\sqrt{10}$ where (a, b) are the solutions of the Pell equations $a^2 - 10b^2 = \pm 1$. If $u \in \mathbb{Z}[\sqrt{10}]$ is a unit, then $u^{-1} = \bar{u}$. Dirichlet's unit theorem describes the structure of the group of units; in our case, $\varepsilon = 3 + \sqrt{10}$ is the fundamental unit and, if $\langle \varepsilon \rangle$ is the multiplicative group generated by ε , the group of units is $\mathbb{Z}[\sqrt{10}]^* = \{\pm 1\} \times \langle \varepsilon \rangle$. Notice that $N(\varepsilon) = -1$. We refer the reader to [1, 2, 3] for further details.

The diophantine equation (3) in \mathbb{Z} , gets transformed into the norm equation

$$N(a + b\sqrt{10}) = a^2 - 10b^2 = -9$$

in the ring $\mathbb{Z}[\sqrt{10}]$. Since the norm is multiplicative, two elements of $\mathbb{Z}[\sqrt{10}]$ with equal norm differ (multiplicatively) by a unit of norm 1; that is, by an even power of ε . Thus, all we need is to find the elements in $\mathbb{Z}[\sqrt{10}]$ of norm -9 .

We shall denote by $\langle \alpha, \beta \rangle$ the ideal of $\mathbb{Z}[\sqrt{10}]$ generated by α and β . The class number of $\mathbb{Z}[\sqrt{10}]$ is two, so that the square of every ideal is a principal ideal. Recall also that the norm of an ideal \mathfrak{a} is the cardinal of the quotient ring $\mathbb{Z}[\sqrt{10}]/\mathfrak{a}$, and the norm of ideals is multiplicative. Moreover, the norm of a principal ideal agrees with the absolute value of the norm of a generator of the ideal. We consider the ideals

$$\mathfrak{P}_1 = \langle 3, 1 + \sqrt{10} \rangle, \quad \mathfrak{P}_2 = \langle 3, 1 - \sqrt{10} \rangle.$$

Lemma 2. *With the above notations, it holds:*

- (i) $\langle 3 \rangle = \mathfrak{P}_1 \cdot \mathfrak{P}_2$;
- (ii) \mathfrak{P}_1 and \mathfrak{P}_2 are prime ideals;
- (iii) $\mathfrak{P}_1^2 = \langle 1 + \sqrt{10} \rangle$ and $\mathfrak{P}_2^2 = \langle 1 - \sqrt{10} \rangle$.
- (iv) Moreover, $\langle a + b\sqrt{10} \rangle \cdot \langle a - b\sqrt{10} \rangle = \langle 9 \rangle$ if and only if, for some unit $u \in \mathbb{Z}[10]^*$, one has

$$a + b\sqrt{10} \in \{3u, (1 - \sqrt{10})u, (1 + \sqrt{10})u\}.$$

Proof. (i) On the one hand, we have

$$\mathfrak{P}_1 \mathfrak{P}_2 = \langle 9, 3(1 + \sqrt{10}), 3(1 - \sqrt{10}), -9 \rangle \subseteq \langle 3 \rangle.$$

On the other hand, $3 = 9 - 3(1 + \sqrt{10}) - 3(1 - \sqrt{10}) \in \mathfrak{P}_1 \mathfrak{P}_2$. Hence, we obtain the equality $\langle 3 \rangle = \mathfrak{P}_1 \mathfrak{P}_2$.

(ii) A routine checking shows that the map $f: \mathbb{Z}[\sqrt{10}] \rightarrow \mathbb{Z}/3\mathbb{Z}$ defined by $f(a + b\sqrt{10}) = [a - b]$ is a surjective ring homomorphism. Clearly $f(3) = [3] = [0]$ and $f(1 + \sqrt{10}) = [0]$. Thus, $\mathfrak{P}_1 \subseteq \text{Ker } f$. Also, if $a + b\sqrt{10} \in \text{Ker } f$, then $[a - b] = [0]$. That implies $a = b + 3c$ for some integer c and then

$$a + b\sqrt{10} = b + 3c + b\sqrt{10} = 3c + b(1 + \sqrt{10}) \in \mathfrak{P}_1.$$

Therefore, $\text{Ker } f = \mathfrak{P}_1$. Since $\mathbb{Z}[\sqrt{10}]/\mathfrak{P}_1 \simeq \mathbb{Z}/3\mathbb{Z}$ is a field, it follows that \mathfrak{P}_1 is a maximal ideal and, in particular, it is a prime ideal. Mutatis mutandis we see that \mathfrak{P}_2 is a prime ideal.

(iii) We have

$$\mathfrak{P}_1^2 = \langle 3, 1 + \sqrt{10} \rangle^2 = \langle 3^2, 3(1 + \sqrt{10}), (1 + \sqrt{10})^2 \rangle \subseteq \langle 1 + \sqrt{10} \rangle,$$

and $1 + \sqrt{10} = (-2) \cdot 9 - 3(1 + \sqrt{10}) + 2(1 + \sqrt{10})^2 \in \mathfrak{P}_1^2$. Thus, $\mathfrak{P}_1^2 = \langle 1 + \sqrt{10} \rangle$. Analogously, for the conjugate ideal we get $\mathfrak{P}_2^2 = \langle 1 - \sqrt{10} \rangle$.

(iv) From the equalities

$$\langle a + b\sqrt{10} \rangle \cdot \langle a - b\sqrt{10} \rangle = \langle 9 \rangle = \langle 3 \rangle \cdot \langle 3 \rangle = \mathfrak{P}_1^2 \mathfrak{P}_2^2,$$

by using that $\mathbb{Z}[\sqrt{10}]$ is a Dedekind domain and the fact that the norm is multiplicative on ideals, it follows that $\langle a + b\sqrt{10} \rangle$ must be one among of the ideals

$$\mathfrak{P}_1\mathfrak{P}_2, \quad \mathfrak{P}_1^2, \quad \mathfrak{P}_2^2.$$

Case $\langle a + b\sqrt{10} \rangle = \mathfrak{P}_1\mathfrak{P}_2 = \langle 3 \rangle$. Then, we get $a + b\sqrt{10} = 3u$ for some unit u .

Case $\langle a + b\sqrt{10} \rangle = \mathfrak{P}_1^2 = \langle 1 + \sqrt{10} \rangle$. Then, we get $a + b\sqrt{10} = (1 + \sqrt{10})u$ for some unit u .

Case $\langle a + b\sqrt{10} \rangle = \mathfrak{P}_2^2 = \langle 1 - \sqrt{10} \rangle$. Then, we get $a + b\sqrt{10} = (1 - \sqrt{10})u$ for some unit u . \square

Finally, we can exhibit all positive integers solutions of $a^2 - 10b^2 = -9$. For a quadratic number $\alpha = c + d\sqrt{10}$ in $\mathbb{Z}[\sqrt{10}]$, we shall say that (c, d) are the coordinates of α . Recall that $\varepsilon = 3 + \sqrt{10}$ and has norm -1 .

Proposition 1. *The positive integer solutions (a, b) of $a^2 - 10b^2 = -9$ are the coordinates of:*

- (i) $3\varepsilon^n$ for every odd positive integer n ;
- (ii) $-(1 - \sqrt{10})\varepsilon^n$, for every even positive integer n ;
- (iii) $(1 + \sqrt{10})\varepsilon^n$, for every even positive integer n .

Proof. First, we check that the quadratic numbers of type (i), (ii) or (iii) give positive solutions of $a^2 - 10b^2 = -9$.

(i) It is clear that $3\varepsilon^n$ has positive coordinates; since n is odd, $N(3\varepsilon^n) = 9 \cdot (-1) = -9$. (ii) Observe that $-(1 - \sqrt{10})\varepsilon^2 = 41 + 13\sqrt{10}$ has positive coordinates, and the successive multiplications by $\varepsilon^2 = 19 + 6\sqrt{10}$ also produce quadratic numbers with positive coordinates. Moreover, n being even, one has $N(-(1 - \sqrt{10})\varepsilon^n) = (-9) \cdot 1 = -9$. (iii) Certainly, $(1 + \sqrt{10})\varepsilon^n$ has positive coordinates for every positive integer n ; moreover, if n is even, then $N((1 + \sqrt{10})\varepsilon^n) = (-9) \cdot 1 = -9$.

Reciprocally, we claim that any positive solution of our equation must arise from one of these types (i), (ii) or (iii).

Let (a, b) denote a positive solution of $a^2 - 10b^2 = -9$. Then, $\langle a + b\sqrt{10} \rangle \cdot \langle a - b\sqrt{10} \rangle = \langle 9 \rangle$ and, due to Lemma 2 (iv), we must have

$$a + b\sqrt{10} \in \{3u, (1 - \sqrt{10})u, (1 + \sqrt{10})u\}$$

with u a unit of norm -1 , 1 , and 1 , respectively. The units of norm -1 are precisely $\pm\varepsilon^n$ for odd integers n . Since $\varepsilon = 3 + \sqrt{10} > 0$ satisfies $\varepsilon^{-1} = -\bar{\varepsilon} = -3 + \sqrt{10} > 0$, the units with positive coordinates of norm -1 are precisely ε^n for odd positive integers n . The claim follows. \square

We have obtained that the positive solutions of $a^2 - 10b^2 = -9$ are the coordinates of the quadratic numbers in the sequence $a_n + b_n\sqrt{10}$:

$$\begin{array}{lll} 3\varepsilon, & -(1 - \sqrt{10})\varepsilon^2, & (1 + \sqrt{10})\varepsilon^2, \\ 3\varepsilon^3, & -(1 - \sqrt{10})\varepsilon^4, & (1 + \sqrt{10})\varepsilon^4, \\ 3\varepsilon^5, & -(1 - \sqrt{10})\varepsilon^6, & (1 + \sqrt{10})\varepsilon^6, \\ \vdots & \vdots & \vdots \end{array}$$

With $\varphi = \varepsilon^2 = 19 + 6\sqrt{10}$, each row above is of the form

$$3\varepsilon\varphi^{n-1}, \quad -(1 - \sqrt{10})\varphi^n, \quad (1 + \sqrt{10})\varphi^n, \quad n \geq 1.$$

The $(n + 3)$ th term $a_{n+3} + b_{n+3}\sqrt{10}$ is obtained from the term $a_n + b_n\sqrt{10}$ by the formula

$$a_{n+3} + b_{n+3}\sqrt{10} = (a_n + b_n\sqrt{10})\varphi = (19a_n + 60b_n) + (6a_n + 19b_n)\sqrt{10},$$

with initial values

$$\begin{aligned} a_1 + b_1\sqrt{10} &= 3\varepsilon = 9 + 3\sqrt{10}, \\ a_2 + b_2\sqrt{10} &= -(1 - \sqrt{10})\varphi = 41 + 13\sqrt{10}, \\ a_3 + b_3\sqrt{10} &= (1 + \sqrt{10})\varphi = 79 + 25\sqrt{10}. \end{aligned}$$

Therefore, the positive integer solutions of (3) are the pairs (a_n, b_n) obtained by the recurrence

$$a_{n+3} = 19a_n + 60b_n, \quad b_{n+3} = 6a_n + 19b_n \quad (4)$$

with initial values

$$(a_1, b_1) = (9, 3), \quad (a_2, b_2) = (41, 13), \quad (a_3, b_3) = (79, 25).$$

Recall that we are interested only on the odd positive solutions (a_n, b_n) . Pleasantly, notice that the initial values are formed by odd integers and that if a_n and b_n are odd, then a_{n+3} and b_{n+3} are odd too. Hence a_n and b_n are odd for all n . Rewriting $a_n = 2x_n + 1$ and $b_n = 2y_n + 1$, we finally get the following result.

Proposition 2. *The set \mathcal{S} consists of couples (x_n, y_n) given by the recursion with initial values*

$$(x_1, y_1) = (4, 1), \quad (x_2, y_2) = (20, 6), \quad (x_3, y_3) = (39, 12),$$

and, for $n \geq 1$,

$$x_{n+3} = 19x_n + 60y_n + 39, \quad y_{n+3} = 6x_n + 19y_n + 12. \quad (5)$$

(4, 1),	(20, 6),
(39, 12),	(175, 55),
(779, 246),	(1500, 474),
(6664, 2107),	(29600, 9360),
(56979, 18018),	(253075, 80029),
(1124039, 355452),	(2163720, 684228),
(9610204, 3039013),	(42683900, 13497834),
(82164399, 25982664),	(364934695, 115402483),
(1620864179, 512562258),	(3120083460, 986657022),
(13857908224, 4382255359),	(61550154920, 19463867988),
(118481007099, 37466984190),	(526235577835, 166410301177),
(2337285022799, 739114421304),	(4499158186320, 1422758742216),
(19983094049524, 6319209189385),	(88755280711460, 28066884141582),

Table 1: The first terms of \mathcal{S} , in green those belonging in \mathcal{C}

We think of \mathcal{S} as a set or as a sequence indistinctly. Table 1 shows the first terms of \mathcal{S} . We display in green color the solutions in \mathcal{C} ; these are the solutions we are interested in; that is, those (x_n, y_n) satisfying $x_n + 1$ has one more digit than $y_n + 1$ which correspond to the solutions of our problem originated by the WhatsApp message.

The above recurrence allows to obtain a direct formula for x_n and y_n .

Proposition 3. For $k \in \{1, 2, 3\}$, let

$$A_k = \frac{1}{2} \left(\frac{x_k}{\sqrt{10}} + y_k + \frac{10 + \sqrt{10}}{20} \right).$$

Then, for $n \geq 1$,

$$x_{3n+k} = \lfloor A_k \sqrt{10} \varphi^n \rfloor, \quad y_{3n+k} = \lfloor A_k \varphi^n \rfloor.$$

Proof. For $n \geq 1$ and $k \in \{1, 2, 3\}$, we have

$$\begin{bmatrix} x_{3n+k} \\ y_{3n+k} \\ 1 \end{bmatrix} = \begin{bmatrix} 19 & 60 & 39 \\ 6 & 19 & 12 \\ 0 & 0 & 1 \end{bmatrix}^n \cdot \begin{bmatrix} x_k \\ y_k \\ 1 \end{bmatrix} = P \begin{bmatrix} \varphi^n & 0 & 0 \\ 0 & \varphi^{-n} & 0 \\ 0 & 0 & 1 \end{bmatrix} P^{-1} \cdot \begin{bmatrix} x_k \\ y_k \\ 1 \end{bmatrix},$$

where $P = \begin{bmatrix} \sqrt{10} & -\sqrt{10} & -1 \\ 1 & 1 & -1 \\ 0 & 0 & 2 \end{bmatrix}$ and $\varphi = \varepsilon^2 = 19 + 6\sqrt{10}$. Hence, if

$$B_k = \frac{1}{2} \left(\frac{-x_k}{\sqrt{10}} + y_k + \frac{10 - \sqrt{10}}{20} \right),$$

we have

$$\begin{aligned}x_{3n+k} &= A_k \sqrt{10} \varphi^n + B_k \sqrt{10} \varphi^{-n} - \frac{1}{2}, \\y_{3n+k} &= A_k \varphi^n + B_k \varphi^{-n} - \frac{1}{2}.\end{aligned}\tag{6}$$

From the fact that $|B_k \varphi^{-1}| < |B_k \sqrt{10} \varphi^{-1}| < 1/2$, and $\varphi^{-1} < 1$, we obtain

$$x_{3n+k} = \lfloor A_k \sqrt{10} \varphi^n - 1/2 \rfloor, \quad y_{3n+k} = \lfloor A_k \varphi^n - 1/2 \rfloor.$$

Now, for z a real number, $\lfloor z - 1/2 \rfloor = \lfloor z \rfloor$. Thus,

$$x_{3n+k} = \lfloor A_k \sqrt{10} \varphi^n \rfloor, \quad y_{3n+k} = \lfloor A_k \varphi^n \rfloor.$$

□

Proposition 4. For $(x_n, y_n) \in \mathcal{S}$, it holds

$$\lim_n \frac{y_n + 1}{x_n + 1} = \lim_n \frac{x_n}{y_n} = \frac{1}{\sqrt{10}}.$$

Moreover, y_n/x_n is strictly increasing and $(y_n + 1)/(x_n + 1)$ is strictly decreasing.

Proof. As $\varphi = 19 + 6\sqrt{10} > 1$, from the formulas (6) it is clear that, for each $k \in \{1, 2, 3\}$,

$$\lim_n \frac{y_{3n+k}}{x_{3n+k}} = \frac{1}{\sqrt{10}}.$$

Every pair $(x_n, y_n) \in \mathcal{S}$ is captured in one of the three sequences (x_{3n+k}, y_{3n+k}) , thus

$$\lim_n \frac{y_n}{x_n} = \frac{1}{\sqrt{10}}.$$

Moreover,

$$\lim_n \frac{y_n + 1}{x_n + 1} = \lim_n \frac{y_n/x_n + 1/x_n}{1 + 1/x_n} = \lim_n \frac{y_n}{x_n} = \frac{1}{\sqrt{10}}.$$

To prove that y_n/x_n is increasing, we use the recursive definition of (x_n, y_n) . For $n \geq 1$, the sign of $y_{n+1}/x_{n+1} - y_n/x_n$ is the sign of

$$\gamma_n = x_n y_{n+1} - x_{n+1} y_n.$$

To show that y_n/x_n is increasing it is sufficient to proof that $\gamma_n > 0$ for all n . Now, $\gamma_1 = 4$, $\gamma_2 = 6$ and $\gamma_3 = 45$. For $n \geq 4$,

$$\begin{aligned}\gamma_n &= (19x_{n-3} + 60y_{n-3} + 39)(6x_{n-2} + 19y_{n-2} + 12) \\ &\quad - (19x_{n-2} + 60y_{n-2} + 39)(6x_{n-3} + 19y_{n-3} + 12) \\ &= \gamma_{n-3} + 6(x_{n-2} - x_{n-3}) + 21(y_{n-2} - y_{n-3}).\end{aligned}$$

The sequences x_n and y_n are increasing, and by induction hypothesis, $\gamma_{n-3} > 0$. Therefore $\gamma_n > 0$ and the sequence y_n/x_n is strictly increasing.

Now, let $e_n = 1/\sqrt{10} - y_n/x_n$. Observe that $e_n - e_{n+1} > 0$ for all n , and

$$\frac{y_n + 1}{x_n + 1} - \frac{y_n}{x_n} = \frac{1}{x_n + 1} \left(1 - \frac{y_n}{x_n} \right) = \frac{1}{x_n + 1} \left(1 + e_n - \frac{1}{\sqrt{10}} \right).$$

Therefore,

$$\begin{aligned} \frac{y_n + 1}{x_n + 1} - \frac{y_{n+1} + 1}{x_{n+1} + 1} &= \frac{y_n}{x_n} + \frac{1}{x_n + 1} \left(1 + e_n - \frac{1}{\sqrt{10}} \right) \\ &\quad - \frac{y_{n+1}}{x_{n+1}} - \frac{1}{x_{n+1} + 1} \left(1 + e_{n+1} - \frac{1}{\sqrt{10}} \right) \\ &> \frac{y_n}{x_n} - \frac{y_{n+1}}{x_{n+1}} + \frac{1}{x_{n+1} + 1} (e_n - e_{n+1}) > 0. \end{aligned}$$

Hence, $(y_n + 1)/(x_n + 1)$ is strictly decreasing. \square

3. Infinite solutions

If the set \mathcal{C} is infinite, then the sequence $(y_n + 1)/(x_n + 1)$ with $(x_n, y_n) \in \mathcal{C}$ has the same limit as $(y_n + 1)/(x_n + 1)$ with $(x_n, y_n) \in \mathcal{S}$, which is $1/\sqrt{10}$. Thus, it only remains to prove that the set \mathcal{C} is infinite. To this end, we first show that if $(x, y) \in \mathcal{S}$, then neither $x + 1$ nor $y + 1$ is a power of 10.

Proposition 5. *If $(x, y) \in \mathcal{S}$, then $x + 1 \neq 10^\beta$ and $y + 1 \neq 10^\beta$ for all $\beta \geq 1$.*

Proof. The strategy of the proof is as follows. For an integer $m \geq 2$, we consider the sequence obtained by taking the terms of the sequence \mathcal{S} modulus m :

$$\mathcal{S}_m = \{(x_n \bmod m, y_n \bmod m) : \text{for } (x_n, y_n) \in \mathcal{S}\}.$$

Since \mathcal{S} is defined by a recurrence, the sequences \mathcal{S}_m are periodic for all m . We denote by $\ell(m)$ the length period of \mathcal{S}_m . Assuming that $x_n + 1$ or $y_n + 1$ are powers of 10, we choose appropriate m to derive a contradiction.

Assume $x_n + 1 = 10^\beta$. By inspection, we can (and do) assume that $n \geq 5$, which implies $x_n > 1000$ and $\beta \geq 3$. We take the modulus $m = 8$. Since $10^\beta \equiv 0 \pmod{8}$, we have $x_n = 10^\beta - 1 \equiv 7 \pmod{8}$. The condition $10y_n(y_n + 1) = x_n(x_n + 1)$, becomes $2y_n(y_n + 1) \equiv 6 \cdot 7 \equiv 2 \pmod{8}$, which has no solution. Therefore, $x_n + 1 \neq 10^\beta$.

Assume $y_n + 1 = 10^\beta$. First, we take the modulus $m = 9$. The first 12 terms of \mathcal{S}_9 are

$$\begin{aligned} &(4, 1), (2, 6), (3, 3), (4, 1), (5, 3), (6, 6), (4, 1), (8, 0), (0, 0), \\ &(4, 1), (2, 6), (3, 3), \end{aligned}$$

so that \mathcal{S}_9 has period $\ell(m) = 9$. We must have $y_n = 10^\beta - 1 \equiv 0 \pmod{9}$ and then $x_n(x_n + 1) = 10y_n(y_n + 1) \equiv 0 \pmod{9}$. This forces the two conditions: $x_n \equiv 0, 8 \pmod{9}$ and $n \equiv 0, 8 \pmod{9}$.

Now, we take $m = 10$. The 33 first terms of \mathcal{S}_{10} are

$$\begin{aligned} &(4, 1), (0, 6), (9, 2), (5, 5), (9, 6), (0, 4), (4, 7), (0, 0), (9, 8), (5, 9), \\ &(9, 2), (0, 8), (4, 3), (0, 4), (9, 4), (5, 3), (9, 8), (0, 2), (4, 9), (0, 8), \\ &(9, 0), (5, 7), (9, 4), (0, 6), (4, 5), (0, 2), (9, 6), (5, 1), (9, 0), (0, 0), \\ &(4, 1), (0, 6), (9, 2), \end{aligned}$$

so that \mathcal{S}_{10} has period $\ell(m) = 30$. Thus, we have $y_n = 10^\beta - 1 \equiv 9 \pmod{10}$, $x_n(x_n + 1) \equiv 0 \pmod{10}$, and $x_n \equiv 0, 4, 5, 9 \pmod{10}$. The pairs $(0, 9)$ and $(9, 9)$ are not in \mathcal{S}_{10} . If $(4, 9)$ is a n -term of \mathcal{S}_{10} then $n \equiv 19 \pmod{30}$, and if $(5, 9)$ is a n -term of \mathcal{S}_{10} then $n \equiv 10 \pmod{30}$.

Gluing the information mod $m = 9$ and mod $m = 10$, we see that none of the four systems

$$n \equiv g \pmod{9}, g \in \{0, 8\}, \quad n \equiv h \pmod{30}, h \in \{10, 19\}$$

has a solution because in the four cases $\gcd(9, 30) = 3$ does not divide $g - h$. Therefore, $y_n + 1 \neq 10^\beta$. \square

As a consequence of Proposition 5, we can precise the condition on digits of Lemma 1.

Corollary 1. *Let $(x, y) \in \mathcal{S}$. Then, one has:*

- (i) $\delta(x) = \delta(x + 1)$ and $\delta(y) = \delta(y + 1)$;
- (ii) $\delta(x) = \delta(y)$ or $\delta(x) = \delta(y) + 1$.

Moreover, $\mathcal{C} = \{(x, y) \in \mathcal{S} : \delta(x) = \delta(y) + 1\}$.

Proof. (i) If $\delta(x) = \beta$, then $10^\beta \leq x < 10^{\beta+1}$ and $x + 1 \leq 10^{\beta+1}$. By Proposition 5, we have $x + 1 \neq 10^{\beta+1}$. Therefore, $x + 1 < 10^{\beta+1}$ and we have $\delta(x + 1) = \beta = \delta(x)$. Analogously, $\delta(y + 1) = \delta(y)$.

(ii) As in the proof of Lemma 1, put $r = \delta(x + 1)$ and $s = \delta(y + 1)$. By (i), $\delta(x) = r$ and $\delta(y) = s$. Then, since $(x, y) \in \mathcal{S}$, then $\delta(x(x + 1)) \in \{2r, 2r + 1\}$ and $\delta(y(y + 1)) \in \{2s, 2s - 1\} = \{2r - 1, 2r\}$. It follows $r = s$ or $r = s + 1$, that is, $\delta(x) = \delta(y)$ or $\delta(x) = \delta(y) + 1$.

Finally, by (i) and Lemma 1, we have $\mathcal{C} = \{(x, y) \in \mathcal{S} : \delta(x) = \delta(y) + 1\}$. \square

Now, we can prove that the set \mathcal{C} is infinite. In fact, we shall prove that in each of the three sequences (x_{3n+k}, y_{3n+k}) there are no more than two consecutive terms which are not in \mathcal{C} . The next two remarks will be useful.

Remark The number $\varphi = \epsilon^2 = 19 + 6\sqrt{10}$ satisfies the conditions $37 < \varphi < 38$ and $1441 < \varphi^2 < 1442$. Thus, for each positive real number x , we have

$$\delta(x) + 1 \leq \delta(x\varphi) \leq \delta(x) + 2, \quad \delta(x) + 3 \leq \delta(x\varphi^2) \leq \delta(x) + 4. \quad (7)$$

Remark Due to Proposition 3, we know that $x_{3n+k} = \lfloor A_k \sqrt{10} \varphi^n \rfloor$. Let $\delta(x_{3n+k}) = \beta$. We have $A_k \sqrt{10} \varphi^n < 1 + x_{3n+k} \leq 10^{\beta+1}$. By Proposition 5, we have $x_{3n+k} + 1 \neq 10^{\beta+1}$. Therefore,

$$10^\beta \leq x_{3n+k} < A_k \sqrt{10} \varphi^n < x_{3n+k} + 1 < 10^{\beta+1}.$$

Thus, $\delta(x_{3n+k}) = \delta(A_k \sqrt{10} \varphi^n)$. Analogously, one gets $\delta(y_{3n+k}) = \delta(A_k \varphi^n)$.

Proposition 6. For $k \in \{1, 2, 3\}$, the sequence (x_{3n+k}, y_{3n+k}) has not three consecutive terms in $\mathcal{S} \setminus \mathcal{C}$. In particular, the set \mathcal{C} is infinite.

Proof. Consider one of the three sequences; that is, fix k . Rename the index of the terms and assume that we have consecutive elements such that $(x_n, y_n) \in \mathcal{C}$ and $(x_{n+1}, y_{n+1}), (x_{n+2}, y_{n+2}) \in \mathcal{S} \setminus \mathcal{C}$. We want to prove that $(x_{n+3}, y_{n+3}) \in \mathcal{C}$.

For every $m \geq 1$, denote $p_m = A_k \sqrt{10} \varphi^m$ and $q_m = A_k \varphi^m$. By using the last Remark, we have

$$\delta(p_n) = \delta(q_n) + 1, \quad \delta(p_{n+1}) = \delta(q_{n+1}), \quad \delta(p_{n+2}) = \delta(q_{n+2}),$$

and we want to prove $\delta(p_{n+3}) = \delta(q_{n+3}) + 1$. Let $\beta = \delta(p_n)$. Then, $\delta(q_n) = \beta - 1$. By (7), one has

$$\delta(p_{n+1}) = \delta(p_n \varphi) \in \{\beta + 1, \beta + 2\}, \quad \delta(q_{n+1}) = \delta(q_n \varphi) \in \{\beta, \beta + 1\}.$$

Since $\delta(p_{n+1}) = \delta(q_{n+1})$, we must have $\delta(p_{n+1}) = \delta(q_{n+1}) = \beta + 1$.

As for (p_{n+2}, q_{n+2}) , by (7), we have

$$\delta(p_{n+2}) = \delta(p_{n+1} \varphi) \in \{\beta + 2, \beta + 3\}, \quad \delta(q_{n+2}) = \delta(q_n \varphi^2) \in \{\beta + 3, \beta + 4\}.$$

Hence, $\delta(p_{n+2}) = \delta(q_{n+2}) = \beta + 3$.

As for (p_{n+3}, q_{n+3}) , by (7), we obtain

$$\delta(p_{n+3}) = \delta(p_{n+2} \varphi) \in \{\beta + 4, \beta + 5\}, \quad \delta(q_{n+3}) = \delta(q_{n+2} \varphi) \in \{\beta + 4, \beta + 5\}.$$

Now, on one hand we have, that the relation $q_n < 10^\beta \leq p_n$ and $\varphi < 38$ implies

$$q_{n+3} = q_n \varphi^3 < 10^\beta \cdot 38^3, \quad \delta(q_{n+3}) \leq \delta(10^\beta \cdot 38^3) = \beta + 4.$$

Hence, $\delta(q_{n+3}) = \beta + 4$. On the other,

$$p_{n+3} = q_{n+3} \sqrt{10} = q_{n+2} \varphi \sqrt{10} \geq 10^{\beta+3} \varphi \sqrt{10} > 10^{\beta+3} \cdot 37 \cdot \sqrt{10}.$$

Then, $\delta(p_{n+3}) \geq \delta(10^{\beta+3} \cdot 37 \cdot \sqrt{10}) = \beta + 5$. Hence, $\delta(p_{n+3}) = \beta + 5$. Thus, we conclude $\delta(p_{n+3}) = \beta + 5 = \delta(q_{n+3}) + 1$. \square

4. Conclusion

We summarize here our enquires on the WhtasApp question raised in the introduction. The set \mathcal{S} of positive solutions of the diophantine equation $x(x+1) = 10y(y+1)$ is the set of terms of the sequence (x_n, y_n) defined by

$$(x_1, y_1) = (4, 1), \quad (x_2, y_2) = (20, 6), \quad (x_3, y_3) = (39, 12),$$

and, for $n \geq 1$,

$$x_{n+3} = 19x_n + 60y_n + 39, \quad y_{n+3} = 6x_n + 19y_n + 12.$$

Explicitly, for $k \in \{1, 2, 3\}$, if

$$A_k = \frac{1}{2} \left(\frac{x_k}{\sqrt{10}} + y_k + \frac{10 + \sqrt{10}}{20} \right), \quad \text{and} \quad \varphi = 19 + 6\sqrt{10},$$

then, for $n \geq 1$,

$$x_{3n+k} = \lfloor A_k \sqrt{10} \varphi^n \rfloor, \quad y_{3n+k} = \lfloor A_k \varphi^n \rfloor.$$

The set \mathcal{C} of solutions that concatenate well with regard to our WhastApp problem is formed by the terms $(x_n, y_n) \in \mathcal{S}$ such that x_n has one more digit than y_n . The set \mathcal{C} has infinitely many elements, and the sequence of quotients $(y_n + 1)/(x_n + 1)$ with $(x_n, y_n) \in \mathcal{C}$ is strictly decreasing with limit $1/\sqrt{10}$.

Acknowledgement

We would like to thanks Xavier de Cabo, the sender of the WhatsApp message.

References

- [1] I. Niven, H. S. Zuckerman, and H. L. Montgomery, *An Introduction to the Theory of Numbers*, John Wiley & Sons, 1991.
- [2] T. Ono, *An Introduction to Algebraic Number Theory*, Plenum Press, 1990.
- [3] P. Stevenhagen, *Number Rings*,
<http://websites.math.leidenuniv.nl/algebra/ant.pdf>