

ZEROS OF RANKIN–SELBERG L -FUNCTIONS IN FAMILIES

PETER HUMPHRIES AND JESSE THORNER

ABSTRACT. Let \mathfrak{F}_n be the set of all cuspidal automorphic representations π of GL_n with unitary central character over a number field F . We prove the first unconditional zero density estimate for the set $\mathcal{S} = \{L(s, \pi \times \pi') : \pi \in \mathfrak{F}_n\}$ of Rankin–Selberg L -functions, where $\pi' \in \mathfrak{F}_{n'}$ is fixed. We use this density estimate to establish

- (i) a hybrid-aspect subconvexity bound at $s = \frac{1}{2}$ for almost all $L(s, \pi \times \pi') \in \mathcal{S}$,
- (ii) a strong on-average form of effective multiplicity one for almost all $\pi \in \mathfrak{F}_n$, and
- (iii) a positive level of distribution for $L(s, \pi \times \tilde{\pi})$, in the sense of Bombieri–Vinogradov, for each $\pi \in \mathfrak{F}_n$.

1. INTRODUCTION AND STATEMENT OF THE MAIN RESULT

Let \mathbb{A}_F be the ring of adèles over a number field F with absolute norm $N = N_{F/\mathbb{Q}}$ and absolute discriminant D_F . Let \mathfrak{F}_n be the set of cuspidal automorphic representations $\pi = \bigotimes_v \pi_v$ of $\mathrm{GL}_n(\mathbb{A}_F)$, where the (restricted) tensor product runs over all places of F and π is normalized so that its central character is trivial on the diagonally embedded copy of the positive reals. Let \mathfrak{q}_π be the arithmetic conductor of π , $C(\pi) \geq 1$ the analytic conductor of π (see (3.4)), and $\mathfrak{F}_n(Q) = \{\pi \in \mathfrak{F}_n : C(\pi) \leq Q\}$. The analytic conductor $C(\pi)$ is a useful measure for the arithmetic and spectral complexity of π . Our normalization for the central characters ensures that $|\mathfrak{F}_n(Q)|$ is finite.

Given $\pi \in \mathfrak{F}_n$ and $\pi' \in \mathfrak{F}_{n'}$, let $L(s, \pi \times \pi')$ be the associated Rankin–Selberg L -function, and let $\tilde{\pi} \in \mathfrak{F}_n$ and $\tilde{\pi}' \in \mathfrak{F}_{n'}$ be the contragredient representations. When $\pi' \in \{\tilde{\pi}, \tilde{\pi}'\}$, work of Brumley [23, Appendix] and the authors [24] shows that there exists an effectively computable constant $c_1 = c_1(n, n') > 0$ such that $L(s, \pi \times \pi')$ has a “standard” zero-free region of the shape

$$(1.1) \quad \mathrm{Re}(s) \geq 1 - \frac{c_1}{\log(C(\pi)C(\pi')(|\mathrm{Im}(s)| + 3)^{[F:\mathbb{Q}]})}$$

apart from at most one real simple zero. This is comparable to the classical zero-free region for Dirichlet L -functions. Brumley ([8] and [32, Appendix]) established a much narrower zero-free region for all choices of π and π' . The generalized Riemann hypothesis (GRH) asserts that $L(s, \pi \times \pi') \neq 0$ for $\mathrm{Re}(s) > \frac{1}{2}$. Zeros near the line $\mathrm{Re}(s) = 1$ are typically most damaging in applications, but even a zero-free region of the shape $\mathrm{Re}(s) \geq 1 - \delta$ for some constant $\delta = \delta(n, n', [F : \mathbb{Q}]) > 0$ would be sufficient to obtain many spectacular arithmetic consequences.

Since such strong zero-free regions for Rankin–Selberg L -functions remain out of reach, it is useful to show that zeros near the line $\mathrm{Re}(s) = 1$ must be “sparse”. A suitable quantitative formulation can serve as a proxy for a zero-free region of the shape $\mathrm{Re}(s) \geq 1 - \delta$. Famous

2020 *Mathematics Subject Classification.* 11F66 (primary); 11F12, 11F67 (secondary).

Key words and phrases. Automorphic forms, Rankin–Selberg L -functions, zero density estimate, subconvexity.

consequences of this philosophy include Hoheisel's proof [22] that $p_{n+1} - p_n \ll p_n^{1-1/33000}$ (where p_n is the n -th prime) and Linnik's proof [34] that if $\gcd(a, q) = 1$, then there exists an absolute, effectively computable constant $B > 0$ and a prime $p \leq q^B$ such that $p \equiv a \pmod{q}$.

To quantify our notion of "sparse", we define for $\sigma \geq 0$ and $T \geq 1$ the quantity

$$N_{\pi \times \pi'}(\sigma, T) = |\{\rho = \beta + i\gamma : L(\rho, \pi \times \pi') = 0, \beta \geq \sigma, |\gamma| \leq T\}|.$$

Note that $N_{\pi \times \pi'}(\frac{1}{2}, T)$ is roughly $T \log(C(\pi)C(\pi')T)$ via the argument principle and the functional equation, and GRH can be restated as $N_{\pi \times \pi'}(\sigma, T) = 0$ for all $\sigma > \frac{1}{2}$. The zero density estimate

$$(1.2) \quad N_{\pi \times \pi'}(\sigma, T) \ll_{n, n', [F: \mathbb{Q}]} (C(\pi)C(\pi')T^{[F: \mathbb{Q}]})^{10^7(n'n)^4(1-\sigma)}.$$

follows from work of Soundararajan and Thorner [45, Corollary 2.6]. Therefore, while an arbitrary Rankin–Selberg L -function $L(s, \pi \times \pi')$ is not yet known to have the standard zero-free region (1.1), the bound (1.2) ensures that the number of zeros in the region (1.1) is $O_{n, n', [F: \mathbb{Q}]}(1)$.

Let $\mathcal{S} \subseteq \mathfrak{F}_n$, and let $\mathcal{S}(Q) = \{\pi \in \mathcal{S} : C(\pi) \leq Q\}$. In this article, we seek a strong averaged form of (1.2), namely

$$(1.3) \quad \sum_{\pi \in \mathcal{S}(Q)} N_{\pi \times \pi'}(\sigma, T) \ll_{n, n', [F: \mathbb{Q}], \varepsilon} (Q|\mathcal{S}(Q)|C(\pi')T^{[F: \mathbb{Q}]})^{A(1-\sigma)+\varepsilon},$$

where $A = A(n, n', [F: \mathbb{Q}]) > 0$ is a constant and $\varepsilon > 0$. The bound (1.3) follows from the works of Brumley, Thorner, and Zaman under at least one of the following hypotheses:¹

- $\pi' \in \mathfrak{F}_1$ is trivial [47, Theorem 1.2],
- $\max\{n, n'\} \leq 4$ [11, Theorem 1.3], or
- π' and each $\pi \in \mathcal{S}(Q)$ satisfy certain unproven partial progress towards the generalized Ramanujan conjecture (GRC) [11, Hypothesis 1.1 and Theorem 1.3].²

Here, we prove the first completely unconditional zero density estimate of the form (1.3).

Theorem 1.1. *Let $n, n' \geq 1$ and $\varepsilon > 0$. Let $\mathcal{S} \subseteq \mathfrak{F}_n$ and $\mathcal{S}(Q) = \{\pi \in \mathcal{S} : C(\pi) \leq Q\}$. If $0 \leq \sigma \leq 1$, $\pi' \in \mathfrak{F}_{n'}$, and $Q, T \geq 1$, then*

$$\sum_{\pi \in \mathcal{S}(Q)} N_{\pi \times \pi'}(\sigma, T) \ll_{n, n', [F: \mathbb{Q}], \varepsilon} (|\mathcal{S}(Q)|^4 (C(\pi')QT^{[F: \mathbb{Q}]})^{6.15 \max\{n^2, n'\}})^{1-\sigma+\varepsilon}.$$

Theorem 1.1 is nontrivial when

$$(1.4) \quad \delta_{\mathcal{S}} = \liminf_{Q \rightarrow \infty} \frac{\log |\mathcal{S}(Q)|}{\log Q} > 0.$$

This is important because in applications, it is usually convenient to bound Q by a power of $|\mathcal{S}(Q)|$ or vice versa. When $\mathcal{S} = \mathfrak{F}_n$, we have the bounds

$$(1.5) \quad Q^{n+1} \ll_{n, F} |\mathfrak{F}_n(Q)| \ll_{\varepsilon} D_F^{-n^2} Q^{2n+\varepsilon}.$$

The upper bound in (1.5) was proved by Brumley, Thorner, and Zaman [11, Theorem A.1]. The lower bound in (1.5) follows from work of Brumley and Milićević [10, Theorem 1.1], who computed a constant $c_{n, F} > 0$ such that if $\mathfrak{F}_n^*(Q)$ is the subset of $\pi \in \mathfrak{F}_n(Q)$ that are spherical at the archimedean places of F , then $|\mathfrak{F}_n^*(Q)| \sim c_{n, F} Q^{n+1}$. (The lower bound in

¹In [11, 45], it is assumed that $F = \mathbb{Q}$. Uniformity over $F \neq \mathbb{Q}$ requires minor modifications.

²If θ_n in (3.3) satisfies $\theta_n \leq \frac{1}{4} - \delta_n$ for some $\delta_n > 0$, then each $\pi \in \mathfrak{F}_n$ satisfies [11, Hypothesis 1.1].

(1.5) reflects the conjectured order of growth; see [10].) Together, Theorem 1.1 and (1.5) imply that

$$(1.6) \quad \sum_{\pi \in \mathfrak{F}_n(Q)} N_{\pi \times \pi'}(\sigma, T) \ll_{n, n', F, \varepsilon} (|\mathfrak{F}_n(Q)| C(\pi')^n T^{n[F:\mathbb{Q}]})^{7.1 \max\{n, n'\}(1-\sigma) + \varepsilon}.$$

Furthermore, if $n' = n \geq 3$ and $\pi' \in \mathfrak{F}_{n'}$, then Theorem 1.1 and (1.5) together imply that

$$(1.7) \quad \sum_{\pi \in \mathfrak{F}_n(Q)} N_{\pi \times \pi'}(\sigma, T) \ll_{\varepsilon} (C(\pi') Q T^{[F:\mathbb{Q}]})^{9n^2(1-\sigma) + \varepsilon}.$$

2. APPLICATIONS

We will now describe some applications of Theorem 1.1. In what follows, we write $f \ll_{\nu} g$, $f = O_{\nu}(g)$, and $g \gg_{\nu} f$ to denote that there exists a constant $c > 0$ such that $|f| \leq c|g|$ in the stated range. The implied constant c , which is effectively computable unless otherwise stated, will depend at most on ν , n , n' , and $[F : \mathbb{Q}]$. The expression $f \asymp_{\nu} g$ means that $f \ll_{\nu} g$ and $g \ll_{\nu} f$. We use $\varepsilon > 0$ to denote an arbitrarily small quantity that depends at most on n , n' , and $[F : \mathbb{Q}]$.

2.1. Bounds for Rankin–Selberg L -functions. It is a classical problem for Dirichlet L -functions to find strong bounds on the critical line $\operatorname{Re}(s) = \frac{1}{2}$. The Phragmén–Lindelöf convexity principle shows that if q_{χ} is the conductor of a primitive Dirichlet character χ , then $L(\frac{1}{2}, \chi) \ll q^{1/4}$; improving this bound by replacing $1/4$ with a smaller exponent is known as a *subconvex* bound. The multiplicative version of the classical large sieve inequality combined with an approximate functional equation shows that for almost all χ , we have the bound $L(\frac{1}{2}, \chi) \ll_{\varepsilon} q_{\chi}^{\varepsilon}$ for all $\varepsilon > 0$, consistent with the generalized Lindelöf hypothesis (GLH).

For $\pi \in \mathfrak{F}_2$, GLH predicts that $L(\frac{1}{2}, \pi) \ll_{\varepsilon} C(\pi)^{\varepsilon}$. Michel and Venkatesh [38] proved that there exists a fixed positive $\delta > 0$ such that $L(\frac{1}{2}, \pi) \ll_F C(\pi)^{1/4-\delta}$, the culmination of several decades of research. When $F = \mathbb{Q}$, a sharp mean value estimate for Hecke eigenvalues proved by Deshouillers and Iwaniec [14], in conjunction with the approximate functional equation, implies the bound $L(\frac{1}{2}, \pi) \ll_{\varepsilon} (qT)^{\varepsilon}$ for almost all $\pi \in \mathfrak{F}_2$ of arithmetic conductor q , trivial central character, and archimedean complexity (Laplace eigenvalue or weight squared) lying in the dyadic interval $[T, 2T]$. Note that in this case, $C(\pi) \asymp qT$.

For $\pi \in \mathfrak{F}_n$ with $n \geq 3$, the best uniform result towards the bound $L(\frac{1}{2}, \pi) \ll_{\varepsilon} C(\pi)^{\varepsilon}$ predicted by GLH is that of Soundararajan and Thorner [45, Corollary 2.7], namely

$$(2.1) \quad L(\frac{1}{2}, \pi) \ll C(\pi)^{\frac{1}{4}} (\log C(\pi))^{-1/(10^{17}n^3)}.$$

We mention three results that improve upon (2.1) in an average sense, each having complementary strengths. Jana [29, Theorem 6] extended the GLH-on-average bound of Deshouillers and Iwaniec to the family of cuspidal automorphic representations of $\operatorname{GL}_n(\mathbb{A}_{\mathbb{Q}})$ of arithmetic conductor 1 and growing analytic conductor. Blomer [3, Corollary 5] proved the corresponding result for the family of cuspidal automorphic representations of $\operatorname{GL}_n(\mathbb{A}_{\mathbb{Q}})$ of a large given prime arithmetic conductor q , trivial central character, and whose archimedean components are principal series representations confined to a compact subset of the unitary dual. Thorner and Zaman [47, Theorem 1.3] proved that there exists a constant $c_2 = c_2(n, [F : \mathbb{Q}]) > 0$ such that if $\varepsilon > 0$, then

$$(2.2) \quad |\{\pi \in \mathfrak{F}_n(Q) : |L(\frac{1}{2}, \pi)| \geq c_2 C(\pi)^{\frac{1}{4} - \frac{\varepsilon}{10^{16}n^3}}\}| \ll_F |\mathfrak{F}_n(Q)|^{\varepsilon}.$$

Unlike the preceding results, (2.2) is uniform in both the arithmetic conductor and spectral aspects and holds over number fields other than \mathbb{Q} , but the savings over (2.1) is not comparable to GLH on average.

Given $\pi \in \mathfrak{F}_n$ and $\pi' \in \mathfrak{F}_{n'}$, Soundararajan and Thorner [45, Corollary 2.7] proved when $F = \mathbb{Q}$ that if $C(\pi \times \pi')$ is the analytic conductor of $L(s, \pi \times \pi')$, then

$$(2.3) \quad L\left(\frac{1}{2}, \pi \times \pi'\right) \ll |L\left(\frac{3}{2}, \pi \times \pi'\right)|^2 C(\pi \times \pi')^{\frac{1}{4}} (\log C(\pi \times \pi'))^{-1/(10^{17}(n'n)^3)}.$$

As of now, the best general upper bound for $|L(\frac{3}{2}, \pi \times \pi')|^2$ is larger than any fixed power of $\log C(\pi \times \pi')$ [33, Theorem 2]. The factor of $|L(\frac{3}{2}, \pi \times \pi')|^2$ can be removed under certain partial progress toward GRC. The bound $L(\frac{1}{2}, \pi \times \pi') \ll_{\varepsilon} C(\pi \times \pi')^{\varepsilon}$ is predicted by GLH.

In order to improve (2.3) on average with uniformity in π and π' , one might first try to mimic the approach that worked well for Dirichlet L -functions and GL_2 L -functions using trace formulae, approximate functional equations, the spectral large sieve, Voronoï summation, etc. While such methods have seen great success for $\mathrm{GL}_n \times \mathrm{GL}_{n'}$ with $n, n' \in \{1, 2\}$, suitably uniform and flexible versions of these tools do not appear to be available yet in the general setting. The special case where $|n - n'| \leq 1$ exhibits some nice structural properties, lending itself to approaches via period integrals that completely avoids the aforementioned tools. To describe work in this direction, let $F = \mathbb{Q}$, $\mathcal{F}_n \subseteq \mathfrak{F}_n$ be the subset of cuspidal automorphic representations of $\mathrm{GL}_n(\mathbb{A}_{\mathbb{Q}})$ of arithmetic conductor 1, and $\mathcal{F}_n(Q) = \{\pi \in \mathcal{F}_n : C(\pi) \leq Q\}$. It follows from work of Jana [30, Corollary 2.2] that if $\varepsilon > 0$, $\pi' \in \mathcal{F}_n$, and the spectral parameters of π' have real part at least $-\frac{1}{n^2+1}$ (which is far stronger than the best known unconditional lower bound $-\frac{1}{2} + \frac{1}{n^2+1}$ due to Luo, Rudnick, and Sarnak [37]), then

$$(2.4) \quad \sum_{\pi \in \mathcal{F}_n(Q)} |L(\frac{1}{2}, \pi \times \pi')|^2 \ll_{\pi', \varepsilon} |\mathcal{F}_n(Q)|^{1+\varepsilon}.$$

Therefore, for fixed $\pi' \in \mathcal{F}_n$, the generalized Lindelöf hypothesis for $L(\frac{1}{2}, \pi \times \pi')$ holds on average over the $\pi \in \mathcal{F}_n$. Using Chebyshev's inequality, we conclude that for all $\delta > 0$, there exists a constant $c_{\pi', \delta} > 0$ such that

$$(2.5) \quad |\{\pi \in \mathcal{F}_n(Q) : |L(\frac{1}{2}, \pi \times \pi')| \geq c_{\pi', \delta} C(\pi \times \pi')^{2\delta}\}| \ll_{\pi', \delta} |\mathcal{F}_n(Q)|^{1-\delta}.$$

See also the work of Blomer [2, Theorem 2], which proves a variant of (2.4) for families of ‘‘spectrally close’’ Hecke–Maaß newforms on $\mathrm{SL}_n(\mathbb{Z})$.

Along the same lines as (2.2), we use (1.6) to prove the following result.

Theorem 2.1. *Let $n, n' \geq 1$ and $Q \geq 1$. If $\varepsilon > 0$ and $\pi' \in \mathfrak{F}_{n'}$, then*

$$|\{\pi \in \mathfrak{F}_n(Q) : |L(\frac{1}{2}, \pi \times \pi')| \geq C(\pi \times \pi')^{\frac{1}{4} - \frac{\varepsilon}{10^{10} \max\{n, n'\}}} \}| \ll_{\varepsilon} (C(\pi')^n |\mathfrak{F}_n(Q)|)^{\varepsilon}.$$

Remark 2.2. Given a subset $\mathcal{S} \subseteq \mathfrak{F}_n$, a similar result can be proved for $\pi \in \mathcal{S}(Q)$ using Theorem 1.1. Such a result would depend effectively on $\delta_{\mathcal{S}}$ in (1.4).

In contrast with Jana's work in (2.5), Theorem 2.1 provides a smaller power-saving improvement over (2.3), but the improvement is uniform in the arithmetic conductor and spectral aspects as well as in π' . The exceptional set in Theorem 2.1 is a much smaller than in (2.5). Theorem 2.1 removes the requirements that $n = n'$, that $q_{\pi} = q_{\pi'} = 1$, and that the spectral parameters of π' have real part at least $-\frac{1}{n^2+1}$. Finally, Theorem 2.1 is proved over any number field, while [30] is only proved over \mathbb{Q} .

2.2. Effective multiplicity one. Let $\pi = \bigotimes_v \pi_v$ and $\pi' = \bigotimes_v \pi'_v$ be cuspidal automorphic representations in $\mathfrak{F}_n(Q)$. Under the assumption of GRH for $L(s, \pi \times \tilde{\pi})$ and $L(s, \pi \times \pi')$ and that $\pi_{\mathfrak{p}}$ and $\pi'_{\mathfrak{p}}$ are tempered for all prime ideals $\mathfrak{p} \mid \mathfrak{q}_{\pi} \mathfrak{q}_{\pi'}$, it is known that if $Y = (\log Q)^2$ and $\pi_{\mathfrak{p}} \cong \pi'_{\mathfrak{p}}$ for all $\mathfrak{p} \nmid \mathfrak{q}_{\pi} \mathfrak{q}_{\pi'}$ with $N\mathfrak{p} \ll Y$, then $\pi = \pi'$ [25, Proposition 5.22]. Brumley [8], improving on work of Moreno [40], proved that there exists a constant $B_n > 0$ such that this result holds unconditionally with $Y = Q^{B_n}$. This result makes effective the multiplicity one theorems of Jacquet and Shalika [28, Theorem 4.8] and Piatetski-Shapiro [43]. Any fixed $B_n > 2n$ suffices [35].

When $n = 2$, we have an average result that nearly achieves what GRH predicts. Specifically, let \mathfrak{F}_2^b be the subset of $\pi \in \mathfrak{F}_2$ with squarefree conductor and trivial central character, and let $\pi' \in \mathfrak{F}_2^b$. For all $\varepsilon > 0$, there exists an effectively computable constant $N_\varepsilon > 0$, depending at most on ε and $[F : \mathbb{Q}]$, such that

$$(2.6) \quad |\{\pi \in \mathfrak{F}_2^b(Q) : \pi_{\mathfrak{p}} \cong \pi'_{\mathfrak{p}} \text{ for all } \mathfrak{p} \nmid \mathfrak{q}_{\pi} \text{ with } N\mathfrak{p} \leq (\log Q)^{N_\varepsilon}\}| \ll_\varepsilon Q^\varepsilon.$$

In particular, the implied constant does not depend on π' . This was proved by Duke and Kowalski [15, Theorem 3] when $F = \mathbb{Q}$ in a stronger form under the assumption of GRC. See Brumley's Ph.D. thesis [7, Corollary 5.2.2] for a proof that does not use GRC.

If $\pi \in \mathfrak{F}_2$ and $\tilde{\pi} \in \mathfrak{F}_2$ is the contragredient, then $L(s, \pi \times \tilde{\pi}) = \zeta_F(s) L(s, \pi, \text{Ad})$, where $\zeta_F(s)$ is the Dedekind zeta function of F and Ad is the adjoint square lift from a representation of $\text{GL}_2(\mathbb{A}_F)$ to a representation of $\text{GL}_3(\mathbb{A}_F)$. The arguments in [7, 15] rely on two key results:

- (1) Gelbart and Jacquet [18] proved that if $\pi \in \mathfrak{F}_2$, then $L(s, \pi, \text{Ad})$ is the L -function of an automorphic representation of $\text{GL}_3(\mathbb{A}_F)$, denoted $\text{Ad } \pi$, complete with a criterion by which one can determine whether $\text{Ad } \pi$ is cuspidal (and lies in \mathfrak{F}_3).
- (2) At most $O_\varepsilon(Q^{1/2+\varepsilon})$ representations $\pi \in \mathfrak{F}_2(Q)$ have the same adjoint lift. Also, Ramakrishnan ([15, Appendix] and [44]) proved that $\text{Ad} : \mathfrak{F}_2^b \rightarrow \mathfrak{F}_3$ is injective.

In attempting to generalize the strategy of Duke and Kowalski to GL_n for $n \geq 3$, one encounters some deep open problems. If $\pi \in \mathfrak{F}_n$, then for $\text{Re}(s)$ sufficiently large, $L(s, \pi \times \tilde{\pi})$ factors as $\zeta_F(s) L(s, \pi, \text{Ad})$, where Ad is the adjoint square lift from GL_n to GL_{n^2-1} . Apart from some special cases, the following obstacles arise:

- (1) The adjoint lift is not yet known to be automorphic for $n \geq 3$, and if it were, there is no known criterion for cuspidality.
- (2) Let $\mathcal{H}_n \subseteq \mathfrak{F}_n(Q)$ be the subset of $\pi \in \mathfrak{F}_n(Q)$ such that $\text{Ad } \pi \in \mathfrak{F}_{n^2-1}$. It is not known how many $\pi \in \mathcal{H}_n$ have the same adjoint lift.

Despite these setbacks, we can use (1.7) (more specifically, Corollary 7.1 below) to prove a GL_n analogue of (2.6) when π' depends mildly on Q .

Theorem 2.3. *Let $n \geq 3$ and $Q \geq 1$. There exists an absolute, effectively computable constant $c_3 > 0$ such that if $0 < \varepsilon < 1$ and $\pi' \in \mathfrak{F}_n((\log Q)^{c_3/(n^2[F:\mathbb{Q}]^2)})$, then*

$$|\{\pi \in \mathfrak{F}_n(Q) : \pi_{\mathfrak{p}} \cong \pi'_{\mathfrak{p}} \text{ for all } \mathfrak{p} \nmid \mathfrak{q}_{\pi} \mathfrak{q}_{\pi'} \text{ with } N\mathfrak{p} \leq (\log Q)^{41n^2/\varepsilon}\}| \ll_\varepsilon Q^\varepsilon.$$

Remark 2.4. Given a subset $\mathcal{S} \subseteq \mathfrak{F}_n$, a similar result can be proved for $\pi \in \mathcal{S}(Q)$ using Theorem 1.1. Such a result would depend effectively on $\delta_{\mathcal{S}}$ in (1.4).

The proof is very flexible. For example, if $\varepsilon > 0$ and $C(\pi') \ll_\varepsilon Q^{c_3\varepsilon^2/(n^2[F:\mathbb{Q}]^2)}$, then the same proof with minor changes in choices of parameters produces the bound

$$|\{\pi \in \mathfrak{F}_n(Q) : \pi_{\mathfrak{p}} \cong \pi'_{\mathfrak{p}} \text{ for all } \mathfrak{p} \nmid \mathfrak{q}_{\pi} \mathfrak{q}_{\pi'} \text{ with } N\mathfrak{p} \leq Q^\varepsilon\}| \ll_\varepsilon Q^\varepsilon.$$

While the number of \mathfrak{p} for which one needs to check that $\pi_{\mathfrak{p}} \cong \pi'_{\mathfrak{p}}$ is larger, the range of $C(\pi')$ is greatly extended, and the threshold $N\mathfrak{p} \leq Q^\varepsilon$ (reminiscent of Vinogradov's conjecture on the size of the least quadratic nonresidue) still greatly improves on the unconditional range $N\mathfrak{p} \ll_\varepsilon Q^{2n+\varepsilon}$ from [35].

2.3. Automorphic level of distribution. Let $\Lambda(m)$ be the von Mangoldt function, equal to $\log p$ if m is a power of a prime p and zero otherwise. The celebrated Bombieri–Vinogradov theorem states that if $\theta < \frac{1}{2}$ is fixed, then for all $A > 0$, we have

$$(2.7) \quad \sum_{q \leq x^\theta} \max_{\gcd(a,q)=1} \max_{y \leq x} \left| \sum_{\substack{m \leq y \\ m \equiv a \pmod{q}}} \Lambda(m) - \frac{y}{\varphi(q)} \right| \ll_A \frac{x}{(\log x)^A}.$$

This may be viewed as an average form of GRH for Dirichlet L -functions. As part of his proof [6], Bombieri proved a strong form of the zero density estimate in Theorem 1.1 for Dirichlet L -functions. We call any θ for which (2.7) holds a *level of distribution* for the primes. Elliott and Halberstam conjectured that any fixed $\theta < 1$ is a level of distribution for the primes.

Number theorists have proved several interesting extensions and variations of (2.7). For example, Murty and Murty [42] proved that primes in the Chebotarev density theorem have a positive level of distribution. To describe a different direction for automorphic representations over \mathbb{Q} , we let $n \geq 2$ and consider $\pi \in \mathfrak{F}_n$ with conductor q_π . Let $\Lambda(m)$ be the von Mangoldt function, and define the numbers $a_\pi(m)$ by

$$-\frac{L'}{L}(s, \pi) = \sum_p \sum_{k=1}^{\infty} \frac{\sum_{j=1}^n \alpha_{j,\pi}(p)^k \log p}{p^{ks}} = \sum_{n=1}^{\infty} \frac{a_\pi(m) \Lambda(m)}{m^s}, \quad \operatorname{Re}(s) > 1.$$

Note that $a_\pi(p) = \lambda_\pi(p)$. For fixed $\theta < \frac{1}{n^2-2}$, Wong [49, Theorem 9] proved that if π satisfies GRC and $L(s, \pi \times (\tilde{\pi} \otimes \chi))$ has no Landau–Siegel zero for all Dirichlet characters χ , then for any $A > 0$,

$$(2.8) \quad \sum_{q \leq x^\theta} \max_{\gcd(a,q)=1} \max_{y \leq x} \left| \sum_{\substack{m \leq y \\ \gcd(m, q_\pi)=1 \\ m \equiv a \pmod{q}}} |a_\pi(m)|^2 \Lambda(m) - \frac{y}{\varphi(q)} \right| \ll_{A,\pi} \frac{x}{(\log x)^A}.$$

This conditionally endows $L(s, \pi \times \tilde{\pi})$ with a positive level of distribution θ . The hypotheses for (2.8) hold for π attached to non-CM holomorphic cuspidal newforms on congruence subgroups of $\mathrm{SL}_2(\mathbb{Z})$.

Let $\pi \in \mathfrak{F}_n$. Using (1.7), we unconditionally endow $L(s, \pi \times \tilde{\pi})$ with a notion of positive level of distribution. In particular, we avoid recourse to unproven progress toward GRC or the absence of Landau–Siegel zeros.

Theorem 2.5. *Let $F = \mathbb{Q}$ and $\pi \in \mathfrak{F}_n$. Fix $\theta < 1/(9n^3)$. If $A > 0$, then*

$$\sum_{\substack{q \leq x^\theta \\ \gcd(q, q_\pi)=1}} \max_{\gcd(a,q)=1} \max_{y \leq x} \left| \sum_{\substack{m \leq y \\ m \equiv a \pmod{q}}} |a_\pi(m)|^2 \Lambda(m) - \frac{y}{\varphi(q)} \right| \ll_{A,\pi} \frac{x}{(\log x)^A}.$$

The implied constants are ineffective.

Remark 2.6. One can prove an analogue of Theorem 2.5 with $F \neq \mathbb{Q}$, replacing residue classes modulo q with ray classes modulo \mathfrak{q} . We restrict to $F = \mathbb{Q}$ for notational simplicity.

Overview of the paper. In Section 3, we recall basic properties of standard L -functions and Rankin–Selberg L -functions that we will use in our proofs. In Section 4, we prove a large sieve inequality for the Dirichlet coefficients of $L(s, \pi \times \pi')^{-1}$ and a corollary on mean values of Dirichlet polynomials, which we use in our proof of Theorem 1.1 in Section 5. We then prove Theorem 2.1 in Section 6, Theorem 2.3 in Section 7, and Theorem 2.5 in Section 8.

Acknowledgements. We thank Gergely Harcos, Rizwanur Khan, and the anonymous referee for helpful comments.

3. PROPERTIES OF L -FUNCTIONS

We recall some standard facts about L -functions arising from automorphic representations and their Rankin–Selberg convolutions. See [8, 19, 27, 39, 45].

3.1. Standard L -functions. Given $\pi \in \mathfrak{F}_n$, let $\tilde{\pi} \in \mathfrak{F}_n$ be the contragredient representation and \mathfrak{q}_π be the conductor of π . We express π as a restricted tensor product $\bigotimes_v \pi_v$ of smooth admissible representations of $\mathrm{GL}_n(F_v)$, where v varies over places of F . When v is a nonarchimedean place corresponding with a prime ideal \mathfrak{p} , then the local L -function $L(s, \pi_{\mathfrak{p}})$ is defined in terms of the Satake parameters $A_\pi(\mathfrak{p}) = \{\alpha_{1,\pi}(\mathfrak{p}), \dots, \alpha_{n,\pi}(\mathfrak{p})\}$ by

$$(3.1) \quad L(s, \pi_{\mathfrak{p}}) = \prod_{j=1}^n (1 - \alpha_{j,\pi}(\mathfrak{p}) \mathrm{N}\mathfrak{p}^{-s})^{-1} = \sum_{k=0}^{\infty} \frac{\lambda_\pi(\mathfrak{p}^k)}{\mathrm{N}\mathfrak{p}^{ks}}.$$

We have $\alpha_{j,\pi}(\mathfrak{p}) \neq 0$ for all j whenever $\mathfrak{p} \nmid \mathfrak{q}_\pi$, and when $\mathfrak{p} \mid \mathfrak{q}_\pi$, it might be the case that there exist j such that $\alpha_{j,\pi}(\mathfrak{p}) = 0$. The standard L -function $L(s, \pi)$ associated to π is of the form

$$L(s, \pi) = \prod_{\mathfrak{p}} L(s, \pi_{\mathfrak{p}}) = \sum_{\mathfrak{n}} \frac{\lambda_\pi(\mathfrak{n})}{\mathrm{N}\mathfrak{n}^s}.$$

The Euler product and Dirichlet series converge absolutely when $\mathrm{Re}(s) > 1$.

At each archimedean place v of F , there are n Langlands parameters $\mu_{j,\pi}(v) \in \mathbb{C}$ such that

$$L(s, \pi_\infty) = \prod_{v|\infty} \prod_{j=1}^n \Gamma_v(s + \mu_{j,\pi}(v)), \quad \Gamma_v(s) := \begin{cases} \pi^{-s/2} \Gamma(s/2) & \text{if } F_v = \mathbb{R}, \\ 2(2\pi)^{-s} \Gamma(s) & \text{if } F_v = \mathbb{C}. \end{cases}$$

By combining the work in [4, 5, 37, 41], we know that there exists

$$(3.2) \quad 0 \leq \theta_n \leq \begin{cases} 0 & \text{if } n = 1, \\ 7/64 & \text{if } n = 2, \\ 5/14 & \text{if } n = 3, \\ 9/22 & \text{if } n = 4, \\ 1/2 - 1/(n^2 + 1) & \text{if } n \geq 5 \end{cases}$$

such that

$$(3.3) \quad |\alpha_{j,\pi}(\mathfrak{p})| \leq \mathrm{N}\mathfrak{p}^{\theta_n} \quad \text{and} \quad \mathrm{Re}(\mu_{j,\pi}(v)) \geq -\theta_n.$$

GRC asserts that in (3.2), one may take $\theta_n = 0$. We have $\mathfrak{q}_\pi = \mathfrak{q}_{\tilde{\pi}}$, and for each \mathfrak{p} and each v , we have the equalities of sets $\{\alpha_{j,\tilde{\pi}}(\mathfrak{p})\} = \overline{\{\alpha_{j,\pi}(\mathfrak{p})\}}$ and $\{\mu_{j,\tilde{\pi}}(v)\} = \overline{\{\mu_{j,\pi}(v)\}}$.

Let r_π be the order of the pole of $L(s, \pi)$ at $s = 1$. The completed L -function

$$\Lambda(s, \pi) = (s(s-1))^{r_\pi} (D_F^n \mathrm{N}\mathfrak{q}_\pi)^{s/2} L(s, \pi) L(s, \pi_\infty)$$

is entire of order 1, and there exists a complex number $W(\pi)$ of modulus 1 such that for all $s \in \mathbb{C}$, we have the functional equation $\Lambda(s, \pi) = W(\pi)\Lambda(1-s, \tilde{\pi})$. Let $d(v) = 1$ if $F_v = \mathbb{R}$ and $d(v) = 2$ if $F_v = \mathbb{C}$. The analytic conductor of π [26] is given by

$$(3.4) \quad C(\pi, t) := D_F^n N_{\mathfrak{q}_\pi} \prod_{v|\infty} \prod_{j=1}^n (3 + |it + \mu_{j,\pi}(v)|^{d(v)}), \quad C(\pi) := C(\pi, 0).$$

Since $\Lambda(s, \pi)$ is entire of order 1, there exist complex numbers a_π and b_π such that

$$\Lambda(s, \pi) = e^{a_\pi + b_\pi s} \prod_{\Lambda(\rho, \pi) = 0} \left(1 - \frac{s}{\rho}\right) e^{s/\rho}.$$

The zeros ρ in the above Hadamard product are the nontrivial zeros of $L(s, \pi)$, and the zeros of $L(s, \pi)$ that arise as poles of $s^{r_\pi} L(s, \pi_\infty)$ are the trivial zeros.

3.2. Rankin–Selberg L -functions. Let $\pi \in \mathfrak{F}_n$ and $\pi' \in \mathfrak{F}_{n'}$. At each prime ideal \mathfrak{p} , Jacquet, Piatetski-Shapiro, and Shalika [27] associate to $\pi_{\mathfrak{p}}$ and $\pi'_{\mathfrak{p}}$ a local Rankin–Selberg L -function

$$(3.5) \quad L(s, \pi_{\mathfrak{p}} \times \pi'_{\mathfrak{p}}) = \prod_{j=1}^n \prod_{j'=1}^{n'} (1 - \alpha_{j,j',\pi \times \pi'}(\mathfrak{p}) N_{\mathfrak{p}}^{-s})^{-1} = \sum_{k=0}^{\infty} \frac{\lambda_{\pi \times \pi'}(\mathfrak{p}^k)}{N_{\mathfrak{p}}^{ks}}$$

and a local conductor $\mathfrak{q}_{\pi_{\mathfrak{p}} \times \pi'_{\mathfrak{p}}}$. If $\mathfrak{p} \nmid \mathfrak{q}_{\pi} \mathfrak{q}_{\pi'}$, then we have the equality of sets

$$(3.6) \quad \{\alpha_{j,j',\pi \times \pi'}(\mathfrak{p})\} = \{\alpha_{j,\pi}(\mathfrak{p}) \alpha_{j',\pi'}(\mathfrak{p})\}.$$

The Rankin–Selberg L -function $L(s, \pi \times \pi')$ associated to π and π' and its arithmetic conductor are

$$L(s, \pi \times \pi') = \prod_{\mathfrak{p}} L(s, \pi_{\mathfrak{p}} \times \pi'_{\mathfrak{p}}) = \sum_{\mathfrak{n}} \frac{\lambda_{\pi \times \pi'}(\mathfrak{n})}{N_{\mathfrak{n}}^s}, \quad \mathfrak{q}_{\pi \times \pi'} = \prod_{\mathfrak{p}} \mathfrak{q}_{\pi_{\mathfrak{p}} \times \pi'_{\mathfrak{p}}}.$$

At an archimedean place v of F , Jacquet, Piatetski-Shapiro, and Shalika associate $n'n$ complex Langlands parameters $\mu_{j,j',\pi \times \pi'}(v)$ to π_v and π'_v , from which one defines

$$L(s, \pi_\infty \times \pi'_\infty) = \prod_{v|\infty} \prod_{j=1}^n \prod_{j'=1}^{n'} \Gamma_v(s + \mu_{j,j',\pi \times \pi'}(v)).$$

Using the explicit descriptions of $\alpha_{j,j',\pi \times \pi'}(\mathfrak{p})$ and $\mu_{j,j',\pi \times \pi'}(v)$ in [23, 45], one sees that

$$(3.7) \quad |\alpha_{j,j',\pi \times \pi'}(\mathfrak{p})| \leq N_{\mathfrak{p}}^{\theta_n + \theta_{n'}}, \quad \operatorname{Re}(\mu_{j,j',\pi \times \pi'}(v)) \geq -\theta_n - \theta_{n'}.$$

Let $r_{\pi \times \pi'} = -\operatorname{ord}_{s=1} L(s, \pi \times \pi')$. By our normalization for the central characters of π and π' , we have that $r_{\pi \times \pi'} = 0$ if and only if $\pi \neq \tilde{\pi}'$, and $r_{\pi \times \tilde{\pi}'} = 1$ otherwise. The function

$$(3.8) \quad \Lambda(s, \pi \times \pi') = (s(s-1))^{r_{\pi \times \pi'}} (D_F^{n'n} N_{\mathfrak{q}_{\pi \times \pi'}})^{s/2} L(s, \pi \times \pi') L(s, \pi_\infty \times \pi'_\infty)$$

is entire of order 1, and there exists a complex number $W(\pi \times \pi')$ of modulus 1 such that $\Lambda(s, \pi \times \pi')$ satisfies the functional equation $\Lambda(s, \pi \times \pi') = W(\pi \times \pi') \Lambda(1-s, \tilde{\pi} \times \tilde{\pi}')$. As with $L(s, \pi)$, the analytic conductor of $L(s, \pi \times \pi')$ is given by

$$(3.9) \quad C(\pi \times \pi', t) := D_F^{n'n} N_{\mathfrak{q}_{\pi \times \pi'}} \prod_{v|\infty} \prod_{j=1}^n \prod_{j'=1}^{n'} (3 + |it + \mu_{j,j',\pi \times \pi'}(v)|^{d(v)}), \quad C(\pi \times \pi') := C(0, \pi \times \pi').$$

The combined work of Bushnell and Henniart [13] and Brumley [23, Appendix] yields

$$(3.10) \quad C(\pi \times \pi', t) \ll C(\pi \times \pi')(3 + |t|)^{[F:\mathbb{Q}]n'n}, \quad C(\pi \times \pi') \ll C(\pi)^{n'} C(\pi')^n.$$

Since $\Lambda(s, \pi \times \pi')$ is entire of order 1, there exist complex numbers $a_{\pi \times \pi'}$ and $b_{\pi \times \pi'}$ such that the Hadamard factorization

$$(3.11) \quad \Lambda(s, \pi \times \pi') = e^{a_{\pi \times \pi'} + b_{\pi \times \pi'} s} \prod_{\Lambda(\rho, \pi \times \pi')=0} \left(1 - \frac{s}{\rho}\right) e^{s/\rho}$$

holds. The zeros ρ in (3.11) are the nontrivial zeros of $L(s, \pi \times \pi')$, and the zeros of $L(s, \pi \times \pi')$ that arise as poles of $s^{r_{\pi \times \pi'}} L(s, \pi_\infty \times \pi'_\infty)$ are the trivial zeros.

It follows from work of Li [33, Theorem 2] (with minor adjustments when $F \neq \mathbb{Q}$) that there exists an absolute and effectively computable constant $c_4 > 0$, which we assume to be sufficiently large for future convenience, such that

$$(3.12) \quad \lim_{\sigma_0 \rightarrow \sigma} (\sigma_0 - 1)^{r_{\pi \times \pi'}} L(\sigma_0, \pi \times \pi') \ll \exp\left(c_4 n' n [F:\mathbb{Q}] \frac{\log C(\pi \times \pi')}{\log \log C(\pi \times \pi')}\right), \quad \sigma \in [1, 3].$$

We can change π' to $\pi' \otimes |\det|^{it}$; at the archimedean places, this has the effect of adding it to each $\mu_{j,j',\pi \times \pi'}(v)$. We then apply functional equation, the Phragmén–Lindelöf convexity principle, and (3.10) to obtain for all $\sigma \geq 0$

$$(3.13) \quad \begin{aligned} & \lim_{\sigma_0 \rightarrow \sigma} \left(\frac{\sigma_0 + it - 1}{\sigma_0 + it + 1}\right)^{r_{\pi \times \pi'}} L(\sigma_0 + it, \pi \times \pi') \\ & \ll_\varepsilon C(\pi \times \pi', t)^{\frac{\max\{1-\sigma, 0\}}{2} + \frac{\varepsilon}{n'n[F:\mathbb{Q}]}} \\ & \ll_\varepsilon (C(\pi)^{n'} C(\pi')^n (3 + |t|)^{n'n[F:\mathbb{Q}]})^{\frac{\max\{1-\sigma, 0\}}{2} + \frac{\varepsilon}{n'n[F:\mathbb{Q}]}}. \end{aligned}$$

Lemma 3.1. *If $\pi \in \mathfrak{F}_n$, $X \geq 3$, and $\varepsilon > 0$, then $\sum_{N\mathfrak{n} \leq X} \lambda_{\pi \times \tilde{\pi}}(\mathfrak{n})/N\mathfrak{n} \ll_\varepsilon C(\pi)^\varepsilon \log X$.*

Proof. Since $\lambda_{\pi \times \tilde{\pi}}(\mathfrak{n}) \geq 0$ for all \mathfrak{n} by [21, Lemma a], we observe by (3.12) that

$$\sum_{N\mathfrak{n} \leq X} \frac{\lambda_{\pi \times \tilde{\pi}}(\mathfrak{n})}{N\mathfrak{n}} \leq e \sum_{\mathfrak{n}} \frac{\lambda_{\pi \times \tilde{\pi}}(\mathfrak{n})}{N\mathfrak{n}^{1+\frac{1}{\log X}}} \ll (\log X) \operatorname{Res}_{s=1} L(s, \pi \times \tilde{\pi}).$$

The desired bounded now follows from (3.13) with $\sigma = 1$, $t = 0$, and $\pi' = \tilde{\pi}$. \square

Lemma 3.2. *Let $J \geq 1$ be an integer. For all $j \in \{1, \dots, J\}$, let $t_j \in \mathbb{R}$; n_j, n'_j be positive integers; and $\pi_j \in \mathfrak{F}_{n_j}$ and $\pi'_j \in \mathfrak{F}_{n'_j}$. Let*

$$D(s) = \prod_{j=1}^J L(s + it_j, \pi_j \times \tilde{\pi}'_j), \quad \Delta(s) = \prod_{j=1}^J \Lambda(s + it_j, \pi_j \times \tilde{\pi}'_j), \quad Q = \prod_{j=1}^J C(\pi_j)^{n'_j} C(\pi'_j)^{n_j}.$$

Let $R = -\operatorname{ord}_{s=1} D(s)$. If $1 < \sigma < 2$ and the \mathfrak{n} -th Dirichlet coefficient of $-\frac{D'}{D}(s)$ is nonnegative when $\gcd(\mathfrak{n}, \prod_{j=1}^J \mathfrak{q}_{\pi_j} \mathfrak{q}_{\pi'_j}) = \mathcal{O}_F$, then

$$\sum_{\Delta(\rho)=0} \operatorname{Re}\left(\frac{1}{\sigma - \rho}\right) < \frac{R}{\sigma - 1} + \sum_{\substack{1 \leq j \leq J \\ \pi_j = \pi'_j, t_j \neq 0}} \frac{\sigma - 1}{(\sigma - 1)^2 + t_j^2} + O(\log Q).$$

Proof. Let $1 < \sigma < 2$ and

$$D_\infty(s) = \prod_{j=1}^J L(s, (\pi_j)_\infty \times (\tilde{\pi}'_j)_\infty), \quad \mathfrak{q}_D = \prod_{j=1}^J \mathfrak{q}_{\pi_j \times \tilde{\pi}'_j}.$$

By comparing the logarithmic derivative of $\Delta(s)$ with the logarithmic derivative of its Hadamard factorization

$$\Delta(s) = e^{a_D + b_D s} \prod_{\Delta(\rho)=0} \left(1 - \frac{s}{\rho}\right) e^{s/\rho},$$

we find that

$$\sum_{\Delta(\rho)=0} \left(\frac{1}{\sigma - \rho} + \frac{1}{\rho}\right) + b_D = \frac{D'}{D}(\sigma) + \frac{\log N\mathfrak{q}_D}{2} + \frac{R}{\sigma - 1} + \sum_{\substack{1 \leq j \leq J \\ \pi_j = \pi'_j}} \left(\frac{1}{\sigma + it_j - 1} + \frac{1}{\sigma + it_j}\right) + \frac{D'_\infty}{D_\infty}(\sigma).$$

Since $\operatorname{Re}(b_D) = -\sum_{\Delta(\rho)=0} \operatorname{Re}(\rho^{-1})$ [25, Proposition 5.7(3)], we take real parts and obtain

$$\sum_{\Delta(\rho)=0} \operatorname{Re}\left(\frac{1}{\sigma - \rho}\right) = \operatorname{Re}\left(\frac{D'}{D}(s) + \frac{\log N\mathfrak{q}_D}{2} + \frac{R}{\sigma - 1} + \sum_{\substack{1 \leq j \leq J \\ \pi_j = \pi'_j \\ t_j \neq 0}} \frac{\sigma - 1}{(\sigma - 1)^2 + t_j^2} + \frac{D'_\infty}{D_\infty}(s)\right) + O(1).$$

By (3.3), (3.7), and (3.10), the bound $\operatorname{Re}\left(\frac{D'}{D}(\sigma)\right) \ll \log Q$ (resp. $\frac{D'_\infty}{D_\infty}(\sigma) \ll \log Q$) follows from our hypothesis on the Dirichlet coefficients of $-\frac{D'}{D}(s)$ (resp. Stirling's formula). \square

3.3. Rankin–Selberg combinatorics. A partition $\mu = (\mu_i)_{i=1}^\infty$ is a sequence of nonincreasing nonnegative integers $\mu_1 \geq \mu_2 \geq \dots$ with only finitely many nonzero entries. For a partition μ , let $\ell(\mu)$ be the number of nonzero μ_i , and let $|\mu| = \sum_{i=1}^\infty \mu_i$. For a set $\{\alpha_1, \dots, \alpha_n\}$ of real numbers and a partition μ with $\ell(\mu) \leq n$, let $s_\mu(\{\alpha_1, \dots, \alpha_n\})$ be the Schur polynomial $\det[(\alpha_i^{\lambda(j)+n-j})_{ij}] / \det[(\alpha_i^{n-j})_{ij}]$ associated to μ . If $|\mu| = 0$, then $s_\mu(\{\alpha_1, \dots, \alpha_n\})$ is identically one. By convention, if $\ell(\mu) > n$, then $s_\mu(\{\alpha_1, \dots, \alpha_n\})$ is identically zero.

Let $\pi \in \mathfrak{F}_n$ and $\pi' \in \mathfrak{F}_{n'}$. By (3.5), (3.6), and Cauchy's identity [12, Theorem 38.1], we have

$$\sum_{k=0}^\infty \frac{\lambda_{\pi \times \pi'}(\mathfrak{p}^k)}{N\mathfrak{p}^{ks}} = L(s, \pi_\mathfrak{p} \times \pi'_\mathfrak{p}) = \sum_{\mu} \frac{s_\mu(A_\pi(\mathfrak{p}))s_\mu(A_{\pi'}(\mathfrak{p}))}{N\mathfrak{p}^{s|\mu|}}, \quad \mathfrak{p} \nmid \mathfrak{q}_\pi \mathfrak{q}_{\pi'},$$

where the sum ranges over all partitions. This yields

$$\lambda_{\pi \times \pi'}(\mathfrak{p}^k) = \sum_{|\mu|=k} s_\mu(A_\pi(\mathfrak{p}))s_\mu(A_{\pi'}(\mathfrak{p})), \quad \mathfrak{p} \nmid \mathfrak{q}_\pi \mathfrak{q}_{\pi'}.$$

For an integral ideal \mathfrak{n} with factorization $\mathfrak{n} = \prod_{\mathfrak{p}} \mathfrak{p}^{\operatorname{ord}_{\mathfrak{p}}(\mathfrak{n})}$ (with $\operatorname{ord}_{\mathfrak{p}}(\mathfrak{n}) = 0$ for all but finitely many \mathfrak{p}), the multiplicativity of $\lambda_{\pi \times \pi'}(\mathfrak{n})$ tells us that if $\gcd(\mathfrak{n}, \mathfrak{q}_\pi \mathfrak{q}_{\pi'}) = \mathcal{O}_F$, then

$$(3.14) \quad \lambda_{\pi \times \pi'}(\mathfrak{n}) = \prod_{\mathfrak{p}} \lambda_{\pi \times \pi'}(\mathfrak{p}^{\operatorname{ord}_{\mathfrak{p}}(\mathfrak{n})}) = \sum_{(\mu_{\mathfrak{p}})_{\mathfrak{p}} \in \underline{\mu}[\mathfrak{n}]} \prod_{\mathfrak{p}} s_{\mu_{\mathfrak{p}}}(A_\pi(\mathfrak{p}))s_{\mu_{\mathfrak{p}}}(A_{\pi'}(\mathfrak{p})),$$

where $(\mu_{\mathfrak{p}})_{\mathfrak{p}}$ denotes a sequence of partitions indexed by prime ideals and

$$(3.15) \quad \underline{\mu}[\mathfrak{n}] := \{(\mu_{\mathfrak{p}})_{\mathfrak{p}} : |\mu_{\mathfrak{p}}| = \operatorname{ord}_{\mathfrak{p}}(\mathfrak{n}) \text{ for all } \mathfrak{p}\}.$$

Define the numbers $\mu_{\pi \times \pi'}(\mathbf{n})$ on unramified prime powers by

$$\sum_{k=0}^{\infty} \frac{\mu_{\pi \times \pi'}(\mathfrak{p}^k)}{N\mathfrak{p}^{ks}} = L(s, \pi_{\mathfrak{p}} \times \pi'_{\mathfrak{p}})^{-1} = \prod_{j=1}^n \prod_{j'=1}^{n'} (1 - \alpha_{j,\pi}(\mathfrak{p}) \alpha_{j',\pi'}(\mathfrak{p}) N\mathfrak{p}^{-s}), \quad \mathfrak{p} \nmid \mathfrak{q}_{\pi} \mathfrak{q}_{\pi'}.$$

By multiplicativity, this defines $\mu_{\pi \times \pi'}(\mathbf{n})$ when $\gcd(\mathbf{n}, \mathfrak{q}_{\pi} \mathfrak{q}_{\pi'}) = \mathcal{O}_F$. For a partition $\mu = (\mu_i)_{i=1}^{\infty}$, let $\mu^* = (\mu_i^*)_{i=1}^{\infty}$ be the dual partition defined by $\mu_i^* = |\{j: \mu_j \geq i\}|$. It follows from the dual Cauchy identity [12, Chapter 38] and (3.6) that

$$\sum_{k=0}^{\infty} \frac{\mu_{\pi \times \pi'}(\mathfrak{p}^k)}{N\mathfrak{p}^{ks}} = \sum_{\mu} \frac{s_{\mu}(A_{\pi}(\mathfrak{p})) s_{\mu^*}(-A_{\pi'}(\mathfrak{p}))}{N\mathfrak{p}^{|\mu|s}}, \quad \mathfrak{p} \nmid \mathfrak{q}_{\pi} \mathfrak{q}_{\pi'},$$

where $-A_{\pi'}(\mathfrak{p}) = \{-\alpha_{1,\pi'}(\mathfrak{p}), \dots, -\alpha_{n,\pi'}(\mathfrak{p})\}$. Hence we have

$$(3.16) \quad \mu_{\pi \times \pi'}(\mathbf{n}) = \sum_{(\mu_{\mathfrak{p}})_{\mathfrak{p}} \in \underline{\mu}[\mathbf{n}]} \prod_{\mathfrak{p}} s_{\mu_{\mathfrak{p}}}(A_{\pi}(\mathfrak{p})) s_{\mu_{\mathfrak{p}}^*}(-A_{\pi'}(\mathfrak{p})), \quad \gcd(\mathbf{n}, \mathfrak{q}_{\pi} \mathfrak{q}_{\pi'}) = \mathcal{O}_F.$$

Lemma 3.3. *If $\gcd(\mathbf{n}, \mathfrak{q}_{\pi} \mathfrak{q}_{\pi'}) = \mathcal{O}_F$, then we have $|\mu_{\pi \times \pi'}(\mathbf{n})| \leq \frac{1}{2}(\lambda_{\pi \times \tilde{\pi}}(\mathbf{n}) + \lambda_{\pi' \times \tilde{\pi}'}(\mathbf{n}))$.*

Proof. We apply the inequality of arithmetic and geometric means to (3.16):

$$|\mu_{\pi \times \pi'}(\mathbf{n})| \leq \frac{1}{2} \left(\sum_{(\mu_{\mathfrak{p}})_{\mathfrak{p}} \in \underline{\mu}[\mathbf{n}]} \left| \prod_{\mathfrak{p}} s_{\mu_{\mathfrak{p}}}(A_{\pi}(\mathfrak{p})) \right|^2 + \sum_{(\mu_{\mathfrak{p}})_{\mathfrak{p}} \in \underline{\mu}[\mathbf{n}]} \left| \prod_{\mathfrak{p}} s_{\mu_{\mathfrak{p}}^*}(-A_{\pi'}(\mathfrak{p})) \right|^2 \right).$$

The first sum equals $\lambda_{\pi \times \tilde{\pi}}(\mathbf{n})$ by (3.14). For the second sum, note that since $|\mu| = |\mu^*|$, we have $(\mu_{\mathfrak{p}})_{\mathfrak{p}} \in \underline{\mu}[\mathbf{n}]$ if and only if $(\mu_{\mathfrak{p}}^*)_{\mathfrak{p}} \in \underline{\mu}[\mathbf{n}]$. Hence by rearranging, we see that

$$(3.17) \quad \sum_{(\mu_{\mathfrak{p}})_{\mathfrak{p}} \in \underline{\mu}[\mathbf{n}]} \left| \prod_{\mathfrak{p}} s_{\mu_{\mathfrak{p}}^*}(-A_{\pi'}(\mathfrak{p})) \right|^2 = \sum_{(\mu_{\mathfrak{p}})_{\mathfrak{p}} \in \underline{\mu}[\mathbf{n}]} \left| \prod_{\mathfrak{p}} s_{\mu_{\mathfrak{p}}}(-A_{\pi'}(\mathfrak{p})) \right|^2 = \lambda_{\pi' \times \tilde{\pi}'}(\mathbf{n}),$$

where the last equality holds because $\alpha_{j,\pi'}(\mathfrak{p}) \overline{\alpha_{j',\pi'}(\mathfrak{p})} = (-\alpha_{j,\pi'}(\mathfrak{p})) \overline{(-\alpha_{j',\pi'}(\mathfrak{p}))}$. \square

4. A NEW MEAN VALUE ESTIMATE

Our proof of Theorem 1.1 uses the following new mean value estimate for the Dirichlet coefficients of $L(s, \pi \times \pi')$ and $L(s, \pi \times \pi')^{-1}$. Let $\mathcal{S} \subseteq \mathfrak{F}_n$, and let $\mathcal{S}(Q) = \{\pi \in \mathcal{S}: C(\pi) \leq Q\}$.

Theorem 4.1. *Let b be a complex-valued function supported on the integral ideals of \mathcal{O}_F . Let $n, n' \geq 1$, and let $\pi' \in \mathfrak{F}_{n'}$. Let $Q, T \geq 1$, $\varepsilon > 0$, and $x \geq 1$. Both*

$$\sum_{\pi \in \mathcal{S}(Q)} \left| \sum_{\substack{N\mathfrak{n} \in (x, xe^{1/T}] \\ \gcd(\mathfrak{n}, \mathfrak{q}_{\pi} \mathfrak{q}_{\pi'}) = \mathcal{O}_F}} \mu_{\pi \times \pi'}(\mathfrak{n}) b(\mathfrak{n}) \right|^2 \quad \text{and} \quad \sum_{\pi \in \mathcal{S}(Q)} \left| \sum_{\substack{N\mathfrak{n} \in (x, xe^{1/T}] \\ \gcd(\mathfrak{n}, \mathfrak{q}_{\pi} \mathfrak{q}_{\pi'}) = \mathcal{O}_F}} \lambda_{\pi \times \pi'}(\mathfrak{n}) b(\mathfrak{n}) \right|^2$$

are

$$\ll_{\varepsilon} Q^{\varepsilon} \left(\frac{x}{T} + Q^{4n^2\theta_n + n} T^{\frac{1}{2}n^2[F:\mathbb{Q}] + \varepsilon} |\mathcal{S}(Q)| \right) \sum_{\substack{N\mathfrak{n} \in (x, xe^{1/T}] \\ \gcd(\mathfrak{n}, \mathfrak{q}_{\pi'}) = \mathcal{O}_F}} \lambda_{\pi' \times \tilde{\pi}'}(\mathfrak{n}) |b(\mathfrak{n})|^2,$$

where $\theta_n \in [0, \frac{1}{2} - \frac{1}{n^2+1}]$ is the best known exponent towards GRC for $\pi \in \mathcal{S}$.

Remark 4.2. If $\pi' = \mathbb{1}$ and $\mathcal{S} = \mathfrak{F}_n$, then Theorem 4.1 recovers [47, Theorem 1.1]³:

$$\sum_{\pi \in \mathfrak{F}_n(Q)} \left| \sum_{\substack{N\mathbf{n} \in (x, xe^{1/T}] \\ \gcd(\mathbf{n}, \mathbf{q}_\pi) = \mathcal{O}_F}} \lambda_\pi(\mathbf{n}) b(\mathbf{n}) \right|^2 \ll_\varepsilon Q^\varepsilon \left(\frac{x}{T} + Q^{4n^2\theta_n + n} T^{\frac{1}{2}n^2[F:\mathbb{Q}] + \varepsilon} |\mathfrak{F}_n(Q)| \right) \sum_{\substack{N\mathbf{n} \in (x, xe^{1/T}] \\ \gcd(\mathbf{n}, \mathbf{q}_{\pi'}) = \mathcal{O}_F}} |b(\mathbf{n})|^2.$$

One could try to prove Theorem 4.1 starting with this, replacing $b(\mathbf{n})$ with $\lambda_{\pi'}(\mathbf{n})\mu_F(\mathbf{n})b(\mathbf{n})$ (where $\mu_F(\mathbf{n})$ is the \mathbf{n} -th Dirichlet coefficient of $\zeta_F(s)^{-1}$), and try to recover a version of Theorem 4.1 with $\mu_{\pi \times \pi'}(\mathbf{n})$ replaced by $\lambda_\pi(\mathbf{n})\lambda_{\pi'}(\mathbf{n})\mu_F(\mathbf{n})$. Note that $\mu_{\pi \times \pi'}(\mathbf{n}) = \lambda_\pi(\mathbf{n})\lambda_{\pi'}(\mathbf{n})\mu_F(\mathbf{n})$ when \mathbf{n} is squarefree and coprime to $\mathbf{q}_\pi\mathbf{q}_{\pi'}$. Otherwise, equality is not guaranteed. If one wants to approximate $L(s, \pi \times \pi')$ with $\sum_{\mathbf{n} \text{ squarefree}} \lambda_\pi(\mathbf{n})\lambda_{\pi'}(\mathbf{n})N\mathbf{n}^{-s}$ and extend into the critical strip when $\pi, \pi' \in \mathfrak{F}_n$ and $n \geq 5$, then one must have progress towards GRC well beyond what is known unconditionally [9, 15]. Such progress would then be a hypothesis for Theorem 1.1.

Theorem 4.1 provides the first nontrivial unconditional mean value estimates of large sieve type for the Dirichlet coefficients $\lambda_{\pi \times \pi'}(\mathbf{n})$ or $\mu_{\pi \times \pi'}(\mathbf{n})$ for arbitrary n and n' . Theorem 4.1 follows from a more general result, Proposition 4.3 below, for sequences of products of Schur polynomials evaluated on the set $A_\pi(\mathfrak{p})$ of Satake parameters of π at \mathfrak{p} . We begin with a mean value estimate for the Satake parameters of π as $\pi \in \mathcal{S}$ varies. Let

$$(4.1) \quad a : \bigcup_{x < N\mathbf{n} \leq xe^{1/T}} \underline{\mu}[\mathbf{n}] \rightarrow \mathbb{C}, \quad \alpha : \mathcal{S}(Q) \rightarrow \mathbb{C}$$

be functions that are not identically zero. Their ℓ^2 norms $\|a\|_2$ and $\|\alpha\|_2$ are defined by

$$\|a\|_2 = \left(\sum_{x < N\mathbf{n} \leq xe^{1/T}} \sum_{(\mu_{\mathfrak{p}})_{\mathfrak{p}} \in \underline{\mu}[\mathbf{n}]} |a((\mu_{\mathfrak{p}})_{\mathfrak{p}})|^2 \right)^{1/2}, \quad \|\alpha\|_2 = \left(\sum_{\pi \in \mathcal{S}(Q)} |\alpha(\pi)|^2 \right)^{1/2}.$$

For convenience, we define $\mathbf{1}_{(\mathbf{n}, \mathbf{q})}$ to equal one when $\gcd(\mathbf{n}, \mathbf{q}) = \mathcal{O}_F$ and zero otherwise.

Proposition 4.3. *Let $x \geq 1$ and $Q, T \geq 1$. Define*

$$C(Q, T, x) := \sup_{\|a\|_2 \neq 0} \frac{1}{\|a\|_2^2} \sum_{\pi \in \mathcal{S}(Q)} \left| \sum_{\substack{x < N\mathbf{n} \leq xe^{1/T} \\ \gcd(\mathbf{n}, \mathbf{q}_\pi) = \mathcal{O}_F}} \sum_{(\mu_{\mathfrak{p}})_{\mathfrak{p}} \in \underline{\mu}[\mathbf{n}]} \left[\prod_{\mathfrak{p}} s_{\mu_{\mathfrak{p}}}(A_\pi(\mathfrak{p})) \right] a((\mu_{\mathfrak{p}})_{\mathfrak{p}}) \right|^2.$$

We have the bound $C(Q, T, x) \ll_\varepsilon Q^\varepsilon (T^{-1}x + Q^{4n^2\theta_n + n} T^{\frac{n^2[F:\mathbb{Q}]}{2} + \varepsilon} |\mathcal{S}(Q)|)$.

Proof. We observe that

$$(4.2) \quad C(Q, T, x) = \sup_{\|a\|_2 = 1} \sum_{\pi \in \mathcal{S}(Q)} \left| \sum_{x < N\mathbf{n} \leq xe^{1/T}} \sum_{(\mu_{\mathfrak{p}})_{\mathfrak{p}} \in \underline{\mu}[\mathbf{n}]} \left[\prod_{\mathfrak{p}} s_{\mu_{\mathfrak{p}}}(A_\pi(\mathfrak{p})) \right] \mathbf{1}_{(\mathbf{n}, \mathbf{q}_\pi)} a((\mu_{\mathfrak{p}})_{\mathfrak{p}}) \right|^2.$$

By the duality principle for bilinear forms, (4.2) is bounded by the supremum over the functions $\alpha : \mathcal{S}(Q) \rightarrow \mathbb{C}$ such that $\|\alpha\|_2 = 1$ of

$$(4.3) \quad \sum_{x < N\mathbf{n} \leq xe^{1/T}} \sum_{(\mu_{\mathfrak{p}})_{\mathfrak{p}} \in \underline{\mu}[\mathbf{n}]} \left| \sum_{\pi \in \mathcal{S}(Q)} \left[\prod_{\mathfrak{p}} s_{\mu_{\mathfrak{p}}}(A_\pi(\mathfrak{p})) \right] \mathbf{1}_{(\mathbf{n}, \mathbf{q}_\pi)} \alpha(\pi) \right|^2.$$

³The factor $Q^{4n^2\theta_n + n}$ fixes a minor error in the proof of [47, Theorem 1.1], which led to a factor of $Q^{n^2 + n}$ instead.

Let ϕ be a fixed smooth test function, supported in a compact subset of $[-2, 2]$, such that $\phi(t) = 1$ for $t \in [0, 1]$ and $\phi(t) \in [0, 1)$ otherwise. Then (4.3) is at most

$$(4.4) \quad \sum_{\mathfrak{n}} \sum_{(\mu_{\mathfrak{p}})_{\mathfrak{p}} \in \underline{\mu}[\mathfrak{n}]} \left| \sum_{\pi \in \mathcal{S}(Q)} \left[\prod_{\mathfrak{p}} s_{\mu_{\mathfrak{p}}}(A_{\pi}(\mathfrak{p})) \right] \mathbf{1}_{(n, q_{\pi})} \alpha(\pi) \right|^2 \phi \left(T \log \frac{N\mathfrak{n}}{x} \right).$$

We expand the square, interchange the order of summation, and apply (3.14) to find that (4.4) equals

$$(4.5) \quad \begin{aligned} & \sum_{\pi, \pi' \in \mathcal{S}(Q)} \alpha(\pi) \overline{\alpha(\pi')} \sum_{\mathfrak{n}} \sum_{(\mu_{\mathfrak{p}})_{\mathfrak{p}} \in \underline{\mu}[\mathfrak{n}]} \left[\prod_{\mathfrak{p}} s_{\mu_{\mathfrak{p}}}(A_{\pi}(\mathfrak{p})) \right] \overline{\left[\prod_{\mathfrak{p}} s_{\mu_{\mathfrak{p}}}(A_{\pi'}(\mathfrak{p})) \right]} \mathbf{1}_{(n, q_{\pi})} \mathbf{1}_{(n, q_{\pi'})} \phi \left(T \log \frac{N\mathfrak{n}}{x} \right) \\ &= \sum_{\pi, \pi' \in \mathcal{S}(Q)} \alpha(\pi) \overline{\alpha(\pi')} \sum_{\gcd(n, q_{\pi} q_{\pi'}) = \mathcal{O}_F} \lambda_{\pi \times \tilde{\pi}'}(\mathfrak{n}) \phi \left(T \log \frac{N\mathfrak{n}}{x} \right). \end{aligned}$$

Let $\kappa_{\pi \times \pi'} = \text{Res}_{s=1} L(s, \pi \times \pi') \prod_{\mathfrak{p} | q_{\pi} q_{\pi'}} L(s, \pi_{\mathfrak{p}} \times \pi'_{\mathfrak{p}})^{-1}$. Note that $\kappa_{\pi \times \pi'} \geq 0$, with equality if and only if $\pi' \neq \tilde{\pi}$. Since $|\alpha_{j, j', \pi \times \pi'}(\mathfrak{p})| \leq N\mathfrak{p}$, we have the bound

$$\prod_{\mathfrak{p} | q_{\pi} q_{\pi'}} |L(1, \pi_{\mathfrak{p}} \times \pi'_{\mathfrak{p}})|^{-1} = \prod_{\mathfrak{p} | q_{\pi} q_{\pi'}} \prod_{j=1}^n \prod_{j'=1}^{n'} \left| 1 - \frac{\alpha_{j, j', \pi \times \pi'}(\mathfrak{p})}{N\mathfrak{p}} \right| \leq \prod_{\mathfrak{p} | q_{\pi} q_{\pi'}} 2^{n'n}.$$

Since $|\{\mathfrak{p} : \mathfrak{p} \mid \mathfrak{n}\}| \ll (\log N\mathfrak{n}) / \log \log N\mathfrak{n}$ [48, Lemma 1.13b], it follows from (3.12) that

$$(4.6) \quad \kappa_{\pi \times \tilde{\pi}} \ll_{\varepsilon} C(\pi)^{\varepsilon}.$$

Let $\hat{\phi}(s) = \int_{\mathbb{R}} \phi(y) e^{sy} dy$. It follows from a standard contour integral calculation using (3.7) and (3.13) that (4.5) equals

$$(4.7) \quad \begin{aligned} & \sum_{\pi, \pi' \in \mathcal{S}(Q)} \alpha(\pi) \overline{\alpha(\pi')} \left(\frac{1}{2\pi i T} \int_{3-i\infty}^{3+i\infty} \frac{L(s, \pi \times \tilde{\pi}')}{\prod_{\mathfrak{p} | q_{\pi} q_{\pi'}} L(s, \pi_{\mathfrak{p}} \times \tilde{\pi}'_{\mathfrak{p}})} x^s \hat{\phi}(s/T) ds \right) \\ &= \sum_{\pi, \pi' \in \mathcal{S}(Q)} \alpha(\pi) \overline{\alpha(\pi')} \left(\kappa_{\pi \times \tilde{\pi}'} x \frac{\hat{\phi}(\frac{1}{T})}{T} + \frac{1}{2\pi i T} \int_{\frac{1}{4 \log(\varepsilon x)} - i\infty}^{\frac{1}{4 \log(\varepsilon x)} + i\infty} \frac{L(s, \pi \times \tilde{\pi}')}{\prod_{\mathfrak{p} | q_{\pi} q_{\pi'}} L(s, \pi_{\mathfrak{p}} \times \tilde{\pi}'_{\mathfrak{p}})} x^s \hat{\phi}(s/T) ds \right) \\ &= \sum_{\pi, \pi' \in \mathcal{S}(Q)} \alpha(\pi) \overline{\alpha(\pi')} \left(\kappa_{\pi \times \tilde{\pi}'} x \frac{\hat{\phi}(\frac{1}{T})}{T} + O_{\phi, \varepsilon} \left(Q^{4n^2 \theta_n + n + \varepsilon} T^{\frac{1}{2} n^2 [F:\mathbb{Q}] + \varepsilon} \right) \right). \end{aligned}$$

Recall that $\kappa_{\pi \times \tilde{\pi}'} > 0$ when $\pi = \pi'$, and $\kappa_{\pi \times \tilde{\pi}'} = 0$ otherwise. Since $\|\alpha\|_2 = 1$ and ϕ is fixed, it follows from the inequality of arithmetic and geometric means that (4.7) equals

$$\begin{aligned} & \frac{\hat{\phi}(\frac{1}{T})}{T} x \sum_{\pi \in \mathcal{S}(Q)} |\alpha(\pi)|^2 \kappa_{\pi \times \tilde{\pi}'} + O_{\phi} \left(Q^{4n^2 \theta_n + n + \varepsilon} T^{\frac{1}{2} n^2 [F:\mathbb{Q}] + \varepsilon} |\mathcal{S}(Q)| \right) \\ & \ll_{\phi, \varepsilon} \frac{x}{T} \max_{\pi \in \mathcal{S}(Q)} \kappa_{\pi \times \tilde{\pi}'} + Q^{4n^2 \theta_n + n + \varepsilon} T^{\frac{1}{2} n^2 [F:\mathbb{Q}] + \varepsilon} |\mathcal{S}(Q)|. \end{aligned}$$

We estimate the maximum using (4.6), and the desired result follows. \square

We use Proposition 4.3 to prove Theorem 4.1.

Proof of Theorem 4.1. For the sum involving $\mu_{\pi \times \pi'}(\mathbf{n})$, we apply Proposition 4.3 with

$$a((\mu_{\mathfrak{p}})_{\mathfrak{p}}) = b\left(\prod_{\mathfrak{p}} N\mathfrak{p}^{|\mu_{\mathfrak{p}}|}\right) \mathbf{1}_{(\prod_{\mathfrak{p}} N\mathfrak{p}^{|\mu_{\mathfrak{p}}|}, q_{\pi'})} \prod_{\mathfrak{p}} s_{\mu_{\mathfrak{p}}}^*(-A_{\pi'}(\mathfrak{p})).$$

If $(\mu_{\mathfrak{p}})_{\mathfrak{p}} \in \underline{\mu}[\mathbf{n}]$, then by (3.15), we have that $|\mu_{\mathfrak{p}}| = \text{ord}_{\mathfrak{p}}(\mathbf{n})$ and

$$a((\mu_{\mathfrak{p}})_{\mathfrak{p}}) = b(\mathbf{n}) \mathbf{1}_{(n, q_{\pi'})} \prod_{\mathfrak{p}} s_{\mu_{\mathfrak{p}}}^*(-A_{\pi'}(\mathfrak{p})).$$

By (3.16), the left-hand side of Proposition 4.3 becomes

$$\begin{aligned} & \sum_{\pi \in \mathcal{S}(Q)} \left| \sum_{\substack{x < N\mathbf{n} \leq xe^{1/T} \\ \gcd(n, q_{\pi}) = \mathcal{O}_F}} \sum_{(\mu_{\mathfrak{p}})_{\mathfrak{p}} \in \underline{\mu}[\mathbf{n}]} \left[\prod_{\mathfrak{p}} s_{\mu_{\mathfrak{p}}}(A_{\pi}(\mathfrak{p})) \right] a((\mu_{\mathfrak{p}})_{\mathfrak{p}}) \right|^2 \\ &= \sum_{\pi \in \mathcal{S}(Q)} \left| \sum_{\substack{x < N\mathbf{n} \leq xe^{1/T} \\ \gcd(n, q_{\pi}) = \mathcal{O}_F}} b(\mathbf{n}) \mathbf{1}_{(n, q_{\pi'})} \sum_{(\mu_{\mathfrak{p}})_{\mathfrak{p}} \in \underline{\mu}[\mathbf{n}]} \prod_{\mathfrak{p}} s_{\mu_{\mathfrak{p}}}(A_{\pi}(\mathfrak{p})) s_{\mu_{\mathfrak{p}}}^*(-A_{\pi'}(\mathfrak{p})) \right|^2 \\ &= \sum_{\pi \in \mathcal{S}(Q)} \left| \sum_{\substack{x < N\mathbf{n} \leq xe^{1/T} \\ \gcd(n, q_{\pi} q_{\pi'}) = \mathcal{O}_F}} \mu_{\pi \times \pi'}(\mathbf{n}) b(\mathbf{n}) \right|^2. \end{aligned}$$

Similarly, the right-hand side of Proposition 4.3 becomes

$$Q^{\varepsilon} \left(\frac{x}{T} + Q^{4n^2\theta_n + n} T^{\frac{1}{2}n^2[F:\mathbb{Q}] + \varepsilon} |\mathcal{S}(Q)| \right) \sum_{\substack{N\mathbf{n} \in (x, xe^{1/T}] \\ \gcd(n, q_{\pi'}) = \mathcal{O}_F}} |b(\mathbf{n})|^2 \sum_{(\mu_{\mathfrak{p}})_{\mathfrak{p}} \in \underline{\mu}[\mathbf{n}]} \left| \prod_{\mathfrak{p}} s_{\mu_{\mathfrak{p}}}^*(-A_{\pi'}(\mathfrak{p})) \right|^2.$$

The desired result now follows from (3.17). Apart from the new choice of

$$a((\mu_{\mathfrak{p}})_{\mathfrak{p}}) = b\left(\prod_{\mathfrak{p}} N\mathfrak{p}^{|\mu_{\mathfrak{p}}|}\right) \mathbf{1}_{(\prod_{\mathfrak{p}} N\mathfrak{p}^{|\mu_{\mathfrak{p}}|}, q_{\pi'})} \prod_{\mathfrak{p}} s_{\mu_{\mathfrak{p}}}(A_{\pi'}(\mathfrak{p})),$$

the result for the sum involving $\lambda_{\pi \times \pi'}(\mathbf{n})$ is handled in the same manner. \square

Corollary 4.4. *Let $\pi' \in \mathfrak{F}_{\mathbf{n}'}$, $Q, T \geq 1$, and $\varepsilon > 0$. If $Y \geq e$ and*

$$X \geq Q^{4n^2\theta_n + n + \varepsilon} T^{\frac{1}{2}n^2[F:\mathbb{Q}] + 1 + \varepsilon} |\mathcal{S}(Q)|, \quad \log Y \asymp_{\varepsilon} \log X,$$

then

$$\begin{aligned} & \sum_{\pi \in \mathcal{S}(Q)} \int_{-T}^T \left| \sum_{\substack{N\mathbf{n} > X \\ \gcd(n, q_{\pi} q_{\pi'}) = \mathcal{O}_F}} \frac{\mu_{\pi \times \pi'}(\mathbf{n})}{N\mathbf{n}^{1 + \frac{1}{\log Y} + iv}} \right|^2 dv \ll_{\varepsilon} C(\pi')^{\varepsilon} Q^{\varepsilon} \log X, \\ & \sum_{\pi \in \mathcal{S}(Q)} \int_{-T}^T \left| \sum_{\substack{N\mathbf{n} \leq X \\ \gcd(n, q_{\pi} q_{\pi'}) = \mathcal{O}_F}} \frac{\mu_{\pi \times \pi'}(\mathbf{n})}{N\mathbf{n}^{\frac{1}{2} + iv}} \right|^2 dv \ll_{\varepsilon} C(\pi')^{\varepsilon} Q^{\varepsilon} X \log X. \end{aligned}$$

Proof. We prove the first result; the second result is proved completely analogously. A formal generalization of [17, Theorem 1] to number fields tells us that if $c(\mathbf{n})$ is a complex-valued function supported on the integral ideals of \mathcal{O}_F with $\sum_{\mathbf{n}} |c(\mathbf{n})| < \infty$, then

$$\int_{-T}^T \left| \sum_{\mathbf{n}} \frac{c(\mathbf{n})}{N\mathbf{n}^{it}} \right|^2 dt \ll T^2 \int_0^{\infty} \left| \sum_{N\mathbf{n} \in (x, xe^{1/T}]} c(\mathbf{n}) \right|^2 \frac{dx}{x}.$$

We choose $b(\mathbf{n}) = \mathbf{Nn}^{-1-\frac{1}{\log Y}}$ if $\mathbf{Nn} > X$ and $b(\mathbf{n}) = 0$ otherwise, which leads to

$$\sum_{\pi \in \mathcal{S}(Q)} \int_{-T}^T \left| \sum_{\substack{\mathbf{Nn} > X \\ \gcd(\mathbf{n}, \mathfrak{q}_\pi \mathfrak{q}_{\pi'}) = \mathcal{O}_F}} \frac{\mu_{\pi \times \pi'}(\mathbf{n})}{\mathbf{Nn}^{1+\frac{1}{\log Y}+it}} \right|^2 dt \ll T^2 \int_0^\infty \sum_{\pi \in \mathcal{S}(Q)} \left| \sum_{\mathbf{Nn} \in (x, x\epsilon^{1/T})} \mu_{\pi \times \pi'}(\mathbf{n}) b(\mathbf{n}) \right|^2 \frac{dx}{x}.$$

Theorem 4.1 and the fact that $\lambda_{\pi' \times \tilde{\pi}'}(\mathbf{n}) \geq 0$ for all \mathbf{n} imply that the above display is

$$\ll_\varepsilon Q^\varepsilon \sum_{\substack{\mathbf{Nn} > X \\ \gcd(\mathbf{n}, \mathfrak{q}_{\pi'}) = \mathcal{O}_F}} \frac{\lambda_{\pi' \times \tilde{\pi}'}(\mathbf{n})}{\mathbf{Nn}^{1+\frac{2}{\log Y}}} \left(1 + \frac{Q^{4n^2\theta_n+n} T^{\frac{1}{2}n^2[F:\mathbb{Q}]+1+\varepsilon} |\mathcal{S}(Q)|}{\mathbf{Nn}} \right) \ll_\varepsilon Q^\varepsilon \sum_{\mathbf{Nn} > X} \frac{\lambda_{\pi' \times \tilde{\pi}'}(\mathbf{n})}{\mathbf{Nn}^{1+\frac{2}{\log Y}}}.$$

The desired result now follows from Lemma 3.1 and our choices of X and Y . \square

5. PROOF OF THEOREM 1.1

Let $n, n' \geq 1$, $\varepsilon > 0$, $T \geq 3$, $Q \geq 3$, $\pi \in \mathcal{S}(Q)$, and $\pi' \in \mathfrak{F}_{n'}$. Define

$$\begin{aligned} X &:= C(\pi')^\varepsilon Q^{4n^2\theta_n+n+\varepsilon} T^{\frac{1}{2}n^2[F:\mathbb{Q}]+1+\varepsilon} |\mathcal{S}(Q)|, \\ Y &:= \left((C(\pi')^{\frac{n}{2}} Q^{\frac{n'}{2}} (C(\pi')Q)^{2n'n \max\{0, \theta_n + \theta_{n'} - \frac{1}{2}\}} |\mathcal{S}(Q)| T^{\frac{n'n[F:\mathbb{Q}]}{2}+1})^{1+\frac{\varepsilon}{4n'n[F:\mathbb{Q}]}} X \right)^{\frac{1}{3-2\sigma}}, \\ L^{\text{wr}}(s, \pi \times \pi') &:= L(s, \pi \times \pi') \prod_{\mathfrak{p} | \mathfrak{q}_\pi \mathfrak{q}_{\pi'}} L(s, \pi_{\mathfrak{p}} \times \pi'_{\mathfrak{p}})^{-1}, \\ (5.1) \quad M_X(s, \pi \times \pi') &:= \sum_{\substack{\mathbf{Nn} \leq X \\ \gcd(\mathbf{n}, \mathfrak{q}_\pi \mathfrak{q}_{\pi'}) = \mathcal{O}_F}} \frac{\mu_{\pi \times \pi'}(\mathbf{n})}{\mathbf{Nn}^s}, \\ LM_X(s, \pi \times \pi') &:= L^{\text{wr}}(s, \pi \times \pi') M_X(s, \pi \times \pi'). \end{aligned}$$

Note that $\log(C(\pi)C(\pi')T) \ll \log X \asymp_\varepsilon \log Y$. We assume that $T \geq 2(\log Y)^2$ since the proof does not change appreciably otherwise. In this section, ε might vary from line to line, and terms of size $(C(\pi')QT^{[F:\mathbb{Q}]})^\varepsilon |\mathcal{S}(Q)|^\varepsilon$ and polynomials in $\log(C(\pi')QT^{[F:\mathbb{Q}]}) |\mathcal{S}(Q)|$ might be bounded by X^ε without mention.

If $t \in [-T, T]$, then

$$|\{\rho = \beta + i\gamma: \beta \geq 0, |\gamma - t| \leq 1\}| \ll \log C(\pi \times \pi', t) \ll \log(C(\pi)C(\pi')T)$$

by [25, Proposition 5.7]. We observe that the rectangle $[\sigma, 1] \times [-T, T]$ is covered by $O(T)$ boxes of the form $[\sigma, 1] \times [y, y + 2(\log Y)^2]$, each containing $O((\log(C(\pi)C(\pi')T))^3)$ zeros. If we write $n_{\pi \times \pi'}$ for the number of such boxes containing at least one zero ρ of $L(s, \pi \times \pi')$, then

$$N_{\pi \times \pi'}(\sigma, T) \ll (\log(C(\pi')QT))^{-3} n_{\pi \times \pi'}.$$

If $\pi' = \tilde{\pi}'$, then it suffices to assume that $|\rho - 1| > 1$ since there are $O(\log(C(\pi')QT))$ zeros ρ such that $|\rho - 1| \leq 1$.

Let $\rho = \beta + i\gamma$ be a zero of $L(s, \pi \times \pi')$, in which case $LM_X(\rho, \pi \times \pi') = 0$. We compute

$$\begin{aligned}
e^{-1/Y} &= \frac{1}{2\pi i} \int_{1-\beta+\frac{1}{\log Y}-i\infty}^{1-\beta+\frac{1}{\log Y}+i\infty} (1 - LM_X(\rho + w, \pi \times \pi'))\Gamma(w)Y^w dw \\
(5.2) \quad &+ \frac{1}{2\pi i} \int_{\frac{1}{2}-\beta-i\infty}^{\frac{1}{2}-\beta+i\infty} LM_X(\rho + w, \pi \times \pi')\Gamma(w)Y^w dw \\
&+ \kappa_{\pi \times \pi'}\Gamma(1 - \rho)Y^{1-\rho} \sum_{\substack{N\mathfrak{n} \leq X \\ \gcd(\mathfrak{n}, \mathfrak{q}_{\pi \times \pi'}) = \mathcal{O}_F}} \frac{\mu_{\pi \times \pi'}(\mathfrak{n})}{N\mathfrak{n}}.
\end{aligned}$$

It follows from (3.13) and Lemmata 3.1 and 3.3 that if $\operatorname{Re}(w) = 1 - \beta + \frac{1}{\log Y}$, then

$$\begin{aligned}
(5.3) \quad &|1 - LM_X(\rho + w, \pi \times \pi')| \\
&= |L^w(\rho + w, \pi \times \pi')| \cdot |L^w(\rho + w, \pi \times \pi')^{-1} - M_X(\rho + w, \pi \times \pi')| \\
&\ll_{\varepsilon} C(\pi \times \pi', |\gamma + \operatorname{Im}(w)| + 1)^{\varepsilon} \sum_{\substack{N\mathfrak{n} > X \\ \gcd(\mathfrak{n}, \mathfrak{q}_{\pi \times \pi'}) = \mathcal{O}_F}} \frac{|\mu_{\pi \times \pi'}(\mathfrak{n})|}{N\mathfrak{n}^{1+\frac{1}{\log Y}}} \\
&\ll_{\varepsilon} C(\pi \times \pi', |\gamma + \operatorname{Im}(w)| + 1)^{\varepsilon} \sum_{\substack{N\mathfrak{n} > X \\ \gcd(\mathfrak{n}, \mathfrak{q}_{\pi \times \pi'}) = \mathcal{O}_F}} \frac{\lambda_{\pi \times \pi}(\mathfrak{n}) + \lambda_{\pi' \times \pi'}(\mathfrak{n})}{N\mathfrak{n}^{1+\frac{1}{\log Y}}} \\
&\ll_{\varepsilon} C(\pi)^{\varepsilon} C(\pi')^{\varepsilon} (|\gamma + \operatorname{Im}(w)| + 1)^{\varepsilon} \log X.
\end{aligned}$$

Therefore, we deduce from Stirling's formula that

$$\begin{aligned}
&\left| \frac{1}{2\pi i} \int_{1-\beta+\frac{1}{\log Y}-i\infty}^{1-\beta+\frac{1}{\log Y}+i\infty} (1 - LM_X(\rho + w, \pi \times \pi'))\Gamma(w)Y^w dw \right. \\
&\quad \left. - \frac{1}{2\pi i} \int_{1-\beta+\frac{1}{\log Y}-i(\log Y)^2}^{1-\beta+\frac{1}{\log Y}+i(\log Y)^2} (1 - LM_X(\rho + w, \pi \times \pi'))\Gamma(w)Y^w dw \right| \ll \frac{1}{Y}.
\end{aligned}$$

The other terms in (5.2) are handled similarly. Since $e^{-1/Y} = 1 + O(1/Y)$, it follows that

$$\begin{aligned}
(5.4) \quad &1 \ll \frac{1}{2\pi i} \int_{1-\beta+\frac{1}{\log Y}-i(\log Y)^2}^{1-\beta+\frac{1}{\log Y}+i(\log Y)^2} (1 - LM_X(\rho + w, \pi \times \pi'))\Gamma(w)Y^w dw \\
&+ \frac{1}{2\pi i} \int_{\frac{1}{2}-\beta-i(\log Y)^2}^{\frac{1}{2}-\beta+i(\log Y)^2} LM_X(\rho + w, \pi \times \pi')\Gamma(w)Y^w dw + \kappa_{\pi \times \pi'} Y^{1-\sigma} \frac{\log X}{\max\{1, |\gamma|^3\}}.
\end{aligned}$$

Write the first integral in (5.4) as I_1 and the second as I_2 . A simple optimization calculation shows that if $c \geq 1$ and $c^{-1} \leq |I_1| + |I_2|$, then $c^{-1} \leq 2c(|I_1|^2 + |I_2|^2)$. We estimate $|I_1|^2$ using the Cauchy–Schwarz inequality, thus obtaining the bound

$$\begin{aligned}
1 &\ll (\log Y)^2 Y^{2(1-\sigma)} \int_{\gamma-(\log Y)^2}^{\gamma+(\log Y)^2} |1 - LM_X(1 + \frac{1}{\log Y} + iv, \pi \times \pi')|^2 dv \\
&+ Y^{\frac{1}{2}-\sigma} \int_{\gamma-(\log Y)^2}^{\gamma+(\log Y)^2} |LM_X(\frac{1}{2} + iv, \pi \times \pi')| dv + \kappa_{\pi \times \pi'} Y^{1-\sigma} \frac{\log X}{\max\{1, |\gamma|^3\}}.
\end{aligned}$$

Since $T \geq 2(\log Y)^2$ by hypothesis, we conclude that

$$(5.5) \quad \begin{aligned} n_{\pi \times \pi'} &\ll (\log Y)^2 Y^{2(1-\sigma)} \int_{-T}^T |1 - L(1 + \frac{1}{\log Y} + iv) M_X(1 + \frac{1}{\log Y} + iv)|^2 dv \\ &+ Y^{\frac{1}{2}-\sigma} \int_{-T}^T |L(\frac{1}{2} + iv) M_X(\frac{1}{2} + iv)| dv + \kappa_{\pi \times \pi'} Y^{1-\sigma} \log X + 1. \end{aligned}$$

Note that $\kappa_{\pi \times \pi'} = 0$ unless $\pi' = \tilde{\pi}$, which occurs for at most one $\pi \in \mathcal{S}(Q)$. When $\kappa_{\pi \times \pi'} \neq 0$, it satisfies $\kappa_{\pi \times \pi'} = \kappa_{\pi \times \tilde{\pi}} \ll_{\varepsilon} C(\pi)^{\varepsilon}$ per (4.6). Therefore, summing (5.5) over $\pi \in \mathcal{S}(Q)$, we find that

$$(5.6) \quad \begin{aligned} \sum_{\pi \in \mathcal{S}(Q)} N_{\pi \times \pi'}(\sigma, T) &\ll (\log C(\pi) C(\pi') T)^3 \sum_{\pi \in \mathcal{S}(Q)} n_{\pi \times \pi'} \\ &\ll_{\varepsilon} X^{\varepsilon} \left(Y^{2(1-\sigma)} \int_{-T}^T \sum_{\pi \in \mathcal{S}(Q)} |1 - LM_X(1 + \frac{1}{\log Y} + iv, \pi \times \pi')|^2 dv \right. \\ &\quad \left. + Y^{\frac{1}{2}-\sigma} \int_{-T}^T \sum_{\pi \in \mathcal{S}(Q)} |LM_X(\frac{1}{2} + iv, \pi \times \pi')| dv + Y^{1-\sigma} \right). \end{aligned}$$

For the first integral in (5.6), we deduce from (3.13) that

$$\begin{aligned} &|1 - LM_X(1 + \frac{1}{\log Y} + it, \pi \times \pi')|^2 \\ &= |L^{\text{ur}}(1 + \frac{1}{\log Y} + it, \pi \times \pi')|^2 \left| \frac{1}{L^{\text{ur}}(1 + \frac{1}{\log Y} + it, \pi \times \pi')} - M_X(1 + \frac{1}{\log Y} + it, \pi \times \pi') \right|^2 \\ &\ll_{\varepsilon} X^{\varepsilon} \left| \sum_{\substack{N\mathbf{n} > X \\ \gcd(\mathbf{n}, \mathfrak{q}_{\pi} \mathfrak{q}_{\pi'}) = \mathcal{O}_F}} \frac{\mu_{\pi \times \pi'}(\mathbf{n})}{N\mathbf{n}^{1+\frac{1}{\log Y}}} \right|^2. \end{aligned}$$

For the second integral in (5.6), it follows from the Cauchy–Schwarz inequality that

$$\begin{aligned} &\int_{-T}^T \sum_{\pi \in \mathcal{S}(Q)} |LM_X(\frac{1}{2} + iv, \pi \times \pi')| dv \\ &\leq \left(\int_{-T}^T \sum_{\pi \in \mathcal{S}(Q)} |L^{\text{ur}}(\frac{1}{2} + iv, \pi \times \pi')|^2 dv \right)^{\frac{1}{2}} \left(\int_{-T}^T \sum_{\pi \in \mathcal{S}(Q)} \left| \sum_{\substack{N\mathbf{n} \leq X \\ \gcd(\mathbf{n}, \mathfrak{q}_{\pi} \mathfrak{q}_{\pi'}) = \mathcal{O}_F}} \frac{\mu_{\pi \times \pi'}(\mathbf{n})}{N\mathbf{n}^{\frac{1}{2}+iv}} \right|^2 dv \right)^{\frac{1}{2}}. \end{aligned}$$

We bound the second moment of $L^{\text{ur}}(\frac{1}{2} + iv, \pi \times \pi')$ trivially as

$$\begin{aligned} &\int_{-T}^T \sum_{\pi \in \mathcal{S}(Q)} |L^{\text{ur}}(\frac{1}{2} + iv, \pi \times \pi')|^2 dv \\ &\ll_{\varepsilon} (QC(\pi'))^{2n' \max\{0, \theta_n + \theta_{n'} - \frac{1}{2}\} + \varepsilon} Q^{\frac{n'}{2}} C(\pi')^{\frac{n'}{2}} T^{\frac{n' n [F:\mathbb{Q}]}{2} + 1 + \varepsilon} |\mathcal{S}(Q)| \ll_{\varepsilon} \frac{Y^{3-2\sigma}}{X} \end{aligned}$$

using (3.7) (to bound the ramified Euler factors) and (3.13). In summary, we find that

$$\begin{aligned} \sum_{\pi \in \mathcal{S}(Q)} N_{\pi \times \pi'}(\sigma, T) &\ll_{\varepsilon} X^{\varepsilon} \left(Y^{2(1-\sigma)} \int_{-T}^T \sum_{\pi \in \mathcal{S}(Q)} \left| \sum_{\substack{N\mathfrak{n} > X \\ \gcd(\mathfrak{n}, \mathfrak{q}_{\pi} \mathfrak{q}_{\pi'}) = \mathcal{O}_F}} \frac{\mu_{\pi \times \pi'}(\mathfrak{n})}{N\mathfrak{n}^{1 + \frac{1}{\log Y} + iv}} \right|^2 dv \right. \\ &\quad \left. + Y^{\frac{1}{2} - \sigma} \left(\frac{Y^{3-2\sigma}}{X} \int_{-T}^T \sum_{\pi \in \mathcal{S}(Q)} \left| \sum_{\substack{N\mathfrak{n} \leq X \\ \gcd(\mathfrak{n}, \mathfrak{q}_{\pi} \mathfrak{q}_{\pi'}) = \mathcal{O}_F}} \frac{\mu_{\pi \times \pi'}(\mathfrak{n})}{N\mathfrak{n}^{\frac{1}{2} + iv}} \right|^2 dv \right)^{\frac{1}{2}} + Y^{1-\sigma} \right). \end{aligned}$$

We bound the two v -integrals using Corollary 4.4 and (5.1), thus obtaining

$$\sum_{\pi \in \mathcal{S}(Q)} N_{\pi \times \pi'}(\sigma, T) \ll_{\varepsilon} Y^{2(1-\sigma)} X^{\varepsilon}.$$

Theorem 1.1 now follows from (5.1), the bounds for θ_n and $\theta_{n'}$ in (3.2), and (1.5).

6. BOUNDS FOR RANKIN–SELBERG L -FUNCTIONS

In this section, we prove Theorem 2.1. Let $\pi' \in \mathfrak{F}_{n'}$ and $Q \geq 1$. In (1.6), we rescale ε to $\varepsilon/72$ and define $\alpha = \varepsilon/(7.2 \max\{n, n'\})$ so that

$$(6.1) \quad |\{\pi \in \mathfrak{F}_n(Q) : N_{\pi \times \pi'}(1 - \alpha, 6) = 0\}| \leq \sum_{\pi \in \mathfrak{F}_n(Q)} N_{\pi \times \pi'}(1 - \alpha, 6) \ll_{F, \varepsilon} (C(\pi')^n |\mathfrak{F}_n(Q)|)^{\varepsilon}.$$

Let $\pi \in \mathfrak{F}_n(Q)$. Proceeding as in the proof of [45, Theorem 1.1], we obtain⁴

$$\log |L(\tfrac{1}{2}, \pi \times \pi')| \leq \left(\frac{1}{4} - \frac{\alpha}{10^9} \right) \log C(\pi \times \pi') + 2 \log |L(\tfrac{3}{2}, \pi \times \pi')| + \frac{\alpha}{10^7} N_{\pi \times \pi'}(1 - \alpha, 6) + O(1).$$

By (3.12) and (3.13), we find that there exists an effectively computable constant $c_5 = c_5(n, n', [F : \mathbb{Q}], \varepsilon) > 0$ such that if $C(\pi \times \pi') \geq c_5$, then

$$(6.2) \quad \log |L(\tfrac{1}{2}, \pi \times \pi')| \leq \left(\frac{1}{4} - \frac{9\alpha}{10^{10}} \right) \log C(\pi \times \pi') + \frac{\alpha}{10^7} N_{\pi \times \pi'}(1 - \alpha, 6).$$

Theorem 2.1 now follows from (6.1) and (6.2).

7. EFFECTIVE MULTIPLICITY ONE ON AVERAGE

In this section, we prove Theorem 2.3. We consider the family of cuspidal automorphic representations $\mathfrak{F}_n(Q)$ over a number field F . Recall our convention that implied constants are allowed to depend on $n \geq 3$, $[F : \mathbb{Q}]$, and ε unless specifically mentioned otherwise.

To prove Theorem 2.3, we use Theorem 1.1 to build large zero-free regions for almost all L -functions $L(s, \pi \times \pi')$ with $\pi \in \mathfrak{F}_n$ varying. If $\pi' \in \{\tilde{\pi}, \tilde{\pi}'\}$, then $L(s, \pi \times \pi')$ has the standard zero-free region (1.1) apart from a possible Landau–Siegel zero, which must be both real and simple. In all other cases, only Brumley’s much narrower zero-free region is known ([8], [32, Appendix]). We can use Theorem 1.1 to establish a much stronger zero-free region for $L(s, \pi \times \pi')$ for all $\pi \in \mathfrak{F}_n(Q)$ with very few exceptions. Here is a simple example.

Corollary 7.1. *Let $\varepsilon > 0$, $n \geq 3$, and $\pi' \in \mathfrak{F}_n(Q)$. For all $\pi \in \mathfrak{F}_n(Q)$ with $O_{\varepsilon}(Q^{\varepsilon})$ exceptions, the L -function $L(s, \pi \times \pi')$ is nonzero in the region*

$$\operatorname{Re}(s) \geq 1 - \frac{\varepsilon}{20n^2}, \quad |\operatorname{Im}(s)| \leq \log Q.$$

⁴Small modifications are required to work over $F \neq \mathbb{Q}$, none of which substantially changes the proof.

Proof. This follows from (1.7) (once we rescale ε to $\varepsilon/21$) with $C(\pi') \leq Q$, $T = \log Q$, and $\sigma = 1 - \varepsilon/(20n^2)$. \square

For $\pi_1, \pi_2 \in \mathfrak{F}_n$ define the numbers $\Lambda_{\pi_1 \times \pi_2}(\mathbf{n})$ by the Dirichlet series identity (for $\operatorname{Re}(s) > 1$)

$$(7.1) \quad \sum_{\mathbf{n}} \frac{\Lambda_{\pi_1 \times \pi_2}(\mathbf{n})}{\mathbf{Nn}^s} = -\frac{L'}{L}(s, \pi_1 \times \pi_2) = \sum_{\mathfrak{p}} \sum_{k=1}^{\infty} \frac{\sum_{j=1}^n \sum_{j'=1}^n \alpha_{j,j',\pi_1 \times \pi_2}(\mathfrak{p})^k \log \mathbf{Np}}{\mathbf{Np}^{ks}}.$$

Lemma 7.2. *Let $\pi' \in \mathfrak{F}_n$. There exist absolute and effectively computable constants $c_6 \in (0, 1)$; $c_7, c_8, c_9, c_{10} \geq 1$; and $c_{11} \in (0, 1)$ such that the following are true.*

(1) *The L -function $L(s, \pi' \times \tilde{\pi}')$ has at most one zero, say β_1 , in the region*

$$\operatorname{Re}(s) \geq 1 - \frac{c_6}{\log(C(\pi')^n (|\operatorname{Im}(s)| + e)^{n^2[F:\mathbb{Q}]})}.$$

If β_1 exists, then it must be real and simple and satisfy $\beta_1 \leq 1 - C(\pi')^{-c_7 n}$.

(2) *If $A \geq c_8$, $\log \log C(\pi') \geq c_9 n^4 [F:\mathbb{Q}]^2$, and $x \geq C(\pi')^{c_{10} A^2 n^3 [F:\mathbb{Q}] \log(en[F:\mathbb{Q}])}$, then*

$$\sum_{\frac{x}{2} < \mathbf{Nn} \leq x} \Lambda_{\pi' \times \tilde{\pi}'}(\mathbf{n}) = \begin{cases} \frac{x}{2}(1 - \xi^{\beta_1 - 1})(1 + O(e^{-c_{11}A})) & \text{if } \beta_1 \text{ exists,} \\ \frac{x}{2}(1 + O(e^{-c_{11}A})) & \text{otherwise,} \end{cases}$$

where $\xi \in [\frac{x}{2}, x]$ satisfies $x^{\beta_1} - (\frac{x}{2})^{\beta_1} = \beta_1 \frac{x}{2} \xi^{\beta_1 - 1}$ and the implied constants are absolute and effectively computable.

Proof. This is [24, Theorem 2.1] with $\delta = 0$ and x replaced with $x/2$. \square

Lemma 7.2 informs the following choices of parameters that we will use throughout the rest of this section. Let $0 < \varepsilon < 1$, let Q be sufficiently large with respect to ε , and let

$$(7.2) \quad \begin{aligned} c_7 &\geq \frac{c_8^2 c_{10}}{3240}, & A &= \sqrt{\frac{3240 c_7}{c_{10} \varepsilon}}, & B &= \frac{41 n^2}{\varepsilon}, & x &= \frac{1}{2} (\log Q)^B, \\ \pi &\in \mathfrak{F}_n(Q), & \pi' &\in \mathfrak{F}_n\left(x^{\frac{1}{A^2 c_{10} n^4 [F:\mathbb{Q}]^2}}\right) = \mathfrak{F}_n\left(\left(2^{-\frac{\varepsilon}{41 n^2}} \log Q\right)^{\frac{41}{3240 c_7 n^2 [F:\mathbb{Q}]^2}}\right). \end{aligned}$$

Corollary 7.3. *Under the notation and hypotheses of (7.2), there holds*

$$\sum_{\substack{\frac{x}{2} < \mathbf{Nn} \leq x \\ \gcd(\mathbf{n}, \mathfrak{q}_{\pi} \mathfrak{q}_{\pi'}) = \mathcal{O}_F}} \Lambda_{\pi' \times \tilde{\pi}'}(\mathbf{n}) \gg_{\varepsilon} x^{1 - \frac{2c_7}{A^2 c_{10} n^4 [F:\mathbb{Q}]^2}}.$$

Proof. Let $\pi, \pi' \in \mathfrak{F}_n(Q)$. In Lemma 7.2, the lower bound on $C(\pi')$ only serves to ensure that the implied constants are absolute. This was pertinent in [24], but it is not pertinent here. Thus, we may replace the two conditions $\log \log C(\pi') \geq c_9 n^4 [F:\mathbb{Q}]^2$ and $x \geq C(\pi')^{A^2 c_{10} n^3 [F:\mathbb{Q}] \log(en[F:\mathbb{Q}])}$ with the single condition $C(\pi') \ll x^{1/(A^2 c_{10} n^4 [F:\mathbb{Q}]^2)}$. We want to refine Lemma 7.2 so that one only sums over \mathbf{n} such that $\gcd(\mathbf{n}, \mathfrak{q}_{\pi} \mathfrak{q}_{\pi'}) = \mathcal{O}_F$. Using (3.3) and (3.7), we find that the contribution to Lemma 7.2(2) arising from \mathbf{n} such that $\gcd(\mathbf{n}, \mathfrak{q}_{\pi} \mathfrak{q}_{\pi'}) \neq \mathcal{O}_F$ is

$$\sum_{\substack{\frac{x}{2} < \mathbf{Nn} \leq x \\ \gcd(\mathbf{n}, \mathfrak{q}_{\pi} \mathfrak{q}_{\pi'}) \neq \mathcal{O}_F}} \Lambda_{\pi' \times \tilde{\pi}'}(\mathbf{n}) \ll \sum_{\mathfrak{p} | \mathfrak{q}_{\pi} \mathfrak{q}_{\pi'}} \sum_{\substack{\log(x/2) < j \leq \frac{\log x}{\log \mathbf{Np}}} \mathbf{Np}^{j(1 - \frac{2}{n^2 + 1})} \ll \frac{\log Q}{\log \log Q} x^{1 - \frac{2}{n^2 + 1}},$$

which is $\ll x^{1-\frac{1}{n^2+1}}$. In the worst case, where β_1 in Lemma 7.2(1) exists, it follows from Lemma 7.2(2) and the above discussion that if $C(\pi') \ll_{n,[F:\mathbb{Q}]} x^{1/(A^2 c_{10} n^3 [F:\mathbb{Q}] \log(en[F:\mathbb{Q}]))}$, then

$$\sum_{\substack{x < N\mathfrak{n} \leq 2x \\ \gcd(\mathfrak{n}, \mathfrak{q}_\pi \mathfrak{q}_{\pi'}) = \mathcal{O}_F}} \Lambda_{\pi' \times \tilde{\pi}'}(\mathfrak{n}) \gg x(1 - \xi^{\beta_1 - 1}) \gg_{n,[F:\mathbb{Q}]} x \min\{1, (1 - \beta_1) \log C(\pi')\} \gg \frac{x \log C(\pi')}{C(\pi')^{c_7 n}}.$$

The desired result now follows. \square

Lemma 7.4. *Let $n \geq 3$. Assume the notation and hypotheses in (7.2). Let Φ be a fixed smooth function supported on a compact subset of $[\frac{1}{4}, 2]$ such that $0 \leq \Phi(t) \leq 1$ for all $t \in [\frac{1}{4}, 2]$ and $\Phi(t) = 1$ for $t \in (\frac{1}{2}, 1]$. If $L(s, \pi \times \pi')$ is entire and does not vanish in the region $\operatorname{Re}(s) \geq 1 - \varepsilon/(20n^2)$ and $|\operatorname{Im}(s)| \leq \log Q$, then*

$$\sum_{\gcd(\mathfrak{n}, \mathfrak{q}_\pi \mathfrak{q}_{\pi'}) = \mathcal{O}_F} \Lambda_{\pi \times \tilde{\pi}'}(\mathfrak{n}) \Phi(N\mathfrak{n}/x) \ll_B x^{1+\frac{2}{B}-\frac{\varepsilon}{20n^2}}.$$

Proof. Writing $\widehat{\Phi}(s) = \int_0^\infty \Phi(t)t^{s-1}dt$ for the Mellin transform of Φ (which is an entire function of s), we observe via Mellin inversion that

$$\sum_{\gcd(\mathfrak{n}, \mathfrak{q}_\pi \mathfrak{q}_{\pi'}) = \mathcal{O}_F} \Lambda_{\pi \times \tilde{\pi}'}(\mathfrak{n}) \Phi(N\mathfrak{n}/x) = \frac{1}{2\pi i} \int_{2-i\infty}^{2+i\infty} -\frac{(L^{\text{ur}})'}{(L^{\text{ur}})}(s, \pi \times \tilde{\pi}') \widehat{\Phi}(s) x^s ds,$$

where $L^{\text{ur}}(s, \pi \times \tilde{\pi}') = L(s, \pi \times \tilde{\pi}') \prod_{\mathfrak{p}|\mathfrak{q}_\pi \mathfrak{q}_{\pi'}} L(s, \pi_{\mathfrak{p}} \times \tilde{\pi}'_{\mathfrak{p}})^{-1}$. A standard contour integral calculation using the argument principle shows that the above display equals

$$- \sum_{L^{\text{ur}}(\rho, \pi \times \tilde{\pi}') = 0} \widehat{\Phi}(\rho) x^\rho,$$

where ρ ranges over all zeros of $L^{\text{ur}}(s, \pi \times \tilde{\pi}')$.

Since Φ is compactly supported and $\widehat{\Phi}$ is entire, it follows that for any $R \geq 2$, we have $|\widehat{\Phi}(s)| \ll_R \min\{1, |s|^{-R} e^{\operatorname{Re}(s)}\}$. Note that the reciprocals of the Euler factors of $L(s, \pi \times \tilde{\pi}')$ at prime ideals $\mathfrak{p} \mid \mathfrak{q}_\pi \mathfrak{q}_{\pi'}$ and all of the trivial zeros of $L(s, \pi \times \tilde{\pi}')$ have real part no larger than $1 - \frac{2}{n^2+1}$ per (3.3) and (3.7). Since for any $t \in \mathbb{R}$ there are $\ll \log Q + \log(|t| + 2)$ zeros $\rho = \beta + i\gamma$ of $L^{\text{ur}}(s, \pi \times \tilde{\pi}')$ that satisfy $0 < \beta < 1$ and $|\gamma - t| \leq 1$, we find for any $T \geq 1$, $R \geq 2$, and $\sigma_0 \geq 1 - \frac{2}{n^2+1}$ that

$$(7.3) \quad \sum_{L^{\text{ur}}(\rho, \pi \times \tilde{\pi}') = 0} \widehat{\Phi}(\rho) x^\rho = \sum_{\substack{L^{\text{ur}}(\rho, \pi \times \tilde{\pi}') = 0 \\ \beta \geq \sigma_0 \\ |\gamma| \leq T}} \widehat{\Phi}(\rho) x^\rho + O(T^{1-R} \log(QT)x + T \log(QT)x^{\sigma_0}).$$

We choose $T = \log Q = (2x)^{1/B}$ and $\sigma_0 = 1 - \varepsilon/(20n^2)$, in which case our hypotheses imply that the sum over zeros on the right-hand side of (7.3) is empty and

$$\left| \sum_{L^{\text{ur}}(\rho, \pi \times \tilde{\pi}') = 0} \widehat{\Phi}(\rho) x^\rho \right| \ll_B x^{1+\frac{2-R}{B}} + x^{\frac{2}{B}+\sigma_0}.$$

The desired result follows from choosing $R = \max\{B(1 - \sigma_0), 3\}$. \square

Proof of Theorem 2.3. Suppose to the contrary that $\pi \neq \pi'$ and $\pi_{\mathfrak{p}} \cong \pi'_{\mathfrak{p}}$ for all $\mathfrak{p} \nmid \mathfrak{q}_\pi \mathfrak{q}_{\pi'}$ with $N\mathfrak{p} \leq 2x$. Then $A_\pi(\mathfrak{p}) = A_{\pi'}(\mathfrak{p})$ for all $\mathfrak{p} \nmid \mathfrak{q}_\pi \mathfrak{q}_{\pi'}$ with $N\mathfrak{p} \leq 2x$. By (3.6) and (7.1), it

follows that $\Lambda_{\pi \times \tilde{\pi}'}(\mathbf{n}) = \Lambda_{\pi' \times \tilde{\pi}'}(\mathbf{n})$ for all \mathbf{n} such that $\gcd(\mathbf{n}, \mathfrak{q}_\pi \mathfrak{q}_{\pi'}) = \mathcal{O}_F$ and $N\mathbf{n} \leq x$. Since $\Lambda_{\pi' \times \tilde{\pi}'}(\mathbf{n}) \geq 0$ for all such \mathbf{n} , the same holds for $\Lambda_{\pi \times \tilde{\pi}'}(\mathbf{n})$. It follows that

$$\sum_{\substack{\frac{x}{2} < N\mathbf{n} \leq x \\ \gcd(\mathbf{n}, \mathfrak{q}_\pi \mathfrak{q}_{\pi'}) = \mathcal{O}_F}} \Lambda_{\pi' \times \tilde{\pi}'}(\mathbf{n}) = \sum_{\substack{\frac{x}{2} < N\mathbf{n} \leq x \\ \gcd(\mathbf{n}, \mathfrak{q}_\pi \mathfrak{q}_{\pi'}) = \mathcal{O}_F}} \Lambda_{\pi \times \tilde{\pi}'}(\mathbf{n}).$$

By (7.2), we have that

$$(7.4) \quad C(\pi') \ll x^{1/(A^2 c_{10} n^4 [F:\mathbb{Q}]^2)}.$$

Therefore, if $L(s, \pi \times \tilde{\pi}') \neq 0$ in the region $\{s \in \mathbb{C} : \operatorname{Re}(s) \geq 1 - \varepsilon/(20n^2), |\operatorname{Im}(s)| \leq \log Q\}$, then by Corollary 7.3 and Lemma 7.4, we have that

$$x^{1 - \frac{2c_7}{A^2 c_{10} n^3 [F:\mathbb{Q}]^2}} \ll \sum_{\substack{\frac{x}{2} < N\mathbf{n} \leq x \\ \gcd(\mathbf{n}, \mathfrak{q}_\pi \mathfrak{q}_{\pi'}) = \mathcal{O}_F}} \Lambda_{\pi' \times \tilde{\pi}'}(\mathbf{n}) = \sum_{\substack{\frac{x}{2} < N\mathbf{n} \leq x \\ \gcd(\mathbf{n}, \mathfrak{q}_\pi \mathfrak{q}_{\pi'}) = \mathcal{O}_F}} \Lambda_{\pi \times \tilde{\pi}'}(\mathbf{n}) \ll_B x^{1 + \frac{2}{B} - \frac{\varepsilon}{20n^2}}.$$

This implies that

$$1 - \frac{2c_7}{A^2 c_{10} n^3 [F:\mathbb{Q}]^2} \leq 1 + \frac{2}{B} - \frac{\varepsilon}{20n^2},$$

which contradicts our choices of A and B in (7.2). The desired result now follows from Corollary 7.1. \square

8. AUTOMORPHIC LEVEL OF DISTRIBUTION

In this section, we prove Theorem 2.5. In what follows, let $F = \mathbb{Q}$. If $n = 2$, then $L(s, \pi \times \tilde{\pi})$ is the L -function of an isobaric automorphic representation of $\operatorname{GL}_4(\mathbb{A}_{\mathbb{Q}})$ whose cuspidal constituents have rank at most 3. Since the L -function any Dirichlet character twist of any cuspidal constituents of rank 2 or 3 have no Landau–Siegel zero by [1, 21], a stronger result than Theorem 2.5 follows from a minor variation of the proof of [31, Theorem 1.1]. Therefore, we may restrict our consideration to $n \geq 3$.

Let $\pi \in \mathfrak{F}_n$ have arithmetic conductor q_π , and let χ be a primitive Dirichlet character modulo q . We will allow π to be fixed, so for notational compactness, we introduce $L_\chi(s) := L(s, \pi \times (\tilde{\pi} \otimes \chi))$ and $L_1(s) := L(s, \pi \times \tilde{\pi})$. If q and q_π are coprime, then $L_\chi(s)$ is entire if and only if χ is nontrivial, since $\tilde{\pi} \otimes \chi \neq \tilde{\pi}$, which is clear by comparing the arithmetic conductors of $q_{\tilde{\pi} \otimes \chi}$ and $q_{\tilde{\pi}}$. If χ is trivial, then $L_\chi(s) = L_1(s)$. We also define $\Lambda_\chi(s) := \Lambda(s, \pi \times (\tilde{\pi} \otimes \chi))$ and $\Lambda_1(s) := \Lambda(s, \pi \times \tilde{\pi})$.

8.1. Preliminaries. We define $a_\pi(m)$ and $a_{\pi \times \tilde{\pi}}(m)$ by the Dirichlet series identities

$$\sum_{n=1}^{\infty} \frac{a_\pi(m) \Lambda(m)}{m^s} = -\frac{L'}{L}(s, \pi), \quad \sum_{n=1}^{\infty} \frac{a_{\pi \times \tilde{\pi}}(m) \Lambda(m)}{m^s} = -\frac{L'}{L}(s, \pi \times \tilde{\pi}), \quad \operatorname{Re}(s) > 1.$$

The local calculations in [36, Lemma 2.1] show that if $\chi \pmod{q}$ is a primitive Dirichlet character and $\gcd(q, q_\pi) = 1$, then for all primes p , we have that

$$(8.1) \quad L(s, \pi_p \times (\tilde{\pi} \otimes \chi)_p) = \prod_{j=1}^n \prod_{j'=1}^n (1 - \alpha_{j, j', \pi \times \tilde{\pi}}(p) \chi(p) p^{-s})^{-1}.$$

It follows from (3.6) and (8.1) that if $\gcd(m, q_\pi) = 1$, then $a_{\pi \times (\tilde{\pi} \otimes \chi)}(m) = |a_\pi(m)|^2 \chi(m)$. We now provide a convenient expression for $-L'_\chi(s)/L_\chi(s)$.

Lemma 8.1. *Let $\pi \in \mathfrak{F}_n$. Let $\psi \pmod{q}$ be a Dirichlet character with q such that $\gcd(q, q_\pi) = 1$ and χ be the primitive Dirichlet character that induces ψ . Let $\delta(\chi) = 1$ if χ is trivial and $\delta(\chi) = 0$ otherwise. There exists a function $H_\pi(s; \chi, \psi)$ such that in the region $\operatorname{Re}(s) \geq 1 - \frac{1}{n^2}$,*

- (1) $H_\pi(s; \chi, \psi)$ is holomorphic,
- (2) $|H_\pi(s; \chi, \psi)| \ll_\pi \log(q(3 + |\operatorname{Im}(s)|))$, and
- (3) we have the identity

$$\sum_{m=1}^{\infty} \frac{|a_\pi(m)|^2 \psi(m) \Lambda(m)}{m^s} = \frac{\delta(\chi)}{s-1} - \frac{\Lambda'_\chi}{\Lambda_\chi}(s) + H_\pi(s; \chi, \psi).$$

Proof. Suppose first that ψ is primitive, in which case $\psi = \chi$ and the function $H_\pi(s; \chi, \psi)$ is

$$\begin{aligned} \frac{\log q_{\pi \times (\tilde{\pi} \otimes \chi)}}{2} + \frac{L'}{L}(s, \pi_\infty \times (\tilde{\pi} \otimes \chi)_\infty) + \frac{\delta(\chi)}{s} + \sum_{p|q\pi q} \frac{L'}{L}(s, \pi_p \times (\tilde{\pi}' \otimes \chi)_p) \\ - \sum_{p|q\pi q} \sum_{\substack{1 \leq j \leq n \\ 1 \leq j' \leq n}} \frac{\alpha_{j,\pi}(p) \overline{\alpha_{j',\pi'}(p)} \chi(p) \log p}{p^s - \alpha_{j,\pi}(p) \alpha_{j',\pi}(p) \chi(p)}. \end{aligned}$$

This is holomorphic and bounded as claimed for $\operatorname{Re}(s) \geq 1 - \frac{1}{n^2}$ by (3.7), (3.10), and Stirling's formula. If ψ is not primitive and χ is the primitive character that induces ψ , then in the same region, we have

$$\left| \sum_{m=1}^{\infty} \frac{|a_\pi(m)|^2 \psi(m) \Lambda(m)}{m^s} - \sum_{m=1}^{\infty} \frac{|a_\pi(m)|^2 \chi(m) \Lambda(m)}{m^s} \right| \leq \sum_{p|q} \sum_{k \geq 1} \frac{|a_\pi(p^k)|^2 \log p}{p^{k \operatorname{Re}(s)}},$$

which is bounded as desired using (3.7) again. \square

We will follow the following zero-free region for $L_\chi(s)$ and Siegel-type bound on any real exceptional zeros. We will prove this theorem later.

Theorem 8.2. *Let $Q \geq 3$ and $\pi \in \mathfrak{F}_n$. There exists an effectively computable constant $c_{12} = c_{12}(\pi) > 0$ such that for all primitive Dirichlet characters $\chi \pmod{q}$ with $q \leq Q$ and $\gcd(q, q_\pi) = 1$ with at most one exception, the L -function $L(s, \pi \times (\tilde{\pi} \otimes \chi))$ does not vanish in the region*

$$\operatorname{Re}(s) \geq 1 - \frac{c_{12}}{\log(Q(|\operatorname{Im}(s)| + 3))}.$$

If the exceptional character $\chi_1 \pmod{q_1}$ exists, then $L(s, \pi \times (\tilde{\pi} \otimes \chi_1))$ has at most one zero, say β_1 , in this region; β_1 is both real and simple; and χ_1 must be quadratic. Moreover, for all $\varepsilon > 0$, there exists an ineffective constant $c_\pi(\varepsilon) > 0$ such that $\beta_1 \leq 1 - c_\pi(\varepsilon) q_1^{-\varepsilon}$.

8.2. Proof of Theorem 2.5. We follow Gallagher's proof of the Bombieri–Vinogradov theorem in [16], with $n \geq 2$ and $\pi \in \mathfrak{F}_n$ fixed at the onset. Note that the function

$$\psi_k(y; q, a) := \frac{1}{k!} \sum_{\substack{m \leq y \\ m \equiv a \pmod{q}}} |a_\pi(m)|^2 \Lambda(m) \left(\log \frac{y}{m} \right)^k$$

is monotonically increasing as a function of y for each $k \geq 0$. Thus, if $0 < \lambda \leq 1$, then

$$\frac{1}{\lambda} \int_{e^{-\lambda y}}^y \psi_{k-1}(t; q, a) \frac{dt}{t} \leq \psi_{k-1}(y; q, a) \leq \frac{1}{\lambda} \int_y^{e^{\lambda y}} \psi_{k-1}(t; q, a) \frac{dt}{t}.$$

The integrals, which equal $\psi_k(y; q, a) - \psi_k(e^{-\lambda}y; q, a)$ and $\psi_k(e^{\lambda}y; q, a) - \psi_k(y; q, a)$ respectively, both have the same asymptotic expansion, namely

$$(\lambda + O(\lambda^2)) \frac{y}{\varphi(q)} + O(\max_{y \leq ex} |r_k(y; q, a)|), \quad r_k(y; q, a) := \psi_k(y; q, a) - \frac{y}{\varphi(q)},$$

where φ is Euler's totient function. Thus, we have the bounds

$$\max_{y \leq x} |r_{k-1}(y; q, a)| \ll \frac{\lambda x}{\varphi(q)} + \frac{1}{\lambda} \max_{y \leq ex} |r_k(y; q, a)|$$

and

$$\sum_{\substack{q \leq x^\theta \\ \gcd(q, q_\pi)=1}} \max_{\gcd(a, q)=1} \max_{y \leq x} |r_{k-1}(y; q, a)| \ll \lambda x \log x + \frac{1}{\lambda} \sum_{\substack{q \leq x^\theta \\ \gcd(q, q_\pi)=1}} \max_{\gcd(a, q)=1} \max_{y \leq ex} |r_k(y; q, a)|.$$

It follows by induction on k that

$$\sum_{\substack{q \leq x^\theta \\ \gcd(q, q_\pi)=1}} \max_{\gcd(a, q)=1} \max_{y \leq x} |r_0(y; q, a)| \ll_k \lambda x \log x + \frac{1}{\lambda^{2^k-1}} \sum_{\substack{q \leq x^\theta \\ \gcd(q, q_\pi)=1}} \max_{\gcd(a, q)=1} \max_{y \leq ex} |r_k(y; q, a)|.$$

Proposition 8.3. *If $\theta < 1/(9n^3)$ is fixed and $k = 9n^2 + 1$, then for any $B > 0$, we have*

$$\sum_{\substack{q \leq x^\theta \\ \gcd(q, q_\pi)=1}} \max_{\gcd(a, q)=1} \max_{y \leq ex} |r_k(y; q, a)| \ll_{\pi, B} \frac{x}{(\log x)^B}.$$

Proposition 8.3 implies Theorem 2.5. It follows from Proposition 8.3 that

$$\sum_{\substack{q \leq x^\theta \\ \gcd(q, q_\pi)=1}} \max_{\gcd(a, q)=1} \max_{y \leq x} |r_0(y; q, a)| \ll_{\pi, B} \lambda x \log x + \frac{1}{\lambda^{2^k-1}} \frac{x}{(\log x)^B}.$$

To finish, we choose $B = 2^k(A+1) - 1$ and $\lambda = (\log x)^{-(B+1)/2^k}$. \square

8.3. Proof of Proposition 8.3. Let $\pi \in \mathfrak{F}_n$, $k = 9n^2 + 1$, $Q = x^\theta$ for some fixed $0 < \theta < 1/(9n^3)$, and $q \leq Q$. We have the decomposition

$$\frac{1}{k!} \sum_{\substack{m \leq y \\ m \equiv a \pmod{q}}} |a_\pi(m)|^2 \Lambda(m) \left(\log \frac{y}{m} \right)^k = \frac{1}{k!} \sum_{\psi \pmod{q}} \frac{\bar{\psi}(a)}{\varphi(q)} \sum_{y \leq m} |a_\pi(m)|^2 \Lambda(m) \psi(m) \left(\log \frac{y}{m} \right)^k.$$

By Lemma 8.1 and Mellin inversion, this equals

$$\frac{1}{\varphi(q)} \sum_{\psi \pmod{q}} \sum_{\substack{\chi \text{ primitive} \\ \chi \text{ induces } \psi}} \frac{\bar{\chi}(a)}{2\pi i} \int_{3-i\infty}^{3+i\infty} \left(-\frac{\Lambda'_\chi(s)}{\Lambda_\chi(s)} + H_\pi(s; \chi, \psi) \right) \frac{y^s}{s^{k+1}} ds.$$

Since a primitive character $\chi \pmod{q}$ induces characters to moduli that are a multiple of q , it follows from the bound $\sum_{f \leq Q, q|f} \varphi(f)^{-1} \ll \frac{\log Q}{\varphi(q)}$ that

$$(8.2) \quad \begin{aligned} & \sum_{\substack{q \leq Q \\ \gcd(q, q_\pi)=1}} \max_{\gcd(a, q)=1} \max_{y \leq x} \left| \frac{1}{k!} \sum_{\substack{m \leq y \\ m \equiv a \pmod{q}}} |a_\pi(m)|^2 \Lambda(m) \left(\log \frac{y}{m} \right)^k - \frac{y}{\varphi(q)} \right| \\ & \ll \sum_{\substack{q \leq Q \\ \gcd(q, q_\pi)=1}} \frac{\log Q}{\varphi(q)} \sum_{\substack{\chi \pmod{q} \\ \chi \text{ primitive}}} \max_{y \leq x} \left| \frac{1}{2\pi i} \int_{3-i\infty}^{3+i\infty} \left(-\frac{\Lambda'_\chi(s)}{\Lambda_\chi(s)} + H_\pi(s; \chi, \chi) \right) \frac{y^s}{s^{k+1}} ds \right|. \end{aligned}$$

Observe that by Lemma 8.1 and our range of θ , (8.2) equals

$$(8.3) \quad \begin{aligned} & \sum_{\substack{q \leq Q \\ \gcd(q, q_\pi)=1}} \frac{\log Q}{\varphi(q)} \sum_{\substack{\chi \pmod{q} \\ \chi \text{ primitive}}} \max_{y \leq x} \left| - \sum_{L_\chi(\rho)=0} \frac{y^\rho}{\rho^{k+1}} + \frac{1}{2\pi i} \int_{1-\frac{1}{\pi^2}-i\infty}^{1-\frac{1}{\pi^2}+i\infty} H_\pi(s; \chi, \chi) \frac{y^s}{s^{k+1}} ds \right| \\ & \ll_{\pi, B} \log Q \sum_{\substack{q \leq Q \\ \gcd(q, q_\pi)=1}} \frac{1}{\varphi(q)} \sum_{\substack{\chi \pmod{q} \\ \chi \text{ primitive}}} \sum_{L_\chi(\rho)=0} \frac{y^\beta}{|\rho|^{k+1}} + \frac{x}{(\log x)^B}, \end{aligned}$$

where $\rho = \beta + i\gamma$ ranges over the nontrivial zeros of $L_\chi(s)$. In light of the bounds $\frac{1}{q} \leq \frac{1}{\varphi(q)} \ll \frac{\log(eq)}{q}$, we dyadically decompose $[1, Q]$ into $O(\log Q)$ subintervals and find that (8.3) is

$$(8.4) \quad \begin{aligned} & \ll_{\pi, B} x(\log Q)^2 \sup_{3 \leq R \leq Q} \sum_{\substack{q \leq R \\ \gcd(q, q_\pi)=1}} \frac{1}{\varphi(q)} \sum_{\substack{\chi \pmod{q} \\ \chi \text{ primitive}}} \sum_{\rho=\beta+i\gamma} \frac{x^{\beta-1}}{|\rho|^k} + \frac{x}{(\log x)^B} \\ & \ll_{\pi, B} x(\log Q)^3 \sup_{3 \leq R \leq x^\theta} \frac{1}{R} \sum_{\substack{q \leq R \\ \gcd(q, q_\pi)=1}} \sum_{\substack{\chi \pmod{q} \\ \chi \text{ primitive}}} \sum_{\rho=\beta+i\gamma} \frac{x^{\beta-1}}{|\rho|^k} + \frac{x}{(\log x)^B} \\ & \ll_{\pi, B} x(\log x)^3 \sup_{3 \leq R \leq x^\theta} \frac{1}{R} \left(\frac{x^{\beta_1-1}}{\beta_1^k} + \sum_{\substack{q \leq R \\ \gcd(q, q_\pi)=1}} \sum_{\substack{\chi \pmod{q} \\ \chi \text{ primitive}}} \sum_{\rho=\beta+i\gamma \neq \beta_1} \frac{x^{\beta-1}}{|\rho|^k} \right) + \frac{x}{(\log x)^B}. \end{aligned}$$

where β_1 is the exceptional zero in Theorem 8.2. The term x^{β_1-1}/β_1^k is omitted if β_1 does not exist.

If β_1 exists as in Theorem 8.2 and the supremum is achieved when $R \leq (\log x)^{4B}$, then we apply Theorem 8.2 with $\varepsilon = \frac{1}{8B}$ and conclude that the contribution from such a zero is absorbed in our error term. If the supremum is achieved when $R > (\log x)^{4B}$, then contribution from β_1 is trivially absorbed in our error term. Hence (8.4) is

$$(8.5) \quad \ll_{\pi, B} x(\log x)^3 \sup_{3 \leq R \leq x^\theta} \frac{1}{R} \sum_{\substack{q \leq R \\ \gcd(q, q_\pi)=1}} \sum_{\substack{\chi \pmod{q} \\ \chi \text{ primitive}}} \sum_{\rho=\beta+i\gamma \neq \beta_1} \frac{x^{\beta-1}}{|\rho|^k} + \frac{x}{(\log x)^B}.$$

Let us now consider the zeros ρ with $|\rho| < \frac{1}{4}$. The number of such zeros is $\ll R^2 \log R$. From the consideration of the corresponding zeros $1 - \rho$ of $L_{\overline{\chi}}(s)$, we deduce that $|\rho| \gg x^{-1/(4k)}$. Thus, the contribution from these zeros is $\ll Rx^{\frac{1}{4} + \frac{k}{4k}} \log R \ll Qx^{\frac{1}{2}} \log Q \ll_B$

$x(\log x)^{-B}$. Define $T_0 = 0$ and $T_j = 2^{j-1}$ for $j \geq 1$. The above discussion shows that (8.5) is

$$(8.6) \quad \ll_{\pi, B} x(\log x)^3 \sup_{3 \leq R \leq x^\theta} \frac{1}{R} \sum_{\substack{q \leq R \\ \gcd(q, q_\pi) = 1}} \sum_{\substack{\chi \pmod{q} \\ \chi \text{ primitive}}} \sum_{\substack{\rho = \beta + i\gamma \neq \beta_1 \\ |\rho| > \frac{1}{4}}} \frac{x^{\beta-1}}{|\rho|^k} + \frac{x}{(\log x)^B}$$

$$\ll_{\pi, B} x(\log x)^3 \sup_{3 \leq R \leq x^\theta} \frac{1}{R} \sum_{j=1}^{\infty} \sum_{\substack{q \leq R \\ \gcd(q, q_\pi) = 1}} \sum_{\substack{\chi \pmod{q} \\ \chi \text{ primitive}}} \sum'_{\substack{\rho = \beta + i\gamma \neq \beta_1 \\ |\rho| \geq \frac{1}{4} \\ T_{j-1} \leq |\gamma| \leq T_j}} \frac{x^{\beta-1}}{|\rho|^k} + \frac{x}{(\log x)^B}.$$

If $|\rho| \geq \frac{1}{4}$ and $T_{j-1} \leq |\gamma| \leq T_j$, then $|\rho| \geq \max\{T_{j-1}, \frac{1}{4}\} \geq T_j/4$ and $|\rho| \gg |\gamma| + 3$. Therefore, if $\delta = \min\{1 - 9n^3\theta, \frac{1}{2}\}$, then

$$x^{\beta-1} |\rho|^{-k} \ll T_j^{-\frac{1}{2}} (|\gamma| + 1)^{-\frac{1}{2}} x^{-(1-\beta)\delta} (x^{1-\delta} T_j^{k-1})^{-(1-\beta)} \ll T_j^{-\frac{1}{2}} (|\gamma| + 1)^{-\frac{1}{2}} x^{-(1-\beta)\delta} (R^{\frac{1-\delta}{\theta}} T_j^{k-1})^{\beta-1}.$$

Since $\rho = \beta + i\gamma \neq \beta_1$, it follows from Theorem 8.2 that

$$(|\gamma| + 1)^{-\frac{1}{2}} x^{-(1-\beta)\delta} \leq e^{-\delta\eta_\pi(x, R)}, \quad \eta_\pi(x, R) := \inf_{t \geq 3} \left[c_\pi \frac{\log x}{\log(Rt)} + \log t \right],$$

so (8.6) is

$$\ll_{\pi, B} x(\log x)^3 \sup_{3 \leq R \leq x^\theta} \frac{e^{-\delta\eta_\pi(x, R)}}{R} \sum_{j=1}^{\infty} T_j^{-\frac{1}{2}} \sum_{\substack{q \leq R \\ \gcd(q, q_\pi) = 1}} \sum_{\substack{\chi \pmod{q} \\ \chi \text{ primitive}}} \sum_{\substack{\rho = \beta + i\gamma \neq \beta_1 \\ |\gamma| \leq T_j}} (R^{\frac{1-\delta}{\theta}} T_j^{k-1})^{\beta-1} + \frac{x}{(\log x)^B}.$$

Define

$$N_\pi^*(\sigma, T, R) = \sum_{\substack{q \leq R \\ \gcd(q, q_\pi) = 1}} \sum_{\substack{\chi \pmod{q} \\ \chi \text{ primitive}}} |\{\rho = \beta + i\gamma : \beta \geq \sigma, |\gamma| \leq T, L_\chi(\rho) = 0\}|.$$

By (3.10), there exists an effectively computable constant $c_{13} = c_{13}(n) > 0$ such that

$$\{\tilde{\pi} \otimes \chi : \chi \pmod{q_\chi} \text{ primitive}, \gcd(q_\chi, q_\pi) = 1, q_\chi \leq Q\} \subseteq \mathfrak{F}_n(c_{13}C(\pi)Q^n).$$

Thus, by (1.7), we have $N_\pi^*(\sigma, T, R) \ll_{\pi, \varepsilon} (R^n T)^{9n^2(1-\sigma)+\varepsilon}$. Partial summation yields

$$(8.7) \quad \sum_{\substack{q \leq R \\ \gcd(q, q_\pi) = 1}} \sum_{\substack{\chi \pmod{q} \\ \chi \text{ primitive}}} \sum_{\substack{\rho = \beta + i\gamma \neq \beta_1 \\ |\gamma| \leq T_j}} (R^{\frac{1-\delta}{\theta}} T_j^{k-1})^{-(1-\beta)}$$

$$\ll (R^{\frac{1-\delta}{\theta}} T_j^{k-1})^{-1} N_\pi^*(0, T_j, R) + \log(RT_j) \int_0^1 (R^{\frac{1-\delta}{\theta}} T_j^{k-1})^{-\sigma} N_\pi^*(1-\sigma, T_j, R) d\sigma$$

$$\ll_{\pi, \varepsilon} (RT_j)^\varepsilon \left((R^{\frac{1-\delta}{\theta}} T_j^{k-1})^{-1} RT_j + \int_0^1 (R^{\frac{1-\delta}{\theta}} T_j^{k-1})^{-\sigma} (R^n T_j)^{9n^2\sigma} d\sigma \right).$$

Our choices for θ , k , and δ ensure that (8.7) is $\ll_{\pi, \varepsilon} (RT_j)^\varepsilon$. Thus, (8.6) is

$$(8.8) \quad \ll_{\pi, B, \varepsilon} x(\log x)^3 \sup_{3 \leq R \leq x^\theta} e^{-\delta\eta_\pi(x, R)} \frac{R^\varepsilon}{R} \sum_{j=1}^{\infty} T_j^{-\frac{1}{2}+\varepsilon} + \frac{x}{(\log x)^B}$$

$$\ll_{\pi, B, \varepsilon} x(\log x)^3 \sup_{3 \leq R \leq x^\theta} e^{-\delta\eta_\pi(x, R)} \frac{R^\varepsilon}{R} + \frac{x}{(\log x)^B}.$$

A small calculation (cf. [46, Section 4]) shows that there is a constant $c_{14} = c_{14}(\pi) > 0$ such that $e^{-\delta\eta_\pi(x,R)} \ll_{\pi,B,\varepsilon} \exp(-c_{14}\frac{\log x}{\log R}) + \exp(-c_{14}\sqrt{\log x})$, and Theorem 2.5 follows.

8.4. Proof of Theorem 8.2. Although the proof of Theorem 8.2 contains only standard techniques, such zero-free regions for $L_\chi(s)$ are new even in the case when χ is trivial. The case when χ is trivial was only recently handled unconditionally in [24].

Lemma 8.4. [24, Theorem 2.1(1)] *Let $\pi \in \mathfrak{F}_n$. There exists an absolute and effectively computable constant $c_{15} > 0$ such that $L_1(s)$ has at most one zero, say β_1 , in the region $\operatorname{Re}(s) \geq 1 - c_{15}/\log(C(\pi)^n(|\operatorname{Im}(s)| + e)^{n^2})$. If β_1 exists, then it must be real and simple, and there exists an absolute and effectively computable constant c_{16} such that $\beta_1 \leq 1 - C(\pi)^{-c_{16}n}$.*

We prove the corresponding result for $L_\chi(s)$. The ideas in [24] inform our approach here

Lemma 8.5. *Let $\pi \in \mathfrak{F}_n$. Let $\chi \pmod{q}$ be a nontrivial primitive Dirichlet character such that $\gcd(q, q_\pi) = 1$. There exists an effectively computable constant $c_{17} = c_{17}(n) > 0$ such that $L_\chi(s)$ has at most one zero, say β_1 , in the region*

$$(8.9) \quad \operatorname{Re}(s) \geq 1 - \frac{c_{17}}{\log(qC(\pi)(|\operatorname{Im}(s)| + 3))}.$$

If the exceptional zero β_1 exists, then it is real and simple, and χ is quadratic.

Proof. Let $\chi \pmod{q}$ be a primitive nontrivial Dirichlet character, let ψ be the primitive character that induces χ^2 , and let $\beta + i\gamma$ be a nontrivial zero of $L_\chi(s)$. Define $\Pi_\chi = \pi \boxplus \pi \otimes \chi \cdot | \cdot |^{it} \boxplus \pi \otimes \bar{\chi} \cdot | \cdot |^{-it}$, and define

$$(8.10) \quad D(s) = L(s, \Pi_\chi \times \tilde{\Pi}_\chi) = L_1(s)^3 L_\chi(s + i\gamma)^2 L_{\bar{\chi}}(s - i\gamma)^2 L_\psi(s + 2i\gamma) L_{\bar{\psi}}(s - 2i\gamma).$$

The factor $L_1(s)^3$ has a pole of order 3 at $s = 1$, and the hypothesis that $\gcd(q, q_\pi) = 1$ ensures that $L_\chi(s + i\gamma)^2 L_{\bar{\chi}}(s - i\gamma)^2$ is entire. If ψ is complex, then $L_\psi(s + 2i\gamma) L_{\bar{\psi}}(s - 2i\gamma)$ is entire; otherwise, it has poles of order 1 at $s = 1 \pm 2i\gamma$. The additional poles when ψ is real require some additional casework when γ is close to zero. For notational compactness, let $\mathcal{Q}_\gamma = qC(\pi)(|\gamma| + 3)$. Note that $-D'(s)$ has nonnegative Dirichlet coefficients per [21, Lemma a].

The functional equation for $L_\chi(s)$ together with the fact that $L(s, \pi \times \tilde{\pi})$ is a self-dual L -function (even if $L(s, \pi)$ itself is not self-dual) implies that if ρ is a zero of $L_\chi(s)$, then $\bar{\rho}$ is a zero of $L_{\bar{\chi}}(s)$. Thus, we have that

$$(8.11) \quad \operatorname{ord}_{s=\beta} D(s) \geq 4.$$

Let ω denote a nontrivial zero of $D(s)$, $\delta(\psi) = 1$ if ψ is trivial, and $\delta(\psi) = 0$ otherwise. We apply Lemma 3.2 and (3.10) to (8.10), concluding that if $1 < \sigma < 2$, then

$$(8.12) \quad \sum_{\omega} \operatorname{Re}\left(\frac{1}{\sigma - \omega}\right) < \frac{3}{\sigma - 1} + \delta(\psi) \frac{2(\sigma - 1)}{(\sigma - 1)^2 + 4\gamma^2} + c_{18} \log \mathcal{Q}_\gamma,$$

where $c_{18} = c_{18}(n) > 1$ is a suitable implied constant. Since $\beta < 1$ is, by hypothesis, one of the zeros in the sum in (8.12), we have by (8.11) and nonnegativity that

$$(8.13) \quad \frac{4}{\sigma - \beta} < \frac{3}{\sigma - 1} + \delta(\psi) \frac{2(\sigma - 1)}{4\gamma^2 + (\sigma - 1)^2} + c_{18} \log \mathcal{Q}_\gamma.$$

Case 1: Either χ is real and $|\gamma| \geq \frac{1}{7c_{22} \log \mathcal{Q}_0}$, or χ is complex. If $\sigma = 1 + \frac{1}{5c_{18} \log \mathcal{Q}_\gamma}$, then

$$\delta(\psi) \frac{2(\sigma - 1)}{4\gamma^2 + (\sigma - 1)^2} \leq \frac{490}{149} c_{18} \log \mathcal{Q}_\gamma.$$

Thus, (8.13) becomes

$$\frac{4}{1 + \frac{1}{5c_{18} \log \mathcal{Q}_\gamma} - \beta} \leq \frac{2874}{149} c_{18} \log \mathcal{Q}_\gamma.$$

Upon solving for β , we conclude that $\beta \leq 1 - \frac{1}{136c_{18} \mathcal{Q}_\gamma}$.

Case 2: χ is real and $\gamma = 0$. We start at (8.12) with $\delta(\psi) = 1$, $\gamma = 0$, and $\sigma = 1 + \frac{1}{3c_{18} \log \mathcal{Q}_0}$. If there are N zeros β (with multiplicity) of $D(s)$ such that $\beta \geq 1 - \frac{1}{96c_{18} \log \mathcal{Q}_0}$, then by (8.12),

$$\frac{32}{11} c_{18} N \log \mathcal{Q}_0 = \frac{N}{\sigma - (1 - \frac{1}{96c_{18} \log \mathcal{Q}_0})} \leq \frac{5}{\sigma - 1} + c_{18} \log \mathcal{Q}_0 = 16c_{18} \log \mathcal{Q}_0.$$

It follows that (since N is an integer) $N \leq \lfloor \frac{11}{2} \rfloor = 5$. By the bound (8.11), $L_\chi(s)$ has at most one real zero β , necessarily simple, satisfying $\beta \geq 1 - \frac{1}{96c_{18} \log \mathcal{Q}_0}$.

Case 3: χ is quadratic and $0 < |\gamma| < \frac{1}{7c_{22} \log \mathcal{Q}_0}$. We apply Lemma 3.2 to $L_1(s)L_\chi(s)$. The only singularity is a simple pole at $s = 1$. Both $L_1(s)$ and $L_\chi(s)$ are self-dual, so if $\beta + i\gamma$ is a nontrivial zero of $L_1(s)L_\chi(s) = 0$, then so is $\beta - i\gamma$. By (8.1), the m -th Dirichlet coefficient of $-\frac{L'_1}{L_1}(s) - \frac{L'_\chi}{L_\chi}(s)$ is $(1 + \chi(m))a_{\pi \times \tilde{\pi}}(m)\Lambda(m) \geq 0$, so Lemma 3.2 yields

$$(8.14) \quad 2 \frac{\sigma - \beta}{(\sigma - \beta)^2 + \gamma^2} \leq \frac{1}{\sigma - 1} + c_{18} \log \mathcal{Q}_\gamma.$$

If $\sigma = 1 + \frac{1}{2c_{18} \log \mathcal{Q}_\gamma}$ and $0 < |\gamma| < \frac{1}{7c_{18} \log \mathcal{Q}_0}$ in (8.14), then $\beta \leq 1 - \frac{1}{8c_{18} \log \mathcal{Q}_\gamma}$. \square

We now prove a Siegel-type bound for β_1 in Lemma 8.5 (if it exists) using the ideas of Hoffstein and Lockhart [20]. This is new for all $\pi \in \mathfrak{F}_n$ with $n \geq 3$. We begin with an auxiliary calculation. Let $\chi \pmod{q}$ and $\chi' \pmod{q'}$ be distinct nontrivial quadratic Dirichlet characters such that $\gcd(q'q, q_\pi) = 1$, and let ψ be the primitive character that induces $\chi'\chi$ (whose conductor is necessarily coprime to q_π). These coprimality restrictions ensure that $\pi \neq \pi \otimes \chi$, $\pi \neq \pi \otimes \chi'$, $\pi \neq \pi \otimes \psi$, $q_{\pi \otimes \chi} = q_\pi q^n$, $q_{\pi \otimes \chi'} = q_\pi (q')^n$, and $q_{\pi \otimes \psi} = q_\psi q^n$. We also have that $q_\psi | q_\chi q_{\chi'}$. Let

$$(8.15) \quad L(s, \Pi^*) = L_1(s)L_\chi(s)L_{\chi'}(s)L_\psi(s).$$

By the above discussion, $L(s, \Pi^*)$ is holomorphic apart from a simple pole at $s = 1$. It follows from (8.1) that if $k \geq 1$, then the p^k -th Dirichlet coefficient of $\log L(s, \Pi^*)$ equals

$$k^{-1} a_{\pi \times \tilde{\pi}}(p^k) (1 + \chi(p^k) + \chi'(p^k) + \psi(p^k)) \geq 0.$$

The nonnegativity of $1 + \chi(p^k) + \chi'(p^k) + \psi(p^k)$ follows from the fact that this sum is a Dirichlet coefficient of the Dedekind zeta function of a biquadratic field. Upon exponentiating, we find that the m -th Dirichlet coefficient $\lambda_{\Pi^*}(m)$ of $L(s, \Pi^*)$ is nonnegative.

Lemma 8.6. *If $\pi \in \mathfrak{F}_n$ and χ is a primitive nontrivial real Dirichlet character, then $L_\chi(1) > 0$ and the Dirichlet coefficients of $L_1(s)L_\chi(s)$ are nonnegative.*

Proof. Let K be the quadratic field associated to χ . If π_{BC} is the base change of π to an automorphic representation of $\text{GL}_n(\mathbb{A}_K)$, then $L(s, \pi_{\text{BC}} \times \tilde{\pi}_{\text{BC}}) = L_1(s)L_\chi(s)$. Since $L(s, \pi_{\text{BC}} \times \tilde{\pi}_{\text{BC}})$ is holomorphic on $\mathbb{C} - \{1\}$ apart from a simple pole at $s = 1$, the same holds for $L_1(s)L_\chi(s)$. Since $\gcd(q, q_\pi) = 1$ (hence $L_\chi(s)$ is entire), the residue of $L(s, \pi_{\text{BC}} \times \tilde{\pi}_{\text{BC}})$ at $s = 1$, which is positive, equals $L_\chi(1)\text{Res}_{s=1}L_1(s)$. Since $\text{Res}_{s=1}L_1(s) > 0$, it follows that $L_\chi(1) > 0$. The Dirichlet coefficients of $L_1(s)L_\chi(s)$ are nonnegative per Case 3 in the proof of Lemma 8.5. \square

Let $0 < \varepsilon < 1$, and let $\beta \in (1 - \varepsilon, 1)$. By (3.13) and the above discussion, we have

$$(8.16) \quad \begin{aligned} L\left(\frac{1}{2} + it, \Pi^*\right) &\ll_{\pi, \varepsilon} (q'q)^{\frac{n^2}{2} + \varepsilon} (3 + |t|)^{n^2 + \varepsilon}, \\ \text{Res}_{s=1-\beta} L(s + \beta, \Pi^*) x^s \Gamma(s) &\ll_{\pi, \varepsilon} L_\chi(1) (q'q)^\varepsilon (1 - \beta)^{-1} x^{1-\beta}. \end{aligned}$$

If $x \geq 3$, then since $\lambda_{\Pi^*}(1) = 1$, we use (8.16) to deduce that

$$(8.17) \quad \begin{aligned} \frac{1}{e} &\leq \sum_{m=1}^{\infty} \frac{\lambda_{\Pi^*}(m)}{m^\beta} e^{-\frac{m}{x}} \\ &= \frac{1}{2\pi i} \int_{3-i\infty}^{3+i\infty} L(s + \beta, \Pi^*) x^s \Gamma(s) ds \\ &= \text{Res}_{s=1-\beta} L(s + \beta, \Pi^*) x^s \Gamma(s) + L(\beta, \Pi^*) + \frac{1}{2\pi i} \int_{\frac{1}{2}-\beta-i\infty}^{\frac{1}{2}-\beta+i\infty} L(s + \beta, \Pi^*) x^s \Gamma(s) ds \\ &\ll_{\pi, \varepsilon} L_\chi(1) (q'q)^\varepsilon (1 - \beta)^{-1} x^{1-\beta} + L(\beta, \Pi^*) + (q'q)^{\frac{n^2}{2} + \varepsilon} x^{\frac{1}{2}-\beta}. \end{aligned}$$

Proposition 8.7. *Recall the notation and hypotheses of Lemma 8.5. If β_1 exists for a primitive character $\chi \pmod{q}$ such that $\gcd(q, q_\pi) = 1$, then for all $\varepsilon > 0$, there exists an (ineffective) constant $c'_\pi(\varepsilon) > 0$ such that $L_\chi(1) \geq c'_\pi(\varepsilon) q^{-\varepsilon}$.*

Proof. It suffices to take $0 < \varepsilon < 1$. Let $\chi \pmod{q}$ and $\chi' \pmod{q'}$ be primitive quadratic Dirichlet characters with, let ψ be the primitive character that induces $\chi'\chi$, and recall the definition of $L(s, \Pi^*)$ from (8.15). Our proof consists of two cases.

First, suppose that there exists no primitive quadratic Dirichlet character ω such that $L_\omega(s)$ does not vanish for $s \in (1 - \frac{\varepsilon}{2}, 1)$. It then follows that there exists a constant $c_{19} = c_{19}(\pi) > 0$ such that $L_1(s)L_\chi(s) \neq 0$ in the interval $s \in (1 - c_{19}/\log q, 1)$. Since the Dirichlet coefficients of $L_1(s)L_\chi(s)$ are nonnegative (using Lemma 8.6) and the residue of $L_1(s)L_\chi(s)$ at $s = 1$ is $L_\chi(1)\text{Res}_{s=1}L_1(s)$, it follows from [20, Proposition 1.1] and (3.13) that $L_\chi(1)\text{Res}_{s=1}L_1(s) \gg_\pi (\log q)^{-1}$. Since each term the left-hand side is positive (using Lemma 8.6), the desired result follows.

Second, suppose that there exists $\chi' \pmod{q'}$ and $\beta \in (1 - \frac{\varepsilon}{2}, 1)$ such that $L_{\chi'}(\beta) = 0$. We may assume that q' is minimal, subject to this condition. Now, let χ be arbitrary. If $q < q'$, then the minimality of q' ensures that $L_\chi(s) \neq 0$ for $s \in (1 - \frac{\varepsilon}{2}, 1)$, and the preceding case implies the desired result. Suppose now that $q \geq q'$. If $L_\chi(s)$ has no real zero within a distance of $c_{17}/\log(3q'qC(\pi))$ of $s = 1$, then since $q \geq q'$, $L_\chi(s)$ has no real zero within a distance of $\frac{1}{2}c_{17}/\log(3qC(\pi))$ of $s = 1$. Again, the desired result follows by the preceding case. Finally, suppose that $L_\chi(s)$ has a real zero within a distance of $c_{17}/\log(3q'qC(\pi))$ of $s = 1$. At this stage, we assume that $\chi \neq \chi'$.

It follows from analysis nearly identical to the second case in Lemma 8.5 (with $L(s, \Pi^*)$ replacing $D(s)$) that $L(s, \Pi^*)$ has at most one real zero within distance $c_{17}/\log(3q'qC(\pi))$

from 1. Since we have supposed that $L_\chi(s)$ has a real zero within distance $c_{17}/\log(3q'qC(\pi))$ of $s = 1$, the above discussion indicates that this is the sole real zero for $L(s, \Pi^*)$ within a distance of $c_{17}/\log(3q'qC(\pi))$ of $s = 1$. It follows that the zero β of $L_{\chi'}(s)$ must satisfy

$$\beta \leq 1 - \frac{c_{17}}{\log(3q'qC(\pi))}, \quad \beta \in \left(1 - \frac{\varepsilon}{2}, 1\right).$$

Since $L_{\chi'}(\beta) = 0$, it follows that $L(\beta, \Pi^*) = 0$. Using (8.17), the above bounds on β , and the fact that $q \geq q'$, we find that

$$1 \ll_{\pi, \varepsilon} L_\chi(1)q^{2\varepsilon}x^{1-\beta} + q^{n^2+2\varepsilon}x^{\frac{1}{2}-\beta} \ll_{\pi, \chi', \varepsilon} L_\chi(1)q^{2\varepsilon}x^{\frac{\varepsilon}{2}} + q^{n^2+2\varepsilon}x^{\frac{\varepsilon}{2}-\frac{1}{2}}$$

Choosing $x = q^{2n^2}/L_\chi(1)^2$ (which is at least 3 by (3.10) and (3.13)) and solving for $L_\chi(1)$, we find that for all $\chi \neq \chi'$ and all $0 < \varepsilon < 1$, there exists a constant $d_\pi(\varepsilon) > 0$ such that $L_\chi(1) \geq d_\pi(\varepsilon)q^{-\varepsilon(n^2+2)/(1-\varepsilon)}$. Upon rescaling ε to $\varepsilon/(n^2+2+\varepsilon)$, we have $L_\chi(1) \geq d_\pi(\varepsilon/(n^2+2+\varepsilon))q^{-\varepsilon}$. As long as $\chi \neq \chi'$, the constant $d_\pi(\varepsilon/(n^2+2+\varepsilon))$ is effective. Once we decrease $d_\pi(\varepsilon/(n^2+2+\varepsilon))$ suitably to account for the case where $\chi = \chi'$, the claimed result holds for arbitrary χ . \square

Corollary 8.8. *Recall the notation and hypotheses of Lemma 8.5. If β_1 exists, then for all $\varepsilon > 0$, there exists an (ineffective) constant $c_\pi(\varepsilon) > 0$ such that $\beta_1 \leq 1 - c_\pi(\varepsilon)q^{-\varepsilon}$.*

Proof. If β_1 exists, then there exists $\sigma \in [\beta_1, 1]$ such that $L'_\chi(\sigma)(1 - \beta_1) = L_\chi(1) \geq c'_\pi(\frac{\varepsilon}{2})q^{-\frac{\varepsilon}{2}}$ by Proposition 8.7 and the mean value theorem. The result follows once we establish the bound $L'_\chi(\sigma) \ll_{\pi, \varepsilon} q^{\frac{\varepsilon}{2}}$ for $\sigma \in [1 - b_n/\log(3qC(\pi)), 1]$, where $b_n > 0$ is a suitable constant depending at most on n . To prove this, we observe via Cauchy's integral formula that

$$L'_\chi(1) = \frac{1}{2\pi i} \int_{|z-1|=\frac{1}{\log q}} \frac{L_\chi(z)}{(z-1)^2} dz \ll (\log q) \max_{|\xi-1| \leq \frac{1}{\log q}} |L_\chi(\xi)|,$$

in which case the desired bound follows from (3.13). \square

We will show that among the primitive characters $\chi \pmod{q}$ with $q \leq Q$, we encounter very few with the property that $L_\chi(s)$ has an exceptional zero.

Lemma 8.9. *Let $Q \geq 3$. There exists an effectively computable constant $c_{20} = c_{20}(n) > 0$ such that there is at most one real nontrivial primitive Dirichlet character $\chi_1 \pmod{q_1}$ with $q_1 \leq Q$ such that $L_{\chi_1}(s)$ has a real zero β_1 satisfying $\beta_1 > 1 - c_{20}/\log(C(\pi)Q)$.*

Proof. Suppose to the contrary that $\chi \pmod{q}$ and $\chi' \pmod{q'}$ are two distinct such characters with $q, q' \leq Q$. Let $\Pi = \pi \boxplus \pi \otimes \chi \boxplus \pi \otimes \chi'$, let ψ be the primitive character that induces $\chi'\chi$, and let

$$F(s) = L(s, \Pi \times \tilde{\Pi}) = L_1(s)^3 L_\chi(s)^2 L_{\chi'}(s)^2 L_\psi(s)^2.$$

By [21, Lemma a], the Dirichlet coefficients of $-\frac{F'}{F}(s)$ are nonnegative. By Lemma 3.2, there exists a constant $c_{21} = c_{21}(n) \geq 1$ such that if ω runs through the nontrivial zeros of $F(s)$ and $1 < \sigma < 2$, then

$$(8.18) \quad \sum_{\omega} \operatorname{Re} \left(\frac{1}{\sigma - \omega} \right) < \frac{3}{\sigma - 1} + c_{21} \log(C(\pi)Q).$$

If $\sigma = 1 + \frac{1}{2c_{21} \log(C(\pi)Q)}$ and M is the number (necessarily an integer) of real zeros (counting multiplicity) of $F(s)$ that are at least $1 - \frac{1}{14c_{21} \log(C(\pi)Q)}$, then it follows from (8.18) that

$$\frac{M}{1 + \frac{1}{2c_{21} \log(C(\pi)Q)} - \left(1 - \frac{1}{14c_{21} \log(C(\pi)Q)}\right)} < \frac{3}{\left(1 + \frac{1}{2c_{21} \log(C(\pi)Q)}\right) - 1} + c_{21} \log(C(\pi)Q).$$

This implies that $M \leq 3$. But if $L_\chi(s)$ and $L_{\chi'}(s)$ both have real zeros that are larger than $1 - \frac{1}{14c_{21} \log(C(\pi)Q)}$, then $F(s)$ has 4 such zeros, a contradiction. The lemma now follows. \square

Proof of Theorem 8.2. This follows from Corollary 8.8 and Lemmata 8.4, 8.5, and 8.9. \square

REFERENCES

- [1] W. D. Banks. Twisted symmetric-square L -functions and the nonexistence of Siegel zeros on $GL(3)$. *Duke Math. J.*, 87(2):343–353, 1997.
- [2] V. Blomer. Period integrals and Rankin–Selberg L -functions on $GL(n)$. *Geom. Funct. Anal.*, 22(3):608–620, 2012.
- [3] V. Blomer. Density theorems for $GL(n)$. *Invent. Math.*, 232(2):683–711, 2023.
- [4] V. Blomer and F. Brumley. On the Ramanujan conjecture over number fields. *Ann. of Math. (2)*, 174(1):581–605, 2011.
- [5] V. Blomer and F. Brumley. Non-vanishing of L -functions, the Ramanujan conjecture, and families of Hecke characters. *Canad. J. Math.*, 65(1):22–51, 2013.
- [6] E. Bombieri. On the large sieve. *Mathematika*, 12:201–225, 1965.
- [7] F. Brumley. *Distinguishing cusp forms on the general linear group*. ProQuest LLC, Ann Arbor, MI, 2004. Thesis (Ph.D.)—Princeton University.
- [8] F. Brumley. Effective multiplicity one on GL_N and narrow zero-free regions for Rankin–Selberg L -functions. *Amer. J. Math.*, 128(6):1455–1474, 2006.
- [9] F. Brumley. Second order average estimates on local data of cusp forms. *Arch. Math. (Basel)*, 87(1):19–32, 2006.
- [10] F. Brumley and D. Milićević. Counting cusp forms by analytic conductor. *Ann. Sci. Éc. Norm. Supér. (4)*. Accepted for publication.
- [11] F. Brumley, J. Thorner, and A. Zaman. Zeros of Rankin–Selberg L -functions at the edge of the critical strip. *J. Eur. Math. Soc. (JEMS)*, 24(5):1471–1541, 2022. With an appendix by Colin J. Bushnell and Guy Henniart.
- [12] D. Bump. *Lie groups*, volume 225 of *Graduate Texts in Mathematics*. Springer, New York, second edition, 2013.
- [13] C. J. Bushnell and G. Henniart. An upper bound on conductors for pairs. *J. Number Theory*, 65(2):183–196, 1997.
- [14] J.-M. Deshouillers and H. Iwaniec. Kloosterman sums and Fourier coefficients of cusp forms. *Invent. Math.*, 70(2):219–288, 1982/83.
- [15] W. Duke and E. Kowalski. A problem of Linnik for elliptic curves and mean-value estimates for automorphic representations. *Invent. Math.*, 139(1):1–39, 2000. With an appendix by Dinakar Ramakrishnan.
- [16] P. X. Gallagher. Bombieri’s mean value theorem. *Mathematika*, 15:1–6, 1968.
- [17] P. X. Gallagher. A large sieve density estimate near $\sigma = 1$. *Invent. Math.*, 11:329–339, 1970.
- [18] S. Gelbart and H. Jacquet. A relation between automorphic representations of $GL(2)$ and $GL(3)$. *Ann. Sci. École Norm. Sup. (4)*, 11(4):471–542, 1978.
- [19] R. Godement and H. Jacquet. *Zeta functions of simple algebras*. Lecture Notes in Mathematics, Vol. 260. Springer-Verlag, Berlin-New York, 1972.
- [20] J. Hoffstein and P. Lockhart. Coefficients of Maass forms and the Siegel zero. *Ann. of Math. (2)*, 140(1):161–181, 1994. With an appendix by Dorian Goldfeld, Jeffrey Hoffstein and Daniel Lieman.
- [21] J. Hoffstein and D. Ramakrishnan. Siegel zeros and cusp forms. *Int. Math. Res. Not. IMRN*, 1995(6):279–308, 1995.
- [22] G. Hoheisel. Primzahl probleme in der Analysis. *S.-B. Preuss. Akad. Wiss.*, 8:580–588, 1930.

- [23] P. Humphries. Standard zero-free regions for Rankin-Selberg L -functions via sieve theory. *Math. Z.*, 292(3–4):1105–1122, 2019. With an appendix by Farrell Brumley.
- [24] P. Humphries and J. Thorner. Towards a GL_n variant of the Hoheisel phenomenon. *Trans. Amer. Math. Soc.*, 375(3):1801–1824, 2022.
- [25] H. Iwaniec and E. Kowalski. *Analytic number theory*, volume 53 of *American Mathematical Society Colloquium Publications*. American Mathematical Society, Providence, RI, 2004.
- [26] H. Iwaniec and P. Sarnak. Perspectives on the analytic theory of L -functions. *Geom. Funct. Anal.*, (Special Volume, Part II):705–741, 2000. GAFA 2000 (Tel Aviv, 1999).
- [27] H. Jacquet, I. I. Piatetskii-Shapiro, and J. A. Shalika. Rankin-Selberg convolutions. *Amer. J. Math.*, 105(2):367–464, 1983.
- [28] H. Jacquet and J. A. Shalika. On Euler products and the classification of automorphic representations. I. *Amer. J. Math.*, 103(3):499–558, 1981.
- [29] S. Jana. Applications of analytic newvectors for $GL(n)$. *Math. Ann.*, 380(3-4):915–952, 2021.
- [30] S. Jana. The second moment of $GL(n) \times GL(n)$ Rankin-Selberg L -functions. *Forum Math. Sigma*, 10:Paper No. e47, 39, 2022.
- [31] Y. Jiang, G. Lü, J. Thorner, and Z. Wang. A Bombieri-Vinogradov theorem for higher-rank groups. *Int. Math. Res. Not. IMRN*, 2023(1):482–535, 2023.
- [32] E. Lapid. On the Harish-Chandra Schwartz space of $G(F)\backslash G(\mathbb{A})$. In *Automorphic representations and L -functions*, volume 22 of *Tata Inst. Fundam. Res. Stud. Math.*, pages 335–377. Tata Inst. Fund. Res., Mumbai, 2013. With an appendix by Farrell Brumley.
- [33] X. Li. Upper bounds on L -functions at the edge of the critical strip. *Int. Math. Res. Not. IMRN*, 2010(4):727–755, 2010.
- [34] U. V. Linnik. On the least prime in an arithmetic progression. *Rec. Math. [Mat. Sbornik] N.S.*, 15(57):139–178, 347–368, 1944.
- [35] J. Liu and Y. Wang. A theorem on analytic strong multiplicity one. *J. Number Theory*, 129(8):1874–1882, 2009.
- [36] W. Luo, Z. Rudnick, and P. Sarnak. On Selberg’s eigenvalue conjecture. *Geom. Funct. Anal.*, 5(2):387–401, 1995.
- [37] W. Luo, Z. Rudnick, and P. Sarnak. On the generalized Ramanujan conjecture for $GL(n)$. In *Automorphic forms, automorphic representations, and arithmetic (Fort Worth, TX, 1996)*, volume 66 of *Proc. Sympos. Pure Math.*, pages 301–310. Amer. Math. Soc., Providence, RI, 1999.
- [38] P. Michel and A. Venkatesh. The subconvexity problem for GL_2 . *Publ. Math. Inst. Hautes Études Sci.*, 111(111):171–271, 2010.
- [39] C. Mœglin and J.-L. Waldspurger. Le spectre résiduel de $GL(n)$. *Ann. Sci. École Norm. Sup. (4)*, 22(4):605–674, 1989.
- [40] C. J. Moreno. Analytic proof of the strong multiplicity one theorem. *Amer. J. Math.*, 107(1):163–206, 1985.
- [41] W. Müller and B. Speh. Absolute convergence of the spectral side of the Arthur trace formula for GL_n . *Geom. Funct. Anal.*, 14(1):58–93, 2004. With an appendix by E. M. Lapid.
- [42] M. R. Murty and V. K. Murty. A variant of the Bombieri-Vinogradov theorem. In *Number theory (Montreal, Que., 1985)*, volume 7 of *CMS Conf. Proc.*, pages 243–272. Amer. Math. Soc., Providence, RI, 1987.
- [43] I. I. Piatetski-Shapiro. Multiplicity one theorems. In *Automorphic forms, representations and L -functions (Proc. Sympos. Pure Math., Oregon State Univ., Corvallis, Ore., 1977), Part 1*, Proc. Sympos. Pure Math., XXXIII, pages 209–212. Amer. Math. Soc., Providence, R.I., 1979.
- [44] D. Ramakrishnan. Modularity of the Rankin-Selberg L -series, and multiplicity one for $SL(2)$. *Ann. of Math. (2)*, 152(1):45–111, 2000.
- [45] K. Soundararajan and J. Thorner. Weak subconvexity without a Ramanujan hypothesis. *Duke Math. J.*, 168:1231–1268, 2019. With an appendix by Farrell Brumley.
- [46] J. Thorner and A. Zaman. A unified and improved Chebotarev density theorem. *Algebra Number Theory*, 13(5):1039–1068, 2019.
- [47] J. Thorner and A. Zaman. An unconditional GL_n large sieve. *Adv. Math.*, 378:Paper No. 107529, 24, 2021.
- [48] A. Weiss. The least prime ideal. *J. Reine Angew. Math.*, 338:56–94, 1983.

- [49] P.-J. Wong. Bombieri–Vinogradov theorems for modular forms and applications. *Mathematika*, 66(1):200–229, 2020.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF VIRGINIA, CHARLOTTESVILLE, VA 22904, USA

Email address: `pclhumphries@gmail.com`

URL: `https://sites.google.com/view/peterhumphries/`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ILLINOIS, URBANA, IL 61801, USA

Email address: `jesse.thorner@gmail.com`