

SOME  $q$ -SUPERCONGRUENCES FROM RAHMAN'S SUMMATION

YUDONG LIU AND XIAOXIA WANG\*

ABSTRACT. Inspired by the recent work on  $q$ -congruences and the quadratic summation formula of Rahman, we provide some new  $q$ -supercongruences. By taking  $q \rightarrow 1$  in one of our results, we obtain a new Ramanujan-type supercongruence, which has the same right-hand side as Van Hamme's (G.2) supercongruence for  $p \equiv 1 \pmod{4}$ . We also formulate some related challenging conjectures on supercongruences and  $q$ -supercongruences.

## 1. INTRODUCTION

In 1913, Ramanujan announced the following infinite sum in his first letter to Hardy (cf. [1, p. 25, Eq. (2)]),

$$\sum_{k=0}^{\infty} (8k+1) \frac{\left(\frac{1}{4}\right)_k^4}{k!^4} = \frac{2\sqrt{2}}{\sqrt{\pi} \Gamma\left(\frac{3}{4}\right)^2}, \quad (1.1)$$

where  $(a)_n = a(a+1)\cdots(a+n-1)$  denotes the Pochhammer symbol and  $\Gamma(x)$  is the Gamma function. Later, Hardy [15, p. 495] gave a proof of this infinite summation. Recently, Chen and Chu [2, Eq. (4)] reproved it by giving its  $q$ -analogue as follows:

$$\sum_{k=0}^{\infty} [8k+1] \frac{(q; q^4)_k^4}{(q^4; q^4)_k^4} q^{2k} = \frac{(q^5, q^3, q^3, q^3; q^4)_{\infty}}{(q^2, q^4, q^4, q^4; q^4)_{\infty}}. \quad (1.2)$$

They also proposed a similar  $q$ -infinite summation [2, Example(8)]:

$$\sum_{k=0}^{\infty} [6k+1] \frac{(q; q^4)_k (q; q^2)_k^3}{(q^2; q^2)_k (q^4; q^4)_k^3} q^{k^2+k} = \frac{(q^5, q^3, q^3, q^3; q^4)_{\infty}}{(q^2, q^4, q^4, q^4; q^4)_{\infty}}, \quad (1.3)$$

whose right-hand side is the same as (1.2). According to [5, p. 353, Eq. (I.36)],

$$\lim_{q \rightarrow 1^-} \frac{(q; q)_{\infty}}{(q^x; q)_{\infty}} (1-q)^{1-x} = \Gamma(x). \quad (1.4)$$

---

2010 *Mathematics Subject Classification.* Primary 33D15; Secondary 11A07, 11B65.

*Key words and phrases.* Basic hypergeometric series; supercongruences;  $q$ -congruences; cyclotomic polynomial; Rahman's summation formula;

This work is supported by National Natural Science Foundations of China (11661032).

\* Corresponding author.

Letting  $q \rightarrow 1^-$  in (1.3), Chen and Chu obtained the following Ramanujan-type  $\pi$ -formula:

$$\sum_{k=0}^{\infty} (6k+1) \frac{\left(\frac{1}{2}\right)_k^3 \left(\frac{1}{4}\right)_k}{4^k k!^4} = \frac{2\sqrt{2}}{\sqrt{\pi} \Gamma\left(\frac{3}{4}\right)^2}. \quad (1.5)$$

In 1997, Van Hamme [29] conjectured that (1.1) has a nice  $p$ -adic analogue:

$$\sum_{k=0}^{(p-1)/4} (8k+1) \frac{\left(\frac{1}{4}\right)_k^4}{k!^4} \equiv p \frac{\Gamma_p\left(\frac{1}{2}\right)\Gamma_p\left(\frac{1}{4}\right)}{\Gamma_p\left(\frac{3}{4}\right)} \pmod{p^3}, \quad \text{if } p \equiv 1 \pmod{4}, \quad (1.6)$$

where  $p$  is an odd prime and  $\Gamma_p(x)$  is the  $p$ -adic Gamma function [21]. In 2015, Swisher proved this supercongruence (1.6). Not long ago, the authors [19] reproved it by giving the following  $q$ -analogue:

$$\sum_{k=0}^m [8k+1] \frac{(q; q^4)_k^4}{(q^4; q^4)_k^4} q^{2k} \equiv \frac{(q^2; q^4)_{(n-1)/4}}{(q^4; q^4)_{(n-1)/4}} [n] q^{(1-n)/4} \pmod{[n]\Phi_n(q)^2}, \quad (1.7)$$

where  $m = (n-1)/4$  or  $n-1$ . Recently, Guo and Schlosser [9, Theorems 2] proved that, for any even integer  $d \geq 4$  and positive integer  $n$  with  $n \equiv -1 \pmod{d}$ ,

$$\sum_{k=0}^{n-1} [2dk+1] \frac{(q; q^d)_k^d}{(q^d; q^d)_k^d} q^{d(d-3)k/2} \equiv 0 \pmod{\Phi_n(q)^2}. \quad (1.8)$$

Setting  $d = 4$  in the above  $q$ -supercongruence, we have

$$\sum_{k=0}^{n-1} [8k+1] \frac{(q; q^4)_k^4}{(q^4; q^4)_k^4} q^{2k} \equiv 0 \pmod{\Phi_n(q)^2} \quad \text{for } n \equiv 3 \pmod{4}.$$

Here and in what follows,  $(a; q)_n = (1-a)(1-aq)\cdots(1-aq^{n-1})$  is the  $q$ -shifted factorial. For convenience, we write  $(a_1, a_2, \dots, a_m; q)_n = (a_1; q)_n (a_2; q)_n \cdots (a_m; q)_n$  for  $n \geq 0$  or  $n = \infty$ . Moreover,  $[n] := [n]_q = (1-q^n)/(1-q)$  denotes the  $q$ -integer. Furthermore,  $\Phi_n(q)$  is the  $n$ -th cyclotomic polynomial in  $q$ , which is defined as

$$\Phi_n(q) = \prod_{\substack{1 \leq k \leq n \\ \gcd(n, k) = 1}} (q - \zeta^k),$$

where  $\zeta$  is an  $n$ -th primitive root of unity.

In the past few years, Ramanujan's and Ramanujan-type  $\pi$ -formulas have been generalized to the  $q$ -world by many authors with various methods (see, for example, [2, 3, 8, 12, 13, 16, 17, 25, 28]). At the same time, the Ramanujan-type (super)congruences and their  $q$ -analogues have gradually aroused the research interest of many authors (see [4, 6, 7, 9–11, 14, 18, 23, 26, 27, 30, 31]). Nowadays, the 'creative microscoping' method recently introduced by Guo and Zudilin [14] makes the research of  $q$ -supercongruences much easier than before.

In this paper, we shall give two  $q$ -supercongruences on truncated forms of (1.3) via 'creative microscoping' method as follows.

**Theorem 1.** *Let  $n \equiv 1 \pmod{4}$  be a positive integer. Then, modulo  $[n]\Phi_n(q)^2$ ,*

$$\begin{aligned} & \sum_{k=0}^{(n-1)/2} [6k+1] \frac{(q; q^4)_k (q; q^2)_k^3}{(q^2; q^2)_k (q^4; q^4)_k^3} q^{k^2+k} \\ & \equiv \begin{cases} \frac{(q^2; q^4)_{\frac{(n-1)}{4}}}{(q^4; q^4)_{\frac{(n-1)}{4}}} [n] q^{(1-n)/4}, & \text{if } n \equiv 1 \pmod{4}; \\ 0, & \text{if } n \equiv 3 \pmod{4}, \end{cases} \end{aligned} \quad (1.9)$$

$$\begin{aligned} & \sum_{k=0}^{n-1} [6k+1] \frac{(q; q^4)_k (q; q^2)_k^3}{(q^2; q^2)_k (q^4; q^4)_k^3} q^{k^2+k} \\ & \equiv \begin{cases} \frac{(q^2; q^4)_{\frac{(n-1)}{4}}}{(q^4; q^4)_{\frac{(n-1)}{4}}} [n] q^{(1-n)/4}, & \text{if } n \equiv 1 \pmod{4}; \\ 0, & \text{if } n \equiv 3 \pmod{4}. \end{cases} \end{aligned} \quad (1.10)$$

Let  $n = p \equiv 1 \pmod{4}$  be a prime and let  $q \rightarrow 1$  in (1.9). We have

$$\sum_{k=0}^{(p-1)/2} (6k+1) \frac{(\frac{1}{2})_k^3 (\frac{1}{4})_k}{4^k k!^4} \equiv \frac{(\frac{1}{2})_{(p-1)/4}}{(1)_{(p-1)/4}} p \pmod{p^3}. \quad (1.11)$$

Notice that, the authors [19] proved the following congruence holds for prime  $p \equiv 1 \pmod{4}$ ,

$$\frac{(\frac{1}{2})_{(p-1)/4}}{(1)_{(p-1)/4}} p \equiv \frac{\Gamma_p(\frac{1}{2})\Gamma_p(\frac{1}{4})}{\Gamma_p(\frac{3}{4})} p \pmod{p^3}. \quad (1.12)$$

From (1.9), we immediately deduce that

$$\sum_{k=0}^{(p-1)/2} (6k+1) \frac{(\frac{1}{2})_k^3 (\frac{1}{4})_k}{4^k k!^4} \equiv \begin{cases} \frac{\Gamma_p(\frac{1}{2})\Gamma_p(\frac{1}{4})}{\Gamma_p(\frac{3}{4})} p \pmod{p^3}, & \text{if } p \equiv 1 \pmod{4}, \\ 0 \pmod{p^3}, & \text{if } p \equiv 3 \pmod{4}. \end{cases} \quad (1.13)$$

It is a remarkable fact that the infinite Ramanujan-type  $\pi$ -formulas (1.1) and (1.5) not only have the same summation, but also have similar  $p$ -adic analogues. It is easy to see that the following supercongruences from (1.10) hold true:

$$\sum_{k=0}^{p-1} (6k+1) \frac{(\frac{1}{2})_k^3 (\frac{1}{4})_k}{4^k k!^4} \equiv \begin{cases} \frac{\Gamma_p(\frac{1}{2})\Gamma_p(\frac{1}{4})}{\Gamma_p(\frac{3}{4})} p \pmod{p^3}, & \text{if } p \equiv 1 \pmod{4}, \\ 0 \pmod{p^3}, & \text{if } p \equiv 3 \pmod{4}. \end{cases}$$

Motivated by (1.7) and Theorem 1, we would like to propose the following challenging conjecture.

**Conjecture 1.** *Let  $n \equiv 1 \pmod{4}$  be a positive integer. Then*

$$\begin{aligned} \sum_{k=0}^{(n-1)/2} [6k+1] \frac{(q; q^4)_k (q; q^2)_k^3}{(q^2; q^2)_k (q^4; q^4)_k^3} q^{k^2+k} &\equiv \sum_{k=0}^{(n-1)/4} [8k+1] \frac{(q; q^4)_k^4}{(q^4; q^4)_k^4} q^{2k} \pmod{[n]\Phi_n(q)^3}, \\ \sum_{k=0}^{n-1} [6k+1] \frac{(q; q^4)_k (q; q^2)_k^3}{(q^2; q^2)_k (q^4; q^4)_k^3} q^{k^2+k} &\equiv \sum_{k=0}^{n-1} [8k+1] \frac{(q; q^4)_k^4}{(q^4; q^4)_k^4} q^{2k} \pmod{[n]\Phi_n(q)^4}. \end{aligned}$$

The rest of the paper is organized as follows. We shall prove Theorem 1 in the next section with the help of Rahman's summation formula. Generalizations of Theorem 1 will be given in Section 3. Finally, in Section 4, we will propose similar  $q$ -congruences, a Ramanujan-type  $\pi$ -series and a related conjecture on  $q$ -supercongruences.

## 2. PROOF OF THEOREM 1

We need the following lemmas in our proof of Theorems 1.

**Lemma 1.** *Let  $n, d$  be positive integers with  $n > 1$  and  $n \equiv 1 \pmod{2d}$ . Then*

$$\sum_{k=0}^{(n-1)/d} [3dk+1] \frac{(q; q^{2d})_k (q^{d-1}; q^d)_k (aq, q/a; q^d)_k}{(q^d; q^d)_k (q^{d+2}; q^{2d})_k (aq^{2d}, q^{2d}/a; q^{2d})_k} q^{d(k^2+k)/2} \equiv 0 \pmod{\Phi_n(q)}. \quad (2.1)$$

*Proof.* We will make use of a quadratic summation formula of Rahman [24, Eq. (1.8)], stated as follow:

$$\sum_{k=0}^{\infty} \frac{1 - aq^{3k}}{1 - a} \frac{(a; q^2)_k (qa/bd; q)_k (b, d; q)_k}{(q; q)_k (qbd; q^2)_k (aq^2/b, aq^2/d; q^2)_k} q^{(k^2+k)/2} = \frac{(aq^2, qb, qd, aq^2/bd; q^2)_{\infty}}{(q, q^2a/b, q^2a/d, qbd; q^2)_{\infty}}. \quad (2.2)$$

By letting  $a = b = d = q^{\frac{1}{2}}$  and  $q \rightarrow q^2$  in (2.2), Chen and Chu [2, Eq. (4)] proved (1.2). Here, setting  $q \rightarrow q^d$ ,  $a = q^{1-n}$ ,  $b = aq$  and  $d = q/a$  in (2.2), then for  $n \equiv 1 \pmod{2d}$ , we have

$$\begin{aligned} &\sum_{k=0}^{(n-1)/2d} \frac{1 - q^{3dk+1-n}}{1 - q^{1-n}} \frac{(q^{1-n}; q^{2d})_k (q^{d-1-n}; q^d)_k (aq, q/a; q^d)_k}{(q^d; q^d)_k (q^{d+2}; q^{2d})_k (aq^{2d-n}, q^{2d-n}/a; q^{2d})_k} q^{d(k^2+k)/2} \\ &= \frac{(q^{2d-1-n}, q^{2d+1-n}, q^{d+1}/a, aq^{d+1}; q^{2d})_{\infty}}{(q^d, q^{d+2}, q^{2d-n}/a, aq^{2d-n}; q^{2d})_{\infty}}. \end{aligned} \quad (2.3)$$

It is easy to check that  $(q^{2d+1-n}; q^{2d})_{\infty} = 0$  for  $n \equiv 1 \pmod{2d}$ , thus the right-hand side of the above  $q$ -congruence is equal to 0. Since  $q^n \equiv 1 \pmod{\Phi_n(q)}$  and for  $(n-1)/2d < k \leq (n-1)/d$ ,  $(q; q^{2d})_k / ((q^d; q^d)_k (q^{d+2}; q^{2d})_k) \equiv 0 \pmod{\Phi_n(q)}$ , we get

$$\sum_{k=0}^{(n-1)/d} [3dk+1] \frac{(q; q^{2d})_k (q^{d-1}; q^d)_k (aq, q/a; q^d)_k}{(q^d; q^d)_k (q^{d+2}; q^{2d})_k (aq^{2d}, q^{2d}/a; q^{2d})_k} q^{d(k^2+k)/2} \equiv 0 \pmod{\Phi_n(q)},$$

as desired.  $\square$

**Lemma 2.** *Let  $n, d$  be positive integers with  $n > 1$  and  $n \equiv -1 \pmod{2d}$ . Then*

$$\sum_{k=0}^{(n+1)/d-1} [3dk+1] \frac{(q; q^{2d})_k (q^{d-1}; q^2)_k (aq, q/a; q^2)_k}{(q^d; q^d)_k (q^{d+2}; q^{2d})_k (aq^{2d}, q^{2d}/a; q^{2d})_k} q^{d(k^2+k)/2} \equiv 0 \pmod{\Phi_n(q)}. \quad (2.4)$$

*Proof.* Take same parameters transformation in (2.2) as the proof of Lemma 1, the left-hand side of (2.2) is truncated at  $(n+1)/d - 1$  place and the right-hand side is same to (2.3), which is equal to 0 for  $n \equiv -1 \pmod{2d}$ . Then applying  $q^n \equiv 1 \pmod{\Phi_n(q)}$ , we immediately obtain (2.4).  $\square$

In order to prove Theorem 1, we also need to establish the following parametric generalization.

**Theorem 2.** *Let  $n$  be a positive odd integer. Then, modulo  $\Phi_n(q)(1 - aq^n)(a - q^n)$ ,*

$$\begin{aligned} & \sum_{k=0}^{(n-1)/2} [6k+1] \frac{(q; q^4)_k (q; q^2)_k (aq, q/a; q^2)_k}{(q^2; q^2)_k (q^4; q^4)_k (aq^4, q^4/a; q^4)_k} q^{k^2+k} \\ & \equiv \begin{cases} \frac{(q^2; q^4)_{(n-1)/4} [n] q^{(1-n)/4}}{(q^4; q^4)_{(n-1)/4}}, & \text{if } n \equiv 1 \pmod{4}, \\ 0, & \text{if } n \equiv 3 \pmod{4}. \end{cases} \end{aligned} \quad (2.5)$$

*Proof.* For  $a = q^n$  or  $a = q^{-n}$ , the left-hand side of (2.5) is equal to

$$\begin{aligned} & \sum_{k=0}^{(n-1)/2} [6k+1] \frac{(q; q^4)_k (q; q^2)_k (q^{1-n}, q^{1+n}; q^2)_k}{(q^2; q^2)_k (q^4; q^4)_k (q^{4-n}, q^{4+n}; q^4)_k} q^{k^2+k} \\ & = \frac{(q^5, q^3, q^{3-n}, q^{3+n}; q^4)_\infty}{(q^2, q^4, q^{4-n}, q^{4+n}; q^4)_\infty}, \end{aligned}$$

which is Rahman's summation formula (2.2) with the parameter substitutions  $q \mapsto q^2, a = q, b = q^{1-n}$  and  $d = q^{1+n}$ . It's easy to see that

$$\frac{(q^5, q^3, q^{3-n}, q^{3+n}; q^4)_\infty}{(q^2, q^4, q^{4-n}, q^{4+n}; q^4)_\infty} = \begin{cases} \frac{(q^2; q^4)_{(n-1)/4} [n] q^{(1-n)/4}}{(q^4; q^4)_{(n-1)/4}}, & \text{if } n \equiv 1 \pmod{4}, \\ 0, & \text{if } n \equiv 3 \pmod{4}. \end{cases}$$

This proves the truth of  $q$ -congruences (2.5) modulo  $(1 - aq^n)(a - q^n)$ . Finally, since  $\Phi_n(q), a - q^n$  and  $1 - aq^n$  are pairwise relatively prime polynomials, applying Lemmas 1 and 2 with  $d = 2$ , we finish the proof of Theorem 2.  $\square$

*Proof of Theorem 1.* For  $k$  in the range  $0 \leq k \leq (n-1)/2$ , the limits of the denominators on both sides of (2.5) related to  $a$  are relatively prime to  $\Phi_n(q)$  when  $a \rightarrow 1$ . On the other hand, the limit  $(1 - aq^n)(a - q^n)$  as  $a \rightarrow 1$  contains the factor  $\Phi_n(q)^2$ . Therefore, the limiting case  $a \rightarrow 1$  of (2.5) leads to (1.9) are true modulo  $\Phi_n(q)^3$ . Observing that

$(q; q^4)_k (q; q^2)_k^3 / ((q^2; q^2)_k (q^4; q^4)_k^3) \equiv 0 \pmod{\Phi_n(q)^3}$  for  $k$  in the range  $(n-1)/2 \leq k \leq (n-1)$ , we immediately get

$$\begin{aligned} & \sum_{k=0}^{n-1} [6k+1] \frac{(q; q^4)_k (q; q^2)_k^3}{(q^2; q^2)_k (q^4; q^4)_k^3} q^{k^2+k} \\ & \equiv \begin{cases} \frac{(q^2; q^4)_{(n-1)/4} [n] q^{(1-n)/4}}{(q^4; q^4)_{(n-1)/4}}, & \text{if } n \equiv 1 \pmod{4} \\ 0, & \text{if } n \equiv 3 \pmod{4} \end{cases} \pmod{\Phi_n(q)^3}. \end{aligned} \quad (2.6)$$

It remains to prove that the following congruences also hold modulo  $[n]$ , i.e.,

$$\begin{aligned} & \sum_{k=0}^m [6k+1] \frac{(q; q^4)_k (q; q^2)_k^3}{(q^2; q^2)_k (q^4; q^4)_k^3} q^{k^2+k} \\ & \equiv \begin{cases} \frac{(q^2; q^4)_{(n-1)/4} [n] q^{(1-n)/4}}{(q^4; q^4)_{(n-1)/4}}, & \text{if } n \equiv 1 \pmod{4}, \\ 0, & \text{if } n \equiv 3 \pmod{4}. \end{cases} \pmod{[n]}, \end{aligned} \quad (2.7)$$

where  $m$  equals  $(n-1)/2$  or  $n-1$ . For  $n > 1$ , let  $\zeta \neq 1$  be an  $n$ -th unity root, not necessarily primitive. Then  $\zeta$  must be a primitive  $m_1$ -th root of unity with  $m_1 | n$ . Let  $c_q(k)$  denote the  $k$ -th term on the left-hand side in (2.5), i.e.,

$$c_q(k) = [6k+1] \frac{(q; q^4)_k (q; q^2)_k (aq, q/a; q^2)_k}{(q^2; q^2)_k (q^4; q^4)_k (aq^4, q^4/a; q^4)_k} q^{k^2+k}. \quad (2.8)$$

Applying Lemmas 1 and 2 with  $d = 2$ ,  $n = m_1$ , and the fact that  $(q; q^2)_k / (q^2; q^2)_k \equiv 0 \pmod{\Phi_{m_1}(q)}$  for  $(m_1-1)/2 < k \leq m_1-1$ , we have

$$\sum_{k=0}^{(m_1-1)/2} c_\zeta(k) = \sum_{k=0}^{m_1-1} c_\zeta(k) = 0.$$

For  $0 \leq k \leq m_1-1$ , the following limit can be calculated:

$$\lim_{q \rightarrow \zeta} \frac{c_q(\ell m_1 + k)}{c_q(\ell m_1)} = c_\zeta(k).$$

Thus, we get

$$\sum_{k=0}^{(n-1)/2} c_\zeta(k) = \sum_{\ell=0}^{(n/m_1-3)/2} c_\zeta(\ell m_1) \sum_{k=0}^{m_1-1} c_\zeta(k) + \sum_{k=0}^{(m_1-1)/2} c_\zeta((n-m_1)/2 + k) = 0, \quad (2.9)$$

and

$$\sum_{k=0}^{n-1} c_\zeta(k) = \sum_{\ell=0}^{n/m_1-1} \sum_{k=0}^{m_1-1} c_\zeta(\ell m_1 + k) = \sum_{\ell=0}^{n/m_1-1} c_\zeta(\ell m_1) \sum_{k=0}^{m_1-1} c_\zeta(k) = 0. \quad (2.10)$$

Noting that

$$\prod_{m_1|n, m_1>1} \Phi_{m_1}(q) = [n],$$

we establish (2.7). Since the least common multiple of  $[n]$  and  $\Phi_n(q)^3$  is  $[n]\Phi_n(q)^2$ , we complete the proof of the theorem.  $\square$

### 3. GENERALIZATIONS OF THEOREM 1

In this section, we first shall give a generalization of  $q$ -supercongruences (1.9) for  $n \equiv 1 \pmod{4}$  modulo  $\Phi_n(q)^3$ .

**Theorem 3.** *Let  $d, n > 1$  be integers with  $n \equiv 1 \pmod{2d}$ . Then*

$$\begin{aligned} & \sum_{k=0}^{(n-1)/d} [3dk+1] \frac{(q; q^{2d})_k (q^{d-1}; q^d)_k (q; q^d)_k^2}{(q^d; q^d)_k (q^{d+2}; q^{2d})_k (q^{2d}; q^{2d})_k^2} q^{d(k^2+k)/2} \\ & \equiv \frac{(q^d; q^{2d})_{(n-1)/(2d)}}{(q^{d+2}; q^{2d})_{(n-1)/(2d)}} [n] q^{-(n-1)(d-1)/2d} \pmod{\Phi_n(q)^3}, \end{aligned} \quad (3.1)$$

$$\begin{aligned} & \sum_{k=0}^{n-1} [3dk+1] \frac{(q; q^{2d})_k (q^{d-1}; q^d)_k (q; q^d)_k^2}{(q^d; q^d)_k (q^{d+2}; q^{2d})_k (q^{2d}; q^{2d})_k^2} q^{d(k^2+k)/2} \\ & \equiv \frac{(q^d; q^{2d})_{(n-1)/(2d)}}{(q^{d+2}; q^{2d})_{(n-1)/(2d)}} [n] q^{-(n-1)(d-1)/2d} \pmod{\Phi_n(q)^3}. \end{aligned} \quad (3.2)$$

Similarly, in order to prove Theorem 3, we first establish the following parametric generalization of (3.1).

**Theorem 4.** *Let  $d, n > 1$  be positive integers with  $n \equiv 1 \pmod{2d}$ . Then, modulo  $\Phi_n(q)(1 - aq^n)(a - q^n)$*

$$\begin{aligned} & \sum_{k=0}^{(n-1)/d} [3dk+1] \frac{(q; q^{2d})_k (q^{d-1}; q^d)_k (aq, q/a; q^d)_k}{(q^d; q^d)_k (q^{d+2}; q^{2d})_k (aq^{2d}, q^{2d}/a; q^{2d})_k} q^{d(k^2+k)/2} \\ & \equiv \frac{(q^d; q^{2d})_{(n-1)/(2d)}}{(q^{d+2}; q^{2d})_{(n-1)/(2d)}} [n] q^{-(n-1)(d-1)/2d}. \end{aligned} \quad (3.3)$$

*Proof.* For  $a = q^n$  or  $a = q^{-n}$ , the left-hand side of (3.3) is equal to

$$\sum_{k=0}^{(n-1)/d} [3dk+1] \frac{(q; q^{2d})_k (q^{d-1}; q^d)_k (q^{1+n}, q^{1-n}; q^d)_k}{(q^d; q^d)_k (q^{d+2}; q^{2d})_k (q^{2d+n}, q^{2d-n}; q^{2d})_k} q^{d(k^2+k)/2}, \quad (3.4)$$

which is the left-hand side of (2.2) with the parameters replaced by  $q \mapsto q^d$ ,  $a = q$ ,  $b = q^{1-n}$  and  $d = q^{1+n}$ . Then, the right-hand side of (2.2) can be written as

$$\frac{(q^{2d-1}, q^{2d+1}, q^{d+1-n}, q^{d+1+n}; q^{2d})_\infty}{(q^d, q^{d+2}, q^{2d-n}, q^{2d+n}; q^{2d})_\infty} = \frac{(q^d; q^{2d})_{(n-1)/(2d)}}{(q^{d+2}; q^{2d})_{(n-1)/(2d)}} [n] q^{-(n-1)(d-1)/2d}.$$

This proves that the  $q$ -congruence (3.3) holds modulo  $1 - aq^n$  and  $a - q^n$ . Moreover, applying Lemma 1, we can prove that (3.3) is also true modulo  $\Phi_n(q)$ .  $\square$

*Proof of Theorem 3.* Similarly as before, letting  $a \rightarrow 1$ , we can see the denominator of (3.3) is relatively to  $\Phi_n(q)$ . Also, the limit of  $(1 - aq^n)(a - q^n)$  as  $a \rightarrow 1$  has factor  $\Phi_n(q)^2$ . This means that (3.1) modulo  $\Phi_n(q)^3$  holds true. On the other hand, since  $(q; q^{2d})_k (q; q^d)_k^2 / ((q^d; q^d)_k (q^{2d}; q^{2d})_k^2) \equiv 0 \pmod{\Phi_n(q)^3}$  for  $k$  in the range  $(n-1)/d \leq k \leq (n-1)$ , the  $q$ -supercongruence (3.2) directly follows from (3.1).  $\square$

We also have the following generalization of (1.9) modulo  $\Phi_n(q)^2$  for  $n \equiv 3 \pmod{4}$ .

**Theorem 5.** *Let  $d$  and  $n > 1$  be positive integers with  $n \equiv d + 1 \pmod{2d}$ . Then*

$$\sum_{k=0}^{(n-1)/d} [3dk + 1] \frac{(q; q^{2d})_k (q^{d-1}; q^d)_k (q; q^d)_k^2}{(q^d; q^d)_k (q^{d+2}; q^{2d})_k (q^{2d}; q^{2d})_k^2} q^{d(k^2+k)/2} \equiv 0 \pmod{\Phi_n(q)^2} \quad (3.5)$$

*Proof.* The proof is the same as that of Theorem 4 and we need to establish the following one more parameter extension:

$$\sum_{k=0}^{(n-1)/d} [3dk + 1] \frac{(q; q^{2d})_k (q^{d-1}; q^d)_k (aq, q/a; q^d)_k}{(q^d; q^d)_k (q^{d+2}; q^{2d})_k (aq^{2d}, q^{2d}/a; q^{2d})_k} q^{d(k^2+k)/2} \equiv 0 \pmod{(a - q^n)(1 - aq^n)}. \quad (3.6)$$

Noticing that, for  $n \equiv d + 1 \pmod{2d}$ ,

$$\frac{(q^{2d+1}, q^{2d-1}, q^{d+1-n}, q^{d+1+n}; q^{2d})_\infty}{(q^d, q^{d+2}, q^{2d-n}, q^{2d+n}; q^{2d})_\infty} = 0,$$

we immediately obtain (3.6) by taking  $q \mapsto q^d$ ,  $a = q$ ,  $b = q^{1-n}$  and  $d = q^{1+n}$  in Rahman's summation (2.2). Then letting  $a \rightarrow 1$ , we are led to (3.5).  $\square$

Letting  $n = p$  be an odd prime,  $d = 4$  and  $q \rightarrow 1$  in Theorem 5, we get the following result.

**Corollary 6.** *Let  $p \equiv 5 \pmod{8}$  be a prime. Then*

$$\sum_{k=0}^{(p-1)/4} (12k + 1) \frac{\left(\frac{1}{4}\right)_k^2 \left(\frac{1}{8}\right)_k}{4^k k!^3} \equiv 0 \pmod{p^2}. \quad (3.7)$$

By the numerical calculations, we believe that the following stronger version is also true.



**Conjecture 2.** *Let  $p \equiv 5 \pmod{8}$  be a prime. Then*

$$\sum_{k=0}^{p-1} (12k+1) \frac{\left(\frac{1}{4}\right)_k^2 \left(\frac{1}{8}\right)_k}{4^k k!^3} \equiv 0 \pmod{p^3}.$$

#### 4. OTHER CONCLUSIONS FROM RAHMAN'S SUMMATION

In this section, we present similar  $q$ -supercongruences and a  $\pi$ -formula from Rahman's quadratic summation formula (2.2).

**Theorem 7.** *Let  $n$  and  $d \neq 1$  be positive integers. Then, module  $\Phi_n(q)^2$ ,*

$$\begin{aligned} & \sum_{k=0}^{(n-1)/d} [3dk-1] \frac{(q^{-1}; q^{2d})_k (q^{d-3}; q^d)_k (q; q^d)_k^2}{(q^d; q^d)_k (q^{d+2}; q^{2d})_k (q^{2d-2}; q^{2d})_k^2} q^{d(k^2+k)/2+1} \\ & \equiv \begin{cases} -\frac{(q^d; q^{2d})_{(n-1)/(2d)} (q^{2d-1}; q^{2d})_{(n-1)/(2d)}}{(q^3; q^{2d})_{(n-1)/(2d)} (q^{d+2}; q^{2d})_{(n-1)/(2d)}} q^{(3-d)(n-1)/2d}, & \text{if } n \equiv 1 \pmod{2d}, \\ 0, & \text{if } n \equiv d+1 \pmod{2d}. \end{cases} \end{aligned} \quad (4.1)$$

*Proof.* We first establish the following congruence module  $(a - q^n)(1 - aq^n)$ :

$$\begin{aligned} & \sum_{k=0}^{(n-1)/d} [3dk-1] \frac{(q^{-1}; q^{2d})_k (q^{d-3}; q^d)_k (q/a; q^d)_k (aq; q^d)_k}{(q^d; q^d)_k (q^{d+2}; q^{2d})_k (aq^{2d-2}; q^{2d})_k (q^{2d-2}/a; q^{2d})_k} q^{d(k^2+k)/2+1} \\ & \equiv \begin{cases} -\frac{(q^d; q^{2d})_{(n-1)/(2d)} (q^{2d-1}; q^{2d})_{(n-1)/(2d)}}{(q^3; q^{2d})_{(n-1)/(2d)} (q^{d+2}; q^{2d})_{(n-1)/(2d)}} q^{(3-d)(n-1)/2d}, & \text{if } n \equiv 1 \pmod{2d}, \\ 0, & \text{if } n \equiv d+1 \pmod{2d}. \end{cases} \end{aligned} \quad (4.2)$$

Letting  $q \mapsto q^d$ ,  $a = q^{-1}$ ,  $b = q^{1-n}$  and  $d = q^{1+n}$  in (2.2), we obtain

$$\begin{aligned} & \sum_{k=0}^{(n-1)/d} [3dk-1] \frac{(q^{-1}; q^{2d})_k (q^{d-3}; q^d)_k (q^{1-n}; q^d)_k (q^{1+n}; q^d)_k}{(q^d; q^d)_k (q^{d+2}; q^{2d})_k (q^{2d-2+n}; q^{2d})_k (q^{2d-2-n}; q^{2d})_k} q^{d(k^2+k)/2+1} \\ & = -\frac{(q^{2d-1}, q^{2d-3}, q^{d+1-n}, q^{d+1+n}; q^{2d})_\infty}{(q^d, q^{d+2}, q^{2d-2-n}, q^{2d-2+n}; q^{2d})_\infty} \\ & = \begin{cases} -\frac{(q^d; q^{2d})_{(n-1)/(2d)} (q^{2d-1}; q^{2d})_{(n-1)/(2d)}}{(q^3; q^{2d})_{(n-1)/(2d)} (q^{d+2}; q^{2d})_{(n-1)/(2d)}} q^{(3-d)(n-1)/2d}, & \text{if } n \equiv 1 \pmod{2d}, \\ 0, & \text{if } n \equiv d+1 \pmod{2d}. \end{cases} \end{aligned}$$

This proves that the  $q$ -congruences generalization (4.2) with one more parameter hold true. The proof then follows by taking limits as  $a \rightarrow 1$  in (4.2).  $\square$

Specially, when  $d = 2$  in Theorem 7, modulo  $\Phi_n(q)^2$ , we have

$$\begin{aligned} & \sum_{k=0}^{(n-1)/2} [6k-1] \frac{(q^{-1}; q^4)_k (q^{-1}; q^2)_k (q; q^2)_k^2}{(q^2; q^2)_k (q^4; q^4)_k (q^2; q^4)_k^2} q^{k^2+k+1} \\ & \equiv \begin{cases} -\frac{(q^2; q^4)_{(n-1)/4}}{(q^4; q^4)_{(n-1)/4}} q^{(n-1)/4}, & \text{if } n \equiv 1 \pmod{4}, \\ 0, & \text{if } n \equiv 3 \pmod{4}. \end{cases} \end{aligned} \quad (4.3)$$

Furthermore, we believe that the following stronger  $q$ -supercongruences are true.

**Conjecture 3.** *Let  $n$  be a positive odd integer. Then, modulo  $[n]\Phi_n(q)^2$ ,*

$$\begin{aligned} & \sum_{k=0}^{(n-1)/2} [6k-1] \frac{(q^{-1}; q^4)_k (q^{-1}; q^2)_k (q; q^2)_k^2}{(q^2; q^2)_k (q^4; q^4)_k (q^2; q^4)_k^2} q^{k^2+k+1} \\ & \equiv \begin{cases} -\frac{(q^2; q^4)_{(n-1)/4}}{(q^4; q^4)_{(n-1)/4}} q^{(n-1)/4}, & \text{if } n \equiv 1 \pmod{4}, \\ 0, & \text{if } n \equiv 3 \pmod{4}. \end{cases} \end{aligned} \quad (4.4)$$

Letting  $n = p$  be a prime,  $q \rightarrow 1$  in the above  $q$ -supercongruences, and applying (1.12) again, we have

$$\sum_{k=0}^{(p-1)/2} (6k-1) \frac{(-\frac{1}{2})_k (-\frac{1}{4})_k}{4^k k!^2} \equiv \begin{cases} -\frac{\Gamma_p(\frac{1}{2})\Gamma_p(\frac{1}{4})}{\Gamma_p(\frac{3}{4})} \pmod{p^3}, & \text{if } p \equiv 1 \pmod{4}, \\ 0 \pmod{p^3}, & \text{if } p \equiv 3 \pmod{4}. \end{cases} \quad (4.5)$$

Meanwhile, replacing  $q$  by  $q^2$ ,  $a = q^{-1}$  and  $b = d = q$  in (2.2), we get an infinity summation formula:

$$\sum_{k=0}^{\infty} [6k-1] \frac{(q^{-1}; q^4)_k (q^{-1}; q^2)_k (q; q^2)_k^2}{(q^2; q^2)_k (q^4; q^4)_k (q^2; q^4)_k^2} q^{k^2+k+1} = -\frac{(q^3, q^3, q^3, q; q^4)_{\infty}}{(q^2, q^2, q^2, q^4; q^4)_{\infty}}. \quad (4.6)$$

Applying (1.4) and noting that  $\Gamma(x)\Gamma(1-x) = \pi/\sin(\pi x)$ , then letting  $q \rightarrow 1^-$  in (4.6), we shall obtain the following Ramanujan-type summation of  $\pi$  as

$$\begin{aligned} \sum_{k=0}^{\infty} (6k-1) \frac{(-\frac{1}{2})_k (-\frac{1}{4})_k}{4^k k!^2} &= -\frac{\Gamma(\frac{1}{2})^3 \Gamma(1)}{\Gamma(\frac{3}{4})^3 \Gamma(\frac{1}{4})} \\ &= -\frac{\sqrt{2\pi}}{2\Gamma^2(\frac{3}{4})}. \end{aligned} \quad (4.7)$$

Thus, the supercongruence (4.5) may be deemed a nice  $p$ -adic analogue of (4.7).

## REFERENCES

- [1] B.C. Berndt and R.A. Rankin, *Ramanujan, Letters and Commentary*, History of Mathematics **9**, Amer. Math. Soc., Providence, RI; London Math. Soc., London, 1995.
- [2] X. Chen and W. Chu, Hidden  $q$ -analogues of Ramanujan-like  $\pi$ -series, *Ramanujan J.* (2020) <https://doi.org/10.1007/s11139-019-00192-7>
- [3] W. Chu,  $q$ -Series reciprocities and further  $\pi$ -formula, *Kodai Math.* **41** (3) (2018), 512–530.
- [4] O. Gorodetsky,  $q$ -Congruences, with applications to supercongruences and the cyclic sieving phenomenon, *Int. J. Number Theory* **15** (9)(2019), 1919–1968.
- [5] G. Gasper and M. Rahman, *Basic hypergeometric series*, second edition, Encyclopedia of Mathematics and Its Applications 96, Cambridge University Press, Cambridge, 2004.
- [6] V.J.W. Guo,  $q$ -Analogues of Dwork-type supercongruences, *J. Math. Anal. Appl.* **487** (2020), Art. 124022.
- [7] V.J.W. Guo, Proof of some  $q$ -supercongruences modulo the fourth power of a cyclotomic polynomial, *Results Math.* **75** (2020), Art. 77.
- [8] V.J.W. Guo,  $q$ -Analogues of three Ramanujan-type formulas for  $1/\pi$ , *Ramanujan J.* **52** (1) (2020), 123–132.
- [9] V.J.W. Guo and M.J. Schlosser, Some new  $q$ -congruences for truncated basic hypergeometric series: even powers, *Results Math.* **75** (2020), Art. 1.
- [10] V.J.W. Guo and M.J. Schlosser, A family of  $q$ -hypergeometric congruences modulo the fourth power of a cyclotomic polynomial, *Israel J. Math.* (to appear).
- [11] V.J.W. Guo and M.J. Schlosser, Some  $q$ -supercongruences from transformation formulas for basic hypergeometric series, *Constr. Approx.* (to appear).
- [12] V.J.W. Guo and J.-C. Liu,  $q$ -Analogues of two Ramanujan-type formulas for  $1/\pi$ , *J. Difference Equ. Appl.* **24** (8) (2018), 1368–1373.
- [13] V.J.W. Guo and W. Zudilin, Ramanujan-type formulae for  $1/\pi$ :  $q$ -analogues, *Integral Transform Spec. Funct.* **29** (7) (2018), 505–513.
- [14] V.J.W. Guo and W. Zudilin, A  $q$ -microscope for supercongruences, *Adv. Math.* **346** (2019), 329–358.
- [15] G.H. Hardy, A chapter from Ramanujan's note-book, *Proc. Cambridge Philos. Soc.* **21** (2) (1923), 492–503.
- [16] Q.-H. Hou, C. Krattenthaler and Z.-W. Sun, On  $q$ -analogues of some series for  $\pi$  and  $\pi^2$ , *Proc. Amer. Math. Soc.* **147** (5) (2019), 1953–1961.
- [17] J. Guillera, Hypergeometric identities for 10 extended Ramanujan-type series, *Ramanujan J.* **15** (2008), 219–234.
- [18] J.-C. Liu and F. Petrov, Congruences on sums of  $q$ -binomial coefficient, *Adv. Appl. Math.* **116** (2020), Art. 102003.
- [19] Y. Liu and X. Wang,  $q$ -Analogues of the (G.2) supercongruence of Van Hamme, preprint, August 2020; arXiv: 2008.01417.
- [20] D. McCarthy and R. Osburn, A  $p$ -adic analogue of a formula of Ramanujan. *Arch. Math. (Basel)* **91** (6) (2008), 492–504.
- [21] Y. Morita, A  $p$ -adic supercongruence of the  $\Gamma$  function, *J. Fac. Sci. Univ. Tokyo* **22** (1975), 255–266.
- [22] E. Mortenson, A  $p$ -adic supercongruence conjecture of van Hamme, *Proc. Amer. Math. Soc.* **136** (12) (2008), 4321–4328.
- [23] H.-X. Ni and H. Pan, On a conjectured  $q$ -congruence of Guo and Zeng, *Int. J. Number Theory* **14** (6) (2018), 1699–1707.
- [24] M. Rahman, Some quadratic and cubic summation formulas for basic hypergeometric series. *Canad J. Math.* **45** (2) (1993), 394–411.
- [25] Z.-W. Sun, Two  $q$ -analogues of Euler's formula  $\zeta(2) = \pi^2/6$ , *Colloq. Math.* **158** (2) (2019), 313–320.

- [26] R. Tauraso, Some  $q$ -analogs of congruences for central binomial sums, *Colloq. Math.* **133** (2013), 133–143.
- [27] X. Wang and M. Yue, A  $q$ -analogue of the (A.2) supercongruence of Van Hamme for any prime  $p \equiv 3 \pmod{4}$ , *Int. J. Number Theory* **16** (6) (2020), 1325–1335.
- [28] C. Wei,  $q$ -Analogues of several  $\pi$ -formulas, *Proc. Amer. Math. Soc.* **148** (2020), 2287–2296.
- [29] L. Van Hamme, Some conjectures concerning partial sums of generalized hypergeometric series, in:  *$p$ -Adic Functional Analysis* (Nijmegen, 1996), Lecture Notes in Pure and Appl. Math. **192**, Dekker, New York (1997), 223–236.
- [30] W. Zudilin, Ramanujan-type supercongruences, *J. Number Theory* **129** (8) (2009), 1848–1857.
- [31] W. Zudilin, Congruences for  $q$ -binomial coefficients, *Ann. Combin.* **23** (2019), 1123–1135.

DEPARTMENT OF APPLIED MATHEMATICS, SHANGHAI UNIVERSITY, SHANGHAI 200444, P. R. CHINA  
Email address: lydshdx@163.com (Y. Liu), xiaoxiawang@shu.edu.cn (X. Wang)