

# Prescribing oscillation behavior of solutions to the heat equation on $\mathbb{R}^n$ via the initial data and its average integral\*

Dong-Ho Tsai

March 12, 2021

## Abstract

Motivated by a classical stabilization result for solution to the Cauchy problem of the heat equation  $\partial_t u = \Delta u$  on  $\mathbb{R}^n$ , we consider its oscillation behavior with radial initial data  $\varphi(x) = \varphi(|x|) \in C^0(\mathbb{R}^n) \cap L^\infty(\mathbb{R}^n)$ . Given four arbitrary finite numbers  $r < \alpha < \beta < s$ , one can construct a radial  $\varphi \in C^0(\mathbb{R}^n) \cap L^\infty(\mathbb{R}^n)$  so that  $\varphi$  together with its corresponding solution  $u(x, t)$  satisfy the oscillation behavior:

$$\begin{aligned} \liminf_{\tau \rightarrow \infty} \varphi(\tau) = r < \liminf_{t \rightarrow \infty} u(0, t) = \alpha \\ < \limsup_{t \rightarrow \infty} u(0, t) = \beta < \limsup_{\tau \rightarrow \infty} \varphi(\tau) = s. \end{aligned}$$

Similarly, given  $p < \alpha < \beta < q$  with  $p+q = \alpha+\beta$ , one can find a radial  $\varphi \in C^0(\mathbb{R}^n) \cap L^\infty(\mathbb{R}^n)$  so that its average integral  $H(\tau)$  and  $u(x, t)$  satisfy

$$\begin{aligned} \liminf_{\tau \rightarrow \infty} H(\tau) = p < \liminf_{t \rightarrow \infty} u(0, t) = \alpha \\ < \limsup_{t \rightarrow \infty} u(0, t) = \beta < \limsup_{\tau \rightarrow \infty} H(\tau) = q. \end{aligned}$$

Here  $H(\tau)$ ,  $\tau \in (0, \infty)$ , is given by

$$H(\tau) = \frac{1}{|B(0, \tau)|} \int_{B(0, \tau)} \varphi(y) dy, \quad \tau \in (0, \infty)$$

and  $B(0, \tau)$  is the open ball with radius  $\tau > 0$  centered at the origin of  $\mathbb{R}^n$ .

## 1 Introduction.

This article is a continuation of our previous ones [TN, CT] and proves several new interesting results. Consider the initial value problem

$$\begin{cases} u_t(x, t) = \Delta u(x, t), & x \in \mathbb{R}^n, \quad t > 0, \\ u(x, 0) = \varphi(x), & x \in \mathbb{R}^n, \end{cases} \quad (1)$$

where  $\varphi : \mathbb{R}^n \rightarrow \mathbb{R}$  is a given continuous function. It is known that if  $\varphi : \mathbb{R}^n \rightarrow \mathbb{R}$  is a **continuous bounded** function, then the function given by convolution integral

$$u(x, t) = \frac{1}{(4\pi t)^{n/2}} \int_{\mathbb{R}^n} e^{-\frac{|x-y|^2}{4t}} \varphi(y) dy, \quad x \in \mathbb{R}^n, \quad t > 0 \quad (2)$$

---

\*Mathematics Subject Classification: 35K05, 35K15.

is a smooth solution of the heat equation on  $\mathbb{R}^n \times (0, \infty)$  with  $\lim_{(x,t) \rightarrow (x_0,0)} u(x,t) = \varphi(x_0)$  for any  $x_0 \in \mathbb{R}^n$ . Due to the example by Tychonoff in 1935 (see the book [J], Chapter 7), it is known that the Cauchy problem for the heat equation (1) has no unique solution on  $\mathbb{R}^n \times (0, \infty)$  even if the initial data  $\varphi(x)$  is bounded (unless we impose certain growth condition of  $u(x,t)$  as  $|x| \rightarrow \infty$  for  $t > 0$ ). From now on, when we say  $u(x,t)$  is "the solution" of the heat equation with  $u(x,0) = \varphi(x)$ ,  $x \in \mathbb{R}^n$ , we always mean that it is the solution given by the convolution integral (2). As a consequence, if  $|\varphi| \leq M$  on  $\mathbb{R}^n$  for some constant  $M > 0$ , we also have  $|u| \leq M$  on  $\mathbb{R}^n \times [0, \infty)$ .

In this paper, we shall always assume that  $\varphi \in C^0(\mathbb{R}^n) \cap L^\infty(\mathbb{R}^n)$  and use  $H(\tau)$ ,  $\tau \in [0, \infty)$ , to denote the **average integral** of  $\varphi$  over the open ball  $B(0, \tau) \subset \mathbb{R}^n$  centered at  $x = 0$  with radius  $\tau > 0$ , i.e.

$$H(\tau) = \frac{1}{|B(0, \tau)|} \int_{B(0, \tau)} \varphi(y) dy = \frac{1}{\omega(n) \tau^n} \int_{B(0, \tau)} \varphi(y) dy, \quad \tau \in (0, \infty), \quad (3)$$

where  $\omega(n) = |B(0, 1)|$  is the volume of the unit ball in  $\mathbb{R}^n$ . By continuity, if we define  $H(0) = \varphi(0)$ , then  $H \in C^0[0, \infty) \cap L^\infty[0, \infty)$ .

The role played by  $H(\tau)$  in the Cauchy problem (1) is the following beautiful **stabilization** result:

**Theorem 1** (See [E, K, RE].) *Assume  $\varphi \in C^0(\mathbb{R}^n) \cap L^\infty(\mathbb{R}^n)$  and  $u(x,t)$  is the solution of the heat equation (1) given by (2). Then*

$$\lim_{t \rightarrow \infty} u(0, t) = 0 \quad \text{if and only if} \quad \lim_{\tau \rightarrow \infty} \left( \frac{1}{|B(0, \tau)|} \int_{B(0, \tau)} \varphi(y) dy \right) = 0. \quad (4)$$

Moreover,

$$\lim_{t \rightarrow \infty} \left( \sup_{x \in \mathbb{R}^n} |u(x, t)| \right) = 0 \quad (5)$$

if and only if

$$\lim_{\tau \rightarrow \infty} \left( \sup_{x \in \mathbb{R}^n} \left| \frac{1}{|B(x, \tau)|} \int_{B(x, \tau)} \varphi(y) dy \right| \right) = 0. \quad (6)$$

Here  $|B(x, \tau)|$  is the volume of the open ball in  $\mathbb{R}^n$  centered at  $x \in \mathbb{R}^n$  with radius  $\tau > 0$ .

**Remark 2** *Similar result holds if we replace the limit value 0 in (4) by any real number  $c$ .*

**Remark 3** *As demonstrated in [CT], Theorem 1 fails if the initial data  $\varphi$  is **unbounded**. In this paper, we will always assume  $\varphi \in C^0(\mathbb{R}^n) \cap L^\infty(\mathbb{R}^n)$ .*

**Remark 4** *Since  $|\varphi| \leq M$  on  $\mathbb{R}^n$  for some constant  $M > 0$ , the convolution solution  $u(x, t)$  satisfies the following gradient estimate*

$$|\nabla_x u(x, t)| \leq \frac{c(n) M}{\sqrt{t}}, \quad \forall (x, t) \in \mathbb{R}^n \times (0, \infty), \quad (7)$$

where  $c(n)$  is a constant depending only on  $n$ . Hence if we have  $\lim_{t \rightarrow \infty} u(0, t) = 0$ , we also have  $\lim_{t \rightarrow \infty} u(x, t) = 0$  for all  $x \in \mathbb{R}^n$  and the convergence is uniform in  $x \in K$  where  $K \subset \mathbb{R}^n$  is any compact set.

By Theorem 1 and Remark 2, the  $\lim_{\tau \rightarrow \infty} H(\tau)$  does not exist if and only if  $\lim_{t \rightarrow \infty} u(0, t)$  does not exist. In this case, we have the following comparison result concerning the oscillation of  $u(0, t)$  and  $H(\tau)$  as  $t \rightarrow \infty$  and  $\tau \rightarrow \infty$  (see [CT]), namely

$$\liminf_{\tau \rightarrow \infty} H(\tau) \leq \liminf_{t \rightarrow \infty} u(0, t) < \limsup_{t \rightarrow \infty} u(0, t) \leq \limsup_{\tau \rightarrow \infty} H(\tau). \quad (8)$$

The key point in the proof of (8) is to use the **representation formula** (see Lemma 4 in [CT]):

$$u(0, t) = \frac{2\omega(n)}{\pi^{n/2}} \int_0^\infty e^{-z^2} z^{n+1} H(\sqrt{4tz}) dz, \quad \forall t \in (0, \infty) \quad (9)$$

together with the classical Fatou's Lemma. Same as in [CT], we shall use  $p \leq \alpha < \beta \leq q$  to denote the four finite limit values in (8) in this paper.

By analogy, can we also compare the oscillation of  $u(0, t)$ ,  $t \in [0, \infty)$ , with the oscillation of the initial data  $\varphi(x)$ ? Since  $\varphi(x)$  is a function defined on  $\mathbb{R}^n$ , it would be quite tricky unless we assume  $\varphi(x) = \varphi(\tau)$ ,  $\tau = |x| \in [0, \infty)$ ,  $x \in \mathbb{R}^n$ , is a **radial function**. With this, both  $u(0, t)$  and  $\varphi(\tau)$  are one-variable functions defined on  $[0, \infty)$  and we can compare their oscillations as  $t \rightarrow \infty$  and  $\tau \rightarrow \infty$  respectively. From now on, we shall assume that  $\varphi(x) = \varphi(|x|)$ ,  $x \in \mathbb{R}^n$ , is a radial function.

Clearly, the limit  $\lim_{\tau \rightarrow \infty} \varphi(\tau) = 0$  will imply  $\lim_{t \rightarrow \infty} u(0, t) = 0$ , but **not** conversely. For example, the solution to the one-dimensional heat equation  $u_t = u_{xx}$  with radial initial data  $\varphi(x) = \cos x$ ,  $x \in [0, \infty)$ , is given by  $u(x, t) = e^{-t} \cos x$ , which satisfies  $\lim_{t \rightarrow \infty} u(0, t) = 0$ . But we have  $\liminf_{x \rightarrow \infty} \varphi(x) = -1$ ,  $\limsup_{x \rightarrow \infty} \varphi(x) = 1$ .

For radial  $\varphi \in C^0[0, \infty) \cap L^\infty[0, \infty)$ , the identities (2) (at  $x = 0$ ) and (3) will become

$$u(0, t) = \frac{n\omega(n)}{\pi^{n/2}} \int_0^\infty e^{-z^2} z^{n-1} \varphi(\sqrt{4tz}) dz, \quad \forall t \in (0, \infty) \quad (10)$$

and

$$H(\tau) = \frac{n}{\tau^n} \int_0^\tau \varphi(r) r^{n-1} dr, \quad \tau \in (0, \infty), \quad H(0) = \varphi(0). \quad (11)$$

In particular, we note the similarity between (10) and (9).

Let  $r = \liminf_{\tau \rightarrow \infty} \varphi(\tau)$  and  $s = \limsup_{\tau \rightarrow \infty} \varphi(\tau)$ . We have the following general inequalities among the six numbers  $\alpha$ ,  $\beta$ ,  $p$ ,  $q$ ,  $r$ ,  $s$ :

$$r \leq p \leq \alpha \leq \beta \leq q \leq s, \quad (12)$$

where the first and last inequalities in (12) can be derived from the identity  $H(\tau) = \frac{n}{\tau^n} \int_0^\tau \varphi(r) r^{n-1} dr$  and the definition of  $\liminf$  and  $\limsup$ . Note that Theorem 1 says that  $\alpha = \beta$  if and only if  $p = q$ . However,  $\alpha = \beta$  does not necessarily imply  $r = s$  (as the above simple example shows), i.e. the stabilization of  $u(0, t)$  as  $t \rightarrow \infty$  does not imply the stabilization of  $\varphi(\tau)$  as  $\tau \rightarrow \infty$ .

In this paper we are interested in prescribing the oscillation values  $\alpha < \beta$  of  $u(0, t)$  as  $t \rightarrow \infty$  by choosing suitable radial  $\varphi(\tau)$  in (10) or by choosing suitable average integral  $H(\tau)$  in (9) (by (21) below, one can determine  $\varphi(\tau)$  as long as  $H(\tau)$  is chosen). There are many choices for such  $\varphi(\tau)$  and  $H(\tau)$ . The interesting results here are that one can also prescribe the oscillation values of  $\varphi(\tau)$  and  $H(\tau)$  as  $\tau \rightarrow \infty$ .

More precisely, the results in this paper are:

1. To prescribe, as  $\tau \rightarrow \infty$  and  $t \rightarrow \infty$ , the oscillation behavior of  $H(\tau)$  and  $u(0, t)$  for four arbitrary finite numbers  $p \leq \alpha < \beta \leq q$  satisfying the **symmetry condition**

$$p + q = \alpha + \beta. \quad (13)$$

2. To prescribe, as  $\tau \rightarrow \infty$  and  $t \rightarrow \infty$ , the oscillation behavior of  $\varphi(\tau)$  and  $u(0, t)$  for four arbitrary finite numbers  $r \leq \alpha \leq \beta \leq s$ .

See Theorem 5 and Theorem 8 in the next section.

## 2 Main results and their proofs.

### 2.1 Prescribing $H(\tau)$ and $u(0, t)$ .

The discussion of prescribing  $H(\tau)$  and  $u(0, t)$  is motivated by Theorem 1. Denote the four limit values in (8) as  $p, \alpha, \beta, q$  respectively. In [CT], with the help of the formula (9), we have constructed two bounded radial functions  $\varphi(x) = \varphi(|x|)$ ,  $x \in \mathbb{R}^n$ , so that the following are satisfied:

- (a).  $p = -1 < \alpha = -\sqrt{A^2 + B^2} < \beta = \sqrt{A^2 + B^2} < q = 1$ . Here  $A, B$  are the values of certain convergent improper integrals satisfying  $0 < A^2 + B^2 < 1$ .
- (b).  $p = \alpha < \beta = q$  for arbitrary two finite numbers  $\alpha < \beta$ .

In the above two examples, we have the identity

$$p + q = \alpha + \beta, \quad \text{where } \alpha < \beta. \quad (14)$$

Our new result in the following is that, as long as the symmetry condition (14) is satisfied, we can prescribe them. For four numbers  $p \leq \alpha < \beta \leq q$  to satisfy (14), we must have either  $p < \alpha < \beta < q$  or  $p = \alpha < \beta = q$ . Since the second case has been done, it suffices to look at the first case. We have:

**Theorem 5** *Let  $p < \alpha < \beta < q$  be four **arbitrary** finite numbers satisfying the identity  $p + q = \alpha + \beta$ . One can find a **continuous bounded radial function**  $\varphi(x) = \varphi(|x|)$ ,  $x \in \mathbb{R}^n$ , so that its average integral  $H(\tau)$  and the convolution solution (2) of the problem (1) satisfy*

$$\liminf_{\tau \rightarrow \infty} H(\tau) = p < \liminf_{t \rightarrow \infty} u(0, t) = \alpha < \limsup_{t \rightarrow \infty} u(0, t) = \beta < \limsup_{\tau \rightarrow \infty} H(\tau) = q. \quad (15)$$

More precisely, we can choose the initial data  $\varphi(x) = \varphi(|x|)$ ,  $x \in \mathbb{R}^n$ , as

$$\varphi(\tau) = \frac{q-p}{2} \sin(m \log(\tau+1)) + \frac{q-p}{2} \frac{m\tau}{n(\tau+1)} \cos(m \log(\tau+1)) + \frac{q+p}{2}, \quad (16)$$

where  $\tau \in [0, \infty)$  and  $m \in (0, \infty)$  is a number satisfying the identity

$$\begin{aligned} & \left( \frac{2\omega(n)}{\pi^{n/2}} \int_0^\infty e^{-z^2} z^{n+1} \cos(m \log z) dz \right)^2 + \left( \frac{2\omega(n)}{\pi^{n/2}} \int_0^\infty e^{-z^2} z^{n+1} \sin(m \log z) dz \right)^2 \\ & = \left( \frac{\beta - \alpha}{q - p} \right)^2 \in (0, 1). \end{aligned} \quad (17)$$

**Remark 6** *Without the condition  $p + q = \alpha + \beta$ , we **do not know** how to achieve the result (15) in general. However, we can construct some specific example with  $p + q \neq \alpha + \beta$ . See Section 3.1.*

**Remark 7** *Note that if the initial data  $\varphi(x) = \varphi(|x|)$  is radial, the convolution solution  $u(x, t)$  of the heat equation will also be radial in  $x \in \mathbb{R}^n$  for each fixed time  $t > 0$ .*

### 2.2 Prescribing $\varphi(\tau)$ and $u(0, t)$ .

It is due to Theorem 1 that we study the oscillation behavior of  $H(\tau)$  and its effect on  $u(0, t)$ . Instead, since  $\varphi(\tau)$  is the initial data, it is perhaps more natural to study the oscillation relation between  $\varphi(\tau)$  and  $u(0, t)$ . However, there is a major difference here. Since we always assume that  $\varphi(\tau) \in C^0[0, \infty) \cap L^\infty[0, \infty)$  is continuous and bounded, its average integral function  $H(\tau)$  will also be continuous and bounded and satisfy  $H'(\tau) = O(1/\tau)$  as  $\tau \rightarrow \infty$ , which we call it a **slow**

**oscillation** if we have  $p < q$  (see (22) and Remark 14 below). Any such oscillation will cause  $u(0, t)$  to have a slow oscillation too (by Theorem 1 we have  $p < q$  if and only if  $\alpha < \beta$ ).

On the other hand,  $\varphi(\tau) \in C^0[0, \infty) \cap L^\infty[0, \infty)$  may not have a slow oscillation (i.e. it may not satisfy  $\varphi'(\tau) = O(1/\tau)$  as  $\tau \rightarrow \infty$  even if it is differentiable) and so its oscillation (with  $r < s$ ) may not pass to  $u(0, t)$ . In fact, for any  $2\pi$ -**periodic** regular oscillation function  $\varphi(\tau)$ ,  $\tau \in [0, \infty)$ , the Cauchy problem (1) with  $u(x, 0) = \varphi(|x|)$ ,  $x \in \mathbb{R}^n$ , will have the convergence  $\lim_{t \rightarrow \infty} u(0, t) = \frac{1}{2\pi} \int_0^{2\pi} \varphi(\tau) d\tau$  (see Lemma 17 below). However, due to the similarity between the two representation formulas (9) and (10), if  $\varphi(\tau)$  has a slow oscillation as  $\tau \rightarrow \infty$  with  $r < s$ , then it has the chance of passing to the function  $u(0, t)$  as  $t \rightarrow \infty$  (see Lemma 18 below).

The major result for prescribing  $\varphi(\tau)$  and  $u(0, t)$  is the following:

**Theorem 8** *Let  $r < \alpha < \beta < s$  be four **arbitrary** finite numbers. One can find a **continuous bounded radial function**  $\varphi(x) = \varphi(|x|)$ ,  $x \in \mathbb{R}^n$ , so that it and the convolution solution (2) of the problem (1) satisfy*

$$\liminf_{\tau \rightarrow \infty} \varphi(\tau) = r < \liminf_{t \rightarrow \infty} u(0, t) = \alpha < \limsup_{t \rightarrow \infty} u(0, t) = \beta < \limsup_{\tau \rightarrow \infty} \varphi(\tau) = s. \quad (18)$$

More precisely, we can choose the initial data  $\varphi(x) = \varphi(|x|)$ ,  $x \in \mathbb{R}^n$ , as

$$\varphi(\tau) = [C_1 \sin(m \log(\tau + 1)) + C_2] + \psi(\tau), \quad \tau \in [0, \infty), \quad (19)$$

for some suitable constants  $m > 0$ ,  $C_1 > 0$ ,  $C_2$  depending on  $r$ ,  $\alpha$ ,  $\beta$ ,  $s$  and some  $2\pi$ -periodic continuous nonzero function  $\psi(\tau)$  satisfying  $\frac{1}{2\pi} \int_0^{2\pi} \psi(\tau) d\tau = 0$ .

**Remark 9** *Theorem 8 is actually valid for four arbitrary finite numbers  $r \leq \alpha \leq \beta \leq s$ . We will deal with the remaining cases in Theorem 20 below.*

**Remark 10** *This is a comparison between Theorem 5 and Theorem 8. For any initial data  $\varphi \in C^0(\mathbb{R}^n) \cap L^\infty(\mathbb{R}^n)$ , no matter it is radial or not, the function  $H(\tau)$  is always a slow oscillation function as  $\tau \rightarrow \infty$ . Therefore, the function space for  $H(\tau)$  is smaller than the function space for  $\varphi(\tau)$ , which may explain why we have more general result in Theorem 8. Note that the function  $\varphi(\tau)$  in (19) does not have a slow oscillation as  $\tau \rightarrow \infty$  due to the term  $\psi(\tau)$ .*

### 2.2.1 Proof of Theorem 5.

The proof is an interesting modification of the argument in Section 3.1.1 of [CT], which can achieve much more general result. Some motivation should be explained at the beginning. First, we choose  $\varphi(x) = \varphi(|x|)$  to be a radial function in order to simplify the computation. We think a non-radial function can also be found as long as we can overcome the computational complexity. Second, we choose the average integral function  $H(\tau)$  first and then go back to find its corresponding  $\varphi(|x|)$ . The function  $H(\tau)$  has to be **oscillatory** so that we have  $\liminf_{\tau \rightarrow \infty} H(\tau) < \limsup_{\tau \rightarrow \infty} H(\tau)$ . Moreover, since we require  $\varphi$  to be bounded and continuous (in view of Theorem 1), its average integral  $H(\tau)$  will satisfy the derivative estimate  $H'(\tau) = O(1/\tau)$  as  $\tau \rightarrow \infty$  (see (22) below). Therefore, a natural choice is roughly like the function  $\sin(\log \tau)$ ,  $\tau \in (0, \infty)$ . After suitable modification of the function  $\sin(\log \tau)$ , the proof can be achieved.

Given  $\varphi(x) = \varphi(|x|) \in C^0[0, \infty) \cap L^\infty[0, \infty)$ , its average integral function

$$H(\tau) = \frac{1}{\omega(n) \tau^n} \int_{B(0, \tau)} \varphi(y) dy = \frac{n}{\tau^n} \int_0^\tau \varphi(r) r^{n-1} dr, \quad \tau \in (0, \infty), \quad H(0) = \varphi(0), \quad (20)$$

is a bounded continuous function on  $[0, \infty)$ , differentiable in  $(0, \infty)$ , and satisfies the identity

$$H'(\tau) = -\frac{n}{\tau} H(\tau) + \frac{n}{\tau} \varphi(\tau), \quad \forall \tau \in (0, \infty). \quad (21)$$

Hence it has the asymptotic behavior

$$H'(\tau) = O\left(\frac{1}{\tau}\right) \quad \text{as } \tau \rightarrow \infty. \quad (22)$$

Moreover, the radial function  $\varphi(\tau)$  can be determined from  $H(\tau)$  by the identity (21).

Let  $m \in (0, \infty)$  be a fixed number. Motivated by our principle stated above, we temporarily choose  $H(\tau)$  (will modify it later on) to be equal to

$$H(\tau) = \frac{n}{\tau^n} \int_0^\tau \varphi(r) r^{n-1} dr = \sin(m \log(\tau + 1)), \quad \tau \in (0, \infty), \quad H(0) = 0, \quad (23)$$

which, by (21), gives

$$\varphi(\tau) = \sin(m \log(\tau + 1)) + \frac{m\tau}{n(\tau + 1)} \cos(m \log(\tau + 1)), \quad \tau \in (0, \infty). \quad (24)$$

That is, if we choose  $\varphi(\tau)$  to be the function given by (24) and define  $\varphi(0) = 0$ , then the function  $\varphi(x) = \varphi(|x|)$ ,  $x \in \mathbb{R}^n$ , will be a radial function defined on  $\mathbb{R}^n$ , lying in the space  $C^0(\mathbb{R}^n) \cap L^\infty(\mathbb{R}^n)$ , and its average integral is given by  $H(\tau) = \sin(m \log(\tau + 1))$  for all  $\tau \in [0, \infty)$ . In particular, we have  $p = -1$ ,  $q = 1$ .

To find the values of  $\alpha$ ,  $\beta$ , we use the representation formula (9). We have

$$\begin{aligned} u(0, t) &= \frac{2\omega(n)}{\pi^{n/2}} \int_0^\infty e^{-z^2} z^{n+1} \sin\left(m \log\left(\sqrt{4t}z + 1\right)\right) dz \\ &= \begin{cases} \left[ \frac{2\omega(n)}{\pi^{n/2}} \int_0^\infty e^{-z^2} z^{n+1} \cos\left(m \log\left(z + \frac{1}{\sqrt{4t}}\right)\right) dz \right] \sin(m \log \sqrt{4t}) \\ + \left[ \frac{2\omega(n)}{\pi^{n/2}} \int_0^\infty e^{-z^2} z^{n+1} \sin\left(m \log\left(z + \frac{1}{\sqrt{4t}}\right)\right) dz \right] \cos(m \log \sqrt{4t}), \end{cases} \quad t \in (0, \infty) \end{aligned} \quad (25)$$

and the Lebesgue Dominated Convergence Theorem implies

$$\begin{aligned} &\lim_{t \rightarrow \infty} \frac{2\omega(n)}{\pi^{n/2}} \int_0^\infty e^{-z^2} z^{n+1} \cos\left(m \log\left(z + \frac{1}{\sqrt{4t}}\right)\right) dz \\ &= \frac{2\omega(n)}{\pi^{n/2}} \int_0^\infty e^{-z^2} z^{n+1} \cos(m \log z) dz := A(m) \in (-1, 1) \end{aligned} \quad (26)$$

and

$$\begin{aligned} &\lim_{t \rightarrow \infty} \frac{2\omega(n)}{\pi^{n/2}} \int_0^\infty e^{-z^2} z^{n+1} \sin\left(m \log\left(z + \frac{1}{\sqrt{4t}}\right)\right) dz \\ &= \frac{2\omega(n)}{\pi^{n/2}} \int_0^\infty e^{-z^2} z^{n+1} \sin(m \log z) dz := B(m) \in (-1, 1), \end{aligned} \quad (27)$$

where we can use the identity  $\frac{2\omega(n)}{\pi^{n/2}} \int_0^\infty e^{-z^2} z^{n+1} dz = 1$  to know that  $A(m), B(m) \in (-1, 1)$ . By Hölder inequality, the constant  $A(m)$  satisfies the estimate

$$\begin{aligned} A^2(m) &= \left( \frac{2\omega(n)}{\pi^{n/2}} \int_0^\infty \sqrt{e^{-z^2} z^{n+1}} \sqrt{e^{-z^2} z^{n+1}} \cos(m \log z) dz \right)^2 \\ &< \left( \frac{2\omega(n)}{\pi^{n/2}} \int_0^\infty e^{-z^2} z^{n+1} dz \right) \left( \frac{2\omega(n)}{\pi^{n/2}} \int_0^\infty e^{-z^2} z^{n+1} \cos^2(m \log z) dz \right) \\ &= \frac{2\omega(n)}{\pi^{n/2}} \int_0^\infty e^{-z^2} z^{n+1} \cos^2(m \log z) dz \end{aligned}$$

and similarly

$$B^2(m) < \frac{2\omega(n)}{\pi^{n/2}} \int_0^\infty e^{-z^2} z^{n+1} \sin^2(m \log z) dz$$

and we conclude

$$0 < A^2(m) + B^2(m) < \frac{2\omega(n)}{\pi^{n/2}} \int_0^\infty e^{-z^2} z^{n+1} [\cos^2(m \log z) + \sin^2(m \log z)] dz = 1. \quad (28)$$

Therefore, we have the convergence

$$\lim_{t \rightarrow \infty} \left| u(0, t) - \left[ A(m) \sin\left(m \log \sqrt{4t}\right) + B(m) \cos\left(m \log \sqrt{4t}\right) \right] \right| = 0, \quad (29)$$

which gives the conclusion

$$\begin{aligned} -1 &= \liminf_{\tau \rightarrow \infty} H(\tau) < -\sqrt{A^2(m) + B^2(m)} = \liminf_{t \rightarrow \infty} u(0, t) \\ &< \limsup_{t \rightarrow \infty} u(0, t) = \sqrt{A^2(m) + B^2(m)} < \limsup_{\tau \rightarrow \infty} H(\tau) = 1 \end{aligned} \quad (30)$$

for fixed  $m \in (0, \infty)$ .

The idea next is to view  $m \in (0, \infty)$  as a variable. We observe the following:

**Lemma 11** *We have*

$$\lim_{m \rightarrow \infty} A(m) = \lim_{m \rightarrow \infty} B(m) = 0, \quad m \in (0, \infty). \quad (31)$$

**Proof.** For any  $\varepsilon > 0$ , one can find small number  $\delta > 0$  and large number  $M > 0$ , both are independent of  $m \in (0, \infty)$ , such that

$$\left| \int_0^\delta e^{-z^2} z^{n+1} \cos(m \log z) dz \right| + \left| \int_M^\infty e^{-z^2} z^{n+1} \cos(m \log z) dz \right| < \varepsilon. \quad (32)$$

On the other hand, by the change of variables, we have

$$\int_\delta^M e^{-z^2} z^{n+1} \cos(m \log z) dz = \int_{\log \delta}^{\log M} F(x) \cos(mx) dx, \quad x = \log z,$$

where  $F(x) = e^{-e^{2x} + (n+2)x}$ . The familiar **Riemann-Lebesgue lemma** in analysis (or just use integration by parts) implies

$$\lim_{m \rightarrow \infty} \int_{\log \delta}^{\log M} F(x) \cos(mx) dx = 0. \quad (33)$$

which, together with (32), implies  $\lim_{m \rightarrow \infty} A(m) = 0$ . The proof of  $\lim_{m \rightarrow \infty} B(m) = 0$  is similar.  $\square$

Since estimate (32) is valid for all  $m \in (0, \infty)$ , the two improper integrals

$$\int_0^\infty e^{-z^2} z^{n+1} \cos(m \log z) dz, \quad \int_0^\infty e^{-z^2} z^{n+1} \sin(m \log z) dz \quad (34)$$

**converge uniformly** with respect to  $m \in (0, \infty)$ . As a consequence, both  $A(m)$  and  $B(m)$  are **continuous functions** of  $m \in (0, \infty)$ , with

$$\lim_{m \rightarrow \infty} A(m) = \lim_{m \rightarrow \infty} B(m) = 0, \quad \lim_{m \rightarrow 0} A(m) = 1, \quad \lim_{m \rightarrow 0} B(m) = 0, \quad (35)$$

where the last two limits in (35) are due to the Lebesgue Dominated Convergence Theorem. Moreover, the intermediate value theorem implies the existence of a number  $m \in (0, \infty)$  satisfying

$$\sqrt{A^2(m) + B^2(m)} = \frac{\beta - \alpha}{q - p} \in (0, 1). \quad (36)$$

Now we choose  $H(\tau)$  as

$$H(\tau) = \frac{q-p}{2} \sin(m \log(\tau+1)) + \frac{q+p}{2}, \quad H(0) = \frac{q+p}{2}, \quad \tau \in [0, \infty), \quad (37)$$

where  $m \in (0, \infty)$  is the number satisfying (36). By (23) and (24), its corresponding initial bounded radial function  $\varphi(x) = \varphi(|x|)$ ,  $x \in \mathbb{R}^n$ , is given by

$$\varphi(\tau) = \begin{cases} \frac{q-p}{2} \sin(m \log(\tau+1)) \\ + \frac{q-p}{2} \frac{m\tau}{n(\tau+1)} \cos(m \log(\tau+1)) + \frac{q+p}{2}, \end{cases} \quad \varphi(0) = \frac{q+p}{2}, \quad \tau \in [0, \infty). \quad (38)$$

Similar to (29), we have

$$\lim_{t \rightarrow \infty} \left| u(0, t) - \left\{ \frac{q-p}{2} \left[ A(m) \sin(m \log \sqrt{4t}) + B(m) \cos(m \log \sqrt{4t}) \right] + \frac{q+p}{2} \right\} \right| = 0, \quad (39)$$

which gives

$$\liminf_{t \rightarrow \infty} u(0, t) = -\frac{q-p}{2} \sqrt{A^2(m) + B^2(m)} + \frac{q+p}{2} = -\frac{\beta - \alpha}{2} + \frac{\beta + \alpha}{2} = \alpha$$

and

$$\limsup_{t \rightarrow \infty} u(0, t) = \frac{q-p}{2} \sqrt{A^2(m) + B^2(m)} + \frac{q+p}{2} = \frac{\beta - \alpha}{2} + \frac{\beta + \alpha}{2} = \beta.$$

Since we clearly have  $\liminf_{\tau \rightarrow \infty} H(\tau) = p$ ,  $\limsup_{\tau \rightarrow \infty} H(\tau) = q$ , the proof of Theorem 5 is now complete.  $\square$

**Remark 12** *The proof of Theorem 5 also reveals the following interesting observation. For fixed  $q - p > 0$ , if we have small  $\beta - \alpha$ , then by (36) and (31), we will have large  $m \in (0, \infty)$  and the function  $\varphi(\tau)$  in (38) will tend to be unbounded due to the term  $m\tau / (n(\tau+1))$ . This may suggest that we cannot find a bounded radial function  $\varphi(\tau)$  satisfying  $p < \alpha = \beta < q$ . This matches with the result in Theorem 1, which implies that if we have  $p < \alpha = \beta < q$ , then the initial data  $\varphi \in C^0(\mathbb{R}^n)$  must be unbounded.*

**Remark 13** *By (7), for a given initial data  $\varphi \in C^0(\mathbb{R}^n) \cap L^\infty(\mathbb{R}^n)$ , if we have*

$$\liminf_{t \rightarrow \infty} u(0, t) = \alpha < \limsup_{t \rightarrow \infty} u(0, t) = \beta,$$

*then for any fixed  $x \in \mathbb{R}^n$  we also have*

$$\liminf_{t \rightarrow \infty} u(x, t) = \alpha < \limsup_{t \rightarrow \infty} u(x, t) = \beta. \quad (40)$$

**Remark 14** *It has been shown in [CT] (see its equations (16) and (37)) that both functions  $H(\tau)$  and  $u(0, t)$  satisfy*

$$H'(\tau) = O\left(\frac{1}{\tau}\right) \quad \text{as } \tau \rightarrow \infty, \quad u_t(0, t) = O\left(\frac{1}{t}\right) \quad \text{as } t \rightarrow \infty, \quad (41)$$

*as long as the initial data  $\varphi$  lies in the space  $C^0(\mathbb{R}^n) \cap L^\infty(\mathbb{R}^n)$  (no matter it is radial or not). By (41), we may say that any oscillation in  $H(\tau)$  (with  $p < q$ ) or in  $u(0, t)$  (with  $\alpha < \beta$ ) is a **slow oscillation**.*



As a consequence of Theorem 5, we have the following two corollaries, which say that we can **prescribe** the oscillation of  $H(\tau)$  and  $u(0, t)$  for **three** arbitrary different numbers.

**Corollary 15** *For any three different numbers, denoted as  $p < \alpha < \beta$ , one can find a continuous bounded radial function  $\varphi(x) = \varphi(|x|)$ ,  $x \in \mathbb{R}^n$ , so that its average integral and the solution (2) of the problem (1) satisfies*

$$\liminf_{\tau \rightarrow \infty} H(\tau) = p < \liminf_{t \rightarrow \infty} u(0, t) = \alpha < \limsup_{t \rightarrow \infty} u(0, t) = \beta < \limsup_{\tau \rightarrow \infty} H(\tau) = q, \quad (42)$$

where  $q = \alpha + \beta - p$ . Similarly, for any three different numbers, denoted as  $\alpha < \beta < q$ , one can find a bounded radial function satisfying (42), where now  $p = \alpha + \beta - q$ .

**Corollary 16** *For any three different numbers, denoted as  $p < \alpha < q$ , with  $\alpha < (p + q)/2$ , the same result as in Corollary 15 holds, where now  $\beta = p + q - \alpha$ . Similarly, for any three different numbers, denoted as  $p < \beta < q$ , with  $\beta > (p + q)/2$ , the same result as in Corollary 15 holds, where now  $\alpha = p + q - \beta$ .*

## 2.2.2 Proof of Theorem 8.

To prove Theorem 8, we first need the following:

**Lemma 17** *Assume  $\varphi(\tau)$  is a  $2\pi$ -**periodic** radial function defined on  $\tau \in [0, \infty)$ . We have the convergence*

$$\lim_{t \rightarrow \infty} u(0, t) = \lim_{\tau \rightarrow \infty} H(\tau) = \lim_{\tau \rightarrow \infty} \frac{n}{\tau^n} \int_0^\tau \varphi(r) r^{n-1} dr = \frac{1}{2\pi} \int_0^{2\pi} \varphi(\tau) d\tau. \quad (43)$$

**Proof.** Since  $\varphi \in C^0[0, \infty) \cap L^\infty[0, \infty)$ , by Remark 2 in Theorem 1, it suffices to prove the identity for  $H(\tau)$ . The proof is quite straightforward for the case  $n = 1$  (see Remark 6 in [TN]), but may need a trick for  $n > 1$ . For large  $\tau > 0$ , we can express it as  $\tau = 2m\pi + R$  for some  $m \in \mathbb{N}$  and  $R \in [0, 2\pi)$ , with the understanding that both  $m$  and  $R$  depend on  $\tau$  and as  $\tau \rightarrow \infty$  we have  $m \rightarrow \infty$ , hence we obtain

$$\begin{aligned} & \frac{n}{\tau^n} \int_0^\tau \varphi(r) r^{n-1} dr \\ &= \frac{n}{(2m\pi + R)^n} \left( \int_0^{2\pi} \varphi(r) r^{n-1} dr + \int_{2\pi}^{4\pi} \varphi(r) r^{n-1} dr + \int_{4\pi}^{6\pi} \varphi(r) r^{n-1} dr \right. \\ & \quad \left. + \cdots + \int_{2(m-1)\pi}^{2m\pi} \varphi(r) r^{n-1} dr + \int_{2m\pi}^{2m\pi+R} \varphi(r) r^{n-1} dr \right). \end{aligned} \quad (44)$$

If we do the change of variables

$$\int_{2\pi}^{4\pi} \varphi(r) r^{n-1} dr = \int_0^{2\pi} \varphi(s) (s + 2\pi)^{n-1} ds, \quad r = s + 2\pi,$$

and

$$\int_{4\pi}^{6\pi} \varphi(r) r^{n-1} dr = \int_0^{2\pi} \varphi(s) (s + 4\pi)^{n-1} ds, \quad r = s + 4\pi,$$

..., etc., (44) becomes

$$\begin{aligned} & \frac{n}{\tau^n} \int_0^\tau \varphi(r) r^{n-1} dr \\ &= \frac{n}{(2m\pi + R)^n} \left( \int_0^{2\pi} \varphi(s) [s^{n-1} + (s + 2\pi)^{n-1} + (s + 4\pi)^{n-1} + \cdots + (s + 2(m-1)\pi)^{n-1}] ds \right. \\ & \quad \left. + \int_0^R \varphi(s) (s + 2m\pi)^{n-1} ds \right) \\ & := I + II. \end{aligned} \quad (45)$$

For the second term  $II$  in (45), we have

$$\lim_{m \rightarrow \infty} |II| \leq \lim_{m \rightarrow \infty} \left( \frac{n}{(2m\pi + R)^n} \cdot 2\pi \max_{s \in [0, 2\pi]} |\varphi(s)| \cdot (2\pi + 2m\pi)^{n-1} \right) = 0. \quad (46)$$

To estimate the first term  $I$  in (45), we first look at the integral

$$\begin{aligned} & \frac{n}{(2m\pi)^n} \int_0^{2\pi} \varphi(s) [s^{n-1} + (s + 2\pi)^{n-1} + (s + 4\pi)^{n-1} + \cdots + (s + 2(m-1)\pi)^{n-1}] ds \\ &= \frac{1}{2\pi} \int_0^{2\pi} \varphi(s) \underbrace{\frac{n}{m^n} \frac{1}{(2\pi)^{n-1}} [s^{n-1} + (s + 2\pi)^{n-1} + (s + 4\pi)^{n-1} + \cdots + (s + 2(m-1)\pi)^{n-1}]}_{\dots} ds, \end{aligned}$$

where the underlined term  $\underbrace{\dots}$  is equal to

$$\begin{aligned} & \underbrace{\dots} \\ &= \frac{n}{m^n} \left[ \left( \frac{s}{2\pi} \right)^{n-1} + \left( \frac{s}{2\pi} + 1 \right)^{n-1} + \left( \frac{s}{2\pi} + 2 \right)^{n-1} + \cdots + \left( \frac{s}{2\pi} + (m-1) \right)^{n-1} \right] \end{aligned}$$

and for  $s \in [0, 2\pi]$  we have

$$\begin{aligned} & \frac{n}{m^n} (0^{n-1} + 1^{n-1} + 2^{n-1} + \cdots + (m-1)^{n-1}) \\ & \leq \underbrace{\dots} \leq \frac{n}{m^n} (1^{n-1} + 2^{n-1} + 3^{n-1} + \cdots + m^{n-1}). \end{aligned} \quad (47)$$

By the inequality

$$\begin{aligned} & \frac{n}{m^n} (0^{n-1} + 1^{n-1} + 2^{n-1} + \cdots + (m-1)^{n-1}) \\ & < \frac{n}{m^n} \int_0^m x^{n-1} dx = 1 < \frac{n}{m^n} (1^{n-1} + 2^{n-1} + 3^{n-1} + \cdots + m^{n-1}) \end{aligned} \quad (48)$$

with

$$\begin{aligned} & \lim_{m \rightarrow \infty} \left[ \frac{n}{m^n} (1^{n-1} + 2^{n-1} + 3^{n-1} + \cdots + m^{n-1}) - \frac{n}{m^n} (0^{n-1} + 1^{n-1} + 2^{n-1} + \cdots + (m-1)^{n-1}) \right] \\ &= \lim_{m \rightarrow \infty} \left( \frac{n}{m^n} m^{n-1} \right) = 0, \end{aligned}$$

we must have

$$\begin{cases} \lim_{m \rightarrow \infty} \frac{n}{m^n} (0^{n-1} + 1^{n-1} + 2^{n-1} + \cdots + (m-1)^{n-1}) = 1 \\ \lim_{m \rightarrow \infty} \frac{n}{m^n} (1^{n-1} + 2^{n-1} + 3^{n-1} + \cdots + m^{n-1}) = 1 \end{cases}$$

and hence  $\lim_{m \rightarrow \infty} \underbrace{\dots} = 1$ , which implies

$$\lim_{m \rightarrow \infty} \frac{n}{(2m\pi)^n} \int_0^{2m\pi} \varphi(r) r^{n-1} dr = \frac{1}{2\pi} \int_0^{2\pi} \varphi(s) \left( \lim_{m \rightarrow \infty} \underbrace{\dots} \right) ds = \frac{1}{2\pi} \int_0^{2\pi} \varphi(s) ds. \quad (49)$$

Finally we have

$$\lim_{m \rightarrow \infty} I = \lim_{m \rightarrow \infty} \left( \frac{(2m\pi)^n}{(2m\pi + R)^n} \frac{n}{(2m\pi)^n} \int_0^{2m\pi} \varphi(r) r^{n-1} dr \right) = \frac{1}{2\pi} \int_0^{2\pi} \varphi(s) ds.$$

The proof is done.  $\square$

As we have said in the paragraph before Theorem 8, if  $\varphi(\tau)$  has a slow oscillation as  $\tau \rightarrow \infty$  with  $r < s$ , then it has the chance of passing to the function  $u(0, t)$ . The following gives a simple way to find a slow oscillation function  $\varphi(\tau)$  on  $[0, \infty)$  which will cause  $\alpha < \beta$ .

**Lemma 18** Let  $g(\tau)$  be a  $2\pi$ -periodic non-constant  $C^2$  function defined on  $[0, \infty)$  and let  $G(\tau) = g(\log(\tau + 1))$ ,  $\tau \in [0, \infty)$ . Then the radial function  $\varphi(\tau) = \frac{\tau}{n}G'(\tau) + G(\tau)$ ,  $\tau \in [0, \infty)$ , has a slow oscillation as  $\tau \rightarrow \infty$ , lies in the space  $C^0[0, \infty) \cap L^\infty[0, \infty)$ , and its average integral  $H(\tau)$ ,  $\tau \in [0, \infty)$ , satisfies  $H(\tau) = g(\log(\tau + 1))$ ,  $\tau \in [0, \infty)$ , with

$$p = \liminf_{\tau \rightarrow \infty} g(\tau) < \limsup_{\tau \rightarrow \infty} g(\tau) = q, \quad (50)$$

which, by Theorem 1, will also imply  $\alpha < \beta$ .

**Proof.** By

$$\begin{cases} \varphi(\tau) = \frac{\tau}{n} \frac{g'(\log(\tau+1))}{\tau+1} + g(\log(\tau+1)), & \varphi(0) = g(0), \\ \varphi'(\tau) = \left(1 + \frac{1}{n}\right) \frac{g'(\log(\tau+1))}{\tau+1} + \frac{\tau[g''(\log(\tau+1)) - g'(\log(\tau+1))]}{n(\tau+1)^2}, & \tau \in [0, \infty), \end{cases}$$

we see that  $\varphi$  is a slow oscillation function on  $[0, \infty)$ , lying in the space  $C^0[0, \infty) \cap L^\infty[0, \infty)$ . Now

$$\begin{aligned} H(\tau) &= \frac{n}{\tau^n} \int_0^\tau \varphi(r) r^{n-1} dr = \frac{n}{\tau^n} \int_0^\tau \left( \frac{r}{n} G'(r) + G(r) \right) r^{n-1} dr \\ &= \frac{1}{\tau^n} \int_0^\tau \frac{d}{dr} (r^n G(r)) dr = G(\tau) = g(\log(\tau + 1)), \quad \tau \in (0, \infty) \end{aligned}$$

and  $H(0) = G(0) = g(0)$ . Therefore, the inequality (50) follows.  $\square$

With the help of Lemma 17, we are ready to prove Theorem 8. For any  $r < \alpha < \beta < s$ , we divide the proof into two cases.

**Case 1:**  $r + s = \alpha + \beta$ ,  $r < \alpha < \beta < s$ .

In this case we can use the representation formula (10) and, similar to (37), choose

$$\varphi(\tau) = \frac{s-r}{2} \sin(m \log(\tau + 1)) + \frac{s+r}{2}, \quad \varphi(0) = \frac{s+r}{2}, \quad \tau \in [0, \infty), \quad (51)$$

and perform the same argument as in the proof of Theorem 5 to prescribe them. More precisely, here we need to choose  $m \in (0, \infty)$  to satisfy  $\sqrt{\tilde{A}^2(m) + \tilde{B}^2(m)} = \frac{\beta-\alpha}{s-r} \in (0, 1)$ , where now

$$\begin{cases} \tilde{A}(m) = \frac{n\omega(n)}{\pi^{n/2}} \int_0^\infty e^{-z^2} z^{n-1} \cos(m \log z) dz \in (-1, 1), \\ \tilde{B}(m) = \frac{n\omega(n)}{\pi^{n/2}} \int_0^\infty e^{-z^2} z^{n-1} \sin(m \log z) dz \in (-1, 1), \end{cases} \quad (52)$$

with

$$0 < \tilde{A}^2(m) + \tilde{B}^2(m) < \frac{n\omega(n)}{\pi^{n/2}} \int_0^\infty e^{-z^2} z^{n-1} [\cos^2(m \log z) + \sin^2(m \log z)] dz = 1. \quad (53)$$

Note that in this case the chosen function  $\varphi(\tau)$ , like  $H(\tau)$  in (37), has a **slow oscillation** on  $[0, \infty)$ .

**Case 2:**  $r + s \neq \alpha + \beta$ ,  $r < \alpha < \beta < s$ .

In this case, we may assume  $r + s > \alpha + \beta$  (the treatment for the case  $r + s < \alpha + \beta$  is similar). We will choose  $\varphi(\tau) \in C^0[0, \infty) \cap L^\infty[0, \infty)$  to be the **sum** of two functions  $\varphi_1(\tau) + \varphi_2(\tau)$ , where  $\varphi_1(\tau)$  has a **slow oscillation**, but  $\varphi_2(\tau)$  has a **regular oscillation** (which is  $2\pi$ -periodic).

Let  $\lambda$  be the number satisfying  $r + \lambda = \alpha + \beta$ . Since  $r < \alpha < \beta < s$  and  $r + s > \alpha + \beta$ , we have  $\beta < \lambda < s$ . We can write  $\lambda$  as  $\lambda = \varepsilon + \delta$  for some small  $\varepsilon > 0$  and  $\delta \in \mathbb{R}$  so that we have

$$r + \varepsilon < \alpha < \beta < \delta < s. \quad (54)$$

Now we have  $(r + \varepsilon) + \delta = \alpha + \beta$  and by Case 1, we can find a radial  $\varphi_1 \in C^0[0, \infty) \cap L^\infty[0, \infty)$  such that it and its convolution solution  $u_1(x, t)$  satisfy

$$\liminf_{\tau \rightarrow \infty} \varphi_1(\tau) = r + \varepsilon < \liminf_{t \rightarrow \infty} u_1(0, t) = \alpha < \limsup_{t \rightarrow \infty} u_1(0, t) = \beta < \limsup_{\tau \rightarrow \infty} \varphi_1(\tau) = \delta. \quad (55)$$

Next let  $\varphi_2(\tau)$  be a  $2\pi$ -periodic continuous function on  $\tau \in [0, \infty)$  which satisfies

$$\frac{1}{2\pi} \int_0^{2\pi} \varphi_2(\tau) d\tau = 0, \quad \min_{\tau \in [0, 2\pi]} \varphi_2(\tau) = -\varepsilon < 0, \quad \max_{\tau \in [0, 2\pi]} \varphi_2(\tau) = s - \delta > 0. \quad (56)$$

Such a function  $\varphi_2$  clearly exists and, together with its convolution solution  $u_2(x, t)$ , they satisfy

$$\liminf_{\tau \rightarrow \infty} \varphi_2(\tau) = -\varepsilon < \lim_{t \rightarrow \infty} u_2(0, t) = \frac{1}{2\pi} \int_0^{2\pi} \varphi_2(\tau) d\tau = 0 < \limsup_{\tau \rightarrow \infty} \varphi_2(\tau) = s - \delta, \quad (57)$$

due to the result in Lemma 17. Finally, we set  $\varphi(\tau) = \varphi_1(\tau) + \varphi_2(\tau)$ ,  $\tau \in [0, \infty)$ . We have  $\varphi \in C^0[0, \infty) \cap L^\infty[0, \infty)$  and its convolution solution  $u(x, t)$  satisfy  $u(x, t) = u_1(x, t) + u_2(x, t)$  for all  $(x, t) \in \mathbb{R}^n \times [0, \infty)$ . Since  $u_2(0, t) \rightarrow 0$  as  $t \rightarrow \infty$ , we have

$$\begin{cases} \liminf_{t \rightarrow \infty} u(0, t) = \liminf_{t \rightarrow \infty} u_1(0, t) = \alpha \\ \limsup_{t \rightarrow \infty} u(0, t) = \limsup_{t \rightarrow \infty} u_1(0, t) = \beta. \end{cases} \quad (58)$$

It remains to look at  $\varphi(\tau)$  for  $\tau \rightarrow \infty$ . By (51) we know  $\varphi_1(\tau)$  is a **slow oscillation** function on  $[0, \infty)$  with the form

$$\varphi_1(\tau) = \frac{\delta - (r + \varepsilon)}{2} \sin(m \log(\tau + 1)) + \frac{\delta + (r + \varepsilon)}{2}, \quad \varphi_1(0) = \frac{\delta + (r + \varepsilon)}{2}, \quad \tau \in [0, \infty) \quad (59)$$

for some fixed  $m \in (0, \infty)$  satisfying  $\sqrt{\tilde{A}^2(m) + \tilde{B}^2(m)} = \frac{\beta - \alpha}{\delta - (r + \varepsilon)} \in (0, 1)$ . On the other hand,  $\varphi_2(\tau)$  is a **regular oscillation** function ( **$2\pi$ -periodic function**) on  $[0, \infty)$ , which oscillates between  $-\varepsilon$  and  $s - \delta$ . It is not difficult to see that there exists a sequence  $\tau_j \rightarrow \infty$  so that

$$\sin(m \log(\tau_j + 1)) \rightarrow -1 \quad \text{and} \quad \varphi_2(\tau_j) \rightarrow -\varepsilon = \min_{\tau \in [0, 2\pi]} \varphi_2(\tau) < 0 \quad (60)$$

as  $j \rightarrow \infty$ . Similarly, there exists a sequence  $\tau_k \rightarrow \infty$  so that

$$\sin(m \log(\tau_k + 1)) \rightarrow 1 \quad \text{and} \quad \varphi_2(\tau_k) \rightarrow s - \delta = \max_{\tau \in [0, 2\pi]} \varphi_2(\tau) > 0 \quad (61)$$

as  $k \rightarrow \infty$ . By (59), (61), and (60), we have

$$\begin{aligned} \liminf_{\tau \rightarrow \infty} \varphi(\tau) &= \liminf_{\tau \rightarrow \infty} (\varphi_1(\tau) + \varphi_2(\tau)) \\ &= \liminf_{\tau \rightarrow \infty} \varphi_1(\tau) + \liminf_{\tau \rightarrow \infty} \varphi_2(\tau) = (r + \varepsilon) - \varepsilon = r \end{aligned} \quad (62)$$

and

$$\begin{aligned} \limsup_{\tau \rightarrow \infty} \varphi(\tau) &= \limsup_{\tau \rightarrow \infty} (\varphi_1(\tau) + \varphi_2(\tau)) \\ &= \limsup_{\tau \rightarrow \infty} \varphi_1(\tau) + \limsup_{\tau \rightarrow \infty} \varphi_2(\tau) = \delta + (s - \delta) = s. \end{aligned} \quad (63)$$

The proof is done due to (58), (62), and (63).

Combining Case 1 and Case 2, the proof of Theorem 8 is now complete.  $\square$

**Remark 19** This is to explain the existence of a sequence  $\tau_k \rightarrow \infty$  so that (61) holds. We first note that, for given  $\varepsilon > 0$ , there exists  $\delta > 0$  (depending only on  $\varepsilon$ ), such that if  $m \log(\tau + 1)$  (here  $m > 0$  is fixed) lies in the interval  $(2k\pi + \frac{\pi}{2} - \delta, 2k\pi + \frac{\pi}{2} + \delta)$  for **some**  $k \in \mathbb{N}$ , then we have  $|\sin(m \log(\tau + 1)) - 1| < \varepsilon$ , which is equivalent for  $\tau + 1$  to lie in the interval

$$\tau + 1 \in \left( \exp\left(\frac{2k\pi + \frac{\pi}{2} - \delta}{m}\right), \exp\left(\frac{2k\pi + \frac{\pi}{2} + \delta}{m}\right) \right). \quad (64)$$

We can choose  $k \in \mathbb{N}$  sufficiently large so that the length of the interval in (64) is sufficiently large. By this observation, we clearly have (61). The reason for (60) to hold is similar.

### 2.3 The remaining cases not covered by Theorem 8.

For prescribing general  $r \leq \alpha \leq \beta \leq s$ , the remaining cases not covered by Theorem 8 are discussed in the following:

**Theorem 20** For any four arbitrary finite numbers  $r, \alpha, \beta, s$  in one of the following cases:

$$\begin{cases} (1). r = \alpha < \beta < s, & (2). r < \alpha = \beta < s, & (3). r < \alpha < \beta = s, \\ (4). r = \alpha = \beta < s, & (5). r = \alpha < \beta = s, & (6). r < \alpha = \beta = s, \end{cases} \quad (65)$$

one can choose a suitable radial initial data  $\varphi(x) = \varphi(|x|) \in C^0[0, \infty) \cap L^\infty[0, \infty)$  satisfying

$$\liminf_{\tau \rightarrow \infty} \varphi(\tau) = r, \quad \limsup_{\tau \rightarrow \infty} \varphi(\tau) = s \quad (66)$$

and its corresponding convolution solution  $u(x, t)$  in (2) satisfies

$$\liminf_{t \rightarrow \infty} u(0, t) = \alpha, \quad \limsup_{t \rightarrow \infty} u(0, t) = \beta. \quad (67)$$

**Remark 21** The case  $r = \alpha = \beta = s$  is trivial. Just choose  $\varphi(x)$  to be a constant function.

**Proof.** (2). For  $r < \alpha = \beta < s$ , we can choose  $\varphi(\tau)$ ,  $\tau \in [0, \infty)$ , to be any  $2\pi$ -periodic function with average value  $\alpha = \beta$ , maximum value  $s$ , minimum value  $r$ . By Lemma 17, it can be achieved.

(5). For  $r = \alpha < \beta = s$ , we choose  $\varphi(\tau)$  to be the **extremely slow oscillation function**

$$\varphi(\tau) = \frac{\beta - \alpha}{2} \sin[\log(\log(\tau + 2))] + \frac{\beta + \alpha}{2}, \quad \tau \in [0, \infty), \quad (68)$$

which has

$$\liminf_{\tau \rightarrow \infty} \varphi(\tau) = \alpha, \quad \limsup_{\tau \rightarrow \infty} \varphi(\tau) = \beta. \quad (69)$$

By (10), its corresponding  $u(0, t)$  is given by

$$u(0, t) = \frac{\beta - \alpha}{2} \frac{n\omega(n)}{\pi^{n/2}} \int_0^\infty e^{-z^2} z^{n-1} \sin\left[\log\left(\log\left(\sqrt{4t}z + 2\right)\right)\right] dz + \frac{\beta + \alpha}{2}, \quad (70)$$

where one can write

$$\sin\left[\log\left(\log\left(\sqrt{4t}z + 2\right)\right)\right] = \sin\left[\log\left(\log\sqrt{4t}\right) + g(t, z)\right],$$

with

$$g(t, z) = \log\left(1 + \frac{1}{\log\sqrt{4t}} \log\left(z + \frac{2}{\sqrt{4t}}\right)\right), \quad t, z \in (0, \infty).$$

We see that  $\lim_{t \rightarrow \infty} g(t, z) = 0$  for fixed  $z \in (0, \infty)$ . Hence the Lebesgue Dominated Convergence Theorem implies

$$\lim_{t \rightarrow \infty} \left| u(0, t) - \left[ \frac{\beta - \alpha}{2} \sin \left( \log \left( \log \sqrt{4t} \right) \right) + \frac{\beta + \alpha}{2} \right] \right| = 0 \quad (71)$$

and so

$$\liminf_{t \rightarrow \infty} u(0, t) = \alpha, \quad \limsup_{t \rightarrow \infty} u(0, t) = \beta. \quad (72)$$

(4). For  $r = \alpha = \beta < s$ , we choose  $\varphi \in C^0[0, \infty) \cap L^\infty[0, \infty)$  with  $\liminf_{\tau \rightarrow \infty} \varphi(\tau) = r = \alpha$ ,  $\limsup_{\tau \rightarrow \infty} \varphi(\tau) = s$ , and for most  $\tau \in [0, \infty)$  the value of  $\varphi(\tau)$  is equal to  $\alpha$  and for  $\varphi(\tau)$  not equal to  $\alpha$ , it looks like a thin bump with height  $s - \alpha$  and the supports of these bumps are spreading further and further apart. By this choice of  $\varphi$  we have

$$\lim_{\tau \rightarrow \infty} H(\tau) = \lim_{\tau \rightarrow \infty} \frac{n}{\tau^n} \int_0^\tau \varphi(r) r^{n-1} dr = \alpha,$$

which implies  $\lim_{t \rightarrow \infty} u(0, t) = \alpha$  due to Theorem 1. Hence, the case  $r = \alpha = \beta < s$  is achieved.

(6). For  $r < \alpha = \beta = s$ , the construction of  $\varphi$  is similar to that in (4) except that we reverse the role of  $r$  and  $s$ .

(1). For  $r = \alpha < \beta < s$ , we first choose  $\varphi_1(\tau)$  to be the function given by (68), which, together with its corresponding  $u_1(0, t)$ , will satisfy (69) and (72). Next, we choose  $\varphi_2(\tau)$ . Let  $\varepsilon > 0$  be a fixed small number and for each  $m \in \mathbb{N}$ , let

$$\tau_m = \exp \left( \exp \left( 2m\pi + \frac{\pi}{2} \right) \right) - 2, \quad \tilde{\tau}_m = \exp \left( \exp \left( 2m\pi + \frac{3\pi}{2} \right) \right) - 2,$$

where we note that  $\varphi_1(\tilde{\tau}_m) = \alpha$ ,  $\varphi_1(\tau_m) = \beta$ . We require  $\varphi_2(\tau)$  to be a nonnegative function satisfying

$$\varphi_2(\tau) = \begin{cases} \frac{s-\beta}{\varepsilon} (\tau - (\tau_m - \varepsilon)), & \tau \in [\tau_m - \varepsilon, \tau_m] \\ -\frac{s-\beta}{\varepsilon} (\tau - (\tau_m + \varepsilon)), & \tau \in [\tau_m, \tau_m + \varepsilon] \\ 0, & \text{otherwise, } \tau \in [0, \infty). \end{cases}$$

It satisfies  $\liminf_{\tau \rightarrow \infty} \varphi_2(\tau) = 0$  and  $\limsup_{\tau \rightarrow \infty} \varphi_2(\tau) = s - \beta$  and

$$\lim_{\tau \rightarrow \infty} H_2(\tau) = \lim_{\tau \rightarrow \infty} \frac{n}{\tau^n} \int_0^\tau \varphi_2(r) r^{n-1} dr = 0.$$

Hence its corresponding  $u_2(0, t)$  satisfies  $\lim_{t \rightarrow \infty} u_2(0, t) = 0$  due to Theorem 1. Now set  $\varphi = \varphi_1 + \varphi_2 \in C^0[0, \infty) \cap L^\infty[0, \infty)$ . We will have  $u(0, t) = u_1(0, t) + u_2(0, t)$  with

$$\begin{cases} \liminf_{t \rightarrow \infty} u(0, t) = \liminf_{t \rightarrow \infty} u_1(0, t) = \alpha \\ \limsup_{t \rightarrow \infty} u(0, t) = \limsup_{t \rightarrow \infty} u_1(0, t) = \beta. \end{cases}$$

Also, by  $\varphi(\tilde{\tau}_m) = \varphi_1(\tilde{\tau}_m) + \varphi_2(\tilde{\tau}_m) = \alpha + 0 = \alpha$  and  $\varphi(\tau_m) = \varphi_1(\tau_m) + \varphi_2(\tau_m) = \beta + (s - \beta) = s$ , we have

$$\begin{cases} \liminf_{\tau \rightarrow \infty} \varphi(\tau) = \liminf_{\tau \rightarrow \infty} (\varphi_1(\tau) + \varphi_2(\tau)) = \alpha \\ \limsup_{\tau \rightarrow \infty} \varphi(\tau) = \limsup_{\tau \rightarrow \infty} (\varphi_1(\tau) + \varphi_2(\tau)) = \beta + (s - \beta) = s. \end{cases}$$

(3) For  $r < \alpha < \beta = s$ , the construction of  $\varphi$  is similar to that in (1).

The proof of Theorem 20 is now complete.  $\square$

### 3 Some side issues.

#### 3.1 An example of prescribing four different numbers not satisfying (13) in Theorem 5.

This is related to Theorem 5. Until now, we still do not know how to **prescribe** the oscillation of  $H(\tau)$  and  $u(0, t)$  for four arbitrary numbers  $p < \alpha < \beta < q$  **not** satisfying the condition (13). This will be an interesting problem to explore. However, it is not difficult to construct a particular example not satisfying (13). We have:

**Lemma 22** *There exists a radial function  $\varphi \in C^0[0, \infty) \cap L^\infty[0, \infty)$  so that its average integral  $H(\tau)$  and the convolution solution (2) of the problem (1) satisfy  $p < \alpha < \beta < q$  with  $p + q \neq \alpha + \beta$ .*

**Proof.** The idea of breaking the symmetry, unlike (23), is to choose  $H(\tau)$  to be equal to the sum of **several** (at least two) different slow oscillation functions. For simplicity, we only look at the case  $n = 1$  (the construction for  $n > 1$  is similar, but we cannot find numerical values involving a general variable  $n$ ) and require  $H(\tau)$  to be equal to the sum of **two** slow oscillation functions, given by

$$\begin{aligned} H(\tau) &= \frac{n}{\tau^n} \int_0^\tau \varphi(r) r^{n-1} dr = \frac{1}{\tau} \int_0^\tau \varphi(r) dr \quad (n = 1) \\ &= \sin(\log(\tau + 1)) + \sin(2 \log(\tau + 1)), \quad \tau \in (0, \infty), \quad H(0) = 0. \end{aligned} \quad (73)$$

Similar to (24), one can find the corresponding initial radial function  $\varphi \in C^0[0, \infty) \cap L^\infty[0, \infty)$ . With the help of Maple software, we can evaluate

$$\begin{cases} p = \liminf_{\tau \rightarrow \infty} H(\tau) = \min_{x \in [0, 2\pi]} (\sin x + \sin 2x) \approx -1.760172593 \\ q = \limsup_{\tau \rightarrow \infty} H(\tau) = \max_{x \in [0, 2\pi]} (\sin x + \sin 2x) \approx 1.760172593. \end{cases}$$

By the representation formula  $u(0, t) = \frac{4}{\sqrt{\pi}} \int_0^\infty e^{-z^2} z^2 H(\sqrt{4t}z) dz$ ,  $t \in (0, \infty)$ , we have

$$u(0, t) = \begin{cases} A(t) \sin(\log \sqrt{4t}) + B(t) \cos(\log \sqrt{4t}) \\ + C(t) \sin(2 \log \sqrt{4t}) + D(t) \cos(2 \log \sqrt{4t}), \quad t \in (0, \infty), \end{cases}$$

where

$$\begin{cases} A(t) = \frac{4}{\sqrt{\pi}} \int_0^\infty e^{-z^2} z^2 \cos\left(\log\left(z + \frac{1}{\sqrt{4t}}\right)\right) dz \\ B(t) = \frac{4}{\sqrt{\pi}} \int_0^\infty e^{-z^2} z^2 \sin\left(\log\left(z + \frac{1}{\sqrt{4t}}\right)\right) dz \\ C(t) = \frac{4}{\sqrt{\pi}} \int_0^\infty e^{-z^2} z^2 \cos\left(2 \log\left(z + \frac{1}{\sqrt{4t}}\right)\right) dz \\ D(t) = \frac{4}{\sqrt{\pi}} \int_0^\infty e^{-z^2} z^2 \sin\left(2 \log\left(z + \frac{1}{\sqrt{4t}}\right)\right) dz, \quad t \in (0, \infty). \end{cases}$$

As  $t \rightarrow \infty$ , we have

$$\begin{cases} \lim_{t \rightarrow \infty} A(t) = A \approx 0.892253317, \quad \lim_{t \rightarrow \infty} B(t) = B \approx 0.030945895 \\ \lim_{t \rightarrow \infty} C(t) = C \approx 0.649173672, \quad \lim_{t \rightarrow \infty} D(t) = D \approx 0.099535090, \end{cases}$$

and so

$$\begin{aligned} &\lim_{t \rightarrow \infty} \left| u(0, t) - \left[ A \sin(\log \sqrt{4t}) + B \cos(\log \sqrt{4t}) + C \sin(2 \log \sqrt{4t}) + D \cos(2 \log \sqrt{4t}) \right] \right| \\ &= 0, \end{aligned}$$

which gives

$$\begin{cases} \alpha = \liminf_{t \rightarrow \infty} u(0, t) = \min_{x \in [0, 2\pi]} (A \sin x + B \cos x + C \sin 2x + D \cos 2x) \approx -1.369211837 \\ \beta = \limsup_{t \rightarrow \infty} u(0, t) = \max_{x \in [0, 2\pi]} (A \sin x + B \cos x + C \sin 2x + D \cos 2x) \approx 1.328017886 \end{cases}$$

and we find that  $p + q \neq \alpha + \beta$ .  $\square$

**Remark 23** *We have used different software to evaluate the above numerical values of  $p$ ,  $\alpha$ ,  $\beta$ ,  $q$  and obtain the same values.*

### 3.2 The behavior of $u(x, t)$ as $|x| \rightarrow \infty$ .

In Theorem 5 and Theorem 8 we only look at the behavior of  $u(x, t)$  for fixed  $x$  with  $t \rightarrow \infty$ . We may as well look at the behavior of  $u(x, t)$  for fixed  $t$  with  $|x| \rightarrow \infty$ . In the following lemma, we discuss such property for a particular radial initial data. We choose  $\varphi(x) = \varphi(|x|) \in C^0[0, \infty) \cap L^\infty[0, \infty)$  to be the function in (24) with  $m = 1$ . Its average integral  $H(\tau)$  is equal to  $\sin(\log(\tau + 1))$ ,  $\tau \in [0, \infty)$ , which is our typical slow oscillation function.

**Lemma 24** *Let  $u(x, t)$  be the solution of the problem (1) given by (2) with  $u(x, 0) = \varphi(|x|)$ ,  $x \in \mathbb{R}^n$ , where  $\varphi(\tau)$  is given by*

$$\varphi(\tau) = \sin(\log(\tau + 1)) + \frac{\tau}{n(\tau + 1)} \cos(\log(\tau + 1)), \quad \tau \in [0, \infty). \quad (74)$$

Then for each fixed  $t > 0$  we have

$$\lim_{|x| \rightarrow \infty} \left| u(x, t) - \left( \sin(\log|x|) + \frac{1}{n} \cos(\log|x|) \right) \right| = 0. \quad (75)$$

**Remark 25** *Note that for fixed  $t > 0$ , we have*

$$\tilde{\alpha} := \liminf_{|x| \rightarrow \infty} u(x, t) = \liminf_{\tau \rightarrow \infty} \varphi(\tau) = r = -\sqrt{1 + \frac{1}{n^2}} < \liminf_{\tau \rightarrow \infty} H(\tau) = p = -1$$

and

$$\tilde{\beta} := \limsup_{|x| \rightarrow \infty} u(x, t) = \limsup_{\tau \rightarrow \infty} \varphi(\tau) = s = \sqrt{1 + \frac{1}{n^2}} > \limsup_{\tau \rightarrow \infty} H(\tau) = q = 1.$$

Thus the **space oscillation** of  $u(x, t)$  is, in general, not necessarily bounded by the oscillation of  $H(\tau)$  (however, it is always bounded by the oscillation of  $\varphi(\tau)$  due to the maximum principle). This is different from the behavior for **time oscillation**, where we always have  $r \leq p \leq \alpha \leq \beta \leq q \leq s$ .

**Proof.** By Remark 7, we know that  $u(x, t)$  is **radial** in  $x \in \mathbb{R}^n$  for each  $t > 0$ . Therefore, it suffices to look at the oscillation behavior of  $u(x, t)$  for  $|x| \rightarrow \infty$ . The representation formulas (9) and (10) cannot be used here since they are valid only for  $u(0, t)$ . Instead, we use the convolution formula (2) to get

$$u(x, t) = \frac{1}{(4\pi t)^{n/2}} \int_{\mathbb{R}^n} e^{-\frac{|x-y|^2}{4t}} \varphi(|y|) dy = \frac{1}{\pi^{n/2}} \int_{\mathbb{R}^n} e^{-|z|^2} \varphi\left(|x + \sqrt{4tz}|\right) dz, \quad (76)$$

where by (74) we have

$$\varphi\left(|x + \sqrt{4tz}|\right) = \begin{cases} \sin(\log(|x + \sqrt{4tz}| + 1)) \\ + \frac{|x + \sqrt{4tz}|}{n(|x + \sqrt{4tz}| + 1)} \cos(\log(|x + \sqrt{4tz}| + 1)). \end{cases}$$



By the Lebesgue Dominated Convergence Theorem and the identity

$$\sin \left( \log \left( \left| x + \sqrt{4tz} \right| + 1 \right) \right) = \begin{cases} \sin (\log |x|) \cos \left( \log \left( \frac{|x + \sqrt{4tz}| + 1}{|x|} \right) \right) \\ + \cos (\log |x|) \sin \left( \log \left( \frac{|x + \sqrt{4tz}| + 1}{|x|} \right) \right), \end{cases}$$

where

$$\lim_{|x| \rightarrow \infty} \frac{|x + \sqrt{4tz}| + 1}{|x|} = 1, \quad \text{for fixed } t \text{ and } z,$$

we have

$$\lim_{|x| \rightarrow \infty} \left| \sin \left( \log \left( \left| x + \sqrt{4tz} \right| + 1 \right) \right) - \sin (\log |x|) \right| = 0$$

and so

$$\lim_{|x| \rightarrow \infty} \left| \frac{1}{\pi^{n/2}} \int_{\mathbb{R}^n} e^{-|z|^2} \sin \left( \log \left( \left| x + \sqrt{4tz} \right| + 1 \right) \right) dz - \sin (\log |x|) \right| = 0. \quad (77)$$

Similarly, we have

$$\lim_{|x| \rightarrow \infty} \left| \frac{1}{\pi^{n/2}} \int_{\mathbb{R}^n} e^{-|z|^2} \left( \frac{|x + \sqrt{4tz}|}{n(|x + \sqrt{4tz}| + 1)} \cos \left( \log \left( \left| x + \sqrt{4tz} \right| + 1 \right) \right) \right) dz - \frac{1}{n} \cos (\log |x|) \right| = 0 \quad (78)$$

and (75) follows. The proof is done.  $\square$

**Acknowledgement.** Research supported by NCTS (National Center for Theoretical Sciences) and MoST (Ministry of Science and Technology) of Taiwan with grant number 108-2115-M-007-013-MY2.

## References

- [CE] P. Collet, J.-P. Eckmann, *Space-time behavior in problems of hydrodynamic type: A case study*, **Nonlinearity**, **5** (1992) 1265-1302.
- [CT] M.-S. Chang, D.-H. Tsai, *On the oscillation behavior of solutions to the heat equation on  $\mathbb{R}^n$* , **Journal of Differential Equations**, **268** (2020) 2040-2062.
- [E] S.D. Eidel'man, *Parabolic System*, North-Holland, Amsterdam, 1969.
- [J] F. John, *Partial Differential Equations, 4th edition*, **Applied Mathematical Sciences**, v. **1**, Springer-Verlag, 1982.
- [K] S. Kamin, *On stabilization of solutions of the Cauchy problem for parabolic equations*, **Proc. Roy. Soc. Edinburgh Sect. A**, **76** (1976) 43-53.
- [N] W.-M. Ni, *The mathematics of diffusion*, **CBMS-NSF Regional Conference Series in Applied Math.**, v. **82**, **SIAM** (2011).
- [NT] M. Nara, M. Taniguchi, *The condition on the stability of stationary lines in a curvature flow in the whole plane*, **J. Diff. Eq.** **237** (2007) 61-76.
- [RE] V.D. Repnikov, S.D. Eidel'man, *A new proof of the theorem on the stabilization of the solution of the Cauchy problem for the heat equation*, **Math. USSR Sb.**, **2** (1967) 135-139.

[TN] D.-H. Tsai, C.-H. Nien, *On the oscillation behavior of solutions to the one-dimensional heat equation*, **Discrete & Continuous Dynamical Systems-A**, vol. **39**, no. **7** (2019) 4073-4089.

Dong-Ho Tsai  
Department of Mathematics  
National Tsing Hua University  
Hsinchu 30013, TAIWAN  
E-mail: [dhtsai@math.nthu.edu.tw](mailto:dhtsai@math.nthu.edu.tw)