

AUTOMORPHISMS OF FREE BRAIDED ASSOCIATIVE ALGEBRAS IN TWO VARIABLES

Riza Mutalip¹, Altyngul Naurazbekova², and Ualbai Umirbaev³

ABSTRACT. We describe the groups of automorphisms of two generated free braided associative algebras with involutive diagonal braidings over a field of characteristic $\neq 2$. Depending on the form of the diagonal involutive braiding, five different automorphism groups arise as automorphism groups of two generated free braided associative algebras. This list covers all three automorphism groups that arise in the case of quantum planes [1].

Mathematics Subject Classification (2010): 16W20, 16T20, 16T25.

Key words: Yang-Baxter equation, braiding, free associative algebra, automorphism.

1. INTRODUCTION

It is well known [7, 8, 11, 14] that all automorphisms of the polynomial algebra $K[x_1, x_2]$ and the free associative algebra $K\langle x_1, x_2 \rangle$ in two variables over an arbitrary field K are tame. Moreover [7, 14], the automorphism groups of the algebras $K[x_1, x_2]$ and $K\langle x_1, x_2 \rangle$ are isomorphic, that is

$$\text{Aut}K[x_1, x_2] \cong \text{Aut}K\langle x_1, x_2 \rangle.$$

It is also known that all automorphisms of two generated free Poisson algebras over a field of characteristic zero [15] and two generated free right-symmetric algebras over an arbitrary field [10] are tame. P. Cohn [6] proved that all automorphisms of free Lie algebras of a finite rank over a field are tame. An analogue of this theorem is true for free algebras of any homogeneous Schreier variety of algebras [13]. Recall that the varieties of all non-associative algebras [12], commutative and anti-commutative algebras [17], Lie algebras [18, 22] and Lie superalgebras [16, 19] are Schreier.

Let $q = (q_{ij}) \in M_n(K)$ be a matrix of order $n \geq 2$ whose elements $q_{ij} \in K^*$ satisfy the relations $q_{ii} = q_{ij}q_{ji} = 1$ for all $1 \leq i, j \leq n$. Denote by $O_q = K_q[x_1, \dots, x_n]$ the associative algebra with unity generated by the elements x_1, \dots, x_n which satisfy the relations

$$x_i x_j = q_{ij} x_j x_i, \quad 1 \leq i, j \leq n.$$

The algebra O_q is called a *quantum polynomial algebra* [5]. In the case $n = 2$ the algebra O_q is called a *quantum plane*.

¹L.N. Gumilyov Eurasian National University, Nur-Sultan, Kazakhstan, e-mail: mutalipriza@yahoo.com

²L.N. Gumilyov Eurasian National University, Nur-Sultan, Kazakhstan, e-mail: altyngul.82@mail.ru

³Wayne State University, Detroit, MI 48202, USA, and Institute of Mathematics and Modeling, Almaty, Kazakhstan, e-mail: umirbaev@wayne.edu

The automorphism groups of the quantum polynomial algebras were described by J. Alev and M. Chamarie [1] (see also [5]). All automorphisms of the quantum plane O_q with $q_{12} \neq \pm 1$ are *toric*, i.e. are isomorphic to the multiplicative group $K^* \times K^*$. Moreover, the automorphism group $\text{Aut}(O_q)$ of the quantum plane O_q is isomorphic to $\text{Aut}K\langle x_1, x_2 \rangle$ if $q_{12} = 1$ and is isomorphic to the semidirect product $(K^* \times K^*) \rtimes \mathbb{Z}_2$ if $q_{12} = -1$. Automorphisms of the field of fractions of quantum polynomial algebras were studied in [2, 4, 3].

A *braided space* is a linear space over an arbitrary field K with a linear map $\tau : V \otimes V \rightarrow V \otimes V$ that satisfies the Yang–Baxter equation or the *braid relation*

$$(1) \quad (\tau \otimes \text{id})(\text{id} \otimes \tau)(\tau \otimes \text{id}) = (\text{id} \otimes \tau)(\tau \otimes \text{id})(\text{id} \otimes \tau).$$

The linear map τ will be called a *braiding* of the space V .

If x_1, x_2, \dots, x_n is a basis of a linear space V , then for arbitrary parameters $q_{is} \in K$, $1 \leq i, s \leq n$, the linear map τ defined by

$$(2) \quad \tau : x_i \otimes x_s \mapsto q_{is} \cdot x_s \otimes x_i$$

is a braiding and is called a *diagonal braiding*. Denote by $\tau = (q_{ij}) \in M_n(K)$ this diagonal braiding.

A braiding $\tau : V \otimes V \rightarrow V \otimes V$ is called *involutive* if $\tau^2 = \text{id}$. The diagonal braiding $\tau = (q_{ij}) \in M_n(K)$ is involutive if and only if $q_{ij}q_{ji} = 1$ for all $1 \leq i, j \leq n$. In particular, $q_{ii} = \pm 1$ for all i . The best known example of an involutive braiding is the *ordinary flip* θ defined by $\theta(x \otimes y) = y \otimes x$ for all $x, y \in V$.

It is well known [9] that every braiding $\tau : V \otimes V \rightarrow V \otimes V$ of a vector space V can be uniquely extended to a braiding τ' of the tensor algebra

$$T(V) = \bigoplus_{i=0}^{\infty} V^{\otimes i}.$$

Moreover, there is a canonical braided structure of the Hopf algebra on $T(V)$, which plays an important role in quantum Lie theory (see, for example [9, 20]). Analogues of these results for free nonassociative algebras are also true [21]. If x_1, x_2, \dots, x_n is a basis of the linear space V then $T(V)$ is isomorphic to the free associative algebra $K\langle x_1, x_2, \dots, x_n \rangle$ over K freely generated by x_1, x_2, \dots, x_n .

In this paper we describe the groups of automorphisms of free braided associative algebras in two variables $A_\tau = (K\langle x_1, x_2 \rangle, \tau)$ with a diagonal involutive braiding $\tau = (q_{11}, q_{12}, q_{21}, q_{22})$ over a field of characteristic $\neq 2$. If $q_{ij} = 1$ for all i, j then $\tau = \theta$ is the ordinary flip and the group of automorphisms $\text{Aut}A_\tau$ of A_τ coincides with the group of all automorphisms $\text{Aut}K\langle x_1, x_2 \rangle$ of the free associative algebra $K\langle x_1, x_2 \rangle$. If $q_{ij} = -1$ for all i, j then the group of automorphisms $\text{Aut}A_\tau$ coincides with the group of all odd automorphisms G_{odd} (full definitions are given in Section 3) of the free associative algebra $K\langle x_1, x_2 \rangle$. Furthermore, $\text{Aut}A_\tau$ is isomorphic to the toric group $K^* \times K^*$ if $q_{12} \neq \pm 1$ or $q_{12} = -1, q_{11}q_{22} = -1$, and isomorphic to the group $(K^* \times K^*) \rtimes \mathbb{Z}_2$ if $q_{11} = q_{22}, q_{12} = q_{21}, q_{11}q_{12} = -1$. If $q_{12} = 1$ and $q_{11}q_{22} = -1$ then $\text{Aut}A_\tau$ is isomorphic to the subgroup of triangular automorphisms of $K\langle x_1, x_2 \rangle$ of the form

$$\{\varphi \in \text{Aut}K\langle x_1, x_2 \rangle \mid \varphi(x_1) = \alpha x_1 + g(x_2^2), \varphi(x_2) = \beta x_2, \alpha, \beta \in K^*, g(x) \in K[x]\}.$$

The paper is organized as follows. In Section 2 some necessary definitions and facts on free braided associative algebras are given. In Section 3 we classify involutive diagonal braidings on the free associative algebra $K\langle x_1, x_2 \rangle$ up to isomorphism. In Section 4 we prove the main result of the paper on the description of all groups of automorphisms of the two generated free braided associative algebras over a field of characteristic $\neq 2$ with an involutive diagonal braiding .

2. BRAIDED ALGEBRAS

A *braid monoid* B_n [9] is an associative monoid generated by braids s_1, s_2, \dots, s_{n-1} subject to the relations

$$(3) \quad s_k s_{k+1} s_k = s_{k+1} s_k s_{k+1}, \quad s_i s_j = s_j s_i, \quad 1 \leq k < n-1, \quad |i-j| > 1.$$

The group with generators s_1, s_2, \dots, s_{n-1} and defining relations (3) is called the *Artin braid group*.

Let V be a linear space with a braiding $\tau : V \otimes V \rightarrow V \otimes V$. Consider the linear maps

$$\tau_i = \text{id}^{\otimes(i-1)} \otimes \tau \otimes \text{id}^{\otimes(n-i-1)} : V^{\otimes n} \rightarrow V^{\otimes n}, \quad 1 \leq i < n.$$

By (1), the maps τ_i satisfy all defining relations (3) of the braid monoid, i.e.,

$$\tau_i \tau_{i+1} \tau_i = \tau_{i+1} \tau_i \tau_{i+1}, \quad 1 \leq i < n-1; \quad \tau_i \tau_j = \tau_j \tau_i, \quad |i-j| > 1.$$

Put $[k; k] = 1$ and

$$[m; k] = \tau_{k-1} \tau_{k-2} \cdots \tau_{m+1} \tau_m, \quad [k; m] = \tau_m \tau_{m+1} \cdots \tau_{k-2} \tau_{k-1}, \quad m < k.$$

Consider a map $\nu_r^{k,n} : V^{\otimes n} \rightarrow V^{\otimes n}$, $k \leq r < n$, defined as a superposition of the τ_i 's:

$$(4) \quad \nu_r^{k,n} = [k; r+1][k+1; r+2] \cdots [k+n-r-1; n].$$

It is proven in [9], that

$$\nu_r^{k,n} = [n; r][n-1; r-1] \cdots [n-r+k; k].$$

Let V and V' be spaces with braidings τ and τ' , respectively. A linear map $\varphi : V \rightarrow V'$ is called a *homomorphism of braided spaces* if

$$\tau(\varphi \otimes \varphi) = (\varphi \otimes \varphi)\tau'.$$

An algebra R with a multiplication $\mathbf{m} : R \otimes R \rightarrow R$ is called a *braided algebra* if R is a braided space and

$$(\mathbf{m} \otimes \text{id})\tau = \tau_2 \tau_1 (\text{id} \otimes \mathbf{m}), \quad (\text{id} \otimes \mathbf{m})\tau = \tau_1 \tau_2 (\mathbf{m} \otimes \text{id}).$$

As above, in these formulas we use the so-called ‘‘exponential notation’’ for actions of the operators, that is, the operators in a superposition act from the left to the right.

A *homomorphism of braided algebras* is a linear map that is both a homomorphism of algebras and braided spaces.

Let V be a linear space and

$$T(V) = \bigoplus_{i=0}^{\infty} V^{\otimes i}$$

3

be its tensor algebra. The product in the tensor algebra $T(V)$ will be denoted by \mathbf{m} , that is

$$\mathbf{m} : T(V) \otimes' T(V) \longrightarrow T(V),$$

where the sign \otimes' is the same tensor product \otimes with one additional function that separates the two tensors to which the product \mathbf{m} is applied. For example, $(u \otimes' v)\mathbf{m} = u \otimes v$. Very often we write $v_1 v_2 \dots v_k$ instead of $v_1 \otimes v_2 \otimes \dots \otimes v_k \in V^{\otimes k}$.

We fix a linear basis $X = \{x_i | i \in I\}$ of V . Then $T(V) = K\langle X \rangle$ is the free associative algebra over K with a free set of generators X . The set of all associative words X^* in the alphabet X forms a linear basis for $K\langle X \rangle$. If $w = x_{i_1} x_{i_2} \dots x_{i_k} \in X^*$ then the length k of w will be denoted by $d(w)$. Notice that $w \in V^{\otimes k}$. Set $\text{mdeg}(x_i) = \varepsilon_i$, where $\varepsilon_1, \dots, \varepsilon_n$ is the standard basis for Z^n . The *multidegree* of the word w is defined by

$$\text{mdeg}(v) = \text{mdeg}(x_{i_1}) + \text{mdeg}(x_{i_2}) + \dots + \text{mdeg}(x_{i_s}).$$

It is well known [9] that every braiding τ has a unique extension τ' on the free associative algebra $K\langle X \rangle$ such that $K\langle X \rangle$ is a braided algebra. For any $0 \leq r \leq n$ denote by θ_r the linear map

$$\theta_r : V^{\otimes n} \rightarrow V^{\otimes r} \otimes' V^{\otimes(n-r)}$$

defined by

$$(z_1 z_2 \dots z_n) \theta_r = z_1 z_2 \dots z_r \otimes' z_{r+1} \dots z_n, \quad z_i \in X.$$

The braiding τ' is defined [9] by

$$(5) \quad (u \otimes' v) \tau' = (u \otimes v) \nu_r^{1,n} \theta_{n-r}, \quad u \in V^{\otimes r}, \quad v \in V^{\otimes(n-r)}.$$

In particular, $(1 \otimes' v) \tau' = v \otimes' 1$ and $(u \otimes' 1) \tau' = 1 \otimes' u$ by this definition.

Lemma 1. *Let V be a vector space over a field K with a linear basis $X = \{x_1, x_2, \dots, x_n\}$ and let τ be a diagonal braiding of V defined by (2). Let $u, v \in X^*$, $\text{mdeg}(u) = (s_1, s_2, \dots, s_n)$, and $\text{mdeg}(v) = (t_1, t_2, \dots, t_n)$. Then*

$$(u \otimes' v) \tau' = \prod_{i,j} q_{ij}^{s_i t_j} (v \otimes' u).$$

Proof. Set $s = s_1 + \dots + s_n = d(u)$ and $t = t_1 + \dots + t_n = d(v)$. We prove the statement of the lemma by induction on t . If $t = 0$, then $v = 1$ and $(u \otimes' 1) \tau' = 1 \otimes' u$. Suppose that $t \geq 1$ and $v = v' x_h$. Set $\text{mdeg}(v') = (l_1, l_2, \dots, l_n) = (t_1, t_2, \dots, t_{h-1}, t_h - 1, t_{h+1}, \dots, t_n)$. By (5) and (4), we have

$$\begin{aligned} (u \otimes' v) \tau' &= (u \otimes' v' x_h) \tau' = (u \otimes v' \otimes x_h) \nu_s^{1,s+t} \theta_t \\ &= (u \otimes v' \otimes x_h) [1, s+1] [2, s+2] \dots [t-1, s+t-1] [t, s+t] \theta_t \\ &= (u \otimes v' \otimes x_h) \nu_s^{1,s+t-1} (\tau_{s+t-1} \tau_{s+t-2} \dots \tau_t) \theta_t \\ &= (((u \otimes v') \nu_s^{1,s+t-1} \theta_{t-1}) \otimes x_h) (\tau_{s+t-1} \tau_{s+t-2} \dots \tau_t) \theta_t \\ &= (((u \otimes' v') \tau') \otimes x_h) (\tau_{s+t-1} \tau_{s+t-2} \dots \tau_t) \theta_t. \end{aligned}$$

By the induction proposition, we get

$$\begin{aligned} (u \otimes' v)\tau' &= (u \otimes' v' \otimes x_h)\tau' = \prod_{i,j} q_{ij}^{s_i l_j} (v' \otimes u \otimes x_h)(\tau_{s+t-1}\tau_{s+t-2}\cdots\tau_t)\theta_t \\ &= \prod_{i,j} q_{ij}^{s_i l_j} q_{ih}^{s_i} (v' \otimes x_h \otimes u)\theta_t = \prod_{i,j} q_{ij}^{s_i l_j} q_{ih}^{s_i} (v \otimes u)\theta_t = \prod_{i,j} q_{ij}^{s_i t_j} (v \otimes' u). \quad \square \end{aligned}$$

3. DIAGONAL BRAIDINGS ON $K\langle x_1, x_2 \rangle$

Let V be a two-dimensional vector space over a field K with a linear basis x_1, x_2 . Denote by

$$\tau = (q_{11}, q_{12}, q_{21}, q_{22})$$

the diagonal braiding of V defined by (2).

Let $A = K\langle x_1, x_2 \rangle = T(V)$ be the free associative algebra of the vector space V . Then A is a free braided associative algebra with the braiding τ' defined in Section 2. Denote this extension by the same symbol τ and denote by $A_\tau = (A, \tau)$ the free associative algebra A with braiding τ .

By interchanging the variables x_1 and x_2 , from τ we can get another diagonal braiding, which will be denoted by

$$\tau^* = (q_{22}, q_{21}, q_{12}, q_{11}).$$

Proposition 1. *Let $A_\tau = (K\langle x_1, x_2 \rangle, \tau)$ and $A_\sigma = (K\langle x_1, x_2 \rangle, \sigma)$ be free braided associative algebras in two generators x_1, x_2 over an arbitrary field K with diagonal braidings τ and σ , respectively. Then $A_\tau \cong A_\sigma$ if and only if $\sigma = \tau$ or $\sigma = \tau^*$.*

Proof. Let $\varphi : A_\tau \rightarrow A_\sigma$ be an isomorphism and $\varphi(x_1) = f_1$, $\varphi(x_2) = f_2$. Denote by $L(f_1) = \alpha_1 x_1 + \alpha_2 x_2$ and $L(f_2) = \beta_1 x_1 + \beta_2 x_2$ the linear parts of f_1 and f_2 , respectively. Since φ is an automorphism, we have

$$(6) \quad \det \begin{bmatrix} \alpha_1 & \alpha_2 \\ \beta_1 & \beta_2 \end{bmatrix} \neq 0$$

and

$$((x_i \otimes x_j)\tau)(\varphi \otimes \varphi) = ((x_i \otimes x_j)(\varphi \otimes \varphi))\sigma, \quad 1 \leq i, j \leq 2.$$

Let $\tau = (q_{11}, q_{12}, q_{21}, q_{22})$ and $\sigma = (p_{11}, p_{12}, p_{21}, p_{22})$. Then

$$(7) \quad q_{ij}(\varphi(x_j) \otimes \varphi(x_i)) = (\varphi(x_i) \otimes \varphi(x_j))\sigma, \quad 1 \leq i, j \leq 2.$$

Consequently, for $i = 1$ and $j = 2$ we have

$$L(q_{12}(\varphi(x_2) \otimes \varphi(x_1))) = L((\varphi(x_1) \otimes \varphi(x_2))\sigma),$$

that is

$$q_{12}(\beta_1 x_1 + \beta_2 x_2) \otimes (\alpha_1 x_1 + \alpha_2 x_2) = ((\alpha_1 x_1 + \alpha_2 x_2) \otimes (\beta_1 x_1 + \beta_2 x_2))\sigma.$$

Hence

$$\begin{aligned} & q_{12}(\alpha_1 \beta_1 (x_1 \otimes x_1) + \alpha_2 \beta_1 (x_1 \otimes x_2) + \alpha_1 \beta_2 (x_2 \otimes x_1) + \alpha_2 \beta_2 (x_2 \otimes x_2)) \\ &= p_{11} \alpha_1 \beta_1 (x_1 \otimes x_1) + p_{12} \alpha_1 \beta_2 (x_2 \otimes x_1) + p_{21} \alpha_2 \beta_1 (x_1 \otimes x_2) + p_{22} \alpha_2 \beta_2 (x_2 \otimes x_2). \end{aligned}$$

Comparing the coefficients of the terms $x_1 \otimes x_1$, $x_1 \otimes x_2$, $x_2 \otimes x_1$, $x_2 \otimes x_2$ in this equality, we get

$$(8) \quad (p_{11} - q_{12})\alpha_1\beta_1 = (p_{21} - q_{12})\alpha_2\beta_1 = (p_{12} - q_{12})\alpha_1\beta_2 = (p_{22} - q_{12})\alpha_2\beta_2 = 0.$$

Applying (7) for the remaining values of i and j , we similarly obtain that

$$(9) \quad (p_{11} - q_{11})\alpha_1^2 = (p_{21} - q_{11})\alpha_1\alpha_2 = (p_{12} - q_{11})\alpha_1\alpha_2 = (p_{22} - q_{11})\alpha_2^2 = 0,$$

$$(10) \quad (p_{11} - q_{21})\alpha_1\beta_1 = (p_{21} - q_{21})\alpha_1\beta_2 = (p_{12} - q_{21})\alpha_2\beta_1 = (p_{22} - q_{21})\alpha_2\beta_2 = 0,$$

$$(11) \quad (p_{11} - q_{22})\beta_1^2 = (p_{21} - q_{22})\beta_1\beta_2 = (p_{12} - q_{22})\beta_1\beta_2 = (p_{22} - q_{22})\beta_2^2 = 0.$$

By (6) $\alpha_1\beta_2 \neq 0$ or $\alpha_2\beta_1 \neq 0$. If $\alpha_1\beta_2 \neq 0$ then it follows from (8) – (11) that

$$p_{12} = q_{12}, p_{11} = q_{11}, p_{21} = q_{21}, p_{22} = q_{22}.$$

If $\alpha_2\beta_1 \neq 0$ then, similarly, we get

$$p_{21} = q_{12}, p_{22} = q_{11}, p_{12} = q_{21}, p_{11} = q_{22}.$$

Consequently, if $A_\tau \cong A_\sigma$, then $\sigma = \tau$ or $\sigma = \tau^*$.

Let $\sigma = \tau^*$. Consider the automorphism φ of the free associative algebra $K\langle x_1, x_2 \rangle$ given by $\varphi(x_1) = x_2$ and $\varphi(x_2) = x_1$. To prove that $\varphi : A_\tau \rightarrow A_\sigma$ is an isomorphism of braided algebras, it is sufficient to check that $\tau(\varphi \otimes \varphi) = (\varphi \otimes \varphi)\tau^*$. Let $u, v \in X^*$, $s_i = \deg_{x_i} u$, $t_i = \deg_{x_i} v$, $i \in \{1, 2\}$. Note that $\deg_{x_1} \varphi(u) = s_2$, $\deg_{x_2} \varphi(u) = s_1$, $\deg_{x_1} \varphi(v) = t_2$, $\deg_{x_2} \varphi(v) = t_1$. By Lemma 1, we have

$$((u \otimes v)\tau)(\varphi \otimes \varphi) = q_{11}^{s_1 t_1} q_{12}^{s_1 t_2} q_{21}^{s_2 t_1} q_{22}^{s_2 t_2} (\varphi(v) \otimes \varphi(u))$$

and

$$((u \otimes v)(\varphi \otimes \varphi))\tau^* = q_{11}^{s_1 t_1} q_{12}^{s_1 t_2} q_{21}^{s_2 t_1} q_{22}^{s_2 t_2} (\varphi(v) \otimes \varphi(u)).$$

Hence

$$((u \otimes v)\tau)(\varphi \otimes \varphi) = ((u \otimes v)(\varphi \otimes \varphi))\tau^*$$

and $A_\tau \cong A_\sigma$. \square

Notice that the braiding τ from (2) is involutive if and only if

$$q_{ij}q_{ji} = 1$$

for all $1 \leq i, j \leq n$. Consequently, in the case of two variables the braiding τ from (6) is involutive if and only if

$$q_{11}^2 = q_{12}q_{21} = q_{22}^2 = 1.$$

In particular, $q_{11} = \pm 1$ and $q_{22} = \pm 1$. Note that if τ is involutive then τ^* is also involutive.

Corollary 1. *Let $A_\tau = (K\langle x_1, x_2 \rangle, \tau)$ be a free braided associative algebra in two variables x_1, x_2 over a field K of characteristic $p \neq 2$ with an involutive diagonal braiding $\tau = (q_{11}, q_{12}, q_{21}, q_{22})$. Then either $q_{12} \neq \pm 1$ or A_τ is isomorphic to A_σ , where σ is equal to one of the following vectors:*

$$(1, 1, 1, 1), (-1, -1, -1, -1), (1, 1, 1, -1), (-1, 1, 1, -1), (1, -1, -1, 1), (1, -1, -1, -1).$$

Proof. If $q_{12} = 1$ then τ is involutive if only if it is equal to one of the following vectors:

$$(1, 1, 1, 1), (1, 1, 1, -1), (-1, 1, 1, 1), (-1, 1, 1, -1).$$

By Proposition 1, $A_{(-1,1,1,1)} \cong A_{(1,1,1,-1)}$.

If $q_{12} = -1$ then τ is equal to one of the following vectors:

$$(1, -1, -1, 1), (1, -1, -1, -1), (-1, -1, -1, 1), (-1, -1, -1, -1).$$

By Proposition 1, $A_{(1,-1,-1,-1)} \cong A_{(-1,-1,-1,1)}$. \square

4. AUTOMORPHISMS OF $A_\tau = (K\langle x_1, x_2 \rangle, \tau)$

Let $K\langle X \rangle = K\langle x_1, x_2, \dots, x_n \rangle$ be the free associative algebra over K with the set of free generators $X = \{x_1, x_2, \dots, x_n\}$. Denote by $\varphi = (f_1, f_2, \dots, f_n)$ an automorphism of the algebra $K\langle X \rangle$ such that $\varphi(x_1) = f_1, \varphi(x_2) = f_2, \dots, \varphi(x_n) = f_n$. Denote by $\text{Aut}K\langle X \rangle$ the set of all automorphisms of the algebra $K\langle X \rangle$ over K . An automorphism of the form

$$(x_1, \dots, x_{i-1}, \alpha x_i + f, x_{i+1}, \dots, x_n),$$

where $0 \neq \alpha \in K$, $f \in K\langle X \setminus \{x_i\} \rangle$, is called *elementary*. An automorphism φ is called *tame* if it is a product of elementary automorphisms.

Consider the natural grading

$$K\langle X \rangle = A_0 \oplus A_1 \oplus \dots \oplus A_m \oplus \dots,$$

where $A_0 = K$, $A_1 = Kx_1 + Kx_2 + \dots + Kx_n$, and A_m is the linear span of words of length m . Every nonzero element $f \in K\langle X \rangle$ can be uniquely represented in the form

$$f = f_0 + f_1 + \dots + f_{m-1} + f_m,$$

where $f_i \in A_i$, $f_m \neq 0$. Then f_m is called *the highest homogeneous part* of f and is denoted by \bar{f} . Set also $\deg f = m$. The degree of an automorphism $\varphi = (f_1, f_2, \dots, f_n)$ is defined by

$$\deg \varphi = \deg f_1 + \deg f_2 + \dots + \deg f_n.$$

An *elementary transformation* of an n -tuple $\varphi = (f_1, f_2, \dots, f_n)$ changes only one element f_i for some i with an element of the form $\alpha f_i + g$, where $0 \neq \alpha \in K$ and $g \in K\langle f_j | j \neq i \rangle$, and leaves unchanged all other elements f_j , where $j \neq i$. We write $\varphi \rightarrow \psi$ if ψ is obtained from φ by one elementary transformation. An automorphism φ is called *elementarily reducible* if there exists an automorphism ψ such that $\varphi \rightarrow \psi$ and $\deg \psi < \deg \varphi$.

Every automorphism of the free associative algebra $A = K\langle x_1, x_2 \rangle$ in two variables over a field K is tame [7, 14]. In the next lemma we formulate a property of automorphisms of $A = K\langle x_1, x_2 \rangle$ that we need in the future.

Lemma 2. *Let $A = K\langle x_1, x_2 \rangle$ be the free associative algebra in two variables x_1, x_2 . If $\varphi = (f_1, f_2)$ is an automorphism of the algebra A with degree ≥ 3 , then*

$$\bar{f}_1 = \alpha(ax_1 + bx_2)^r, \quad \bar{f}_2 = \beta(ax_1 + bx_2)^s,$$

where $r + s \geq 3$, $r \mid s$ or $s \mid r$, $\alpha, \beta \in K^*$, $a, b \in K$, and $(a, b) \neq (0, 0)$.

Proof. Let $\deg f_1 = r$ and $\deg f_2 = s$. The property $r \mid s$ or $s \mid r$ is explicitly formulated in [7, 14]. Without loss of generality, we can assume that $r \leq s$. Prove the remaining statement of the lemma by induction on $\deg \varphi = r + s$.

Every automorphism of A with degree ≥ 3 is elementarily reducible [7, 14]. Therefore, there is an elementary transformation $(f_1, f_2) \rightarrow (f_1, \gamma f_2 - g(f_1)) = (f_1, f'_2)$, where $\gamma \neq 0$, $g(f_1) \in K\langle f_1 \rangle$, such that $\deg f'_2 = \deg(\gamma f_2 - g(f_1)) < s$. If $\deg(f_1, f'_2) \geq 3$ then by the induction proposition we get $\overline{f_1} = \alpha(ax_1 + bx_2)^r$. Notice that this is true even if $\deg(f_1, f'_2) = 2$. Consequently,

$$\overline{f_2} = 1/\gamma \overline{g(f_1)} = \beta(ax_1 + bx_2)^s$$

for some $\beta \in K^*$. \square

We fix the grading

$$(12) \quad A = B_0 \oplus B_1$$

of the algebra A , where B_0 is the linear span of all monomials of even length and B_1 is the linear span of all monomials of odd length.

An automorphism $\varphi = (f_1, f_2)$ of A is called *odd* if $f_1, f_2 \in B_1$. The set of all odd automorphisms forms a group and this group will be denoted by G_{odd} .

Define the following subgroups of the group of automorphisms $A = \text{Aut}K\langle x_1, x_2 \rangle$:

- (1) $G_1 = \text{Aut}K\langle x_1, x_2 \rangle$ is the group of all automorphisms of A ;
- (2) $G_2 = G_{\text{odd}}$ is the group of all odd automorphisms of A ;
- (3) $G_3 = \{\varphi \in G_1 \mid \varphi = (\alpha_1 x_1, \beta_2 x_2) \text{ or } \varphi = (\beta_1 x_2, \alpha_2 x_1), \alpha_1, \beta_2, \alpha_2, \beta_1 \in K^*\}$;
- (4) $G_4 = \{\varphi \in G_1 \mid \varphi = (\alpha_1 x_1 + g(x_2^2), \beta_2 x_2), \alpha_1, \beta_2 \in K^*, g(x) \in K[x]\}$;
- (5) $G_5 = \{\varphi \in G_1 \mid \varphi = (\alpha_1 x_1, \beta_2 x_2), \alpha_1, \beta_2 \in K^*\}$ is the group of all *toric* automorphisms of $K\langle x_1, x_2 \rangle$.

Let $A_\tau = (A, \tau)$ be the free braided associative algebra in two generators x_1, x_2 over a field K of arbitrary characteristic $\neq 2$ with an involutive diagonal braiding $\tau = (q_{11}, q_{12}, q_{21}, q_{22})$. Notice that an automorphism $\varphi \in \text{Aut}A$ of the algebra A is an automorphism of A_τ if and only if

$$(13) \quad \tau(\varphi \otimes \varphi) = (\varphi \otimes \varphi)\tau.$$

We identify the group of automorphisms $\text{Aut}A_\tau$ of A_τ with the corresponding subgroup of $\text{Aut}K\langle x_1, x_2 \rangle$.

Lemma 3. *If $\tau = (1, 1, 1, 1)$ then $\text{Aut}A_\tau = G_1$.*

Proof. For any $u, v \in X^*$ we have $(u \otimes v)\tau = v \otimes u$ by Lemma 1. Then

$$((u \otimes v)\tau)(\varphi \otimes \varphi) = \varphi(v) \otimes \varphi(u) = (\varphi(u) \otimes \varphi(v))\tau = ((u \otimes v)(\varphi \otimes \varphi))\tau$$

for any $\varphi \in G_1$ and any $u, v \in X^*$. Consequently, (13) is true for all $\varphi \in G_1$. This means that $G_\tau = G_1$. \square

Lemma 4. *If $\tau = (-1, 1, 1, -1)$ or $\tau = (1, -1, -1, 1)$ then $\text{Aut}A_\tau = G_3$.*

Proof. Let $\varphi = (f_1, f_2)$ be an automorphism of the algebra A_τ with degree ≥ 3 . By Lemma 2, we have

$$(14) \quad \overline{f_1} = \gamma_1(ax_1 + bx_2)^{s_1}, \overline{f_2} = \gamma_2(ax_1 + bx_2)^{s_2},$$

where $s_1 + s_2 \geq 3$, $s_1 \mid s_2$, or $s_2 \mid s_1$, $\gamma_1, \gamma_2 \in K^*$, $a, b \in K$ and $(a, b) \neq 0$. By (13), we have

$$\overline{((x_i \otimes x_j)\tau)(\varphi \otimes \varphi)} = \overline{((x_i \otimes x_j)(\varphi \otimes \varphi))\tau}, \quad i, j \in \{1, 2\}.$$

Hence

$$\overline{q_{ij}\varphi(x_j) \otimes \varphi(x_i)} = \overline{(\varphi(x_i) \otimes \varphi(x_j))\tau}, \quad i, j \in \{1, 2\}.$$

By (14), we have

$$q_{ij}\gamma_i\gamma_j(ax_1 + bx_2)^{s_j} \otimes (ax_1 + bx_2)^{s_i} = \gamma_i\gamma_j((ax_1 + bx_2)^{s_i} \otimes (ax_1 + bx_2)^{s_j})\tau, \quad i, j \in \{1, 2\}.$$

Using Lemma 1 and comparing the coefficients of the terms $x_1^{s_j} \otimes x_1^{s_i}$, $x_2^{s_j} \otimes x_1^{s_i}$, $x_1^{s_j} \otimes x_2^{s_i}$, $x_2^{s_j} \otimes x_2^{s_i}$ in this equality, we obtain

$$(q_{ij} - q_{11}^{s_i s_j})a^{s_i + s_j} = (q_{ij} - q_{12}^{s_i s_j})a^{s_i}b^{s_j} = (q_{ij} - q_{21}^{s_i s_j})a^{s_j}b^{s_i} = (q_{ij} - q_{22}^{s_i s_j})b^{s_i + s_j} = 0, \quad i, j \in \{1, 2\}.$$

Varying the values of $i, j \in \{1, 2\}$, we get

$$(15) \quad (q_{11} - q_{11}^{s_1^2})a^{2s_1} = (q_{11} - q_{12}^{s_1^2})a^{s_1}b^{s_1} = (q_{11} - q_{21}^{s_1^2})a^{s_1}b^{s_1} = (q_{11} - q_{22}^{s_1^2})b^{2s_1} = 0,$$

$$(16) \quad (q_{12} - q_{11}^{s_1 s_2})a^{s_1 + s_2} = (q_{12} - q_{12}^{s_1 s_2})a^{s_1}b^{s_2} = (q_{12} - q_{21}^{s_1 s_2})a^{s_2}b^{s_1} = (q_{12} - q_{22}^{s_1 s_2})b^{s_1 + s_2} = 0,$$

$$(17) \quad (q_{21} - q_{11}^{s_1 s_2})a^{s_1 + s_2} = (q_{21} - q_{12}^{s_1 s_2})a^{s_2}b^{s_1} = (q_{21} - q_{21}^{s_1 s_2})a^{s_1}b^{s_2} = (q_{21} - q_{22}^{s_1 s_2})b^{s_1 + s_2} = 0,$$

$$(18) \quad (q_{22} - q_{11}^{s_2^2})a^{2s_2} = (q_{22} - q_{12}^{s_2^2})a^{s_2}b^{s_2} = (q_{22} - q_{21}^{s_2^2})a^{s_2}b^{s_2} = (q_{22} - q_{22}^{s_2^2})b^{2s_2} = 0.$$

If $\tau = (-1, 1, 1, -1)$ then (15) – (18) implies that

$$\begin{aligned} & (-1 - (-1)^{s_1^2})a^{2s_1} = (-1 - (-1)^{s_1^2})b^{2s_1} \\ & = (1 - (-1)^{s_1 s_2})a^{s_1 + s_2} = (1 - (-1)^{s_1 s_2})b^{s_1 + s_2} \\ & = (-1 - (-1)^{s_2^2})a^{2s_2} = (-1 - (-1)^{s_2^2})b^{2s_2} = 0. \end{aligned}$$

It is easy to check that over a field of characteristic $\neq 2$, these equations give $a = b = 0$.

If $\tau = (1, -1, -1, 1)$ then (16) gives

$$(-1 - 1^{s_1 s_2})a^{s_1 + s_2} = (-1 - 1^{s_1 s_2})b^{s_1 + s_2} = 0$$

and, consequently, $a = b = 0$.

Therefore, if $\tau = (-1, 1, 1, -1)$ or $\tau = (1, -1, -1, 1)$, then A_τ does not have automorphisms of degree ≥ 3 . Consequently,

$$\varphi = (\alpha_1 x_1 + \beta_1 x_2 + \gamma_1, \alpha_2 x_1 + \beta_2 x_2 + \gamma_2), \quad \alpha_i, \beta_i, \gamma_i \in K.$$

By (13), we have

$$((x_i \otimes x_j)\tau)\varphi \otimes \varphi = ((x_i \otimes x_j)\varphi \otimes \varphi)\tau, \quad i, j \in \{1, 2\}.$$

Hence

$$\begin{aligned} & q_{ij}(\alpha_j x_1 + \beta_j x_2 + \gamma_j) \otimes (\alpha_i x_1 + \beta_i x_2 + \gamma_i) \\ & = ((\alpha_i x_1 + \beta_i x_2 + \gamma_i) \otimes (\alpha_j x_1 + \beta_j x_2 + \gamma_j))\tau, \quad i, j \in \{1, 2\}. \end{aligned}$$

Comparing the coefficients in the terms $x_1 \otimes x_1$, $x_1 \otimes x_2$, $x_2 \otimes x_1$, $x_2 \otimes x_2$, x_1 , x_2 , and 1 in this equality, we get

$$(q_{ij} - q_{11})\alpha_i \alpha_j = (q_{ij} - q_{12})\alpha_i \beta_j = (q_{ij} - q_{21})\alpha_j \beta_i = (q_{ij} - q_{22})\beta_i \beta_j$$

$$= (q_{ij} - 1)(\alpha_i \gamma_j + \alpha_j \gamma_i) = (q_{ij} - 1)(\beta_i \gamma_j + \beta_j \gamma_i) = (q_{ij} - 1)\gamma_i \gamma_j = 0, \quad i, j \in \{1, 2\}.$$

Varying the values of $i, j \in \{1, 2\}$, we have

$$(19) \quad (q_{11} - q_{12})\alpha_1 \beta_1 = (q_{11} - q_{21})\alpha_1 \beta_1 = (q_{11} - q_{22})\beta_1^2 = (q_{11} - 1)\alpha_1 \gamma_1 \\ = (q_{11} - 1)\beta_1 \gamma_1 = (q_{11} - 1)\gamma_1^2 = 0,$$

$$(20) \quad (q_{12} - q_{11})\alpha_1 \alpha_2 = (q_{12} - q_{21})\alpha_2 \beta_1 = (q_{12} - q_{22})\beta_1 \beta_2 \\ = (q_{12} - 1)(\alpha_1 \gamma_2 + \alpha_2 \gamma_1) = (q_{12} - 1)(\beta_1 \gamma_2 + \beta_2 \gamma_1) = (q_{12} - 1)\gamma_1 \gamma_2 = 0,$$

$$(21) \quad (q_{21} - q_{11})\alpha_1 \alpha_2 = (q_{21} - q_{12})\alpha_2 \beta_1 = (q_{21} - q_{22})\beta_1 \beta_2 \\ = (q_{21} - 1)(\alpha_2 \gamma_1 + \alpha_1 \gamma_2) = (q_{21} - 1)(\beta_2 \gamma_1 + \beta_1 \gamma_2) = (q_{21} - 1)\gamma_2 \gamma_1 = 0,$$

$$(22) \quad (q_{22} - q_{11})\alpha_2^2 = (q_{22} - q_{12})\alpha_2 \beta_2 = (q_{22} - q_{21})\alpha_2 \beta_2 = (q_{22} - 1)\alpha_2 \gamma_2 \\ = (q_{22} - 1)\beta_2 \gamma_2 = (q_{22} - 1)\gamma_2^2 = 0.$$

If $\tau = (-1, 1, 1, -1)$ or $\tau = (1, -1, -1, 1)$, it follows from (19) – (22) that

$$\alpha_1 \beta_1 = \alpha_1 \alpha_2 = \beta_1 \beta_2 = \alpha_2 \beta_2 = \gamma_1^2 = \gamma_2^2 = 0$$

or

$$\alpha_1 \beta_1 = \alpha_1 \alpha_2 = \beta_1 \beta_2 = \alpha_2 \beta_2 = \alpha_1 \gamma_2 + \alpha_2 \gamma_1 = \beta_1 \gamma_2 + \beta_2 \gamma_1 = \gamma_1 \gamma_2 = 0,$$

respectively. Using this and (6), we get

$$\alpha_1 \neq 0, \beta_2 \neq 0, \beta_1 = \alpha_2 = \gamma_1 = \gamma_2 = 0$$

or

$$\beta_1 \neq 0, \alpha_2 \neq 0, \alpha_1 = \beta_2 = \gamma_1 = \gamma_2 = 0.$$

This means that

$$\varphi = (\alpha_1 x_1, \beta_2 x_2) \text{ or } \varphi = (\beta_1 x_2, \alpha_2 x_1).$$

Using Lemma 1, it is easy to check that each of these automorphisms is an automorphism of A_τ . Consequently, $\text{Aut}A_\tau = G_3$. \square

Lemma 5. *If $\tau = (-1, -1, -1, -1)$ then $\text{Aut}A_\tau = G_2$.*

Proof. Let $u, v \in X^*$, $s = d(u)$, and $t = d(v)$. If $\tau = (-1, -1, -1, -1)$ then, by Lemma 1,

$$(u \otimes v)\tau = (-1)^{st}(v \otimes u).$$

Notice that $(-1)^{st} = -1$ if and only if both s and t are odd. This implies that

$$(23) \quad (f \otimes g)\tau = (-1)^{ij}(g \otimes f)$$

for any homogeneous elements $f \in B_i$ and $g \in B_j$ with respect to the grading (12).

Let $\varphi \in G_2 = G_{\text{odd}}$ be an arbitrary odd automorphism of $A = K\langle x_1, x_2 \rangle$. Notice that $\varphi(B_i) \subseteq B_i$. Using (23), we get

$$(f \otimes g)\tau(\varphi \otimes \varphi) = (-1)^{ij}(g \otimes f)(\varphi \otimes \varphi) \\ = (-1)^{ij}(\varphi(g) \otimes \varphi(f)) = (\varphi(f) \otimes \varphi(g))\tau = (f \otimes g)(\varphi \otimes \varphi)\tau,$$

that is, (13) holds for φ . Consequently, $\varphi \in \text{Aut}A_\tau$ and $G_2 \subseteq \text{Aut}A_\tau$.

Let $\varphi = (f_1, f_2)$ be an arbitrary automorphism of the algebra A_τ . We prove that $\varphi \in G_2$ by induction on $\deg(\varphi)$. Let $\deg(\varphi) = 2$, that is,

$$\varphi = (\alpha_1x_1 + \beta_1x_2 + \gamma_1, \alpha_2x_1 + \beta_2x_2 + \gamma_2), \quad \alpha_i, \beta_i, \gamma_i \in K.$$

It follows from (19) – (22) that $\gamma_1 = \gamma_2 = 0$. Consequently,

$$\varphi = (\alpha_1x_1 + \beta_1x_2, \alpha_2x_1 + \beta_2x_2) \in G_2.$$

Suppose that $\deg(\varphi) \geq 3$, $\deg f_1 = s_1$, and $\deg f_2 = s_2$. Then (15) – (18) imply that

$$\begin{aligned} (-1 - (-1)^{s_i s_j}) a^{s_i + s_j} &= (-1 - (-1)^{s_i s_j}) a^{s_i} b^{s_j} \\ &= (-1 - (-1)^{s_i s_j}) a^{s_j} b^{s_i} = (-1 - (-1)^{s_i s_j}) b^{s_i + s_j} = 0, \quad i, j \in \{1, 2\}. \end{aligned}$$

If $s_1 s_2$ is even then these equations give $a = b = 0$ over a field of characteristic $\neq 2$. Consequently, s_1 and s_2 are both odd. Suppose that $s_1 | s_2$ for definiteness by Lemma 2. Set $t = s_2 / s_1$. Then there exists an odd automorphism $\lambda = (x_1, x_2 + \beta x_1^t)$ such that $\deg(\varphi\lambda) < \deg(\varphi)$. Since $\lambda \in G_2 \subseteq A_\tau$, it follows that $\varphi\lambda \in A_\tau$. By the induction proposition, $\varphi\lambda \in G_2$. Consequently, $\varphi \in G_2$. \square

Lemma 6. *If $\tau = (1, 1, 1, -1)$ then $\text{Aut}_K A_\tau = G_4$.*

Proof. Consider the grading

$$A = C_0 \oplus C_1$$

of the algebra A , where C_0 is the linear span of all monomials of even degree in x_2 and C_1 is the linear span of all monomials of odd degree in x_2 .

Let $u, v \in X^*$, $\text{mdeg}(u) = (s, s')$, $\text{mdeg}(v) = (t, t')$. Lemma 1 gives

$$(u \otimes v)\tau = (-1)^{s't'}(v \otimes u).$$

Notice that $(-1)^{s't'} = -1$ if and only if both s and t are odd. This implies that

$$(24) \quad (f \otimes g)\tau = (-1)^{ij}(g \otimes f)$$

for any homogeneous elements $f \in C_i$ and $g \in C_j$.

Let $\varphi \in G_4$. Notice that $\varphi(C_i) \subseteq C_i$. Using (24), we easily get

$$\begin{aligned} (f \otimes g)\tau(\varphi \otimes \varphi) &= (-1)^{ij}(g \otimes f)(\varphi \otimes \varphi) \\ &= (-1)^{ij}(\varphi(g) \otimes \varphi(f)) = (\varphi(f) \otimes \varphi(g))\tau = (f \otimes g)(\varphi \otimes \varphi)\tau. \end{aligned}$$

Consequently, $\varphi \in \text{Aut}A_\tau$ and $G_4 \subseteq \text{Aut}A_\tau$.

Let $\varphi = (f_1, f_2)$ be an arbitrary automorphism of the algebra A_τ . We prove by induction on $\deg \varphi$ that $\varphi \in G_4$. Let $\deg(\varphi) = 2$, i.e.,

$$\varphi = (\alpha_1x_1 + \beta_1x_2 + \gamma_1, \alpha_2x_1 + \beta_2x_2 + \gamma_2), \quad \alpha_i, \beta_i, \gamma_i \in K.$$

It follows from (19) – (22) that

$$\beta_1^2 = \beta_1\beta_2 = \alpha_2^2 = \alpha_2\beta_2 = \alpha_2\gamma_2 = \beta_2\gamma_2 = \gamma_2^2 = 0.$$

Using this and (6), we get

$$\alpha_1 \neq 0, \beta_2 \neq 0, \beta_1 = \alpha_2 = \gamma_2 = 0.$$

This means that

$$\varphi = (\alpha_1 x_1 + \gamma_1, \beta_2 x_2) \in G_4.$$

Suppose that $\deg(\varphi) \geq 3$, $\deg f_1 = s_1$, and $\deg f_2 = s_2$. As in the proof of Lemma 4, (15) – (18) imply that

$$(-1 - 1^{s_2^2})a^{2s_2} = (1 - (-1)^{s_1^2})b^{2s_1} = (1 - (-1)^{s_1 s_2})b^{s_1 + s_2} = (-1 - (-1)^{s_2^2})b^{2s_2} = 0.$$

From this we immediately get $a = 0$ over a field of characteristic $\neq 2$. Then $b \neq 0$. This is possible only if $(s_1, s_2) = (2k, 2l - 1)$ for some positive integers k, l . By Lemma 2, $s_2 | s_1$ and $t = s_1/s_2 = 2k/(2l - 1)$ is even. Consequently, there exists $\lambda = (x_1 + \beta x_2^t, x_2) \in G_4$ such that $\deg(\varphi\lambda) < \deg(\varphi)$. Since $\lambda \in G_4 \subseteq A_\tau$, it follows that $\varphi\lambda \in A_\tau$. By the induction proposition, $\varphi\lambda \in G_4$. Consequently, $\varphi \in G_4$. \square

Lemma 7. *If $\tau = (1, -1, -1, -1)$ then $\text{Aut}A_\tau = G_5$.*

Proof. Let $\varphi = (f_1, f_2)$ be an automorphism of the algebra A_τ . If $\deg(\varphi) \geq 3$, $\deg f_1 = s_1$, and $\deg f_2 = s_2$, then (15) – (18) imply that

$$\begin{aligned} (-1 - 1^{s_1 s_2})a^{s_1 + s_2} &= (1 - (-1)^{s_1^2})b^{2s_1} \\ &= (-1 - (-1)^{s_1 s_2})b^{s_1 + s_2} = (-1 - (-1)^{s_2^2})b^{2s_2} = 0. \end{aligned}$$

It is easy to check that over a field of characteristic $\neq 2$, these equations give $a = b = 0$. Therefore, A_τ does not have any automorphisms with degree ≥ 3 .

Suppose that

$$\varphi = (\alpha_1 x_1 + \beta_1 x_2 + \gamma_1, \alpha_2 x_1 + \beta_2 x_2 + \gamma_2), \quad \alpha_i, \beta_i, \gamma_i \in K.$$

It follows from (19) – (22) that

$$\alpha_1 \beta_1 = \beta_1^2 = \alpha_1 \alpha_2 = \alpha_1 \gamma_2 + \alpha_2 \gamma_1 = \beta_1 \gamma_2 + \beta_2 \gamma_1 = \gamma_1 \gamma_2 = \alpha_2^2 = \alpha_2 \gamma_2 = \beta_2 \gamma_2 = \gamma_2^2 = 0.$$

Using this and (6), we get

$$\alpha_1 \neq 0, \beta_2 \neq 0, \beta_1 = \alpha_2 = \gamma_1 = \gamma_2 = 0.$$

This means that

$$\varphi = (\alpha_1 x_1, \beta_2 x_2) \in G_5.$$

Using Lemma 1, it is easy to check that each element of G_5 is an automorphism of A_τ . Consequently, $\text{Aut}A_\tau = G_5$. \square

Lemma 8. *If $\tau = (q_{11}, q_{12}, q_{21}, q_{22})$ with $q_{12} \neq \pm 1$ then $\text{Aut}A_\tau = G_5$.*

Proof. Let $\tau = (q_{11}, q_{12}, q_{21}, q_{22})$ with $q_{12} \neq \pm 1$. We have $q_{11}^2 = q_{22}^2 = q_{12}q_{21} = 1$ since $\tau^2 = \text{id}$. Consequently, $q_{11} = \pm 1$, $q_{22} = \pm 1$, $q_{21} \neq q_{12}$, $q_{12} \neq \pm 1$.

Let $\varphi = (f_1, f_2)$ be an automorphism of the algebra A_τ . If $\deg(\varphi) \geq 3$, $\deg f_1 = s_1$, and $\deg f_2 = s_2$, then over a field of characteristic $\neq 2$, (15) – (18) give $a = b = 0$. Therefore, A_τ does not have any automorphisms with degree ≥ 3 .

Suppose that

$$\varphi = (\alpha_1 x_1 + \beta_1 x_2 + \gamma_1, \alpha_2 x_1 + \beta_2 x_2 + \gamma_2), \quad \alpha_i, \beta_i, \gamma_i \in K.$$

It follows from (19) – (22) that

$$\alpha_1 \beta_1 = \alpha_1 \alpha_2 = \alpha_2 \beta_1 = \beta_1 \beta_2 = \alpha_2 \beta_2 = \alpha_2 \gamma_1 + \alpha_1 \gamma_2 = \beta_2 \gamma_1 + \beta_1 \gamma_2 = \gamma_1 \gamma_2 = 0.$$

Using this and (6), we get

$$\alpha_1 \neq 0, \beta_2 \neq 0, \beta_1 = \alpha_2 = \gamma_1 = \gamma_2 = 0.$$

This means that

$$\varphi = (\alpha_1 x_1, \beta_2 x_2) \in G_5.$$

Using Lemma 1, it is easy to check that each element of G_5 is an automorphism of A_τ . Consequently, $\text{Aut}A_\tau = G_5$. \square

Theorem 1. *Let $A_\tau = \langle K\langle x_1, x_2 \rangle, \tau \rangle$ be a free braided associative algebra in two generators x_1, x_2 over a field K of arbitrary characteristic $\neq 2$ with an involutive diagonal braiding $\tau = (q_{11}, q_{12}, q_{21}, q_{22})$. Then*

- (1) $\text{Aut}A_\tau = \text{Aut}K\langle x_1, x_2 \rangle$ if $q_{ij} = 1$ for all i, j ;
- (2) $\text{Aut}A_\tau = G_{\text{odd}}$ if $q_{ij} = -1$ for all i, j , where G_{odd} is the group of all odd automorphisms of $K\langle x_1, x_2 \rangle$;
- (3) $\text{Aut}A_\tau \cong (K^* \times K^*) \rtimes \mathbb{Z}_2$ if $q_{11} = q_{22}, q_{12} = q_{21}$, and $q_{11}q_{12} = -1$;
- (4) $\text{Aut}A_\tau \cong G_4$ if $q_{12} = 1$ and $q_{11}q_{22} = -1$;
- (5) $\text{Aut}A_\tau \cong K^* \times K^*$ if $q_{12} \neq \pm 1$ or $q_{12} = -1, q_{11}q_{22} = -1$.

Proof. The statements (1) and (2) are proven in Lemmas 3 and 5, respectively. Let $q_{11} = q_{22}, q_{12} = q_{21}$, and $q_{11}q_{12} = -1$. Using Proposition 1, we can assume that $\tau = (-1, 1, 1, -1)$ or $\tau = (1, -1, -1, 1)$. By Lemma 4, $\text{Aut}A_\tau \cong G_3$. Notice that

$$G_3 \cong (K^* \times K^*) \rtimes \mathbb{Z}_2,$$

where \mathbb{Z}_2 is the subgroup of G_3 generated by (x_2, x_1) .

If $q_{12} = 1$ and $q_{11}q_{22} = -1$ then we can assume that $\tau = (1, 1, 1, -1)$ by Proposition 1. By Lemma 6, $\text{Aut}A_\tau \cong G_4$. If $q_{12} \neq \pm 1$ then $\text{Aut}A_\tau = G_5 \cong K^* \times K^*$ by Lemma 8. If $q_{12} = -1$ and $q_{11}q_{22} = -1$ then we can assume that $\tau = (1, -1, -1, -1)$ by Proposition 1. Then Lemma 7 gives $\text{Aut}A_\tau = G_5$ again. \square

Since $\text{Aut}K\langle x_1, x_2 \rangle \cong \text{Aut}K[x_1, x_2]$, the groups $\text{Aut}K\langle x_1, x_2 \rangle$, $K^* \times K^*$, and $(K^* \times K^*) \rtimes \mathbb{Z}_2$ are the automorphism groups of quantum planes [1].

Acknowledgments

This work was supported by the grant of the Ministry of Education and Science of the Republic of Kazakhstan (project AP08052290).

REFERENCES

- [1] Alev, J., Chamarie, M. (1992). Dérivations et automorphismes de quelques algèbres quantiques. *Comm. Algebra* 20(6):1787–1802.
- [2] Alev, J., Dumas, F. (1997). Invariants du corps de Weyl sous l'action de groupes finis. (French. English summary) [Invariants of the Weyl field under the action of finite groups] *Comm. Algebra* 25(5):1655–1672.
- [3] Artamonov, V.A., Cohn, P.M. (1999). The skew field of rational functions on the quantum plane. (English summary) *Algebra*, 11. *J. Math. Sci. (New York)* 93(6):824–829.
- [4] Artamonov, V.A. (2000). Automorphisms of the skew field of rational quantum functions. (Russian) *Mat. Sb.* 191(12):3–26; translation in *Sb. Math.* 191(11–12):1749–1771.
- [5] Artamonov, V.A. (2002). Automorphisms and derivations of quantum polynomials. (English summary) *Recent advances in Lie theory (Vigo, 2000)*, 109–120, *Res. Exp. Math.*, 25, Heldermann.

- [6] Cohn, P.M. (1964). Subalgebras of free associative algebras, Proc. London Math. Soc. (3)14:618–632.
- [7] Czerniakiewicz, A.J. (1971,1972). Automorphisms of a free associative algebra of rank 2. I, II. Trans. Amer. Math. Soc. 160:393–401; 171:309–315.
- [8] Jung, H.W.E. (1942). Über ganze birationale Transformationen der Ebene. J. reine angew. Math. 184:161–174.
- [9] Kharchenko, V. (2015). Quantum Lie theory. A multilinear approach, Lecture Notes in Mathematics, vol. 2150, Springer, Cham, xiii+302 pp.
- [10] Kozybaev, D., Makar-Limanov, L., Umirbaev, U. (2008). The Freiheitssatz and the automorphisms of free right-symmetric algebras. Asian-Eur. J. Math. 1(2):243–254.
- [11] van der Kulk, W. (1953). On polynomial rings in two variables. Nieuw Archief voor Wisk. (3)1:33–41.
- [12] Kurosh, A.G. (1947). Nonassociative free algebras and free products of algebras. Mat. Sb. 20:239–262.
- [13] Lewin, J. (1968). On Schreier varieties of linear algebras. Trans. Amer. Math. Soc. 132:553–562.
- [14] Makar-Limanov, L.G. (1970). The automorphisms of the free algebra of two generators. Funktsional. Anal. i Prilozhen. 4(3):107–108.
- [15] Makar-Limanov, L., Turusbekova, U., Umirbaev, U. (2009). Automorphisms and derivations of free Poisson algebras in two variables. J. Algebra 322:3318–3330.
- [16] Mikhalev, A.A. (1985). Subalgebras of free colored Lie superalgebras. Mat. Zametki 37(5):653–661.
- [17] Shirshov, A.I. (1954). Subalgebras of free commutative and free anticommutative algebras. (Russian) Mat. Sbornik N.S. 34(76):81–88.
- [18] Shirshov, A.I. (1953). Subalgebras of free Lie algebras. (Russian) Mat. Sbornik 33(75):441–452.
- [19] Shtern, A.S. (1986). Free Lie superalgebras. Sibirsk. Mat. Zh. 27(1):170–174.
- [20] Takeuchi, M. (2000). Survey of braided Hopf algebras, in: New Trends in Hopf Algebra Theory, in: Contemp. Math. 267:301–324, AMS, Providence, RI.
- [21] Umirbaev, U., Kharchenko, V. (2017). Free braided nonassociative Hopf algebras and Sabinin τ -algebras. J. Algebra 492:130–156.
- [22] Witt, E. (1956). Die Unterringe der freien Lieschen Rings. Math. Z. 64:195–216.