From Short-Range to Contact Interactions in Two-dimensional Many-Body Quantum Systems

Marcel Griesemer[∗] and Michael Hofacker† Fachbereich Mathematik, Universität Stuttgart, D-70569 Stuttgart, Germany

Abstract

Quantum systems composed of N distinct particles in \mathbb{R}^2 with two-body contact interactions of TMS type are shown to arise as limits - in the norm resolvent sense - of Schrödinger operators with suitably rescaled pair potentials.

1 Introduction

Many-particle quantum systems with short-range interactions are often described by simplified models with zero-range (contact) interactions. Effective models of low energy nuclear physics, the Lieb-Liniger model for the 1d Bose gas, and the Fermi polaron model for an impurity within an ideal Fermi gas are well-known models of this type [25, 14, 18]. The understanding of these models is that zero-range interactions provide an idealized description of short-ranged interparticle forces whose details are not known or considered irrelevant. In the present paper we fully justify this idealization for 2d many-particle systems with (spin-independent) two-body forces. We prove norm resolvent convergence for suitably rescaled Schrödinger operators towards TMS Hamiltonians [6]. This mathematically justifies the use of TMS Hamiltonians for describing contact interactions, and, moreover, provides a new way of defining them.

In the Hilbert space

$$
\mathscr{H} := L^2\left(\mathbb{R}^{2N}, d(x_1,\ldots,x_N)\right),
$$

we consider Schrödinger operators of the form

$$
H_{\varepsilon} := \sum_{i=1}^{N} (-\Delta_{x_i}/m_i) - \sum_{\sigma \in \mathcal{I}} g_{\varepsilon,\sigma} V_{\sigma,\varepsilon}(x_j - x_i), \qquad \varepsilon > 0,
$$
 (1.1)

where $m_i > 0$ is the mass of the *i*th particle, I denotes the set of all pairs $\sigma = (i, j)$, $i < j$, and

$$
V_{\sigma,\varepsilon}(r) := \varepsilon^{-2} V_{\sigma}(r/\varepsilon), \qquad \varepsilon > 0.
$$
\n(1.2)

We assume $V_{\sigma} \in L^1 \cap L^2(\mathbb{R}^2)$, $V_{\sigma}(-r) = V_{\sigma}(r)$ as well as some mild decay of $V_{\sigma}(r)$ as $|r| \to \infty$. For H_{ε} to have a limit as $\varepsilon \to 0$, the coupling constant $g_{\varepsilon,\sigma}$ must have an asymptotic behavior, as $\varepsilon \to 0$, adjusted to the space dimension. For $d = 2$ we may set

$$
-\frac{1}{g_{\varepsilon,\sigma}} = \mu_{\sigma} \left[a_{\sigma} \ln(\varepsilon) + b_{\sigma} \right] + o(1)
$$
\n(1.3)

[∗]marcel.griesemer@mathematik.uni-stuttgart.de

[†]michael.hofacker@mathematik.uni-stuttgart.de

with $a_{\sigma}, b_{\sigma} \in \mathbb{R}$, $a_{\sigma} > 0$. Here the reduced mass μ_{σ} of the pair $\sigma = (i, j)$ has been factored out. We assume that

$$
a_{\sigma} \ge \frac{1}{2\pi} \int V_{\sigma}(r) \, \mathrm{d}r. \tag{1.4}
$$

If equality holds in (1.4), we will see that the two-body interaction V_{σ} gives rise to a non-trivial contact interaction, whose strength is determined by the parameter b_{σ} . In the case of inequality there is no contribution from V_σ . This is well-known for the case of $N=1$ particles, where our problem was solved long ago [2]. We use $\mathcal{J} \subset \mathcal{I}$ to denote the subset of pairs for which equality holds in (1.4) .

Our main result asserts norm resolvent convergence of H_{ε} as $\varepsilon \to 0$. In addition, we give an explicit expression for the resolvent of the limiting Hamiltonian, a characterization of the domain in terms of so-called TMS boundary conditions at the collision planes $\{x \in \mathbb{R}^{2N} \mid x_i = x_j\}$, and we show, in the case of equal masses $m_j = 1$, that our limiting Hamiltonian agrees with the Hamiltonian H_{*β*} (for suitable $\beta = (\beta_{\sigma})_{\sigma \in \mathcal{I}}$) of Dell'Antonio et al. (see [6, Eqs. (5.3)-(5.4)]). To describe these results in detail, we need some auxiliary spaces and operators.

Let

$$
\mathfrak{X} := \bigoplus_{\sigma \in \mathcal{J}} \mathfrak{X}_{\sigma},
$$
\n
$$
\mathfrak{X}_{\sigma=(i,j)} := L^2 \left(\mathbb{R}^{2(N-1)}, \mathbf{d}\left(R, x_1, \dots, \widehat{x}_i \dots, \widehat{x}_j \dots, x_N\right) \right),
$$
\n
$$
(1.5)
$$

where the hat, as in \hat{x}_i , indicates omission of a variable. Let $H_0 = \sum_{i=1}^N$ $\sum_{i=1}$ (- Δ_{x_i}/m_i) on $D(H_0)$ = $H^2(\mathbb{R}^{2N})$ and let

$$
T_{\sigma}:D(T_{\sigma})\to\mathfrak{X}_{\sigma}
$$

denote the H_0 -bounded trace operator, which, for smooth functions ψ , is given by

$$
(T_{\sigma}\psi)(R, x_1, ... \hat{x}_i ... \hat{x}_j ... , x_N) = \psi(x_1, ..., x_N)|_{x_i = x_j = R}, \qquad \sigma = (i, j). \tag{1.6}
$$

Then $G(z)_{\sigma} = T_{\sigma}R_0(z)$, with $R_0(z) = (H_0 + z)^{-1}$, is a bounded operator from \mathscr{H} to \mathfrak{X}_{σ} . This operator is closely related to the Green's function of $H_0 + z$, see, e.g., (4.52), below, and the letter *G* serves to remind us of the connection. Let *T* and $G(z) = TR_0(z)$ denote the operator-valued vectors with components T_{σ} and $G(z)_{\sigma}$, respectively. Our main result is the following:^{*}

Theorem 1.1. Suppose, for all $\sigma \in \mathcal{I}$, that $V_{\sigma} \in L^1 \cap L^2(\mathbb{R}^2)$, $V_{\sigma}(-r) = V_{\sigma}(r)$, and there exists *some* $s > 0$ *such that* $\int |r|^{2s} |V_{\sigma}(r)| dr < \infty$ *. Let* H_{ε} *be defined by* (1.1)*-*(1.4)*. Then, as* $\varepsilon \to 0$ *, H^ε converges in norm resolvent sense to a self-adjoint, semibounded operator H. The resolvent* $of H, for z \in \rho(H_0) \cap \rho(H), obeys$

$$
(H+z)^{-1} = R_0(z) + G(\overline{z})^* \Theta(z)^{-1} G(z)
$$
\n(1.7)

with some invertible and unbounded operator $\Theta(z) : D \subset \mathfrak{X} \to \mathfrak{X}$ *, whose domain is independent of z.*

Remarks:

1. Θ(*z*) is an operator matrix with entries $\Theta(z)_{\sigma\nu} : \mathfrak{X}_{\nu} \to \mathfrak{X}_{\sigma}$. For $\sigma \neq \nu$ these entries are bounded operators given by $\Theta(z)_{\sigma\nu} = -T_{\sigma}G(\overline{z})^*_{\nu} = -T_{\sigma}(T_{\nu}R_0(\overline{z}))^*$. The unbounded operators $\Theta(z)_{\sigma\sigma}$ are given explicitly in Equation (4.32), below.

[∗]Note our definition of the resolvent set at the end of this section.

2. Antisymmetric wave functions from $H^2(\mathbb{R}^{2N})$ belong to the kernel of all trace operators *T*_σ. Hence for identical particles, by (1.7), $(H + z)^{-1} = R_0(z)$ on the subspace $\mathscr{H}_{\text{anti}} \subset$ $\mathscr H$ of antisymmetric wave functions. In fact, there are no contact interactions among identical, spin-aligned fermions in space dimensions $d \geq 2$ [10]. Theorem 1.1 implies $\lim_{\varepsilon \to 0} (H_{\varepsilon} + z)^{-1} = R_0(z)$ on $\mathscr{H}_{\text{anti}}$ and its proof gives a rate of convergence [10].

The main novelty in Theorem 1.1, compared to previous results of similar type, is that convergence is established in norm resolvent sense. Norm resolvent convergence implies convergence of the spectrum, which is not true for the weaker strong resolvent convergence [24]. In the present case, where $\sigma(H_{\varepsilon}) = [\Sigma_{\varepsilon}, \infty)$, the norm resolvent convergence implies $\sigma(H) = [\Sigma, \infty)$ with $\Sigma = \lim_{\varepsilon \to 0} \Sigma_{\varepsilon}$.

The analog of Theorem 1.1 in the case $N = 1$ is well-known and true without a condition of the type (1.4). But if $N > 1$ this condition is necessary: if (1.4) is not satisfied for some pair σ , then one may expect strong resolvent convergence at best. To see this, consider a twoparticle system with pair potential *V* and reduced mass $\mu = 1$. If $0 < a < \frac{1}{2\pi} \int V(r) dr$ and $-g_{\varepsilon}^{-1} = a \ln(\varepsilon) + b$, then the Hamiltonian in the center of mass frame has a negative eigenvalue running towards $-\infty$ as $\varepsilon \to 0$ [23]. Due to the center of mass motion, the Hamiltonian H_{ε} then has essential spectrum filling the entire real axis in the limit $\varepsilon \to 0$. This is not compatible with norm resolvent convergence towards a semibounded Hamiltonian.

By the following corollary, domain vectors of *H* satisfy the so-called TMS boundary conditions at the collision planes $\{x \in \mathbb{R}^{2N} \mid x_i = x_j\}.$

Corollary 1.2. *Under the assumptions of Theorem 1.1, a vector* $\psi \in \mathcal{H}$ *belongs to* $D(H)$ *if and only if the following holds: For some (and hence all)* $z \in \rho(H_0) \cap \rho(H)$ *there exist* $\psi_0 \in D(H_0)$ $and w \in D$ *such that*

$$
\psi = \psi_0 + G(\overline{z})^* w \tag{1.8}
$$

and

$$
T\psi_0 = \Theta(z)w.\tag{1.9}
$$

In this case,

$$
(H+z)\psi = (H_0+z)\psi_0.
$$
\n(1.10)

The vectors ψ_0 *and w are uniquely determined by* $\psi \in D(H)$ *and* $z \in \rho(H_0) \cap \rho(H)$ *.*

Choosing $w = 0$ in Corollary 1.2, we see that $D(H_0) \cap \text{Ker } T \subset D(H)$ and that $H = H_0$ on $D(H_0) \cap \text{Ker } T$. Thus *H* is a self-adjoint extension of $H_0 \restriction \text{Ker } T$. It is well-known that a Krein formula, like (1.7), is the characteristic equation for the resolvent of all such extensions [19, 20].

Recently, in a study of the 2d stochastic heat equation, Gu, Quastel and Tsai have derived a result very similar to Theorem 1.1 for *N* identical particles [12]. In [12] the two-body potentials are compactly supported smooth functions and convergence in strong resolvent sense is established. In the case of bosons, Gu et al. give a formula for the resolvent that is similar to the one in Theorem 1.1 and the subsequent remark. In one space dimension, results similar to Theorem 1.1 for $N = 3$ quantum particles are obtained in [3], and for arbitrary $N \ge 1$ in [11]. The well-known Lieb-Liniger model with repulsive *δ*-interactions is derived from a trapped 3d Bose gas with two-body potentials in [22]. TMS Hamiltonians like *H* in Theorem 1.1 have also been described as resolvent limits of *N*-body Hamiltonians, where the regularized two-body contact interaction is an integral operator, rather than a potential, and the regularization is achieved by an ultraviolet cutoff [6, 7, 9] or a reversed heat flow [8]. In these cases the convergence is easier to establish than in the case studied here. Nevertheless, all previous approximation results of this kind in 2d with $N \geq 2$ particles establish *strong* resolvent convergence only. In three space dimensions, TMS Hamiltonians for $N \geq 3$ bosons are symmetric but not self-adjoint, and all

(non-trivial) self-adjoint extensions are unbounded below [17, 16]. A possible solution to this problem in the case of $N = 3$ bosons is worked out in the recent paper [4], where a semi-bounded self-adjoint Hamiltonian containing a three-body contact interaction is constructed.

To prove Theorem 1.1, we further develop the methods and tools introduced in our previous paper [11]. The key elements along with some auxiliary spaces and tools are described in Section 2. Sections 3 and 4 provide all preparations needed for the proof of Theorem 1.1, which is given in Section 5. In addition, we derive a lower bound on $\sigma(H)$ in Section 5. In Section 6 we compute the quadratic form of the Hamiltonian *H* and we show it agrees, in the case of equal masses, with a quadratic form derived by Dell'Antonio et al. [6], and which, in recent years, has become a standard starting point for investigations of TMS Hamiltonians.

We conclude this introduction with some remarks on our notations and with the proof of Corollary 1.2. In this paper the resolvent set $\rho(H)$ of a closed operator *H* is defined as the set of $z \in \mathbb{C}$ for which $H + z : D(H) \subset \mathcal{H} \to \mathcal{H}$ is a bijection. This differs by a minus sign from the conventional definition and it means that the spectrum $\sigma(H)$ is the complement of $-\rho(H)$. The L^2 -norm will be denoted by $\|\cdot\|$, without index, while all other norms carry the space as an index, as e.g. in $||V||_{L^1}$.

Proof of Corollary 1.2. In this proof, $T: D(H_0) \to \mathfrak{X}$ and $G(z): \mathcal{H} \to \mathfrak{X}$ are the operatorvalued vectors defined in terms of the components T_{σ} and $G(z)_{\sigma}$. We know from [5] and Proposition 5.1 that a point $z \in \rho(H_0)$ belongs to $\rho(H)$ if and only if $\Theta(z)$ has a bounded inverse.

Suppose $\psi \in D(H)$ and $z \in \rho(H_0) \cap \rho(H)$. Define $\varphi := (H + z)\psi$, $\psi_0 := R_0(z)\varphi$, and $w := \Theta(z)^{-1} T \psi_0$. Then, by (1.7),

$$
\psi = (H + z)^{-1} \varphi = R_0(z)\varphi + G(\overline{z})^* \Theta(z)^{-1} G(z)\varphi
$$

= $\psi_0 + G(\overline{z})^* w$.

Conversely, if (1.8) and (1.9) hold for some $z \in \rho(H_0) \cap \rho(H)$, let $\varphi := (H_0 + z)\psi_0$. Then $T\psi_0 = TR_0(z)\varphi = G(z)\varphi$ and $w = \Theta(z)^{-1}T\psi_0 = \Theta(z)^{-1}G(z)\varphi$. Hence, by (1.8) and (1.7),

$$
\psi = \psi_0 + G(\overline{z})^* w = R_0(z)\varphi + G(\overline{z})^* \Theta(z)^{-1} G(z)\varphi
$$

=
$$
(H + z)^{-1} \varphi \in D(H).
$$

It follows that $(H + z)\psi = \varphi = (H_0 + z)\psi_0$. In particular, $\psi_0 = (H_0 + z)^{-1}(H + z)\psi$ is uniquely determined by ψ , and, since $\Theta(z)$ is invertible, it follows from (1.9) that $w \in D$ is unique as well. 口

2 Auxiliary operators and strategy of the proof

The proof of Theorem (1.1) starts with the new expression (2.9), below, for H_{ε} , which allows for the explicit representation of the resolvent in terms of a generalized Konno-Kuroda formula, see Equation (2.11) . We now describe the various operators occurring in (2.9) and (2.11) . To this end, we need the auxiliary Hilbert spaces

$$
\widetilde{\mathfrak{X}} := \bigoplus_{\sigma \in \mathcal{I}} \widetilde{\mathfrak{X}}_{\sigma},
$$
\n
$$
\widetilde{\mathfrak{X}}_{\sigma=(i,j)} := L^2 \left(\mathbb{R}^{2N}, \mathrm{d}\left(r, R, x_1, \dots \widehat{x}_i \dots \widehat{x}_j \dots, x_N\right) \right).
$$
\n(2.1)

The integration variables r and R in (2.1) correspond to the relative and center of mass coordinates

$$
r = r_{\sigma} := x_j - x_i,
$$
 $R = R_{\sigma} := \frac{m_i x_i + m_j x_j}{m_i + m_j}$ (2.2)

of the particles constituting the pair $\sigma = (i, j)$. This change of coordinates is implemented unitarily by the operator $\mathscr{K}_{(i,j)} : \mathscr{H} \to \widetilde{\mathfrak{X}}_{(i,j)}$ defined by

$$
\left(\mathcal{K}_{(i,j)}\Psi\right)(r, R, x_1, \dots \hat{x}_i \dots \hat{x}_j \dots, x_N)
$$

:= $\Psi\left(x_1, \dots, x_{i-1}, R - \frac{m_j r}{m_i + m_j}, x_{i+1}, \dots, x_{j-1}, R + \frac{m_i r}{m_i + m_j}, x_{j+1}, \dots, x_N\right).$ (2.3)

The adjoint thereof is the operator $\mathscr{K}^*_{(i,j)} : \widetilde{\mathfrak{X}}_{(i,j)} \to \mathscr{H}$ with

$$
(\mathcal{K}_{(i,j)}^* \Psi)(x_1, ..., x_N) = \Psi\left(x_j - x_i, \frac{m_i x_i + m_j x_j}{m_i + m_j}, x_1, ... \hat{x}_i ... \hat{x}_j ..., x_N\right).
$$
 (2.4)

Let $U_{\varepsilon} \in \mathscr{L}(L^2(\mathbb{R}^2))$ denote the unitary rescaling that, on $\mathscr{L}(\widetilde{\mathfrak{X}}_{(i,j)})$, is given by

$$
(U_{\varepsilon}\Psi)(r,\underline{X}) := \varepsilon \Psi(\varepsilon r,\underline{X}), \qquad \underline{X} = (R,x_1,...\widehat{x}_i...\widehat{x}_j...,x_N), \qquad (2.5)
$$

and let

$$
v_{\sigma}(r) = |V_{\sigma}(r)|^{1/2},
$$

$$
u_{\sigma}(r) = J_{\sigma}|V_{\sigma}(r)|^{1/2}, \qquad J_{\sigma} := \text{sgn}(V_{\sigma}),
$$

so that $V_{\sigma} = u_{\sigma}v_{\sigma}$. Then, in terms of the above operators, we define new operators $A_{\varepsilon,\sigma}, B_{\varepsilon,\sigma}$: $D(A_{\varepsilon,\sigma}) \subseteq \mathscr{H} \to \widetilde{\mathfrak{X}}_{\sigma}$ by

$$
A_{\varepsilon,\sigma} := (v_{\sigma} \otimes 1) \varepsilon^{-1} U_{\varepsilon} \mathscr{K}_{\sigma}, \tag{2.6}
$$

$$
B_{\varepsilon,\sigma} := (u_{\sigma} \otimes 1) \,\varepsilon^{-1} U_{\varepsilon} \mathscr{K}_{\sigma} = J_{\sigma} A_{\varepsilon,\sigma},\tag{2.7}
$$

where $D(A_{\varepsilon,\sigma})$ is determined by the domain of the multiplication operator $v_{\sigma} \otimes 1$. Obviously, $A_{\varepsilon,\sigma}$ and $B_{\varepsilon,\sigma}$ are densely defined and closed. These operators allow us to write the two-body interaction in the form

$$
V_{\sigma,\varepsilon}(x_j - x_i) = \mathcal{K}_{\sigma}^*(V_{\sigma,\varepsilon} \otimes 1)\mathcal{K}_{\sigma} = A_{\varepsilon,\sigma}^* B_{\varepsilon,\sigma}
$$
\n(2.8)

and therefore

$$
H_{\varepsilon} = H_0 - \sum_{\sigma \in \mathcal{I}} g_{\varepsilon,\sigma} A_{\varepsilon,\sigma}^* B_{\varepsilon,\sigma}, \qquad \varepsilon > 0.
$$
 (2.9)

Since $V_{\sigma} \in L^2(\mathbb{R}^2)$, it is clear from (2.8) that $A^*_{\varepsilon,\sigma}B_{\varepsilon,\sigma}$ as well as $A^*_{\varepsilon,\sigma}A_{\varepsilon,\sigma}$ is infinitesimally *H*₀-bounded. So H_{ε} is self-adjoint on $D(H_0)$, and moreover, the hypotheses of Corollary B.2 are satisfied. This means that

$$
\Lambda_{\varepsilon}(z)_{\sigma\nu} = g_{\varepsilon,\sigma}^{-1} \delta_{\sigma,\nu} - B_{\varepsilon,\sigma} R_0(z) A_{\varepsilon,\nu}^*, \qquad \sigma, \nu \in \mathcal{I}
$$
\n(2.10)

are bounded operators and that $\Lambda_{\varepsilon}(z)$ has a bounded inverse if and only if $z \in \rho(H_{\varepsilon})$. Then

$$
(H_{\varepsilon} + z)^{-1} = R_0(z) + \sum_{\sigma,\nu \in \mathcal{I}} (A_{\varepsilon,\sigma} R_0(\overline{z}))^* (\Lambda_{\varepsilon}(z)^{-1})_{\sigma\nu} B_{\varepsilon,\nu} R_0(z).
$$
 (2.11)

In Section 3 we prove existence of the limit

$$
S(z)_{\sigma} = \lim_{\varepsilon \to 0} A_{\varepsilon,\sigma} R_0(z), \qquad \sigma \in \mathcal{I}
$$
 (2.12)

for some, and hence for all $z \in \rho(H_0)$. This works in all dimensions $d \in \{1,2,3\}$. Obviously, $\lim_{\varepsilon \to 0} B_{\varepsilon,\sigma} R_0(z) = J_{\sigma} S(z)_{\sigma}$. We will see that

$$
S(z)_{\sigma} \Psi = v_{\sigma} \otimes G(z)_{\sigma} \Psi,
$$
\n(2.13)

where $G(z)_{\sigma} = T_{\sigma}R_0(z)$ and T_{σ} is the trace operator introduced in (1.6).

The hard part, which is non-trivial even for $N = 1$, is the convergence of $\Lambda_{\varepsilon}(z)^{-1}$ in space dimensions $d \geq 2$. The limit is, in general, not the inverse of an operator. Our analysis is based on the decomposition

$$
\Lambda_{\varepsilon}(z)=\Lambda_{\varepsilon}(z)_{\rm diag}+\Lambda_{\varepsilon}(z)_{\rm off}
$$

into diagonal and off-diagonal parts of the operator matrix (2.10). We then show that both $(\Lambda_{\varepsilon}(z)_{\text{diag}})^{-1}$ and $(\Lambda_{\varepsilon}(z)_{\text{diag}})^{-1}\Lambda_{\varepsilon}(z)_{\text{off}}$ have limits as $\varepsilon \to 0$. The diagonal parts $B_{\varepsilon,\sigma}R_0(z)A_{\varepsilon,\sigma}^*$ contain a divergent contribution that must be cancelled by the divergence of $g_{\varepsilon,\sigma}^{-1}$ as $\varepsilon \to 0$. It turns out that $(g_{\varepsilon,\sigma}^{-1} - B_{\varepsilon,\sigma} R_0(z) A_{\varepsilon,\sigma}^*)^{-1}$ has a vanishing limit unless $\int V_{\sigma}(r) dr = 2\pi a_{\sigma}$. In the end, we arrive at

$$
\lim_{\varepsilon \to 0} (\Lambda_{\varepsilon}(z)^{-1})_{\sigma \nu} = \begin{cases} \frac{|u_{\sigma}\rangle \langle v_{\nu}|}{\langle u_{\sigma} | v_{\sigma} \rangle \langle u_{\nu} | v_{\nu} \rangle} \otimes (\Theta(z)^{-1})_{\sigma \nu} & \text{if } \sigma, \nu \in \mathcal{J} \\ 0 & \text{else} \end{cases}
$$
(2.14)

for large enough $z > 0$, with some closed and invertible operator $\Theta(z)$ in the Hilbert space \mathfrak{X} . Combining (2.14) with (2.12) and (2.13), it follows that the expression on the right hand side of (2.11) has the limit (1.7) . It is then a standard argument to show that (1.7) defines the resolvent of a self-adjoint operator.

3 The limit of $A_{\varepsilon,\sigma}R_0(z)$

Let $\varepsilon > 0$, $z \in \rho(H_0)$ and $\sigma \in \mathcal{I}$. Then the operator $A_{\varepsilon,\sigma} R_0(z)$ is bounded because, by (2.6), $A_{\varepsilon,\sigma}$ is closed and $D(H_0) \subset D(A_{\varepsilon,\sigma})$. In this section we prove that

$$
S(z)_{\sigma} = \lim_{\varepsilon \to 0} A_{\varepsilon,\sigma} R_0(z), \qquad z \in \rho(H_0)
$$
\n(3.1)

exists in the norm of $\mathscr{L}(\mathscr{H}, \widetilde{\mathfrak{X}}_{\sigma})$. In fact, we have an explicit expression for $S(z)_{\sigma}$ in terms of the trace T_{σ} defined in (1.6). Clearly,

$$
T_{\sigma} = \tau \mathcal{K}_{\sigma},\tag{3.2}
$$

where $\mathscr{K}_{\sigma} \in \mathscr{L}(\mathscr{H}, \widetilde{\mathfrak{X}}_{\sigma})$ is the change of coordinates defined in (2.3) and, for smooth functions *ψ* with compact support,

$$
(\tau\psi) (R, x_3, ..., x_N) := \psi(r, R, x_3, ..., x_N)|_{r=0}.
$$

We shall use (3.2) for the definition of T_{σ} , where the trace τ is defined by

$$
\widehat{\tau\psi}(\underline{P}) := \frac{1}{2\pi} \int_{\mathbb{R}^2} \widehat{\psi}(p, \underline{P}) \, dp,\tag{3.3}
$$

 (p, P) being conjugate to $(r, R, x_3, ..., x_N)$, and

$$
D(\tau) = \left\{ \psi \in L^{2}(\mathbb{R}^{2N}) \middle| \int \left(\int \left| \widehat{\psi}(p, \underline{P}) \right| \, \mathrm{d}p \right)^{2} \mathrm{d} \underline{P} < \infty \right\}.
$$

For lack of a good reference, we state and prove the following result about *τ* :

Lemma 3.1. For all $s > 1$, $H^s(\mathbb{R}^{2N}) \subset D(\tau)$ and $\tau : H^s(\mathbb{R}^{2N}) \to H^{s-1}(\mathbb{R}^{2N-2})$ is a bounded *operator.*

Remark: From the Definition (3.2) of T_{σ} , from $\mathscr{K}_{\sigma} H^2(\mathbb{R}^{2N}) = H^2(\mathbb{R}^{2N})$, and from Lemma 3.1, it follows that $H^2(\mathbb{R}^{2N}) \subset D(T_\sigma)$.

Proof. We leave it to the reader to check that $H^s(\mathbb{R}^{2N}) \subset D(\tau)$. For $\psi \in H^s(\mathbb{R}^{2N})$, we have, by Definition (3.3) and by Cauchy-Schwarz inequality,

$$
|\widehat{\tau\psi}(P)|^2 \le \frac{1}{4\pi^2} \int |\widehat{\psi}(p,P)|^2 (1+|p|^2+|P|^2)^s \, \mathrm{d}p \cdot \int (1+|p|^2+|P|^2)^{-s} \, \mathrm{d}p,\tag{3.4}
$$

where

$$
\frac{1}{4\pi^2} \int (1+|p|^2+|\underline{P}|^2)^{-s} \, \mathrm{d}p = C_s (1+|\underline{P}|^2)^{1-s},\tag{3.5}
$$

and $C_s < \infty$ because $s > 1$. From (3.4) and (3.5), it follows that $\|\tau\psi\|_{H^{s-1}} \leq C_s^{1/2} \|\psi\|_{H^s}$. \Box

In view of $A_{\varepsilon,\sigma} = [(\varepsilon^{-1}v_{\sigma}U_{\varepsilon}) \otimes 1] \mathscr{K}_{\sigma}$, proving (3.1) is a problem in $\mathscr{L}(L^2(\mathbb{R}^2))$, which we solve in the following lemma. For a given pair σ , let $V = V_{\sigma}$, $v = v_{\sigma}$ and let $|v\rangle \langle \delta| : H^2(\mathbb{R}^2) \to L^2(\mathbb{R}^2)$ denote the rank-one operator given by

$$
|v\rangle\langle\delta|\,\psi:=\psi(0)v,
$$

where $\psi(0) := (2\pi)^{-1} \int \widehat{\psi}(p) dp$. Then the following holds:

Lemma 3.2. *If* $\int (1 + |r|^{2s}) |V(r)| dr < \infty$ *for some* $s \in (0, 1)$ *, then*

$$
v\varepsilon^{-1}U_{\varepsilon} \to |v\rangle \langle \delta|, \qquad (\varepsilon \downarrow 0)
$$

in the norm of $\mathscr{L}(H^2(\mathbb{R}^2), L^2(\mathbb{R}^2))$ *and the rate of convergence is at least* $O(\varepsilon^s)$ *.*

Proof. Due to the Sobolev embedding $H^2(\mathbb{R}^2) \hookrightarrow C^{0,s}(\mathbb{R}^2)$, valid for $s \in (0,1)$, we have that

$$
\left| \left(v \varepsilon^{-1} U_{\varepsilon} \psi - |v\rangle \langle \delta | \psi \right) (r) \right| = |v(r)| \, |\psi(\varepsilon r) - \psi(0)|
$$

$$
\leq |v(r)| \, c_s |\varepsilon r|^s \, ||\psi||_{H^2}.
$$

This implies the statement of the lemma.

We now come to the proof of (3.1) :

Proposition 3.3. *If* $z \in \rho(H_0)$ *and* $\int (1 + |r|^{2s}) |V_\sigma(r)| dr < \infty$ *for some* $s \in (0,1)$ *, then, as ε* ↓ 0*,*

$$
||A_{\varepsilon,\sigma}R_0(z)-S(z)_{\sigma}||=O(\varepsilon^s),
$$

where $S(z)_{\sigma} \in \mathscr{L}(\mathscr{H}, \widetilde{\mathfrak{X}}_{\sigma})$ *is given by*

$$
S(z)_{\sigma}\psi = v_{\sigma} \otimes (G(z)_{\sigma}\psi) \tag{3.6}
$$

 $and G(z)_{\sigma} = T_{\sigma}R_0(z) \in \mathcal{L}(\mathcal{H}, \mathfrak{X}_{\sigma}).$

Proof. By (2.6), we have that $A_{\varepsilon,\sigma}R_0(z) = (v_{\sigma} \otimes 1)\varepsilon^{-1}U_{\varepsilon}\mathscr{K}_{\sigma}R_0(z)$. Comparing this with $S(z)_{\sigma}\psi = v_{\sigma} \otimes (\tau \mathscr{K}_{\sigma}R_0(z)\psi) = (v_{\sigma}) \langle \delta | \otimes 1 \rangle \mathscr{K}_{\sigma}R_0(z)\psi$, we see that the proposition follows from Lemma 3.2 since \mathscr{K}_{σ} defines a bounded operator in $H^2(\mathbb{R}^{2N})$. \Box

4 Convergence of $\Lambda_{\varepsilon}(z)^{-1}$

The goal of this section is to show that the limit $\lim_{\varepsilon \to 0} \Lambda_{\varepsilon}(z)^{-1}$ exists for all large enough $z > 0$, provided that, for all pairs σ , $V_{\sigma} \in L^1 \cap L^2(\mathbb{R}^2)$ and $\int |r|^{2s} |V_{\sigma}(r)| dr < \infty$ for some $s > 0$. As explained at the end of Section 2, this proof is based on a decomposition of $\Lambda_{\varepsilon}(z)$ into diagonal and off-diagonal parts that we now define.

Recall from (2.10) that, for given $\varepsilon, z > 0$, the operator matrix $\Lambda_{\varepsilon}(z) \in \mathscr{L}(\tilde{\mathfrak{X}})$ has the components

$$
\Lambda_{\varepsilon}(z)_{\sigma\nu} = g_{\varepsilon,\sigma}^{-1} \delta_{\sigma,\nu} - \phi_{\varepsilon}(z)_{\sigma\nu}, \qquad \sigma, \nu \in \mathcal{I},
$$

where

$$
\phi_{\varepsilon}(z)_{\sigma\nu}:=B_{\varepsilon,\sigma}R_0(z)A_{\varepsilon,\nu}^*
$$

defines a bounded operator $\phi_{\varepsilon}(z)_{\sigma\nu} \in \mathscr{L}(\tilde{\mathfrak{X}}_{\nu}, \tilde{\mathfrak{X}}_{\sigma})$ for all pairs $\sigma, \nu \in \mathcal{I}$. Using the definitions (2.7) and (2.6) for $B_{\varepsilon,\sigma}$ and $A_{\varepsilon,\nu}$, respectively, leads to the explicit formula

$$
\phi_{\varepsilon}(z)_{\sigma\nu} = \varepsilon^{-2} \left(u_{\sigma} \otimes 1 \right) U_{\varepsilon} \mathcal{K}_{\sigma} R_0(z) \mathcal{K}_{\nu}^* U_{\varepsilon}^*(v_{\nu} \otimes 1). \tag{4.1}
$$

We shall see that, for fixed $z > 0$, the off-diagonal operators $\phi_{\varepsilon}(z)_{\sigma\nu}$ ($\sigma \neq \nu$) are uniformly bounded in $\varepsilon > 0$, whereas the specific behavior of the coupling constant $g_{\varepsilon,\sigma}$ (see (1.3)) cancels the singular part in the diagonal operators $\phi_{\varepsilon}(z)_{\sigma\sigma}$ if and only if $a_{\sigma} = \int V_{\sigma}(r) dr/(2\pi) > 0$. Therefore, we write the operator matrix $\Lambda_{\varepsilon}(z)$ in the form

$$
\Lambda_{\varepsilon}(z) = \Lambda_{\varepsilon}(z)_{\text{diag}} + \Lambda_{\varepsilon}(z)_{\text{off}},\tag{4.2}
$$

where the diagonal part $\Lambda_{\varepsilon}(z)_{\text{diag}}$ and the off-diagonal part $\Lambda_{\varepsilon}(z)_{\text{off}}$ are given by

$$
\left(\Lambda_{\varepsilon}(z)_{\text{diag}}\right)_{\sigma\nu} = \left(g_{\varepsilon,\sigma}^{-1} - \phi_{\varepsilon}(z)_{\sigma\sigma}\right)\delta_{\sigma\nu}, \qquad \sigma,\nu \in \mathcal{I}, \tag{4.3}
$$

and

$$
(\Lambda_{\varepsilon}(z)_{\text{off}})_{\sigma\nu} = \phi_{\varepsilon}(z)_{\sigma\nu} (\delta_{\sigma\nu} - 1), \qquad \sigma, \nu \in \mathcal{I}, \qquad (4.4)
$$

respectively. Section 4.1 is devoted to $\Lambda_{\varepsilon}(z)_{\text{diag}}$ and Section 4.2 is devoted to $\Lambda_{\varepsilon}(z)_{\text{off}}$.

4.1 The limit of $(\Lambda_{\varepsilon}(z)_{\text{diag}})^{-1}$

We first show that $\Lambda_{\varepsilon}(z)_{\text{diag}}$ is invertible for small enough $\varepsilon > 0$ and large enough $z > 0$ and then we compute the limit $\lim_{\varepsilon \to 0} (\Lambda_{\varepsilon}(z)_{\text{diag}})^{-1}$. This can be done for each component $\Lambda_{\varepsilon}(z)_{\sigma\sigma} = g_{\varepsilon,\sigma}^{-1} - \phi_{\varepsilon}(z)_{\sigma\sigma}$ separately and without loss of generality we may choose $\sigma = (1,2)$. In the following, $v := v_{(1,2)}$, $\mathscr{K} := \mathscr{K}_{(1,2)}$, $\mu := \mu_{(1,2)}$, $\phi_{\varepsilon}(z) := \phi_{\varepsilon}(z)_{(1,2)(1,2)}$, etc.

We begin by computing the kernel of $\phi_{\varepsilon}(z)$. To this end, we note that

$$
\mathcal{K}R_0(z) = (\widetilde{H}_0 + z)^{-1}\mathcal{K},\tag{4.5}
$$

where H_0 is the free Hamiltonian H_0 expressed in the relative and center of mass coordinates of the pair $(1, 2)$ (cf. (2.2)), i.e.

$$
\widetilde{H}_0 = -\frac{\Delta_r}{\mu} - \frac{\Delta_R}{m_1 + m_2} + \sum_{i=3}^N \left(-\frac{\Delta_{x_i}}{m_i} \right). \tag{4.6}
$$

Starting from the explicit formula (4.1), the identity (4.5) and $\mathscr{K} \mathscr{K}^* = 1$ imply that

$$
\phi_{\varepsilon}(z) = \varepsilon^{-2} \left(u \otimes 1 \right) U_{\varepsilon} (\widetilde{H}_0 + z)^{-1} U_{\varepsilon}^*(v \otimes 1).
$$

Hence, after a Fourier transform in $(R, x_3, ..., x_N)$, $\phi_{\varepsilon}(z)$ acts pointwise in the conjugate variable $P = (P, p_3, ..., p_N)$ by the operator

$$
\phi_{\varepsilon}(z,\underline{P}) = \mu \varepsilon^{-2} u U_{\varepsilon}(-\Delta_r + \mu(z+Q))^{-1} U_{\varepsilon}^* v
$$

= $\mu u (-\Delta_r + \varepsilon^2 \mu(z+Q))^{-1} v \in \mathcal{L}(L^2(\mathbb{R}^2)),$ (4.7)

where

$$
Q := \frac{P^2}{m_1 + m_2} + \sum_{i=3}^{N} \frac{p_i^2}{m_i},\tag{4.8}
$$

and the scaling property of the Laplace operator with respect to the unitary scaling U_{ε} has been used in (4.7). As the resolvent in (4.7) is divergent in the limit $\varepsilon \downarrow 0$, the first step in the analysis of $\lim_{\varepsilon \to 0} (\Lambda_{\varepsilon}(z)_{\text{diag}})^{-1}$ is to study the asymptotic behavior of $\phi_{\varepsilon}(z, \underline{P})$ for small $\varepsilon > 0$.

A similar but less specific version of the following lemma can be found in [23, Proposition 3.2].

Lemma 4.1. Let $V \in L^1 \cap L^2(\mathbb{R}^2)$ satisfy $\int |r|^{2s} |V(r)| dr < \infty$ for some $s \in (0, 2)$. Then there *exists a constant* $C = C(s, V) > 0$ *such that*

$$
\forall \alpha > 0: \qquad \left\| u \left(-\Delta + \alpha \right)^{-1} v + \left(2\pi \right)^{-1} \left[\left(\ln \left(\sqrt{\alpha}/2 \right) + \gamma \right) |u\rangle \langle v| + L \right] \right\|_{\text{HS}} \leq C \alpha^{s/2},\tag{4.9}
$$

$$
\forall \alpha \ge 1: \qquad \left\| u \left(-\Delta + \alpha \right)^{-1} v \right\|_{\text{HS}} \le C, \tag{4.10}
$$

where $\|\cdot\|_{\text{HS}}$ *denotes the Hilbert-Schmidt norm in* $L^2(\mathbb{R}^2)$, γ *is the Euler–Mascheroni constant* and *L* is a Hilbert-Schmidt operator in $L^2(\mathbb{R}^2)$ that is defined in terms of the kernel

$$
u(r) \ln(|r - r'|) v(r'), \qquad r \neq r'.
$$

Proof. First, we observe that $u(-\Delta + \alpha)^{-1}v$ is associated with the integral kernel

$$
u(r) G_{\alpha}(r - r') v(r'), \qquad (4.11)
$$

where $G_{\alpha} = G_{\alpha}^2$ denotes the Green's function of $-\Delta + \alpha : H^2(\mathbb{R}^2) \to L^2(\mathbb{R}^2)$ (we refer to the appendix for details about G_{α}). By [2, Chapter I.5], we have the explicit description

$$
G_{\alpha}(x) = G_1(\sqrt{\alpha}x) = \frac{i}{4}H_0^{(1)}(i\sqrt{\alpha}|x|), \qquad 0 \neq x \in \mathbb{R}^2, \qquad (4.12)
$$

where $H_0^{(1)}$ denotes the Hankel function of first kind and order zero. It is known (see e.g. [1, Chapter 9.1]) that

$$
iH_0^{(1)}(ir) = -2\pi^{-1} \left(\ln \left(r/2 \right) + \gamma \right) + O(r^2 |\ln(r)|), \qquad (r \downarrow 0). \tag{4.13}
$$

As G_1 is smooth in $\mathbb{R}^2 \setminus \{0\}$ and exponentially decaying as $|x| \to \infty$ (see Lemma A.2), we conclude from (4.12) and (4.13) that there is some constant $\lambda = \lambda(s) > 0$ such that

$$
\left|G_1(x) + (2\pi)^{-1} \left(\ln(|x|/2) + \gamma\right)\right| \le \lambda |x|^s, \qquad x \in \mathbb{R}^2 \setminus \{0\}.
$$

Using again (4.12), this implies that

$$
\left| G_{\alpha}(x) + (2\pi)^{-1} \left(\ln \left(\sqrt{\alpha} |x|/2 \right) + \gamma \right) \right| \leq \lambda \alpha^{s/2} |x|^s
$$

uniformly in $\alpha > 0$ and $x \in \mathbb{R}^2 \setminus \{0\}$. Therefore, (4.9) follows from

$$
\|u(-\Delta + \alpha)^{-1} v + (2\pi)^{-1} \left[(\ln(\sqrt{\alpha}/2) + \gamma) |u\rangle\langle v| + L \right] \|_{\text{HS}}^2
$$

= $\int \text{d}r \, \text{d}r' |u(r)|^2 |G_{\alpha}(r - r') + (2\pi)^{-1} \left[\ln(\sqrt{\alpha}|r - r'|/2) + \gamma \right]|^2 |v(r')|^2$
 $\leq \lambda^2 \alpha^s \int \text{d}r \, \text{d}r' |V(r)| |r - r'|^{2s} |V(r')|$
 $\leq 2^{2s+1} \lambda^2 \alpha^s \|V\|_{L^1} \int |r|^{2s} |V(r)| \, \text{d}r,$

where the elementary inequality $(a + b)^{2s} \leq 2^{2s}(a^{2s} + b^{2s})$ $(a, b > 0)$ was used for the last inequality. To show that *L* indeed defines a Hilbert-Schmidt operator, we note that $||L||_{\text{HS}}^2$ = $I_1 + I_2$, where

$$
I_1 = \int_{|r| \le 1} dr \ln(|r|)^2 \int dr' \, |V(r+r')||V(r')| \le 2\pi ||V||^2 \int_0^1 x \ln(x)^2 dx < \infty
$$

by the Cauchy-Schwarz inequality and

$$
I_2 = \int \mathrm{d}r \, \ln(|r|)^2 \int \mathrm{d}r' \, |V(r+r')||V(r')|
$$

\n
$$
\leq \sup_{x \geq 1} \left(x^{-2s} \ln(x)^2 \right) \int \mathrm{d}r \, \mathrm{d}r' \, |r|^{2s} |V(r+r')||V(r')|
$$

\n
$$
= \sup_{x \geq 1} \left(x^{-2s} \ln(x)^2 \right) \int \mathrm{d}r \, \mathrm{d}r' \, |r-r'|^{2s} |V(r)||V(r')|
$$

\n
$$
\leq 2^{2s+1} \sup_{x \geq 1} \left(x^{-2s} \ln(x)^2 \right) ||V||_{L^1} \int |r|^{2s} |V(r)| \, \mathrm{d}r < \infty.
$$

It remains to prove (4.10). To this end, we note that for fixed $x \neq 0$ the function $\alpha \to G_{\alpha}(x)$ is strictly monotonically decreasing in $\alpha > 0$ (cf. Lemma A.1 (v)). Therefore, the Cauchy-Schwarz inequality yields for $\alpha \geq 1$ the uniform estimate

$$
\|u(-\Delta + \alpha)^{-1}v\|_{\text{HS}}^2 \le \int \mathrm{d}r \,\mathrm{d}r' \, |V(r)| \, |G_1(r - r')|^2 \, |V(r')|
$$

\$\leq \|V\|^2 \|G_1\|^2 < \infty\$,

where we used that $G_1 \in L^2(\mathbb{R}^2)$ (cf. Lemma A.1 (iv)).

In the next lemma we show that $g_{\varepsilon}^{-1} - \phi_{\varepsilon}(z)$ is invertible for all large enough $z > 0$ and small enough $\varepsilon > 0$. It is here, where the asymptotics (1.3) of g_{ε} , i.e.

$$
g_{\varepsilon}^{-1} = -\mu(a\ln(\varepsilon) + b) + o(1), \qquad (\varepsilon \downarrow 0) \tag{4.14}
$$

for some $a > 0$, will play a crucial role.

Lemma 4.2. *Let the hypotheses of Lemma 4.1 be satisfied. Then there exist* ε_0 , $z_0 > 0$ *such that* $g_{\varepsilon}^{-1} - \phi_{\varepsilon}(z, \underline{P})$ *is invertible for all* \underline{P} *,* $\varepsilon \in (0, \varepsilon_0)$ *and* $z > z_0$ *. Moreover,*

$$
\left\| \left(g_{\varepsilon}^{-1} - \phi_{\varepsilon}(z, \underline{P}) \right)^{-1} \right\| \leq \widetilde{C} \begin{cases} |\ln(\varepsilon)|^{-1} & \text{if } f \ V(r) \, \mathrm{d}r/(2\pi) < a \\ \max \{ |\ln(\varepsilon)|^{-1}, \ln(\mu(z+Q))^{-1} \} & \text{if } f \ V(r) \, \mathrm{d}r/(2\pi) = a, \end{cases} \tag{4.15}
$$

where the constant $\widetilde{C} > 0$ *is independent of* \underline{P} *,* $z > z_0$ *and* $\varepsilon \in (0, \varepsilon_0)$ *.*

 \Box

Proof. Let $C > 0$ denote the constant from Lemma 4.1. By (4.14), we find $\varepsilon_0 \in (0,1)$ such that $|g_{\varepsilon}\mu C| \leq 1/2$ and $|g_{\varepsilon}| \leq \lambda |\ln(\varepsilon)|^{-1}$ for all $\varepsilon \in (0, \varepsilon_0)$, where the constant $\lambda > 0$ is independent of $\varepsilon \in (0, \varepsilon_0)$.

If $\alpha_{\varepsilon} := \varepsilon^2 \mu(z + Q) \ge 1$, then it follows from (4.7) and (4.10) that

$$
||g_{\varepsilon}\phi_{\varepsilon}(z,\underline{P})|| \leq |g_{\varepsilon}|\mu||u(-\Delta + \alpha_{\varepsilon})^{-1}v|| \leq |g_{\varepsilon}\mu C| \leq \frac{1}{2}.
$$

Hence, $g_{\varepsilon}^{-1} - \phi_{\varepsilon}(z, \underline{P})$ is invertible and

$$
\left\| \left(g_{\varepsilon}^{-1} - \phi_{\varepsilon}(z, \underline{P}) \right)^{-1} \right\| = \left\| g_{\varepsilon} \left(1 - g_{\varepsilon} \phi_{\varepsilon}(z, \underline{P}) \right)^{-1} \right\| \leq 2\lambda |\ln(\varepsilon)|^{-1}
$$

for all $\varepsilon \in (0, \varepsilon_0)$. This establishes (4.15) in the case of $\alpha_{\varepsilon} \geq 1$.

In all the following, $\alpha_{\varepsilon} = \varepsilon^2 \mu(z + Q) < 1$. Then, it follows from (4.7) and (4.9) that

$$
\phi_{\varepsilon}(z,\underline{P}) = -\frac{\mu}{4\pi} \ln(\alpha_{\varepsilon}) \left| u \right\rangle \left\langle v \right| + O(1),\tag{4.16}
$$

where the norm of the operator $O(1)$ is uniformly bounded in P , $z > 0$ and $\varepsilon \in (0, \varepsilon_0)$ as long as α_{ε} < 1. To determine the inverse of $g_{\varepsilon}^{-1} - \phi_{\varepsilon}(z, \underline{P})$, we apply the identity (cf. [2, Chapter I.1, Eq. (1.3.47)])

$$
(1 + R + \beta \left| \tilde{\eta} \right\rangle \left\langle \eta \right|)^{-1} = (1 + R)^{-1} - \frac{(1 + R)^{-1} \left| \tilde{\eta} \right\rangle \left\langle \eta \right| (1 + R)^{-1}}{\beta^{-1} + \left\langle \eta \right| (1 + R)^{-1} \tilde{\eta} \rangle},\tag{4.17}
$$

which is valid under the assumption that $\beta \in \mathbb{C}$, $\tilde{\eta}, \eta \in \mathcal{H}$, R , $(1 + R)^{-1} \in \mathscr{L}(\mathcal{H})$ and β^{-1} + $\langle \eta | (1+R)^{-1} \tilde{\eta} \rangle \neq 0$ in some separable Hilbert space H. This identity (with $R = 0$) and $|g_{\varepsilon}| = O(|\ln(\varepsilon)|^{-1})$ yield

$$
\left(g_{\varepsilon}^{-1} + \frac{\mu}{4\pi} \ln(\alpha_{\varepsilon}) \ket{u} \bra{v}\right)^{-1} = g_{\varepsilon} \left(1 + \frac{g_{\varepsilon} \mu}{4\pi} \ln(\alpha_{\varepsilon}) \ket{u} \bra{v}\right)^{-1} = g_{\varepsilon}^{2} f(z, \underline{P}, \varepsilon)^{-1} \ket{u} \bra{v} + O(\ln(\varepsilon))^{-1},
$$
(4.18)

provided that

$$
f(z, \underline{P}, \varepsilon) := -\frac{4\pi}{\mu \ln(\alpha_{\varepsilon})} - g_{\varepsilon} \int V(r) dr \neq 0.
$$

To derive a lower bound on $f(z, P, \varepsilon)$, we consider the two cases $\int V(r) dr/(2\pi) < a$ and $\int V(r) dr/(2\pi) = a$ separately. In both cases we assume that $z > z_0$ and $\varepsilon \in (0, \varepsilon_0)$ for some $z_0 \geq \mu^{-1}$ and $\varepsilon_0 \in (0,1)$, which we will fix below. This implies that $\mu(z+Q) \geq 1$ and hence

$$
0 < -\ln(\alpha_{\varepsilon}) \le 2|\ln(\varepsilon)|. \tag{4.19}
$$

If $\int V(r) dr/(2\pi) < a$, then we use (4.14) and (4.19) to derive the lower bound

$$
f(z, \underline{P}, \varepsilon) \ge (\mu |\ln(\varepsilon)|)^{-1} \left(2\pi - \frac{1}{a} \int V(r) dr\right) + O(\ln(\varepsilon)^{-2}) \qquad (\varepsilon \downarrow 0). \tag{4.20}
$$

Hence, there exists $\varepsilon_0 \in (0,1)$ such that $f(z, \underline{P}, \varepsilon) > 0$ for all $\underline{P}, z > z_0 \ge \mu^{-1}$ and $\varepsilon \in (0,\varepsilon_0)$. Moreover, inserting (4.20) in (4.18) and using $g_{\varepsilon}^2 = O(\ln(\varepsilon)^{-2})$ shows that

$$
\left\| \left(g_{\varepsilon}^{-1} + \frac{\mu}{4\pi} \ln(\alpha_{\varepsilon}) |u\rangle \langle v| \right)^{-1} \right\| = O(|\ln(\varepsilon)|^{-1}) \qquad (\varepsilon \downarrow 0).
$$

In view of (4.16), this proves (4.15) for $\int V(r) dr/(2\pi) < a$ and $\alpha_{\varepsilon} < 1$.

If $\int V(r) dr/(2\pi) = a$, then we again use (4.14) and (4.19) to derive a more refined version of (4.20):

$$
|\ln(\varepsilon)|^2 |f(z, \underline{P}, \varepsilon)| = \frac{|\ln(\varepsilon)|^2}{|\ln(\alpha_{\varepsilon})|} \left| 4\pi \mu^{-1} + g_{\varepsilon} \ln(\alpha_{\varepsilon}) \int V(r) dr \right|
$$

\n
$$
\geq \frac{|\ln(\varepsilon)|}{2\mu} \left| 4\pi + g_{\varepsilon} \mu \ln(\alpha_{\varepsilon}) \int V(r) dr \right|
$$

\n
$$
= \frac{|\ln(\varepsilon)|}{2\mu} \left| 4\pi - \frac{\int V(r) dr}{a \ln(\varepsilon)} \left(2 \ln(\varepsilon) + \ln(\mu(z+Q)) \right) + O\left(\frac{|\ln(\alpha_{\varepsilon})|}{\ln(\varepsilon)^2} \right) \right|
$$

\n
$$
= \pi \mu^{-1} \ln(\mu(z+Q)) + O(1),
$$

where the remainder $O(1)$ is uniformly bounded in \underline{P} , $z > \mu^{-1}$ and $\varepsilon \in (0, \varepsilon_0)$. By choosing $z_0 \geq \mu^{-1}$ large enough, we conclude that $|f(z, \underline{P}, \varepsilon)| \geq \pi \ln(\mu(z+Q))/(2\mu \ln(\varepsilon)^2) > 0$ for all \underline{P} , $z > z_0$ and $\varepsilon \in (0, \varepsilon_0)$. With the help of this bound, the desired estimate for the operator norm of $(g_{\varepsilon}^{-1} - \phi_{\varepsilon}(z, \underline{P}))^{-1}$ can be obtained similarly to the case $\int V(r) dr/(2\pi) < a$. \Box

An immediate consequence of Lemma 4.2 is that $g_{\varepsilon}^{-1} - \phi_{\varepsilon}(z)$ is invertible for all small enough $\varepsilon > 0$ and large enough $z > 0$. Yet, we shall see that the limit $\varepsilon \to 0$ of the inverse operator essentially depends on the leading order of the coupling constant g_{ε} or, more precisely, on *a*. If $\int V(r) dr/(2\pi) < a$, then it follows from (4.15) that $\lim_{\varepsilon \to 0} (g_{\varepsilon}^{-1} - \phi_{\varepsilon}(z))^{-1}$ vanishes for $z > z_0 > 0$, while for $\int V(r) dr/(2\pi) = a$ this limit will turn out to be non-trivial. To prove this, we first recall that (4.7) and (4.11) imply that for fixed P the operator $\phi_{\varepsilon}(z, P)$ is associated with the kernel

$$
\mu u(r) G_{\varepsilon^2 \mu(z+Q)}(r-r') v(r').
$$

In the notation of [2, Chapter I.5] for the one-particle case, $(-g_{\varepsilon}) \cdot \phi_{\varepsilon}(z, P)$ agrees with the operator $B_{\varepsilon}(k)$ for $k^2 := -\mu(z+Q) < 0$, $\lambda_1 := 1/a$ and $\lambda_2 := -b/a^2$. In view of [2, Chapter I.5, Eq. (5.61) ^{*}, we thus expect that

$$
\lim_{\varepsilon \to 0} \left(g_{\varepsilon}^{-1} - \phi_{\varepsilon}(z) \right)^{-1} = \frac{|u\rangle \langle v|}{\langle u | v \rangle^2} \otimes D(z)^{-1},\tag{4.21}
$$

where $D(z)$ is an (unbounded) operator in \mathfrak{X}_{σ} , which, after passing to Fourier space, acts as the multiplication operator

$$
D(z, \underline{P}) := \frac{\mu}{2\pi} \left(\ln \left(\frac{\sqrt{\mu(z+Q)}}{2} \right) + \gamma + 2\pi\alpha \right), \tag{4.22}
$$

$$
\alpha := -\frac{b}{\langle u | v \rangle} + \frac{\langle v | Lu \rangle}{2\pi \langle u | v \rangle^2}.
$$
\n(4.23)

Here, *γ* denotes the Euler–Mascheroni constant and *L* is the Hilbert-Schmidt operator defined in Lemma 4.1. We now turn to the proof of (4.21):

Proposition 4.3. Let $\int V(r) dr/(2\pi) = a$ and let the hypotheses of Lemma 4.1 be satisfied. *Then* $D(z)$ *is invertible and* (4.21) *holds true for all large enough* $z > 0$ *.*

Proof. Clearly, $D(z)$ is invertible if ln $(\sqrt{\mu z}/2) + \gamma + 2\pi\alpha > 0$, i.e. $z > z_1 := 4\mu^{-1} \exp(-4\pi\alpha - 2\gamma)$. Let ε_0 , $z_0 > 0$ be chosen as in Lemma 4.2 and, for the rest of this proof, let $z > \max(z_0, z_1)$ be fixed. Then (4.21) is equivalent to the statement that

$$
\lim_{\varepsilon \to 0} \sup_{\underline{P}} \left\| \left(g_{\varepsilon}^{-1} - \phi_{\varepsilon}(z, \underline{P}) \right)^{-1} - D(z, \underline{P})^{-1} \frac{|u\rangle \langle v|}{\langle u | v \rangle^2} \right\| = 0 \tag{4.24}
$$

[∗]There is a typo in [2, Chapter I.5, Eq. (5.61)]: The term in curly braces is to be inverted.

with $\|\cdot\|$ denoting the operator norm in $\mathscr{L}(L^2(\mathbb{R}^2))$.

The idea of the following proof is that both operators in the difference vanish as $|P| \to \infty$, while for $|P| \leq K$ the arguments from the one-particle case still work. To make this more explicit, we fix $\delta > 0$. Then it follows from Lemma 4.2 and (4.22) that

$$
\left\| \left(g_{\varepsilon}^{-1} - \phi_{\varepsilon}(z, \underline{P}) \right)^{-1} \right\| + \left\| D(z, \underline{P})^{-1} \frac{|u\rangle \langle v|}{\langle u | v \rangle^2} \right\| < \delta,
$$

provided that $\varepsilon > 0$ is sufficiently small and $|P|$, and hence *Q*, are sufficiently large. To prove (4.24), it therefore suffices to show that for any $K > 0$ there exists $\varepsilon_K > 0$ such that

$$
\sup_{\underline{P}: Q \le K} \left\| \left(g_{\varepsilon}^{-1} - \phi_{\varepsilon}(z, \underline{P}) \right)^{-1} - D(z, \underline{P})^{-1} \frac{|u\rangle \langle v|}{\langle u | v \rangle^2} \right\| < \delta \tag{4.25}
$$

holds for all $\varepsilon \in (0, \varepsilon_K)$.

Let $K > 0$ be fixed. Then (4.9) implies that, for some constant $C = C(s, V) > 0$,

$$
\left\| \phi_{\varepsilon}(z,\underline{P}) + \frac{\mu}{2\pi} \left[\left(\ln \left(\frac{\varepsilon}{2} \sqrt{\mu(z+Q)} \right) + \gamma \right) |u\rangle \langle v| + L \right] \right\|_{\text{HS}} \leq C \mu^{1+s/2} (z+K)^{s/2} \varepsilon^s
$$

uniformly in \underline{P} with $Q = Q(\underline{P}) \leq K$. Combining this with the asymptotic behavior of g_{ε} in the limit $\varepsilon \to 0$ (see (4.14)), we see that

$$
(-g_{\varepsilon}) \cdot \phi_{\varepsilon}(z, \underline{P})
$$

= $-\frac{1}{2\pi a} |u\rangle\langle v| - \frac{1}{2\pi a \ln(\varepsilon)} \left[\left\{ \ln \left(\frac{\sqrt{\mu(z+Q)}}{2} \right) + \gamma - \frac{b}{a} \right\} |u\rangle\langle v| + L \right] + o\left(\frac{1}{|\ln(\varepsilon)|} \right)$ (4.26)

is valid in Hilbert-Schmidt norm uniformly in P with $Q = Q(P) \leq K$. As this coincides with the expansion of $B_{\varepsilon}(k)$ from the one-particle case (cf. [2, Chapter I.5, Eq. (5.56)]), the proof of (4.25) now follows the line of arguments from that case. For the convenience of the reader, we spell out the details in the following.

To start with, we derive from (4.26) the equation

$$
\left(g_{\varepsilon}^{-1} - \phi_{\varepsilon}(z, \underline{P})\right)^{-1} = g_{\varepsilon}\left(1 + \beta(Q) \left|u\right\rangle \langle v| + R_{\varepsilon}\right)^{-1},\tag{4.27}
$$

where

$$
\beta(Q) := -\frac{1}{2\pi a} \left(1 + \frac{\tilde{\beta}(Q)}{\ln(\varepsilon)} \right),
$$

$$
\tilde{\beta}(Q) := \ln \left(\frac{\sqrt{\mu(z+Q)}}{2} \right) + \gamma - \frac{b}{a},
$$

$$
R_{\varepsilon} := -\frac{1}{2\pi a \ln(\varepsilon)} L + o\left(|\ln(\varepsilon)|^{-1} \right).
$$

The expansion of R_{ε} holds uniformly in *P* with $Q = Q(\underline{P}) \leq K$. To obtain an expression for the inverse on the right side of (4.27) , we now apply the identity (4.17) . This yields that

$$
\left(1+\beta(Q)|u\rangle\langle v|+R_{\varepsilon}\right)^{-1} = \left(1+R_{\varepsilon}\right)^{-1} - \frac{\left(1+R_{\varepsilon}\right)^{-1}|u\rangle\langle v|\left(1+R_{\varepsilon}\right)^{-1}}{\beta(Q)^{-1}+\langle v|\left(1+R_{\varepsilon}\right)^{-1}u\rangle} \tag{4.28}
$$

as operators in $\mathscr{L}(L^2(\mathbb{R}^2))$. By the definitions of $\beta(Q)$ and R_{ε} , and by $2\pi a = \langle u | v \rangle$,

$$
\beta(Q)^{-1} = -\langle u \, | \, v \rangle \left(1 - \frac{\tilde{\beta}(Q)}{\ln(\varepsilon)} \right) + O\left(\ln(\varepsilon)^{-2} \right), \qquad Q \le K, \tag{4.29}
$$

and

$$
\left\langle v \left| (1 + R_{\varepsilon})^{-1} u \right\rangle = \left\langle u \left| v \right\rangle + \frac{\left\langle v \left| Lu \right\rangle}{2\pi a \ln(\varepsilon)} + o \left(\left| \ln(\varepsilon) \right|^{-1} \right) \right\rangle, \tag{4.30}
$$

where $||R_{\varepsilon}|| = O(|\ln(\varepsilon)|^{-1})$ was used. Note that the sum of (4.29) and (4.30) is of order $1/|\ln(\varepsilon)|$ because $\langle u | v \rangle$ cancels. It follows that the second term in (4.28) is of order $|\ln(\varepsilon)|$. Hence we may ignore the first summand, $(1+R_\varepsilon)^{-1}$, in Equation (4.28) and we may replace the numerator in the second summand by $|u\rangle\langle v|$. It follows that, uniformly in *P* with $Q = Q(P) \le K$,

$$
\left(1+\beta(Q) \left|u\right\rangle\left\langle v\right| + R_{\varepsilon}\right)^{-1} = -\frac{\ln(\varepsilon)}{\langle u \left|v\right\rangle} \left[\ln\left(\frac{\sqrt{\mu(z+Q)}}{2}\right) + \gamma + 2\pi\alpha\right]^{-1} \left|u\right\rangle\left\langle v\right| + o\left(\left|\ln(\varepsilon)\right|\right)
$$

with *α* defined by (4.23). Finally, it follows from (4.27) and from the asymptotics of g_{ε} with $a = \langle u | v \rangle / (2\pi)$ that (4.25) with $D(z, P)$ from (4.22) holds true for all sufficiently small $\varepsilon > 0$. This proves the lemma. \Box

For later convenience, we now collect the conclusions of this section. To this end, we need to reinstall the index σ , which we dropped at the beginning of the section. Let $\sigma \in \mathcal{I}$ be a fixed pair. If $V_{\sigma} \in L^1 \cap L^2(\mathbb{R}^2)$ and $\int |r|^{2s} |V_{\sigma}(r)| dr < \infty$ for some $s \in (0, 2)$, then the analogues of Lemma 4.2 and Proposition 4.3 show that $\Lambda_{\varepsilon}(z)_{\sigma\sigma} = g_{\varepsilon,\sigma}^{-1} - \phi_{\varepsilon}(z)_{\sigma\sigma}$ is invertible for all sufficiently small $\varepsilon > 0$ and sufficiently large $z > 0$ and

$$
\lim_{\varepsilon \to 0} (\Lambda_{\varepsilon}(z)_{\sigma\sigma})^{-1} = \begin{cases} 0 & \text{if } \int V_{\sigma}(r) \, \mathrm{d}r/(2\pi) < a_{\sigma} \\ \frac{|u_{\sigma}\rangle \langle v_{\sigma}|}{\langle u_{\sigma} | v_{\sigma}\rangle^{2}} \otimes (\Theta(z)_{\sigma\sigma})^{-1} & \text{if } \int V_{\sigma}(r) \, \mathrm{d}r/(2\pi) = a_{\sigma}, \end{cases}
$$
(4.31)

where $\Theta(z)_{\sigma\sigma}$ is an invertible operator in \mathfrak{X}_{σ} . For $\sigma = (1,2)$, $\Theta(z)_{\sigma\sigma} = D(z)$ with $D(z)$ defined by (4.22) and (4.23). For general pairs $\sigma = (i, j)$, the operator $\Theta(z)_{\sigma\sigma}$ acts pointwise in $P_{\sigma} = (P, p_1, \ldots, \hat{p}_i \ldots \hat{p}_j \ldots, p_N)$ by multiplication with

$$
\Theta(z, \underline{P}_{\sigma})_{\sigma\sigma} = \frac{\mu_{\sigma}}{4\pi} \left[\ln \left(z + \frac{P^2}{m_i + m_j} + \sum_{\substack{n=1 \ n \neq i,j}}^N \frac{p_n^2}{m_n} \right) + \frac{\beta_{\sigma}}{\pi} \right], \qquad \sigma = (i, j), \qquad (4.32)
$$

where

$$
\beta_{\sigma} = 2\pi \left(\ln(\sqrt{\mu_{\sigma}}/2) + \gamma + 2\pi \alpha_{\sigma} \right),\tag{4.33}
$$

$$
\alpha_{\sigma} = -\frac{b_{\sigma}}{\langle u_{\sigma} | v_{\sigma} \rangle} + \frac{\langle v_{\sigma} | Lu_{\sigma} \rangle}{2\pi \langle u_{\sigma} | v_{\sigma} \rangle^{2}}.
$$
\n(4.34)

4.2 Analysis of $\Lambda_{\epsilon}(z)_{\text{off}}$

If $N > 2$, then $\Lambda_{\varepsilon}(z)$ has an off-diagonal part $\Lambda_{\varepsilon}(z)_{\text{off}}$ defined by Eq. (4.4). We will see in this section that $\Lambda_{\varepsilon}(z)_{\text{off}}$ is uniformly bounded in $\varepsilon > 0$ and $z > 0$ and that, upon introducing a cutoff, it has a limit as $\varepsilon \downarrow 0$. These results will allow us in Section 4.3 to prove existence of $\lim_{\varepsilon \to 0} \Lambda_{\varepsilon}(z)^{-1}$ for sufficiently large $z > 0$. With this goal in mind, the results of this section are formulated for $z > 0$ only, although most of them still hold in a slightly modified way for general $z \in \rho(H_0)$.

4.2.1 Uniform boundedness of $\Lambda_{\varepsilon}(z)_{\text{off}}$ in $\varepsilon > 0$ and $z > 0$

Recall from Eqs. (4.4) and (4.1) that the non-vanishing components of $\Lambda_{\varepsilon}(z)_{\text{off}}$ are given by

$$
\Lambda_{\varepsilon}(z)_{\sigma\nu} = -\varepsilon^{-2} \left(u_{\sigma} \otimes 1 \right) U_{\varepsilon} \mathscr{K}_{\sigma} R_0(z) \mathscr{K}_{\nu}^* U_{\varepsilon}^*(v_{\nu} \otimes 1), \qquad \sigma, \nu \in \mathcal{I}, \sigma \neq \nu. \tag{4.35}
$$

We will prove in this section, among other things, that for $\sigma \neq \nu$ the norm $\|\Lambda_{\varepsilon}(z)_{\sigma\nu}\|$ is uniformly bounded in $\varepsilon > 0$ and $z > 0$, the main result being Proposition 4.4 below. The presence of the distinct coordinate transformations \mathcal{K}_{σ} and \mathcal{K}_{ν} in (4.35) makes the proof very technical and somewhat tedious. Since the tools used in this proof are not needed anymore in the sequel, it is possible and advisable to take Proposition 4.4 for granted and to skip the proof at first reading. With this said, we now start developing the tools for proving Proposition 4.4. To estimate the norm of $\Lambda_{\varepsilon}(z)_{\sigma\nu}$, we compute its integral kernel in terms of the Green's function $G_{z,m}$ of H_0 , where the vector $\underline{m} := (m_1, ..., m_N)$ collects the masses of the *N* particles. By a simple scaling argument,

$$
G_{z,\underline{m}}(x_1,...,x_N) = \left(\prod_{i=1}^N m_i\right) G_z^{2N}(\sqrt{m_1}x_1,...,\sqrt{m_N}x_N),\tag{4.36}
$$

where G_z^{2N} denotes the usual Green's function of $-\Delta + z : H^2(\mathbb{R}^{2N}) \to L^2(\mathbb{R}^{2N})$. As the choice of the pair σ is immaterial for our estimates, we can assume that $\sigma = (1, 2)$ and $\nu = (k, l) \neq (1, 2)$. Now, with U_{ε} , $\mathscr{K}_{(1,2)}$ and $\mathscr{K}_{(k,l)}^*$ defined by (2.5), (2.3) and (2.4), respectively, we find that (4.35) defines an integral operator and, for $\psi \in \widetilde{\mathfrak{X}}_{(k,l)},$

$$
\left(\Lambda_{\varepsilon}(z)_{(1,2)(k,l)}\,\psi\right)(r,R,x_3,...,x_N) \n= -\varepsilon^{-2}u_{(1,2)}(r)\int dx'_1\cdots dx'_N\,G_{z,\underline{m}}\left(R - \frac{\varepsilon m_2r}{m_1 + m_2} - x'_1, R + \frac{\varepsilon m_1r}{m_1 + m_2} - x'_2, x_3 - x'_3, ..., x_N - x'_N\right) \n\cdot v_{(k,l)}\left(\frac{x'_l - x'_k}{\varepsilon}\right)\,\psi\left(\frac{x'_l - x'_k}{\varepsilon}, \frac{m_kx'_k + m_lx'_l}{m_k + m_l}, x'_1, \dots, \widehat{x'_k}...\widehat{x'_l}...\,x'_N\right) \n= -u_{(1,2)}(r)\int dx'_1\cdots dx'_N\,dr'\,dR'\,G_{z,\underline{m}}\left(R - \varepsilon c_{21}r - x'_1, R + \varepsilon c_{12}r - x'_2, x_3 - x'_3, ..., x_N - x'_N\right) \n\cdot v_{(k,l)}\left(r'\right)\,\psi\left(r', R', x'_1, \dots, \widehat{x'_k}...\widehat{x'_l}...\,x'_N\right)\,\delta\left(x'_k - R' + \varepsilon c_{lk}r'\right)\delta\left(x'_l - R' - \varepsilon c_{kl}r'\right),\n\tag{4.37}
$$

where

$$
c_{ij} := \frac{m_i}{m_i + m_j}, \qquad i, j = 1, ..., N.
$$

The second equation of (4.37) was obtained by the substitution

$$
r' := \frac{x'_l - x'_k}{\varepsilon}, \qquad R' := \frac{m_k x'_k + m_l x'_l}{m_k + m_l},
$$

and two more integrations were introduced that are compensated by delta distributions. In the following, we distinguish two cases: In the first case $\sigma = (1, 2)$ and $\nu = (k, l)$ have one particle in common, so $k \in \{1,2\}$ and $l \geq 3$. In the second case σ and ν are composed of distinct particles, which means that $3 \leq k < l \leq N$.

If $\sigma = (1, 2)$ and $\nu = (1, l)$ with $l \geq 3$, then, after the evaluation of the delta distributions in x'_1 and x'_l , (4.37) shows that the operator $\Lambda_{\varepsilon}(z)_{\sigma\nu}$ simply acts by convolution in $(x_3, \ldots, \hat{x}_l, \ldots, x_N)$. Consequently, the explicit formula (4.36) for $G_{z,m}$ and Lemma A.1 (*vi*) imply that $\Lambda_{\varepsilon}(z)_{\sigma\nu}$ acts pointwise in the conjugate variable $\underline{p}_{(1,l)} := (p_3, \ldots, \widehat{p}_l, \ldots, p_N)$ by the operator $\Lambda_{\varepsilon}(z, \underline{p}_{(1,l)})_{\sigma\nu}$ that has the kernel

$$
-u_{\sigma}(r)\left(m_1m_2m_l G_{z+Q_{\nu}}^6(X_{\sigma\nu,\varepsilon})\right)v_{\nu}(r'), \qquad \sigma = (1,2), \nu = (1,l), l \ge 3,
$$
 (4.38)

where

$$
X_{(1,2)(1,l),\varepsilon} := \left(\begin{array}{c} \sqrt{m_1} (R - R' - \varepsilon (c_{21}r - c_{l1}r')) \\ \sqrt{m_2} (R - x'_2 + \varepsilon c_{12}r) \\ \sqrt{m_l} (x_l - R' - \varepsilon c_{1l}r') \end{array} \right) \in \mathbb{R}^6
$$

and

$$
Q_{(k,l)} := \sum_{\substack{n=3 \ n \neq k,l}}^N \frac{p_n^2}{m_n}, \qquad \nu = (k,l) \in \mathcal{I}.
$$

Similar considerations for $\sigma = (1, 2)$ and $\nu = (2, l)$ with $l \geq 3$ show that $\Lambda_{\varepsilon}(z)_{\sigma\nu}$ acts pointwise in $\underline{p}_{(2,l)} := (p_3, \ldots \widehat{p}_l \ldots, p_N)$ by the operator $\Lambda_{\varepsilon}(z, \underline{p}_{(2,l)})_{\sigma\nu}$ with kernel

$$
-u_{\sigma}(r)\left(m_1m_2m_l G_{z+Q_{\nu}}^6(X_{\sigma\nu,\varepsilon})\right)v_{\nu}(r'), \qquad \sigma = (1,2), \nu = (2,l), l \ge 3,
$$
 (4.39)

where

$$
X_{(1,2)(2,l),\varepsilon} := \left(\sqrt{\frac{m_1}(R - x_1' - \varepsilon c_{21}r)}{\sqrt{m_2}(R - R' + \varepsilon (c_{12}r + c_{l2}r'))} \right) \in \mathbb{R}^6.
$$

By inspection, the kernels (4.38) and (4.39) only differ by the permutations $x'_1 \leftrightarrow x'_2$, $m_1 \leftrightarrow m_2$, $v_{(1,l)} \leftrightarrow v_{(2,l)}$ and the reflection $r \to -r$, which will allow us to analyze them simultaneously.

So far, we have considered all operators that appear in the case of $N \leq 3$ particles. Let now $N > 3$, $\sigma = (1, 2)$ and $\nu = (k, l)$ with $3 \leq k < l \leq N$. Then, after the evaluation of the delta distributions in x'_k and x'_l , it follows from (4.37) that $\Lambda_{\varepsilon}(z)_{\sigma\nu}$ acts by convolution in $(x_3, \ldots, \widehat{x_k} \ldots \widehat{x_l} \ldots, x_N)$. Hence, the action of $\Lambda_{\varepsilon}(z)_{\sigma\nu}$ is pointwise in the conjugate variables $p_{(k,l)} := (p_3, \ldots \widehat{p_k} \ldots \widehat{p_l} \ldots p_N)$ by the operator $\Lambda_{\varepsilon}(z, \underline{p}_{(k,l)})_{\sigma\nu}$ with kernel

$$
-u_{\sigma}(r)\left(m_1m_2m_km_l\,G_{z+Q_{\nu}}^8(X_{\sigma\nu,\varepsilon})\right)v_{\nu}(r'), \qquad \sigma = (1,2), \nu = (k,l), 3 \le k < l \le N, \tag{4.40}
$$

where

$$
X_{(1,2)(k,l),\varepsilon} := \left(\begin{array}{c} \sqrt{m_1} (R - x_1' - \varepsilon c_{21}r) \\ \sqrt{m_2} (R - x_2' + \varepsilon c_{12}r) \\ \sqrt{m_k} (x_k - R' + \varepsilon c_{lk}r') \\ \sqrt{m_l} (x_l - R' - \varepsilon c_{kl}r') \end{array} \right) \in \mathbb{R}^8.
$$

Besides the bound for the norm of $\Lambda_{\varepsilon}(z)_{\sigma\nu}$, we shall also estimate the error caused by cutting off the potential outside some ball of radius $h > 0$ in the following proposition. This, in turn, will reduce the proof of our convergence result (see Proposition 4.8) to the case of compactly supported potentials. For any pair $\sigma \in \mathcal{I}$ and $h > 0$, let

$$
V_{\sigma}^{h}(x) := \begin{cases} V_{\sigma}(x) & \text{if } |x| \le h \\ 0 & \text{else} \end{cases}
$$

and let $\Lambda_{\varepsilon}^{h}(z)_{\sigma\nu}$ denote the operator $\Lambda_{\varepsilon}(z)_{\sigma\nu}$, where the potentials V_{σ} and V_{ν} are replaced by V_{σ}^{h} and V^h_ν , respectively. This means that the kernel of $\Lambda^h_\varepsilon(z)_{\sigma\nu}$ emerges from the kernel of $\Lambda_\varepsilon(z)_{\sigma\nu}$ by replacing u_{σ} and v_{ν} with $u_{\sigma}^h := \text{sgn}(V_{\sigma}) |V_{\sigma}^h|^{1/2}$ and $v_{\nu}^h := |V_{\nu}^h|^{1/2}$, respectively.

Proposition 4.4. For all pairs $\sigma = (i, j)$, $\nu = (k, l)$ with $\sigma \neq \nu$ the operator $\Lambda_{\varepsilon}(z)_{\sigma\nu}$ is uniformly *bounded in* $z > 0$ *and* $\varepsilon > 0$ *. Explicitly, it holds that*

$$
\|\Lambda_{\varepsilon}(z)_{\sigma\nu}\| \le C(\sigma,\nu) \|V_{\sigma}\|_{L^{1}}^{1/2} \|V_{\nu}\|_{L^{1}}^{1/2},\tag{4.41}
$$

where $C(\sigma,\nu) := (4\sqrt{2})^{-1} m_i m_j m_k m_l / \min(m_i, m_j, m_k, m_l)^3$. Furthermore, for any $h > 0$,

$$
\|\Lambda_{\varepsilon}(z)_{\sigma\nu} - \Lambda_{\varepsilon}^{h}(z)_{\sigma\nu}\| \le C(\sigma,\nu) \left(\|V_{\sigma}\|_{L^{1}} \|V_{\nu}\|_{L^{1}} - \|V_{\sigma}^{h}\|_{L^{1}} \|V_{\nu}^{h}\|_{L^{1}} \right)^{1/2}.
$$
 (4.42)

Proof. Without loss of generality, we may assume that $\sigma = (1, 2)$ and then we have to establish (4.41) and (4.42) for all pairs $\nu \neq (1,2)$. We first consider the case $\nu = (1,l)$ with $l \geq 3$.

The proofs of (4.41) and (4.42) are similar: It follows from (4.38) that in both cases we have to estimate the norm of an operator that, for fixed $\underline{p}_{(1,l)}$, is given by a kernel of the form

$$
m_1m_2m_l W(r,r') G_{z+Q}^6(X_\varepsilon),
$$

where $X_{\varepsilon} := X_{(1,2)(1,l),\varepsilon}$ and $Q := Q_{(1,l)}$ for short. Explicitly, we have $W(r,r') = u_{(1,2)}(r)v_{(1,l)}(r')$ in the case of (4.41) and $W(r,r') = u_{(1,2)}(r)v_{(1,l)}(r') - u_{(1,2)}^h(r)v_{(1,l)}^h(r')$ in the case of (4.42). Therefore, we only demonstrate the desired estimate in the case of (4.41).

For $\psi \in L^2(\mathbb{R}^2 \times \mathbb{R}^2 \times \mathbb{R}^2)$, the Cauchy-Schwarz inequality in the *r'*-integration yields

$$
(m_1 m_2 m_l)^{-2} || \Lambda_{\varepsilon}(z, \underline{p}_{(1,l)})_{(1,2)(1,l)} \psi ||^2
$$

= $\int \mathrm{d}r \, \mathrm{d}R \, \mathrm{d}x_l \left| \int \mathrm{d}r' \, \mathrm{d}R' \, \mathrm{d}x'_2 \, W(r, r') \, G_{z+Q}^6(X_{\varepsilon}) \, \psi(r', R', x'_2) \right|^2$
= $\int \mathrm{d}r \, \mathrm{d}R \, \mathrm{d}x_l \left| \int \mathrm{d}r' \, W(r, r') \int \mathrm{d}R' \, \mathrm{d}x'_2 \, G_{z+Q}^6(X_{\varepsilon}) \, \psi(r', R', x'_2) \right|^2$
 $\leq \int \mathrm{d}r \, \mathrm{d}R \, \mathrm{d}x_l \left\{ \int \mathrm{d}r' \, W(r, r')^2 \right\} \int \mathrm{d}r' \left| \int \mathrm{d}R' \, \mathrm{d}x'_2 \, G_{z+Q}^6(X_{\varepsilon}) \, \psi(r', R', x'_2) \right|^2$
 $\leq \left\{ \int \mathrm{d}r \, \mathrm{d}r' \, W(r, r')^2 \right\} \cdot \sup_{r \in \mathbb{R}^2} \left(\int \mathrm{d}r' \mathrm{d}R \, \mathrm{d}x_l \left| \int \mathrm{d}R' \, \mathrm{d}x'_2 \, G_{z+Q}^6(X_{\varepsilon}) \, \psi(r', R', x'_2) \right|^2 \right), \quad (4.43)$

where

$$
\int \mathrm{d}r \, \mathrm{d}r' \, W(r, r')^2 = \|V_{(1,2)}\|_{L^1} \|V_{(1,l)}\|_{L^1}.
$$
\n(4.44)

For a further estimate of (4.43), we fix $r \in \mathbb{R}^2$. Then triangle inequality and the sequence of substitutions $R' + \varepsilon (c_{21}r - c_{l1}r') \rightarrow R', x_l + \varepsilon c_{21}r - \varepsilon r' \rightarrow x_l, x_2' - \varepsilon c_{12}r \rightarrow x_2'$ in the first step and the monotonicity of the Green's function w.r.t. z and m_1, m_2, m_l (cf. Lemma A.1 (v)) in the second step yield that

$$
\int d\mathbf{r}' dR d\mathbf{x}_l \left| \int dR' d\mathbf{x}_2' G_{z+Q}^6(X_{\varepsilon}) \psi(\mathbf{r}', R', \mathbf{x}_2') \right|^2
$$
\n
$$
= \int d\mathbf{r}' dR d\mathbf{x}_l \left(\int dR' d\mathbf{x}_2' G_{z+Q}^6(\sqrt{m_1}(R - R'), \sqrt{m_2}(R - \mathbf{x}_2'), \sqrt{m_l}(x_l - R')) \widetilde{\psi}(\mathbf{r}', R', \mathbf{x}_2') \right)^2
$$
\n
$$
\leq \int d\mathbf{r}' \left\| F \widetilde{\psi}(\mathbf{r}', \cdot) \right\|^2, \tag{4.45}
$$

where $\widetilde{\psi} \in L^2(\mathbb{R}^2 \times \mathbb{R}^2 \times \mathbb{R}^2)$ is given by

$$
\widetilde{\psi}(r', R', x_2') := |\psi(r', R' - \varepsilon (c_{21}r - c_{l1}r'), x_2' + \varepsilon c_{12}r)|
$$

and $F: L^2(\mathbb{R}^2 \times \mathbb{R}^2, d(R, x_2)) \to L^2(\mathbb{R}^2 \times \mathbb{R}^2, d(R, x_l))$ is defined by the kernel

$$
G_z^6(\sqrt{m}(R - R'), \sqrt{m}(R - x'_2), \sqrt{m}(x_l - R'))
$$
, $m = \min(m_1, m_2, m_l)$.

By Lemma A.3, *F* is bounded and $||F|| \leq (4\sqrt{2}m^2)^{-1}$. Using this together with the fact that $\|\psi\| = \|\tilde{\psi}\|$, we obtain from (4.45) that

$$
\int dr' dR \, dx_l \left| \int dR' \, dx_2' \, G_{z+Q}^6 \left(X_\varepsilon \right) \, \psi(r', R', x_2') \right|^2 \leq (32 \min(m_1, m_2, m_l)^4)^{-1} \|\psi\|^2, \tag{4.46}
$$

where the right side is independent of $r \in \mathbb{R}^2$ and $\underline{p}_{(1,l)}$. Hence, (4.41) for $\sigma = (1,2)$ and $\nu = (1,l)$ now follows by combining (4.43) , (4.44) and (4.46) .

As already mentioned, the bound (4.42) is obtained in the same manner with the only difference that $W(r, r') = u_{\sigma}(r)v_{\nu}(r') - u_{\sigma}^h(r)v_{\nu}^h(r')$ in this case. Then, (4.44) has to be replaced by the identity

$$
\int \! \mathrm{d}r \, \mathrm{d}r' \, W(r,r')^2 = \|V_\sigma\|_{L^1} \|V_\nu\|_{L^1} - \|V_\sigma^h\|_{L^1} \|V_\nu^h\|_{L^1},
$$

which is a consequence of the identities $u_{\sigma} \cdot u_{\sigma}^h = |V_{\sigma}^h|$ and $v_{\nu} \cdot v_{\nu}^h = |V_{\nu}^h|$. This completes the proof of (4.41) and (4.42) if $\sigma = (1, 2)$ and $\nu = (1, l)$ with $l \geq 3$.

If $\sigma = (1,2)$ and $\nu = (2,l)$ with $l \geq 3$, then the kernel of $\Lambda_{\varepsilon}(z)_{\sigma\nu}$ only differs from the kernel of $\Lambda_{\varepsilon}(z)$ _(1,2)(_{1,*l*)} by the permutations $x'_1 \leftrightarrow x'_2$, $m_1 \leftrightarrow m_2$, $v_{(1,l)} \leftrightarrow v_{(2,l)}$ and the (unitary) reflection $r \to -r$. Therefore, the above proof also works in this case.

If $\sigma = (1, 2)$ and $\nu = (k, l)$ with $3 \leq k < l \leq N$, then the operator $\Lambda_{\varepsilon}(z)_{\sigma\nu}$ acts pointwise in $p_{(k,l)} = (p_3, \ldots \widehat{p_k} \ldots \widehat{p_l} \ldots p_N)$ by the operator $\Lambda_{\varepsilon}(z, \underline{p}_{(k,l)})_{\sigma\nu}$ that is defined by the kernel (4.40). In this case, the above estimates have to be slightly adjusted. The role of the operator F is now played by the operator $B \in \mathscr{L}(L^2(\mathbb{R}^2 \times \mathbb{R}^2 \times \mathbb{R}^2))$ from Lemma A.3. As the bounds for $||F||$ and for $||B||$ in Lemma A.3 differ by a factor of m^{-1} , while there is one mass factor more in front of the Green's function for $\nu = (k, l)$ with $3 \le k \le l \le N$ than for $\nu = (1, l)$ with $l \ge 3$, we again obtain (4.41) and (4.42) with $C(\sigma, \nu)$ given in the statement of the proposition. \Box

4.2.2 The off-diagonal limit operators

This section is a preparation for the next one, where we shall be concerned with the convergence, as $\varepsilon \to 0$, of

$$
\Lambda_{\varepsilon}(z)_{\sigma\nu} = -B_{\varepsilon,\sigma}R_0(z)A_{\varepsilon,\nu}^*, \qquad \sigma \neq \nu. \tag{4.47}
$$

From Proposition 3.3 we know that

$$
R_0(z)A_{\varepsilon,\nu}^* \to S(z)_{\nu}^* = G(z)_{\nu}^* \langle v_{\nu} |, \qquad (\varepsilon \downarrow 0) \tag{4.48}
$$

in $\mathscr{L}(\widetilde{\mathfrak{X}}_{\nu},\mathscr{H}),$ and

$$
B_{\varepsilon,\sigma} \to |u_{\sigma}\rangle T_{\sigma}, \qquad (\varepsilon \downarrow 0) \tag{4.49}
$$

in $\mathscr{L}(H^2(\mathbb{R}^{2N}), \widetilde{\mathfrak{X}}_{\sigma})$. Here, $|u_{\sigma}\rangle : \mathfrak{X}_{\sigma} \to \widetilde{\mathfrak{X}}_{\sigma}$ is defined by $|u_{\sigma}\rangle \psi = u_{\sigma} \otimes \psi$ and $\langle v_{\nu}| : \widetilde{\mathfrak{X}}_{\nu} \to \mathfrak{X}_{\nu}$ is the adjoint of $|v_\nu\rangle$. The negative of the formal composition of the limits in (4.48) and (4.49) is the operator

$$
|u_{\sigma}\rangle\,\langle v_{\nu}|\otimes\Theta(z)_{\sigma\nu}
$$

with

$$
\Theta(z)_{\sigma\nu} := -T_{\sigma} G(z)^{*}_{\nu}, \qquad \sigma \neq \nu. \tag{4.50}
$$

In the remainder of this section, we show that (4.50) defines an element $\Theta(z)_{\sigma\nu} \in \mathscr{L}(\mathfrak{X}_{\nu}, \mathfrak{X}_{\sigma})$. We begin by computing representations of T_{σ} and $G(z)^{*}_{\nu}$ in Fourier space. Using the definitions (3.2), (3.3) and (2.3) of T_{σ} , τ and \mathscr{K}_{σ} , respectively, we find for $\sigma = (i, j)$ and $\psi \in D(T_{\sigma})$ that

$$
\widehat{(T_{(i,j)}\psi)}\left(P, p_1, \ldots \widehat{p_i} \ldots \widehat{p_j} \ldots, p_N\right) = \frac{1}{2\pi} \int dp \, \widehat{\mathcal{K}_{(i,j)}\psi}\left(p, P, p_1, \ldots \widehat{p_i} \ldots \widehat{p_j} \ldots, p_N\right)
$$
\n
$$
= \frac{1}{2\pi} \int dp \, \widehat{\psi}\left(p_1, \ldots, p_{i-1}, \frac{m_i P}{m_i + m_j} - p, p_{i+1}, \ldots, p_{j-1}, \frac{m_j P}{m_i + m_j} + p, p_{j+1}, \ldots, p_N\right). \tag{4.51}
$$

To compute the Fourier transform of $G(z)_{\nu}^*w$ for $z > 0$, $\nu = (k, l)$, and $w \in \mathfrak{X}_{\nu}$, we use $G(z)_{\nu} = T_{\nu}R_0(z)$ in combination with (4.51) and the substitution

$$
P := p_k + p_l, \quad p := \frac{m_k p_l - m_l p_k}{m_k + m_l}.
$$

After a straightforward computation, we find that $\langle w | G(z)_{\nu} \psi \rangle = \langle G(z)_{\nu}^* w | \psi \rangle$ for any $\psi \in \mathcal{H}$, where

$$
\widehat{(G(z)_{\nu}^*w)}(p_1, ..., p_N) = \frac{1}{2\pi} \left(z + \sum_{n=1}^N \frac{p_n^2}{m_n} \right)^{-1} \widehat{w}(p_k + p_l, p_1, ... \widehat{p}_k ... \widehat{p}_l ..., p_N).
$$
 (4.52)

With these identities at hand, we now come to the main result of this section:

Proposition 4.5. Let $\sigma = (i, j) \neq (k, l) = \nu$ and $z > 0$ be given. Then $\Theta(z)_{\sigma\nu} = -T_{\sigma}G(z)_{\nu}^*$ *defines a bounded operator and* $\|\Theta(z)_{\sigma\nu}\| \leq \max(m_i, m_j, m_k, m_l)/4$.

In the proof of Proposition 4.5 we will need the following result taken from [6, Lemma 3.1]. For the convenience of the reader, we give a short proof with a different constant here.

Lemma 4.6. *For all* $f, g \in L^2(\mathbb{R}^2)$ *,*

$$
\int \frac{|f(x)||g(x')|}{|x|^2 + |x'|^2} dx dx' \le \pi^2 ||f|| ||g||. \tag{4.53}
$$

Proof. Let *K* denote the integral operator in $L^2(\mathbb{R}^2)$ that is defined by the kernel $K(x; x') =$ $(|x|^2 + |x'|^2)^{-1}$. Using the Schur test with $h(x) = |x|^{-1}$, it is straightforward to verify that *K* defines a bounded operator and that

$$
||K|| \le \underset{x' \in \mathbb{R}^2}{\text{ess sup}} \left(|x'| \int_{\mathbb{R}^2} \frac{1}{|x|^2 + |x'|^2} \frac{1}{|x|} \, \mathrm{d}x \right) = \pi^2.
$$

Hence, $\langle |f|, K |g| \rangle \leq \pi^2 ||f|| ||g||$, which establishes (4.53).

Proof of Proposition 4.5. Without loss of generality, we may assume that $\sigma = (1, 2)$. Moreover, it suffices to consider the two cases $\nu = (1,3)$ and $\nu = (3,4)$ corresponding to pairs with one common particle and no common particle, respectively.

To show that $\text{Ran } G(z)_{\nu}^* \subset D(T_{\sigma})$, or equivalently, that $\text{Ran }\mathscr{K}_{\sigma}G(z)_{\nu}^* \subset D(\tau)$, we have to verify that, for all $w \in \mathfrak{X}_{\nu}$,

$$
\widehat{\phi}(P, p_3, ..., p_N) := \frac{1}{2\pi} \int dp \left| \widehat{G(z)_{\nu}^* w} \left(\frac{m_1 P}{m_1 + m_2} - p, \frac{m_2 P}{m_1 + m_2} + p, p_3, ..., p_N \right) \right|
$$

defines an L^2 -function. To this end, by the Riesz-Lemma, it suffices to show that

$$
\int dP \, dp_3 \cdots dp_N \, |(\widehat{\psi}\widehat{\phi})(P, p_3, ..., p_N)| \le \text{const.} ||\psi||
$$

for all $\psi \in \mathfrak{X}_{\sigma}$. The substitution

$$
P := p_1 + p_2, \quad p := \frac{m_1 p_2 - m_2 p_1}{m_1 + m_2}
$$

and the identity (4.52) for $G(z)_{\nu}^*w$ show that

$$
\int dP \, dp_3 \cdots dp_N |\hat{\psi}(P, p_3, ..., p_N)||\hat{\phi}(P, p_3, ..., p_N)|
$$
\n
$$
= \frac{1}{2\pi} \int dp_1 \cdots dp_N |\hat{\psi}(p_1 + p_2, p_3, ..., p_N)| |\widehat{G(z)}_p^* w(p_1, ..., p_N)|
$$
\n
$$
\leq \frac{\max(m_2, m_l)}{4\pi^2} \int dp_1 \cdots dp_N \frac{|\hat{\psi}(p_1 + p_2, p_3, ..., p_N)||\hat{w}(p_k + p_l, p_1, ... \hat{p}_k ... \hat{p}_l ..., p_N)|}{p_2^2 + p_l^2}.
$$
\n(4.54)

If $\nu = (k, l) = (1, 3)$, let $p := (p_4, ..., p_N)$ and set

$$
f(p_3, \underline{p}) := \left(\int dp_1 \, |\widehat{\psi}(p_1, p_3, \underline{p})|^2 \right)^{1/2}
$$

$$
g(p_2, \underline{p}) := \left(\int dp_1 \, |\widehat{w}(p_1, p_2, \underline{p})|^2 \right)^{1/2}.
$$

Then $f(\cdot, \underline{p}), g(\cdot, \underline{p}) \in L^2(\mathbb{R}^2)$ for almost all \underline{p} . Hence, after applying the Cauchy-Schwarz inequality in the p_1 -integration, it follows from Lemma 4.6 that

$$
\int d\underline{p} \, dp_2 \, dp_3 \, dp_1 \, \frac{|\widehat{\psi}(p_1 + p_2, p_3, \underline{p})| \, |\widehat{w}(p_1 + p_3, p_2, \underline{p})|}{p_2^2 + p_3^2} \le \int d\underline{p} \, dp_2 \, dp_3 \, \frac{|f(p_3, \underline{p})| \, |g(p_2, \underline{p})|}{p_2^2 + p_3^2} \le \pi^2 ||\psi|| ||w||. \tag{4.55}
$$

If $\nu = (k, l) = (3, 4)$, we set $p := (p_5, ..., p_N)$ and

$$
f(p_4, \underline{p}) := \left(\int dp_1 dp_3 \, |\hat{\psi}(p_1, p_3, p_4, \underline{p})|^2 \right)^{1/2}
$$

$$
g(p_2, \underline{p}) := \left(\int dp_1 dp_3 \, |\hat{w}(p_3, p_1, p_2, \underline{p})|^2 \right)^{1/2}.
$$

Using the Cauchy-Schwarz inequality in the (p_1, p_3) -integration, we conclude from Lemma 4.6 that

$$
\int d\underline{p} dp_2 dp_4 dp_1 dp_3 \frac{|\widehat{\psi}(p_1 + p_2, p_3, p_4, \underline{p})| |\widehat{w}(p_3 + p_4, p_1, p_2, \underline{p})|}{p_2^2 + p_4^2}
$$

\n
$$
\leq \int d\underline{p} dp_2 dp_4 \frac{|f(p_4, \underline{p})| |g(p_2, \underline{p})|}{p_2^2 + p_4^2} \leq \pi^2 ||\psi|| ||w||. \tag{4.56}
$$

From (4.54), (4.55) and (4.56), it follows that $\hat{\phi} \in L^2$ and hence $\text{Ran } G(z)_{\nu}^* \subset D(T_{\sigma})$. Furthermore, the above estimates now show that

$$
|\langle \psi | T_{\sigma} G(z)_{\nu}^* w \rangle| \le \frac{\max(m_1, m_2, m_3, m_4)}{4} ||\psi|| ||w||.
$$

Therefore, $\Theta(z)_{\sigma\nu} = -T_{\sigma}G(z)^{*}_{\nu}$ is a bounded operator, whose norm satisfies the desired estimate. \Box

4.2.3 Cutoff functions and convergence

The goal of this section is to prove Proposition 4.8 on the convergence, as $\varepsilon \to 0$, of $\Lambda_{\varepsilon}(z)_{\sigma\nu}$ with a suitable space cutoff $\chi_{\sigma\nu,c}$ defined below. In the analysis of $\lim_{\varepsilon\to 0} \Lambda_{\varepsilon}(z)^{-1}$ in Section 4.3, this cutoff will be removed again.

We begin by motivating the cutoff: By (4.49), $\lim_{\varepsilon \to 0} B_{\varepsilon,\sigma}$ exists in $\mathscr{L}(H^2(\mathbb{R}^{2N}), \widetilde{\mathfrak{X}}_{\sigma})$. Here, the Sobolev index 2 could be reduced to $1 + \delta$ for some $\delta > 0$ but not further because of the presence of the trace operator T_{σ} in (4.49). Moreover, by (4.48), $\lim_{\varepsilon \to 0} R_0(z) A_{\varepsilon,\nu}^*$ exists in $\mathscr{L}(\widetilde{\mathfrak{X}}_{\nu},\mathscr{H})$, or perhaps in $\mathscr{L}(\widetilde{\mathfrak{X}}_{\nu},H^{1-\delta}(\mathbb{R}^{2N}))$, which is not enough to prove that $\Lambda_{\varepsilon}(z)_{\sigma\nu}$ given by (4.47) has a limit as $\varepsilon \to 0$. The space cutoff to be introduced below removes the singularity due to T^*_{ν} . More explicitly, for $\nu = (1, 2)$, $v = v_{\nu}$ and $\widetilde{R}_0(z) = (\widetilde{H}_0 + z)^{-1}$ (see (4.5) and (4.6)), the operator $\widetilde{R}_0(z) \varepsilon^{-1} U^*_{\varepsilon}(v \otimes 1)$ has the kernel

$$
G_{z,\underleftarrow{\widetilde{m}}}(r-\varepsilon r', R-R', x_3-x'_3, ..., x_N-x'_N) v(r'),
$$

where $\tilde{m} := (\mu, m_1 + m_2, m_3, ..., m_N)$. Hence, if *v* has compact support and $|r| \geq c > 0$ is enforced by a space cutoff χ_c , then, for sufficiently small $\varepsilon > 0$, the singularity of $G_{z,\tilde{m}}$ at the origin is avoided and, by Lemma A.2, this means that the kernel becomes a smooth function of $(r, R, x_3, ..., x_N)$.

Let $\chi \in C^{\infty}(\mathbb{R}, [0,1])$ be a function with $\chi(x) = 0$ for $x \le 1$ and $\chi(x) = 1$ for $x \ge 2$ and set $\chi_c(r) := \chi(|r|/c)$ for $r \in \mathbb{R}^2$ and $c > 0$.

Figure 1: The cutoff function *χ*

Then we have the following result:

Lemma 4.7. *Assume that* $v \in L^2(\mathbb{R}^2)$ *has compact support* supp $(v) \subset \{x \in \mathbb{R}^2 \mid |x| \leq h\}$ *for some* $h > 0$ *and let* $c, z > 0$ *. Then, for all* $n \in \mathbb{N}_0$ *, the limit*

$$
\lim_{\varepsilon \to 0} (\chi_c \otimes 1) \widetilde{R}_0(z) \varepsilon^{-1} U_{\varepsilon}^*(v \otimes 1) = (\chi_c \otimes 1) \mathscr{K} S(z)^*
$$

exists in $\mathscr{L}(L^2(\mathbb{R}^{2N}), H^n(\mathbb{R}^{2N}))$ *, where* $\mathscr{K} = \mathscr{K}_{(1,2)}$ *and* $S(z) = S(z)_{(1,2)}$ *.*

Proof. Clearly, the lemma will follow if we show that

$$
\lim_{\varepsilon \to 0} \left(\partial^{\alpha} \chi_c \right) \widetilde{R}_0(z) \varepsilon^{-1} U_{\varepsilon}^*(v \otimes 1) = \left(\partial^{\alpha} \chi_c \right) \mathcal{K} S(z)^* \qquad \text{in } \mathcal{L}(L^2(\mathbb{R}^{2N}), H^n(\mathbb{R}^{2N})) \tag{4.57}
$$

for all $n \in \mathbb{N}_0$ and all multi-indices $\alpha \in \mathbb{N}_0^2$, where $\partial^\alpha \chi_c = (\partial^\alpha \chi_c) \otimes 1$ for short. Using (4.5) in combination with the adjoint of (2.6) and (4.48), it follows that

$$
\lim_{\varepsilon \to 0} \widetilde{R}_0(z) \varepsilon^{-1} U_{\varepsilon}^*(v \otimes 1) = \mathcal{K} S(z)^* \qquad \text{in } \mathcal{L}(L^2(\mathbb{R}^{2N})),
$$

so (4.57) holds for $n = 0$.

We now proceed by induction and assume that (4.57) holds for some $n \in \mathbb{N}_0$ and all $\alpha \in \mathbb{N}_0^2$. To show that (4.57) also holds for $n + 1$, we are going to use that

$$
\eta \widetilde{R}_0(z) = \widetilde{R}_0(z) \left[\widetilde{H}_0, \eta \right] \widetilde{R}_0(z) + \widetilde{R}_0(z) \eta, \qquad \eta = \partial^{\alpha} \chi_c, \qquad (4.58)
$$

where [·*,* ·] denotes the commutator and

$$
\lim_{\varepsilon \to 0} \widetilde{R}_0(z) \eta \,\varepsilon^{-1} U_{\varepsilon}^*(v \otimes 1) = 0 \qquad \qquad \text{in } \mathscr{L}(L^2(\mathbb{R}^{2N}), H^{n+1}(\mathbb{R}^{2N})) \tag{4.59}
$$

because $\eta U_{\varepsilon}^*(v \otimes 1) = 0$ for $\varepsilon h < c$. Let $D_k : H^1(\mathbb{R}^2) \to L^2(\mathbb{R}^2)$ denote the partial derivative $D_k \varphi := \partial_k \varphi$ for $k \in \{1, 2\}$. From the definition (4.6) of H_0 and from $\Delta_r = D_1 D_1 + D_2 D_2$, it follows that

$$
\mu\left[\widetilde{H}_0,\eta\right] = \left[-\Delta_r,\eta\right] = -\sum_{k=1}^2 \left(D_k(\partial_k \eta) + (\partial_k \eta)D_k\right) = \left(\Delta_r \eta\right) - 2\sum_{k=1}^2 D_k(\partial_k \eta),\tag{4.60}
$$

where $\mu = \mu_{(1,2)}$ is a reduced mass. Equations (4.58) and (4.60) imply that

$$
\eta \widetilde{R}_0(z) = \widetilde{R}_0(z)\eta + \mu^{-1} \widetilde{R}_0(z)(\Delta_r \eta) \widetilde{R}_0(z) - 2\mu^{-1} \sum_{k=1}^2 D_k \widetilde{R}_0(z)(\partial_k \eta) \widetilde{R}_0(z).
$$
 (4.61)

After multiplying (4.61) with $\varepsilon^{-1}U^*_{\varepsilon}(v\otimes 1)$ from the right, we get from (4.59) combined with the induction hypothesis and the fact that $\widetilde{R}_0(z)$ and $D_k \widetilde{R}_0(z)$ belong to $\mathscr{L}(H^n(\mathbb{R}^{2N}), H^{n+1}(\mathbb{R}^{2N}))$ that $\lim_{\varepsilon \to 0} \eta \widetilde{R}_0(z) \varepsilon^{-1} U_{\varepsilon}^*(v \otimes 1)$ exists in $\mathscr{L}(L^2(\mathbb{R}^{2N}), H^{n+1}(\mathbb{R}^{2N}))$. Furthermore, the limit operator has to be the one in (4.57) because convergence in $\mathscr{L}(L^2(\mathbb{R}^{2N}), H^{n+1}(\mathbb{R}^{2N}))$ implies convergence in $\mathscr{L}(L^2(\mathbb{R}^{2N}), H^n(\mathbb{R}^{2N}))$. Hence, (4.57) holds with $n+1$ in place of *n* and the proof is complete. П

With the help of Lemma 4.7 we can now prove convergence of the regularized operators $(1 \otimes \chi_{\sigma\nu,c})\Lambda_{\varepsilon}(z)_{\sigma\nu}$ for pairs $\sigma = (i,j) \neq (k,l) = \nu$, and a cutoff function $\chi_{\sigma\nu,c}$ defined by

$$
\chi_{(i,j)(k,l),c}(R,x_1,...\hat{x}_i...\hat{x}_j...,x_N) := \begin{cases} \chi_c(x_l - R) & \text{if } k \in \{i,j\}, l \notin \{i,j\}, \\ \chi_c(x_k - R) & \text{if } k \notin \{i,j\}, l \in \{i,j\}, \\ \chi_c(x_l - x_k) & \text{if } k, l \notin \{i,j\}. \end{cases}
$$
(4.62)

Proposition 4.8. *For all pairs* $\sigma, \nu \in \mathcal{I}, \sigma \neq \nu$ *and* $c, z > 0$ *the limit*

$$
\lim_{\varepsilon \to 0} (1 \otimes \chi_{\sigma \nu, c}) \Lambda_{\varepsilon}(z)_{\sigma \nu} = (1 \otimes \chi_{\sigma \nu, c}) \Lambda(z)_{\sigma \nu}
$$
\n(4.63)

exists in $\mathscr{L}(\widetilde{\mathfrak{X}}_{\nu}, \widetilde{\mathfrak{X}}_{\sigma})$ *, where*

$$
\Lambda(z)_{\sigma\nu} := |u_{\sigma}\rangle \langle v_{\nu}| \otimes \Theta(z)_{\sigma\nu} \tag{4.64}
$$

 $and \Theta(z)_{\sigma\nu} = -T_{\sigma}G(z)^{*}_{\nu} \in \mathscr{L}(\mathfrak{X}_{\nu}, \mathfrak{X}_{\sigma}).$

Proof. Assume, for the moment, that for all $h > 0$,

$$
\lim_{\varepsilon \to 0} (1 \otimes \chi_{\sigma \nu, c}) \Lambda_{\varepsilon}^{h}(z)_{\sigma \nu} = (1 \otimes \chi_{\sigma \nu, c}) \Lambda^{h}(z)_{\sigma \nu}, \tag{4.65}
$$

where $\Lambda_{\varepsilon}^{h}(z)_{\sigma\nu}$ was introduced in Section 4.2.1 and $\Lambda^{h}(z)_{\sigma\nu} := |u_{\sigma}^{h}\rangle \langle v_{\nu}^{h}| \otimes \Theta(z)_{\sigma\nu}$. By Proposition 4.5, $\Theta(z)_{\sigma\nu} = -T_{\sigma}G(z)_{\nu}^*$ defines a bounded operator and $\|\Theta(z)_{\sigma\nu}\| \leq \widetilde{C}(\sigma,\nu)$, so the definition (4.64) and the corresponding identity for $\Lambda^h(z)_{\sigma\nu}$ imply that

$$
\|\Lambda(z)_{\sigma\nu} - \Lambda^h(z)_{\sigma\nu}\| \leq \left\| |u_{\sigma}\rangle \langle v_{\nu}| - |u_{\sigma}^h\rangle \langle v_{\nu}^h| \right\| \|\Theta(z)_{\sigma\nu}\|
$$

$$
\leq \widetilde{C}(\sigma, \nu) \left(\|V_{\sigma}\|_{L^1} \|V_{\nu}\|_{L^1} - \|V_{\sigma}^h\|_{L^1} \|V_{\nu}^h\|_{L^1} \right)^{1/2}, \tag{4.66}
$$

which is the equivalent of (4.42) for $\varepsilon = 0$. Now, the general case of (4.63), where u_{σ} and v_{ν} do not have compact support, follows from (4.65) , (4.42) and (4.66) by a simple $\delta/3$ -argument.

It remains to prove (4.65). As the choice of the pair ν is immaterial for the following arguments, it suffices to consider the case $\nu = (1, 2)$ only. Moreover, we may assume that $\supp(u_{\sigma}) \cup \supp(v_{\nu}) \subset \{x \in \mathbb{R}^2 \mid |x| \leq h\}$, i.e. $u_{\sigma}^h = u_{\sigma}$ and $v_{\nu}^h = v_{\nu}$. Then (4.35) becomes

$$
\Lambda_{\varepsilon}^{h}(z)_{\sigma\nu} = -\varepsilon^{-2} \left(u_{\sigma} \otimes 1 \right) U_{\varepsilon} \mathscr{K}_{\sigma} R_{0}(z) \mathscr{K}_{\nu}^{*} U_{\varepsilon}^{*}(v_{\nu} \otimes 1). \tag{4.67}
$$

From Lemma 3.2, combined with Eqs. (3.2) and (3.3), it follows that

$$
\lim_{\varepsilon \to 0} \varepsilon^{-1} \left(u_{\sigma} \otimes 1 \right) U_{\varepsilon} \mathscr{K}_{\sigma} = \left(|u_{\sigma} \rangle \langle \delta | \otimes 1 \rangle \mathscr{K}_{\sigma} = |u_{\sigma} \rangle T_{\sigma} \qquad \text{in } \mathscr{L}(H^{2}(\mathbb{R}^{2N}), \widetilde{\mathfrak{X}}_{\sigma}). \tag{4.68}
$$

Next, setting

$$
\chi_{\nu,c}(x_1,...,x_N) := \chi_c(|x_2 - x_1|), \qquad \nu = (1,2),
$$

the identities $R_0(z)\mathscr{K}_\nu^* = \mathscr{K}_\nu^* \widetilde{R}_0(z)$, $\chi_{\nu,c}\mathscr{K}_\nu^* = \mathscr{K}_\nu^*(\chi_c \otimes 1)$ and Lemma 4.7 imply that

$$
\lim_{\varepsilon \to 0} \chi_{\nu,c} R_0(z) \mathscr{K}_{\nu}^* \varepsilon^{-1} U_{\varepsilon}^*(v_\nu \otimes 1) = \chi_{\nu,c} S(z)_{\nu}^* \qquad \text{in } \mathscr{L}(\widetilde{\mathfrak{X}}_{\nu}, H^n(\mathbb{R}^{2N})) \tag{4.69}
$$

for any $n \in \mathbb{N}_0$. Hence, we conclude from (4.68) and (4.69) that

$$
\lim_{\varepsilon \to 0} -\varepsilon^{-2} \left(u_{\sigma} \otimes 1 \right) U_{\varepsilon} \mathscr{K}_{\sigma} \chi_{\nu,c} R_0(z) \mathscr{K}_{\nu}^* U_{\varepsilon}^*(v_{\nu} \otimes 1) = - \left| u_{\sigma} \right\rangle T_{\sigma} \chi_{\nu,c} S(z)_{\nu}^* \tag{4.70}
$$

in $\mathscr{L}(\widetilde{\mathfrak{X}}_{\nu}, \widetilde{\mathfrak{X}}_{\sigma})$. To complete the proof of (4.65), we have to move the cutoff to the left on both sides of (4.70). We start by considering the limit operators. For a given $\psi \in \widetilde{\mathfrak{X}}_{\nu}$, let $\widetilde{\psi} := S(z)_{\nu}^*\psi$. Then (4.69) shows that $\chi_{\nu,c}\widetilde{\psi} \in H^n(\mathbb{R}^{2N})$ for any $n \in \mathbb{N}$, so it follows from a standard Sobolev embedding theorem (see e.g. [15, Theorem 8.8]) that $\chi_{\nu,c}\tilde{\psi}$ defines a smooth function. In particular, $\chi_{\nu,c}\tilde{\psi} \in D(T_{\sigma})$ and $T_{\sigma}(\chi_{\nu,c}\tilde{\psi})$ is explicitly given by (1.6). Now, a direct calculation, using the defining relations for $\chi_{\nu,c}$ and $\chi_{\sigma\nu,c}$, yields that

$$
T_{\sigma}(\chi_{\nu,c}\widetilde{\psi})=\chi_{\sigma\nu,c}T_{\sigma}\widetilde{\psi},
$$

where, by (3.6) and Proposition 4.5, $\tilde{\psi} = S(z)_{\nu}^* \psi = G(z)_{\nu}^* \langle v_{\nu} | \psi \in D(T_{\sigma})$. Therefore,

$$
- |u_{\sigma}\rangle T_{\sigma}\chi_{\nu,c}S(z)_{\nu}^*\psi = (|u_{\sigma}\rangle \langle v_{\nu}| \otimes [\chi_{\sigma\nu,c}\Theta(z)_{\sigma\nu}])\psi,
$$

so the limit operator in (4.70) agrees with the desired limit operator in (4.65).

It remains to show that the left side of (4.70) agrees with the left side of (4.65) with $\Lambda_{\varepsilon}^{h}(z)_{\sigma\nu}$ given by (4.67). If $\sigma = (k, l)$ with $3 \leq k < l \leq N$, then this follows immediately from $U_{\varepsilon} \mathscr{K}_{\sigma} \chi_{\nu, c} =$ $(1 \otimes \chi_{\sigma\nu,c})U_{\varepsilon}\mathscr{K}_{\sigma}$. If $\sigma = (1,l)$ with $l \geq 3$, then, by some abuse of notation,

$$
U_{\varepsilon}\mathscr{K}_{\sigma}\chi_{\nu,c} = \chi_c \left(x_2 - R + \frac{\varepsilon m_l r}{m_1 + m_l}\right) U_{\varepsilon}\mathscr{K}_{\sigma}.
$$
 (4.71)

For the purpose of computing the limit in (4.70) , the right side of (4.71) may be replaced by $\chi_c(|x_2 - R|)U_\varepsilon \mathscr{K}_\sigma$ because χ_c is Lipschitz, u_σ has compact support and, by Proposition 4.4, $\|\Lambda_{\varepsilon}^{h}(z)_{\sigma\nu}\|$ is uniformly bounded in $\varepsilon > 0$. Again, (4.65) follows. The remaining case $\sigma = (2, l)$ with $l \geq 3$ is treated similarly.

4.3 Convergence of $\Lambda_{\varepsilon}(z)^{-1}$

In Sections 4.1 and 4.2 we have seen that $(\Lambda_{\varepsilon}(z)_{\text{diag}})^{-1}$ and a regularized version of $\Lambda_{\varepsilon}(z)_{\text{off}}$ have limits as $\varepsilon \to 0$. This will now allow us to prove invertibility of $\Lambda_{\varepsilon}(z)$ for small $\varepsilon > 0$ and large enough $z > 0$, and existence of $\lim_{\varepsilon \to 0} \Lambda_{\varepsilon}(z)^{-1}$. We claim that

$$
\lim_{\varepsilon \to 0} (\Lambda_{\varepsilon}(z)^{-1})_{\sigma \nu} = \begin{cases} \frac{|u_{\sigma}\rangle \langle v_{\nu}|}{\langle u_{\sigma} | v_{\sigma}\rangle \langle u_{\nu} | v_{\nu}\rangle} \otimes (\Theta(z)^{-1})_{\sigma \nu} & \text{if } \sigma, \nu \in \mathcal{J} \\ 0 & \text{else,} \end{cases}
$$
(4.72)

where $\Theta(z) = (\Theta(z)_{\sigma\nu})_{\sigma,\nu\in\mathcal{J}}$ is invertible in the reduced Hilbert space \mathfrak{X} defined by (1.5).

To prove (4.72) , we introduce some auxiliary operators. Let $\Pi : \widetilde{\mathfrak{X}} \to L^2(\mathbb{R}^2, dr) \otimes \mathfrak{X}$ denote the matrix operator defined by

$$
\Pi_{\sigma\nu} := \begin{cases} \delta_{\sigma\nu} & \text{if } \sigma, \nu \in \mathcal{J} \\ 0 & \text{if } \sigma \in \mathcal{J}, \nu \in \mathcal{I} \setminus \mathcal{J} \end{cases}
$$

and let $U = (U_{\sigma\nu})_{\sigma,\nu \in \mathcal{J}}$, where $U_{\sigma\nu} \in \mathscr{L}(L^2(\mathbb{R}^2))$ is given by

$$
U_{\sigma\nu} := \frac{|u_{\sigma}\rangle \langle v_{\nu}|}{\langle u_{\sigma} | v_{\sigma}\rangle \langle u_{\nu} | v_{\nu}\rangle}.
$$

Let $\Theta(z)_{\text{diag}}$ and $\Theta(z)_{\text{off}}$ denote the operators in $\mathfrak X$ defined in terms of the components $\Theta(z)_{\sigma\sigma}$ and $\Theta(z)_{\sigma\nu}$, $\sigma \neq \nu$, that we have introduced in Sections 4.1 and 4.2.2, respectively, and let

$$
\Theta(z) := \Theta(z)_{\text{diag}} + \Theta(z)_{\text{off}}.
$$

Proposition 4.5 implies that $\Theta(z)_{\text{off}} \in \mathscr{L}(\mathfrak{X})$, while, by (4.32), $\Theta(z)_{\text{diag}}$ is unbounded in \mathfrak{X} . With the help of Π and U , Equation (4.31) can now be written as

$$
\lim_{\varepsilon \to 0} (\Lambda_{\varepsilon}(z)_{\text{diag}})^{-1} = \Pi^* \left(U \circ (\Theta(z)_{\text{diag}})^{-1} \right) \Pi. \tag{4.73}
$$

Here, the operator product $A \circ B$ is defined by $(A \circ B)_{ij} = A_{ij} \otimes B_{ij}$, a notation inspired by the Hadamard-Schur product of matrices.

The following proposition proves (4.72) in these new notations:

Proposition 4.9. *There exist* ε_0 , $z_0 > 0$ *such that the operator matrix* $\Lambda_{\varepsilon}(z)$ *is invertible in* $\widetilde{\mathfrak{X}}$ *for all* $\varepsilon \in (0, \varepsilon_0)$ *and* $z > z_0$ *, and then*

$$
\lim_{\varepsilon \to 0} \Lambda_{\varepsilon}(z)^{-1} = \Pi^*(U \circ \Theta(z)^{-1})\Pi,
$$
\n(4.74)

where $\Theta(z)$ *is a closed and invertible operator in* \mathfrak{X} *.*

Proof. Recall from (4.2)-(4.4) that $\Lambda_{\varepsilon}(z) = \Lambda_{\varepsilon}(z)_{\text{diag}} + \Lambda_{\varepsilon}(z)_{\text{off}}$, where $\Lambda_{\varepsilon}(z)_{\text{diag}}$ and $\Lambda_{\varepsilon}(z)_{\text{off}}$ is the diagonal and off-diagonal part of the operator-valued matrix $\Lambda_{\varepsilon}(z) = (g_{\varepsilon,\sigma}^{-1} \delta_{\sigma\nu} - \phi_{\varepsilon}(z)_{\sigma\nu})_{\sigma,\nu \in \mathcal{I}},$ respectively. By Proposition 4.4, $\|\Lambda_{\varepsilon}(z)_{\text{off}}\| \leq C_{\text{off}} < \infty$ holds uniformly in $\varepsilon, z > 0$. Furthermore, (4.73) and the definition (4.32) of $\Theta(z, \underline{P}_{\sigma})_{\sigma\sigma}$ imply that $\Lambda_{\varepsilon}(z)_{\text{diag}}$ is invertible and $\|(\Lambda_{\varepsilon}(z)_{\text{diag}})^{-1}\| \leq (2C_{\text{off}})^{-1}$, provided that $z > z_0$ and $\varepsilon \in (0, \varepsilon_0)$ for sufficiently large $z_0 > 0$ and sufficiently small $\varepsilon_0 > 0$. We conclude that $\Lambda_{\varepsilon}(z)$ is invertible and

$$
\Lambda_{\varepsilon}(z)^{-1} = \left(1 + (\Lambda_{\varepsilon}(z)_{\text{diag}})^{-1} \Lambda_{\varepsilon}(z)_{\text{off}}\right)^{-1} (\Lambda_{\varepsilon}(z)_{\text{diag}})^{-1}, \qquad z > z_0, \ \varepsilon \in (0, \varepsilon_0). \tag{4.75}
$$

Next, we claim that $\lim_{\varepsilon \to 0} (\Lambda_{\varepsilon}(z)_{\text{diag}})^{-1} \Lambda_{\varepsilon}(z)_{\text{off}}$ exists for all large enough $z > 0$, which, by (4.3), is equivalent to the assertion that $\lim_{\varepsilon\to 0} (g_{\varepsilon,\sigma}^{-1} - \phi_{\varepsilon}(z)_{\sigma\sigma})^{-1}\Lambda_{\varepsilon}(z)_{\sigma\nu}$ exists for all pairs $\sigma, \nu \in I$, $\sigma \neq \nu$. Without loss of generality, we may assume that $\sigma = (1, 2) \neq (k, l) = \nu$ and in the following we drop the index σ in the diagonal contributions, i.e. $\phi_{\varepsilon}(z) := \phi_{\varepsilon}(z)_{\sigma\sigma}$, $V := V_{\sigma}$, $g_{\varepsilon} := g_{\varepsilon,\sigma}$, $a := a_{\sigma}$ etc. If $\int V(r) dr/(2\pi) < a$, then it follows from Lemma 4.2 and from the fact that, by Proposition 4.4, $\|\Lambda_{\varepsilon}(z)_{\sigma\nu}\| \leq C(\sigma,\nu)$ is uniformly bounded in $\varepsilon, z > 0$ that, for large enough $z > 0$,

$$
\lim_{\varepsilon \to 0} \left(g_{\varepsilon}^{-1} - \phi_{\varepsilon}(z) \right)^{-1} \Lambda_{\varepsilon}(z)_{\sigma \nu} = 0, \qquad \nu \neq (1, 2).
$$

If $\int V(r) dr = 2\pi a$, then it is more subtle to prove existence of the limit $\varepsilon \to 0$. We claim that, for all large enough $z > 0$,

$$
\lim_{\varepsilon \to 0} \left(g_{\varepsilon}^{-1} - \phi_{\varepsilon}(z) \right)^{-1} \Lambda_{\varepsilon}(z)_{\sigma \nu} = \left(\frac{|u\rangle \langle v|}{\langle u | v \rangle^2} \otimes D(z)^{-1} \right) \Lambda(z)_{\sigma \nu}, \qquad \nu \neq (1, 2), \qquad (4.76)
$$

where $D(z, \underline{P})$ is defined by (4.22) with $\alpha = \alpha_{\sigma}$ and $\Lambda(z)_{\sigma\nu} \in \mathscr{L}(\widetilde{\mathfrak{X}}_{\nu}, \widetilde{\mathfrak{X}}_{\sigma})$ is defined by (4.64). To prove this, we start by showing that large momenta may be discarded: As in Section 4.1, let $P = (P, p_3, ..., p_N)$ be conjugate to $(R, x_3, ..., x_N)$ and let $\eta_K(K > 0)$ denote multiplication with

the characteristic function of the ball $\{P \mid |P| \leq K\}$ in Fourier space. Now, observe that (4.8) implies $Q \ge \lambda |P|^2$ with $\lambda := 1/\max(m_1 + m_2, m_3, ..., m_N) > 0$, so for fixed $\delta > 0$ and $z > z_0$, Lemma 4.2 gives

$$
\left\| \left(g_{\varepsilon}^{-1} - \phi_{\varepsilon}(z) \right)^{-1} (1 \otimes (1 - \eta_K)) \Lambda_{\varepsilon}(z)_{\sigma\nu} \right\| \leq \tilde{C} \max \left\{ |\ln(\varepsilon)|^{-1}, \ln(\mu(z + \lambda K^2))^{-1} \right\} C(\sigma, \nu) < \delta/3,
$$
\n(4.77)

provided that $K > 0$ is large enough and $\varepsilon > 0$ is small enough. Furthermore, using that, by Proposition 4.3,

$$
\lim_{\varepsilon \to 0} \left(g_{\varepsilon}^{-1} - \phi_{\varepsilon}(z) \right)^{-1} = \frac{|u\rangle \langle v|}{\langle u | v \rangle^2} \otimes D(z)^{-1}, \tag{4.78}
$$

similar estimates also show that

$$
\left\| \left(\frac{|u\rangle\langle v|}{\langle u|v\rangle^2} \otimes D(z)^{-1} \right) (1 \otimes (1 - \eta_K)) \Lambda(z)_{\sigma\nu} \right\| < \delta/3 \tag{4.79}
$$

for large enough $K > 0$. In view of (4.77) and (4.79), it is clear that (4.76) will follow if we can show that

$$
\lim_{\varepsilon \to 0} \left(g_{\varepsilon}^{-1} - \phi_{\varepsilon}(z) \right)^{-1} (1 \otimes \eta_K) \Lambda_{\varepsilon}(z)_{\sigma\nu} = \left(\frac{|u\rangle \langle v|}{\langle u | v \rangle^2} \otimes D(z)^{-1} \right) (1 \otimes \eta_K) \Lambda(z)_{\sigma\nu} \tag{4.80}
$$

for any fixed $K > 0$, $\nu \neq (1, 2)$ and large enough $z > 0$. To further simplify (4.80), we decompose

$$
\eta_K = \eta_K \chi_{\sigma\nu,c} + \eta_K (1 - \chi_{\sigma\nu,c}), \qquad c > 0,
$$
\n(4.81)

where $\chi_{\sigma\nu,c} = \chi_{(1,2)\nu,c}$ is defined by (4.62). Furthermore, for $\nu = (k,l) \neq (1,2) = \sigma$, we set

$$
\gamma_{\sigma\nu}(R, x_3, ..., x_N) := \begin{cases} |R - x_l|^{1/2} & \text{if } k \in \{1, 2\}, l \ge 3\\ |x_k - x_l|^{1/2} & \text{if } 3 \le k < l \le N \end{cases}
$$

and we note that the definition of $\chi_{\sigma\nu,c}$ implies that

$$
\|\eta_K(1 - \chi_{\sigma\nu,c})\| = \|(1 - \chi_{\sigma\nu,c})\eta_K\| \le \|(1 - \chi_{\sigma\nu,c})\gamma_{\sigma\nu}\| \|\gamma_{\sigma\nu}^{-1}(-\Delta' + 1)^{-1}(-\Delta' + 1)\eta_K\|
$$

$$
\le (2c)^{1/2} \|\gamma_{\sigma\nu}^{-1}(-\Delta' + 1)^{-1}\| (K^2 + 1), \tag{4.82}
$$

where

$$
-\Delta' := -\Delta_R + \sum_{i=3}^N (-\Delta_{x_i}).
$$

 $\text{As } | \cdot |^{-1/2} \in L^2(\mathbb{R}^2) + L^{\infty}(\mathbb{R}^2)$, a standard result (see e.g. [24, Theorem 11.1]) yields that $\|\gamma_{\sigma\nu}^{-1}(-\Delta'+1)^{-1}\| < \infty$, so the right side of (4.82) vanishes as $c \to 0$. Hence, with the help of (4.81) and another $\delta/3$ -argument, we see that the proof of (4.80) can be reduced to the claim that, for any fixed $c, K > 0, \nu \neq (1, 2)$ and large enough $z > 0$,

$$
\lim_{\varepsilon \to 0} \left(g_{\varepsilon}^{-1} - \phi_{\varepsilon}(z) \right)^{-1} (1 \otimes (\eta_K \chi_{\sigma \nu,c})) \Lambda_{\varepsilon}(z)_{\sigma \nu} = \left(\frac{|u\rangle \langle v|}{\langle u | v \rangle^{2}} \otimes D(z)^{-1} \right) (1 \otimes (\eta_K \chi_{\sigma \nu,c})) \Lambda(z)_{\sigma \nu}.
$$

However, this is immediate from (4.78) and Proposition 4.8, so (4.80) and thus (4.76) are established.

Since the choice $\sigma = (1, 2)$ in the analysis above was immaterial, we conclude that, for all pairs $\sigma, \nu \in \mathcal{I}, \sigma \neq \nu$,

$$
\lim_{\varepsilon \to 0} \left(g_{\varepsilon,\sigma}^{-1} - \phi_{\varepsilon}(z)_{\sigma\sigma} \right)^{-1} \Lambda_{\varepsilon}(z)_{\sigma\nu} = \begin{cases} \left(\frac{|u_{\sigma}\rangle \langle v_{\sigma}|}{\langle u_{\sigma} | v_{\sigma} \rangle^{2}} \otimes (\Theta(z)_{\sigma\sigma})^{-1} \right) \Lambda(z)_{\sigma\nu} & \text{if } \sigma \in \mathcal{J} \\ 0 & \text{else.} \end{cases}
$$
(4.83)

Let the operator $\Lambda(z)_{\text{off}}$ in $\tilde{\mathfrak{X}}$ be defined by $(\Lambda(z)_{\text{off}})_{\sigma\nu} = \Lambda(z)_{\sigma\nu} (1 - \delta_{\sigma\nu})$. Then, in the notation of (4.73), (4.83) takes the form

$$
\lim_{\varepsilon \to 0} (\Lambda_{\varepsilon}(z)_{\text{diag}})^{-1} \Lambda_{\varepsilon}(z)_{\text{off}} = \Pi^* \left(U \circ (\Theta(z)_{\text{diag}})^{-1} \right) \Pi \Lambda(z)_{\text{off}} =: L(z). \tag{4.84}
$$

From (4.75), (4.73) and (4.84), it follows that

$$
\lim_{\varepsilon \to 0} \Lambda_{\varepsilon}(z)^{-1} = (1 + L(z))^{-1} \Pi^* \left(U \circ (\Theta(z)_{\text{diag}})^{-1} \right) \Pi,
$$
\n(4.85)

provided that $z > z_0$ with $z_0 > 0$ large enough.

To simplify the right side of (4.85) , we first note that the inverse $(1 + L(z))^{-1}$ is only needed on Ran Π^* , which, in view of (4.84), is left invariant by $L(z)$. Explicitly, we have that

$$
(1 + L(z))^{-1} \Pi^* = \Pi^* \left(1 + \left[U \circ (\Theta(z)_{\text{diag}})^{-1} \right] \Pi \Lambda(z)_{\text{off}} \Pi^* \right)^{-1}.
$$
 (4.86)

Secondly, by Proposition 4.8, we have the factorization property

$$
(\Lambda(z)_{\text{off}})_{\sigma\nu} = \Lambda(z)_{\sigma\nu} = |u_{\sigma}\rangle \langle v_{\nu} | \otimes \Theta(z)_{\sigma\nu}, \qquad \sigma \neq \nu,
$$
 (4.87)

where $\Theta(z)_{\sigma\nu} \in \mathscr{L}(\mathfrak{X}_{\nu}, \mathfrak{X}_{\sigma})$ for $\sigma \neq \nu$. From (4.87), it follows that

$$
\left[U \circ (\Theta(z)_{\text{diag}})^{-1}\right] \Pi \Lambda(z)_{\text{off}} \Pi^* = \tilde{U} \circ \left[(\Theta(z)_{\text{diag}})^{-1} \Theta(z)_{\text{off}} \right],\tag{4.88}
$$

where the components of $U = (U_{\sigma\nu})_{\sigma,\nu \in \mathcal{J}}$ are defined by

$$
\widetilde{U}_{\sigma\nu} := \frac{|u_{\sigma}\rangle\langle v_{\nu}|}{\langle u_{\sigma} | v_{\sigma}\rangle}
$$

and the identity $U_{\sigma\nu} = U_{\sigma\lambda} u_{\lambda} \rangle \langle v_{\nu} |$ was used. Furthermore, a direct computation, using the identity $(\tilde{U} \circ A)(\tilde{U} \circ B) = (\tilde{U} \circ (AB))$, shows that, on Ran II,

$$
\left(1+\widetilde{U}\circ\left[\left(\Theta(z)_{\text{diag}}\right)^{-1}\Theta(z)_{\text{off}}\right]\right)^{-1}=1-\widetilde{U}\circ\left[\Theta(z)^{-1}\Theta(z)_{\text{off}}\right].\tag{4.89}
$$

 $\text{Indeed, } \Theta(z)^{-1} = (1 + (\Theta(z)_{\text{diag}})^{-1} \Theta(z)_{\text{off}})^{-1} (\Theta(z)_{\text{diag}})^{-1}$ exists for large enough $z > 0$, since our analysis in Section 4.1 shows that $\lim_{z\to\infty} (\Theta(z)_{\text{diag}})^{-1} = 0$ and, by Proposition 4.5, $\|\Theta(z)_{\sigma\nu}\|$ with $\sigma \neq \nu$ and hence $\|\Theta(z)_{\text{off}}\|$ are uniformly bounded in $z > 0$. Now, (4.85) combined with (4.86), (4.88) and (4.89) yields

$$
\lim_{\varepsilon \to 0} \Lambda_{\varepsilon}(z)^{-1} = \Pi^* \left(1 + \tilde{U} \circ \left[(\Theta(z)_{\text{diag}})^{-1} \Theta(z)_{\text{off}} \right] \right)^{-1} \left(U \circ (\Theta(z)_{\text{diag}})^{-1} \right) \Pi
$$
\n
$$
= \Pi^* \left(1 - \tilde{U} \circ \left[\Theta(z)^{-1} \Theta(z)_{\text{off}} \right] \right) \left(U \circ (\Theta(z)_{\text{diag}})^{-1} \right) \Pi
$$
\n
$$
= \Pi^* \left(U \circ \left[(\Theta(z)_{\text{diag}})^{-1} - \Theta(z)^{-1} \Theta(z)_{\text{off}} (\Theta(z)_{\text{diag}})^{-1} \right] \right) \Pi
$$
\n
$$
= \Pi^* \left(U \circ \Theta(z)^{-1} \right) \Pi,
$$

where the last equation follows from the second resolvent identity and the second to last equation \Box used $U_{\sigma\lambda}U_{\lambda\nu}=U_{\sigma\nu}$.

5 Proof of Theorem 1.1. and lower bound on $\sigma(H)$

Recall from Corollary B.2 that for any $z \in \rho(H_0)$ we have that $z \in \rho(H_\varepsilon)$ if and only if $\Lambda_\varepsilon(z)$ is invertible in $\widetilde{\mathfrak{X}}$ and in this case the resolvent $(H_{\varepsilon}+z)^{-1}$ can be expressed by the Konno-Kuroda formula

$$
(H_{\varepsilon} + z)^{-1} = R_0(z) + \sum_{\sigma,\nu \in \mathcal{I}} (A_{\varepsilon,\sigma} R_0(\overline{z}))^* (\Lambda_{\varepsilon}(z)^{-1})_{\sigma\nu} J_{\nu} A_{\varepsilon,\nu} R_0(z).
$$
(5.1)

In virtue of Proposition 4.9, there exist ε_0 , $z_0 > 0$ such that $\Lambda_{\varepsilon}(z)$ is invertible for all $z > z_0$ and $\varepsilon \in (0, \varepsilon_0)$. This implies that $(z_0, \infty) \subset \rho(H_\varepsilon) \cap \rho(H_0)$ for any $\varepsilon \in (0, \varepsilon_0)$ and $(H_\varepsilon + z)^{-1}$ is then given by (5.1). Moreover, Proposition 4.9 also shows that $\lim_{\varepsilon \to 0} \Lambda_{\varepsilon}(z)^{-1}$ exists for any $z > z_0$ and, by the componentwise version (4.72) of (4.74),

$$
\lim_{\varepsilon \to 0} (\Lambda_{\varepsilon}(z)^{-1})_{\sigma \nu} = \begin{cases} \frac{|u_{\sigma}\rangle \langle v_{\nu}|}{\langle u_{\sigma} | v_{\sigma} \rangle \langle u_{\nu} | v_{\nu} \rangle} \otimes (\Theta(z)^{-1})_{\sigma \nu} & \text{if } \sigma, \nu \in \mathcal{J} \\ 0 & \text{else.} \end{cases}
$$

Here, $\Theta(z)$ is a closed and invertible operator in the reduced Hilbert space $\mathfrak X$ defined by (1.5). Since $J_{\nu} = \text{sgn}(V_{\nu})$ is bounded and since, by Proposition 3.3, $\lim_{\varepsilon \to 0} A_{\varepsilon,\nu} R_0(z) = S(z)_{\nu}$ exists for all *v*, we conclude that we can take the limit $\varepsilon \to 0$ on the right side of (5.1). We find that $\lim_{\varepsilon \to 0} (H_{\varepsilon} + z)^{-1} = R(z)$ for all $z > z_0$, where

$$
R(z) := R_0(z) + \sum_{\sigma,\nu \in \mathcal{J}} S(\overline{z})^*_{\sigma} \left(\frac{|u_{\sigma}\rangle \langle v_{\nu}|}{\langle u_{\sigma} | v_{\sigma}\rangle \langle u_{\nu} | v_{\nu}\rangle} \otimes (\Theta(z)^{-1})_{\sigma\nu} \right) J_{\nu} S(z)_{\nu}.
$$
 (5.2)

Expression (5.2) can be simplified as follows: By Proposition 3.3, we have that $S(z)_{\sigma}\psi =$ $v_{\sigma} \otimes (G(z)_{\sigma} \psi)$ with $G(z)_{\sigma} \in \mathscr{L}(\mathscr{H}, \mathfrak{X}_{\sigma})$ and, similarly, $J_{\nu}S(z)_{\nu}\psi = u_{\nu} \otimes (G(z)_{\nu}\psi)$. Hence, (5.2) takes the form

$$
R(z) = R_0(z) + \sum_{\sigma,\nu \in \mathcal{J}} G(\overline{z})^*_{\sigma} (\Theta(z)^{-1})_{\sigma\nu} G(z)_{\nu} = R_0(z) + G(\overline{z})^* \Theta(z)^{-1} G(z), \tag{5.3}
$$

where $G(z) \in \mathscr{L}(\mathscr{H}, \mathfrak{X})$ is given by $G(z)\psi = (G(z)_{\sigma}\psi)_{\sigma \in \mathcal{J}}$. This expression for $R(z)$ agrees with the right side of (1.7). Moreover, it also shows that $R(z)$ only depends on V_{σ} ($\sigma \in \mathcal{J}$) via the parameters β_{σ} that are defined by (4.33).

To prove Theorem 1.1 for $z > z_0$, it remains to show that $R(z)$ indeed defines the resolvent $(H+z)^{-1}$ of a self-adjoint operator *H*. This follows from the Trotter-Kato Theorem (see e.g. [7, Theorem 5 and $[21,$ Theorem VIII.22^{\dagger}, provided that we can show that Ran $R(z)$ is dense in \mathscr{H} . Let $T: D(H_0) \to \mathfrak{X}$ be given by $T\psi = (T_\sigma \psi)_{\sigma \in \mathcal{J}}$, where the trace $T_\sigma: D(T_\sigma) \subset \mathscr{H} \to \mathfrak{X}_\sigma$ is defined by (3.2) on the domain $D(T_{\sigma}) \supset D(H_0)$. For compactly supported, smooth functions ψ , $T_{\sigma}\psi$ agrees with (1.6), so Ker *T* contains the dense set $C_0^{\infty}(\mathbb{R}^{2N} \setminus \Gamma) \subset \mathcal{H}$, where

$$
\Gamma := \{(x_1, ..., x_N) \in \mathbb{R}^{2N} \mid \exists i \neq j \text{ with } x_i = x_j\}
$$

denotes the union of the contact planes of the *N* particles. Let $\phi \in (\text{Ran } R(z))^{\perp}$. Then, for any $\psi \in \text{Ker } T$, it follows from (5.3) and $G(z) = TR_0(z)$ that

$$
0 = \langle \phi | R(z) (H_0 + z) \psi \rangle = \langle \phi | \psi \rangle + \langle \phi | G(\overline{z})^* \Theta(z)^{-1} T \psi \rangle = \langle \phi | \psi \rangle.
$$

As Ker $T \subset \mathcal{H}$ is dense, this implies that $\phi = 0$, i.e. Ran $R(z) \subset \mathcal{H}$ is dense. Now, by the Trotter-Kato Theorem, there exists a self-adjoint operator *H* such that $R(z) = (H + z)^{-1}$

[†]In [21] existence of $\lim_{\varepsilon\to 0}(H_{\varepsilon}+z)^{-1}$ in two points *z* with $\pm \text{Im}(z) > 0$ is assumed, but the proof can be adapted to the case where the limit exists for all *z* from a non-empty open interval.

for $z > z_0$ and, moreover, $H_\varepsilon \to H$ in the norm resolvent sense. This completes the proof of Theorem 1.1 for $z > z_0$.

To prove Theorem 1.1 for all $z \in \rho(H) \cap \rho(H_0)$, it suffices to verify the hypotheses of [5, Theorem 2.19]. To this end, we first need to define $\Theta(z)$ for all $z \in \rho(H_0) = \mathbb{C} \setminus (-\infty, 0]$. For the diagonal parts $\Theta(z)_{\sigma\sigma}$, this is achieved by (4.32) with ln denoting the principal branch of the logarithm. To define the off-diagonal parts for $z \in \rho(H_0)$, we use $\Theta(z)_{\sigma\nu} := -T_\sigma G(\overline{z})^*_{\nu}$ for $\sigma \neq \nu$, which agrees with (4.50) for $z > 0$, but, a priori, may be an unbounded operator for other values of $z \in \rho(H_0)$. From $G(z)_{\nu} = T_{\nu}R_0(z)$ and from the first resolvent identity, we get

$$
G(\overline{z})_{\nu}^* = G(\overline{w})_{\nu}^* + (w - z)R_0(z)G(\overline{w})_{\nu}^*, \qquad z, w \in \rho(H_0). \tag{5.4}
$$

Choosing $w \in (0, \infty)$, we know from Proposition 4.5 that $\text{Ran } G(\overline{w})^*_{\nu} \subset D(T_{\sigma})$ and that $T_{\sigma}G(\overline{w})^*_{\nu}$ is a bounded operator. Likewise, $T_{\sigma}R_0(z)$ is a bounded operator and hence, by (5.4) , $\Theta(z)_{\sigma\nu} =$ $-T_{\sigma}G(\overline{z})^*_{\nu}$ is a bounded operator defined on \mathfrak{X}_{ν} for all $z \in \rho(H_0)$. Moreover,

$$
\Theta(z)_{\sigma\nu} = \Theta(w)_{\sigma\nu} + (z - w)G(z)_{\sigma}G(\overline{w})^*_{\nu}, \qquad \sigma \neq \nu, \quad z, w \in \rho(H_0). \tag{5.5}
$$

By Proposition 5.1, below, the extended operator $\Theta(z)$ satisfies the hypotheses of [5, Theorem 2.19], so the proof of Theorem 1.1 is complete. \square

Proposition 5.1. *Let* $w, z \in \rho(H_0)$ *. Then the operator* $\Theta(z)$ *has the following properties:*

- *(i)* $\Theta(z)^* = \Theta(\overline{z})$
- *(ii)* $\Theta(z) = \Theta(w) + (z w)G(z)G(\overline{w})^*$
- *(iii)* $0 \in \rho(\Theta(z))$ *for some* $z \in \rho(H_0)$

Remark. (*ii*) implies that $D = D(\Theta(z))$ is independent of $z \in \rho(H_0)$.

Proof. Property (*iii*) has already been verified in Proposition 4.9. To prove (*ii*), we first recall that $\Theta(z)$ and $G(z)$ have the components $\Theta(z)_{\sigma\nu}$ ($\sigma, \nu \in \mathcal{J}$) and $G(z)_{\sigma}$ ($\sigma \in \mathcal{J}$), respectively. Hence, we have to verify that

$$
\Theta(z)_{\sigma\nu} = \Theta(w)_{\sigma\nu} + (z - w)G(z)_{\sigma}G(\overline{w})^*_{\nu}, \qquad \sigma, \nu \in \mathcal{J}.
$$
 (5.6)

If $\sigma \neq \nu$, then this has been shown in (5.5). In the case of $\sigma = \nu$, we assume that $\sigma = (1,2)$ for notational simplicity. Equations (3.2) and (4.5) imply that

$$
G(z)_{\sigma} G(\overline{w})_{\sigma}^* = \tau \mathscr{K}_{\sigma} R_0(z) (\tau \mathscr{K}_{\sigma} R_0(\overline{w}))^* = \tau (\widetilde{H}_0 + z)^{-1} \left(\tau (\widetilde{H}_0 + \overline{w})^{-1} \right)^*,
$$

where H_0 is defined by (4.6). Using now the definition (3.3) of τ together with a Fourier transform in $(R, x_3, ..., x_N)$, we find that $(z-w)G(z)_{\sigma}G(\overline{w})_{\sigma}^*$ acts pointwise in $\underline{P} = (P, p_3, ..., p_N)$ by multiplication with

$$
\frac{z-w}{4\pi^2} \int\limits_{\mathbb{R}^2} \left(\frac{p^2}{\mu_{\sigma}} + Q + z\right)^{-1} \left(\frac{p^2}{\mu_{\sigma}} + Q + w\right)^{-1} dp = \frac{\mu_{\sigma}}{4\pi} \left[\ln(z+Q) - \ln(w+Q)\right],\tag{5.7}
$$

where $Q > 0$ is defined by (4.8). Now (5.6), for $\sigma = \nu = (1, 2)$, follows from (5.7) and from the definition (4.32) of $\Theta(z, \underline{P}_{\sigma})_{\sigma\sigma}$.

To prove (*i*), we use the fact that $\Theta(z)$ does not depend on the particular choices of V_{σ} , a_{σ} and b_{σ} as long as the parameters β_{σ} and the set $\mathcal J$ remain unchanged. For $\sigma \in \mathcal J$ we choose $V_{\sigma} > 0$, $a_{\sigma} = \int V_{\sigma}(r) dr/(2\pi)$ and b_{σ} to solve (4.33) and (4.34) for the given values of β_{σ} . For $\sigma \in \mathcal{I} \setminus \mathcal{J}$ we set $V_{\sigma} = 0$. Moreover, we choose $z \in (z_0, \infty)$ so that (4.74) holds. Then $u_{\sigma} = v_{\sigma}$ for all pairs $\sigma \in \mathcal{I}$ and hence $\Lambda_{\varepsilon}(z)$ is self-adjoint for all $\varepsilon > 0$. Now, it follows from (4.74), or more directly from (4.72) , that $\Theta(z)^{-1}$ is self-adjoint, too. Therefore, $\Theta(z)$ is self-adjoint for $z \in (z_0, \infty)$ and (*i*) for general $z \in \rho(H_0)$ now easily follows from (*ii*). \Box Our results on $\Theta(z)_{\sigma\nu}$ imply the following lower bound on $\sigma(H)$ (see also [6, 7]).

Proposition 5.2. Let *H* denote the Hamiltonian from Theorem 1.1. Then, with $N_{\mathcal{J}} := |\mathcal{J}|$, $\mu^- := \min_{\sigma \in \mathcal{J}} \mu_{\sigma}, \ \beta^- := \min_{\sigma \in \mathcal{J}} \beta_{\sigma} \text{ and } m := \max_{i=1,\dots,N} m_i, \text{ it holds that}$

$$
\inf \sigma(H) \ge -\exp\left(\frac{\pi m}{\mu^{-}}(N_{\mathcal{J}} - 1) - \frac{\beta^{-}}{\pi}\right). \tag{5.8}
$$

Proof. Without restriction, we can assume that $N_{\mathcal{J}} \geq 1$ because for $H = H_0$ the bound (5.8) is obvious. From [5, Theorem 2.19] and from Proposition 5.1, we know that a point $z \in \rho(H_0)$ belongs to $\rho(H)$ if and only if $\Theta(z)$ is invertible in \mathfrak{X} . Since $\Theta(z)$ is self-adjoint for $z > 0$, it suffices to show that $\Theta(z)$ is bounded from below by a positive constant for $z > z_0 :=$ $\exp(\pi m(N_{\mathcal{J}}-1)/\mu^{-}-\beta^{-}/\pi)$. To this end, we use the definition (4.32) of $\Theta(z)_{\sigma\sigma}$ and, for $\sigma \neq \nu$, we use the bound $\|\Theta(z)_{\sigma\nu}\| \leq m/4$ from Proposition 4.5. As $z > z_0$ implies that $\ln(z) + \beta^-/\pi > 0$, we find that, for all $w = (w_\sigma)_{\sigma \in \mathcal{J}} \in D(\Theta(z)),$

$$
\langle w | \Theta(z)w \rangle = \sum_{\sigma \in \mathcal{J}} \langle w_{\sigma} | \Theta(z)_{\sigma\sigma} w_{\sigma} \rangle + \sum_{\substack{\sigma, \nu \in \mathcal{J} \\ \sigma \neq \nu}} \langle w_{\sigma} | \Theta(z)_{\sigma\nu} w_{\nu} \rangle
$$

\n
$$
\geq \frac{\mu^{-}}{4\pi} \left(\ln(z) + \frac{\beta^{-}}{\pi} \right) \sum_{\sigma \in \mathcal{J}} ||w_{\sigma}||^{2} - \frac{m}{4} \sum_{\substack{\sigma, \nu \in \mathcal{J} \\ \sigma \neq \nu}} ||w_{\sigma}|| ||w_{\nu}||
$$

\n
$$
\geq \frac{\mu^{-}}{4\pi} \left(\ln(z) + \frac{\beta^{-}}{\pi} \right) ||w||^{2} - \frac{m}{8} \sum_{\substack{\sigma, \nu \in \mathcal{J} \\ \sigma \neq \nu}} (||w_{\sigma}||^{2} + ||w_{\nu}||^{2})
$$

\n
$$
= \left[\frac{\mu^{-}}{4\pi} \left(\ln(z) + \frac{\beta^{-}}{\pi} \right) - \frac{m}{4} \left(N_{\mathcal{J}} - 1 \right) \right] ||w||^{2}.
$$

The expression in brackets is positive for $z > z_0 = \exp(\pi m (N_{\mathcal{J}} - 1)/\mu^- - \beta^-/\pi)$, which proves $(5.8).$ □

6 The quadratic form of the Hamiltonian

In this section we determine the quadratic form of *H* and we show, in the case of *N* particles of mass one, that it agrees with a quadratic form introduced in [6]. This proves that *H* is the TMS Hamiltonian of Dell'Antonio et al. (see [6, Eqs. (5.3), (5.4)]).

We start by deriving an explicit formula for the quadratic form of H restricted to $D(H)$:

Lemma 6.1. *Let H* denote the self-adjoint operator from Theorem 1.1 and let $z \in \rho(H_0) \cap \rho(H)$. *Then, for any* $\psi \in D(H)$ *, it holds that*

$$
\langle \psi | H\psi \rangle = \langle \psi - G(z)^* w | (H_0 + z)(\psi - G(\overline{z})^* w) \rangle + \langle w | \Theta(z) w \rangle - z ||\psi||^2, \tag{6.1}
$$

where $w = (w_{\sigma})_{\sigma \in \mathcal{J}} \in D(\Theta(z))$ *is uniquely determined by* (1.8) *and* (1.9)*.*

Proof. By (1.10), we have that $(H + z)\psi = (H_0 + z)\psi_0$, which yields that

$$
\langle \psi | (H+z)\psi \rangle = \langle \psi | (H_0+z)\psi_0 \rangle
$$

=
$$
\langle \psi - G(z)^* w | (H_0+z)\psi_0 \rangle + \langle w | G(z)(H_0+z)\psi_0 \rangle.
$$
 (6.2)

By the definition of $G(z)$ and by (1.9), $G(z)(H_0 + z)\psi_0 = T\psi_0 = \Theta(z)w$, so it follows from (1.8) and (6.2) that

$$
\langle \psi | (H+z)\psi \rangle = \langle \psi - G(z)^* w | (H_0+z) (\psi - G(\overline{z})^* w) \rangle + \langle w | \Theta(z) w \rangle,
$$

29

which proves (6.1) .

 \Box

Next, we are going to determine an explicit description of the closure of the quadratic form from Lemma 6.1. From (4.32), it follows that $\Theta(z)_{\sigma\sigma} \geq c_1 \ln(z) + c_2$ in operator sense, where $c_1 > 0$ and $c_2 \in \mathbb{R}$ depend on the pair $\sigma \in \mathcal{J}$ but not on $z \in (0, \infty)$. Combining this with the fact that $\|\Theta(z)_{\text{off}}\|$ is uniformly bounded in $z \in (0, \infty)$, we see that $\Theta(z) \geq c > 0$ for sufficiently large $z \in \rho(H) \cap (0, \infty)$. For such z, we introduce a quadratic form q with domain

$$
D(q) := \left\{ \psi \in \mathcal{H} \, \middle| \, \exists w \in D(\Theta(z)^{1/2}) : \psi - G(z)^* w \in H^1(\mathbb{R}^{2N}) \right\}
$$

by

$$
q(\psi) := \|(H_0 + z)^{1/2}(\psi - G(z)^*w)\|^2 + \|\Theta(z)^{1/2}w\|^2 - z\|\psi\|^2,
$$
\n(6.3)

which agrees with the right side of (6.1) if $\psi \in D(H) \subset D(q)$. Similarly to [6, Lemma 3.2], one verifies that $G(z)^*w \notin H^1(\mathbb{R}^{2N})$ for any $w \in D(\Theta(z)^{1/2}) \setminus \{0\}$, so $w = w^{\psi}$ is uniquely determined by $\psi \in D(q)$ and, in particular, q is well-defined. Furthermore, it is not hard to see that *q* is bounded from below and closed. Hence, there exists a unique self-adjoint operator H_q associated with *q*. We are going to show that $H_q = H$. Let $\psi \in D(H_q)$ be fixed. Then, by the definitions of *q* and H_q , for all $\phi \in D(q)$,

$$
\langle (H_q + z)\psi | \phi \rangle = \langle (H_0 + z)^{1/2} (\psi - G(z)^* w^{\psi}) | (H_0 + z)^{1/2} (\phi - G(z)^* w^{\phi}) \rangle + \langle \Theta(z)^{1/2} w^{\psi} | \Theta(z)^{1/2} w^{\phi} \rangle.
$$
 (6.4)

Choosing $w^{\phi} = 0$, wee see that any $\phi \in H^1(\mathbb{R}^{2N})$ belongs to $D(q)$, and hence, for $\phi \in H^1(\mathbb{R}^{2N})$, Eq. (6.4) becomes

$$
\langle (H_q + z)\psi | \phi \rangle = \langle (H_0 + z)^{1/2} (\psi - G(z)^* w^{\psi}) | (H_0 + z)^{1/2} \phi \rangle.
$$
 (6.5)

This equation shows that $\psi_0 := \psi - G(z)^* w^{\psi} \in D(H_0)$, and that

$$
(H_q + z)\psi = (H_0 + z)\psi_0.
$$
\n(6.6)

We have thus verified condition (1.8) of Corollary 1.2. It remains to check condition (1.9) .

Given $w \in D(\Theta(z)^{1/2})$, we choose $\phi := G(z)^*w$, so that $\phi - G(z)^*w = 0 \in H^1(\mathbb{R}^{2N})$. It follows that $\phi \in D(q)$, and from Eqs. (6.4) and (6.6), we see that, for all $w \in D(\Theta(z)^{1/2})$,

$$
\langle \Theta(z)^{1/2} w^{\psi} | \Theta(z)^{1/2} w \rangle = \langle (H_q + z) \psi | \phi \rangle
$$

=
$$
\langle (H_0 + z) \psi_0 | G(z)^* w \rangle
$$

=
$$
\langle G(z) (H_0 + z) \psi_0 | w \rangle = \langle T \psi_0 | w \rangle.
$$

This equation implies that $w^{\psi} \in D(\Theta(z))$ and that $\Theta(z)w^{\psi} = T\psi_0$, which is condition (1.9). Corollary 1.2 now shows that $\psi \in D(H)$ and, in view of Eq. (6.6), that $H_q \subset H$. From the self-adjointness of H_q and H , we conclude that $H_q = H$.

In the case of $m_i = 1$ for $i = 1, ..., N$, we are going to show that *q* agrees with a quadratic form introduced in [6, Eqs. (2.13)-(2.16)]. For given $\beta = (\beta_{\sigma})_{\sigma \in \mathcal{I}}$, the quadratic form in [6] is denoted by F_β and, in our notation, it is defined by

$$
\mathbf{F}_{\beta}(\psi) := \|\nabla(\psi - G_z^{2N} * \xi^{\psi})\|^2 + z \|\psi - G_z^{2N} * \xi^{\psi}\|^2 + \Phi_{\beta}^{z,1}(\xi^{\psi}) + \Phi^{z,2}(\xi^{\psi}) - z \|\psi\|^2 \tag{6.7}
$$

on the domain

$$
D(\mathbf{F}_{\beta}) = \left\{ \psi \in \mathcal{H} \, \middle| \, \exists \xi^{\psi} \in D(\Phi_{\beta}^{z,1}) : \psi - G_z^{2N} * \xi^{\psi} \in H^1(\mathbb{R}^{2N}) \right\}.
$$

The right side of (6.7) is independent of the particular choice of $z \in (0, \infty)$ and $\xi^{\psi} = (\xi_{\sigma})_{\sigma \in \mathcal{J}}$ is a collection of "charges" $\xi_{\sigma} \in L^2(\mathbb{R}^{2(N-1)})$, which are uniquely determined by $\psi \in D(\mathcal{F}_{\beta})$. For $\sigma = (i, j)$, ξ_{σ} may be interpreted as a function on the hyperplane $x_i = x_j$ and the convolution $G_z^{2N} * \xi^{\psi}$ is to be understood in the sense that

$$
\widehat{G_z^{2N} * \xi^{\psi}}(p_1, ..., p_N) = \sum_{\sigma = (i,j)\in\mathcal{J}} \left(z + \sum_{n=1}^N p_n^2 \right)^{-1} \widehat{\xi_{\sigma}} \left(p_1, ..., p_{i-1}, \frac{p_i + p_j}{\sqrt{2}}, ..., \widehat{p_j} ..., p_N \right) \tag{6.8}
$$

(cf. the proof of [6, Lemma 3.2 (*a*)]). Moreover, by [6, Theorem 3.3], F_β is bounded from below and closed on $D(F_\beta)$.

Assuming that $m_i = 1$ for $i = 1, ..., N$ and that $z \in \rho(H) \cap (0, \infty)$ is so large that $\Theta(z) \geq c > 0$, we now write the various contributions to the right side of (6.3) in a form that will allow us to compare them to their counterparts in (6.7). Recall from Section 5 that $G(z) \in \mathscr{L}(\mathscr{H}, \mathfrak{X})$ is given by $G(z)\psi = (G(z)_{\sigma}\psi)_{\sigma \in \mathcal{J}}$ with $G(z)_{\sigma} = T_{\sigma}R_0(z)$. For a given $w = (w_{\sigma})_{\sigma \in \mathcal{J}} \in \mathfrak{X}$, this implies that

$$
G(z)^* w = \sum_{\sigma \in \mathcal{J}} G(z)^*_{\sigma} w_{\sigma}, \tag{6.9}
$$

where, by (4.52) for $\sigma = (i, j)$,

$$
\widehat{\left(G(z)_{\sigma}^* w_{\sigma}\right)}\left(p_1, \dots, p_N\right) = \frac{1}{2\pi} \left(z + \sum_{n=1}^N p_n^2\right)^{-1} \widehat{w_{\sigma}}\left(p_i + p_j, p_1, \dots \widehat{p}_i \dots \widehat{p}_j \dots, p_N\right). \tag{6.10}
$$

From (6.8) , (6.9) and (6.10) , we infer that

$$
G(z)^* w = G_z^{2N} * \xi, \qquad w = (w_\sigma)_{\sigma \in \mathcal{J}}, \xi = (\xi_\sigma)_{\sigma \in \mathcal{J}}, \qquad (6.11)
$$

where ξ_{σ} agrees with w_{σ} up to a permutation of the arguments and rescaling:

$$
\xi_{\sigma}(x_1, \ldots \hat{x}_j, \ldots, x_N) = \frac{1}{4\pi} w_{\sigma} \left(\frac{x_i}{\sqrt{2}}, x_1, \ldots, \hat{x}_i, \ldots, \hat{x}_j, \ldots, x_N \right), \qquad \sigma = (i, j), \qquad (6.12)
$$

which, after Fourier transform, is equivalent to

$$
\widehat{\xi_{\sigma}}(p_1, \ldots \widehat{p}_j \ldots, p_N) = \frac{1}{2\pi} \widehat{w_{\sigma}} \left(\sqrt{2}p_i, p_1, \ldots \widehat{p}_i \ldots \widehat{p}_j \ldots, p_N \right), \qquad \sigma = (i, j). \tag{6.13}
$$

Conversely, given $\xi_{\sigma} \in L^2(\mathbb{R}^{2(N-1)})$ for all $\sigma \in \mathcal{J}$, the identity (6.12) uniquely determines $w = (w_{\sigma})_{\sigma \in \mathcal{J}} \in \mathfrak{X}$ and again (6.11) holds true. In particular, it follows that

$$
||(H_0 + z)^{1/2}(\psi - G(z)^*w)||^2 = ||\nabla(\psi - G_z^{2N} * \xi)||^2 + z||\psi - G_z^{2N} * \xi||^2,
$$

provided that $\psi - G(z)^*w \in H^1(\mathbb{R}^{2N})$. The third and the fourth terms in (6.7) correspond to the diagonal and off-diagonal contributions, respectively, in

$$
\|\Theta(z)^{1/2}w\|^2 = \sum_{\sigma \in \mathcal{J}} \|(\Theta(z)_{\sigma\sigma})^{1/2}w_{\sigma}\|^2 + \sum_{\substack{\sigma,\nu \in \mathcal{J} \\ \sigma \neq \nu}} \langle w_{\sigma} | \Theta(z)_{\sigma\nu} w_{\nu} \rangle.
$$

We begin with the diagonal ones. Recall that for $\sigma = (i, j)$ the operator $\Theta(z)_{\sigma\sigma}$ acts pointwise in $\underline{P}_{\sigma} = (P, p_1, \ldots, \widehat{p}_i, \ldots, p_N)$ by multiplication with $\Theta(z, \underline{P}_{\sigma})_{\sigma\sigma}$ defined by (4.32). For given $w = (w_{\sigma})_{\sigma \in \mathcal{J}} \in D(\Theta(z)^{1/2})$, it follows that

$$
\sum_{\sigma \in \mathcal{J}} \|(\Theta(z)_{\sigma\sigma})^{1/2} w_{\sigma}\|^2 = \sum_{\sigma \in \mathcal{J}} \int d\underline{P}_{\sigma} \Theta(z, \underline{P}_{\sigma})_{\sigma\sigma} |\widehat{w_{\sigma}} (\underline{P}_{\sigma})|^2,
$$

which agrees with $\Phi_{\beta}^{z,1}(\xi)$ defined by [6, Eq. (2.15)], where $\beta = (\beta_{\sigma})_{\sigma \in \mathcal{J}}$ and $\xi \in D(\Phi_{\beta}^{z,1})$ are given by (4.33) and (6.13), respectively. Conversely, given $\xi \in D(\Phi_{\beta}^{z,1})$, a straightforward computation shows that (6.13) defines a vector $w \in D(\Theta(z)^{1/2})$ and that $\Phi_{\beta}^{z,1}(\xi)$ $\sum_{\sigma \in \mathcal{J}} ||(\Theta(z)_{\sigma\sigma})^{1/2} w_{\sigma}||^2$. In particular, we see that $w \in D(\Theta(z)^{1/2})$ if and only if (6.13) defines a vector $\xi \in D(\Phi_{\beta}^{z,1})$ and hence, using (6.11), it follows that $D(q) = D(\mathbf{F}_{\beta})$. It remains to examine the contributions of the off-diagonal operators $\Theta(z)_{\sigma\nu} \in \mathscr{L}(\mathfrak{X}_{\nu}, \mathfrak{X}_{\sigma})$. Here, the key is the identity (cf. Eq. (4.50))

$$
\Theta(z)_{\sigma\nu} = -T_{\sigma}G(z)^{*}_{\nu}, \qquad \sigma \neq \nu. \tag{6.14}
$$

Recall from (4.51) that, for $\sigma = (i, j)$ and $\psi \in D(T_{\sigma})$,

$$
(\widehat{T_{\sigma}\psi}) (P, p_1, \ldots \widehat{p_i} \ldots \widehat{p_j} \ldots, p_N) = \frac{1}{2\pi} \int dp \, \widehat{\psi} \Big(p_1, \ldots, p_{i-1}, \frac{P}{2} - p, p_{i+1}, \ldots, p_{j-1}, \frac{P}{2} + p, p_{j+1}, \ldots, p_N \Big) \,.
$$
 (6.15)

Now, using first (6.14), (6.15) and the substitution

$$
P := p_i + p_j
$$
, $p := \frac{p_j - p_i}{2}$,

and inserting the identity (6.10) for $G(z)_{\nu}^* w_{\nu}$ thereafter, we obtain that

$$
\sum_{\substack{\sigma,\nu \in \mathcal{J} \\ \sigma \neq \nu}} \langle w_{\sigma} | \Theta(z)_{\sigma\nu} w_{\nu} \rangle = -(2\pi)^{-2} \sum_{\substack{\sigma,\nu \in \mathcal{J} \\ \sigma \neq \nu}} \int d p_1 \cdots d p_N \, \overline{\hat{w}_{\sigma}} \left(p_i + p_j, p_1, \ldots \hat{p}_i \ldots \hat{p}_j \ldots, p_N \right) \cdot \left(z + \sum_{n=1}^N p_n^2 \right)^{-1} \widehat{w}_{\nu} \left(p_k + p_l, p_1, \ldots \hat{p}_k \ldots \hat{p}_l \ldots, p_N \right), \tag{6.16}
$$

where the sum runs over all $\sigma = (i, j) \neq (k, l) = \nu$. In view of (6.13), (6.16) coincides with the expression for $\Phi^{z,2}(\xi)$ from [6, Eq. (2.16)]. This completes the proof that $q = F_\beta$.

Appendix

A Properties of the Green's function

This appendix collects facts and estimates on the Green's function of $-\Delta+z : H^2(\mathbb{R}^d) \to L^2(\mathbb{R}^d)$.

For $d \in \mathbb{N}$ and $z \in \mathbb{C}$ with $\text{Re}(z) > 0$, let the function $G_z^d : \mathbb{R}^d \to \mathbb{C}$ be defined by

$$
G_z^d(x) := \int_0^\infty (4\pi t)^{-d/2} \exp\left(-\frac{x^2}{4t} - zt\right) dt.
$$
 (A.1)

Note that G_z^d has a singularity at $x = 0$ unless $d = 1$. The proof of the following lemma is left to the reader.

Lemma A.1. *Let* $d \in \mathbb{N}$ *and* $z \in \mathbb{C}$ *with* $\text{Re}(z) > 0$ *be given. Then* G_z^d *defined by* (A.1) *satisfies the following properties:*

- *(i)* G_z^d ∈ $L^1(\mathbb{R}^d)$ *and* $||G_z^d||_{L^1}$ ≤ Re(*z*)⁻¹
- *(ii)* The Fourier transform of G_z^d is given by $G_z^d(p) = (2\pi)^{-d/2}(p^2 + z)^{-1}$
- *(iii)* G_z^d *is the Green's function of* $-\Delta + z$ *, i.e.* $(-\Delta + z)^{-1}f = G_z^d * f$ *holds for all* $f \in L^2(\mathbb{R}^d)$
- (iv) $G_z^d \in L^2(\mathbb{R}^d)$ *if and only if* $d \in \{1, 2, 3\}$
- *(v)* G_z^d *is spherically symmetric, i.e.* $G_z^d(x)$ *only depends on* |*x*|*, and for* $z \in (0, \infty)$ *it is positive and strictly monotonically decreasing both as a function of* $|x|$ and z
- *(vi)* Let $d_1, d_2 \in \mathbb{N}$ with $d_1 + d_2 = d$ and decompose $x = (x_1, x_2) \in \mathbb{R}^{d_1} \times \mathbb{R}^{d_2}$. If $x_1 \neq 0$ or $d_1 = 1$, then $G_z^d(x_1, \cdot) \in L^1(\mathbb{R}^{d_2})$ and the Fourier transform with respect to x_2 is

$$
\widehat{G}_z^d(x_1, p_2) = (2\pi)^{-d_2/2} G_{z+p_2^2}^{d_1}(x_1).
$$

In particular, we have that

$$
\int_{\mathbb{R}^{d_2}} G_z^d(x_1, x_2) \, dx_2 = G_z^{d_1}(x_1).
$$

It follows from (A.1) that $G_z^d(x) \to \infty$ as $|x| \to 0$ in dimensions $d \geq 2$. However, the next lemma shows that, apart from $x = 0$, G_z^d is smooth and exponentially decaying as $|x| \to \infty$:

Lemma A.2. Let $d \in \mathbb{N}$ and $z \in \mathbb{C}$ with $\text{Re}(z) > 0$. Then G_z^d is smooth in $\mathbb{R}^d \setminus \{0\}$ and, for *any multi-index* $\alpha \in \mathbb{N}_0^d$, $\partial^{\alpha} G_z^d$ *is exponentially decaying as* $|x| \to \infty$ *.*

Proof. First, we claim that, for all $c > 0$, $k \in \mathbb{R}$, $l \geq 0$ and $z' \in (0, \sqrt{\text{Re}(z)})$,

$$
\sup_{|x|\geq c} \left(\int_0^\infty t^k |x|^l \exp\left(-\frac{x^2}{4t} - \text{Re}(z)t + z'|x|\right) dt \right) < \infty.
$$
 (A.2)

To prove this, we choose $\delta > 0$ so small that $z' \leq (1 - \delta)\sqrt{\text{Re}(z)}$. Then the estimate

$$
\exp\left(-\frac{x^2}{4t} - \text{Re}(z)t + z'|x|\right) \le \exp\left(-\frac{x^2}{4t} - \text{Re}(z)t + (1 - \delta)\sqrt{\text{Re}(z)}|x|\right)
$$

$$
= \exp\left(-(1 - \delta)\left(\frac{|x|}{2\sqrt{t}} - \sqrt{\text{Re}(z)t}\right)^2 - \delta\left(\frac{x^2}{4t} + \text{Re}(z)t\right)\right)
$$

$$
\le \exp\left(-\frac{\delta x^2}{8t} - \frac{\delta c^2}{8t} - \delta \text{Re}(z)t\right)
$$

 $\text{holds for all } t > 0 \text{ and } |x| \geq c.$ With the help of $C_{l,\delta} := \sup_{s \geq 0} (s^{l/2} \exp(-\delta s)) < \infty$, (A.2) now follows from

$$
\sup_{|x|\geq c} \left(\int_{0}^{\infty} t^{k} |x|^l \exp\left(-\frac{x^2}{4t} - \text{Re}(z)t + z'|x| \right) dt \right)
$$

\n
$$
\leq 8^{l/2} \sup_{|x|\geq c} \left(\int_{0}^{\infty} t^{k+l/2} \left(\frac{x^2}{8t} \right)^{l/2} \exp\left(-\frac{\delta x^2}{8t} - \frac{\delta c^2}{8t} - \delta \text{ Re}(z)t \right) dt \right)
$$

\n
$$
\leq 8^{l/2} C_{l,\delta} \int_{0}^{\infty} t^{k+l/2} \exp\left(-\frac{\delta c^2}{8t} - \delta \text{ Re}(z)t \right) dt < \infty,
$$

where the last integral is finite because the term $\exp(-\delta c^2/(8t))$ cancels a possible divergence at $t = 0$. (A.2) allows us to change the order of differentiation and integration in the representation $(A.1)$ of G_z^d . We find that, for $x \neq 0$, $\partial^{\alpha} G_z^d(x)$ is a sum of terms of the form

$$
\int_{0}^{\infty} (4\pi t)^{-d/2} \frac{x^{\alpha'}}{(2t)^n} \exp\left(-\frac{x^2}{4t} - zt\right) dt,
$$

where $0 \le n \le |\alpha|$ and $\alpha' \in \mathbb{N}_0^d$ is a multi-index with $|\alpha'| \le |\alpha|$. In view of $(A.2)$, all of these terms are exponentially decaying as $|x| \to \infty$. \Box

In the proof of Proposition 4.4 two integral operators F and B depending on G_z^d were introduced. We close this section with the bypassed proof of their boundedness:

Lemma A.3. *Let* $z, m > 0$. *Then the operator* $F: L^2(\mathbb{R}^2 \times \mathbb{R}^2) \to L^2(\mathbb{R}^2 \times \mathbb{R}^2)$ *that is defined in terms of the kernel*

$$
K_F(x, y; x', y') := G_z^6(\sqrt{m}(x - x'), \sqrt{m}(x - y'), \sqrt{m}(y - x'))
$$

is bounded with $||F|| \leq (4\sqrt{2}m^2)^{-1}$ *. Similarly, the kernel*

$$
K_B(x, y, w; x', y', w') := G_z^8(\sqrt{m}(x - y'), \sqrt{m}(x - w'), \sqrt{m}(y - x'), \sqrt{m}(w - x'))
$$

defines an operator $B \in \mathcal{L}(L^2(\mathbb{R}^2 \times \mathbb{R}^2 \times \mathbb{R}^2))$ *with* $||B|| \leq (4\sqrt{2}m^3)^{-1}$ *.*

Proof. The bounds for $||F||$ and $||B||$ are both based on the Schur test. In the case of *F*, the Schur test, in a general version, yields that

$$
||F|| \le \underset{x,y}{\text{ess sup}} \left(h(x,y)^{-1} \int dx' dy' h(x',y') K_F(x,y;x',y') \right) \tag{A.3}
$$

for any measurable test function $h : \mathbb{R}^2 \times \mathbb{R}^2 \to (0, \infty)$, provided that the right side is finite. We choose the test function $h(x, y) := |x - y|^{-1}$, so after scaling we may assume that $m = 1$. To evaluate the right side of (A.3), we insert the integral representation (A.1) of the Green's function G_z^6 and we substitute $x - y' \rightarrow y'$. This results in the estimate

$$
\begin{split}\n&\underset{x,y}{\text{ess sup}}\left(|x-y|\int \mathrm{d}x' \, \mathrm{d}y' \, |x'-y'|^{-1} \, G_z^6 \left(x-x', x-y', y-x'\right)\right) \\
&= \underset{x,y}{\text{ess sup}}\left(|x-y|\int \mathrm{d}x' \int_0^\infty \mathrm{d}t \, (4\pi t)^{-3} \exp\left(-\frac{1}{4t}\left((x-x')^2+(y-x')^2\right)-zt\right) \\
&\cdot \int \mathrm{d}y' \, |y'+x'-x|^{-1} \exp\left(-\frac{|y'|^2}{4t}\right)\right).\n\end{split} \tag{A.4}
$$

Using a rearrangement inequality (see [15, Theorem 3.4]), the last integral is bounded by

$$
\int dy' |y' + x' - x|^{-1} \exp\left(-\frac{|y'|^2}{4t}\right) \le \int dy' |y'|^{-1} \exp\left(-\frac{|y'|^2}{4t}\right)
$$

$$
= 2\pi \int_0^\infty ds \exp\left(-\frac{s^2}{4t}\right) = \sqrt{4\pi^3 t}.
$$
 (A.5)

For the remaining integral, we use $(x - x')^2 + (y - x')^2 = \frac{1}{2}$ 2 $((2x'-x-y)^2 + (x-y)^2)$ together with the substitution $2x' - x - y \rightarrow 2x'$. This yields that

$$
\pi \operatorname{ess} \sup_{x,y} \left(|x - y| \int dx' \int_0^{\infty} dt \, (4\pi t)^{-5/2} \exp\left(-\frac{1}{4t} \left((x - x')^2 + (y - x')^2 \right) - zt \right) \right)
$$
\n
$$
\leq \pi \operatorname{ess} \sup_{x,y} \left(|x - y| \int dx' \int_0^{\infty} dt \, (4\pi t)^{-5/2} \exp\left(-\frac{|x'|^2}{2t} - \frac{(x - y)^2}{8t} \right) \right)
$$
\n
$$
= (16\sqrt{\pi})^{-1} \operatorname{ess} \sup_{x,y} \left(|x - y| \int_0^{\infty} dt \, t^{-3/2} \exp\left(-\frac{(x - y)^2}{8t} \right) \right)
$$
\n
$$
= (4\sqrt{2\pi})^{-1} \int_0^{\infty} dt \, t^{-3/2} \exp\left(-t^{-1} \right). \tag{A.6}
$$

In the last equality, the ess sup cancels after the scaling $t \to (x - y)^2 t/8$. The remaining integral can be explicitly computed with the help of the substitution $\tilde{t} := t^{-1}$. We find that

$$
\int_{0}^{\infty} dt \ t^{-3/2} \exp\left(-t^{-1}\right) = \int_{0}^{\infty} d\tilde{t} \ \tilde{t}^{-1/2} \exp\left(-\tilde{t}\right) = \Gamma(1/2) = \sqrt{\pi},\tag{A.7}
$$

where Γ denotes the Gamma function. Using $(A.5)$, $(A.6)$ and $(A.7)$ to bound the right side of (A.4), the Schur test yields that *F* is bounded and its norm satisfies the desired estimate.

The proof in case of the operator *B* is similar, but for the convenience of the reader we go into details here. First, applying the Schur test with the test function $h(y, w) = |y - w|^{-1}$ gives

$$
||B|| \le \underset{x,y,w}{\text{ess sup}} \left(|y-w| \int \! \mathrm{d}x' \, \mathrm{d}y' \, \mathrm{d}w' \, |y'-w'|^{-1} K_B(x,y,w;x',y',w') \right),
$$

provided that the right side is finite. After scaling, we may again assume that $m = 1$. Next, inserting the integral formula (A.1) for G_z^8 and substituting $x - y' \rightarrow y'$, it follows that

$$
||B|| \le \underset{x,y,w}{\text{ess sup}} \left(|y-w| \int dx' \, dw' \int_0^\infty dt \, (4\pi t)^{-4} \exp\left(-\frac{1}{4t} \left[(x-w')^2 + (y-x')^2 + (w-x')^2 \right] \right) \cdot \int dy' \, |x-y'-w'|^{-1} \exp\left(-\frac{|y'|^2}{4t} \right) \right). \tag{A.8}
$$

Similarly to (A.5), a rearrangement inequality now yields that

$$
\int dy' |x - y' - w'|^{-1} \exp\left(-\frac{|y'|^2}{4t}\right) \le \sqrt{4\pi^3 t}.
$$
 (A.9)

In the remaining integral, we use $(y - x')^2 + (w - x')^2 = \frac{1}{2}$ 2 $((2x'-w-y)^2+(y-w)^2)$ in combination with the substitutions $2x' - w - y \rightarrow 2x'$ and $x - w' \rightarrow w'$. This leads to

$$
\pi \operatorname{ess} \sup_{x,y,w} \left(|y-w| \int dx' dw' \int_0^{\infty} dt \ (4\pi t)^{-7/2} \exp\left(-\frac{1}{4t} \left[(x-w')^2 + (y-x')^2 + (w-x')^2 \right] \right) \right)
$$
\n
$$
= \pi \operatorname{ess} \sup_{x,y,w} \left(|y-w| \int dx' dw' \int_0^{\infty} dt \ (4\pi t)^{-7/2} \exp\left(-\frac{1}{4t} \left(|w'|^2 + 2|x'|^2 \right) - \frac{(y-w)^2}{8t} \right) \right)
$$
\n
$$
= (16\sqrt{\pi})^{-1} \operatorname{ess} \sup_{y,w} \left(|y-w| \int_0^{\infty} dt \ t^{-3/2} \exp\left(-\frac{(y-w)^2}{8t}\right) \right)
$$
\n
$$
= (4\sqrt{2\pi})^{-1} \int_0^{\infty} dt \ t^{-3/2} \exp\left(-t^{-1}\right), \tag{A.10}
$$

where the ess sup vanishes after the scaling $t \to (y-w)^2 t/8$. By (A.7), the last integral in (A.10) is equal to $\sqrt{\pi}$, so the desired estimate for $||B||$ follows from (A.8), (A.9) and (A.10). \Box

B Konno-Kuroda formula

In this section we sketch the proof of the Konno-Kuroda resolvent identity, see [13, Eq. (2.3)], for operators of the type (B.1). The main difference between Theorem B.1 below and [13] is that we do not assume that $\phi(z)$ defined by (B.2) extends to a compact operator for some (and hence all) $z \in \rho(H_0)$.

Let H and $\widetilde{\mathfrak{X}}$ be arbitrary complex Hilbert spaces, let $H_0 \geq 0$ be a self-adjoint operator in *H* and let *A* : *D*(*A*) ⊂ *H* → **X** be densely defined and closed with *D*(*A*) ⊃ *D*(*H*₀). Let $J \in \mathscr{L}(\widetilde{\mathfrak{X}})$ be a self-adjoint isometry and let $B = JA$. Suppose that $BD(H_0) \subset D(A^*)$ and that *A*[∗]*A* and *A*[∗]*B* are *H*₀-bounded with relative bound less than one. Then

$$
H = H_0 - A^*B \tag{B.1}
$$

is self-adjoint on $D(H_0)$ by the Kato-Rellich Theorem. For $z \in \rho(H_0)$, let $R_0(z) := (H_0 + z)^{-1}$ and let $\phi(z) : D(A^*) \subset \widetilde{\mathfrak{X}} \to \widetilde{\mathfrak{X}}$ be defined by

$$
\phi(z) := BR_0(z)A^*.
$$
\n(B.2)

Note that $D(A^*) \subset \tilde{\mathfrak{X}}$ is dense because *A* is closed. The resolvent $(H + z)^{-1}$ and the operator $\phi(z)$ are related by the following theorem:

Theorem B.1. Let the above hypotheses be satisfied and let $z \in \rho(H_0)$. Then $\phi(z)$ is a bounded *operator. The operator* $1 - \phi(z)$ *is invertible if and only if* $z \in \rho(H_0) \cap \rho(H)$ *, and then*

$$
(H+z)^{-1} = R_0(z) + R_0(z)A^*(1 - \phi(z))^{-1}BR_0(z)
$$
 (B.3)

$$
(1 - \phi(z))^{-1} = 1 + B(H + z)^{-1}A^*.
$$
 (B.4)

Remark. Note that $(1 - \phi(z))^{-1}$ leaves $D(A^*)$ invariant. This follows from (B.4) and from the assumption $BD(H_0) \subset D(A^*)$.

Proof.

Step 1. $AR_0(c)^{1/2}$ is bounded for $c > 0$, and $A(H + c)^{-1/2}$ is bounded for $c > 0$ large enough.

As A^*A is H_0 -bounded with relative bound less than one, the operator $H_0 - A^*A$ is bounded from below. This implies that, for all $\psi \in D(H_0)$ and all $c > 0$,

$$
||A\psi||^2 \le ||(H_0 + c)^{1/2}\psi||^2 + C||\psi||^2
$$

with some constant $C > 0$. Due to the fact that $D(H_0)$ is dense in $D(H_0^{1/2})$ $\binom{1}{0}$ and A is closed, this bound extends to all $\psi \in D(H_0^{1/2})$ $\binom{1}{0}$ by an approximation argument. In particular, $D(A) \supset$ $D(H_0^{1/2}$ $\binom{1}{0}$ and the first statement of Step 1 follows. The second statement is a consequence of the first and the fact that *H* and H_0 have equivalent form norms, which implies that $D(H^{1/2}) =$ $D(H_0^{1/2})$ $\binom{1}{0}$.

Step 2. If $z \in \rho(H_0)$, then $\phi(z)$ is a bounded operator, and if $z \in \rho(H)$, then

$$
M(z) := B(H+z)^{-1}A^*
$$

is a bounded operator too.

This easily follows from Step 1 and from the first resolvent identity.

Step 3. If $z \in \rho(H_0) \cap \rho(H)$, then $(1 - \phi(z))$ is invertible and $1 + M(z) = (1 - \phi(z))^{-1}$.

Both $\phi(z)$ and $M(z)$ leave $D(A^*)$ invariant and on this subspace, by straightforward computations using the second resolvent identity, $(1 - \phi(z))(1 + M(z)) = 1 = (1 + M(z))(1 - \phi(z)).$ *Step 4.* If $z \in \rho(H_0)$ and $1 - \phi(z)$ is invertible, then $z \in \rho(H)$, $(1 - \phi(z))^{-1}$ leaves $D(A^*)$ invariant and (B.3) holds.

By Step 3, $(1 - \phi(i))^{-1} = 1 + M(i)$, which leaves $D(A^*)$ invariant. Now suppose that $z \in \rho(H_0)$ and that $1 - \phi(z)$ has a bounded inverse. Then

$$
(1 - \phi(z))^{-1} = (1 - \phi(i))^{-1} + (1 - \phi(i))^{-1}(\phi(z) - \phi(i))(1 - \phi(z))^{-1}
$$

= $(1 - \phi(i))^{-1} + (i - z)(1 - \phi(i))^{-1}BR_0(i)(AR_0(\bar{z}))^*(1 - \phi(z))^{-1}.$

Since $BR_0(i): \mathcal{H} \to D(A^*)$, it follows that $(1 - \phi(z))^{-1}$ leaves $D(A^*)$ invariant as well, and

$$
R(z) := R_0(z) + R_0(z)A^*(1 - \phi(z))^{-1}BR_0(z)
$$

is well-defined. Now it is a matter of straightforward computations to show that $(H+z)R(z) = 1$ on $\mathscr H$ and that $R(z)(H+z) = 1$ on $D(H)$. \Box

Assume that $\widetilde{\mathfrak{X}} = \bigoplus_{i=1}^{I} \widetilde{\mathfrak{X}}_i$, where $I \in \mathbb{N}$ and $\widetilde{\mathfrak{X}}_i$ are complex Hilbert spaces for $i = 1, ..., I$. Then we consider operators of the more general form

$$
H = H_0 - \sum_{i=1}^{I} g_i A_i^* J_i A_i,
$$
 (B.5)

where $0 \neq g_i \in \mathbb{R}$, $A_i: D(A_i) \subset \mathscr{H} \to \tilde{\mathfrak{X}}_i$ are densely defined and closed with $D(A_i) \supset D(H_0)$ and *J_i* are self-adjoint isometries in $\tilde{\mathfrak{X}}_i$ for $i = 1, ..., I$. Suppose that $J_iA_iD(H_0) \subset D(A_i^*)$ and that $A_i^* A_i$ and $A_i^* J_i A_i$ are infinitesimally H_0 -bounded for $i = 1, ..., I$, which ensures that *H* is self-adjoint on $D(H_0)$. Now, observe that operators of the form $(B.5)$ can be reduced to the form (B.1), where $A: D(A) \subset \mathscr{H} \to \tilde{\mathfrak{X}}$ is the closure of the operator $A_0: D(H_0) \to \tilde{\mathfrak{X}}$ defined by $A_0\psi = (|g_i|^{1/2}A_i\psi)_{i=1}^I$ and $J \in \mathscr{L}(\tilde{\mathfrak{X}})$ is defined by $J(\psi_i)_{i=1}^I = (sgn(g_i)J_i\psi_i)_{i=1}^I$. Hence, Theorem B.1 implies the following corollary:

Corollary B.2. Let the above hypotheses be satisfied and let $z \in \rho(H_0)$. Then

$$
\Lambda(z)_{ij} := g_i^{-1} \delta_{ij} - J_i A_i R_0(z) A_j^*, \qquad i, j = 1, ..., N
$$

defines a bounded operator $\Lambda(z) = (\Lambda(z)_{ij})_{i,j=1}^I$ *in* $\widetilde{\mathfrak{X}}$ *. Moreover,* $\Lambda(z)$ *is invertible if and only if* $z \in \rho(H_0) \cap \rho(H)$ *, and then*

$$
(H+z)^{-1} = R_0(z) + \sum_{i,j=1}^{I} (A_i R_0(\overline{z}))^* (\Lambda(z)^{-1})_{ij} J_j A_j R_0(z),
$$

$$
\Lambda(z)^{-1} = (g_i \delta_{ij} + g_i g_j J_i A_i (H+z)^{-1} A_j^*)_{i,j=1}^{I}.
$$

Acknowledgement. Initially, Ulrich Linden was involved in the project as well. He was the first of us to sit down and do some preliminary computations and estimates in the special case of a 2d Bose gas. This helped tremendously to get the project off the ground. - M. H. was supported partially by the *Deutsche Forschungsgemeinschaft (DFG)* through the Research Training Group 1838: *Spectral Theory and Dynamics of Quantum Systems*.

References

- [1] Milton Abramowitz and Irene A. Stegun. *Handbook of mathematical functions with formulas, graphs, and mathematical tables*, volume 55 of *National Bureau of Standards Applied Mathematics Series*. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C., 1964.
- [2] S. Albeverio, F. Gesztesy, R. Høegh-Krohn, and H. Holden. *Solvable models in quantum mechanics*. AMS Chelsea Publishing, Providence, RI, second edition, 2005. With an appendix by Pavel Exner.
- [3] Giulia Basti, Claudio Cacciapuoti, Domenico Finco, and Alessandro Teta. The threebody problem in dimension one: from short-range to contact interactions. *J. Math. Phys.*, 59(7):072104, 18, 2018.
- [4] Giulia Basti, Claudio Cacciapuoti, Domenico Finco, and Alessandro Teta. Three-body hamiltonian with regularized zero-range interactions in dimension three, 2021.
- [5] Claudio Cacciapuoti, Davide Fermi, and Andrea Posilicano. On inverses of Krein's Qfunctions. *Rend. Mat. Appl.(7)*, 39:229–240, 2018.
- [6] G. F. Dell'Antonio, R. Figari, and A. Teta. Hamiltonians for systems of *N* particles interacting through point interactions. *Ann. Inst. H. Poincaré Phys. Théor.*, 60(3):253–290, 1994.
- [7] J. Dimock and S. G. Rajeev. Multi-particle Schrödinger operators with point interactions in the plane. *J. Phys. A*, 37(39):9157–9173, 2004.
- [8] Fatih Erman and O Teoman Turgut. A many-body problem with point interactions on twodimensional manifolds. *Journal of Physics A: Mathematical and Theoretical*, 46(5):055401, 2013.
- [9] M. Griesemer and U. Linden. Spectral Theory of the Fermi Polaron. *Ann. Henri Poincaré*, 20(6):1931–1967, 2019.
- [10] Marcel Griesemer and Michael Hofacker. On the weakness of short-range interactions in Fermi gases. *in preparation*, 2021.
- [11] Marcel Griesemer, Michael Hofacker, and Ulrich Linden. From short-range to contact interactions in the 1d Bose gas. *Math. Phys. Anal. Geom.*, 23(2):Paper No. 19, 28, 2020.
- [12] Yu Gu, Jeremy Quastel, and Li-Cheng Tsai. Moments of the 2d she at criticality. *Probability and Mathematical Physics*, 2(1):179–219, 2021.
- [13] Reiji Konno and S. T. Kuroda. On the finiteness of perturbed eigenvalues. *J. Fac. Sci. Univ. Tokyo Sect. I*, 13:55–63 (1966), 1966.
- [14] E. H. Lieb and W. Liniger. Exact analysis of an interacting bose gas i. the general solution and the ground state. *Phys. Rev. 130*, 39:1605–1616, 1963.
- [15] Elliott H. Lieb and Michael Loss. *Analysis*, volume 14 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, second edition, 2001.
- [16] R. A. Minlos and L. D. Faddeev. Comment on the problem of three particles with point interactions. *Soviet Physics JETP*, 14:1315–1316, 1962.
- [17] R. A. Minlos and L. D. Faddeev. On the point interaction for a three-particle system in quantum mechanics. *Soviet Physics Dokl.*, 6:1072–1074, 1962.
- [18] Meera M. Parish and Jesper Levinsen. Highly polarized fermi gases in two dimensions. *Phys. Rev. A*, 87:033616, Mar 2013.
- [19] Andrea Posilicano. A Krein-like formula for singular perturbations of self-adjoint operators and applications. *J. Funct. Anal.*, 183(1):109–147, 2001.
- [20] Andrea Posilicano. Self-adjoint extensions of restrictions. *Oper. Matrices*, 2(4):483–506, 2008.
- [21] M. Reed and B. Simon. *Methods of modern mathematical physics. I*. Academic Press, Inc. [Harcourt Brace Jovanovich, Publishers], New York, second edition, 1980. Functional analysis.
- [22] Robert Seiringer and Jun Yin. The Lieb-Liniger model as a limit of dilute bosons in three dimensions. *Comm. Math. Phys.*, 284(2):459–479, 2008.
- [23] Barry Simon. The bound state of weakly coupled Schrödinger operators in one and two dimensions. *Ann. Physics*, 97(2):279–288, 1976.
- [24] Gerald Teschl. *Mathematical methods in quantum mechanics*, volume 99 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2009. With applications to Schrödinger operators.
- [25] U Van Kolck. Effective field theory of short-range forces. *Nuclear Physics A*, 645(2):273– 302, 1999.