

# WHEN IS A REDUCTIVE GROUP SCHEME LINEAR?

PHILIPPE GILLE

ABSTRACT. We show that a reductive group scheme over a base scheme  $S$  admits a faithful linear representation if and only if its radical torus is isotrivial; that is, it splits after a finite étale cover.

*to the 75-th anniversary of Gopal Prasad*

*Keywords:* Reductive group schemes, representations, tori, resolution property.

*MSC 2000:* 14L15, 20G35

## 1. INTRODUCTION

Let  $S$  be a scheme. Let  $\mathfrak{G}$  be an  $S$ -group scheme. It is natural to ask whether  $\mathfrak{G}$  is linear; that is, there exists a group monomorphism  $\mathfrak{G} \rightarrow \mathrm{GL}(\mathcal{E})$  where  $\mathcal{E}$  is a locally free  $\mathcal{O}_S$ -module of finite rank. In particular,  $\mathfrak{G}$  admits a faithful representation on  $\mathcal{E}$ . This holds for affine algebraic groups over a field [4, II, §2.3.3].

In the case  $S$  is locally noetherian and  $G$  is of multiplicative type of constant type and of finite type, Grothendieck has shown that  $G$  is linear if and if  $G$  is isotrivial, i.e.  $G$  is split by a finite étale extension of  $S$  [11, XI.4.6]. In particular there exist non linear tori of rank 2 over the local ring (at a node) of a nodal algebraic curve (*ibid*, X.1.6). Firstly we extend that criterion over an arbitrary base by using Azumaya and separable  $\mathcal{O}_S$ -algebras (Theorem 3.3).

Secondly we deal with the case  $\mathfrak{G}$  reductive; that is,  $\mathfrak{G}$  is smooth affine with reductive (connected) geometric fibers. In this case a faithful representation is necessarily a closed immersion [11, XVI.1.5]. Positive results on the linearity question are due to M. Raynaud [11, VI<sub>B</sub>] and R. Thomason [13, 3.1] which is essentially the implication (i)  $\implies$  (ii) in the theorem below.

We can restrict our attention to the case when  $\mathfrak{G}$  is of constant type (recall that the type is a locally constant function on  $S$ ); this implies that there exists a Chevalley  $\mathbb{Z}$ -group scheme  $G$  such that  $\mathfrak{G}$  is locally isomorphic to  $G_S$  for the étale topology [11, XXII.2.3, 2.5]. A short version of our main result is the following.

---

*Date:* April 23, 2021.

The author is supported by the project ANR Geolie, ANR-15-CE 40-0012, (The French National Research Agency).

**Theorem 1.1.** *The following are equivalent:*

- (i) *The radical torus  $\text{rad}(\mathfrak{G})$  is isotrivial;*
- (ii)  *$\mathfrak{G}$  is linear.*

*Furthermore if  $S$  is affine, the above are equivalent to*

- (ii') *there exists a closed immersion  $i : \mathfrak{G} \rightarrow \text{GL}_n$  with  $n \geq 1$  which is a homomorphism.*

We recall that  $\text{rad}(\mathfrak{G})$  is the maximal central subtorus of  $\mathfrak{G}$  [11, XXIV.4.3.6] and that (i) means that  $\text{rad}(\mathfrak{G})$  splits after passing to a finite étale cover  $S' \rightarrow S$ . In the noetherian setting, a variant of the implication (i)  $\implies$  (ii) has been shown by Margaux who furthermore provided an  $\text{Aut}(\mathfrak{G})$ -equivariant representation [9]. Note that condition (i) depends only on the quasi-split form of  $\mathfrak{G}$  and also that it is always satisfied if  $\mathfrak{G}$  is semisimple or if  $\text{rad}(\mathfrak{G})$  is of rank one. Furthermore, if  $S$  is a semilocal scheme, Demazure's characterization of isotrivial group schemes [11, XXIV.3.5] permits us to deduce that  $\mathfrak{G}$  is isotrivial if and only if  $\mathfrak{G}$  is linear, see section 5.

Finally for  $S = \text{Spec}(R)$  with  $R$  noetherian, we complete Thomason's approach by showing that linearity for  $\mathfrak{G}$  is equivalent to the resolution property (Th. 6.2).

**Acknowledgements.** I thank Vladimir Chernousov, Laurent Moret-Bailly, Erhard Neher, Arturo Pianzola, and Anastasia Stavrova for their valuable suggestions. I thank the referee for a simplification of the proof of Proposition 3.2.

## 2. DEFINITIONS AND BASIC FACTS

**2.1. Notation.** We use mainly the terminology and notation of Grothendieck-Dieudonné [5, §9.4 and 9.6] which agrees with that of Demazure-Grothendieck used in [11, Exp. I.4].

(a) Let  $S$  be a scheme and let  $\mathcal{E}$  be a quasi-coherent sheaf over  $S$ . For each morphism  $f : T \rightarrow S$ , we denote by  $\mathcal{E}_T = f^*(\mathcal{E})$  the inverse image of  $\mathcal{E}$  by the morphism  $f$ . We denote by  $\mathbf{V}(\mathcal{E})$  the affine  $S$ -scheme defined by  $\mathbf{V}(\mathcal{E}) = \text{Spec}(\text{Sym}^\bullet(\mathcal{E}))$ ; it represents the  $S$ -functor  $Y \mapsto \text{Hom}_{\mathcal{O}_Y}(\mathcal{E}_Y, \mathcal{O}_Y)$  [5, 9.4.9].

(b) We assume now that  $\mathcal{E}$  is locally free of finite rank and denote by  $\mathcal{E}^\vee$  its dual. In this case the affine  $S$ -scheme  $\mathbf{V}(\mathcal{E})$  is of finite presentation (ibid, 9.4.11); also the  $S$ -functor  $Y \mapsto H^0(Y, \mathcal{E}_Y) = \text{Hom}_{\mathcal{O}_Y}(\mathcal{O}_Y, \mathcal{E}_Y)$  is representable by the affine  $S$ -scheme  $\mathbf{V}(\mathcal{E}^\vee)$  which is also denoted by  $\mathbf{W}(\mathcal{E})$  [11, I.4.6].

The above applies to the locally free quasi-coherent sheaf  $\mathcal{E}nd(\mathcal{E}) = \mathcal{E}^\vee \otimes_{\mathcal{O}_S} \mathcal{E}$  over  $S$  so that we can consider the affine  $S$ -scheme  $\mathbf{V}(\mathcal{E}nd(\mathcal{E}))$  which is an  $S$ -functor in associative and unital algebras [5, 9.6.2]. Now we consider the  $S$ -functor  $Y \mapsto \text{Aut}_{\mathcal{O}_Y}(\mathcal{E}_Y)$ . It is representable by an open  $S$ -subscheme of  $\mathbf{V}(\mathcal{E}nd(\mathcal{E}))$  which is denoted by  $\text{GL}(\mathcal{E})$  (*loc. cit.*, 9.6.4).

(c) If  $\mathcal{B}$  is a locally free  $\mathcal{O}_S$ -algebra (unital, associative) of finite rank, we recall that the functor of invertible elements of  $\mathcal{B}$  is representable by an affine  $S$ -group scheme which is denoted by  $\mathrm{GL}_1(\mathcal{B})$  [3, 2.4.2.1].

For separable and Azumaya algebras, we refer to [8]. Note that in [3, §2.5.1], separable algebras are supposed furthermore to be locally free of finite rank.

If  $\mathcal{B}$  is a separable  $\mathcal{O}_S$ -algebra which is a locally free  $\mathcal{O}_S$ -algebra of finite rank, then  $\mathrm{GL}_1(\mathcal{B})$  it is a reductive  $S$ -group scheme [3, 3.1.0.50].

(d) We use the theory and terminology of tori and multiplicative group schemes of [11]; see also Oesterlé's survey [10].

**2.2. Finite étale covers.** The next lemma is a consequence of the equivalence of categories describing finite étale  $\mathcal{O}_S$ -algebra of rank  $N$  [3, §2.5.2]; it admits a simple direct proof.

**Lemma 2.1.** *Let  $N$  be a positive integer and let  $\mathcal{C}$  be a finite étale  $\mathcal{O}_S$ -algebra of rank  $N$ . Then there exists a finite étale cover  $T$  of  $S$  of degree  $N!$  such that  $\mathcal{C} \otimes_{\mathcal{O}_S} \mathcal{O}_T \xrightarrow{\sim} (\mathcal{O}_T)^N$ .*

*Proof.* We proceed by induction on  $N$ , the case  $N = 1$  being obvious. We put  $S' = \mathrm{Spec}(\mathcal{O}_{\mathcal{C}})$ , this is a finite étale cover of  $S$  of degree  $N$ . Since the diagonal map  $S' \rightarrow S' \times_S S'$  is closed and open [6, 4.17.4.2] there exists a decomposition  $\mathcal{C} \otimes_{\mathcal{O}_S} \mathcal{O}'_S = \mathcal{O}_{S'} \times \mathcal{C}'$  where  $\mathcal{C}'$  is a finite étale  $\mathcal{O}_{S'}$ -algebra of rank  $N - 1$ . Applying the induction process to  $\mathcal{C}'$  provides a finite étale cover  $T$  of  $S'$  of degree  $(N - 1)!$  such that  $\mathcal{C}' \otimes_{\mathcal{O}_{S'}} \mathcal{O}_T \cong (\mathcal{O}_T)^{N-1}$ . Thus  $\mathcal{C} \otimes_{\mathcal{O}_S} \mathcal{O}_T = \mathcal{O}_T \times (\mathcal{O}_T)^{N-1}$  and  $T$  is a finite étale cover of  $S$  of degree  $N! = N \times (N - 1)!$ .  $\square$

**2.3. Isotriviality.** [11, XXIV.4] Let  $\mathcal{H}$  be a fppf  $S$ -sheaf in groups and let  $\mathcal{X}$  be a  $\mathcal{H}$ -torsor. We say that  $\mathcal{X}$  is *isotrivial* if there exists a finite étale cover  $S'$  of  $S$  which trivializes  $\mathcal{X}$ ; that is, satisfying  $\mathcal{X}(S') \neq \emptyset$ .

The notion of locally isotrivial (with respect to the Zariski topology) is then clear and there is also the following variant of *semilocally isotrivial*.

We say that  $\mathcal{X}$  is *semilocally isotrivial* if for each subset  $\{s_1, \dots, s_n\}$  of points of  $S$  contained in an affine open subset of  $S$ , there exists an open subscheme  $U$  of  $S$  containing  $s_1, \dots, s_n$  such that  $\mathcal{X} \times_S U$  is isotrivial over  $U$ .

A reductive  $S$ -group scheme  $\mathfrak{G}$  is *isotrivial* if it is split by a finite étale cover  $S'$  of  $S$ . An isotrivial reductive  $S$ -group scheme  $\mathfrak{G}$  is necessarily of constant type. If  $\mathfrak{G}$  is of constant type with underlying Chevalley group scheme  $G$ ,  $\mathfrak{G}$  is isotrivial if and only if the  $\mathrm{Aut}(G)$ -torsor  $\mathrm{Isom}(G_S, \mathfrak{G})$  is isotrivial.

**2.4. Rank one tori.** The simplest case is that of  $G = \mathbb{G}_{m,S}$ , the split  $S$ -torus of rank 1. The  $S$ -functor  $S' \mapsto \mathrm{Hom}_{S'-gp}(\mathbb{G}_{m,S'}, \mathbb{G}_{m,S'})$  is representable by the constant  $S$ -group scheme  $\mathbb{Z}_S$  [11, VIII.1.5]. It follows that the  $S$ -functor  $S' \mapsto \mathrm{Isom}_{S'-gp}(\mathbb{G}_{m,S'}, \mathbb{G}_{m,S'})$  is representable by the constant  $S$ -group scheme  $(\mathbb{Z}/2\mathbb{Z})_S = \mathrm{Aut}_{S-gp}(\mathbb{Z}_S)$ . On the other hand,  $(\mathbb{Z}/2\mathbb{Z})_S$  is the automorphism group of the split

étale cover  $S \sqcup S \rightarrow S$  of degree 2. By definition an  $S$ -torus of rank one is a form of  $\mathbb{G}_m$  for the fpqc topology; in the other hand, a degree 2 étale cover is a form of  $S \sqcup S$  for the finite étale topology (see for example Lemma 2.1) and a fortiori for the fpqc topology.

According to the faithfully flat descent technique (e.g. [11, XXIV.1.17]), there is then an equivalence of categories between the groupoid of rank one tori over  $S$  (resp. the groupoid of degree 2 étale covers of  $S$ ) and the groupoid of  $(\mathbb{Z}/2\mathbb{Z})_S$ -torsors.

More precisely one associates to an  $S$ -torus  $T$  of rank 1 (resp. to an étale cover  $E$  of degree 2) the  $(\mathbb{Z}/2\mathbb{Z})_S$ -torsor  $\text{Isom}_{gr}(\mathbb{G}_m, T)$  (resp.  $\text{Isom}(S \sqcup S, E)$ ). The inverse map is given by twisting the split object by a given  $(\mathbb{Z}/2\mathbb{Z})_S$ -torsor.

Let  $T$  be an  $S$ -torus of rank one and let  $S' \rightarrow T$  be its associated étale cover of degree 2. Then  $T$  splits after base change to  $S'$  and so is isotrivial. Furthermore we claim that  $T$  is isomorphic to the quotient  $Q = R_{S'/S}(\mathbb{G}_{m,S'})/\mathbb{G}_m$  where  $R_{S'/S}(\mathbb{G}_{m,S'})$  stands for the Weil restriction. We consider the  $(\mathbb{Z}/2\mathbb{Z})_S$ -equivariant exact sequences

$$1 \rightarrow \mathbb{G}_{m,S} \xrightarrow{\Delta} \mathbb{G}_{m,S} \times \mathbb{G}_{m,S} \xrightarrow{\Pi} \mathbb{G}_{m,S} \rightarrow 1$$

where  $\Pi(x, y) = xy^{-1}$  and the  $\mathbb{Z}/2\mathbb{Z}$ -action on  $\mathbb{G}_{m,S} \times \mathbb{G}_{m,S}$  is by  $(x, y) \mapsto (y, x)$ . The  $\mathbb{Z}/2\mathbb{Z}$ -action on the last factor is then by  $x \mapsto x^{-1}$ . Twisting this sequence by the  $(\mathbb{Z}/2\mathbb{Z})_S$ -torsor  $\text{Isom}(S \sqcup S, S')$  yields the desired exact sequence

$$1 \rightarrow \mathbb{G}_m \rightarrow R_{S'/S}(\mathbb{G}_{m,S'}) \rightarrow Q \rightarrow 1$$

where the identification of the second term is left to the reader (it is similar to that of the proof of [11, XXIX.3.13]).

**2.5. Linear representations.** Let  $\mathfrak{G}$  be an  $S$ -group scheme. We say that  $\mathfrak{G}$  is *linear* if there exists a locally free  $\mathcal{O}_S$ -module  $\mathcal{E}$  of finite rank and a group homomorphism  $\mathfrak{G} \rightarrow \text{GL}(\mathcal{E})$  which is a monomorphism.

The notion of locally linear  $S$ -group scheme is then clear and there is also the following variant of *semilocally linear*.

We say that  $\mathfrak{G}$  is *semilocally linear* if for each subset  $\{s_1, \dots, s_n\}$  of points of  $S$  contained in an affine open subset of  $S$ , there exists an open subscheme  $U$  of  $S$  containing  $s_1, \dots, s_n$  such that  $\mathfrak{G} \times_S U$  is linear over  $U$ .

**Lemma 2.2.** *Assume that  $S = \text{Spec}(R)$  and let  $\mathcal{E}$  be a locally free  $\mathcal{O}_S$ -module of finite rank. Then  $\text{GL}(\mathcal{E})$  embeds as a closed  $S$ -subgroup scheme in  $\text{GL}_n$  for some  $n \geq 1$ .*

*Proof.* Since the rank of  $\mathcal{E}$  is a locally constant function [5, Ch. 0, 5.4.1], we can assume that  $\mathcal{E}$  is locally free of constant rank  $r$ . Then  $E = H^0(R, \mathcal{E})$  is a locally free  $R$ -module of rank  $r$ , so is finitely generated projective [12, Tag 00NX]. It follows that there exists an integer  $n \geq 1$  and a decomposition  $R^n = E \oplus E'$ . The homomorphism  $\text{GL}(\mathcal{E}) \rightarrow \text{GL}_n$  is a closed immersion.  $\square$

**Lemma 2.3.** *Let  $\mathfrak{G}$  be an  $S$ -group scheme and let  $S'$  be a finite locally free cover of  $S$ . Then  $\mathfrak{G}$  is linear if and only if  $\mathfrak{G} \times_S S'$  is linear.*

*Proof.* We denote by  $p : S' \rightarrow S$  the structure map. If  $\mathfrak{G}$  is linear, then  $\mathfrak{G} \times_S S'$  is linear. Conversely we assume that there exists a monomorphism  $i : \mathfrak{G} \times_S S' \rightarrow \mathrm{GL}(\mathcal{E}')$  where  $\mathcal{E}'$  is a locally free  $\mathcal{O}_{S'}$ -module of finite rank. We put  $\mathcal{E} = p_*(\mathcal{E}')$ , this is a locally free  $\mathcal{O}_S$ -module of finite rank. We consider the sequence of  $S$ -functors in  $S$ -groups

$$\mathfrak{G} \rightarrow R_{S'/S}(\mathfrak{G} \times_S S') \xrightarrow{R_{S'/S}(i)} R_{S'/S}(\mathrm{GL}(\mathcal{E}')) \rightarrow \mathrm{GL}(\mathcal{E})$$

where  $R_{S'/S}$  stands for the Weil restriction and the first map is the diagonal map which is a monomorphism. Since the Weil restriction for  $S'/S$  transforms monomorphisms into monomorphisms, the map  $R_{S'/S}(i)$  is also a monomorphism and so is the last map since  $R_{S'/S}(\mathrm{GL}(\mathcal{E}'))(T) \subset \mathrm{GL}(\mathcal{E})(T)$  corresponds to automorphisms of  $\mathcal{E} \otimes_{\mathcal{O}_S} \mathcal{O}_T$  which are  $\mathcal{O}_{S'} \otimes_{\mathcal{O}_S} \mathcal{O}_T$ -linear. Since all maps are monomorphisms, we conclude that  $\mathfrak{G}$  is linear.  $\square$

### 3. TORI AND GROUP OF MULTIPLICATIVE TYPE

**3.1. Maximal tori of linear groups.** Let  $\mathcal{A}$  be an Azumaya  $\mathcal{O}_S$ -algebra. We consider the reductive  $S$ -group scheme  $\mathrm{GL}_1(\mathcal{A})$ . Let  $\mathcal{B} \subset \mathcal{A}$  be a separable  $\mathcal{O}_S$ -subalgebra of  $\mathcal{A}$  which is a locally free  $\mathcal{O}_S$ -module of finite rank and which is locally a direct summand of  $\mathcal{A}$  as  $\mathcal{O}_S$ -module. We get a monomorphism of reductive  $S$ -group schemes  $\mathrm{GL}_1(\mathcal{B}) \rightarrow \mathrm{GL}_1(\mathcal{A})$ . In particular, if  $\mathcal{B}$  is commutative,  $\mathcal{B}$  is a finite étale  $\mathcal{O}_S$ -algebra and  $\mathrm{GL}_1(\mathcal{B})$  is a torus. We come now to Grothendieck's definition of maximal étale subalgebras.

**Definition 3.1.** [7, Déf. 5.6]. *We say that a finite étale  $\mathcal{O}_S$ -subalgebra  $\mathcal{C} \subset \mathcal{A}$  is maximal if  $\mathcal{C}$  is locally a direct summand of  $\mathcal{A}$  as  $\mathcal{O}_S$ -module and if the rank of  $\mathcal{C} \otimes_{\mathcal{O}_S} \kappa(s)$  is the degree of  $\mathcal{A}_s \otimes_{\mathcal{O}_S} \kappa(s)$  for each  $s \in S$ .*

If  $\mathcal{C} \subset \mathcal{A}$  is maximal finite étale  $\mathcal{O}_S$ -subalgebra of  $\mathcal{A}$ , then the torus  $\mathrm{GL}_1(\mathcal{C})$  is a maximal torus of  $\mathrm{GL}_1(\mathcal{A})$  since it is the case on geometric fibers. According to [7, §7.5], all maximal  $S$ -tori of  $\mathrm{GL}_1(\mathcal{A})$  occur in that manner; this is part (3) of the following enlarged statement.

**Proposition 3.2.** *Let  $\mathfrak{S}$  be a subgroup scheme of multiplicative type of  $\mathrm{GL}_1(\mathcal{A})$  and put  $\mathcal{B} = \mathcal{A}^{\mathfrak{S}}$ , the centralizer subalgebra of  $\mathfrak{S}$ .*

- (1)  $\mathcal{B}$  is a separable  $\mathcal{O}_S$ -algebra which is locally a direct summand of  $\mathcal{A}$  as  $\mathcal{O}_S$ -module.
- (2) Let  $\mathcal{C}$  be the center of  $\mathcal{B}$ ; this is a finite étale  $\mathcal{O}_S$ -algebra of positive rank which is locally a direct summand of  $\mathcal{B}$  (and  $\mathcal{A}$ ) as  $\mathcal{O}_S$ -module. We have the closed immersions

$$\mathfrak{S} \subset \mathrm{GL}_1(\mathcal{C}) \subset \mathrm{GL}_1(\mathcal{A}).$$

- (3) If  $\mathfrak{S}$  is a maximal torus of  $\mathrm{GL}_1(\mathcal{A})$ , then  $\mathfrak{S} = \mathrm{GL}_1(\mathcal{C})$  and  $\mathcal{C}$  is a maximal finite étale  $\mathcal{O}_S$ -subalgebra of  $\mathcal{A}$ .

(4) If  $\mathfrak{S}$  is of constant type, then  $\mathfrak{S}$  is isotrivial.

*Proof.* According to [11, IX.2.5], the map  $\mathfrak{S} \rightarrow \mathrm{GL}_1(\mathcal{A})$  is a closed immersion so that  $\mathfrak{S}$  is of finite type over  $S$ . As a preliminary observation we notice that (1), (2), (3) are local for the étale topology. We can assume that  $S = \mathrm{Spec}(R)$ ,  $\mathcal{A} = \mathrm{M}_n(R)$  and that  $\mathfrak{S} = D(M)$  for  $M$  an abelian group. Note that  $M$  is finitely generated since  $\mathfrak{S}$  is of finite type (*ibid*, VII.2.1.b).

(1) We consider the  $M$ -grading  $R^n = \bigoplus_{m \in M} R_m^n$ . The  $R$ -modules  $(R_m^n)_{m \in M}$  are finitely generated projective so locally free of finite rank. There is a finite subset  $M' \subset M$  such that  $R^n = \bigoplus_{m \in M'} R_m^n$ . Then  $\mathcal{B} = \prod_{m \in M'} \mathrm{End}_R(R_m^n)$  and each  $\mathrm{End}_R(R_m^n)$  is a separable  $R$ -algebra which is locally free of finite rank [8, III, example 2.8]. Since a product of separable algebras is a separable algebra (*ibid*, III, proposition 1.7), it follows that  $\mathcal{B}$  is a separable  $R$ -algebra. Furthermore  $\mathcal{B}$  is locally a direct summand of  $\mathrm{M}_n(R)$  as  $R$ -module.

(2) Let  $\mathcal{C}$  be the center of  $\mathcal{B}$ . We have  $R$ -monomorphisms of groups  $\mathfrak{S} \subset \mathrm{GL}_1(\mathcal{C}) \subset \mathrm{GL}_1(\mathcal{A})$ .

(3) We assume that  $\mathfrak{S}$  is a maximal torus of  $\mathrm{GL}_1(\mathcal{A})$  and want to establish that  $\mathfrak{S} = \mathrm{GL}_1(\mathcal{C})$ . So  $\mathrm{GL}_1(\mathcal{C})$  is an  $R$ -torus containing  $\mathfrak{T}$  and since maximality holds also in the naive sense [11, XII.1.4], we conclude that  $\mathfrak{T} = \mathrm{GL}_1(\mathcal{C})$ .

(4) We come back to the initial setting (i.e. without localizing). We want to show that  $\mathfrak{S}$  is isotrivial. According to [5, ch.0, 5.4.1], for each integer  $l \geq 0$ ,  $S_l = \{s \in S \mid \mathrm{rank}(\mathcal{C}_{\kappa(s)}) = l\}$  is an open subset of  $S$  so that we have a decomposition in clopen subschemes  $S = \coprod_{l \geq 0} S_l$ . Without loss of generality we can assume

that  $\mathcal{C}$  is locally free of rank  $l$ . Lemma 2.1 provides a finite étale cover  $S'$  of  $S$  such that  $\mathcal{C} \otimes_{\mathcal{O}_S} \mathcal{O}_{S'} \cong (\mathcal{O}_{S'})^l$ . Hence  $\mathrm{GL}_1(\mathcal{C}) \times_S S' \cong (\mathbb{G}_m)^l \times_S S'$ . It follows that  $\mathfrak{S} \times_S S'$  is a subgroup  $S'$ -scheme of  $(\mathbb{G}_m)^l \times_S S'$  of multiplicative type. According to [11, IX.2.11.(i)] there exists a partition in clopen subsets  $S' = \sqcup_{i \in I} S'_i$  such that each  $\mathfrak{S} \times_S S'_i$  is diagonalizable. Since  $\mathfrak{S} \times_S S'$  is of constant type we conclude that  $\mathfrak{S} \times_S S'$  is diagonalizable.  $\square$

### 3.2. Characterization of isotrivial groups of multiplicative type.

**Theorem 3.3.** *Let  $\mathfrak{S}$  be an  $S$ -group scheme of multiplicative type of finite type and of constant type. Then the following are equivalent:*

- (i)  $\mathfrak{S}$  is isotrivial;
- (ii)  $\mathfrak{S}$  is linear;
- (iii)  $\mathfrak{S}$  is an  $S$ -subgroup scheme of an  $S$ -group scheme  $\mathrm{GL}_1(\mathcal{A})$  where  $\mathcal{A}$  is an Azumaya  $\mathcal{O}_S$ -algebra.

*Proof.* (i)  $\implies$  (ii). This follows from Lemma 2.3.

(ii)  $\implies$  (iii). By definition we have a monomorphism  $\mathfrak{S} \rightarrow \mathrm{GL}(\mathcal{E})$  where  $\mathcal{E}$  is a locally free  $\mathcal{O}_S$ -module of finite rank. Since  $\mathrm{End}_{\mathcal{O}_S}(\mathcal{E})$  is an Azumaya  $\mathcal{O}_S$ -algebra, we get (iii).

(iii)  $\implies$  (i). This follows of Proposition 3.2.(4).  $\square$

**Examples 3.4.** (a) Grothendieck constructed a scheme  $S$  and an  $S$ -torus  $\mathfrak{T}$  which is locally trivial of rank 2 but which is not isotrivial [11, §X.1.6] (e.g.  $S$  consists of two copies of the projective line over a field pinched at 0 and  $\infty$ ). Theorem 3.3 shows that such an  $S$ -torus is not linear.

(b) Also there exists a local ring  $R$  and an  $R$ -torus  $\mathfrak{T}$  of rank 2 which is not isotrivial [11, §1.6]; the ring  $R$  can be taken as the local ring of an algebraic curve at a double point. Theorem 3.3 shows that such an  $R$ -torus is not linear.

#### 4. REDUCTIVE CASE

For stating the complete version of our main result, we need more notation. As in the introduction,  $\mathfrak{G}$  is a reductive  $S$ -group scheme of constant type and  $G$  is the underlying Chevalley  $\mathbb{Z}$ -group scheme. We denote by  $\mathrm{Aut}(G)$  the automorphism group scheme of  $G$  and we have an exact sequence of  $\mathbb{Z}$ -group schemes [11, th. XXIV.1.3]

$$1 \rightarrow G_{ad} \rightarrow \mathrm{Aut}(G) \rightarrow \mathrm{Out}(G) \rightarrow 1.$$

We remind the reader of the representability of the fppf sheaf  $\underline{\mathrm{Isom}}(G_S, \mathfrak{G})$  by a  $\mathrm{Aut}(G)_S$ -torsor  $\mathrm{Isom}(G_S, \mathfrak{G})$  defined in [11, XXIV.1.8]. The contracted product

$$\mathrm{Isomext}(G_S, \mathfrak{G}) := \mathrm{Isom}(G_S, \mathfrak{G}) \wedge^{\mathrm{Aut}(G)_S} \mathrm{Out}(G)_S$$

is an  $\mathrm{Out}(G)_S$ -torsor (*ibid*, 1.10) which encodes the isomorphism class of the quasi-split form of  $\mathfrak{G}$ .

**Theorem 4.1.** *The following are equivalent:*

- (i) *The torus  $\mathrm{rad}(\mathfrak{G})$  is isotrivial;*
- (ii) *the  $\mathrm{Out}(G)_S$ -torsor  $\mathrm{Isomext}(G_S, \mathfrak{G})$  is isotrivial;*
- (iii)  *$\mathfrak{G}$  is linear;*
- (iv)  *$\mathrm{rad}(\mathfrak{G})$  is linear.*

Furthermore if  $S$  is affine, we can take a faithful linear representation in some  $\mathrm{GL}_n$  for (iii) and (iv).

*Proof.* (i)  $\implies$  (ii). We assume that  $\mathrm{rad}(\mathfrak{G})$  is isotrivial and want to show that the  $\mathrm{Out}(G)_S$ -torsor  $\mathcal{F} = \mathrm{Isomext}(G_S, \mathfrak{G})$  is isotrivial. In other words, we want to show that there exists a finite étale cover  $S'$  of  $S$  such that  $\mathfrak{G} \times_S S'$  is an inner form of  $G$ .

Without loss of generality we can assume that  $\mathrm{rad}(\mathfrak{G})$  is a split torus. We quote now [11, XXIV.2.16] for the Chevalley group  $G$  over  $\mathbb{Z}$  which introduces the  $\mathbb{Z}$ -group scheme

$$H = \ker(\mathrm{Aut}(G) \rightarrow \mathrm{Aut}(\mathrm{rad}(G)));$$

furthermore there is an equivalence of categories between the category of  $H$ -torsors over  $S$  and the category of pairs  $(\mathfrak{M}, \phi)$  where  $\mathfrak{M}$  is an  $S$ -form of  $G$  and  $\phi : \text{rad}(G)_S \xrightarrow{\sim} \text{rad}(\mathfrak{M})$ .

Since  $\text{rad}(\mathfrak{G})$  is split, we choose an isomorphism  $\phi : \text{rad}(G)_S \xrightarrow{\sim} \text{rad}(\mathfrak{G})$  and consider an  $H$ -torsor  $\mathfrak{P}$  mapping to an object isomorphic to  $(\mathfrak{G}, \phi)$ . Furthermore the quoted reference provides an exact sequence of  $\mathbb{Z}$ -group schemes

$$1 \rightarrow G_{ad} \rightarrow H \xrightarrow{p} F \rightarrow 1$$

where  $F$  is finite étale over  $\mathbb{Z}$ , so is constant. We denote by  $S' = \mathfrak{M} \wedge^{H_S} F_S$  the contracted product of  $\mathfrak{M}$  and  $F_S$  with respect to  $H_S$ ; this is an  $F$ -torsor over  $S$ , hence is a finite étale cover of  $S$ . It follows that  $\mathfrak{P} \times_S S'$  admits a reduction to a  $G_{ad, S'}$ -torsor  $\mathfrak{Q}'$ . Since the map  $G_{ad} \rightarrow H \rightarrow \text{Aut}(G)$  is the canonical map, we conclude that  $\mathfrak{G}_{S'} \cong {}^{\mathfrak{Q}'}G$  is an inner form of  $G$ .

(ii)  $\implies$  (iii). Our assumption is that there exists a finite étale cover  $S'/S$  which splits the  $\text{Out}(G)$ -torsor  $\text{Isomext}(G_S, \mathfrak{G})$ . Lemma 2.3 permits us to replace  $S$  by  $S'$ , so we can assume that the  $\text{Out}(G)$ -torsor  $\text{Isomext}(G_S, \mathfrak{G})$  is trivial; that is,  $\mathfrak{G}$  is an inner form of  $G$ . There exists a  $G_{ad}$ -torsor  $\mathfrak{Q}$  over  $S$  such that  $\mathfrak{G} \cong {}^{\mathfrak{Q}}G$ . Since  $G \rtimes G_{ad}$  is defined over  $\mathbb{Z}$ , it admits a faithful representation  $\rho : G \rtimes G_{ad} \rightarrow \text{GL}_n$  over  $\mathbb{Z}$  [2, §1.4.5]. The map  $\rho$  is then  $G_{ad}$ -equivariant and can be twisted by the  $G_{ad}$ -torsor  $\mathfrak{Q}$ . We obtain a faithful representation  ${}^{\mathfrak{Q}}G \rtimes {}^{\mathfrak{Q}}G_{ad} \rightarrow {}^{\mathfrak{Q}}\text{GL}_n = \text{GL}(\mathcal{E})$  where  $\mathcal{E}$  is the locally free  $\mathcal{O}_S$ -module of rank  $n$  which is the twist of  $(\mathcal{O}_S)^n$  by the  $\text{GL}_n$ -torsor  $\mathfrak{Q} \wedge^{G_{ad}} \text{GL}_n$ . Thus  ${}^{\mathfrak{Q}}G$  is linear.

(iii)  $\implies$  (iv). Obvious.

(iv)  $\implies$  (i). Since  $\text{rad}(\mathfrak{G})$  is a form of  $\text{rad}(G)$ , it is of constant rank and Theorem 3.3 shows that  $\text{rad}(G)$  is isotrivial.

Finally the refinement for  $S$  affine follows from Lemma 2.2.  $\square$

**Corollary 4.2.** *Under the assumptions of  $\mathfrak{G}$ , let  $\mathfrak{G}^{qs}$  be the quasi-split form of  $\mathfrak{G}$ . Then  $\mathfrak{G}$  is linear if and only if  $\mathfrak{G}^{qs}$  is linear.*

The next corollary slightly generalizes a result by Thomason [13, cor. 3.2].

**Corollary 4.3.** *Assume that either*

- (i)  *$S$  is locally noetherian and geometrically unibranch (e.g. normal);*
- (ii)  *$\text{rad}(\mathfrak{G})$  is of rank  $\leq 1$  (in particular if  $G$  is semisimple).*

*Then  $\mathfrak{G}$  is linear.*

*Proof.* In case (i), the torus  $\text{rad}(\mathfrak{G})$  is isotrivial [11, X.5.16]. In case (ii), we have  $\text{rad}(G) = 1$  or  $\mathbb{G}_m$  (since  $G$  is split), so that  $\text{rad}(\mathfrak{G})$  is split by a quadratic étale cover of  $S$  (§2.4), hence is isotrivial. Hence Theorem 1.1 implies that  $\mathfrak{G}$  is linear.  $\square$

The next corollary extends Demazure's characterization of locally isotrivial reductive group schemes [11, XXIV.4.1.5].



**Corollary 4.4.** *The following are equivalent:*

- (i)  $\mathfrak{G}$  is locally (resp. semilocally) isotrivial;
- (ii) The torus  $\text{rad}(\mathfrak{G})$  is locally (resp. semilocally) isotrivial;
- (iii) the  $\text{Out}(G)_S$ -torsor  $\text{Isomext}(G_S, \mathfrak{G})$  is locally (resp. semilocally) isotrivial;
- (iv)  $\mathfrak{G}$  is locally (resp. semilocally) linear;
- (v)  $\text{rad}(\mathfrak{G})$  is locally (resp. semilocally) linear.

*Proof.* In view of Theorem 4.1, it remains to establish the equivalence (i)  $\iff$  (ii). Since this is precisely the quoted result [11, XXIV.3.5], the proof is complete.  $\square$

**Corollary 4.5.** *Let  $\mathfrak{H}$  be a reductive  $S$ -subgroup scheme of  $\mathfrak{G}$ . If  $\mathfrak{G}$  is locally (resp. semilocally) isotrivial, then  $\mathfrak{H}$  is locally (resp. semilocally) isotrivial.*

*Proof.* Corollary 4.4 shows that  $\mathfrak{G}$  is locally (resp. semilocally) linear and so is  $\mathfrak{H}$ . Therefore  $\mathfrak{H}$  is locally (resp. semilocally) isotrivial.  $\square$

## 5. THE SEMILOCAL CASE

We assume that  $S = \text{Spec}(R)$  where  $R$  is a semilocal ring and continue to assume that the reductive  $S$ -group scheme  $\mathfrak{G}$  is of constant type. We remind the reader that  $\mathfrak{G}$  admits a maximal torus (Grothendieck, [11, XIV.3.20 and footnote]).

**Corollary 5.1.** *Let  $\mathfrak{T}$  be a maximal torus of  $\mathfrak{G}$ . The following are equivalent:*

- (i)  $\mathfrak{G}$  is isotrivial;
- (ii) The torus  $\text{rad}(\mathfrak{G})$  is isotrivial;
- (iii) the  $\text{Out}(G)_S$ -torsor  $\text{Isomext}(G_S, \mathfrak{G})$  is isotrivial;
- (iv)  $\mathfrak{G}$  is linear;
- (v)  $\text{rad}(\mathfrak{G})$  is linear;
- (vi)  $\mathfrak{T}$  is linear;
- (vii)  $\mathfrak{T}$  is isotrivial.

*Proof.* From Corollary 4.4, we have the equivalences (i)  $\iff$  (ii)  $\iff$  (iii)  $\iff$  (iv)  $\iff$  (v). On the other hand, the equivalence (vi)  $\iff$  (vii) holds according to Theorem 3.3. Now we observe that the implications (iv)  $\implies$  (vi) and (vi)  $\implies$  (v) are obvious so the proof is complete.  $\square$

## 6. EQUIVARIANT RESOLUTION PROPERTY

**Definition 6.1.** *Let  $\mathfrak{G}$  be a flat group scheme over  $S$  acting on a locally noetherian  $S$ -scheme  $\mathfrak{X}$ . One says that  $(\mathfrak{G}, S, \mathfrak{X})$  has the resolution property (RE) for short) if for every coherent  $\mathfrak{G}$ -module  $\mathcal{F}$  on  $\mathfrak{X}$ , there is a locally free coherent  $\mathfrak{G}$ -module (i.e. a  $\mathfrak{G}$ -vector bundle  $\mathcal{E}$ ) and a  $\mathfrak{G}$ -equivariant epimorphism  $\mathcal{E} \rightarrow \mathcal{F} \rightarrow 0$ .*

We strengthen Thomason's results.

**Theorem 6.2.** *Let  $\mathfrak{G}$  be a reductive  $S$ -group scheme. We assume that  $S$  is separated noetherian and that  $(1, S, S)$  satisfies the resolution property, e.g.  $S$  is affine or regular or admits an ample family of line bundles. Then the following are equivalent:*

- (i)  $\mathfrak{G}$  is linear;
- (ii)  $\text{rad}(\mathfrak{G})$  is isotrivial;
- (iii)  $\mathfrak{G}$  satisfies (RE).

*Proof.* (i)  $\iff$  (ii). This is a special case of Theorem 4.1.

(ii)  $\implies$  (iii). This is Thomason's result [13, Theorem 2.18].

(iii)  $\implies$  (i). This is Thomason's result [13, Theorem 3.1], see also [11, VI<sub>B</sub>.13.5].  $\square$

**Remark 6.3.** Example 3.4.(b) is an example of a local noetherian ring  $R$  and of a rank two non-isotrivial torus  $\mathfrak{T}$ . Theorem 6.2 shows that  $\mathfrak{T}$  does not satisfy (RE). This answers a question of Thomason [13, §2.3].

#### REFERENCES

- [1] A. Borel, *Linear algebraic groups*, Graduate Texts in Mathematics **126** (2nd ed.), Berlin, New York: Springer-Verlag.
- [2] F. Bruhat, J. Tits, *Groupes réductifs sur un corps local : II. Schémas en groupes. Existence d'une donnée radicielle valuée*, Publications Mathématiques de l'IHÉS **60** (1984), 5-184.
- [3] B. Calmès, J. Fasel, *Groupes classiques*, Autour des schémas en groupes, vol II, Panoramas et Synthèses **46** (2015), 1-133.
- [4] M. Demazure et P. Gabriel, *Groupes algébriques*, Masson (1970).
- [5] A. Grothendieck, J.-A. Dieudonné, *Eléments de géométrie algébrique. I*, Grundlehren der Mathematischen Wissenschaften 166; Springer-Verlag, Berlin, 1971.
- [6] A. Grothendieck (avec la collaboration de J. Dieudonné), *Eléments de Géométrie Algébrique IV*, Publications mathématiques de l'I.H.É.S. no 20, 24, 28 and 32 (1964 - 1967).
- [7] A. Grothendieck, *Le groupe de Brauer. I. Algèbres d'Azumaya et interprétations diverses*, Dix exposés sur la cohomologie des schémas, 46-66, Adv. Stud. Pure Math., 3, North-Holland, Amsterdam, 1968.
- [8] M.-A. Knus, M. Ojanguren, *Théorie de la Descente et Algèbres d'Azumaya*, Lecture Notes in Mathematics **389** (1974), Springer
- [9] B. Margaux, *Formal torsors under reductive group schemes*, Rev. Un. Mat. Argentina **60** (2019), 217-224.
- [10] J. Oesterlé, *Schémas en groupes de type multiplicatif*, Autour des schémas en groupes. Vol. I, 63-91, Panor. Synthèses, 42/43, Soc. Math. France, Paris, 2014.
- [11] *Séminaire de Géométrie algébrique de l'I.H.É.S., 1963-1964, Schémas en groupes, dirigé par M. Demazure et A. Grothendieck*, Lecture Notes in Math. 151-153. Springer (1970).
- [12] Stacks project, <https://stacks.math.columbia.edu>
- [13] R. W. Thomason, *Equivariant resolution, linearization, and Hilbert's fourteenth problem over arbitrary base schemes*, Adv. in Math. **65** (1987), 16-34.

P. GILLE, INSTITUT CAMILLE JORDAN - UNIVERSITÉ CLAUDE BERNARD LYON 1 43 BOULEVARD DU 11 NOVEMBRE 1918, 69622 VILLEURBANNE CEDEX - FRANCE

*Email address:* [gille@math.univ-lyon1.fr](mailto:gille@math.univ-lyon1.fr)