## A MINKOWSKI INEQUALITY FOR HOROWITZ-MYERS GEON

AGHIL ALAEE AND PEI-KEN HUNG

Abstract. We prove a sharp inequality for toroidal hypersurfaces in three and four dimensional Horowitz-Myers geon. This extend previous results on Minkowski inequality in the static spacetime to toroidal surfaces in asymptotically hyperbolic manifold with flat toroidal conformal infinity.

#### <span id="page-0-0"></span>1. Introduction

The classical Minkowski inequality for a closed convex hypersurface  $\Sigma$  with induced metric  $\gamma$  in  $\mathbb{R}^n$  reads

(1.1) 
$$
\int_{\Sigma} H \, \mathrm{d} \mathrm{vol}_{\gamma} \geq |\mathbb{S}^{n-1}|^{\frac{1}{n-1}} |\Sigma|^{\frac{n-2}{n-1}}.
$$

Moreover, the rigidity holds if and only if  $\Sigma$  is isometric to the  $(n-1)$ -sphere. The inequality [\(1.1\)](#page-0-0) has been improved to include star-shaped surfaces [\[6,](#page-19-0) [7\]](#page-19-1). One can replace the star-shaped condition by an outward minimizing condition using the weak inverse mean curvature flow of Huisken and Ilmanen [\[10\]](#page-19-2). It is an open problem whether [\(1.1\)](#page-0-0) holds for general mean convex hypersurfaces. This inequality has been generalized to different settings. In [\[4\]](#page-19-3), Brendle, Wang and the second author extended this inequality to convex, star-shaped hypersurface  $\Sigma$  with induced metric  $\gamma$  in the Anti-de-Sitter Schwarzschild space with horizon  $\Sigma_H$  and static potential  $\phi$ 

$$
(1.2) \qquad \int_{\Sigma} \phi H \, \mathrm{d} \mathrm{vol}_{\gamma} - n(n-1) \int_{\Omega} \phi \, \mathrm{d} \mathrm{vol}_{g} \ge (n-1) |\mathbb{S}^{n-1}|^{\frac{1}{n-1}} \left( |\Sigma|^{\frac{n-2}{n-1}} - |\Sigma_H|^{\frac{n-2}{n-1}} \right),
$$

where  $\Omega$  is a bounded region with boundary  $\partial \Omega = \Sigma \cup \Sigma_H$ . Moreover, the equality holds if and only if  $\Sigma$  is a coordinate sphere. For bounded region with outward minimizing boundary  $\Sigma$  in the Schwarzschild space with mass  $m$ , Wei [\[11\]](#page-19-4) proved that

<span id="page-0-1"></span>(1.3) 
$$
\frac{1}{(n-1)|\mathbb{S}^{n-1}|}\int_{\Sigma}\phi H \, \mathrm{dvol}_{\gamma} \ge \left(\frac{|\Sigma|}{|\mathbb{S}^{n-1}|}\right)^{\frac{n-2}{n-1}} - 2m.
$$

with rigidity for coordinate sphere. Recently, McCormick [\[9\]](#page-19-5) showed that the inequality [\(1.3\)](#page-0-1) holds for asymptotically flat static spacetime of dimension  $3 \leq n \leq 7$ . Note that all of these inequalities are for static manifolds with spherical infinity.

In 1999, Horowitz and Myers [\[8\]](#page-19-6) discovered a complete static spacetime  $(N^{n+1}, -\phi^2 dt^2 + g)$  with negative cosmological constant  $-n$ , that is a solution of the following static equation

<span id="page-0-2"></span>(1.4) 
$$
g\Delta_g\phi + \phi \text{Ric}_g - \text{Hess}_g\phi = 0, \qquad R_g = -n(n-1)
$$

where Ric<sub>g</sub>,  $R_g$ ,  $\Delta_g$ , and Hess<sub>g</sub> are Ricci curvature, scalar curvature, Laplace-Beltrami operator, and the hessian with respect to metric  $g$ , respectively. The time constant slice of this spacetime

Date: March 16, 2021.

A. Alaee acknowledges the support of the AMS-Simons travel grant.

## $2\,$   $\,$  ALAEE AND HUNG

is an asymptotically hyperbolic manifold with flat toroidal conformal infinity and it is called the Horowitz-Myers geon  $(M, q, \phi)$  [\[12\]](#page-19-7). The metric takes the form

(1.5) 
$$
g = ds^2 + \left(\frac{d\phi}{ds}\right)^2 d\xi^2 + \phi^2 \sum_{i=3}^n (d\theta^i)^2, \qquad \phi(s) := \cosh^{\frac{2}{n}} \left(\frac{ns}{2}\right),
$$

where  $s \in [0, \infty)$  is the geodesic coordinate from central torus,  $\xi \in [0, 4\pi/n]$ , and  $\theta^i \in [0, a_i]$  for  $0 < a_3 \leq \cdots \leq a_n$ . The period of  $\xi$  is chosen such that, after a coordinate change, the metric is smooth up to the central torus  $\{s = 0\}$ . The function  $\phi(s)$  satisfies the equation

<span id="page-1-5"></span>(1.6) 
$$
\frac{d\phi}{ds} = \phi (1 - \phi^{-n})^{1/2}.
$$

The total mass of the Horowitz-Myers geon is

<span id="page-1-0"></span>(1.7) 
$$
m = -\frac{4\pi}{n} \prod_{i=3}^{n} a_i.
$$

They postulated that this may be the ground state for a conjectural nonsupersymmetric AdS/CFT correspondence [\[8\]](#page-19-6). More precisely, let  $(M, g)$  be an asymptotically hyperbolic manifold with flat toroidal conformal infinity and scalar curvature  $R(g) \geq -n(n-1)$ . Then the total mass is at least equal to the mass of the Horowitz-Myers geon. Moreover, the rigidity holds if and only if  $(M, g)$  is isometric to the Horowitz-Myers geon. The Horowitz-Myers conjecture follows from Conjecture 3 of [\[8\]](#page-19-6). Progress on the Horowitz-Myers conjecture has been very limited. For small perturbations of Horowitz-Myers the conjecture is true by Constable and Myers [\[5\]](#page-19-8) and Woolgar proved the rigidity of this conjecture in 3 dimensions [\[12\]](#page-19-7). Recently, in [\[1\]](#page-19-9), there are more supporting evidence in the validity of this conjecture.

In this paper, we investigate a Minkowski type inequality for toroidal hypersurfaces  $\Sigma$  with induced metric  $\gamma$  in the Horowitz-Myers geon  $(M, g, \phi)$ . We consider a Minkowski-type quantity

<span id="page-1-4"></span>(1.8) 
$$
Q(\Sigma) := n(n-1) \int_{\Omega} \phi \, \mathrm{dvol}_g - \int_{\Sigma} \phi H \, \mathrm{dvol}_{\gamma}.
$$

Here H is the mean curvature of  $\Sigma$  in M and  $\Omega$  is the bonded region enclosed by  $\Sigma$ . Applying the Divergence Theorem and the static equation [\(1.4\)](#page-0-2),  $Q(\Sigma)$  can be rewritten as

<span id="page-1-6"></span>(1.9) 
$$
Q(\Sigma) = \int_{\Sigma} (n-1)g(\nabla \phi, \nu) - \phi H \, \text{dvol}_{\gamma}.
$$

We aim to prove that under certain convexity assumption,

<span id="page-1-1"></span>
$$
(1.10)\t\t Q(\Sigma) \le -2^{-1}nm,
$$

where  $m$  is the mass defined in  $(1.7)$ . The main results are

<span id="page-1-3"></span>**Theorem 1.1.** Let  $n = 3$  and  $\Sigma \subset M$  be a graph over  $T^2$ . Suppose that the second fundamental form  $\tilde{h}_{ab}$  of  $\Sigma$  that corresponds to the conformal metric  $\tilde{g} = \phi^2 g$  is non-negative definite and that

<span id="page-1-2"></span>
$$
\min_{\Sigma} \phi > 1.
$$

Then [\(1.10\)](#page-1-1) holds and the equality is achieved if and only if  $\Sigma$  is given by a coordinate torus  $s \equiv$ constant.

<span id="page-2-1"></span>**Theorem 1.2.** Let  $n = 4$  and  $\Sigma \subset M$  be a graph over  $T^3$ . Assume  $\Sigma$  is symmetric along the  $\xi$ direction. Suppose that the second fundamental form  $\bar{h}_{ab}$  of  $\Sigma$  that corresponds to the conformal metric  $\bar{g} = \left(\phi \frac{d\phi}{ds}\right)^2 g$  is non-negative definite and that

$$
\min_{\Sigma} \phi^4 \ge 1 + \frac{2}{\sqrt{3}}.
$$

Then [\(1.10\)](#page-1-1) holds and the equality is achieved if and only if  $\Sigma$  is given by a coordinate torus  $s \equiv$ constant.

**Remark 1.3.** The assumption [\(1.11\)](#page-1-2) in Theorem [1.1](#page-1-3) is to avoid the coordinate singularity at  $\{s \equiv 0\}$ . The assumption  $(1.12)$  in Theorem [1.2](#page-2-1) is to ensure the flow  $(4.1)$  admits a long-time solution. As long as the solution of [\(4.1\)](#page-10-0) exists for all time, the result of Theorem [1.2](#page-2-1) holds. See Lemma [4.1](#page-10-1) for more detail.

The structure of this article is as follows. In Section [2](#page-2-2) we discuss the geometry of graphs and prove the monotinicity of the  $Q$  in  $(1.8)$  under a weighted normal flow. In Section [3,](#page-5-0) we prove the global existence of the flow in dimensions 3 and prove Theorem [1.1.](#page-1-3) Finally, in Section [4,](#page-10-2) we prove the global existence of the flow in dimensions 4 for axially symmetric graphs and prove Theorem [1.2.](#page-2-1)

## <span id="page-2-0"></span>2. GEOMETRY OF GRAPHS

<span id="page-2-2"></span>In this section, we investigate the geometry of graph  $\Sigma$  in  $(M, g, \phi)$ . First, we define the function

<span id="page-2-3"></span>(2.1) 
$$
q(s) := \int_0^s \frac{ds'}{\phi(s')}.
$$

Using q as a coordinate, the conformal metric  $g' := \phi^{-2}g$  is asymptotic to a flat metric on  $\mathbb{R} \times T^{n-1}$ 

(2.2) 
$$
g' = dq^2 + \left(\phi^{-1}\frac{d\phi}{ds}\right)^2 d\xi^2 + \sum_{i=3}^n (d\theta^i)^2.
$$

We will further consider other conformal metrics

(2.3) 
$$
\tilde{g} = \phi^2 g, \ \bar{g} = \left(\phi \frac{d\phi}{ds}\right)^2 g.
$$

The flows we consider, [\(3.1\)](#page-5-1) and [\(4.1\)](#page-10-0), are unit normal flows in  $\tilde{q}$  and  $\bar{q}$  respectively.

We use  $\nabla$  and  $R_{\alpha\beta\lambda\mu}$  to denote the Levi-Civita connection and the Riemannian curvature tensor of g respectively. For a hypersurface  $\Sigma$ , we use  $\{x^a\}$ ,  $a = 1, 2, \ldots, n-1$  as a local coordinate. The induced metric and the second fundamental form are denoted by  $\gamma_{ab}$  and  $h_{ab}$  respectively. We write D for the Levi-Civita connection of  $\gamma$ . We use  $dvol_q$  and  $dvol_\gamma$  to denote the volume form of g and  $\gamma$  respectively. The tensors and connections induced by the conformal metrics  $g', \tilde{g}$  or  $\bar{g}$  will carry the corresponding accents. For instance, the second fundamental form with respect to  $g'$  is written as  $h'_{ab}$ . Also, the index in Weingarten tensor is raised accordingly. For example,  $\tilde{h}^b_a = \tilde{\gamma}^{bc} \tilde{h}_{ca}$ .

In the Appendix [A,](#page-14-0) we calculate the curvatures and the second fundamental forms with respect to the metric

$$
dq^{2} + \Psi(q)^{2} d\xi^{2} + \sum_{i=3}^{n} (d\theta^{i})^{2},
$$

for a general positive function  $\Psi(q)$ . The metric g' corresponds to  $\Psi = \phi^{-1} \frac{d\phi}{ds}$ , which is the only case we will use. We further record the curvatures and the second fundamental forms under a conformal transformation  $\check{g} = e^{2\psi} g'$ . The metrics g,  $\tilde{g}$  and  $\bar{g}$  correspond to  $\psi = \log \phi$ ,  $2 \log \phi$  and  $2 \log \phi + \log \frac{d\phi}{ds}$ respectively.

Let  $\Sigma$  be a graph given by  $s = v(\xi, \theta^i)$  for some smooth function v defined on  $T^{n-1}$ . Define the height function in the  $q$  coordinate as

$$
u(\xi, \theta^i) = q(v(\xi, \theta^i)).
$$

Define the slope of  $\Sigma$  by

<span id="page-3-2"></span>(2.4) 
$$
\rho^2 = 1 + \left(\phi \left/ \frac{d\phi}{ds}\right)^2 \left(\frac{\partial u}{\partial \xi}\right)^2 + \delta^{ij} \frac{\partial u}{\partial \theta^i} \frac{\partial u}{\partial \theta^j}.
$$

Let H be the mean curvature of  $\Sigma$  with respect to g. A direct computation using [\(A.11\)](#page-18-0) with  $\Psi = \phi^{-1} \frac{d\phi}{ds}, \psi = \log \phi$  and [\(1.6\)](#page-1-5) shows

$$
H = -\rho \phi^{-1} \Delta' u + (2^{-1}n)\rho^{-1} \phi^{1-n} \left(\frac{d\phi}{ds}\right)^{-1} \delta^{ij} \frac{\partial u}{\partial \theta^i} \frac{\partial u}{\partial \theta^j} + (n-1)\rho^{-1} \phi^{-1} \frac{d\phi}{ds} + (2^{-1}n)\rho^{-1} \phi^{1-n} \left(\frac{d\phi}{ds}\right)^{-1}
$$

.

Here  $\Delta'$  is the Laplace-Beltrami operator of  $\gamma'$ . Together with

<span id="page-3-0"></span>
$$
d\text{vol}_{\gamma} = \rho \phi^{n-2} \frac{d\phi}{ds} d\xi \wedge d\theta^3 \wedge \dots \wedge d\theta^n,
$$

we derive from [\(1.9\)](#page-1-6) that

<span id="page-3-3"></span>(2.5) 
$$
Q(\Sigma) + 2^{-1}nm = \int_{T^{n-1}} \rho^2 \phi^{n-2} \frac{d\phi}{ds} \Delta' u - (2^{-1}n) \delta^{ij} \frac{\partial u}{\partial \theta^i} \frac{\partial u}{\partial \theta^j} d\xi \wedge d\theta^3 \wedge \cdots \wedge d\theta^n.
$$

Let p be a function defined on M to be determined. Let  $F : [0, T_0) \times T^{n-1} \to M$  be a family of embeddings that satisfies

$$
\frac{\partial F}{\partial t} = p\nu,
$$

where  $\nu$  is the unit outward normal. Denote by  $\Sigma_t$  the image of  $F(t, \cdot)$  and by  $\gamma_t$  the induced metric.

#### <span id="page-3-1"></span>Lemma 2.1.

- For  $n = 3$ , by taking  $p = \phi^{-1}$ ,  $Q(\Sigma_t)$  is monotone non-decreasing along the flow [\(2.6\)](#page-3-0).
- For  $n = 4$ , assume  $\Sigma_0$  is symmetric along the  $\xi$  direction. Then by taking  $p = (\phi \frac{d\phi}{ds})^{-1}$ ,  $Q(\Sigma_t)$  is monotone non-decreasing along the flow [\(2.6\)](#page-3-0).

Furthermore, in the two cases above,  $Q(\Sigma_t)$  is strictly increasing unless  $\Sigma_0$  is a coordinate torus.

Proof. We compute

(2.7) 
$$
\frac{d}{dt}Q(\Sigma_t) = n(n-1)\int_{\Sigma_t} p\phi \,d\mathrm{vol}_{\gamma_t} - \int_{\Sigma_t} \left(\frac{\partial \phi}{\partial t}H + p\phi H^2 + \phi \frac{\partial H}{\partial t}\right) \,d\mathrm{vol}_{\gamma_t}.
$$

Using the first variation of area and the Gauss equation,

(2.8) 
$$
\frac{\partial H}{\partial t} = -\Delta_{\gamma t} p - p|h|^2 - p\text{Ric}(\nu, \nu),
$$

(2.9) 
$$
n(n-1) - H^2 + |h|^2 = -R_{\gamma_t} - 2\text{Ric}(\nu, \nu).
$$

Hence

(2.10) 
$$
\frac{d}{dt}Q(\Sigma_t) = \int_{\Sigma_t} p\phi \left[ -R_{\gamma_t} - 2\mathrm{Ric}(\nu,\nu) \right] - pHg(\nabla\phi,\nu) + p\phi \mathrm{Ric}(\nu,\nu) + \phi \Delta_{\gamma_t} p \, d\mathrm{vol}_{\gamma_t}.
$$

Using the static equation [\(1.4\)](#page-0-2),

(2.11) 
$$
\Delta_{\gamma t} \phi = \Delta_g \phi - \nabla^2 \phi(\nu, \nu) - Hg(\nabla \phi, \nu) = -\phi \text{Ric}(\nu, \nu) - Hg(\nabla \phi, \nu),
$$

we obtain

(2.12)  
\n
$$
\frac{d}{dt}Q(\Sigma_t) = \int_{\Sigma_t} -p\phi R_{\gamma_t} + p\Delta_{\gamma_t}\phi + \phi \Delta_{\gamma_t} p \,d\mathrm{vol}_{\gamma_t}
$$
\n
$$
= \int_{\Sigma_t} -p\phi R_{\gamma_t} - 2\gamma_t (D\phi, Dp) \,d\mathrm{vol}_{\gamma_t}.
$$

For  $n=3$ , by choosing  $p=\phi^{-1}$ , we obtain from the Gauss-Bonnet Theorem that

(2.13) 
$$
\frac{d}{dt}Q(\Sigma_t) = \int_{\Sigma_t} -R_{\gamma_t} + 2\phi^{-2}\gamma_t(D\phi, D\phi) d\mathrm{vol}_{\gamma_t} \geq 0.
$$

Moreover,  $\frac{d}{dt}Q(\Sigma_t) = 0$  if and only if  $\phi$  is a constant on  $\Sigma_t$ . This implies  $\Sigma_t$  is a coordinate torus. By reversing the flow  $(2.6)$ ,  $\Sigma_0$  is also a coordinate torus.

For  $n = 4$ , we assume that  $\Sigma_0$  is symmetric along the  $\xi$  direction. Hence  $\Sigma_t$  preserves the same symmetry. Denote by  $\gamma_t$  the induced metric on  $\Sigma_t = \Sigma_t \cap {\xi \equiv \text{constant}}$ . Then

$$
\gamma_t = \mathring{\gamma}_t + \left(\frac{d\phi}{ds}\right)^2 d\xi^2.
$$

The scalar curvature of  $\gamma_t$  and  $\mathring{\gamma}_t$  are related through

$$
R_{\gamma_t} = R_{\dot{\gamma}_t} - 2 \left( \frac{d\phi}{ds} \right)^{-1} \Delta_{\dot{\gamma}_t} \left( \frac{d\phi}{ds} \right).
$$

Assuming  $p$  is also symmetric along the  $\xi$  direction, we get

$$
\frac{d}{dt}Q(\Sigma_t) = \frac{4\pi}{n} \int_{\mathring{\Sigma}_t} -p\phi \frac{d\phi}{ds} R_{\dot{\gamma}_t} + 2p\phi \Delta_{\dot{\gamma}_t} \left(\frac{d\phi}{ds}\right) - 2\frac{d\phi}{ds} \dot{\gamma}_t (D\phi, Dp) d\mathrm{vol}_{\dot{\gamma}_t}.
$$

Taking  $p = \left(\phi \frac{d\phi}{ds}\right)^{-1}$ , we deduce from the Gauss-Bonnet Theorem that

$$
\frac{d}{dt}Q(\Sigma_t) = \frac{4\pi}{n} \int_{\hat{\Sigma}_t} -R_{\hat{\gamma}_t} + 2\left(\frac{d\phi}{ds}\right)^{-1} \Delta_{\hat{\gamma}_t} \left(\frac{d\phi}{ds}\right) - 2\frac{d\phi}{ds}\hat{\gamma}_t \left(D\phi, D\left(\phi\frac{d\phi}{ds}\right)^{-1}\right) d\text{vol}_{\hat{\gamma}_t}
$$
\n
$$
= \frac{4\pi}{n} \int_{\hat{\Sigma}_t} \left(-2\frac{d}{ds}\left(\frac{d\phi}{ds}\right)^{-1} \frac{d^2\phi}{ds^2} - 2\left(\frac{d\phi}{ds}\right)^2 \frac{d}{ds}\left(\phi\frac{d\phi}{ds}\right)^{-1}\right) \cdot \hat{\gamma}_t(Dv_t, Dv_t) d\text{vol}_{\hat{\gamma}_t}.
$$

From  $(1.6)$  and  $n = 4$ , we derive

$$
-2\frac{d}{ds}\left(\frac{d\phi}{ds}\right)^{-1}\frac{d^2\phi}{ds^2} - 2\left(\frac{d\phi}{ds}\right)^2\frac{d}{ds}\left(\phi\frac{d\phi}{ds}\right)^{-1} = (1 - \phi^{-4})^{-1}(6 + 2\phi^{-8}) \ge 0.
$$

Thus  $Q(\Sigma_t)$  is monotone non-decreasing along the flow. Using the same argument for the 3 dimensional case,  $\frac{d}{dt}Q(\Sigma_t) = 0$  if and only if  $\Sigma_0$  is a coordinate sphere.

## <span id="page-5-0"></span> $6$   $\,$  ALAEE AND HUNG  $\,$

## <span id="page-5-1"></span>3. Three Dimensional Case

In this section, we fix  $n = 3$  and prove Theorem [1.1.](#page-1-3) Recall that  $F : [0, T_0) \times T^{n-1} \to M$  solves

(3.1) 
$$
\frac{\partial F}{\partial t} = \phi^{-1} \nu.
$$

Here we take  $T_0 \in (0,\infty]$  to be the largest number such that  $(3.1)$  has a smooth graphical solution in  $t \in [0, T_0)$ .

Equation [\(3.1\)](#page-5-1) can be viewed as the unit normal flow with respect to the conformal metric  $\tilde{g} = \phi^2 g$ . Recall that  $\tilde{h}_{ab}$  is the the second fundamental form with respect to  $\tilde{g}$ . Our main assumption is

<span id="page-5-2"></span>(3.2) 
$$
\tilde{h}_{ab} \text{ is non-negative definite for } \Sigma_0 \text{ and } \min_{\Sigma_0} \phi > 1.
$$

We list two lemmas which allow us to prove Theorem [1.1.](#page-1-3)

<span id="page-5-4"></span>**Lemma 3.1.** Suppose  $T_0 = \infty$ . Then

<span id="page-5-8"></span>(3.3) 
$$
\limsup_{t \to \infty} Q(\Sigma_t) \leq -\frac{3m}{2}.
$$

<span id="page-5-3"></span>**Lemma 3.2.** Under the assumption [\(3.2\)](#page-5-2), we have  $T_0 = \infty$ .

*Proof of Theorem [1.1.](#page-1-3)* Let  $\Sigma_t$  be the solution of [\(3.1\)](#page-5-1) starting from  $\Sigma_0 = \Sigma$ . Combining Lemmas [2.1,](#page-3-1) [3.2](#page-5-3) and [3.1,](#page-5-4)

$$
Q(\Sigma) \le \limsup_{t \to \infty} Q(\Sigma_t) \le -\frac{3m}{2}.
$$

Furthermore, the equality implies  $Q(\Sigma_t)$  is a constant along the flow. And Lemma [2.1](#page-3-1) implies  $\Sigma$  is a coordinate torus.

We prove Lemmas [3.1](#page-5-4) and Lemma [3.2](#page-5-3) in the rest of this section. We adapt the convention that C denotes a large constant depending on  $\Sigma_0$ . The value of C may change from line to line.

Denote by  $v_t(\xi, \theta)$  the height function of  $\Sigma_t$  and  $u_t = q(v_t)$ . Then  $v_t$  solves the equation

(3.4) 
$$
\frac{\partial v_t}{\partial t} = \rho \phi(v_t)^{-1}.
$$

Here  $\rho$  is the slope of  $\Sigma_t$  defined in [\(2.4\)](#page-3-2). We start with the  $C^0$  and  $C^1$  estimates for  $v_t$  and  $u_t$ .

<span id="page-5-7"></span>**Lemma 3.3.** There exists a constant C depending on  $\Sigma_0$  such that for all  $t \in [0, T_0)$ ,

<span id="page-5-6"></span>(3.5) 
$$
|\phi(v_t) - t| \leq C, \ \phi(v_t) \geq 1 + C^{-1}(t+1).
$$

*Proof.* At the minimum point of  $v_t$ , we have  $\rho = 1$ . Together with  $(1.6)$ ,

<span id="page-5-5"></span>
$$
\frac{d}{dt} \min_{T^2} \phi(v_t) \ge \left(1 - (\min_{T^2} \phi(v_t))^{-3}\right)^{1/2}.
$$

Let  $\Phi(t, a)$  be the solution to the ODE

$$
\frac{\partial \Phi}{\partial t} = (1 - \Phi^{-3})^{1/2}, \ \Phi(0, a) = a.
$$

Through the ODE comparison,

$$
\min_{T^2} \phi(v_t) \ge \Phi(t, \min_{T^2} \phi(v_0)).
$$

From now on, we take  $a = \min_{T^2} \phi(v_0)$ . By the assumption [\(3.2\)](#page-5-2),  $a > 1$ . Because  $\Phi(t, a)$  is increasing in t, we have  $\frac{\partial \Phi}{\partial t} \geq (1 - a^{-3})^{1/2}$ . Through integration,  $\Phi(t, a) \geq (1 - a^{-3})^{1/2}t + a$ . This implies  $\phi(v_t) \geq 1 + C^{-1}(t+1)$  for C large enough. Furthermore, using the equation of  $\Phi$  again,

$$
\left|\frac{\partial \Phi}{\partial t} - 1\right| \le \frac{C}{(1+t)^3}.
$$

Thus  $|\Phi(t, a) - t| \leq C$  and  $\min_{T^2} \phi(v_t) \geq t - C$ . With a similar argument, we can also derive

$$
\max_{T^2} \phi(v_t) \le t + C.
$$

The proof is finished.  $\Box$ 

For the notation simplicity, we later denote  $\phi(v_t)$  and  $\frac{d\phi}{ds}(v_t)$  by  $\phi$  and  $\frac{d\phi}{ds}$  respectively. Recall that

$$
\rho^2 = 1 + \left(\frac{\partial u_t}{\partial \theta}\right)^2 + \left(\phi \left/\frac{d\phi}{ds}\right)^2 \left(\frac{\partial u_t}{\partial \xi}\right)^2.
$$

<span id="page-6-0"></span>**Lemma 3.4.** There exists a constant C depending on  $\Sigma_0$  such that for  $t \in [0, T_0)$ ,

(3.6) 
$$
\max_{\Sigma_t} (\rho^2 - 1) \le (C^{-1}t + 1)^{-4} \max_{\Sigma_0} (\rho^2 - 1).
$$

*Proof.* Using  $(1.6)$  and  $(3.4)$ , we have

<span id="page-6-1"></span>
$$
\frac{\partial^2 u_t}{\partial x^a \partial t} = \phi^{-2} \frac{\partial \rho}{\partial x^a} - 2\rho \phi^{-1} (1 - \phi^{-3})^{1/2} \frac{\partial u_t}{\partial x^a}
$$

and

$$
\frac{\partial}{\partial t} \left( \phi \bigg/ \frac{d\phi}{ds} \right)^2 = -3\rho \phi^{-4} (1 - \phi^{-3})^{-3/2}.
$$

It follows that

$$
\frac{\partial}{\partial t}(\rho^2 - 1) = 2\phi^{-2} \left( \frac{\partial u_t}{\partial \theta} \frac{\partial \rho}{\partial \theta} + \left( \phi \left/ \frac{d\phi}{ds} \right)^2 \frac{\partial u_t}{\partial \xi} \frac{\partial \rho}{\partial \xi} \right) - 3\rho \phi^{-4} (1 - \phi^{-3})^{3/2} \left( \frac{\partial u_t}{\partial \xi} \right)^2 - 4\rho \phi^{-1} (1 - \phi^{-3})^{1/2} (\rho^2 - 1).
$$

At the maximum point of  $\rho^2$ ,  $d\rho = 0$  and

$$
\frac{\partial}{\partial t}(\rho^2 - 1) \le -4\rho\phi^{-1}(1 - \phi^{-3})^{1/2}(\rho^2 - 1).
$$

Using [\(3.5\)](#page-5-6), there exists a constant C such that for all  $t \in [0, T_0)$ ,

$$
4\rho\phi^{-1}(1-\phi^{-3})^{1/2} \ge 4\phi^{-1}(1-\phi^{-3})^{1/2} \ge 4(t+C)^{-1}.
$$

Therefore,

$$
\frac{d}{dt} \max_{\Sigma_t} (\rho^2 - 1) \leq -4(t + C)^{-1} \max_{\Sigma_t} (\rho^2 - 1).
$$

By the ODE comparison,

$$
\max_{\Sigma_t}(\rho^2 - 1) \le (C^{-1}t + 1)^{-4} \max_{\Sigma_0}(\rho^2 - 1).
$$

The proof is finished.  $\square$ 

Recall that  $g' = \phi^{-2}g$  is a conformal metric which asymptotic to the flat metric on  $\mathbb{R} \times T^2$ . Let  $\gamma'_t$ be the induced metric of g' on  $\Sigma_t$  and D' be the Levi-Civita connection of  $\gamma'_t$ . Lemmas [3.3](#page-5-7) and [3.4](#page-6-0) imply, provided  $T_0 = \infty$ ,  $\gamma'_t$  converges the the flat metric  $d\xi^2 + d\theta^2$  in  $C^0$ . Next, we use the second fundamental form  $h'_{ab}$  to bound the hessian of  $u_t$  from below.

<span id="page-7-1"></span>**Lemma 3.5.** Suppose  $T_0 = \infty$ . Then there exists a constant C depending on  $\Sigma_0$  such that for  $t \geq C+1$ ,

$$
D'_{b}(D')^{a}u_{t} \geq -\left(C(t-C)^{-2}\log t\right)\delta_{b}^{a}.
$$

Proof. The flow [\(3.1\)](#page-5-1) can be rewritten as

$$
\frac{\partial F}{\partial t} = \phi^{-2} \nu'.
$$

Here  $\nu' = \phi \nu$  is the unit normal vector with respect to g'. The evolution equation of  $(h')_b^a$  is given by

$$
\frac{\partial}{\partial t}(h')^a_b = -\phi^{-2}(h')^c_b(h')^a_c - D'_b(D')^a\phi^{-2} - \phi^{-2}(dx^a)_{\alpha}(\nu')^{\beta} \left(\frac{\partial}{\partial x^b}\right)^{\lambda} (\nu')^{\mu} (R')^{\alpha}_{\beta\lambda\mu}.
$$

Here  $(R')_{\alpha\beta\lambda\mu}$  is the Riemannian curvature tensor of g'. In the estimates below, the norms are computed with respect to  $\gamma'_t$ . From [\(A.3\)](#page-14-1) and [\(1.6\)](#page-1-5), the only non-zero component of R' is

$$
R'_{q\xi q\xi} = 3\phi^{-1}(1 - \phi^{-3}).
$$

Together with  $(3.5)$ , for t large enough,

$$
\left| -\phi^{-2} (dx^a)_{\alpha} (\nu')^{\beta} \left( \frac{\partial}{\partial x^b} \right)^{\lambda} (\nu')^{\mu} (R')^{\alpha}_{\beta \lambda \mu} \right| \leq C(t - C)^{-3}.
$$

Using  $(1.6)$  and  $(2.1)$ ,

$$
-D'_b(D')^a\phi^{-2} = 2\phi^{-1}(1-\phi^{-3})^{1/2}D'_b(D')^au_t + (-2+5\phi^{-3})D'_bu_t(D')^au_t.
$$

The relation between  $h'_{ab}$  and  $D'_b D'_a u_t$  is given in Corollary [A.3](#page-16-0) with  $\Psi = \phi^{-1} \frac{d\phi}{ds}$ . Using [\(1.6\)](#page-1-5), we derive

<span id="page-7-0"></span>(3.7) 
$$
D'_b(D')_au_t = -\rho^{-1}h'_{ab} + \frac{3}{2}\phi^{-2}(1-\phi^{-3})^{1/2}D'_b\xi D'_a\xi.
$$

Hence

$$
-D'_b(D')^a\phi^{-2} = -2\rho^{-1}\phi^{-1}(1-\phi^{-3})^{1/2}(h')^a_b + (-2+5\phi^{-3})D'_b u_t(D')^a u_t + 3\phi^{-3}(1-\phi^{-3})D'_b \xi(D')^a \xi.
$$

From  $(3.5)$  and  $(3.6)$ , for t large enough, we have

$$
|D'_b u_t (D')^a u_t| \le C(t - C)^{-4}
$$

and

$$
|3\phi^{-3}(1-\phi^{-3})D'_b\xi(D')^a\xi| \le (t-C)^{-3}.
$$

Putting the above together, we get

$$
\left| \frac{\partial}{\partial t} (h')^a_b + 2\rho^{-1} \phi^{-1} (1 - \phi^{-3})^{1/2} (h')^a_b + \phi^{-2} (h')^c_b (h')^a_c \right| \le C(t - C)^{-3}.
$$

Let  $w_t$  be the maximum eigenvalue of  $(h')_a^b$ . Using again [\(3.5\)](#page-5-6) and [\(3.6\)](#page-6-1), for t large enough, we have  $2\rho^{-1}\phi^{-1}(1-\phi^{-3})^{1/2} \geq 2(t+C)^{-1}$ . Therefore,

$$
\frac{dw_t}{dt} \le -2(t+C)^{-1}w_t + C(t-C)^{-3}.
$$

From the ODE comparison,  $w_t \leq C(t-C)^{-2} \log t$  for t large enough. The assertion then follows from the view of  $(3.7)$ .

Proof of Lemma [3.1.](#page-5-4) From  $(2.5)$ ,

$$
Q(\Sigma_t) + \frac{3m}{2} = \int_{T^2} \rho^2 \phi \frac{d\phi}{ds} \Delta' u_t - \frac{3}{2} \left(\frac{\partial u_t}{\partial \theta}\right)^2 d\xi \wedge d\theta.
$$

From Lemma [3.4,](#page-6-0)  $\left(\frac{\partial u_t}{\partial \theta}\right)^2 \leq C(t-C)^{-4}$ . It remains to prove

$$
\limsup_{t \to \infty} \int_{T^2} \rho^2 \phi \frac{d\phi}{ds} \Delta' u_t d\xi \wedge d\theta \le 0.
$$

Recall that  $\gamma'_t$  is the induced metric of  $\Sigma$  from g'. Because

$$
d\text{vol}_{\gamma'_t} = \rho \phi^{-1} \frac{d\phi}{ds} d\xi \wedge d\theta,
$$

$$
\int_{T^2} \rho^2 \phi \frac{d\phi}{ds} \Delta' u_t d\xi \wedge d\theta = \int_{\Sigma_t} \rho \phi^2 \Delta' u_t d\mathrm{vol}_{\gamma'_t} = \underbrace{\int_{\Sigma_t} (\rho - 1) \phi^2 \Delta' u_t d\mathrm{vol}_{\gamma'_t}}_{\text{I}} + \underbrace{\int_{\Sigma_t} \phi^2 \Delta' u_t d\mathrm{vol}_{\gamma'_t}}_{\text{II}}.
$$

From Lemmas [3.3,](#page-5-7) [3.4](#page-6-0) and [3.5,](#page-7-1)  $\phi^2 \le C(t+C)^2$ ,  $|\rho - 1| \le C(t-C)^{-4}$  and

$$
\int_{\Sigma_t} |\Delta' u_t| \, d\text{vol}_{\gamma'_t} = 2 \int_{\Sigma_t} (\Delta' u_t)_- \, d\text{vol}_{\gamma'_t} \le C(t - C)^{-2} \log t.
$$

Therefore the first term I goes to zero. Through integration by parts,

$$
\mathrm{II} = -\int_{\Sigma_t} 2\phi \frac{d\phi}{ds}|D'u_t|^2_{\gamma'_t}\,d\mathrm{vol}_{\gamma'_t} \leq 0.
$$

Thus the assertion  $(3.3)$  follows.

Lastly, we prove Lemma [3.2.](#page-5-3) Recall that  $\tilde{g} = \phi^2 g$  and that  $\tilde{h}^a_b$  is the Weingarten tensor with respect to  $\tilde{g}$ . The flow [\(3.1\)](#page-5-1) is a unit normal flow with respect to  $\tilde{g}$ . The evolution equation of  $\tilde{h}^a_b$  is given by

(3.8) 
$$
\frac{\partial}{\partial t} \tilde{h}_b^a = -\tilde{h}_b^c \tilde{h}_c^a - (dx^a)_{\alpha} \tilde{\nu}^{\beta} \left(\frac{\partial}{\partial x^b}\right)^{\lambda} \tilde{\nu}^{\mu} \tilde{R}^{\alpha}_{\ \beta \lambda \mu}.
$$

<span id="page-8-1"></span>**Lemma 3.6.** Suppose the assumption [\(3.2\)](#page-5-2) holds. Then there exists a continuous function  $C(t)$ defined on  $[0, \infty)$  such that for all  $t \in [0, T_0)$ ,

<span id="page-8-0"></span>
$$
|\tilde{h}_b^a|_{\tilde{\gamma}_t} \leq |\tilde{h}_b^a|_{\tilde{\gamma}_0} + C(t).
$$

*Proof.* Using [\(A.9\)](#page-16-1) with  $\Psi = \phi^{-1} \frac{d\phi}{ds}$ ,  $\psi = 2 \log \phi$  and [\(1.6\)](#page-1-5), we derive that non-zero components of  $R_{\alpha\beta\lambda\mu}$  are

$$
\widetilde{R}_{q\xi q\xi} = -\phi^6 (2 + \phi^{-3})(1 - \phi^{-3}), \n\widetilde{R}_{q\theta q\theta} = -\phi^6 (2 + \phi^{-3}), \n\widetilde{R}_{\xi \theta \xi \theta} = -\phi^6 (4 - \phi^{-3})(1 - \phi^{-3}).
$$

In particular, the sectional curvature of  $\tilde{g}$  is non-positive. From the view of [\(3.8\)](#page-8-0),  $\tilde{h}^a_b$  remains non-negative definite under the assumption [\(3.2\)](#page-5-2). From [\(3.5\)](#page-5-6), there exists a continuous function  $C(t)$ such that for any  $t \in [0, T_0)$ ,

$$
\left| (dx^a)_{\alpha} \tilde{\nu}^{\beta} \left( \frac{\partial}{\partial x^b} \right)^{\lambda} \tilde{\nu}^{\mu} \tilde{R}^{\alpha}_{ \; \; \beta \lambda \mu } \right| \leq C(t).
$$

Applying the ODE comparison to [\(3.8\)](#page-8-0), we have an upper bound for the maximum eigenvalue of  $\tilde{h}^a_b$ in any finite time. The proof is finished.

*Proof of Lemma [3.2.](#page-5-3)* Recall that  $v_t$  is the height function of  $\Sigma_t$  in the s coordinate and  $u_t = q(v_t)$ . It suffices to show that for any  $T < \infty$ , there exists constants  $C_j(T)$ ,  $j = 0, 1, \ldots$  such that for all  $0\leq t< T$ 

$$
\left|\frac{\partial^j u_t}{\partial x^{a_1} \partial x^{a_2} \dots \partial x^{a_j}}\right| \leq C_j(T).
$$

Lemma [3.3](#page-5-7) and Lemma [3.4](#page-6-0) provide such bounds for the case  $j = 0$  and  $j = 1$  respectively. From the view of  $(3.7)$  and Lemma [A.5,](#page-17-0) Lemma [3.6](#page-8-1) shows the case  $j = 2$ . Furthermore, by differentiating [\(3.7\)](#page-7-0), it suffices to show that for all  $j \geq 1$ ,

<span id="page-9-0"></span>
$$
(3.9) \t\t |\widetilde{D}^j \widetilde{h}|_{\widetilde{\gamma}} \le C_{j+2}(T).
$$

Here  $\tilde{D}$  is the Levi-Civita connection of  $\tilde{\gamma}$ . Compute

$$
\frac{\partial}{\partial t}\tilde{\gamma}(\widetilde{D}^j\tilde{h},\widetilde{D}^j\tilde{h})=\underbrace{\left(\frac{\partial}{\partial t}\tilde{\gamma}\right)(\widetilde{D}^j\tilde{h},\widetilde{D}^j\tilde{h})}_{I}+\underbrace{\tilde{\gamma}\left(\left[\frac{\partial}{\partial t},\widetilde{D}^j\right]\tilde{h},\widetilde{D}^j\tilde{h}\right)}_{II}+\underbrace{\tilde{\gamma}\left(\widetilde{D}^j\frac{\partial}{\partial t}\tilde{h},\widetilde{D}^j\tilde{h}\right)}_{III}.
$$

Because  $\frac{\partial}{\partial t} \tilde{\gamma}_{ab} = 2\tilde{h}_{ab},$ 

$$
\mathbf{I} = \widetilde{D}^j \widetilde{h} * \widetilde{D}^j \widetilde{h} * \widetilde{h}.
$$

For any tensor N, we have the commutation relation  $[\frac{\partial}{\partial t}, \tilde{D}^j]N = \sum_{i=0}^{j-1} \tilde{D}^{j-i}\tilde{h} * \tilde{D}^iN$ . Hence

$$
II = \sum_{i=0}^{j} \widetilde{D}^{j} \widetilde{h} * \widetilde{D}^{j-i} \widetilde{h} * \widetilde{D}^{i} \widetilde{h}.
$$

Finally, from [\(3.8\)](#page-8-0),

$$
III = \sum_{i=0}^{j} \widetilde{D}^{j} \widetilde{h} * \widetilde{D}^{j-i} \widetilde{h} * \widetilde{D}^{i} \widetilde{h} + \widetilde{D}^{j} K * \widetilde{D}^{j} \widetilde{h}.
$$

Here

$$
K_{ab} = \left(\frac{\partial}{\partial x^a}\right)^{\alpha} \tilde{\nu}^{\beta} \left(\frac{\partial}{\partial x^b}\right)^{\lambda} \tilde{\nu}^{\mu} \tilde{R}_{\alpha\beta\lambda\mu}.
$$

Let

$$
\omega_j(t) = \max_{\Sigma_t} |\widetilde{D}^j \widetilde{h}|_{\widetilde{\gamma}}^2.
$$

We use induction and assume  $\omega_i(t) \leq C_i(T)$  for  $0 \leq i \leq j-1$  and  $t \in [0, T)$ . By Lemma [B.2,](#page-18-1)  $|\widetilde{D}^i K|_{\widetilde{\gamma}}$ is bounded for  $0 \leq i \leq j$ . We then deduce

$$
\frac{d}{dt}\omega_j \le C\omega_j + C.
$$

<span id="page-10-2"></span>From the ODE comparison,  $\omega_i$  remains bounded in  $t \in [0, T)$ . Thus [\(3.9\)](#page-9-0) follows.

## <span id="page-10-0"></span>4. FOUR-DIMENSIONAL WITH SYMMETRY IN  $\xi$

In this section, we fix  $n = 4$  and prove Theorem [1.2.](#page-2-1) Recall that  $F : [0, T_0) \times T^{n-1} \to M$  solves

(4.1) 
$$
\frac{\partial F}{\partial t} = \left(\phi \frac{d\phi}{ds}\right)^{-1} \nu.
$$

Here we take  $T_0 \in (0,\infty]$  to be the largest number such that [\(4.1\)](#page-10-0) has a smooth graphical solution in  $t \in [0, T_0)$ .

Equation [\(4.1\)](#page-10-0) can be viewed as the unit normal flow with respect to the conformal metric  $\bar{g} =$  $\left(\phi \frac{d\phi}{ds}\right)^2 g$ . Recall that  $\bar{h}_{ab}$  is the the second fundamental form with respect to  $\bar{g}$ . Our main assumption is

<span id="page-10-3"></span>(4.2) 
$$
\bar{h}_{ab} \text{ is non-negative definite for } \Sigma_0 \text{ and } \min_{\Sigma_0} \phi^4 \ge 1 + \frac{2}{\sqrt{3}}.
$$

We list two lemmas which allow us to prove Theorem [1.2.](#page-2-1)

<span id="page-10-1"></span>**Lemma 4.1.** Suppose  $T_0 = \infty$ . Then

<span id="page-10-5"></span>(4.3) 
$$
\limsup_{t \to \infty} Q(\Sigma_t) \leq -2m.
$$

<span id="page-10-4"></span>**Lemma 4.2.** Under the assumption [\(4.2\)](#page-10-3), we have  $T_0 = \infty$ .

Proof of Theorem [1.2.](#page-2-1) Let  $\Sigma_t$  be the solution of [\(4.1\)](#page-10-0) starting from  $\Sigma_0 = \Sigma$ . Combining Lemmas [2.1,](#page-3-1) [4.2](#page-10-4) and [4.1,](#page-10-1)

$$
Q(\Sigma) \le \limsup_{t \to \infty} Q(\Sigma_t) \le -2m.
$$

Furthermore, the equality implies  $Q(\Sigma_t)$  is a constant along the flow. And Lemma [2.1](#page-3-1) implies  $\Sigma$  is a coordinate torus.

We prove Lemmas [4.1](#page-10-1) and Lemma [4.2](#page-10-4) in the rest of this section. We again adapt the convention that C denotes a large constant depending on  $\Sigma_0$ . The value of C may change from line to line.

Denote by  $v_t(\xi, \theta)$  the height function of  $\Sigma_t$  and  $u_t = q(v_t)$ . Then  $v_t$  solves the equation

(4.4) 
$$
\frac{\partial v_t}{\partial t} = \rho \left( \phi(v_t) \frac{d\phi}{ds}(v_t) \right)^{-1}
$$

Here  $\rho$  is the slope of  $\Sigma_t$  defined in [\(2.4\)](#page-3-2). Because  $\Sigma_t$  are symmetric along the  $\xi$  direction, the slope  $\rho$  is given by

.

$$
\rho^2 = 1 + \delta^{ij} \frac{\partial u_t}{\partial \theta^i} \frac{\partial u_t}{\partial \theta^j}.
$$

We start with the  $C^0$  and  $C^1$  estimates for  $v_t$  and  $u_t$ .

<span id="page-11-1"></span>**Lemma 4.3.** There exists a constant C depending on  $\Sigma_0$  such that for all  $t \in [0, T_0)$ ,

<span id="page-11-0"></span>(4.5) 
$$
|\phi^2(v_t) - 2t| \le C.
$$

*Proof.* At the maximum point of  $v_t$ , we have  $\rho = 1$ . Together with  $(1.6)$ ,

$$
\frac{d}{dt} \max_{T^2} \phi^2(v_t) \le 2.
$$

Similarly,

$$
\frac{d}{dt}\min_{T^2} \phi^2(v_t) \ge 2.
$$

Thus the assertion follows by taking  $C = \max_{T^2} \phi^2$  $(v_0).$ 

<span id="page-11-2"></span>**Lemma 4.4.** There exists constant C depending on 
$$
\Sigma_0
$$
 such that for  $t \in [0, T_0)$ ,

(4.6) 
$$
\max_{\Sigma_t} (\rho^2 - 1) \le (C^{-1}t + 1)^{-3} \max_{\Sigma_0} (\rho^2 - 1).
$$

Proof. From

$$
\frac{\partial^2 u_t}{\partial x^a \partial t} = \phi^{-2} \left( \frac{d\phi}{ds} \right)^{-1} \frac{\partial \rho}{\partial x^a} + \rho \phi^{-2} (-3 + \phi^{-4}) (1 - \phi^{-4})^{-1} \frac{\partial u_t}{\partial x^a},
$$

we deduce

$$
\frac{\partial}{\partial t}(\rho^2 - 1) = 2\phi^{-2} \left(\frac{d\phi}{ds}\right)^{-1} \delta^{ij} \frac{\partial u_t}{\partial \theta^i} \frac{\partial \rho}{\partial \theta^j} + \rho \phi^{-2}(-3 + \phi^{-4})(1 - \phi^{-4})^{-1}(\rho^2 - 1).
$$

At the maximum point of  $\rho^2 - 1$ ,  $d\rho = 0$  and

$$
\frac{\partial}{\partial t}(\rho^2 - 1) \leq \rho \phi^{-2}(-3 + \phi^{-4})(1 - \phi^{-4})^{-1}(\rho^2 - 1).
$$

Using [\(4.5\)](#page-11-0), there exists a constant C such that for all  $t \in [0, T_0)$ ,

$$
\rho\phi^{-2}(-3+\phi^{-4})(1-\phi^{-4})^{-1} \ge \phi^{-2}(-3+\phi^{-4})(1-\phi^{-4})^{-1} \ge -3(t+C)^{-1}.
$$

Hence

$$
\frac{d}{dt} \max_{\Sigma_t} (\rho^2 - 1) \leq -3(t + C)^{-1} \max_{\Sigma_t} (\rho^2 - 1).
$$

By the ODE comparison,

$$
\max_{\Sigma_t}(\rho^2 - 1) \le (C^{-1}t + 1)^{-3} \max_{\Sigma_0}(\rho^2 - 1).
$$

The proof is finished.  $\square$ 

Recall that  $g' = \phi^{-2}g$  is a conformal metric which asymptotic to the flat metric on  $\mathbb{R} \times T^3$ . Let  $\gamma'_t$ be the induced metric of g' on  $\Sigma_t$  and D' be the Levi-Civita connection of  $\gamma'_t$ . Lemmas [4.3](#page-11-1) and [4.4](#page-11-2) imply, provided  $T_0 = \infty$ ,  $\gamma'_t$  converges the the flat metric  $d\xi^2 + d\theta^2$  in  $C^0$ . Next, we use the second fundamental form  $h'_{ab}$  to bound the hessian of  $u_t$  from below.

**Lemma 4.5.** Suppose  $T_0 = \infty$ . Then there exists a constant C depending on  $\Sigma_0$  such that for  $t \geq C+1$ ,

$$
D'_b(D')^a u_t \ge -\left(C(t-C)^{-3/2}\log t\right)\delta_b^a.
$$

Proof. The flow [\(4.1\)](#page-10-0) can be rewritten as

$$
\frac{\partial F}{\partial t} = \left(\phi^2 \frac{d\phi}{ds}\right)^{-1} \nu'
$$

.

Here  $\nu' = \phi \nu$  is the unit normal vector with respect to g'. Let  $G = \phi^2 \frac{d\phi}{ds}$ . Then the evolution equation of  $(h')_b^a$  is given by

$$
\frac{\partial}{\partial t}(h')^a_b = -G^{-1}(h')^c_b(h')^a_c - D'_b(D')^a G^{-1} - G^{-1}(dx^a)_{\alpha}(\nu')^{\beta} \left(\frac{\partial}{\partial x^b}\right)^{\lambda} (\nu')^{\mu} (R')^{\alpha}_{\beta\lambda\mu}.
$$

Here  $(R')_{\alpha\beta\lambda\mu}$  is the Riemannian curvature tensor of g'. In the estimates below, the norms are computed with respect to  $\gamma'_t$ . From [\(A.3\)](#page-14-1) and [\(1.6\)](#page-1-5), the only non-zero component of R' is

$$
R'_{q\xi q\xi} = 6\phi^{-2}(1 - \phi^{-4}).
$$

Together with  $(4.5)$ , for t large enough,

$$
\left| -G^{-1} (dx^a)_{\alpha} (\nu')^{\beta} \left( \frac{\partial}{\partial x^b} \right)^{\lambda} (\nu')^{\mu} (R')^{\alpha}_{\beta \lambda \mu} \right| \leq C (t - C)^{-5/2}.
$$

Using [\(2.1\)](#page-2-3),

$$
-D'_{b}(D')^{a}G^{-1} = G^{-2}\frac{dG}{dq}D'_{b}(D')^{a}u_{t} + \frac{d}{dq}\left(G^{-2}\frac{dG}{dq}\right)D'_{b}u_{t}(D')^{a}u_{t}
$$

The relation between  $h'_{ab}$  and  $D'_b D'_a u_t$  is given in Corollary [A.3](#page-16-0) with  $\Psi = \phi^{-1} \frac{d\phi}{ds}$ . Using [\(1.6\)](#page-1-5), we derive

<span id="page-12-0"></span>(4.7) 
$$
D'_b D'_a u_t = -\rho^{-1} h'_{ab} + 2\phi^{-3} (1 - \phi^{-4})^{1/2} D'_b \xi D'_a \xi.
$$

Together with

$$
G^{-2}\frac{dG}{dq} = \phi^{-2}(3 - \phi^{-4})(1 - \phi^{-4})^{-1},
$$

$$
\frac{d}{dq}\left(G^{-2}\frac{dG}{dq}\right) = -2\phi^{-1}(3 + \phi^{-8})(1 - \phi^{-4})^{-3/2},
$$

we derive

$$
-D'_{b}(D')^{a}G^{-1} = -\rho^{-1}\phi^{-2}(3-\phi^{-4})(1-\phi^{-4})^{-1}(h')^{a}_{b} - 2\phi^{-1}(3+\phi^{-8})(1-\phi^{-4})^{-3/2}D'_{b}u_{t}(D')^{a}u_{t}
$$
  
+  $2\phi^{-5}(3-\phi^{-4})(1-\phi^{-4})^{-1/2}D'_{b}\xi(D')^{a}\xi.$ 

From  $(3.5)$  and  $(3.6)$ , for t large enough,

$$
|2\phi^{-5}(3-\phi^{-4})(1-\phi^{-4})^{-1/2}D_{b}'\xi(D')^{a}\xi| \leq C(t-C)^{-5/2}.
$$

Putting the above together, we derive

$$
\left| \frac{\partial}{\partial t} (h')_b^a + \rho^{-1} \phi^{-2} (3 - \phi^{-4}) (1 - \phi^{-4})^{-1} (h')_b^a + G^{-1} (h')_b^c (h')_c^a
$$
  
+  $2\phi^{-1} (3 + \phi^{-8}) (1 - \phi^{-4})^{-3/2} D'_b u_t (D')^a u_t \right| \le C(t - C)^{-5/2}.$ 

Let  $w_t$  be the maximum eigenvalue of  $(h')_a^b$ . From [\(3.5\)](#page-5-6) and [\(3.6\)](#page-6-1), for t large enough,

$$
\rho^{-1}\phi^{-2}(3-\phi^{-4})(1-\phi^{-4}) \ge \frac{3}{2}(t+C)^{-1}.
$$

.

We derive

$$
\frac{dw_t}{dt} \le -\frac{3}{2}(t+C)^{-1}w_t + C(t-C)^{-5/2}.
$$

From the ODE comparison,  $w_t \leq C(t-C)^{-3/2} \log t$  for t large enough. Hence the assertion follows  $(4.7).$  $(4.7).$ 

*Proof of Lemma [4.1.](#page-10-1)* From  $(2.5)$ ,

$$
Q(\Sigma_t) + 2m = \int_{T^3} \rho^2 \phi \frac{d\phi}{ds} \Delta' u_t - 2\delta^{ij} \frac{\partial u_t}{\partial \theta^i} \frac{\partial u_t}{\partial \theta^j} d\xi \wedge d\theta^3 \wedge d\theta^4.
$$

From Lemma [4.4,](#page-11-2)  $\delta^{ij} \frac{\partial u_t}{\partial \theta^i} \frac{\partial u_t}{\partial \theta^j} \leq C(t-C)^{-3}$ . It remains to prove

$$
\limsup_{t \to \infty} \int_{T^3} \rho^2 \phi \frac{d\phi}{ds} \Delta' u_t d\xi \wedge d\theta^3 \wedge d\theta^4 \le 0.
$$

Recall that  $\gamma'_t$  is the induced metric of  $\Sigma$  from g'. Because

$$
d\text{vol}_{\gamma'_t} = \rho \phi^{-1} \frac{d\phi}{ds} d\xi \wedge d\theta^3 \wedge d\theta^4,
$$

$$
\int_{T^3} \rho^2 \phi \frac{d\phi}{ds} \Delta' u_t d\xi \wedge d\theta^3 \wedge d\theta^4 = \int_{\Sigma_t} \rho \phi^2 \Delta' u_t d\mathrm{vol}_{\gamma'_t} = \underbrace{\int_{\Sigma_t} (\rho - 1) \phi^2 \Delta' u_t d\mathrm{vol}_{\gamma'_t}}_{\text{I}} + \underbrace{\int_{\Sigma_t} \phi^2 \Delta' u_t d\mathrm{vol}_{\gamma'_t}}_{\text{II}}.
$$

From Lemmas [4.3,](#page-11-1) [4.4](#page-11-2) and [4.6,](#page-13-0)  $\phi^2 \leq Ct$ ,  $|\rho - 1| \leq Ct^{-3}$  and

$$
\int_{\Sigma_t} |\Delta' u_t| \, d\text{vol}_{\gamma'} = 2 \int_{\Sigma_t} (\Delta' u_t)_- \, d\text{vol}_{\gamma'} \leq C t^{-3/2} \log t.
$$

Hence the first term I goes to zero. Through integration by parts,

$$
II = -\int_{\Sigma_t} 2\phi \frac{d\phi}{ds} |D'u_t|^2_{\gamma'_t} d\mathrm{vol}_{\gamma'} \le 0.
$$

Thus [\(4.3\)](#page-10-5) follows.

Lastly, we prove Lemma [4.2.](#page-10-4) Recall that  $\bar{g} = \left(\phi \frac{d\phi}{ds}\right)^2 g$  and that  $\bar{h}_b^a$  is the Weingarten tensor with respect to  $\bar{g}$ . The flow [\(4.1\)](#page-10-0) is the unit normal flow with respect to  $\bar{g}$ . The evolution equation of  $\bar{h}^a_b$ is given by

(4.8) 
$$
\frac{\partial}{\partial t} \bar{h}_b^a = -\bar{h}_b^c \bar{h}_c^a - (dx^a)_{\alpha} \bar{\nu}^{\beta} \left(\frac{\partial}{\partial x^b}\right)^{\lambda} \bar{\nu}^{\mu} \overline{R}^{\alpha}_{\ \beta \lambda \mu}.
$$

<span id="page-13-0"></span>**Lemma 4.6.** Under the assumption [\(4.2\)](#page-10-3), there exists a continuous function  $C(t)$  defined on  $[0,\infty)$ such that for all  $t \in [0, T_0)$ ,

$$
|\bar{h}_b^a|_{\bar{\gamma}_t} \le |\bar{h}_b^a|_{\bar{\gamma}_0} + C(t)
$$



*Proof.* Using [\(A.9\)](#page-16-1) with  $\Psi = \phi^{-1} \frac{d\phi}{ds}$  and  $\psi = 2 \log \phi + \log \frac{d\phi}{ds}$ , we derive that non-zero components of  $\overline{R}_{\alpha\beta\lambda\mu}$  are

$$
\overline{R}_{q\xi q\xi} = -3\phi^8 (1 - \phi^{-4})^3,
$$
  
\n
$$
\overline{R}_{qiqi} = -\phi^8 (3 - 6\phi^{-4} - \phi^{-8}),
$$
  
\n
$$
\overline{R}_{\xi i\xi i} = -\phi^8 (9 - \phi^{-8}) (1 - \phi^{-4}),
$$
  
\n
$$
\overline{R}_{ijij} = -\phi^8 (1 - \phi^{-4})^2, i \neq j.
$$

Besides  $\overline{R}_{qiqi}$ , other sectional curvatures are apparently non-positive. Actually, the polynomial  $3x^2 6x - 1$  has roots  $x = 1 \pm \frac{2}{\sqrt{2}}$  $\frac{1}{3}$ . Under the assumption [4.2,](#page-10-3) we have  $\phi^4 \geq 1 + \frac{2}{\sqrt{3}}$  $\frac{1}{3}$  along the flow. Hence the term

$$
-(dx^a)_{\alpha}\bar{\nu}^{\beta}\left(\frac{\partial}{\partial x^b}\right)^{\lambda}\bar{\nu}^{\mu}\overline{R}^{\alpha}_{\ \beta\lambda\mu}
$$

is non-negative definite and  $\bar{h}^a_b$  also remains non-negative definite . On the other hand, from [\(4.5\)](#page-11-0), for any  $t \in [0, T_0)$ ,

$$
\left| (dx^a)_{\alpha} \bar{\nu}^{\beta} \left( \frac{\partial}{\partial x^b} \right)^{\lambda} \bar{\nu}^{\mu} \overline{R}^{\alpha}_{\ \beta \lambda \mu} \right| \leq C(t).
$$

for some function  $C(t)$ . Then the assertion follows by the ODE comparison.

<span id="page-14-0"></span>With Lemma [4.6,](#page-13-0) the proof of Lemma [4.2](#page-10-4) is similar to the one for Lemma [3.2.](#page-5-3)

## Appendix A. Curvature and Second Fundamental Form

In this appendix, we record formulas of curvatures and second fundamental forms. Let  $\Psi(q)$  be a smooth and positive function. Consider the metric

(A.1) 
$$
g' = dq^2 + \Psi(q)^2 d\xi^2 + \sum_{i=3}^n (d\theta^i)^2.
$$

The non-zero components of the Christoffel symbols are

<span id="page-14-2"></span>(A.2) 
$$
(\Gamma')_{q\xi}^{\xi} = \Psi^{-1} \frac{d\Psi}{dq}, (\Gamma')_{\xi\xi}^{q} = -\Psi \frac{d\Psi}{dq}.
$$

The only non-zero component of the curvature is

<span id="page-14-1"></span>(A.3) 
$$
R'_{q\xi q\xi} = -\Psi \frac{d^2 \Psi}{dq^2}.
$$

We now consider a hypersurface given by a graph over  $T^{n-1}$ . Let  $u(\xi, \theta^i)$  be a smooth function and consider the map  $F(\xi, \theta^i) = (u(\xi, \theta^i), \xi, \theta^i)$ . Denote the image of F by  $\Sigma$ .

$$
\frac{\partial F}{\partial \xi} = \frac{\partial}{\partial \xi} + \frac{\partial u}{\partial \xi} \frac{\partial}{\partial q}, \ \frac{\partial F}{\partial \theta^i} = \frac{\partial}{\partial \theta^i} + \frac{\partial u}{\partial \theta^i} \frac{\partial}{\partial q}.
$$

Let  $\{x^a\} = \{\xi, \theta^i\}$  and define a metric on  $T^{n-1}$  as

(A.4) 
$$
\hat{\gamma}_{ab} dx^a dx^b = \Psi(u)^2 d\xi^2 + \delta_{ij} d\theta^i d\theta^j.
$$

The induced metric  $\gamma'_{ab}$  on  $\Sigma$  is given by

(A.5) 
$$
\gamma'_{ab} = \hat{\gamma}_{ab} + \frac{\partial u}{\partial x^a} \frac{\partial u}{\partial x^b}.
$$

The normal vector of  $\Sigma$  is given by

<span id="page-15-0"></span>(A.6) 
$$
\nu' = \rho^{-1} \left( \frac{\partial}{\partial q} - \frac{1}{\Psi(u)^2} \frac{\partial u}{\partial \xi} \frac{\partial}{\partial \xi} - \delta^{ij} \frac{\partial u}{\partial \theta^i} \frac{\partial}{\partial \theta^j} \right),
$$

where  $\rho$  is the slope given by

(A.7) 
$$
\rho^2 = 1 + \Psi(u)^{-2} \left(\frac{\partial u}{\partial \xi}\right)^2 + \delta^{ij} \frac{\partial u}{\partial \theta^i} \frac{\partial u}{\partial \theta^j}.
$$

<span id="page-15-1"></span>**Lemma A.1.** The second fundamental form of  $\Sigma$  is given by

$$
h'_{\xi\xi} = -\rho^{-1} \frac{\partial^2 u}{\partial \xi^2} + 2\rho^{-1} \Psi^{-1} \frac{d\Psi}{dq} \left(\frac{\partial u}{\partial \xi}\right)^2 + \rho^{-1} \Psi \frac{d\Psi}{dq},
$$
  
\n
$$
h'_{ij} = -\rho^{-1} \frac{\partial^2 u}{\partial \theta^i \partial \theta^j},
$$
  
\n
$$
h'_{\xi i} = -\rho^{-1} \frac{\partial^2 u}{\partial \xi \partial \theta^i} + \rho^{-1} \Psi^{-1} \frac{d\Psi}{dq} \frac{\partial u}{\partial \xi} \frac{\partial u}{\partial \theta^i}.
$$

*Proof.* Let  $\nabla'$  be the Levi-Civita connection of  $g'$ . From  $(A.2)$ ,

$$
\nabla'_{\frac{\partial F}{\partial \xi}} \frac{\partial F}{\partial \xi} = \left( \frac{\partial^2 u}{\partial \xi^2} - \Psi \frac{d\Psi}{dq} \right) \frac{\partial}{\partial q} + 2\Psi^{-1} \frac{d\Psi}{dq} \frac{\partial u}{\partial \xi} \frac{\partial}{\partial \xi},
$$
  

$$
\nabla'_{\frac{\partial F}{\partial \theta^i}} \frac{\partial F}{\partial \theta^j} = \frac{\partial^2 u}{\partial \theta^i \theta^j} \frac{\partial}{\partial q},
$$
  

$$
\nabla'_{\frac{\partial F}{\partial \theta^i}} \frac{\partial F}{\partial \xi} = \frac{\partial^2 u}{\partial \xi \partial \theta^i} \frac{\partial}{\partial q} + \Psi^{-1} \frac{d\Psi}{dq} \frac{\partial u}{\partial \theta^i} \frac{\partial}{\partial \xi}.
$$

Taking inner product with  $\nu'$  given in [\(A.6\)](#page-15-0), the assertion follows.

We now view u as a function defined on  $\Sigma$  and compute its hessian. Let  $D'$  be the Levi-Civita connection of  $\gamma'$ .

# <span id="page-15-2"></span>Lemma A.2.

$$
D_{\xi}'D_{\xi}'u = \rho^{-2}\frac{\partial \tilde{u}}{\partial \xi^{2}} - 2\rho^{-2}\Psi^{-1}\frac{d\Psi}{dq}\left(\frac{\partial u}{\partial \xi}\right)^{2} - \rho^{-2}\Psi\frac{d\Psi}{dq} + \Psi\frac{d\Psi}{dq},
$$
  
\n
$$
D_{i}'D_{j}'u = \rho^{-2}\frac{\partial^{2}u}{\partial \theta^{i}\partial \theta^{j}},
$$
  
\n
$$
D_{\xi}'D_{i}'u = \rho^{-2}\frac{\partial^{2}u}{\partial \xi\partial \theta^{i}} - \rho^{-2}\Psi^{-1}\frac{d\Psi}{dq}\frac{\partial u}{\partial \xi}\frac{\partial u}{\partial \theta^{i}}.
$$

*Proof.* Let  $\hat{\Gamma}_{ab}^c$  be the Christoffel symbols of  $\hat{\gamma}_{ab}$ . The non-zero components of  $\hat{\Gamma}_{ab}^c$  are

$$
\hat{\Gamma}_{\xi i}^{\xi} = \Psi^{-1} \frac{d\Psi}{dq} \frac{\partial u}{\partial \theta^{i}}, \ \hat{\Gamma}_{\xi \xi}^{i} = -\Psi \frac{d\Psi}{dq} \frac{\partial u}{\partial \theta^{i}}, \ \hat{\Gamma}_{\xi \xi}^{\xi} = \Psi^{-1} \frac{d\Psi}{dq} \frac{\partial u}{\partial \xi}.
$$

For any smooth function  $\tilde{u}$ ,

$$
\hat{D}_{\xi}\hat{D}_{\xi}\tilde{u} = \frac{\partial \tilde{u}}{\partial \xi^{2}} + \Psi \frac{d\Psi}{dq} \delta^{ij} \frac{\partial u}{\partial \theta^{i}} \frac{\partial \tilde{u}}{\partial \theta^{j}} - \Psi^{-1} \frac{d\Psi}{dq} \frac{\partial u}{\partial \xi} \frac{\partial \tilde{u}}{\partial \xi},
$$
\n
$$
\hat{D}_{i}\hat{D}_{j}\tilde{u} = \frac{\partial^{2}\tilde{u}}{\partial \theta^{i}\partial \theta^{j}},
$$
\n
$$
\hat{D}_{\xi}\hat{D}_{i}\tilde{u} = \frac{\partial^{2}\tilde{u}}{\partial \xi\partial \theta^{i}} - \Psi^{-1} \frac{d\Psi}{dq} \frac{\partial u}{\partial \theta^{i}} \frac{\partial \tilde{u}}{\partial \xi}.
$$

Here  $\hat{D}$  is the Levi-Civita connection for  $\hat{\gamma}$ . Because  $\gamma'_{ab} = \hat{\gamma}_{ab} + \frac{\partial u}{\partial x^a} \frac{\partial u}{\partial x^b}$ , we have

$$
D'_a D'_b u = \rho^{-2} \hat{D}_a \hat{D}_b u.
$$

Combining the above, the assertion for  $D'_i D'_j u$  and  $D'_\xi D'_i u$  follows. For  $D'_\xi D'_\xi u$ , we use

$$
\delta^{ij}\frac{\partial u}{\partial \theta^i}\frac{\partial u}{\partial \theta^j} = -\Psi^{-2}\left(\frac{\partial u}{\partial \xi}\right)^2 - 1 + \rho^2
$$

.

Then

$$
D_{\xi}'D_{\xi}'u = \rho^{-2}\hat{D}_{\xi}\hat{D}_{\xi}u = \rho^{-2}\left(\frac{\partial\tilde{u}}{\partial\xi^{2}} + \Psi\frac{d\Psi}{dq}\delta^{ij}\frac{\partial u}{\partial\theta^{i}}\frac{\partial u}{\partial\theta^{j}} - \Psi^{-1}\frac{d\Psi}{dq}\left(\frac{\partial u}{\partial\xi}\right)^{2}\right)
$$
  

$$
= \rho^{-2}\left(\frac{\partial\tilde{u}}{\partial\xi^{2}} - 2\Psi^{-1}\frac{d\Psi}{dq}\left(\frac{\partial u}{\partial\xi}\right)^{2} - \Psi\frac{d\Psi}{dq} + \rho^{2}\Psi\frac{d\Psi}{dq}\right)
$$
  

$$
= \rho^{-2}\frac{\partial\tilde{u}}{\partial\xi^{2}} - 2\rho^{-2}\Psi^{-1}\frac{d\Psi}{dq}\left(\frac{\partial u}{\partial\xi}\right)^{2} - \rho^{-2}\Psi\frac{d\Psi}{dq} + \Psi\frac{d\Psi}{dq}.
$$

From Lemmas [A.1](#page-15-1) and [A.2,](#page-15-2) we obtain the following relation between  $h'_{ab}$  and  $D'_b D'_b u$ .

# <span id="page-16-0"></span>Corollary A.3.

<span id="page-16-2"></span>(A.8) 
$$
h'_{ab} = -\rho D'_a D'_b u + \rho \Psi \frac{d\Psi}{dq} D'_a \xi D'_b \xi.
$$

Next, we record the formulas under a conformal transformation. Let  $\psi(q)$  be a smooth function of q. Define

$$
\check{g} = e^{2\psi}g'.
$$

Let  $\check{R}_{\alpha\beta\lambda\mu}$  be the Riemannian curvature of  $\check{g}$ .

**Lemma A.4.** The non-zero components of  $\check{R}_{\alpha\beta\lambda\mu}$  are

<span id="page-16-1"></span>
$$
\tilde{R}_{q\xi q\xi} = -e^{2\psi} \left( \Psi \frac{d^2 \Psi}{dq^2} + \Psi^2 \frac{d^2 \psi}{dq^2} + \Psi \frac{d\Psi}{dq} \frac{d\psi}{dq} \right),
$$
\n
$$
\tilde{R}_{qiqi} = -e^{2\psi} \frac{d^2 \psi}{dq^2},
$$
\n
$$
\tilde{R}_{\xi i\xi i} = -e^{2\psi} \Psi \frac{d\Psi}{dq} \frac{d\psi}{dq} - e^{2\psi} \Psi^2 \left( \frac{d\psi}{dq} \right)^2,
$$
\n
$$
\tilde{R}_{ijij} = -e^{2\psi} \left( \frac{d\psi}{dq} \right)^2, \quad i \neq j.
$$

 $\Box$ 

*Proof.* As  $\check{g} = e^{2\psi} g'$ , the formula of  $\check{R}_{\alpha\beta\lambda\mu}$  is given by [\[2,](#page-19-10) page 58]

$$
\check{R}_{\alpha\beta\lambda\mu} = e^{2\psi} R'_{\alpha\beta\lambda\mu} - e^{2\psi} (g'_{\alpha\lambda} T_{\beta\mu} + g'_{\beta\mu} T_{\alpha\lambda} - g'_{\alpha\mu} T_{\beta\lambda} - g'_{\beta\lambda} T_{\alpha\mu}),
$$

with  $T_{\alpha\beta} = \nabla'_{\alpha}\nabla'_{\beta}\psi - \nabla'_{\alpha}\psi\nabla'_{\beta}\psi + 2^{-1}|\nabla'\psi|^2_{g'}g'_{\alpha\beta}$ . From [\(A.2\)](#page-14-2), non-zero components of  $T_{\alpha\beta}$  are

$$
T_{qq} = \frac{d^2\psi}{dq^2} - 2^{-1} \left(\frac{d\psi}{dq}\right)^2,
$$
  
\n
$$
T_{\xi\xi} = \Psi \frac{d\Psi}{dq} \frac{d\psi}{dq} + 2^{-1} \Psi^2 \left(\frac{d\psi}{dq}\right)^2,
$$
  
\n
$$
T_{ii} = 2^{-1} \left(\frac{d\psi}{dq}\right)^2.
$$

Combining with  $(A.3)$ , the assertion follows a direct computation.

We turn to the second fundamental form of  $\Sigma$  with respect to  $\check{g}$ . The normal vector and the induced metric are given by  $\check{\nu} = e^{-\psi} \nu'$  and  $\check{\gamma} = e^{2\psi} \gamma'$  respectively. Let  $\check{h}_{ab}$  be the second fundamental form with respect to  $\check{g}$ .

# <span id="page-17-0"></span>Lemma A.5.

<span id="page-17-1"></span>(A.10) 
$$
\check{h}_{ab} = e^{\psi} \left( h'_{ab} + \rho^{-1} \frac{d\psi}{dq} \gamma'_{ab} \right).
$$

*Proof.* Let  $\check{\nabla}$  be the Levi-Civita connection of  $\check{q}$ . From

$$
\tilde{\nabla}_X Y = \nabla'_X Y + g'(\nabla' \psi, X) Y + g'(\nabla' \psi, Y) X - g'(X, Y) \nabla' \psi,
$$
  

$$
\tilde{h}_{ab} = -\tilde{g} \left( \tilde{\nabla}_{\frac{\partial F}{\partial x^a}} \frac{\partial F}{\partial x^b}, \tilde{\nu} \right) = -e^{\psi} g' \left( \nabla'_{\frac{\partial F}{\partial x^a}} \frac{\partial F}{\partial x^b} - \gamma'_{ab} \frac{d\psi}{dq} \frac{\partial}{\partial q}, \nu \right) = e^{\psi} \left( h'_{ab} + \rho^{-1} \frac{d\psi}{dq} \gamma'_{ab} \right).
$$

Using  $(A.8)$ ,  $(A.10)$ , we have

$$
\check{h}_{ab} = e^{\psi} \left( -\rho D_a' D_b' u + \rho \Psi \frac{d\Psi}{dq} D_a' \xi D_b' \xi + \rho^{-1} \frac{d\psi}{dq} \gamma_{ab}' \right).
$$

To get the mean curvature  $\check{H}$ , we need to calculate  $|D'\xi|^2_{\gamma'}$ . Recall that

$$
\gamma'_{ab} = \hat{\gamma}_{ab} + \frac{\partial u}{\partial x^a} \frac{\partial u}{\partial x^b}.
$$

We deduce

$$
(\gamma')^{ab} = \hat{\gamma}^{ab} - \rho^{-2} \hat{\gamma}^{ac} \hat{\gamma}^{bd} \frac{\partial u}{\partial x^c} \frac{\partial u}{\partial x^d}.
$$

Hence

$$
|D'\xi|_{\gamma'}^2 = (\gamma')^{\xi\xi} = \Psi^{-2} - \rho^{-2}\Psi^{-4} \left(\frac{\partial u}{\partial \xi}\right)^2
$$

$$
= \rho^{-2}\Psi^{-2} \left(1 + \delta^{ij}\frac{\partial u}{\partial \theta^i}\frac{\partial u}{\partial \theta^j}\right).
$$

With  $\check{\gamma}^{ab} = e^{-2\psi} (\gamma')^{ab}$ , we obtain

 $\Box$ 

**Corollary A.6.** The mean curvature of  $\Sigma$  with respect to  $\check{g}$  is given by

<span id="page-18-0"></span>(A.11) 
$$
\check{H} = e^{-\psi} \left( -\rho \Delta' u + \rho^{-1} \Psi^{-1} \frac{d\Psi}{dq} \delta^{ij} \frac{\partial u}{\partial \theta^i} \frac{\partial u}{\partial \theta^j} + (n-1)\rho^{-1} \frac{d\psi}{dq} + \rho^{-1} \Psi^{-1} \frac{d\Psi}{dq} \right).
$$

Here  $\Delta'$  is the Laplace-Beltrami operator of  $\gamma'$ .

## Appendix B. Projection

Let M be a n-dimensional manifold with coordinates  $\{y^{\alpha}\}_{\alpha=1}^{n}$ . Let  $g_{\alpha\beta}$  be a Riemannian metric on M and  $\Sigma^{n-1}$  be a hypersurface. We use  $\{x^a\}_{a=1}^n$  to denote a local coordinates of  $\Sigma$ . Denote by  $\nu^{\alpha}$  the unit normal vector of  $\Sigma$ , by  $\gamma_{ab}$  the induced metric on  $\Sigma$  and by  $h_{ab}$  the second fundamental form of  $\Sigma$  with respect to  $\nu^{\alpha}$ . Define the projection tensor

$$
P_{\beta}^{\alpha} = \delta_{\beta}^{\alpha} - \nu_{\beta} \nu^{\alpha}.
$$

Denote by  $\nabla$  and  $D$  the Levi-Civita connection of  $g_{\alpha\beta}$  and  $\gamma_{ab}$  respectively.

<span id="page-18-2"></span>**Lemma B.1.** Let  $\overline{F}_{\alpha}$  be a one-form and  $F_a = P_a^{\alpha} \overline{F}_{\alpha}$  be the projection of  $\overline{F}_{\alpha}$ . Then

(B.2) 
$$
D_a F_b = P_a^{\alpha} P_b^{\beta} \nabla_{\alpha} \overline{F}_{\beta} - \overline{F}_{\alpha} \nu^{\alpha} h_{ab}.
$$

(B.3) 
$$
D_a(F_\alpha \nu^\alpha) = (P_a^\beta \nabla_\beta \overline{F}_\alpha) \nu^\alpha + F_b h_a^b.
$$

Let

(B.4) 
$$
K_{ab} = P_a^{\alpha} \nu^{\beta} P_b^{\lambda} \nu^{\mu} R_{\alpha\beta\lambda\mu}.
$$

<span id="page-18-1"></span>**Lemma B.2.** For any  $m \in \mathbb{N} \cup \{0\}$  there exists a degree  $m+1$  polynomial  $p_m$  such that the following holds. Suppose there exists a constant C such that for all  $0 \le j \le m$ ,

<span id="page-18-3"></span>
$$
|\nabla^j Rm|_g \le C, \ |D^{j-1}h|_{\gamma} \le C.
$$

Then

$$
|D^m K|_{\gamma} \le p_m(C).
$$

*Proof.* For  $m, k \in \mathbb{N} \cup \{0\}$ , let  $L_{m,k}$  be the collection  $4+m-k$  tensors on  $\Sigma$  obtained from projections of  $\nabla^m Rm \cdot \nu^{\otimes k}$ . We adapt the convention that  $L_{m,-1} = 0$  and  $L_{m,k} = 0$  if  $k \geq m+3$ . Note that  $K \in L_{0,2}$ . From Lemma [B.1,](#page-18-2) we have

$$
DL_{m,k} = L_{m+1,k} + L_{m,k-1} * h + L_{m,k+1} * h.
$$

Here we write  $L * h$  for any linear combination formed by contracting L and h by  $\gamma$ . Through induction, we have for all  $m \in \mathbb{N} \cup \{0\},\$ 

$$
D^m L_{0,2} = \sum L_{j,i} * D^{\ell_1} h * \cdots * D^{\ell_k} h,
$$

where the summation goes over

$$
0 \le i \le j+3
$$
,  $0 \le \ell_i$ , and  $j + k + \sum_{i=1}^k \ell_i = m$ .

Under the assumption [\(B.5\)](#page-18-3), each term  $L_{j,i}$  and  $D^{\ell_i}$  above are bounded by C. Each individual summand  $L_{j,i} * D^{\ell_1} h * \cdots * D^{\ell_k} h$  is bounded by  $C^{k+1}$ . Then the assertion then follows.

#### **REFERENCES**

- <span id="page-19-10"></span><span id="page-19-9"></span>[1] H. Barzegar, P. T. Chrusciel, M. Hörzinger, M. Maliborski, and L. Nguyen, On the energy of the Horowitz-Myers metrics, Phys. Rev. D., **101** (2020), 024007.
- [2] A. L. Besse, Einstein manifolds, Classics in Mathematics. Springer-Verlag, Berlin, 1987.
- <span id="page-19-3"></span>[3] S. Brendle, Constant mean curvature surfaces in warped product manifolds, Publ. Math. IHES, 117 (2013), 247–269.
- <span id="page-19-8"></span>[4] S. Brendle, P. K. Hung, and M. T. Wang, A Minkowski Inequality for Hypersurfaces in the Anti-de Sitter-Schwarzschild Manifold., Comm. Pure Appl. Math., 69 (2016), no. 1, 124–144.
- <span id="page-19-0"></span>[5] N. R. Constable, and R. C. Myers, Spin two glueballs, positive energy theorems and the AdS/CFT correspondence, J. High Energy Phys., 037 (1999).
- [6] P. Guan, X. N. Ma, N. Trudinger, and X. Zhu, A form of Alexandrov-Fenchel inequality., Pure Appl. Math. Q., 6 (2010), 999–1012.
- <span id="page-19-6"></span><span id="page-19-1"></span>[7] P. Guan, and J. Li, The quermassintegral inequalities for k-convex starshaped domain., Adv. Math. 221 (2009), 1725-1732.
- [8] G.T. Horowitz and R.C. Myers, The AdS/CFT Correspondence and a New Positive Energy Conjecture for General *Relativity*, Phys. Rev. D, **59** (1999),  $1-12$ .
- <span id="page-19-5"></span>[9] S. McCormick, On a Minkowski-like inequality for asymptotically flat static manifolds, Proc. Am. Math. Soc., 146 (2018), no. 9, 4039–4046.
- <span id="page-19-2"></span>[10] G. Huisken, and T. Ilmanen, The inverse mean curvature flow and the Riemannian Penrose inequality, J. Differential Geom., 59 (2001), 353-437.
- <span id="page-19-4"></span>[11] Yong Wei, On the Minkowski-type inequality for outward minimizing hypersurfaces in Schwarzschild space, Calc. Var. Partial Differ. Equ. 57 (2018), 46.
- <span id="page-19-7"></span>[12] E. Woolgar, The rigid Horowitz-Myers conjecture, J. High Energy Phys., 104 (2017).

#### Aghil Alaee

Department of Mathematics and Computer Science, Clark University, Worcester, MA 01610, USA Center of Mathematical Sciences and Applications, Harvard University, Cambridge, MA 02138, USA Email address: aalaeekhangha@clarku.edu, aghil.alaee@cmsa.fas.harvard.edu

#### Pei-Ken Hung

Department of Mathematics, Massachusetts Institute of Technology, Cambridge, MA 02139, USA Email address: pkhung@mit.edu