

# Control theory approach to continuous-time finite state mean field games

Yurii Averboukh<sup>a</sup>

March 1, 2025

## Abstract

In the paper, we use the equivalent formulation of a finite state mean field game as a control problem with mixed constraints to study the dependence of solutions to finite state mean field game on an initial distribution of players. We introduce the concept of value multifunction of the mean field game that is a mapping assigning to an initial time and an initial distribution a set of expected outcomes of the representative player corresponding to solutions of the mean field game. Using the control reformulation of the mean field game, we give the sufficient condition on a given multifunction to be a value multifunction in the terms of the viability theory. The maximal multifunction (i.e. the mapping assigning to an initial time and distribution the whole set of values corresponding to solutions of the mean field game) is characterized via the backward attainability set for the certain control systems.

**MSC Classification (2020):** 49N80, 91A16, 49J45, 60J27, 34H05 .

**Keywords:** continuous-time finite state mean field game, value multifunction, viability, backward attainability domain, master equation.

## 1 Introduction

The mean field game theory examines systems of identical players those interacts via some external media. It was proposed independently by Lasry, Lions [21], [22] and Huang, Caines, Malhamé [13], [14]. In the paper, we study the continuous-time finite state mean field games that is the infinite player dynamic game under assumptions that the players are similar, when the dynamics of each players is given by a continuous-time finite state Markov chain with transition probabilities depending on distribution of all players and player's actions. The finite state mean field games find various application in the analysis of socio-economic systems and modeling of cybersecurity [15], [16], [17], [18], [20].

The study of the continuous-time finite state mean field games started with papers [3], [11]. In those papers the approach based on the mean field game system consisting of Bellman equation and kynetic equation was developed. Notice that, for the finite state

---

<sup>a</sup>Higher School of Economics, Moscow, Russia, e-mail: [averboukh@gmail.com](mailto:averboukh@gmail.com)

mean field game, the mean field game system is a system of ODEs with mixed (initial and boundary) conditions. The results of the aforementioned papers were extended to the case of finite state mean field games with common noise acting in a finite sequence of times in [8].

The further progress of the theory of finite state mean field games is due to the probabilistic approach [9] and master equation [4], [5], [6], [10]. The master equation first proposed by Lions [23] for the second-order mean field games is used to justify the convergence of feedback equilibria in the finite player games to the solution of the mean field game and to study the dependence of the solution of the mean field game on the initial distribution of players. However, the theory of master equation for the mean field games is developed in the case of unique solution to the mean field game [4], [5], [6], [10].

In the paper, we study the dependence of the solution of the continuous-time finite state mean field game in the general case without any uniqueness assumptions. To this end, we express the mean field game as a finite-dimensional control problem with mixed constraints. To examine the dependence of the solution of the mean field game on initial distribution, we use the concept of value multifunction that is a set-valued mapping assigning to an initial time and an initial distribution of agent a set of expected outcomes of the representative player. The maximal value multifunction comprises all solutions of the mean field game. We find the sufficient condition on a given multifunction to be a value multifunction in the terms of the viability theory. Furthermore, we characterize the maximal value function using the attainability domain for the certain dynamical system in the backward time constructed by the original mean field game. Finally, we show that a graph of smooth solution of the master equation is a value multifunction.

The paper is organized as follows. In Section 2, we introduce the main concepts of the continuous-time finite state mean field games. Section 3 is concerned with the relaxation of the control space. The control problem equivalent to the finite state mean field game is introduced in Section 4. The study of dependence of the solution of the finite state mean field game on the initial distribution is studied in Section 5. Here we introduce the mean field game dynamical system and show that a multifunction satisfying the viability condition for this dynamical system is a value multifunction. Furthermore, we prove that the maximal value multifunction can be found using the attainability domain for the certain dynamical system. The last result of Section 5 gives the link between the the solution of master equation and the value multifunction.

## 2 Mean field game system for finite state space

We consider the mean field game with the finite state space. Without loss of generality, we assume the state space is  $\{1, \dots, d\}$ , where  $d \in \mathbb{N}$ . Distributions on the space  $\{1, \dots, d\}$  lie in the  $d$ -dimensional simplex

$$\Sigma^d \triangleq \{(m_1, \dots, m_d) : m_i \geq 0, \quad m_1 + \dots + m_d = 1\}.$$

In the following, we regard  $m \in \Sigma^d$  as a row-vector and endow  $\Sigma^d$  with the Euclidean metric. Additionally, we embed  $\Sigma^d$  into the Euclidean space of row-vectors denoted below by  $\mathbb{R}^{d*}$ .

Recall that mean field games examine systems of infinitely many similar players. In the finite state case (see [3], [11]), the dynamics of each player is given by the Markov chain with the Kolmogorov matrix depending on the current distribution of agents

$$Q(t, m, u) = (Q_{i,j}(t, m, u))_{i,j=1}^d.$$

Here,  $t$  stands for a time,  $m$  is a distribution of all player,  $u$  is a control of the player chosen from some set  $U$ . We consider the mean field game on the finite time horizon  $[t_0, T]$  and assume that each player tries to maximize

$$\mathbb{E} \left[ \sigma(X(T), m(T)) + \int_{t_0}^T g(t, X(t), m(t), u(t)) dt \right], \quad (1)$$

where  $X$  is the stochastic process describing the motion of the player,  $m(t)$  is the distribution of players at time  $t$ , whereas  $u(t)$  denotes the instantaneous control of the player.

We impose the following conditions on the control space, the Kolmogorov matrix and the payoff functions.

- (C1) The set  $U$  is a metric compact.
- (C2) The functions  $Q_{i,j}$ ,  $\sigma$  and  $g$  are continuous.
- (C3) The dependence of  $Q_{i,j}$  and  $g$  on  $m$  is Lipschitz continuous.

The solution of the mean field game is given by the common solution of two problems. The first one describes the dynamics of the distribution of players, when the second problem is the optimization problem for the representative player. To introduce them, let us describe the strategies used by the players.

We assume the players are informed about current time and state. Since the state space is finite, one can assume that, a policy is now a  $d$ -tuple  $u(\cdot) = (u_i(\cdot))_{i=1}^d$ , where  $u_i(\cdot)$ ,  $i = 1, \dots, d$ , are measurable functions from  $[t_0, T]$  to  $U$ . Below, we assume that the player who occupies the state  $i$  at time  $t$  uses the control  $u_i(t)$ . Therefore, the dynamics of distribution of players obeys the following Kolmogorov equation

$$\frac{d}{dt} m_j(t) = \sum_{i=1}^d m_i(t) Q_{i,j}(t, m(t), u_i(t)), \quad j = 1, \dots, d, \quad m(t_0) = m_0. \quad (2)$$

It is convenient to rewrite this system in the vector form. To this end, given a vector of controls  $u = (u_i)_{i=1}^d \in U^d$  and  $m \in \Sigma^d$ , denote the matrix with the elements  $Q_{i,j}(t, m, u_i)$  by  $\mathcal{Q}(t, m, u)$ . Notice that now  $U^d$  serves as a new control space. Therefore, system (2) can be rewritten as follows:

$$\frac{d}{dt} m(t) = m(t) \mathcal{Q}(t, m(t), u(t)), \quad m(t_0) = m_0. \quad (3)$$

The second problem is the optimization problem for each player. Given a flow of probabilities  $m(\cdot)$ , we obtain the Markov decision problem with the dynamics determined by the Kolmogorov matrix

$$Q(t, m(t), u)$$

and objective function (1). Recall that we use the feedback strategies. Within this framework, the solution of the optimal control problem can be obtained from the dynamic programming. To introduce it for the Markov decision problem, we define the Hamiltonian by the following rule: for  $t \in [0, T]$ ,  $m \in \Sigma^d$ ,  $\phi \in \mathbb{R}^d$ , set

$$H_i(t, m, \phi) \triangleq \max_{u \in U} \left[ \sum_{j=1}^d Q_{i,j}(t, m, u) \phi_j + g(t, i, m, u) \right],$$

$$H(t, m, u) = (H_i(t, m, u))_{i=1}^d.$$

For any flow of probabilities is  $m(\cdot)$ , the value function  $\varphi : [0, T] \rightarrow \mathbb{R}^d$  satisfies the Bellman equation

$$\frac{d}{dt} \varphi(t) = -H(t, m(t), \varphi(t)), \quad \phi(T) = \sigma(i, T), \quad (4)$$

whereas the feedback optimal strategy is computed by the rule

$$\hat{u}_i(t) \in \operatorname{Argmax}_{u \in U} \left[ \sum_{j=1}^d Q_{i,j}(t, m, u) \phi_j(t) + g(t, i, m(t), u) \right]. \quad (5)$$

A solution of the system (3), (4), (5) is a solution of the finite state mean field game.

*Remark 2.1.* To study the dependence of the solution of the mean field game on the initial distribution of players, Lions proposed the concept of the master equation [23]. It was developed for the case of finite state space in [4], [6]. Let us introduce it in the multivalued setting. If  $\phi \in \mathbb{R}^d$ , then put

$$\mathcal{O}(t, m, \phi) \triangleq \left\{ m \mathcal{Q}(t, m, u) : u = (u_i)_{i=1}^d, \right. \\ \left. \sum_{j=1}^d Q_{i,j}(t, m, u_i) \phi_j(t) + g(t, i, m(t), u_i) = H_i(t, m, \phi) \right\}.$$

Since we allow nonuniqueness of the optimal controls, it is worth to write the master equation for the finite state mean field game in the multivalued manner:

$$\frac{\partial}{\partial s} \Phi(s, \mu) + H(t, \mu, \Phi(s, \mu)) \in -\mathcal{O}(s, \mu, \Phi(s, \mu)) \frac{\partial \Phi}{\partial \mu}(s, \mu). \quad (6)$$

Here  $\Phi_i(s, \mu)$  can be interpreted as the optimal outcome of the representative player who starts at the time  $s$  from the state  $i$  under assumptions that the initial distribution of agents is  $\mu$  and all players use an optimal strategy realizing a solution of the MFG;  $\partial \Phi / \partial m$  stands for the derivative of  $\Phi$  w.r.t. to  $m$ . Notice that formally this inclusion can be deduced from the mean field game system (2), (4), (5). Indeed, by construction  $\Phi(s, m(s))$  solves Bellman equation (4), when  $m(\cdot)$  satisfies (2) for the strategy  $\hat{u}$  obeying (5). The last assumption is equivalent to the inclusion

$$\frac{d}{dt} m(t) \in \mathcal{O}(t, m(t), \Phi(t, m(t))). \quad (7)$$

Differentiating formally  $\Phi(t, m(t))$  and using the Bellman equation, we obtain

$$\frac{\partial \Phi(t, m(t))}{\partial t} + H(t, m(t), \Phi(t, m(t))) + \frac{m(t)}{dt} \cdot \frac{\partial \Phi(t, m(t))}{\partial m} = 0.$$

Using this, (7) and plugging in the resulting inclusion only for the initial position  $(s, \mu)$ , we arrive at the master equation in the multivalued form (6).

### 3 Relaxation of the finite state mean field game

In the previous section, we assume that players occupying the same state use the same control. However, in the case when the Hamiltonian is not strictly convex, the optimal control is nonunique. Furthermore, the set of feedback strategies is neither convex nor closed. To overcome this difficulties, we use the relaxed (randomized) feedback strategies [12].

A feedback strategy is a mapping

$$t \mapsto \nu(t) \triangleq (\nu_1(t, du), \dots, \nu_d(t, du))$$

such that

- $\nu_i(t, \cdot)$  is a probability on  $U$ ;
- the dependence  $t \mapsto \int_U \zeta(u) \nu_i(t, du)$  is measurable for any continuous function  $\zeta : U \rightarrow \mathbb{R}$ .

The relaxed strategies imply that the players occupying the state  $i$  at the time  $t$  distribute their controls according to the probability  $\nu_i(t, du)$ .

Thus, for a fixed flow of probabilities  $m(\cdot)$  and a relaxed strategy  $\nu(\cdot)$ , the motion of a representative player is the Markov chain with the Kolmogorov matrix

$$\left( \int_U Q(t, m(t), u) \nu_i(t, du) \right)_{i,j=1}^d.$$

With some abuse of notation, for  $t \in [0, T]$ ,  $m \in \Sigma^d$ ,  $\nu = (\nu_1, \dots, \nu_d) \in (\mathcal{P}(U))^d$ , we set

$$\mathcal{Q}_{i,j}(t, m, \nu) \triangleq \left( \int_U Q_{i,j}(t, m, u) \nu_i(du) \right),$$

$$g_i(t, m, \nu) = g(t, i, m, \nu) \triangleq \int_U g_i(t, i, m, u) \nu_i(du).$$

Furthermore, notice that

$$H_i(t, m, \phi) = \max_{\nu_i \in \mathcal{P}(U)} \left[ \sum_{j=1}^q \int_U Q(t, m, u) \nu_i(du) \cdot \phi_j + g_i(t, m, \nu_i) \right].$$

Finally, we introduce the matrix and vector notation:

$$\mathcal{Q}(t, m, \nu) = (\mathcal{Q}_{i,j}(t, m, \nu))_{i,j \in \overline{1,d}}.$$

$$\begin{aligned}
H(t, m, \phi) &\triangleq (H_1(t, m, \phi), \dots, H_d(t, m, \phi))^d, \\
g(t, m, \nu) &= (g_1(t, m, \nu), \dots, g_d(t, m, \nu))^d, \\
\sigma(m) &\triangleq (\sigma(1, m), \dots, \sigma(d, m))^T.
\end{aligned}$$

Using this notation, we give the following definition.

**Definition 3.1.** Let  $t_0 \in [0, T]$  be an initial time,  $\mu_0 \in \Sigma^d$  be an initial probability. We say that the pair  $(\phi(\cdot), m(\cdot))$  is a solution of the mean field game if there exists a relaxed feedback strategy  $\hat{\nu}(\cdot) = (\hat{\nu}_1(\cdot), \dots, \hat{\nu}_d(\cdot))$  such that

1.  $m(\cdot)$  satisfies the Kolmogorov equation

$$\frac{d}{dt}m(t) = m(t)\mathcal{Q}(t, m(t), \hat{\nu}(t)), \quad m(t_0) = m_0. \quad (8)$$

2.  $\phi(\cdot)$  satisfies Bellman equation (4), i.e.,

$$\frac{d}{dt}\phi(t) = -H(t, m(t), \phi(t)), \quad \phi(T) = \sigma(m(T)) \quad (9)$$

3.  $\hat{\nu}$  is an optimal control, i.e.,

$$\hat{\nu}_i(t) \in \operatorname{Argmax}_{\nu_i \in \mathcal{P}(U)} \left[ \sum_{j=1}^q \int_U \mathcal{Q}(t, m, u) \nu_i(du) \cdot \phi_j + \int_U g(t, i, m, u) \nu_i(du) \right] \quad (10)$$

or, equivalently,

$$\frac{d}{dt}\phi(t) + \mathcal{Q}(t, m(t), \hat{\nu}(t))\phi(t) + g(t, m(t), \hat{\nu}(t)) = 0.$$

**Theorem 3.2.** *Under assumptions (C1)–(C3), there exists at least one solution of the mean field game.*

The proof is by standard fixed point arguments and relies on the convexity and compactness of the set of relaxed controls. Thus, we omit it.

Below we also will use the probabilistic representation of the finite state mean field game. It is convenient to introduce the dynamics of the representative player using generator. Let  $t \in [0, T]$ ,  $i \in \{1, \dots, d\}$ ,  $m \in \Sigma^d$ ,  $\nu \in (\mathcal{P}(U))^d$ . Define the operator  $L_t[m, \nu]$  by the following rule: for  $\psi \in \mathbb{R}^d$ ,

$$(L_t[m, \nu]\psi)(i) \triangleq \sum_{j=1}^d \mathcal{Q}_{i,j}(t, m, \nu)\psi(j).$$

As above we regard,  $(L_t[m, \nu]\psi)(i)$  as a vector in  $\mathbb{R}^d$ . Therefore,  $L_t[m, \nu]\psi$  is a mapping from  $\mathbb{R}^d$  to  $\mathbb{R}^d$ . Given,  $s, r \in [0, T]$ ,  $s < r$ , a flow of probabilities  $m(\cdot)$  and a relaxed feedback strategy  $\eta(\cdot)$ , we say that the 5-tuple  $(\Omega, \mathcal{F}, \{\mathcal{F}_t\}_{t \in [s, r]}, P, X)$ , where  $(\Omega, \mathcal{F}, \{\mathcal{F}_t\}_{t \in [s, r]}, P)$  is a filtered probability space, and  $X$  is  $\{\mathcal{F}_t\}_{t \in [s, r]}$ -adapted stochastic

process taking values in  $\{1, \dots, d\}$ , provides a motion of the representative player if, for any  $\psi \in \mathbb{R}^d$ , the process

$$\psi(X(t)) - \int_s^t (L_\tau[m(\tau), \eta(\tau)]\psi)(X(\tau))d\tau$$

is a  $\{\mathcal{F}_t\}_{t \in [s, r]}$ -martingale.

Notice that, for every initial probability on  $\{1, \dots, d\}$ , at least one motion of the representative player always exists [19]. One can introduce the probabilities  $\mu_i(t) \triangleq P(X(t) = i)$ . If  $\mu_* = (\mu_{*,i})_{i=1}^d$  is the initial distribution at the time  $s$ , then the vector  $\mu(t) = (\mu_i(t))_{i=1}^d$  obeys the Kolmogorov equation

$$\frac{d}{dt}\mu(t) = \mu\mathcal{Q}(t, m(t), \eta(t)), \quad \mu(s) = \mu_*.$$

**Proposition 3.3.** *The pair  $(\phi(\cdot), m(\cdot))$  solves the mean field game with the equilibrium feedback strategy  $\hat{\nu}(\cdot)$ , if and only if (8), (9) holds and  $\hat{\nu}(\cdot)$  is the optimal feedback relaxed strategy at any initial distribution for the Markov decision problem on  $[t_0, T]$*

$$\text{maximize } \mathbb{E} \left[ \sigma(X(T), m(T)) + \int_{t_0}^T g(t, X(t), m(t), \eta(t))dt \right] \quad (11)$$

subject to feedback relaxed  $\eta(\cdot) : [t_0, T] \rightarrow (\mathcal{P}(U))^d$  and corresponding motion of the representative player  $(\Omega, \mathcal{F}, \{\mathcal{F}_t\}_{t \in [t_0, T]}, P, X)$  at any initial position, where  $\mathbb{E}$  stands for the expectation according to the probability  $P$ .

*Proof.* It suffices to prove that the fact that  $\hat{\nu}$  is optimal control for the Markov decision problem on  $[t_0, T]$  with payoff given by (11) subject to a strategy  $\eta$  and a corresponding motion of the representative player  $(\Omega, \mathcal{F}, \{\mathcal{F}_t\}_{t \in [t_0, T]}, P, X)$  is equivalent to (10). This directly follows from the dynamic programming principle.  $\square$

Notice that the expectation of payoff (11) is determined only by  $\eta(\cdot)$ ,  $m(\cdot)$  and the initial distribution  $\mu_0$  by the formula:

$$\begin{aligned} & \mathbb{E} \left[ \sigma(X(T), m(T)) + \int_{t_0}^T g(t, X(t), m(t), \eta(t))dt \right] \\ &= \mu(T)\sigma(m(T)) + \int_{t_0}^T \mu(t)g(t, m(t), \eta(t))dt. \end{aligned} \quad (12)$$

Thus, the optimal feedback strategies do not depend on the concrete choice of the realization of the motion of the representative player.

## 4 Optimal control reformulation

The purpose of the section is to show that the solution of the mean field game is the solution of the following control problem:

$$\text{minimize } J(\phi(\cdot), m(\cdot), \mu(\cdot), \nu(\cdot)) \triangleq \mu_0\phi(t_0) - \mu(T)\sigma(m(T)) + \int_{t_0}^T \mu(t)g(t, m(t), \nu(t))dt \quad (13)$$

subject to  $(\phi(\cdot), m(\cdot), \mu(\cdot), \nu(\cdot))$  satisfying

$$\frac{d}{dt}m(t) = m(t)\mathcal{Q}(t, m(t), \nu(t)), \quad (14)$$

$$\frac{d}{dt}\mu(t) = \mu(t)\mathcal{Q}(t, m(t), \nu(t)), \quad (15)$$

$$\frac{d}{dt}\phi(t) = -H(t, m(t), \phi(t)), \quad (16)$$

$$m(t_0) = m_0, \mu(t_0) = \mu_0, \phi(T) = \sigma(m(T)). \quad (17)$$

Notice that the state vector for this problem is  $(\phi, m, \mu)$ , where  $\phi \in \mathbb{R}^d$ ,  $m, \mu \in \Sigma^d$ , when the control parameter is  $\nu = (\nu_1, \dots, \nu_d) \in (\mathcal{P}(U))^d$ . One may regard the vector  $\phi(t)$  as the upper bound of the rewards;  $m(t)$  stands for the distribution of all players;  $\mu(t)$  describes the evolution of fictitious players.

**Proposition 4.1.** *For every control process  $(\phi(\cdot), m(\cdot), \mu(\cdot), \nu(\cdot))$  satisfying (14), (15), (16), (17)*

$$J(\phi(\cdot), m(\cdot), \mu(\cdot), \nu(\cdot)) \geq 0.$$

*Proof.* First, let  $e^k$  stand for the  $k$ -th coordinate vector of  $\mathbb{R}^{d^*}$ . Notice that, for any  $\mu_0 = (\mu_{0,1}, \dots, \mu_{0,d}) \in \Sigma^d$ ,

$$\mu_0 = \sum_{k=1}^d \mu_{0,k} e^k.$$

Furthermore, let  $[t_0, T] \ni t \mapsto \mu^k(t) \in \Sigma^d$  be a solution of the initial value problem

$$\frac{d}{dt}\mu^k(t) = \mu^k(t)\mathcal{Q}(t, m(t), \nu(t)), \quad \mu^k(t_0) = e^k. \quad (18)$$

The trajectory  $\mu^k(\cdot)$  describes the probability distribution for the representative player who starts at time  $t_0$  from the state  $k$ . Notice that, if  $\mu(\cdot)$  satisfies (15), then

$$\mu(\cdot) = \sum_{k=1}^d \mu_{0,k} \mu^k(\cdot).$$

Put

$$\begin{aligned} J_k(\phi(\cdot), m(\cdot), \mu^k(\cdot), \nu(\cdot)) &\triangleq \phi_k(t_0) - \mu^k(T)\sigma(m(T)) - \int_{t_0}^T \mu^k(t)g(t, m(t), \nu(t))dt \\ &= e^k\phi(t_0) - \mu^k(T)\sigma(m(T)) - \int_{t_0}^T \mu^k(t)g(t, m(t), \nu(t))dt. \end{aligned} \quad (19)$$

Notice that the quantity

$$\mu^k(T)\sigma(m(T)) + \int_{t_0}^T \mu^k(t)g(t, m(t), \nu(t))dt$$



is the reward of the representative player who starts from the state  $k$  and uses the strategy  $\nu$  in the case when the distribution of all players is given by  $m(\cdot)$ . Since  $\phi_k(\cdot)$  is the equal to the maximal expected reward of the representative player, we deduce that

$$J_k(\phi(\cdot), m(\cdot), \mu^k(\cdot), \nu(\cdot)) \geq 0. \quad (20)$$

Furthermore, if  $\mu(\cdot)$  satisfies (15), then

$$J(\phi(\cdot), m(\cdot), \mu(\cdot), \nu(\cdot)) = \sum_{k=1}^d \mu_{0,k} J_k(\phi(\cdot), m(\cdot), \mu^k(\cdot), \nu(\cdot)). \quad (21)$$

This and (20) give the conclusion of the proposition.  $\square$

In the following theorem, we assume that the data of the mean field game  $m_0$  and  $\sigma$  are fixed.

**Theorem 4.2.** *For the function  $t \mapsto (\phi^*(t), m^*(t)) \in \mathbb{R}^d \times \Sigma^d$  and the control  $\nu^*(\cdot)$  the following statements are equivalent.*


- (i) *The pair  $(\phi^*(\cdot), m^*(\cdot))$  solves the mean field game with initial distribution  $m_0$ , whereas  $\nu^*(\cdot)$  is the corresponding equilibrium feedback strategy.*
- (ii) *There exists a flow of probabilities  $\mu^*(\cdot)$  and an initial distribution  $\mu_0^* \in \Sigma^d$  with nonzero coordinates such that  $(\phi^*(\cdot), m^*(\cdot), \mu^*(\cdot), \nu^*(\cdot))$  provides the solution of the optimal control problem (13)–(17).*
- (iii) *For every  $\mu^*(\cdot)$  satisfying (15), the control process  $(\phi^*(\cdot), m^*(\cdot), \mu^*(\cdot), \nu^*(\cdot))$  is the solution of the optimal control problem (13)–(17) with  $\mu_0 = \mu^*(t_0)$ .*
- (iv) *The triple  $(\phi^*(\cdot), m^*(\cdot), \nu^*(\cdot))$  is such that, for some  $\mu_0 \in \Sigma^d$  with nonzero coordinates and  $\mu^*(\cdot)$ , the process  $(\phi^*(\cdot), m^*(\cdot), \mu^*(\cdot), \nu^*(\cdot))$  satisfies (14)–(17) and the equality*

$$J(\phi^*(\cdot), m^*(\cdot), \mu^*(\cdot), \nu^*(\cdot)) = 0.$$

- (v) *For any  $\mu^*(\cdot)$  satisfying (15) and  $\mu_0 = \mu^*(t_0)$ , the control process  $(\phi^*(\cdot), m^*(\cdot), \mu^*(\cdot), \nu(\cdot))$  satisfies (14)–(17) and the following equality holds:*

$$J(\phi^*(\cdot), m^*(\cdot), \mu^*(\cdot), \nu^*(\cdot)) = 0.$$

*Proof.* The scheme of implications proving the desired equivalence can be illustrate by the following commutative diagram.



Obviously, we have that (iii) implies (ii). Analogously, (iv) follows from (v).

Now we prove implication (v) $\Rightarrow$ (iii). Let  $\mu^{*,k}$  satisfies (18) for  $m(\cdot) = m^*(\cdot)$  and  $\nu = \nu^*(\cdot)$ . Condition (v) implies that each value  $J_k(\phi^*(\cdot), m^*(\cdot), \mu^{*,k}(\cdot), \nu^*(\cdot)) = 0$ . Further, let  $\mu_0 = (\mu_{0,1}, \dots, \mu_{0,d})$  be an arbitrary element of  $\Sigma^d$ . The function  $\mu^*(\cdot)$  defined by

$$\mu^*(t) \triangleq \sum_{k=1}^d \mu_{0,k} \mu^{*,k}(t)$$

solves (15) with initial condition  $\mu^*(t_0) = \mu_0$ . Moreover, we have that

$$J(\phi^*(\cdot), m^*(\cdot), \mu^*(\cdot), \nu^*(\cdot)) = \sum_{k=1}^d \mu_{0,k} J_k(\phi^*(\cdot), m^*(\cdot), \mu^{*,k}(\cdot), \nu^*(\cdot)) = 0.$$

This and Proposition 4.1 yield that  $(\phi^*(\cdot), m^*(\cdot), \mu^*(\cdot), \nu^*(\cdot))$  provides the solution of the optimal control problem (13)–(17) for every  $\mu^*(\cdot)$ ,  $\mu_0$  satisfying ODE (15) and initial condition  $\mu^*(t_0) = \mu_0$ . Thus, (v) implies (iii) and, consequently, (ii).

By Proposition 3.3, the statement (i) is equivalent to the following one: equations (14), (16) hold true and  $J_k(\phi^*(\cdot), m^*(\cdot), \mu^{*,k}(\cdot), \nu^*(\cdot)) = 0$ . Indeed, let  $(\Omega, \mathcal{F}, \{\mathcal{F}_t\}_{t \in [t_0, T]}, P, X^k)$  be the motion of the representative player corresponding to the control  $\nu^*(\cdot)$ , flow of probabilities  $m^*(\cdot)$  and initial distribution  $e^k$ . Proposition 3.3 yields that

$$\phi_k(t_0) = \mathbb{E} \left[ \sigma(X^k(T), m(T)) + \int_{t_0}^T g(t, X^k(t), m(t), \nu(t)) dt \right],$$

where  $\mu_i^{*,k} = P(X^k(t) = i)$ . This and equality (12) imply that

$$\phi_k(t_0) = \mu^{*,k}(T) \phi(m^*(T)) + \int_{t_0}^T \mu^{*,k}(t) g(t, m^*(t), \nu^*(t)) dt.$$

Using definition of  $J_k$ , we deduce that optimality of  $\nu(\cdot)$  is equivalent to the equality  $J_k(\phi^*(\cdot), m^*(\cdot), \mu^{*,k}(\cdot), \nu^*(\cdot)) = 0$ .

Therefore, if (i) holds, then one can use (21) and deduce (v). This implies (iv).

Let us prove implication (iv) $\Rightarrow$ (i). Since

$$J(\phi^*(\cdot), m^*(\cdot), \mu^*(\cdot), \nu^*(\cdot)) = \sum_{k=1}^d \mu_{0,k} J_k(\phi^*(\cdot), m^*(\cdot), \mu^{*,k}(\cdot), \nu^*(\cdot))$$

and

$$J_k(\phi^*(\cdot), m^*(\cdot), \mu^{*,k}(\cdot), \nu^*(\cdot)) \geq 0, \quad k = 1, \dots, d,$$

we conclude that  $J_k(\phi^*(\cdot), m^*(\cdot), \mu^{*,k}(\cdot), \nu^*(\cdot)) = 0$ . This together with (14) and (16) is equivalent to (i).

To complete the proof, we prove the implication (ii) $\Rightarrow$ (iv). Assume that  $(\phi^*(\cdot), m^*(\cdot), \mu^*(\cdot), \nu^*(\cdot))$  provides the optimal solution of the control problem (13), (14), (16), (15). From Proposition 4.1, it follows that

$$J(\phi^*(\cdot), m^*(\cdot), \mu^*(\cdot), \nu^*(\cdot)) \geq 0.$$

Furthermore, recall (see Theorem 3.2) that there exists at least one solution of the finite state mean field game  $(\phi^{\natural}(\cdot), m^{\natural}(\cdot))$  for the initial distribution  $m_0$ . Let  $\nu'(\cdot)$  be the corresponding equilibrium strategy. Using the implication (i) $\Rightarrow$ (v) for the solution of the mean field game  $(\phi^{\natural}(\cdot), m^{\natural}(\cdot))$ , we deduce that the infimum of  $J(\phi'(\cdot), m'(\cdot), \mu'(\cdot), \nu'(\cdot))$  over the set of 4-tuples obeying (14)–(16) is not greater than 0. This leads the equality

$$J(\phi^*(\cdot), m^*(\cdot), \mu^*(\cdot), \nu^*(\cdot)) = 0.$$

This means that (ii) implies (iv). □

**Corollary 4.3.** *If  $m_0$  has nonzero-coordinates, then the following statements are equivalent.*

- *The pair  $(\phi^*(\cdot), m^*(\cdot))$  solves the mean field game with initial distribution  $m_0$ , whereas  $\nu^*(\cdot)$  is the corresponding equilibrium feedback strategy.*
- *The process  $(\phi^*(\cdot), m^*(\cdot), \nu^*(\cdot))$  is optimal in the problem*

$$\begin{aligned} \text{minimize } J'(\phi(\cdot), m(\cdot), \nu(\cdot)) &\triangleq m_0\phi(t_0) - m(T)\sigma(m(T)) \\ &\quad - \int_{t_0}^T m(t)g(t, m(t), \nu(t))dt \end{aligned}$$

*subject to (14), (16) and  $m(t_0) = m_0, \phi(T) = \sigma(m(T))$ .*

- *The triple  $(\phi^*(\cdot), m^*(\cdot), \nu^*(\cdot))$  satisfies (14), (16), boundary condition  $m^*(t_0) = m_0, \phi^*(T) = \sigma(m^*(T))$  and the equality  $J'(\phi^*(\cdot), m^*(\cdot), \nu^*(\cdot)) = 0$ .*

*Proof.* The desired equivalence follows from Theorem 4.2 and the fact that, if  $m(\cdot)$  and  $\mu(\cdot)$  satisfying (14) and (15) respectively are such that  $m(t_0) = \mu(t_0)$ , then  $m(\cdot) = \mu(\cdot)$ . □

## 5 Dependence of the solution of the mean field game on the initial distribution

Generally, the solution of the mean field game is nonunique [7]. Thus, it is reasonable to examine the multifunction assigning to an initial time and initial distribution of players a set of expected values of the representative player. In this section, we examine this multifunction using the viability and attainability theories.

**Definition 5.1.** We say that  $\mathcal{V} : [0, T] \times \Sigma^d \rightrightarrows \mathbb{R}^d$  is a value multifunction if, for any  $t_0 \in [0, T], m_0 \in \Sigma^d, \phi_0 \in \mathcal{V}(t_0, m_0)$ , there exists a solution of the mean field game  $(\phi(\cdot), m(\cdot))$  such that

$$\phi(t_0) = \phi_0, \quad m(t_0) = m_0.$$

We also consider the maximal value function that is multivalued mapping assigning to each pair  $(t_0, m(t_0))$  the set of all vectors  $\phi_0$  such that  $\phi_0 = \phi(t_0), m_0 = m(t_0)$  for some solution of the mean field game  $(\phi(\cdot), m(\cdot))$ . Denote the maximal value multifunction by  $\mathcal{W}$ .

First, let us present the sufficient condition for a given multifunction  $\mathcal{V} : [t_0, T] \times \Sigma^d \rightrightarrows \mathbb{R}^d$  to be a value multifunction. To this end, we use the viability property for the dynamics in the space  $\mathbb{R}^d \times \Sigma^d \times \Sigma^d \times \mathbb{R}$  given by equations (14)–(16) and

$$\frac{d}{dt}z(t) = -\mu(t)\mathcal{Q}(t, m(t), \nu(t))\phi(t) - \mu(t)H(t, m(t), \phi(t)) - \mu(t)g(t, m(t), \nu(t)). \quad (22)$$

**Definition 5.2.** We say that the set  $\mathcal{A} \subset [0, T] \times \mathbb{R}^d \times \Sigma^d \times \Sigma^d \times \mathbb{R}$  is viable w.r.t. to the dynamics (14)–(16), (22) if, for every  $s, r \in [0, T]$ ,  $s < r$ ,  $(s, \phi_*, m_*, \mu_*, z_*) \in \mathcal{A}$ , there exists a feedback relaxed control  $\nu(\cdot)$  and the 4-tuple  $(\phi(\cdot), m(\cdot), \mu(\cdot), z(\cdot))$  such that

- $\phi(\cdot), m(\cdot), \mu(\cdot), z(\cdot)$  and  $\nu(\cdot)$  satisfy (14)–(16), (22);
- $\phi(s) = \phi_*, m(s) = m_*, \mu(s) = \mu_*, z(s) = z_*$ ;
- $(r, \phi(r), m(r), \mu(r), z(r)) \in \mathcal{A}$ .

**Theorem 5.3.** Assume that, given a multifunction  $\mathcal{V} : [0, T] \times \Sigma^d \rightrightarrows \mathbb{R}^d$ , one can find a closed set  $\mathcal{A} \subset [0, T] \times \mathbb{R}^d \times \Sigma^d \times \Sigma^d \times \mathbb{R}$  such that

- $\mathcal{A}$  is viable w.r.t. (14)–(16), (22);
- the inclusion  $(T, \phi, m, \mu, z) \in \mathcal{A}$  implies that  $\phi = \sigma(m), z = 0$ ;
- if  $\phi_0 \in \mathcal{V}(t_0, m_0)$ , then  $(t_0, \phi_0, m_0, \mu_0, 0) \in \mathcal{A}$  for some  $\mu_0 \in \Sigma^d$  with nonzero coordinates.

Then,  $\mathcal{V}$  is a value multifunction.

The proof of the Theorem relies on the following lemma.

**Lemma 5.4.** If  $(\phi(\cdot), m(\cdot), \mu(\cdot), z(\cdot), \nu(\cdot))$  obeys (14)–(16), (22), then, for every  $s, r \in [0, T]$ ,  $s < r$ ,

$$z(r) - z(s) = \mu(s)\phi(s) - \mu(r)\phi(r) - \int_s^r \mu(t)g(t, m(t), \nu(t))dt. \quad (23)$$

*Proof.* In fact,

$$\mu(r)\phi(r) - \mu(s)\phi(s) = \int_s^r \frac{d}{dt} [\mu(t)\phi(t)] dt.$$

Since  $\mu(\cdot)$  and  $\phi(\cdot)$  satisfy equations (15) and (16) respectively, we obtain that

$$\frac{d}{dt} [\mu(t)\phi(t)] = \mu(t)\mathcal{Q}(t, m(t), \nu(t))\phi(t) + \mu(t)H(t, m(t), \phi(t)).$$

This implies representation (23). □

*Proof of Theorem 5.3.* Let  $(t_0, \phi_0, m_0, \mu_0)$  be such that

$$(t_0, \phi_0, m_0, \mu_0, 0) \in \mathcal{A},$$

and let  $\mu_0$  have nonzero coordinates. Using the standard viability technique (see [2, Theorem 3.3.4]), we deduce that there exists  $(\phi(\cdot), m(\cdot), \mu(\cdot), z(\cdot), \nu(\cdot))$  satisfying (14)–(16), (22) and boundary conditions

$$m(t_0) = m_0, \quad \phi(t_0) = \phi_0, \quad \mu(t_0) = \mu_0, \quad z(t_0) = 0,$$

$$\phi(T) = \sigma(m(T)), \quad z(T) = 0.$$

Using Lemma 5.4 and definition of the function  $J$  (see (13)), we deduce that

$$\begin{aligned} z(T) &= z(t_0) + \mu(t_0)\phi(t_0) - \mu(T)\phi(T) - \int_{t_0}^T \mu(t)g(t, m(t), \nu(t))dt \\ &= z(t_0) + J(m(\cdot), \phi(\cdot), \mu(\cdot), \nu(\cdot)). \end{aligned}$$

From the boundary conditions, it follows that

$$J(\phi(\cdot), m(\cdot), \mu(\cdot), \nu(\cdot)) = 0.$$

This means that, for  $(\phi(\cdot), m(\cdot), \nu(\cdot))$ , statement (iv) of Theorem 4.2 is in force. By Theorem 4.2  $(\phi(\cdot), m(\cdot))$  is a solution of the finite state mean field game. Thus,  $\mathcal{V}$  is a value multifunction.  $\square$

Now let us characterize the maximal value multifunction  $\mathcal{W}$  in the terms of the attainability domain. Denote

$$\hat{\nu} \triangleq (1/\sqrt{d}, \dots, 1/\sqrt{d}) = \frac{1}{\sqrt{d}}(1, \dots, 1) = \frac{1}{\sqrt{d}} \sum_{i=1}^d e^i.$$

**Theorem 5.5.** *Let  $\mathcal{A}_*$  be the set of 5-tuples  $(s, \phi(s), m(s), \mu(s), z(s))$  such that  $s \in [0, T]$ , when the 4-tuple  $(\phi(\cdot), m(\cdot), \mu(\cdot), z(\cdot))$  satisfies (14)–(16), (22) for some relaxed strategy  $\nu(\cdot)$  and the boundary condition*

$$m(T) \in \Sigma^d, \quad \phi(T) = \sigma(m(T)), \quad \mu(T) \in \Sigma^d, \quad z(T) = 0.$$

*Then, given  $t_0 \in [0, T]$ ,  $m_0 \in \Sigma^d$ ,*

$$\mathcal{W}(t_0, m_0) = \{\phi_0 : (t_0, \phi_0, m_0, \hat{\nu}, 0) \in \mathcal{A}_*\}.$$

*Remark 5.6.* In Theorem 5.5 one can replace  $\hat{\nu}$  with any row-vector with nonzero coordinates.

The proof of Theorem 5.5 uses the following auxiliary statement.

**Lemma 5.7.** *Assume that  $(\phi(\cdot), m(\cdot), \mu(\cdot), z(\cdot), \nu(\cdot))$  satisfies (14)–(16), (22). Then, the function  $t \mapsto z(t)$  is nonincreasing.*

*Proof.* Consider the Markov decision problem on  $[s, r]$  with the dynamics given by the Markov chain with the Kolmogorov matrix

$$\mathcal{Q}(t, m(t), \nu(t))$$

and the reward equal to

$$\mathbb{E} \left[ \phi(X(r), m(r)) + \int_s^r g(t, X(t), m(t), \nu(t)) dt \right]. \quad (24)$$

From dynamic programming principle it follows that the value of this problem at  $t \in [s, r]$  is equal to  $\phi(\cdot)$ .

Let  $(\Omega, \mathcal{F}, \{\mathcal{F}_t\}_{t \in [t_0, T]}, P, X)$  be a motion of the representative player corresponding to the feedback relaxed control  $\nu(\cdot)$ , the flow of probabilities  $m(\cdot)$  and the initial distribution at time  $s$  equal to  $\mu(s)$ . As above, we have that the probability  $P(X(t) = i) = \mu_i(t)$  and, thus, obeys (15). We have that

$$\begin{aligned} \mathbb{E} \left[ \phi(X(r), m(r)) + \int_s^r g(t, X(t), m(t), \nu(t)) dt \right] \\ = \mu(r)\phi(m(r)) + \int_s^r \mu(t)g(t, m(t), \nu(t)) dt. \end{aligned}$$

Since  $\phi(s)$  is the value function for the Markov decision problem with the payoff given by (24) and the dynamics given by the Markov chain  $\mathcal{Q}(t, m(t), \nu(t))$  on  $[s, r]$ ,

$$\mu(s)\phi(s) - \mu(r)\phi(r) - \int_s^r \mu(t)g(t, m(t), \nu(t)) dt \leq 0.$$

Combining this with (23), we obtain that  $z(r) \leq z(s)$  when  $r > s$ .  $\square$

*Proof of Theorem 5.5.* Notice that  $\mathcal{A}_*$  is viable with respect to equations (14)–(16), (22). Hence, by Theorem 5.3 the mapping  $(t_0, m_0) \rightarrow \{\phi_0 : (t_0, \phi_0, m_0, \vartheta, 0) \in \mathcal{A}_*\}$  is a value multifunctions. Hence,

$$\{\phi_0 : (t_0, \phi_0, m_0, \hat{\vartheta}, 0) \in \mathcal{A}_*\} \subset \mathcal{W}(t_0, m_0). \quad (25)$$

Let us prove the opposite inclusion.

Choose  $\phi_0 \in \mathcal{W}(t_0, m_0)$ . This means that there exists a solution of the mean field game  $(\phi(\cdot), m(\cdot))$  such that  $\phi(t_0) = \phi_0$ ,  $m(t_0) = m_0$ . Let  $\nu(\cdot)$  be a equilibrium relaxed feedback strategy corresponding to this solution and let  $\mu(\cdot)$  solve (15) with the initial condition  $\mu(t_0) = \hat{\vartheta}$ . Further, put

$$z(s) \triangleq \mu(s)\phi(s) - \mu(T)\sigma(m(T)) + \int_s^T \mu(t)g(t, m(t), \nu(t)) dt.$$

Notice, that  $z(\cdot)$  satisfies (22) and  $z(t_0) = J(m(\cdot), \phi(\cdot), \mu(\cdot), \nu(\cdot))$ . By Theorem 4.2, we conclude that  $z(t_0) = 0$ . Additionally,  $z(T) = 0$ . Using Lemma 5.7, we obtain that  $z(s) = 0$  for every  $s \in [t_0, T]$ . This implies that  $(t_0, m_0, \phi_0, \hat{\vartheta}, 0) \in \mathcal{A}_*$ . Therefore,

$$\mathcal{W}(t_0, m_0) \subset \{\phi_0 : (t_0, \phi_0, m_0, \hat{\vartheta}, 0) \in \mathcal{A}_*\}.$$

This together with (25) proves the Theorem.  $\square$

Let us complete the section by the fact that any solution of the master equation is a graph of a value multifunction.

Master equation (6) relies on the measurable feedback strategies. However, we primary consider relaxed strategies. Thus, we relax the master equation and arrive at the following:

$$\frac{\partial}{\partial s}\Phi(s, \mu) + H(t, \mu, \Phi(s, \mu)) \in -\text{co } \mathcal{O}(s, \mu, \Phi(s, \mu)) \cdot \frac{\partial \Phi}{\partial \mu}(s, \mu). \quad (26)$$

Here,

$$\text{co } \mathcal{O}(s, m, \phi) = \left\{ m\mathcal{Q}(t, m, \nu) : \right. \\ \left. \mathcal{Q}(t, m, \nu)\phi + g(t, m(t), \nu) = H(t, m, \phi), \quad \nu \in (\mathcal{P}(U))^d \right\}.$$

**Proposition 5.8.** *Let  $\Phi$  be a smooth solution of the master equation in the multivalued form (26). Then, the multifunction*

$$\mathcal{V}(t_0, m_0) \triangleq \{\Phi(t_0, m_0)\}$$

*is a value multifunction.*

*Proof.* Choose  $t_0 \in [0, T]$ ,  $m_0 \in \Sigma^d$ . We shall prove that there exists a solution of the mean field game  $(\phi(\cdot), m(\cdot))$  such that  $\phi(t_0) = \Phi(t_0, m_0)$ ,  $m(t_0) = m_0$ .

Let  $\mathcal{O}^\sharp(s, \mu)$  be the set of row-vectors  $\xi \in \text{co } \mathcal{O}(s, \mu, \Phi(s, \mu))$  such that

$$-\xi \frac{\partial \Phi(s, \mu)}{\partial \mu} = \frac{\partial}{\partial s}\Phi(s, \mu) + H(t, \mu, \Phi(s, \mu)). \quad (27)$$

Notice that  $\mathcal{O}^\sharp(s, \mu)$  is a nonempty convex compact subset of  $\mathbb{R}^{d*}$ . Moreover, the multifunction  $(s, \mu) \mapsto \mathcal{O}^\sharp(s, \mu)$  is upper semicontinuous.

Further, let  $m(\cdot)$  solve the differential inclusion

$$\frac{d}{dt}m(t) \in \mathcal{O}^\sharp(t, m(t)), \quad m(t_0) = m_0. \quad (28)$$

Consider the multivalued mapping

$$t \mapsto \left\{ \nu \in (\mathcal{P}(U))^d : \frac{d}{dt}m(t) = m(t)\mathcal{Q}(t, m(t), \nu), \right. \\ \left. \mathcal{Q}(t, m(t), \nu)\Phi(t, m(t)) + g(t, m(t), \nu) = H(t, m(t), \Phi(t, m(t))) \right\}.$$

By [1, §18.17, Filippov's Implicit Function Theorem], this mapping admits a measurable selector. Denote it by  $\hat{\nu}(\cdot)$ . Thus,

$$\frac{d}{dt}m(t) = m(t)\mathcal{Q}(t, m(t), \hat{\nu}(t)), \quad m(t_0) = m_0. \quad (29)$$

$$\mathcal{Q}(t, m(t), \hat{\nu}(t))\Phi(t, m(t)) + g(t, m(t), \hat{\nu}(t)) = H(t, m(t), \Phi(t, m(t))) \quad (30)$$

Furthermore, (27) and (28) yield the equality

$$-\left(\frac{d}{dt}m(t)\right)\frac{\partial\Phi(s,\mu)}{\partial\mu}\Big|_{s=t,\mu=m(t)}=\frac{\partial}{\partial s}\Phi(s,\mu)\Big|_{s=t,\mu=m(t)}+H(t,m,\Phi(s,m)).\quad (31)$$

Now, set  $\phi(t)\triangleq\Phi(t,m(t))$ . Due to (31), we have that it satisfies the Bellman equation

$$\frac{d}{dt}\phi(t)=-H(t,m(t),\phi(t))\quad (32)$$

and the boundary conditions

$$\phi(T)=\Phi(T,\sigma(m(T)))=\sigma(m(T)),\quad \phi(t_0)=\Phi(t_0,m(t_0)).\quad (33)$$

The fact that  $\hat{\nu}(\cdot)$  is an optimal control for the representative player directly follows from (30). Combining this with (29), (32) and (33), we conclude that the pair  $(\phi(\cdot),m(\cdot))$  is a solution of the mean field game such that  $\phi(t_0)=\Phi(t_0,m(t_0))$ ,  $m(t_0)=m_0$ . Since we choose  $(t_0,m_0)$  arbitrarily, the multifunction  $\mathcal{V}$  is the value multifunction.  $\square$

**Acknowledgment.** The article was prepared in the framework of a research grant funded by the Ministry of Science and Higher Education of the Russian Federation (grant ID: 075-15-2020-928).

## References

- [1] C. D. Aliprantis and K. C. Border. *Infinite Dimensional Analysis: A Hitchhiker's Guide*. Springer, Berlin, Heidelberg, 2006.
- [2] J.-P. Aubin. *Viability theory*. Birkhäuser, Boston, 2009.
- [3] R. Basna, A. Hilbert, and V. N. Kolokoltsov. An approximate nash equilibrium for pure jump markov games of mean-field-type on continuous state space. *Stochastics*, 89(6-7), 2016.
- [4] E. Bayraktar, A. Cecchin, A. Cohen, and F. Delarue. Finite state mean field games with Wright-Fisher common noise. Preprint at ArXiv:1912.06701, 2019.
- [5] E. Bayraktar, A. Cecchin, A. Cohen, and F. Delarue. Finite state mean field games with Wright-Fisher common noise as limits of  $N$ -player weighted games. Preprint at ArXiv:2012.04845, 2020.
- [6] E. Bayraktar and A. Cohen. Analysis of a finite state many player game using its master equation. *SIAM J. Control. Optim.*, 56(5):3538–3568, 2018.
- [7] E. Bayraktar and X. Zhang. On non-uniqueness in mean field games. *Proc. Amer. Math. Soc.*, 148:4091–4106, 2020.



- [8] C. Belak, D. Hoffmann, and F. T. Seifried. Continuous-time mean field games with finite state space and common noise. *SSRN Electronic Journal*, 2019.
- [9] A. Cecchin and M. Fischer. Probabilistic approach to finite state mean field games. *Appl. Math. Opt.*, 81(2):253–300, 2020.
- [10] A. Cecchin and G. Pelino. Convergence, fluctuations and large deviations for finite state mean field games via the master equation. *Stochastic Process. Appl.*, 129:4510–4555, 2019.
- [11] D. A. Gomes, J. Mohr, and R. R. Souza. Continuous time finite state mean field games. *Appl. Math. Opt.*, 68:99–143, 2013.
- [12] X. Guo and O. Hernández-Lerma. *Continuous-Time Markov Decision Processes*. Springer, New York, 2009.
- [13] M. Huang, P. E. Caines, and R. P. Malhamé. Large-population cost-coupled LQG problems with nonuniform agents: individual-mass behavior and decentralized Nash equilibria. *IEEE Trans. Automat. Control*, 52:1560–1571, 2007.
- [14] M. Huang, R. P. Malhamé, and P. E. Caines. Large population stochastic dynamic games: closed-loop McKean-Vlasov systems and the Nash certainty equivalence principle. *Commun. Inf. Syst.*, 6:221–251, 2006.
- [15] S. Katsikas and V. N. Kolokoltsov. Evolutionary, mean-field and pressure-resistance game modelling of networks security. *J. Dyn. Games*, 6(4):315–335, 2019.
- [16] V. Kolokoltsov and A. Bensoussan. Mean-field-game model for botnet defense in cyber-security. *Appl. Math. Opt.*, 74(3):669–692, 2016.
- [17] V. Kolokoltsov and O. Malafeyev. *Many agent games in socio-economic systems: corruption, inspection, coalition building, network growth, security*. Springer Nature, New York, 2019.
- [18] V. Kolokoltsov and W. Yang. Inspection games in a mean field setting. Preprint at ArXiv:1507.08339, 2015.
- [19] V. N. Kolokoltsov. *Nonlinear Markov process and kinetic equations*. Cambridge University Press, Cambridge, 2010.
- [20] V. N. Kolokoltsov and O. A. Malafeyev. Corruption and botnet defense: a mean field game approach. *Int. J. Game Theory*, 47:977–999, 2018.
- [21] J.-M. Lasry and P.-L. Lions. Jeux à champ moyen. I. Le cas stationnaire. *C. R. Math. Acad. Sci. Paris*, 343:619–625, 2006.
- [22] J.-M. Lasry and P.-L. Lions. Jeux à champ moyen. II. Horizon fini et contrôle optimal. *C. R. Math. Acad. Sci. Paris*, 343:679–684, 2006.
- [23] P.-L. Lions. College de France course on mean-field games. College de France, 2007-2011.