

# DISCRIMINANT OF TAUTOLOGICAL BUNDLES ON SYMMETRIC PRODUCTS OF CURVES

ANDREAS KRUG

ABSTRACT. We compute a formula for the discriminant of tautological bundles on symmetric powers of a complex smooth projective curve. It follows that the Bogomolov inequality does not give a new restriction to stability of these tautological bundles. It only rules out tautological bundles which are already known to have the structure sheaf as a destabilising subbundle.

## 1. INTRODUCTION

There is a natural way to produce a vector bundle on the symmetric product of a curve out of a vector bundle on the curve – use the Fourier–Mukai transform along the universal effective divisor.

More precisely, let  $C$  be a smooth curve over an algebraically closed field of characteristic zero, and let  $n \geq 2$ . Then the symmetric product  $C^{(n)} = C^n / \mathfrak{S}_n$  can be identified with the moduli space of effective degree  $n$  divisors on  $C$ , with the universal family  $\Xi \subset C \times C^{(n)}$  given by the image of the embedding

$$C \times C^{(n-1)} \hookrightarrow C \times C^{(n)} \quad , \quad (x, x_1 + \cdots + x_{n-1}) \mapsto (x, x_1 + \cdots + x_{n-1} + x).$$

Let  $p: \Xi \rightarrow C$  and  $q: \Xi \rightarrow C^{(n)}$  be the projections. For a vector bundle  $E$  on  $C$ , the associated *tautological bundle* on  $C^{(n)}$  is  $E^{[n]} := q_* p^* E$ . Since  $q$  is flat and finite of degree  $n$ , this is indeed a vector bundle with  $\text{rank}(E^{[n]}) = n \text{rank}(E)$  and fibres  $E^{[n]}(x_1 + \cdots + x_n) = H^0(E|_{x_1 + \cdots + x_n})$ .

One much-studied question regarding tautological bundles is whether they are stable. Here, by *stability* we mean slope stability with respect to the polarisation  $H = x + C^{(n-1)} \subset C^{(n)}$ . Strengthening earlier results of [AO94, BS92, Mis19, EMLN11, BN13, DP16, BD18], it was proven in [Kru18] that if  $E$  is (semi-)stable with slope  $\mu(E) \notin [-1, n-1]$  (or, for semi-stability,  $\mu(E) \notin (-1, n-1)$ ), then also  $E^{[n]}$  is (semi-)stable.

For  $\mu(E) \in [-1, n-1]$ , the situation seems more complicated. It was observed in [Kru18, Sect. 3] that the above condition for stability is already numerically optimal: For every curve  $C$  and every  $d = 0, \dots, n-2$  there is a line bundle of degree  $d$  on  $C$  such that  $L^{[n]}$  is unstable. There are also line bundles of degree  $-1$  and  $n-1$  such that  $L^{[n]}$  is strictly semi-stable. On the other hand, there are also examples of line bundles with  $d \in (-1, n-1)$  such that  $L^{[n]}$  is stable. Namely, by [BN13],  $L^{[2]}$  is stable for every non-trivial line bundle  $L$  of degree zero. We would like to find a complete answer to the following question:

*Let  $E$  be a (semi-)stable bundle of slope  $\mu(E) \in [-1, n-1]$ . Under which circumstances is  $E^{[n]}$  a (semi-)stable bundle on  $C^{(n)}$ ?*

An obvious first approach to get a meaningful restriction on semi-stability of  $E^{[n]}$  is to use the Bogomolov inequality

$$\int_{C^{(n)}} \Delta(E^{[n]}) H^{n-2} \geq 0$$

where  $\Delta(E^{[n]}) = -2 \operatorname{rank}(E^{[n]}) \operatorname{ch}_2(E^{[n]}) + c_1(E^{[n]})^2$  is the discriminant; see [Bog78], [Miy87], [HL10, Thm. 7.3.1]. In this paper, we compute the relevant intersection number.

**Theorem 1.1.** *Let  $C$  be a smooth projective curve of genus  $g$ , and let  $E$  be a vector bundle on  $C$  of rank  $r$  and degree  $d$ . Then, for every  $n \geq 2$ ,*

$$\int_{C^{(n)}} \Delta(E^{[n]}) H^{n-2} = d^2 - (n-2)dr + (n-1)(g-1)r^2.$$

Dividing the equation of **Theorem 1.1** by  $r^2$ , we see that

$$\int \Delta(E^{[n]}) H^{n-2} \geq 0 \iff \mu^2 - (n-2)\mu + (n-1)(g-1) \geq 0.$$

Hence, the Bogomolov inequality gives

**Corollary 1.2.** *The tautological bundle  $E^{[n]}$  is unstable for every vector bundle  $E$  on  $C$  with*

$$\mu(E) \in \left( \frac{n-2}{2} - \frac{1}{2} \sqrt{(n-2)^2 - 4(n-1)(g-1)}, \frac{n-2}{2} + \frac{1}{2} \sqrt{(n-2)^2 - 4(n-1)(g-1)} \right)$$

For  $g \geq 1$ , the interval of **Corollary 1.2** lies inside the interval  $(g-1, n-g)$ , and it approaches for large  $n$  asymptotically the whole interval  $(g-1, n-g)$ .

However, the instability criterion of **Corollary 1.2** is weaker than one already given in [Kru18, Sect. 3]. There, it was observed that  $\mathcal{O}_{C^{(n)}}$  is a destabilising subbundle of  $E^{[n]}$  for every  $E$  with  $\mu(E) < n-1$  and  $H^0(E) \neq 0$ . By the Riemann-Roch theorem, this implies that  $E^{[n]}$  is always unstable when  $\mu(E) \in (g-1, n-1)$ .

Let now  $n = 2$ , such that  $C^{(2)}$  is a surface. Then for  $d \notin [-1, r]$ , the bundle  $E^{[2]}$  is stable by [Kru18], and we write  $\mathcal{M} = \mathcal{M}_{C^{(2)}}(\operatorname{ch}(E^{[2]}), \det(E^{[2]}))$  for the moduli space of stable sheaves on  $C^{(2)}$  with the same Chern character and determinant. **Theorem 1.1** tells us that  $\int \Delta(E^{[n]})$  grows quadratically with  $d$ . Hence, we can apply the results of [O'G96] to obtain

**Corollary 1.3.** *For  $|d| \gg 0$ , the moduli space  $\mathcal{M}$  is irreducible and generically reduced of the expected dimension.*

Since  $\chi(\mathcal{O}_{C^{(2)}}) = 1 - g + \binom{g}{2} = \frac{1}{2}(g^2 - 3g + 2)$ , the expected dimension of  $\mathcal{M}$  is

$$\Delta(E^{[2]}) - (\operatorname{rank}(E^{[2]})^2 - 1)\chi(\mathcal{O}_X) = d^2 + r^2(g-1) - \frac{1}{2}(4r^2 - 1)(g^2 - 3g + 2).$$

## 2. PROOF OF **THEOREM 1.1**

**2.1. General Conventions.** All our computations are carried out in the Chow rings modulo numerical equivalence. In particular, we will often omit the integral symbol  $\int$  when writing intersection numbers.

Throughout  $C$  will denote a smooth projective curve over a field of characteristic zero, and  $x$  will denote a point of the curve  $C$ . It will not matter which point, since we are doing our computations modulo numerical equivalence. In particular, the notion of slope stability with respect to our polarisation  $H_n = x + C^{(n-1)}$  of  $C^{(n)}$  does not depend on the choice of  $x \in C$ .

**2.2. Some intersection numbers on  $C^n$ .** Let  $\pi_n: C^n \rightarrow C^{(n)}$  be the  $\mathfrak{S}_n$ -quotient morphism. We have  $\tilde{H}_n := \pi_n^* H_n = \sum_{i=1}^n \text{pr}_i^*([x])$  where  $\text{pr}_i: C^n \rightarrow C$  is the projection to the  $i$ -th factor. For  $1 \leq i < j \leq n$ , we consider the *pairwise diagonal*

$$\Delta_{ij} = \{(x_1, \dots, x_n) \in C^n \mid x_i = x_j\}.$$

The *big diagonal* is the sum of all pairwise diagonals

$$\delta_n = \sum_{1 \leq i < j \leq n} \Delta_{ij}.$$

Another divisor on  $C^n$  that we will need is  $\delta'_n = \delta_n - \overline{\text{pr}}_1^* \delta_n = \sum_{j=2}^n \Delta_{1j}$  where

$$\overline{\text{pr}}_1: C^n \rightarrow C^{n-1} \quad , \quad (x_1, x_2, \dots, x_n) \mapsto (x_2, \dots, x_n).$$

We will often omit the index  $n$  in the notation and write  $\pi, H, \tilde{H}, \delta, \delta'$  instead of  $\pi_n, H_n, \tilde{H}_n, \delta_n, \delta'_n$ . For  $I \subset \{1, \dots, n\}$ , we define

$$\eta_I := \prod_{i \in I} \text{pr}_i^*([x]) = [\{(x_1, \dots, x_n) \mid x_i = x \ \forall i \in I\}].$$

Since  $\text{pr}_i^*([x])^2 = 0$  for every  $i = 1, \dots, n$ , we get

$$(1) \quad \tilde{H}^k = k! \sum_{|I|=k} \eta_I \quad \text{for all } k = 1, \dots, n.$$

**Lemma 2.1.** *We have the following intersection numbers on  $C^n$ :*

$$(2) \quad \tilde{H}^n = n! \quad , \quad \delta \tilde{H}^{n-1} = n!(n-1) \quad , \quad \delta^2 \tilde{H}^{n-2} = -n!(g-1) + n!(n-2)n,$$

$$(3) \quad \text{pr}_1^*([x]) \delta' \tilde{H}^{n-2} = (n-1)! \quad , \quad \delta'^2 \tilde{H}^{n-2} = -(n-1)!2(g-1) + (n-1)!(n-2)3.$$

*Proof.* The first equation is just a special case of (1). For the second equation, note that, for  $|I| = n-1$ , we have

$$\Delta_{ij} \eta_I = \begin{cases} 1 & \text{if } |I \cap \{i, j\}| = 1, \\ 0 & \text{if } |I \cap \{i, j\}| = 2. \end{cases}$$

Hence, by (1), we get  $\Delta_{ij} \tilde{H}^{n-1} = (n-1)!2$ . Since there are  $\binom{n}{2}$  pairwise diagonals, we get  $\delta \tilde{H}^{n-1} = \binom{n}{2} (n-1)!2 = n!(n-1)$ .

We have  $\text{pr}_1^*([x]) \Delta_{1j} = \eta_{\{1, j\}}$ . Hence, the only summand of  $\tilde{H}^{n-2}$  which has a non-zero intersection pairing with  $\text{pr}_1^*([x]) \Delta_{1j}$  is  $\eta_{\{2, \dots, n\} \setminus \{j\}}$ . Hence  $\text{pr}_1^*([x]) \Delta_{1j} \tilde{H}^{n-2} = (n-2)!$ , again by (1). The formula  $\text{pr}_1^*([x]) \delta' \tilde{H}^{n-2} = (n-2)!(n-1) = (n-1)!$  follows as  $\delta'$  consists of  $(n-1)$  pairwise diagonals. For the computation of  $\delta'^2 \tilde{H}^{n-2}$ , we first note that

$$(4) \quad \delta'^2 = \sum_{j=2}^n \Delta_{1j}^2 + 2 \sum_{2 \leq i < j \leq n} \Delta_{1i} \Delta_{1j}.$$

The self-intersection of a pairwise diagonal is  $\Delta_{1j}^2 = -2(g-1)\eta_{\{1, j\}}$ . Hence,

$$(5) \quad \Delta_{1j}^2 \tilde{H}^{n-2} = -(n-2)!2(g-1).$$

Furthermore,  $\Delta_{1i} \Delta_{1j} = \Delta_{1ij} = \{(x_1, \dots, x_n) \mid x_1 = x_i = x_j\}$ . For  $|I| = n-2$ , we have

$$\Delta_{1ij} \eta_I = \begin{cases} 1 & \text{if } |I \cap \{1, i, j\}| = 1, \\ 0 & \text{if } |I \cap \{1, i, j\}| \geq 2. \end{cases}$$

Since there are three  $I \subset \{1, \dots, n\}$  with  $|I| = n - 2$  and  $|I \cap \{1, i, j\}| = 1$ , equation (1) yields

$$(6) \quad \Delta_{1ij} \tilde{H}^{n-2} = (n-2)!3.$$

We obtain the formula for  $\delta^2 \tilde{H}^{n-2}$  by combining (4), (5), and (6).

The computation of  $\delta^2 \tilde{H}^{n-2}$  is very similar using that

$$\delta^2 = \sum_{1 \leq i < j \leq n} \Delta_{ij}^2 + 6 \sum_{1 \leq i < j < k \leq n} \Delta_{ijk} + \sum_{\{i,j\} \cap \{k,\ell\} = \emptyset} \Delta_{ij} \Delta_{k,\ell}$$

instead of (4). The only additional ingredient is that, for  $\{i, j\} \cap \{k, \ell\} = \emptyset$  and  $|I| = n - 2$ , we have

$$\Delta_{ij} \Delta_{k\ell} \eta_I = \begin{cases} 1 & \text{if } |I \cap \{i, j\}| = 1 = |I \cap \{k, \ell\}|, \\ 0 & \text{else.} \end{cases} \quad \square$$

**Lemma 2.2.** *Let  $\alpha \in \text{CH}^2(C^{n-2})$ . Then*

$$\int_{C^n} \tilde{H}_n^{n-2} \overline{\text{pr}}_1^* \alpha = (n-2) \int_{C^{n-1}} \tilde{H}_{n-1}^{n-3} \alpha.$$

*Proof.* By projection formula,

$$\int_{C^n} \tilde{H}_n^{n-2} \overline{\text{pr}}_1^* \alpha = \int_{C^{n-1}} \overline{\text{pr}}_{1*} (\tilde{H}_n^{n-2} \overline{\text{pr}}_1^* \alpha) = \int_{C^{n-1}} (\overline{\text{pr}}_{1*} \tilde{H}_n^{n-2}) \alpha.$$

We have  $\overline{\text{pr}}_{1*} \tilde{H}_n^{n-2} = (n-2) \tilde{H}_{n-1}^{n-3}$  as follows from (1) together with

$$\overline{\text{pr}}_{1*} \eta_I = \begin{cases} \eta_{I \setminus \{1\}} & \text{if } 1 \in I, \\ 0 & \text{if } 1 \notin I. \end{cases} \quad \square$$

**2.3. Computation of the discriminant by induction.** The Chern characters of tautological bundles are computed in [Mat65, Sect. 3] in the case that  $E = L$  is a line bundle. So one way to proceed would be to generalise the formula [Mat65, Prop. 6] to higher rank bundles, and then use it to compute  $\Delta(E^{[n]})H^{n-2}$ . Instead, we chose a more direct route to the computation of  $\Delta(E^{[n]})H^{n-2}$ , using the short exact sequence

$$(7) \quad 0 \rightarrow \text{pr}_1^* E(-\delta'_n) \rightarrow \pi_n^* E^{[n]} \rightarrow \overline{\text{pr}}_1^* \pi_{n-1}^* E^{[n-1]} \rightarrow 0.$$

of [Kru18, Prop. 1.4] to proceed by induction on  $n$ .

We consider a vector bundle  $E$  of rank  $r$  and degree  $d$  on the curve  $C$  of genus  $g$ .

**Lemma 2.3.** *For all  $n \geq 2$ , we have*

$$c_1(E^{[n]})^2 H^{n-2} = d^2 - 2dr(n-1) - r^2(g-1) + r^2n(n-2).$$

*Proof.* Using the short exact sequence (7) one deduces by induction the formula

$$c_1(\pi^* E^{[n]}) = d\tilde{H} - r\delta;$$

see [Kru18, Sect. 1.7]. Hence,  $c_1(\pi^* E^{[n]})^2 = d^2 \tilde{H}^2 - 2dr\delta\tilde{H} + r^2\delta^2$ . By (2), we get

$$c_1(\pi^* E^{[n]})^2 \tilde{H}^{n-2} = n!(d^2 - 2dr(n-1) - r^2(g-1) + r^2n(n-2)).$$

Since  $\pi: C^n \rightarrow C^{(n)}$  is of degree  $n!$ , we have  $c_1(E^{[n]})^2 H^{n-2} = \frac{c_1(\pi^* E^{[n]})^2 \tilde{H}^{n-2}}{n!}$ . □

**Lemma 2.4.** *For every  $n \geq 2$ , we have*

$$\mathrm{ch}_2(E^{[n]})H^{n-2} = -\frac{1}{2}(d + (g-1)r - (n-2)r).$$

*Proof.* We have  $\mathrm{ch}(\mathrm{pr}_1^* E) = r + d \mathrm{pr}_1^*([x])$  and  $\mathrm{ch}(\mathcal{O}(-\delta')) = 1 - \delta' + \frac{1}{2}\delta'^2 \pm \dots$ . Hence,  $\mathrm{ch}_2(\mathrm{pr}_1^* E(-\delta')) = -d \mathrm{pr}_1^*([x])\delta' + \frac{1}{2}r\delta'^2$  which by (3) gives

$$(8) \quad \mathrm{ch}_2(\mathrm{pr}_1^* E(-\delta'))\tilde{H}^{n-2} = -(n-1)!d - (n-1)!(g-1)r + (n-1)!(n-2)\frac{3}{2}r$$

We now can prove the assertion, or rather the equivalent formula

$$(9) \quad \mathrm{ch}_2(\pi_n^* E^{[n]})\tilde{H}^{n-2} = -\frac{1}{2}n!(d + (g-1)r - (n-2)r),$$

by induction using the short exact sequence (7). For  $n = 2$ , the sequence gives  $\mathrm{ch}_2(\pi_2^* E^{[2]}) = \mathrm{ch}_2(\mathrm{pr}_1^* E(-\delta'))$ , and we check that the right-hand sides of (8) and (9) are equal for  $n = 2$ . For  $n \geq 3$ , the sequence (7) together with Lemma 2.2 give

$$\begin{aligned} \mathrm{ch}_2(\pi_n^* E^{[n]})\tilde{H}_n^{n-2} &= \mathrm{ch}_2(\mathrm{pr}_1^* E(-\delta'))\tilde{H}_n^{n-2} + \overline{\mathrm{pr}}_1^* \mathrm{ch}_2(\pi_{n-1}^* E^{[n-1]})\tilde{H}_n^{n-2} \\ &= \mathrm{ch}_2(\mathrm{pr}_1^* E(-\delta'))\tilde{H}_n^{n-2} + (n-2) \mathrm{ch}_2(\pi_{n-1}^* E^{[n-1]})\tilde{H}_{n-1}^{n-3}. \end{aligned}$$

Now, the induction step is done by plugging (8) and the  $n-1$  case of (9) into the right-hand side.  $\square$

Recall that the discriminant is defined as

$$\Delta(E^{[n]}) = -2 \mathrm{rank}(E^{[n]}) \mathrm{ch}_2(E^{[n]}) + c_1(E^{[n]})^2 = -2nr \mathrm{ch}_2(E^{[n]}) + c_1(E^{[n]})^2.$$

Hence Lemma 2.3 and Lemma 2.4 together compute  $\Delta(E^{[n]})H^{n-2}$  and prove Theorem 1.1.

#### REFERENCES

- [AO94] Vincenzo Ancona and Giorgio Ottaviani. Stability of special instanton bundles on  $\mathbf{P}^{2n+1}$ . *Trans. Amer. Math. Soc.*, 341(2):677–693, 1994.
- [BD18] Suratno Basu and Krishanu Dan. Stability of secant bundles on the second symmetric power of curves. *Arch. Math. (Basel)*, 110(3):245–249, 2018.
- [BN13] Indranil Biswas and D. S. Nagaraj. Stability of secant bundles on second symmetric power of a curve. In *Commutative algebra and algebraic geometry (CAAG-2010)*, volume 17 of *Ramanujan Math. Soc. Lect. Notes Ser.*, pages 13–18. Ramanujan Math. Soc., Mysore, 2013.
- [Bog78] Fedor A. Bogomolov. Holomorphic tensors and vector bundles on projective manifolds. *Izv. Akad. Nauk SSSR Ser. Mat.*, 42(6):1227–1287, 1439, 1978.
- [BS92] Guntram Bohnhorst and Heinz Spindler. The stability of certain vector bundles on  $\mathbf{P}^n$ . In *Complex algebraic varieties (Bayreuth, 1990)*, volume 1507 of *Lecture Notes in Math.*, pages 39–50. Springer, Berlin, 1992.
- [DP16] Krishanu Dan and Sarbeswar Pal. Semistability of certain bundles on second symmetric power of a curve. *J. Geom. Phys.*, 103:37–42, 2016.
- [EMLN11] A. El Mazouni, F. Laytimi, and D. S. Nagaraj. Secant bundles on second symmetric power of a curve. *J. Ramanujan Math. Soc.*, 26(2):181–194, 2011.
- [HL10] Daniel Huybrechts and Manfred Lehn. *The geometry of moduli spaces of sheaves*. Cambridge Mathematical Library. Cambridge University Press, Cambridge, second edition, 2010.
- [Kru18] Andreas Krug. Stability of tautological bundles on symmetric products of curves. *arXiv:1809.06450*, 2018.
- [Mat65] Arthur Mattuck. Secant bundles on symmetric products. *Amer. J. Math.*, 87:779–797, 1965.
- [Mis19] Ernesto C. Mistretta. On stability of tautological bundles and their total transforms. *Milan J. Math.*, 87(2):273–282, 2019.

- [Miy87] Yoichi Miyaoka. The Chern classes and Kodaira dimension of a minimal variety. In *Algebraic geometry, Sendai, 1985*, volume 10 of *Adv. Stud. Pure Math.*, pages 449–476. North-Holland, Amsterdam, 1987.
- [O’G96] Kieran G. O’Grady. Moduli of vector bundles on projective surfaces: some basic results. *Invent. Math.*, 123(1):141–207, 1996.