

THE ISOMORPHISM PROBLEM FOR SCHUBERT VARIETIES

EDWARD RICHMOND AND WILLIAM SLOFSTRA

ABSTRACT. Schubert varieties in the full flag variety of Kac-Moody type are indexed by elements of the corresponding Weyl group. We give a practical criterion for when two such Schubert varieties (from potentially different flag varieties) are isomorphic, in terms of the Cartan matrix and reduced words for the indexing Weyl group elements. As a corollary, we show that two such Schubert varieties are isomorphic if and only if there is an isomorphism between their integral cohomology rings that preserves the Schubert basis.

1. INTRODUCTION

Kac-Moody flag varieties are central objects of study in geometry, topology, and representation theory. In particular, the finite-type Kac-Moody flag varieties are the usual generalized flag varieties associated to semisimple Lie groups. While generalized flag varieties are finite-dimensional, Kac-Moody flag varieties of non-finite type are infinite-dimensional. However, all Kac-Moody flag varieties can be realized as ind-varieties stratified by finite-dimensional Schubert varieties [Kum87]. Schubert varieties are themselves important examples of algebraic varieties, and their singularities are closely connected with the representation theory of the corresponding Kac-Moody groups and algebras. As a result, their geometry has been closely studied (see, for instance, the surveys [BL00, AB16]).

We are interested in the isomorphism problem for Schubert varieties: when are two Schubert varieties isomorphic as algebraic varieties? This natural geometric question was first raised for Schubert varieties by Develin, Martin, and Reiner [DMR07]. They show that two partition varieties (a subclass of type A Schubert varieties introduced by Ding [Din97]) are isomorphic if and only if their integral cohomology rings are isomorphic. Aside from this, and the case of toric Schubert varieties which we cover below, the question does not seem to have been pursued further, even in type A or other finite types. In this paper, we give a complete solution to the isomorphism problem for Schubert varieties in full flag varieties of any Kac-Moody type over \mathbb{C} .

To describe this solution, it is helpful to recall some basic facts about Schubert varieties. We follow the conventions from [Kum02]. Recall that the starting data for constructing a Kac-Moody Lie algebra (and subsequently its Kac-Moody group, full flag variety, and Schubert varieties) is a **(generalized) Cartan matrix**, which is an integer matrix $A := [A_{st}]_{(s,t) \in S^2}$ indexed by some finite set S , such that for any $s, t \in S$,

- (1) $A_{st} = 2$ if $s = t$,
- (2) $A_{st} \leq 0$ if $s \neq t$, and
- (3) $A_{st} = 0$ if and only if $A_{ts} = 0$.

Let $\mathcal{G} := \mathcal{G}(A)$ and $\mathcal{B} := \mathcal{B}(A)$ denote the Kac-Moody group and Borel subgroup of a Cartan matrix A . The full flag variety corresponding to A is the quotient $\mathcal{X} = \mathcal{X}(A) := \mathcal{G}/\mathcal{B}$. The Weyl group $W := W(A)$ of A is the crystallographic Coxeter group generated by S , and

satisfying relations $(st)^{m_{st}} = e$, where $m_{ss} = 1$, and

$$m_{st} = \begin{cases} 2 & \text{if } A_{st}A_{ts} = 0 \\ 3 & \text{if } A_{st}A_{ts} = 1 \\ 4 & \text{if } A_{st}A_{ts} = 2 \\ 6 & \text{if } A_{st}A_{ts} = 3 \\ \infty & \text{if } A_{st}A_{ts} \geq 4 \end{cases}$$

for all $s, t \in S$. The pair (W, S) forms a Coxeter system, and the elements of S are referred to as the **simple reflections** (or **simple transpositions**) in W . For every $w \in W$, the **Schubert variety** $X(w, A)$ is defined to be the closure of $\mathcal{B}w\mathcal{B}/\mathcal{B}$ in \mathcal{X} . It is well known that $X(w, A)$ is an irreducible finite dimensional complex variety of dimension $\ell(w)$, where $\ell : W \rightarrow \mathbb{Z}_{\geq 0}$ is the length function of the Coxeter system (W, S) . A product $w = s_1 \cdots s_k$ of simple reflections s_1, \dots, s_k in W is a **reduced word** if $k = \ell(w)$. Every element of W can be written as a reduced word.

If $w = s$ is a simple reflection, then $X(w, A) \cong \mathbb{P}^1$, and hence all one-dimensional Schubert varieties are isomorphic, independent of A . The case of two-dimensional Schubert varieties is more interesting:

Example 1.1. *Suppose $X(w, A)$ is a two-dimensional Schubert variety, so $w = st \in S$ for some $s \neq t$. Then $X(w, A)$ is a Zariski-locally-trivial \mathbb{P}^1 -bundle over \mathbb{P}^1 . The cohomology ring $H^*(X(w, A); \mathbb{Z})$ is the free \mathbb{Z} -module generated by the Schubert classes ξ_u for $u \in \{e, s, t, w\}$, where ξ_u has degree $2\ell(u)$ and e is the identity. The ring structure is determined by the relations $\xi_s^2 = 0$, $\xi_s \cdot \xi_t = \xi_w$, and $\xi_t^2 = -A_{st}\xi_w$. From this, it follows that $X(w, A)$ is the Hirzebruch surface Σ_n , where $n = -A_{st}$.*

It is well-known that $\Sigma_n \cong \Sigma_m$ if and only if $m = n$. Hence $X(w, A) \cong X(w', A')$ with $w' = s't'$ if and only if $A_{st} = A'_{s't'}$. In particular, the isomorphism type of $X(w, A)$ depends only on the value of A_{st} , not on A_{ts} .

Let \leq denote the Bruhat order for the Coxeter system (W, S) , and define the **support** of an element $w \in W$ to be the set

$$S(w) := \{s \in S : s \leq w\}.$$

A simple reflection $s \in S$ belongs to $S(w)$ if and only if s appears in some (or equivalently, every) reduced word of w . Inspired by the example of Hirzebruch surfaces, we define:

Definition 1.2. *Let A and A' be Cartan matrices over S and S' respectively. Let $w \in W(A)$ and $w' \in W(A')$. We say the pair (w, A) and (w', A') are **Cartan equivalent** if there is a bijection $\sigma : S(w) \rightarrow S(w')$ such that the following are satisfied:*

- (a) *There are reduced words $w = s_1 \cdots s_k$ and $w' = t_1 \cdots t_k$ such that $\sigma(s_i) = t_i$ for all $1 \leq i \leq k$.*
- (b) *If $ss' \leq w$ for $s \neq s' \in S(w)$, then $A_{ss'} = A'_{\sigma(s)\sigma(s')}$.*

Although it's not immediately obvious from the definition, we show in the next section (see Corollary 2.3) that Cartan equivalence is an equivalence relation (and in particular is symmetric). We also refer to the bijection $\sigma : S(w) \rightarrow S(w')$ in Definition 1.2 as a Cartan equivalence.

Our main result is that, somewhat surprisingly, two Schubert varieties $X(w, A)$ and $X(w', A')$ are isomorphic if and only if (w, A) and (w', A') are Cartan equivalent. As

part of the proof, we give a cohomological characterization of isomorphism as well. Recall that Schubert varieties are stratified by their Schubert subvarieties, and this stratification implies that the Schubert classes form a basis for the integral cohomology ring $H^*(X(w, A)) := H^*(X(w, A); \mathbb{Z})$.

Theorem 1.3. *Let A and A' be Cartan matrices over S and S' respectively. Let $w \in W(A)$ and $w' \in W(A')$. Then the following are equivalent.*

- (1) *The pairs (w, A) and (w', A') are Cartan equivalent.*
- (2) *The Schubert varieties $X(w, A)$ and $X(w', A')$ are algebraically isomorphic.*
- (3) *There is a graded ring isomorphism $\phi : H^*(X(w, A)) \rightarrow H^*(X(w', A'))$ which sends the Schubert basis for $H^*(X(w, A))$ to the Schubert basis for $H^*(X(w', A'))$.*

Theorem 1.3 can be readily applied in many situations. To illustrate this, we give several examples and applications in Section 1.1. We remark that the theorem does not hold if we drop the requirement in part (3) that the isomorphism preserve Schubert bases. Indeed, returning to Example 1.1, two Hirzebruch surfaces Σ_n and Σ_m have isomorphic integral cohomology rings if and only if they are diffeomorphic, which happens if and only if $m = n \pmod{2}$.

Hirzebruch surfaces are also examples of toric varieties. In general, a Schubert variety $X(w, A)$ is a toric variety if and only if $w = s_1 \cdots s_k$ for distinct simple reflections $s_1, \dots, s_k \in S$ [Kar13]. It follows from this that toric Schubert varieties are toric manifolds (smooth compact toric varieties). It is well known that isomorphism classes of toric varieties are determined by the combinatorial data in their associated fans [Ful93]. In addition, a result of Masuda states that toric manifolds are isomorphic if and only if their equivariant cohomology rings are weakly isomorphic [Mas08]. Both of these criteria apply to toric Schubert varieties in particular. It is an open question as to whether toric manifolds are cohomologically rigid, in the sense that any two toric manifolds with isomorphic cohomology rings are diffeomorphic or homeomorphic. We can ask the same question for Schubert varieties:

Question 1.4. *Suppose $X(w, A)$ and $X(w', A')$ are both smooth, and $H^*(X(w, A))$ and $H^*(X(w', A'))$ are isomorphic as graded rings. Are $X(w, A)$ and $X(w', A')$ diffeomorphic or homeomorphic?*

If integral cohomology is replaced with rational cohomology $H^*(X(w, A); \mathbb{Q})$, then Question 1.4 has a negative answer, as all Hirzebruch surfaces have isomorphic cohomology rings over \mathbb{Q} . Another counterexample is provided by the flag varieties of finite types B_n and C_n , since the cohomology rings are isomorphic over \mathbb{Q} , but not over \mathbb{Z} [BS02, Bor53, EG95].

Finally, recall that if A is a Cartan matrix over S , then for any subset $J \subseteq S$ there is a partial flag variety $\mathcal{X}^J := \mathcal{G}(A)/\mathcal{P}(A)_J$, where $\mathcal{P}(A)_J$ is the parabolic subgroup generated by the elements of J . Partial flag varieties are also stratified by Schubert varieties $X^J(w, A)$, where $X^J(w, A)$ is defined as the closure of $\mathcal{B}w\mathcal{P}_J/\mathcal{P}_J$ in \mathcal{X}^J . Partial flag varieties include familiar examples such as the Grassmannians. Next we show the notion of Cartan equivalence is neither necessary nor sufficient for distinguishing Schubert varieties in partial flag varieties.

Example 1.5. *Consider the Cartan matrices of types A_3 and C_2 given by*

$$A_3 = \begin{pmatrix} 2 & -1 & 0 \\ -1 & 2 & -1 \\ 0 & -1 & 2 \end{pmatrix} \quad \text{and} \quad C_2 = \begin{pmatrix} 2 & -2 \\ -1 & 2 \end{pmatrix}$$

over index sets $S = \{s_1, s_2, s_3\}$ and $S' = \{s'_1, s'_2\}$ respectively. First consider the case where $J = \{s_2, s_3\}$ and $J' = \{s'_2\}$ with $w = s_3s_2s_1$ and $w' = s'_1s'_2s'_1$. Then $X^J(w, A_3)$ and $X^{J'}(w', C_2)$ are both isomorphic to the projective space \mathbb{P}^3 . However $|S(w)| = 3$ and $|S(w')| = 2$ and hence (w, A_3) and (w', C_3) cannot be Cartan equivalent.

Conversely, consider $w = s_2s_1s_3s_2$ and $J = \{s_1, s_3\}$. Then $X^\emptyset(w, A_3) = X(w, A_3)$ is singular, whereas $X^J(w, A_3)$ is the Grassmannian $\text{Gr}(2, 4)$, and hence is smooth. Clearly (w, A_3) is Cartan equivalent to itself; however $X^\emptyset(w, A_3) \not\cong X^J(w, A_3)$.

The class of Schubert varieties of partial flag varieties is much broader than the class of Schubert varieties in full flag varieties. It is not clear whether the isomorphism problem for this broader class should have a simple combinatorial solution like Cartan equivalence. We leave this as an open problem.

In proving Theorem 1.3, we need to show that a Cartan equivalence can be constructed from the isomorphism between cohomology rings. To do this, we prove that the Cartan matrix entries A_{st} for $st \leq w$ and the reduced words for w can be recovered from $H^*(X(w), A)$ along with its Schubert basis. This gives a procedure to solve a related problem of independent interest: constructing a presentation of a Schubert variety (as a Schubert variety) solely from geometric data. We outline this procedure in Section 4.1.

1.1. Examples and applications of Theorem 1.3. In this section, we illustrate the potential applications of Theorem 1.3 with several examples. We start with a basic example of how the theorem works:

Example 1.6. Let A be the following Cartan matrix over $S = \{s_1, s_2, s_3, s_4\}$:

$$\begin{pmatrix} 2 & -1 & -1 & 0 \\ -1 & 2 & -3 & 0 \\ -2 & -1 & 2 & -5 \\ 0 & 0 & -3 & 2 \end{pmatrix}$$

Let w be the reduced word $s_2s_3s_2s_1s_4$ in $W(A)$. To illustrate Definition 1.2 part (2), we list all elements $s_i s_j \leq w$ in the corresponding positions within the matrix A and then highlight the relevant data in A needed to determine the Cartan equivalence class of (w, A) .

$$(1) \quad \begin{pmatrix} & & s_1s_4 \\ s_2s_1 & & s_2s_3 & s_2s_4 \\ s_3s_1 & s_3s_2 & & s_3s_4 \\ s_4s_1 & s_4s_2 & & \end{pmatrix} \Rightarrow \begin{pmatrix} 2 & * & * & 0 \\ -1 & 2 & -3 & 0 \\ -2 & -1 & 2 & -5 \\ 0 & 0 & * & 2 \end{pmatrix}$$

The “*” entries correspond to pairs of indices (s_i, s_j) , $i \neq j$, such that $s_i s_j \not\leq w$. Suppose A' is another Cartan matrix over S which agrees with A on all the non-starred entries. By Lemma 2.2, the word $s_2s_3s_2s_1s_4$ will be reduced in $W(A')$ as well, so (w, A) and (w, A') will be Cartan equivalent. By Theorem 1.3, we would then have $X(w, A) \cong X(w', A')$.

More generally, if a Cartan matrix A' contains a submatrix of the form on the right in Equation (1) (where the “*” entries can be any number), then there will be an element $w' \in W(A')$ such that (w', A') is Cartan equivalent to (w, A) , and hence $X(w, A) \cong X(w', A')$.

One of the advantages of Theorem 1.3 is that it makes it easy to determine whether Schubert varieties in different types are isomorphic. For instance:

Example 1.7. Consider the Cartan matrices

$$A_3 = \begin{pmatrix} 2 & -1 & 0 \\ -1 & 2 & -1 \\ 0 & -1 & 2 \end{pmatrix} \quad \text{and} \quad B_3 = \begin{pmatrix} 2 & -1 & 0 \\ -1 & 2 & -1 \\ 0 & -2 & 2 \end{pmatrix}$$

of types A_3 and B_3 over $S = \{s_1, s_2, s_3\}$. Theorem 1.3 implies that

$$X(s_1s_2s_3, A_3) \cong X(s_1s_2s_3, B_3)$$

and

$$X(s_3s_2s_1, A_3) \not\cong X(s_3s_2s_1, B_3).$$

Also note that

$$X(s_2s_1s_3, A_3) \not\cong X(s_1s_3s_2, A_3).$$

Note that every Schubert variety in this example is smooth and has the same Poincaré polynomial. Hence the varieties in this example cannot be distinguished by these properties. In the last example, we have a case where $X(w, A) \not\cong X(w^{-1}, A)$.

For any A is a Cartan matrix indexed by S , and $J \subseteq S$, let $A_J := [A_{st}]_{(s,t) \in J^2}$ denote the induced Cartan matrix over J . The group $W(A_J)$ can be thought of as the subgroup of $W(A)$ generated by J , and is typically denoted by W_J . If $w \in W(A)$, then $w \in W(A_{S(w)})$. Something that shows up in the previous example is that the isomorphism type of $X(w, A)$ depends only on $A_{S(w)}$. In fact, $X(w, A) \cong X(w, A_{S(w)})$. While this follows from Theorem 1.3, it can also be easily proved without it (see for instance [RS16, Lemma 4.8]). We say $w \in W(A)$ is **fully supported** if $S(w) = S$.

Finite and affine type Cartan matrices are classified by Dynkin diagrams, which are graphs with simple edges and decorated multiedges (See Figure 1). The vertex set of the Dynkin diagram is the index set S of the Cartan matrix A , and the edge or multiedge between vertices s and t determines the matrix values A_{st} and A_{ts} . For any Cartan matrix A , we can also consider the Coxeter graph of A , which is the graph with vertex set S , and $m_{st} - 2$ edges between vertices s and t . In finite and affine types, the Coxeter graph is the Dynkin diagram with decorations removed from the multiedges.

A particularly interesting case to look at is finite versus infinite type. Recall that a Cartan matrix A is **simply-laced** if $A_{st} \in \{0, -1\}$ for all $s \neq t \in S$. Equivalently, a Cartan matrix is simply-laced if the exponents m_{st} of the Coxeter relations are either 2 or 3, i.e. the Coxeter graph is simple. Given a Cartan matrix A , it is convenient to define the **simple Coxeter graph** $\Gamma(A) := (S, E)$ to be the graph with vertex set S and edges $(s, t) \in E$ if and only if $A_{st} \neq 0$. In other words, $\Gamma(A)$ is the underlying simple graph of the Coxeter graph of A .

Lemma 1.8. *If $X(w, A) \cong X(w', A')$, then $\Gamma(A_{S(w)}) \cong \Gamma(A_{S(w')})$.*

Proof. Let $\sigma : S(w) \rightarrow S(w')$ be a Cartan equivalence. If $s, t \in S(w)$, then either $st \leq w$ or $ts \leq w$. If $st \leq w$, then $A_{st} = A'_{\sigma(s)\sigma(t)}$, while if $ts \leq w$ then $A_{ts} = A'_{\sigma(t)\sigma(s)}$. Using the fact that $A_{st} = 0$ if and only if $A_{ts} = 0$, we conclude that $A_{st} = 0$ if and only if $A'_{\sigma(s)\sigma(t)} = 0$. Therefore σ is a graph isomorphism. \square

Corollary 1.9. *If $X(w, A)$ is isomorphic to a Schubert variety in a finite type flag variety, then $\Gamma(A_{S(w)})$ is a finite type Coxeter graph. In particular, if $A_{S(w)}$ is simply-laced, then $X(w, A)$ is isomorphic to a Schubert variety of finite type if and only if $A_{S(w)}$ is of finite type.*

Finite type	Dynkin diagram	$\text{Aut}(A)$	$\text{Aut}(\Gamma(A))$
A_n		\mathbb{Z}_2	\mathbb{Z}_2
$B_n (n \geq 3)$		\mathbb{Z}_1	\mathbb{Z}_2
$C_n (n \geq 2)$		\mathbb{Z}_1	\mathbb{Z}_2
D_4		S_3	S_3
$D_n (n \geq 5)$		\mathbb{Z}_2	\mathbb{Z}_2
E_6		\mathbb{Z}_2	\mathbb{Z}_2
E_7		\mathbb{Z}_1	\mathbb{Z}_1
E_8		\mathbb{Z}_1	\mathbb{Z}_1
F_4		\mathbb{Z}_1	\mathbb{Z}_2
G_2		\mathbb{Z}_1	\mathbb{Z}_2

Affine type	Dynkin diagram	$\text{Aut}(A)$	$\text{Aut}(\Gamma(A))$
\tilde{A}_n		$I_2(n)$	$I_2(n)$
$\tilde{B}_n (n \geq 3)$		\mathbb{Z}_1	\mathbb{Z}_2
$\tilde{C}_n (n \geq 2)$		\mathbb{Z}_2	\mathbb{Z}_2
\tilde{D}_4		S_4	S_4
$\tilde{D}_n (n \geq 5)$		$\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2$	$\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2$
\tilde{E}_6		S_3	S_3
\tilde{E}_7		\mathbb{Z}_2	\mathbb{Z}_2
\tilde{E}_8		\mathbb{Z}_1	\mathbb{Z}_1
\tilde{F}_4		\mathbb{Z}_1	\mathbb{Z}_2
\tilde{G}_2		\mathbb{Z}_1	\mathbb{Z}_2
\tilde{A}_2^2		\mathbb{Z}_1	\mathbb{Z}_2
\tilde{A}_{2n}^2		\mathbb{Z}_1	\mathbb{Z}_2
\tilde{A}_{2n+1}^2		\mathbb{Z}_2	\mathbb{Z}_2
\tilde{D}_n^2		\mathbb{Z}_2	\mathbb{Z}_2
\tilde{D}_4^3		\mathbb{Z}_1	\mathbb{Z}_2
\tilde{E}_6^2		\mathbb{Z}_1	\mathbb{Z}_2

FIGURE 1. Dynkin diagrams and automorphism groups of all finite and affine Lie types. S_n denotes the symmetric group, and $I_2(n)$ denotes the dihedral group.

Proof. It is easy to verify from Figure 1 that the underlying simple graph of a finite type Dynkin diagram is also a finite type Dynkin diagram. If $X(w, A) \cong X(w', A')$ where A' has finite type, then by Lemma 1.8, the graph $\Gamma(A_{S(w)})$ is isomorphic to $\Gamma(A'_{S(w)})$ and hence has finite type. If $A_{S(w)}$ is simply-laced, then $\Gamma(A_{S(w)})$ is the Coxeter graph of $A_{S(w)}$, and hence $A_{S(w)}$ is of finite type. \square

Example 1.10. *The Coxeter graph of affine type \tilde{A}_n is a cycle for $n \geq 2$. Hence if $w \in W(\tilde{A}_n)$ is fully supported, then $X(w, \tilde{A}_n)$ is not isomorphic to any Schubert variety of finite type. The same is true for affine types \tilde{D}_n and \tilde{E}_n .*

Example 1.11. *If $A_{S(w)}$ is not simply-laced, then it's possible for $X(w, A_{S(w)})$ to be isomorphic to a finite-type Schubert variety, even if $A_{S(w)}$ is not of finite type. For instance, consider the Cartan matrices*

$$\tilde{A}_1 = \begin{pmatrix} 2 & -2 \\ -2 & 2 \end{pmatrix} \text{ and } C_2 = \begin{pmatrix} 2 & -2 \\ -1 & 2 \end{pmatrix}$$

of affine type \tilde{A}_1 and finite type C_2 over index set $\{s_1, s_2\}$. Then $X(s_1 s_2, \tilde{A}_1) \cong X(s_1 s_2, C_2)$.

Example 1.12. *The criterion in Definition 1.2 simplifies if A is simply-laced, since $A_{st} = A_{ts}$ for all s, t . More generally, suppose that A and A' are symmetric Cartan matrices, in the sense that $A_{st} = A_{ts}$ for all $s, t \in S$,¹ and let $w \in W(A)$, $w' \in W(A')$. If $X(w, A) \cong X(w', A')$, then the Cartan matrices $A_{S(w)}$ and $A'_{S(w')}$ are the same up to permutation of rows and columns. In other words, if $X(w, A)$ and $X(w', A')$ are isomorphic, then the flag varieties $\mathcal{X}(A_{S(w)})$ and $\mathcal{X}(A'_{S(w')}$ are isomorphic, with an isomorphism that identifies $X(w, A)$ and $X(w', A')$.*

We can use Lemma 1.8 for comparison to other classes of Schubert varieties as well.

Example 1.13. *The finite type A Schubert varieties are the best studied class of Schubert varieties. By Lemma 1.8, a Schubert variety $X(w, A)$ is isomorphic to a Schubert variety of finite type A if and only if*

- (1) *the simple Coxeter graph $\Gamma(A_{S(w)})$ is a disjoint union of paths, and*
- (2) *for every $st \leq w$, we have $A_{st} \in \{0, -1\}$.*

We can also use Theorem 1.3 to calculate the isomorphism classes of Schubert varieties in a fixed Kac-Moody flag variety. For any Cartan matrix A and $w \in W(A)$, let

$$\text{Isom}(w, A) := \{w' \in W(A) : X(w, A) \cong X(w', A)\}$$

denote the isomorphism class of $X(w, A)$ within the Kac-Moody flag variety $\mathcal{X}(A)$.

¹Symmetric is not the same as symmetrizable, a common condition imposed on Cartan matrices when studying the representation theory of Kac-Moody Lie algebras.

Example 1.14. *The flag variety $\mathcal{X}(A_3)$ (where A_3 is the Cartan matrix over $S = \{s_1, s_2, s_3\}$ as in Example 1.7) has 14 isomorphism classes of Schubert varieties:*

$\ell(w)$	$\text{Isom}(w, A_3)$
0	$\{1\}$
1	$\{s_1, s_2, s_3\}$
2	$\{s_1s_3\}, \{s_1s_2, s_2s_1, s_2s_3, s_3s_2\}$
3	$\{s_1s_2s_1, s_2s_3s_2\}, \{s_1s_3s_2\}, \{s_2s_1s_3\}, \{s_1s_2s_3, s_3s_2s_1\}$
4	$\{s_1s_2s_3s_2, s_3s_2s_1s_2\}, \{s_2s_1s_2s_3, s_2s_3s_2s_1\}, \{s_2s_1s_3s_2\}$
5	$\{s_2s_1s_2s_3s_2, s_2s_3s_2s_1s_2\}, \{s_3s_2s_1s_2s_3\}$
6	$\{s_3s_2s_1s_3s_2s_3\}$

In types A_4 and A_5 there are 54 and 316 Schubert isomorphism classes respectively.

When $w \in W(A)$ is not fully supported, it's possible for $X(w, A)$ to be isomorphic to $X(w', A)$ where $S(w) \neq S(w')$. For instance, $X(s, A) \cong X(t, A)$ for all $s, t \in S$. So in Example 1.14, $|\text{Isom}(s_1, A_3)| = 3$, and in general $|\text{Isom}(s, A)| = |S|$. However, in Example 1.14 all the fully supported elements w have isomorphism classes of size 1 or 2. In fact, the sets $\text{Isom}(w, A)$ when A is of finite and affine type and w is fully supported are surprisingly small. To explain this, we can bound the size of the sets $\text{Isom}(w, A)$ in terms of the automorphism groups of $\Gamma(A)$. Recall that a **diagram automorphism** of a Cartan matrix A with index set S is a bijection $\sigma : S \rightarrow S$ such that $A_{st} = A_{\sigma(s)\sigma(t)}$. We let $\text{Aut}(A)$ denote the group of diagram automorphisms of A . Diagram automorphisms play an important role in the classification of automorphisms of Kac-Moody groups [CM05, CC93, KW92]. We let $\text{Aut}(\Gamma(A))$ denote the graph automorphism group of $\Gamma(A)$, i.e. the set of bijections $\sigma : S \rightarrow S$ such that $A_{st} = 0$ if and only if $A_{\sigma(s)\sigma(t)} = 0$. Since $\Gamma(A)$ is the Dynkin diagram of the Cartan matrix where every off-diagonal non-zero entry of A is changed to -1 , $\text{Aut}(\Gamma(A))$ is also the automorphism group of some Cartan matrix.

Corollary 1.15. *If A is a Cartan matrix over S and $w \in W(A)$ is fully supported, then*

$$|\text{Isom}(w, A)| \leq |\text{Aut}(\Gamma(A))|.$$

Furthermore, if A is symmetric, then

$$|\text{Isom}(w, A)| \leq |\text{Aut}(A)|.$$

Proof. If $w \in W(A)$ is fully-supported, then a Cartan equivalence $\sigma : S \rightarrow S$ between (w, A) and (w', A) is an element of $\text{Aut}(\Gamma(A))$.

If A is symmetric, and $s \neq t \in S$, then either $st \leq w$, in which case $A_{st} = A_{\sigma(s)\sigma(t)}$, or $ts \leq w$, in which case $A_{st} = A_{ts} = A_{\sigma(t)\sigma(s)} = A_{\sigma(s)\sigma(t)}$. So σ will be in $\text{Aut}(A)$. \square

The automorphisms groups of A for A finite and (untwisted) affine types are well-known, and are shown in Figure 1. As can be verified from the table, if A is finite or affine, then $\Gamma(A)$ is also finite or affine, so $\text{Aut}(\Gamma(A))$ can be determined from the table as well. It follows that for all simple finite and affine types except affine type \tilde{A}_n , $|\text{Isom}(w, A)| \leq 24$ for all fully supported w , and for many simple finite and affine types, the bound is $|\text{Isom}(w, A)| \leq 2$.

1.2. Outline of paper. The remainder of this paper focuses on proving Theorem 1.3. In Section 2, we review some basic facts on Coxeter groups and apply them to Cartan equivalences. In Section 3 we recall the definition of the algebraic structure on Schubert varieties using the Kac-Moody Lie algebra, and prove (1) implies (2). In Section 4, we study the cohomology ring of Schubert varieties, and prove (2) implies (3) and (3) implies (1). An explanation of how to construct a presentation of a Schubert variety from geometric data is given in Subsection 4.1. We use [Kum02] as the primary reference for background material throughout the paper.

1.3. Acknowledgements. The authors thank Dave Anderson, Shrawan Kumar, Kevin Purbhoo, and Alex Yong for helpful conversations. WS was supported by NSERC DG 2018-03968

2. COMBINATORICS OF CARTAN EQUIVALENCE

In this section, we establish some basic combinatorial properties of Cartan equivalence. As usual, a word over alphabet S is a sequence (s_1, \dots, s_k) with $s_i \in S$ for $1 \leq i \leq k$. As mentioned in the introduction, if A is a Cartan matrix over index set S , then (s_1, \dots, s_k) is a reduced word if there is no way to write $s_1 \cdots s_k \in W(A)$ as a product of fewer than k elements of S . For any $w \in W(A)$, let $\text{RW}(w)$ denote the set of reduced words of w . Normally when working with $W(A)$, a word (s_1, \dots, s_k) is written as $s_1 \cdots s_k$, so the expression $s_1 \cdots s_k$ can refer to a word or to an element of $W(A)$ depending on the context. We use the same convention in this paper, but we'll also use the sequence notation for words when there is potential for confusion.

We need the following special case of the subword property for Bruhat order.

Lemma 2.1. *Let A be a Cartan matrix over a finite set S . Let $w \in W(A)$ and suppose $s, t \in S(w)$. If $A_{st} < 0$ then the following are equivalent:*

- $st \leq w$.
- The element s appears before t in some reduced word for w .
- The element s appears before t in any reduced word for w .

Otherwise, if $A_{st} = 0$, then $st = ts \leq w$.

Proof. Special case of [BB05, Theorem 2.2.2]. □

The definition of Cartan equivalence ostensibly requires us to find a reduced expression for w which corresponds to a reduced expression for w' . However, any reduced expression will do:

Lemma 2.2. *Let A and A' be Cartan matrices on S and S' respectively and let $w \in W(A)$. Let (s_1, \dots, s_k) be a reduced word for $w \in W(A)$, and suppose $\sigma : S(w) \rightarrow S'$ is an injection satisfying the condition that $A_{st} = A'_{\sigma(s)\sigma(t)}$ whenever $st \leq w$. Then $(\sigma(s_1), \dots, \sigma(s_k))$ is a reduced word for $w' := \sigma(s_1) \cdots \sigma(s_k) \in W(A')$, and (w, A) is Cartan equivalent to (w', A') . Furthermore, $(t_1, \dots, t_k) \in \text{RW}(w)$ if and only if $(\sigma(t_1), \dots, \sigma(t_k)) \in \text{RW}(w')$.*

Proof. Let m_{st} and m'_{st} be the Coxeter exponents for A and A' respectively. A word (t_1, \dots, t_k) with $t_1, \dots, t_k \in S'$ is non-reduced if and only if it is possible to apply Coxeter relations

$$\underbrace{(s, t, \dots)}_{m'_{st} \text{ times}} = \underbrace{(t, s, \dots)}_{m'_{st} \text{ times}}$$

for simple reflections $s, t \in S'$, to get a word (t'_1, \dots, t'_k) with $t'_i = t'_{i+1}$ for some $1 \leq i < k$. Suppose $(\sigma(s_1), \dots, \sigma(s_k))$ contains an alternating subword $(\sigma(s), \sigma(t), \dots)$ of length $m'_{\sigma(s)\sigma(t)}$. If $m'_{\sigma(s)\sigma(t)} \geq 3$, then $st \leq w$ and $ts \leq w$, and so $A_{st} = A'_{\sigma(s)\sigma(t)}$ and $A_{ts} = A'_{\sigma(t)\sigma(s)}$. If $m'_{\sigma(s)\sigma(t)} = 2$, then $st \leq w$, and hence $A_{st} = A'_{\sigma(s)\sigma(t)} = 0$, and so $A_{ts} = 0 = A'_{\sigma(t)\sigma(s)}$ as well. Thus in either case, $m_{st} = m'_{\sigma(s)\sigma(t)}$, and so any Coxeter relation that can be applied to $(\sigma(s_1), \dots, \sigma(s_k))$ can also be applied to (s_1, \dots, s_k) , giving another reduced word for w . Since (s_1, \dots, s_k) is reduced, we'll never get a word of the form (s'_1, \dots, s'_k) with $s'_i = s'_{i+1}$ by applying Coxeter relations. So the same is true of $(\sigma(s_1), \dots, \sigma(s_k))$, and thus $(\sigma(s_1), \dots, \sigma(s_k))$ is reduced as well. The fact that (w, A) and (w', A') are Cartan equivalent follows immediately.

Similarly, if (s_1, \dots, s_k) contains an alternating subword (s, t, \dots) of length m_{st} , then the same argument shows that $m_{st} = m'_{\sigma(s)\sigma(t)}$, and hence any Coxeter relation that can be applied to (s_1, \dots, s_k) can also be applied to $(\sigma(s_1), \dots, \sigma(s_k))$. Since $\text{RW}(w)$ and $\text{RW}(w')$ are exactly the words we get by applying Coxeter relations to (s_1, \dots, s_k) and $(\sigma(s_1), \dots, \sigma(s_k))$, it follows that $(t_1, \dots, t_k) \in \text{RW}(w)$ if and only if $(\sigma(t_1), \dots, \sigma(t_k)) \in \text{RW}(w')$. \square

Corollary 2.3. *Cartan equivalence is an equivalence relation.*

Proof. Clearly Cartan equivalence is reflexive. Suppose that $\sigma : S(w) \rightarrow S(w')$ is a Cartan equivalence as in Definition 1.2. By definition, there is a reduced word $s_1 \cdots s_n$ for w such that $\sigma(s_1) \cdots \sigma(s_n)$ is a reduced word for w' . Suppose $s't' \leq w'$, where $s' = \sigma(s)$ and $t' = \sigma(t)$. By Lemma 2.1, if $A'_{s't'} \neq 0$, then s' occurs before t' in $\sigma(s_1) \cdots \sigma(s_n)$. So $st \leq w$, and hence $A'_{s't'} = A_{st} = A_{\sigma^{-1}(s')\sigma^{-1}(t')}$. If $A'_{s't'} = 0$, then at least one of st or $ts \leq w$. In both cases $A_{\sigma^{-1}(s')\sigma^{-1}(t')} = A_{st} = 0 = A'_{s't'}$, since if $ts \leq w$, then $A_{ts} = A'_{t's'} = 0$, implying $A_{st} = 0$. So σ^{-1} is also a Cartan equivalence, and Cartan equivalence is symmetric.

For transitivity, suppose $\sigma : S(w) \rightarrow S(w')$ and $\tau : S(w') \rightarrow S(w'')$ are Cartan equivalences from (w, A) to (w', A') and (w', A') to (w'', A'') respectively. Let s_1, \dots, s_n be a reduced word for w . By Lemma 2.2, $\sigma(s_1) \cdots \sigma(s_n)$ is a reduced word for w' , and therefore $\tau(\sigma(s_1)) \cdots \tau(\sigma(s_n))$ is a reduced word for w'' . If $st \leq w$, then $\sigma(s)\sigma(t) \leq w'$. So $A_{st} = A'_{\sigma(s)\sigma(t)} = A''_{\tau(\sigma(s)), \tau(\sigma(t))}$ for all $st \leq w$, and hence $\tau \circ \sigma : S(w) \rightarrow S(w'')$ is a Cartan equivalence between (w, A) and (w'', A'') . \square

Another corollary of Lemma 2.2 is that Cartan equivalences preserve Bruhat intervals:

Corollary 2.4. *Suppose (w, A) and (w', A') are Cartan equivalent under the bijection $\sigma : S(w) \rightarrow S(w')$. For any $v \leq w$, there is $v' \leq w'$ such that (v, A) and (v', A') are Cartan equivalent under an induced bijection $\sigma|_{S(v)} : S(v) \rightarrow \sigma(S(v))$. Furthermore, this correspondence gives a poset isomorphism $\tilde{\sigma} : [e, w] \rightarrow [e, w']$.*

Proof. Fix a reduced word $w = s_1 \cdots s_k$ and let $v \leq w$. Then there is a subsequence (i_1, \dots, i_m) for which $v = s_{i_1} \cdots s_{i_m}$ is a reduced word for v . Let $v' := \sigma(s_{i_1}) \cdots \sigma(s_{i_m})$. It follows from Lemma 2.2 that (v, A) is Cartan equivalent to (v', A') and $v' \leq w'$. Also by Lemma 2.2, the element v' is independent of the choice of reduced word for v . So σ induces a function $\tilde{\sigma} : [e, w] \rightarrow [e, w']$. If $u \leq v$, then there is another subsequence (j_1, \dots, j_k) such that $u = s_{i_{j_1}} \cdots s_{i_{j_k}}$ is a reduced word for u , and hence $\tilde{\sigma}(u) = \sigma(s_{i_{j_1}}) \cdots \sigma(s_{i_{j_k}}) \leq \tilde{\sigma}(v)$, so $\tilde{\sigma}$ is order-preserving. The function $\tilde{\sigma}^{-1} : [e, w'] \rightarrow [e, w]$ induced by σ^{-1} is an inverse to $\tilde{\sigma}$, so $\tilde{\sigma}$ is a bijection. \square

3. CARTAN EQUIVALENCE AND KAC-MOODY LIE ALGEBRAS

In this section we prove that condition (1) implies condition (2) from Theorem 1.3, which we state as its own proposition:

Proposition 3.1. *If (w, A) and (w', A') are Cartan equivalent, then $X(w, A) \cong X(w', A')$ as algebraic varieties.*

The general idea of the proof of Proposition 3.1 is to use the Cartan equivalence between (w, A) and (w', A') to construct a linear map $\pi : V \rightarrow V'$ between certain vector spaces V and V' for which the algebraic structures of $X(w, A)$ and $X(w', A')$ are realized as closed embeddings into $\mathbb{P}(V)$ and $\mathbb{P}(V')$ respectively. Restricting π to $X(w, A)$ will then give an algebraic bijection between $X(w, A)$ and $X(w', A')$. Since these varieties are normal, they will be isomorphic. We begin by recalling the construction of a Kac-Moody Lie algebra $\mathfrak{g}(A)$ from a Cartan matrix A over a finite set S of size n as given in [Kum02] or [Kac90]. Let $\mathfrak{h}(A)$ be a complex vector space of dimension $2n - \text{rank}(A)$. Let $\{h_s\}_{s \in S} \subset \mathfrak{h}(A)$ denote a set of linearly independent vectors with corresponding simple root vectors $\{\alpha_s\}_{s \in S} \subset \mathfrak{h}^*(A)$ satisfying

$$\alpha_t(h_s) = A_{st}.$$

The **Kac-Moody Lie algebra** $\mathfrak{g}(A)$ is the Lie algebra generated by \mathfrak{h} , along with elements $\{e_s\}_{s \in S}$, $\{f_s\}_{s \in S}$, satisfying Lie bracket relations:

- (1) $[\mathfrak{h}, \mathfrak{h}] = 0$,
- (2) $[h, e_s] = \alpha_s(h) e_s$ and $[h, f_s] = -\alpha_s(h) f_s$ for all $h \in \mathfrak{h}(A)$ and $s \in S$,
- (3) $[e_s, f_t] = \delta_{st} h_s$,
- (4) $\text{ad}(e_s)^{1-A_{st}}(e_t) = 0$ for all $s \neq t \in S$, and
- (5) $\text{ad}(f_s)^{1-A_{st}}(f_t) = 0$ for all $s \neq t \in S$.

The algebra $\mathfrak{g}(A)$ has a triangular decomposition

$$\mathfrak{g}(A) = \mathfrak{n}^-(A) \oplus \mathfrak{h}(A) \oplus \mathfrak{n}^+(A),$$

where $\mathfrak{n}^\pm(A)$ are the subalgebras generated by $\{e_s\}_{s \in S}$ and $\{f_s\}_{s \in S}$ respectively. More specifically, the algebra $\mathfrak{n}^+(A)$ is the free Lie algebra generated by $\{e_s\}_{s \in S}$ satisfying the relations in (4) above. Similarly, $\mathfrak{n}^-(A)$ is freely generated by the $\{f_s\}_{s \in S}$ subject to the relations in (5).

In the proof of Proposition 3.1, we want to be able to work with Kac-Moody Lie algebras without specifying a Cartan matrix. To do this, consider the set the variables $\{a_{st}\}_{(s,t) \in S^2}$ and let $R_0 = \mathbb{C}[a_{st} : (s, t) \in S^2]$ denote the polynomial ring generated by these variables. Let \mathcal{R} be the free (associative non-commutative) R_0 -algebra generated by symbols \tilde{f}_s , \tilde{h}_s , and \tilde{e}_s , for $s \in S$. We use \tilde{f}^I to denote an arbitrary non-commutative monomial in the variables $\{\tilde{f}_s\}_{s \in S}$, y^I to denote a non-commutative monomial in the variables $\{\tilde{h}_s\}_{s \in S}$ and $\{\tilde{e}_s\}_{s \in S}$, and x^I to denote a non-commutative monomial in all three families of variables. A general element of \mathcal{R} can then be written as

$$\sum_I g_I(a) x^I,$$

where each $g_I(a)$ is an element of R_0 , and $g_I(a) = 0$ for all but finitely many I . We say an element of \mathcal{R} is **independent of a_{st}** if each monomial in the coefficients $g_I(a)$ do not contain a_{st} as a factor. We say that an element of \mathcal{R} is a **normal form** if it can be written as

$$\sum_{I, J} g_{IJ}(a) \tilde{f}^I y^J.$$

In other words, every monomial is ordered so that all \tilde{f}_s 's precede all \tilde{h}_s 's and \tilde{e}_s 's. Given a Cartan matrix $A = [A_{st}]_{(s,t) \in S^2}$ over S , there is a morphism $\phi(A) : \mathcal{R} \rightarrow \mathcal{U}(\mathfrak{g}(A))$ which sends

$$(\tilde{f}_s, \tilde{h}_s, \tilde{e}_s) \mapsto (f_s, h_s, e_s) \quad \text{and} \quad a_{st} \mapsto A_{st} \text{ for all } (s, t) \in S^2.$$

Note that a_{ss} is a variable in R_0 . However, $\phi(A)(a_{ss}) = 2$ for all Cartan matrices A . Hence the reader can replace a_{ss} with 2 in all the proceeding calculations.

Let $\mathfrak{g}'(A)$ denote the commutator subalgebra of $\mathfrak{g}(A)$. The morphism $\phi(A)$ maps \mathcal{R} surjectively onto $\mathcal{U}(\mathfrak{g}'(A))$, the universal enveloping algebra of $\mathfrak{g}'(A)$, since it is generated by $\{f_s, h_s, e_s \mid s \in S\}$. Given an element $\tau \in \mathcal{R}$, we say that $v \in \mathcal{R}$ is a **normal form** of τ if:

- (a) v is a normal form, and
- (b) $\phi(A)(v) = \phi(A)(\tau)$ for all Cartan matrices A over S^2 .

Using the relations (2) and (3) from the definition of $\mathfrak{g}(A)$, it is clear that every element of \mathcal{R} has a normal form.² Given $\tau \in \mathcal{R}$, we construct a specific normal form $\eta(\tau)$ as follows:

- if $\tilde{e}_s \tilde{f}_t$ or $\tilde{h}_s \tilde{f}_t$ does not occur in any monomial of τ , then set $\eta(\tau) = \tau$.
- Otherwise find the rightmost occurrence of $\tilde{e}_s \tilde{f}_t$ or $\tilde{h}_s \tilde{f}_t$ in each monomial of τ , and either
 - replace $\tilde{e}_s \tilde{f}_t$ with $\tilde{f}_t \tilde{e}_s + \delta_{st} \tilde{h}_s$, or
 - replace $\tilde{h}_s \tilde{f}_t$ with $\tilde{f}_t \tilde{h}_s - a_{st} \tilde{f}_t$.
- Repeat for each monomial in the resulting sum.

We call $\eta(\tau)$ **the normal form** of τ .

Example 3.2. Let $S = \{s_1, s_2, s_3\}$ and $\tau = \tilde{h}_1 \tilde{e}_2 \tilde{e}_3 \tilde{f}_2$. Applying the algorithm for $\eta(\tau)$ yields:

$$\begin{aligned} \tau = \tilde{h}_1 \tilde{e}_2 (\tilde{e}_3 \tilde{f}_2) &\rightarrow \tilde{h}_1 \tilde{e}_2 (\tilde{f}_2 \tilde{e}_3) = \tilde{h}_1 (\tilde{e}_2 \tilde{f}_2) \tilde{e}_3 \\ &\rightarrow \tilde{h}_1 (\tilde{f}_2 \tilde{e}_2 + \tilde{h}_2) \tilde{e}_3 = (\tilde{h}_1 \tilde{f}_2) \tilde{e}_2 \tilde{e}_3 + \tilde{h}_1 \tilde{h}_2 \tilde{e}_3 \\ &\rightarrow (\tilde{f}_2 \tilde{h}_1 - a_{12} \tilde{f}_2) \tilde{e}_2 \tilde{e}_3 + \tilde{h}_1 \tilde{h}_2 \tilde{e}_3 \\ &= \tilde{f}_2 \tilde{h}_1 \tilde{e}_2 \tilde{e}_3 - a_{12} \tilde{f}_2 \tilde{e}_2 \tilde{e}_3 + \tilde{h}_1 \tilde{h}_2 \tilde{e}_3 = \eta(\tau). \end{aligned}$$

Lemma 3.3. Let τ be an element of \mathcal{R} which is independent of a_{st} with $s \neq t$. Suppose that, in every monomial of τ , every \tilde{f}_t occurs to the left of every \tilde{h}_s and \tilde{f}_s . Then $\eta(\tau)$ is also independent of a_{st} .

Proof. An a_{st} is created in the above procedure when we switch

$$\tilde{h}_s \tilde{f}_t \rightarrow \tilde{f}_t \tilde{h}_s - a_{st} \tilde{f}_t$$

and \tilde{h}_s is created whenever we switch

$$\tilde{e}_s \tilde{f}_s \rightarrow \tilde{f}_s \tilde{e}_s + \tilde{h}_s.$$

When the hypothesis of the lemma holds, all \tilde{h}_s 's occur to the right of all \tilde{f}_t 's in every monomial of τ . If we always apply the above steps to the rightmost occurrence of $\tilde{e}_r \tilde{f}_u$ or $\tilde{h}_r \tilde{f}_u$, then we never create a \tilde{h}_s to the left of any \tilde{f}_t . Thus we will never create any a_{st} 's. Since τ is independent of a_{st} , we conclude that $\eta(\tau)$ is also independent of a_{st} . \square

²This is analogous to the isomorphism $\mathcal{U}(\mathfrak{g}(A)) \cong \mathcal{U}(\mathfrak{n}^-(A)) \otimes \mathcal{U}(\mathfrak{h}(A)) \oplus \mathfrak{n}^+(A)$. Working with \mathcal{R} allows us to use this isomorphism without specifying the Cartan matrix A .

Example 3.2 illustrates that if \tilde{f}_t occurs to the right of \tilde{h}_s in τ , then $\eta(\tau)$ may depend on a_{st} even if τ does not.

For the rest of this section, we work with two Cartan matrices A and A' over finite sets S and S' with $|S| = |S'|$. We will also fix a bijection $\sigma : S \rightarrow S'$ and for the sake of notational simplicity, we let $A'_{st} := A'_{\sigma(s)\sigma(t)}$ for all $s, t \in S$ (note that we are not assuming that $A_{st} = A'_{st}$). Let $\mathfrak{g}(A)$ and $\mathfrak{g}(A')$ denote the corresponding Kac-Moody algebras to A and A' . We will use the notation above for the generators of $\mathfrak{g}(A)$, and refer to the generators of $\mathfrak{g}(A')$ by

$$h'_s := h_{\sigma(s)}, \quad e'_s := e_{\sigma(s)}, \quad \text{and} \quad f'_s := f_{\sigma(s)}.$$

Note that we always use $\mathfrak{g}(A')$ instead of \mathfrak{g}' to avoid confusion with the commutator subgroup of $\mathfrak{g} = \mathfrak{g}(A)$.

Recall that a representation for $\mathfrak{g}(A)$ is said to be integrable if e_s and f_s are locally nilpotent for every $s \in S$. Given a dominant integral weight λ , we let $L^{\max}(\lambda) = L^{\max}(\lambda, A)$ denote the maximal integrable $\mathfrak{g}(A)$ module of highest weight λ .

Given two Cartan matrices A and A' over S and S' as above, we write $A \leq A'$ to mean that $A_{st} \leq A'_{st}$ for all $s, t \in S$ (in other words, $|A_{st}| \geq |A'_{st}|$). For the next few lemmas, we assume the following hypotheses:

- Hypothesis 3.4.** (i) A and A' are Cartan matrices over S and S' with $A \leq A'$.
 (ii) $\lambda \in \mathfrak{h}(A)^*$ and $\lambda' \in \mathfrak{h}(A')^*$ are dominant integrable weights such that $\lambda(h_s) \geq \lambda'(h'_s)$ for all $s \in S$.
 (iii) $V = L^{\max}(\lambda, A)$ and $V' = L^{\max}(\lambda', A')$ are the maximal integrable modules with highest weights λ and λ' .
 (iv) ω and ω' are highest weight vectors for V and V' respectively.

Lemma 3.5. *Suppose hypotheses 3.4 hold. Then:*

- (a) *There are surjective Lie algebra morphisms*

$$\psi^+ : \mathfrak{n}^+(A) \rightarrow \mathfrak{n}^+(A') \quad \text{and} \quad \psi^- : \mathfrak{n}^-(A) \rightarrow \mathfrak{n}^-(A')$$

mapping $e_s \mapsto e'_s$ and $f_s \mapsto f'_s$ respectively.

- (b) *There is a surjective $\mathfrak{n}^-(A)$ -module morphism $\pi : V \rightarrow V'$ sending ω to ω' , where V' is regarded as a $\mathfrak{n}^-(A)$ -module using ψ^- .*

Proof. Part (a), the fact that $\psi^+ : \mathfrak{n}^+(A) \rightarrow \mathfrak{n}^+(A')$ is well-defined and surjective follows from the fact that $1 - A_{st} \geq 1 - A'_{st}$ for all for all $s, t \in S$. In particular, the relation $\text{ad}(e'_s)^{1-A_{st}}(e'_t) = 0$ holds in $\mathfrak{n}^+(A')$. Similar argument holds for the map $\psi^- : \mathfrak{n}^-(A) \rightarrow \mathfrak{n}^-(A')$.

For part (b), recall that $L^{\max}(\lambda, A)$ is the quotient of the Verma module (see [Kum02, Definition 2.1.1 and Definition 2.1.5])

$$M(\lambda, A) := \mathcal{U}(\mathfrak{n}^-(A)) \otimes_{\mathbb{C}} \mathbb{C}_{\lambda}$$

by the $\mathfrak{g}(A)$ -module $M^1(\lambda, A)$ generated by $f_s^{\lambda(h_s)+1} v_{\lambda}$, where $s \in S$ and v_{λ} is the cyclic weight vector $1 \otimes 1$ in $M(\lambda, A)$. By [Kum02, Lemma 2.1.6], the action of \mathfrak{n}^+ sends the generators $f_s^{\lambda(h_s)+1} v_{\lambda}$ to zero, so that $M^1(\lambda, A)$ is also the $\mathcal{U}(\mathfrak{n}^-(A))$ -module generated by $f_s^{\lambda(h_s)+1} v_{\lambda}$. By part (a), for any cyclic weight vector $v_{\lambda'}$ of $M(\lambda', A')$, there is a surjective $\mathcal{U}(\mathfrak{n}^-(A))$ -module map $\pi : M(\lambda, A) \rightarrow M(\lambda', A')$ sending $v_{\lambda} \mapsto v_{\lambda'}$. Now

$$\pi(f_s^{\lambda(h_s)+1} v_{\lambda}) = (f'_s)^{\lambda(h_s)+1} v_{\lambda'}$$

belongs to $M^1(\lambda', A')$ for all $s \in S$ since $\lambda(h_s) + 1 \geq \lambda'(h'_s) + 1$. Thus we get an induced morphism $\pi : L^{\max}(\lambda, A) \rightarrow L^{\max}(\lambda', A')$ on the quotients. \square

We now recall the root system of a Kac-Moody algebra $\mathfrak{g}(A)$ and the inversion set of an element $w \in W(A)$. Let $Q = Q(A) := \bigoplus_{s \in S} \mathbb{Z} \alpha_s \subset \mathfrak{h}^*$ denote the root lattice and let $R(A) \subseteq Q$ denote the root system of $\mathfrak{g}(A)$. We can decompose the set $R(A) = R^+(A) \sqcup R^-(A)$ where $R^+(A)$ and $R^-(A)$ denote the subsets of positive and negative roots respectively. We have that $\mathfrak{n}^\pm(A) = \bigoplus_{\alpha \in R^\pm(A)} \mathfrak{g}(A)_\alpha$ where $\mathfrak{g}(A)_\alpha$ is the root space corresponding to α . The Weyl group $W(A)$ (which is generated by S) acts on Q by

$$s(\alpha_t) := \alpha_t - \alpha_t(h_s) \alpha_s$$

for $s, t \in S$. Given $w \in W(A)$, define the **inversion set**

$$I(w) := \{\alpha \in R^+(A) : w^{-1}(\alpha) \in R^-(A)\}.$$

Note that the inversion set is finite of size $\ell(w)$ and lies inside the sublattice $\bigoplus_{s \in S(w)} \mathbb{Z} \alpha_s$.

Lemma 3.6. *Let A and A' be Cartan matrices over S and S' with $w \in W(A)$ and $w' \in W(A')$. Suppose that w is Cartan equivalent to w' under the bijection $\sigma : S(w) \rightarrow S(w')$. Then the induced isomorphism*

$$\sigma : \bigoplus_{s \in S(w)} \mathbb{Z} \alpha_s \rightarrow \bigoplus_{s' \in S(w')} \mathbb{Z} \alpha'_{s'}$$

given by $\alpha_s \mapsto \alpha'_{\sigma(s)}$ identifies $I(w)$ with $I(w')$.

Proof. Let $w = s_1 \cdots s_k$ be a reduced expression. Then $I(w) = \{\beta_1, \dots, \beta_k\}$, where

$$\beta_\ell = s_1 \cdots s_{\ell-1}(\alpha_{s_\ell}).$$

Given $1 \leq \ell \leq k$, we want to show that $\sigma(\beta_\ell)$ belongs to $I(w')$. First note that if $st \leq w$, then $A_{st} = A'_{\sigma(s)\sigma(t)}$ and hence

$$\sigma(s(\alpha_t)) = \sigma(\alpha_t - A_{st}\alpha_s) = \alpha'_{\sigma(t)} - A'_{\sigma(s)\sigma(t)}\alpha'_{\sigma(s)} = \sigma(s)(\alpha'_{\sigma(t)}).$$

In particular, $\sigma(s_{\ell-1}(\alpha_{s_\ell})) = \sigma(s_{\ell-1})(\alpha'_{\sigma(s_\ell)})$. For the purpose of induction, suppose that

$$\sigma(s_m \cdots s_{\ell-1}(\alpha_{s_\ell})) = \sigma(s_m) \cdots \sigma(s_{\ell-1})(\alpha'_{\sigma(s_\ell)}).$$

Let $s_{m-1} = s$ and write $s_m \cdots s_{\ell-1}(\alpha_{s_\ell}) = \sum_{t \in S} c_t \alpha_t$ where $c_t \in \mathbb{Z}$. If $st \not\leq w$, then t does not appear to the right of any s in the reduced expression for w . This implies the coefficient $c_t = 0$ for all $t \in S$ such that $st \not\leq w$. Hence

$$\begin{aligned} \sigma(s_{m-1} s_m \cdots s_{\ell-1}(\alpha_{s_\ell})) &= \sum_{t \in S} c_t \sigma(s_{m-1}(\alpha_t)) = \sum_{t \in S} c_t \sigma(s_{m-1})(\alpha'_{\sigma(t)}) \\ &= \sigma(s_{m-1}) \left(\sum_{t \in S} c_t \alpha'_{\sigma(t)} \right) = \sigma(s_{m-1}) \sigma(s_m \cdots s_{\ell-1}(\alpha_{s_\ell})). \end{aligned}$$

By induction, we have

$$\sigma(\beta_\ell) = \sigma(s_1 \cdots s_{\ell-1}(\alpha_{s_\ell})) = \sigma(s_1) \cdots \sigma(s_{\ell-1})(\alpha'_{\sigma(s_\ell)}).$$

Since $\sigma(s_1) \cdots \sigma(s_k)$ is a reduced expression for w' , we have $\sigma(\beta_\ell) \in I(w')$. \square

For $w \in W(A)$, define

$$\mathfrak{n}^+(A)_w := \bigoplus_{\alpha \in I(w)} \mathfrak{g}(A)_\alpha.$$

Since $I(w)$ is closed under bracket, $\mathfrak{n}^+(A)_w$ is a finite-dimensional nilpotent Lie algebra. We now add two additional hypotheses:

Hypothesis 3.7. (v) Suppose that $w \in W(A)$ such that $A_{st} = A'_{st}$ for all $st \leq w$.
 (vi) Let $w' \in W(A')$ denote the element which is Cartan equivalent to w under the bijection $\sigma : S \rightarrow S'$.

Lemma 3.8. Suppose that hypotheses 3.4 and 3.7 hold. Then ψ^+ induces an isomorphism $\mathfrak{n}^+(A)_w \rightarrow \mathfrak{n}^+(A')_{w'}$.

Proof. Because $\mathfrak{n}^+(A)$ is generated by $\{e_s\}_{s \in S}$, it is spanned by Lie monomials in these same variables. Since each monomial is contained in a root space, we have that $\psi^+(\mathfrak{g}(A)_\alpha) \subseteq \mathfrak{g}(A')_{\sigma(\alpha)}$ where $\sigma : S(w) \rightarrow S(w')$ is given in Lemma 3.6. The lemma now follows from Lemma 3.6. \square

Lemma 3.9. Let $\exp : \mathfrak{g}(A) \rightarrow \mathcal{G}(A)$ denote the exponential map and suppose that hypotheses 3.4 and 3.7 hold. Further suppose that $\lambda(h_s) = \lambda'(h'_s)$ for all $s \in S$. Then

$$\pi(\exp(z)v \cdot \omega) = \exp(\psi^+(z))\sigma(v) \cdot \omega'$$

for all $z \in \mathfrak{n}^+(A)$ and $v \leq w$.

Proof. Let $v = s_1 \dots s_k$ be a reduced expression. If $st \not\leq w$ then $st \not\leq v$, so $\sigma(s_1) \dots \sigma(s_k)$ is also a reduced expression for $\sigma(v)$. By [Kum02, Definition 1.3.2 (5)], the action of s (resp. $\sigma(s)$) on V (resp. V') is given by

$$\exp(f_s)\exp(-e_s)\exp(f_s) \quad (\text{resp.} \quad \exp(f'_s)\exp(-e'_s)\exp(f'_s)).$$

Since V is integrable, for any $s \in S$ and $u \in V$, there is an integer N_0 such that

$$\exp(-e_s) \cdot u = \left(\sum_{k=0}^N \frac{(-e_s)^k}{k!} \right) \cdot u \quad \text{and} \quad \exp(f_s) \cdot u = \left(\sum_{k=0}^N \frac{f_s^k}{k!} \right) \cdot u$$

for all $N \geq N_0$. The same applies to $\exp(-e'_s)$ and $\exp(f'_s)$ when acting on V' . Since V and V' are highest-weight modules, we also have similar expressions for $\exp(z)$ and $\exp(\psi^+(z))$. Thus we can find an integer $N \gg 0$ such that

$$\begin{aligned} \exp(z)v \cdot \omega &= \exp(z)(s_1 \dots s_k) \cdot \omega = \left(\sum_{j=0}^N \frac{z^j}{j!} \right) \cdot \left(\sum_{(l_1, m_1, n_1) \in [N]^3} \frac{f_{s_1}^{l_1} (-e_{s_1})^{m_1} f_{s_1}^{n_1}}{(l_1!)(m_1!)(n_1!)} \right) \dots \\ &\dots \left(\sum_{(l_k, m_k, n_k) \in [N]^3} \frac{f_{s_k}^{l_k} (-e_{s_k})^{m_k} f_{s_k}^{n_k}}{(l_k!)(m_k!)(n_k!)} \right) \cdot \omega \end{aligned}$$

and

$$\begin{aligned} \exp(\psi^+(z))\sigma(v) \cdot \omega' &= \left(\sum_{j=0}^N \frac{\psi^+(z)^j}{j!} \right) \cdot \left(\sum_{(l_1, m_1, n_1) \in [N]^3} \frac{(f'_{s_1})^{l_1} (-e'_{s_1})^{m_1} (f'_{s_1})^{n_1}}{(l_1!)(m_1!)(n_1!)} \right) \dots \\ &\dots \left(\sum_{(l_k, m_k, n_k) \in [N]^3} \frac{(f'_{s_k})^{l_k} (-e'_{s_k})^{m_k} (f'_{s_k})^{n_k}}{(l_k!)(m_k!)(n_k!)} \right) \cdot \omega', \end{aligned}$$

where $[N] := \{0, 1, \dots, N\}$. Since $\mathfrak{n}^+(A)$ is generated by $\{e_s \mid s \in S\}$, the element z is a polynomial in $\{e_s \mid s \in S\}$. Since ψ^+ sends $e_s \mapsto e'_s$, we can thus write down an element Φ in \mathcal{R} such that

$$\exp(z)v \cdot \omega = \phi(A)(\Phi) \cdot \omega \quad \text{and} \quad \exp(\psi^+(z))\sigma(v) \cdot \omega' = \phi(A')(\Phi) \cdot \omega'.$$

Now replace Φ with the normal form

$$\eta(\Phi) = \sum_{I,J} g_{IJ}(a) \tilde{f}^I y^J.$$

Note that $h_s \omega = \lambda(h_s) \omega$ and $e_s \omega = 0$ for all $s \in S$. Hence, for any I in the above sum, we have that $\phi(A)(y^I) \omega = c_I \omega$ for some $c_I \in \mathbb{Z}$. Since $\lambda(h_s) = \lambda'(h'_s)$, we also have $\phi(A')(y^I) \omega' = c_I \omega'$. Letting

$$\eta_0 = \sum_{I,J} c_J g_{IJ}(a) \tilde{f}^I$$

yields

$$\phi(A)(\Phi) \cdot \omega = \phi(A)(\eta_0) \cdot \omega \quad \text{and} \quad \phi(A')(\Phi) \cdot \omega' = \phi(A')(\eta_0) \cdot \omega'.$$

If $st \not\leq w$ then $st \not\leq v$, and this implies that f_t occurs to the left of every f_s in Φ by Lemma 2.1. Since h_s does not occur in Φ , Lemma 3.3 implies that $\eta(\Phi)$ is independent of a_{st} . This also implies that η_0 is independent of a_{st} . On the other hand, if $st \leq w$, then $A_{st} = A'_{st}$. So

$$\eta_0 = \sum_{I,J} c_J g_{IJ}(A) \tilde{f}^I = \sum_{I,J} c_J g_{IJ}(A') \tilde{f}^I.$$

It follows that $\phi(A')(\eta_0) = \psi^-(\phi(A)(\eta_0))$. Finally, the map π is ψ^- -equivariant, so

$$\begin{aligned} \pi(\exp(z)v \cdot \omega) &= \pi(\phi(A)(\Phi) \cdot \omega) = \pi(\phi(A)(\eta_0) \cdot \omega) \\ &= \psi^-(\phi(A)(\eta_0))\pi(\omega) = \phi(A')(\eta_0)\omega' \\ &= \phi(A')(\Phi)\omega' = \exp(\psi^+(z))\sigma(v) \cdot \omega'. \end{aligned}$$

□

Proof of Proposition 3.1. Let A and A' be Cartan matrices over S and S' respectively and that $\sigma : S(w) \rightarrow S(w')$ gives a Cartan equivalence between (w, A) and (w', A') . Without loss of generality, we can assume that $S = S(w)$ and $S' = S(w')$ (for instance, see [RS16, Lemma 4.8]). In particular, hypotheses 3.7 will hold. Assume (now with loss of generality) that $A \leq A'$. By [Kum02, Definition 7.1.19], the stable variety structure on $X(w, A)$ is induced (meaning that there is a closed embedding) by taking the map

$$\mathcal{G}(A)/\mathcal{B}(A) \rightarrow \mathbb{P}(V) : g \mapsto g \cdot \omega,$$

where $\mathcal{B}(A)$ is the Borel subgroup of $\mathcal{G}(A)$, and ω is the highest weight vector of $V = L^{\max}(\lambda, A)$ for a large enough dominant weight λ . By possibly increasing λ , we can assume that the variety structure on $X(w', A')$ is induced by taking the map

$$\mathcal{G}(A')/\mathcal{B}(A') \rightarrow \mathbb{P}(V') : g \mapsto g \cdot \omega',$$

where ω' is the highest weight of $V' = L^{\max}(\lambda', A')$ for a dominant weight λ' with $\lambda'(h_{\sigma(s)}) = \lambda(h_s)$. Hence all parts of Hypothesis 3.4 are satisfied.

Consider the map $\pi : X(w, A) \rightarrow \mathbb{P}(V')$ induced by $\pi : V \rightarrow V'$. By [Kum02, 6.2.E.1], every point of $X(w, A)$ can be written uniquely as $\exp(x)v \cdot \omega$ for some $v \leq w$ and

$x \in \mathfrak{n}^+(A)_v$. By Corollary 2.4 and Lemmas 3.8 and 3.9, the map π induces a bijection $X(w, A) \rightarrow X(w', A')$. Since $X(w', A')$ is a normal variety (see [Kum02, Theorem 8.2.2 (b)]), the map π restricted to $X(w, A)$ must be an algebraic isomorphism.

Now suppose that $A \not\leq A'$. Define \tilde{A} by $\tilde{A}_{st} = \min(A_{st}, A'_{st})$. By Lemma 2.2, there is an element \tilde{w} in $W(\tilde{A})$ such that (\tilde{w}, \tilde{A}) is Cartan equivalent to both (w, A) and (w', A') . We also have $\tilde{A} \leq A$ and $\tilde{A} \leq A'$ and hence the above argument implies $X(w, A) \cong X(\tilde{w}, \tilde{A}) \cong X(w', A')$. \square

4. THE COHOMOLOGY RING OF SCHUBERT VARIETIES

In this section, we finish the proof of Theorem 1.3 by showing that condition (2) implies condition (3), and that condition (3) implies condition (1). Let A be a Cartan matrix over S , and let $w \in W(A)$. Recall that $\mathcal{B} = \mathcal{B}(A)$ is the Borel subgroup of the Kac-Moody group $\mathcal{G}(A)$. The Schubert variety $X(w, A)$ has a stratification given by its decomposition into Schubert cells

$$X(w, A) = \overline{\mathcal{B}w\mathcal{B}}/\mathcal{B} = \bigsqcup_{u \leq w} \mathcal{B}u\mathcal{B}/\mathcal{B}.$$

Let x_v denote the fundamental class of the Schubert subvariety $X(v, A) = \overline{\mathcal{B}v\mathcal{B}}/\mathcal{B} \subseteq X(w, A)$ in the integral homology group $H_{2\ell(v)}(X(w, A)) := H_{2\ell(v)}(X(w, A), \mathbb{Z})$. Equivalently, if we consider $X(v, A)$ as a cycle in the Chow group $\text{Ch}_*(X(w, A))$, then

$$x_v = \text{cl}(X(v, A))$$

where $\text{cl} : \text{Ch}_*(X(w, A)) \rightarrow H_*(X(w, A))$ denotes the cycle map between Chow groups and homology. It is well known that the Schubert homology classes $\{x_v\}_{v \leq w}$ form a \mathbb{Z} -basis of $H_*(X(w, A))$. The corresponding Schubert basis $\{\xi_v\}_{v \leq w}$ in integral cohomology $H^*(X(w, A)) := H^*(X(w, A), \mathbb{Z})$ is defined (using the identification $H^*(X(w, A)) \simeq \text{Hom}_{\mathbb{Z}}(H_*(X(w, A)), \mathbb{Z})$) by

$$\xi_v(x_u) := \delta_{vu}.$$

We now prove that condition (2) implies condition (3) in Theorem 1.3.

Proposition 4.1. *Let A and A' be Cartan matrices with $w \in W(A)$ and $w' \in W(A')$ and suppose that $\phi : X(w, A) \rightarrow X(w', A')$ is an algebraic isomorphism. Then the induced map*

$$\phi^* : H^*(X(w', A')) \rightarrow H^*(X(w, A))$$

is a graded ring isomorphism that identifies Schubert bases.

Proof of Proposition 4.1. Let $\text{Ch}_*(X(w, A))$ and $\text{Ch}_*(X(w', A'))$ denote the Chow groups of $X(w, A)$ and $X(w', A')$. Since ϕ is an algebraic isomorphism, the induced isomorphism of Chow groups $\phi_* : \text{Ch}_*(X(w, A)) \rightarrow \text{Ch}_*(X(w', A'))$ preserves the cone of effective classes. By [FMSS95, Corollary to Theorem 1], any effective class is a nonnegative \mathbb{Z} -linear combination of Schubert cycles.³ Hence the Schubert cycles form the minimal extremal rays of the effective cone, so ϕ_* maps Schubert cycles to Schubert cycles. Since the Schubert classes in cohomology are dual to the Schubert cycles in homology, ϕ^* maps Schubert classes to Schubert classes. \square

To finish the proof of Theorem 1.3, we just need to show that condition (3) implies condition (1):

³This idea effectively goes back to [Hir84]; see also [KN98].

Proposition 4.2. *Let A and A' be Cartan matrices with $w \in W(A)$ and $w' \in W(A')$. Suppose there is graded ring isomorphism*

$$\phi : H^*(X(w, A)) \rightarrow H^*(X(w', A'))$$

which identifies Schubert bases. Then (w, A) is Cartan equivalent to (w', A') .

For the proof of Proposition 4.2, suppose we have two Schubert varieties $X(w, A)$ and $X(w', A')$ with an isomorphism ϕ between cohomology rings as in the proposition. Let $E := \{\xi_v\}_{v \leq w}$ denote the Schubert basis of $H^*(X(w, A))$. As sets, there is a bijection $E \rightarrow [e, w]$ by mapping $\xi_u \mapsto u$. Define $\tilde{S}(w) := E \cap H^2(X(w, A))$. The bijection $E \rightarrow [e, w]$ sends ξ_s to s , and hence identifies $\tilde{S}(w)$ with $S(w)$. We similarly define E' and $\tilde{S}(w')$ for $H^*(X(w', A'))$. By assumption, the isomorphism ϕ restricted to E gives a degree-preserving bijection $\phi : E \rightarrow E'$. Let $\sigma : S(w) \rightarrow S(w')$ be the bijection corresponding to $\phi|_{\tilde{S}(w)}$, so in particular $\phi(\xi_s) = \xi_{\sigma(s)}$ for any $s \in S(w)$. To prove Proposition 4.2, we will show that σ is a Cartan equivalence. For this, we need to show that all of the relevant entries of the Cartan matrix and the reduced word structure of $w \in W(A)$ can be reconstructed from the ring structure of $H^*(X(w, A))$.

Recall the set of roots $R(A) \subseteq Q(A) = \bigoplus_{s \in S} \mathbb{Z}\alpha_s \subseteq \mathfrak{h}^*(A)$. For notational simplicity, we denote $R := R(A)$ and $R^\pm := R^\pm(A)$. A root $\beta \in R$ is said to be a real root if $\beta = v(\alpha_s)$ for some $v \in W(A)$ and $s \in S$. The Weyl group $W(A)$ acts on the dual space $\mathfrak{h}(A)$ by

$$s(h) := h - \alpha_s(h) h_s$$

for $s \in S$ and $h \in \mathfrak{h}(A)$, where $h_s \in \mathfrak{h}(A)$, $s \in S$ are vectors as in Section 3 (so $\alpha_t(h_s) = A_{st}$). The restriction of this action to $\bigoplus_{s \in S} \mathbb{Z}h_s$ is the dual action to the action of $W(A)$ on $Q(A)$. For a real root $\beta = v(\alpha_s)$, the corresponding coroot is $\beta^\vee := v(h_s) \in \mathfrak{h}(A)$, and the corresponding reflection is $s_\beta := vsv^{-1} \in W(A)$. Note that $\alpha_t^\vee = h_t$ and $s_{\alpha_t} = t$ for all $t \in S$. Finally, we pick fundamental coweights $\{\omega_s\}_{s \in S} \subseteq \mathfrak{h}^*(A)$ satisfying the formula

$$\omega_s(h_t) := \delta_{st}.$$

The main computational tool used to prove Proposition 4.2 is Chevalley's (non-equivariant) formula for multiplying Schubert classes by simple Schubert classes. We use here the version from [Kum02, Theorem 11.1.7 (i)].

Proposition 4.3. *(Chevalley's formula) For any $\xi_s \in \tilde{S}(w)$ and $\xi_u \in E$,*

$$\xi_s \cdot \xi_u = \sum \omega_s(u^{-1}(\beta^\vee)) \xi_{s_\beta u}$$

where the sum is over all real roots $\beta \in R^+$ such that $\ell(s_\beta u) = \ell(u) + 1$ and $s_\beta u \leq w$.

Note that if $\ell(s_\beta u) = \ell(u) + 1$, then $u^{-1}(\beta^\vee) \in R^+$ and hence $\omega_s(u^{-1}(\beta^\vee))$ is a nonnegative integer. For any $F \in H^*(X(w, A))$, define **the support of F** as

$$\text{Supp}(F) := \{\xi_v \in E : F(x_v) \neq 0\}.$$

In other words, if we write $F = \sum_{v \leq w} c_v \xi_v$, then $\xi_v \in \text{Supp}(F)$ if and only if $c_v \neq 0$. Let \prec be the partial order on E generated by the covering relations $\xi_u \prec \xi_v$ for u and v such that $\xi_v \in \text{Supp}(\xi_s \xi_u)$ for some $s \in S(w)$. In the next lemma, we show that \prec corresponds to Bruhat order \leq on $[e, w]$:

Lemma 4.4. *$\xi_u \prec \xi_v$ if and only if $u \leq v$.*

Proof. It suffices to consider covering relations. Recall that $u \leq v$ if and only if $v = s_\beta u$ with $\ell(v) = \ell(u) + 1$ for some real root $\beta \in R^+$. Hence $\xi_u \prec \xi_v$ immediately implies $u \leq v$. Conversely, if $u \leq v$ with $\ell(v) = \ell(u) + 1$, then $v = s_\beta u$ where $u^{-1}(\beta^\vee) \in R^+$. So there exists $s \in S(w)$ for which $\omega_s(u^{-1}(\beta^\vee)) > 0$, implying that $\xi_v \in \text{Supp}(\xi_s \cdot \xi_u)$. Hence $\xi_u \prec \xi_v$. \square

Next we show that Cartan matrix entries A_{st} for $st \leq w$ can be recovered from products of simple Schubert classes.

Lemma 4.5. *Let $s, t \in S(w)$ such that $s \neq t$. Then:*

(1) $\text{Supp}(\xi_s \xi_t) \cap \text{Supp}(\xi_t^2) \neq \emptyset$ if and only if $st \leq w$ and $st \neq ts$. In this case,

$$\text{Supp}(\xi_s \xi_t) \cap \text{Supp}(\xi_t^2) = \{\xi_{st}\} \quad \text{and} \quad A_{st} = -\xi_t^2(x_{st}).$$

(2) $\text{Supp}(\xi_s \xi_t) \cap (\text{Supp}(\xi_t^2) \cup \text{Supp}(\xi_s^2)) = \emptyset$ if and only if $st = ts$. In this case, $A_{st} = 0$.

Proof. By Chevalley's formula, we have

$$\xi_t^2 = \sum_{\substack{s': s't \leq w \\ s' \neq t}} \omega_t(t(\alpha_{s'}^\vee)) \xi_{s't} + \sum_{\substack{s'': ts'' \leq w \\ ts'' \neq s''t}} \omega_t(\alpha_{s''}^\vee) \xi_{ts''}.$$

In the second sum, since $t \neq s''$, $\omega_t(\alpha_{s''}^\vee) = 0$. So

$$\xi_t^2 = \sum_{\substack{s': s't \leq w \\ s' \neq t}} \omega_t(\alpha_{s'}^\vee - A_{s't} \alpha_t^\vee) \xi_{s't} = \sum_{\substack{s': s't \leq w \\ s' \neq t}} -A_{s't} \xi_{s't}.$$

Since $A_{s't} = 0$ if and only if $s't = ts'$, we conclude that

$$\text{Supp}(\xi_t^2) = \{\xi_{s't} \mid s' \in S(w) \text{ such that } s't \neq ts' \text{ and } s't \leq w\}.$$

If $s \neq t$, then

$$\begin{aligned} \xi_s \xi_t &= \sum_{\substack{s': s't \leq w \\ s' \neq t}} \omega_s(t(\alpha_{s'}^\vee)) \xi_{s't} + \sum_{\substack{s'': ts'' \leq w \\ ts'' \neq s''t}} \omega_s(\alpha_{s''}^\vee) \xi_{ts''} \\ &= \sum_{\substack{s': s't \leq w \\ s' \neq t}} \omega_s(\alpha_{s'}^\vee - A_{s't} \alpha_t^\vee) \xi_{s't} + \sum_{\substack{s'': ts'' \leq w \\ ts'' \neq s''t}} \omega_s(\alpha_{s''}^\vee) \xi_{ts''} \end{aligned}$$

If $st \leq w$ then the first sum is ξ_{st} , and otherwise the first sum is 0. If $st \neq ts$, and $ts \leq w$, then the second sum is ξ_{ts} , while otherwise it's 0. So we conclude that $\text{Supp}(\xi_s \xi_t) \subseteq \{\xi_{st}, \xi_{ts}\}$ with $\xi_{st} \in \text{Supp}(\xi_s \xi_t)$ if and only if $st \leq w$. This proves part (1). For part (2), note that either $st \leq w$ or $ts \leq w$. So if $\text{Supp}(\xi_t^2) \cup \text{Supp}(\xi_s^2)$ contains neither ξ_{st} or ξ_{ts} , we must have $st = ts$. Conversely, if $st = ts$, then $\xi_{st} = \xi_{ts}$ cannot belong to either $\text{Supp}(\xi_t^2)$ or $\text{Supp}(\xi_s^2)$. Since $\text{Supp}(\xi_s \xi_t)$ will always contain one of ξ_{st} or ξ_{ts} , this proves part (2). \square

Corollary 4.6. *Let $s, t \in S(w)$ such that $s \neq t$. If $st \leq w$, then $A_{st} = A'_{\sigma(s)\sigma(t)}$.*

Proof. First suppose that $s, t \in S(w)$ commute. By Lemma 4.5 part (2), we have

$$\text{Supp}(\xi_s \xi_t) \cap (\text{Supp}(\xi_t^2) \cup \text{Supp}(\xi_s^2)) = \emptyset.$$

Since ϕ is a graded ring isomorphism identifying Schubert bases, we also have that

$$\text{Supp}(\phi(\xi_s)\phi(\xi_t)) \cap (\text{Supp}(\phi(\xi_t)^2) \cup \text{Supp}(\phi(\xi_s)^2)) = \emptyset.$$

This implies that $\sigma(s), \sigma(t)$ also commute and $A_{st} = A'_{\sigma(s)\sigma(t)} = 0$. If s, t do not commute, then Lemma 4.5 part (1) implies

$$\text{Supp}(\xi_s \xi_t) \cap \text{Supp}(\xi_t^2) = \{\xi_{st}\}.$$

Applying the map ϕ to this equation gives

$$\text{Supp}(\phi(\xi_s)\phi(\xi_t)) \cap \text{Supp}(\phi(\xi_t)^2) = \{\phi(\xi_{st})\}.$$

Since this set is nonempty, Lemma 4.5 part (1) also implies that $\sigma(s)\sigma(t) \leq w'$ and $\sigma(s), \sigma(t)$ do not commute. Moreover, we know that

$$\text{Supp}(\phi(\xi_s)\phi(\xi_t)) \cap \text{Supp}(\phi(\xi_t)^2) = \text{Supp}(\xi_{\sigma(s)}\xi_{\sigma(t)}) \cap \text{Supp}(\xi_{\sigma(t)}^2) = \{\xi_{\sigma(s)\sigma(t)}\},$$

so $\phi(\xi_{st}) = \xi_{\sigma(s)\sigma(t)}$. Since A_{st} is the coefficient of ξ_{st} in $-\xi_t^2$, and $A'_{\sigma(s)\sigma(t)}$ is the coefficient of $\xi_{\sigma(s)\sigma(t)}$ in $-\xi_{\sigma(t)}^2 = \phi(-\xi_t^2)$, we conclude that $A_{st} = A'_{\sigma(s)\sigma(t)}$. \square

One consequence of Lemma 2.2 and Corollary 4.6 is that if $w = s_1 \cdots s_k$ is a reduced expression, then $\sigma(s_1) \cdots \sigma(s_k) \in W(A')$ is a reduced expression Cartan equivalent to w . What remains to be shown is that $w' = \sigma(s_1) \cdots \sigma(s_k)$. In order to show this, we next show how to reconstruct the descent sets of elements in $W(A)$ from the ring structure of $H^*(X(w, A))$. For any $J \subseteq S$, let W_J denote Coxeter subgroup generated by J and let W^J denote the set of minimal length coset representatives for the cosets $W(A)/W_J$. The right descent set of $u \in W(A)$ is

$$D_R(u) := \{s \in S : \ell(us) = \ell(u) - 1\}.$$

If $u \leq w$, then $D_R(u) \subseteq S(w)$. It is well known that $s \in D_R(u)$ if and only if $u \notin W^{\{s\}}$, and that $D_R(u)$ is the set of simple reflections s for which there are reduced expressions $u = s_1 \cdots s_k$ with $s_k = s$. For any subset $J \subseteq S(w)$, let H^J be the subring of $H^*(X(w, A))$ generated by $\{\xi_s \mid s \in S(w) \setminus J\}$, and let

$$E^J := \bigcup_{F \in H^J} \text{Supp}(F)$$

be the support set of H^J in E .

Lemma 4.7. *The map $E \rightarrow [e, w]$ given by $\xi_v \mapsto v$ restricts to a bijection $E^J \rightarrow W^J \cap [e, w]$.*

Proof. We first show that $\{v \in [e, w] : \xi_v \in E^J\}$ is a subset of W^J . Suppose $u \in W^J \cap [e, w]$, $s \in S(w) \setminus J$. By Chevalley's formula, if $\xi_v \in S(\xi_s \xi_u)$ then $v = s_\beta u$ for some real root $\beta \in R^+$ with $\ell(v) = \ell(u) + 1$ and $\omega_s(u^{-1}(\beta^\vee)) \neq 0$. Suppose v is of this form. If $v \notin W^J$, then there is some simple reflection $s' \in D_R(v) \cap J$, and consequently there is a reduced expression $v = s_1 \cdots s_k$ for v with $s_k = s' \in J$. But $u = s_1 \cdots \hat{s}_\ell \cdots s_k$, where \hat{s}_ℓ means that s_ℓ is omitted from the product, and $u \in W^J$, so we must have that $\ell = k$, and ultimately $v = us'$. Hence $s_{u^{-1}\beta} = u^{-1}s_\beta u = u^{-1}v = s'$, and since the correspondence between roots and reflections is a bijection, $u^{-1}\beta = \alpha_{s'}$ and $u^{-1}\beta^\vee = \alpha_{s'}^\vee$. Since $s \notin J$, $\omega_s(u^{-1}\beta^\vee) = \omega_s(\alpha_{s'}^\vee) = 0$, so if $v \notin W^J$, then $\xi_v \notin \text{Supp}(\xi_s \xi_u)$.

Because $H^*(X(w, A))$ is graded, $\xi_s \in E^J$ for $s \in S(w)$ if and only if $s \in W^J$. We've shown that if $u \in W^J \cap [e, w]$, $s \in S(W) \setminus J$, and $\xi_v \in \text{Supp}(\xi_s \xi_u)$, then $v \in W^J$, so we conclude by induction on degree that $\{v \in [e, w] : \xi_v \in E^J\} \subset W^J$.

For the converse, we show that $\{\xi_v : v \in W_J \cap [e, w]\}$ is a subset of E^J by induction on length. Indeed, suppose that $v \in W^J \cap [e, w]$, and write $v = s_1 \cdots s_k$. Let $u := s_1 v = s_2 \cdots s_k$. Clearly $u \in W^J$, and by induction we can assume that $\xi_u \in E^J$. Since α_{s_1} is an inversion of v , we have that $\alpha_{s_1} \in R^+ \cap vR^-$. Hence $v^{-1}(\alpha_{s_1}) \in v^{-1}R^+ \cap R^-$ which

implies $v^{-1}(-\alpha_{s_1}) \in v^{-1}R^- \cap R^+$. Now [Kum02, Exercise 1.3.E] says that, since $v \in W^J$, the set $v^{-1}R^- \cap R^+ \subseteq R^+ \setminus R_J^+$, where R_J^+ is the subset of the positive roots in the span of $\{\alpha_{s'} : s' \in J\}$. Thus

$$u^{-1}(\alpha_{s_1}) = u^{-1}s_1(-\alpha_{s_1}) = v^{-1}(-\alpha_{s_1}) \in R^+ \setminus R_J^+.$$

We conclude that there exists $s \notin J$ such that $\omega_s(u^{-1}(\alpha_{s_1}^\vee)) \neq 0$. Consequently $\xi_v \in \text{Supp}(\xi_s \xi_u)$, implying $\xi_v \in E^J$. \square

We now define

$$\tilde{D}_R(\xi_u) := \{\xi_s \in \tilde{S}(w) : \xi_u \notin E^{\{s\}}\}.$$

Lemma 4.8. *For any $u \leq w$, the bijection $\tilde{S}(w) \rightarrow S(w)$ given by $\xi_s \mapsto s$ restricts to a bijection $\tilde{D}_R(\xi_u) \rightarrow D_R(u)$.*

Proof. Since $s \in D_R(u)$ if and only if $u \notin W^{\{s\}}$, the lemma follows immediately from Lemma 4.7. \square

Lemma 4.9. *For every $\xi_s \in \tilde{D}_R(\xi_v)$, there exists a unique $\xi_u \in E^{\{s\}}$ for which $\xi_v \in \text{Supp}(\xi_s \xi_u)$. Moreover, $us = v$.*

Proof. Fix $\xi_s \in \tilde{D}_R(\xi_v)$. By Lemma 4.8, $s \in D_R(v)$ and hence $u = vs \leq v$. The element u is the unique element less than v such that $\ell(v) = \ell(u) + 1$ and $u \in W^{\{s\}}$. Hence Lemma 4.4 and 4.7 imply $\xi_v \in \text{Supp}(\xi_s \xi_u)$. \square

Recall that if $w \in W(A)$, then $\text{RW}(w)$ is the set of reduced words of w . Lemma 4.9 implies we can inductively define $\widetilde{\text{RW}}(\xi_v)$ by setting $\widetilde{\text{RW}}(\xi_\epsilon) := \{\epsilon\}$ where ϵ denotes the empty sequence, and

$$\widetilde{\text{RW}}(\xi_v) := \left\{ (\xi_{s_1}, \dots, \xi_{s_m}, \xi_s) \quad : \quad \begin{array}{l} \xi_s \in \tilde{D}_R(\xi_v), (\xi_{s_1}, \dots, \xi_{s_m}) \in \widetilde{\text{RW}}(\xi_u) \text{ for } u \\ \text{the unique element such that } \xi_u \in E^{\{s\}} \text{ and} \\ \xi_v \in \text{Supp}(\xi_s \xi_u) \end{array} \right\}.$$

Lemma 4.10. *For any $v \in [e, w]$, the bijection $\tilde{S}(w) \rightarrow S(w)$ given by $\xi_s \mapsto s$ induces a bijection between $\widetilde{\text{RW}}(\xi_v)$ and $\text{RW}(v)$.*

Proof. $\text{RW}(v)$ is the set of sequences of the form (s_1, \dots, s_k) where $s_k \in D_R(v)$ and $(s_1, \dots, s_{k-1}) \in \text{RW}(vs_k)$. The lemma follows by induction on length using Lemmas 4.8 and 4.9. \square

Proof of Proposition 4.2. Suppose that $\phi : H^*(X(w, A)) \rightarrow H^*(X(w', A'))$ is a graded ring isomorphism which identifies Schubert classes. In particular, the restricted map $\phi|_{H^2(X(w, A))}$ induces a bijection $\sigma : S(w) \rightarrow S(w')$. Let $w = s_1 \cdots s_k$ be a reduced expression. Lemma 4.10 implies that $(\xi_{s_1}, \dots, \xi_{s_k}) \in \widetilde{\text{RW}}(\xi_w)$. Since ϕ is an isomorphism that identifies Schubert classes, $\phi(\text{Supp}(F)) = \text{Supp}(\phi(F))$ for any $F \in H^*(X(w, A))$. As a result, $\phi(E^J) = (E')^{\sigma(J)}$ for any $J \subset S(w)$ and hence $\phi(\tilde{D}_R(\xi_v)) = \tilde{D}_R(\phi(\xi_v))$. Also we have that $\xi_v \in \text{Supp}(\xi_s \xi_u)$ if and only if

$$\phi(\xi_v) \in \text{Supp}(\phi(\xi_s)\phi(\xi_u)) = \text{Supp}(\xi_{\sigma(s)}\phi(\xi_u)).$$

So Lemma 4.9 implies $(\xi_{\sigma(s_1)}, \dots, \xi_{\sigma(s_k)}) \in \widetilde{\text{RW}}(\phi(\xi_w))$. Since ϕ is graded, $\phi(\xi_w) = \xi_{w'}$, and hence $\sigma(s_1) \cdots \sigma(s_k) \in \text{RW}(w')$ by Lemma 4.10. Finally, Corollary 4.6 says that $A_{st} = A_{\sigma(s)\sigma(t)}$ for all $st \leq w$. Thus w and w' are Cartan equivalent. \square

Together, Propositions 3.1, 4.1, and 4.2 complete the proof of Theorem 1.3.

4.1. Constructing a presentation of a Schubert variety. In section, we describe how to construct a presentation of a Schubert variety from geometric data. Suppose we have a variety X which is isomorphic to a Schubert variety $X(w, A)$ for some $w \in W(A)$, but the element w and matrix A are unknown. We want to find a Cartan matrix A' and $w' \in W(A')$ such that $X \cong X(w', A')$. This can be done if we know the integral cohomology ring $H^*(X; \mathbb{Z})$, and the effective cone X in $H^*(X) = H^*(X; \mathbb{Z})$. Indeed, let E be the generators of the extremal rays of the effective cone. Since X is isomorphic to a Schubert variety, we know that E is the Schubert basis for $H^*(X)$. The proof of Proposition 4.2 then gives a procedure to construct w' and A' from $H^*(X)$ and E . Specifically, for any $F \in H^*(X)$, we can write $F = \sum_{\xi \in E} c_\xi \xi$ for some integers c_ξ and define

$$\text{Supp}(F) := \{\xi \in E \mid c_\xi \neq 0\}.$$

Let $\tilde{S} := E \cap H^2(X)$. We first construct a Cartan matrix A' over the set \tilde{S} using Lemma 4.5. Let $\zeta_1, \zeta_2 \in \tilde{S}$ and consider the following cases (where products denote the product in $H^*(X)$):

- (1) If $\zeta_1 = \zeta_2$, then set $A'_{\zeta_1 \zeta_2} = 2$.
- (2) If $\zeta_1 \neq \zeta_2$ and $\text{Supp}(\zeta_1 \zeta_2) \cap (\text{Supp}(\zeta_1^2) \cup \text{Supp}(\zeta_2^2)) = \emptyset$, then set $A'_{\zeta_1 \zeta_2} = A'_{\zeta_2 \zeta_1} = 0$.
- (3) If $\zeta_1 \neq \zeta_2$ and $\text{Supp}(\zeta_1 \zeta_2) \cap \text{Supp}(\zeta_2^2) = \{\nu\}$, then write $\zeta_2^2 = \sum_{\xi \in E} c_\xi \xi$, and set

$$A'_{\zeta_1 \zeta_2} = -c_\nu.$$

- (4) If $\zeta_1 \neq \zeta_2$, $\text{Supp}(\zeta_1 \zeta_2) \cap \text{Supp}(\zeta_2^2) = \emptyset$, and $\text{Supp}(\zeta_1 \zeta_2) \cap \text{Supp}(\zeta_1^2) \neq \emptyset$, then set $A'_{\zeta_1 \zeta_2}$ to any negative integer.

By Lemma 4.5, exactly one of these cases must hold for each pair $(\zeta_1, \zeta_2) \in \tilde{S}^2$. In addition, if $\xi : S \rightarrow \tilde{S} : s \mapsto \xi_s$, then $A_{st} = A'_{\xi_s \xi_t}$ for all $st \leq w$.

Next, observe that for any $\xi \in E$, we can construct $\tilde{D}_R(\xi)$ and $\widetilde{RW}(\xi)$ strictly in terms of $H^*(X)$ and E , without referring to w or A . Indeed, for any $J \subseteq \tilde{S}$ we can define H^J to be the subalgebra of $H^*(X)$ generated by J , and E^J to be the set $\bigcup_{F \in H^J} \text{Supp}(F) \subseteq E$. We can then define $\tilde{D}_R(\xi)$ and $\widetilde{RW}(\xi)$ as in the previous section, except that instead of making a distinction between ξ_s and s , we write $\zeta \in \tilde{S}$ in place of both of them (so for instance, we'd define $\tilde{D}_R(\xi) = \{\zeta \in \tilde{S} : \xi \notin E^{\{\zeta\}}\}$).

Let ξ be the unique element of E of highest degree, let $(\zeta_1, \dots, \zeta_k) \in \widetilde{RW}(\xi)$, and let $w' := \zeta_1 \cdots \zeta_k \in W(A')$. Then ξ must be ξ_w , and if we write $\zeta_i = \xi_{s_i}$, then $s_1 \cdots s_k$ is a reduced word for w by Lemma 4.10. Thus ξ is a Cartan equivalence between (w, A) and (w', A') , and so $X = X(w, A) \cong X(w', A')$.

REFERENCES

- [AB16] Hiraku Abe and Sara Billey, *Consequences of the Lakshmibai-Sandhya theorem: the ubiquity of permutation patterns in Schubert calculus and related geometry*, Schubert calculus—Osaka 2012, Adv. Stud. Pure Math., vol. 71, Math. Soc. Japan, [Tokyo], 2016, pp. 1–52. MR 3644818
- [BB05] Anders Björner and Francesco Brenti, *Combinatorics of Coxeter groups*, Graduate Texts in Mathematics, vol. 231, Springer, New York, 2005. MR 2133266
- [BL00] Sara Billey and V. Lakshmibai, *Singular loci of Schubert varieties*, Progress in Mathematics, vol. 182, Birkhäuser Boston, Inc., Boston, MA, 2000.
- [Bor53] Armand Borel, *Sur la cohomologie des espaces fibrés principaux et des espaces homogènes de groupes de Lie compacts*, Ann. of Math. (2) **57** (1953), 115–207. MR 51508
- [BS02] Nantel Bergeron and Frank Sottile, *A Pieri-type formula for isotropic flag manifolds*, Trans. Amer. Math. Soc. **354** (2002), no. 7, 2659–2705. MR 1895198

- [CC93] R. W. Carter and Y. Chen, *Automorphisms of affine Kac-Moody groups and related Chevalley groups over rings*, J. Algebra **155** (1993), no. 1, 44–94. MR 1206622
- [CM05] Pierre-Emmanuel Caprace and Bernhard Mühlherr, *Isomorphisms of Kac-Moody groups*, Invent. Math. **161** (2005), no. 2, 361–388. MR 2180452
- [Din97] Kequan Ding, *Rook placements and cellular decomposition of partition varieties*, Discrete Math. **170** (1997), no. 1-3, 107–151. MR 1452940
- [DMR07] Mike Develin, Jeremy L. Martin, and Victor Reiner, *Classification of Ding’s Schubert varieties: finer rook equivalence*, Canad. J. Math. **59** (2007), no. 1, 36–62. MR 2289417
- [EG95] Dan Edidin and William Graham, *Characteristic classes and quadric bundles*, Duke Math. J. **78** (1995), no. 2, 277–299. MR 1333501
- [FMSS95] W. Fulton, R. MacPherson, F. Sottile, and B. Sturmfels, *Intersection theory on spherical varieties*, J. Algebraic Geom. **4** (1995), no. 1, 181–193. MR 1299008
- [Ful93] William Fulton, *Introduction to toric varieties*, Annals of Mathematics Studies, vol. 131, Princeton University Press, Princeton, NJ, 1993, The William H. Roever Lectures in Geometry. MR 1234037
- [Hir84] André Hirschowitz, *Le groupe de Chow équivariant*, C. R. Acad. Sci. Paris Sér. I Math. **298** (1984), no. 5, 87–89. MR 741066
- [Kac90] Victor G. Kac, *Infinite-dimensional Lie algebras*, third ed., Cambridge University Press, Cambridge, 1990. MR 1104219
- [Kar13] Paramasamy Karuppuchamy, *On Schubert varieties*, Comm. Algebra **41** (2013), no. 4, 1365–1368. MR 3044412
- [KN98] Shrawan Kumar and Madhav V. Nori, *Positivity of the cup product in cohomology of flag varieties associated to Kac-Moody groups*, Internat. Math. Res. Notices (1998), no. 14, 757–763. MR 1637105
- [Kum87] Shrawan Kumar, *Demazure character formula in arbitrary Kac-Moody setting*, Invent. Math. **89** (1987), no. 2, 395–423. MR 894387
- [Kum02] ———, *Kac-Moody groups, their flag varieties and representation theory*, Progress in Mathematics, vol. 204, Birkhäuser Boston, Inc., Boston, MA, 2002. MR 1923198
- [KW92] V. G. Kac and S. P. Wang, *On automorphisms of Kac-Moody algebras and groups*, Adv. Math. **92** (1992), no. 2, 129–195. MR 1155464
- [Mas08] Mikiya Masuda, *Equivariant cohomology distinguishes toric manifolds*, Adv. Math. **218** (2008), no. 6, 2005–2012. MR 2431667
- [RS16] Edward Richmond and William Slofstra, *Billey-Postnikov decompositions and the fibre bundle structure of Schubert varieties*, Math. Ann. **366** (2016), no. 1-2, 31–55.

Email address: `edward.richmond@okstate.edu`

Email address: `weslofst@uwaterloo.ca`