

## INTERNAL ABSOLUTE GEOMETRY

KARSTEN BOHLEN

ABSTRACT. We model systems as objects in a certain ambient Grothendieck site with additional structure. We introduce generalized sheaves, called virtual manifolds. These sheaves reconstruct an internal groupoid which models instances of the holonomy groupoid of foliations, but in the broader framework of integrable, almost regular systems. A generalization of the Morita invariant homotopy type to the setting of internal groupoids, furnishes a generalized version of the Lusternik-Schnirelmann invariant for groupoids. This invariant, referred to herein as geometric complexity, computes the number of transitive components of the generalized holonomy groupoid, up to Morita invariant homotopy. We introduce a so-called internal absolute geometry, as a bicategory localization at the Morita homotopies and internal point groupoids, i.e. at the absolute point. We extend the geometric complexity to a well-defined bifunctor on this bicategory localization.

## 1. INTRODUCTION

In the setting of smoothly differentiable functions to any singular foliation one can associate a holonomy groupoid. The construction furnishes a smooth (Lie) groupoid whose orbits correspond precisely to the leaves of the foliation. The holonomy groupoid was studied by C. Ehresmann [9], extended to the regular foliation case by H. E. Winkelkemper [22] and to the setting of singular foliations by J. Pradines [20], as well as recently by C. Debord [7], G. Skandalis and I. Androulidakis [2]. The concept was extended to other settings by various authors. We mention the construction of a holonomy groupoid for Teichmüller and Riemann moduli spaces viewed as stacks, in connection with partially foliated structures based on infinite dimensional Fréchet manifolds in [17]. In general, if one is concerned not only with systems of a strictly smooth nature, the holonomy groupoid may still provide a useful geometric resolution of a foliation like structure. For example in the setting of cognitive systems, the geometry of information spaces is only smooth in a generalized (diffeological) sense, cf. [12]. This necessitates a synthetic approach to the holonomy groupoid and its associated geometry. The setting we want to study therefore is that of faithful subcategories of the category of diffeological spaces. A diffeological space can be viewed as a certain sheaf on the site of euclidean spaces [3]. As such it is a representable stack. It carries the  $D$ -topology generated by the plots [13]. We distill these properties in abstract terms to a suitable ambient category which in particular cases is a subcategory of diffeological spaces (though not by any means itself a "convenient" category). The ambient category consists of generalized systems each possessing a suitable generalization of the orbit structure reconstructing a holonomy groupoid. The behavior of a system should be analyzed via suitable invariants associated to a geometry generated by the candidate holonomy groupoids attached to the system. One such invariant is the Lusternik-Schnirelmann "category" invariant [16]. The invariant measures the minimal number

---

2000 *Mathematics Subject Classification.* Primary 18F40; Secondary 18E99.

*Key words and phrases.* internal absolute geometry, site, groupoid, geometric complexity.

of components that are transitive in a weak sense, sufficient to cover the space. It has properties very much in line to the general complexity functional one would like to associate to a system, cf. [15]. Recently, the invariant has been extended to the case of differentiable stacks [1] and Lie groupoids [5].

As it seems more suitable, we refer to the generalized version of the invariant henceforth as geometric complexity. The classical result of Lusternik-Schnirelmann asserts that a lower bound for the critical points of a functional on the space in question is furnished by the geometric complexity of the space. In the setting of groupoids, the invariant is the least number of point groupoids (transitive components) - up to a Morita invariant homotopy type - required to cover the groupoid. At the end of the paper we define a bicategory localization of the bicategory of internal groupoids that furnishes an absolute point geometry. Thereby, the point groupoids of candidate holonomy groupoids are viewed as a unit, up to a weak equivalence. Since the geometric complexity is invariant with regard to the 2-cells (Morita homotopies) and evaluates to a constant on the point groupoids, it extends to a well-defined bifunctor on the absolute point geometry. There is a formal similarity to the construction of operator  $KK$ -theory, where the role of the absolute point is played by the compact operators. We should mention possible connections to classical index theory. The index of an operator in the classical setting of smooth compact manifolds (with boundary) can be computed by studying the singularities of the zeta function of the Laplacian formed by the operator, see for example [11]. The geometric complexity furnishes in principle a lower bound on the critical points of a functional on the given space. In addition, singularities of the partition function of a system can in principle be controlled by suitable functionals, e.g. the Gibbs-functional in classical mechanics. Thereby a tenuous connection to index theory and operator  $KK$ -theory for systems suggests itself. It could be of interest to study these topics in greater detail. Another motivation for this study is a general property of systems, especially those with an information crosstalk structure, such as the rate distortion manifolds described formally in [12]. Such systems in general may possess a holonomy groupoid whose function is to describe the underlying path space structure, where holonomy encodes the nonlinear interactions of the components. Usually, such systems possess redundancy in that many different paths give rise to equivalent functional behavior and may therefore be identified, whenever one studies the problem of path inference. For example the redundancy of the neural code [8]. In addition, repeated simultaneous cross activation often results in a synergy and strengthening of connections which then act as a unit, as exemplified e.g. by the strong Hebb rule in neurobiology. These units are precisely the transitive components, up to a notion of homotopy. This motivates the introduction of the absolute point geometry. These issues will be studied more in depth in future research.

The overall axiomatics for the construction of the internal absolute geometry, as a bicategory localization, are listed below. The axioms constitute the underlying structure  $(\mathbb{A}, \mathcal{T}, \mathbb{E}, \mathcal{I})$ , where  $\mathbb{A}$  is the ambient category,  $\mathcal{T}$  is a Grothendieck site,  $\mathbb{E}$  is a suitable collection of admissible epimorphisms and  $\mathcal{I}$  is an interval object.

- (1) The site  $\mathcal{T}$  is subcanonical 2.4 and its covers are local 2.6.
- (2) The ambient site  $(\mathbb{A}, \mathcal{T})$  admits effective descent 3.3.
- (3) The triple  $(\mathbb{A}, \mathcal{T}, \mathbb{E})$  forms a site-structure 2.2.
- (4) The site-structure  $(\mathbb{A}, \mathcal{T}, \mathbb{E})$  is diadic 4.1.
- (5) The diadic site-structure  $(\mathbb{A}, \mathcal{T}, \mathbb{E}, \mathcal{I})$  admits filtered homotopic pullbacks 4.4.

## 2. INTERNAL GROUPOIDS

Let  $\mathbb{A} = (\mathbb{A}_0, \mathbb{A}_1)$  denote the ambient category. It is endowed with a site-structure  $(\mathcal{T}, \mathbb{E})$ , where  $\mathcal{T}$  is a subcanonical Grothendieck site and  $\mathbb{E}$  is a singleton-site consisting of admissible epimorphisms. The class  $\mathbb{E}$  consists by definition of the maximal collection of epimorphisms of  $\mathbb{A}$  that are both stable with regard to pullback and  $\mathcal{T}$ -universal, see also [21]. The purpose of this formalism is to work with groupoids internal to the site. Let us recall the meaning of these notions.

**Definition 2.1.** A *site*  $\mathcal{T}$  on an ambient category  $\mathbb{A}$  is a collection of families  $\{(U_i \rightarrow X)_{i \in I}\}_{X \in \mathbb{A}_0}$  of arrows in  $\mathbb{A}_1$  called *covers* between objects of  $\mathbb{A}_0$  called *opens*. A site verifies the following conditions:

- (1) We have that  $\{U' \xrightarrow{\sim} U\} \in \mathcal{T}$ , i.e. all *isomorphisms* are contained in  $\mathcal{T}$ .
- (2) Given  $\{U_i \rightarrow U\}_{i \in I}$  collection of covers in  $\mathcal{T}$ , then for each  $V \rightarrow U \in \mathbb{A}_1$  we have that  $\{U_i *_U V \rightarrow V\}_{i \in I}$  is in  $\mathcal{T}$ , i.e.  $\mathcal{T}$  is closed with regard to *base change*.
- (3) Given  $\{U_i \rightarrow U\}_{i \in I}$  in  $\mathcal{T}$  and  $\{U_{ij} \rightarrow U_i\}_{j \in J}$  in  $\mathcal{T}$  for each  $i \in I$ , we have that the composite  $\{U_{ij} \rightarrow U\}_{(i,j) \in I \times J}$  is also in  $\mathcal{T}$ , i.e.  $\mathcal{T}$  is closed with regard to *intersections*.

If each cover consists of a single map, the site is called *singleton site*.

The condition (3) in particular implies that  $\mathcal{T}$  is a sieve-structure, reflecting the alternative formulation of the three axioms to be found in the literature, in terms of the notion of a covering sieve. We can always consider a single map  $(f_i)_{i \in I}: \bigsqcup_{i \in I} U_i \rightarrow X$  in lieu of the collection of covers  $\{U_i \rightarrow U\}_{i \in I}$ .

**Definition 2.2.** Let  $(\mathbb{A}, \mathcal{T})$  be a site. An epimorphism  $f: S \rightarrow X$  is called  $\tau$ -*universal* if there is a covering family  $(U_i \rightarrow X)_{i \in I}$  and a lift  $U_i \rightarrow S$  for each  $i \in I$  and  $f$  is universal with this property. The dual notion is that of a  $\tau$ -*universal* monomorphism, i.e. a monomorphism  $f: S \rightarrow X$  such that there is a co-covering family  $(S \rightarrow U_i)_{i \in I}$  and an extension  $X \rightarrow U_i$  for each  $i \in I$  and  $f$  is universal with this property. The maximal class of  $\mathcal{T}$ -universal epimorphisms, such that for each epimorphism the pullback along an arbitrary arrow exists, forms a singleton site called *admissible epimorphisms* and is denoted by  $\mathbb{E}$ . The maximal class of  $\mathcal{T}$ -universal monomorphism, such that for each monomorphism the pushout along an arbitrary arrow exists, forms a singleton site called *admissible monomorphisms* and is denoted by  $\mathbb{E}^{\text{op}}$ . We call the triple  $(\mathbb{A}, \mathcal{T}, \mathbb{E})$  a *site-structure*. The triple  $(\mathbb{A}^{\text{op}}, \mathcal{T}^{\text{op}}, \mathbb{E}^{\text{op}})$  is called a *co-site-structure*.

**Notation 2.3.** Throughout, the arrow  $\rightrightarrows$  designates an admissible monomorphism and  $\rightrightarrows$  designates an admissible epimorphism.

**Definition 2.4.** A site  $(\mathbb{A}, \mathcal{T})$  fulfilling the conditions of Proposition 2.5 is called *sub-canonical*.

**Proposition 2.5** ([19], Lem. 2.2). *Let  $(\mathbb{A}, \mathcal{T})$  be a site. The following conditions are equivalent*

- (1) Each cover  $f: U \rightarrow X$  in  $\mathcal{T}$  is a coequalizer of some parallel arrows  $g_1, g_2: Z \rightrightarrows U$ .
- (2) Each cover  $f: U \rightarrow X$  in  $\mathcal{T}$  is the coequalizer of  $\text{pr}_1, \text{pr}_2: U *_f U \rightrightarrows U$ .
- (3) For each  $W \in \mathbb{A}_0$  and  $f: U \rightarrow X$  in  $\mathcal{T}$ , we have a bijection

$$\begin{aligned} \text{Mor}_{\mathbb{A}}(X, W) &\xrightarrow{\sim} \{h \in \text{Mor}_{\mathbb{A}}(U, W) : h \circ \text{pr}_1 = h \circ \text{pr}_2 \text{ in } \text{Mor}_{\mathbb{A}}(U *_f U, W)\}, \\ g &\mapsto g \circ f. \end{aligned}$$

(4) All representable functors  $\text{Mor}_{\mathbb{A}}(-, W)$  on  $\mathbb{A}$  are sheaves.

**Definition 2.6.** Let  $(\mathbb{A}, \mathcal{T})$  be a site. We say that  $f: Y \rightarrow X \in \mathbb{A}_1$  has property  $(P)$  locally if there is a cover  $g: U \rightarrow X$  such that  $\text{pr}_2: Y *_f^g U \rightarrow U$  has property  $(P)$ .

We henceforth assume that the property of being a cover in  $\mathcal{T}$  is a local property.

**Definition 2.7.** An  $\mathbb{A}$ -internal groupoid is a groupoid internal to  $(\mathbb{A}, \mathcal{T}, \mathbb{E})$  with structure

$$\mathcal{G}_1 * \mathcal{G}_1 \xrightarrow{m} \mathcal{G}_1 \xrightarrow{i} \mathcal{G}_1 \xrightarrow{r,s} \mathcal{G}_0 \xrightarrow{u} \mathcal{G}_1.$$

with  $\mathcal{G}_0, \mathcal{G}_1 \in \mathbb{A}_0$  and internal arrows such that  $r \in \mathbb{E}$ .

**Remark 2.8.** • By the structural axioms of a groupoid,  $s \in \mathbb{E} \Leftrightarrow r \in \mathbb{E}$ , since  $r \circ i = s$ . Also  $i$  is in  $\mathbb{E} \cap \mathbb{E}^{op}$ , where  $\mathbb{E}^{op}$  is  $\mathbb{E}$  in the opposite category  $\mathbb{A}^{op}$ . We have  $m \in \mathbb{E}$ ,  $u \in \mathbb{E}^{op}$ .

- By definition of  $\mathbb{E}$  we have  $\mathcal{G} * \mathcal{G} \in \mathbb{A}_0$ .
- The use of elementwise notation for groupoids and actions can be justified by the algorithm described in section 3 of [19].

**Equivalences and essential equivalences.** The *strict arrow* is defined as

$$\begin{array}{ccc} \mathcal{H}_1 & \xrightarrow{f_1} & \mathcal{G}_1 \\ \bullet \downarrow & & \bullet \downarrow \\ \mathcal{H}_0 & \xrightarrow{f_0} & \mathcal{G}_0. \end{array}$$

commutes for  $\bullet = s$  or  $r$  and

$$f(\gamma \cdot \eta) = f(\gamma) \cdot f(\eta), \quad (\gamma, \eta) \in \mathcal{H} * \mathcal{H}, \quad (1)$$

$$f(\gamma^{-1}) = f(\gamma)^{-1}, \quad \gamma \in \mathcal{H}. \quad (2)$$

We denote by  $\mathbb{A}\mathcal{G}$  the strict category of internal groupoids.

**Remark 2.9.** • If  $(\gamma, \eta) \in \mathcal{H} * \mathcal{H}$ , then (1), (2) and  $s = r \circ i$  implies  $(f(\gamma), f(\eta)) \in \mathcal{G} * \mathcal{G}$ .

- A strict isomorphism corresponds to an equivalence of groupoids, i.e.

$$\mathcal{H} \begin{array}{c} \xrightarrow{f} \\ \xleftarrow{g} \end{array} \mathcal{G}$$

such that  $gf \sim_T \text{id}_{\mathcal{H}}$ ,  $fg \sim_{\tilde{T}} \text{id}_{\mathcal{G}}$ . Given strict arrows  $f_1, f_2: \mathcal{H} \rightarrow \mathcal{G}$ , then  $f_1 \sim_T f_2 \Leftrightarrow T: \mathcal{H}_0 \rightarrow \mathcal{G}_1 \in \mathbb{A}_1$  is such that  $T(x): f_1^0(x) \rightarrow f_2^0(x)$  and for each  $\gamma: x \rightarrow y$  in  $\mathcal{H}_1$  we have  $f_2(\eta)T(x) = T(y)f_1(\eta)$ .

**Definition 2.10.** A strict arrow  $\epsilon: \mathcal{H} \rightarrow \mathcal{G}$  is an *essential equivalence*  $\Leftrightarrow \epsilon$  is essentially surjective, i.e.  $r\pi_1: \mathcal{G}_1 * \mathcal{H}_0 \rightarrow \mathcal{G}_0 \in \mathbb{E}$  and  $\epsilon$  is fully faithful, i.e.

$$\begin{array}{ccc} \mathcal{H}_1 & \xrightarrow{\epsilon} & \mathcal{G}_1 \\ s \oplus r \downarrow & & s \oplus r \downarrow \\ \mathcal{H}_0 \times \mathcal{H}_0 & \xrightarrow{\epsilon \times \epsilon} & \mathcal{G}_0 \times \mathcal{G}_0 \end{array}$$

is a pullback.

**Remark 2.11.** An equivalence of internal groupoids furnishes in particular an essential equivalence of internal groupoids.

**The bi-category of bi-bundle correspondences.** A right-action  $Z \circlearrowright \mathcal{G}$  of  $\mathcal{G} \in \mathbb{A}\mathcal{G}_0$  on  $Z \in \mathbb{A}_0$  is a map  $\alpha: Z * \mathcal{G} \rightarrow Z, (z, \gamma) \mapsto z \cdot \gamma$ , defined on the pullback  $Z * \mathcal{G}$  along the actor map  $q: Z \rightarrow \mathcal{G}_0 \in \mathbb{E}$  and  $r: \mathcal{G} \rightarrow \mathcal{G}_0$ . The action verifies the following conditions:

$$q(z\gamma) = q(z), (z, \gamma) \in Z * \mathcal{G}, \quad (3)$$

$$z \cdot (\gamma\eta) = (z\gamma) \cdot \eta, (z, \gamma) \in Z * \mathcal{G}, (\gamma, \eta) \in \mathcal{G} * \mathcal{G}. \quad (4)$$

A left-action is a right-action over  $(\mathbb{A}, \mathcal{T}, \mathbb{E})^{op} = (\mathbb{A}^{op}, \mathcal{T}^{op}, \mathbb{E}^{op})$ .

A *principal right-action* is a right-action  $Z \circlearrowright \mathcal{G}$  which is *free*, i.e.  $\{\gamma \in \mathcal{G} : z \cdot \gamma = z\} = \{\text{id}_{q(z)}\}$  for each  $z \in Z$  and the quotient map  $q_{\mathcal{G}}: Z \rightarrow Z/\mathcal{G}$  is in  $\mathbb{E}$  such that  $Z *_{Z/\mathcal{G}} Z \rightarrow \mathcal{G} \in \mathbb{A}_1$ .

**Example 2.12.** If  $\mathbb{A} = \underline{C}^\infty$  is the category of  $C^\infty$ -functions between  $C^\infty$ -manifolds and  $\mathbb{E}$  consists of the surjective submersions or if  $\mathbb{A} = \underline{C}_{\text{Fr}}^\infty$  is the category of  $C^\infty$ -maps between Fréchet manifolds, then a right action of internal groupoids being principal is equivalent to the action being free and proper, cf. [19].

**Definition 2.13.** A *bi-bundle correspondence* is given by  $\mathcal{H} \circlearrowright Z \circlearrowright \mathcal{G}$  such that the right  $\mathcal{G}$ -action is principal, the actions commute and  $Z/\mathcal{G} \cong \mathcal{H}_0$  in  $\mathbb{A}$ , via the actor map given by the  $\mathcal{H}$ -action. A bi-bundle correspondence is called *bi-bundle equivalence* if the left action is principal and  $\mathcal{H}/Z \cong \mathcal{G}_0$  via the actor map given by the  $\mathcal{G}$ -action.

Bi-bundle correspondences can be composed, i.e. we have a generalized tensor product, denoted by  $\odot$ . The resulting object is in general not an internal object of the ambient category. We state the additional condition of effective descent on the ambient site-structure and prove the bicategory property in the next section, within the broader setting of partial internal groupoids. Throughout the remainder of this section we tacitly assume the effective descent condition. Note that the condition can be slightly weakened, at least in the case for internal groupoids. Sufficient is to assume that the condition of being a cover is a local property and the associated Čech groupoid furnishes a sheaf, according to [19]. Let  $\mathcal{G} \in \mathbb{A}\mathcal{G}_0$ , then the unit arrow is given by the  $\mathcal{G}$ - $\mathcal{G}$  bi-bundle correspondence

$$\begin{array}{ccccc} \mathcal{G} & & \circlearrowright & \mathcal{G} & \circlearrowright & \mathcal{G} \\ \Downarrow & & \swarrow r & & \searrow s & \Downarrow \\ \mathcal{G}_0 & & & & & \mathcal{G}_0 \end{array}$$

Let  $\mathcal{H} \circlearrowright Z \circlearrowright \mathcal{G}$  and  $\mathcal{G} \circlearrowright \tilde{Z} \circlearrowright \mathcal{K}$  be bi-bundle correspondences, define

$$(\mathcal{H} \circlearrowright Z \circlearrowright \mathcal{G}) \odot (\mathcal{G} \circlearrowright \tilde{Z} \circlearrowright \mathcal{K}) := (\mathcal{H} \circlearrowright (Z *_{\mathcal{G}_0} \tilde{Z}) / \mathcal{G} \circlearrowright \mathcal{K}),$$

where  $Z *_{\mathcal{G}_0} \tilde{Z}$  is the pullback

$$\begin{array}{ccc} Z *_{\mathcal{G}_0} \tilde{Z} & \xrightarrow{\text{pr}_1} & Z \\ \text{pr}_2 \downarrow & & \downarrow q_Z \\ \tilde{Z} & \xrightarrow{p_{\tilde{Z}}} & \mathcal{G}_0. \end{array}$$

The right  $\mathcal{G}$ -action on  $Z *_{\mathcal{G}_0} \tilde{Z}$  is defined by

$$(z, w) \cdot \gamma = (z\gamma, \gamma^{-1}w).$$

This furnishes a  $\mathcal{H}\text{-}\mathcal{K}$  bi-bundle structure:

$$\begin{array}{ccccc} \mathcal{H} & \circlearrowleft & (Z *_{\mathcal{G}_0} \tilde{Z})/\mathcal{G} & \circlearrowright & \mathcal{K} \\ \Downarrow & & \swarrow p & & \searrow q \\ \mathcal{H}_0 & & & & \mathcal{K}_0 \end{array}$$

such that the actor maps are well-defined via  $p[z, w] = p_Z(z)$ ,  $q[z, w] = q_{\tilde{Z}}(w)$ . The left-action is given by  $\eta[z, w] = [\eta z, w]$  and the right-action by  $[z, w]\kappa = [z, w\kappa]$ . One checks that these yield well-defined actions and that the property of principality is preserved, under generalized tensor product.

**Remark 2.14.** To any bi-bundle correspondence a right and left unit, up to isomorphism, exists and is given by the self actions. To this end, note that

$$(\mathcal{H} \circlearrowleft Z \circlearrowright \mathcal{G}) \circlearrowleft (\mathcal{G} \circlearrowleft \mathcal{G} \circlearrowright \mathcal{G}) = (\mathcal{H} \circlearrowleft (Z *_{\mathcal{G}_0} \mathcal{G})/\mathcal{G} \circlearrowright \mathcal{G}) \cong (\mathcal{H} \circlearrowleft Z \circlearrowright \mathcal{G}).$$

The last isomorphism is given by  $f: (Z *_{\mathcal{G}_0} \mathcal{G})/\mathcal{G} \rightarrow Z$ ,  $f([z, x]) = z \cdot u(x)$ . By principality of the right  $\mathcal{G}$ -action, this is an isomorphism in  $\mathbb{A}_1$  which is equivariant, i.e.  $f(\eta[z, x]\gamma) = \eta f([z, x])\gamma$ . Therefore,  $f$  furnishes a bi-morphism. Likewise, we obtain a left unit  $\mathcal{H} \circlearrowleft \mathcal{H} \circlearrowright \mathcal{H}$ .

The previous considerations suggest to us a bi-category structure on internal groupoids. We denote by  $\mathbb{A}\mathcal{G}_{\text{bi}}$  the bi-category with objects  $(\mathbb{A}\mathcal{G}_{\text{bi}})_0 = \mathbb{A}\mathcal{G}_0$ , the 1-arrows given by bi-bundle correspondences and the 2-arrows given by isomorphisms of bi-bundle correspondences.

### The bi-category of bi-maps.

**Definition 2.15.** A *bi-map* is a pair of strict arrows

$$\mathcal{H} \xleftarrow{\epsilon} \mathcal{B} \xrightarrow{\varphi} \mathcal{G}$$

such that  $\epsilon$  is an essential equivalence. A bi-map is denoted by  $\langle \epsilon, \varphi \rangle$ . A *bi-equivalence* is a bi-map  $\langle \epsilon, \varphi \rangle$  such that  $\varphi$  is an essential equivalence.

The bi-maps  $\mathcal{H} \xleftarrow{\epsilon} \mathcal{B} \xrightarrow{\varphi} \mathcal{G}$ ,  $\mathcal{H} \xleftarrow{\tilde{\epsilon}} \tilde{\mathcal{B}} \xrightarrow{\tilde{\varphi}} \mathcal{G}$  are *isomorphic* if there is an internal groupoid  $\mathcal{B} \diamond \tilde{\mathcal{B}}$  and essential equivalences  $\mathcal{B} \xleftarrow{\alpha} \mathcal{B} \diamond \tilde{\mathcal{B}} \xrightarrow{\beta} \tilde{\mathcal{B}}$ . such that the following diagram commutes:

$$\begin{array}{ccc} & \mathcal{B} & \\ \epsilon \swarrow & \uparrow \alpha & \searrow \varphi \\ \mathcal{H} & \mathcal{B} \diamond \tilde{\mathcal{B}} & \mathcal{G} \\ \tilde{\epsilon} \swarrow & \downarrow \beta & \searrow \tilde{\varphi} \\ & \tilde{\mathcal{B}} & \end{array}$$

We introduce the bi-category  $\widetilde{\mathbb{A}\mathcal{G}_{\text{bi}}}$  with objects  $(\widetilde{\mathbb{A}\mathcal{G}_{\text{bi}}})_0 = \mathbb{A}\mathcal{G}_0$ , the 1-arrows given by bi-maps and the 2-arrows as isomorphisms between bi-maps.

**Theorem 2.16** ([5]). *There is a canonical isomorphism of bi-categories*

$$\Phi: \mathbb{A}\mathcal{G}_{\text{bi}} \xrightarrow{\sim} \widetilde{\mathbb{A}\mathcal{G}_{\text{bi}}}.$$

**Remark 2.17.** Let  $\mathcal{H} \xrightarrow{\varphi} \mathcal{G} \in \mathbb{A}\mathcal{G}_1$ . Then  $\mathcal{H} \xleftarrow{\text{id}} \mathcal{H} \xrightarrow{\varphi} \mathcal{G}$  is a bi-map. Likewise, we define explicitly a bi-bundle correspondence as follows. Notice

$$Z = (\mathcal{H} *_{\mathcal{H}_0}^r \mathcal{H} *_{\mathcal{G}_0}^r \mathcal{G}) / \mathcal{H} \cong \mathcal{H}_0 *_{\mathcal{G}_0}^r \mathcal{G}.$$

This furnishes the bi-bundle correspondence

$$\begin{array}{ccccc} \mathcal{H} & \circlearrowleft & \mathcal{H}_0 *_{\mathcal{G}_0}^r \mathcal{G} & \circlearrowright & \mathcal{G} \\ \Downarrow & \swarrow \text{spr}_2 & & \searrow \text{pr}_1 & \Downarrow \\ \mathcal{H}_0 & & & & \mathcal{G}_0 \end{array}$$

We obtain bi-functors of inclusion  $I_2$  and  $\tilde{I}_2$ , fitting into the commuting diagram

$$\begin{array}{ccc} \mathbb{A}\mathcal{G} & \xrightarrow{I_2} & \mathbb{A}\mathcal{G}_{\text{bi}} \\ \downarrow \tilde{I}_2 & \swarrow \cong \Phi & \\ \widetilde{\mathbb{A}\mathcal{G}_{\text{bi}}} & & \end{array}$$

### 3. VIRTUAL MANIFOLDS

We describe a reconstruction theorem for internal groupoids resulting from a generalized atlas of germs of partial Morita equivalences, adapted to a subcanonical site structure with effective descent. As a preparatory step we recall the definition of the descent functor from [14, B 1.5]. Let  $(\mathbb{A}, \mathcal{T})$  be a site and  $\mathcal{R} = (\varphi_i: U_i \rightarrow X)_{i \in I}$  a covering family in  $\mathcal{T}$ . We form the pullback

$$\begin{array}{ccc} U_{ij} & \xrightarrow{\psi_{ij}} & U_i \\ \chi_{ij} \downarrow & & \varphi_i \downarrow \\ U_j & \xrightarrow{\varphi_j} & X \end{array}$$

Define the discrete category  $\underline{X}$  by  $\underline{X}_0 = X$ ,  $\underline{X}_1 = \text{id}_X$ . Given  $A_i \in \mathbb{A}^{U_i}$ ,  $A_j \in \mathbb{A}^{U_j}$ , denote the canonical arrows  $f_{ij}: \psi_{ij}^*(A_i) \xrightarrow{\sim} \chi_{ij}^*(A_j)$  in  $\mathbb{A}^{U_{ij}}$ . We record the following *compatibility conditions* for  $f_{ij}$ :

- (a) Denote by  $\Delta_i: U_i \rightarrow U_{ii}$  the canonical diagonal arrow. Then via  $\psi_{ii}\Delta_i = \text{id}_{U_i} = \chi_{ii}\Delta_i$  the compatibility relation is

$$\Delta_i^* \psi_{ii}^*(A_i) \cong A_i \cong \Delta_i^* \chi_{ii}^*(A_i).$$

- (b) There are arrows  $\pi_{ij}: U_{ijk} \rightarrow U_{ij}$ ,  $\pi_{ik}: U_{ijk} \rightarrow U_{ik}$ ,  $\pi_{jk}: U_{ijk} \rightarrow U_{jk}$  as part of the triple pullback:

$$\begin{array}{ccccc} & & U_i & & \\ & \nearrow & & \searrow \varphi_i & \\ U_{ijk} & \longrightarrow & U_j & \xrightarrow{\varphi_j} & X \\ & \searrow & & \nearrow \varphi_k & \\ & & U_k & & \end{array}$$

such that the following diagram commutes:

$$\begin{array}{ccccc} \chi_{ij}^* \psi_{ij}^*(A_i) & \xrightarrow{\pi_{ij}^*(f_{ij})} & \pi_{ij}^* \chi_{ij}^*(A_j) & \xrightarrow{\cong} & \pi_{jk}^* \psi_{jk}^*(A_j) \\ \downarrow \cong & & & & \downarrow \pi_{jk}^*(f_{jk}) \\ \pi_{ik}^* \psi_{ik}^*(A_i) & \xrightarrow{\pi_{ik}^*(f_{ik})} & \pi_{ik}^* \chi_{ik}^*(A_k) & \xrightarrow{\cong} & \pi_{jk}^* \chi_{jk}^*(A_k) \end{array}$$

The isomorphisms in the above diagrams are canonical and arise from commuting squares in  $\mathcal{T}$ .

Define the category  $\underline{\text{Desc}}(\mathbb{A}, \mathcal{R})$  with objects  $\underline{\text{Desc}}(\mathbb{A}, \mathcal{R})_0 := \{(A_i)_{i \in I}, (f_{ij})_{(i,j) \in I \times I}\}$  such that  $(f_{ij})$  fulfills the compatibility conditions (a), (b) above. The arrows between objects  $\underline{\text{Desc}}(\mathbb{A}, \mathcal{R})_1$  are of the form  $\theta: ((A_i), (f_{ij})) \mapsto ((B_i), (g_{ij}))$  such that the following diagram commutes:

$$\begin{array}{ccc} \psi_{ij}^*(A_i) & \xrightarrow{f_{ij}} & \chi_{ij}^*(A_j) \\ \psi_{ij}^*(\theta_i) \downarrow & & \chi_{ij}^*(\theta_j) \downarrow \\ \psi_{ij}^*(B_i) & \xrightarrow{\quad} & \chi_{ij}^*(B_j). \end{array}$$

**Definition 3.1.** Given  $X \in \mathbb{A}_0$  and a covering family  $\mathcal{R}$  of  $X$  in  $\mathcal{T}$ , the *descent functor*  $\Psi_{\mathcal{R}}: \mathbb{A}^X \rightarrow \underline{\text{Desc}}(\mathbb{A}, \mathcal{R})$  is defined as

$$\begin{aligned} (\mathbb{A}^X)_0 \ni A &\mapsto (\varphi_i^*(A), f_{ij}) \in \underline{\text{Desc}}(\mathbb{A}, \mathcal{R})_0, \\ (\mathbb{A}^X)_1 \ni h: A \rightarrow B &\mapsto (\varphi_i^*(h) : i \in I) \in \underline{\text{Desc}}(\mathbb{A}, \mathcal{R})_1. \end{aligned}$$

**Remark 3.2.** Conditions (a) and (b) entail that  $f_{ij}$  are isomorphisms.

**Definition 3.3.** A site  $(\mathbb{A}, \mathcal{T})$  fulfills *effective descent* if for each covering family  $\mathcal{R}$  the descent functor  $\Psi_{\mathcal{R}}$  induces an equivalence of categories  $\mathbb{A}^X \leftrightarrow \underline{\text{Desc}}(\mathbb{A}, \mathcal{R})$ .

Groupoids that are of holonomy type, are generally reconstructed from "local" data, in this case defined by partial Morita equivalence classes that glue properly. We describe the type of sheaf whose global objects are precisely groupoids of holonomy type. To this end we introduce partial groupoids, i.e. internal groupoids whose structural maps are defined only partially.

**Definition 3.4.** A *partial groupoid* is given by the data

$$\begin{array}{ccccc} & & \mathcal{D}_m & & \\ & & \text{pr}_1 \downarrow & \searrow & \\ \mathcal{D}_i & \xrightarrow{i} & \mathcal{L} & \xrightarrow{r} & \mathcal{L}_0 \\ & & \uparrow m & \nearrow & \\ & & \mathcal{L}_1 * \mathcal{L}_1 \supseteq \mathcal{D}_m & & \end{array}$$

and  $\mathcal{L}_0 \rightrightarrows \mathcal{D}_i, \mathcal{D}_r \rightrightarrows \mathcal{L}_1$  are admissible monomorphisms. Likewise  $\Delta_{\mathcal{L}_0} \rightrightarrows \mathcal{D}_m \rightrightarrows \mathcal{L}_1 * \mathcal{L}_1$  are admissible monomorphisms. The following conditions hold:

- (1)  $(s \circ u)|_{\mathcal{L}_0} = (r \circ u)|_{\mathcal{D}_r} = \text{id}_{\mathcal{D}_r}$ ,
- (2)  $m \circ (m \times \text{id}_{\mathcal{L}}) = m \circ (\text{id}_{\mathcal{L}} \times m)$ ,
- (3)  $r \circ m = r \circ \text{pr}_1, s \circ m = s \circ \text{pr}_2$ ,
- (4)  $m(m(-), -) = m(-, m(-))$ ,
- (5)  $r \circ i = s$ ,
- (6)  $m(i(a), a) = \text{id}_{s(a)}, m(a, i(a)) = \text{id}_{r(a)}, a \in \mathcal{D}_i$ .

**Definition 3.5.** Given a site structure  $(\mathbb{A}, \mathcal{T}, \mathbb{E})$ . A *partial right action* is given by the data

$$\begin{array}{ccc} Z & \circlearrowright & \mathcal{L} \\ & \searrow q & \Downarrow \\ & & \mathcal{L}_0 \end{array}$$

where  $\alpha: \mathcal{D}_\alpha \rightarrow Z$ ,  $\alpha(z, \gamma) = z \cdot \gamma$  is internal and  $\mathcal{L}_0 \rightarrow \mathcal{D}_\alpha \rightarrow Z *_q^r \mathcal{L}$  are admissible monomorphisms and the following conditions hold:

- (1)  $q \circ \alpha = q \circ \text{pr}_1$ ,  $(z, \gamma) \in \mathcal{D}_\alpha$ ,
- (2)  $\alpha(z, m(\gamma, \eta)) = m(\alpha(z, \gamma), \eta)$ ,  $(z, \gamma) \in \mathcal{D}_\alpha$ ,  $(\gamma, \eta) \in \mathcal{D}_m$ .

The action is called *locally free* if  $z\gamma = z$  for each  $z$  implies that  $\gamma \in \mathcal{L}$ . It is called *locally proper* if the quotient arrow  $\pi: Z \rightarrow Z/\mathcal{L} \in \mathbb{E}$  and  $Z *_Z/\mathcal{L} Z \rightarrow \mathcal{L} \in \mathbb{A}_1$ .

Consider the situation of two partial bibundle correspondences:

$$\begin{array}{ccccccc} \mathcal{L}_1 & \circlearrowright & Z_1 & \circlearrowright & \mathcal{L} & \circlearrowright & Z_2 & \circlearrowright & \mathcal{L}_2 \\ \Downarrow & \swarrow p_1 & \searrow q_1 & \Downarrow & \swarrow p_2 & \searrow q_2 & \Downarrow \\ \mathcal{L}_1^{(0)} & & & \mathcal{L}_0 & & & \mathcal{L}_2^{(0)} \end{array}$$

i.e. partial right principal bibundles such that  $p_1, p_2 \in \mathbb{E}$ .

**Proposition 3.6.** Let  $(\mathbb{A}, \mathcal{T}, \mathbb{E})$  be a subcanonical site structure that admits effective descent. Denote by  $\mathcal{L}_1 \circlearrowright Z_1 \circlearrowright \mathcal{L}$  and  $\mathcal{L} \circlearrowright Z_2 \circlearrowright \mathcal{L}_2$  partial bibundle correspondences. Then the generalized tensor product  $Z_1 \circlearrowright Z_2$  is an  $\mathbb{A}$ -object.

*Proof.* Since  $p_2 \in \mathbb{E}$  the pullback  $Z_1 * Z_2$  is internal. Define the action  $\Psi$  of  $\mathcal{L}$  on  $Z_1 * Z_2$  via  $\Psi(z_1, z_2, \gamma) = (z_1\gamma, \gamma^{-1}z_2)$ ,  $(z_1, z_2) \in Z_1 * Z_2$ . There is an arrow  $\pi: Z_1 * Z_2 \rightarrow \mathcal{L}_0$ ,  $(z_1, z_2) \mapsto q_1(z_1) = p_2(z_2)$ . The action is locally free. Denote the canonical quotient by  $Q: Z_1 * Z_2 \rightarrow Z_1 * Z_2/\mathcal{L}$ , defined via the equivalence relation  $(z_1, z_2) \sim (z'_1, z'_2) \Leftrightarrow (z_1, z_2) = (z'_1\gamma, \gamma^{-1}z'_2)$ . The arrow  $(z_1, z_2) \mapsto (z_1\gamma, \gamma^{-1}z_2)$  induces an isomorphism of  $Z_1 * Z_2$ . Since  $p_1: Z_1 \rightarrow \mathcal{L}_1^{(0)} =: U_1$  is an admissible epimorphism, we fix  $(\varphi_i: U_i \rightarrow U_1)_{i \in I} \in \mathcal{T}^{U_1}$ ,  $s_i: U_i \rightarrow Z_1 \in \mathbb{A}_1$  such that  $p_1 \circ s_i = \varphi_i$ ,  $i \in I$ . We fix the isomorphism of "undoing the  $\mathcal{L}$ -action on  $Z_1$ "  $\delta: Z_1 *_U Z_1 \xrightarrow{\sim} Z_1 *_\mathcal{L}_0^{q_1, r} \mathcal{L}$  whose inverse is  $(z_1, \gamma) \mapsto (z_1, z_1\gamma)$ , by local properness. The diagram

$$\begin{array}{ccc} Z_1 *_U Z_1 & \xrightarrow[\simeq]{\delta} & Z_1 *_\mathcal{L}_0^{q_1, r} \mathcal{L} \\ q_1 \circ \text{pr}_1 \downarrow & \circlearrowright & \downarrow \text{pr}_2 \\ \mathcal{L}_0 & \xleftarrow[r]{} & \mathcal{L} \end{array}$$

commutes. Furthermore, the relations  $(\text{pr}_2 \circ \delta)(z_1, z'_1 \cdot \gamma) = (\text{pr}_2 \circ \delta)(z_1, z'_1) \cdot \gamma$ ,  $z_1 \cdot (\text{pr}_2 \circ \delta)(z_1, z'_1)^{-1} = z'_1$  and  $(\text{pr}_2 \circ \delta)(z, z) = i(q_1(z))$  hold. We construct the arrow  $\Phi_i: U_i \circlearrowright Z_2 \rightarrow U_i *_\mathcal{L}_0^{q_1 \circ s_i, p_2} Z_2$  defined by  $[(z_1, z_2)] \mapsto (z_1, (\text{pr}_2 \circ \delta \circ s_i)(z_1, z_2) \cdot z_2)$ . It is well-defined by the relations above:

$$\begin{aligned} p_2((\text{pr}_2 \circ \delta \circ s_i)(z_1, z_1) \cdot z_2) &= (r \circ \text{pr}_2 \circ \delta \circ s_i)(z_1, z_1) \\ &= (q_1 \circ s_i \circ \text{pr}_1)(z_1, z_1) = (q_1 \circ s_i)(z_1) \end{aligned}$$

and it can be similarly checked that  $\delta$  is compatible with the  $\mathcal{L}$ -action.

$$\begin{array}{ccccc}
& & Z_1 & \xrightarrow{q_1} & \mathcal{L}_0 \\
& \nearrow s_i & \downarrow p_1 & & \uparrow \\
U_i & \xrightarrow{\varphi_i} & U_1 & & Z_2 \\
& \nwarrow \text{pr}_1 & & \nearrow \text{pr}_2 & \\
& & U_i *_{\mathcal{L}_0}^{q_1 \circ s_i, p_2} Z_2 & \longrightarrow & Z_1 * Z_2 \\
& & \uparrow \Phi_i \simeq & & \downarrow Q \\
& & U_i \odot Z_2 & \longrightarrow & Z_1 \odot Z_2
\end{array}$$

The descent arrows  $\Phi_{ij}: U_{ij} *_{\mathcal{L}_0}^{q_1 \circ s_j, p_2} Z_2 \rightarrow U_{ij} *_{\mathcal{L}_0}^{q_1 \circ s_i, p_2} Z_2$  are defined via  $(z_1, z_2) \mapsto (z_1, (\text{pr}_2 \circ \delta)(s_i(z_1), s_j(z_2)) \cdot z_2)$ . This furnishes the descent datum

$$((U_{ij} *_{\mathcal{L}_0}^{q_1 \circ s_i, p_2} Z_2, U_{ij} *_{\mathcal{L}_0}^{q_1 \circ s_j, p_2} Z_2, \Phi_{ij})_{(i,j) \in I \times I}) \in \underline{\text{Desc}}(\mathbb{A}, \mathcal{R}).$$

By effective descent, this data reconstructs the object  $Z_1 \odot Z_2$ . The representation is unique, i.e. independent of the initial choice of covering family up to internal isomorphism, by subcanonicity. The arrow  $p_1$  has a local representation  $\text{pr}_1: U_i *_{\mathcal{L}_0}^{q_1 \circ s_i, p_2} Z_2 \rightarrow U_i$  by  $[(z_1, z_2)] \mapsto p_1(z_1)$  which makes  $p_1$  locally liftable. Therefore  $p_1$  is an admissible epimorphism. Likewise, the arrow  $\tilde{\text{pr}}_1: U_i *_{\mathcal{L}_0} \mathcal{L} *_{\mathcal{L}_0} Z_2 \rightarrow U_i *_{\mathcal{L}_0} Z_2$  is a local representation of  $Q$ . Therefore  $Q$  is an admissible epimorphism. It follows that  $Z_1 \odot Z_2$  is an internal object.  $\square$

**Proposition 3.7.** *Let  $(\mathbb{A}, \mathcal{T}, \mathbb{E})$  be a subcanonical site structure that admits effective descent. Then, the generalized tensor product  $\odot$  is compatible with partial Morita equivalence, i.e.  $\mathcal{L}_0$  partially Morita equivalent to  $\mathcal{L}_1$  if and only if there is a right-principal partial  $\mathcal{L}_0$ - $\mathcal{L}_1$  bibundle  $Z$  and a right-principal partial  $\mathcal{L}_1$ - $\mathcal{L}_0$  bibundle  $Z^{-1}$  such that*

$$\mathcal{L}_1 \odot Z^{-1} \odot Z \odot \mathcal{L}_1 \cong \mathcal{L}_1 \odot \mathcal{L}_1 \odot \mathcal{L}_1, \quad (5)$$

$$\mathcal{L}_0 \odot Z \odot Z^{-1} \odot \mathcal{L}_0 \cong \mathcal{L}_0 \odot \mathcal{L}_0 \odot \mathcal{L}_0. \quad (6)$$


*Proof.*  $(\Leftarrow)$ : (5) implies that  $q_Z: \mathcal{L}_1 \rightarrow (\mathcal{L}_1)_0$  is an admissible epimorphism, since  $s: \mathcal{L}_1 \rightarrow (\mathcal{L}_1)_0$  is an admissible epimorphism. By (6) the  $\mathcal{L}_0$ -action on  $Z$  is free and transitive on the  $q_Z$ -fibers.

$(\Rightarrow)$ : We prove (5). Note that  $Z/\mathcal{L}_0 \cong (\mathcal{L}_1)_0$  implies that  $Z * Z/\mathcal{L}_0 \cong \mathcal{L}_1$  is a natural isomorphism which is furnished by the fact that  $[x, y] \in Z * Z/\mathcal{L}_0$ ,  $q(x) = q(y)$  implies there is a unique  $\gamma \in \mathcal{L}_1$  such that  $x\gamma = y$ . The proof of (6) proceeds along the same lines.  $\square$

Given a site  $\mathcal{T}$ , let us fix the optic bi-category  $\mathbb{A}\mathcal{S}(\mathcal{T})_{\text{bi}}$  with the objects given by  $\mathbb{A}_0$ , the 1-arrows given by spans from singleton  $\mathcal{T}$ -covers and 2-cells defined by span-arrows of the form:

$$\begin{array}{ccc}
& S_1 & \\
\swarrow & & \searrow \\
A & & B \\
\swarrow & & \searrow \\
& S_2 &
\end{array}$$

Composition is defined via pullbacks, they exist up to isomorphism:



**Definition 3.8.** Let  $A \leftarrow S \rightarrow B$  be a span where the arrows are singleton covers from  $\mathcal{T}$ . A *span-cover* is given by the data  $(A \leftarrow \tilde{S} \rightarrow B, \varphi)$  where  $\varphi: \tilde{S} \rightarrow S$  is a singleton cover which forms a 2-cell in the category  $\mathbb{A}\mathcal{S}(\mathcal{T})_{\text{bi}}$ .

It is straightforward to check the axioms of a Grothendieck site for the collection of span-covers.

**Proposition 3.9.** *Given a site  $(\mathbb{A}, \mathcal{T})$ , then  $\widehat{\mathcal{T}}$  the span-covers formed out of  $\mathcal{T}$  furnish a singleton site on the category consisting of spans as objects and 2-cells as arrows.*

**Definition 3.10.** A *spanoid*  $\mathcal{L}$  is a partial groupoid that is final in the 1-arrows of  $\mathbb{A}\mathcal{S}(\mathcal{T})_{\text{bi}}$ , i.e. there exists a span  $S$  and a unique span-arrow  $\mathcal{L} \rightarrow S$ .

Denote by  $\mathbb{A}\mathcal{P}\mathcal{S}$  the category with sets of partial Morita equivalences between spanoids, associated to coverings, as objects and sets of span 2-cells as arrows between objects, i.e.  $\mathbb{A}\mathcal{P}\mathcal{S}_1 = \mathbb{A}\mathcal{S}(\mathcal{T})_{\text{bi},2}$ . By a virtual manifold over a site structure we mean a generalized presheaf taking values in  $\mathbb{A}\mathcal{P}\mathcal{S}$ , but viewed at first as a Set-valued functor, that fulfills effective descent *horizontally*, i.e. with regard to span-arrows and that fulfills the sheaf-condition *vertically*, i.e. with regard to partial Morita equivalences. This construction is derived from the geometry of foliations where the orbit structure of the symmetries of the system corresponds to the leaves of the foliation, and where (co-)dimensionality within leaves is preserved by Morita equivalence, cf. [2], [7], [20].

**Definition 3.11.** A *generalized presheaf* is a functor  $\mathcal{M}: \mathbb{A}^{op} \rightarrow \mathbb{A}\mathcal{P}\mathcal{S}$  such that  $\mathcal{M}$  fulfills horizontal effective descent with regard to span isomorphisms. Let  $\{\varphi_i: U_i \rightarrow U\}_{i \in I}$  denote a cover in  $\mathcal{T}$ . The following conditions characterize this effective descent:

- (1) For each  $W_{U_i}, Z_{U_i}$  representatives of objects in  $\mathcal{M}(U)_0$  and  $\psi_i: W_{U_i} \rightarrow Z_{U_i}$  span arrows, there is a span arrow  $\psi: W_U \rightarrow Z_U$  such that  $\psi|_{Z_{U_i}} = \psi_i$ .
- (2) For each span arrows  $\psi, \chi: W_U \rightarrow Z_U \in \mathcal{M}(U)_1$  such that  $\psi|_{U_i} = \chi|_{U_i}$  for each  $i$  we have  $\psi = \chi$ .
- (3) Let  $\{U_{ij} := U_i * U_j \rightarrow U_i\}_{(i,j) \in I \times J}$  be the subcover, let  $Z_i \in \mathcal{M}(U)_0$  and  $\varphi_{ij}: Z_{i|ij} \rightarrow Z_{i|ij}$  satisfying the cocycle condition  $\varphi_{ij|ijk} \circ \varphi_{jk|ijk} = \varphi_{ik|ijk}$ . Then there exists  $Z \in \mathcal{M}(U)_0$  and span isomorphisms  $\varphi_i: Z|_i \xrightarrow{\sim} Z_i$  such that  $\varphi_{ij} \circ \varphi_{i|ij} = \varphi_{j|ij}$ .

*Notation:* The category of generalized presheaves with natural transformations as arrows is denoted by  $\underline{\mathcal{P}\mathcal{V}}_{\widehat{\mathcal{T}}}(\mathbb{A})$ .

We record some easily verified properties of span covers and germs of partial Morita equivalences.

**Lemma 3.12.** 1) *Partial Morita equivalences along span isomorphisms have equal germs: Let  $Z_f, Z_g \in \mathbb{A}\mathcal{P}\mathcal{S}_0$  representatives of partial Morita equivalences  $f, g$ , then there are covers  $V_f \rightarrow Z_f, V_g \rightarrow Z_g$  in  $\widehat{\mathcal{T}}$  such that  $f|_{\mathcal{H}_0, V_f, \mathcal{H}_1} \equiv g|_{\mathcal{H}_0, V_g, \mathcal{H}_1}$ .*

- 2) *Germ replacement:* Let  $f$  denote partial Morita equivalence  $\mathcal{L}_1 \sim \mathcal{L}_0$  and  $g$  the partial Morita equivalence  $\mathcal{L}'_1 \sim \mathcal{L}'_0$  between spanoids. Then there is a maximal spanoid  $\tilde{\mathcal{L}}_i \rightrightarrows O_i * O'_i$  such that the restrictions agree  $f|_{\tilde{\mathcal{L}}_1, Z_f, \tilde{\mathcal{L}}_0} \equiv g|_{\tilde{\mathcal{L}}_1, Z_f, \tilde{\mathcal{L}}_0}$ .
- 3) A partial Morita equivalence has a final span cover.

Via a generalized Yoneda embedding  $\hat{j}: \mathbb{A} \hookrightarrow \mathcal{PV}_{\hat{\mathcal{T}}}(\mathbb{A})$  we construct the sheafification of generalized presheaves. We obtain the category  $\mathcal{V}_{\hat{\mathcal{T}}}(\mathbb{A})$  of so-called virtual manifolds and an adjunction  $L \dashv i$ , where  $L$  is the sheafification functor and  $i$  is the inclusion functor  $\mathcal{V}_{\hat{\mathcal{T}}}(\mathbb{A}) \hookrightarrow \mathcal{PV}_{\hat{\mathcal{T}}}(\mathbb{A})$ . The Yoneda embedding for the site  $(\mathbb{AP}\mathcal{S}, \hat{\mathcal{T}})$  is obtained via  $h_Z: \mathbb{AP}\mathcal{S} \rightarrow \text{Set}$ ,  $U \mapsto \text{Mor}_{\mathbb{AP}\mathcal{S}}(U, Z)$ . A 2-cell  $f: Z \rightarrow W$  induces a natural transformation  $h_f: h_Z \rightarrow h_W$  by the associativity of 2-cells. This furnishes the Yoneda embedding  $\hat{j}: \mathbb{AP}\mathcal{S} \rightarrow [\mathbb{AP}\mathcal{S}^{op}, \text{Set}]$ . By Lemma 3.12 1)-2) we can adapt the sheafification in the following fashion. Setting

$$\widehat{W} := \left\{ \widehat{S}(\{Z_i\}) := \varprojlim_{i,j} \left( \prod_{i,j} \hat{j}(Z_i) *_{\hat{j}(Z)} \hat{j}(Z_j) \rightrightarrows \prod_i \hat{j}(Z_i) \right) \rightarrow \hat{j}(Z) : \{Z_i \rightarrow Z\}_{i \in I} \in \hat{\mathcal{T}} \right\}.$$

Define the sheaf  $\mathcal{L}(\mathcal{M})$  via

$$L(\mathcal{M})(U) := \varprojlim_{w: \widehat{S}(Z_U) \rightarrow \hat{j}(Z_U) \in \widehat{W}} \mathcal{PV}_{\hat{\mathcal{T}}}(\mathbb{A})(\widehat{S}(Z_U), \mathcal{M}), \quad U \in \mathbb{A}_0, \quad \mathcal{M} \in \mathcal{PV}_{\hat{\mathcal{T}}}(\mathbb{A})_0.$$

We call  $\mathcal{L}(\mathcal{M})$  a *generalized sheaf*.

**Definition 3.13.** A *generalized sheaf*  $\widehat{\mathcal{M}}$  such that the inclusions of equivalence classes of germs of partial Morita equivalences are admissible monomorphisms is called a *virtual manifold*. The category of virtual manifolds with natural transformations as arrows is denoted by  $\mathcal{V}_{\hat{\mathcal{T}}}(\mathbb{A})$ .

We have thus constructed the localization:

$$\mathcal{PV}_{\hat{\mathcal{T}}}(\mathbb{A})[\widehat{W}^{-1}](\mathcal{M}, \widehat{\mathcal{M}}) := \varprojlim_{\widehat{S}(\mathcal{M}) \rightarrow \mathcal{M} \in \widehat{W}} \mathcal{PV}_{\hat{\mathcal{T}}}(\mathbb{A})(\widehat{S}(\mathcal{M}), \widehat{\mathcal{M}})$$

for  $\mathcal{M} \in \mathcal{PV}_{\hat{\mathcal{T}}}(\mathbb{A})_0$ ,  $\widehat{\mathcal{M}} \in \mathcal{V}_{\hat{\mathcal{T}}}(\mathbb{A})_0$ .

**Proposition 3.14.** *The global objects of virtual manifolds over  $(\mathbb{A}, \mathcal{T}, \mathbb{E})$  are internal groupoids.*

*Proof.* Let  $X \in \mathbb{A}_0$  and denote by  $\mathcal{G} = \widehat{\mathcal{M}}(X)_0$  the global object of  $\widehat{\mathcal{M}} \in \mathcal{V}_{\hat{\mathcal{T}}}(\mathbb{A})$ . As a set  $\mathcal{G}$  consists of equivalence classes of germs of partial Morita equivalences. The structure maps of  $\mathcal{G}$  are defined as follows. The unit inclusion  $u: X \rightarrow \mathcal{G}$  which maps to the identities of the corresponding maximal partial spanoids. The source  $s: \mathcal{G} \rightarrow X$  maps  $[\mathcal{L}_1 \circ Z_f \circ \mathcal{L}_2]_z \mapsto q_f(z)$  and  $r: \mathcal{G} \rightarrow X$  maps  $[\mathcal{L}_1 \circ Z_f \circ \mathcal{L}_2]_z \mapsto p_f(z)$ . The composition is defined by

$$[\mathcal{L}_2 \circ Z_g \circ \mathcal{L}_1]_t \cdot [\mathcal{L}_1 \circ Z_f \circ \mathcal{L}_0]_z = [\mathcal{L}_2 \circ Z_g \circ Z_f \circ \mathcal{L}_0]_{(t,z)}.$$

By Proposition 3.7 the inverse can be defined as

$$i: \mathcal{G} \rightarrow \mathcal{G}, \quad [\mathcal{L}_1 \circ Z_f \circ \mathcal{L}_0]^{-1} = [\mathcal{L}_0 \circ Z_f^{-1} \circ \mathcal{L}_1]$$

and this furnishes the groupoid structure on  $\mathcal{G}$ . By Lemma 3.12, 3) the section functor is internal, hence  $\mathcal{G}$  is an internal span. From the definition of a virtual manifold, the inclusions of the fibers aka sections of the sheaf are admissible monomorphisms. Hence, the structural maps of  $\mathcal{G}$  are internal as well.  $\square$

## 4. MORITA HOMOTOPY TYPE

A homotopy type for internal groupoids that is invariant with regard to Morita equivalences is described. We fix a site-structure  $(\mathbb{A}, \mathcal{T}, \mathbb{E})$ . Let  $\mathcal{I}$  denote the groupoid on two objects, i.e. the interval object. We seek to endow the site structure with an additional *diad* structure, cf. [10].

**Definition 4.1.** We say that  $\mathbb{A}\mathcal{G}_2$  admits a *cocylinder* if the groupoid  $\mathcal{I}$  exists, and the cocylinder is given by coproducts with  $\mathcal{I}$ , i.e. by the data  $((\cdot)^{\mathcal{I}} : \mathbb{A}\mathcal{G} \rightarrow \mathbb{A}\mathcal{G}, e_0, e_1 : (\cdot)^{\mathcal{I}} \rightarrow \text{id}_{\mathbb{A}\mathcal{G}}, t : \text{id}_{\mathbb{A}\mathcal{G}} \rightarrow (\cdot)^{\mathcal{I}})$  where  $e_0, e_1$  and  $t$  are natural transformations such that  $e_0 t = e_1 t = \text{id}_{\mathbb{A}\mathcal{G}}$ . In this case, we refer to  $(\mathbb{A}, \mathcal{T}, \mathbb{E}, \mathcal{I})$  as a *diadic site structure*.

We assume that all site structures considered admit a cocylinder.

**Definition 4.2.** Let  $(\mathbb{A}, \mathcal{T}, \mathbb{E}, \mathcal{I})$  be a diadic site structure. Given  $\varphi, \psi : \mathcal{G} \rightarrow \mathcal{H}$  strict arrows in  $\mathbb{A}\mathcal{G}$ . Then  $\varphi, \psi$  are *right-homotopic* if there is a strict arrow  $H : \mathcal{G} \rightarrow \mathcal{H}^{\mathcal{I}}$  such that  $e_0(\mathcal{H}) \circ H = \varphi$ ,  $e_1(\mathcal{H}) \circ H = \psi$ . Notation:  $\varphi \sim_h \psi$ .

To any co-span of internal groupoids we define an associated homotopy pullback of the  $n$ -th degree. If these homotopy pullbacks fulfill a certain vertical filtration condition we meet the requirements to generalize the construction of the Morita invariant homotopy type constructed for the case of Lie groupoids in [5].

Denote by  $\mathcal{P}_1 := \mathcal{P}_1(\mathcal{J} \xrightarrow{\psi} \mathcal{G} \xleftarrow{\varphi} \mathcal{K})$  the homotopic pullback of co-span internal to  $\mathbb{A}\mathcal{G}$ , i.e. the diagram

$$\begin{array}{ccc} \mathcal{P}_1 & \xrightarrow{\text{pr}_1} & \mathcal{K} \\ \downarrow \text{pr}_2 & & \downarrow \varphi \\ \mathcal{J} & \xrightarrow{\psi} & \mathcal{G} \end{array}$$

where

$$\begin{aligned} \mathcal{P}_{1,0} &:= \{(x, \sigma, y) \in \mathcal{K}_0 \times \mathcal{G}^{\mathcal{I}} \times \mathcal{J}_0 : e_0(\mathcal{G}) \circ \sigma = \varphi(x), e_1(\mathcal{G}) \circ \sigma = \psi(y)\}, \\ \mathcal{P}_{1,1} &:= \{(\kappa, \sigma, \iota) \in \mathcal{K}_1 \times \mathcal{G}^{\mathcal{I}} \times \mathcal{J}_1 : e_0(\mathcal{G}) \circ \sigma = \varphi(s(\kappa)), e_1(\mathcal{G}) \circ \sigma = \psi(r(\iota))\}. \end{aligned}$$

**Remark 4.3.** If  $\varphi$  is an admissible epimorphism, then  $\mathcal{P}_1$  is an internal object and up to homotopy an internal pullback.

The definition of  $\mathcal{P}_{1,1}$  is rewritten in terms of ordinary pullbacks

$$\begin{aligned} \mathcal{P}_{1,1} &= (\mathcal{K}_1 *_{\mathcal{G}_0} \mathcal{G}_1) *^h (\mathcal{J}_1 *_{\mathcal{G}_0} \mathcal{G}_1) \\ &= \mathcal{K}_1 *_{\mathcal{G}_0} \mathcal{G}_1 *^h \mathcal{G}_1 *_{\mathcal{G}_0} \mathcal{J}_1. \end{aligned}$$

Set  $\mathcal{K}_0 *^h \mathcal{G}^{n-1}$  to denote the  $n$ -th homotopy pullback, i.e.  $\mathcal{K}_0 *^h \mathcal{G}_1 *^h \dots *^h \mathcal{G}_1$  iterating the above construction  $n - 1$  times. Setting

$$\begin{aligned} \mathcal{P}_n(\mathcal{K} \xrightarrow{\varphi} \mathcal{G} \xleftarrow{\psi} \mathcal{J})_0 &:= \mathcal{K}_0 *_{\mathcal{G}_0}^h \mathcal{G}^{n-1} *_{\mathcal{G}_0}^h \mathcal{J}_0, \\ \mathcal{P}_n(\mathcal{K} \xrightarrow{\varphi} \mathcal{G} \xleftarrow{\psi} \mathcal{J})_1 &:= \mathcal{K}_1 *_{\mathcal{G}_0}^h \mathcal{G}^{n-1} *_{\mathcal{G}_0}^h \mathcal{J}_1, \end{aligned}$$

we obtain the *n-th homotopy pullback*. The construction is universal up to a stronger version of homotopy. We denote the underlying equivalence relation by  $\sim_n$  and call two strict arrows  $\varphi, \psi$  *strongly homotopic* if there is an  $n \in \mathbb{N}$  such that  $\varphi \sim_n \psi$ . We note that  $\sim_h \equiv \sim_1$ . The vertical composition of homotopic pullbacks is denoted by  $\square$ .

$$\begin{array}{ccc}
\mathcal{P}_n & \xrightarrow{\text{pr}_1} & \mathcal{K} \\
\downarrow \text{pr}_2 & \cong & \downarrow \varphi \\
\mathcal{J} & \xrightarrow{\psi} & \mathcal{G}
\end{array}$$

**Definition 4.4.** The diadic site structure  $(\mathbb{A}, \mathcal{T}, \mathbb{E}, \mathcal{I})$  admits filtered homotopic pullbacks if

$$\mathcal{P}_n(\text{cospan}_1) \square \mathcal{P}_n(\text{cospan}_2) \cong \mathcal{P}_{n+m}(\text{cospan}_1 \cdot \text{cospan}_2)$$

whenever the co-spans  $\text{cospan}_1$  and  $\text{cospan}_2$  are composable over  $\mathbb{A}\mathcal{G}$ .

**Remark 4.5.** The condition can be shown to hold whenever a subdivision structure is present, allowing refinements.

**Definition 4.6.** A strict arrow  $\eta: \mathcal{K} \rightarrow \mathcal{G}$  between internal groupoids is an *essential homotopy equivalence* if there is a strong homotopy equivalence  $h: \mathcal{K} \rightarrow \mathcal{L}$  and an essential equivalence  $\epsilon: \mathcal{L} \rightarrow \mathcal{G}$  such that the following diagram commutes up to strong homotopy:

$$\begin{array}{ccc}
\mathcal{K} & \xrightarrow{\eta} & \mathcal{G} \\
\downarrow h & \nearrow \epsilon & \\
\mathcal{L} & & 
\end{array}$$

**Remark 4.7.** If the groupoids are represented by virtual manifolds  $\widehat{\mathcal{M}}_{\mathcal{G}}$ ,  $\widehat{\mathcal{M}}_{\mathcal{K}}$ , then  $\eta$  induces homotopy equivalences between local sections of  $\widehat{\mathcal{M}}_{\mathcal{G}}$  and  $\widehat{\mathcal{M}}_{\mathcal{K}}$ . In addition, we note that strong homotopy equivalences and essential equivalences are particular instances of essential homotopy equivalences

**Definition 4.8.** An internal groupoid  $\mathcal{K}$  is *Morita homotopy equivalent* to  $\mathcal{G}$  if there is an internal groupoid  $\mathcal{L}$  and essential homotopy equivalences

$$\mathcal{K} \xleftarrow{\eta} \mathcal{L} \xrightarrow{\nu} \mathcal{G}.$$

**Proposition 4.9.** *Morita homotopy is an equivalence relation over internal groupoids.*

*Proof.* The assertion follows mutatis mutandis, given our assumptions, from the considerations of section 3 in [5].  $\square$

Denote by  $\mathbb{A}HG_{\text{bi}}$  the bicategory consisting of internal groupoids as objects, strict arrows as 1-arrows and homotopy classes of strong homotopies as 2-arrows.

**Proposition 4.10.** *As defined above,  $\mathbb{A}HG_{\text{bi}}$  carries a bicategory structure.*

*Proof.* The definition of vertical composition of strong homotopies is well-defined since vertically filtered homotopy pullbacks exist by assumption.  $\square$

**Proposition 4.11.** *The essential homotopy equivalences  $W$  allow a bicalculus of fractions.*

*Proof.* We have to check the conditions stated in [4]. The proof relies essentially on associativity which is furnished by the existence of vertically filtered homotopy pullbacks. The remainder of the argument follows mutatis mutandis the discussion of [5].  $\square$

**Remark 4.12.** The previous results allow us to form the bicategory localization  $\mathbb{A}HG_{\text{bi}}[W^{-1}]$ . The resulting bicategory fulfills the following universal property: There is a bifunctor  $U: \mathbb{A}HG_2 \rightarrow \mathbb{A}HG_2[W^{-1}]$  that sends essential homotopy equivalences ( $\sim_{eh}$ ) to weak generalized equivalences ( $\sim_W$ ) and is universal with that property, i.e. each bifunctor  $F$  which has this property there is a unique bicategory functor  $G$  such that

$$\begin{array}{ccc} \mathbb{A}HG_{\text{bi}} & \xrightarrow{F} & \mathbb{A}HG_{\text{bi}}[W^{-1}] \\ & \searrow U & \uparrow G \\ & & \mathbb{A}HG_{\text{bi}}[W^{-1}]. \end{array}$$

commutes. In addition, denoting by  $\mathbb{A}G_2[E^{-1}]$  the localization at essential equivalences, the following diagram commutes

$$\begin{array}{ccc} \mathbb{A}G_2 & \xrightarrow{U|_{\mathbb{A}G_2}} & \mathbb{A}G_2[E^{-1}] \\ i_{\mathbb{A}G_2} \downarrow & & \downarrow i_{\mathbb{A}G_2[E^{-1}]} \\ \mathbb{A}HG_{\text{bi}} & \xrightarrow{U} & \mathbb{A}HG_{\text{bi}}[W^{-1}]. \end{array}$$

Morita homotopy equivalence amounts to equivalence of objects in the bicategory  $\mathbb{A}HG_{\text{bi}}[W^{-1}]$ .

The 1-arrows of  $\mathbb{A}HG_{\text{bi}}[W^{-1}]$  are given by  $\mathcal{K} \xleftarrow{\eta} \mathcal{J} \xrightarrow{\varphi} \mathcal{G}$  with  $\eta$  denoting an essential homotopy equivalence. The 2-arrows of  $\mathbb{A}HG_{\text{bi}}[W^{-1}]$  take the form:

$$\begin{array}{ccccc} & & \mathcal{C} & & \\ & \eta & \uparrow u & \varphi & \\ \mathcal{K} & & \mathcal{L} & & \mathcal{G} \\ & \Downarrow \eta & \downarrow v & \Downarrow \eta & \\ & & \mathcal{D} & & \end{array}$$

where  $\mathcal{L}$  is an internal groupoid,  $u, v$  are essential homotopy equivalences and the double arrows denote strong homotopies.

Horizontal composition of 1-arrows is defined by

$$(\mathcal{K} \xleftarrow{\eta} \mathcal{J} \xrightarrow{\varphi} \mathcal{G}) \boxplus (\mathcal{G} \xleftarrow{\nu} \mathcal{J}' \xrightarrow{\psi} \mathcal{L}) = \begin{array}{ccccc} & & \mathcal{P}_n & & \\ & \swarrow & \downarrow \Downarrow & \searrow & \\ \mathcal{K} & \xleftarrow{\eta} & \mathcal{J} & \xrightarrow{\varphi} & \mathcal{G} & \xleftarrow{\nu} & \mathcal{J}' & \xrightarrow{\psi} & \mathcal{L} \end{array}$$

Vertical composition of 2-arrows is defined by

$$\begin{array}{ccccc} & & \mathcal{C} & & \\ & \eta & \uparrow u & \varphi & \\ \mathcal{K} & & \mathcal{L} & & \mathcal{G} \\ & \Downarrow \eta & \downarrow v & \Downarrow \eta & \\ & & \mathcal{D} & & \end{array} \boxplus \begin{array}{ccccc} & & \mathcal{D} & & \\ & \eta' & \uparrow u' & \varphi' & \\ \mathcal{K} & & \mathcal{L}' & & \mathcal{G} \\ & \Downarrow \eta' & \downarrow v' & \Downarrow \eta' & \\ & & \mathcal{E} & & \end{array}$$

$$\begin{array}{ccccc}
& & \mathcal{C} & & \\
& & \uparrow u & & \\
& & \mathcal{L} & & \\
& \eta & & \varphi & \\
& \swarrow & & \searrow & \\
= \mathcal{K} & & & & \mathcal{G} \\
& \Downarrow \mathcal{P}_{n''} & & & \Downarrow \mathcal{P}_{n''} \\
& & \mathcal{L}' & & \mathcal{D} \\
& \eta'' & & \varphi'' & \\
& \swarrow & & \searrow & \\
& & \mathcal{E} & & \\
& & \downarrow v' & & 
\end{array}$$

Internal groupoids  $\mathcal{G}$  and  $\mathcal{K}$  are equivalent in  $\mathbb{A}H\mathcal{G}_{\text{bi}}[W^{-1}]$  if there are arrows  $(\eta, \varphi)$  from  $\mathcal{K}$  to  $\mathcal{G}$  and  $(\theta, \psi)$  from  $\mathcal{G}$  to  $\mathcal{K}$  such that  $(\eta, \varphi) \circ (\theta, \psi)$  is Morita homotopic to the identity  $(\text{id}_{\mathcal{G}}, \text{id}_{\mathcal{G}})$  and  $(\theta, \psi) \circ (\eta, \varphi) \sim_W (\text{id}_{\mathcal{K}}, \text{id}_{\mathcal{K}})$ . The resulting homotopy type is Morita invariant.

**Proposition 4.13.** *If  $\mathcal{K} \sim_{\mathcal{M}} \mathcal{G}$  then  $\mathcal{K} \sim_W \mathcal{G}$ .*

## 5. GEOMETRIC COMPLEXITY

We define the geometric complexity bifunctor on the 2-category of virtual manifolds. This is an extension of the classical Lusternik-Schnirelmann invariant, see [16]. Roughly speaking, this invariant measures the number of transitive components of a given geometric object, up to a Morita invariant homotopy type. Throughout this section we fix a diadic site structure  $(\mathbb{A}, \mathcal{T}, \mathbb{E}, \mathcal{I})$  capturing all the previously stated assumptions. We assume that all internal groupoids globalize a virtual manifold structure.

**Definition 5.1.** Let  $(\mathbb{A}, \tau, \mathbb{E})$  be a site-structure. Given an internal group  $G$  and a point  $\bullet$ , the groupoid  $G \times \bullet \rightrightarrows \bullet$  is called an internal *point groupoid*.

Up to Morita equivalence, point groupoids are transitive, cf. [18].

**Proposition 5.2.** *A groupoid  $\mathcal{G}$ , internal to  $(\mathbb{A}, \tau, \mathbb{E})$ , is transitive if and only if  $\mathcal{G}$  is internally Morita equivalent to an internal point groupoid.*

*Proof.* Assuming  $\mathcal{G}$  is transitive, note that  $s \oplus r \in \mathbb{E}$ . Then the map  $r: \mathcal{G}_{x_0} \rightarrow \mathcal{G}_0$  is the pullback of  $s \oplus r$  along  $\mathcal{G}_0 \rightarrow \mathcal{G}_0 \times \mathcal{G}_0$ ,  $x \mapsto (x_0, x)$  and therefore  $r|_{\mathcal{G}_{x_0}} \in \mathbb{E}$ . The inclusion  $\mathcal{G}_{x_0} \rightarrow \mathcal{G}$  induces a Morita equivalence. In order to see this, since  $\mathcal{G}_{x_0} = \mathcal{G}_{\{x_0\}}$ , we need to show that  $r \circ \text{pr}_1: \mathcal{G}_1 *_{\mathcal{G}_0} \{x_0\} \rightarrow \mathcal{G}_0 \in \mathbb{E}$ . The latter arrow however equals  $r|_{\mathcal{G}_{x_0}}$  which yields the assertion. The other direction is immediate by the induced orbit correspondence.  $\square$

Let  $\widehat{\mathcal{M}}$  be a virtual manifold,  $X \in \mathbb{A}_0$  and set  $\mathcal{G} := \widehat{\mathcal{M}}(X)$ . A given cover  $U \rightarrow X$  is invariant if it is closed with regard to the equivalence relation generated by  $r \oplus s: \mathcal{G} \rightarrow X$ . The equivalence classes are called *orbits*. An orbit groupoid  $\widehat{\mathcal{M}}(\mathcal{O})$  for an orbit  $\mathcal{O} \rightarrow X$  is denoted by  $\mathcal{O}^K$ , where  $K$  denotes the associated isotropy group. A *constant bimap* is given by  $\mathcal{H} \xleftarrow{\varphi} \mathcal{B} \xrightarrow{c} \mathcal{G}$  such that the image of the functor  $c$  is contained in an internal point groupoid.

**Definition 5.3.** Let  $U \twoheadrightarrow X$  be an invariant cover. Then  $\widehat{\mathcal{M}}(X)$  is a *weak point groupoid* if the inclusion functor  $i_U: \widehat{\mathcal{M}}(U) \hookrightarrow \widehat{\mathcal{M}}(X)$  is Morita homotopic to a constant bimap, i.e.

$$\begin{array}{ccccc}
 & & \widehat{\mathcal{M}}(U) & & \\
 & \swarrow \text{id} & \uparrow u & \searrow i_U & \\
 \widehat{\mathcal{M}}(U) & \Downarrow n & \mathcal{L} & \Downarrow n' & \widehat{\mathcal{M}}(X) \\
 & \swarrow c & \downarrow v & \searrow c & \\
 & & \widehat{\mathcal{M}}(U') & \longrightarrow & \bullet^K
 \end{array}$$

where  $u, v$  are essential homotopy equivalences.

**Remark 5.4.** In a weak point subgroupoid  $\widehat{\mathcal{M}}(U)$  the constant bimap can be substituted with a bimap taking values in an orbit groupoid by Proposition 5.2.

We adapt to our framework the definitions of geometric complexity, relative geometric complexity and deformations of groupoids from [5] and [1]. The geometric complexity is invariant with regard to Morita homotopy equivalence. We omit the proof of this, since it can be adapted from loc. cit. to our framework with little effort. The relative geometric complexity elucidates the structure of the absolute point geometry, as we will show.

**Definition 5.5.** The *geometric complexity* of a virtual manifold  $\widehat{\mathcal{M}}$  denoted by  $c^{\text{geo}}(\widehat{\mathcal{M}})$  is the least number of weak point subgroupoids required to cover  $\widehat{\mathcal{M}}(X)$ . If a finite number does not suffice  $c^{\text{geo}}(\mathcal{M}) = \infty$ , i.e. the geometric complexity takes values in  $\mathbb{N} \cup \{\infty\} =: \mathbb{N}_\infty$ .

**Proposition 5.6.** Let  $\widehat{\mathcal{M}}$  and  $\widehat{\mathcal{N}}$  be virtual manifolds. If  $\widehat{\mathcal{M}} \sim_W \widehat{\mathcal{N}}$ , then  $c^{\text{geo}}(\widehat{\mathcal{M}}) = c^{\text{geo}}(\widehat{\mathcal{N}})$ .

By definition, on point groupoids  $\bullet^K$  the geometric complexity is constant equal to 1. Recall that the 2-arrows in  $\mathbb{A}HG_{\text{bi}}[W^{-1}]$  correspond precisely to Morita homotopies. Consider all the point groupoids  $S$ , i.e. all groupoids of the form  $\bullet^K$  for  $K$  an internal group. This can be a set, e.g. if we assume  $\mathbb{A}$  to be locally small. We form the bi-category localization  $\mathbb{A}HG_{\text{bi}}[W^{-1}][S^{-1}]$ . The localization functor  $L_S: \mathbb{A}HG_{\text{bi}}[W^{-1}] \rightarrow (\mathbb{A}HG_{\text{bi}}[W^{-1}])[S^{-1}]$  has the following properties:

- 1) It maps  $S$  to the so-called absolute point  $\bullet$ .
- 2) For each bi-category  $\mathcal{D}$  we have an isomorphism

$$\begin{aligned}
 (-) \circ L_S: \text{Fun}_{\text{bi}}((\mathbb{A}HG_{\text{bi}}[W^{-1}])[S^{-1}], \mathcal{D}) &\xrightarrow{\sim} \text{Fun}_{\text{bi}}(\mathbb{A}HG_{\text{bi}}[W^{-1}], \mathcal{D})_S \\
 &\subset \text{Fun}_{\text{bi}}(\mathbb{A}HG_{\text{bi}}[W^{-1}], \mathcal{D}).
 \end{aligned}$$

Given two bi-categories  $\mathcal{C}, \mathcal{D}$ , we denote by  $\text{Fun}_{\text{bi}}(\mathcal{C}, \mathcal{D})$  the bi-category of bi-functors. The objects consist of strict bi-functors from  $\mathcal{C}$  to  $\mathcal{D}$ . The 2-arrows are modifications between natural transformations, 2-*transfors* in the sense of Crans [6]. In addition for given  $S \subset \text{Mor}(\mathcal{C})_1$  we denote by  $\text{Fun}_{\text{bi}}(\mathcal{C}, \mathcal{D})_S$  the functor bi-category consisting of those bi-functors which send elements of  $S$  to equivalences.

Note that the localization does not work in the same way as the previous construction, since  $S$  does clearly not induce a bicalculus of fractions. Instead we employ a general construction of localization which is possible in any (bi-)category, cf. [6]. The localization can be very large in a set-theoretic sense. In practice we will be interested in the

slice categories  $\mathbb{A}H\mathcal{G}_{\text{bi}}[W^{-1}][S^{-1}]/[\mathcal{G}] =: \mathcal{L}_{\mathcal{G}}$ , where  $\mathcal{G}$  is an internal groupoid and  $[\mathcal{G}]$  denotes its class in the localization. We refer to  $\mathcal{L}_{\mathcal{G}}$  as the *locus* at  $\mathcal{G}$ .

Then  $\mathbf{c}^{\text{geo}}$  extends by Proposition 5.6 to a bifunctor such that the following diagram commutes:

$$\begin{array}{ccc} \mathbb{A}\mathcal{G}_2 & \xrightarrow{\tilde{U}} & \mathbb{A}H\mathcal{G}_{\text{bi}}[W^{-1}][S^{-1}] \\ \downarrow \mathbf{c}^{\text{geo}} & \swarrow \mathbf{c}^{\text{geo}} & \\ \underline{N}_{\infty} & & \end{array}$$

The application of the functor  $\mathbf{c}^{\text{geo}}$  to  $[\mathcal{G}]$  corresponds to the evaluation of the absolute point at the locus  $\mathcal{L}_{\mathcal{G}}$ .

This follows from the universal properties:

$$\begin{array}{ccccc} & & \tilde{U} & & \\ & & \text{---} & & \\ \mathbb{A}H\mathcal{G}_{\text{bi}} & \xrightarrow{U} & \mathbb{A}H\mathcal{G}_{\text{bi}}[W^{-1}] & \xrightarrow{L_S} & (\mathbb{A}H\mathcal{G}_{\text{bi}}[W^{-1}])[S^{-1}] \\ \uparrow i & & \uparrow i & & \uparrow \\ \mathbb{A}\mathcal{G}_2 & \xrightarrow{U|_{\mathbb{A}\mathcal{G}_2}} & \mathbb{A}\mathcal{G}_2[E^{-1}] & & \\ & & \tilde{U}|_{\mathbb{A}\mathcal{G}_2} & & \end{array}$$

**Definition 5.7.** The *relative geometric complexity*  $\mathbf{c}^{\text{geo}}(\mathcal{H}, \mathcal{G})$  of a full subgroupoid  $\mathcal{H}$  of  $\mathcal{G}$  is defined as the least number of  $\mathcal{G}$  weak point groupoids required to cover  $\mathcal{H}$ .

Let  $\mathcal{H}$  and  $\mathcal{K}$  be full subgroupoids of  $\mathcal{G}$ .

**Definition 5.8.** A  $\mathcal{G}$ -*deformation* of  $\mathcal{H}$  into  $\mathcal{K}$  is defined by the following commuting diagram:

$$\begin{array}{ccccc} & & \mathcal{H} & & \\ & \swarrow \text{id} & \uparrow u & \searrow i_{\mathcal{H}} & \\ \mathcal{H} & \Downarrow n & \mathcal{L} & \Downarrow n' & \mathcal{G} \\ & \swarrow \epsilon & \downarrow v & \nearrow \varphi & \uparrow \\ & & \tilde{\mathcal{H}} & \longrightarrow & \mathcal{K} \end{array}$$

Since weak point groupoids are exactly those groupoids that are  $\mathcal{G}$ -deformable to an orbit groupoid we have that complexity increases along deformations.

**Proposition 5.9.** Let  $\mathcal{H}, \mathcal{K}$  be full subgroupoids of  $\mathcal{G}$  such that there is a  $\mathcal{G}$ -deformation from  $\mathcal{H}$  to  $\mathcal{K}$ , then  $\mathbf{c}^{\text{geo}}(\mathcal{H}, \mathcal{G}) \leq \mathbf{c}^{\text{geo}}(\mathcal{K}, \mathcal{G})$ .

**Remark 5.10.** It follows that full subgroupoids  $\mathcal{H}$  induce a partial order structure on the loci  $\mathcal{L}_{\mathcal{H}}$  with respect to  $\mathcal{L}_{\mathcal{G}}$ , with the relation given by deformation. Hence for any fixed system  $X$  endowed with a virtual manifold structure, the past evolution of the system to its present level of complexity is represented by the partial order structure of the loci in the absolute point geometry.

## REFERENCES

- [1] S. Alsulami, H. Colman, F. Neumann, *The Lusternik-Schnirelmann category for a differentiable stack*, International Conference on Mathematics and Statistics. Springer, Cham, 2015.
- [2] I. Androulidakis, G. Skandalis, *The holonomy groupoid of a singular foliation*, Journal für die reine und angewandte Mathematik (Crelles Journal), 2009(626), 1-37.
- [3] J. Baez, A. Hoffnung, *Convenient categories of smooth spaces*, Transactions of the American Mathematical Society 363.11 (2011): 5789-5825.
- [4] J. Benabou, *Introduction to bicategories*, In Midwest Category Seminar, volume 42 of LNM, pages 1–77. Springer, 1967.
- [5] H. Colman, *The Lusternik-Schnirelmann category of a Lie groupoid*, Transactions of the American Mathematical Society 362.10 (2010): 5529-5567.
- [6] S. E. Crans, *Localizations of transfors*, K-theory, 28(1), 39-105, 2003.
- [7] C. Debord, *Holonomy Groupoids of singular Foliations*, J. Differential Geometry, 58 (2001), 467-500.
- [8] G. M. Edelman, *The remembered present: a biological theory of consciousness*, Basic Books, 1989.
- [9] C. Ehresmann, *Catégories et structures*, Dunod, Paris, 1965.
- [10] M. Grandis, *Directed algebraic topology. Models of non-reversible worlds*, New Mathematical Monographs, 13. Cambridge University Press, Cambridge, 2009. x+434 pp.
- [11] G. Grubb, R. T. Seeley, *Zeta and eta functions for Atiyah-Patodi-Singer operators*, The Journal of Geometric Analysis 6.1 (1996): 31-77.
- [12] J. F. Glazebrook, R. Wallace, *Rate Distortion Manifolds as Model Spaces for Cognitive Information*, Informatica (03505596) 33.3 (2009).
- [13] P. Iglesias-Zemmour, *Dimension in diffeology*, Indagationes Mathematicae 18.4 (2007): 555-560.
- [14] P. T. Johnstone, *Sketches of an Elephant: A Topos Theory Compendium: Volume 1*, Oxford University Press, 2002.
- [15] G. J. Klir, *Facets of systems science*, Vol. 7, Springer Science & Business Media, 2013.
- [16] L. Lusternik, L. Schnirelmann, *Méthodes topologiques dans les Problèmes Variationnels*, Hermann, Paris, 1934.
- [17] L. Meersseman, *The Teichmüller and Riemann moduli stacks*, Journal de l'École polytechnique—Mathématiques 6 (2019): 879-945.
- [18] I. Moerdijk, J. Mrčun, *Introduction to foliations and Lie groupoids*, 91. Cambridge Studies in Advanced Mathematics. Cambridge University Press, 2003.
- [19] R. Meyer, C. Zhu, *Groupoids in categories with pretopology*, Theory & Applications of Categories, 2015, 30. Jg., Nr. 55.
- [20] J. Pradines, *How to define the differentiable graph of a singular foliation*, Cahiers Topologie Géom. Différentielle Catég. 26 (1985), no. 4, 339–380.
- [21] D. M. Roberts, *Internal categories, anafunctors and localisations*, Theory and Applications of Categories 26 (2012), 788-829.
- [22] H.E. Winkelkemper, *The graph of a foliation*, Ann. Glob. Analysis and Geometry 1(3) (1983) 51-75.

Email address: [karsten.bohlen@mathematik.uni-regensburg.de](mailto:karsten.bohlen@mathematik.uni-regensburg.de)