# PQ-TYPE ADJACENCY POLYTOPES OF JOIN GRAPHS

HIDEFUMI OHSUGI AND AKIYOSHI TSUCHIYA

ABSTRACT. PQ-type adjacency polytopes  $\nabla_G^{PQ}$  are lattice polytopes arising from finite graphs G. There is a connection between  $\nabla_G^{PQ}$  and the engineering problem known as power-flow study, which models the balance of electric power on a network of power generation. In particular, the normalized volume of  $\nabla_G^{PQ}$  plays a central role. In the present paper, we focus the case where G is a join graph. In fact, formulas of the  $h^*$ -polynomial and the normalized volume of  $\nabla_G^{PQ}$  of a join graph G are presented. Moreover, we give explicit formulas of the  $h^*$ -polynomial and the normalized volume of  $\nabla_G^{PQ}$  when G is a complete multipartite graph or a wheel graph.

#### 1. INTRODUCTION

A *lattice polytope*  $\mathscr{P} \subset \mathbb{R}^n$  is a convex polytope all of whose vertices have integer coordinates. Its *normalized volume*,  $Vol(\mathscr{P}) = dim(\mathscr{P})!vol(\mathscr{P})$  where  $vol(\mathscr{P})$  is the relative volume of  $\mathscr{P}$ , is always a positive integer. To compute  $Vol(\mathscr{P})$  is a fundamental but hard problem in polyhedral geometry.

Let *G* be a simple graph on  $[n] := \{1, ..., n\}$  with edge set E(G). The *PV-type adjacency poly*tope  $\nabla_G^{PV}$  of *G* is the lattice polytope which is the convex hull of

$$\{\pm(\mathbf{e}_i-\mathbf{e}_j)\in\mathbb{R}^n:\{i,j\}\in E(G)\},\$$

where  $\mathbf{e}_i$  is the *i*-th unit coordinate vector in  $\mathbb{R}^n$ . The normalized volumes of PV-type adjacency polytopes have attracted much attention. In fact, the normalized volume of a PV-type adjacency polytope gives an upper bound on the number of possible solutions in the Kuramoto equations ([5]), which models the behavior of interacting oscillators ([13]). For several classes of graphs, explicit formulas for the normalized volume of their PV-type adjacency polytopes have been given (e.g., [1, 7, 10]). In particular, we can compute the normalized volume of the PV-type adjacency polytope of a suspension graph by using interior polynomials ([15]). Here interior polynomials are a version of the Tutte polynomials for hypergraphs introduced by Kálmán [11]. On the other hand, the *PQ-type adjacency polytope*  $\nabla_G^{PQ}$  of *G* is the lattice polytope which is the convex hull of

$$\{(\mathbf{e}_i, \mathbf{e}_j) \in \mathbb{R}^{2n} : \{i, j\} \in E(G) \text{ or } i = j\}.$$

There is a connection between PQ-type adjacency polytopes and the engineering problem known as power-flow study, which models the balances of electric power on a network of power generation ([6]). In fact, the normalized volume of a PQ-type adjacency polytope gives an upper bound on the number of possible solutions in the algebraic power-flow equations. Davis and Chen [8] showed that the normalized volume of  $\nabla_G^{PQ}$  can be computed by using sequences of nonnegative integers related to the *Dragon Marriage Problem* [17].

<sup>2010</sup> Mathematics Subject Classification. 05A15, 05C31, 52B12, 52B20.

Key words and phrases. adjacency polytope,  $h^*$ -polynomial, interior polynomial, perfectly matchable set, join graph, root polytope.

In the present paper, we focus on the  $h^*$ -polynomial of a PQ-type adjacency polytope. Here, the  $h^*$ -polynomial  $h^*(\mathscr{P}, x)$  of a lattice polytope  $\mathscr{P}$  is a discrete tool to compute the normalized volume Vol( $\mathscr{P}$ ) (see Section 2). We recall a relation between  $\nabla_G^{PQ}$  and a root polytope. For a bipartite graph H on [n] with edge set E(H), the root polytope  $\mathscr{Q}_H$  of H is the lattice polytope which is the convex hull of

$$\{\mathbf{e}_i + \mathbf{e}_j \in \mathbb{R}^n : \{i, j\} \in E(H)\}.$$

For a positive integer *n*, set  $[\overline{n}] := \{\overline{1}, ..., \overline{n}\}$ . Define D(G) to be the bipartite graph on  $[n] \cup [\overline{n}]$  with edges  $\{i, \overline{i}\}$  for each  $i \in [n]$  and  $\{i, \overline{j}\}$  and  $\{\overline{i}, j\}$  for each edge  $\{i, j\}$  in *G*. It then follows that  $\nabla_G^{PQ}$ is unimodularly equivalent to  $\mathscr{Q}_{D(G)}$  ([8, Lemma 2.4]). On the other hand, it is known [12] that the  $h^*$ -polynomial of the root polytope  $\mathscr{Q}_H$  of a connected bipartite graph *H* coincides with the interior polynomial  $I_H(x)$  of the associated hypergraph of *H*. In particular, the normalized volume of  $\mathscr{Q}_H$ is equal to |HT(H)|, where HT(H) denotes the set of hypertrees of the associated hypergraph of *H*. Therefore, we can compute the  $h^*$ -polynomial and the normalized volume of  $\nabla_G^{PQ}$  of a connected graph *G* by using an interior polynomial and counting hypertrees.

The main result of the present paper is formulas of the  $h^*$ -polynomial and the normalized volume of  $\nabla_G^{PQ}$  of a join graph G. Let  $G_1, \ldots, G_s$  be graphs with  $m_1, \ldots, m_s$  vertices. Suppose that  $G_i$ and  $G_j$  have no common vertices for each  $i \neq j$ . Then the *join*  $G_1 + \cdots + G_s$  of  $G_1, \ldots, G_s$  is obtained from  $G_1 \cup \cdots \cup G_s$  joining each vertex of  $G_i$  to each vertex of  $G_j$  for any  $i \neq j$ . Note that  $G_1 + \cdots + G_s$  (s > 1) is connected and hence so is  $D(G_1 + \cdots + G_s)$ . For example, the complete bipartite graph  $K_{\ell,m}$  is equal to the join  $E_{\ell} + E_m$  where  $E_k$  is the empty graph with k vertices. For complete graphs  $K_{\ell}$  and  $K_m$ , one has  $K_{\ell} + K_m = K_{\ell+m}$ . We can compute the  $h^*$ -polynomial and the normalized volume of the PQ-type adjacency polytope of a join graph by using perfectly matchable set polynomials (see Section 2 for the definition of perfectly matchable set polynomials).

**Theorem 1.1.** Let  $G_1, \ldots, G_s$  be graphs with  $m_1, \ldots, m_s$  vertices. Suppose that  $G_i$  and  $G_j$  have no common vertices for each  $i \neq j$ . Then for the join  $G = G_1 + \cdots + G_s$  with  $m = \sum_{i=1}^s m_i$  vertices, we have

(1) 
$$h^*(\nabla_G^{PQ}, x) = \sum_{i=1}^s h^*(\nabla_{G_i + K_{m-m_i}}^{PQ}, x) - (s-1) \sum_{k=0}^{m-1} {\binom{m-1}{k}}^2 x^k$$

(2) 
$$= \sum_{i=1}^{s} I_{D(G_i+K_{m-m_i})}(x) - (s-1) \sum_{k=0}^{m-1} {\binom{m-1}{k}}^2 x^k$$

(3) 
$$= \sum_{i=1}^{s} p(D(G_i + K_{m-m_i-1}), x) - (s-1) \sum_{k=0}^{m-1} {\binom{m-1}{k}}^2 x^k,$$

where p(H,x) denotes the perfectly matchable set polynomial of a graph H. In particular, one has

$$Vol(\nabla_G^{PQ}) = \sum_{i=1}^{s} Vol(\nabla_{G_i+K_{m-m_i}}^{PQ}) - (s-1) \binom{2(m-1)}{m-1}$$
$$= \sum_{i=1}^{s} |HT(D(G_i+K_{m-m_i}))| - (s-1) \binom{2(m-1)}{m-1}$$
$$= \sum_{i=1}^{s} |PM(D(G_i+K_{m-m_i-1}))| - (s-1) \binom{2(m-1)}{m-1},$$

where PM(H) denotes the set of perfectly matchable sets of a graph H.

By using this theorem, we give explicit formulas of the  $h^*$ -polynomial and the normalized volume of  $\nabla_G^{PQ}$  when *G* is a complete multipartite graph (Corollary 4.4). On the other hand, Theorem 1.1 is not useful for computing the  $h^*$ -polynomial and the normalized volume of  $\nabla_G^{PQ}$  when *G* is a wheel graph  $W_n$ , that is, *G* is the join of a cycle  $C_n$  and  $K_1$ . We give explicit formulas for the  $h^*$ -polynomial and the normalized volume of  $\nabla_{W_n}^{PQ}$  when *G* is a wheel graph  $W_n$ , that is, *G* is the join of a cycle  $C_n$  and  $K_1$ . We give explicit formulas for the  $h^*$ -polynomial and the normalized volume of  $\nabla_{W_n}^{PQ}$  and prove the conjecture [8, Conjecture 4.4] on the normalized volume of  $\nabla_{W_n}^{PQ}$  (Theorem 5.1) by computing the perfectly matchable set polynomial of  $D(C_n)$ .

The paper is organized as follows: After reviewing the definitions and properties of the  $h^*$ -polynomials of lattice polytopes and the interior polynomials of connected bipartite graphs in Section 2, we give a proof of Theorem 1.1 in Section 3. By using Theorem 1.1, explicit formulas of the  $h^*$ -polynomial and the normalized volume of the PQ-type adjacency polytope of a complete multipartite graph are presented in Section 4. Finally, we compute the  $h^*$ -polynomial and the normalized volume of a wheel graph in Section 5.

## 2. PRELIMINARIES

As explained in the previous section, the  $h^*$ -polynomial of  $\nabla_G^{PQ}$  is equal to the interior polynomial of D(G). First, we recall what  $h^*$ -polynomials are. Let  $\mathscr{P} \subset \mathbb{R}^n$  be a lattice polytope of dimension d. Given a positive integer t, we define

$$L_{\mathscr{P}}(t) = |t\mathscr{P} \cap \mathbb{Z}^n|,$$

where  $t\mathscr{P} := \{t\mathbf{x} \in \mathbb{R}^n : \mathbf{x} \in \mathscr{P}\}$ . The study on  $L_{\mathscr{P}}(t)$  originated in Ehrhart [9] who proved that  $L_{\mathscr{P}}(t)$  is a polynomial in t of degree d with the constant term 1. We call  $L_{\mathscr{P}}(t)$  the *Ehrhart polynomial* of  $\mathscr{P}$ . The generating function of the lattice point enumerator, i.e., the formal power series

$$\operatorname{Ehr}_{\mathscr{P}}(x) = 1 + \sum_{k=1}^{\infty} L_{\mathscr{P}}(k) x^{k}$$

is called the *Ehrhart series* of  $\mathcal{P}$ . It is known that it can be expressed as a rational function of the form

$$\operatorname{Ehr}_{\mathscr{P}}(x) = \frac{h^*(\mathscr{P}, x)}{(1-x)^{d+1}},$$

where  $h^*(\mathscr{P}, x)$  is a polynomial in x of degree at most d with nonnegative integer coefficients and it is called the  $h^*$ -polynomial (or the  $\delta$ -polynomial) of  $\mathscr{P}$ . Moreover,

$$h^*(\mathscr{P}, x) = \sum_{i=0}^d h_i^* x^i$$

satisfies  $h_0^* = 1$ ,  $h_1^* = |\mathscr{P} \cap \mathbb{Z}^n| - (d+1)$  and  $h_d^* = |int(\mathscr{P}) \cap \mathbb{Z}^n|$ , where  $int(\mathscr{P})$  is the relative interior of  $\mathscr{P}$ . Furthermore,  $h^*(\mathscr{P}, 1) = \sum_{i=0}^d h_i^*$  is equal to the normalized volume  $Vol(\mathscr{P})$  of  $\mathscr{P}$ . We refer the reader to [4] for the detailed information about Ehrhart polynomials and  $h^*$ -polynomials.

Next, we recall the definition of interior polynomials and their properties. A *hypergraph* is a pair  $\mathscr{H} = (V, E)$ , where  $E = \{e_1, \ldots, e_n\}$  is a finite multiset of non-empty subsets of  $V = \{v_1, \ldots, v_m\}$ . Elements of *V* are called vertices and the elements of *E* are the hyperedges. Then we can associate  $\mathscr{H}$  to a bipartite graph Bip $\mathscr{H}$  on the vertex set  $V \cup E$  with the edge set  $\{\{v_i, e_j\} : v_i \in e_j\}$ . Assume that Bip $\mathscr{H}$  is connected. A *hypertree* in  $\mathscr{H}$  is a function  $f : E \to \mathbb{Z}_{\geq 0}$  such that there exists a

spanning tree  $\Gamma$  of Bip $\mathscr{H}$  whose vertices have degree f(e) + 1 at each  $e \in E$ . Then we say that  $\Gamma$  induces f. Let HT( $\mathscr{H}$ ) denote the set of all hypertrees in  $\mathscr{H}$ . A hyperedge  $e_i \in E$  is said to be *internally inactive* with respect to the hypertree f if there exists j' < j such that  $g: E \to \mathbb{Z}_{\geq 0}$ defined by /

$$g(e_i) = \begin{cases} f(e_i) + 1 & (i = j') \\ f(e_i) - 1 & (i = j) \\ f(e_i) & (\text{otherwise}) \end{cases}$$

is a hypertree. Let  $\overline{\iota}(f)$  be the number of internally inactive hyperedges of f. Then the *interior* polynomial of  $\mathscr{H}$  is the generating function  $I_{\mathscr{H}}(x) = \sum_{f \in \mathrm{HT}(\mathscr{H})} x^{\overline{\iota}(f)}$ . It is known [11, Proposition 6.1] that deg  $I_{\mathscr{H}}(x) \leq \min\{|V|, |E|\} - 1$ . If  $G = \operatorname{Bip}\mathscr{H}$ , then we set  $\operatorname{HT}(G) = \operatorname{HT}(\mathscr{H})$  and  $I_G(x) = I_{\mathcal{H}}(x)$ . The coefficients of  $I_G(x)$  are described as follows.

**Proposition 2.1** ([11, Theorem 3.4]). Let G be a connected bipartite graph on the vertex set  $V_1 \sqcup V_2$ where  $V_1 = \{v_1, \ldots, v_p\}$  and  $|V_2| = q$ . Then the coefficient of  $x^k$  in  $I_G(x)$  is the number of functions  $f: V_1 \to \mathbb{Z}_{>0}$  such that

- (i)  $\sum_{i=1}^{p} f(v_i) = q 1;$ (ii)  $\sum_{v \in V'} f(v) \le |\Gamma_G(V')| 1$  for all  $V' \subset V_1$ , where  $\Gamma_G(S) \subset V_2$  is the set of vertices adjacent to some vertex in S;
- (iii)  $\overline{\iota}(f) = k$ , *i.e.*,  $|\eta_G(f)| = k$  where  $\eta_G(f)$  is the set of vertices  $v_i \in V_1$  satisfying the following condition: there exists j' < j such that the function  $g: V_1 \to \mathbb{Z}_{>0}$  defined by

$$g(v_i) = \begin{cases} f(v_i) + 1 & (i = j') \\ f(v_i) - 1 & (i = j) \\ f(v_i) & (otherwise) \end{cases}$$

satisfies condition (ii) above.

From [8, Lemma 2.4] and [12, Theorems 1.1 and 3.10], we have the following.

**Proposition 2.2.** Let G be a connected graph. Then  $h^*(\nabla_G^{PG}, x) = I_{D(G)}(x)$ . In particular, the normalized volume of  $\nabla_G^{PG}$  is  $I_{D(G)}(1) = |HT(D(G))|$ .

Let G be a finite graph on the vertex set V = [n]. A k-matching of G is a set of k pairwise non-adjacent edges of G. The matching generating polynomial of G is

$$g(G,x) = \sum_{k\geq 0} m_G(k) x^k,$$

where  $m_G(k)$  is the number of k-matchings of G. A k-matching of G is said to be *perfect* if 2k = n. A subset  $S \subset V$  is called a *perfectly matchable set* [2] if the induced subgraph of G on the vertex set S has a perfect matching. Let PM(G,k) be the set of all perfectly matchable sets S of G with |S| = 2k and PM(G) the set of all perfectly matchable sets of G. We regard  $\emptyset$  as a perfectly matchable set and we set  $PM(G,0) = \{\emptyset\}$ . Note that  $|PM(G,k)| \le m_G(k)$  holds in general. We call the polynomial

$$p(G,x) = \sum_{k \ge 0} |\mathsf{PM}(G,k)| x^k$$

the *perfectly matchable set polynomial* (PMS polynomial) of G.

**Example 2.3.** For the cycle  $C_4$  of length 4, we have  $g(C_4, x) = 2x^2 + 4x + 1$  and  $p(C_4, x) = x^2 + 4x + 1$ . On the other hand, if a graph *G* has no even cycles, then we have g(G, x) = p(G, x).

Assume that *G* is a bipartite graph with a bipartition  $[n] = V_1 \sqcup V_2$ . Then let  $\tilde{G}$  be a connected bipartite graph on [n+2] whose edge set is

$$E(\tilde{G}) = E(G) \cup \{\{i, n+1\} : i \in V_1\} \cup \{\{j, n+2\} : j \in V_2 \cup \{n+1\}\}$$

**Proposition 2.4** ([14, Proposition 3.4]). Let G be a bipartite graph. Then we have

$$I_{\widetilde{G}}(x) = p(G, x).$$

The following Lemma is easy to see. However it is useful when we apply Proposition 2.4 to such graphs.

**Lemma 2.5.** Let G be a graph. Then we have

$$D(G+K_1)=D(G).$$

Moreover, we have

**Proposition 2.6.** Let G be a graph with n vertices. Then the  $h^*$ -polynomial of  $\nabla_{G+K_1}^{PQ}$  is p(D(G), x). In particular, the normalized volume of  $\nabla_{G+K_1}^{PQ}$  is |PM(D(G))|.

*Proof.* The assertion follows from Propositions 2.2, 2.4 and Lemma 2.5.

**Remark 2.7.** Given a graph *G*, the number of matchings of *G* is called *Hosoya index* of *G* and denoted by Z(G). From Proposition 2.6, the normalized volume of  $\nabla_{G+K_1}^{PQ}$  is at most Z(D(G)).

Let *G* be a graph on the vertex set [n]. Given a subset  $S \subset [n]$ , we associate the (0,1)-vector  $\rho(S) = \sum_{i \in S} \mathbf{e}_i \in \mathbb{R}^n$ . For example,  $\rho(\emptyset) = \mathbf{0} \in \mathbb{R}^n$ . The convex hull of

$$\{\rho(S): S \in \mathsf{PM}(G,k) \text{ for some } k\}$$

is called a *perfectly matchable subgraph polytope* (PMS polytope) of G. A system of linear inequalities for a PMS polytope of G was given in [2] for bipartite graphs, and in [3] for arbitrary graphs.

**Proposition 2.8** ([2, Theorem 1]). Let G be a bipartite graph on the vertex set  $[n] = V_1 \sqcup V_2$ . Then the PMS polytope of G is the set of all vectors  $(x_1, \ldots, x_n) \in \mathbb{R}^n$  such that

$$0 \le x_i \le 1$$
 for each  $i = 1, 2, ..., n$ ,

$$\sum_{i\in V_1} x_i = \sum_{j\in V_2} x_j,$$

$$\sum_{i \in S} x_i \leq \sum_{j \in \Gamma_G(S)} x_j \text{ for all } S \subset V_1,$$

where  $\Gamma_G(S) \subset V_2$  is the set of vertices adjacent to some vertex in *S*.

#### 3. JOIN GRAPHS

In the present section, we give a proof of Theorem 1.1. Given a graph G and a nonnegative integer k, let  $F_G(k)$  be the set of functions satisfying conditions (i)–(iii) in Proposition 2.1 for D(G).

**Lemma 3.1.** Let  $G_1$  and  $G_2$  be graphs with  $m_1$  and  $m_2$  vertices, respectively. Suppose that  $G_1$  and  $G_2$  have no common vertices. Then for the join  $G = G_1 + G_2$  with  $m = m_1 + m_2$  vertices, we have

(4) 
$$\bigsqcup_{k} F_{G}(k) = \left(\bigsqcup_{k} F_{G_{1}+K_{m_{2}}}(k)\right) \cap \left(\bigsqcup_{k} F_{K_{m_{1}}+G_{2}}(k)\right),$$

(5) 
$$\left(\bigsqcup_{k} F_{G_1+K_{m_2}}(k)\right) \cup \left(\bigsqcup_{k} F_{K_{m_1}+G_2}(k)\right) = \bigsqcup_{k} F_{K_m}(k).$$

*Proof.* Let  $V_1 \sqcup V_2$  with  $V_1 = \{v_1, ..., v_m\}$  and  $V_2 = \{v'_1, ..., v'_m\}$  denote the common vertex set of D(G),  $D(G_1 + K_{m_2})$ ,  $D(K_{m_1} + G_2)$  and  $D(K_m)$  (=  $K_{m,m}$ ). In addition, let  $V_{1,1} = \{v_1, ..., v_{m_1}\}$ ,  $V_{1,2} = \{v_{m_1+1}, ..., v_m\}$ ,  $V_{2,1} = \{v'_1, ..., v'_{m_1}\}$  and  $V_{2,2} = \{v'_{m_1+1}, ..., v'_m\}$ , where  $V_{1,j} \sqcup V_{2,j}$  corresponds to the vertex set of  $D(G_j)$  for j = 1, 2.

**Case 1** ( $D(K_m)$ ). Since  $|\Gamma_{D(K_m)}(V')| - 1 = m - 1$  for all  $V' \subset V_1$ , condition (i) implies condition (ii) in Proposition 2.1.

**Case 2** ( $D(G_1 + K_{m_2})$ ). Suppose that f satisfies condition (i). If  $v \in V' \subset V_1$  for some  $v \in V_{1,2}$ , then condition (ii) holds for V' since deg(v) = m. Thus, if f satisfies condition (i), then condition (ii) in Proposition 2.1 is equivalent to the condition

(6) 
$$\sum_{v \in V'} f(v) \le |\Gamma_{D(G_1)}(V')| + m_2 - 1 \text{ for all } V' \subset V_{1,1}$$

**Case 3**  $(D(K_{m_1} + G_2))$ . By the similar argument as in Case 2, if f satisfies condition (i), then condition (ii) in Proposition 2.1 is equivalent to the condition

(7) 
$$\sum_{v \in V'} f(v) \le |\Gamma_{D(G_2)}(V')| + m_1 - 1 \text{ for all } V' \subset V_{1,2}$$

**Case 4** (D(G)). Suppose that f satisfies condition (i). If  $v, v' \in V' \subset V_1$  for some  $v \in V_{1,1}$  and  $v' \in V_{1,2}$ , then condition (ii) holds for V' since  $\Gamma_{D(G)}(\{v, v'\}) = V_2$ . Hence, if f satisfies condition (i), then condition (ii) in Proposition 2.1 holds if and only if both (6) and (7) hold.

Hence (4) holds.

Since any element in  $F_{G_1+K_{m_2}}(k) \cup F_{K_{m_1}+G_2}(k)$  satisfies condition (i), we have

$$\left(\bigsqcup_{k}F_{G_{1}+K_{m_{2}}}(k)\right)\cup\left(\bigsqcup_{k}F_{K_{m_{1}}+G_{2}}(k)\right)\subset\bigsqcup_{k}F_{K_{m}}(k).$$

Suppose that f satisfies none of (6) and (7). Then

$$\sum_{v \in V'} f(v) \ge |\Gamma_{D(G_1)}(V')| + m_2 \text{ for some } V' \subset V_{1,1},$$
$$\sum_{v \in V''} f(v) \ge |\Gamma_{D(G_2)}(V'')| + m_1 \text{ for some } V'' \subset V_{1,2}.$$

It then follows that  $\sum_{v \in V_1} f(v) \ge m_1 + m_2 = m$ . Hence, if *f* satisfies condition (i), then at least one of (6) or (7) holds. Thus (5) holds.

**Lemma 3.2.** Let  $G_1$  and  $G_2$  be graphs with  $m_1$  and  $m_2$  vertices, respectively. Suppose that  $G_1$  and  $G_2$  have no common vertices. Then for the join  $G = G_1 + G_2$  with  $m = m_1 + m_2$  vertices,  $F_{K_m}(k)$  is decomposed into the disjoint sets

$$F_{K_m}(k) = F_{K_{m_1}+G_2}(k) \sqcup (F_{G_1+K_{m_2}}(k) \setminus F_G(k))$$

as in Fig. 1.



FIGURE 1. Decomposition of  $F_{K_m}(k)$ 

*Proof.* Let  $V_1 \sqcup V_2$  with  $V_1 = \{v_1, ..., v_m\}$  and  $V_2 = \{v'_1, ..., v'_m\}$  denote the common vertex set of D(G),  $D(G_1 + K_{m_2})$ ,  $D(K_{m_1} + G_2)$  and  $D(K_m)$  (=  $K_{m,m}$ ). In addition, let  $V_{1,1} = \{v_1, ..., v_{m_1}\}$ ,  $V_{1,2} = \{v_{m_1+1}, ..., v_m\}$ ,  $V_{2,1} = \{v'_1, ..., v'_{m_1}\}$  and  $V_{2,2} = \{v'_{m_1+1}, ..., v'_m\}$ , where  $V_{1,j} \sqcup V_{2,j}$  corresponds to the vertex set of  $D(G_j)$  for j = 1, 2.

**Claim 1.**  $F_{K_{m_1}+G_2}(k) \subset F_{K_m}(k)$ .

Suppose that f belongs to  $F_{K_{m_1}+G_2}(k) \cup F_{K_m}(k)$ . Then condition (ii) in Proposition 2.1 is independent from the value  $f(v_1)$  since deg $(v_1) = m$ . Thus one can choose j' = 1 for condition (iii) for any j > 1, and hence we have

$$k = |\{v_j \in V_1 : v_j \neq v_1, f(v_j) > 0\}|.$$

Therefore

$$F_{K_{m_1}+G_2}(k) \subset F_{K_m}(k)$$

and  $F_{K_{m_1}+G_2}(\ell) \cap F_{K_m}(k) = \emptyset$  if  $\ell \neq k$ .

**Claim 2.**  $F_G(k) \subset F_{G_1+K_{m_2}}(k)$ .

Suppose that  $f \in F_G(k)$ . Then  $f \in F_{G_1+K_{m_2}}(\ell)$  for some  $\ell$ . Note that  $k = |\eta_{D(G)}(f)|$  and  $\ell = |\eta_{D(G_1+K_{m_2})}(f)|$  (where  $\eta$  is defined in Proposition 2.1). In order to prove  $k = \ell$ , we will show that  $\eta_{D(G)}(f) = \eta_{D(G_1+K_{m_2})}(f)$ . Suppose that  $f(v_j) > 0$  for  $v_j \in V_1$  ( $j \neq 1$ ). If  $j \leq m_1 + 1$ , then  $v_j$  belongs to  $\eta_{D(G)}(f)$  if and only if  $v_j$  belongs to  $\eta_{D(G_1+K_{m_2})}(f)$  since the vertex  $v_{j'}$  in condition (iii) in Proposition 2.1 should be chosen from  $V_{1,1}$ . Let  $j \geq m_1 + 2$ . Since  $v_{m_1+1}$  belongs to  $V_{1,2}$  with

 $m_1 + 1 < j$ , (6) holds for g defined in condition (iii) of Proposition 2.1 where  $v_{j'} = v_{m_1+1}$ . Thus  $v_{j'} = v_{m_1+1}$  satisfies condition (iii) of Proposition 2.1 for  $v_j$  in  $D(G_1 + K_{m_2})$ . Hence  $v_j$  belongs to  $\eta_{D(G_1+K_{m_2})}(f)$ . We now show that any  $v_j$   $(j \ge m_1+2)$  belongs to  $\eta_{D(G)}(f)$ , i.e., there exists j' < j such that  $g: V_1 \to \mathbb{Z}_{\ge 0}$  defined by

(9) 
$$g(v_i) = \begin{cases} f(v_i) + 1 & (i = j') \\ f(v_i) - 1 & (i = j) \\ f(v_i) & (\text{otherwise}) \end{cases}$$

is a hypertree in D(G). Let  $\Gamma$  be a spanning tree of D(G) that induces f. Suppose that  $\Gamma$  does not contain an edge  $e = \{v_k, v'_1\}$  for some  $m_1 < k \le m$ . Then  $\Gamma \cup \{e\}$  has a unique cycle, and the cycle contains  $v_k$ . Let e' be the edge of the cycle that is adjacent to  $v_k$  but different from e. Then  $(\Gamma \cup \{e'\}) \setminus \{e\}$  induces f. Thus we may assume that  $\{\{v_k, v'_1\} : m_1 < k \le m\} \subset \Gamma$ . By the same argument, we may assume that  $\{\{v_k, v'_{m_1+1}\} : 1 \le k \le m_1\}$  is a subset of  $\Gamma$ . Hence

$$S = \{\{v_1, v'_{m_1+1}\}, \dots, \{v_{m_1}, v'_{m_1+1}\}, \{v_{m_1+1}, v'_1\}, \dots, \{v_m, v'_1\}\}$$

is a subset of  $\Gamma$ . Since  $\Gamma$  is spanning,

$$T = \{\{v_{i_2}, v'_2\}, \dots, \{v_{i_{m_1}}, v'_{m_1}\}, \{v_{i_{m_1+2}}, v'_{m_1+2}\}, \dots, \{v_{i_m}, v'_m\}\}$$

is a subset of  $\Gamma$  for some  $1 \le i_2, \ldots, i_{m_1}, i_{m_1+2}, \ldots, i_m \le m$ . Then  $S \cap T = \emptyset$  and  $|S \cup T| = 2m - 2$ . Since  $\Gamma$  has 2m - 1 edges, we have

 $\Gamma = S \sqcup T \sqcup \{e\}$ 

for some edge *e* of D(G). In particular, the degree of each vertex  $v'_i$   $(i \in [m] \setminus \{1, m_1 + 1\})$  in  $\Gamma$  is at most 2. Since  $f(v_j) > 0$ , the degree of  $v_j$  in  $\Gamma$  is at least 2, and hence  $\{v_j, v'_k\}$  is an edge of  $\Gamma$  for some  $k \in [m] \setminus \{1\}$ . Let  $e' = \{v_j, v'_k\}$ . We now construct a spanning tree of D(G) that induces a hypertree of the form (9). Note that, for an edge e'' of D(G),  $\Gamma' = (\Gamma \cup \{e''\}) \setminus \{e'\}$  is a spanning tree of D(G) if  $e'' \notin \Gamma$  and the (unique) cycle of  $\Gamma \cup \{e''\}$  contains e'.

**Case A**  $(k = m_1 + 1)$ . Then  $\Gamma$  contains a path  $(v'_1, v_j, v'_{m_1+1}, v_1)$ . Since  $\Gamma$  has no cycles,  $\{v_1, v'_1\}$  is not an edge of  $\Gamma$ . Hence  $(\Gamma \cup \{v_1, v'_1\}) \setminus \{e'\}$  is a spanning tree of D(G) that induces a hypertree of the form (9) where j' = 1.

**Case B**  $(k \neq m_1 + 1 \text{ and } \deg(v'_k) = 1)$ . If  $2 \leq k \leq m_1$ , then  $(\Gamma \cup \{v_k, v'_k\}) \setminus \{e'\}$  is a spanning tree of D(G) that induces a hypertree of the form (9) where j' = k. If  $m_1 + 2 \leq k \leq m$ , then  $(\Gamma \cup \{v_1, v'_k\}) \setminus \{e'\}$  is a spanning tree of D(G) that induces a hypertree of the form (9) where j' = 1.

**Case** C ( $k \neq m_1 + 1$  and deg( $v'_k$ ) = 2). Then  $\Gamma$  contains an edge { $v_\ell, v'_k$ } for some  $\ell \neq j$ . If  $m_1 < \ell \le m$ , then  $\Gamma$  contains a cycle ( $v'_1, v_j, v'_k, v_\ell, v'_1$ ) of length 4. This is a contradiction. Hence  $\ell \le m_1$ . Then  $\Gamma$  contains a path ( $v'_1, v_j, v'_k, v_\ell, v'_{m_1+1}, v_1$ ). Since  $\Gamma$  has no cycles, { $v_1, v'_1$ } is not an edge of  $\Gamma$ . Hence ( $\Gamma \cup \{v_1, v'_1\}$ ) \ {e'} is a spanning tree of D(G) that induces a hypertree of the form (9) where j' = 1.

Thus we have  $k = \ell$ , and hence

(10) 
$$F_G(k) \subset F_{G_1+K_{m_2}}(k).$$

**Claim 3.**  $(F_{G_1+K_{m_2}}(k) \setminus F_G(k)) \subset F_{K_m}(k).$ 

Suppose that f belongs to  $F_{G_1+K_{m_2}}(k) \setminus F_G(k)$ . Then there exists  $V' \subset V_{1,2}$  such that

$$\sum_{v \in V'} f(v) \ge |\Gamma_{D(G_2)}(V')| + m_1.$$

In particular,  $\sum_{v \in V_{1,2}} f(v) > m_1$ . Since  $\sum_{v \in V_1} f(v) = m - 1$ , we have  $\sum_{v \in V_{1,1}} f(v) < m_2 - 1$ . Hence

$$\sum_{v \in V''} f(v) < m_2 - 1 < |\Gamma_{D(G_1)}(V'')| + m_2 - 1$$

for all  $V'' \subset V_{1,1}$ . Thus, for each  $v_j \neq v_1$  with  $f(v_j) > 0$ ,  $g: V_1 \to \mathbb{Z}_{\geq 0}$  defined by

$$g(v_i) = \begin{cases} f(v_i) + 1 & (i = 1) \\ f(v_i) - 1 & (i = j) \\ f(v_i) & (\text{otherwise}) \end{cases}$$

satisfies

$$\sum_{v \in V''} g(v) \le 1 + \sum_{v \in V''} f(v) \le |\Gamma_{D(G_1)}(V'')| + m_2 - 1$$

for all  $V'' \subset V_{1,1}$ . Therefore we have

$$k = |\{v_j \in V_1 : v_j \neq v_1, f(v_j) > 0\}|_{\mathcal{X}}$$

and hence

(11) 
$$(F_{G_1+K_{m_2}}(k)\setminus F_G(k))\subset F_{K_m}(k).$$

**Claim 4.**  $F_{K_m}(k) = F_{K_{m_1}+G_2}(k) \cup (F_{G_1+K_{m_2}}(k) \setminus F_G(k)).$ From (8) and (11),

$$F_{K_m}(k) \supset F_{K_{m_1}+G_2}(k) \cup (F_{G_1+K_{m_2}}(k) \setminus F_G(k)).$$

Let  $f \in F_{K_m}(k) \setminus F_{K_{m_1}+G_2}(k)$ . Since  $F_{K_{m_1}+G_2}(k') \cap F_{K_m}(k) = \emptyset$  for any  $k' \neq k$ , it follows that  $f \notin F_{K_{m_1}+G_2}(\ell)$  for any  $\ell$ . Hence from (5),  $f \in F_{G_1+K_{m_2}}(\ell)$  for some  $\ell$ . If  $f \in F_G(\ell)$ , then  $f \in F_{K_{m_1}+G_2}(\ell')$  for some  $\ell'$  by (4). This is a contradiction. Thus  $f \in F_{G_1+K_{m_2}}(\ell) \setminus F_G(\ell)$ . From (11), we have

$$f \in F_{G_1 + K_{m_2}}(\ell) \setminus F_G(\ell) \subset F_{K_m}(\ell)$$

Then  $\ell = k$ . It follows that

$$F_{K_m}(k) = F_{K_{m_1}+G_2}(k) \cup (F_{G_1+K_{m_2}}(k) \setminus F_G(k)).$$

Finally, we show that this is a decomposition. Suppose that  $f \in F_{G_1+K_{m_2}}(k) \cap F_{K_{m_1}+G_2}(k)$ . From (4),  $f \in F_G(\ell)$  for some  $\ell$ . Moreover, from (10), we have  $\ell = k$ . Thus

$$F_{G_1+K_{m_2}}(k) \cap F_{K_{m_1}+G_2}(k) \subset F_G(k).$$

Therefore  $F_{K_m}(k)$  is decomposed into the disjoint sets

$$F_{K_m}(k) = F_{K_{m_1}+G_2}(k) \sqcup (F_{G_1+K_{m_2}}(k) \setminus F_G(k))$$

as in Fig. 1.

Now, we are in the position to give a proof of Theorem 1.1.

*Proof of Theorem 1.1.* We prove this by induction on *s*. First we discuss the case when s = 2, i.e.,  $G = G_1 + G_2$ . It is known [12, Example 5.3] that

$$|F_{K_m}(k)| = \binom{m-1}{k}^2.$$

From Lemma 3.2,

$$\binom{m-1}{k}^2 = |F_{K_{m_1}+G_2}(k)| + (|F_{G_1+K_{m_2}}(k)| - |F_G(k)|).$$

Thus we have

$$I_{D(G)}(x) = I_{D(G_1+K_{m_2})}(x) + I_{D(G_2+K_{m_1})}(x) - \sum_{k=0}^{m-1} {\binom{m-1}{k}}^2 x^k.$$

Let s > 2 and assume that the assertion holds for the join of at most s - 1 graphs. Since  $G = (G_1 + \cdots + G_{s-1}) + G_s$ , we have

$$I_{D(G)}(x) = I_{D((G_{1}+\dots+G_{s-1})+K_{m_{s}})}(x) + I_{D(G_{s}+K_{m-m_{s}})}(x) - \sum_{k=0}^{m-1} {\binom{m-1}{k}}^{2} x^{k}$$

$$= \sum_{i=1}^{s-2} I_{D(G_{i}+K_{m-m_{i}})}(x) + I_{D((G_{s-1}+K_{m_{s}})+K_{m-m_{s-1}-m_{s}})}(x)$$

$$-(s-2) \sum_{k=0}^{m-1} {\binom{m-1}{k}}^{2} x^{k} + I_{D(G_{s}+K_{m-m_{s}})}(x) - \sum_{k=0}^{m-1} {\binom{m-1}{k}}^{2} x^{k}$$

$$= \sum_{i=1}^{s} I_{D(G_{i}+K_{m-m_{i}})}(x) - (s-1) \sum_{k=0}^{m-1} {\binom{m-1}{k}}^{2} x^{k}.$$

From Proposition 2.4 and Lemma 2.5, this is equal to equation (3).

# 4. COMPLETE MULTIPARTITE GRAPHS

In this section, applying Theorem 1.1, we give explicit formulas for the  $h^*$ -polynomial and the normalized volume of the PQ-type adjacency polytope of a complete multipartite graph.

Given positive integers  $\ell$  and m, let

$$f_{\ell,m}(x) = \sum_{k=0}^{\ell+m-1} \sum_{\alpha=0}^{k} \binom{\ell-1}{k-\alpha} \binom{m}{\alpha} \sum_{\beta=\alpha}^{k} \binom{\ell+\alpha-1}{\beta} \binom{m-\alpha}{k-\beta} x^{k}.$$

Since

$$\sum_{k=0}^{\ell+m-1} \sum_{\alpha=0}^{k} {\ell-1 \choose k-\alpha} {m \choose \alpha} \sum_{\beta=0}^{k} {\ell+\alpha-1 \choose \beta} {m-\alpha \choose k-\beta} x^{k}$$

$$= \sum_{k=0}^{\ell+m-1} \sum_{\alpha=0}^{k} {\ell-1 \choose k-\alpha} {m \choose \alpha} {\ell+m-1 \choose k} x^{k}$$

$$= \sum_{k=0}^{\ell+m-1} {\ell+m-1 \choose k} \sum_{\alpha=0}^{k} {\ell-1 \choose k-\alpha} {m \choose \alpha} x^{k}$$

$$= \sum_{k=0}^{\ell+m-1} {\ell+m-1 \choose k}^{2} x^{k}$$

holds, we have

$$f_{\ell,m}(x) = \sum_{k=0}^{\ell+m-1} \binom{\ell+m-1}{k}^2 x^k - \sum_{k=1}^{m-1} \sum_{\alpha=1}^k \binom{\ell-1}{k-\alpha} \binom{m}{\alpha} \sum_{\beta=0}^{\alpha-1} \binom{\ell+\alpha-1}{\beta} \binom{m-\alpha}{k-\beta} x^k.$$

The *h*<sup>\*</sup>-polynomial of  $\nabla_G^{PQ}$  for the graph  $G = K_\ell + E_m$  (=  $K_{1,...,1,m}$ ) coincides with  $f_{\ell,m}(x)$ .

**Theorem 4.1.** Let  $G = K_{\ell} + E_m$ . Then we have

$$h^{*}(\nabla_{G}^{\mathrm{PQ}}, x) = f_{\ell,m}(x),$$
  

$$\mathrm{Vol}(\nabla_{G}^{\mathrm{PQ}}) = f_{\ell,m}(1) = \sum_{\alpha=0}^{m} \binom{m}{\alpha} \sum_{\beta=0}^{\ell-1} \binom{\ell+\alpha-1}{\beta} \binom{\ell+m-\alpha-1}{\beta}.$$

*Proof.* Let  $G' = D(K_{\ell-1} + E_m)$ . Since  $G = (K_{\ell-1} + E_m) + K_1$ , from Proposition 2.6, we have

$$h^*(\nabla_G^{\mathrm{PQ}}, x) = p(G', x) = \sum_{k=0}^{\ell+m-1} |\mathrm{PM}(G', k)| x^k.$$

Let  $n = \ell + m - 1$  and let  $[n] \cup [\overline{n}]$  be the vertex set of G'. We decompose [n] into two disjoint set  $V_{1,1} = [\ell - 1]$  and  $V_{1,2} = [n] \setminus [\ell - 1]$  where  $V_{1,1}$  (resp.  $V_{1,2}$ ) corresponds to  $K_{\ell-1}$  (resp.  $E_m$ ). From Proposition 2.8, each |PM(G',k)| is the number of (0,1)-vectors  $(x_1,\ldots,x_n,y_1,\ldots,y_n) \in \mathbb{R}^{2n}$  such that

(12) 
$$\sum_{i=1}^{n} x_i = \sum_{i=1}^{n} y_i = k,$$

(13) 
$$\sum_{i\in S} x_i \leq \sum_{\overline{j}\in \Gamma_{G'}(S)} y_j \text{ for all } S\subset [n].$$

If (12) holds and a subset  $S \subset [n]$  contains an element of  $V_{1,1}$ , then we have  $\Gamma_{G'}(S) = [\overline{n}]$ , and hence

$$\sum_{\overline{j}\in\Gamma_{G'}(S)} y_j = k = \sum_{i=1}^n x_i \ge \sum_{i\in S} x_i.$$

Thus (12) and (13) hold if and only if

(14) 
$$\sum_{i=1}^{n} x_i = \sum_{i=1}^{n} y_i = k,$$

(15) 
$$\sum_{i\in S} x_i \leq \sum_{\overline{j}\in \Gamma_{G'}(S)} y_j \text{ for all } S\subset V_{1,2}.$$

We count such vectors with  $|\{i \in V_{1,2} : x_i = 1\}| = \alpha$  for each  $\alpha = 0, 1, ..., k$ . There are  $\binom{m}{\alpha}$  possibilities for the choice of the subset  $S' = \{i \in V_{1,2} : x_i = 1\} \subset V_{1,2}$ . For each S', there are  $\binom{\ell-1}{k-\alpha}$  possibilities for the choice of the subset  $\{i \in V_{1,1} : x_i = 1\} \subset V_{1,1}$ . Then equations (14) and (15) hold if and only if  $\sum_{i=1}^{n} y_i = k$  and  $\alpha \leq \sum_{\overline{j} \in \Gamma_{G'}(S')} y_j$ . Let  $\beta = \sum_{\overline{j} \in \Gamma_{G'}(S')} y_j$ . Since  $|\Gamma_{G'}(S')| = \ell + \alpha - 1$ , there are  $\binom{\ell+\alpha-1}{\beta}$  possibilities for the choice of the subset  $S'' = \{\overline{j} \in \Gamma_{G'}(S') : y_j = 1\} \subset \Gamma_{G'}(S')$ . For each S'', there are  $\binom{m-\alpha}{k-\beta}$  possibilities for the choice of the subset  $\{\overline{j} \in [\overline{n}] \setminus \Gamma_{G'}(S') : y_j = 1\} \subset [\overline{n}] \setminus \Gamma_{G'}(S')$ . Thus we have  $I_{D(G)}(x) = f_{\ell,m}(x)$ .

Moreover, the normalized volume of  $\nabla_G^{PQ}$  is equal to

$$\begin{split} f_{\ell,m}(1) &= \sum_{k=0}^{\ell+m-1} \sum_{\alpha=0}^{k} \binom{\ell-1}{k-\alpha} \binom{m}{\alpha} \sum_{\beta=\alpha}^{k} \binom{\ell+\alpha-1}{\beta} \binom{m-\alpha}{k-\beta} \\ &= \sum_{\alpha=0}^{m} \binom{m}{\alpha} \sum_{\beta=\alpha}^{\ell+\alpha-1} \binom{\ell+\alpha-1}{\beta} \sum_{k=\beta}^{\ell+\alpha-1} \binom{\ell-1}{k-\alpha} \binom{m-\alpha}{k-\beta} \\ &= \sum_{\alpha=0}^{m} \binom{m}{\alpha} \sum_{\beta=\alpha}^{\ell+\alpha-1} \binom{\ell+\alpha-1}{\beta} \sum_{k=\beta}^{\ell+\alpha-1} \binom{\ell-1}{\ell-1+\alpha-\beta-(k-\beta)} \binom{m-\alpha}{k-\beta} \\ &= \sum_{\alpha=0}^{m} \binom{m}{\alpha} \sum_{\beta=\alpha}^{\ell+\alpha-1} \binom{\ell+\alpha-1}{\beta} \binom{\ell+m-\alpha-1}{\ell+\alpha-\beta-1} \\ &= \sum_{\alpha=0}^{m} \binom{m}{\alpha} \sum_{\beta'=0}^{\ell-1} \binom{\ell+\alpha-1}{\beta'} \binom{\ell+m-\alpha-1}{\beta'}. \end{split}$$

**Remark 4.2.** Let  $G = K_{\ell} + E_m$ . Since

$$\sum_{\alpha=0}^{m} \binom{m}{\alpha} \sum_{\beta=0}^{\ell+\alpha-1} \binom{\ell+\alpha-1}{\beta} \binom{\ell+m-\alpha-1}{\beta} = \binom{2(\ell+m-1)}{\ell+m-1},$$

we have

$$\operatorname{Vol}(\nabla_G^{\operatorname{PQ}}) = \binom{2(\ell+m-1)}{\ell+m-1} - \sum_{\alpha=1}^{m-1} \binom{m}{\alpha} \sum_{\beta=\ell}^{\ell+\alpha-1} \binom{\ell+\alpha-1}{\beta} \binom{\ell+m-\alpha-1}{\beta}.$$

**Example 4.3.** For small  $\ell$  and m,  $f_{\ell,m}(1)$  in Theorem 4.1 is

$$\begin{split} f_{1,m}(1) &= 2^{m}, \\ f_{2,m}(1) &= 2^{m-2} \left( m^{2} + 3m + 8 \right), \\ f_{3,m}(1) &= \frac{2^{m-4}}{(2!)^{2}} \left( m^{4} + 10m^{3} + 59m^{2} + 186m + 384 \right), \\ f_{4,m}(1) &= \frac{2^{m-6}}{(3!)^{2}} \left( m^{6} + 21m^{5} + 229m^{4} + 1563m^{3} + 7762m^{2} + 24984m + 46080 \right), \\ f_{\ell,1}(1) &= \binom{2\ell}{\ell}, \\ f_{\ell,1}(1) &= \binom{2\ell}{\ell}, \\ f_{\ell,2}(1) &= \binom{2(\ell+1)}{\ell+1} - 2, \\ f_{\ell,3}(1) &= \binom{2(\ell+2)}{\ell+2} - (6\ell+6), \\ f_{\ell,4}(1) &= \binom{2(\ell+3)}{\ell+3} - (10\ell^{2} + 24\ell + 20). \end{split}$$

From Theorems 1.1 and 4.1, we have the following.

**Corollary 4.4.** Let G be a complete multipartite graph  $K_{m_1,...,m_s}$ , and let  $m = \sum_{i=1}^{s} m_i$ . Then we have

$$h^{*}(\nabla_{G}^{PQ}, x) = \sum_{i=1}^{s} f_{m-m_{i},m_{i}}(x) - (s-1) \sum_{k=0}^{m-1} {\binom{m-1}{k}}^{2} x^{k},$$
  

$$Vol(\nabla_{G}^{PQ}) = \sum_{i=1}^{s} f_{m-m_{i},m_{i}}(1) - (s-1) {\binom{2(m-1)}{m-1}}.$$

**Example 4.5.** From Corollary 4.4, the normalized volume of  $\nabla_{K_{\ell,m}}^{PQ}$  is

$$f_{\ell,m}(1) + f_{m,\ell}(1) - {2(\ell+m-1) \choose \ell+m-1}.$$

From Example 4.3, we have

$$\begin{aligned} \operatorname{Vol}(\nabla_{K_{1,m}}^{\operatorname{PQ}}) &= 2^{m}, \\ \operatorname{Vol}(\nabla_{K_{2,m}}^{\operatorname{PQ}}) &= 2^{m-2} \left(m^{2} + 3m + 8\right) - 2, \\ \operatorname{Vol}(\nabla_{K_{3,m}}^{\operatorname{PQ}}) &= \frac{2^{m-4}}{(2!)^{2}} \left(m^{4} + 10m^{3} + 59m^{2} + 186m + 384\right) - (6m + 6), \\ \operatorname{Vol}(\nabla_{K_{4,m}}^{\operatorname{PQ}}) &= \frac{2^{m-6}}{(3!)^{2}} \left(m^{6} + 21m^{5} + 229m^{4} + 1563m^{3} + 7762m^{2} + 24984m + 46080\right) \\ &- (10m^{2} + 24m + 20). \end{aligned}$$

The formula for Vol( $\nabla_{K_{2,m}}^{PQ}$ ) coincides with that in [8, Proposition 4.2].

**Example 4.6.** Let *G* be the complete bipartite graph  $K_{2,n-2}$ . Since

$$f_{2,m}(x) = \sum_{k=0}^{m+1} \sum_{\alpha=0}^{k} {\binom{1}{k-\alpha} \binom{m}{\alpha}} \sum_{\beta=\alpha}^{k} {\binom{\alpha+1}{\beta} \binom{m-\alpha}{k-\beta}} x^{k}$$

$$= \sum_{k=0}^{m+1} {\binom{m}{k-1}} k(m-k+1)x^{k} + \sum_{k=0}^{m+1} {\binom{m}{k-1}} x^{k} + \sum_{k=0}^{m+1} {\binom{m}{k}} (k+1)x^{k}$$

$$= \sum_{k=0}^{m} {\binom{m}{k}} k^{2}x^{k} + \sum_{k=0}^{m} {\binom{m}{k}} x^{k+1} + \sum_{k=0}^{m} {\binom{m}{k}} (k+1)x^{k}$$

$$= \sum_{k=0}^{m} {\binom{m}{k}} (x+1+k^{2}+k)x^{k}$$

$$= (x+1)^{m+1} + m((m+1)x+2)x(x+1)^{m-2}$$

and

$$f_{\ell,2}(x) = \sum_{k=0}^{\ell+1} {\ell+1 \choose k}^2 x^k - 2x,$$

we have

$$h^*(\nabla_G^{PQ}, x) = f_{2,n-2}(x) + f_{n-2,2}(x) - \sum_{k=0}^{n-1} \binom{n-1}{k}^2 x^k$$
  
=  $(x+1)^{n-1} + (n-2)((n-1)x+2)x(x+1)^{n-4} - 2x$ .

## 5. WHEEL GRAPHS

For  $n \ge 3$ , the wheel graph  $W_n$  with n + 1 vertices is the join graph  $W_n = C_n + K_1$ . Unfortunately, Theorem 1.1 is not useful for computing the  $h^*$ -polynomial of  $\nabla_{W_n}^{PQ}$ . We will give an explicit formula for the  $h^*$ -polynomial of  $\nabla_{W_n}^{PQ}$  and prove the conjecture [8, Conjecture 4.4] on the normalized volume of  $\nabla_{W_n}^{PQ}$  by using Proposition 2.6 on  $\nabla_{G+K_1}^{PQ}$ .

Let

$$\gamma(n,x) = \begin{cases} 1 & \text{if } n = 0, \\ \frac{(1+\sqrt{1+8x})^n + (1-\sqrt{1+8x})^n}{2^n} & \text{if } n \text{ is odd,} \\ \frac{(1+\sqrt{1+8x})^n + (1-\sqrt{1+8x})^n}{2^n} - 2x^{\frac{n}{2}} & \text{otherwise.} \end{cases}$$

For  $n \ge 3$ ,

$$\frac{(1+\sqrt{1+8x})^n + (1-\sqrt{1+8x})^n}{2^n} = g(C_n, 2x),$$

where  $g(C_n, x)$  is the matching generating polynomial of  $C_n$ . Moreover, for  $n \ge 3$ , it is known [16, Example 4.5] that,  $\gamma(n, x)$  is the  $\gamma$ -polynomial of the PV-type adjacency polytope  $\nabla_{W_n}^{PV}$  of  $W_n$ . (Note that  $\nabla_{W_n}^{PV}$  is called *the symmetric edge polytope of type A* of  $W_n$  in [16].) The  $h^*$ -polynomial of  $\nabla_{W_n}^{PQ}$  is described by this function as follows.

**Theorem 5.1.** The  $h^*$ -polynomial of  $\nabla_{W_n}^{PQ}$  is

$$\sum_{k=0}^{n} \binom{n}{k} \gamma(n-k,x) x^{k}$$

$$= \left(\frac{1+2x+\sqrt{1+8x}}{2}\right)^{n} + \left(\frac{1+2x-\sqrt{1+8x}}{2}\right)^{n} + x^{n} - (x+\sqrt{x})^{n} - (x-\sqrt{x})^{n}.$$
PO

Moreover, the normalized volume of  $\nabla_{W_n}^{PQ}$  is  $3^n - 2^n + 1$ .

*Proof.* Since  $W_n = C_n + K_1$ , the  $h^*$ -polynomial of  $\nabla_{W_n}^{PQ}$  is

$$p(D(C_n), x) = \sum_{\ell=0}^n |\mathrm{PM}(D(C_n), \ell)| x^{\ell}$$

by Proposition 2.6. From Proposition 2.8,  $|PM(D(C_n), \ell)|$  is equal to the number of (0, 1)-vectors  $(x_1, \ldots, x_n, y_1, \ldots, y_n) \in \mathbb{R}^{2n}$  such that

(16) 
$$\sum_{i=1}^{n} x_i = \sum_{j=1}^{n} y_j = \ell,$$

(17) 
$$\sum_{i \in S} x_i \leq \sum_{\overline{j} \in \Gamma_{G'}(S)} y_j \text{ for all } S \subset [n],$$

where  $V = [n] \cup [\overline{n}]$  is the set of vertices of  $D(C_n)$ . Let  $C_n = (1, 2, ..., n, 1)$ . Given a subset  $T \subset [n]$ and an integer  $\ell \in [n]$ , let  $PM_{T,\ell}$  denote the set of all (0,1) vectors  $(x_1, ..., x_n, y_1, ..., y_n) \in \mathbb{R}^{2n}$ satisfying (16), (17), and  $T = \{i \in [n] : x_i = y_i = 1\}$ . Note that  $PM_{T,\ell} = \emptyset$  if  $\ell < |T|$ . Let  $T \subset [n]$  with |T| = k. We will show that  $\sum_{\ell=k}^{n} |PM_{T,\ell}| x^{\ell} = \gamma(n-k, x) x^k$ .

**Case 1** (k = n). It is easy to see that  $PM_{T,n} = \{(1, ..., 1)\}$  and  $PM_{T,\ell} = \emptyset$  if  $\ell \neq n$ . Note that  $\gamma(n-k,x)x^k = x^n$  if k = n. Thus  $\sum_{\ell=k}^n |PM_{T,\ell}|x^\ell = \gamma(n-k,x)x^k$ .

**Case 2** (k = n - 1). Let  $T = [n] \setminus \{i\}$  where  $i \in [n]$ . It then follows that  $PM_{T,n-1} = \{(1, \dots, 1) - \mathbf{e}_i - \mathbf{e}_{n+i}\}$  and  $PM_{T,\ell} = \emptyset$  if  $\ell \neq n-1$ . Note that  $\gamma(n-k,x)x^k = x^{n-1}$  if k = n-1. Thus  $\sum_{\ell=k}^n |PM_{T,\ell}|x^\ell = \gamma(n-k,x)x^k$ .

**Case 3** (k = n - 2). Let  $T = [n] \setminus \{i, j\}$  where  $1 \le i < j \le n$ . Since (16) holds, each element of  $PM_{T,\ell}$  is

$$\alpha_1 = (1,\ldots,1) - \mathbf{e}_i - \mathbf{e}_j - \mathbf{e}_{n+i} - \mathbf{e}_{n+j}$$

if  $\ell = n - 2$ , and is one of

$$\alpha_2 = (1,\ldots,1) - \mathbf{e}_i - \mathbf{e}_{n+j}, \alpha_3 = (1,\ldots,1) - \mathbf{e}_j - \mathbf{e}_{n+i},$$

if  $\ell = n - 1$ . Then each  $\alpha_i$  corresponds to a perfectly matchable set. In fact, a matching of  $D(C_n)$  which corresponds to  $\alpha_1$ ,  $\alpha_2$ ,  $\alpha_3$  is

$$\{\{s,\bar{s}\}: s \in [n] \setminus \{i,j\}\}, \\ \{\{s,\bar{s}\}: 1 \le s \le i-1 \text{ or } j+1 \le s \le n\} \cup \{\{s+1,\bar{s}\}: i \le s \le j-1\}, \\ \{\{s,\bar{s}\}: 1 \le s \le i-1 \text{ or } j+1 \le s \le n\} \cup \{\{s,\bar{s}+1\}: i \le s \le j-1\}, \\ 15 \end{cases}$$

respectively. Thus

$$PM_{T,\ell} = \begin{cases} \{\alpha_1\} & \text{if } \ell = n-2, \\ \{\alpha_2, \alpha_3\} & \text{if } \ell = n-1, \\ \emptyset & \text{otherwise.} \end{cases}$$

Note that  $\gamma(n-k,x)x^k = (2x+1)x^{n-2} = x^{n-2} + 2x^{n-1}$  if k = n-2. Thus  $\sum_{\ell=k}^n |PM_{T,\ell}|x^\ell = \gamma(n-k,x)x^k$ .

**Case 4**  $(k \le n-3)$ . Suppose that  $T' = \{p, p+1, \dots, q\} \subset T$  and  $p-1, q+1 \notin T$ . Since  $|T| \le n-3$ , we have  $n-q+p-1 \ge 3$ . Let  $U = \{p, p+1, \dots, q, \overline{p}, \overline{p+1}, \dots, \overline{q}\}$ . We will show that there exists a perfectly matchable set *S* of  $D(C_{n-q+p-1})$  such that

$$\rho(S) = (x_1, \dots, x_{p-1}, x_{q+1}, \dots, x_n, y_1, \dots, y_{p-1}, y_{q+1}, \dots, y_n)$$

for any  $(x_1,...,x_n,y_1,...,y_n) \in PM_{T,\ell}$ . Here  $C_{n-q+p-1} = (1,...,p-1,q+1,...,n,1)$  and the vertex set of  $D(C_{n-q+p-1})$  is  $\{1,...,p-1,q+1,...,n\} \cup \{\overline{1},...,\overline{p-1},\overline{q+1},...,\overline{n}\}$ . Let *M* be a matching of  $D(C_n)$  which corresponds to  $(x_1,...,x_n,y_1,...,y_n)$ .

**Case 4.1** (either  $x_{p-1} = x_{q+1} = 0$  or  $y_{p-1} = y_{q+1} = 0$ ). Exchanging [n] and  $[\overline{n}]$  if needed, we may assume that  $y_{p-1} = y_{q+1} = 0$ . Then, for the subset  $T' \subset [n]$ ,

$$\sum_{i \in T'} x_i = \sum_{\overline{j} \in \Gamma_{G'}(T')} y_j = |T'|.$$

Hence the matching M is the union of a perfect matching of the induced subgraph  $D(C_n)_U$  of  $D(C_n)$  and a matching M' of the induced subgraph  $D(C_n)_{V\setminus U}$  of  $D(C_n)$ . Then one can regard M' as a matching of  $D(C_{n-q+p-1})$  since  $D(C_n)_{V\setminus U}$  is a subgraph of  $D(C_{n-q+p-1})$ .

**Case 4.2**  $(x_{p-1} = y_{q+1} \neq y_{p-1} = x_{q+1})$ . Exchanging [n] and  $[\overline{n}]$  if needed, we may assume that  $x_{p-1} = y_{q+1} = 1$  and  $y_{p-1} = x_{q+1} = 0$ . If  $e = \{q+2, \overline{q+1}\}$  belongs to the matching M, then  $M \setminus \{e\}$  is the union of a perfect matching of  $D(C_n)_U$  and a matching M' of  $D(C_n)_{V\setminus U}$  by the same argument in Case 4.1. Suppose that  $e = \{q, \overline{q+1}\}$  belongs to the matching M. It then follows that M is the union of  $\{\{p-1, \overline{p}\}, \dots, \{q-1, \overline{q}\}, \{q, \overline{q+1}\}\}$  and a matching M' of  $D(C_n)_{V\setminus U}$ . Thus one can regard  $M' \cup \{\{p-1, \overline{q+1}\}\}$  as a matching of  $D(C_{n-q+p-1})$ .

Thus there exists a perfectly matchable set  $S_1$  of  $D(C_{n-q+p-1})$  such that

$$\rho(S_1) = (x_1, \dots, x_{p-1}, x_{q+1}, \dots, x_n, y_1, \dots, y_{p-1}, y_{q+1}, \dots, y_n)$$

for  $(x_1, \ldots, x_n, y_1, \ldots, y_n)$  if  $T' = \{p, p+1, \ldots, q\} \subset T$  and  $p-1, q+1 \notin T$ . If, in addition,  $T'' = \{p', p'+1, \ldots, q'\} \subset T$  and  $p'-1, q'+1 \notin T$  for some p' > q+1, then there exists a perfectly matchable set  $S_2$  of  $D(C_{n-(q-p+1)-(q'-p'+1)})$  such that

$$\rho(S_2) = (x_1, \dots, x_{p-1}, x_{q+1}, \dots, x_{p'-1}, x_{q'+1}, \dots, x_n, y_1, \dots, y_{p-1}, y_{q+1}, \dots, y_{p'-1}, y_{q'+1}, \dots, y_n)$$

by the same argument as above. Repeating the above argument, it follows that there exists a perfectly matchable set *S* of  $D(C_{n-k})$  such that  $\rho(S) = (x_{i_1}, \dots, x_{i_{n-k}}, y_{i_1}, \dots, y_{i_{n-k}})$  where  $[n] \setminus T = \{i_1, \dots, i_{n-k}\}$  and  $x_{i_r} + y_{i_r} \leq 1$  for all *r*. Then there exists a perfectly matchable set *S'* of  $C_{n-k}$  such that  $|S'| = 2(\ell - k)$  and  $\rho(S') = (x_{i_1} + y_{i_1}, \dots, x_{i_{n-k}} + y_{i_{n-k}})$ . The matching corresponding to *S'* is

not unique if and only if n - k is even and  $\rho(S') = (1, ..., 1)$ . There exist two matchings for the perfectly matchable set *S* of  $D(C_{n-k})$  exactly when

$$(x_{i_1},\ldots,x_{i_{n-k}},y_{i_1},\ldots,y_{i_{n-k}}) = (1,0,\ldots,1,0,0,1,\ldots,0,1), (0,1,\ldots,0,1,1,0,\ldots,1,0).$$

Conversely, for each matching M of  $C_{n-k}$ , there exist  $2^{|M|}$  vectors  $(x_{i_1}, \ldots, x_{i_{n-k}}, y_{i_1}, \ldots, y_{i_{n-k}})$  where  $x_{i_r} + y_{i_r} \leq 1$  for all r and associated with a perfectly matchable set of  $D(C_{n-k})$  since there are two possibilities  $(x_{i_p}, x_{i_{p+1}}, y_{i_p}, y_{i_{p+1}}) = (1, 0, 0, 1), (0, 1, 1, 0)$  for each  $\{i_p, i_{p+1}\} \in M$ . Hence we have

$$|PM_{T,\ell}| = \begin{cases} 2^{\ell-k} m_{C_{n-k}}(\ell-k) - 2 & \text{if } \ell - k = (n-k)/2, \\ 2^{\ell-k} m_{C_{n-k}}(\ell-k) & \text{otherwise.} \end{cases}$$

Recall that  $m_G(k)$  is the number of k-matchings of a graph G. Thus

$$\sum_{\ell=k}^{n} |\mathbf{PM}_{T,\ell}| x^{\ell} = \sum_{\ell=k}^{n} 2^{\ell-k} m_{C_{n-k}} (\ell-k) x^{\ell} - 2x^{(n+k)/2}$$
$$= \left( \sum_{\ell=k}^{n} m_{C_{n-k}} (\ell-k) (2x)^{\ell-k} - 2x^{(n-k)/2} \right) x^{k}$$
$$= \gamma(n-k,x) x^{k}$$

if n - k is even, and

$$\begin{split} \sum_{\ell=k}^{n} |\mathbf{PM}_{T,\ell}| x^{\ell} &= \sum_{\ell=k}^{n} 2^{\ell-k} m_{C_{n-k}}(\ell-k) x^{\ell} \\ &= \left( \sum_{\ell=k}^{n} m_{C_{n-k}}(\ell-k) (2x)^{\ell-k} \right) x^{k} \\ &= \gamma(n-k,x) x^{k} \end{split}$$

if n - k is odd.

Therefore the  $h^*$ -polynomial of  $\nabla_{W_n}^{PQ}$  is

$$\sum_{T \subset [n]} \sum_{\ell=|T|}^{n} |\mathrm{PM}_{T,\ell}| x^{\ell} = \sum_{T \subset [n]} \gamma(n-|T|) x^{|T|} = \sum_{k=0}^{n} \binom{n}{k} \gamma(n-k,x) x^{k}.$$

Moreover

$$\begin{split} &\sum_{k=0}^{n} \binom{n}{k} \gamma(n-k,x) x^{k} \\ &= x^{n} + \sum_{k=0}^{n} \binom{n}{k} \left(\frac{1+\sqrt{1+8x}}{2}\right)^{n-k} x^{k} + \sum_{k=0}^{n} \binom{n}{k} \left(\frac{1-\sqrt{1+8x}}{2}\right)^{n-k} x^{k} \\ &- \sum_{\substack{k \ge 0 \\ k \equiv n \pmod{2}}} 2\binom{n}{k} \sqrt{x}^{n-k} x^{k} \\ &= \left(\frac{1+2x+\sqrt{1+8x}}{2}\right)^{n} + \left(\frac{1+2x-\sqrt{1+8x}}{2}\right)^{n} + x^{n} - (x+\sqrt{x})^{n} - (x-\sqrt{x})^{n}. \end{split}$$

In particular, substituting x = 1, we have

$$\sum_{k=0}^{n} \binom{n}{k} \gamma(n-k,1) = 3^{n} - 2^{n} + 1$$

### ACKNOWLEDGEMENT

The authors were partially supported by JSPS KAKENHI 18H01134, 19K14505, and 19J00312.

### REFERENCES

- F. Ardila, M. Back, S, Hoşten, J. Pfeifle and K. Seashore, Root Polytopes and Growth Series of Root Lattices, SIAM J. Discrete Math. 25 (2011), 360–378.
- [2] E. Balas and W.R. Pulleyblank, The perfectly matchable subgraph polytopes of a bipartite graph, *Networks* **13** (1983), 495–516.
- [3] E. Balas and W.R. Pulleyblank, The perfectly matchable subgraph polytope of an arbitrary graph, *Combinatorica* **9** (1989), 321–337.
- [4] M. Beck and S. Robins, "Computing the continuous discretely", Undergraduate Texts in Mathematics, Springer, second edition, 2015.
- [5] T. Chen, R. Davis and D. Mehta, Counting equilibria of the Kuramoto model using birationally invariant intersection index, *SIAM J. Appl. Algebra Geometry* **2** (2018), 489–507.
- [6] T. Chen and D. Mehta, On the network topology dependent solution count of the algebraic load flow equations, *IEEE Transactions on Power Systems* **33** (2018), 1451–1460.
- [7] A. D'Alì, E. Delucchi and M. Michałek, Many faces of symmetric edge polytopes, preprint, arXiv: 1910.05193
- [8] R. Davis and T. Chen, Computing volumes of adjacency polytopes via draconian sequences, preprint, arXiv:2007.11051.
- [9] E. Ehrhart, "Polynomês Arithmétiques et Méthode des Polyédres en Combinatorie", Birkhäuser, Boston/ Basel/ Stuttgart, 1977.
- [10] A. Higashitani, K. Jochemko and M. Michałek, Arithmetic aspects of symmetric edge polytopes, *Mathematika*, 65 (2019), 763–784.
- [11] T. Kálmán, A version of Tutte's polynomial for hypergraphs, Adv. Math. 244 (2013), 823–873.
- [12] T. Kálmán and A. Postnikov, Root polytopes, Tutte polynomials, and a duality theorem for bipartite graphs, *Proc. Lond. Math. Soc.* **114** (2017), 561–588.
- [13] Y. Kuramoto, Self-entrainment of a population of coupled non-linear oscillators, in International Symposium on Mathematical Problems in Theoretical Physics (Kyoto Univ., Kyoto, 1975), 1975, pp. 420–422. Lecture Notes in Phys., 39.
- [14] H. Ohsugi and A. Tsuchiya, Reflexive polytopes arising from bipartite graphs with  $\gamma$ -positivity associated to interior polynomials, *Selecta Math.* (*N.S.*) **26** (2020), 59.
- [15] H. Ohsugi and A. Tsuchiya, The  $h^*$ -polynomials of locally anti-blocking lattice polytopes and their  $\gamma$ -positivity, *Discrete Comput. Geom.* (2020), published online.
- [16] H. Ohsugi and A. Tsuchiya, Symmetric edge polytopes and matching generating polynomials, preprint, arXiv:2008.08621.
- [17] A. Postnikov, Permutohedra, associahedra, and beyond, Int. Math. Res. Notices 6 (2009), 1026–1106.

HIDEFUMI OHSUGI, DEPARTMENT OF MATHEMATICAL SCIENCES, SCHOOL OF SCIENCE, KWANSEI GAKUIN UNIVERSITY, SANDA, HYOGO 669-1337, JAPAN

Email address: ohsugi@kwansei.ac.jp

AKIYOSHI TSUCHIYA, GRADUATE SCHOOL OF MATHEMATICAL SCIENCES, UNIVERSITY OF TOKYO, KOMABA, MEGURO-KU, TOKYO 153-8914, JAPAN

Email address: akiyoshi@ms.u-tokyo.ac.jp