

Parameterized discrete uniformization theorems and curvature flows for polyhedral surfaces, II

Xu Xu, Chao Zheng

October 5, 2021

Abstract

This paper investigates the combinatorial α -curvature for vertex scaling of piecewise hyperbolic metrics on polyhedral surfaces, which is a parameterized generalization of the classical combinatorial curvature. A discrete uniformization theorem for combinatorial α -curvature is established, which generalizes Gu-Guo-Luo-Sun-Wu's discrete uniformization theorem for classical combinatorial curvature [16]. We further introduce combinatorial α -Yamabe flow and combinatorial α -Calabi flow for vertex scaling to find piecewise hyperbolic metrics with prescribed combinatorial α -curvatures. To handle the potential singularities along the combinatorial curvature flows, we do surgery along the flows by edge flipping. Using the discrete conformal theory established by Gu-Guo-Luo-Sun-Wu [16], we prove the longtime existence and convergence of combinatorial α -Yamabe flow and combinatorial α -Calabi flow with surgery, which provide effective algorithms for finding piecewise hyperbolic metrics with prescribed combinatorial α -curvatures.

Keywords: Discrete uniformization theorem; Piecewise hyperbolic metric; Combinatorial curvature; Combinatorial Yamabe flow; Combinatorial Calabi flow

1 Introduction

This is a continuation of [27], in which the first author established a parameterized discrete uniformization theorem for combinatorial α -curvature for vertex scaling of piecewise linear metrics (PL metrics for short in the following) and studied the longtime behavior of combinatorial α -Yamabe flow and combinatorial α -Calabi flow for vertex scaling of PL metrics by surgery. In this paper, we establish the counterpart results for combinatorial α -curvature for vertex scaling of piecewise hyperbolic metrics (PH metrics for short in the following) on surfaces.

Suppose S is a connected closed surface and V is a finite subset of S with $|V| = N$, (S, V) is called a marked surface. A PH metric on the marked surface (S, V) is a

hyperbolic metric on S with conic singularities contained in V . Suppose $\mathcal{T} = (V, E, F)$ is a triangulation of (S, V) , where V, E, F represent the sets of vertices, edges and faces respectively, (S, V, \mathcal{T}) is called a triangulated surface. Denote a vertex, an edge and a face in the triangulation by $v_i, \{v_i v_j\}, \Delta v_i v_j v_k$ respectively, where i, j, k are natural numbers. A PH metric on a triangulated surface (S, V, \mathcal{T}) defines a map $d : E \rightarrow (0, +\infty)$ such that for any triangle $\Delta v_i v_j v_k \in F$, d_{ij}, d_{ik}, d_{jk} satisfy the triangle inequalities, where $d_{st} := d(\{v_s v_t\})$ for $\{v_s v_t\} \in E$. Conversely, by gluing hyperbolic triangles isometrically along the edges in pair, we obtain PH metrics on a triangulated surface (S, V, \mathcal{T}) , which give rise to PH metrics on the marked surface (S, V) . The classical combinatorial curvature $K : V \rightarrow (-\infty, 2\pi)$ is used to describe the conic singularities of PH metrics at the vertices. Suppose (S, V, \mathcal{T}) is a triangulated surface with a PH metric d , the classical combinatorial curvature for d at $v_i \in V$ is defined to be

$$K_i = 2\pi - \sum_{\Delta v_i v_j v_k \in F} \alpha_i^{jk}, \quad (1)$$

where α_i^{jk} is the inner angle at the vertex v_i in the hyperbolic triangle $\Delta v_i v_j v_k \in F$. Note that the combinatorial curvature for the PH metric d is independent of the geometric triangulations of d on (S, V) .

The vertex scaling for PL metric on polyhedral surfaces is a discrete analogy of the conformal change in Riemannian geometry, which was introduced physically by Róček-Williams [23] and mathematically by Luo [19] independently. Motivated by the vertex scaling for PL metrics on polyhedral surfaces, Bobenko-Pinkall-Springborn [1] introduced the following definition of vertex scaling for PH metrics on triangulated surfaces.

Definition 1.1 ([1]). Suppose (S, V, \mathcal{T}) is a connected closed triangulated surface with a PH metric $d : E \rightarrow (0, +\infty)$ and $u : V \rightarrow \mathbb{R}$ is a function defined on the vertices. The vertex scaling of PH metric d by u is defined to be the PH metric $u * d$ such that

$$\sinh \frac{(u * d)_{ij}}{2} = \sinh \frac{d_{ij}}{2} e^{u_i + u_j} \quad (2)$$

for any edge $\{v_i v_j\} \in E$. The function $u : V \rightarrow \mathbb{R}$ is called a discrete conformal factor.

As mentioned in [14], the classical combinatorial curvature K is scaling invariant in the Euclidean background geometry and does not approximate the Gauss curvature on smooth surface as the triangulation of the surface becomes finer and finer. To overcome these disadvantages, Ge and the first author [14] introduced the combinatorial α -curvature for Thurston's Euclidean circle packing metrics. After that, there are lots of studies on combinatorial α -curvature on surfaces and 3-manifolds [4, 8, 9, 11–14, 25–27]. Following [14], we introduce the following combinatorial α -curvature for vertex scaling of PH metrics on triangulated surfaces.

Definition 1.2. Suppose (S, V, \mathcal{T}) is a connected closed triangulated surface with a PH metric d_0 , $\alpha \in \mathbb{R}$ is a constant and $u : V \rightarrow \mathbb{R}$ is a discrete conformal factor for d_0 . The combinatorial α -curvature at vertex $v_i \in V$ is defined to be

$$R_{\alpha,i} = \frac{K_i}{w_i^\alpha}, \quad (3)$$

where $w_i = e^{u_i}$ and K_i is the classical combinatorial curvature at v_i defined by (1).

Remark 1. $R_{\alpha,i}$ is a parameterized generalization of K_i . Specially, if $\alpha = 0$, $R_{0,i}$ is the classical combinatorial curvature K_i . Similar to the classical combinatorial curvature K , the combinatorial α -curvature R_α in Definition 1.2 is independent of the geometric triangulation of the marked surface (S, V) with a PH metric. Therefore, the combinatorial α -curvature is well-defined for PH metrics discrete conformal to d_0 .

One of the remarkable theorems proved by Bobenko-Pinkall-Springborn [1] is that on a triangulated surface with a PH metric, the classical combinatorial curvature determines the discrete conformal factor. We prove the following global rigidity for the combinatorial α -curvature R_α with respect to discrete conformal factors, which is a parameterized generalization of Bobenko-Pinkall-Springborn's global rigidity.

Theorem 1.3. Suppose (S, V, \mathcal{T}) is a connected closed triangulated surface with a PH metric d_0 , $\alpha \in \mathbb{R}$ is a constant and \overline{R} is a function defined on the vertices. If $\alpha \overline{R} \leq 0$, there exists at most one discrete conformal factor u such that the combinatorial α -curvature of $u * d_0$ is \overline{R} .

An important problem on polyhedral surfaces is about the existence of discrete conformal factors with prescribed combinatorial curvatures. Recently, Gu-Luo-Sun-Wu [17], Gu-Guo-Luo-Sun-Wu [16] and Springborn [22] gave nice answers to this problem. As a corollary, Gu-Luo-Sun-Wu [17] and Gu-Guo-Luo-Sun-Wu [16] proved the Euclidean and hyperbolic discrete uniformization theorem for the classical combinatorial curvature on closed marked surfaces respectively. We prove the following result on the existence of discrete conformal factor for PH metrics with prescribed combinatorial α -curvature.

Theorem 1.4. Suppose (S, V) is a connected closed marked surface with a PH metric d_0 , α is a constant and $\overline{\mathbf{F}}$ is a function defined on the vertices. Then there exists a PH metric in the discrete conformal class $\mathcal{D}(d_0)$ with combinatorial α -curvature $\overline{\mathbf{F}}$ if one of the following three conditions is satisfied

- (1) $\alpha > 0$, $\chi(S) < 0$, $\overline{\mathbf{F}} \leq 0$;
- (2) $\alpha < 0$, $\overline{\mathbf{F}} > 0$;

(3) (Gu-Guo-Luo-Sun-Wu [16]) $\alpha = 0$, $\bar{\mathbf{F}} \in (-\infty, 2\pi)$, $\sum_{i=1}^N \bar{\mathbf{F}}_i > 2\pi\chi(S)$.

Here $\mathcal{D}(d_0)$ is the discrete conformal class introduced by Gu-Guo-Luo-Sun-Wu [16]. See also Definition 3.1.

As a corollary of Theorem 1.4, we have the following result on the existence of PH metrics with constant combinatorial α -curvature on connected closed marked surfaces, which is a parameterized generalization of the hyperbolic discrete uniformization theorem obtained by Gu-Guo-Luo-Sun-Wu for the classical combinatorial curvature [16].

Corollary 1.5. Suppose (S, V) is a connected closed marked surface with a PH metric d_0 and α is a constant.

- (1) If $\alpha > 0$ and $\chi(S) < 0$, for any nonpositive constant $\bar{\mathbf{F}}$, there exists a unique PH metric in $\mathcal{D}(d_0)$ with combinatorial α -curvature $\bar{\mathbf{F}}$.
- (2) If $\alpha < 0$, for any positive constant $\bar{\mathbf{F}}$, there exists a unique PH metric in $\mathcal{D}(d_0)$ with combinatorial α -curvature $\bar{\mathbf{F}}$.
- (3) (Gu-Guo-Luo-Sun-Wu [16]) If $\alpha = 0$, for any constant $\bar{\mathbf{F}} \in (\frac{2\pi\chi(S)}{N}, 2\pi)$, there exists a unique PH metric in $\mathcal{D}(d_0)$ with combinatorial α -curvature $\bar{\mathbf{F}}$.

Another important problem on polyhedral surfaces is to find effective algorithms searching for polyhedral metrics (PL metrics and PH metrics) with prescribed combinatorial curvatures, which has lots of applications in practical activities [29]. Chow-Luo's combinatorial Ricci flows for circle packings [2] and Luo's combinatorial Yamabe flow for vertex scaling [19] provide effective algorithms to find polyhedral metrics with prescribed classical combinatorial curvature. Motivated by [2, 12–14, 19, 27], we introduce the following combinatorial α -Yamabe flow for vertex scaling of PH metrics, which is a parameterized generalization of Luo's combinatorial Yamabe flow in [19].

Definition 1.6. Suppose (S, V, \mathcal{T}) is a connected closed triangulated surface with a PH metric d_0 and $\alpha \in \mathbb{R}$ is a constant. The combinatorial α -Yamabe flow is defined to be

$$\begin{cases} \frac{dw_i}{dt} = -R_{\alpha,i}w_i, \\ w_i(0) = 1, \end{cases} \quad (4)$$

where $w_i = e^{u_i}$ and $u : V \rightarrow \mathbb{R}$ is a discrete conformal factor for d_0 .

Remark 2. When $\alpha = 0$, the combinatorial 0-Yamabe flow (4) is Luo's combinatorial Yamabe flow in [19]. One can further generalize the combinatorial α -Yamabe flow (4) to the following form to find PH metrics with prescribed combinatorial α -curvature \bar{R}

$$\frac{dw_i}{dt} = (\bar{R}_i - R_{\alpha,i})w_i. \quad (5)$$

If $\bar{R}_i = 0$ for all $i \in V$, (5) is reduced to (4).

Combinatorial Calabi flow is another effective combinatorial curvature flow to find polyhedral metrics with prescribed combinatorial curvature. The combinatorial Calabi flow for Thurston's circle packing metrics and for vertex scaling of PL metrics was introduced by Ge [5, 6] and the combinatorial Calabi flow for vertex scaling of PH metrics was introduced by Zhu and the first author [30]. Motivated by [5–7, 9, 12, 14, 27, 30], we introduce the following combinatorial α -Calabi flow for vertex scaling of PH metrics.

Definition 1.7. Suppose (S, V, \mathcal{T}) is a connected closed triangulated surface with a PH metric d_0 and $\alpha \in \mathbb{R}$ is a constant. The combinatorial α -Calabi flow is defined to be

$$\begin{cases} \frac{dw_i}{dt} = \Delta_\alpha^\mathcal{T} R_{\alpha,i} w_i, \\ w_i(0) = 1, \end{cases} \quad (6)$$

where $w_i = e^{u_i}$, $u : V \rightarrow \mathbb{R}$ is a discrete conformal factor for d_0 and the discrete α -Laplace operator $\Delta_\alpha^\mathcal{T}$ is defined by

$$\Delta_\alpha^\mathcal{T} f_i = -\frac{1}{w_i^\alpha} \sum_{j \in V} \frac{\partial K_i}{\partial u_j} f_j$$

for $f : V \rightarrow \mathbb{R}$.

Remark 3. When $\alpha = 0$, the combinatorial 0-Calabi flow (6) is the combinatorial Calabi flow introduced by Zhu and the first author [30]. One can also use the following modified combinatorial α -Calabi flow to study the prescribed combinatorial α -curvature problem

$$\frac{dw_i}{dt} = \Delta_\alpha^\mathcal{T} (R_\alpha - \bar{R})_i w_i, \quad (7)$$

where \bar{R} is a function defined on the vertices. If $\bar{R}_i = 0$ for all $i \in V$, (7) is reduced to (6).

Along the combinatorial α -flows (combinatorial α -Yamabe flow and combinatorial α -Calabi flow), singularities may develop, which correspond to that some triangles degenerate or discrete conformal factors tend to infinity. To handle the potential singularities along the combinatorial α -flows, we do the following surgery on combinatorial α -flows by flipping, the idea of which comes from [16, 17, 19]. Recall that a Delaunay triangulation \mathcal{T} for a PH metric d on a marked surface (S, V) is a geometric triangulation of (S, V, d) such that for any adjacent triangles $\Delta v_i v_j v_k, \Delta v_i v_j v_l \in F$ sharing an edge $\{v_i v_j\} \in E$,

$$\alpha_l^{ij} + \alpha_k^{ij} \leq \alpha_i^{jk} + \alpha_j^{ik} + \alpha_i^{jl} + \alpha_j^{il}. \quad (8)$$

Please refer to [16, 18] for more information on Delaunay triangulations for PH metrics on marked surfaces. There is always a Delaunay triangulation \mathcal{T} for a marked surface (S, V) with a PH metric d [16]. Along the combinatorial α -flows on a triangulated surface

(S, V, \mathcal{T}) , if \mathcal{T} is Delaunay in $u(t) * d_0$ for $t \in [0, T]$ and not Delaunay in $u(t) * d_0$ for $t \in (T, T + \epsilon)$, $\epsilon > 0$, there exists an edge $\{v_i v_j\} \in E$ such that $\alpha_i^{jk} + \alpha_j^{ik} + \alpha_i^{jl} + \alpha_j^{il} \geq \alpha_i^{ij} + \alpha_k^{ij}$ for $t \in [0, T]$ and $\alpha_i^{jk} + \alpha_j^{ik} + \alpha_i^{jl} + \alpha_j^{il} < \alpha_i^{ij} + \alpha_k^{ij}$ for $t \in (T, T + \epsilon)$. Then we replace the triangulation \mathcal{T} by a new triangulation \mathcal{T}' at time $t = T$ via replacing two triangles $\Delta v_i v_j v_k$ and $\Delta v_i v_j v_l$ adjacent to $\{v_i v_j\}$ by two new triangles $\Delta v_i v_k v_l$ and $\Delta v_j v_k v_l$. This is called a **surgery by flipping** on the triangulation \mathcal{T} , which is also an isometry of (S, V) with PH metric $u(T) * d_0$. After the surgery by flipping at time $t = T$, we run the combinatorial α -flows on (S, V, \mathcal{T}') with initial metric $u(T) * d_0$. Whenever the Delaunay condition (8) is not satisfied along the combinatorial α -flows, we do surgery by flipping along the combinatorial α -flows.

Theorem 1.8. Suppose (S, V) is a connected closed marked surface with a PH metric d_0 , α is a constant and $\bar{\mathbf{F}}$ is a function defined on the vertices. If one of the conditions (1)(2)(3) in Theorem 1.4 is satisfied, the solution of combinatorial α -Yamabe flow and combinatorial α -Calabi flow with surgery exist for all time and converge exponentially fast.

Remark 4. If $\alpha = 0$, the longtime existence and convergence of hyperbolic combinatorial Yamabe flow (0-Yamabe flow) with surgery was proved by Gu-Guo-Luo-Sun-Wu [16] and the longtime existence and convergence of hyperbolic combinatorial Calabi flow (0-Calabi flow) with surgery was proved by Zhu and the first author [30]. Theorem 1.8 is a parameterized generalization of these results.

The paper is organized as follows. In Section 2, we study the combinatorial α -curvature on a marked surface with a fixed triangulation and prove Theorem 1.3. We also discuss the stability of combinatorial α -flows on triangulated surfaces in Section 2. In Section 3, we allow the triangulation on the marked surface to be changed by edge flipping and prove Theorem 1.4 and Theorem 1.8.

Acknowledgements

The authors thank Dr. Tianqi Wu for communications on the details of [16, 17]. The research of the first author is supported by the Fundamental Research Funds for the Central Universities under grant no. 2042020kf0199.

2 Combinatorial α -curvature and combinatorial α -flows on a marked surface with a fixed triangulation

2.1 Rigidity of combinatorial α -curvature on triangulated surfaces

Suppose (S, V) is a marked surface with a fixed triangulation \mathcal{T} , $\Delta v_i v_j v_k \in F$ is a triangle and d_{ij}, d_{ik}, d_{jk} defines a PH metric on $\Delta v_i v_j v_k$. The admissible space $\mathcal{U}_{ijk}^{H, \mathcal{T}}(d)$ of discrete conformal factors for the triangle $\Delta v_i v_j v_k$ with a PH metric d_{ij}, d_{ik}, d_{jk} is defined to be the set of discrete conformal factors $(u_i, u_j, u_k) \in \mathbb{R}^3$ such that the triangle with edge lengths given by (2) exists in 2-dimensional hyperbolic space \mathbb{H}^2 , i.e.

$$\mathcal{U}_{ijk}^{H, \mathcal{T}}(d) = \{(u_i, u_j, u_k) \in \mathbb{R}^3 \mid (u * d)_{rs} + (u * d)_{rt} > (u * d)_{ts}, \{r, s, t\} = \{i, j, k\}\}. \quad (9)$$

Then the admissible space $\mathcal{U}^{H, \mathcal{T}}(d)$ for (S, V, \mathcal{T}) with a PH metric d could be written as $\mathcal{U}^{H, \mathcal{T}}(d) = \bigcap_{\Delta v_i v_j v_k \in F} \mathcal{U}_{ijk}^{H, \mathcal{T}}(d)$. Here and in the following, we do not distinguish $\mathcal{U}_{ijk}^{H, \mathcal{T}}(d)$ and its natural preimage in \mathbb{R}^N if it causes no confusion.

For a triangle $\Delta v_i v_j v_k \in F$, we denote α_i^{jk} as the inner angle opposite to the edge $\{v_j v_k\}$. Based on Leibon's work [18] on volume of some generalized hyperbolic tetrahedron in \mathbb{H}^3 , Bobenko-Pinkall-Springborn [1] proved the following result.

Lemma 2.1 ([1]). Suppose (S, V, \mathcal{T}) is a connected closed triangulated surface with a PH metric d .

- (1) The admissible space $\mathcal{U}_{ijk}^{H, \mathcal{T}}(d)$ is nonempty and simply connected with analytical boundaries in \mathbb{R}^3 .
- (2) The matrix $\frac{\partial(\alpha_i^{jk}, \alpha_j^{ik}, \alpha_k^{ij})}{\partial(u_i, u_j, u_k)}$ is symmetric and negative definite on $\mathcal{U}_{ijk}^{H, \mathcal{T}}(d)$, which implies the matrix L is symmetric and positive definite on $\mathcal{U}^{H, \mathcal{T}}(d)$, where

$$L = (L_{ij})_{N \times N} = \frac{\partial(K_1, \dots, K_N)}{\partial(u_1, \dots, u_N)} = \begin{pmatrix} \frac{\partial K_1}{\partial u_1} & \dots & \frac{\partial K_1}{\partial u_N} \\ \vdots & \ddots & \vdots \\ \frac{\partial K_N}{\partial u_1} & \dots & \frac{\partial K_N}{\partial u_N} \end{pmatrix}. \quad (10)$$

- (3) The inner angles $\alpha_i^{jk}, \alpha_j^{ik}, \alpha_k^{ij}$ defined for $(u_i, u_j, u_k) \in \mathcal{U}_{ijk}^{H, \mathcal{T}}(d)$ could be extended by constants to be continuous functions $\tilde{\alpha}_i^{jk}, \tilde{\alpha}_j^{ik}, \tilde{\alpha}_k^{ij}$ defined on \mathbb{R}^3 .

One can also refer to [28] for a new proof of Lemma 2.1. We further have the following decomposition for the matrix L .

Lemma 2.2. Suppose (S, V, \mathcal{T}) is a connected closed triangulated surface with a PH metric d . The positive definite matrix L defined by (10) could be decomposed to be

$$L = L_A + L_B,$$

where $L_A = \text{diag}\{A_1, \dots, A_N\}$ is a diagonal matrix with

$$A_i = \frac{\partial}{\partial u_i} \left(\sum_{\Delta v_i v_j v_k \in F} \text{Area}(\Delta v_i v_j v_k) \right) = \sum_{j \sim i} B_{ij} (\cosh d_{ij} - 1) \quad (11)$$

and L_B is a symmetric matrix defined by

$$(L_B)_{ij} = \begin{cases} \sum_{k \sim i} B_{ik}, & j = i, \\ -B_{ij}, & j \sim i, \\ 0, & \text{otherwise,} \end{cases}$$

with

$$B_{ij} = \frac{\partial \alpha_i^{jk}}{\partial u_j} + \frac{\partial \alpha_i^{jl}}{\partial u_j} = \frac{1}{\cosh^2 \frac{d_{ij}}{2}} \left(\tan \frac{\alpha_i^{jk} + \alpha_j^{ik} - \alpha_k^{ij}}{2} + \tan \frac{\alpha_i^{jl} + \alpha_j^{il} - \alpha_l^{ij}}{2} \right). \quad (12)$$

Here $\Delta v_i v_j v_k$ and $\Delta v_i v_j v_l$ are two adjacent triangles sharing an edge $\{v_i v_j\}$. Furthermore, if the triangulation \mathcal{T} is Delaunay in the PH metric d on (S, V) , then $A_i > 0$, $B_{ij} \geq 0$, which implies L_A is positive definite and L_B is positive semi-definite.

The proof of Lemma 2.2 is too long and deviates from the theme of this paper, so we defer it to Appendix A. For Thurston's hyperbolic circle packing metrics on surfaces, Ge and the first author [10] obtained a decomposition similar to Lemma 2.2.

By Lemma 2.1, the Ricci energy function

$$F_{ijk}(u_i, u_j, u_k) = \int^{(u_i, u_j, u_k)} \alpha_i^{jk} du_i + \alpha_j^{ik} du_j + \alpha_k^{ij} du_k$$

for the triangle $\Delta v_i v_j v_k$ is a well-defined locally strictly concave function of $(u_i, u_j, u_k) \in \mathcal{U}_{ijk}^{H, \mathcal{T}}(d)$. $F_{ijk}(u_i, u_j, u_k)$ was introduced by Bobenko-Pinkall-Springborn [1] using the volume function of a generalized hyperbolic tetrahedron. Here we construct $F_{ijk}(u_i, u_j, u_k)$ directly. Based on Leibon's work on the volume of a generalized hyperbolic tetrahedron [15], Bobenko-Pinkall-Springborn [1] further observed that $F_{ijk}(u_i, u_j, u_k)$ could be extended to be a globally defined concave function on \mathbb{R}^3 . In the following, we use Luo's generalization [20] of Bobenko-Pinkall-Springborn's extension to extend the function $F_{ijk}(u_i, u_j, u_k)$.

Definition 2.3 ([20], Definition 2.3). A differential 1-form $w = \sum_{i=1}^n a_i(x) dx^i$ in an open subset $U \subset \mathbb{R}^n$ is said to be continuous if each $a_i(x)$ is continuous on U . A continuous differential 1-form w is called closed if $\int_{\partial\tau} w = 0$ for each triangle $\tau \subset U$.

Theorem 2.4 ([20], Corollary 2.6). Suppose $X \subset \mathbb{R}^n$ is an open convex set and $A \subset X$ is an open subset of X bounded by a real analytic codimension-1 submanifold in X . If

$w = \sum_{i=1}^n a_i(x) dx_i$ is a continuous closed 1-form on A so that $P(x) = \int_a^x w$ is locally convex on A and each a_i can be extended continuous to X by constant functions to a function \tilde{a}_i on X , then $\tilde{P}(x) = \int_a^x \sum_{i=1}^n \tilde{a}_i(x) dx_i$ is a C^1 -smooth convex function on X extending P .

Combining Lemma 2.1 and Theorem 2.4, $F_{ijk}(u_i, u_j, u_k)$ defined on $\mathcal{U}_{ijk}^{H,T}(d)$ could be extended to be

$$\tilde{F}_{ijk}(u_i, u_j, u_k) = \int^{(u_i, u_j, u_k)} \tilde{\alpha}_i^{jk} du_i + \tilde{\alpha}_j^{ik} du_j + \tilde{\alpha}_k^{ij} du_k, \quad (13)$$

which is a C^1 -smooth concave function defined on \mathbb{R}^3 .

Using the extension \tilde{F}_{ijk} of the Ricci energy function F_{ijk} , we can prove the following rigidity for the extended combinatorial α -curvature $\tilde{R}_\alpha = \frac{\tilde{K}_i}{w_i^\alpha} = \frac{1}{w_i^\alpha} (2\pi - \sum_{\Delta v_i v_j v_k} \tilde{\alpha}_i^{jk})$, which is a generalization of Theorem 1.3.

Theorem 2.5. Suppose (S, V, \mathcal{T}) is a connected closed triangulated surface with a PH metric d_0 , $\alpha \in \mathbb{R}$ is a constant and \bar{R} is a function defined on the vertices with $\alpha \bar{R} \leq 0$. If there exist $u_A \in \mathcal{U}^{H,T}(d_0)$ and $u_B \in \mathbb{R}^N$ with the same extended combinatorial α -curvature \bar{R} , then $u_A = u_B$.

Proof. Define the following Ricci energy $P(u)$ for \bar{R} by

$$P(u) = - \sum_{\Delta v_i v_j v_k \in F} F_{ijk}(u_i, u_j, u_k) + \int^u \sum_{i=1}^N (2\pi - \bar{R}_i w_i^\alpha) du_i.$$

By direct calculations, we have

$$\nabla_{u_i} P(u) = - \sum_{\Delta v_i v_j v_k \in F} \alpha_i^{jk} + 2\pi - \bar{R}_i w_i^\alpha = K_i - \bar{R}_i w_i^\alpha,$$

and

$$\text{Hess}_u P = L - \alpha \begin{pmatrix} \bar{R}_1 w_1^\alpha & & \\ & \ddots & \\ & & \bar{R}_N w_N^\alpha \end{pmatrix}.$$

Combining Lemma 2.1 and the condition $\alpha \bar{R} \leq 0$, $\text{Hess}_u P$ is positive definite and P is locally strictly convex on $\mathcal{U}^{H,T}(d_0)$.

By the extension $\tilde{F}_{ijk}(u_i, u_j, u_k)$ of $F_{ijk}(u_i, u_j, u_k)$ in (13), the Ricci energy function $P(u)$ defined on $\mathcal{U}^{H,T}(d_0)$ could be extended to be

$$\tilde{P}(u) = - \sum_{\Delta v_i v_j v_k \in F} \tilde{F}_{ijk}(u_i, u_j, u_k) + \int^u \sum_{i=1}^N (2\pi - \bar{R}_i w_i^\alpha) du_i,$$

which is a C^1 -smooth convex function defined on \mathbb{R}^N and locally strictly convex on $\mathcal{U}^{H,\mathcal{T}}(d_0) \subset \mathbb{R}^N$ with $\nabla_{u_i} \tilde{P} = \tilde{K}_i - \bar{R}_i w_i^\alpha$.

Set

$$f(t) = \tilde{P}((1-t)u_A + tu_B), \quad t \in [0, 1].$$

Then $f(t)$ is a C^1 smooth convex function of $t \in [0, 1]$ with

$$f'(t) = \sum_{i=1}^N \nabla_{u_i} \tilde{P}|_{(1-t)u_A + tu_B} \cdot (u_{B,i} - u_{A,i}) = \sum_{i=1}^N (\tilde{K}_i - \bar{R}_i w_i^\alpha)|_{(1-t)u_A + tu_B} \cdot (u_{B,i} - u_{A,i}).$$

By the assumption that $\tilde{R}_\alpha(u_A) = \tilde{R}_\alpha(u_B) = \bar{R}$, we have $f'(0) = f'(1) = 0$, which implies $f'(t) \equiv 0$ by the convexity of $f(t)$. Note that u_A is in the open subset $\mathcal{U}^{H,\mathcal{T}}(d_0)$ of \mathbb{R}^N , where $P(u)$ is smooth and $\text{Hess}_u P$ is positive definite, there exists $0 < \epsilon < 1$ such that $(1-t)u_A + tu_B \in \mathcal{U}^{H,\mathcal{T}}(d_0)$ and $f(t)$ is smooth and strictly convex for $t \in [0, \epsilon]$. By $f'(t) \equiv 0$ for $t \in [0, 1]$, we have $f''(t) \equiv 0$ for $t \in [0, \epsilon]$. Combining with

$$f''(t) = (u_B - u_A)^T \cdot \text{Hess}_u P|_{(1-t)u_A + tu_B} \cdot (u_B - u_A)$$

for $t \in [0, \epsilon]$ and $\text{Hess}_u P$ is positive definite for $u \in \mathcal{U}^{H,\mathcal{T}}(d_0)$, this implies $u_A = u_B$. \square

2.2 Combinatorial α -flows on triangulated surfaces

The combinatorial α -Yamabe flow (5) and the combinatorial α -Calabi flow (7) are ODE systems with smooth coefficients. Therefore, the solutions always exist locally around the initial time $t = 0$. We further have the following result on the longtime existence and convergence for the solutions of combinatorial α -flows.

Theorem 2.6. Suppose (S, V, \mathcal{T}) is a connected closed triangulated surface with a PH metric d_0 , $\alpha \in \mathbb{R}$ is a constant and \bar{R} is a function defined on V . If the solutions of combinatorial α -flows converge, there exist PH metrics with combinatorial α -curvature \bar{R} . Furthermore, if there is a PH metric $d^* = u^* * d_0$ with combinatorial α -curvature \bar{R} and $\alpha \bar{R} \leq 0$, there exists a constant $\delta > 0$ such that if $\|R_\alpha(u(0)) - R_\alpha(u^*)\| < \delta$, then the solutions of combinatorial α -flows exist for all time and converge exponentially fast to u^* .

Proof. Suppose $w(t)$ is a solution of the combinatorial α -Yamabe flow (5), then $u(t) = \ln w(t)$ is a solution of the flow $\frac{du_i}{dt} = \bar{R}_i - R_{\alpha,i}$. If $w^* := w(+\infty) = \lim_{t \rightarrow +\infty} w(t)$ exists, then $u^* := u(+\infty) = \ln w(+\infty)$ exists in $\mathcal{U}^{H,\mathcal{T}}(d_0)$ and $R_\alpha(u^*) = \lim_{t \rightarrow +\infty} R_\alpha(u(t))$ exists. Furthermore, there exists a sequence $\xi_n \in (n, n+1)$ such that

$$u_i(n+1) - u_i(n) = u'_i(\xi_n) = \bar{R}_i - R_{\alpha,i}(u(\xi_n)) \rightarrow 0 \text{ as } n \rightarrow +\infty,$$

which implies $R_{\alpha,i}(u^*) = \overline{R}_i$ for all $i \in V$ and $u^* * d_0$ is a PH metric with combinatorial α -curvature \overline{R} . Similarly, if the combinatorial α -Calabi flow (7) converges, there exists a PH metric with combinatorial α -curvature \overline{R} .

Suppose there exists a PH metric $d^* = u^* * d_0$ with combinatorial α -curvature \overline{R} . For the combinatorial α -Yamabe flow (5), set $\Gamma(u) = \overline{R} - R_\alpha$. By direct calculations, we have

$$D\Gamma|_{u=u^*} = -\Sigma^{-\alpha}L + \alpha\Lambda = -\Sigma^{-\frac{\alpha}{2}}(\Sigma^{-\frac{\alpha}{2}}L\Sigma^{-\frac{\alpha}{2}} - \alpha\Lambda)\Sigma^{\frac{\alpha}{2}},$$

where $\Sigma = \text{diag}\{w_1, w_2, \dots, w_N\}$, $\Lambda = \text{diag}\{\overline{R}_1, \overline{R}_2, \dots, \overline{R}_N\}$. By the condition $\alpha\overline{R} \leq 0$, $D\Gamma|_{u=u^*}$ has N negative eigenvalues, which implies u^* is a local attractor of (5). Then the conclusion follows from Lyapunov Stability Theorem ([21], Chapter 5).

Similarly, for the combinatorial α -Calabi flow (7), set $\Gamma(u) = \Delta_\alpha^\top(R_\alpha - \overline{R})$. By direct calculations, we have

$$\begin{aligned} D\Gamma|_{u=u^*} &= -\Sigma^{-\alpha}L\Sigma^{-\alpha}L + \alpha\Sigma^{-\alpha}L\Lambda \\ &= -\Sigma^{-\frac{\alpha}{2}}\left(\Sigma^{-\frac{\alpha}{2}}L\Sigma^{-\alpha}L\Sigma^{-\frac{\alpha}{2}} - \alpha\Sigma^{-\frac{\alpha}{2}}L\Sigma^{-\frac{\alpha}{2}}\Lambda\right)\Sigma^{\frac{\alpha}{2}} \\ &= -\Sigma^{-\frac{\alpha}{2}}(Q^2 - \alpha Q\Lambda)\Sigma^{\frac{\alpha}{2}} \\ &= -\Sigma^{-\frac{\alpha}{2}}Q^{\frac{1}{2}}\left(Q^2 + Q^{\frac{1}{2}}(-\alpha\Lambda)Q^{\frac{1}{2}}\right)Q^{-\frac{1}{2}}\Sigma^{\frac{\alpha}{2}}, \end{aligned}$$

where $Q = \Sigma^{-\frac{\alpha}{2}}L\Sigma^{-\frac{\alpha}{2}}$ is a symmetric and positive definite matrix. By the condition $\alpha\overline{R} \leq 0$, $D\Gamma|_{u=u^*}$ has N negative eigenvalues, which implies u^* is a local attractor of (5). Then the conclusion follows from Lyapunov Stability Theorem ([21], Chapter 5). \square

Theorem 2.6 gives the longtime existence and convergence for the solutions of combinatorial α -flows for initial PH metrics with small initial energy. But for general initial PH metrics, the combinatorial α -flows may develop singularities, including that some triangles degenerate and the conformal factors tend to infinity along the corresponding combinatorial α -flows. To handle the possible singularities along the combinatorial α -flows, we can do surgery by flipping on the α -flows.

3 Combinatorial α -curvature and combinatorial α -flows on discrete Riemannian surfaces

3.1 Gu-Guo-Luo-Sun-Wu's work on discrete uniformization theorem

To analyze the behavior of the combinatorial α -flows with surgery, we need the discrete conformal theory for PH metrics established by Gu-Guo-Luo-Sun-Wu [16]. In this subsection, we briefly recall the theory. Please refer to [16] for more details.

Definition 3.1. ([16], Definition 1) Two PH metrics d, d' on a connected closed marked surface (S, V) are discrete conformal if there exist a sequence of PH metrics $d_1 = d, d_2, \dots, d_m = d'$ on (S, V) and triangulations $\mathcal{T}_1, \dots, \mathcal{T}_m$ of (S, V) satisfying

- (a) each \mathcal{T}_i is Delaunay in d_i ,
- (b) if $\mathcal{T}_i = \mathcal{T}_{i+1}$, there exists a function $u : V \rightarrow \mathbb{R}$, called a conformal factor, so that if e is an edge in \mathcal{T}_i with end points v and v' , then the lengths $x_{d_i}(e)$ and $x_{d_{i+1}}(e)$ of e in metrics d_i and d_{i+1} are related by

$$\sinh \frac{x_{d_{i+1}}(e)}{2} = \sinh \frac{x_{d_i}(e)}{2} e^{u(v)+u(v')},$$

- (c) if $\mathcal{T}_i \neq \mathcal{T}_{i+1}$, then (S, d_i) is isometric to (S, d_{i+1}) by an isometry homotopic to the identity in (S, V) .

The discrete conformal class of a PH metric d on (S, V) is called a discrete Riemannian surface, which is denoted by $\mathcal{D}(d)$ [16].

Lemma 3.2. ([16], Proposition 16) If \mathcal{T} and \mathcal{T}' are Delaunay triangulations of a PH metric d on a closed marked surface (S, V) , there exist a sequence of Delaunay triangulations $\mathcal{T}_1 = \mathcal{T}, \mathcal{T}_2, \dots, \mathcal{T}_k = \mathcal{T}'$ of d so that \mathcal{T}_{i+1} is obtained from \mathcal{T}_i by a diagonal switch.

The diagonal switch in Lemma 3.2 is the surgery by flipping described in the introduction. Using the new definition of discrete conformality, Gu-Guo-Luo-Sun-Wu [16] proved the following discrete conformal uniformization theorem for vertex scaling of PH metrics.

Theorem 3.3. ([16], Theorem 3) Suppose (S, V) is a connected closed marked surface and d is a PH metric on (S, V) . Then for any $K^* : V \rightarrow (-\infty, 2\pi)$ with $\sum_{v \in V} K^*(v) > 2\pi\chi(S)$, there exists a unique PH metric d' on (S, V) so that d' is discrete conformal to d and the discrete curvature of d' is K^* . Furthermore, the discrete Yamabe flow with surgery associated to curvature K^* having initial value d converges to d' linearly fast.

Denote the Teichmüller space of all PH metrics on (S, V) by $T_{hp}(S, V)$ and decorated Teichmüller space of all equivalence class of decorated hyperbolic metrics on $S - V$ by $T_D(S - V)$. To prove Theorem 3.3, Gu-Guo-Luo-Sun-Wu proved the following result.

Theorem 3.4. ([16], Theorem 22, Corollary 24) There is a C^1 -diffeomorphism $\mathbf{A} : T_{hp}(S, V) \rightarrow T_D(S, V)$ between $T_{hp}(S, V)$ and $T_D(S, V)$. Furthermore, the space $\mathcal{D}(d) \subset T_{hp}(S, V)$ of all equivalence classes of PH metrics discrete conformal to d is C^1 -diffeomorphism to $\{p\} \times \mathbb{R}_{>0}^V$ under the diffeomorphism \mathbf{A} , where p is the unique hyperbolic metric on $S - V$ determined by the PH metric d on (S, V) .

Theorem 3.4 implies that the union of the admissible spaces $\mathcal{U}_D^{H,\mathcal{T}}(d')$ of conformal factors such that \mathcal{T} is Delaunay for $d' \in \mathcal{D}(d)$ is \mathbb{R}^n .

Set $u_i = \ln w_i$, $1 \leq i \leq n$, for $w = (w_1, w_2, \dots, w_n) \in \mathbb{R}_{>0}^n$ and $\mathbf{P} = \{x \in (-\infty, 2\pi)^n \mid \sum_{i=1}^n x_i > 2\pi\chi(S)\}$. Using the map \mathbf{A} , Gu-Guo-Luo-Sun-Wu [16] defined the curvature map

$$\begin{aligned} \mathbf{F} : \mathbb{R}^n &\rightarrow \mathbf{P} \\ u &\mapsto K_{\mathbf{A}^{-1}(p,w(-2u))} \end{aligned} \tag{14}$$

and proved the following property of \mathbf{F} .

Proposition 3.5. ([16]) There exists a C^2 -smooth strictly convex function $W : \mathbb{R}^n \rightarrow \mathbb{R}$ defined by $W(u) = \int_0^u \sum_{i=1}^N \mathbf{F}_i du_i$ so that its gradient ∇W is \mathbf{F} .

Remark 5. Proposition 3.5 implies that \mathbf{F} defined on \mathbb{R}^n is a C^1 -extension of the combinatorial curvature K defined on the space of conformal factors $\mathcal{U}_D^{H,\mathcal{T}}(d')$ for $d' \in \mathcal{D}(d)$. Furthermore, the discrete conformal factor u is uniquely determined by the curvature map \mathbf{F} and then determined by the PH metric in the discrete conformal class $\mathcal{D}(d)$ [16].

Note that the discrete α -Laplace operator $\Delta_\alpha^\mathcal{T}$ is independent of the Delaunay triangulations of a PH metric, then the discrete α -Laplace operator $\Delta_\alpha^\mathcal{T}$ could be extend to the following operator Δ_α defined on \mathbb{R}^n , which is the space of discrete conformal factors for the discrete conformal class $\mathcal{D}(d)$.

Definition 3.6. Suppose (S, V) is a connected closed marked surface with a PH metric d_0 . For a function $f : V \rightarrow \mathbb{R}$ on the vertices, the hyperbolic discrete α -Laplace operator of $d \in \mathcal{D}(d_0)$ on (S, V) is defined to be the map

$$\begin{aligned} \Delta_\alpha : \mathbb{R}^V &\rightarrow \mathbb{R}^V \\ f &\mapsto \Delta_\alpha f, \end{aligned}$$

where the value of $\Delta_\alpha f$ at vertex v_i is

$$\Delta_\alpha f_i = -\frac{1}{w_i^\alpha} \sum_{j \in V} \frac{\partial \mathbf{F}_i}{\partial u_j} f_j = -\frac{1}{w_i^\alpha} (\tilde{L}f)_i \tag{15}$$

with $\tilde{L}_{ij} = \frac{\partial \mathbf{F}_i}{\partial u_j}$ being an extension of $L_{ij} = \frac{\partial K_i}{\partial u_j}$.

Remark 6. By Remark 5, the discrete conformal factor $u \in \mathbb{R}^N$ is uniquely determined by the PH metric $d \in \mathcal{D}(d_0)$ and the discrete α -Laplace operator in Definition 3.6 is well-defined. By Proposition 3.5, Δ_α is continuous and piecewise smooth on \mathbb{R}^n as a matrix-valued function of u . By Lemma 2.2, we have

$$\Delta_\alpha f_i = \sum_{j \sim i} \frac{B_{ij}}{w_i^\alpha} (f_j - f_i) - \frac{A_i}{w_i^\alpha} f_i \tag{16}$$

with $A_i > 0$ and $B_{ij} \geq 0$.

3.2 Rigidity of combinatorial α -curvature on discrete Riemannian surfaces

Motivated by the relationship of \mathbf{F} defined on a discrete Riemannian surface and the combinatorial curvature K defined on a triangulated surface, we introduce the following definition of combinatorial α -curvature \mathbf{F}_α for PH metrics on a discrete Riemannian surface, which is an extension of the combinatorial α -curvature R_α defined on a triangulated surface.

Definition 3.7. Suppose (S, V) is a connected closed marked surface with a PH metric d_0 , $\alpha \in \mathbb{R}$ is a constant and \mathbf{F} is the curvature map defined by (14). The combinatorial α -curvature for $d \in \mathcal{D}(d_0)$ is defined to be

$$\mathbf{F}_{\alpha,i} = \frac{\mathbf{F}_i}{w_i^\alpha}, \quad (17)$$

where $w_i = e^{u_i}$ and $u : V \rightarrow \mathbb{R}$ is the unique discrete conformal factor determined by the PH metric $d \in \mathcal{D}(d_0)$.

By Remark 5, the combinatorial α -curvature \mathbf{F}_α in Definition 3.7 is well-defined for $d \in \mathcal{D}(d_0)$. Denote the space of conformal factors by $\mathcal{U}(d)$ for a PH metric d on the marked surface (S, V) , which is \mathbb{R}^N by Theorem 3.4. Analogous to Theorem 1.3, we have the following global rigidity for the combinatorial α -curvature \mathbf{F}_α with respect to discrete conformal factors on discrete Riemannian surfaces.

Theorem 3.8. Suppose (S, V) is a closed marked surface with a PH metric d_0 , $\alpha \in \mathbb{R}$ is a constant and \bar{F} is a function defined on V . If $\alpha \bar{F} \leq 0$, there exists at most one discrete conformal factor $u^* \in \mathcal{U}(d_0)$ such that $\mathbf{A}^{-1}(p, w(-2u^*)) \in \mathcal{D}(d_0)$ has the combinatorial α -curvature \bar{F} .

Proof. By Proposition 3.5, define the following Ricci energy function

$$W_\alpha(u) = W(u) - \int_0^u \sum_{i=1}^N \bar{F}_i w_i^\alpha du_i = \int_0^u \sum_{i=1}^N \mathbf{F}_i du_i - \int_0^u \sum_{i=1}^N \bar{F}_i w_i^\alpha du_i.$$

Then $W_\alpha(u)$ is a C^2 -smooth function defined on \mathbb{R}^n with $\nabla_{u_i} W_\alpha = \nabla_{u_i} W - \bar{F}_i w_i^\alpha = \mathbf{F}_i - \bar{F}_i w_i^\alpha$. Therefore, for $u^* \in \mathcal{U}(d)$, $\mathbf{A}^{-1}(p, w(-2u^*)) \in \mathcal{D}(d)$ has the combinatorial α -curvature \bar{F} if and only if $\nabla_{u_i} W_\alpha(u^*) = 0$. By direct calculations, we have

$$\text{Hess}W_\alpha = \tilde{L} - \alpha \begin{pmatrix} \bar{F}_1 w_1^\alpha & & \\ & \ddots & \\ & & \bar{F}_N w_N^\alpha \end{pmatrix}.$$

Due to $\alpha\bar{F} \leq 0$, $\text{Hess}W_\alpha$ is positive definite and W_α is strictly convex on \mathbb{R}^n by Proposition 3.5. Then the rigidity follows from the following well-known result from analysis.

Lemma 3.9. If $W : \Omega \rightarrow \mathbb{R}$ is a C^1 -smooth strictly convex function on an open convex set $\Omega \subset \mathbb{R}^m$, then its gradient $\nabla W : \Omega \rightarrow \mathbb{R}^m$ is an embedding. □

3.3 Combinatorial α -Yamabe flow with surgery

Using the discrete conformal theory established by Gu-Guo-Luo-Sun-Wu [16], the combinatorial α -Yamabe flow with surgery can be defined as follows.

Definition 3.10. Suppose (S, V) is a connected closed marked surface with a PH metric d_0 and $\alpha \in \mathbb{R}$ is a constant. The combinatorial α -Yamabe flow with surgery is defined to be

$$\begin{cases} \frac{du_i}{dt} = -\mathbf{F}_{\alpha,i}, \\ u_i(0) = 0. \end{cases} \quad (18)$$

One can modify the combinatorial α -Yamabe flow with surgery (18) to the following form

$$\frac{du_i}{dt} = \bar{\mathbf{F}}_i - \mathbf{F}_{\alpha,i} \quad (19)$$

to study prescribed curvature problems, where $\bar{\mathbf{F}}$ is a function defined on V .

Theorem 3.11. Suppose (S, V) is a connected closed marked surface with a PH metric d_0 , $\alpha \in \mathbb{R}$ is a constant and $\bar{\mathbf{F}} : V \rightarrow \mathbb{R}$ is a function defined on V with $\alpha\bar{\mathbf{F}} \leq 0$. Then there exists a PH metric in the discrete conformal class $\mathcal{D}(d_0)$ with combinatorial α -curvature $\bar{\mathbf{F}}$ if and only if the solution of modified combinatorial α -Yamabe flow with surgery (19) exists for all time and converges exponentially fast to a PH metric with combinatorial α -curvature $\bar{\mathbf{F}}$.

Proof. Suppose the solution $u(t)$ of the modified combinatorial α -Yamabe flow with surgery (19) converges to $u^* \in \mathcal{U}(d_0)$ as $t \rightarrow +\infty$, then $\mathbf{F}_\alpha(u^*) = \lim_{t \rightarrow +\infty} \mathbf{F}_\alpha(u(t))$ by the C^1 -smoothness of \mathbf{F}_α . Furthermore, there exists a sequence $\xi_n \in (n, n+1)$ such that

$$u_i(n+1) - u_i(n) = u'_i(\xi_n) = \bar{\mathbf{F}}_i - \mathbf{F}_{\alpha,i}(u(\xi_n)) \rightarrow 0, \text{ as } n \rightarrow +\infty,$$

which implies $\mathbf{F}_{\alpha,i}(u^*) = \lim_{n \rightarrow +\infty} \mathbf{F}_{\alpha,i}(u(\xi_n)) = \bar{\mathbf{F}}_i$ for all $v_i \in V$ and $u^* \in \mathcal{U}(d_0)$ is a discrete conformal factor with combinatorial α -curvature $\bar{\mathbf{F}}$.

Conversely, suppose there exists a PH metric u^* in the discrete conformal class $\mathcal{D}(d_0)$ with combinatorial α -curvature $\bar{\mathbf{F}}$. Set

$$W_\alpha(u) = W(u) - \int_{u^*}^u \sum_{i=1}^N \bar{\mathbf{F}}_i w_i^\alpha du_i = \int_{u^*}^u \sum_{i=1}^N \mathbf{F}_i du_i - \int_{u^*}^u \sum_{i=1}^N \bar{\mathbf{F}}_i w_i^\alpha du_i.$$

Then $W_\alpha(u)$ is a C^2 -smooth strictly convex function defined on \mathbb{R}^n by Proposition 3.5 and the condition $\alpha\bar{\mathbf{F}} \leq 0$. Furthermore, $W_\alpha(u^*) = 0$, $\nabla W_\alpha(u^*) = 0$, which implies $W_\alpha(u) \geq W_\alpha(u^*) = 0$ and $\lim_{u \rightarrow \infty} W_\alpha(u) = +\infty$ by the convexity of $W_\alpha(u)$. One can also refer to Lemma 4.6 in [13] for a proof for this fact. By direct calculations, we have

$$\frac{dW_\alpha(u(t))}{dt} = \sum_{i=1}^N \frac{\partial W_\alpha}{\partial u_i} \frac{du_i}{dt} = \sum_{i=1}^N (\mathbf{F}_{\alpha,i} - \bar{\mathbf{F}}_i) w_i^\alpha (\bar{\mathbf{F}}_i - \mathbf{F}_{\alpha,i}) = - \sum_{i=1}^N (\bar{\mathbf{F}}_i - \mathbf{F}_{\alpha,i})^2 w_i^\alpha \leq 0,$$

which implies $0 \leq W_\alpha(u(t)) \leq W_\alpha(u(0))$. Therefore, $\{u(t)\} \subset\subset \mathbb{R}^N$ by $\lim_{u \rightarrow \infty} W_\alpha(u) = +\infty$, which implies the solution of modified combinatorial α -Yamabe flow with surgery (19) exists for all time and $W_\alpha(u(t))$ converges. Moreover, there exists a sequence $\xi_n \in (n, n+1)$ such that as $n \rightarrow +\infty$,

$$W_\alpha(u(n+1)) - W_\alpha(u(n)) = (W_\alpha(u(t)))'|_{\xi_n} = \nabla W_\alpha \cdot \frac{du_i}{dt}|_{\xi_n} = - \sum_{i=1}^N (\bar{\mathbf{F}}_i - \mathbf{F}_{\alpha,i})^2 w_i^\alpha|_{\xi_n} \rightarrow 0.$$

Then $\lim_{n \rightarrow +\infty} \mathbf{F}_{\alpha,i}(u(\xi_n)) = \bar{\mathbf{F}}_i = \mathbf{F}_{\alpha,i}(u^*)$ for all $v_i \in V$. By $\{u(t)\} \subset\subset \mathbb{R}^N$, there exists $\bar{u} \in \mathbb{R}^N$ and a subsequence of $\{u(\xi_n)\}$, still denoted as $\{u(\xi_n)\}$ for simplicity, such that $\lim_{n \rightarrow \infty} u(\xi_n) = \bar{u}$, which implies $\mathbf{F}_{\alpha,i}(\bar{u}) = \lim_{n \rightarrow +\infty} \mathbf{F}_{\alpha,i}(u(\xi_n)) = \mathbf{F}_{\alpha,i}(u^*)$. This further implies $\bar{u} = u^*$ by Theorem 3.8. Therefore, $\lim_{n \rightarrow \infty} u(\xi_n) = u^*$.

Set $\Gamma(u) = \bar{\mathbf{F}} - \mathbf{F}_\alpha$, then $D\Gamma|_{u=u^*}$ has N negative eigenvalues, which implies that u^* is a local attractor of (19). Then the conclusion follows from Lyapunov Stability Theorem ([21], Chapter 5). \square

3.4 Existence of PH metrics with the prescribed α -curvature

Set $M_{\alpha,i} = \mathbf{F}_{\alpha,i} - \bar{\mathbf{F}}_i$. Then the modified combinatorial Yamabe flow with surgery (19) could be written as

$$\frac{dw_i}{dt} = -M_{\alpha,i}.$$

We have the following evolution equation for $M_{\alpha,i}$ along the modified combinatorial α -Yamabe flow with surgery (19).

Lemma 3.12. Along the modified combinatorial α -Yamabe flow with surgery (19), $M_{\alpha,i}$ evolves according to

$$\frac{dM_{\alpha,i}}{dt} = \sum_{j \sim i} \frac{B_{ij}}{w_i^\alpha} (M_{\alpha,j} - M_{\alpha,i}) - \frac{A_i}{w_i^\alpha} M_{\alpha,i} + \alpha M_{\alpha,i} (M_{\alpha,i} + \bar{\mathbf{F}}_i), \quad (20)$$

where $A_i > 0$, $B_{ij} \geq 0$.

Proof. By the chain rules,

$$\begin{aligned} \frac{dM_{\alpha,i}}{dt} &= \sum_{j=1}^N \frac{\partial \mathbf{F}_{\alpha,i}}{\partial u_j} \frac{du_j}{dt} \\ &= \sum_{j=1}^N \left(\frac{1}{w_i^\alpha} \frac{\partial \mathbf{F}_i}{\partial u_j} - \frac{\mathbf{F}_i}{w_i^\alpha} \alpha \delta_{ij} \right) \cdot (-M_{\alpha,j}) \\ &= (\Delta_\alpha M_\alpha)_i + \alpha M_{\alpha,i} (M_{\alpha,i} + \bar{\mathbf{F}}_i) \\ &= \sum_{j \sim i} \frac{B_{ij}}{w_i^\alpha} (M_{\alpha,j} - M_{\alpha,i}) - \frac{A_i}{w_i^\alpha} M_{\alpha,i} + \alpha M_{\alpha,i} (M_{\alpha,i} + \bar{\mathbf{F}}_i), \end{aligned}$$

where (16) is used in the last line. $A_i > 0$ and $B_{ij} \geq 0$ follows from Lemma 2.2. \square

To analyze the longtime behavior of the combinatorial α -Yamabe flow with surgery, we need the following discrete maximum principle obtained in [14].

Theorem 3.13. (Maximum Principle) Let $f : V \times [0, T] \rightarrow \mathbb{R}$ be a C^1 function such that

$$\frac{\partial f_i}{\partial t} \geq \sum_{j \sim i} a_{ij} (f_j - f_i) + \Phi_i(f_i), \quad \forall (v_i, t) \in V \times [0, T],$$

where $a_{ij} \geq 0$ and $\Phi_i : \mathbb{R} \rightarrow \mathbb{R}$ is a local Lipschitz function. Suppose there exists $C_1 \in \mathbb{R}$ such that $f_i(0) \geq C_1$ for all $v_i \in V$. Let φ be the solution to the associated ODE

$$\begin{cases} \frac{d\varphi}{dt} = \Phi_i(\varphi) \\ \varphi(0) = C_1, \end{cases}$$

then $f_i(t) \geq \varphi(t)$ for all $(v_i, t) \in V \times [0, T]$ such that $\varphi(t)$ exists.

Similarly, suppose $f : V \times [0, T] \rightarrow \mathbb{R}$ be a C^1 function such that

$$\frac{\partial f_i}{\partial t} \leq \sum_{j \sim i} a_{ij} (f_j - f_i) + \Phi_i(f_i), \quad \forall (v_i, t) \in V \times [0, T].$$

Suppose there exists $C_2 \in \mathbb{R}$ such that $f_i(0) \leq C_2$ for all $v_i \in V$. Let ψ be the solution to the associated ODE

$$\begin{cases} \frac{d\psi}{dt} = \Phi_i(\psi) \\ \psi(0) = C_2, \end{cases}$$

then $f_i(t) \leq \psi(t)$ for all $(v_i, t) \in V \times [0, T)$ such that $\psi(t)$ exists.

Remark 7. In Theorem 3.13, $\sum_{j \sim i} a_{ij}(f_j - f_i)$ is actually the classical discrete Laplace operator Δf_i used in [3]. In order to avoid confusion with the α -Laplace operator Δ_α in Definition 3.6, here we use the notation $\sum_{j \sim i} a_{ij}(f_j - f_i)$ rather than Δf_i .

As a direct application of Theorem 3.13 to (20), we have the following property for M_α .

Corollary 3.14. If $M_{\alpha,i}(0) \leq 0$ for all $v_i \in V$, then $M_{\alpha,i}(t) \leq 0$ for all $v_i \in V$ along the modified combinatorial α -Yamabe flow (19). If $M_{\alpha,i}(0) \geq 0$ for all $v_i \in V$, then $M_{\alpha,i}(t) \geq 0$ for all $v_i \in V$ along the modified combinatorial α -Yamabe flow (19).

As another application of Theorem 3.13, we have the following existence of PH metrics with prescribed combinatorial α -curvature.

Lemma 3.15. Suppose (S, V) is a connected closed marked surface with a PH metric d_0 , $\alpha > 0$ is a constant and $\bar{\mathbf{F}} : V \rightarrow \mathbb{R}$ is a negative constant function defined on V . If there exists $u^* \in \mathcal{U}(d_0)$ such that $\bar{\mathbf{F}} < \mathbf{F}_{\alpha,i}(u^*) < 0$ for all $v_i \in V$, then there exists a PH metric in the discrete conformal class $\mathcal{D}(d_0)$ with combinatorial α -curvature $\bar{\mathbf{F}}$.

Proof. Taking u^* as the initial value of the modified combinatorial α -Yamabe flow with surgery (19). We shall prove that the solution of (19) with initial value $u(0) = u^*$ exists for all time and converges to a discrete conformal factor with combinatorial α -curvature $\bar{\mathbf{F}}$, which implies the existence of a PH metric in the discrete conformal class $\mathcal{D}(d_0)$ with combinatorial α -curvature $\bar{\mathbf{F}}$.

Set

$$\begin{aligned} \mathbf{F}_{\alpha,\max}(u(0)) &= \max_{i \in V} \mathbf{F}_{\alpha,i}(u^*), \quad M_{\alpha,\max} = \max_{i \in V} M_{\alpha,i}. \\ \mathbf{F}_{\alpha,\min}(u(0)) &= \min_{i \in V} \mathbf{F}_{\alpha,i}(u^*), \quad M_{\alpha,\min} = \min_{i \in V} M_{\alpha,i}. \end{aligned}$$

Due to $\bar{\mathbf{F}} < \mathbf{F}_{\alpha,\min}(u(0))$, we have $M_{\alpha,i}(0) > 0$ for all $v_i \in V$, which implies $M_{\alpha,i}(t) \geq 0$ for all $v_i \in V$ by Corollary 3.14. Combining $A_i > 0$, $M_{\alpha,i}(t) \geq 0$ and (20), we have

$$\frac{dM_{\alpha,i}}{dt} \leq \sum_{j \sim i} \frac{B_{ij}}{w_i^\alpha} (M_{\alpha,j} - M_{\alpha,i}) + \alpha M_{\alpha,i} (M_{\alpha,i} + \bar{\mathbf{F}}),$$

which implies

$$0 \leq M_{\alpha,i}(t) \leq \frac{\bar{\mathbf{F}}}{-1 + \left(\frac{\bar{\mathbf{F}}}{M_{\alpha,\max}(0)} + 1\right)e^{-\alpha \bar{\mathbf{F}}t}} \leq \frac{\bar{\mathbf{F}} M_{\alpha,\max}(u(0))}{\mathbf{F}_{\alpha,\max}(u(0))} e^{\alpha \bar{\mathbf{F}}t} \quad (21)$$

by Theorem 3.13. Combining $\frac{du_i}{dt} = -M_{\alpha,i}$, $\alpha > 0$ and $\bar{\mathbf{F}} < 0$, the estimation (21) further implies $\{u(t)\} \subset \subset \mathbb{R}^N$. Therefore, the solution of modified combinatorial α -Yamabe flow

with surgery (19) exists for all time and converges, which implies the existence of a PH metric in the conformal class of $\mathcal{D}(d_0)$ with combinatorial α -curvature $\overline{\mathbf{F}}$. \square

Remark 8. Lemma 3.15 still holds under the condition that $\alpha < 0$, $\overline{\mathbf{F}}$ is a positive constant and $0 < \mathbf{F}_{\alpha,i}(u^*) < \overline{\mathbf{F}}$ for all $v_i \in V$, the proof of which is the same as that for Lemma 3.15.

To handle more general cases, we need the following technique lemma.

Lemma 3.16. Suppose $\triangle v_i v_j v_k \in F$ is a hyperbolic triangle with discrete hyperbolic metrics d_{ij}, d_{jk}, d_{ki} . If the discrete conformal factors u_i, u_j, u_k for the hyperbolic triangle are uniformly bounded from below, i.e. $\exists C$ such that $u_i, u_j, u_k \in [C, +\infty)$. Then for any $\epsilon > 0$, there exists a number $\tilde{C} = \tilde{C}(d_{ij}, d_{jk}, d_{ki}, C)$ such that if $u_i > \tilde{C}$, the inner angle α_i^{jk} is smaller than ϵ .

Proof. We use the trick from [12] to prove the lemma. Due to

$$\sinh \frac{l_{ij}}{2} = \sinh \frac{d_{ij}}{2} e^{u_i+u_j} \geq \sinh \frac{d_{ij}}{2} e^{u_i+C},$$

we have $\sinh \frac{l_{ij}}{2} \rightarrow +\infty$, $\sinh \frac{l_{ik}}{2} \rightarrow +\infty$ uniformly as $u_i \rightarrow +\infty$, where $l_{rs} = (u * d)_{rs}$, $\{r, s, t\} = \{i, j, k\}$. By the hyperbolic cosine law,

$$\begin{aligned} \cos \alpha_i^{jk} &= \frac{\cosh l_{ij} \cosh l_{ik} - \cosh l_{jk}}{\sinh l_{ij} \sinh l_{ik}} \\ &= \frac{\cosh(l_{ij} + l_{ik}) + \cosh(l_{ij} - l_{ik}) - 2 \cosh l_{jk}}{\cosh(l_{ij} + l_{ik}) - \cosh(l_{ij} - l_{ik})} \\ &= \frac{1 + \lambda - 2\mu}{1 - \lambda}, \end{aligned}$$

where $\lambda = \frac{\cosh(l_{ij}-l_{ik})}{\cosh(l_{ij}+l_{ik})}$ and $\mu = \frac{\cosh l_{jk}}{\cosh(l_{ij}+l_{ik})}$. To prove $\alpha_i^{jk} \rightarrow 0$ uniformly, we just need to prove $\lambda, \mu \rightarrow 0$ uniformly as $u_i \rightarrow +\infty$. Note that

$$\begin{aligned} 0 < \lambda &< \frac{\cosh l_{ij}}{\cosh(l_{ij} + l_{ik})} < \frac{1}{\cosh l_{ik}} \leq \frac{1}{\sinh \frac{d_{ik}}{2} e^{u_i+C}}, \\ 0 < \mu &= \frac{1}{(1 + 2 \sinh^2 \frac{l_{ij}}{2})(1 + 2 \sinh^2 \frac{l_{ik}}{2})} + \frac{\sinh^2 \frac{d_{jk}}{2}}{\sinh^2 \frac{d_{ij}}{2} \sinh^2 \frac{d_{ik}}{2} e^{4u_i}}, \end{aligned}$$

we have $\lambda, \mu \rightarrow 0$ uniformly as $u_i \rightarrow +\infty$. Therefore, $\alpha_i^{jk} \rightarrow 0$ uniformly as $u_i \rightarrow +\infty$. \square

As an application of Lemma 3.16, we have the following result.

Lemma 3.17. Suppose (S, V) is a connected closed marked surface with a PH metric d_0 , $\alpha > 0$ is a constant and $\bar{\mathbf{F}}$ is a non-positive constant. If there exists $u^* \in \mathcal{U}(d_0)$ such that $\mathbf{F}_\alpha(u^*)$ is a negative constant, then there exists a PH metric in the discrete conformal class $\mathcal{D}(d_0)$ with combinatorial α -curvature $\bar{\mathbf{F}}$.

Proof. Because $\mathbf{F}_\alpha(u^*)$ and $\bar{\mathbf{F}}$ are both constants, then

$$\bar{\mathbf{F}} = \mathbf{F}_\alpha(u^*) < 0 \quad \text{or} \quad \bar{\mathbf{F}} < \mathbf{F}_\alpha(u^*) < 0 \quad \text{or} \quad \mathbf{F}_\alpha(u^*) < \bar{\mathbf{F}} \leq 0.$$

In the case that $\bar{\mathbf{F}} = \mathbf{F}_\alpha(u^*) < 0$, $\mathbf{A}^{-1}(p, w(-2u^*)) \in \mathcal{D}(d_0)$ is a PH metric with combinatorial α -curvature $\bar{\mathbf{F}}$.

In the case that $\bar{\mathbf{F}} < \mathbf{F}_\alpha(u^*) < 0$, the conclusion follows from Lemma 3.15.

In the case that $\mathbf{F}_\alpha(u^*) < \bar{\mathbf{F}} \leq 0$, we take u^* as the initial value of the modified combinatorial α -Yamabe flow with surgery (19). Then $M_\alpha(0) < 0$ is a constant. By Corollary 3.14, $M_{\alpha,i}(t) \leq 0$ for all $v_i \in V$, which implies $\mathbf{F}_{\alpha,i}(u(t)) \leq \bar{\mathbf{F}} \leq 0$ for all $v_i \in V$. Combining $A_i > 0$, $M_{\alpha,i}(t) \leq 0$ and (20), we have

$$\frac{dM_{\alpha,i}}{dt} \geq \sum_{j \sim i} \frac{B_{ij}}{w_i^\alpha} (M_{\alpha,j} - M_{\alpha,i}) + \alpha M_{\alpha,i} (M_{\alpha,i} + \bar{\mathbf{F}}).$$

Therefore, by Theorem 3.13,

$$\begin{aligned} \frac{\bar{\mathbf{F}}}{-1 + \left(\frac{\bar{\mathbf{F}}}{M_\alpha(0)} + 1\right)e^{-\alpha\bar{\mathbf{F}}t}} &\leq M_{\alpha,i}(t) \leq 0, \quad \text{if } \bar{\mathbf{F}} < 0, \\ \frac{M_\alpha(0)}{-\alpha M_\alpha(0)t + 1} &\leq M_{\alpha,i}(t) \leq 0, \quad \text{if } \bar{\mathbf{F}} = 0. \end{aligned} \tag{22}$$

In the case that $\bar{\mathbf{F}} < 0$, by the the estimation (22) and $\frac{du_i}{dt} = -M_{\alpha,i}(t)$, we have $\{u(t)\} \subset \subset \mathbb{R}^N$. In the case that $\bar{\mathbf{F}} = 0$, we have $\frac{du_i}{dt} = -M_{\alpha,i}(t) \geq 0$, which implies $u_i(t)$ is uniformly bounded from below for all $v_i \in V$. We claim that $u(t)$ is bounded from above. If not, then there exists at least one vertex $v_i \in V$ such that $\lim_{t \rightarrow +\infty} u_i(t) = +\infty$. For the vertex v_i , by Lemma 3.16, there exists $C_i > 0$ such that whenever $u_i(t) > C_i$, the inner angle α_i^{jk} is smaller than $\frac{\pi}{N_i}$, where N_i is the degree at the vertex i . Thus $K_i = 2\pi - \sum_{\Delta v_i v_j v_k \in F} \alpha_i^{jk} > \pi > 0$ and then $\mathbf{F}_{\alpha,i}(u(t)) > 0$, which contradicts $\mathbf{F}_{\alpha,i}(u(t)) \leq 0$. Therefore, $u_i(t)$ is bounded from above. This completes the proof of the claim. As a consequence, the solution of modified combinatorial α -Yamabe flow with surgery (19) exists for all time and converges, which implies the existence of a PH metric in the discrete conformal class $\mathcal{D}(d_0)$ with combinatorial α -curvature $\bar{\mathbf{F}}$. \square

Remark 9. Lemma 3.17 still holds under the condition that $\alpha < 0$, $\bar{\mathbf{F}}$ is a positive constant and $\mathbf{F}_\alpha(u^*)$ is a positive constant. As we do not have a result paralleling to Lemma 3.16 under the condition that u_i is bounded from above for all $v_i \in V$, the result in Lemma 3.17 can not be generalized to the case that $\alpha < 0$, $\bar{\mathbf{F}}$ is a nonnegative constant and $\mathbf{F}_\alpha(u^*)$ is a positive constant.

As a direct corollary of Lemma 3.15, Lemma 3.17, Remark 8 and Remark 9, we have the following result.

Corollary 3.18. Suppose (S, V) is a connected closed marked surface with a PH metric d_0 and α is a constant.

- (1) If $\alpha > 0$ and there exists $u^* \in \mathcal{U}(d_0)$ such that $\mathbf{F}_{\alpha,i}(u^*) < 0$ for all $v_i \in V$, then for any nonpositive constant $\bar{\mathbf{F}}$, there exists a PH metric in $\mathcal{D}(d_0)$ with combinatorial α -curvature $\bar{\mathbf{F}}$;
- (2) If $\alpha < 0$ and there exists $u^* \in \mathcal{U}(d_0)$ such that $\mathbf{F}_{\alpha,i}(u^*) > 0$ for all $v_i \in V$, then for any positive constant $\bar{\mathbf{F}}$, there exists a PH metric in $\mathcal{D}(d_0)$ with combinatorial α -curvature $\bar{\mathbf{F}}$.

The following result shows that the constant condition on $\bar{\mathbf{F}}$ in Corollary 3.18 could be removed.

Theorem 3.19. Suppose (S, V) is a connected closed marked surface with a PH metric d_0 , α is a constant and $\bar{\mathbf{F}} : V \rightarrow \mathbb{R}$ is a function defined on V . If there exists $u^* \in \mathcal{U}(d_0)$ such that one of the following conditions is satisfied :

- (1) $\alpha > 0$, $\bar{\mathbf{F}}_i \leq 0$, $\mathbf{F}_{\alpha,i}(u^*) < 0$ for all $v_i \in V$,
- (2) $\alpha < 0$, $\bar{\mathbf{F}}_i > 0$, $\mathbf{F}_{\alpha,i}(u^*) > 0$ for all $v_i \in V$,

then there exists a PH metric in the discrete conformal class $\mathcal{D}(d_0)$ with combinatorial α -curvature $\bar{\mathbf{F}}$.

Proof. In the case that $\alpha > 0$, $\bar{\mathbf{F}} \leq 0$, $\mathbf{F}_{\alpha,i}(u^*) < 0$, set

$$\bar{\mathbf{F}}_{\max} = \max_{v_i \in V} \bar{\mathbf{F}}_i, \quad \bar{\mathbf{F}}_{\min} = \min_{v_i \in V} \bar{\mathbf{F}}_i.$$

By Corollary 3.18, there exists $\bar{u} \in \mathcal{U}(d_0)$ with negative constant combinatorial α -curvature $\mathbf{F}_\alpha(\bar{u}) < \bar{\mathbf{F}}_{\min} < 0$. Taking \bar{u} as the initial value of the modified combinatorial α -Yamabe

flow with surgery (19), then $M_{\alpha,i}(0) < 0$ for all $v_i \in V$, which implies $M_{\alpha,i}(t) \leq 0$ for all $v_i \in V$ by Corollary 3.14. Combining $\alpha > 0$, $\bar{\mathbf{F}} \leq 0$, $A_i > 0$, $M_{\alpha,i}(t) \leq 0$ and (20), we have

$$\begin{aligned} \frac{dM_{\alpha,i}}{dt} &\geq \sum_{j \sim i} \frac{B_{ij}}{w_i^\alpha} (M_{\alpha,j} - M_{\alpha,i}) + \alpha M_{\alpha,i} (M_{\alpha,i} + \bar{\mathbf{F}}_i) \\ &\geq \sum_{j \sim i} \frac{B_{ij}}{w_i^\alpha} (M_{\alpha,j} - M_{\alpha,i}) + \alpha M_{\alpha,i} (M_{\alpha,i} + \bar{\mathbf{F}}_{\max}). \end{aligned}$$

Therefore, by Theorem 3.13,

$$\begin{aligned} \frac{\bar{\mathbf{F}}_{\max}}{-1 + \left(\frac{\bar{\mathbf{F}}_{\max}}{M_{\alpha,\min}(0)} + 1\right)e^{-\alpha\bar{\mathbf{F}}_{\max}t}} &\leq M_{\alpha,i}(t) \leq 0, \text{ if } \bar{\mathbf{F}}_{\max} < 0, \\ \frac{M_{\alpha,\min}(0)}{-\alpha M_{\alpha,\min}(0)t + 1} &\leq M_{\alpha,i}(t) \leq 0, \text{ if } \bar{\mathbf{F}}_{\max} = 0. \end{aligned}$$

Following the arguments in the proof of Lemma 3.17, the solution of the modified combinatorial α -Yamabe flow with surgery (19) stays in a compact subset of \mathbb{R}^N . Therefore, the solution of (19) exists for all time and converges, which implies the existence of a PH metric in the discrete conformal class of $\mathcal{D}(d_0)$ with combinatorial α -curvature $\bar{\mathbf{F}}$.

The same arguments apply to the case that $\alpha < 0$, $\bar{\mathbf{F}} > 0$, $\mathbf{F}_{\alpha,i}(u(0)) > 0$. We omit the details here. \square

Proof of Theorem 1.4: In the case of $\alpha > 0$, $\chi(S) < 0$, $\bar{\mathbf{F}} \leq 0$, by Theorem 3.3, there exists $\bar{u} \in \mathcal{U}(d_0)$ such that $\sum_i^N K_i^*(\bar{u}) > 2\pi\chi(S)$ and $K_i^*(\bar{u}) < 0$ for all $v_i \in V$. Then $\mathbf{F}_{\alpha,i}(u^*) < 0$ for all $v_i \in V$. By Theorem 3.19, there exists a PH metric in the discrete conformal class $\mathcal{D}(d_0)$ with non-positive combinatorial α -curvature $\bar{\mathbf{F}}$.

In the case of $\alpha < 0$, $\bar{\mathbf{F}} > 0$, by Theorem 3.3, there exists $u^* \in \mathcal{U}(d_0)$ such that $\mathbf{F}_{\alpha,i}(u^*) > 0$ for all $v_i \in V$. By Theorem 3.19, there exists a PH metric in the discrete conformal class $\mathcal{D}(d_0)$ with positive combinatorial α -curvature $\bar{\mathbf{F}}$.

The case of $\alpha = 0$ follows from Theorem 3.3. \square

As a corollary of Theorem 3.11 and Theorem 1.4, we have the following longtime existence and convergence for the solution of modified combinatorial α -Yamabe flow (19), which is the first part of Theorem 1.8.

Theorem 3.20. Suppose (S, V) is a connected closed marked surface with a PH metric d_0 , α is a constant and $\bar{\mathbf{F}} : V \rightarrow \mathbb{R}$ is a function defined on V . If one of the conditions (1)(2)(3) in Theorem 1.4 is satisfied, then the solution of modified combinatorial α -Yamabe flow with surgery (19) exists for all time and converges exponentially fast to a PH metric with combinatorial α -curvature $\bar{\mathbf{F}}$.

3.5 Combinatorial α -Calabi flow with surgery

Analogy to Definition 3.10 for combinatorial α -Yamabe flow with surgery, we define the following combinatorial α -Calabi flow with surgery.

Definition 3.21. Suppose (S, V) is a connected closed marked surface with a PH metric d_0 and $\alpha \in \mathbb{R}$ is a constant. The combinatorial α -Calabi flow with surgery is defined to be

$$\begin{cases} \frac{du_i}{dt} = \Delta_\alpha \mathbf{F}_{\alpha,i}, \\ u_i(0) = 0, \end{cases} \quad (23)$$

where Δ_α is the α -Laplace operator defined by (15).

One can use the following modified combinatorial α -Calabi flow with surgery to study prescribed combinatorial α -curvature problem on polyhedral surfaces

$$\frac{du_i}{dt} = \Delta_\alpha (\mathbf{F}_\alpha - \bar{\mathbf{F}})_i, \quad (24)$$

where $\bar{\mathbf{F}}$ is a function defined on V .

Theorem 3.22. Suppose (S, V) is a connected closed marked surface with a PH metric d_0 , $\alpha \in \mathbb{R}$ is a constant and $\bar{\mathbf{F}} : V \rightarrow \mathbb{R}$ is a function defined on V with $\alpha \bar{\mathbf{F}} \leq 0$. Then there exists a PH metric in the discrete conformal class $\mathcal{D}(d_0)$ with combinatorial α -curvature $\bar{\mathbf{F}}$ if and only if the solution of modified combinatorial α -Calabi flow with surgery (24) exists for all time and converges exponentially fast to a PH metric with combinatorial α -curvature $\bar{\mathbf{F}}$.

Proof. Suppose the solution $u(t)$ of the modified combinatorial α -Calabi flow with surgery (24) converges to $u^* \in \mathcal{U}(d_0)$ as $t \rightarrow +\infty$, then $\mathbf{F}_\alpha(u^*) = \lim_{t \rightarrow +\infty} \mathbf{F}_\alpha(u(t))$ by the C^1 -smoothness of \mathbf{F} . Furthermore, there exists a sequence $\xi_n \in (n, n+1)$ such that

$$u_i(n+1) - u_i(n) = u'_i(\xi_n) = \Delta_\alpha (\mathbf{F}_\alpha(u(\xi_n)) - \bar{\mathbf{F}})_i \rightarrow 0, \text{ as } n \rightarrow +\infty,$$

which implies $\mathbf{F}_{\alpha,i}(u^*) = \lim_{n \rightarrow +\infty} \mathbf{F}_{\alpha,i}(u(\xi_n)) = \bar{\mathbf{F}}_i$ for all $v_i \in V$ and $u^* \in \mathcal{U}(d_0)$ corresponds to a PH metric with combinatorial α -curvature $\bar{\mathbf{F}}$.

Conversely, suppose there exists a PH metric in the conformal class $\mathcal{D}(d_0)$ with combinatorial α -curvature $\bar{\mathbf{F}}$ and u^* is the corresponding discrete conformal factor in $\mathcal{U}(d_0)$. Set

$$W_\alpha(u) = W(u) - \int_{u^*}^u \sum_{i=1}^N \bar{\mathbf{F}}_i w_i^\alpha du_i = \int_{u^*}^u \sum_{i=1}^N \mathbf{F}_i du_i - \int_{u^*}^u \sum_{i=1}^N \bar{\mathbf{F}}_i w_i^\alpha du_i.$$

Then $W_\alpha(u)$ is a C^2 -smooth strictly convex function defined on \mathbb{R}^n by Proposition 3.5 and $\alpha\bar{\mathbf{F}} \leq 0$. Furthermore, $W_\alpha(u^*) = 0$, $\nabla W_\alpha(u^*) = 0$ and $\text{Hess}W_\alpha > 0$, which implies $W_\alpha(u) \geq W_\alpha(u^*) = 0$ and $\lim_{u \rightarrow \infty} W_\alpha(u) = +\infty$. By direct calculations, we have

$$\frac{dW_\alpha(u(t))}{dt} = \sum_{i=1}^N \frac{\partial W_\alpha}{\partial u_i} \frac{du_i}{dt} = \sum_{i=1}^N (\mathbf{F}_\alpha - \bar{\mathbf{F}})_i w_i^\alpha \Delta_\alpha (\mathbf{F}_\alpha - \bar{\mathbf{F}})_i = -(\mathbf{F}_\alpha - \bar{\mathbf{F}})^T \cdot \tilde{L} \cdot (\mathbf{F}_\alpha - \bar{\mathbf{F}}) \leq 0$$

by Proposition 3.5, which implies $0 \leq W_\alpha(u(t)) \leq W_\alpha(u(0))$. Thus $\{u(t)\} \subset \subset \mathbb{R}^N$ by $\lim_{u \rightarrow \infty} W_\alpha(u) = +\infty$, which implies the solution of the modified combinatorial α -Calabi flow with surgery (24) exists for all time and $W_\alpha(u(t))$ converges. Moreover, there exists a sequence $\xi_n \in (n, n+1)$ such that as $n \rightarrow +\infty$,

$$\begin{aligned} W_\alpha(u(n+1)) - W_\alpha(u(n)) &= (W_\alpha(u(t)))'|_{\xi_n} = \nabla W_\alpha \cdot \frac{du_i}{dt} \Big|_{\xi_n} \\ &= \sum_{i=1}^N (\mathbf{F}_\alpha - \bar{\mathbf{F}})_i w_i^\alpha \Delta_\alpha (\mathbf{F}_\alpha - \bar{\mathbf{F}})_i \Big|_{\xi_n} = -(\mathbf{F}_\alpha - \bar{\mathbf{F}})^T \cdot \tilde{L} \cdot (\mathbf{F}_\alpha - \bar{\mathbf{F}}) \Big|_{\xi_n} \rightarrow 0. \end{aligned}$$

Then $\lim_{n \rightarrow +\infty} \mathbf{F}_{\alpha,i}(u(\xi_n)) = \bar{\mathbf{F}}_i = \mathbf{F}_{\alpha,i}(u^*)$ for all $v_i \in V$, which implies $\lim_{n \rightarrow +\infty} u(\xi_n) = u^*$ by Theorem 3.8.

Similar to the proof of Theorem 2.6, set $\Gamma(u) = \Delta_\alpha(\mathbf{F}_\alpha - \bar{\mathbf{F}})$, then $D\Gamma|_{u=u^*}$ has N negative eigenvalues, which implies that u^* is a local attractor of (24). Then the conclusion follows by Lyapunov Stability Theorem ([21], Chapter 5). \square

As a corollary of Theorem 3.22 and Theorem 1.4, we have the following longtime existence and convergence for the solution of modified combinatorial α -Calabi flow with surgery (24), which is the second part of Theorem 1.8.

Theorem 3.23. Suppose (S, V) is a connected closed marked surface with a PH metric d_0 , α is a constant and $\bar{\mathbf{F}} : V \rightarrow \mathbb{R}$ is a function defined on V . If one of the conditions (1)(2)(3) in Theorem 1.4 is satisfied, then the solution of modified combinatorial α -Calabi flow with surgery (24) exists for all time and converges exponentially fast to a PH metric in $\mathcal{D}(d_0)$ with combinatorial α -curvature $\bar{\mathbf{F}}$.

A Proof of Lemma 2.2

We use $j \sim i$ to denote that the vertices v_i and v_j are adjacent. By Lemma 2.1, $\frac{\partial \alpha_i^{jk}}{\partial u_j} = \frac{\partial \alpha_j^{ik}}{\partial u_i}$ and $\frac{\partial K_i}{\partial u_j} = \frac{\partial K_j}{\partial u_i}$.

(1) If $j \approx i$ and $j \neq i$, then $\frac{\partial K_i}{\partial u_j} = 0$.

(2) If $j \sim i$, then

$$\frac{\partial K_i}{\partial u_j} = \frac{\partial(2\pi - \sum_{\Delta v_i v_j v_k \in F} \alpha_i^{jk})}{\partial u_j} = - \sum_{\Delta v_i v_j v_k \in F} \frac{\partial \alpha_i^{jk}}{\partial u_j} = -\left(\frac{\partial \alpha_i^{jk}}{\partial u_j} + \frac{\partial \alpha_i^{jl}}{\partial u_j}\right),$$

where $\Delta v_i v_j v_k$ and $\Delta v_i v_j v_l$ are adjacent triangles sharing a common edge $\{v_i v_j\}$.

Set $B_{ij} = \frac{\partial \alpha_i^{jk}}{\partial u_j} + \frac{\partial \alpha_i^{jl}}{\partial u_j}$, then $\frac{\partial K_i}{\partial u_j} = -B_{ij}$.

(3) If $j = i$, then

$$\begin{aligned} \frac{\partial K_i}{\partial u_i} &= \sum_{\Delta v_i v_j v_k \in F} \frac{\partial \left(\alpha_j^{ik} + \alpha_k^{ij} + \text{Area}(\Delta v_i v_j v_k) \right)}{\partial u_i} \\ &= \sum_{\Delta v_i v_j v_k \in F} \left(\frac{\partial \alpha_i^{jk}}{\partial u_j} + \frac{\partial \alpha_i^{jk}}{\partial u_k} \right) + \sum_{\Delta v_i v_j v_k \in F} \frac{\partial \text{Area}(\Delta v_i v_j v_k)}{\partial u_i} \\ &= \sum_{j \sim i} \left(\frac{\partial \alpha_i^{jk}}{\partial u_j} + \frac{\partial \alpha_i^{jl}}{\partial u_j} \right) + \frac{\partial}{\partial u_i} \left(\sum_{\Delta v_i v_j v_k \in F} \text{Area}(\Delta v_i v_j v_k) \right). \end{aligned}$$

Set $A_i = \frac{\partial}{\partial u_i} \left(\sum_{\Delta v_i v_j v_k \in F} \text{Area}(\Delta v_i v_j v_k) \right)$, then

$$\frac{\partial K_i}{\partial u_i} = A_i + \sum_{j \sim i} B_{ij}. \quad (25)$$

Set $L_A = \text{diag}\{A_1, \dots, A_N\}$ and $L_B = ((L_B)_{ij})_{N \times N}$, where

$$(L_B)_{ij} = \begin{cases} \sum_{k \sim i} B_{ik}, & j = i, \\ -B_{ij}, & j \sim i, \\ 0, & j \not\sim i, j \neq i. \end{cases}$$

Then $L = L_A + L_B$.

To get the expression of B_{ij} in (12), we need the following lemma.

Lemma A.1. Suppose $\Delta v_i v_j v_k$ is a hyperbolic triangle with a PH metric d_0 and u is a discrete conformal factor for the triangle. Denote $d_{st} := (u * d_0)_{st}$ as the edge length of $\{st\}$ in the triangle $\Delta v_i v_j v_k$ and $\alpha_i^{jk}, \alpha_j^{ik}, \alpha_k^{ij}$ as the inner angles facing $\{jk\}, \{ik\}, \{ij\}$ respectively. Then

$$\frac{\partial \alpha_i^{jk}}{\partial u_j} = \frac{1}{\cosh^2 \frac{d_{ij}}{2}} \tan \frac{\alpha_i^{jk} + \alpha_j^{ik} - \alpha_k^{ij}}{2}.$$

Proof. By the derivative cosine law (see Lemma A1 in [2] for example),

$$\frac{\partial \alpha_i^{jk}}{\partial d_{jk}} = \frac{\sinh d_{jk}}{A}, \quad \frac{\partial \alpha_i^{jk}}{\partial d_{ik}} = \frac{-\sinh d_{jk} \cos \alpha_k^{ij}}{A}, \quad \frac{\partial \alpha_i^{jk}}{\partial d_{ij}} = \frac{-\sinh d_{jk} \cos \alpha_j^{ik}}{A},$$

where $A = \sinh d_{ik} \sinh d_{ij} \sin \alpha_i^{jk}$. By (2), we have

$$\frac{\partial d_{jk}}{\partial u_i} = 0, \quad \frac{\partial d_{jk}}{\partial u_j} = \frac{\partial d_{jk}}{\partial u_k} = 2 \tanh \frac{d_{jk}}{2}.$$

According to the chain rules,

$$\begin{aligned} \frac{\partial \alpha_i^{jk}}{\partial u_j} &= \frac{\partial \alpha_i^{jk}}{\partial d_{jk}} \frac{\partial d_{jk}}{\partial u_j} + \frac{\partial \alpha_i^{jk}}{\partial d_{ik}} \frac{\partial d_{ik}}{\partial u_j} + \frac{\partial \alpha_i^{jk}}{\partial d_{ij}} \frac{\partial d_{ij}}{\partial u_j} \\ &= \frac{2 \sinh d_{jk}}{A} \tanh \frac{d_{jk}}{2} - \frac{2 \sinh d_{jk} \cos \alpha_j^{ik}}{A} \tanh \frac{d_{ij}}{2} \\ &= \frac{2(\cosh d_{jk} + \cosh d_{ik} - \cosh d_{ij} - 1)}{A(\cosh d_{ij} + 1)}. \end{aligned} \quad (26)$$

By the hyperbolic cosine law expressing d_{ik} in terms of α_i^{jk} , α_j^{ik} , α_k^{ij} , we have

$$\begin{aligned} \cosh^2 \frac{d_{ik}}{2} &= \frac{1}{2}(\cosh d_{ik} + 1) \\ &= \frac{1}{2} \left(\frac{\cos \alpha_j^{ik} + \cos \alpha_k^{ij} \cos \alpha_i^{jk}}{\sin \alpha_k^{ij} \sin \alpha_i^{jk}} + 1 \right) \\ &= \frac{\cos \frac{\alpha_j^{ik} + \alpha_k^{ij} - \alpha_i^{jk}}{2} \cos \frac{\alpha_j^{ik} - \alpha_k^{ij} + \alpha_i^{jk}}{2}}{\sin \alpha_k^{ij} \sin \alpha_i^{jk}}, \end{aligned} \quad (27)$$

which implies

$$\begin{aligned} &\cosh \frac{d_{jk}}{2} \cosh \frac{d_{ik}}{2} \sin \alpha_k^{ij} \\ &= \sqrt{\frac{\cos \frac{\alpha_i^{jk} + \alpha_j^{ik} - \alpha_k^{ij}}{2} \cos \frac{\alpha_i^{jk} - \alpha_j^{ik} + \alpha_k^{ij}}{2}}{\sin \alpha_j^{ik} \sin \alpha_k^{ij}}} \cdot \sqrt{\frac{\cos \frac{\alpha_j^{ik} + \alpha_k^{ij} - \alpha_i^{jk}}{2} \cos \frac{\alpha_j^{ik} - \alpha_k^{ij} + \alpha_i^{jk}}{2}}{\sin \alpha_k^{ij} \sin \alpha_i^{jk}}} \sin \alpha_k^{ij} \\ &= \cosh \frac{d_{ij}}{2} \cos \frac{\alpha_i^{jk} + \alpha_j^{ik} - \alpha_k^{ij}}{2}, \end{aligned} \quad (28)$$

where (27) is used in the last line. Recall the following formula obtained by Gu-Guo-Luo-Sun-Wu ([16] Lemma 11)

$$2 \sin \frac{\alpha_i^{jk} + \alpha_j^{ik} - \alpha_k^{ij}}{2} \cdot \cosh \frac{d_{ij}}{2} = \frac{\sinh^2 \frac{d_{jk}}{2} + \sinh^2 \frac{d_{ik}}{2} - \sinh^2 \frac{d_{ij}}{2}}{\sinh \frac{d_{jk}}{2} \sinh \frac{d_{ik}}{2}}. \quad (29)$$

Then (26) implies

$$\begin{aligned}
\frac{\partial \alpha_i^{jk}}{\partial u_j} &= \frac{2(\cosh d_{jk} + \cosh d_{ik} - \cosh d_{ij} - 1)}{\sinh d_{ik} \sinh d_{jk} \sin \alpha_k^{ij} (\cosh d_{ij} + 1)} \\
&= \frac{4(\sinh^2 \frac{d_{jk}}{2} + \sinh^2 \frac{d_{ik}}{2} - \sinh^2 \frac{d_{ij}}{2})}{\sinh d_{ik} \sinh d_{jk} \sin \alpha_k^{ij} (\cosh d_{ij} + 1)} \\
&= \frac{8 \sinh \frac{d_{jk}}{2} \sinh \frac{d_{ik}}{2} \cosh \frac{d_{ij}}{2}}{\sinh d_{ik} \sinh d_{jk} \sin \alpha_k^{ij} (\cosh d_{ij} + 1)} \sin \frac{\alpha_i^{jk} + \alpha_j^{ik} - \alpha_k^{ij}}{2} \\
&= \frac{1}{\cosh \frac{d_{ij}}{2} \cosh \frac{d_{jk}}{2} \cosh \frac{d_{ik}}{2} \sin \alpha_k^{ij}} \sin \frac{\alpha_i^{jk} + \alpha_j^{ik} - \alpha_k^{ij}}{2} \\
&= \frac{1}{\cosh^2 \frac{d_{ij}}{2}} \tan \frac{\alpha_i^{jk} + \alpha_j^{ik} - \alpha_k^{ij}}{2},
\end{aligned}$$

where (29) is used in the third line and (28) is used in the last line. \square

Lemma A.2. Suppose (S, V, \mathcal{T}) is a connected closed triangulated surface with a PH metric d_0 . $d := u * d_0$ is a PH metric defined by a function $u : V \rightarrow \mathbb{R}$. Then the triangulation \mathcal{T} is Delaunay in d if and only if $B_{ij} \geq 0$ for any $\{v_i v_j\} \in E$, which implies L_B is positive semi-definite. Furthermore,

$$A_i = \sum_{j \sim i} B_{ij} (\cosh d_{ij} - 1), \quad (30)$$

which is nonnegative if the triangulation \mathcal{T} is Delaunay in d .

Proof. By Lemma A.1, for $j \sim i$,

$$\begin{aligned}
B_{ij} &= \frac{\partial \alpha_i^{jk}}{\partial u_j} + \frac{\partial \alpha_i^{jl}}{\partial u_j} \\
&= \frac{1}{\cosh^2 \frac{d_{ij}}{2}} \left(\tan \frac{\alpha_i^{jk} + \alpha_j^{ik} - \alpha_k^{ij}}{2} + \tan \frac{\alpha_i^{jl} + \alpha_j^{il} - \alpha_l^{ij}}{2} \right) \\
&= \frac{1}{\cosh^2 \frac{d_{ij}}{2}} \cdot \frac{\sin \frac{\alpha_i^{jk} + \alpha_j^{ik} - \alpha_k^{ij} + \alpha_i^{jl} + \alpha_j^{il} - \alpha_l^{ij}}{2}}{\sin \frac{\pi - \alpha_i^{jk} - \alpha_j^{ik} + \alpha_k^{ij}}{2} \sin \frac{\pi - \alpha_i^{jl} - \alpha_j^{il} + \alpha_l^{ij}}{2}}.
\end{aligned} \quad (31)$$

Note that the triangulation \mathcal{T} is Delaunay if and only if $\alpha_i^{jk} + \alpha_j^{ik} + \alpha_i^{jl} + \alpha_j^{il} \geq \alpha_l^{ij} + \alpha_k^{ij}$ [16, 18], which is equivalent to $B_{ij} \geq 0$.

For any $x \in \mathbb{R}^N$, we have

$$x^T L_B x = \sum_{i,j=1}^N (L_B)_{ij} x_i x_j = \sum_{i \sim j} (-B_{ij} x_i x_j) + \sum_{i \sim j} x_i^2 B_{ij} = \frac{1}{2} \sum_{i \sim j} B_{ij} (x_i - x_j)^2 \geq 0,$$

which implies L_B is positive semi-definite. One can also refer to Lemma 3.10 in [2] for the positive semi-definiteness of L_B .

Recall the following formula obtained by Glickenstein-Thomas ([15] Proposition 9)

$$\frac{\partial \text{Area}(\Delta v_i v_j v_k)}{\partial u_k} = \frac{\partial \alpha_i^{jk}}{\partial u_k} (\cosh d_{ik} - 1) + \frac{\partial \alpha_j^{ik}}{\partial u_k} (\cosh d_{jk} - 1).$$

Then

$$\begin{aligned} A_i &= \sum_{\Delta v_i v_j v_k \in F} \frac{\partial \text{Area}(\Delta v_i v_j v_k)}{\partial u_i} \\ &= \sum_{\Delta v_i v_j v_k \in F} \frac{\partial \alpha_i^{jk}}{\partial u_j} (\cosh d_{ij} - 1) + \sum_{\Delta v_i v_j v_k \in F} \frac{\partial \alpha_i^{jk}}{\partial u_k} (\cosh d_{ik} - 1) \\ &= \sum_{j \sim i} \left(\frac{\partial \alpha_i^{jk}}{\partial u_j} + \frac{\partial \alpha_i^{jl}}{\partial u_j} \right) (\cosh d_{ij} - 1) \\ &= \sum_{j \sim i} B_{ij} (\cosh d_{ij} - 1). \end{aligned}$$

Therefore, $B_{ij} \geq 0$ implies $A_i \geq 0$. □

Remark 10. Submitting (12) into (30) gives

$$A_i = 2 \sum_{j \sim i} \tanh^2 \frac{d_{ij}}{2} \cdot \left(\tan \frac{\alpha_i^{jk} + \alpha_j^{ik} - \alpha_k^{ij}}{2} + \tan \frac{\alpha_i^{jl} + \alpha_j^{il} - \alpha_l^{ij}}{2} \right). \quad (32)$$

Then the decomposition $L = L_A + L_B$ with A_i given by (32) and B_{ij} given by (12) is equivalent to the formula obtained by Bobenko-Pinkall-Springborn ([1], Proposition 6.1.7).

We further have the following result on the sign of A_i under the Delaunay condition, which completes the proof of Lemma 2.2.

Lemma A.3. Suppose (S, V, \mathcal{T}) is a connected closed triangulated surface with a PH metric d_0 . $d := u * d_0$ is a PH metric defined by a function $u : V \rightarrow \mathbb{R}$. If the triangulation \mathcal{T} is Delaunay in d , then $A_i > 0$ for all $v_i \in V$, which implies L_A is positive definite.

Proof. Combining (25) and (30) gives

$$\frac{\partial K_i}{\partial u_i} = A_i + \sum_{j \sim i} B_{ij} = \sum_{j \sim i} B_{ij} \cosh d_{ij}. \quad (33)$$

For a single triangle $\Delta v_i v_j v_k$, by direct calculations, we have

$$\frac{\partial \alpha_i^{jk}}{\partial u_i} = \frac{\cosh^2 d_{ik} + \cosh^2 d_{ij} - 2 \cosh d_{jk} \cosh d_{ik} \cosh d_{ij} + (1 - \cosh d_{jk})(\cosh d_{ik} + \cosh d_{ij})}{A(1 + \cosh d_{ik})(1 + \cosh d_{ij})}, \quad (34)$$

where $A = \sinh d_{ik} \sinh d_{ij} \sin \alpha_i^{jk}$. Note that

$$\begin{aligned} \sin^2 \alpha_i^{jk} &= 1 - \left(\frac{\cosh d_{ik} \cosh d_{ij} - \cosh d_{jk}}{\sinh d_{ik} \sinh d_{ij}} \right)^2 \\ &= \frac{1}{\sinh^2 d_{ik} \sinh^2 d_{ij}} (-\cosh^2 d_{jk} - \cosh^2 d_{ik} - \cosh^2 d_{ij} + 1 + 2 \cosh d_{jk} \cosh d_{ik} \cosh d_{ij}), \end{aligned}$$

which implies

$$\begin{aligned} & -\cosh^2 d_{ik} - \cosh^2 d_{ij} + 2 \cosh d_{jk} \cosh d_{ik} \cosh d_{ij} + (\cosh d_{jk} - 1)(\cosh d_{ik} + \cosh d_{ij}) \\ &= \sinh^2 d_{ik} \sinh^2 d_{ij} \sin^2 \alpha_i^{jk} + \sinh^2 d_{jk} + (\cosh d_{jk} - 1)(\cosh d_{ik} + \cosh d_{ij}) \\ &> 0, \end{aligned}$$

we have $\frac{\partial \alpha_i^{jk}}{\partial u_i} < 0$ by (34). Then

$$\frac{\partial K_i}{\partial u_i} = \frac{\partial(2\pi - \sum_{\Delta v_i v_j v_k \in F} \alpha_i^{jk})}{\partial u_i} = - \sum_{\Delta v_i v_j v_k \in F} \frac{\partial \alpha_i^{jk}}{\partial u_i} > 0,$$

which implies $\sum_{j \sim i} B_{ij} > 0$ by (33) and $B_{ij} \geq 0$. Therefore, $A_i = \sum_{j \sim i} B_{ij}(\cosh d_{ij} - 1) > 0$ by (30). \square

After the paper was finished, Dr. Tianqi Wu and Dr. Xiaoping Zhu told us that there was a similar proof of Lemma 2.2 in [24]. We thank Dr. Tianqi Wu and Dr. Xiaoping Zhu for informing us this and communications on related topics. However, our proof is different. So we present our proof here.

References

- [1] A. Bobenko, U. Pinkall, B. Springborn, *Discrete conformal maps and ideal hyperbolic polyhedra*. *Geom. Topol.* 19 (2015), no. 4, 2155-2215.
- [2] B. Chow, F. Luo, *Combinatorial Ricci flows on surfaces*, *J. Differential Geom.*, Volume 63, No. 1 (2003), 97-129.
- [3] F. R. K. Chung, *Spectral graph theory*, *CBMS Regional Conference Series in Mathematics*, 92. American Mathematical Society, Providence, RI, 1997.
- [4] S. Dai, H. Ge, *Discrete Yamabe flows with R-curvature revisited*. *J. Math. Anal. Appl.* 484 (2020), no. 1, 123681, 11 pp.

- [5] H. Ge, *Combinatorial methods and geometric equations*, Thesis (Ph.D.)-Peking University, Beijing. 2012.
- [6] H. Ge, *Combinatorial Calabi flows on surfaces*, Trans. Amer. Math. Soc. 370 (2018), no. 2, 1377-1391.
- [7] H. Ge, B. Hua, *On combinatorial Calabi flow with hyperbolic circle patterns*, Adv. Math. 333 (2018), 523-538.
- [8] H. Ge, W. Jiang, *On the deformation of inversive distance circle packings, III*. J. Funct. Anal. 272 (2017), no. 9, 3596-3609.
- [9] H. Ge, X. Xu, *Discrete quasi-Einstein metrics and combinatorial curvature flows in 3-dimension*, Adv. Math. 267 (2014), 470-497.
- [10] H. Ge, X. Xu, *2-dimensional combinatorial Calabi flow in hyperbolic background geometry*, Differential Geom. Appl. 47 (2016) 86-98.
- [11] H. Ge, X. Xu, *α -curvatures and α -flows on low dimensional triangulated manifolds*. Calc. Var. Partial Differential Equations 55 (2016), no. 1, Art. 12, 16 pp.
- [12] H. Ge; X. Xu, *A discrete Ricci flow on surfaces with hyperbolic background geometry*, Int. Math. Res. Not. IMRN 2017, no. 11, 3510-3527.
- [13] H. Ge, X. Xu, *On a combinatorial curvature for surfaces with inversive distance circle packing metrics*. J. Funct. Anal. 275 (2018), no. 3, 523-558.
- [14] H. Ge, X. Xu, *A combinatorial Yamabe problem on two and three dimensional manifolds*, Calc. Var. Partial Differential Equations 60 (2021), no. 1, 20.
- [15] D. Glickenstein, J. Thomas, *Duality structures and discrete conformal variations of piecewise constant curvature surfaces*, Adv. Math. 320 (2017), 250-278.
- [16] X. D. Gu, R. Guo, F. Luo, J. Sun, T. Wu, *A discrete uniformization theorem for polyhedral surfaces II*, J. Differential Geom. 109 (2018), no. 3, 431-466.
- [17] X. D. Gu, F. Luo, J. Sun, T. Wu, *A discrete uniformization theorem for polyhedral surfaces*, J. Differential Geom. 109 (2018), no. 2, 223-256.
- [18] G. Leibon, *Characterizing the Delaunay decompositions of compact hyperbolic surface*, Geom. Topol. 6 (2002), 361-391.
- [19] F. Luo, *Combinatorial Yamabe flows on surfaces*, Commun. Contemp. Math. 6 (2004), no. 5, 765-780.
- [20] F. Luo, *Rigidity of polyhedral surfaces, III*, Geom. Topol. 15 (2011), 2299-2319.
- [21] L.S. Pontryagin, *Ordinary differential equations*, Addison-Wesley Publishing Company Inc., Reading, 1962.
- [22] B. Springborn, *Ideal hyperbolic polyhedra and discrete uniformization*. Discrete Comput. Geom. 64 (2020), no. 1, 63-108.
- [23] M. Röcek, R. M. Williams, *The quantization of Regge calculus*. Z. Phys. C 21 (1984), no. 4, 371-381.
- [24] T. Wu, X. Zhu, *The convergence of discrete uniformizations for closed surfaces*, arXiv:2008.06744v2 [math.GT].
- [25] X. Xu, *Rigidity of inversive distance circle packings revisited*, Adv. Math. 332 (2018), 476-509.
- [26] X. Xu, *On the global rigidity of sphere packings on 3-dimensional manifolds*, J. Differential Geom. 115 (2020), no. 1, 175-193.
- [27] X. Xu, *Parameterized discrete uniformization theorems and curvature flows for polyhedral surfaces, I*, arXiv:1806.04516v1 [math.GT].
- [28] X. Xu, C. Zheng, *A new proof for global rigidity of vertex scaling on polyhedral surfaces*, accepted by Asian J. Math. 2021.

- [29] W. Zeng, X. Gu, *Ricci flow for shape analysis and surface registration*, Springer Briefs in Mathematics. Springer, New York (2013).
- [30] X. Zhu, X. Xu, *Combinatorial Calabi flow with surgery on surfaces*, Calc. Var. Partial Differential Equations 58 (2019), no. 6, Paper No. 195, 20 pp.

(Xu Xu) School of Mathematics and Statistics, Wuhan University, Wuhan 430072, P.R. China

E-mail: xuxu2@whu.edu.cn

(Chao Zheng) School of Mathematics and Statistics, Wuhan University, Wuhan 430072, P.R. China

E-mail: 2019202010023@whu.edu.cn