
ON THE VALIDITY OF CLASSICAL PARTITION FUNCTION.

A PREPRINT

Sushil K. Singh
 Department of Physics
 SGTB Khalsa College
 University of Delhi

Savinder Kaur*
 Department of Physics
 SGTB Khalsa College
 University of Delhi

April 6, 2021

ABSTRACT

The partition function for a system of non-interacting N -particles can be found by summing over all the states of the system. The classical partition function for an ideal gas differs from Bosonic or Fermionic partition function in the classical regime. Students find it difficult to follow the differences arising out of incorrect counting by the classical partition function by missing out on the indistinguishability of particles and Fermi-Bose statistics. We present a pedagogical computer-based experiment to probe and demonstrate the key differences in the partition functions (i) Q , for the system of distinguishable and indistinguishable particles (ii) B , for Bosons and (ii) F , for Fermions without formally using the single-particle partition function.

Keywords Partition Function · Gibbs Gas · Fermions and Bosons

1 Introduction

The quintessential role that partition function (PF) plays in finding properties of a system is well established. The indistinguishability of identical particles has been a central point in the understanding of thermodynamic properties of the system. The thermodynamics function for a classical system can be obtained by defining the appropriate partition function

$$Q_N^* \equiv \frac{1}{A} Q_N$$

where A is a normalization factor and $Q_N = \int e^{\beta E_s}$ is the PF for indistinguishable N -particles. The conclusion that entropy should be an additive function for macroscopic objects, led Gibbs to propose that the constant A should be proportional to $N!$ in order not to overcount configurations of identical particles which are identical except for how the particles are labeled [1, 2]. Though, the idea is totally unintelligible within classical mechanics, because interchanging particles does lead to classically different states, much has been deliberated on the appropriate normalization factor A [3–6]. For most purposes, classical physics suffices to describe a system of N -particles under laboratory conditions and the indistinguishability is often incorporated by a judicious division by $N!$ [8, 9]. The factor $N!$, although correctly accounts for all states in which no orbital is occupied by more than one atom, fails for multiple occupancy [5]. We are tempted to summarize this discussion by saying that the number of quantum states for identical Fermions n_F and that for identical Bosons n_B are given by $n_F = n_B = n_C/N!$ where n_C is the number of states for identical classical atoms. The division by $N!$ overcounts Fermi states and undercounts Bose states with the consequence that the PFs [5, 6]

$$B_N > Q_N^* > F_N$$

where B_N and F_N represent the PF for N non-interacting Bosons and Fermions respectively. Though the topic has been explored in detail [4–6], the subtlety of the subject demands exploration and implementation in the computational lab. Despite the significance of this result, a pedagogical solution is not explicitly present in the literature. Our goal is to perform a *computer based experiment* (written in open source software Python) to analyse the differences between PFs, Q_N , B_N , Q_N^* , F_N and the state functions which are usually overlooked in textbooks.

*sk_savinder2005@yahoo.co.in

2 Canonical Ensemble

Consider a system of N -particles occupying a volume V maintained at a constant temperature T , through thermal contact with a heat bath. A macrostate for the system can be realized by a large number of microstates with *equal a priori probabilities*. The microstate is defined as any possible individual configuration in the phase space. The system resides in any of these microstates at an instant t while continually switching from one microstate to another. The more the number of available microstates the more the system spends time in the corresponding macrostate. The state functions are then expected to be "averaged" over these microstates through which the system passes [7]. An entirely different approach would be to imagine all these microstates to be occupied by *copies* of the system at the same instant. The average behavior of the system in this collection or *ensemble* would be identical to the time-averaged behavior of the given system. Statistically, the system evolves towards a macrostate with the largest number of microstates and spends an "overwhelmingly" large fraction of its time in this macrostate [1, 7, 8].

2.1 The "Canonical" Distribution Function

For the system in equilibrium with a thermal reservoir at temperature T , the Boltzmann factor $\exp(-\beta E_s)$ determines the "canonical" probability distribution P_s for state s

$$P_s = \frac{e^{-\beta E_s}}{\sum_r^{\text{all states}} e^{-\beta E_r}} \quad (1)$$

where s, r refer to complete orthonormal set of energy eigenstates of the system. The properties of the reservoir appear only in the scalar factor $\beta = 1/k_B T$ where k_B is the Boltzmann's constant.

2.2 The Partition Function

The general recipe to extract information about the various macroscopic properties of the given system is to find the quantity $Q_N(V, T)$ called the "sum-over-states" (German: Zustandssumme) or the partition function

$$Q_N(V, T) = \sum_s^{\text{all states}} e^{-\beta E_s} \quad (2)$$

The dependence on N and V comes through the energy eigenvalues E_s . In most physical cases the energy levels *accessible* to a system are degenerate, that is, one has a group of states, g_s in number, all belonging to the same energy value E_s . In such cases it is more useful to write the partition function as

$$Q_N(V, T) = \sum_s^{\text{energy states}} g_s e^{-\beta E_s} \quad (3)$$

The partition function counts the number of states available to the system.

The connection with thermodynamics is made by the Helmholtz Free energy ($H = E - TS$)

$$H_N(V, T) = -\frac{1}{\beta} \ln Q_N \quad (4)$$

which is used to determine all other state functions of the system.

3 Computer Based Experiment (CBE)

Initially, we consider a system of *distinguishable* N -particles in which each particle can access n number of single-particle energy levels ϵ_r . There are $M = n^N$ possible combination of energy states available to the system. Each state of the system is characterized by energy E_s ($s = 1, \dots, M$)

$$E_s = \epsilon_{r_1}^s + \epsilon_{r_2}^s + \dots + \epsilon_{r_N}^s \quad (5)$$

where $\epsilon_{r_m}^s$ ($m = 1, \dots, N$ and $r = 1, \dots, n$) denotes energy of single-particle state when the m^{th} particle is in its r^{th} level while the system is in the s^{th} state. The PF Q_N can be written as

$$Q_N = \sum_{s=1}^M e^{-\beta \sum_{m=1}^N \epsilon_{r_m}^s} = \sum_{s=1}^M \left(\prod_{m=1}^N e^{-\beta \epsilon_{r_m}^s} \right) \quad (6)$$

For *distinguishable* particles it is often useful to define a single-particle PF $q = \sum_r e^{-\beta\epsilon_r}$ where ϵ_r belongs to the complete orthonormal set of single-particle observable energies. The N -particle PF can then be found as $Q_N = q^N$. However, this single-particle PF q is not a relevant quantity when we are dealing with quantum statistics, that is, Fermi-Dirac or Bose-Einstein statistics. The Fermionic and Bosonic particles are indistinguishable and are not independent because of the symmetry requirements of the wave functions [4, 8]. We thus approach the problem without using the single-particle PF. We numerically calculate PFs for an N -particle system with each particle being able to access n energy levels. All the three cases (a) the distinguishable system (b) the Bosonic system and (c) the Fermionic system will be considered to find the differences arising in the number of available states.

To illustrate the *CBE*, consider 2 identical *distinguishable* particles each of which can access the 3 energy levels labelled 0 (ϵ_0), 1 (ϵ_1) and 2 (ϵ_2). There are $3^2 = 9$ available states $S_p(i, j)$ to the system where p refers to a state in which the particle 1 is in the i^{th} energy level and particle 2 is in the j^{th} energy level.

$S_p(i, j)$	(i) ($\epsilon_0, \epsilon_1, \epsilon_2$)		(ii) (0, 1, 2)	
	E_s	g_s	E_s	g_s
$S_1(0, 0)$	$E_0 = 2\epsilon_0$	1	0	1
$S_2(0, 1), S_3(1, 0)$	$E_1 = \epsilon_0 + \epsilon_1$	2	1	2
$S_4(0, 2), S_5(2, 0)$	$E_2 = \epsilon_0 + \epsilon_2$	2	2	3
$S_6(1, 1)$	$E_3 = 2\epsilon_1$	1	3	2
$S_7(1, 2), S_8(2, 1)$	$E_4 = \epsilon_1 + \epsilon_2$	2	4	1
$S_9(2, 2)$	$E_5 = 2\epsilon_2$	1		

Table 1: Distinguishable Particles : $S_p(i, j)$ observable states for 2-particle 3-level system, E_s energy and g_s degeneracy for levels (i) ($\epsilon_0, \epsilon_1, \epsilon_2$) & (ii) in particular ($\epsilon_0 = 0, \epsilon_1 = 1, \epsilon_2 = 2$)

The Table 1 shows 9 possible states of the system ($p = 1, 2 \dots 9$). States with label 2 and 3 i.e. $S_2(0, 1)$ and $S_3(1, 0)$ in which the particles just exchange their energy levels will be degenerate. Thus energy with label 1 i.e. E_1 has degeneracy $g_1 = 2$. States with same energy are listed together and thus there are only 6 distinct energy E_s of the system (label s is 0 to 5) with degeneracy g_s . Table 1 also lists the allowed energy states when single-particle energy levels $\epsilon_0 = 0, \epsilon_1 = 1$ and $\epsilon_2 = 2$. Apart from the exchange degeneracy, the choice of ϵ 's also leads to additional degeneracy in E_2 and E_3 .

$S_p(i, j)$	(i) ($\epsilon_0, \epsilon_1, \epsilon_2$)		(ii) (0, 1, 2)	
	E_s	g_s	E_s	g_s
$S_1(0, 0)$	$E_0 = 2\epsilon_0$	1	0	1
$S_2(0, 1) = S_3(1, 0)$	$E_1 = \epsilon_0 + \epsilon_1$	1	1	1
$S_4(0, 2) = S_5(2, 0)$	$E_2 = \epsilon_0 + \epsilon_2$	1	2	2
$S_6(1, 1)$	$E_3 = 2\epsilon_1$	1		
$S_7(1, 2) = S_8(2, 1)$	$E_4 = \epsilon_1 + \epsilon_2$	1	3	1
$S_9(2, 2)$	$E_5 = 2\epsilon_2$	1	4	1

Table 2: Same as Table 1 but for Bosons

FD (ϵ_1, ϵ_2)	(i) ($\epsilon_0, \epsilon_1, \epsilon_2$)		(ii) (0, 1, 2)	
	E_s	g_s	E_s	g_s
$S_1(0, 0)$	\times	\times	\times	\times
$S_2(0, 1) = S_3(1, 0)$	$E_1 = \epsilon_0 + \epsilon_1$	1	1	1
$S_4(0, 2) = S_5(2, 0)$	$E_2 = \epsilon_0 + \epsilon_2$	1	2	1
$S_6(1, 1)$	\times	\times	\times	\times
$S_7(1, 2) = S_8(2, 1)$	$E_4 = \epsilon_1 + \epsilon_2$	1	3	1
$S_9(2, 2)$	\times	\times	\times	\times

Table 3: Same as Table 1 but for Fermions

The Table 2 lists states available for the same system but with indistinguishable Bosonic particles along with energy eigen-states E_s and their degeneracies g_s . The symbols have the same meaning as in Table 1. Since the Bosons are indistinguishable, states $S_2(0, 1)$ and $S_3(1, 0)$ are now the same state with just two different labels. This has to be counted only once as exchange of particle for indistinguishable particles is meaningless. Thus the degeneracy g_s for corresponding energy E_1 is now counted as $g_1 = 1$ and not 2. Thus all the exchange degeneracy are removed. However, the choice of $\epsilon_0 = 0, \epsilon_1 = 1$ and $\epsilon_2 = 2$ leads to a degeneracy in energy states E_2 and E_3 .

The Table 3 lists states available for the same system but with indistinguishable Fermions along with energy eigen-states E_s and their degeneracies g_s . Fermions are indistinguishable particles and have additional restriction that no two particles occupy the same single-particle state (Pauli Exclusion principle). Thus the exchange degeneracy are again removed and the Pauli Exclusion principle disqualifies some states (S_1, S_6 & S_9) as being observable. Degeneracy may arise due to choice of ϵ for a larger system.

The PF Q, B and F for the three systems are then evaluated. The PF for the Gibbs gas is also evaluated as $Q^* = Q/N!$. The program is provided in the Appendix A.

4 Results

4.1 Partition Function

The PFs for 2-particle 3-level distinguishable system Q_N , Gibbs gas Q_N^* , Boson B_N and Fermion F_N are plotted with respect to dimensionless temperature T/T_0 where $T_0 = \epsilon_0/k_B$ in Figure 1. At high T/T_0 values (i) $Q \rightarrow 9$ for




Figure 1: 2-particle 3-level Partition Function




Figure 2: 3-particle 10-level Partition Function

distinguishable particles, (ii) $B \rightarrow 6$ for Bosonic particles, (iii) $F \rightarrow 3$ for Fermionic system and (iv) $Q^* \rightarrow 4.5$ for Gibbs gas. They reflect the available energy states as can be verified from the respective Tables 1, 2 and 3. Further, for all temperatures the partition function for Gibbs gas follows the inequality $B_N > Q_N^* > F_N$ [5, 6].

In figures 2 – 6 we consider a modest 3-particles and 10-level system which has 10^3 accessible energy states. Computationally, we need finite particles and levels to study the behaviour as we scale up the system. The 1000 available states are adequate to show that the $1/N!$ term is justified as we choose larger number of particles.

The PF for 3-particle 10-level distinguishable system, Gibbs gas, Boson and Fermion are shown in Figure 2. Though the system is still small, the factor of $1/N!$ used to scale down the distinguishable system PF Q to that for a Gibbs gas Q^* seems to predict values in-between those of B and F .

4.2 State Functions

For the 3-particle 10-level system (with $\epsilon_0 = 1, \epsilon_1 = 2, \dots$ and $\epsilon_9 = 10$) the non-dimensional state functions $H/\epsilon_0, S, E/\epsilon_0$ and C_v/ϵ_0 per particle are presented with T/T_0 in Figures 3, 4, 5 and 6 respectively. Though the system is still




Figure 3: The Helmholtz Free Energy




Figure 4: The Entropy

small, the factor of $1/N!$ used to scale down the distinguishable system partition function Q to that for a Gibbs gas Q^* seems to predict values of the state functions close to those for B and F . We see that the logarithmic dependence of the state function on the partition function bring the Gibbs gas closer to Boson and Fermion gas [2].




Figure 5: The Energy




Figure 6: The Specific heat

5 Conclusions

We find that (a) the PFs Q_N^* , B_N and F_N are different and (b) by taking large number of particles with even larger number of energy states, the state functions for Bosons and Fermions approach those for the classical case because the state functions have logarithmic dependence on the PF.

The advantage of this approach is that (i) students by ignoring the single-particle PF understand states available to the system itself (ii) the PF correctly counts the available states and provides an interpretation to the PF. (iii) Even though the PFs Q_N , B_N and F_N differ the state functions can be described by using the $Q_N^* = 1/N! \times Q_N$

References

- [1] A J Berlinsky and A B Harris, *Statistical Mechanics: An Introductory Graduate Course*, Springer Nature, Switzerland, 2019
- [2] L D Landau and E M Lifshitz, *Statistical Physics*, Addison-Wesley, Reading, MA, 1958
- [3] H A Buchdahl, *Remark on the Factor $1/N!$ in the Partition Function*, Am. J. Phys. **42** : 51-53, 1974
- [4] D I Ford, *A note on the partition function for systems of Independent Particles*, Am. J. Phys. **39** : 215-220, 1971
- [5] H Kroemer, *How incorrect is the classical partition function for the ideal gas?*, Am. J. Phys. **48** : 962, 1980
- [6] R. Baierlein, *The fraction of "all different" combinations: Justifying the semi-classical partition function*, Am. J. Phys. **65**(4) : 314-316, 1997
- [7] R K Pathria and P D Beale, *Statistical Mechanics*, Elsevier, United States, 2011
- [8] D A McQuarrie, *Statistical Mechanics*, Harper and Row, New York, 1976
- [9] C Kittle and H Kroemer, *Thermal Physics*, Freeman, San Francisco, 1980