

Obliquely propagating ion-acoustic shock waves in degenerate quantum plasma

M.K. Islam^{1,*}, S. Biswas^{1,**}, N.A. Chowdhury^{2,***}, A. Mannan^{1,†}, M. Salahuddin^{1,‡}, and A.A. Mamun^{1,§}

¹ Department of Physics, Jahangirnagar University, Savar, Dhaka-1342, Bangladesh

² Plasma Physics Division, Atomic Energy Centre, Dhaka-1000, Bangladesh

e-mail: *islam.stu2018@juniv.edu, **shawonbiswas440@gmail.com, ***nurealam1743phy@gmail.com,

†abdulmannan@juniv.edu, ‡su_2960@juniv.edu, §mamun_phys@juniv.edu

arXiv:2104.02121v1 [physics.plasm-ph] 5 Apr 2021

Abstract

A theoretical investigation has been carried out on the propagation of nonlinear ion-acoustic shock waves (IASHWs) in a collisionless magnetized degenerate quantum plasma system composed of inertial non-relativistic positively charged light and heavy ions, inertialess ultra-relativistically degenerate electrons and positrons. The reductive perturbation method has been employed to drive the Burgers' equation. It has been observed that under consideration, our plasma model supports only positive potential shock structure. It is also found that the amplitude and steepness of the IASHWs have been significantly modified by the variation of ion kinematic viscosity, oblique angle, number density, and charge state of the plasma species. The results of our present investigation will be helpful for understanding the propagation of IASHWs in white dwarfs and neutron stars.

Keywords: Shock waves; Degenerate quantum plasma; White dwarfs; Neutron stars.

1. Introduction

The existence of light (viz., ^1H [1, 2, 3, 4], ^4He [1, 5, 6, 7, 8], ^{12}C [7, 8], and ^{16}O [7, 8, 9], etc.) and heavy (viz., ^{56}Fe [10, 11], ^{85}Rb [11, 12], and ^{96}Mo [11, 12], etc.) ions in astrophysical compact objects (viz., white dwarfs and neutron stars) has received a substantial attention to investigate ion-acoustic (IA) waves (IAWs) in degenerate quantum plasma (DQP). The particle number density in the white dwarfs (i.e., $6 \times 10^{29} \text{cm}^{-3}$) and neutron stars (i.e., $6 \times 10^{36} \text{cm}^{-3}$) is extremely high [1, 9, 11, 13]. This extreme number density of these degenerate particles dictates to follow the Heisenberg's uncertainty principle, and according to the uncertainty principle when the position of these particles is confined then the momentum of these particles tends to very large. This excessive momentum leads to generate extreme outward degenerate pressure which is counterbalanced by the inward gravitational compression. The dynamics of these particles is mathematically modeled under two categories, namely, non-relativistic and ultra-relativistic limits by Chandrasekhar [1]. The pressure of non-relativistic light and heavy ions is expressed as $P_j = K_j N_j^\alpha$; where $\alpha = 5/3$, $K_j = 3\pi\hbar^2/5m_j$, $j = l$ for light ion, $j = h$ for heavy ion, and \hbar is the Planck constant [14, 15, 16]. The degenerate pressure of ultra-relativistic electrons and positrons can be expressed as $P_s = K_s N_s^{7/3}$; where $s = e$ for electron, $s = p$ for positron, $\gamma_s = 4/3$, $K_s = 3\hbar c/4$, and c is the speed of light [14, 15, 16].

The presence of positrons in white dwarfs and neutron stars has been extensively discussed in the Refs. [17, 18, 19, 20]. Sultana and Schlickeiser [16] investigated IA solitary waves in DQP containing degenerate electrons, light ions, and inertial mobile non-degenerate heavy ions. Gill *et al.* [21] investigated the IA shock waves (IASHWs) in relativistic DQP composed of

electrons, positrons and ions, and found that the height of the potential is maximum for the lower positron density. Ata-ur-Rahman *et al.* [22] considered an unmagnetized DQP containing inertial ions, and inertialess electrons and positrons, and demonstrated that the amplitude of the IAWs decreases with the increase of positron number density. Hossen *et al.* [23] studied IASHWs in a four-component plasma system having inertialess electrons and positrons, and inertial heavy and light ions, and reported that the amplitude of the shock profile decreases with positron number density. Mamun and Shukla [24] investigated electrostatic solitary waves propagating in ultra-relativistic plasma medium consisting of degenerate electrons and cold mobile ions, and reported that the wave amplitude increases with the increase of ion number density.

The strong magnetic field (i.e., about 1 Mega Gauss) in white dwarfs was predicted by Blackett [25] and observed by Zeeman spectroscopy [26, 27]. El-Taibany *et al.* [28] analyzed solitary waves in a magnetized degenerate electron-positron plasma, and found that the wave amplitude increases with the oblique angle which is the angle between the external magnetic field and the direction of wave propagation. Shaukat [29] investigated IA solitary waves in the presence of magnetic field, and observed that the solitary wave amplitude increases with increasing oblique angle.

The energy dissipation of the shock wave, which is governed by the Burgers' equation [30], may arise due to the kinematic viscosity of the medium. Hafez *et al.* [31] studied IASHWs in weakly relativistic plasma containing electrons, positrons, and ions, and noticed that the steepness of the IASHWs decreases with the increase in the value of viscosity of plasma species but the amplitude of the IASHWs does not change. Abdelwahed *et al.* [32] analyzed IASHWs in a pair-ion plasma, and also found

that the shock steepness decreases with increasing ion viscosity.

Recently, Saini *et al.* [33] investigated heavy nucleus acoustic periodic waves in DQP. Haider [34] examined the shock profiles in the presence of degenerate inertial ions, and inertialess electrons and positrons. Atteya *et al.* [35] studied IASHWs in a DQP which contains ion fluids, degenerate electrons, and stationary heavy ions. To the best of authors' knowledge, still no one investigated IASHWs in a magnetized DQP having inertialess ultra-relativistically degenerate electrons and positrons, and inertial positively charged non-relativistic light and heavy ions. In this manuscript, we will derive the Burgers' equation, and will also use associated solution to examine the basic features of IASHW in DQP.

The manuscript is organized in the preceding way: The governing equations are described in section 2. The Burgers' equation and associated shock solution are presented in section 3. The results and discussion are presented in section 4. A brief conclusion is presented in section 5.

2. Governing Equations

We consider a magnetized DQP system consisting of inertial positively charged light ion (mass m_l ; charge eZ_l ; number density N_l), positively charged heavy ion (mass m_h ; charge eZ_h , number density N_h), inertialess electron (mass m_e ; charge $-e$, number density N_e), and positron (mass m_p ; charge e ; number density N_p); where Z_l (Z_h) is the charge state of the light (heavy) ion. An uniform external magnetic field \mathbf{B} is existing in the direction of z -axis ($\mathbf{B} = B_0\hat{z}$ and \hat{z} is the unit vector). The propagation of IAWs in DQP system is governed by the following equations:

$$\frac{\partial N_h}{\partial T} + \tilde{\nabla} \cdot (N_h U_h) = 0, \quad (1)$$

$$\frac{\partial U_h}{\partial T} + (U_h \cdot \tilde{\nabla})U_h = -\frac{Z_h e}{m_h} \nabla \tilde{\Phi} + \frac{Z_h e B_0}{m_h} (U_h \times \hat{z}) - \frac{1}{m_h N_h} \tilde{\nabla} P_h + \tilde{\eta}_h \tilde{\nabla}^2 U_h, \quad (2)$$

$$\frac{\partial N_l}{\partial T} + \tilde{\nabla} \cdot (N_l U_l) = 0, \quad (3)$$

$$\frac{\partial U_l}{\partial T} + (U_l \cdot \tilde{\nabla})U_l = -\frac{Z_l e}{m_l} \tilde{\nabla} \tilde{\Phi} + \frac{Z_l e B_0}{m_l} (U_l \times \hat{z}) - \frac{1}{m_l N_l} \tilde{\nabla} P_l + \tilde{\eta}_l \tilde{\nabla}^2 U_l, \quad (4)$$

$$\tilde{\nabla}^2 \tilde{\Phi} = 4\pi e(N_e - N_p - Z_l N_l - Z_h N_h), \quad (5)$$

where U_l (U_h) is the fluid speed of light (heavy) ion; $\tilde{\Phi}$ is the electrostatic wave potential; P_l (P_h) is the pressure for light (heavy) ion; $\tilde{\eta}_l = \mu/m_l N_l$ ($\tilde{\eta}_h = \mu/m_h N_h$) is the kinematic viscosity for light (heavy) ion. The degenerate pressure equations for electrons and positrons can be expressed, respectively, as

$$\tilde{\nabla} \tilde{\Phi} - \frac{1}{e N_e} \tilde{\nabla} P_e = 0, \quad (6)$$

$$\tilde{\nabla} \tilde{\Phi} + \frac{1}{e N_p} \tilde{\nabla} P_p = 0. \quad (7)$$

Now, we have introduced the normalizing parameters: $n_h \rightarrow N_h/n_{h0}$; $n_l \rightarrow N_l/n_{l0}$; $n_e \rightarrow N_e/n_{e0}$; $n_p \rightarrow N_p/n_{p0}$; $u_h \rightarrow U_h/C_h$; $u_l \rightarrow U_l/C_h$; $\phi \rightarrow e\tilde{\Phi}/m_e c^2$; $t \rightarrow \omega_{ph} T$; $\nabla \rightarrow \lambda_{Dh}^{-1} \tilde{\nabla}$; $\eta \rightarrow \tilde{\eta}/\omega_{ph} \lambda_{Dh}^2$ [where IAWs speed $C_h = (Z_h m_e c^2/m_h)^{1/2}$; plasma frequency $\omega_{ph} = (4\pi Z_h^2 e^2 n_{h0}/m_h)^{1/2}$; the Debye length $\lambda_{Dh} = (m_e c^2/4\pi Z_h e^2 n_{h0})^{1/2}$, and for simplicity we have considered $\eta = \eta_l = \eta_h$]. At equilibrium, the charge neutrality condition can be written as $n_{e0} = n_{p0} + Z_l n_{l0} + Z_h n_{h0}$. By using these normalizing parameters, Eqs. (1)-(5) can be expressed in the normalized form

$$\frac{\partial n_h}{\partial t} + \nabla \cdot (n_h u_h) = 0, \quad (8)$$

$$\frac{\partial}{\partial t}(u_h) + (u_h \cdot \nabla)u_h = -\nabla \phi + \Omega_{ch}(u_h \times \hat{z}) - \frac{\mu_1 K_1}{n_h} \nabla n_h^\alpha + \eta \nabla^2 u_h, \quad (9)$$

$$\frac{\partial n_l}{\partial t} + \nabla \cdot (n_l u_l) = 0, \quad (10)$$

$$\frac{\partial}{\partial t}(u_l) + (u_l \cdot \nabla)u_l = -\mu_2 \nabla \phi + \mu_2 \Omega_{cl}(u_l \times \hat{z}) - \frac{\mu_3 K_2}{n_l} \nabla n_l^\alpha + \eta \nabla^2 u_l, \quad (11)$$

$$\nabla^2 \phi = (1 + \mu_4 + \mu_5)n_e - \mu_4 n_p - \mu_5 n_l - n_h, \quad (12)$$

where $\Omega_{ch} = \omega_{ch}/\omega_{ph}$, $\mu_1 = 1/Z_h$, $\mu_2 = Z_l m_h/Z_h m_l$, $\mu_3 = m_h/Z_h m_e$, $\mu_4 = n_{p0}/Z_h n_{h0}$, $\mu_5 = Z_l n_{l0}/Z_h n_{h0}$, $K_1 = n_{h0}^{\alpha-1} K_h/m_e c^2$, and $K_2 = n_{l0}^{\alpha-1} K_l/m_e c^2$. Now, by normalizing and integrating Eqs. (6) and (7), the number densities of the inertialess electrons and positrons can be obtained in terms of electrostatic potential ϕ , respectively, as

$$n_e = \left[1 + \frac{\gamma_e - 1}{\gamma_e K_3} \phi \right]^{\frac{1}{\gamma_e - 1}}, \quad (13)$$

$$n_p = \left[1 - \frac{\gamma_p - 1}{\gamma_p K_4} \phi \right]^{\frac{1}{\gamma_p - 1}}, \quad (14)$$

where $K_3 = n_{e0}^{\gamma_e-1} K_e/m_e c^2$ and $K_4 = n_{p0}^{\gamma_p-1} K_p/m_e c^2$. By expanding the right hand side of Eqs. (13) and (14) up to second order in ϕ , and substituting in Eq. (12), we get

$$\nabla^2 \phi + \mu_5 n_l + n_h = 1 + \mu_5 + \beta_1 \phi + \beta_2 \phi^2 + \dots, \quad (15)$$

where

$$\beta_1 = \frac{[\gamma_p K_4 (1 + \mu_4 + \mu_5) + \mu_4 \gamma_e K_3]}{\gamma_e \gamma_p K_3 K_4},$$

$$\beta_2 = \frac{[(\gamma_p K_4)^2 (1 + \mu_4 + \mu_5) (2 - \gamma_e) - \mu_4 (\gamma_e K_3)^2 (2 - \gamma_p)]}{2(\gamma_e \gamma_p K_3 K_4)^2}.$$

3. Derivation of the Burgers' Equation

To study IASHWs, we derive Burgers' equation by employing the reductive perturbation method (RPM) [36, 37], and the stretched coordinates for independent variables can be written as [38, 39, 40]

$$\xi = \epsilon(l_x x + l_y y + l_z z - v_p t), \quad (16)$$

$$\tau = \epsilon^2 t, \quad (17)$$

where v_p is the phase speed and ϵ is a smallness parameter measuring the weakness of the dissipation ($0 < \epsilon < 1$). The l_x , l_y , and l_z are the directional cosines of \mathbf{k} (wave vector) along x , y , and z -axes, respectively (i.e., $l_x^2 + l_y^2 + l_z^2 = 1$). Then, the dependent variables can be expressed in power series of ϵ as [39]

$$n_h = 1 + \epsilon n_h^{(1)} + \epsilon^2 n_h^{(2)} + \dots, \quad (18)$$

$$n_l = 1 + \epsilon n_l^{(1)} + \epsilon^2 n_l^{(2)} + \dots, \quad (19)$$

$$u_{hx,y} = \epsilon^2 u_{hx,y}^{(1)} + \epsilon^3 u_{hx,y}^{(2)} + \dots, \quad (20)$$

$$u_{lx,y} = \epsilon^2 u_{lx,y}^{(1)} + \epsilon^3 u_{lx,y}^{(2)} + \dots, \quad (21)$$

$$u_{hz} = \epsilon u_{hz}^{(1)} + \epsilon^2 u_{hz}^{(2)} + \dots, \quad (22)$$

$$u_{lz} = \epsilon u_{lz}^{(1)} + \epsilon^2 u_{lz}^{(2)} + \dots, \quad (23)$$

$$\phi = \epsilon \phi^{(1)} + \epsilon^2 \phi^{(2)} + \dots. \quad (24)$$

Now, by substituting Eqs. (16)-(24) into Eqs. (8)-(11) and (15), and collecting the terms containing ϵ , the first-order equations reduce to

$$n_h^{(1)} = \frac{l_z^2}{(v_p^2 - \alpha \mu_1 l_z^2 K_1)} \phi^{(1)}, \quad (25)$$

$$u_{hz}^{(1)} = \frac{v_p l_z}{(v_p^2 - \alpha \mu_1 l_z^2 K_1)} \phi^{(1)}, \quad (26)$$

$$n_l^{(1)} = \frac{\mu_2 l_z^2}{(v_p^2 - \alpha \mu_3 l_z^2 K_2)} \phi^{(1)}, \quad (27)$$

$$u_{lz}^{(1)} = \frac{\mu_2 v_p l_z}{(v_p^2 - \alpha \mu_3 l_z^2 K_2)} \phi^{(1)}. \quad (28)$$

Now, the phase speed of IASHWs can be written as

$$v_p = v_{p+} = l_z \sqrt{\frac{a + \sqrt{a^2 - 4\beta_1 b}}{2\beta_1}}, \quad (29)$$

$$v_p = v_{p-} = l_z \sqrt{\frac{a - \sqrt{a^2 - 4\beta_1 b}}{2\beta_1}}, \quad (30)$$

where $a = \alpha \mu_1 \beta_1 K_1 + \alpha \mu_3 \beta_1 K_2 + \mu_2 \mu_5 + 1$ and $b = \alpha^2 \mu_1 \mu_3 \beta_1 K_1 K_2 - \alpha \mu_1 \mu_2 \mu_5 K_1 - \alpha \mu_3 K_2$. The x and y -components of the first-order momentum equations can be manifested as

$$u_{hx}^{(1)} = -\frac{l_y v_p^2}{\Omega_{ch}(v_p^2 - \alpha \mu_1 l_z^2 K_1)} \frac{\partial \phi^{(1)}}{\partial \xi}, \quad (31)$$

$$u_{hy}^{(1)} = \frac{l_x v_p^2}{\Omega_{ch}(v_p^2 - \alpha \mu_1 l_z^2 K_1)} \frac{\partial \phi^{(1)}}{\partial \xi}, \quad (32)$$

$$u_{lx}^{(1)} = -\frac{l_y v_p^2}{\Omega_{ch}(v_p^2 - \alpha \mu_3 l_z^2 K_2)} \frac{\partial \phi^{(1)}}{\partial \xi}, \quad (33)$$

$$u_{ly}^{(1)} = \frac{l_x v_p^2}{\Omega_{ch}(v_p^2 - \alpha \mu_3 l_z^2 K_2)} \frac{\partial \phi^{(1)}}{\partial \xi}. \quad (34)$$

Now, by taking the next higher-order terms, the equation of continuity, momentum equation, and Poisson's equation can be

written as

$$\begin{aligned} \frac{\partial n_h^{(1)}}{\partial \tau} - v_p \frac{\partial n_h^{(2)}}{\partial \xi} + l_x \frac{\partial u_{hx}^{(1)}}{\partial \xi} + l_y \frac{\partial u_{hy}^{(1)}}{\partial \xi} + l_z \frac{\partial u_{hz}^{(2)}}{\partial \xi} \\ + l_z \frac{\partial}{\partial \xi} (n_h^{(1)} u_{hz}^{(1)}) = 0, \end{aligned} \quad (35)$$

$$\begin{aligned} \frac{\partial u_{hz}^{(1)}}{\partial \tau} - v_p \frac{\partial u_{hz}^{(2)}}{\partial \xi} + l_z u_{hz}^{(1)} \frac{\partial u_{hz}^{(1)}}{\partial \xi} + \mu_1 l_z \frac{\partial \phi^{(2)}}{\partial \xi} \\ + \alpha \mu_1 l_z K_1 \left[\frac{\partial n_h^{(2)}}{\partial \xi} + \frac{(\alpha - 2)}{2} \frac{\partial n_h^{(1)^2}}{\partial \xi} \right] - \eta \frac{\partial^2 u_{hz}^{(1)}}{\partial \xi^2} = 0, \end{aligned} \quad (36)$$

$$\begin{aligned} \frac{\partial n_l^{(1)}}{\partial \tau} - v_p \frac{\partial n_l^{(2)}}{\partial \xi} + l_x \frac{\partial u_{lx}^{(1)}}{\partial \xi} + l_y \frac{\partial u_{ly}^{(1)}}{\partial \xi} + l_z \frac{\partial u_{lz}^{(2)}}{\partial \xi} \\ + l_z \frac{\partial}{\partial \xi} (n_l^{(1)} u_{lz}^{(1)}) = 0, \end{aligned} \quad (37)$$

$$\begin{aligned} \frac{\partial u_{lz}^{(1)}}{\partial \tau} - v_p \frac{\partial u_{lz}^{(2)}}{\partial \xi} + l_z u_{lz}^{(1)} \frac{\partial u_{lz}^{(1)}}{\partial \xi} + \mu_2 l_z \frac{\partial \phi^{(2)}}{\partial \xi} \\ + \alpha \mu_3 l_z K_2 \left[\frac{\partial n_l^{(2)}}{\partial \xi} + \frac{(\alpha - 2)}{2} \frac{\partial n_l^{(1)^2}}{\partial \xi} \right] - \eta \frac{\partial^2 u_{lz}^{(1)}}{\partial \xi^2} = 0, \end{aligned} \quad (38)$$

$$\mu_5 n_l^{(2)} + n_h^{(2)} = \beta_1 \phi^{(2)} + \beta_2 \phi^{(1)^2}. \quad (39)$$

Finally, the next higher-order terms of Eqs. (8)-(11) and (15), with the help of Eqs. (25)-(39), can provide the Burgers' equation

$$\frac{\partial \Phi}{\partial \tau} + A \Phi \frac{\partial \Phi}{\partial \xi} = B \frac{\partial^2 \Phi}{\partial \xi^2}, \quad (40)$$

where $\Phi = \phi^{(1)}$ for simplicity. In Eq. (32), the nonlinear coefficient A and dissipative coefficient B are given

$$A = A'(A'' - 2\beta_2), \quad (41)$$

$$B = \frac{\eta}{2}, \quad (42)$$

where

$$\begin{aligned} A' &= \frac{v_p^4 + \alpha^2 \mu_1 \mu_3 l_z^4 K_1 K_2 - \alpha l_z^2 v_p^2 (\mu_1 K_1 + \mu_3 K_2)}{2 v_p l_z^2 [(v_p^2 - \alpha \mu_3 l_z^2 K_2)^2 + \mu_2 \mu_5 (v_p^2 - \alpha \mu_1 l_z^2 K_1)^2]}, \\ A'' &= \frac{l_z^4 \{ [3v_p^2 + \mu_1 l_z^2 K_1 \alpha (\alpha - 2)] + \mu_2^2 \mu_5 [3v_p^2 + \mu_3 l_z^2 K_2 \alpha (\alpha - 2)] \}}{\{(v_p^2 - \alpha \mu_1 l_z^2 K_1)(v_p^2 - \alpha \mu_3 l_z^2 K_2)\}^3}, \end{aligned}$$

Now, we look for stationary shock wave solution of this Burgers' equation by considering $\zeta = \xi - U_0 \tau$ (where ζ is a new space variable and U_0 is the speed of the ion fluid). These allow us to write the stationary shock wave solution as

$$\Phi = \Phi_0 \left[1 - \tanh \left(\frac{\zeta}{\Delta} \right) \right], \quad (43)$$

where the amplitude Φ_0 and width Δ are, respectively, given by

$$\Phi_0 = \frac{U_0}{A}, \quad \text{and} \quad \Delta = \frac{2B}{U_0}. \quad (44)$$

It is clear from Eqs. (43) and (44) that the IASHWs exist, which are formed due to the balance between nonlinearity and dissipation, because $B > 0$ and the IASHWs with $\Phi > 0$ ($\Phi < 0$) exist if $A > 0$ ($A < 0$) because $U_0 > 0$.




Figure 1: Plot of the nonlinear coefficient A vs μ_5 when $\alpha = 5/3$, $\gamma_e = 4/3$, $\gamma_p = 4/3$, $\delta = 20^\circ$, $\eta = 0.3$, $Z_h = 37$, $Z_l = 6$, $n_{h0} = 7 \times 10^{29} \text{cm}^{-3}$, $n_{l0} = 5 \times 10^{30} \text{cm}^{-3}$, $n_{p0} = 10^{31} \text{cm}^{-3}$, and $v_p = v_{p+}$.




Figure 3: Plot of Φ vs ζ for different values of η when $\alpha = 5/3$, $\gamma_e = 4/3$, $\gamma_p = 4/3$, $\delta = 20^\circ$, $Z_h = 37$, $Z_l = 6$, $n_{h0} = 7 \times 10^{29} \text{cm}^{-3}$, $n_{l0} = 5 \times 10^{30} \text{cm}^{-3}$, $n_{p0} = 10^{31} \text{cm}^{-3}$, $U_0 = 0.05$, and $v_p = v_{p+}$.




Figure 2: Plot of Φ vs ζ for different values of δ when $\alpha = 5/3$, $\gamma_e = 4/3$, $\gamma_p = 4/3$, $\eta = 0.3$, $Z_h = 37$, $Z_l = 6$, $n_{h0} = 7 \times 10^{29} \text{cm}^{-3}$, $n_{l0} = 5 \times 10^{30} \text{cm}^{-3}$, $n_{p0} = 10^{31} \text{cm}^{-3}$, $U_0 = 0.05$, and $v_p = v_{p+}$.




Figure 4: Plot of Φ vs ζ for different values of Z_l when $\alpha = 5/3$, $\gamma_e = 4/3$, $\gamma_p = 4/3$, $\delta = 20^\circ$, $\eta = 0.3$, $Z_h = 37$, $n_{h0} = 7 \times 10^{29} \text{cm}^{-3}$, $n_{l0} = 5 \times 10^{30} \text{cm}^{-3}$, $n_{p0} = 10^{31} \text{cm}^{-3}$, $U_0 = 0.05$, and $v_p = v_{p+}$.

4. Results and Discussion

Our investigation is reasonably valid for degenerate cold plasma systems (viz., white dwarfs and neutron stars) in which light (viz., ^1_1H [1, 2, 3, 4], ^4_2He [1, 5, 6, 7, 8], $^{12}_6\text{C}$ [7, 8], and $^{16}_8\text{O}$ [7, 8, 9], etc.) and heavy (viz., $^{56}_{26}\text{Fe}$ [10, 11], $^{85}_{37}\text{Rb}$ [11, 12], and $^{96}_{42}\text{Mo}$ [11, 12], etc.) ions can exist. For our numerical analysis, we have considered the range of the plasma parameters as $Z_l = 6 \sim 13$, $Z_h = 35 \sim 65$, $n_{l0} = 3 \times 10^{30} \text{cm}^{-3} \sim 7 \times 10^{30} \text{cm}^{-3}$, $n_{h0} = 4 \times 10^{29} \text{cm}^{-3} \sim 10 \times 10^{29} \text{cm}^{-3}$, and $n_{p0} = 1 \times 10^{31} \text{cm}^{-3} \sim 10 \times 10^{31} \text{cm}^{-3}$.

It is obvious from Eq. (40) that the wave potential becomes infinite when nonlinear coefficient is equal to zero (i.e., $A = 0$), and in that case, the validity of reductive perturbation method breaks down. The positive (i.e., $\Phi > 0$) and negative (i.e., $\Phi < 0$) electrostatic potentials can exist corresponding to the values of $A > 0$ and $A < 0$, respectively. It is clear from Fig. 1 that our plasma model supports only positive potential shock structure (i.e., $\Phi > 0$) associated with $A > 0$ under consideration of non-relativistic light and heavy ions (i.e., $\alpha = 5/3$), and ultra-relativistic degenerate electrons and positrons (i.e., $\gamma_e = \gamma_p = 4/3$).

The oblique angle δ indicates the angle between the direc-

tion of the propagation of IASHW and the direction of existing external magnetic field which is parallel with z -axes. The variation of electrostatic shock potential structure (i.e., $\Phi > 0$) associated with $A > 0$ with δ under consideration of non-relativistic light and heavy ions (i.e., $\alpha = 5/3$) and ultra-relativistic degenerate electrons and positrons (i.e., $\gamma_e = \gamma_p = 4/3$) can be observed in Fig. 2. It is obvious from this figure that the electrostatic shock potential (i.e., $\Phi > 0$) associated with $A > 0$ increases with the increase of δ . Physically, the interaction between the electrostatic shock potential associated with $A > 0$ and the external magnetic field is clearly increased with the increase in δ .

Figure 3 describes the effects of the kinematic viscosity of the positively charged non-relativistic light and heavy ions (via η) on the electrostatic shock potential structure (i.e., $\Phi > 0$) associated with $A > 0$. We have noticed that the amplitude of the electrostatic shock profile (i.e., $\Phi > 0$) associated with $A > 0$ is independent to the variation of the ion kinematic viscosity but the steepness of the electrostatic shock profile (i.e., $\Phi > 0$) associated with $A > 0$ is rigorously dependent to the variation of ion kinematic viscosity. The steepness of the electrostatic shock potential structure (i.e., $\Phi > 0$) associated with $A > 0$




Figure 5: Plot of Φ vs ζ for different values of Z_h when $\alpha = 5/3$; $\gamma_e = 4/3$, $\gamma_p = 4/3$, $\delta = 20^\circ$, $\eta = 0.3$, $Z_l = 6$, $n_{h0} = 7 \times 10^{29} \text{cm}^{-3}$, $n_{l0} = 5 \times 10^{30} \text{cm}^{-3}$, $n_{p0} = 10^{31} \text{cm}^{-3}$, $U_0 = 0.05$, and $v_p = v_{p+}$.




Figure 7: Plot of Φ vs ζ for different values of n_{h0} when $\alpha = 5/3$; $\gamma_e = 4/3$, $\gamma_p = 4/3$, $\delta = 20^\circ$, $\eta = 0.3$, $Z_h = 37$, $Z_l = 6$, $n_{l0} = 5 \times 10^{30} \text{cm}^{-3}$, $n_{p0} = 10^{31} \text{cm}^{-3}$, $U_0 = 0.05$, and $v_p = v_{p+}$.




Figure 6: Plot of Φ vs ζ for different values n_{l0} when $\alpha = 5/3$; $\gamma_e = 4/3$, $\gamma_p = 4/3$, $\delta = 20^\circ$, $\eta = 0.3$, $Z_h = 37$, $Z_l = 6$, $n_{h0} = 7 \times 10^{29} \text{cm}^{-3}$, $n_{p0} = 10^{31} \text{cm}^{-3}$, $U_0 = 0.05$, and $v_p = v_{p+}$.




Figure 8: Plot of Φ vs ζ for different values of n_{p0} when $\alpha = 5/3$; $\gamma_e = 4/3$, $\gamma_p = 4/3$, $\delta = 20^\circ$, $\eta = 0.3$, $Z_h = 37$, $Z_l = 6$, $n_{h0} = 7 \times 10^{29} \text{cm}^{-3}$, $n_{l0} = 5 \times 10^{30} \text{cm}^{-3}$, $U_0 = 0.05$, and $v_p = v_{p+}$.

decreases with η , and this result agrees with the result of Refs. [31, 32].

The amplitude of the positive electrostatic shock structure (i.e., $\Phi > 0$) associated with $A > 0$ is so much sensitive to the change of the charge state of non-relativistic light and heavy ions. Figures 4 and 5 show that the amplitude of the positive shock structure (i.e., $\Phi > 0$) associated with $A > 0$ increases with the charge state of non-relativistic light and heavy ions. Similarly, the increasing number of non-relativistic light and heavy ions enhances the positive electrostatic shock structure (i.e., $\Phi > 0$) associated with $A > 0$. It can easily demonstrate from Figs. 6 and 7 that as we increase n_{l0} and n_{h0} , the amplitude of the positive electrostatic shock structure (i.e., $\Phi > 0$) associated with $A > 0$ increases. Physically, the charge state and number density of non-relativistic light and heavy ions can control the dynamics of the DQP system in the similar way.

We have studied the characteristics of IASHWs for different values of positron number density (via n_{p0}) under consideration of non-relativistic light and heavy ions (i.e., $\alpha = 5/3$) and ultra-relativistic degenerate electrons and positrons (i.e., $\gamma_e = \gamma_p = 4/3$) in Fig. 8. It can be highlighted from this figure that the amplitude of electrostatic shock structure (i.e., $\Phi > 0$)

associated with $A > 0$ decreases with positron number density, and this finding is analogous to the result of Refs. [21, 22].

5. Conclusion

We have studied the basic characteristics of IASHWs in an extremely dense DQP containing non-relativistic light and heavy ions, and inertialess ultra-relativistic degenerate electrons and positrons in the presence of external magnetic field. The RPM [41] has been utilized to derive Burgers' equation. The results that have been found from our present investigation can be summarized as follows:

- Our plasma model supports only positive potential shock structure (i.e., $\Phi > 0$) associated with $A > 0$ under consideration of non-relativistic light and heavy ions (i.e., $\alpha = 5/3$), and ultra-relativistic degenerate electrons and positrons (i.e., $\gamma_e = \gamma_p = 4/3$).
- The electrostatic shock potential (i.e., $\Phi > 0$) associated with $A > 0$ increases with the increase in δ .
- The steepness of the positive potential shock structure (i.e., $\Phi > 0$) associated with $A > 0$ decreases with η .

- The amplitude of electrostatic shock structure (i.e., $\Phi > 0$) associated with $A > 0$ is to be found to increase with the charge state and number density of non-relativistic light and heavy ions.
- The increasing positron number density decreases the height of the positive shock profile.

It may be noted that the self-gravitational effect of the plasma species is important to be considered in our governing equations but beyond the scope of our present work. Overall, the outcomes from our present investigation will be helpful to understand the IASHWs in white dwarfs and neutron stars.

References

- [1] S. Chandrasekhar, *Astrophys. J.* **74**, 81 (1931).
- [2] S. Chandrasekhar, *The Observatory* **57**, 373 (1934).
- [3] R.S. Fletcher, *et al.*, *Phys. Rev. Lett.* **96**, 105003 (2006).
- [4] T.C. Killian, *Nature (London)* **441**, 297 (2006).
- [5] H.M. Van Horn, *Science* **252**, 384 (1991).
- [6] R.H. Fowler, *J. Astrophys. Astron.* **15**, 105 (1994).
- [7] D. Koester and G. Chanmugam, *Rep. Prog. Phys.* **53**, 837 (1990).
- [8] D. Koester, *Astron. Astrophys. Rev.* **11**, 33 (2002).
- [9] S. Chandrasekhar, *Philos. Mag.* **11**, 592 (1931).
- [10] A. Vanderburg, *et al.*, *Nature* **526**, 546 (2015).
- [11] N. Chamel and P. Haensel, *Living Rev. Relativ.* **11**, 10 (2008).
- [12] A. Witze, *Nature* **510**, 196 (1990).
- [13] S. Chandrasekhar and R.F. Tooper, *Astrophys. J.* **139**, 1396 (1964).
- [14] P.K. Shukla, *et al.*, *Phys. Rev. E* **84**, 026405 (2011).
- [15] W.F. El-Taibany, *et al.*, *Adv. space res.* **50**, 101 (2012).
- [16] S. Sultana and R. Schlickeiser, *Astrophys. Space Sci.* **363**, 103 (2018).
- [17] B. Zhang and J. Gil, *Astrophys. J.* **631**, 143 (2005).
- [18] P.A. Sturrock, *Astrophys. J.* **164** (1971) 529.
- [19] A.K. Harding and A. G. Muslimov, *Astrophys. J.* **508**, 328 (1998).
- [20] A.K. Harding and D. Lai, *Rep. Prog. Phys.* **69**, 2631 (2006).
- [21] T.S. Gill, *et al.*, *J. Phys.* **208**, 012040 (2010).
- [22] Ata-Ur-Rahman, *et al.*, *J. Plasma Phys.* **79**, 817 (2013).
- [23] M.R. Hossen, *et al.*, *Plasma Phys. Rep.* **43**, 1189 (2017).
- [24] A.A. Mamun and P.K. Shukla, *Phys. Plasmas* **17**, 104504 (2010).
- [25] P.M.S. Blackett, *Nature* **159**, 658 (1947).
- [26] J. Liebert, *et al.*, *Astrophys. J.* **214**, 457 (1977).
- [27] F. Euchner, *et al.*, *Astron. Astrophys.* **390**, 633 (2002).
- [28] W.F. El-Taibany, *et al.*, *Adv. Space Res.* **50**, 101 (2012).
- [29] M.I. Shaikat, *Phys. Plasmas* **24**, 102301 (2017).
- [30] J.M. Burgers, *Elsevier* **1**, 171 (1948).
- [31] M.G. Hafez, *et al.*, *Plasma Phys. Rep.* **43**, 499 (2017).
- [32] H.G. Abdelwahed, *et al.*, *J. Exp. Theor. Phys.* **122**, 1111 (2016).
- [33] N.S. Saini, *et al.*, *Wave Random Complex.* DOI: 10.1080/17455030.2020.1798561.
- [34] M. M. Haider, *Z. Naturforsch. A* **71**, 1131 (2016).
- [35] A. Atteya, *et al.*, *Eur. Phys. J. Plus* **132**, 1 (2017).
- [36] M.H. Rahman, *et al.*, *Phys. Plasmas* **25**, 102118 (2018); N.A. Chowdhury, *et al.*, *Phys. plasmas* **24**, 113701 (2017); M.H. Rahman, *et al.*, *Chin. J. Phys.* **56**, 2061 (2018); N.A. Chowdhury, *et al.*, *Plasma Phys. Rep.* **45**, 459 (2019); S. Jahan, *et al.*, *Plasma Phys. Rep.* **46**, 90 (2020).
- [37] S.K. Paul, *et al.*, *Pramana J. Phys.* **94**, 58 (2020); N.A. Chowdhury, *et al.*, *Chaos* **27**, 093105 (2017); N.A. Chowdhury, *et al.*, *Contrib. Plasma Phys.* **58**, 870 (2018); M. Hassan, *et al.*, *Commun. Theor. Phys.* **71**, 1017 (2019); D.M.S. Zaman, *et al.*, *High Temp.* **58**, 789 (2020).
- [38] A. A. Mamun, *Phys. Scr.* **59**, 454 (1999).
- [39] M. M. Hossen, *et al.*, *High Energy Density Phys.* **24**, 9 (2017).
- [40] H. Washimi, T. Taniuti, *Phys. Rev. Lett.* **17**, 996 (1966).
- [41] S. Jahan, *et al.*, *Commun. Theor. Phys.* **71**, 327 (2019); N. Ahmed, *et al.*, *Chaos* **28**, 123107 (2018); N.A. Chowdhury, *et al.*, *Vacuum* **147**, 31 (2018); R.K. Shikha, *et al.*, *Eur. Phys. J. D* **73**, 177 (2019); S. Banik, *et al.*, *Eur. Phys. J. D* **75**, 43 (2021).