Tuning symplectic integrators is easy and worthwhile

Robert I McLachlan*

School of Fundamental Sciences, Massey University, New Zealand

(Dated: April 22, 2021)

Many applications in computational physics that use numerical integrators based on splitting and composition can benefit from the development of optimized algorithms and from choosing the best ordering of terms. The cost in programming and execution time is minimal, while the performance improvements can be large.

PACS numbers: Valid PACS appear here

I. INTRODUCTION

Symplectic numerical integration by splitting the Hamiltonian and composing the flows of the associated vector fields has become an extremely widely used technique in computational science, especially computational physics and chemistry. A splitting into two parts, $H = H_1 + H_2$, together with the 3-term composition

$$e^{\frac{1}{2}hX_1}e^{hX_2}e^{\frac{1}{2}hX_1}$$

is the most common and is often all that is needed. It is variously called the leapfrog, Störmer–Verlet, or Strang splitting method. (Here X_i is the Hamiltonian vector field associated with Hamiltonian H_i , e^{hX} is the time-*h* flow of the vector field X, and h is the time step.) For example, it is the basic building block of the Hamiltonian Monte Carlo method.

Some Hamiltonians can only be written as the sum of more than two explicitly integrable terms, say $H = \sum_{i=1}^{n} H_i$. In addition, some applications need order higher than two to achieve the required accuracy for a given amount of computational effort. Methods of all orders exist, but are progressively more expensive. Optimized methods have been develop that can significantly reduce discretization errors at fixed cost [1].

However, an informal survey of the current literature suggests that unoptimized composition methods of order 4, 6, and 8 [2] are in common use in cosmology, celestial mechanics, quantum mechanics, quantum statistical mechanics, solid state physics, kinetic theory, plasma physics, molecular dynamics, optics, neural networks, and fluid mechanics [3–27]. Computations in these fields could benefit from experience gained in numerical analysis to reduce errors and error growth at little cost either in programming or execution time.

In this note we review four such techniques.

II. OPTIMIZED METHODS FOR MULTI-TERM SPLITTINGS

Let the vector field $X = \sum_{i=1}^{n}$ be split into *n* parts, each explicitly integrable and let

$$\chi_h = \mathrm{e}^{hX_n} \circ \cdots \circ \mathrm{e}^{hX_1}$$

be a first-order integrator for X. We define the adjoint χ_h^* of χ_h by $\chi_h^* = (\chi^{-1})_{-h}$; that is,

$$\chi_h^* = \mathrm{e}^{hX_1} \circ \cdots \circ \mathrm{e}^{hX_n}.$$

Let

$$\psi_h(\alpha) = \chi_{\alpha_s h} \circ \chi^*_{\alpha_{(s-1)}h} \circ \cdots \circ \chi_{\alpha_2 h} \circ \chi^*_{\alpha_1 h}.$$
(1)

The second-order leapfrog method is $S_{2,h} = \psi_h(\frac{1}{2}, \frac{1}{2})$. Note that when n = 2, this reduces to a composition that alternates steps of X_1 and X_2 . Methods of this type designed for n = 2 and for arbitrary n have the same order [28], although the optimal coefficients may not be the same. Nevertheless, coefficients optimised for the method (1) can reduce the error significantly.

In particular, the minimal-s methods formed by recursively increasing the order from k to k + 2 by $S_{k,ch} \circ S_{k,(1-2c)h} \circ S_{k,ch}$ with $c = 1/(2 - 2^{1/(2k+1)})$ have very large error constants and poor stability and should be avoided. The 3-stage method with k = 2, called S₃4, will be used as a reference method here. (The notation indicates that it has order 4 and uses work equivalent to 3 leapfrog steps.) Its large substeps, 1.35h, -1.70h, and 1.35h, contribute to its large error constants and poor stability.

The optimized method that we will use for the purpose of illustration here is called BM₆4 and was introduced by Blanes and Moan [29]. It has s = 12 and $\alpha_1 = \alpha_{12} =$ 0.0792036964311957, $\alpha_3 = \alpha_{10} = 0.2228614958676077$, $\alpha_5 = \alpha_8 = 0.3246481886897062$, $\alpha_7 = \alpha_6 =$ 0.1096884778767498, $\alpha_9 = \alpha_4 = -0.3667132690474257$, and $\alpha_{11} = \alpha_2 = 0.1303114101821663$. BM₆4 has errors are around 500 times smaller than those of S₃4 and also allows the use of larger time steps.

^{*}Electronic address: r.mclachlan@massey.ac.nz

III. ORDER MATTERS

The ordering of the *n* terms affects the errors. It can also affect the cost slightly, as there is an opportunity to collect two steps of X_1 and of X_n into single steps. There is no easy way to anticipate the best ordering. Fasso [30] studied this question in detail for the splitting of the rigid body Hamiltonian into three terms, finding that the ordering affected the error by a factor of 10, 100, or more, and that the best ordering depended in a non-obvious way on the moments of inertia of the body.

If n is not too large, one approach is to try all permutations of the n terms with a typical initial condition and choose the ordering with the smallest local truncation error.

In Table I we give the results of numerical experiments testing the influence of the term ordering on the error of the optimized 4th order integrator BM₆4. Linear systems in \mathbb{R}^{16} with random coefficients are integrated for time 1 with time-step 0.1. The difference between the best and worst orders is significant and increases with n. The principal error in 4th order symmetric integrators contains $\mathcal{O}(n^3)$ terms, each a triple commutator of the terms of the ODE. Some of these terms might cancel each other; the experiment suggests that the effect of using the best ordering scales like n^c with c in the region 1.5 to 2.5. The effect in these unstructured systems is less that that observed by Fasso [30] for the (more highly structured) rigid body.

IV. COMPENSATED SUMMATION REDUCES ROUND-OFF ERROR

Round-off error is not often a dominant source of error, but in long integrations it does accumulate and can sometimes dominate the truncation error. It can be drastically reduced at minuscule cost (both in programming and execution) by the technique of compensated summation [31]. In each update $x_{n+1} = x_n + hy_n$, the increment hy_n is computed to full precision, but x_n is stored to higher precision by storing a pair of full precision numbers. Compensated summation reduces roundoff errors by a factor $||hy_n||/||x_n||$.

Computing hy_n to full precision is important and may need attention, especially if the flow e^{hX} involves special functions. One non-trivial example that we will use below involves $H = \frac{1}{2}q^np^2$, an instance of Channell's observation that any monomial Hamiltonian is explicitly integrable in terms of elementary functions [32]. First, the solution of Hamilton's equations should be presented algebraically with the update separated out:

$$\begin{aligned} \dot{q} &= q^{n}p, \qquad q(0) = q_{0} \\ \dot{p} &= -\frac{n}{2}q^{n-1}p, \quad p(0) = p_{0} \\ \Rightarrow q(t) &= q_{0} \left(1 + \delta\right)^{\frac{2}{2-n}} \\ &= q_{0} + q_{0} \left((1 + \delta)^{\frac{2}{2-n}} - 1\right) \quad (n \neq 2) \\ p(t) &= p_{0} \left(1 + \delta\right)^{\frac{n}{n-2}} \\ &= p_{0} + p_{0} \left((1 + \delta)^{\frac{n}{n-2}} - 1\right), \\ \delta &= t \left(1 - \frac{n}{2}\right) q_{0}^{n-1} p_{0} \end{aligned}$$

Second, the update should be computed to full precision. In this case this can be achieved using the functions $\operatorname{expm1}(z) = e^{z} - 1$ and $\log 1p(z) = \log(1+z)$ provided in most mathematics libraries:

$$q(t) = q_0 + q_0 \operatorname{expm1}\left(\frac{2}{2-n}\log p(\delta)\right)$$

$$p(t) = p_0 + p_0 \operatorname{expm1}\left(\frac{n}{n-2}\log p(\delta)\right)$$
(2)

Similarly for n = 2, we have

$$q(t) = e^{p_0 q_0 t} q_0 = q_0 + q_0 \text{expm1}(p_0 q_0 t)$$

$$p(t) = e^{-p_0 q_0 t} p_0 = q_0 + p_0 \text{expm1}(-p_0 q_0 t)$$
(3)

These formulations are faster and more accurate than the direct implementation using powers.

Secondly, "Brouwer's Law" states that roundoff errors should accumulate like a random walk, leading to squareroot growth in time [33]. Linear error growth of roundoff errors, which is often observed in momenta that would be conserved exactly in exact arithmetic, is sign of a failure of Brouwer's Law and an indication of systematic bias in the floating point arithmetic. Techniques for detecting and correcting this bias are found in the literature [34– 40].

V. METHODS WITH PROCESSING

Methods of the form $\rho \circ \psi \circ \rho^{-1}$ are called "processed" or "corrected" methods [41]. The basic method ψ is called the kernel, and ρ the processor. Processed methods were put to spectacular use by Wisdom [42], in which the stored values of a very long solar system simulation (based on leapfrog with a 2-term splitting) were processed years after the original calculation to dramatically reduce errors. (Or to put it another way, to reveal the true, underlying errors.) Methods with processing have been optimized for many different cases of splitting: the general case (considered in this note); near-integrable systems (as in the Wisdom–Holman example); and systems based on splitting into kinetic and potential parts. The processor can either be a method of the same class

	antisy	mmetri	ic case	general case		
n	$\min R$	$\operatorname{med} R$	$\max R$	$\min R$	$\operatorname{med} R$	$\max R$
3	1.16	1.70	4.05	1.20	1.90	3.95
4	1.66	2.57	5.32	1.98	3.40	7.68
5	2.23	3.46	7.64	2.80	5.41	11.39
6	3.46	5.00	9.71	4.38	9.29	18.55
7	4.80	6.36	15.62	6.41	13.35	24.22

TABLE I: Linear systems $\dot{X} = (\sum_{i=1}^{n} A_i)X$, $0 \leq t \leq 1$, $X \in \mathbb{R}^{16 \times 16}$, X(0) = I, are integrated using an optimized 4th order splitting method. In each trial, all n! orderings of A_1, \ldots, A_n are tested and the ratio R between the greatest and the least global error of all orderings at t = 1 is computed. The minimum, median, and maximum ratio R over 50 trials is reported. On the left ('antisymmetric case'), each A_i is antisymmetric with independent normally distributed entries, modelling periodic systems with global errors growing linearly in time; on the right ('general case'), each A_i has independent normally distributed entries, modelling systems with sensitive dependence on initial conditions.

as the kernel, or calculated more cheaply by other methods. Either way, if output is taken every m steps, then only $\rho \circ \psi^m \circ \rho^{-1}$ is calculated. Observables that are invariant under conjugation by the diffeomorphism ρ (such as invariant phase space structures, Lyapunov exponents, periods of periodic orbits) can be computed without applying ρ at all. The error can now be reduced relative to the non-corrected case because only those error terms in the kernel that cannot be removed by a corrected need to be minimized.

In the examples of linear systems in Table I, it reduce errors by a factor of about 1.4. Greater improvements have been reported at higher order, for nearly integrable systems, and for systems with Hamiltonians of the form kinetic plus potential [41].

VI. EXAMPLE: PARTICLES NEAR BLACK HOLES

We consider two examples of explicit symplectic integrators based on splitting and composition. Both have four-dimensional (reduced) space space. The first tracks charged particles near a Schwarzschild black hole [24] and has a Hamiltonian separable into 4 terms; the second tracks charged particles near a Reissner–Nordström antide Sitter black hole [25] and has a Hamiltonian separable into 6 terms. In both cases the Hamiltonians are of the form $V(q) + \sum_{i=1}^{n-1} T_i(q, p)$ where each kinetic energy term T_i is a monomial in q and p and hence explicitly integrable in terms of elementary functions [32] as in Eqs (2,3).

The optimized method can reduce energy errors by a factor of up to 10^4 . In all cases in the best ordering of terms the potential term was applied first, but the ordering of the remaining kinetic energy terms had to be determined experimentally.



FIG. 1: Energy error in integration of a particle near a Schwarzschild black hole over proper time $\tau = 10^8$ with time step $\Delta \tau = 1$ by two 4th order symplectic integrators, S₃4 (top) and BM₆4 with compensated summation (bottom), the latter showing smaller errors and no energy drift.

geometry	method	least	greatest	ratio
Sahwarzashild	BM_64	4.0×10^{-12}	1.2×10^{-10}	30
Schwarzschlid	S_34	3.9×10^{-9}	3.0×10^{-8}	8
P. N. odS	$BM_{6}4$	1.3×10^{-11}	3.1×10^{-10}	24
n-n aus	S_34	2.9×10^{-9}	6.8×10^{-8}	23

TABLE II: Energy errors in particle tracking near black holes in two cases by each of two different methods. The Schwarzschild case integrates a quasiperiodic orbit ('Orbit 1' from [24, Fig. 1] with 4-part splitting the R–N adS case integrates a chaotic orbit [25, Fig. 2(c)] with 6-part splitting. The least and the greatest energy errors over the different orderings of the terms, and their ratio, are given. The reference energy level is -0.5 and the energy error is measured in the sup ($\|\cdot\|_{\infty}$) norm; the time step is 1 in all cases.

In addition, the use of compensated summation drastically reduces the roundoff error to the point that it is invisible in Figure 1. On closer inspection, it is not accumulating linearly, and the roundoff error after 10^8 steps is about 2×10^{-16} . In constrast, the method S₃4 without compensated summation shows a linear drift in energy of about 10^{-9} over the same time interval.

In the Schwarzschild example, the optimized 4th order processed method P_{64} of [41] has similar errors to BM_{64} .

VII. DISCUSSION

While second-order Strang splitting is sufficient for many applications, when higher accuracy is required, the optimized higher order methods are preferred. We know of no case in which the method S_34 is preferred over BM₆4; the latter generally has errors orders of magnitude smaller and can be used with larger time steps. Compensated summation, and checking that roundoff errors are unbiased, costs almost nothing in programming or execution time. Despite this, the method S_34 without compensated summation remains widely used.

Possibly one reason for this situation is that these algorithms are so simple and flexible, and are used in such diverse applications in computational science, that they are often implemented by hand afresh for each new application. In contrast, numerical methods that are more complicated, such as the finite element method on un-

- S. Blanes, F. Casas, and A. Murua, arXiv preprint arXiv:0812.0377 (2008).
- [2] H. Yoshida, Physics letters A **150**, 262 (1990).
- [3] G. Biondini and J. Oregero, Studies in Applied Mathematics 145, 325 (2020).
- [4] A. Bravetti, M. Seri, M. Vermeeren, and F. Zadra, Celestial Mechanics and Dynamical Astronomy 132, 1 (2020).
- [5] J. Budd, Y. van Gennip, and J. Latz, GAMM-Mitteilungen 44, e202100004 (2021).
- [6] J. Cai and H. Zhang, Applied Mathematics Letters 102, 106158 (2020).
- [7] S. Y. Cho, S. Boscarino, G. Russo, and S.-B. Yun, Journal of Computational Physics p. 110281 (2021).
- [8] C. Deng, X. Wu, and E. Liang, Monthly Notices of the Royal Astronomical Society 496, 2946 (2020).
- [9] P. K. Faehrmann, M. Steudtner, R. Kueng, M. Kieferova, and J. Eisert, arXiv preprint arXiv:2101.07808 (2021).
- [10] T. E. Faver, R. H. Goodman, and J. D. Wright, Zeitschrift f
 ür angewandte Mathematik und Physik 71, 1 (2020).
- [11] D. G. Figueroa, A. Florio, F. Torrenti, and W. Valkenburg, arXiv preprint arXiv:2006.15122 (2020).
- [12] C. Klein and N. Stoilov, Studies in Applied Mathematics 145, 36 (2020).
- [13] K. D. Lozanov and M. A. Amin, Journal of Cosmology and Astroparticle Physics **2020**, 058 (2020).
- [14] G. Mancini, S. Del Galdo, B. Chandramouli, M. Pagliai, and V. Barone, Journal of Chemical Theory and Computation 16, 5747 (2020).
- [15] V. Manwadkar, A. A. Trani, and N. W. Leigh, Monthly Notices of the Royal Astronomical Society 497, 3694 (2020).
- [16] H. Ohno, JOSA A **37**, 411 (2020).
- [17] K. Seki and S. Yunoki, PRX Quantum 2, 010333 (2021).
- [18] C. Sheng, J. Shen, T. Tang, L.-L. Wang, and H. Yuan, SIAM Journal on Numerical Analysis 58, 2435 (2020).
- [19] H. Song, L. Vogt-Maranto, R. Wiscons, A. J. Matzger, and M. E. Tuckerman, The Journal of Physical Chemistry Letters 11, 9751 (2020).
- [20] E. Stern, Y. Alexahin, A. Burov, and V. Shiltsev, Journal of Instrumentation 16, P03045 (2021).
- [21] Y. Tong, S. Xiong, X. He, G. Pan, and B. Zhu, Journal of Computational Physics p. 110325 (2021).

structured meshes, have passed into packages and opensource platforms that incorporate advances as they become available.

Acknowledgments

The author thanks Ying Wang and Sergio Blanes for discussions.

- [22] L. Wang, K. Nitadori, and J. Makino, Monthly Notices of the Royal Astronomical Society 493, 3398 (2020).
- [23] Z. Wang, W. Fu, Y. Zhang, and H. Zhao, Physical Review Letters **124**, 186401 (2020).
- [24] Y. Wang, W. Sun, F. Liu, and X. Wu, The Astrophysical Journal 907, 66 (2021).
- [25] Y. Wang, W. Sun, F. Liu, and X. Wu, arXiv preprint arXiv:2103.12272 (2021).
- [26] Y. Wu, H. Li, and J. Hou, Computational Materials Science 190, 110273 (2021).
- [27] J. Xiao and H. Qin, Plasma Science and Technology (2021).
- [28] R. I. McLachlan, SIAM Journal on Scientific Computing 16, 151 (1995).
- [29] S. Blanes and P. C. Moan, Journal of Computational and Applied Mathematics 142, 313 (2002).
- [30] F. Fassò, Journal of computational physics 189, 527 (2003).
- [31] N. J. Higham, Accuracy and stability of numerical algorithms (SIAM, 2002).
- [32] P. Channell, AT-6: ATN-86-6, Los Alamos National Laboratory (1986).
- [33] E. Hairer, R. I. McLachlan, and A. Razakarivony, BIT Numerical Mathematics 48, 231 (2008).
- [34] M. Antonana, J. Makazaga, and A. Murua, Numerical Algorithms 78, 63 (2018).
- [35] R. I. McLachlan, K. Modin, and O. Verdier, Physical Review E 89, 061301 (2014).
- [36] H. Rein and D. S. Spiegel, Monthly Notices of the Royal Astronomical Society 446, 1424 (2015).
- [37] M. Rodríguez and R. Barrio, Applied Numerical Mathematics 62, 1014 (2012).
- [38] J. Seyrich and G. Lukes-Gerakopoulos, Physical Review D 86, 124013 (2012).
- [39] L. Symes, R. McLachlan, and P. Blakie, Physical Review E 93, 053309 (2016).
- [40] G. Vilmart, Journal of computational physics 227, 7083 (2008).
- [41] S. Blanes, F. Casas, and A. Murua, SIAM Journal on Scientific Computing 27, 1817 (2006).
- [42] J. Wisdom, M. Holman, and J. Touma, Fields Institute Communications 10, 217 (1996).