

A NON-HYPERELLIPTIC CURVE WITH TORSION CERESA CYCLE MODULO ALGEBRAIC EQUIVALENCE

ARNAUD BEAUVILLE AND CHAD SCHOEN

ABSTRACT. We exhibit a non-hyperelliptic curve C of genus 3 such that the class of the Ceresa cycle $[C] - [C^-]$ in JC modulo algebraic equivalence is torsion.

1. INTRODUCTION

Let C be a complex curve of genus $g \geq 2$, and p a point of C . We embed C into its Jacobian J by the Abel-Jacobi map $x \mapsto [x] - [p]$; we denote by C^- the image of C under the involution $(-1)_J : a \mapsto -a$ of J . The *Ceresa class* is the class $\mathfrak{z}(C) := [C] - [C^-]$ in the group $A_1(J)$ of 1-cycles on J modulo algebraic equivalence (it is independent of the choice of p). Since $(-1)_J$ acts trivially on $H^p(J, \mathbb{Z})$ for p even, $\mathfrak{z}(C)$ belongs to the *Griffiths group* $G(J)$, the kernel of the cycle class map $A_1(J) \rightarrow H^{2g-2}(J, \mathbb{Z})$.

Ceresa classes have played a prominent role in the study of Griffiths groups, especially in the development of techniques for showing that a given element is non-zero [C, C-P, H]. In addition they played an important role in showing that $G(J)$ can have infinite rank [N]. As the conjectures of Bloch and Beilinson were developed and are studied $\mathfrak{z}(C)$ appears repeatedly [Bl, BST], [Z, §1.5], always as an element of infinite order.

When C is hyperelliptic, $\mathfrak{z}(C) = 0$; in fact $C - C^-$ is zero as a cycle when p is a Weierstrass point. In this note we will exhibit what we believe to be the first example of a non-hyperelliptic curve C with $\mathfrak{z}(C) = 0$ in $A_1(J) \otimes \mathbb{Q}$. The curve C has genus 3, and admits an automorphism σ of order 9, such that the quotient variety $J/\langle\sigma\rangle$ is uniruled. This implies that the Griffiths group of a resolution of $J/\langle\sigma\rangle$ is torsion; going back to J gives the result.

2. MAIN RESULT

Theorem. *Let $C \subset \mathbb{P}^2$ be the genus 3 curve defined by $X^4 + XZ^3 + Y^3Z = 0$. Then $\mathfrak{z}(C) = 0$ in $A_1(J) \otimes \mathbb{Q}$.*

Proof : Let ζ be a primitive 9-th root of unity. We consider the automorphism σ of C defined by $\sigma(X, Y, Z) = (X, \zeta^2 Y, \zeta^3 Z)$. We use the fixed point $p = (0, 0, 1)$ to embed C in its Jacobian J , so that the action of σ on J preserves C and C^- . We denote by V the quotient variety $J/\langle\sigma\rangle$, and by $\pi : J \rightarrow V$ the quotient map. Let $F \subset J$ be the subset of elements with nontrivial stabilizer; the singular locus $\text{Sing } V$ of V is $\pi(F)$. We put $J^\circ := J \setminus F$ and $V^\circ := V \setminus \text{Sing } V$.

Lemma 1. *Sing V is finite; the points $\pi(x)$ for $x \in \text{Ker}(1_J - \sigma)$ are non-canonical singularities.*

Proof : The space $T_0(J)$ is canonically identified with $H^0(C, K_C)^*$. The elements of $H^0(C, K_C)$ are of the form $L \frac{XdZ - ZdX}{Y^2Z}$, with $L \in H^0(\mathbb{P}^2, \mathcal{O}_{\mathbb{P}^2}(1))$ [B-K, §9.3, Corollary of Theorem 1]. It follows that the eigenvalues of σ on $H^0(C, K_C)$ are $\zeta^5, \zeta^7, \zeta^8$, and those on $T_0(J) = H^0(C, K_C)^*$ are ζ, ζ^2, ζ^4 .

Thanks to E. Colombo and B. van Geemen for their crucial input. The second named author thanks S. Katz and M. Reid for helpful discussions.

Therefore $\text{Ker}(1_J - \sigma^d)$ is finite for $0 < d < 9$, so F is finite. Since $1 + 2 + 4 < 9$, Reid's criterion [R, Theorem 3.1] implies that the singular points $\pi(x)$ for $x \in \text{Ker}(1_J - \sigma)$ are not canonical. ■

Lemma 2. *The variety V is uniruled.*

Proof : Let $\rho : \tilde{V} \rightarrow V$ be a resolution of singularities; it suffices to prove that \tilde{V} has Kodaira dimension $-\infty$ [M]. Suppose this is not the case: there exist an integer $r \geq 1$ and a nonzero section $\tilde{\omega}$ of $K_{\tilde{V}}^r$. By restriction to $\rho^{-1}(V^\circ) \cong V^\circ$, we get a section ω of $K_{V^\circ}^r$, whose pull back under π is a nonzero section of $K_{V^\circ}^r$; therefore ω is a generator of $K_{V^\circ}^r$, hence extends to a generator of the reflexive sheaf $K_V^{[r]}$ (with the notation of [R]). By construction this generator remains regular on \tilde{V} , which means that the singularities of V are canonical [R, Proposition 1.2], contradicting Lemma 1. ■

Lemma 3. *Let X be a uniruled smooth projective threefold. The Griffiths group $G(X)$ is torsion.*

Proof : There exists a smooth projective surface S and a dominant rational map $S \times \mathbb{P}^1 \dashrightarrow X$. After blowing up some points and some smooth curves in $S \times \mathbb{P}^1$, we get a smooth projective threefold W birational to $S \times \mathbb{P}^1$, and a generically finite morphism $f : W \rightarrow X$. Since the Griffiths group is a stably birational invariant (see [V, Proposition 2.30]), we have $G(W) = G(S) = 0$. For $z \in G(X)$, we have $(\deg f)z = f_* f^*(z) = 0$, hence $G(X)$ is annihilated by $\deg f$. ■

Remark.— One can actually deduce from [Bl-S, Theorem 1 (ii)] that $G(X) = 0$ — but we will not need this fact.

Proof of the Theorem: We can choose the resolution $\rho : \tilde{V} \rightarrow V$ so that $E := \rho^{-1}(\text{Sing}V)$ is a normal crossing divisor, whose irreducible components are smooth and *rational* [Fuj, Corollary of Theorem 1].

Let \tilde{C} and \tilde{C}^- be the images in V of C and C^- , and let \tilde{C} and \tilde{C}^- be their proper transforms in \tilde{V} . We have $[\tilde{C}] - [\tilde{C}^-] = \frac{1}{9}\pi_*([C] - [C^-]) = 0$ in $H^4(V^\circ, \mathbb{Q})$. Now we have an exact sequence [D, Corollaire 8.2.8]

$$H^2(\tilde{E}, \mathbb{Q}) \xrightarrow{i_*} H^4(\tilde{V}, \mathbb{Q}) \rightarrow H^4(V^\circ, \mathbb{Q}),$$

where \tilde{E} is the normalization of E , and i the composition $\tilde{E} \rightarrow E \hookrightarrow \tilde{V}$. Therefore we have $[\tilde{C}] - [\tilde{C}^-] = i_* z$ in $H^4(\tilde{V}, \mathbb{Q})$ for some class $z \in H^2(\tilde{E}, \mathbb{Q})$. Since the components of \tilde{E} are rational, z is the class of an element \mathfrak{z} of $A_1(\tilde{E}) \otimes \mathbb{Q}$. Then $[\tilde{C}] - [\tilde{C}^-] - i_* \mathfrak{z} \in A_1(\tilde{V}) \otimes \mathbb{Q}$ is homologous to zero, hence equal to zero by Lemma 3. Restricting to $\tilde{V} \setminus E \cong V^\circ$, we get $[\tilde{C}] - [\tilde{C}^-] = 0$ in $A_1(V^\circ \otimes \mathbb{Q})$, hence $[C] - [C^-] = \pi^*([\tilde{C}] - [\tilde{C}^-]) = 0$ in $A_1(J^\circ) \otimes \mathbb{Q}$. But the restriction map $A_1(J) \rightarrow A_1(J^\circ)$ is an isomorphism [Ful, Example 10.3.4], hence the Theorem. ■

3. COMPLEMENTS

Corollary 1. *Let Θ be a Theta divisor on J . We have $[C] = \frac{[\Theta]^2}{2}$ in $A_1(J) \otimes \mathbb{Q}$ (Poincaré formula).*

Proof : Indeed for any genus 3 curve C we have $[\Theta]^2 = [C] + [C^-]$ in $A_1(J) \otimes \mathbb{Q}$ (if p, q are two distinct points of C , the intersection of Θ with its translate by $[p] - [q]$ is the union of a translate of C and a translate of C^- — see for instance [Mu, Lecture IV]). Thus the corollary is equivalent to the theorem. ■

Recall that the *modified diagonal cycle* $\Gamma(C, p)$, first considered in [G-S], is the element $\Gamma(C, p)$ of $A_1(C^3)$ defined as follows. We denote by $[x, x, x], [x, x, p], [x, p, p]$ etc. the classes in $A_1(C \times C \times C)$ of the image of C by the maps $x \mapsto (x, x, x)$, $x \mapsto (x, x, p)$, $x \mapsto (x, p, p)$ etc. Then:

$$\Gamma(C, p) := [x, x, x] - [x, x, p] - [x, p, x] - [p, x, x] + [x, p, p] + [p, x, p] + [p, p, x].$$

By [FLV, Remark 3.4], we have

Corollary 2. $\Gamma(C, p) = 0$ in $A_1(C^3) \otimes \mathbb{Q}$.

Finally let us mention the result of [B3]: the class of $[C] - [C^-]$ in the intermediate Jacobian $\mathcal{J}_1(J)$ is torsion. It can be also deduced from our theorem, though the proof in [B3] is more direct.

In [BLLS] the authors construct a genus 7 curve with the same property, and suggest that the corresponding Ceresa cycle should be torsion modulo algebraic equivalence (Remark 1.2).

REFERENCES

- [B1] A. Beauville: *Sur l'anneau de Chow d'une variété abélienne*. Math. Ann. **273** (1986), 647-651.
- [B2] A. Beauville: *Algebraic cycles on Jacobian varieties*. Compositio Math. **140** (2004), 683-688.
- [B3] A. Beauville: *A non-hyperelliptic curve with torsion Ceresa class*. C.R. Acad. Sci. Paris, to appear; preprint arXiv:2105.07160.
- [BLLS] D. Bisogno, W. Li, D. Litt, P. Srinivasan: *Group-theoretic Johnson classes and non-hyperelliptic curves with torsion Ceresa class*. Preprint arXiv:2004.06146.
- [Bl] S. Bloch: *Algebraic cycles and values of L-functions*. J. reine angew. Math. **350** (1984), 94-107.
- [Bl-S] S. Bloch, V. Srinivas: *Remarks on correspondences and algebraic cycles*. Amer. J. Math. **105** (1983), no. 5, 1235-1253.
- [B-K] E. Brieskorn, H. Knörrer: *Plane algebraic curves*. Birkhäuser Verlag, Basel, 1986.
- [BST] J. Buhler, C. Schoen, J. Top: *Cycles, L-functions and triple products of elliptic curves*. J. Reine Angew. Math. **492** (1997), 93-133.
- [C] G. Ceresa: *C is not algebraically equivalent to C⁻ in its Jacobian*. Ann. of Math. **117** (1983), no. 2, 285-291.
- [C-vG] E. Colombo, B. van Geemen: *Note on curves in a Jacobian*. Compositio Math. **88** (1993), 333-353.
- [C-P] A. Collino, G. P. Pirola: *The Griffiths infinitesimal invariant for a curve in its Jacobian*. Duke Math J. **78** (1995), 59-88.
- [D] P. Deligne: *Théorie de Hodge, III*. Publ. Math. IHES **44** (1974), 5-77.
- [Fuj] A. Fujiki: *On resolutions of cyclic quotient singularities*. Publ. Res. Inst. Math. Sci. **10** (1974/75), no. 1, 293-328.
- [Ful] W. Fulton: *Intersection theory*. Ergebnisse der Math. **2**. Springer-Verlag, Berlin, 1984.
- [FLV] L. Fu, R. Laterveer, C. Vial: *Multiplicative Chow-Künneth decompositions and varieties of cohomological K3 type*. Ann. Mat. Pura Appl. **200** (2021), no. 5, p. 2085-2126.
- [G-S] B. Gross, C. Schoen: *The modified diagonal cycle on the triple product of a pointed curve*. Ann. Inst. Fourier **45** (1995), 649-679.
- [H] B. Harris: *Homological versus algebraic equivalence in a Jacobian*. Proc. Nat. Acad. Sci. U.S.A. **80** (1983), no. 4, 1157-1158.
- [M] Y. Miyaoka: *On the Kodaira dimension of minimal threefolds*. Math. Ann. **281** (1988), 325-332.
- [Mu] D. Mumford: *Curves and their Jacobians*. The red book of varieties and schemes, Lecture Notes in Math. **1358**. Springer-Verlag, Berlin, 1999.
- [N] M. Nori: *Cycles on the generic abelian threefold*. Proc. Indian Acad. Sci. Math. Sci. **99** (1989), no. 3, 191-196.
- [R] M. Reid: *Canonical 3-folds*. Journées de Géométrie Algébrique d'Angers, pp. 273-310, Sijthoff & Noordhoff, 1980.
- [V] C. Voisin: *Birational invariants and decomposition of the diagonal*. Birational Geometry of Hypersurfaces, pp. 3-71; Lecture Notes of the UMI **26**, Springer, 2019.
- [Z] S-W Zhang: *Gross-Schoen cycles and dualising sheaves*. Invent. math. **179** (2010), no. 1, 1-73.

UNIVERSITÉ CÔTE D'AZUR, CNRS – LABORATOIRE J.-A. DIEUDONNÉ, PARC VALROSE, F-06108 NICE CEDEX 2, FRANCE
 Email address: arnaud.beauville@unice.fr

DEPARTMENT OF MATHEMATICS, DUKE UNIVERSITY BOX 90320, DURHAM NC 27708-0320, USA
 Email address: schoen@math.duke.edu