

Strangeness content of the pion in the U(3) Nambu-Jona-Lasinio model

Fábio L. Braghin

Instituto de Física, Federal University of Goiás,
Av. Esperança, s/n, 74690-900, Goiânia, GO, Brazil

May 2, 2022

Abstract

The Nambu-Jona-Lasinio model is considered with flavor-dependent coupling constants $G_{ij} [(\bar{\psi}\lambda_i\psi)(\bar{\psi}\lambda_j\psi) + (\bar{\psi}i\gamma_5\lambda_i\psi)(\bar{\psi}i\gamma_5\lambda_j\psi)]$ for $i, j = 0, 1 \dots N_f^2 - 1$, and $N_f = 3$. A self consistent calculation of quark effective masses and coupling constants is performed making the strange quark effective mass to vary considerably. Quantum mechanical mixings between up, down and strange constituent quarks yields a strangeness content of the light u and d quarks constituent and of the pions. Different estimates for the strangeness contribution for the pion mass are provided.

1 Introduction

Hadron structure involves a large number of degrees of freedom and intrincated phenomena. Consequently their description with analytical methods in Quantum Chromodynamics presents many difficulties that usually require some approximative schemes being also possible to resort to effective models usually valid within a range of a variable, usually energy, associated to some physical scale. In spite of the limitations of an effective model, when compared to first principles calculations, one might expect that improvements eventually may produce a framework hoppefully comparable to effective field theories (EFT) if fundamental properties of QCD are taken into account by introducing the correct degrees of freedom in a suitable and correct way. Besides that, effective models can show very clearly the main connections between observables and the corresponding relevant degrees of freedom. The quark-level Nambu-Jona-Lasinio model (NJL) [1, 2, 3] captures some important features of quark dynamics and it has shown to be appropriate to describe several aspects of hadrons whenever Dynamical Chiral Symmetry Breaking (DChSB) plays an important role. It provides a framework, in general, consistent with the constituent quark model [4]. Constituent, or dressed, quark masses are obtained with contributions of the chiral condensates that, added to the masses originated from the Higgs boson, provides the correct scale of magnitude of hadrons masses [5, 6]. Usually, DChSB is only produced as long as coupling constants are minimally strong and this imposes restrictions in coupling constant of the NJL model. A strongly interacting gluon cloud can be seen as to give rise to (at least part of) the NJL coupling constant [7, 8, 9, 10] and, among other possible related mechanisms,

trace anomaly might also be involved in that [11]. Other similar successful calculations with contact-interactions, usually vector-current interactions inspired in large gluon effective mass induced interaction, have also been done [12, 13], although there might have a (at least partial) relation between these models because of the Fierz transformations.

Light hadron spectrum is mostly described quite well by NJL-type models in agreement with the quark model for which therefore, in addition to valence quarks, their corresponding quark-antiquark condensates produce important contributions. Indeed, other partons, besides the valence quarks, were shown to be needed to describe with precision hadron masses, spin and other properties in different approaches, for few examples [5, 17, 14, 15, 16]. Among these partons, strange sea quarks might yield a strangeness content of the non-strange light hadron sector by starting with the nucleon properties. Results for electromagnetic properties of the nucleon suggest a strangeness content (s-content) to be of the order of 5% [18, 20, 19, 21, 22, 23, 24]. The strange, up and down sea quark densities, however, were found to be nearly the same [21]. Among the light hadrons, pions and kaons are quasi-Goldstone bosons of the DChSB and it becomes of special interest to understand further their detailed structure. Current investigations in different facilities, for example EIC, JLab and AMBERS-CERN, are planned to investigate the (parton structure of the) pion and the kaon. Earlier estimates about the strangeness content of the pion by means of mesons loops, that reduce mesons masses [25], an extremely small contribution to their masses was found [26]. In the present work this subject is investigated in the framework of the NJL model.

Quark-antiquark polarization for the NJL-model was found to provide flavor dependent corrections to the NJL coupling constant [27]. Being an effective model for QCD, it is reasonable to expect that all the (free) parameters of the model might be traced back to degrees of freedom of QCD. At the QCD level the quark current masses are the only parameters containing flavor symmetry breaking and therefore, in an effective model, contributions of these parameters should be expected for all the free parameters of the effective model, similarly to the underlying ideas for an effective field theory (EFT) [28, 29, 30]. Preliminary perturbative estimation for pseudoscalar and scalar light mesons masses, in the absence of mixing-type interactions, showed that flavor-dependent coupling constants change resulting mesons masses slightly less than the flavor-dependence of quark effective masses and the sizeable corrections improve description of several observables. In the present work, a self consistent calculation of coupling constants and effective masses will be done. In a self consistent calculation the dynamics of different quark flavors become directly interdependent and each of the quarks, u,d and s, develops components due to the others, as it will be shown explicitly. The resulting s-content of u and d constituent quarks can be understood in terms of the quantum mixings because of the eigenstates, with corresponding interactions, of two different flavor-U(N) representations for quarks and quark-antiquark mesons [31, 5]. Although fundamental up-down-strange quark mixings are given in terms of the Cabibbo angle for the Cabibbo-Kobayashi-Maskawa (CKM) matrix [32, 17], the nature of the polarization process and the amplification of the quark masses due to DChSB make less clear a direct and unambiguous association of the mixing found in this work with the corresponding *amplified* Cabibbo angles. The method employed to derive these coupling constants made possible to derive form factors for constituent quarks interacting with quark-antiquark light mesons in a dynamical way, as well as average quadratic radii, and also a whole effective field theory for large N_c QCD with corrections due to quark current masses [33, 34].

Therefore, in this work, the strangeness content of the u and d constituent quarks and of the neutral and charged pion masses will be analyzed in the NJL model with flavor-dependent coupling constants. Different estimations for the strangeness contribution for the up and down quarks effective masses and for the neutral and charged pion masses will be provided. The work is organized as follows. In the next section the whole framework will be reminded with particular attention to the definition of the coupling constants and of logics of the self consistent calculation of G_{ij} and quark effective masses will be presented. The bound state

equation, a Bethe-Salpeter equation at the Born level, i.e. with constant amplitude, (BSE) for the quark-antiquark pseudoscalar mesons will be also briefly reminded. Numerical results will be presented in the following section for sets of coupling constants generated by three different gluon propagators. Results will be compared with a calculation for the flavor-independent NJL model with a coupling constant of reference G_0 . The neutral pion and kaon masses, or conversely the charged pion and kaon masses, will be used to fix the set of parameters with which further observables are also presented to assess the overall predictions of the model within the self consistent calculation. After the self consistent calculation, the strange quark effective mass will be freely varied, by keeping the self consistency of the up and down constituent quark effective masses and the coupling constants, such as to understand the role of the quark-antiquark strange condensate on the up and down quark effective masses and pion masses. Particular values for mass differences will be also presented such that different strange quark contributions for the up and down quark and the pion masses are envisaged. In the last section there is a Summary.

2 Masses and coupling constants: a self consistent analysis

The generating functional of the NJL model with flavor dependent corrections to the coupling constants can be written as:

$$Z[\eta, \bar{\eta}] = \int D_{\psi, \bar{\psi}} \exp \left[i \int_x \left\{ \bar{\psi}(i\not{\partial} - m_f)\psi + G_{ij} [(\bar{\psi}\lambda_i\psi)(\bar{\psi}\lambda_j\psi) + (\bar{\psi}i\gamma_5\lambda_i\psi)(\bar{\psi}i\gamma_5\lambda_j\psi)] + L_s \right\} \right], \quad (1)$$

where $D_{\psi, \bar{\psi}} = D[\psi, \bar{\psi}]$ is the functional measure, $\int_x = \int d^4x$, the subscript $f=u,d,s$ is used for the flavor $SU(3)$ fundamental representation, $i, j = 0, \dots, N_f^2 - 1$ is used for flavor indices in the adjoint representation, being $N_f = 3$ the number of flavors, and λ_i are the flavor Gell-Mann matrices with $\lambda_0 = \sqrt{2/3}I$. Quark sources are encoded in $L_s = \bar{\eta}\psi + \bar{\psi}\eta$. The coupling constant $G_{ij} = (G_0 + \tilde{G}_{ij})$ have a component due to gluon dynamics that is calculated for the case of a global color model in which quark-antiquark interaction is mediated by a non perturbative gluon exchange. By means of the background field method in the very long wavelength limit they were found to be given by:

$$\tilde{G}_{ij} = d_2 N_c (\alpha g^2)^2 \text{Tr}_D \text{Tr}_F \int \frac{d^4k}{(2\pi)^4} S_{0f}(k) R(k) i\gamma_5 \lambda_i S_{0f}(-k) R(-k) i\gamma_5 \lambda_j, \quad (2)$$

where Tr_D, Tr_F are the traces in Dirac and flavor indices, $\alpha = 4/9$, g^2 is the running quark-gluon coupling constant, $d_n = \frac{(-1)^n}{2^n}$, $S_{0f}(k)$ is the Fourier transform of the effective quark propagator $S_{0f}(x-y)$ which account for the DChSB by means of the quark effective mass M_f or M_f^* as discussed below. In this last equation $R(k) = 2(R_T(k) + R_L(k))$, where $R_T(k)$ and $R_L(k)$ are transversal and longitudinal components of an effective gluon propagator in a covariant gauge. Other types of contributions, due to gauge boson dynamics and confinement, proportional to delta functions, $\delta(p^2)$, provide smaller or vanishing contributions [27]. The following important properties, due to CP and electromagnetic U(1) invariances, hold:

$$G_{ij} = G_{ji}, \quad G_{22} = G_{11}, \quad G_{55} = G_{44}, \quad G_{77} = G_{66}. \quad (3)$$

The mixing type interactions $G_{i \neq j}$ are proportional to quark effective mass differences and therefore they have considerably smaller numerical values. Within the usual approach for the NJL, scalar and pseudoscalar, S_i, P_i , auxiliary fields are introduced by means of an unit integral multiplied in the generating functional with the corresponding shifts with quark currents that make possible the integration of the quark field. In the limit of degenerate quark effective masses, $M_u = M_d = M_s$, the coupling constants reduce to a single

constant that, as discussed below, will be normalized to be the coupling constant of reference G_0 such that: $G_{ij} \rightarrow G_0 \delta_{ij}$ and the standard treatment of the model can be done. In this case the quark effective masses of constituent quarks are obtained with the contribution of the scalar-quark-antiquark condensate, $M_f = m_s + \bar{S}_f$, where \bar{S}_f are the solutions of the auxiliary field gap equations. Gap equations might be found as saddle point equations for the scalar and pseudoscalar auxiliary fields S_i, P_i and non trivial solutions should emerge for the (neutral) scalar fields S_0, S_3, S_8 . For the coupling constant of reference G_0 these equations can be written as:

$$(G1) \quad M_f - m_f = G_0 \text{Tr}(S_{0,f}(0)). \quad (4)$$

The gap equations for the model (1) however receive corrections from the flavor dependent coupling constants. The final values of the coupling constants G_{ij} therefore are obtained from a self consistent calculation with the corrected gap equations as discussed below such that:

$$G_{ij} = G_{ij}(M_u^*, M_d^*, M_s^*). \quad (5)$$

In these equations for G_{ij} one has the first type of mixing interactions, i.e. $G_{i \neq j}$, that are numerically much smaller than the diagonal ones G_{ii} because they depend on the differences between quark effective masses and they will not be considered in most part of this work.

Corrections to the coupling constant from polarization, however, might produce spurious increasing values of the NJL coupling constant that is a free parameters of the model. To make possible comparisons of numerical results from different choices of the gluon propagator for G_{ij} with results from a coupling constant of reference, $G_0 = 10 \text{GeV}^{-2}$, a normalization procedure will be adopted after the calculation of the integrals in eqs. (5). Since polarization process should produce corrections to an initial NJL -coupling constant, say G_0 the following resulting complete coupling constant should be obtained to compute observables:

$$G_{ij}^{comp} = (G_0 + G_{ij}) \bar{G}_0, \quad (6)$$

where G_{ij} is obtained by eq. (2) and \bar{G}_0 is a renormalization factor that brings the resulting value to a value of reference whenever the symmetric limit is reached, i.e. $G_{ij}(m^*) = G_0 \delta_{ij} = 10 \text{GeV}^{-2}$, being, in that limit, $m^* = m_u^* = m_d^* = m_s^*$. Besides that, the different effective gluon propagator with the running quark-gluon coupling constant, defined below, have different normalizations and it becomes important to normalize all the results by a common factor to make possible to understand the role of each of the variables in the set of parameters and effective gluon propagator. By choosing, for example, the charged pion mass to be a fitted parameter/observable for $G_{11}^{comp} = 10 \text{ GeV}^{-2}$, the following normalization, written in the main text, can be used:

$$G_{i=j}^n \equiv G_{i=j}^{comp} = 10 \times \frac{G_{sym} \delta_{ij} + G_{ij}}{G_{sym} \delta_{ij} + G_{11}}. \quad (7)$$

Because the coupling constants G_{11} is almost equal to G_{33} , it makes basically no difference to adopt neutral or charged pion mass to be a fitted parameter. In the flavor-symmetric calculation for the NJL model, polarization effect contributes with around 20% – 30% of the original value of G_0 [35, 36] and the above normalization is compatible with these results. For the mixing type interactions $G_{i \neq j}$ a similar reasoning is adopted, being that in the flavor symmetric limit and in the original NJL model $G_{i \neq j}^n = 0$, so that one can write:

$$G_{i \neq j}^n = 10 \times \frac{G_{ij}}{G_{sym}}. \quad (8)$$

This normalization is compatible with the one for diagonal G_{ii} although it is somewhat arbitrary. This normalization (7) is different from the one considered in the perturbative investigation [27] and, although numerical results for G_{ij} are somewhat similar to the ones presented in the perturbative case just mentioned, it was found to be more appropriated for the convergence of the self consistent numerical calculations.

These corrections for the NJL-coupling constants can re-arrange quark effective masses. Restricting to the diagonal generators, $i, j = 0, 3, 8$, the coupling constants for the diagonal flavor singlet quark currents, G_{ff} ($f = u, d, s$), can be defined in the following way:

$$G_{ij}^n(\bar{\psi}\lambda_i\psi)(\bar{\psi}\lambda_j\psi) = 2 G_{f_1f_2}(\bar{\psi}\lambda_{f_1}\psi)(\bar{\psi}\lambda_{f_2}\psi), \quad (9)$$

where λ_{f_1} are three single-entry matrices obtained from combinations of diagonal Gell-Mann matrices with a single non zero matrix element that are: $\lambda_u = 1e_{11}$, $\lambda_d = 1e_{22}$ and $\lambda_s = 1e_{33}$, where e_{ii} is a diagonal matrix element. The following relations between the coupling constants G_{ii} and G_{ff} in the absence of the (numerically smaller) mixing-type interactions, $G_{i \neq j} = G_{f_1 \neq f_2} = 0$, are obtained:

$$\begin{aligned} 2G_{uu} &= 2\frac{G_{00}^n}{3} + G_{33}^n + x_s\frac{G_{88}^n}{3}, \\ 2G_{dd} &= 2\frac{G_{00}^n}{3} + G_{33}^n + x_s\frac{G_{88}^n}{3}, \\ 2G_{ss} &= 2\frac{G_{00}^n}{3} + 4x_s\frac{G_{88}^n}{3}, \end{aligned} \quad (10)$$

where x_s is an *ad hoc* parameter to control the strength of G_{88} . For equal quark masses the flavor independent coupling constants reduce to an unique constant $G_{ij} = G_{sym}\delta_{ij}$ and $G_{f_1f_2} = G_{sym}\delta_{f_1f_2}$. Note that for the diagonal interactions $i, j = 0, 3, 8$ one has $G_{uu} = G_{dd}$. Two cases for the strangeness content of the coupling constants will be considered by introducing a parameter x_s in G_{88} that provide the a contributions from the asymmetry of strange to up and down quark dynamics. Therefore an *ad doc* parameter x_s , that controls its strength will be introduced by multiplying G_{88} and it will be made variable to test with more details the contribution of the strangeness in the u and d sector. This parameter can be set $x_s = 1$ at any time, without loss of generality, in which case one can expect to reach a physical point that describes mesons masses. Also, it can be used to make the flavor-breaking content of G_{88} to be suppressed whenever $x_s G_{88} = 10 \text{ GeV}^{-2}$, that is the value of reference for the flavor symmetric point. The following cases will be considered into the equations of G_{ff} as written above:

$$\begin{aligned} (M2) \quad x_s G_{88} &= 10 \text{ GeV}^{-2}, \\ (M3) \quad x_s G_{88} &= G_{88} \text{ GeV}^{-2}, \end{aligned} \quad (11)$$

In the first case, $M2$, the role of strangeness does not take into account the flavor- asymmetry interaction G_{88} which is obtained from the eighth flavor generator $\lambda_8/2$. The second case, $M3$, is obtained with a more realistic account of the strange quark content.

The gap equations for the flavor dependent coupling constants in the absence of mixing-type interactions can be written as:

$$(G2) \quad M_f^* - m_f = G_{ff} \text{Tr} (S_{0,f}(0)), \quad (12)$$

where Tr includes traces in color, flavor and Dirac indices and momentum integral, and $S_{0,f}(x - y)$ is the quark propagator in terms of the quark effective masses M_f^* .

2.1 Mesons bound state equation

Pseudoscalar auxiliary fields for the composite quark-antiquark states can describe pseudoscalar mesons. In particular for the case of the pseudoscalar mesons, the two point Green's function have pole at a time-like momentum $P^2 = P_0^2 = -M_{PS}^2$, at zero tridimensional (Euclidean) momentum $\vec{P} = 0$, where M_{PS} is the mass of the pseudoscalar meson. The NJL- model condition for the quark-antiquark pseudoscalar BSE can be written as:

$$1 - 2G_{ij}I_{f_1f_2}^{ij}(P_0^2 = -M_{PS}^2, \vec{P}^2 = 0) = 0, \quad (13)$$

where

$$I_{f_1f_2}^{ij}(P_0, \vec{P}) = \text{Tr}_{D,F,C} \int \frac{d^4k}{(2\pi)^4} \lambda_i i\gamma_5 S_{0,f_1}(k + P/2) \lambda_j i\gamma_5 S_{0,f_2}(k - P/2), \quad (14)$$

where the different flavor indices for the case of the pion bound states are the following: π^0 with $i, j = 3$ and $f_1, f_2 = u, d$ and π^\pm with $i, j = 1, 2$ and $f_1, f_2 = u, d$. Kaons and some of the scalar mesons were discussed in [27].

In particular for the case of the pions and kaons, one can write the following reduced equation:

$$(M_{PS}^2 - (M_{f_1}^* - M_{f_2}^*)^2)G_{ij}I_2^{f_1f_2} = \frac{G_{ij}}{2} \left(\frac{m_{f_1}}{G_{f_1f_1}M_{f_1}^*} + \frac{m_{f_2}}{G_{f_2f_2}M_{f_2}^*} \right) + 1 - \frac{1}{2} \left(\frac{G_{ij}}{G_{f_1f_1}} + \frac{G_{ij}}{G_{f_2f_2}} \right). \quad (15)$$

In this equation there is a logarithmic divergent integral given by:

$$I_2^{f_1f_2} = \int \frac{d^4k}{(2\pi)^2} \frac{1}{R_{f_1}} \frac{1}{R_{f_2}}, \quad (16)$$

where $R_f = k^2 + M_f^{*2}$ in Euclidean momentum space. These integrals are solved with the same 3-dim cutoff Λ of the gap equations,

3 Numerical results

Flavor dependent coupling constants were calculated by considering three different effective gluon propagators each of the two different sets of current quark masses and UV cutoff S and V_3 . These effective gluon propagators will be labeled by: $_{2,5}$ and $_6$. They incorporate the quark-gluon running coupling constant g^2 as shown below. The first effective gluon propagator ($_2$) is a transversal one extracted from Schwinger Dyson equations calculations [37, 38]. It can be written as:

$$(S_2, V_{3_2}) : D_2(k) = g^2 R_T(k) = \frac{8\pi^2}{\omega^4} D e^{-k^2/\omega^2} + \frac{8\pi^2 \gamma_m E(k^2)}{\ln \left[\tau + (1 + k^2/\Lambda_{QCD}^2)^2 \right]}, \quad (17)$$

where $\gamma_m = 12/(33 - 2N_f)$, $N_f = 4$, $\Lambda_{QCD} = 0.234 \text{ GeV}$, $\tau = e^2 - 1$, $E(k^2) = [1 - \exp(-k^2/[4m_t^2])]/k^2$, $m_t = 0.5 \text{ GeV}$, $D = 0.55^3/\omega$ (GeV^2) and $\omega = 0.5 \text{ GeV}$.

The second type of effective gluon propagator is based in a longitudinal effective confining parameterization [39] that can be written as:

$$(S_{5,6}, V_{3_{5,6}}) : D_{\alpha=5,6}(k) = g^2 R_{L,\alpha}(k) = \frac{K_F}{(k^2 + M_\alpha^2)^2}, \quad (18)$$

where $K_F = (0.5\sqrt{2}\pi)^2/0.6 \text{ GeV}^2$, as considered in previous works [34] to describe several mesons-constituent quark effective coupling constants and form factors, with, however, either a constant effective gluon mass ($M_5 = 0.8 \text{ GeV}$) or a running effective mass given by: $M_6 = M_6(k^2) = \frac{0.5}{1+k^2/\omega_6^2} \text{ GeV}$ for $\omega_6 = 1 \text{ GeV}$.

The sets of (free) parameters, that reproduce the neutral pion and kaon masses after the self consistent calculation, are given in Table (1): current quark masses m_u, m_d, m_s and the ultraviolet (UV) cutoff Λ . In this Table the resulting effective masses from the gap equation (G1), for G_0 , are also presented. It is important to emphasize that the only role of the effective gluon propagator is to produce numerical results for G_{ij} . The sets of parameters S and V_3 yield the same overall behavior of results when M_s^* is varied, therefore figures will be exhibited only for S . Having obtained these fittings from the self consistent calculation, for different sets of G_{ij} , kaons are neglected and the investigation of the contribution of the strangeness is done. For this, the free-variation of the effective mass, M_s^* , will be done, by keeping the effective masses of the up and down quarks calculated self consistently. The values of the mesons masses at the physical point, obtained from the sets of parameters of Table (1), will be shown below in Table (2). The strange quark current mass, m_s , is not really relevant for the pion observables, but it helps to define the physical point, where kaon masses are obtained, and to keep track of the value of the strange quark-antiquark condensate.

Table 1: Sets of parameters: Lagrangian quark masses, ultraviolet cutoff and the quark effective masses obtained from an initial NJL-gap equation (G1) for $G_0 = 10 \text{ GeV}^{-2}$.

set of parameters	m_u MeV	m_d MeV	m_s MeV	Λ MeV	M_u MeV	M_d MeV	M_s MeV
S	3	7	133	680	405	415	612
V_3	3	7	133	685	422	431	625

In figures (1) and (2) results for the self consistent calculation for the up and strange flavor-dependent coupling constants, G_{uu} and G_{ss} , are presented as functions of the (freely-varied) strange quark effective mass M_s^* for the sets S_2, S_5 and S_6 and for the parameterizations $M2$ and $M3$. All the resulting G_{ff} are normalized in the flavor symmetric point according to eq. (7), i.e. $G_{ff}(M^* = m^*) = 10 \text{ GeV}^{-2}$ when $M_u^* = M_d^* = M_s^* = m^*$. Results are sounder physically for $M_s^* \geq m_s \simeq 0.133 \text{ GeV}$, below this value, M_s^* represents nearly a variable strange quark current mass in the absence of self-consistency. Result for the down quark is the same as G_{uu} , according to eqs. (10). The different coupling constants G_{ij} obtained for the different effective gluon propagators (S_2, S_5 and S_6) may produce quite different numerical results (in particular for $M3$) although the overall behavior is basically the same. The behavior for small and large strange quark effective mass limits are quite different depending on the set S_2, S_5 or S_6 . For lower values of M_s^* , the sets S_2 and S_6 that have larger variations than S_5 . For larger strange quark effective masses the quantities G_{ff} become smaller and tend to reach finite definite values at $M_s^* \rightarrow \infty$ that tend to be nearly independent of the gluon propagator. Note that the strange quark effective mass that produces correct values for the kaon masses is around $0.550 - 0.580 \text{ GeV}$ (for $M3$), shown in the Table 4) below, for all the sets S_2, S_5, S_6 and for $V3_2, V3_5, V3_6$.




Figure 1: The up quark coupling constant G_{uu} , eq. (10), as a function of M_s^* , arbitrarily varied. Up and down quark masses are obtained self consistently from their gap equations (G2).




Figure 2: The strange quark coupling constant G_{ss} , eq. (10), as a function of M_s^* , arbitrarily varied. Up and down quark effective masses are obtained self consistently from their gap equations (G2).

In figure (3) the up quark effective mass, as self consistent solution to the gap ($G2$), eq. (12), is presented as a function of the strange quark effective mass M_s^* that is made to vary freely. Again, the figure has a clear meaning for $M_s^* > m_s$. The point in which all the cases coincide is the symmetric point $G_{ij} = G_{sym}\delta_{ij}$ due to the normalization adopted.

The self consistency is implemented for both cases $M2$ and $M3$ that controls the strangeness dependence of G_{88} . The "physical value" of M_s^* , i.e., the value that reproduces the correct kaons masses, being solution of the gap equation $G2$, is around $0.500 - 0.600\text{GeV}$ depending on set $S2, S5, S6$, as presented in Table (2). This self consistence procedure lowers the values of the quark effective masses. It is interesting to note that both limits, zero and very large strange quark effective mass, might be, in different ways, somehow associated to absence of strangeness in the up and down quark dynamics.




Figure 3: Self consistent solutions for the up quark effective mass, M_u^* , obtained from the gap equation ($G2$) (12), as a function of M_s^* that is arbitrarily varied.

In figure (4) the difference between the self consistent solutions of down and up quark effective masses, $M_d^* - M_u^*$, is presented as a function of the strange quark effective mass M_s^* that is freely varied. Again it is important to stress that the quantity $M_s^* \geq m_s$ corresponds to varying the strange chiral condensate arbitrarily. It can be seen that the parameterizations $M3$ (smaller symbols) yield much larger variation of $M_d^* - M_u^*$ mainly for the case of smaller strange quark masses. The behavior with M_s^* is the opposite of the individual quark effective masses M_u^*, M_d^* . The difference in the results between the sets $S2, S5, S6$ (i.e. effective gluon propagator) reaches around only 1 MeV either for $M2$ or $M3$, that is of the order of 10% of the effective mass difference.




Figure 4: Self consistent values for the down and up quarks effective mass difference ($M_d^* - M_u^*$) as a function of M_s^* , arbitrarily varied, both of them obtained from the gap equations ($G2$) eqs. (12).

In figure (5) the neutral pion mass as a function of the strange quark effective mass is exhibited for the self consistent values of M_u^*, M_d^* and coupling constants and for M_s^* freely varying. Because of the normalization adopted, the point in which all the cases coincide is the symmetric point for which $G_{ij} = G_{sym}\delta_{ij}$. The same sets of parameters $S2, S5$ and $S6$ were considered for the two parameterizations $M2$ and $M3$. Both limits of strange quark mass going to zero and going to infinite are well defined, although some points were left out of the figure to emphasize the behavior for $M_s^* > m_s$, i.e. for the strange quark condensate. It is seen that the variation of the pion mass with the strange quark effective mass is larger for smaller strange quark masses, i.e. smaller or vanishing strange quark condensate.




Figure 5: M_{π_0} as a function of M_s^* , arbitrarily varied. All the other parameters M_u^*, M_d^* and coupling constants are obtained self consistently.

Finally, in the figure (6) the mass difference of charged and neutral pions, $\Delta M_\pi = M_{\pi^\pm} - M_{\pi^0}$, is exhibited as a function of the strange quark effective mass, arbitrarily varied. Whereas parameterizations $M3$, smaller symbols, provide small values of ΔM_π for smaller M_s^* , for large strange quark masses parameterizations $M3$ tends however to produce an increase considerably larger than $M2$. $M2$ ($M3$) makes the mass difference to reach a maximum value close to $0.15 - 0.20\text{MeV}$ ($0.15 - 0.27\text{MeV}$) for $M_s^* > 0.8\text{GeV}$.




Figure 6: $\Delta M_\pi = M_{\pi^\pm} - M_{\pi^0}$ as a function of M_s^* , arbitrarily varied. All the other parameters M_u^*, M_d^* and coupling constants are obtained self consistently.

3.1 Other observables

In this section some of the observables calculated with the resulting quark and mesons masses and coupling constants are described and their values are displayed in Table II.

The quark-antiquark scalar condensate, chiral condensate, is defined as:

$$\langle (\bar{q}q)_f \rangle \equiv -Tr(S_{0,f}(k)), \quad (19)$$

and therefore is directly calculated by means of the solutions for the gap equations for the three flavors. Values in the Table correspond to the final self consistent solution of the (G2), i.e. eq. (12). These values improve initial estimation when calculated for G_0 and M_f .

The quark-meson, pion or kaon, coupling constants, or correspondingly the normalization of the field, obtained from the residue of the pole of the vertex can be written as [2, 3]:

$$G_{qqPS} = \left(\frac{\partial \Pi_{ij}(P^2)}{\partial P_0^2} \right)_{P_0^2 \equiv -M_{ps}^2}^{-2}, \quad (20)$$

where the values were calculated at the physical mesons masses $P_0^2 = -M_{ps}^2$, and the polarization tensor was written in eq. (14), $\Pi_{ij}(P^2) = I_{f_1 f_2}^{ij}(P_0^2, \vec{P}^2)$.

The weak decay constant of charged mesons, pion and kaon, $F_{ps} = F_\pi, F_K$, can be calculated as [2, 3]:

$$F_{ps} = \frac{N_c}{4} G_{qqPS} \int \frac{d^4 q}{(2\pi)^4} Tr_{F,D} (\gamma_\mu \gamma_5 \lambda_i S_{f_1}(q + P/2) \lambda_j S_{f_2}(q - P/2)) \Big|_{P_0^2 = -M_{ps}^2}, \quad (21)$$

where f_1, f_2 correspond to the quark/antiquark of the meson and i, j are the associated flavor indices as discussed for eq. (14).

The pseudoscalar mesons mixing that is responsible for the eta-eta' mass difference can also be addressed by means of the flavor-dependent coupling constants due to the mixing-type interaction G_{08} . The auxiliary fields can be introduced by means of functional delta functions in the generating functional [40, 41], for the case of the pseudoscalar fields one can write:

$$1 = \int D[P_i] \delta(P_i - G_{ik} j_{ps}^k), \quad (22)$$

where $j_{ps}^k = \bar{\psi} \lambda^k i \gamma_5 \psi$, $i, k = 0, 3, 8$ provides the needed components to describe the mesons η, η' and π_0 , and where the fields dimensions are properly taken into account. This method neglects possible non factorizations [42] which, nevertheless, can be expected to be small. This definition is completely compatible with usual auxiliary fields when mixing interactions are neglected. For the mixing-type interactions $G_{i \neq j}$ for $i, j = 0, 3, 8$ one can neglect the neutral pion field P_3 since its mixings interactions, G_{03} and G_{38} , are still smaller than G_{08} . By considering resulting auxiliary field and corresponding mesons masses, written in the adjoint representation, $M_{ii}^2 P_i^2$, one can write, in general, the following quadratic terms from the pseudoscalar auxiliary fields with the mixing interactions G_{ij} :

$$\mathcal{L}_{mix} = -\frac{M_{88}^2}{2} P_8^2 - \frac{M_{00}^2}{2} P_0^2 + 2G_{08}^n \bar{G}_{08} P_0 P_8 + \mathcal{O}(P_3, P_3^2) \dots \quad (23)$$

where M_{ii}^2 include the contributions from $G_{i=j}$ derived above, and

$$\bar{G}_{08} = \frac{2}{G_{00}^n \left(G_{88}^n - \frac{G_{08}^{n,2}}{G_{00}^n} \right)}, \quad (24)$$

where the mixing terms $G_{i \neq j}$ are exclusively obtained from the one-loop polarization. By performing the usual rotation to mass eigenstates η, η' , according to the convention from [43], it can be written:

$$\begin{aligned} |\eta\rangle &= \cos \theta_{ps} |P_8\rangle - \sin \theta_{ps} |P_0\rangle, \\ |\eta'\rangle &= \sin \theta_{ps} |P_8\rangle + \cos \theta_{ps} |P_0\rangle. \end{aligned} \quad (25)$$

Although one needs two angles to describe both masses, η, η' [44], in this work only the mass difference will be calculated being directly due to the mixing-type interaction G_{08} . Therefore resulting mixing angle θ_{ps} - as calculated for the mass difference - turns out to be, basically, a composition of the usual two mixing angles θ_0, θ_8 . By calculating \mathcal{L}_{int} in this mass eingestates basis, and comparing to the above 0 – 8 mixing, the following $\eta - \eta'$ mixing angle is obtained:

$$\theta_{ps} = \frac{1}{2} \arcsin \left(\frac{8G_{08}^n \bar{G}_{08}}{(M_\eta^2 - M_{\eta'}^2)} \right). \quad (26)$$

This equation provides numerical results similar to the equation used in [27] being however more complete. As commented above, the normalization of the coupling constants $G_{i \neq j}$ is more ambiguous than the diagonal ones and this has direct consequences in the estimations of θ_{ps} .

In Table (2) several observables calculated for the sets of parameters shown above (S and V_3 for the three different gluon propagators 2,5 and 6 and G_0) are exhibited. The neutral pion and kaon masses were fitted to values close to the experimental value, $M_{\pi^0} = 135$ MeV and $M_{K^0} = 498$ MeV. For the case of the

pion, there are two estimates, one for the set $M2$ and the other from $M3$. The flavor-dependent coupling constants, however, tends to lower the mesons masses with respect to their values calculated with G_0 . All the other observables, are obtained from the more complete calculation with $M3$. Self consistency, and a more complete account of strangeness in the coupling constants by means of $M3$, lead to lower values of the mesons masses and quark effective masses and the need of a larger value of the UV cutoff. The initial fit of the neutral pion mass, for the value of reference for the coupling constant $G_0 = 10 \text{ GeV}^{-2}$, was found to be $M_{\pi^0} = 136.4 - 137.1 \text{ MeV}$. The final value with the flavor-dependent coupling constants and strange effective mass close to value that reproduce a physical point, for $M3$ goes to $M_{\pi^0} = 133 - 135 \text{ GeV}$, whereas for $M2$ it goes to $M_{\pi^0} = 135.0 - 136.3 \text{ GeV}$. Although the sets $M2$ pin down the correct (expected) value the idea is to show the effect of the mixing by comparing with the more complete result from $M3$. The estimation for the charged pion and kaon masses, M_{π^\pm}, M_{K^\pm} , are in quite good agreement with experimental or expected values. Note that, electromagnetic effects are not taken into account, and the expected mass differences due to strong interactions effects have opposite signs and they are respectively $M_{\pi^\pm} - M_{\pi^0} \simeq 0.1 \text{ MeV}$ and $M_{K^\pm} - M_{K^0} \simeq -5.3 \text{ MeV}$ [45, 46, 47]. The kaon masses are exhibited for the sake of completeness to show the entire set of observables used to fit parameters in the self consistent part of the calculation and the prediction for the neutral-charged meson mass difference.

The values of the charged pion and kaon decay constants F_π, F_K are not far from the experimental/expected values (e.v.) although the choice made for the fitting yielded a much better value for the kaon decay constant than F_π , differently from usual results in the literature, see e.g. in [12] and references therein. These results show an improvement with respect to the standard NJL model treatment. The up, down and strange quarks condensates, $\langle \bar{u}u \rangle, \langle \bar{d}d \rangle$ and $\langle \bar{s}s \rangle$, and the pion (kaon)-quark coupling constants $G_{\pi qq}$ ($G_{K qq}$) are also presented. The last observable shown in the Table is the pseudoscalar mesons mixing angle θ_{ps} that provides the $\eta - \eta'$ mixing that is usually defined for other types of effective interactions [48, 49]. It emerges due to the mixing-interaction coupling constant G_{08} and it makes possible to reproduce the mass difference between $M_\eta = 548 \text{ MeV}$ and $M_{\eta'} = 958 \text{ MeV}$ [43]. These values still eventually may change further by different values of the UV cutoff and current quark masses besides effects from the mixing-type interactions. Although the values of all the observables are not as close as they could be to the experimental or expected values they are somewhat improved with respect to the flavor independent calculation which is represented in the Table by the column for the set of parameters with G_0 . It is important to stress that, the coupling constants of reference is slightly larger than usual values and this makes the values of the condensates to be larger than they should. Large values of G_0 , however, favor the convergence of the self consistent solutions for masses and coupling constants. A criterium for analyzing the s-content of the pion is the probability of finding a sea s-quark in it, denoted by Pr-s-content π . It can be approximately defined by means of the change in the pion normalization, $Z_\pi = G_{qq\pi}^2$, with respect to the calculation with G_0 . The most relevant reason for this change in the normalization is the variation of the strange quark effective mass (condensate). However, due to the self consistency of the problem, there might have other much smaller contributions due to up and down quarks. These values are considerably larger than the estimation from ref. [26] based on meson loops.

Finally, in the last two last lines the reduced chi-square for each of the set of parameters for two different situations are presented always for calculation with $M3$. Firstly the chiral condensates are taken into account, resulting in ten observables being two fitted observables, and secondly if the condensates are neglected, it provides seven observables with two fitted observables. The e.v. value for pseudoscalar mixing angle and quark-antiquark condensates were taken to be the average value of those shown in the last column of the Table. These two different estimations of the chi-square were done because, although the resulting values of the chiral condensates are improved with respect to the flavor-independent calculation, the deviations of their (corrected) values are still large with respect to the e.v. and this makes χ_{red}^2 to be

very large. Another source of increase of the chi-squared are the values of F_π . Although the pion decay constant was not really in agreement with experimental value, the important point is that the relative results $F_K - F_\pi$ is slightly improved with respect to standard NJL, that corresponds to the set of parameters for G_0 . As discussed above, the relatively large value of G_0 may responsible for these discrepancies, and further terms in the gap equations, eventually due to higher order interactions, vector interactions or mixing-type interactions, may also be needed to pin down the corrected e.v. The numerical values of quark masses and meson masses needed to calculate the entries of Tables (2) and (4) are displayed in Table (3).

Table 2: Numerical results for some observables of the pion and the kaon. Where it has not been indicated, only the more complete set [M3] was considered. The estimation of the strangeness content of the pion is Pr.s-content π . e.v. refers to experimental or expected values.

Observable	S2	S5	S6	G_0	V_3-2	V_3-5	V_3-6	V_3-G0	e.v.
M_{π^0} (MeV) [M2]	135.0	135.3	135.1	136.4	135.5	136.1	135.6	137.1	135 [43]
M_{π^0} (MeV) [M3]	133.5	134.2	133.7	136.4	134.15	134.9	134.4	137.1	
M_{π^\pm} (MeV) [M2]	135.2	135.4	135.3	136.7	135.7	136.2	135.8	137.4	
M_{π^\pm} (MeV) [M3]	133.7	134.4	133.9	136.7	134.4	135.0	134.5	137.4	
$M_{\pi^\pm} - M_{\pi^0}$ (MeV) [M3]	0.2	0.2	0.2	0.3	0.1	0.1	0.1	0.3	0.1 [43, 45]
M_{K^0} (MeV) [M3]	498.5	498.5	498.5	499	498	498	499	498	498 [43]
M_{K^\pm} (MeV) [M3]	490	491	493	490	486	487	488	490	494 [43]
F_π (MeV)	99	99	99	102	100	101	101	103	92
F_K (MeV)	111	111	111	112	112	112	111	113	111
$(-\langle \bar{u}u \rangle)^{1/3}$ MeV	331	334	332	343	336	338	336	347	240-260 [50, 51]
$(-\langle \bar{d}d \rangle)^{1/3}$ MeV	333	335	333	344	338	339	338	349	240-260 [50, 51]
$(-\langle \bar{s}s \rangle)^{1/3}$ MeV	348	353	349	366	352	356	353	369	290-300 [52]
$G_{qq\pi}$	3.3	3.2	3.2	3.4	3.3	3.3	3.3	3.6	
G_{qqK}	3.8	3.8	3.8	4.2	3.9	4.0	3.9	4.3	
θ_{ps}	-3.7	-2.7	-3.6	0.0	- 3.4	2.6	3.4	0.0	(-11°)-(-24°) [43]
Pr.s-content π	6%	10%	10%	0	16%	16%	16%	0	
χ_{red}^2 (with $\langle \bar{q}q \rangle$)	103	110	103	146	122	129	123	164	
χ_{red}^2 (without $\langle \bar{q}q \rangle$)	29	30	27	51	37	41	39	56	

3.2 Strangeness contribution for the masses of the u, d constituent/dressed quarks and pion

In the Table 3 further numerical results obtained from the change in M_s^* are displayed. In the upper part of the Table there are particular values for the effective quark effective masses M_u^*, M_d^* calculated self consistently (for M2 and M3). The values obtained from complete self consistent calculation - that reproduces the neutral and charged pion and kaon masses - are identified by G2. Both calculations for M2 and M3 correspond to different ways of taking into account G_{88} as responsible for a strange-up/down asymmetry. The resulting up and down quark effective masses when $M_s^* \rightarrow 0$, $M_s^* \rightarrow m_s$ and $M_s \rightarrow \infty$ are also shown. Note that the self consistent calculation of $M_{ch.L}^*$, that is a flavor symmetric limit with degenerate quark masses, for the chiral limit, $m_u = m_d = m_s = 0$, provides a lower value for the quark

effective masses than the value for $M_f^*(M_s^* \rightarrow 0)$ (f=u and d). Note also that self consistent calculation for $M_{u,d}^*$ can easily provide values lower than their value in the chiral limit. Values of the neutral pion masses are shown in the same limits of $M2$ and $M3$ - self consistent results identified by (G2) - and also for $M_s^* = 0$, $M_s^* = m_s$ and $M_s^* \rightarrow \infty$. The analysis is basically the same as that for the up and down constituent or dressed quarks above, being however that in the chiral limit $M_\pi^{ch.L} = 0$ since the pion is a Goldstone boson. The behavior of the charged pion mass is basically the same as the neutral pion mass as shown above.

Table 3: Numerical results for up and down quark effective masses and for the neutral pion mass, several of them obtained by varying freely the strange quark effective mass, M_s^* , and some of them obtained fully self consistently, identified by G2: M_f^* and $M_{\pi 0}$ for (M2) and (M3) and M^* (ch.lim.).

Observable/M3	S2	S5	S6	S-G ₀	V ₃ -2	V ₃ -5	V ₃ -6	V ₃ -G0	e.v.
M_u^* M3-(G2) (MeV)	367	377	368	405	385	393	387	422	
M_u^* M2-(G2) (MeV)	386	394	387	405	401	406	402	422	
$M_u^*(M_s^* \rightarrow 0)$ MeV	618	512	595	405	651	537	626	422	
$M_u^*(M_s^* \rightarrow m_{0,s})$ MeV	563	491	547	405	579	508	563	422	
$M_u^*(M_s^* \rightarrow \infty)$ MeV	290	310	295	405	300	316	304	422	
M_d^* M3-(G2) (MeV)	375	384	378	415	394	402	395	431	
M_d^* M2-(G2) (MeV)	396	399	396	415	410	415	411	431	
$M_d^*(M_s^* \rightarrow 0)$ MeV	625	520	602	415	657	544	632	431	
$M_d^*(M_s^* \rightarrow m_{0,s})$ MeV	555	491	541	415	585	514	568	431	
$M_d^*(M_s^* \rightarrow \infty)$ MeV	305	320	305	415	314	328	316	431	
M_s^* M3-(G2) (MeV)	555	567	558	612	566	581	569	625	
M_s^* M2-(G2) (MeV)	560	570	563	612	600	595	604	625	
ch.lim. $M_{ch.L}^*$ (MeV)	381	381	381	381	415	415	415	415	
$M_{\pi 0}$ M3-(G2) (MeV)	133.5	134.2	133.7	136.7	134.2	134.9	134.4	137.4	0.135 [43, 45]
$M_{\pi 0}$ M2-(G2) (MeV)	135.0	135.3	135.1	136.7	135.5	136.1	135.6	137.4	
$M_{\pi 0}(M_s^* \rightarrow 0)$ MeV	158	147	156	136.7	162	149	159	137.4	
$M_{\pi 0}(M_s^* \rightarrow m_{0,s})$ MeV	150	144	149	136.7	150	144	149	137.4	
$M_{\pi 0}(M_s^* \rightarrow \infty)$ MeV	129	129	129	136.7	129	129	129	137.4	

Different ways of defining strangeness and flavor asymmetry content of the constituent, or dressed, up and down quarks and of the pion can be envisaged. It becomes useful to define particular differences of values that might correspond to variations with specific meanings. From here on, these quantities will be referred as to $T_{ff} = M_f^*, M_\pi$ and also G_{ff} . Furthermore these quantities defined below can also apply to the coupling constants G_{ff} that present basically the same behavior of the up, down quark effective masses when varying M_s^* . We will make use of the following differences of a quantity T_{ff} to characterize a specific s-content of the up and down quarks and of the pion:

$$\Delta_s^{2,3} = T_{ff}(M3) - T_{ff}(M2), \quad (27)$$

$$\Delta_s^0 = T_{ff}(M_s'^*) - T_{ff}(M_s^* = 0), \quad (28)$$

$$\Delta_s^{m_0} = T_{ff}(M_s'^*) - T_{ff}(M_s^* = m_s), \quad (29)$$

$$\Delta_s^\infty = T_{ff}(M_s'^*) - T_{ff}(M_s^* \rightarrow \infty). \quad (30)$$

First, note that the difference between the two curves, $M2$ and $M3$, can be considered as a first measure of the effect of the flavor- asymmetry (for the strange quark) for sea quarks in the coupling constants due to coupling G_{88} . This quantity $\Delta_s^{2,3}$ is defined at the physical point.

Deviations with respect to the limit in which the strange quark effective mass is zero is encoded in Δ_s^0 , and it could be interpreted as the overall contribution of the strange quark effective mass to the observable T_{ff} . The deviation of the quantity T_{ff} with respect to the point where strange quark condensate goes to zero was defined as $\Delta_s^{m_0}$. This is an effective measure of the strange quark condensate in the constituent/dressed quark or pion. These mass differences, however, are not fully extracted in physical points in the sense that in these limits, $M_s^* = 0$ and $M_s^* = m_s$, there are no non trivial solutions for the full self consistent problem and kaons are not bound. The mass difference Δ_s^∞ provides a dynamical way to measure the shift on the value of T_{ff} due to the the strange quark effective mass. In its definition one considers the limit in which the strange quark effective mass goes to infinite $T_{ff}(M_s^* \rightarrow \infty)$. Curiously this is a well definite limit with interesting physical appeal, since in this limit the strange quark degrees of freedom should be frozen.

The same reasoning done above for mass differences, with T_{ff} , applies for the G_{uu} and G_{ss} because these coupling constants present a very similar behavior with the change in M_s^* .

Some mass differences for the up and down quarks, calculated according to eqs. (27-30), are exhibited in Table (4). Two different sources of changes in the up and down quark effective masses can be immediately identified. Firstly the shift in effective masses due to the explicit chiral symmetry breaking by mean of the quark current masses, and secondly the shift in the quark effective masses due to the flavor-dependent coupling constants by means of the self consistent procedure. By denoting the quark effective mass in the chiral limit ($m_0 = 0$) by $M_{ch.L.}$ these two mass shifts can be written respectively as:

$$\Delta_{chL}^f \simeq M_f - M_{ch.L.}, \quad (31)$$

$$\Delta_{G_{ij}}^f \simeq M_f^* - M_f, \quad (32)$$

where M_f are solutions for the first gap equation $G1$ and M_f^* are solutions for the second gap equations $G2$. This second quantity, $\Delta_{G_{ij}}^f$ for the up and down quarks, is mainly due to strangeness content of the coupling constant and, less importantly, also correspond to providing a u-content (d-content) for the d (u) quark effective mass. The numerical values for the neutral pion mass differences, eqs. (27-30), are also shown in Table (4), and they present the same relative behavior of the mass differences of up and down quarks.

Table 4: Numerical results for up and down quark effective mass differences and neutral pion mass differences defined in eqs. (27-30). Results for the charged pion mass are very similar and proportional as shown in Figure (6).

Observable [M3]	S2	S5	S6	S- G_0	V_3 -2	V_3 -5	V_3 -6	V_3 -G0
Δ_u^{chL} (MeV)	24	24	24	24	7	7	7	7
Δ_d^{chL} (MeV)	34	34	34	34	16	16	16	16
$\Delta_{G_{ij}}$ (MeV)	- 38	-28	-37	0	-37	- 29	-35	0
$\Delta_{G_{ij}}$ (MeV)	-40	-31	-37	0	-37	-29	-36	0
$\Delta_s^{2,3}(M_u^*(s))$ (MeV)	-19	-17	-19	0	-16	-13	-15	0
$\Delta_s^0(M_u^*(s))$ (MeV)	-251	-135	-227	0	-266	- 144	-239	0
$\Delta_s^{m_0}(M_u^*(s))$ (MeV)	-194	-114	-179	0	-194	-115	-176	0
$\Delta_s^\infty(M_u^*(s))$ (MeV)	77	67	73	0	85	77	83	0
$\Delta_s^{2,3}(M_\pi^*(s))$ (MeV)	-1.5	-1.1	-1.4	0	-1.3	-1.2	-1.2	0
$\Delta_s^0(M_\pi^*(s))$ (MeV)	-24	-13	-22	0	-28	-14	-25	0
$\Delta_s^{m_0}(M_\pi^*(s))$ (MeV)	-16	-10	-16	0	-16	-9	-14	0
$\Delta_s^\infty(M_\pi^*(s))$ (MeV)	6	5	4	0	5	6	5	0

4 Summary

A self consistent calculation for the quark effective masses and flavor-dependent coupling constants of the NJL model was employed to investigate the role of the strange quark effective mass on the up and down constituent quarks and on the pion masses. A fully self consistent calculation was done to fit the parameters of the model to reproduce neutral (or charged) pion and kaon masses. In this step, several observables were calculated to assess the reliability of the model for different sets of parameters. The strong value of the coupling constant of reference, $G_0 = 10 \text{ GeV}^{-2}$, in one hand helps with the convergence of the self consistent calculation but in the other hand induces considerable large values of the quark-antiquark condensates. Flavor dependent coupling constants, nevertheless, improve their values with respect to the ones obtained with G_0 . In general, results for light pseudoscalar mesons observables from Table (2) show the flavor dependent coupling constants help to improve predictions of the NJL model. Quark effective masses get lower and they require a larger cutoff to make possible the description of light mesons masses. Because of the quantum mixing it was possible to estimate the s-content of the up and down constituent/dressed quarks as well as of the pion even in the absence of mixing-type interactions. The different ways of extracting the strangeness contributions for the up and down quark effective masses and pion masses were analyzed in this work. The mass difference $\Delta_s^{2,3}$ is an interesting quantity since it provides the change in the pion (or up and down effective) mass when exchanging the value of the interaction G_{88} , obtained without any flavor-dependence or mixing effect (M2), with $G_{88} = 10\text{GeV}^{-2}$, by a value that takes into account the strange sea quark dynamics, (M3) with $G_{88} = G_{88}(M_s^*)$. The mass difference Δ_s^∞ corresponds to more dynamical criterium to define a strangeness contribution for the pion mass (or up and down effective masses) such that the deviation of the observables are taken with respect to the limit in which the strange quark mass goes to infinite, i.e. it becomes a frozen degree of freedom. The same analysis is valid for the flavor dependent coupling constants G_{ff} . It is important to note that the mixing investigated in the present work is responsible for the strangeness in the up and down quark sector corresponding to a different mechanism from the quark mixing induced by the Higgs field that gives rise to the CKM matrix. Although these two mechanisms should add to each other, polarization process with the self consistent procedure does not

necessarily lead to a simple rotation of dressed quarks but it might involve some other transformation, like a dilation. This can be associated not only to the DChSB but also to the chosen normalization that reflects the strength of the flavor-dependent corrections to G_0 , i.e. $\Delta G_{ij} = G_{ij} - G_0$. Besides that, due to the self consistent character of the calculation, results contain implicitly not only mixings of the type U_{us} and U_{ds} , like the CKM matrix, but also it contains effective mixings that can be labeled by U_{us}, U_{ds} and U_{ud} . Although the NJL-model does not exhibit confinement of quark and gluons, it provides an interesting and quite appropriate effective way of investigating aspects of hadron structure and dynamics, and it is not clear whether or how confinement would imply modification(s) in the different mixings interactions and mechanisms. A complete account of the flavor dependent - NJL model with mixing type interactions, $G_{i \neq j}$ and $G_{f_1 \neq f_2}$, will be reported in another paper.

Acknowledgements

F.L.B. thanks short discussions with B. El Bennich, C.D. Roberts and G.I. Krein. The author is member of INCT-FNA, Proc. 464898/2014-5 and he acknowledges partial support from CNPq-312072/2018-0 and CNPq-421480/2018-1.

References

- [1] Y. Nambu, G. Jona-Lasinio, Dynamical Model of Elementary Particles Based on an Analogy with Superconductivity I, Phys. Rev. **122**, 345 (1961).
- [2] S. P. Klevansky, The Nambu-Jona-Lasinio model of quantum chromodynamics, Rev. Mod. Phys. **64**, 649 (1992).
- [3] U. Vogl, W. Weise, The Nambu and Jona-Lasinio model, Progr. in Part. and Nucl. Phys. **27**, 195 (1991).
- [4] M. Lavelle, D. McMullan, *Constituent quarks from QCD*, Phys. Rept. **279**, 1 (1997). E. de Rafael, *The Constituent Chiral Quark Model revisited*, Phys. Lett. **B 703**, 60 (2011).
- [5] S. Weinberg, The Quantum Theory of Fields, Cambridge (1996).
- [6] S.J. Brodsky, C.D. Roberts, R. Schrock, P.C. Tandy, New perspectives on the quark condensate, Phys. Rev. C **82**, 022201 (2010).
- [7] J. L. Cortés, J. Gamboa, L. Velásquez, A Nambu-Jona-Lasinio like model from QCD at low energies, Phys. Lett. B **432**, 397 (1998).
- [8] P. Costa, O. Oliveira, P.J.Silva, What does low energy physics tell us about the zero momentum gluon propagator, Phys. Lett. B **695**, 454 (2011).
- [9] K.-I. Kondo, Toward a first-principle derivation of confinement and chiral-symmetry-breaking crossover transitions in QCD, Phys. Rev. D **82**, 065024 (2010).
- [10] E. Barros Jr, F.L. Braghin, Eighth order quark interaction with gluon condensate $\langle A^2 \rangle$, Phys. Rev. D **88**, 034011 (2013).
- [11] Y. B. Yang, *et al*, Proton Mass Decomposition from the QCD Energy Momentum Tensor, Phys. Rev. Lett. **121**, 212001 (2018).

- [12] F. E. Serna, B. El-Bennich, G. Krein, Charmed mesons with a symmetry-preserving contact interaction Phys. Rev. D96, 014013 (2017).
- [13] P.L. Yin, *et al*, Masses of positive- and negative-parity hadron ground-states, including those with heavy quark, arXiv:2102.12568v1.
- [14] D.F. Geesaman, P.E. Reimer, The sea of quarks and antiquarks in the nucleon, Rep. Prog. Phys. 82, 046301 (2019)
- [15] J. Dove *et al*, The asymmetry of antimatter in the proton, Nature 590, 561 (2021).
- [16] EMC, A measurement of the spin asymmetry and determination of the structure function g_1 in deep inelastic muon proton scattering, Phys. Lett. B 206, 364 (1988).
- [17] M.D. Schwartz, Quantum Field Theory and the Standard Model, Cambridge (2014). T.P. Cheng, L.F. Li, Gauge Theory of Elementary Particles, Oxford, (1984).
- [18] D. S. Armstrong, R. D. McKeown, Parity-Violating Electron Scattering and the Electric and Magnetic Strange Form Factors of the Nucleon, Ann.Rev.Nucl.Part.Sci. 62 (2012) 337-359.
- [19] R. D. Young, *et al*, Extracting Nucleon Strange and Anapole Form Factors from World Data, Phys. Rev. Lett. 97, 102002 (2006).
- [20] D.B. Leinweber, *et al*, Precise Determination of the Strangeness Magnetic Moment of the Nucleon, Phys. Rev. Lett. 94, 212001 (2005). D.B. Leinweber, *et al*, Strange Electric Form Factor of the Proton, Phys. Rev. Lett. 97, 022001 (2006)
- [21] M. Aaboud, G. Aad, ATLAS Collaboration, Precision measurement and interpretation of inclusive W^+ , W^- and Z/γ production cross sections with the ATLAS detector, Eur. Phys. Journ. 77, 367 (2017).
- [22] H. Abdolmaleki, *et al*, Probing the strange content of the proton with charm production in charged current at LHeC, DESY Report 19-107, arXiv:1907.01014v2,
- [23] G.S. Bali *et al*, A lattice study of the strangeness content of the nucleon, Progress in Particle and Nuclear Physics 67, 467 (2012).
- [24] R.D. Young, Strangeness in the proton, Nature 544, 419 (2017).
- [25] M.A. Pichowsky, S. Walawalkar, S. Capstick, Meson-loop contributions for the $\rho - \omega$ mass splitting and ρ charge radius, Phys. Rev. D60, 054030 (1999).
- [26] I.C. Cloet, C.D. Roberts, Form Factors and Dyson-Schwinger Equations, arXiv:0811.2018v1; LIGHT CONE 2008 Proceedings of Science, <http://pos.sissa.it/>, Mulhouse, France
- [27] F.L. Braghin, Flavor-dependent $U(3)$ Nambu–Jona-Lasinio coupling constant, Phys. Rev. D103, 094028 (2021).
- [28] S. Weinberg, Phenomenological Lagrangians, Physica (Amsterdam) 96A, 327 (1979).
- [29] J. Gasser, H. Leutwyler, Chiral perturbation theory to one loop, Ann. Phys. (N.Y.) 158, 142 (1984).
- [30] D. B. Kaplan, in Lectures on Effective Field Theory ICTP-SAIFR (2016).
- [31] J.J. Sakurai, S. F. Tuan, Modern Quantum Mechanics, Addison-Wesley (1985).
- [32] N. Cabibbo, Unitary symmetry and leptonic decays, Phys. Rev. Lett. 10, 531 (1963). L.L. Chau, W.-Y. Keung, Comments on the Parametrization of the Kobayashi-Maskawa Matrix, Phys. Rev. Lett. 53, 1802 (1984)

- [33] F.L. Braghin, Quark and pion effective couplings from polarization effects, *Eur. Phys. J. A* 52: 134 (2016) .
- [34] F.L. Braghin, Pion Constituent Quark Couplings strong form factors: A dynamical approach, *Phys. Rev. D* **99**, 014001 (2019).
- [35] A. Paulo Jr., F.L. Braghin, Vacuum polarization corrections to low energy quark effective couplings, *Phys. Rev. D* **90** 014049 (2014).
- [36] F.L. Braghin, SU(2) Higher-order effective quark interactions from polarization, *Phys. Lett. B* 761, 424 (2016).
- [37] A. Bashir, et al., Collective Perspective on Advances in Dyson–Schwinger Equation QCD, *Commun. Theor. Phys.* 58,79 (2012). I.C. Cloet, C.D. Roberts, Explanation and prediction of observables using continuum strong QCD, *Prog. Part. Nucl. Phys.* 77, 1 (2014).
- [38] D. Binosi, L. Chang, J. Papavassiliou, C.D. Roberts, Bridging a gap between continuum-QCD and ab initio predictions of hadron observables, *Phys. Lett. B* **742**, 183 (2015) and references therein.
- [39] J. M. Cornwall, Entropy, confinement, and chiral symmetry breaking, *Phys. Rev. D* 83, 076001 (2011).
- [40] H. Reinhardt, R. Alkofer, Instanton-induced flavour mixing in mesons, *Phys. Lett. B* 207, 482 (1988).
- [41] A.A. Osipov, B. Hiller, A.H. Blin, J. da Providência, Effects of eight-quark interactions on the hadronic vacuum and mass spectra of light mesons, *Annals of Phys.* 322, 2021 (2007).
- [42] F.L. Braghin, F.S. Navarra, Factorization breaking of four-quark condensates in the Nambu–Jona-Lasinio model, *Phys. Rev. D* 91, 074008 (2015).
- [43] M. Tanabashi et al. (Particle Data Group), *Phys. Rev. D* 98, (2018) 030001.
- [44] G. S. Bali, V. Braun, S. Collins, A. Schafer, J. Simeth, Masses and decay constants of the η and η' mesons from lattice QCD, arXiv:2106.05398v1 [hep-lat], and references therein.
- [45] J. Gasser, H. Leutwyler, Quark masses, *Phys. Rept.* 87, 77 (1982).
- [46] J. F. Donoghue, A.F. Perez, The Electromagnetic Mass Differences of Pions and Kaons, *Phys. Rev. D* 55, 7075 (1997). J.F. Donoghue, Light quark masses and chiral symmetry, *Annu. Rev. Nucl. Part. Sci.* 39, 1 (1989).
- [47] S. Basak *et al*, Lattice computation of the electromagnetic contributions to kaon and pion masses, *Phys. Rev. D* 99, 034503 (2019). D. Giusti, *et al*, Leading isospin-breaking corrections to pion, kaon and charmed-meson masses with Twisted-Mass fermions *Phys. Rev. D* 95, 114505 (2017).
- [48] E. Witten, *Nucl. Phys. B* 156, 269 (1979); G. Veneziano, *Nucl. Phys. B* 159, 213 (1979). G. 't Hooft, *Phys. Rev. D* 14, 3432 (1976); Erratum: *ibid D* 18, 2199 (1978).
- [49] C. Rosenzweig, J. Schechter and C. G. Trahern, *Phys. Rev. D* 21, 3388 (1980). R. Alkofer and I. Zahed, *Mod. Phys. Lett. A* 4, 1737 (1989); R. Alkofer and I. Zahed, *Phys. Lett. B* 238, 149 (1990).
- [50] M. Jamin, Flavour-symmetry breaking of the quark condensate and chiral corrections to the Gell-Mann-Oakes-Renner relation *Phys. Lett. B* 538, 71 (2002).
- [51] S. Aoki, FLAG, Review of lattice results concerning low energy particle physics, *Eur. Phys. Journ. C* 80, 113 (2020).
- [52] C.T.H. Davies *et al*, Determination of the quark condensate from heavy-light current-current correlators in full lattice QCD, *Phys. Rev. D* 100, 034506 (2019).