

On the quaternion transformation and field equations in curved space-time

B. C. Chanyal

*Department of Physics, G. B. Pant University of Agriculture & Technology
Pantnagar-263145 (Uttarakhand) India. Email: bcchanyal@gmail.com*

Abstract

In this paper, we use four-dimensional quaternionic algebra to describing space-time geometry in curvature form. The transformation relations of a quaternionic variable are established with the help of basis transformations of quaternion algebra. We deduced the quaternionic covariant derivative that explains how the quaternion components vary with scalar and vector fields. The quaternionic metric tensor and geodesic equation are also discussed to describing the quaternionic line element in curved space-time. Moreover, we discussed an expression for the Riemannian Christoffel curvature tensor in terms of the quaternionic metric tensor. We have deduced the quaternionic Einstein's field-like equation which shows an equivalence between quaternionic matter and geometry.

Keywords: quaternion, curvature space-time, Riemannian Christoffel curvature tensor, Einstein field equation

PACS: 02.40.Hw, 02.40.Ky, 4.20.-q

Mathematics Subject Classification 2020: 12Hxx, 35Q76, 83C10

1 Introduction

The theory of the physical world has two fundamental components: matter and its interactions. The fundamental forces or interactions can be described as a field in which particles interact with one another. These fundamental interactions broke down into four fundamental forces of nature [1]: the gravitational force due to the attraction between massive bodies, the electromagnetic force due to the force between electrically charged particles, the weak force responsible for the radioactive decay of atoms, and the strong force that binding fundamental particles together. Gravitational force is the weakest force experiences as attractive. Because it is a very long-range force, the dominant force on the macroscopic scale is the reason for the formation, shape, and orbit of the astronomical bodies. This force is responsible for an object in space to move in a circular orbit. In Newtonian physics, the concept of absolute space and time

is important for the dynamics of an object but they are not relative to each other. Further, Albert Einstein developed the theory of relativity, therefore, the traditional Newtonian concept of absolute space and time has been replaced by the concept of space-time. Time is considered as the fourth dimension in the four-spaces theory of special and general relativity. The theory which explains the gravitation is named 'General Theory of Relativity' (GTR) [2]. In GTR, gravity does not act as a force but significance the space-time curvature where the source of the curved space-time is energy and matter. The Einstein field equations also elaborate that the mass and energy are responsible for the curvature of space-time, and the geometry of space-time is responsible for the motion of matter. Since, the description of space-time geometry should be non-Euclidean geometry, such that the space-time path becomes curved in presence of the gravitational field. In this case, the path of motion of particles in such a field or non-Euclidean space-time shows curvilinear instead of straight. Objects in nature have basic properties that are independent of the manual selection of any specific coordinate system. The basic assumption of physics is that the laws of nature are stated by equations, and should be valid for all frames of reference.

The theory of relativity required the 'covariant form' that is independent of the coordinate system. The covariant form remains the same in all co-ordinate systems whether the reference frame will be changed or not. The laws of physics must be expressed by an equation that remains covariant for all coordinate systems. The tensor has such properties that show covariant form in all coordinate systems. If all the equations, related to the general theory of relativity, are written in the tensor form, then it will transform easily from one reference frame to another. Therefore, in mathematics, the tensor calculus is the best tool to explain the theory of curvature as ordinary calculus. Besides, for a flat space-time, the Pythagoras theorem is valid perfectly but for a curved space-time, this theorem no longer holds and uses a 'metric' to measure the distance. Moreover, in GTR, the metric tensor behaves as the gravitational potential of Newtonian gravitation. The Riemannian Christoffel's curvature tensor captures the notion of parallel transport. It gives the idea about the connections of the parallel transported vector on a curve space-time. Further, Riemannian curvature is required for all the changes in the tangent vectors when we transport them around a curved space. The Riemann-Christoffel tensor is the only tensor that can be constructed from the second derivative of the metric tensor [3]. The contraction of the Riemann tensor gives Ricci curvature and scalar curvature. Ricci curvature explains the concept in which mass converges and diverges in time corresponding to the part of the curvature of space-time. Scalar curvature gives the single real number which represents the quantity of Riemannian manifold volume of geodesic ball differs from the normal ball in Euclidean space. The equation of motion of free particles is recognized by geodesic equation followed the analogous to Newton's equation of motion which clarifies the equation for the acceleration of particles. The source of the curvature of space-time is generalized by the energy and momentum of the field expressed as an energy-momentum tensor.

The study of algebraic structure in curvature space-time has an important role to describe the natural world. In mathematical algebraic structure, there are four types of division algebra [4], i.e., the real algebra, complex algebra, quaternion algebra (Hamilton algebra)[5], and octonion algebra (Cayley algebra)[6][7][8]. Real numbers are the numbers, which are used normally without any imaginary number. The numbers written as the mixture of a real number and the imaginary number are renowned by complex numbers. Quaternions are the extension of a complex number with non-commutative property, used to label the rotations in three-dimension. Cayley algebra is the extension of the quaternions with non-commutative and non-associative as well. There are many applications of quaternionic algebra in various branches of theoretical and computational physics [9][10][11][12][13]. In electrodynamics, Maxwell's equations in the presence of magnetic monopoles [14] and the classical wave equations of motion [15] have been constructed in terms of quaternionic algebra. The quaternionic form of quantum electrodynamics has been discussed [16] [17]. Chanyal [18][19] proposed the quaternionic covariant theory of four-dimensional particle dyons in the form of relativistic quantum mechanics and also focused on the quantized Dirac-Maxwell equations for dyons. Recently, in magneto-hydrodynamics, the quaternionic dual fields equations for dyonic cold plasma have been analyzed [20]. Further, it has been developed Dirac-Maxwell, Bernoulli, and Navier Stokes like equations for dyonic fluid-plasma in the generalized quaternionic field [21]. Currently, a new approach to Dirac's relativistic field equation for rotating free particle has been investigated in quaternionic form [22]. Beyond the quaternionic algebra, many authors [23][24][25][26][27][28][29][30][31][32][33][34][35] have studied the role of higher dimensional hypercomplex division algebras in various fields of modern physics. In GTR, Edmonds [36] discussed the quaternion wave equation in curved space-time by using the curvilinear coordinate system in relativistic quantum theory. The quaternionic form of curvature quantum theory fills the gap between quantum and gravity theory. In the same way, Weng [37] studied the electromagnetic and gravitational field equations in complex curved space with the help of quaternions and octonions. Keeping in view the Riemannian space and its connection with quaternions, we discussed the Einstein field-like equation in quaternionic curvilinear form. Starting with quaternionic basis transformation from one frame to another, we define the transformation of scalar and vector field components of a quaternion curvature variable. From the quaternionic covariant derivative, we expressed the Christoffel symbol, quaternionic metric tensor, and quaternionic geodesic equation. We also formulated the quaternionic Riemannian tensor that keeps track of how much scalar and vector components of quaternion changed if we propagate parallel along with a small parallelogram. Interestingly, if quaternionic Riemannian Christoffel curvature is zero then the quaternionic curved space-time is converted into flat space-time. Further, the quaternionic Ricci tensor is an important contraction of the quaternionic Riemannian Christoffel tensor which explains the changes in four space-time when an object parallels transport along a geodesic. We have discussed the quaternionic form of the Einstein field-like-equation in a compact form which shows

that the quaternionic value of matter-energy is equivalent to the quaternionic geometry.

2 Preliminaries

A tensor is the generalized form of a vector, defined by simply the arrangement of numbers, or functions that transform according to certain rules under a change of coordinates [38] [39]. According to Einstein summation convention, if any same index appears twice in a term then that index stands for the sum of all terms at the complete range of values. For example, we can write the two forms of a vector \mathbf{u} as

$$\mathbf{u} = \sum_i u^i e_i, \quad \mathbf{u} = \sum_j u_j e^j \quad \forall (i, j = 1, 2, 3), \quad (2.1)$$

where i and j are the repeated indices or summation indices while u^i and u_j are the vector components in contravariant and covariant form, respectively. The scalar product of two vectors can be represented in terms of indices with their components, i.e.,

$$\mathbf{a} \cdot \mathbf{b} = \left(\sum_i a_i e_i \right) \cdot \left(\sum_j b_j e_j \right) = \sum_{ij} a_i b_j (e_i \cdot e_j) = \sum_{ij} a_i b_j \delta_{ij} = \sum_i a_i b_i, \quad (2.2)$$

where δ_{ij} is the Kronecker delta symbol defined as $\delta_{ij} = 1$, for $(i = j)$ and $\delta_{ij} = 0$, for $(i \neq j)$. Also, the vector product becomes

$$[\mathbf{a} \times \mathbf{b}]_i = \sum_{jk} \epsilon_{ijk} a_j b_k. \quad (2.3)$$

Here ϵ_{ijk} is the Levi-Civita symbol with three indices having value $\epsilon_{ijk} = +1$ for cyclic permutation, $\epsilon_{ijk} = -1$ for non-cyclic permutation, and $\epsilon_{ijk} = 0$ for any two repeated indices. The tensors recognized by their order viz. vectors are first-order tensors, dyadics are second-order tensors, triadics are third-order tensors and tetradics are fourth-order tensors. By adding any two tensors of the same rank it gives the tensor of the same order, i.e.,

$$\mathbf{A}_{st}^r = \mathbf{B}_{st}^r + \mathbf{C}_{st}^r, \quad (2.4)$$

where \mathbf{A}_{st}^r , \mathbf{B}_{st}^r and \mathbf{C}_{st}^r are the tensors of same rank, i.e. three-rank tensor. The product of two tensors is given by

$$\mathbf{B}_s^r \mathbf{C}_{uv}^t = \mathcal{A}_{suv}^{rt}. \quad (2.5)$$

In the above representation, \mathbf{B}_s^r is tensor of rank two having indices r and s ; \mathbf{C}_{uv}^t is a tensor of rank three having indices t, u and v , while \mathcal{A}_{suv}^{rt} is a tensor

of rank five. If a magnitude or scalar quantity ϕ transform from one reference frame to another, it remains invariant, such that [40]

$$\phi' = \phi. \quad (2.6)$$

This invariant quantity may also be known as contravariant tensor of rank zero or covariant tensor of rank zero. On the other hand, a vector quantity can be transformed as

$$v'^r \mapsto v^s, \implies v'^r = \frac{\partial x'^r}{\partial x^s} v^s. \quad (2.7)$$

Equation (2.7) signifies that the components of vector v^s transformed to components of vector v'^r when the coordinate x^s transformed to x'^r . We can also define the contravariant tensor components of rank two as

$$A'^ru \mapsto A^{st}, \implies A'^ru = \frac{\partial x'^r}{\partial x^s} \frac{\partial x'^u}{\partial x^t} A^{st}. \quad (2.8)$$

Similarly, we can transform the tensors for higher rank.

3 The quaternionic algebra

In mathematics, a complex number over the real algebra \mathbb{R} having the imaginary number i , is defined by the algebraic extension of a normal real number. Any complex number $Z \in \mathbb{C}$ can be expressed in the form of its basis $(1, i)$ as

$$Z = \xi_1 + i\xi_2, \quad \forall (\xi_1, \xi_2) \in \mathbb{R}^2 \text{ and } Z \in \mathbb{C}. \quad (3.1)$$

If $Re(Z) = 0$ then the complex number is called purely imaginary. In the same way, the quaternion is a number system that extends the complex number in a form of algebra having some properties over addition and multiplication. W. R. Hamilton [5], extended the complex number in terms of four-dimensional norm-division algebra (quaternion) over real algebra \mathbb{R} . The quaternions has four unit elements (e_0, e_1, e_2, e_3) called basis elements, in which e_0 is the scalar unit and e_1, e_2, e_3 are the imaginary units. A quaternion $\mathbb{H} \in \mathbb{Q}$ can be written as

$$\mathbb{H} = e_0w + e_1x + e_2y + e_3z = e_0w + \sum_{j=1}^3 e_j r_j, \quad (3.2)$$

where w, x, y, z are the real numbers. The quaternion may also be composed of the scalar and vector parts as

$$\mathbb{H} = S_{\mathbb{H}} + V_{\mathbb{H}}. \quad (3.3)$$

Here e_0w is the scalar part of quaternion denoted by $S_{\mathbb{H}}$ and $(e_1x + e_2y + e_3z)$ is the vector part of quaternion denoted by $V_{\mathbb{H}}$. If scalar part is zero in equation (3.3), then

$$\mathbb{H} \mapsto V_{\mathbb{H}} = e_1x + e_2y + e_3z, \quad (3.4)$$

is known as right quaternion or pure quaternion. Although any quaternion can be seen as a vector in a four-dimensional vector space, it is usually referred to as the pure quaternion as a vector. Conditionally, if the vector part is zero then

$$\mathbb{H} \rightarrow S_{\mathbb{H}} = e_0 w, \quad (3.5)$$

is known as scalar quaternion. The addition of two quaternions will be

$$\begin{aligned} \mathbb{A} + \mathbb{B} &= (e_0 A_0 + e_1 A_1 + e_2 A_2 + e_3 A_3) + (e_0 B_0 + e_1 B_1 + e_2 B_2 + e_3 B_3) \\ &= e_0 (A_0 + B_0) + e_1 (A_1 + B_1) + e_2 (A_2 + B_2) + e_3 (A_3 + B_3) \\ &= e_0 (H_0) + e_j (H_j) \quad \forall (j = 1, 2, 3), \end{aligned} \quad (3.6)$$

where $H_0 \sim (A_0 + B_0)$ is the scalar part of quaternion while $H_j \sim (A_j + B_j)$ is the vector part of quaternion. Therefore, the equation (3.6) shows the closure property concerning quaternionic addition. The quaternionic algebra also satisfies the associative and commutative properties of addition. Further, the multiplication properties of two quaternions can be expressed by

$$\begin{aligned} \mathbb{A} \circ \mathbb{B} &= (e_0 A_0 + e_1 A_1 + e_2 A_2 + e_3 A_3) \circ (e_0 B_0 + e_1 B_1 + e_2 B_2 + e_3 B_3) \\ &= e_0 P_0 + e_1 P_1 + e_2 P_2 + e_3 P_3 = \mathbb{P} \in \mathbb{Q}, \end{aligned} \quad (3.7)$$

where ' \circ ' is a symbol used for the quaternionic multiplication and the components of \mathbb{P} given in equation (3.7) are written by

$$\begin{aligned} P_0 &:= (A_0 B_0 - A_1 B_1 - A_2 B_2 - A_3 B_3) && \text{(coefficient of } e_0) \\ P_1 &:= (A_0 B_1 + A_1 B_0 + A_2 B_3 - A_3 B_2) && \text{(coefficient of } e_1) \\ P_2 &:= (A_0 B_2 + A_2 B_0 + A_3 B_1 - A_1 B_3) && \text{(coefficient of } e_2) \\ P_3 &:= (A_0 B_3 + A_3 B_0 + A_1 B_2 - A_2 B_1) && \text{(coefficient of } e_3). \end{aligned} \quad (3.8)$$

Here the quaternionic multiplication follows the given rules for basis (e_0, e_1, e_2, e_3) , i.e.,

$$\begin{aligned} e_0^2 &= 1, \quad e_i^2 = -1, \quad e_0 e_i = e_i e_0 = e_i, \\ e_i e_j &= -\delta_{ij} e_0 + \epsilon_{ijk} e_k, \quad \forall (i, j, k = 1, 2, 3). \end{aligned} \quad (3.9)$$

All indices in Levi-Civita symbol ϵ_{ijk} are antisymmetric and satisfy $e_i \times e_j = \sum_k \epsilon_{ijk} e_k$. Thus, equation (3.7) can be written in compact form in terms of ordinary dot and cross product as,

$$\mathbb{A} \circ \mathbb{B} = e_0 (A_0 B_0 - \vec{A} \cdot \vec{B}) + e_j \left[A_0 \vec{B} + B_0 \vec{A} + (\vec{A} \times \vec{B})_j \right]. \quad (3.10)$$

In order to check the non-commutative property, we may write

$$\begin{aligned} \mathbb{B} \circ \mathbb{A} &= (e_0 B_0 + e_1 B_1 + e_2 B_2 + e_3 B_3) \circ (e_0 A_0 + e_1 A_1 + e_2 A_2 + e_3 A_3) \\ &= e_0 (B_0 A_0 - B_1 A_1 - B_2 A_2 - B_3 A_3) + e_1 (B_0 A_1 + B_1 A_0 + B_2 A_3 - B_3 A_2) \\ &\quad + e_2 (B_0 A_2 + B_2 A_0 + B_3 A_1 - B_1 A_3) + e_3 (B_0 A_3 + B_3 A_0 + B_1 A_2 - B_2 A_1) \\ &= e_0 (B_0 A_0 - \vec{B} \cdot \vec{A}) + e_j \left[B_0 \vec{A} + A_0 \vec{B} + (\vec{B} \times \vec{A})_j \right]. \\ &\neq \mathbb{A} \circ \mathbb{B}. \end{aligned} \quad (3.11)$$

From equation (3.11) it is clear that the product of two quaternions is non-commutative because the product of two vectors is always non-commutative, so that $\vec{A} \times \vec{B} = -\vec{B} \times \vec{A}$. On the other hand, the multiplication of quaternions satisfy the associative property i.e., $(\mathbb{A} \circ \mathbb{B}) \circ \mathbb{C} = \mathbb{A} \circ (\mathbb{B} \circ \mathbb{C})$. Further, the quaternionic conjugate of equation (3.2) can be expressed as

$$\mathbb{H}^* = e_0 w - (e_1 x + e_2 y + e_3 z), \quad (3.12)$$

and the product of two quaternions is given by [41]

$$\mathbb{A} \cdot \mathbb{B} = -\frac{1}{2}(\mathbb{A} \circ \mathbb{B}^* + \mathbb{B} \circ \mathbb{A}^*) = -\frac{1}{2}(\mathbb{A}^* \circ \mathbb{B} + \mathbb{B}^* \circ \mathbb{A}). \quad (3.13)$$

The norm of a quaternion \mathbb{H} denoted by $|\mathbb{H}|$ can be represented as

$$|\mathbb{H}| = \sqrt{\mathbb{H} \circ \mathbb{H}^*} = \sqrt{w^2 + x^2 + y^2 + z^2}. \quad (3.14)$$

Now, the inverse of a quaternion can be defined by

$$\mathbb{H}^{-1} = \frac{\mathbb{H}^*}{|\mathbb{H}|^2} \equiv \frac{e_0 w - e_1 x - e_2 y - e_3 z}{w^2 + x^2 + y^2 + z^2}. \quad (3.15)$$

Moreover, the quotient of two vectors is also known as quaternion which can be represented by [42]

$$\mathbb{H} = \frac{\alpha}{\beta}, \quad (3.16)$$

where α and β are the two vectors. The another approach on a ‘‘Tensor of a quaternion’’ ($\mathbf{T}_{\mathbb{H}}$) is given by Hamilton [42], which can be written by

$$(\mathbf{T}_{\mathbb{H}})^2 = \mathbb{H} \circ \mathbb{H}^* = w^2 + x^2 + y^2 + z^2, \quad (3.17)$$

so that $T_{\mathbb{H}} = \sqrt{(w^2 + x^2 + y^2 + z^2)}$. The tensor of a quaternion is similar to the norm of a quaternion, it will be always positive or real number [43]. If the scalar part is zero then $\mathbf{T}(\mathbf{V}_{\mathbb{H}}) = \sqrt{(x^2 + y^2 + z^2)}$ is known as the tensor of a vector quaternion and if vector part is zero then $\mathbf{T}(\mathbf{S}_{\mathbb{H}}) = w$ is known as the tensor of a scalar quaternion. Further, the ‘versor’ of a quaternion which indicates direction, is a unit quaternion known as a normalized quaternion ($\mathbf{U}_{\mathbb{H}}$), so that

$$\mathbf{U}_{\mathbb{H}} = \frac{\mathbb{H}}{|\mathbb{H}|} \equiv \frac{e_0 w + e_1 x + e_2 y + e_3 z}{\sqrt{(w^2 + x^2 + y^2 + z^2)}}. \quad (3.18)$$

The scalar and vector parts of a quaternionic versor can be expressed by $\mathbf{S}(\mathbf{U}_{\mathbb{H}}) = w(w^2 + x^2 + y^2 + z^2)^{-1/2}$ and $\mathbf{V}(\mathbf{U}_{\mathbb{H}}) = (e_1 x + e_2 y + e_3 z)(w^2 + x^2 + y^2 + z^2)^{-1/2}$. A quaternion can also be written in terms of tensor of a quaternion and versor of a quaternion as [42]

$$\mathbb{H} = \mathbf{T}_{\mathbb{H}} \mathbf{U}_{\mathbb{H}} = e_0 w + e_1 x + e_2 y + e_3 z. \quad (3.19)$$

4 Transformation of quaternionic basis elements

If a system of coordinates rotates from its original frame of reference then there will be a change in unit vectors to original unit vectors. This change of unit vectors represents the transformation equations. Here, we introduce the transformation equations for the 2-dimensional system in which the equations are used for the transformation of coordinates from one plane to another plane, i.e.,

$$x' = x \cos \phi + y \sin \phi, \quad y' = -x \sin \phi + y \cos \phi, \quad (4.1)$$

which gives

$$\frac{\partial x'}{\partial x} = \cos \phi, \quad \frac{\partial x'}{\partial y} = \sin \phi, \quad \frac{\partial y'}{\partial x} = -\sin \phi, \quad \frac{\partial y'}{\partial y} = \cos \phi. \quad (4.2)$$

Thus, the transformation of basis elements can be written in the matrix form

$$\begin{pmatrix} e'_1 \\ e'_2 \end{pmatrix} \mapsto \begin{pmatrix} \cos \phi & \sin \phi \\ -\sin \phi & \cos \phi \end{pmatrix} \begin{pmatrix} e_1 \\ e_2 \end{pmatrix}, \quad (4.3)$$

which leads to following linear transformation equations

$$\begin{aligned} e'_1 &\mapsto \frac{\partial x'}{\partial x} e_1 + \frac{\partial x'}{\partial y} e_2, \\ e'_2 &\mapsto \frac{\partial y'}{\partial x} e_1 + \frac{\partial y'}{\partial y} e_2, \end{aligned} \quad (4.4)$$

where e_1 and e_2 are the unit vectors corresponding to x and y - axis in XY -plane, while e'_1 and e'_2 are the unit vectors corresponding to x' and y' - axis in $X'Y'$ - plane, respectively. ϕ is the angle between these two planes. Like 2-D transformation, we may extend the transformation relations of basis elements for 3-D system. Thus, the 3-D transformation matrix (D) can be written by

$$D = \begin{pmatrix} \cos \psi \cos \phi - \sin \psi \sin \phi \cos \theta & -\cos \psi \sin \phi - \sin \psi \cos \theta \cos \phi & \sin \psi \sin \theta \\ \sin \psi \cos \phi + \cos \psi \sin \phi \cos \theta & -\sin \psi \sin \phi + \cos \psi \cos \theta \cos \phi & -\cos \psi \sin \theta \\ \sin \theta \sin \phi & \sin \theta \cos \phi & \cos \theta \end{pmatrix}, \quad (4.5)$$

where ψ, θ, ϕ are the three independent parameters known as *Euler angles* which represents the rotation of axis in new frame of reference with respect to original axis. Let (e_1, e_2, e_3) are pure quaternionic unit vectors along x, y, z - axis in S -frame while (e'_1, e'_2, e'_3) are unit vectors for x', y', z' - axis in S' - frame, then the linear transformation relations for pure quaternionic unit vectors become,

$$\begin{aligned} e'_1 &\mapsto \frac{\partial x'}{\partial x} e_1 + \frac{\partial x'}{\partial y} e_2 + \frac{\partial x'}{\partial z} e_3, \\ e'_2 &\mapsto \frac{\partial y'}{\partial x} e_1 + \frac{\partial y'}{\partial y} e_2 + \frac{\partial y'}{\partial z} e_3, \\ e'_3 &\mapsto \frac{\partial z'}{\partial x} e_1 + \frac{\partial z'}{\partial y} e_2 + \frac{\partial z'}{\partial z} e_3, \end{aligned} \quad (4.6)$$

where the derivatives become

$$\begin{aligned}
\frac{\partial x'}{\partial x} &= \cos \psi \cos \phi - \sin \psi \sin \phi \cos \theta, & \frac{\partial x'}{\partial y} &= -\cos \psi \sin \phi - \sin \psi \cos \theta \cos \phi, \\
\frac{\partial x'}{\partial z} &= \sin \psi \sin \theta, & \frac{\partial y'}{\partial x} &= \sin \psi \cos \phi + \cos \psi \sin \phi \cos \theta, \\
\frac{\partial y'}{\partial y} &= -\sin \psi \sin \phi + \cos \psi \cos \theta \cos \phi, & \frac{\partial y'}{\partial z} &= -\cos \psi \sin \theta, \\
\frac{\partial z'}{\partial x} &= \sin \theta \sin \phi, & \frac{\partial z'}{\partial y} &= \sin \theta \cos \phi, & \frac{\partial z'}{\partial z} &= \cos \theta.
\end{aligned}$$

Interestingly, these transformation equations satisfy the properties of vector algebra. The cross product of two vectors is always non-commutative for pure quaternionic space. Moreover, the properties of pure quaternionic unit elements (e_1, e_2, e_3) corresponding to vector algebra basis can be represented as

$$\begin{aligned}
e'_1 \circ e'_1 &= \left(\frac{\partial x'}{\partial x} e_1 + \frac{\partial x'}{\partial y} e_2 + \frac{\partial x'}{\partial z} e_3 \right) \circ \left(\frac{\partial x'}{\partial x} e_1 + \frac{\partial x'}{\partial y} e_2 + \frac{\partial x'}{\partial z} e_3 \right) \\
&= - \left(\frac{\partial x'}{\partial x} \right)^2 - \left(\frac{\partial x'}{\partial y} \right)^2 - \left(\frac{\partial x'}{\partial z} \right)^2 \\
&= - (\cos \psi \cos \phi - \sin \psi \sin \phi \cos \theta)^2 - (-\cos \psi \sin \phi - \sin \psi \cos \theta \cos \phi)^2 \\
&\quad - (\sin \psi \sin \theta)^2 = -1.
\end{aligned} \tag{4.7}$$

and

$$e'_2 \circ e'_2 = -1, \quad e'_3 \circ e'_3 = -1. \tag{4.8}$$

On the other hand

$$\begin{aligned}
e'_1 \circ e'_2 &= \left(\frac{\partial x'}{\partial x} e_1 + \frac{\partial x'}{\partial y} e_2 + \frac{\partial x'}{\partial z} e_3 \right) \circ \left(\frac{\partial y'}{\partial x} e_1 + \frac{\partial y'}{\partial y} e_2 + \frac{\partial y'}{\partial z} e_3 \right) \\
&= - \frac{\partial x'}{\partial x} \frac{\partial y'}{\partial x} - \frac{\partial x'}{\partial y} \frac{\partial y'}{\partial y} - \frac{\partial x'}{\partial z} \frac{\partial y'}{\partial z} - \left(\frac{\partial x'}{\partial z} \frac{\partial y'}{\partial y} - \frac{\partial x'}{\partial y} \frac{\partial y'}{\partial z} \right) e_1 \\
&\quad - \left(\frac{\partial x'}{\partial x} \frac{\partial y'}{\partial z} - \frac{\partial x'}{\partial z} \frac{\partial y'}{\partial x} \right) e_2 - \left(\frac{\partial x'}{\partial y} \frac{\partial y'}{\partial x} - \frac{\partial x'}{\partial x} \frac{\partial y'}{\partial y} \right) e_3
\end{aligned}$$

$$\begin{aligned}
&= -(\cos \psi \cos \phi - \sin \psi \sin \phi \cos \theta)(\sin \psi \cos \phi + \cos \psi \sin \phi \cos \theta) \\
&- (-\cos \psi \sin \phi - \sin \psi \cos \theta \cos \phi)(-\sin \psi \sin \phi + \cos \psi \cos \theta \cos \phi) \\
&- (\sin \psi \sin \theta)(-\cos \psi \sin \theta) - e_1[(\sin \psi \sin \theta)(-\sin \psi \sin \phi + \cos \psi \cos \theta \cos \phi) \\
&+ (-\cos \psi \sin \phi - \sin \psi \cos \theta \cos \phi)(-\cos \psi \sin \theta)] \\
&- e_2[(\cos \psi \cos \phi - \sin \psi \sin \phi \cos \theta)(-\cos \psi \sin \theta) \\
&+ (\sin \psi \sin \theta)(\sin \psi \cos \phi + \cos \psi \sin \phi \cos \theta)] \\
&- e_3[(-\cos \psi \sin \phi - \sin \psi \cos \theta \cos \phi)(\sin \psi \cos \phi + \cos \psi \sin \phi \cos \theta) \\
&+ (\cos \psi \cos \phi - \sin \psi \sin \phi \cos \theta)(-\sin \psi \sin \phi + \cos \psi \cos \theta \cos \phi)] \\
&= e'_3, \tag{4.9}
\end{aligned}$$

and

$$\begin{aligned}
e'_2 \circ e'_1 &= \left(\frac{\partial y'}{\partial x} e_1 + \frac{\partial y'}{\partial y} e_2 + \frac{\partial y'}{\partial z} e_3 \right) \circ \left(\frac{\partial x'}{\partial x} e_1 + \frac{\partial x'}{\partial y} e_2 + \frac{\partial x'}{\partial z} e_3 \right) \\
&= -\frac{\partial y'}{\partial x} \frac{\partial x'}{\partial x} - \frac{\partial y'}{\partial y} \frac{\partial x'}{\partial y} - \frac{\partial y'}{\partial z} \frac{\partial x'}{\partial z} + \left(\frac{\partial y'}{\partial y} \frac{\partial x'}{\partial z} - \frac{\partial y'}{\partial z} \frac{\partial x'}{\partial y} \right) e_1 \\
&+ \left(\frac{\partial y'}{\partial z} \frac{\partial x'}{\partial x} - \frac{\partial y'}{\partial x} \frac{\partial x'}{\partial z} \right) e_2 + \left(\frac{\partial y'}{\partial x} \frac{\partial x'}{\partial y} - \frac{\partial y'}{\partial y} \frac{\partial x'}{\partial x} \right) e_3 \\
&= -(\sin \psi \cos \phi + \cos \psi \sin \phi \cos \theta)(\cos \psi \cos \phi - \sin \psi \sin \phi \cos \theta) \\
&- (-\sin \psi \sin \phi + \cos \psi \cos \theta \cos \phi)(-\cos \psi \sin \phi - \sin \psi \cos \theta \cos \phi) \\
&- (-\cos \psi \sin \theta)(\sin \psi \sin \theta) + e_1[(\sin \psi \sin \theta)(-\sin \psi \sin \phi + \cos \psi \cos \theta \cos \phi) \\
&- (-\cos \psi \sin \phi - \sin \psi \cos \theta \cos \phi)(-\cos \psi \sin \theta)] \\
&+ e_2[(\cos \psi \cos \phi - \sin \psi \sin \phi \cos \theta)(-\cos \psi \sin \theta) \\
&- (\sin \psi \sin \theta)(\sin \psi \cos \phi + \cos \psi \sin \phi \cos \theta)] \\
&+ e_3[(-\cos \psi \sin \phi - \sin \psi \cos \theta \cos \phi)(\sin \psi \cos \phi + \cos \psi \sin \phi \cos \theta) \\
&- (\cos \psi \cos \phi - \sin \psi \sin \phi \cos \theta)(-\sin \psi \sin \phi + \cos \psi \cos \theta \cos \phi)] \\
&= -e'_3. \tag{4.10}
\end{aligned}$$

Similarly, we obtain

$$\begin{aligned}
e'_2 \circ e'_3 &= e'_1, & e'_3 \circ e'_2 &= -e'_1, \\
e'_3 \circ e'_1 &= e'_2, & e'_1 \circ e'_3 &= -e'_2.
\end{aligned} \tag{4.11}$$

Now, we easily can extend the pure-quaternion to quaternion by adding the scalar unit element e_0 . It should be noticed that the scalar unit element becomes invariant under transformation given in equation (2.6) but it plays an important role with unit elements e_1, e_2, e_3 in quaternionic transformation. Myszkowski [44] has given the idea of 4-D transformation similar to the 3-D transformation. Thus, we assume the transformation equations for quaternionic basis elements

as

$$\begin{aligned}
e'_0 &\mapsto e_0, \\
e'_1 &\mapsto \frac{\partial x'}{\partial t} e_0 + \frac{\partial x'}{\partial x} e_1 + \frac{\partial x'}{\partial y} e_2 + \frac{\partial x'}{\partial z} e_3, \\
e'_2 &\mapsto \frac{\partial y'}{\partial t} e_0 + \frac{\partial y'}{\partial x} e_1 + \frac{\partial y'}{\partial y} e_2 + \frac{\partial y'}{\partial z} e_3, \\
e'_3 &\mapsto \frac{\partial z'}{\partial t} e_0 + \frac{\partial z'}{\partial x} e_1 + \frac{\partial z'}{\partial y} e_2 + \frac{\partial z'}{\partial z} e_3.
\end{aligned} \tag{4.12}$$

Here, we focused on space-time structure for quaternionic transformation, the quaternionic properties will also be satisfied as similar to the 3-D rotation. Now, in the next section, we shall interpret these relations for Riemannian geometry.

5 Quaternionic approach on Riemannian geometry

5.1 Quaternion transformation

In Riemannian geometry, the motion of object takes place in curvature space-time in which 4-dimensional frame of reference rotate along the curve. The structure of quaternionic space-time coordinates can be written as $P^\mu \equiv (P^0, P^j) \simeq (P^\xi, P^1, P^2, P^3)$. The time coordinate may refer corresponding to e_0 i.e. $t \mapsto P^\xi$, and the spatial coordinates may refer corresponding to e_j i.e. $(x, y, z) \mapsto (P^1, P^2, P^3)$ where $j = 1, 2, 3$. Now, the quaternionic elements may transform as

$$e'_0 = e_0, \quad e'_i = \frac{\partial P'^i}{\partial P^\mu} e_\mu, \quad \forall (i = 1, 2, 3). \tag{5.1}$$

The transformation of scalar and vector field components of a quaternion variable (\mathbb{H}) from S to S' -frame can be expressed as

$$H^\xi = H'^\xi + \frac{\partial P^\xi}{\partial P'^i} H'^i \quad (\text{coefficient of } e_0), \tag{5.2}$$

$$H^j = \frac{\partial P^j}{\partial P'^i} H'^i \quad (\text{coefficient of } e_j). \tag{5.3}$$

In the above transformation relation (5.2), we assume that the effect of H'^ξ on H^ξ is linear because it shows the simple transformation on flat space-time. So, we can neglect the component H'^ξ to get the quaternionic transformation (Q-transformation) in curved space-time. Therefore, we have

$$\mathbb{H}' \mapsto \mathbb{H} = \left(\frac{\partial P^\xi}{\partial P'^i} H'^i, \frac{\partial P^j}{\partial P'^i} H'^i \right). \tag{5.4}$$

Equation (5.4) represents the Q-transformation in contravariant form, we can also write the above transformation relations in quaternionic covariant form as

$$H_\xi = \frac{\partial P'^i}{\partial P^\xi} H'_i, \quad H_j = \frac{\partial P'^i}{\partial P^j} H'_i. \quad (5.5)$$

Now, we can write the transformation of quaternionic differential operator $\square \simeq \{\partial_\phi, \partial_j\}$ as

$$\partial_\phi = \frac{\partial P^\phi}{\partial P'^k} \partial'_k, \quad \partial_j = \frac{\partial P^j}{\partial P'^k} \partial'_k, \quad \forall (j, k = 1, 2, 3). \quad (5.6)$$

Since the components of a quaternion can also represent in form of coordinates, then we can write the transformation of quaternionic coordinates [36],

$$dP^\xi = \frac{\partial P^\xi}{\partial P'^i} dP'^i, \quad dP^j = \frac{\partial P^j}{\partial P'^i} dP'^i, \quad \forall (i, j = 1, 2, 3). \quad (5.7)$$

5.2 Quaternionic covariant derivative

In the quaternionic form of curvature space-time, the scalar field derivative of equation (5.5) can be written as

$$H_{;\phi}^\xi = H_{,\phi}^\xi + \left\{ \begin{array}{c} \xi \\ \phi\phi \end{array} \right\} H^\phi, \quad (5.8)$$

where the Christoffel symbol is denoted by $\left\{ \begin{array}{c} \xi \\ \phi\phi \end{array} \right\} = \frac{\partial}{\partial P^\phi} \left(\frac{\partial P^\xi}{\partial P^\phi} \right)$. Similarly, we have

$$H_{\xi;\phi} = H_{\xi,\phi} - \left\{ \begin{array}{c} \phi \\ \phi\xi \end{array} \right\} H_\phi. \quad (5.9)$$

Here the role of Christoffel symbol is very important for Riemannian space-time. If $\left\{ \begin{array}{c} \xi \\ \phi\phi \end{array} \right\} \rightarrow 0$, then the quaternionic curvature covariant derivative transforms into the quaternionic usual partial derivative which indicates the linear transformation. Correspondingly, applying quaternionic derivative ∂_ϕ on H^j , we obtain

$$H_{;\phi}^j = H_{,\phi}^j + \left\{ \begin{array}{c} j \\ \phi\phi \end{array} \right\} H^\phi.$$

As such, we also can operate the quaternionic derivative ∂_k ($k = 1, 2, 3$) on given equation (5.5), so that

$$H_{;k}^\xi = H_{,k}^\xi + \left\{ \begin{array}{c} \xi \\ km \end{array} \right\} H^m, \quad (5.10)$$

$$H_{;k}^j = H_{,k}^j + \left\{ \begin{array}{c} j \\ km \end{array} \right\} H^m, \quad (5.11)$$

where $j, k, m = 1, 2, 3$. Equations (5.10) and (5.11) are represented scalar to vector and vector to vector field transformations, respectively, for generalized quaternionic fields [42]. It should be remarked that through quaternionic derivative, the generalized covariant derivative gives enlargement to the rank of quaternionic tensors which is important to provide more information about the transformed coordinate. In more general the quaternionic tensor derivatives are

$$T_{\xi\xi;\phi} = T_{\xi\xi,\phi} - \left\{ \begin{matrix} \eta \\ \phi\xi \end{matrix} \right\} T_{\xi\eta} - \left\{ \begin{matrix} \eta \\ \phi\xi \end{matrix} \right\} T_{\eta\xi}, \quad (5.12)$$

$$T_{\xi\xi;k} = T_{\xi\xi,k} - \left\{ \begin{matrix} l \\ k\xi \end{matrix} \right\} T_{\xi l} - \left\{ \begin{matrix} l \\ k\xi \end{matrix} \right\} T_{l\xi}, \quad (5.13)$$

$$T_{\xi n;\phi} = T_{\xi n,\phi} - \left\{ \begin{matrix} \eta \\ \phi n \end{matrix} \right\} T_{\xi\eta} - \left\{ \begin{matrix} \eta \\ \phi\xi \end{matrix} \right\} T_{\eta n}, \quad (5.14)$$

$$T_{\xi n;k} = T_{\xi n,k} - \left\{ \begin{matrix} l \\ kn \end{matrix} \right\} T_{\xi l} - \left\{ \begin{matrix} l \\ kn \end{matrix} \right\} T_{ln}, \quad (5.15)$$

$$T_{j\xi;\phi} = T_{j\xi,\phi} - \left\{ \begin{matrix} \eta \\ \phi\xi \end{matrix} \right\} T_{j\eta} - \left\{ \begin{matrix} \eta \\ \phi j \end{matrix} \right\} T_{\eta\xi}, \quad (5.16)$$

$$T_{j\xi;k} = T_{j\xi,k} - \left\{ \begin{matrix} l \\ k\xi \end{matrix} \right\} T_{jl} - \left\{ \begin{matrix} l \\ kj \end{matrix} \right\} T_{l\xi}, \quad (5.17)$$

$$T_{jn;\phi} = T_{jn,\phi} - \left\{ \begin{matrix} \eta \\ \phi n \end{matrix} \right\} T_{j\eta} - \left\{ \begin{matrix} \eta \\ \phi j \end{matrix} \right\} T_{\eta n}, \quad (5.18)$$

$$T_{jn;k} = T_{jn,k} - \left\{ \begin{matrix} l \\ kn \end{matrix} \right\} T_{jl} - \left\{ \begin{matrix} l \\ kj \end{matrix} \right\} T_{ln}, \quad (5.19)$$

where the right-hand side of equations (5.12)-(5.19), the first term shows the tensorial transformation of derivative of quaternionic tensor and the last two terms show the Christoffel symbols which tell us how the geodesic path changes from point to point.

5.3 Quaternionic metric tensor

To expressing the quaternionic metric tensor in terms of the Christoffel symbol, we can use the covariant derivative of quaternionic tensor. Therefore, one can write

$$g_{\xi\xi;\phi} = g_{\xi\xi,\phi} - \left\{ \begin{matrix} \eta \\ \phi\xi \end{matrix} \right\} g_{\xi\eta} - \left\{ \begin{matrix} \eta \\ \phi\xi \end{matrix} \right\} g_{\eta\xi}. \quad (5.20)$$

where the line element becomes $ds = \sqrt{-g_{\mu\nu}(x) dx^\mu dx^\nu}$. Under the covariant differentiation, the quaternionic metric tensor is constant i.e.,

$$\begin{aligned} 0 &= g_{\xi\xi,\phi} - \left\{ \begin{matrix} \eta \\ \phi\xi \end{matrix} \right\} g_{\xi\eta} - \left\{ \begin{matrix} \eta \\ \phi\xi \end{matrix} \right\} g_{\eta\xi} \\ g_{\xi\xi,\phi} &= \left\{ \begin{matrix} \eta \\ \phi\xi \end{matrix} \right\} g_{\xi\eta} + \left\{ \begin{matrix} \eta \\ \phi\xi \end{matrix} \right\} g_{\eta\xi}. \end{aligned} \quad (5.21)$$

Here the metric tensor $g'_{im} = \frac{\partial P^\mu}{\partial P'^i} \frac{\partial P^\nu}{\partial P'^m} \Omega_{\mu\nu}$, where $\Omega_{\mu\nu}$ is the four-dimensional Minkowski metric define by

$$\Omega_{\mu\nu} = \begin{pmatrix} -1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix},$$

and the Christoffel symbol satisfy the following properties

$$\left\{ \begin{matrix} i \\ kl \end{matrix} \right\} = \left\{ \begin{matrix} i \\ lk \end{matrix} \right\}, \quad \left\{ \begin{matrix} m \\ il \end{matrix} \right\} g_{mk} = \left\{ \begin{matrix} k \\ il \end{matrix} \right\}. \quad (5.22)$$

Now, the value of Christoffel symbol corresponding to quaternionic scalar field can be written as,

$$\left\{ \begin{matrix} \eta \\ \xi\xi \end{matrix} \right\} = \frac{1}{2} g^{\eta\phi} (g_{\phi\xi,\xi} + g_{\xi\phi,\xi} - g_{\xi\xi,\phi}), \quad (5.23)$$

where $g^{\eta\phi}$ is the inverse matrix of $g_{\eta\phi}$. Correspondingly, in quaternionic pure vectorial field, the Christoffel symbol can be played by

$$\left\{ \begin{matrix} l \\ jn \end{matrix} \right\} = \frac{1}{2} g^{lk} (g_{kj,n} + g_{nk,j} - g_{jn,k}), \quad (5.24)$$

and the other diversified quaternionic scalar and vector fields show a mixed form of quaternionic Christoffel symbol in terms of the metric tensor, i.e.

$$\left\{ \begin{matrix} l \\ \xi\xi \end{matrix} \right\} = \frac{1}{2} g^{lk} (g_{k\xi,\xi} + g_{\xi k,\xi} - g_{\xi\xi,k}), \quad (5.25)$$

$$\left\{ \begin{matrix} \eta \\ \xi n \end{matrix} \right\} = \frac{1}{2} g^{\eta\phi} (g_{\phi\xi,n} + g_{n\phi,\xi} - g_{\xi n,\phi}), \quad (5.26)$$

$$\left\{ \begin{matrix} l \\ \xi n \end{matrix} \right\} = \frac{1}{2} g^{lk} (g_{k\xi,n} + g_{nk,\xi} - g_{\xi n,k}), \quad (5.27)$$

$$\left\{ \begin{matrix} \eta \\ jn \end{matrix} \right\} = \frac{1}{2} g^{\eta\phi} (g_{\phi j,n} + g_{n\phi,j} - g_{jn,\phi}), \quad (5.28)$$

where (ξ, η, ϕ) are used for generalized quaternionic scalar field variables while (i, j, k, l, m, n) are the generalized quaternionic vector field variables.

5.4 Quaternionic geodesic equation

In general theory of relativity, geodesic equation gives the replacement of linear space-time to curved space-time. In other words, a particle moving in a curvature space-time is always follow the path of geodesic. The geodesic path in curvature space-time structure can be described in terms of four-vector form

with the principle of least action, i.e. $dH^\mu = 0$, see ref.[3]. In the pure quaternionic scalar field,

$$\begin{aligned} H_{;\phi}^\xi &\equiv H_{,\phi}^\xi + \left\{ \begin{array}{c} \xi \\ \phi\phi \end{array} \right\} H^\phi = 0 \\ \Rightarrow dH^\xi + \left\{ \begin{array}{c} \xi \\ \phi\phi \end{array} \right\} dP^\phi H^\phi &= 0. \end{aligned} \quad (5.29)$$

Now, dividing equation (5.29) by ds (e.g. a scalar parameter of motion as proper time) and substituting $H^\xi = \frac{dP^\xi}{ds}$, $H^\phi = \frac{dP^\phi}{ds}$, we get

$$\frac{d^2 P^\xi}{ds^2} + \left\{ \begin{array}{c} \xi \\ \phi\phi \end{array} \right\} \frac{dP^\phi}{ds} \frac{dP^\phi}{ds} = 0. \quad (5.30)$$

Similarly, for quaternionic vector fields, we get

$$\frac{d^2 P^j}{ds^2} + \left\{ \begin{array}{c} j \\ km \end{array} \right\} \frac{dP^k}{ds} \frac{dP^m}{ds} = 0, \quad (5.31)$$

and for a mixed fields,

$$\begin{aligned} \frac{d^2 P^j}{ds^2} + \left\{ \begin{array}{c} j \\ \phi\phi \end{array} \right\} \frac{dP^\phi}{ds} \frac{dP^\phi}{ds} &= 0, \\ \frac{d^2 P^\xi}{ds^2} + \left\{ \begin{array}{c} \xi \\ km \end{array} \right\} \frac{dP^k}{ds} \frac{dP^m}{ds} &= 0. \end{aligned} \quad (5.32)$$

Equations (5.30)-(5.32) represent the non-linear equations called the quaternionic geodesic equations that arise due to the effect of Christoffel symbols. In quaternionic formulation, we also can emphasize that the space-time curvature path is not only followed by vector components but also followed by the scalar components and mixed components of a quaternion variable. If the Christoffel symbol is zero, then the quaternionic geodesic equations of motion lead to

$$\frac{d^2 P^\mu}{ds^2} = 0, \quad \forall (\mu = 0, 1, 2, 3). \quad (5.33)$$

This implies that the generalized quaternionic acceleration will be zero i.e. the uniform velocity of the particle moving in a straight line (or non-Riemannian space-time).

5.5 Quaternionic Riemannian Christoffel curvature tensor

The Riemannian Christoffel curvature tensor is a four-indices tensor, can be obtained by subtraction of two covariant derivatives of quaternionic tensors in which indices are interchanged, such that

$$T_{\xi\xi;\phi} - T_{\xi\phi;\xi} = \left(R_{\xi\xi\phi}^\eta \right) H_\eta, \quad (5.34)$$

where

$$R_{\xi\xi\phi}^{\eta} = \partial_{\xi} \left(\left\{ \begin{matrix} \eta \\ \phi\xi \end{matrix} \right\} \right) - \partial_{\phi} \left(\left\{ \begin{matrix} \eta \\ \xi\xi \end{matrix} \right\} \right) + \left\{ \begin{matrix} \varphi \\ \phi\xi \end{matrix} \right\} \left\{ \begin{matrix} \eta \\ \varphi\xi \end{matrix} \right\} - \left\{ \begin{matrix} \varphi \\ \xi\xi \end{matrix} \right\} \left\{ \begin{matrix} \eta \\ \varphi\phi \end{matrix} \right\} \quad (5.35)$$

is the Riemannian Christoffel curvature tensor for purely quaternionic scalar field. It is a four-indices tensor in curved space-time which describes the curvature of manifolds. Further, we can obtain the Riemannian Christoffel curvature tensor to other quaternionic components given in equation (5.13)-(5.19), i.e.,

$$T_{\xi\xi;k} - T_{\xi k;\xi} = (R_{\xi\xi k}^l) H_l, \quad (5.36)$$

$$T_{\xi n;\phi} - T_{\xi\phi;n} = (R_{\xi n\phi}^{\eta}) H_{\eta}, \quad (5.37)$$

$$T_{\xi n;k} - T_{\xi k;n} = (R_{\xi n k}^l) H_l, \quad (5.38)$$

$$T_{j\xi;\phi} - T_{j\phi;\xi} = (R_{j\xi\phi}^{\eta}) H_{\eta}, \quad (5.39)$$

$$T_{j\xi;k} - T_{jk;\xi} = (R_{j\xi k}^l) H_l, \quad (5.40)$$

$$T_{jn;\phi} - T_{j\phi;n} = (R_{jn\phi}^{\eta}) H_{\eta}, \quad (5.41)$$

$$T_{jn;k} - T_{jk;n} = (R_{jn k}^l) H_l, \quad (5.42)$$

where the Riemannian Christoffel curvature tensors lead to the following way as

$$R_{\xi\xi k}^l = \partial_{\xi} \left(\left\{ \begin{matrix} l \\ k\xi \end{matrix} \right\} \right) - \partial_k \left(\left\{ \begin{matrix} l \\ \xi\xi \end{matrix} \right\} \right) + \left\{ \begin{matrix} i \\ k\xi \end{matrix} \right\} \left\{ \begin{matrix} l \\ i\xi \end{matrix} \right\} - \left\{ \begin{matrix} i \\ \xi\xi \end{matrix} \right\} \left\{ \begin{matrix} l \\ ik \end{matrix} \right\}, \quad (5.43)$$

$$R_{\xi n\phi}^{\eta} = \partial_n \left(\left\{ \begin{matrix} \eta \\ \phi\xi \end{matrix} \right\} \right) - \partial_{\phi} \left(\left\{ \begin{matrix} \eta \\ \xi n \end{matrix} \right\} \right) + \left\{ \begin{matrix} \varphi \\ \phi\xi \end{matrix} \right\} \left\{ \begin{matrix} \eta \\ \varphi n \end{matrix} \right\} - \left\{ \begin{matrix} \varphi \\ n\xi \end{matrix} \right\} \left\{ \begin{matrix} \eta \\ \varphi\phi \end{matrix} \right\}, \quad (5.44)$$

$$R_{\xi n k}^l = \partial_n \left(\left\{ \begin{matrix} l \\ k\xi \end{matrix} \right\} \right) - \partial_k \left(\left\{ \begin{matrix} l \\ \xi n \end{matrix} \right\} \right) + \left\{ \begin{matrix} i \\ k\xi \end{matrix} \right\} \left\{ \begin{matrix} l \\ in \end{matrix} \right\} - \left\{ \begin{matrix} i \\ n\xi \end{matrix} \right\} \left\{ \begin{matrix} l \\ ik \end{matrix} \right\}, \quad (5.45)$$

$$R_{j\xi\phi}^{\eta} = \partial_{\xi} \left(\left\{ \begin{matrix} \eta \\ \phi j \end{matrix} \right\} \right) - \partial_{\phi} \left(\left\{ \begin{matrix} \eta \\ j\xi \end{matrix} \right\} \right) + \left\{ \begin{matrix} \varphi \\ \phi j \end{matrix} \right\} \left\{ \begin{matrix} \eta \\ \varphi\xi \end{matrix} \right\} - \left\{ \begin{matrix} \varphi \\ \xi j \end{matrix} \right\} \left\{ \begin{matrix} \eta \\ \varphi\phi \end{matrix} \right\}, \quad (5.46)$$

$$R_{j\xi k}^l = \partial_{\xi} \left(\left\{ \begin{matrix} l \\ k j \end{matrix} \right\} \right) - \partial_k \left(\left\{ \begin{matrix} l \\ j\xi \end{matrix} \right\} \right) + \left\{ \begin{matrix} i \\ k j \end{matrix} \right\} \left\{ \begin{matrix} l \\ i\xi \end{matrix} \right\} - \left\{ \begin{matrix} i \\ \xi j \end{matrix} \right\} \left\{ \begin{matrix} l \\ ik \end{matrix} \right\}, \quad (5.47)$$

$$R_{jn\phi}^{\eta} = \partial_n \left(\left\{ \begin{matrix} \eta \\ \phi j \end{matrix} \right\} \right) - \partial_{\phi} \left(\left\{ \begin{matrix} \eta \\ j n \end{matrix} \right\} \right) + \left\{ \begin{matrix} \varphi \\ \phi j \end{matrix} \right\} \left\{ \begin{matrix} \eta \\ \varphi n \end{matrix} \right\} - \left\{ \begin{matrix} \varphi \\ n j \end{matrix} \right\} \left\{ \begin{matrix} \eta \\ \varphi\phi \end{matrix} \right\}, \quad (5.48)$$

$$R_{jn k}^l = \partial_n \left(\left\{ \begin{matrix} l \\ k j \end{matrix} \right\} \right) - \partial_k \left(\left\{ \begin{matrix} l \\ j n \end{matrix} \right\} \right) + \left\{ \begin{matrix} i \\ k j \end{matrix} \right\} \left\{ \begin{matrix} l \\ in \end{matrix} \right\} - \left\{ \begin{matrix} i \\ n j \end{matrix} \right\} \left\{ \begin{matrix} l \\ ik \end{matrix} \right\}. \quad (5.49)$$

Here, the quaternionic Riemannian tensor keeps track that how much scalar and vector components of quaternion change when we propagate parallel along with a small parallelogram. If the value of quaternionic Riemannian Christoffel curvature is zero then the quaternionic curved space-time is converted into flat space-time. On the other hand, the quaternionic Ricci tensor is an important contraction of quaternionic Riemannian Christoffel tensor which explains the volume changes when an object parallels transport along a geodesic. In this case

$$\begin{aligned}
g_{h\eta}R_{\xi n\phi}^{\eta} &= R_{h\xi n\phi} \\
&= \partial_n \left(g_{h\eta} \begin{Bmatrix} \eta \\ \phi\xi \end{Bmatrix} \right) - \partial_{\phi} \left(g_{h\eta} \begin{Bmatrix} \eta \\ \xi n \end{Bmatrix} \right) + g_{h\eta} \begin{Bmatrix} \varphi \\ \phi\xi \end{Bmatrix} \begin{Bmatrix} \eta \\ \varphi n \end{Bmatrix} \\
&\quad - g_{h\eta} \begin{Bmatrix} \varphi \\ n\xi \end{Bmatrix} \begin{Bmatrix} \eta \\ \varphi\phi \end{Bmatrix}, \tag{5.50}
\end{aligned}$$

which can be simplified in terms of a metric tensor as

$$\begin{aligned}
R_{h\xi n\phi} &= \frac{1}{2} \left[\frac{\partial^2 g_{h\phi}}{\partial P^n \partial P^{\xi}} + \frac{\partial^2 g_{\xi n}}{\partial P^h \partial P^{\phi}} - \frac{\partial^2 g_{\xi\phi}}{\partial P^n \partial P^h} - \frac{\partial^2 g_{nh}}{\partial P^{\phi} \partial P^{\xi}} \right] \\
&\quad + g_{\eta h} \begin{Bmatrix} \varphi \\ \phi\xi \end{Bmatrix} \begin{Bmatrix} \eta \\ \phi n \end{Bmatrix} - g_{\eta h} \begin{Bmatrix} \varphi \\ n\xi \end{Bmatrix} \begin{Bmatrix} \eta \\ \varphi\phi \end{Bmatrix}. \tag{5.51}
\end{aligned}$$

Now, contracting equation (5.51) with the metric tensor as $g^{hn}R_{h\xi n\phi} = R_{\xi\phi}$, where $R_{\xi\phi}$ is a quaternionic form of Ricci tensor in scalar field, so that

$$R_{\xi\phi} = \partial_n \left(\begin{Bmatrix} n \\ \phi\xi \end{Bmatrix} \right) - \partial_{\phi} \left(\begin{Bmatrix} n \\ \xi n \end{Bmatrix} \right) + \begin{Bmatrix} \varphi \\ \phi\xi \end{Bmatrix} \begin{Bmatrix} n \\ \varphi n \end{Bmatrix} - \begin{Bmatrix} \varphi \\ n\xi \end{Bmatrix} \begin{Bmatrix} n \\ \varphi\phi \end{Bmatrix}. \tag{5.52}$$

Similarly, to involving quaternionic vector field in addition to scalar field, also in pure vector field the Ricci tensors become

$$R_{\xi k} = \partial_n \left(\begin{Bmatrix} n \\ k\xi \end{Bmatrix} \right) - \partial_k \left(\begin{Bmatrix} n \\ \xi n \end{Bmatrix} \right) + \begin{Bmatrix} i \\ k\xi \end{Bmatrix} \begin{Bmatrix} n \\ in \end{Bmatrix} - \begin{Bmatrix} i \\ n\xi \end{Bmatrix} \begin{Bmatrix} n \\ ik \end{Bmatrix}, \tag{5.53}$$

$$R_{jk} = \partial_n \left(\begin{Bmatrix} n \\ kj \end{Bmatrix} \right) - \partial_k \left(\begin{Bmatrix} n \\ jn \end{Bmatrix} \right) + \begin{Bmatrix} i \\ kj \end{Bmatrix} \begin{Bmatrix} n \\ in \end{Bmatrix} - \begin{Bmatrix} i \\ nj \end{Bmatrix} \begin{Bmatrix} n \\ ik \end{Bmatrix}. \tag{5.54}$$

We should notice that the Riemannian tensor ($R_{h\xi n\phi}$) is symmetric to first with third indices and second with fourth indices [3]. Thus the Ricci tensor is also symmetric. The quaternionic form of Ricci tensor can further contract with metric tensor and obtain scalar curvature tensor (R) of rank zero, such that $g^{\xi\phi}R_{\xi\phi} = R$.

5.6 Quaternionic Einstein-field like equation

To express the quaternionic Einstein field like equation, let us start with the Poisson equation as

$$\nabla^2\Phi = 4\pi G\rho, \quad (5.55)$$

where Φ is the Newtonian gravitational potential, G is gravitational constant and ρ is the mass density. Moreover, in metric tensor form [45], we have $\nabla^2\Phi = -\frac{1}{2}\nabla^2g_{00}$. Thus

$$\nabla^2g_{00} = -8\pi G\rho. \quad (5.56)$$

It should be noticed that in equation (5.56) we introduced the quaternionic form of scalar metric tensor ($g_{\xi\xi}$) in place of Newtonian potential (Φ) and quaternionic mass density ($T_{\xi\xi}$) in place of (ρ). Generally, the quaternionic energy-momentum tensor can be written as

$$T^{\mu\nu} \simeq \begin{bmatrix} T^{\xi\xi} & T^{\xi 1} & T^{\xi 2} & T^{\xi 3} \\ T^{1\xi} & T^{11} & T^{12} & T^{13} \\ T^{2\xi} & T^{21} & T^{22} & T^{23} \\ T^{3\xi} & T^{31} & T^{32} & T^{33} \end{bmatrix}. \quad (5.57)$$

Therefore, equation (5.56) leads to $\nabla^2g_{\xi\xi} = -8\pi GT_{\xi\xi}$. Since the quaternionic Ricci tensor can also be written in the form of a double derivative of the quaternionic metric tensor. So, we get

$$R_{\xi\xi} = -8\pi GT_{\xi\xi}. \quad (5.58)$$

Further, the rest components of quaternionic Riemannian Christoffel curvature tensors can express as

$$\begin{aligned} R_{\xi 1} &= -8\pi GT_{\xi 1}, & R_{\xi 2} &= -8\pi GT_{\xi 2}, & R_{\xi 3} &= -8\pi GT_{\xi 3}, \\ R_{11} &= -8\pi GT_{11}, & R_{22} &= -8\pi GT_{22}, & R_{33} &= -8\pi GT_{33}, \\ R_{12} &= -8\pi GT_{12}, & R_{23} &= -8\pi GT_{23}, & R_{31} &= -8\pi GT_{31}, \end{aligned} \quad (5.59)$$

where the other components i.e. $R_{1\xi}, R_{2\xi}, R_{3\xi}, R_{21}, R_{32}, R_{13}$ are symmetrical to $R_{\xi 1}, R_{\xi 2}, R_{\xi 3}, R_{12}, R_{23}, R_{31}$, respectively. Therefore, we may write a generalized quaternionic form of Einstein field-like equation as $R_{\mu\nu} = -8\pi GT_{\mu\nu}$. Since the quaternionic energy-momentum tensor is conserved quantity but quaternionic Ricci tensor is not, so it should be divergence-free. Thus, by using Bianchi identity $R_{\alpha\mu\beta\gamma;\sigma} + R_{\alpha\mu\gamma\sigma;\beta} + R_{\alpha\mu\sigma\beta;\gamma} = 0$, the quaternionic Einstein field-like equation can be written as $R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}R = -8\pi GT_{\mu\nu}$. Here, we can see that the right-hand part shows quaternionic energy and matter, while the left-hand part shows the quaternionic geometry. Thus the quaternionic formalism is perfectly fitted to describe a new approach in the dynamics of a particle in curved space-time.

6 Conclusion

In Riemannian geometry, the curvature of space-time is directly connected to the energy and momentum of whatever matter and radiation are present. Generally, the space-time can be shown using simple experiments following the free-fall trajectories of different test particles, the result of transporting space-time vectors that can denote a particle's velocity (time-like vectors) will vary with the particle's trajectory. The origin of the idea of this space-time curvature is a mathematical concept. Thus the four-dimensional curvature space-time has been visualized by a quaternionic algebra. The transformation rules for two quaternionic frames have been discussed. Interestingly, we discussed that the effect of curvature space-time arises due to the quaternionic Christoffel symbol. Also, it is concluded that the curved path of freely moving particles is effectively explained by the quaternionic geodesic equation. The Christoffel symbol has been established in terms of the derivative of quaternionic metric tensor while the gravitational potential in tensorial form is represented by the metric tensor. It has been confirmed that the volume element of an object obeying the geodesic path changes due to the contraction of the quaternionic Riemannian tensor while the contraction of the quaternionic Ricci tensor gives a scalar quantity representing the magnitude of change in volume of the object. We also have been discussed the quaternionic Einstein field-like equation for gravitation that relates the geometry of quaternionic space-time with the distribution of matter within it. The simplification of quaternionic formalism will be achieved by the approximation as quaternionic flat space-time with a small deviation, which leads to the linearized Einstein field equation. These quaternionic equations may be used to study the phenomena such as gravitational waves.

References

- [1] Zhang T (2010) Fundamental Elements and Interactions of Nature: A Classical Unification Theory. *Prog Phys* 2 (2010): 36
- [2] Beiser A (2003) *Concepts of Modern Physics*. (6th Ed) McGraw-Hill, New York
- [3] Weinberg S (1972) *Gravitation and Cosmology*. John Wiley & Sons, New York
- [4] Dickson L E (1919) On Quaternion and their Generalization and the History of the eight square theorem. *Ann Math* 20: 153
- [5] Hamilton W R (1969) *Elements of Quaternions*. (Vol. I & II) Chelsea Publishing, New York
- [6] Cayley A (1845) On certain results relating to quaternions. *Phil Mag* 26: 210

- [7] Graves R P (1975) Life of Sir William Rowan Hamilton. (Vol 3) Arno Press, New York
- [8] Baez J C (2001) The octonions. Bull Am Math Soc 39: 145
- [9] Finklestein D, Jauch J M, Schiminovich S, Speiser D (1963) Principle of general quaternion covariance. J Math Phys 4: 788
- [10] Adler S L (1995) Quaternion Quantum Mechanics and Quantum Fields. Oxford University Press, New York
- [11] Bisht P S, Negi O P S, Rajput B S (1991) Quaternion Gauge theory of dyonic fields. Prog Theor Phys 85: 157
- [12] Leo S D (1996) Quaternions and Special Relativity. J Math Phys 37: 2955
- [13] Imaeda K (1979) Quaternionic formulation of tachyons, superluminal transformations and a complex space-time. Nuovo Cimento B50: 271
- [14] Negi O P S, Rajput B S (1983) Quaternionic formulation for electromagnetic field equations. Lett Nuovo Cimento 37: 325
- [15] Majerniček V (1999) Quaternionic formulation of the classical fields. Adv Appl Cliff Algebra 9: 119
- [16] Arbab A I (2011) A Quaternionic Quantum Mechanics. Appl Phys Res 3: 160
- [17] Giardino S (2016) Quaternionic particle in a relativistic box. Found Phys 46: 473
- [18] Chanyal B C (2017) A relativistic quantum theory of dyons wave propagation. Canadian J Phys 95: 1200
- [19] Chanyal B C (2018) A new development in quantum field equations of dyons. Canadian J Phys 96: 1192
- [20] Chanyal B C, Pathak M (2018) Quaternionic Approach to Dual Magneto-hydrodynamics of Dyonic cold plasma. Adv High Energy Phys 2018 Article ID 7843730: 13 pp
- [21] Chanyal B C (2019) Quaternionic approach on the Dirac Maxwell, Bernoulli and Navier Stokes equations for dyonic fluid plasma. Int J Mod Phys A 34: 1950202
- [22] Chanyal B C, Karnatak S (2020) A comparative study of quaternionic rotational Dirac equation and its interpretation. Int J Geom Methods Mod Phys 17: 2050018
- [23] Demir S, Tanisli M, Kansu M E (2015) Octonic massless field equations. Int J Mod Phys A 30: 1550084

- [24] Demir S, Tanisli M, Tolan T (2013) Octonic gravitational field equations. *Int J Mod Phys A* 28: 1350112
- [25] Mironov V L, Mironov S V (2009) Octonic representation of electromagnetic field equations. *J Math Phys* 50: 12901
- [26] Mironov V L, Mironov S V (2015) Sedeonic equations of massive fields. *Int J Theor Phys* 54: 153
- [27] Weng ZH (2015) Some properties of dark matter field in the complex octonion space. *Int J Mod Phys A* 30: 1550212
- [28] Weng ZH (2019) Four interactions in the sedenion curved spaces. *Int J Geom Methods Mod Phys* 16: 1950019
- [29] Chanyal B C, Bisht P S, Negi O P S (2010) Generalized octonion electrodynamics. *Int J Theor Phys* 49: 1333
- [30] Chanyal B C, Bisht P S, Negi O P S (2011) Generalized split-octonion electrodynamics. *Int J Theor Phys* 50: 1919
- [31] Chanyal B C, Sharma V K, Negi O P S (2015) Octonionic gravi-electromagnetism and dark matter. *Int J Theor Phys* 54: 3516
- [32] Chanyal B C (2015) Octonion generalization of Pauli and Dirac matrices. *Int J Geom Methods Mod Phys* 12: 1550007
- [33] Chanyal B C, Chanyal S K, Bektas O, Yuce S (2016) A new approach on electromagnetism with dual number coefficient octonion algebra. *Int J Geom Methods Mod Phys* 13: 1630013
- [34] Chanyal B C (2017) Split octonion reformulation for electromagnetic Chiral media of massive dyons. *Commun Theor Phys* 68: 701
- [35] Chanyal B C (2014) Sedenion unified theory of gravi-electromagnetism. *Indian J Phys* 88: 1197
- [36] Edmonds J D (1974) Quaternion Wave Equations in Curved Space-Time. *Int J Theor Phys* 10: 115
- [37] Weng ZH (2016) Forces in the complex octonion curved space. *Int J Geom Methods Mod Phys* 13: 1650076
- [38] Heinbockel J H (1996) Introduction to tensor calculus and continuum mechanics. Old Dominion University, Virginia
- [39] Ferkinghoff-Borg J (2007) Introduction to vector and tensor analysis. University of Phoenix, Arizona
- [40] Narlikar J V (1978) General Relativity and Cosmology. Macmillan India Press, Madras

- [41] Rastall P (1964) Quaternions in relativity. *Rev Mod Phys* 36(3): 820
- [42] Hamilton W R (1853) *Lectures on Quaternions*. Hodge and Smith, Dublin
- [43] Weinberger P (2014) Hamilton and the square root of minus one. *Philos Mag* 94: 1263
- [44] Myszkowski M (2019) A new perspective on space-time 4D rotations and the $SO(4)$ transformation group. *Results Phys* 13: 1
- [45] Gron O, Hervik S (2007) *Einstein's General Theory of Relativity: With Modern Applications in Cosmology*. Springer Publication, New York