

# The tree world

---

**Markos Maniatis,**

*Centro de Ciencias Exactas & Departamento de Ciencias Básicas, Universidad del Bío-Bío,  
Chillán, Chile*

*E-mail:* [maniatis8@gmail.com](mailto:maniatis8@gmail.com)

ABSTRACT: Tree amplitudes of any gauge theory and gravity can be factorized into primitive three-particle amplitudes by the BCFW recursion relations. We show that the amplitudes at any perturbation order are given by tree amplitudes with additional pairs of particles in the forward limit. As an example we compute the four-gluon all-plus helicity amplitude at next-to-leading order.

---

## Contents

<b>1</b>	<b>Introduction</b>	<b>1</b>
<b>2</b>	<b>Tree amplitudes</b>	<b>2</b>
<b>3</b>	<b>Tree amplitudes forming higher orders</b>	<b>4</b>
<b>4</b>	<b>Example four-gluon amplitude</b>	<b>7</b>
4.1	NLO all-plus four-gluon amplitude	8
4.2	Sketch of the NNLO four-gluon amplitude	13
<b>5</b>	<b>Conclusions</b>	<b>14</b>
<b>A</b>	<b>Spinor helicity conventions</b>	<b>15</b>

---

## 1 Introduction

It is well known that the computation of scattering amplitudes via Feynman diagrams has a number of drawbacks: First of all, Feynman diagrams are plagued with gauge redundancies, that is, single diagrams are in general not gauge invariant and only the sum of diagrams (or certain subsets) restore gauge invariance. For instance the three-gluon vertex violates gauge invariance in general and another four-gluon vertex has to be introduced along with “ghost” particles. Feynman diagrams are in general even not Lorentz covariant, and only the product with polarization “tensors” form Lorentz-invariant expressions. Moreover, in the diagrams momentum conservation is violated, with internal lines described by virtual particles.

Of course, Feynman diagrams are nothing more than a mathematical prescription to generate amplitudes. However, this is also done in a redundant way: Every Feynman diagram to a certain perturbation order is of the same order as the final result. This means that most contributions cancel each other. Maybe the most striking example in this respect is the Parke-Taylor formula for the tree level scattering amplitude of  $n$  external gluons, two carrying negative helicity and the rest positive helicities (maximally helicity violating amplitude). Suppressing the color factor, the correct result was conjectured [1],

$$A_n(1^+, \dots, i^-, \dots, j^-, \dots, n^+) = \frac{\langle ij \rangle}{\langle 12 \rangle \langle 23 \rangle \cdots \langle n1 \rangle}, \quad (1.1)$$

where the expressions  $\langle ab \rangle$  denote scalar products in the helicity formalism - see appendix A for a very brief recap and the conventions used. In terms of Feynman diagrams the computation of this amplitude requires for  $n = 5$  external gluons to calculate 25 diagrams, and for  $n = 6$  external gluons already 220 diagrams.

A milestone in the program to deduce scattering amplitudes from *first principles*, that is, Lorentz invariance, locality, unitarity, and gauge groups, are the BCFW recursion relations [2, 3]. Via analytic continuation, or deformation, of the momenta of external particles it has been shown that in gauge theories and gravity the tree amplitudes can be factorized into on-shell *primitive* three-particle amplitudes. For instance the Parke-Taylor formula (1.1) can be proven by the BCFW recursion relations.

However, the BCFW recursion relations rely on tree amplitudes, which of course form only an unphysical subset of the scattering amplitude. There has been spent a lot of effort to go beyond leading order. In particular at next-to-leading order it has been shown that via unitary cuts the amplitudes can be computed [4–8].

A great step towards the derivation of amplitudes at any perturbation order is the Feynman-tree theorem [9, 10]. In a Feynman diagram at any perturbation order a loop can be *opened* by cutting the propagators which form the loop in all different ways. Recursively, all loops can be opened and finally there remain only tree amplitudes. This approach has been modified and applied to different amplitudes [11–13] and combined with a subsequent application of the BCFW recursion relations [14–16]. However, this approach is dedicated to Feynman diagrams based on loops, that is, virtual particles. As to be expected, there is no direct translation from Feynman diagrams opening the loops and on-shell amplitudes at higher perturbation orders [17].

Here we want to propose an approach relying on on-shell amplitudes: We argue that in the tree on-shell amplitudes we inevitably have to take pairs of particles in the forward limit into account. Since the particles in the forward limit have opposite helicities and momenta, they are physically indistinguishable from the leading order on-shell tree amplitudes. We study on-shell tree diagrams adding systematically pairs of particles in the forward limit. We can then apply the BCFW recursion relations to these tree amplitudes taking particles in the forward limit into account. We show that we get the on-shell amplitudes at any perturbation order by a factorization into primitive three-point on-shell amplitudes. This is in particular very attractive since the three-point primitives are fixed by Lorentz-invariance, that is, little group scaling and the gauge group.

We apply the proposed method to the four-gluon all-plus helicity amplitude at next-to-leading order. The four-point amplitude is an excellent framework to study the proposed methods. Eventually we sketch how at next-to-next-to leading order the four-gluon amplitude can be computed taking particles in the forward limit into account.

## 2 Tree amplitudes

We start considering an arbitrary  $n$ -point on-shell tree amplitude,

$$A_n(1^{h_1}, 2^{h_2}, \dots, n^{h_n}). \tag{2.1}$$

We note that we are considering here the complete amplitude and not a color-ordered subamplitude. The momenta and helicities of the external massless particles are denoted by  $p_i$  and  $h_i$ , with  $p_i$  written simply as the index  $i$ , with  $i = 1, \dots, n$ . The convention here is that all momenta are outgoing.

From the BCFW recursion relations we know that we can factorize the tree amplitude in terms of *primitive* three-point amplitudes. This requires in every factorizing recursion step to *deform* the momenta of the external particles,

$$\hat{p}_i = p_i + z r_i \quad (2.2)$$

with complex momenta  $r_i$  and one common complex number  $z$ . This analytic continuation of the external momenta is done such that on-shellness as well as momentum conservation are maintained. Note that the recursion relations may give boundary terms. Here we shall consider only theories where the boundary terms do not appear. As has been shown this is the case for gauge theories as well as gravity.

As an example we may think of an QCD amplitude. Since it is an on-shell tree amplitude it is trivalent. Let us remind ourselves that it is a peculiarity of Feynman diagrams that we have to consider the four-gluon vertex in QCD. Any  $n$ -point trivalent tree amplitude, with  $n \geq 3$  has the following number of couplings, propagators, and subamplitudes,

external particles	couplings	propagators	subamplitudes	
$n$	$n-2$	$n-3$	$(2n-5)!!$	(2.3)

Here we have as usual denoted by  $1!! = 1$ ,  $3!! = 3 \cdot 1$ ,  $5!! = 5 \cdot 3 \cdot 1$ ,  $\dots$ . The number of couplings equals the number of *primitive* three-point amplitudes which built the subamplitudes. An example is the  $n = 6$  gluon amplitude with all-plus helicities, consisting of  $(2n - 5)!! = 105$  trivalent subamplitudes, one of them is

$$(2.4)$$

Suppose that the particles obey an underlying color structure, that is, they fulfill the Jacobi identity. Then we know from the color-kinematics relations [18] that we can write the trivalent subamplitude in terms of subamplitudes of *multi-peripheral* form or equivalently denoted by *half-ladder* form. The subamplitude (2.4) can then be written as minus the sum of the two subamplitudes,

$$(2.5)$$

But for the moment we keep the amplitudes completely general, that is, we do not assume that they are written in multi-peripheral form.

What about the three-point primitive amplitudes from which all tree amplitudes are formed? The kinematics of these primitive amplitudes is completely determined by little-group scaling, that is Lorentz invariance. An interaction of three scalars can not have



order to do this we have to deform the momenta in general, complying with BCFW. However, the amplitude  $A_n(1^{h_1}, \dots, n^{h_n})$  is indistinguishable from amplitudes with additional pairs of particles in the forward limit: For instance adding one pair of particles with opposite momenta and opposite helicities, the amplitude  $A_{n+2}^{\text{pair}}(1^{h_1}, \dots, n^{h_n}, l_1^{h_{l_1}}, -l_1^{-h_{l_1}})$  is indistinguishable from the amplitude  $A_n(1^{h_1}, \dots, n^{h_n})$ . The pair of additional particles are *balanced* and together form a vacuum state. Therefore we find that along with the amplitude  $A_n(1^{h_1}, \dots, n^{h_n})$  we have also to take into account the amplitudes with  $k = 1, 2, 3, \dots$  additional pairs of particles in the forward limit:

$$A_{n+2k}^{\text{pairs}}(1^{h_1}, \dots, n^{h_n}, l_1^{h_{l_1}}, -l_1^{-h_{l_1}}, \dots, l_k^{h_{l_k}}, -l_k^{-h_{l_k}}). \quad (3.1)$$

With every additional pair we add two primitive  $A_3$  amplitudes, increase the number of couplings by two, that is, every additional pair in the forward limit increases the order by one unit. The forward momenta  $l_i$  and  $-l_i$ ,  $i = 1, \dots, k$  are opposite but arbitrary, therefore we have to integrate over their phase space. For the amplitude with  $k$  additional pairs in the forward limit we get

$$(-1)^{2(h_{l_1} + \dots + h_{l_k})} \int_{l_1, \dots, l_k} A_{n+2k}^{\text{pairs}}(1^{h_1}, \dots, n^{h_n}, l_1^{h_{l_1}}, -l_1^{-h_{l_1}}, \dots, l_k^{h_{l_k}}, -l_k^{-h_{l_k}}), \quad (3.2)$$

where we abbreviate the integrations as

$$\int_{l_i} = \int \frac{d^D l_i}{(2\pi)^D} 2\pi \delta^{(+)}(l_i). \quad (3.3)$$

As usual we write  $\delta^{(+)}(l_i) = \theta(l_{i0})\delta(l_i^2)$ . We have to perform the calculation in general  $D = 4 - 2\epsilon$  dimensions in order to regularize the in general singular integrals. The prefactor in (3.2) gives a factor of  $-1$  for every additional fermion pair.

We emphasize that all the subamplitudes which appear on the right-hand side of (3.2) with  $k$  additional pairs are trivalent or, as we will see, products of trivalent amplitudes. Any loop of on-shell amplitudes would violate momentum conservation. The number of additional pairs  $k$  corresponds to subamplitudes at  $N^k\text{LO}$ . However, we have to take into account also split - or cut - subamplitudes, where propagators are replaced by further pairs in the forward limit; without increasing the perturbation order.

Let us consider as an example the six-gluon maximally helicity violating amplitude, that is two gluons carry negative, all remaining ones carry plus helicity. One of the trivalent subamplitudes at leading order (LO) of  $A_6(1^-, 2^-, 3^+, 4^+, 5^+, 6^+)$  is

$$\begin{array}{ccccccc}
 & 2^- & 3^+ & 4^+ & 5^+ & & \\
 & | & | & | & | & & \\
 1^- & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & 6^+ \\
 & & & & & & & & & & 
 \end{array} \quad (3.4)$$

Obviously, we can not split this leading-order amplitude by replacing a propagator by a vacuum pair, for instance,

$$\begin{array}{c}
2^- \quad 3^+ \\
| \quad | \\
1^- \text{---} \bullet \text{---} \bullet \text{---} -l_1^- \quad l_1^+ \text{---} \bullet \text{---} \bullet \text{---} 6^+ \\
4^+ \quad 5^+
\end{array} \quad (3.5)$$

This is not possible since we have to satisfy both, on-shellness and momentum conservation but  $l_1^2 = (p_1 + p_2 + p_3)^2 \neq 0$  for general external momenta.

Now let us go beyond LO and add one pair in the forward limit, a subamplitude of  $A_{6+2}^{1 \text{ pair}}(1^-, 2^-, 3^+, 4^+, 5^+, 6^+, -l_1^-, l_1^+)$  is

$$\begin{array}{c}
2^- \quad 3^+ \quad -l_1^- \quad 4^+ \quad 5^+ \quad l_1^+ \\
| \quad | \quad | \quad | \quad | \quad | \\
1^- \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} 6^+
\end{array} \quad (3.6)$$

The number of couplings is increased by two corresponding to an amplitude at NLO, corresponding to  $k = 1$ . At the same order we have also to consider the split subamplitude  $A_{6+2+2}^{1 \text{ pair}, 1 \text{ split}}(1^-, 2^-, 3^+, 4^+, 5^+, 6^+, -l_1^-, l_1^+, -l_2^-, l_2^+)$ , for instance,

$$\begin{array}{c}
2^- \quad 3^+ \quad -l_1^- \quad 4^+ \quad \quad \quad 5^+ \quad l_1^+ \\
| \quad | \quad | \quad | \quad \quad \quad | \quad | \\
1^- \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} -l_2^- \quad l_2^+ \text{---} \bullet \text{---} \bullet \text{---} 6^+
\end{array} \quad (3.7)$$

This decomposition in two *partial* amplitudes, denoted with the superscript “1 split”, does neither violate momentum conservation nor on-shellness. Alternatively we can replace also different propagators - as long as we do not split into a vanishing  $A_3$  subamplitude. We see that at NLO we can at most replace one propagator, that is split once into two partial amplitudes, respecting on-shellness and momentum conservation. In general the number of splittings of the subamplitudes can not exceed the number of additional pairs in the forward limit added.

For an arbitrary order  $k$ , that is, amplitudes at  $N^k\text{LO}$ , we have to add  $k$  pairs of external forward particles. After the addition of these  $k$  pairs of external particles we can, without changing the order  $k$ , replace in the resulting amplitudes propagators by additional pairs in the forward limit. By every such cut of a propagator we split the amplitude into separate parts. Obviously there is an upper limit of propagators we can replace: The maximal number of propagators we may replace by particles in the forward limit is equal to the number of pairs of additional external particles  $k$ . This comes from the fact that we have to fulfill simultaneously on-shellness and momentum conservation. We get therefore at a certain perturbation order  $k$  exactly  $k$  pairs of external particles in the forward limit with afterwards  $0, \dots, k$  replaced propagators. The subamplitudes split into a number of parts given by the number of replaced propagators plus one. We thus get the composition of tree sub and partial amplitudes at order  $N^k\text{LO}$ :

$$A_n^{\text{N}^k\text{LO}}(1^{h_1}, \dots, n^{h_n}) = (-1)^{2(h_{l_1} + \dots + h_{l_k})} \int_{l_1, \dots, l_k} A_{n+2k}^{k \text{ pairs}}(1^{h_1}, \dots, n^{h_n}, l_1^{h_{l_1}}, -l_1^{-h_{l_1}}, \dots, l_k^{h_{l_k}}, -l_k^{-h_{l_k}}) \quad (3.8)$$

with

$$A_{n+2k}^{k \text{ pairs}} = A_{n+2k}^{k \text{ pairs}, 0 \text{ split}} + (-1)^{2h'_1} \int_{l'_1} A_{n+2k+2}^{k \text{ pairs}, 1 \text{ split}} + \dots$$

$$+ (-1)^{2(h'_1 + \dots + h'_k)} \int_{l'_1, \dots, l'_k} A_{n+2k+2k}^{k \text{ pairs}, k \text{ splits}}$$

and partial amplitudes  $i = 1, \dots, k$ ,

$$A_{n+2k+2i}^{k \text{ pairs}, i \text{ splits}}(1^{h_1}, \dots, n^{h_n}, l_1^{h_{l_1}}, -l_1^{-h_{l_1}}, \dots, l_k^{h_{l_k}}, -l_k^{-h_{l_k}}, l_1^{h'_1}, -l_1^{-h'_1}, \dots, l_i^{h'_i}, -l_i^{-h'_i}).$$

Similar to the pairs added to the trivalent amplitude we have to integrate over the phase space of the momenta ( $l'_i$ ) of the cut propagators. Let us illustrate this in an example - the all-plus four-gluon amplitude. The leading order ( $k = 0$ ) amplitude with  $n = 4$ , that is,  $A_4$  consists of  $(2n - 5)!! = 3$  subamplitudes with coupling order  $n - 2 = 2$ , that is,  $g^2$ . At the next-to-leading order ( $k = 1$ ), we have to add one external pair of forward particles in order to increase the number of couplings by two. For  $k = 1$  we can either cut no propagator at all or at most cut one propagator in the resulting subamplitude. Without any propagator replacements we get one trivalent amplitude and with one cut we get the subamplitude split into two. At next-to-next-to leading order, that is, N<sup>k</sup>L0 with  $k = 2$ , we have to add exactly two pairs of forward particles. In these amplitudes we may either cut no propagator or cut one or maximally two, yielding trivalent subamplitudes unsplit, split into two or into three parts.

With (3.8) we have written the amplitudes to any perturbation order in terms of trivalent subamplitudes or products of trivalent partial amplitudes. We see that we get split amplitudes, that is, products of tree amplitudes, where the number of maximal splittings equals the perturbation order. Since all subamplitudes are tree amplitudes we can apply the BCFW recursion relations, eventually express scattering amplitudes in terms of primitive  $A_3$  amplitudes. From the splittings we get the different color structures: By color kinematics every trivalent subamplitude gives a trace of color generators and we see that we get the known products of traces for higher order scattering amplitudes.

## 4 Example four-gluon amplitude

In this section we want to show how the four-gluon interaction is determined in terms of tree amplitudes. With respect to an increasing perturbation order we have to add systematically, as outlined in the previous sections, particle pairs in the forward limit.

The four-gluon amplitude provides an excellent framework to study the methods. It is the lowest-point amplitude complying with on-shellness and momentum conservation for massless particles. The simplest one is the all-plus helicity amplitude, but all remaining ones can be studied in a similar way.

At leading order, the amplitude vanishes for all plus helicities for any number of external gluons,

$$A_n^{\text{LO}}(1^+, 2^+, \dots, n^+) = 0. \quad (4.1)$$

The usual proof is given with the help of polarization “tensors”. Not resorting to polarization “tensors” we can easily see that the partial amplitudes (4.1) have to vanish from the recursive application of BCFW. First of all we can assume that the amplitudes are in multi-peripheral form by the application of color-kinematics relations [18]. By a convenient relabeling of the external momenta we can apply BCFW as indicated in the following amplitude,

$$\begin{array}{ccccccc}
 & \hat{2}^+ & \hat{3}^+ & & (n-1)^+ & & \\
 & | & | & & | & & \\
 1^+ & \text{---} & \hat{P}_{12} & \text{---} & \dots & \text{---} & n^+ \\
 & \cdot & \cdot & & \cdot & & \\
 & - & + & & & & \\
 & | & | & & | & & \\
 & \cdot & \cdot & & \cdot & & 
 \end{array} \quad (4.2)$$

We choose a  $|2, 3\rangle$  shift with the helicities as indicated and the propagator given by  $P_{12} = 1/(p_1 + p_2)^2$ . The opposite choice of helicities for the particles connected to the propagator  $\hat{P}_{12}$  is excluded because it is directly proportional to a vanishing all-plus amplitude  $A_3$ . However with further recursions of BCFW we can not avoid to get one vanishing all-plus amplitude  $A_3$  and therefore the all-plus  $n$ -gluon amplitude vanishes at leading order.

The situation changes if we consider the amplitude beyond leading order, employing (3.8), for instance at next-to-leading order we have to consider subamplitudes of the form  $A_{n+2}^{\text{pair}}(1, \dots, n, -l_1^-, l_1^+)$  with a pair of gluons added in the forward limit. Applying color-kinematics relations, we get the subamplitudes in multi-peripheral form, where we have the extra pair of external particles with momentum, respectively helicity  $-l_1^-$ , and  $l_1^+$  in all possible positions. Inevitably, we encounter now one external gluon with negative helicity. Then, in the chain of  $A_3$  primitive amplitudes, similar to (4.2) we can choose the helicities of the internal lines such that there never appears an all-plus amplitude  $A_3$  and the argument above does not apply.

In supersymmetry, we immediately see that the all-plus amplitudes have to vanish at all orders, because for every external additional pair of bosons, at any position, we have to add an amplitude where the bosons are replaced by fermions. The corresponding factor  $(-1)^{2h_i}$  in (3.8) ensures that both contributions enter with opposite signs and therefore the sum of partial amplitudes to all orders of all-plus (or equivalently all minus) amplitudes vanishes in supersymmetry.

#### 4.1 NLO all-plus four-gluon amplitude

The conventional method is to compute the amplitude via Feynman diagrams. An alternative approach [5, 8] is based on unitary cuts applied to an expansion of the amplitude in terms of scalar integrals. In [16] the all-plus four-gluon amplitude has been calculated employing the Feynman-tree theorem. Here we want to show the method in a changed manner explicitly starting from primitive  $A_3$  amplitudes integrating over additional pairs of particles in the forward limit.

As has been argued before, the factor  $(-1)$  in (3.8) ensures that the  $n$ -gluon amplitude vanishes to all orders in supersymmetry (SYM), in particular in  $\mathcal{N} = 1$  and  $\mathcal{N} = 4$  SYM.

From the vanishing amplitude, a simple context between the additional pairs of bosons and fermions can be deduced [20]: In a supersymmetric  $n$ -gluon amplitude we have to consider additional pairs of scalars, fermions and gluons in general. Here we want to calculate the next-to-leading order amplitude so we have to consider one additional pair of forward particles. In  $\mathcal{N} = 4$  SYM we have at next-to-leading order for the vanishing amplitude one additional pair of gluons, ( $g$ ), paired with 4 fermions, ( $f$ ), and 3 complex scalars, ( $s$ ):

$$A_{n+2}^{\mathcal{N}=4, 1 \text{ pair}} = A_{n+2}^{1 \text{ pair } (g)} - 4 A_{n+2}^{1 \text{ pair } (f)} + 3 A_{n+2}^{1 \text{ pair } (s)}. \quad (4.3)$$

In  $\mathcal{N} = 1$  SYM we have at next-to-leading order for the vanishing amplitude one additional pair of gluons, ( $g$ ), paired with 1 fermion, ( $f$ ):

$$A_{n+2}^{\mathcal{N}=1, 1 \text{ pair}} = A_{n+2}^{1 \text{ pair } (g)} - A_{n+2}^{1 \text{ pair } (f)}. \quad (4.4)$$

Since the amplitudes (4.3) and (4.4) vanish for all-plus helicity gluons we see that

$$A_{n+2}^{1 \text{ pair } (g)} = A_{n+2}^{1 \text{ pair } (s)}. \quad (4.5)$$

The task now is to calculate the amplitude (3.8) with one additional pair of scalars in the forward limit:

$$A_4^{\text{NLO}}(1^+, 2^+, 3^+, 4^+) = \int_{l_1} A_{4+2}^{1 \text{ pair } (s)}(1^+, 2^+, 3^+, 4^+, -l_1, l_1). \quad (4.6)$$

The subamplitude  $A_{4+2}^{1 \text{ pair}}$  has one contribution without splitting and one partial amplitude split once (3.8), where we suppress from now on the the superscript ( $s$ ):

$$\begin{aligned} A_{4+2}^{1 \text{ pair}}(1^+, 2^+, 3^+, 4^+, -l_1, l_1) = \\ A_{4+2}^{1 \text{ pair}, 0 \text{ splits}}(1^+, 2^+, 3^+, 4^+, -l_1, l_1) + \int_{l_2} A_{4+2+2}^{1 \text{ pair}, 1 \text{ split}}(1^+, 2^+, 3^+, 4^+, -l_1, l_1, -l_2, l_2). \end{aligned} \quad (4.7)$$

Eventually, at NLO we find the expression

$$\begin{aligned} A_4^{\text{NLO}}(1^+, 2^+, 3^+, 4^+) = A_4^{1 \text{ pair}, 0 \text{ splits}} + A_4^{1 \text{ pair}, 1 \text{ split}} \\ \int_{l_1} A_{4+2}^{1 \text{ pair}, 0 \text{ splits}}(1^+, 2^+, 3^+, 4^+, -l_1, l_1) + \int_{l_1, l_2} A_{4+2+2}^{1 \text{ pair}, 1 \text{ split}}(1^+, 2^+, 3^+, 4^+, -l_1, l_1, -l_2, l_2). \end{aligned} \quad (4.8)$$

A subamplitude of the first term on the r.h.s. of (4.8) can be graphically represented as

$$\begin{array}{ccccccc} & 1^+ & 2^+ & 3^+ & 4^+ & & \\ & \downarrow & \downarrow & \downarrow & \downarrow & & \\ -l_1 & \cdots & \bullet & \cdots & \bullet & \cdots & l_1 \end{array} \quad (4.9)$$

The gluon is drawn as a full line and the complex scalar as a dashed line. A subamplitude with one splitting, that is, with one propagator replaced by a pair in the forward limit corresponds to the partial amplitude in the second term on the r.h.s. of (4.8),

$$\begin{array}{ccc}
 1^+ & 2^+ & \\
 \downarrow & \downarrow & \\
 -l_1 & \text{---} \bullet \text{---} \bullet \text{---} & -l_2
 \end{array}
 \quad
 \begin{array}{ccc}
 3^+ & 4^+ & \\
 \downarrow & \downarrow & \\
 l_2 & \text{---} \bullet \text{---} \bullet \text{---} & l_1
 \end{array}
 \tag{4.10}$$

Note that a subamplitude with one additional pair, as shown in (4.9) can only be cut in the propagator as shown in (4.10) since all other propagators cut yield an isolated (unshifted)  $A_3$  amplitude. We recall that the primitive amplitudes  $A_3$  is only non-vanishing for BCFW shifted, that is, deformed, momenta. Let us assume that the gluons are color ordered. Then we have for the in total  $4!$  permutations of the external gluons only to take the cyclic permutations  $1 \rightarrow 2 \rightarrow 3 \rightarrow 4 \rightarrow 1$  into account. Following (4.8) we have to integrate over the phase space of the particle pairs in the forward limit.

We encounter the problem that the phase-space integration has to be performed in general  $D$  dimensions but the helicity formalism is given in 4 dimensions. We follow here the solution as outlined in [21]: The  $D = 4 - 2\epsilon$  dimensional momenta  $l_i$  are split into its four-dimensional and the remaining  $\mu = (-2\epsilon)$  dimensional part. The external momenta are kept four dimensional. Suppose that  $p$  is an external four-dimensional momentum, then we find for a propagator with momentum  $(l_i - p)^2 \rightarrow (l_i - p)^2 - \mu^2$ . The integration measure transforms as  $d^D l_i / (2\pi)^D \rightarrow d^{-2\epsilon} \mu / (2\pi)^{-2\epsilon} d^4 l_i / (2\pi)^4$ . We get a four-dimensional phase space integration with an artificial mass parameter  $\mu$  which appears in the phase space momenta in the integrand.

With view on (4.8), respectively, (4.9) and (4.10) we have to calculate the six-point and four-point amplitudes. In this respect we follow closely [22] where tree amplitudes for up to four gluons and two complex massive scalars have been computed. We start the composition of the amplitude with the primitive  $A_3$  amplitude and glue following BCFW the higher-point amplitudes together. The primitive amplitude of one gluon with  $\pm$  helicity and two massive scalars is

$$\begin{array}{ccc}
 2^\pm & & \\
 \downarrow & & \\
 1 & \text{---} \bullet \text{---} & 3
 \end{array}
 \tag{4.11}$$

Since the scalar is massive, the expressions in (2.6) have to be modified. Depending on the helicity of the gluon, the primitive three-point amplitudes read [22, 23]

$$A_3(1, 2^+, 3) = \frac{\langle q|1|2\rangle}{\langle q|2\rangle}, \quad A_3(1, 2^-, 3) = -\frac{\langle 2|1|q\rangle}{[q|2]}. \tag{4.12}$$

Here  $q$  denotes an arbitrary null vector, linearly independent of the momenta of the particles 1 and 2. Glueing two primitive subamplitudes (4.11) together, we get the  $A_4(1, 2^+, 3^+, 4)$  amplitude by one BCFW recursion where we shift the two massless gluon momenta, that is, a  $|2, 3\rangle$  shift,

$$|\hat{2}\rangle = |2\rangle + z|3\rangle, \quad |\hat{3}\rangle = |3\rangle, \quad |\hat{3}\rangle = |3\rangle - z|2\rangle, \quad |\hat{2}\rangle = |2\rangle, \tag{4.13}$$

$$\begin{array}{c}
\hat{2}^+ \quad \hat{3}^+ \\
| \quad | \\
1 \text{ --- } \bullet \text{ --- } \bullet \text{ --- } 4 \\
\quad \quad \quad | \\
\quad \quad \quad 1/\hat{P}^2
\end{array} \tag{4.14}$$

and find with the propagator momentum  $P^2 = (p_1 + p_2)^2$ ,

$$\begin{aligned}
A_4(1, 2^+, 3^+, 4) &= A_3(\hat{2}^+, 1, -\hat{P}) \frac{1}{P^2 - \mu^2} A_3(\hat{P}, \hat{3}^+, 4) = -\frac{\langle \hat{3} | \hat{P} | \hat{2} \rangle}{\langle \hat{3} \hat{2} \rangle} \frac{1}{P^2 - \mu^2} \frac{\langle \hat{2} | 4 | \hat{3} \rangle}{\langle \hat{2} \hat{3} \rangle} \\
&= \frac{\mu^2}{P^2 - \mu^2} \frac{[23]}{\langle 23 \rangle}. \tag{4.15}
\end{aligned}$$

Glueing another  $A_3$  amplitude to the  $A_4$  amplitude we get the  $A_5(1, 2^+, 3^+, 4^+, 5)$  amplitude with a  $|2, 3\rangle$  shift,

$$\begin{array}{c}
\hat{2}^+ \quad \hat{3}^+ \quad 4^+ \\
| \quad | \quad | \\
1 \text{ --- } \bullet \text{ --- } \bullet \text{ --- } \bullet \text{ --- } 5 \\
\quad \quad \quad | \\
\quad \quad \quad 1/\hat{P}^2
\end{array} \tag{4.16}$$

$$A_5(1, 2^+, 3^+, 4^+, 5) = A_3(1, \hat{2}^+, \hat{P}) \frac{1}{P^2 - \mu^2} A_4(-\hat{P}, \hat{3}^+, 4^+, 5). \tag{4.17}$$

Here we plug in our results for  $A_3$ , (4.11), and  $A_4$ , (4.15). Similar we glue together via BCFW the six point amplitude

$$A_6(-l_1, 1^+, 2^+, 3^+, 4^+, l_1) = A_3(-l_1, \hat{1}^+, -\hat{P}) \frac{1}{P^2 - \mu^2} A_5(\hat{P}, \hat{2}^+, 3^+, 4^+, l_1). \tag{4.18}$$

Only one factorization is possible employing a  $[1, 2)$  shift. We eventually find

$$\begin{aligned}
A_6(-l_1, 1^+, 2^+, 3^+, 4^+, l_1) &= \\
&= \frac{\mu^4}{[(p_1 - l_1)^2 - \mu^2][(p_1 + p_2 - l_1)^2 - \mu^2][(p_1 + p_2 + p_3 - l_1)^2 - \mu^2]} \frac{[12][34]}{\langle 12 \rangle \langle 34 \rangle}. \tag{4.19}
\end{aligned}$$

With these preparations we get for the first term on the r.h.s of (4.8) with (4.19)

$$\begin{aligned}
A_4^{\text{one pair, 0 splits}} &= \frac{[12][34]}{\langle 12 \rangle \langle 34 \rangle} \\
&= \int \frac{d^{-2\epsilon} \mu}{(2\pi)^{-2\epsilon}} \mu^4 \int \frac{d^4 l_1}{(2\pi)^3} \frac{\delta^{(+)}(l_1^2 - \mu^2)}{[(p_1 - l_1)^2 - \mu^2][(p_1 + p_2 - l_1)^2 - \mu^2][(p_1 + p_2 + p_3 - l_1)^2 - \mu^2]}. \tag{4.20}
\end{aligned}$$

For the second term on the r.h.s. of (4.8) with one additional pair, split into two parts,

$$\begin{aligned}
A_4^{\text{one pair, 1 split}} &= \\
&= \int \frac{d^{-2\epsilon} \mu}{(2\pi)^{-2\epsilon}} \mu^4 \int \frac{d^4 l_1}{(2\pi)^3} \frac{d^4 l_2}{(2\pi)^3} \delta^{(+)}(l_1^2 - \mu^2) \delta^{(+)}(l_2^2 - \mu^2) A_4(-l_1, 1^+, 2^+, -l_2) A_4(l_2, 3^+, 4^+, l_1). \tag{4.21}
\end{aligned}$$

Here we plug in our results for the two amplitudes from (4.15) and get

$$A_4^{\text{one pair, 1 split}} = \frac{[12][34]}{\langle 12 \rangle \langle 34 \rangle} \int \frac{d^{-2\epsilon} \mu}{(2\pi)^{-2\epsilon} \mu^4} \int \frac{d^4 l_1}{(2\pi)^3} \frac{d^4 l_2}{(2\pi)^3} \frac{\delta^{(+)}(l_1^2 - \mu^2) \delta^{(+)}(l_2^2 - \mu^2)}{[(l_1 - p_1)^2 - \mu^2][(l_2 - p_3)^2 - \mu^2]}. \quad (4.22)$$

This phase space integral is known from the Cutkosky cutting rules and equals twice the imaginary part of a box integral which vanishes. All other permutations of the external momenta give also vanishing contributions. Especially the kinematic factor

$$\frac{[12][34]}{\langle 12 \rangle \langle 34 \rangle} \quad (4.23)$$

is invariant under all permutations. This can be seen for any cyclic permutation  $1 \rightarrow 2 \rightarrow 3 \rightarrow 4$ , using momentum conservation and  $[12] = \langle 41 \rangle [12] / \langle 41 \rangle = -\langle 43 \rangle [32] / \langle 41 \rangle$  and similar  $[34] = -\langle 21 \rangle [14] / \langle 23 \rangle$ . Besides, the expression (4.23) is invariant under  $1 \leftrightarrow 2$ , under  $3 \leftrightarrow 4$ , and under reflections, which gives invariance under all permutations.

It remains to calculate (4.20). We go conveniently back to a phase space integral in  $D$  dimensions,

$$A_4^{1 \text{ pair, 0 splits}} = \frac{[12][34]}{\langle 12 \rangle \langle 34 \rangle} \int \frac{d^D l_1}{(2\pi)^{D-1}} \frac{\delta^{(+)}(l_1^2)}{(p_1 - l_1)^2 (p_1 + p_2 - l_1)^2 (p_1 + p_2 + p_3 - l_1)^2}. \quad (4.24)$$

The integral can be carried out directly with the help of Schwinger parameters [16, 17]

$$\frac{i}{x + i\epsilon} \rightarrow \int_0^\infty da e^{iax}, \quad 2\pi\delta(x) \rightarrow \int_{-\infty}^\infty da e^{iax}. \quad (4.25)$$

We now integrate in the following order: First over the common Schwinger parameter, then over the parameter corresponding to the delta function, and then over the momentum  $l_1$  in  $D$  dimensions. For the integral we get

$$I = \frac{-i\pi^{\frac{D}{2}}}{2(2\pi)^D} \Gamma\left(2 - \frac{D}{2}\right) \int_0^\infty d\alpha_2 d\alpha_3 d\alpha_4 [s \alpha_3 (\alpha_2 + \alpha_3 + \alpha_4 - 1) - t \alpha_2 \alpha_4 - i\epsilon]^{\frac{D}{2}-2}. \quad (4.26)$$

with the Mandelstam variables  $s = 2p_1 p_2$  and  $t = 2p_2 p_3$ . We then substitute  $\alpha_3 \rightarrow \alpha_3 \alpha_4$ , we integrate over  $\alpha_2$ , followed by  $\alpha_4$ , and finally expand about  $D = 4$  dimensions. Eventually, we integrate over  $\alpha_3$  and get for the integral

$$I = \frac{-i}{32\pi^2} \frac{1}{6} \left[ \frac{s}{s+t} + \frac{st}{(s+t)^2} \log\left(\frac{s}{-t}\right) \right] + \mathcal{O}(\epsilon). \quad (4.27)$$

For the color-ordered amplitude we have to consider all cyclic permutations and we see that we get two equal contributions (4.27) and two contributions with the exchange  $s \leftrightarrow t$ . In the sum the brackets (4.27) cancel exactly and we get the known result

$$A_4^{\text{NLO}}(1^+, 2^+, 3^+, 4^+) = \frac{-i}{16\pi^2} \frac{1}{6} \frac{[12][34]}{\langle 12 \rangle \langle 34 \rangle} + \mathcal{O}(\epsilon). \quad (4.28)$$

## 4.2 Sketch of the NNLO four-gluon amplitude

Here we want to briefly sketch, how we may proceed to calculate the all-plus four-gluon amplitude at NNLO with the help of (3.8), that is, with two additional pairs of forward particles. With two additional forward pairs we inevitably encounter two gluons with negative helicity and the supersymmetry relations mentioned in the last section do not hold. Therefore we construct the amplitude directly from the three-gluon primitive amplitude (2.6).

Adding two pairs in the forward limit we have different contributions depending on the kind where the paired momenta appear, for instance,

$$\begin{array}{ccccccccccc}
 & & -l_1^- & 2^+ & l_1^+ & 3^+ & -l_2^- & 4^+ & & & \\
 & & | & | & | & | & | & | & & & \\
 1^+ & \text{---} & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \text{---} & l_2^+ & \\
 \end{array} \tag{4.29}$$

and contributions, where the vacuum pair momenta do not appear separated, for instance,

$$\begin{array}{ccccccccccc}
 & & -l_1^- & 2^+ & -l_2^- & 3^+ & l_1^+ & 4^+ & & & \\
 & & | & | & | & | & | & | & & & \\
 1^+ & \text{---} & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \text{---} & l_2^+ & \\
 \end{array} \tag{4.30}$$

The latter class would, in terms of Feynman diagrams correspond to non-planar contributions, which we can easily see connecting the lines of opposite momenta.

For the unsplit subamplitudes (3.8) we get maximally helicity violating amplitudes which are given by the Parke-Taylor formula (1.1), [1]. Following (3.8) we have for the NNLO calculation to consider two additional pairs and up to two splittings of the subamplitudes, for instance a subamplitude with one splitting - originating from the subamplitude (4.30) reads

$$\begin{array}{ccccccccccc}
 & & -l_1^- & 2^+ & & & -l_2^- & 3^+ & l_1^+ & 4^+ & \\
 & & | & | & & & | & | & | & | & \\
 1^+ & \text{---} & \bullet & \bullet & \text{---} & -l_3^- & l_3^+ & \text{---} & \bullet & \bullet & \bullet & \bullet & \text{---} & l_2^+ \\
 \end{array} \tag{4.31}$$

However, this partial amplitudes have to vanish because we do only have three minus helicities available and have to distribute them to two partial amplitudes. Since all  $n$ -gluon tree amplitudes with less than two minus helicities vanish, at least one partial amplitude has to vanish. Similar, for two splittings we have in total three partial amplitudes and four pairs in the forward limit, that is, we have in total four gluons with minus helicity available - however we would need at least six minus helicities.

Altogether we see that the NNLO four-gluon all-plus helicity gluon amplitude is given by

$$A_4^{NNLO}(1^+, 2^+, 3^+, 4^+) = \int_{l_1, l_2} A_{4+4}^{2 \text{ pairs}, 0 \text{ splits}}(1^+, 2^+, 3^+, 4^+, -l_1^{-h_1}, l_1^{+h_1}, -l_2^{-h_1}, l_2^{h_1}). \tag{4.32}$$

The subamplitudes of (4.32) are all maximally helicity violating amplitudes (1.1). We have to consider all permutations of the positions of the two additional pairs. This gives contributions with different denominators in (1.1). Moreover, the integration has to be done in  $D$  dimensions. One possibility would be to use the four-dimensional helicity scheme as applied in the last section, where the external momenta are kept four dimensional and only the forward pairs are kept in  $D = 4 - 2\epsilon$  dimensions. It is left for future work to do this calculation and show that the known result can be reproduced at NNLO.

## 5 Conclusions

The main point of this paper is the statement that for the computation of on-shell amplitudes we have to take pairs of particles in the forward limit into account. These contributions are indistinguishable from the leading order amplitudes and give higher order contributions. Over the phase-space of the momenta of the particles in the forward limit has to be integrated. Additional pairs of particles in the forward limit appear by the cut of propagators. It has been shown by adding systematically vacuum pairs in the forward limit that on-shell amplitudes are generated order by order in terms of trivalent tree subamplitudes.

Applying the BCFW recursion relations we can decompose all the trivalent subamplitudes in terms of primitive  $A_3$  amplitudes. These primitive amplitudes in turn are fixed by little-group scaling and the gauge group.

In an explicit example we have shown the calculation of the four-gluon all-plus helicity amplitude at next-to-leading. Starting with a primitive  $A_3$  amplitude we have first glued together the higher-point amplitudes, taking additional pairs of particles in the forward limit into account. Eventually we have performed the phase space integrations and reproduced the known result.

We have briefly sketched how at NNLO the all-plus four-gluon amplitude can be constructed. The explicit calculation at this order is left for future work. It would be very interesting to see this computation in detail, especially with respect to the phase-space integrals of the gluons in the forward limit.

With the proposed method we get the scattering amplitudes at any order based on first principles, that is, Lorentz invariance, locality, unitarity and the underlying gauge groups. We are not facing any gauge redundancies, virtual particles, ghosts, or polarization “tensors”. All amplitudes are eventually built by on-shell tree amplitudes.

The forward limit is nevertheless in general singular. Therefore the calculation has to be regularized where we have here used dimensional regularization. Let us mention that the renormalization program is not changed at all compared to the conventional approach to calculate scattering amplitudes via Feynman diagrams.

We expect that massive particles do not cause principal concerns. However, obviously the calculation will be more complicated. This can be seen directly from the helicity formalism, see also appendix A: The spinors follow from the two-by-two matrix defined as  $p = p_\mu(\sigma^\mu)$ , which for massive particles is no longer of rank one, but will have full rank. The decomposition in terms of one pair of spinors therefore has to be replaced by two pairs

of spinor vectors. For instance in [23] it has been shown how the helicity formalism has to be extended for the massive case. It would be interesting to see explicit calculations of amplitudes based on the here proposed methods with massive particles.

Let us mention eventually that we find it quite striking that the perturbative expansion of scattering amplitudes can be deduced from first principles.

## Acknowledgments

The project was supported in part by the project Fondecyt with No. 1200641.

## A Spinor helicity conventions

We follow the conventions of the Weyl formalism as outlined in [24]. We write the contravariant four-vector  $(p^\mu) = (E, \mathbf{p})^T$ , the metric is  $(\eta_{\mu\nu}) = \text{diag}(-1, +1, +1, +1)$ , and we arrange the unit matrix and the three Pauli matrices into one vector  $(\sigma^\mu) = (\mathbb{1}_2, \boldsymbol{\sigma})$ . Then the  $2 \times 2$  matrix is defined

$$p = p_\mu(\sigma^\mu) = \begin{pmatrix} -E + p_3 & p_1 - ip_2 \\ p_1 + ip_2 & -E - p_3 \end{pmatrix}. \quad (\text{A.1})$$

For massless particles we have  $\det(p) = p^\mu p_\mu = m^2 = 0$ , that is,  $p$  has rank 1 and can be written as a product of two-component vectors, called spinors,

$$p_{a\dot{b}} = -|p\rangle_a \langle p|_{\dot{b}} \quad (\text{A.2})$$

with indices  $a, \dot{a}, b, \dot{b} \in \{1, 2\}$ . The minus sign is convention. The massive case corresponds to a matrix  $p$  with full rank and has for instance been discussed in [23]. We note that the dotted and undotted indices are simply different indices. In order to form Lorentz-invariant scalar products, antisymmetric  $\epsilon$  symbols are introduced,

$$\epsilon^{ab} = \epsilon^{\dot{a}\dot{b}} = -\epsilon_{ab} = -\epsilon_{\dot{a}\dot{b}} = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}. \quad (\text{A.3})$$

The  $\epsilon$  symbols obey

$$\epsilon_{ab}\epsilon^{bc} = \delta_a^c, \quad \epsilon_{\dot{a}\dot{b}}\epsilon^{\dot{b}\dot{c}} = \delta_{\dot{a}}^{\dot{c}}. \quad (\text{A.4})$$

Defining the spinors with upper indices

$$|p\rangle^a \equiv \epsilon^{ab}|p\rangle_b, \quad |p\rangle^{\dot{a}} \equiv \epsilon^{\dot{a}\dot{b}}\langle p|_{\dot{b}}. \quad (\text{A.5})$$

it can be shown that the following scalar products are Lorentz-invariant,

$$\langle pq \rangle \equiv \langle p|_{\dot{a}} \cdot |q\rangle^{\dot{a}} = \epsilon_{\dot{a}\dot{b}}|p\rangle^{\dot{b}} \cdot |q\rangle^{\dot{a}}, \quad [pq] \equiv [p]^a \cdot |q\rangle_a = \epsilon^{ab}|p\rangle_b \cdot |q\rangle_a. \quad (\text{A.6})$$

We set

$$p_{a\dot{b}}\epsilon^{ab}\epsilon^{\dot{a}\dot{b}} = -|p\rangle_a \langle p|_{\dot{b}}\epsilon^{ab}\epsilon^{\dot{a}\dot{b}} = -|p\rangle^{\dot{a}}[p]^b \equiv p^{\dot{a}b} \quad (\text{A.7})$$

and get

$$p_{a\dot{a}}\dot{q}^{\dot{a}a} = -\langle pq\rangle[pq] = (p+q)^2 = 2pq. \quad (\text{A.8})$$

Usually, the following scalar products are also defined,

$$\langle p|k|q\rangle \equiv \langle p|_{\dot{a}}k^{\dot{a}b}|q\rangle_b = -\langle pk\rangle[kq], \quad [p|k|q\rangle \equiv [p]^a k_{a\dot{b}}|q\rangle^{\dot{b}} = -[pk]\langle kq\rangle. \quad (\text{A.9})$$

Let us also mention some equations useful in the explicit calculation - see [24] for more details:

$$\langle ab\rangle = -\langle ba\rangle, \quad [ab] = -[ba] \quad (\text{A.10})$$

and the Shouten identities

$$\langle ab\rangle\langle cd\rangle + \langle ac\rangle\langle db\rangle + \langle ad\rangle\langle bc\rangle = 0, \quad [ab][cd] + [ac][db] + [ad][bc] = 0. \quad (\text{A.11})$$

For real momenta we have  $[p]^a = (|p\rangle^{\dot{a}})^*$  and  $\langle p|_{\dot{a}} = (|p\rangle_a)^*$  justifying the dot notation of the indices. Also useful is the context

$$|-p\rangle = -|p\rangle, \quad |-p] = +|p]. \quad (\text{A.12})$$

## References

- [1] S. J. Parke and T. R. Taylor, *An Amplitude for  $n$  Gluon Scattering*, *Phys. Rev. Lett.* **56** (1986) 2459.
- [2] R. Britto, F. Cachazo and B. Feng, *New recursion relations for tree amplitudes of gluons*, *Nucl. Phys. B* **715** (2005) 499 [[hep-th/0412308](#)].
- [3] R. Britto, F. Cachazo, B. Feng and E. Witten, *Direct proof of tree-level recursion relation in Yang-Mills theory*, *Phys. Rev. Lett.* **94** (2005) 181602 [[hep-th/0501052](#)].
- [4] W. L. van Neerven and J. A. M. Vermaseren, *Large Loop Integrals*, *Phys. Lett. B* **137** (1984) 241.
- [5] Z. Bern, L. J. Dixon, D. C. Dunbar and D. A. Kosower, *One loop  $n$  point gauge theory amplitudes, unitarity and collinear limits*, *Nucl. Phys. B* **425** (1994) 217 [[hep-ph/9403226](#)].
- [6] S. J. Bidder, N. E. J. Bjerrum-Bohr, D. C. Dunbar and W. B. Perkins, *One-loop gluon scattering amplitudes in theories with  $N < 4$  supersymmetries*, *Phys. Lett. B* **612** (2005) 75 [[hep-th/0502028](#)].
- [7] C. Anastasiou, R. Britto, B. Feng, Z. Kunszt and P. Mastrolia,  *$D$ -dimensional unitarity cut method*, *Phys. Lett. B* **645** (2007) 213 [[hep-ph/0609191](#)].
- [8] W. T. Giele, Z. Kunszt and K. Melnikov, *Full one-loop amplitudes from tree amplitudes*, *JHEP* **04** (2008) 049 [[0801.2237](#)].
- [9] R. P. Feynman, *Quantum theory of gravitation*, *Acta Phys. Polon.* **24** (1963) 697.
- [10] R. P. Feynman, *Selected papers of Richard Feynman: Closed Loop And Tree Diagrams*, *World Scientific, Singapore* (2000).
- [11] S. Catani, T. Gleisberg, F. Krauss, G. Rodrigo and J.-C. Winter, *From loops to trees by-passing Feynman's theorem*, *JHEP* **09** (2008) 065 [[0804.3170](#)].

- [12] I. Bierenbaum, S. Catani, P. Draggiotis and G. Rodrigo, *A Tree-Loop Duality Relation at Two Loops and Beyond*, *JHEP* **10** (2010) 073 [[1007.0194](#)].
- [13] C. Baadsgaard, N. E. J. Bjerrum-Bohr, J. L. Bourjaily, S. Caron-Huot, P. H. Damgaard and B. Feng, *New Representations of the Perturbative S-Matrix*, *Phys. Rev. Lett.* **116** (2016) 061601 [[1509.02169](#)].
- [14] M. Maniatis, *Scattering amplitudes abandoning virtual particles*, [1511.03574](#).
- [15] M. Maniatis, *Application of the Feynman-tree theorem together with BCFW recursion relations*, *Int. J. Mod. Phys. A* **33** (2018) 1850042 [[1609.00377](#)].
- [16] M. Maniatis, *Application of Britto-Cachazo-Feng-Witten recursion relations and the Feynman-tree theorem to the four-gluon amplitude with all plus helicities*, *Phys. Rev. D* **100** (2019) 096022 [[1906.10821](#)].
- [17] S. Caron-Huot, *Loops and trees*, *JHEP* **05** (2011) 080 [[1007.3224](#)].
- [18] V. Del Duca, L. J. Dixon and F. Maltoni, *New color decompositions for gauge amplitudes at tree and loop level*, *Nucl. Phys. B* **571** (2000) 51 [[hep-ph/9910563](#)].
- [19] P. Benincasa and F. Cachazo, *Consistency Conditions on the S-Matrix of Massless Particles*, [0705.4305](#).
- [20] Z. Bern, L. J. Dixon, D. C. Dunbar and D. A. Kosower, *One loop selfdual and N=4 superYang-Mills*, *Phys. Lett. B* **394** (1997) 105 [[hep-th/9611127](#)].
- [21] Z. Bern and D. A. Kosower, *The Computation of loop amplitudes in gauge theories*, *Nucl. Phys. B* **379** (1992) 451.
- [22] S. D. Badger, E. W. N. Glover, V. V. Khoze and P. Svrcek, *Recursion relations for gauge theory amplitudes with massive particles*, *JHEP* **07** (2005) 025 [[hep-th/0504159](#)].
- [23] N. Arkani-Hamed, T.-C. Huang and Y.-t. Huang, *Scattering Amplitudes For All Masses and Spins*, [1709.04891](#).
- [24] H. Elvang and Y.-t. Huang, *Scattering Amplitudes*, [1308.1697](#).