

Probability Distribution on Full Rooted Trees

Yuta Nakahara*, Shota Saito†, Akira Kamatsuka‡, and Toshiyasu Matsushima§

*Center for Data Science, Waseda University, Tokyo, 169-8050, Japan, yuta.nakahara@aoni.waseda.jp

†Faculty of Informatics, Gunma University, Gunma 371-8510, Japan shota.s@gunma-u.ac.jp

‡Dept. of Information Science, Shonan Institute of Technology, Kanagawa, 251-8511, Japan kamatsuka@info.shonan-it.ac.jp

§Dept. of Applied Math., Waseda University, Tokyo, 169-8555, Japan, toshimat@waseda.jp

Abstract—The recursive and hierarchical structure of full rooted trees is applicable to represent statistical models in various areas, such as data compression, image processing, and machine learning. In most of these cases, the full rooted tree is not a random variable; as such, model selection to avoid overfitting becomes problematic. A method to solve this problem is to assume a prior distribution on the full rooted trees. This enables overfitting to be avoided based on the Bayes decision theory. For example, by assigning a low prior probability to a complex model, the maximum a posteriori estimator prevents overfitting. Furthermore, overfitting can be avoided by averaging all the models weighted by their posteriors. In this paper, we propose a probability distribution on a set of full rooted trees. Its parametric representation is suitable for calculating the properties of our distribution using recursive functions, such as the mode, expectation, and posterior distribution. Although such distributions have been proposed in previous studies, they are only applicable to specific applications. Therefore, we extract their mathematically essential components and derive new generalized methods to calculate the expectation, posterior distribution, etc.

I. INTRODUCTION

A. Objective of this study

In this paper, we propose a discrete probability distribution on a set with a recursive and hierarchical structure, i.e., a finite set of full rooted trees. Mathematically, a tree is defined as a connected graph without cycles (see, e.g., [1]). A rooted tree is a tree that has one node known as a root node, and a full rooted tree is a rooted tree in which each inner node has the same number of child nodes. Subsequently, we can define a finite set of subtrees of a full rooted tree. This full rooted tree, which contains all the subtrees in the finite set, is denoted as a base tree herein.

A trivial method to define a probability distribution on this set is to assign occurrence probabilities to all subtrees and regard these values as parameters. In other words, we can define the categorical distribution on the finite set of subtrees of the base tree. However, this definition requires the same number of parameters as the subtrees, which increases in a doubly exponential order of the depth of the base tree.

Therefore, we propose an efficient parametric representation of the probability distribution on a set of subtrees. It is suitable for the recursive structure of full rooted trees and allows the number of parameters to be reduced. Moreover, it enables us to calculate its mode, expectation, posterior distribution, etc., using recursive functions. Therefore, it is efficient from a computational viewpoint. Furthermore, we

expect these recursive functions to be effective as a subroutine of the variational Bayesian method and the Markov chain Monte Carlo method in hierarchical Bayesian modeling (see, e.g., [2]).

However, our distribution has already been proposed independently in source coding and machine learning, as will be detailed in the next subsection. The novelty of our study is the extraction of the essence from the previous discussion, which depends on the applicational objects, and its representation as a clear mathematical theory. Hence, we derived new generalized recursive algorithms to calculate the expectation, posterior distribution, etc., which could not be derived in previous studies pertaining to real-world applications.

B. Examples of applications

Full rooted trees are utilized in various fields of study. For example, for text compression in information theory, a full rooted tree represents a set of contexts, which are strings of the most recent symbols at each time point, and it is known as a context tree [3]. In image processing, it represents a variable block-size segmentation, and it is known as quadtree block partitioning [4]. In machine learning, it represents a nonlinear function that comprises many conditional branches and is known as a decision tree [5]. In most of these studies, the rooted tree is not a random variable and serves as an index of a statistical model or function, i.e., one full rooted tree τ corresponds to one statistical model $p(x; \tau)$ or one function $f_\tau(x)$.

Their recursive and hierarchical structures are suitable for representing complex statistical models or functional structures. For example, the expansion of the leaf nodes represents an increase in the contexts of a context tree [3], a division of a block on the image in quadtree partitioning [4], and the addition of a conditional branch in the decision tree [5]. Such expressive capability and extensibility of full rooted trees render them widely applicable in various fields.

However, such hierarchical expressive capability causes a problem in tree selection, i.e., the selection of one statistical model or function. This is because the optimal tree under the criterion of the likelihood or squared loss for training data is inevitably the deepest one, thereby resulting in overfitting. Therefore, most previous studies applied a stopping rule for node expansion [4], [5], introduced a normalization term into the objective function [6], or averaged the statistical models or

the functions with some weights [3], [6], [7]. However, these algorithmic modifications are heuristic at times.

A theoretical method to solve this problem is to consider the full rooted tree as a random variable and assuming a prior distribution on it. An appropriate prior distribution provides a unified method for selecting one full rooted tree or combining them based on the Bayes decision theory (see, e.g., [8]). Although the Bayes decision theory is typically applied to statistical models with unknown continuous parameters, it is also applicable to statistical models with unknown discrete random variables such as full rooted trees (see, e.g., [9]). By assigning a high prior probability to a shallow tree and a low prior probability to a deep tree, we can avoid overfitting the statistical model.

As mentioned above, most previous studies regard the full rooted tree as a non-stochastic variable. However, few studies adopted the above mentioned approach. In terms of text compression, the complete Bayesian interpretation of context tree weighting method was first investigated by the authors of [10]. Not only the theory, but also the associated algorithm has been improved over the decade it was first investigated (see, e.g., [11]). Moreover, similar results obtained from rich real data analysis have been reported recently [12], [13] (Note that the prior form reported in [12], [13] is extremely restricted and cannot be updated as a posterior, in contrast to that reported in [10], [11]). In image processing, the author of [14] were the first to regard the quadtree as a stochastic model, and its optimal estimation was derived under the Bayes criteria. In machine learning, the authors of [15] redefined the decision tree as a stochastic generative model and improved most tree weighting methods (e.g., [7]).

However, these studies depend on specific data or generative models. This might have been the reason that more than 25 years had passed before the first study [10] pertaining to text compression was applied to image processing [14] and machine learning [15]. Therefore, we separated the mathematically essential component of the discussion from the modifiable component based on specific data or the generative model. This theoretically expands the potential application of probability distributions on full rooted trees. Subsequently, we derived new generalized methods to evaluate the characteristics of the probability distribution on full rooted trees, which have not been performed in previous studies. More precisely, only Theorems 1 and 3 and Corollary 2 has been used in previous studies. Meanwhile, the other methods expand the possibility of the applying the probability distribution on full rooted trees.

C. Organization of this paper

The reminder of this paper is organized as follows: In Section II, we present the notations used herein. In Section III, we define the prior on full rooted trees. In Section IV, we describe the algorithms for calculating the properties of the proposed distribution, e.g., a marginal distribution for each node, an efficient calculation of the expectation, mode, and the




Fig. 1. The notations for the rooted trees.

posterior distribution. In Section V, we propose some future work. In Section VI, we conclude the paper.

II. NOTATIONS USED FOR FULL ROOTED TREES

In this section, we define notations for the rooted trees. They are shown in Fig. 1. Let $k \in \mathbb{N}$ denote the maximum number of child nodes and $d_{\max} \in \mathbb{N}$ denote the maximum depth. Let $\tau_p = (\mathcal{V}_p, \mathcal{E}_p)$ denote the perfect¹ k -ary rooted tree whose depth is d_{\max} and root node is v_λ . \mathcal{V}_p and \mathcal{E}_p denote the set of the nodes and edges of it, respectively. Then, let $\mathcal{I}_p \subset \mathcal{V}_p$, and $\mathcal{L}_p \subset \mathcal{V}_p$ denote the set of the inner nodes and the leaf nodes of τ_p , respectively. For each node $v \in \mathcal{V}_p$, let $d_v \in \mathbb{N}$ denote the depth of $v \in \mathcal{V}_p$ and $\text{Ch}_p(v) \subset \mathcal{V}_p$ denote the set of child nodes of v on τ_p . Notations about the relation between two nodes $v, v' \in \mathcal{V}_p$ are as follows. Let $v \succ v'$ denote that v is an ancestor node of v' , (v' is a descendant node of v), $v \succeq v'$ denote that v is an ancestor node of v' or v' itself, (v' is a descendant node of v or v itself), $\text{An}(v) := \{v' \in \mathcal{V}_p \mid v' \succ v\}$, and $\text{Dep}(v) := \{v' \in \mathcal{V}_p \mid v \succ v'\}$.

Subsequently, we consider rooted subtrees of τ_p in which their root nodes are the same as v_λ and all inner nodes have exactly k children. They are called full rooted subtrees and τ_p is called a base tree. Let \mathcal{T} denote the set of all full rooted subtrees of τ_p . Let \mathcal{V}_τ and \mathcal{E}_τ denote the set of the nodes and the edges of $\tau \in \mathcal{T}$, respectively. Let $\mathcal{I}_\tau \subset \mathcal{V}_\tau$, and $\mathcal{L}_\tau \subset \mathcal{V}_\tau$ denote the set of the inner nodes and the leaf nodes of $\tau \in \mathcal{T}$, respectively.

III. DEFINITION OF PROBABILITY DISTRIBUTION ON FULL ROOTED SUBTREES

In this section, we define a probability distribution on full rooted subtrees \mathcal{T} . Let T denote the random variable on \mathcal{T} , and τ denote its realized value.

Definition 1: For $(\alpha_v)_{v \in \mathcal{V}_p} \in [0, 1]^{|\mathcal{V}_p|}$, we define probability distribution $p(\tau)$ on \mathcal{T} as below.

$$p(\tau) := \prod_{v \in \mathcal{I}_\tau} \alpha_v \prod_{v' \in \mathcal{L}_\tau} (1 - \alpha_{v'}), \quad (1)$$

where $\alpha_v = 0$ for $v \in \mathcal{L}_p$.

Intuitively, α_v represents the probability that v has child nodes under the condition that v is contained in the tree. It will be proved as a theoretical fact in Remark 2.

¹“Perfect” means that all inner nodes have exactly k children and all leaf nodes have the same depth.




Fig. 2. An example of the probability distribution on full rooted subtrees. Here, $k = 2$ and $d_{\max} = 2$. The parameters of the distribution are in the upper right figure. When $k = 2$ and $d_{\max} = 2$, $|\mathcal{T}| = 5$. The probability of each full rooted subtree is calculated under the graph of it.

Example 1: An example of the probability distribution on full rooted subtrees for $k = 2$ and $d_{\max} = 2$ is shown in Fig. 2.

Theorem 1: The quantity $p(\tau)$ defined as in (1) fulfills the condition of the probability distribution, that is, $\sum_{\tau \in \mathcal{T}} p(\tau) = 1$.

Example 2: Before the proof of Theorem 1 for the general case, we describe an example where $d_{\max} = 2$ and $k = 2$ (see Fig. 2). First, we factorize the sum as below.

$$\begin{aligned}
& \sum_{\tau \in \mathcal{T}} p(\tau) \\
&= (1 - \alpha_{v_\lambda}) \\
&\quad + \alpha_{v_\lambda}(1 - \alpha_{v_0})(1 - \alpha_{v_1}) \\
&\quad + \alpha_{v_\lambda}\alpha_{v_0}(1 - \alpha_{v_{00}})(1 - \alpha_{v_{01}})(1 - \alpha_{v_1}) \\
&\quad + \alpha_{v_\lambda}(1 - \alpha_{v_0})\alpha_{v_1}(1 - \alpha_{v_{10}})(1 - \alpha_{v_{11}}) \\
&\quad + \alpha_{v_\lambda}\alpha_{v_0}(1 - \alpha_{v_{00}})(1 - \alpha_{v_{01}})\alpha_{v_1}(1 - \alpha_{v_{10}})(1 - \alpha_{v_{11}}) \quad (2) \\
&= (1 - \alpha_{v_\lambda}) \\
&\quad + \alpha_{v_\lambda} \left\{ (1 - \alpha_{v_0})(1 - \alpha_{v_1}) \right. \\
&\quad \quad + \alpha_{v_0}(1 - \alpha_{v_{00}})(1 - \alpha_{v_{01}})(1 - \alpha_{v_1}) \\
&\quad \quad + (1 - \alpha_{v_0})\alpha_{v_1}(1 - \alpha_{v_{10}})(1 - \alpha_{v_{11}}) \\
&\quad \quad \left. + \alpha_{v_0}(1 - \alpha_{v_{00}})(1 - \alpha_{v_{01}}) \right. \\
&\quad \quad \quad \times \alpha_{v_1}(1 - \alpha_{v_{10}})(1 - \alpha_{v_{11}}) \left. \right\} \quad (3) \\
&= (1 - \alpha_{v_\lambda}) \\
&\quad + \alpha_{v_\lambda} \left\{ (1 - \alpha_{v_0})[(1 - \alpha_{v_1}) + \alpha_{v_1}(1 - \alpha_{v_{10}})(1 - \alpha_{v_{11}})] \right. \\
&\quad \quad \left. + \alpha_{v_0}(1 - \alpha_{v_{00}})(1 - \alpha_{v_{01}}) \right\}
\end{aligned}$$

$$\begin{aligned}
& \times [(1 - \alpha_{v_1}) + \alpha_{v_1}(1 - \alpha_{v_{10}})(1 - \alpha_{v_{11}})] \left. \right\} \quad (4) \\
&= (1 - \alpha_{v_\lambda}) \\
&\quad + \alpha_{v_\lambda} \left\{ [(1 - \alpha_{v_0}) + \alpha_{v_0}(1 - \alpha_{v_{00}})(1 - \alpha_{v_{01}})] \right. \\
&\quad \quad \times [(1 - \alpha_{v_1}) + \alpha_{v_1}(1 - \alpha_{v_{10}})(1 - \alpha_{v_{11}})] \left. \right\}. \quad (5)
\end{aligned}$$

Here, $\alpha_{v_{00}} = \alpha_{v_{01}} = \alpha_{v_{10}} = \alpha_{v_{11}} = 0$ since $v_{00}, v_{01}, v_{10}, v_{11} \in \mathcal{L}_p$. Then,

$$(5) = (1 - \alpha_{v_\lambda}) + \alpha_{v_\lambda} \left\{ [(1 - \alpha_{v_0}) + \alpha_{v_0}] \cdot [(1 - \alpha_{v_1}) + \alpha_{v_1}] \right\} \quad (6)$$

$$= (1 - \alpha_{v_\lambda}) + \alpha_{v_\lambda} \quad (7)$$

$$= 1. \quad (8)$$

The general proof of Theorem 1 is in the following. We will first prove Lemma 1, which is the essential lemma since it is not used only in the proof of Theorem 1 but also in the proof of other theorems later.

Lemma 1: Let $F : \mathcal{T} \rightarrow \mathbb{R}$ be a real-valued function on the set \mathcal{T} of the full rooted subtrees of the base tree τ_p . If F has the form

$$F(\tau) = \prod_{v \in \mathcal{I}_\tau} G(v) \prod_{v' \in \mathcal{L}_\tau} H(v'), \quad (9)$$

where $G : \mathcal{V}_p \rightarrow \mathbb{R}$ and $H : \mathcal{V}_p \rightarrow \mathbb{R}$ are real-valued functions on \mathcal{V}_p , then the summation $\sum_{\tau \in \mathcal{T}} F(\tau)$ can be recursively decomposed as follows.

$$\sum_{\tau \in \mathcal{T}} F(\tau) = \phi(v_\lambda), \quad (10)$$




Fig. 3. The example of trees in $\{[v_\lambda]\}$ and $\mathcal{T} \setminus \{[v_\lambda]\}$, where $k = 2$. The left side shows the structure of the tree in $\{[v_\lambda]\}$. There is only one tree $[v_\lambda]$. The right side shows the structure of the trees in $\mathcal{T} \setminus \{[v_\lambda]\}$. All of them have the root node v_λ as its inner node. The other structure is determined by choosing subtrees from $\mathcal{T}^{v_{\lambda 0}}$ and $\mathcal{T}^{v_{\lambda 1}}$.

where $\phi : \mathcal{V}_p \rightarrow \mathbb{R}$ is defined as below.

$$\phi(v) := \begin{cases} H(v), & v \in \mathcal{L}_p, \\ H(v) + G(v) \prod_{v' \in \text{Ch}_p(v)} \phi(v'), & v \in \mathcal{I}_p. \end{cases} \quad (11)$$

Proof: Let $[v_\lambda]$ denote the tree that consists of only the root node v_λ of the base tree τ_p . Then, the cases of the sum is divided as follows.

$$\sum_{\tau \in \mathcal{T}} F(\tau) = \sum_{\tau \in \mathcal{T}} \prod_{v \in \mathcal{I}_\tau} G(v) \prod_{v' \in \mathcal{L}_\tau} H(v') \quad (12)$$

$$= \prod_{v \in \mathcal{I}_{[v_\lambda]}} G(v) \prod_{v' \in \mathcal{L}_{[v_\lambda]}} H(v') + \sum_{\tau \in \mathcal{T} \setminus \{[v_\lambda]\}} \prod_{v \in \mathcal{I}_\tau} G(v) \prod_{v' \in \mathcal{L}_\tau} H(v') \quad (13)$$

$$= H(v_\lambda) + \sum_{\tau \in \mathcal{T} \setminus \{[v_\lambda]\}} \prod_{v \in \mathcal{I}_\tau} G(v) \prod_{v' \in \mathcal{L}_\tau} H(v') \quad (14)$$

$$= H(v_\lambda) + G(v_\lambda) \sum_{\tau \in \mathcal{T} \setminus \{[v_\lambda]\}} \prod_{v \in \mathcal{I}_\tau \setminus \{v_\lambda\}} G(v) \prod_{v' \in \mathcal{L}_\tau} H(v'), \quad (15)$$

where (14) is because $[v_\lambda]$ has no inner node and its leaf node is only v_λ ; (15) is because every tree in $\mathcal{T} \setminus [v_\lambda]$ has v_λ and the corresponding factor $G(v_\lambda)$.

We have already pointed out that each tree $\tau \in \mathcal{T} \setminus \{[v_\lambda]\}$ contains v_λ as its inner node. The other structure of τ is determined by the shape of k subtrees whose root nodes are the child nodes of v_λ (see Fig. 3). We index them in an appropriate order. Then, let $v_{\lambda i}$ denote the i -th child node of v_λ for $i \in \{0, 1, \dots, k-1\}$, i.e. $\{v_{\lambda 0}, \dots, v_{\lambda k-1}\} = \text{Ch}_p(v_\lambda)$. Let $\mathcal{T}^{v_{\lambda i}}$ denote the set of subtrees whose root node is $v_{\lambda i}$. Then, there is a natural bijection from $\mathcal{T} \setminus \{[v_\lambda]\}$ to $\mathcal{T}^{v_{\lambda 0}} \times \dots \times \mathcal{T}^{v_{\lambda k-1}}$. Therefore, the summation of (15) is

further factorized. Consequently, we have

$$(15) = H(v_\lambda) + G(v_\lambda) \times \sum_{(\tau_0, \dots, \tau_{k-1}) \in \mathcal{T}^{v_{\lambda 0}} \times \dots \times \mathcal{T}^{v_{\lambda k-1}}} \left[\prod_{v \in \mathcal{I}_{\tau_0}} G(v) \prod_{v' \in \mathcal{L}_{\tau_0}} H(v') \dots \prod_{v \in \mathcal{I}_{\tau_{k-1}}} G(v) \prod_{v' \in \mathcal{L}_{\tau_{k-1}}} H(v') \right] \quad (16)$$

$$= H(v_\lambda) + G(v_\lambda)$$

$$\times \sum_{\tau_0 \in \mathcal{T}^{v_{\lambda 0}}} \dots \sum_{\tau_{k-1} \in \mathcal{T}^{v_{\lambda k-1}}} \left[\prod_{v \in \mathcal{I}_{\tau_0}} G(v) \prod_{v' \in \mathcal{L}_{\tau_0}} H(v') \dots \prod_{v \in \mathcal{I}_{\tau_{k-1}}} G(v) \prod_{v' \in \mathcal{L}_{\tau_{k-1}}} H(v') \right] \quad (17)$$

$$= H(v_\lambda) + G(v_\lambda) \prod_{i=0}^{k-1} \sum_{\tau \in \mathcal{T}^{v_{\lambda i}}} \prod_{v \in \mathcal{I}_\tau} G(v) \prod_{v' \in \mathcal{L}_\tau} H(v'). \quad (18)$$

Then, from (12) and (18), we have

$$\underbrace{\sum_{\tau \in \mathcal{T}} \prod_{v \in \mathcal{I}_\tau} G(v) \prod_{v' \in \mathcal{L}_\tau} H(v')}_{(a)} = H(v_\lambda) + G(v_\lambda) \underbrace{\prod_{i=0}^{k-1} \sum_{\tau \in \mathcal{T}^{v_{\lambda i}}} \prod_{v \in \mathcal{I}_\tau} G(v) \prod_{v' \in \mathcal{L}_\tau} H(v')}_{(b)}. \quad (19)$$

The underbraced parts (a) and (b) have the same structure except for the depth of the root node of the subtree. Therefore, (b) can be decomposed in a similar manner from (12) to (18). We can continue this decomposition to the leaf nodes.

Then, let \mathcal{T}^v denote the set of subtrees whose root node is $v \in \mathcal{V}_p$ in general, i.e., we define a notion similar to $\mathcal{T}^{v_{\lambda i}}$ for not only $v_{\lambda 0}, v_{\lambda 1}, \dots, v_{\lambda k-1}$ but also any other nodes $v \in \mathcal{V}_p$. Finally, we have an alternative definition of $\phi(v) : \mathcal{V}_p \rightarrow \mathbb{R}$, which is equivalent to (11).

$$\phi(v) := \sum_{\tau \in \mathcal{T}^v} \prod_{v' \in \mathcal{I}_\tau} G(v') \prod_{v'' \in \mathcal{L}_\tau} H(v''), \quad (20)$$

The equivalence is confirmed by substituting it into both sides of (19). Therefore, Lemma 1 is proved. \blacksquare

Then, the proof of Theorem 1 is as follows.

Proof: Using Lemma 1, we can divide the cases of the sum and factorize the common terms of $\sum_{\tau \in \mathcal{T}} p(\tau)$ in the following recursive manner.

$$\sum_{\tau \in \mathcal{T}} p(\tau) = \phi(v_\lambda), \quad (21)$$

where

$$\phi(v) := \begin{cases} 1 - \alpha_v, & v \in \mathcal{L}_p, \\ (1 - \alpha_v) + \alpha_v \prod_{v' \in \text{Ch}_p(v)} \phi(v'), & v \in \mathcal{I}_p. \end{cases} \quad (22)$$

Then, we prove $\phi(v) = 1$ for any node $v \in \mathcal{V}_p$ by structural induction. For any leaf node $v \in \mathcal{L}_p$, $\alpha_v = 0$ from Definition 1. Therefore,

$$\phi(v) = 1 - \alpha_v = 1, \quad v \in \mathcal{L}_p. \quad (23)$$

For any inner node $v \in \mathcal{I}_p$, assuming $\phi(v') = 1$ as the induction hypothesis for any descendant nodes $v' \in \text{De}_p(v)$,

$$\phi(v) = (1 - \alpha_v) + \alpha_v \prod_{v' \in \text{Ch}_p(v)} \phi(v') \quad (24)$$

$$= (1 - \alpha_v) + \alpha_v \prod_{v' \in \text{Ch}_p(v)} 1 \quad (25)$$

$$= 1. \quad (26)$$

Therefore, $\sum_{\tau \in \mathcal{T}} p(\tau) = \phi(v_\lambda) = 1$ since v_λ is also in \mathcal{V}_p . ■

Remark 1: Although Theorem 1 is also proved in [14] and [15], we extract the essential part of them as Lemma 1. In [12], [13], a restricted case of Theorem 1 is proved, in which α_v has a common value for all $v \in \mathcal{I}_p$.

IV. PROPERTIES OF PROBABILITY DISTRIBUTION ON FULL ROOTED SUBTREES

In this section, we describe properties of the probability distribution on full rooted subtrees and methods to calculate them. All the proofs are in Appendix A.

A. Probability of events on nodes

At the beginning, we explain why $v \in \mathcal{V}_T$ is a probabilistic event. We consider any $v \in \mathcal{V}_p$ is given and fixed. After that, a full rooted subtree is randomly chosen according to the probability distribution proposed in Section III. Then, T sometimes contains v and sometimes not, depending on its realized value τ . Therefore, $v \in \mathcal{V}_T$ is a probabilistic event on $p(\tau)$. In the following, we describe probabilities of such events.

Theorem 2: For any $v \in \mathcal{V}_p$, we have the following:

$$\Pr\{v \in \mathcal{V}_T\} = \prod_{v' \in \text{An}(v)} \alpha_{v'}, \quad (27)$$

$$\Pr\{v \in \mathcal{I}_T\} = \alpha_v \prod_{v' \in \text{An}(v)} \alpha_{v'}, \quad (28)$$

$$\Pr\{v \in \mathcal{L}_T\} = (1 - \alpha_v) \prod_{v' \in \text{An}(v)} \alpha_{v'}. \quad (29)$$

Example 3: Let us consider $p(\tau)$ shown in Fig. 2. Trivially, $\Pr\{v_{01} \in \mathcal{V}_T\}$, $\Pr\{v_1 \in \mathcal{I}_T\}$, and $\Pr\{v_0 \in \mathcal{L}_T\}$ are calculated as

$$\Pr\{v_{01} \in \mathcal{V}_T\} = p(\tau_2) + p(\tau_4) = 0.28, \quad (30)$$

$$\Pr\{v_1 \in \mathcal{I}_T\} = p(\tau_3) + p(\tau_4) = 0.56, \quad (31)$$

$$\Pr\{v_0 \in \mathcal{L}_T\} = p(\tau_1) + p(\tau_3) = 0.42. \quad (32)$$

The same probabilities are also given by

$$\Pr\{v_{01} \in \mathcal{V}_T\} = \alpha_{v_\lambda} \alpha_{v_0} = 0.28, \quad (33)$$

$$\Pr\{v_1 \in \mathcal{I}_T\} = \alpha_{v_\lambda} \alpha_{v_1} = 0.56, \quad (34)$$

$$\Pr\{v_0 \in \mathcal{L}_T\} = \alpha_{v_\lambda} (1 - \alpha_{v_1}) = 0.42. \quad (35)$$

Remark 2: Probabilities of many other events on nodes are derived from Theorem 2. For example,

$$\Pr\{v \in \mathcal{I}_T \mid v \in \mathcal{V}_T\} = \frac{\Pr\{v \in \mathcal{I}_T \wedge v \in \mathcal{V}_T\}}{\Pr\{v \in \mathcal{V}_T\}} \quad (36)$$

$$= \frac{\Pr\{v \in \mathcal{I}_T\}}{\Pr\{v \in \mathcal{V}_T\}} \quad (37)$$

$$= \alpha_v. \quad (38)$$

B. Mode

We describe an algorithm to find the mode of $p(\tau)$ with $O(k^{d_{\max}+1})$ computational cost.² Note that, the size of search space \mathcal{T} is of the order of $\Omega(2^{k^{d_{\max}-2}})$ in general.³ First, replacing all the sum in the proof of Lemma 1 for the max, we can derive the following recursive expression of $\max_{\tau \in \mathcal{T}} p(\tau)$.

Proposition 1:

$$\max_{\tau \in \mathcal{T}} p(\tau) = \psi(v_\lambda), \quad (39)$$

where

$$\psi(v) := \begin{cases} 1 - \alpha_v = 1, & v \in \mathcal{L}_p, \\ \max \left\{ 1 - \alpha_v, \alpha_v \prod_{v' \in \text{Ch}_p(v)} \psi(v') \right\}, & v \in \mathcal{I}_p. \end{cases} \quad (40)$$

Example 4: On $p(\tau)$ shown in Fig. 2, the maximum probability is $p(\tau_3) = 0.336$. It is also calculated as follows.

$$\max\{1 - \alpha_{v_\lambda}, \alpha_{v_\lambda} \max\{1 - \alpha_{v_0}, \alpha_{v_0} \max\{1 - \alpha_{v_1}, \alpha_{v_1}\}\}\} \quad (41)$$

$$= \max\{0.3, 0.7 \max\{0.6, 0.4\} \max\{0.2, 0.8\}\} \quad (42)$$

$$= \max\{0.3, 0.336\} = 0.336. \quad (43)$$

In addition, we define a flag variable $\delta_v \in \{0, 1\}$ as follows.

Definition 2: For any $v \in \mathcal{V}_p$, we define

$$\delta_v := \begin{cases} 1, & 1 - \alpha_v < \alpha_v \prod_{v' \in \text{Ch}_p(v)} \psi(v'), \\ 0, & \text{otherwise.} \end{cases} \quad (44)$$

We can calculate $\psi(v)$ and δ_v simultaneously. Then, the mode of $p(\tau)$ is given by the following proposition.

Proposition 2: $\arg \max_{\tau \in \mathcal{T}} p(\tau)$ is identified as the tree that satisfies

$$v \in \mathcal{I}_\tau \Rightarrow \delta_v = 1, \quad (45)$$


$$v \in \mathcal{L}_\tau \Rightarrow \delta_v = 0. \quad (46)$$

Then, the following theorem holds.

Theorem 3: The mode of $p(\tau)$ can be found by backtracking search from v_λ after the calculation of $\psi(v)$ and δ_v . It is detailed in Algorithm 1 in Appendix B.

Remark 3: In [12], [13], Papageorgiou et al. proposed the same algorithm as Algorithm 1 as well as an algorithm to find multiple most likely trees on the background of text compression.

Flag variable calculation



Backtracking




Fig. 4. An example of the mode calculation. The parameters are the same as those in Fig. 2 and shown in the lower right figure. Figures on the upper side show the process of the calculation of the flag variable δ_v , which is determined from leaf nodes in order. Figures on the lower side show the process of backtracking. If $\delta_v = 1$, expand the edge. If $\delta_v = 0$, stop the expansion.

Example 5: See Fig. 4. The parameters are the same as those in Fig. 2. The mode τ_3 is found by the proposed algorithm.

C. Expectation

Let $f : \mathcal{T} \rightarrow \mathbb{R}$ denote a real-valued function on \mathcal{T} . Here, we discuss sufficient conditions of f , under which the following expectation can be calculated efficiently with $O(k^{d_{\max}+1})$ cost.

$$\mathbb{E}[f(T)] := \sum_{\tau \in \mathcal{T}} f(\tau) p(\tau). \quad (47)$$

Note that the size of \mathcal{T} is of the order of $\Omega(2^{k^{d_{\max}-2}})$ in general.

Condition 1: There exist $g : \mathcal{V}_p \rightarrow \mathbb{R}$ and $h : \mathcal{V}_p \rightarrow \mathbb{R}$ such that

$$f(\tau) = \prod_{v \in \mathcal{I}_\tau} g(v) \prod_{v' \in \mathcal{L}_\tau} h(v'). \quad (48)$$

Theorem 4: Under Condition 1, we define a recursive function $\phi : \mathcal{V}_p \rightarrow \mathbb{R}$ as

$$\phi(v) := \begin{cases} h(v), & v \in \mathcal{L}_p, \\ (1 - \alpha_v)h(v) + \alpha_v g(v) \prod_{v' \in \text{Ch}_p(v)} \phi(v'), & v \in \mathcal{I}_p. \end{cases} \quad (49)$$

Then, we can calculate $\mathbb{E}[f(T)]$ as $\mathbb{E}[f(T)] = \phi(v_\lambda)$.

Example 6: Theorem 2 can be regarded examples of Theorem 4.

² $O(\cdot)$ denotes the Big-O notation, i.e., $f(n) = O(g(n))$ means that $\exists k > 0, \exists n_0 > 0, \forall n > n_0, |f(n)| \leq k \cdot g(n)$.

³ $\Omega(\cdot)$ denote the Big-Omega notation in complexity theory, i.e., $f(n) = \Omega(g(n))$ means that $\exists k > 0, \exists n_0 > 0, \forall n > n_0, f(n) \geq k \cdot g(n)$. $|\mathcal{T}| = \Omega(2^{k^{d_{\max}-2}})$ is proved by substituting $G(v) \equiv H(v) \equiv 1$ in Lemma 1.

Condition 2: There exist $g : \mathcal{V}_p \rightarrow \mathbb{R}$ and $h : \mathcal{V}_p \rightarrow \mathbb{R}$ such that

$$f(\tau) = \sum_{v \in \mathcal{I}_\tau} g(v) + \sum_{v' \in \mathcal{L}_\tau} h(v'). \quad (50)$$

Theorem 5: Under Condition 2, we define a recursive function $\xi : \mathcal{V}_p \rightarrow \mathbb{R}$ as

$$\xi(v) := \begin{cases} h(v), & v \in \mathcal{L}_p, \\ (1 - \alpha_v)h(v) + \alpha_v \left(g(v) + \sum_{v' \in \text{Ch}_p(v)} \xi(v') \right), & v \in \mathcal{I}_p. \end{cases} \quad (51)$$

Then, we can calculate $\mathbb{E}[f(T)]$ as $\mathbb{E}[f(T)] = \xi(v_\lambda)$.

Remark 4: Theorem 5 is useful to calculate the Shannon entropy of $p(\tau)$. It is described in Section IV-D.

D. Shannon entropy

Corollary 1: Substituting $g(v) = -\log \alpha_v$ and $h(v) = -\log(1 - \alpha_v)$ into (51), the Shannon entropy $H[T] := -\sum_{\tau \in \mathcal{T}} p(\tau) \log p(\tau)$ can be recursively calculated as follows.

$$H[T] = \xi(v_\lambda), \quad (52)$$

where

$$\xi(v) := \begin{cases} 0, & v \in \mathcal{L}_p, \\ -(1 - \alpha_v) \log(1 - \alpha_v) + \alpha_v \left(-\log \alpha_v + \sum_{v' \in \text{Ch}_p(v)} \xi(v') \right), & v \in \mathcal{I}_p. \end{cases} \quad (53)$$

Remark 5: Kullback-Leibler divergence between two tree distributions $p(\tau)$ and $p'(\tau)$ can be calculated in a similar

manner to Corollary 1. This fact may be useful for Variational Bayesian inference, in which the Kullback-Leibler divergence is minimized. This is a future work.

E. Conjugate prior of $p(\tau|\alpha)$

Here, we consider that $\alpha_v \in [0, 1]$ is also a realized value of a random variable. Let α denote $\{\alpha_v\}_{v \in \mathcal{V}_p}$, and we describe $p(\tau)$ as $p(\tau|\alpha)$ to emphasize the dependency of α in the following theorem. Then, a conjugate prior for $p(\tau|\alpha)$ is as follows.

Theorem 6: The following probability distribution is a conjugate prior for $p(\tau|\alpha)$.

$$p(\alpha) := \prod_{v \in \mathcal{V}_p} \text{Beta}(\alpha_v | \beta_v, \gamma_v), \quad (54)$$

where $\text{Beta}(\cdot | \beta_v, \gamma_v)$ denotes the probability density function of the beta distribution whose parameters are β_v and γ_v . More precisely,

$$p(\alpha|\tau) = \prod_{v \in \mathcal{V}_p} \text{Beta}(\alpha_v | \beta_{v|\tau}, \gamma_{v|\tau}), \quad (55)$$

where

$$\beta_{v|\tau} := \begin{cases} \beta_v + 1, & v \in \mathcal{I}_\tau, \\ \beta_v, & \text{otherwise,} \end{cases} \quad (56)$$

$$\gamma_{v|\tau} := \begin{cases} \gamma_v + 1, & v \in \mathcal{L}_\tau, \\ \gamma_v, & \text{otherwise.} \end{cases} \quad (57)$$

F. $p(\tau)$ as conjugate prior

We define another random variable X on a set \mathcal{X} and assume X depends on T , i.e., it follows a distribution $p(x|\tau)$. Here, we discuss a sufficient condition of $p(x|\tau)$, under which $p(\tau)$ becomes a conjugate prior for it and we can efficiently calculate the posterior $p(\tau|x)$.

Condition 3: There exist two functions $g : \mathcal{V}_p \times \mathcal{X} \rightarrow \mathbb{R}$ and $h : \mathcal{V}_p \times \mathcal{X} \rightarrow \mathbb{R}$, and $p(x|\tau)$ has the following form.

$$p(x|\tau) = \prod_{v \in \mathcal{I}_\tau} g(x, v) \prod_{v' \in \mathcal{L}_\tau} h(x, v'). \quad (58)$$

Note that g and h are not necessarily probability density functions.

Example 7: For given $\mu_1, \mu_2 \in \mathbb{R}$ and $\sigma_1, \sigma_2 \in \mathbb{R}_{>0}$, let $\mathcal{N}(x|\mu_1, \sigma_1^2)$ and $\mathcal{N}(x|\mu_2, \sigma_2^2)$ denote the probability density functions of the normal distributions governed by them. Let $\mathbf{x} := (x_v)_{v \in \mathcal{V}_p}$. If we assume

$$g(x, v) = \mathcal{N}(x_v | \mu_1, \sigma_1^2), \quad (59)$$

$$h(x, v) = \prod_{v' \in \text{Dep}(v) \cup \{v\}} \mathcal{N}(x_{v'} | \mu_2, \sigma_2^2), \quad (60)$$

we can construct $p(\mathbf{x}|\tau)$ that satisfies Condition 3. In other words, the elements of the $|\mathcal{V}_p|$ dimensional vector \mathbf{x} follows the mixture of two normal distributions and either of the two is chosen by τ .

Theorem 7: Under Condition 3, we define $q(x|v)$ and $\alpha_{v|x}$ as follows.

$$q(x|v) := \begin{cases} h(x, v), & v \in \mathcal{L}_p, \\ (1 - \alpha_v)h(x, v) \\ + \alpha_v g(x, v) \prod_{v' \in \text{Ch}_p(v)} q(x|v'), & v \in \mathcal{I}_p, \end{cases} \quad (61)$$

$$\alpha_{v|x} := \begin{cases} \alpha_v, & v \in \mathcal{L}_p, \\ \frac{\alpha_v g(x, v) \prod_{v' \in \text{Ch}_p(v)} q(x|v')}{q(x|v)}, & v \in \mathcal{I}_p. \end{cases} \quad (62)$$

Note that $\alpha_v = 0$ for $v \in \mathcal{L}_p$ (see Definition 1). Then, the posterior $p(\tau|x)$ is represented as follows.

$$p(\tau|x) = \prod_{v \in \mathcal{I}_\tau} \alpha_{v|x} \prod_{v' \in \mathcal{L}_\tau} (1 - \alpha_{v'|x}). \quad (63)$$

It should be noted that the calculation of $q(x|v)$ and $\alpha_{v|x}$ requires $O(k^{d_{\max}+1})$ cost while it requires $\Omega(2^{k^{d_{\max}-2}})$ cost in general.

Moreover, if we assume the following condition stronger than Condition 3, we can calculate the posterior $p(\tau|x)$ more efficiently with $O(d_{\max})$ cost.

Condition 4: In addition to Condition 3, we assume that there exists a path from v_λ to a leaf node $v_{\text{end}} \in \mathcal{L}_p$ and

$$g(x, v) \equiv 1, \quad (64)$$

$$h(x, v) := h'(x, v)^{I\{v \succeq v_{\text{end}}\}}, \quad (65)$$

where $I\{\cdot\}$ denotes the indicator function. In other words, only $h(x, v)$ on the path from v_λ to v_{end} takes a value different of 1.

Corollary 2: Under Condition 4, $q(x|v)$ and $\alpha_{v|x}$ are calculated as follows, more efficiently than (61) and (62).

$$q(x|v) = \begin{cases} h'(x, v), & v = v_{\text{end}}, \\ (1 - \alpha_v)h'(x, v) + \alpha_v q(x|v_{\text{ch}}), & v \succ v_{\text{end}}, \\ 1, & \text{otherwise,} \end{cases} \quad (66)$$

$$\alpha_{v|x} = \begin{cases} \alpha_v, & v \not\succ v_{\text{end}}, \\ \frac{\alpha_v q(x|v_{\text{ch}})}{q(x|v)}, & v \succ v_{\text{end}}, \end{cases} \quad (67)$$

where v_{ch} is a child node of v on the path from v_λ to v_{end} . Note that we need not calculate $q(x|v)$ for $v \not\succ v_{\text{end}}$ to update the posterior and it costs only $O(d_{\max})$.

Remark 6: Condition 4 is effective to represent a generation of sequential data x_1, x_2, \dots, x_N , in which there exists a path from root node v_λ to a leaf node $v_{\text{end}}^n \in \mathcal{L}_p$ for each $n \in \{1, 2, \dots, N\}$. (v_{end}^n and $v_{\text{end}}^{n'}$ may different each other for $n \neq n'$.) The remarkable previous studies using Corollary 2 are [10]–[15] (In [12], [13], only (66) is used but (67) is not). In other words, they treat only the case under Condition 4. The other theorems in this paper have potential applications to broader fields of study.

V. FUTURE WORK

Since the present study is a theoretical study, the theorems derived will be applied in future studies. Theorems 1 and 3 and Corollary 2 have been used in previous studies [10]–[15]. Therefore, the other theorems can be applied.

In this study, we did not use approximative algorithms such as the variational Bayes or Markov chain Monte Carlo method. Such algorithms are required for learning hierarchical models that contain the probability distribution on full rooted subtrees. The methods proposed herein may serve as a subroutine. The expansion of our methods to approximative algorithms is another future work.

In this study, the class of trees is restricted to that of full trees, in which every inner node has the same number of child nodes. Hence, another the generalization of the class to that of any rooted tree can be considered in future studies.

VI. CONCLUSION

In this paper, we discuss the probability distribution on full rooted subtrees. Although such a distribution has been used in many fields of studies, such as information theory [10]–[13], image processing [14], and machine learning [15], it depends significantly on the specific applications and data generative models. By contrast, we discussed it theoretically, collectively, and independently from a specific data generative model. Subsequently, we derived new generalized methods to evaluate the characteristics of the probability distribution on full rooted subtrees, which have not been performed in previous studies. The derived methods are efficient for calculating the events on the nodes, the mode, the expectation, the Shannon entropy, and the posterior distribution for full rooted subtrees. Therefore, this study expands the possibility of the applying the probability distribution on full rooted subtrees.

ACKNOWLEDGMENT

This work was supported by JSPS KAKENHI Grant Numbers JP17K06446, JP19K04914, and JP19K14989.

REFERENCES

- [1] R. Kenneth, *Discrete Mathematics and Its Applications*, 7th ed. McGraw-Hill Science, 2011.
- [2] C. Bishop, *Pattern Recognition and Machine Learning*. Springer, January 2006. [Online]. Available: <https://www.microsoft.com/en-us/research/publication/pattern-recognition-machine-learning/>
- [3] F. M. J. Willems, Y. M. Shtarkov, and T. J. Tjalkens, “The context-tree weighting method: basic properties,” *IEEE Transactions on Information Theory*, vol. 41, no. 3, pp. 653–664, 1995.
- [4] G. J. Sullivan, J. Ohm, W. Han, and T. Wiegand, “Overview of the high efficiency video coding (HEVC) standard,” *IEEE Transactions on Circuits and Systems for Video Technology*, vol. 22, no. 12, pp. 1649–1668, Dec 2012.
- [5] L. Breiman, J. Friedman, C. J. Stone, and R. A. Olshen, *Classification and regression trees*. CRC press, 1984.
- [6] T. Chen and C. Guestrin, “XGBoost: A scalable tree boosting system,” in *Proceedings of the 22nd ACM SIGKDD International Conference on Knowledge Discovery and Data Mining*, ser. KDD ’16. New York, NY, USA: ACM, 2016, pp. 785–794. [Online]. Available: <http://doi.acm.org/10.1145/2939672.2939785>
- [7] L. Breiman, “Random forests,” *Machine Learning*, vol. 45, no. 1, pp. 5–32, 2001. [Online]. Available: <http://dx.doi.org/10.1023/A%3A1010933404324>

- [8] J. O. Berger, *Statistical Decision Theory and Bayesian Analysis*. Springer Science & Business Media, 2013.
- [9] T. Matsushima, H. Inazumi, and S. Hirasawa, “A class of distortionless codes designed by Bayes decision theory,” *IEEE Transactions on Information Theory*, vol. 37, no. 5, pp. 1288–1293, Sep. 1991.
- [10] T. Matsushima and S. Hirasawa, “A Bayes coding algorithm using context tree,” in *Proceedings of 1994 IEEE International Symposium on Information Theory*, 1994, p. 386.
- [11] T. Matsushima and S. Hirasawa, “Reducing the space complexity of a Bayes coding algorithm using an expanded context tree,” in *2009 IEEE International Symposium on Information Theory*, June 2009, pp. 719–723.
- [12] I. Papageorgiou, I. Kontoyiannis, L. Mertzanis, A. Panotopoulou, and M. Skoularidou, “Revisiting context-tree weighting for bayesian inference,” in *2021 IEEE International Symposium on Information Theory (ISIT)*, 2021, pp. 2906–2911.
- [13] I. Kontoyiannis, L. Mertzanis, A. Panotopoulou, I. Papageorgiou, and M. Skoularidou, “Bayesian context trees: Modelling and exact inference for discrete time series,” arXiv, 2020. [Online]. Available: <https://arxiv.org/abs/2007.14900>
- [14] Y. Nakahara and T. Matsushima, “A stochastic model for block segmentation of images based on the quadtree and the Bayes code for it,” *Entropy*, vol. 23, no. 8, 2021. [Online]. Available: <https://www.mdpi.com/1099-4300/23/8/991>
- [15] N. Dobashi, S. Saito, Y. Nakahara, and T. Matsushima, “Meta-tree random forest: Probabilistic data-generative model and Bayes optimal prediction,” *Entropy*, vol. 23, no. 6, 2021. [Online]. Available: <https://www.mdpi.com/1099-4300/23/6/768>

APPENDIX A

A. Proof of Theorem 2

Proof: First, we prove (27). Let $I\{\cdot\}$ denote the indicator function. Then, $\Pr\{v \in \mathcal{V}_T\}$ is expressed as

$$\Pr\{v \in \mathcal{V}_T\} = \sum_{\tau \in \mathcal{T}} I\{v \in \mathcal{V}_\tau\} p(\tau). \quad (68)$$

Here, $v \in \mathcal{V}_\tau$ is equivalent that all the leaf nodes is not a ancestor node of v . Then,

$$\Pr\{v \in \mathcal{V}_T\} = \sum_{\tau \in \mathcal{T}} \prod_{v' \in \mathcal{L}_\tau} I\{v' \neq v\} p(\tau) \quad (69)$$

$$= \sum_{\tau \in \mathcal{T}} \prod_{v' \in \mathcal{I}_\tau} \alpha_{v'} \prod_{v'' \in \mathcal{L}_\tau} I\{v'' \neq v\} (1 - \alpha_{v''}). \quad (70)$$

Therefore, using Lemma 1,

$$\Pr\{v \in \mathcal{V}_T\} = \phi_v(v_\lambda), \quad (71)$$

where

$$\phi_v(v') := \begin{cases} I\{v' \neq v\} (1 - \alpha_{v'}), & v' \in \mathcal{L}_p, \\ I\{v' \neq v\} (1 - \alpha_{v'}) \\ + \alpha_{v'} \prod_{v'' \in \text{Ch}_p(v')} \phi_v(v''), & v' \in \mathcal{I}_p. \end{cases} \quad (72)$$

We further transform this function.

If $v' \neq v$, then $I\{v' \neq v\} = 1$ and consequently,

$$\phi_v(v') = \begin{cases} (1 - \alpha_{v'}), & v' \in \mathcal{L}_p, \\ (1 - \alpha_{v'}) + \alpha_{v'} \prod_{v'' \in \text{Ch}_p(v')} \phi_v(v''), & v' \in \mathcal{I}_p. \end{cases} \quad (73)$$

It has the same form as (22), and every child node $v'' \in \text{Ch}_p(v')$ also satisfies $v'' \neq v$. Therefore, $\phi_v(v') = 1$ for $v' \neq v$.

If $v' \succ v$, v' cannot be in \mathcal{L}_p and has only one child node in $\text{An}(v) \cup \{v\}$. Let v'_{ch} denote it. Then, $\phi_v(v'') = 1$ for the other child nodes $v'' \in \text{Ch}_p(v') \setminus \{v'_{\text{ch}}\}$. Therefore, (72) is represented as follows.

$$\phi_v(v') = \alpha_{v'} \phi_v(v'_{\text{ch}}). \quad (74)$$

Therefore, expanding $\phi_v(v_\lambda)$,

$$\Pr\{v \in \mathcal{V}_T\} = \prod_{v' \in \text{An}(v)} \alpha_{v'}. \quad (75)$$

Next, we prove (28). It is proved in a similar manner to the proof of (27) since

$$\Pr\{v \in \mathcal{I}_T\} = \sum_{\tau \in \mathcal{T}} \prod_{v' \in \mathcal{L}_\tau} I\{v' \not\prec v\} p(\tau). \quad (76)$$

Lastly, we prove (29). We have

$$\Pr\{v \in \mathcal{L}_T\} = \Pr\{v \in \mathcal{V}_T\} - \Pr\{v \in \mathcal{I}_T\}. \quad (77)$$

Therefore, (29) follows from (27) and (28). \blacksquare

B. Proof of Theorem 4

Proof: Substituting (48) into (47), $\mathbb{E}[f(T)]$ can be represented as follows.

$$\mathbb{E}[f(T)] = \sum_{\tau \in \mathcal{T}} \prod_{v \in \mathcal{I}_\tau} \alpha_v g(v) \prod_{v' \in \mathcal{L}_\tau} (1 - \alpha_{v'}) h(v'). \quad (78)$$

Then, using Lemma 1, Theorem 4 straightforwardly follows. \blacksquare

C. Proof of Theorem 5

Proof: First, we switch the order of the summation as follows.

$$\mathbb{E}[f(T)] = \sum_{\tau \in \mathcal{T}} p(\tau) \left(\sum_{v \in \mathcal{I}_\tau} g(v) + \sum_{v' \in \mathcal{L}_\tau} h(v') \right) \quad (79)$$

$$= \sum_{\tau \in \mathcal{T}} p(\tau) \sum_{v \in \mathcal{V}_p} I\{v \in \mathcal{I}_\tau\} g(v) + \sum_{\tau \in \mathcal{T}} p(\tau) \sum_{v' \in \mathcal{V}_p} I\{v' \in \mathcal{L}_\tau\} h(v') \quad (80)$$

$$= \sum_{v \in \mathcal{V}_p} g(v) \sum_{\tau \in \mathcal{T}} p(\tau) I\{v \in \mathcal{I}_\tau\} + \sum_{v' \in \mathcal{V}_p} h(v') \sum_{\tau \in \mathcal{T}} p(\tau) I\{v' \in \mathcal{L}_\tau\} \quad (81)$$

$$= \sum_{v \in \mathcal{V}_p} g(v) \Pr\{v \in \mathcal{I}_T\} + \sum_{v' \in \mathcal{V}_p} h(v') \Pr\{v' \in \mathcal{L}_T\} \quad (82)$$

$$= \sum_{v \in \mathcal{V}_p} g(v) \alpha_v \prod_{v' \in \text{An}(v)} \alpha_{v'} + \sum_{v \in \mathcal{V}_p} h(v) (1 - \alpha_v) \prod_{v' \in \text{An}(v)} \alpha_{v'} \quad (83)$$

$$= \sum_{v \in \mathcal{V}_p} (\alpha_v g(v) + (1 - \alpha_v) h(v)) \prod_{v' \in \text{An}(v)} \alpha_{v'}, \quad (84)$$

where (83) is because of Theorem 2.

Next, we decompose the right-hand side of (84) until it has the same form as (51). Since $\mathcal{V}_p = \{v_\lambda\} \cup \text{Dep}(v_\lambda)$,

$$(84) = \alpha_{v_\lambda} g(v_\lambda) + (1 - \alpha_{v_\lambda}) h(v_\lambda) + \sum_{v \in \text{Dep}(v_\lambda)} (\alpha_v g(v) + (1 - \alpha_v) h(v)) \prod_{v' \in \text{An}(v)} \alpha_{v'}. \quad (85)$$

For any $v \in \text{Dep}(v_\lambda)$, $\text{An}(v)$ contains v_λ . Therefore,

$$(85) = (1 - \alpha_{v_\lambda}) h(v_\lambda) + \alpha_{v_\lambda} \left(g(v_\lambda) + \sum_{v \in \text{Dep}(v_\lambda)} (\alpha_v g(v) + (1 - \alpha_v) h(v)) \prod_{v': v_\lambda \succ v' \succ v} \alpha_{v'} \right). \quad (86)$$

Further, since $\text{Dep}(v_\lambda) = \bigcup_{v \in \text{Ch}_p(v_\lambda)} (\text{Dep}(v) \cup \{v\})$,

$$(86) = (1 - \alpha_{v_\lambda}) h(v_\lambda) + \alpha_{v_\lambda} \left(g(v_\lambda) + \sum_{v \in \text{Ch}_p(v_\lambda)} \left[\sum_{v' \in \text{Dep}(v) \cup \{v\}} (\alpha_{v'} g(v') + (1 - \alpha_{v'}) h(v')) \prod_{v'': v \succeq v'' \succ v'} \alpha_{v''} \right] \right). \quad (87)$$

Comparing (84) and (87), we have

$$\underbrace{\sum_{v \in \mathcal{V}_p} (\alpha_v g(v) + (1 - \alpha_v) h(v)) \prod_{v' \in \text{An}(v)} \alpha_{v'}}_{(a)} = (1 - \alpha_{v_\lambda}) h(v_\lambda) + \alpha_{v_\lambda} \left(g(v_\lambda) + \sum_{v \in \text{Ch}_p(v_\lambda)} \underbrace{\left[\sum_{v' \in \text{Dep}(v) \cup \{v\}} (\alpha_{v'} g(v') + (1 - \alpha_{v'}) h(v')) \prod_{v'': v \succeq v'' \succ v'} \alpha_{v''} \right]}_{(b)} \right). \quad (88)$$

The underbraced parts (a) and (b) have the same structure. Therefore, (b) can be decomposed in a similar manner from (84) to (87). We can continue this decomposition to the leaf nodes.

Finally, we have an alternative definition of $\xi(v) : \mathcal{V}_p \rightarrow \mathbb{R}$, which is equivalent to (51).

$$\xi(v) := \sum_{v' \in \text{Dep}(v) \cup \{v\}} (\alpha_{v'} g(v') + (1 - \alpha_{v'}) h(v')) \prod_{v'': v \succeq v'' \succ v'} \alpha_{v''} \quad (89)$$

The equivalence is confirmed by substituting it into both sides of (88). Therefore, Theorem 5 is proved. ■

D. Proof of Theorem 6

Proof: By the Bayes theorem, we have

$$p(\alpha|\tau) \propto p(\tau|\alpha)p(\alpha) \quad (90)$$

$$= \prod_{v \in \mathcal{I}_\tau} \alpha_v \prod_{v' \in \mathcal{L}_\tau} (1 - \alpha_{v'}) \prod_{v'' \in \mathcal{V}_p} \text{Beta}(v''|\beta_{v''}, \gamma_{v''}) \quad (91)$$

$$= \prod_{v \in \mathcal{I}_\tau} \alpha_v \text{Beta}(v|\beta_v, \gamma_v) \times \prod_{v' \in \mathcal{L}_\tau} (1 - \alpha_{v'}) \text{Beta}(v'|\beta_{v'}, \gamma_{v'}) \times \prod_{v'' \in \mathcal{V}_p \setminus \mathcal{V}_\tau} \text{Beta}(v''|\beta_{v''}, \gamma_{v''}) \quad (92)$$

$$\propto \prod_{v \in \mathcal{V}_p} \text{Beta}(\alpha_v|\beta_{v|\tau}, \gamma_{v|\tau}), \quad (93)$$

where we used the conjugate property between the Bernoulli distribution and the Beta distribution for each term and

$$\beta_{v|\tau} := \begin{cases} \beta_v + 1, & v \in \mathcal{I}_\tau, \\ \beta_v, & \text{otherwise,} \end{cases} \quad (94)$$

$$\gamma_{v|\tau} := \begin{cases} \gamma_v + 1, & v \in \mathcal{L}_\tau, \\ \gamma_v, & \text{otherwise.} \end{cases} \quad (95)$$

E. Proof of Theorem 7

Proof: We prove (63) from the right-hand side to the left.

$$\begin{aligned} & \prod_{v \in \mathcal{I}_\tau} \alpha_{v|x} \prod_{v' \in \mathcal{L}_\tau} (1 - \alpha_{v'|x}) \\ &= \prod_{v \in \mathcal{I}_\tau} \alpha_{v|x} \prod_{v' \in \mathcal{L}_\tau \cap \mathcal{L}_p} (1 - \alpha_{v'|x}) \prod_{v'' \in \mathcal{L}_\tau \cap \mathcal{I}_p} (1 - \alpha_{v''|x}). \end{aligned} \quad (96)$$

In the following, we transform each of the above products in order. First, the first product is transformed by substituting (62) as follows.

$$\prod_{v \in \mathcal{I}_\tau} \alpha_{v|x} = \prod_{v \in \mathcal{I}_\tau} \frac{\alpha_v g(x, v) \prod_{v' \in \text{Ch}_p(v)} q(x|v')}{q(x|v)}. \quad (97)$$

Next, the second product is transformed as follows.

$$\prod_{v \in \mathcal{L}_\tau \cap \mathcal{L}_p} (1 - \alpha_{v|x}) = \prod_{v \in \mathcal{L}_\tau \cap \mathcal{L}_p} (1 - \alpha_v) \quad (98)$$

$$= \prod_{v \in \mathcal{L}_\tau \cap \mathcal{L}_p} \frac{(1 - \alpha_v) h(x, v)}{q(x|v)}, \quad (99)$$

where (98) is because of (62) and (99) is because $q(x|v) = h(x, v)$ for $v \in \mathcal{L}_p$.

Lastly, the third product is transformed as follows.

$$\begin{aligned} & \prod_{v \in \mathcal{L}_\tau \cap \mathcal{I}_p} (1 - \alpha_{v|x}) \\ &= \prod_{v \in \mathcal{L}_\tau \cap \mathcal{I}_p} \left(1 - \frac{\alpha_v g(x, v) \prod_{v' \in \text{Ch}_p(v)} q(x|v')}{q(x|v)} \right) \end{aligned} \quad (100)$$

$$= \prod_{v \in \mathcal{L}_\tau \cap \mathcal{I}_p} \frac{q(x|v) - \alpha_v g(x, v) \prod_{v' \in \text{Ch}_p(v)} q(x|v')}{q(x|v)} \quad (101)$$

$$= \prod_{v \in \mathcal{L}_\tau \cap \mathcal{I}_p} \frac{(1 - \alpha_v) h(x, v)}{q(x|v)}, \quad (102)$$

where (100) is because of (62) and (102) is because of substitution of (61) into $q(x|v)$ at the numerator.

Therefore, we can combine (97), (99) and (102). Then,

$$\begin{aligned} (96) &= \prod_{v \in \mathcal{I}_\tau} \frac{\alpha_v g(x, v) \prod_{v' \in \text{Ch}_p(v)} q(x|v')}{q(x|v)} \\ &\quad \times \prod_{v' \in \mathcal{L}_\tau} \frac{(1 - \alpha_{v'}) h(x, v')}{q(x|v')}. \end{aligned} \quad (103)$$

Here, (103) is a telescoping product, i.e., $q(x|v)$ appears at once in each of the denominator and the numerator. Therefore, we can cancel them except for $q(x|v_\lambda)$. Then,

$$(103) = \frac{1}{q(x|v_\lambda)} \prod_{v \in \mathcal{I}_\tau} \alpha_v g(x, v) \prod_{v' \in \mathcal{L}_\tau} (1 - \alpha_{v'}) h(x, v') \quad (104)$$

$$\begin{aligned} &= \frac{1}{q(x|v_\lambda)} \left(\prod_{v \in \mathcal{I}_\tau} g(x, v) \prod_{v' \in \mathcal{L}_\tau} h(x, v') \right) \\ &\quad \times \left(\prod_{v \in \mathcal{I}_\tau} \alpha_v \prod_{v' \in \mathcal{L}_\tau} (1 - \alpha_{v'}) \right) \end{aligned} \quad (105)$$

$$= \frac{p(x|\tau)p(\tau)}{q(x|v_\lambda)}, \quad (106)$$

where we used (58) and Definition 1.

In addition, because of Theorem 4,

$$q(x|v_\lambda) = \mathbb{E}[p(x|T)] = \sum_{\tau \in \mathcal{T}} p(x|\tau)p(\tau) = p(x). \quad (107)$$

Therefore,

$$(106) = \frac{p(x|\tau)p(\tau)}{p(x)} = p(\tau|x). \quad (108)$$

Then, Theorem 7 holds. ■

F. Proof of Corollary 2

Proof: We will prove only $q(x|v) = 1$ for $v \not\prec v_{\text{end}}$. Then, (66) and (67) are straightforwardly derived by substituting it with (64) and (65) into (61) and (62).

For $v \not\prec v_{\text{end}}$, substituting (64) and (65) into (61),

$$q(x|v) = \begin{cases} 1, & v \in \mathcal{L}_p, \\ (1 - \alpha_v) + \alpha_v \prod_{v' \in \text{Ch}_p(v)} q(x|v'), & v \in \mathcal{I}_p. \end{cases} \quad (109)$$

Since this has the same form as (73), $q(x|v) = 1$ for $v \not\preceq v_{\text{end}}$ is derived in a similar manner. ■

APPENDIX B

PSEUDOCODE TO CALCULATE MODE OF $p(\tau)$

Algorithm 1 Calculation of mode of $p(\tau)$

Input: $\{\alpha_v\}_{v \in \mathcal{V}_p}$
Output: $\tau^* = \arg \max_{\tau} p(\tau)$

```

1: function FLAG_CALCULATION( $v$ )           ▷ Subroutine
2:   if  $v \in \mathcal{L}_p$  then
3:      $\delta_v \leftarrow 0$ 
4:   return 1
5:   else if  $v \in \mathcal{I}_p$  then
6:      $\text{tmp} \leftarrow \prod_{v' \in \text{Ch}_p(v)} \text{FLAG\_CALCULATION}(v')$ 
7:     if  $1 - \alpha_v < \alpha_v \cdot \text{tmp}$  then
8:        $\delta_v \leftarrow 1$ 
9:     return  $\alpha_v \cdot \text{tmp}$ 
10:   else
11:      $\delta_v \leftarrow 0$ 
12:   return  $1 - \alpha_v$ 
13:   end if
14: end if
15: end function

16:
17: function BACKTRACKING( $v, \mathcal{V}, \mathcal{E}$ )         ▷ Subroutine
18:   if  $\delta_v = 0$  then
19:     return
20:   else if  $\delta_v = 1$  then
21:      $\mathcal{V} \leftarrow \mathcal{V} \cup \text{Ch}_p(v)$ 
22:      $\mathcal{E} \leftarrow \mathcal{E} \cup \bigcup_{v' \in \text{Ch}_p(v)} (v, v')$ 
23:     for all  $v' \in \text{Ch}_p(v)$  do
24:       BACKTRACKING( $v', \mathcal{V}, \mathcal{E}$ )
25:     end for
26:   return
27: end if
28: end function

29:
30: procedure                               ▷ The main procedure
31:   FLAG_CALCULATION( $v_\lambda$ )
32:    $\mathcal{V} \leftarrow \emptyset$ 
33:    $\mathcal{E} \leftarrow \emptyset$ 
34:   BACKTRACKING( $v_\lambda, \mathcal{V}, \mathcal{E}$ )
35:   return  $\tau^* = (\mathcal{V}, \mathcal{E})$ 
36: end procedure

```
