

Anti-Ramsey problems in the generalized Petersen graphs for cycles

Huiqing LIU^{1*} Mei LU^{2†} Shunzhe ZHANG^{1‡}

¹Hubei Key Laboratory of Applied Mathematics, Faculty of Mathematics and Statistics, Hubei University, Wuhan 430062, China

²Department of Mathematical Sciences, Tsinghua University, Beijing 100084, China.

Abstract

The anti-Ramsey number $Ar(G, H)$ is the maximum number of colors in an edge-coloring of G with no rainbow copy of H . In this paper, we determine the exact anti-Ramsey number in the generalized Petersen graph $P_{n,k}$ for cycles C_d , where $1 \leq k \leq \lfloor \frac{n-1}{2} \rfloor$ and $5 \leq d \leq 6$. We also give an algorithm to obtain the upper bound or lower bound of anti-Ramsey number.

Keywords: anti-Ramsey number; rainbow; generalized Petersen graph; cycle

1 Introduction

Let $G = (V(G), E(G))$ be a graph. We denote by $E(v)$ the set of all edges incident with v in G , and by C_d a cycle of order d . Let $M(G)$ be the maximum number of vertex-disjoint cycles with length at least 2 in G .

An edge-coloring of a graph G is a mapping $\phi : E(G) \rightarrow \{1, 2, \dots, c\}$, where $\{1, 2, \dots, c\}$ is a set of colors. An edge colored graph is called *rainbow* if all the colors on the edges are distinct. A *rainbow copy* of a graph H in an edge-colored graph G is a subgraph of G isomorphic to H such that the coloring restricted to H is rainbow. Given

*Partially supported by NNSFC (No. 11971158), email: hql_2008@163.com;

†Partially supported by NNSFC (No. 12171272), email: lumei@mail.tsinghua.edu.cn;

‡Partially supported by Youth Project Funds of Hubei Provincial Department of Education, email: shunzhezhang@hubu.edu.cn.

two graphs G and H , the *anti-Ramsey number* $Ar(G, H)$ is the maximum number of colors in an edge-coloring of G which has no rainbow copy of H . If there is no copy of H in G , then $Ar(G, H) = |E(G)|$.

The study of anti-Ramsey theory was initiated by Erdős, Simonovits and Sós in 1973 [4] and considered in the classical case when $G = K_n$. They presented a close relationship between the anti-Ramsey number and Turán number. Since then, plentiful results were researched for a variety of graphs H , including cycles [1, 2, 12, 13, 18, 19], cliques [4, 17], trees [9, 11], and matchings [7, 16]. Some other graphs were also considered as the host graphs in anti-Ramsey problems, such as hypergraphs [6], hypercubes [3], complete split graphs [5, 14], and triangulations [8, 10, 15].

For cycles, Erdős, Simonovits and Sós [4] showed that $Ar(K_n, C_3) = n - 1$. Alon [1] showed that $Ar(K_n, C_4) = \lfloor \frac{4n}{3} \rfloor - 1$. Montellano-Ballesteros and Neumann-Lara [18] proved that $Ar(K_n, C_d) = (\frac{d-2}{2} + \frac{1}{d-1})n + O(1)$ for $3 \leq d \leq n$. In 2015, Hornák, Jendrol, Schiermeyer and Soták [8] investigated the anti-Ramsey numbers for cycles in plane triangulations. Lan, Shi and Song [15] gave the upper and lower bounds of the anti-Ramsey number $Ar(W_n, C_d)$ for $n \geq d - 1$ and $d \geq 6$. Recently, Xu, Lu and Liu [19] completely solved the anti-Ramsey numbers of cycles in wheel graphs and determined some exact values of $Ar(G, C_d)$ when the host graph G is $P_m \times P_n$, $P_m \times C_n$, $C_m \times C_n$ and cyclic Cayley graph, respectively.

Here, we are interested in the anti-Ramsey problems in generalized Petersen graphs for cycles. Let $n \geq 3$ and k be integers such that $1 \leq k \leq n - 1$. The generalized Petersen graph $P_{n,k}$ is a graph of order $2n$ with $V(P_{n,k}) = \{u_i, v_i : 0 \leq i \leq n - 1\}$ and edge set $E(P_{n,k}) = \{u_i u_{i+1}, u_i v_i, v_i v_{i+k} : 0 \leq i \leq n - 1\}$, where subscripts are taken modulo n . Let $O_n = \{u_0, u_1, \dots, u_{n-1}\}$ and $I_n = \{v_0, v_1, \dots, v_{n-1}\}$. Then $V(P_{n,k}) = O_n \cup I_n$. The subgraph induced by O_n is called the outer rim, while the subgraph induced by I_n is called the inner rim. A spoke of $P_{n,k}$ is an edge of the form $u_i v_i$ for some $0 \leq i \leq n - 1$. Note that $P_{n,k}$ is isomorphic to $P_{n,n-k}$. Hence we can always assume that $1 \leq k \leq \lfloor \frac{n-1}{2} \rfloor$.

The generalized Petersen graph has been extensively investigated as many nice structural and algorithmic properties.

Proposition 1.1 (i) $P_{n,k}$ contains C_5 if and only if $n \in \{3, 5\}$ or $k \in \{2, \frac{n}{5}, \frac{2n}{5}, \frac{n-1}{2}\}$;
(ii) $P_{n,k}$ contains C_6 if and only if $k \in \{1, 3, \frac{n}{6}, \frac{n-1}{3}, \frac{n+1}{3}, \frac{n-2}{2}\}$.

Proof. Note that $V(P_{n,k}) = O_n \cup I_n$. By the definition of $P_{n,k}$, $|V(C_5) \cap O_n| \in \{0, 2, 3, 5\}$ and $|V(C_6) \cap O_n| \in \{0, 2, 3, 4, 6\}$. Obviously, $P_{n,1}$ contains C_5 for $n \in \{3, 5\}$ and C_6 for $n \geq 3$. So, in the following, we can assume $k \geq 2$.

Let $C_t = w_1 w_2 \cdots w_t w_1$ be a cycle of length t in $P_{n,k}$. If $V(C_t) \subseteq O_n$, then $t = n$. If $|V(C_t) \cap O_n| = 0$, i.e., $V(C_t) \subseteq I_n$, then we can assume $w_1 = v_0$ by symmetry, and then $C_t = v_0 v_k v_{2k} \cdots v_{(t-1)k} v_0$, which implies $tk \equiv 0 \pmod{n}$ and $ik \not\equiv 0 \pmod{n}$ for $1 \leq i \leq t-1$. Combining this with $1 \leq k \leq \lfloor \frac{n-1}{2} \rfloor$, we have $k \in \{\frac{n}{5}, \frac{2n}{5}\}$ for $t = 5$ and $k = \frac{n}{6}$ for $t = 6$. Thus we may assume $2 \leq |V(C_t) \cap O_n| \leq t-1$. Without loss of generality, we assume $w_1 = u_0$ and $w_2 = u_1$.

(i) Note that $|V(C_5) \cap O_n| \in \{2, 3\}$. If $|V(C_5) \cap O_n| = 2$, then $|V(C_5) \cap I_n| = 3$, $w_5 = v_0$ and $w_3 = v_1$. Let $w_4 = v_s$, then $v_1 v_s, v_s v_0 \in E(P_{n,k})$, which implies $s-1 = k$ and $n-s = k$. So $k = \frac{n-1}{2}$.

If $|V(C_5) \cap O_n| = 3$, that is $|V(C_5) \cap I_n| = 2$, then we can assume that $w_3 = u_2$, and $w_4 = v_2, w_5 = v_0$, which implies $v_2 v_0 \in E(P_{n,k})$. Thus $k = 2$.

(ii) Note that $|V(C_6) \cap O_n| \in \{2, 3, 4\}$ and $N_{I_n}(v_i) = \{v_{i+k}, v_{n-k+i}\}$ for $0 \leq i \leq n-1$.

If $|V(C_6) \cap O_n| = 2$, then $w_3 = v_1, w_6 = v_0, w_4 = v_{k+1}$ (resp. $w_4 = v_{n-k+1}$) and $w_5 = v_{2k+1}$ (resp. $w_5 = v_{n-2k+1}$), which implies $3k+1 \equiv 0 \pmod{n}$ (resp. $n-3k+1 \equiv 0 \pmod{n}$), i.e., $k = \frac{n-1}{3}$ (resp. $k = \frac{n+1}{3}$).

If $|V(C_6) \cap O_n| = 3$, say $w_3 = u_2$, then $w_4 = v_2$, and $w_5 = v_{k+2}$ (resp. $w_5 = v_{n-k+2}$), which implies $2k+2 \equiv 0 \pmod{n}$ (resp. $n-2k+2 \equiv 0 \pmod{n}$), i.e., $k = \frac{n-2}{2}$.

If $|V(C_6) \cap O_n| = 4$, then we can assume that $w_3 = u_2$ and $w_4 = u_3$. Thus $w_5 = v_3$, which implies $k = 3$ as $w_6 = v_0$. ■

In this paper, we determine the exact anti-Ramsey number in $P_{n,k}$ for C_d , where $1 \leq k \leq \lfloor \frac{n-1}{2} \rfloor$ and $5 \leq d \leq 6$. Our results are as follows.

Theorem 1.2 For $1 \leq k \leq \lfloor \frac{n-1}{2} \rfloor$, we have

$$Ar(P_{n,k}, C_5) = \begin{cases} 7, & \text{if } (n, k) = (3, 1); \\ 13, & \text{if } (n, k) = (5, 1); \\ 10, & \text{if } (n, k) = (5, 2); \\ 22, & \text{if } (n, k) = (10, 2); \\ \lfloor \frac{7n}{3} \rfloor, & \text{if } n \geq 6, n \neq 10 \text{ and } k = 2, \text{ or } k = \frac{n-1}{2} \geq 3; \\ \frac{14n}{5}, & \text{if } k \geq 3 \text{ and } k \in \{\frac{n}{5}, \frac{2n}{5}\}; \\ 3n, & \text{otherwise.} \end{cases}$$

Theorem 1.3 For $n \geq 3$, we have

$$Ar(P_{n,1}, C_6) = \begin{cases} 7, & \text{if } n = 3; \\ 9, & \text{if } n = 4; \\ 14, & \text{if } n = 6; \\ \lfloor \frac{5n}{2} \rfloor, & \text{if } n \geq 5, n \neq 6; \end{cases}$$

Theorem 1.4 For $n \geq 5$, we have

$$Ar(P_{n,2}, C_6) = \begin{cases} 11, & \text{if } n = 5; \\ 14, & \text{if } n = 6; \\ 17, & \text{if } n = 7; \\ 34, & \text{if } n = 12; \\ 3n, & \text{otherwise.} \end{cases}$$

Theorem 1.5 For $n \geq 7$, we have

$$Ar(P_{n,3}, C_6) = \begin{cases} 17, & \text{if } n = 8; \\ 22, & \text{if } n = 10; \\ 42, & \text{if } n = 18; \\ \lfloor \frac{5n}{2} \rfloor, & \text{otherwise.} \end{cases}$$

Theorem 1.6 For $4 \leq k \leq \lfloor \frac{n-1}{2} \rfloor$, we have

$$Ar(P_{n,k}, C_6) = \begin{cases} \lfloor \frac{5n}{2} \rfloor, & \text{if } k \in \{ \frac{n-1}{3}, \frac{n+1}{3} \}; \\ \lfloor \frac{7n}{3} \rfloor, & \text{if } k = \frac{n-2}{2}; \\ \frac{17n}{6}, & \text{if } k = \frac{n}{6}; \\ 3n, & \text{otherwise.} \end{cases}$$

The rest of the paper is organized as follows. In Section 2, we give some notations and lemmas which will be used in the proof of main results. The exact values of $Ar(P_{n,k}, C_5)$ and $Ar(P_{n,k}, C_6)$ for $1 \leq k \leq \lfloor \frac{n-1}{2} \rfloor$ are determined in Sections 3 and 4, respectively.

2 Preliminaries

A hypergraph is a pair $H = (V, \mathcal{F})$, where V is a finite set of vertices and \mathcal{F} is a family of subsets of V such that for every $F \in \mathcal{F}$, $F \neq \emptyset$ and $V = \bigcup_{F \in \mathcal{F}} F$. If $|F| = 1$, we call F a loop. The rank of H is defined as $r(H) = \max_{F \in \mathcal{F}} |F|$. If $r(H) = 2$, then H is a graph. For every $v \in V$, denote $F_H(v) = \{F \in \mathcal{F} | v \in F\}$. Let $s(H) = \max\{|F_i \cap F_j| : F_i, F_j \in \mathcal{F}\}$.

Let G be a simple graph with $E(G) = \{e_1, e_2, \dots, e_m\}$, and let $\Psi = \{G_1, \dots, G_l\}$ with $E(G) = \bigcup_{i=1}^l E(G_i)$, where G_i is a subgraph of G for $1 \leq i \leq l$. A hypergraph $H_{G,\Psi}$ is constructed in [19] with $V(H_{G,\Psi}) = \{x_1, x_2, \dots, x_l\}$ and $E(H_{G,\Psi}) = \{F_1, F_2, \dots, F_m\}$, where each x_i corresponds to G_i in Ψ ($1 \leq i \leq l$) and $x_i \in F_j$ if and only if $e_j \in E(G_i)$ ($1 \leq j \leq m$).

Given a graph G and $\Psi = \{G_1, \dots, G_l\}$, where G_i is a subgraph of G for $1 \leq i \leq l$. For an edge-coloring of G , $H \in \Psi$ is called *rainbow- Ψ graph* if the coloring restricted to H is rainbow. Let $Ar_G(\Psi)$ denote the maximum number of colors in a coloring of the edges of G such that there is no rainbow- Ψ graph of G . Obviously, if $\Psi = \{G_1, \dots, G_l\}$ is the set of all copies of H in G , then $Ar(G, H) = Ar_G(\Psi)$.

The following two lemmas will be useful in the proof of our main results.

Lemma 2.1 [19] *Let $G = (V, E)$ be a simple graph. Let $\Psi = \{G_1, \dots, G_l\}$ such that $E(G) = \cup_{i=1}^l E(G_i)$, where G_i is a subgraph of G for $1 \leq i \leq l$. Suppose $H_{G, \Psi}$ is a graph. Then*

$$Ar_G(\Psi) = |E(G)| - |\Psi| + M(H_{G, \Psi}).$$

Lemma 2.2 [19] *Let $G = (V, E)$ be a simple graph. Let $\Psi = \{G_1, \dots, G_l\}$ such that $E(G) = \cup_{i=1}^l E(G_i)$, where G_i is a subgraph of G for $1 \leq i \leq l$. Suppose $H_{G, \Psi}$ is a hypergraph with $r(H_{G, \Psi}) = r$ and $s(H_{G, \Psi}) = s$, then*

$$Ar_G(\Psi) \leq |E(G)| - \left\lceil \frac{2l}{r+s} \right\rceil.$$

Let $H = (V, \mathcal{F})$ be a hypergraph with $|V(H)| = l$ and $\mathcal{F} = \{F_1, F_2, \dots, F_m\}$. Given an integer h with $0 \leq h \leq m-1$. Let $\{\mathcal{F}_1, \mathcal{F}_2, \dots, \mathcal{F}_{m-h}\}$ be an $(m-h)$ -partition of \mathcal{F} with $|\mathcal{F}_1| \geq |\mathcal{F}_2| \geq \dots \geq |\mathcal{F}_{m-h}| \geq 1$. For $1 \leq i \leq m-h$, we denote $L_i := \{v \in F_t \cap F_j \mid F_t, F_j \in \mathcal{F}_i, t \neq j\}$ if $|\mathcal{F}_i| \geq 2$, and \emptyset otherwise. Then $|L_i|$ is the number of vertices which appear simultaneously in at least two edges in \mathcal{F}_i when $|\mathcal{F}_i| \geq 2$. If each vertex contained in the edges of \mathcal{F}_i appears simultaneously in only two edges of \mathcal{F}_i , then \mathcal{F}_i is called a *barrier*. Obviously, $|\cup_{i=1}^{m-h} L_i| \leq l$. If $|\cup_{i=1}^{m-h} L_i| < l$ for any $(m-h)$ -partition of \mathcal{F} , then we say $H \in \mathcal{P}_{h,l}$ and call that H satisfies the property $\mathcal{P}_{h,l}$. Clearly, if $H \notin \mathcal{P}_{h,l}$, then there exists a $(m-h)$ -partition $\{\mathcal{F}_1, \mathcal{F}_2, \dots, \mathcal{F}_{m-h}\}$ of $E(H)$ such that $|\cup_{i=1}^{m-h} L_i| = l$, that is, $\cup_{i=1}^{m-h} L_i = V$.

Note that for each $1 \leq i \leq m-h$,

$$|L_i| \leq \begin{cases} \lfloor \frac{r(H)|\mathcal{F}_i|}{2} \rfloor, & \text{if } |\mathcal{F}_i| \geq 3 \\ s(H), & \text{if } |\mathcal{F}_i| = 2 \\ 0, & \text{otherwise.} \end{cases} \quad (1)$$

Hence, if $|\mathcal{F}_i| \geq 3$ and $r(H)$ is even, then the equality in (1) holds if and only if \mathcal{F}_i is a barrier.

Lemma 2.3 *Let $H = (V, \mathcal{F})$ be a hypergraph with $|V(H)| \geq 7$ and $\mathcal{F} = \{F_1, F_2, \dots, F_m\}$ with $m \geq 4$. If $r(H) = 3$ and $s(H) = 2$, then $H \in \mathcal{P}_{3,7}$.*

Proof. Let $\{\mathcal{F}_1, \mathcal{F}_2, \dots, \mathcal{F}_{m-3}\}$ be an $(m-3)$ -partition of \mathcal{F} with $|\mathcal{F}_1| \geq |\mathcal{F}_2| \geq \dots \geq |\mathcal{F}_{m-3}| \geq 1$. Then $2 \leq |\mathcal{F}_1| \leq 4$ and $\sum_{i=1}^{m-3} (|\mathcal{F}_i| - 1) = 3$. Since $r(H) = 3$ and $s(H) = 2$, we have $|L_i| \leq 2(|\mathcal{F}_i| - 1)$ for $1 \leq i \leq m-3$ by (1). So $|\bigcup_{i=1}^{m-h} L_i| \leq \sum_{i=1}^{m-3} |L_i| \leq 2 \sum_{i=1}^{m-3} (|\mathcal{F}_i| - 1) = 6 < 7$, and hence $H \in \mathcal{P}_{3,7}$. ■

Let $G = (V, E)$ be a (hyper)graph. Let ϕ be an edge-coloring of G with the color set $K = \{c_1, c_2, \dots, c_t\}$. Note that coloring here need not be proper. Suppose $\phi^{-1}(c_i) \neq \emptyset$ for each $c_i \in K$. Let $E(c_i) = \{e \in E(G) \mid \phi(e) = c_i\}$ and $D(c_i) = |E(c_i)| - 1$ for $c_i \in K$. Set $D(G) = \sum_{c_i \in K} D(c_i)$. Then $D(c_i)$ is the repeated number of the color c_i and $D(G)$ is the repeated number of all colors in G . Obviously, $|K| = |E(G)| - D(G)$. In the following, we will give a necessary and sufficient condition for the bounds of $Ar_G(\Psi)$.

Lemma 2.4 *Let $G = (V, E)$ be a simple graph with $|E(G)| = m$. Let $\Psi = \{G_1, \dots, G_l\}$ such that $E(G) = \bigcup_{i=1}^l E(G_i)$, where G_i is a subgraph of G for $1 \leq i \leq l$. Suppose h is an integer with $0 \leq h \leq m-1$. Then $H_{G,\Psi} \in \mathcal{P}_{h,l}$ if and only if $Ar_G(\Psi) \leq |E(G)| - (h+1)$.*

Proof. Suppose that $E(G) = \{e_1, \dots, e_m\}$, $V(H_{G,\Psi}) = \{x_1, \dots, x_l\}$ and $E(H_{G,\Psi}) = \{F_1, F_2, \dots, F_m\}$, where each x_i corresponds to G_i in Ψ for $1 \leq i \leq l$ and each F_j corresponds to e_j for $1 \leq j \leq m$.

Let ϕ be an edge-coloring of G with the color set $K = \{c_1, c_2, \dots, c_{m-h}\}$ such that $\phi^{-1}(c_i) \neq \emptyset$. Consider the edge-coloring ϕ^* of $H_{G,\Psi}$: $\phi^*(F_j) = \phi(e_j)$ for $1 \leq j \leq m$. Then $D_\phi(G) = D_{\phi^*}(H_{G,\Psi}) = h$. Denote $\mathcal{F}_i = \{F \in E(H_{G,\Psi}) \mid \phi^*(F) = c_i\}$, $1 \leq i \leq m-h$. Then $\{\mathcal{F}_1, \mathcal{F}_2, \dots, \mathcal{F}_{m-h}\}$ is a $(m-h)$ -partition of $E(H_{G,\Psi})$.

If $H_{G,\Psi} \in \mathcal{P}_{h,l}$, then $|\bigcup_{i=1}^{m-h} L_i| < l$, and thus $\bigcup_{i=1}^{m-h} L_i \subset V(H_{G,\Psi})$. Let $x_p \in V(H_{G,\Psi}) \setminus (\bigcup_{i=1}^{m-h} L_i)$. Then each edge of $F_{H_{G,\Psi}}(x_p)$ is colored with distinct colors in $H_{G,\Psi}$, and then every edge of $E(G_p)$ is colored with distinct colors in G . It follows that there exists a rainbow G_p in any edge-coloring of G by using $m-h = |E(G)| - D_\phi(G)$ colors. So $Ar_G(\Psi) \leq (|E(G)| - D_\phi(G)) - 1 = |E(G)| - (h+1)$.

Conversely, we suppose that $H_{G,\Psi} \notin \mathcal{P}_{h,l}$. Then there exists a $(m-h)$ -partition $\{\mathcal{F}'_1, \mathcal{F}'_2, \dots, \mathcal{F}'_{m-h}\}$ of $E(H_{G,\Psi})$ such that $\bigcup_{i=1}^{m-h} L'_i = V(H_{G,\Psi})$. Let ϕ' be an edge-coloring of $H_{G,\Psi}$ such that $\phi'^{-1}(c_i) = \mathcal{F}'_i$ for $1 \leq i \leq m-h$. Then $D_{\phi'}(H_{G,\Psi}) = h$.

Now we give an edge-coloring ϕ'' of G as follows: $\phi''(e_j) = \phi'(F_j)$, where each F_j corresponds to e_j ($1 \leq j \leq m$). Then $D_{\phi''}(G) = D_{\phi'}(H_{G,\Psi}) = h$. Note that, for any $x_p \in V(H_{G,\Psi})$, $x_p \in L'_i$ for some i ($1 \leq i \leq m-h$), and hence there are at least two edges of $F_{H_{G,\Psi}}(x_p)$ colored with the same color in $H_{G,\Psi}$, which implies at least two edges in each G_p are colored with the same color. So, G contains no rainbow- Ψ graph of G . Therefore, $Ar_G(\Psi) \geq |E(G)| - D_{\phi''}(G) = |E(G)| - h$. ■

In order to obtain the bounds of $Ar_G(\Psi)$, we need to check that whether a hypergraph $H_{G,\Psi}$ satisfies the property $\mathcal{P}_{h,l}$ or not by Lemma 2.4. So we give the following algorithm.

Algorithm 2.5 *Determine a hypergraph $H_{G,\Psi}$ with the property $\mathcal{P}_{h,l}$ or not.*

Input: A hypergraph $H_{G,\Psi} = (V, \mathcal{F})$ with $|V| = l$ and $\mathcal{F} = \{F_1, F_2, \dots, F_m\}$, an integer h with $0 \leq h \leq m - 1$.

Output: $H_{G,\Psi} \in \mathcal{P}_{h,l}$ or $H_{G,\Psi} \notin \mathcal{P}_{h,l}$.

```

1: set  $Q := \emptyset$ 
2: for each decreasing positive integer sequence  $(f_1, f_2, \dots, f_{m-h})$  such that
    $\sum_{i=1}^{m-h} f_i = m$  do
3:   append  $(f_1, f_2, \dots, f_{m-h})$  to  $Q$ 
4: end for
5: while  $Q$  is nonempty do
6:   consider a sequence  $(f_1, f_2, \dots, f_{m-h})$  of  $Q$ 
7:   for each  $(m-h)$ -partition  $\{\mathcal{F}_1, \mathcal{F}_2, \dots, \mathcal{F}_{m-h}\}$  of  $\mathcal{F}$  such that  $|\mathcal{F}_t| = f_t$ ,
      $1 \leq t \leq m-h$  do
8:     set  $i := 1$ 
9:     for  $i < m-h+1$  do
10:      if  $f_i \geq 2$  then
11:         $L_i := \{v \in F_t \cap F_j \mid F_t, F_j \in \mathcal{F}_i \text{ with } t \neq j\}$ 
12:      else
13:         $L_i := \emptyset$ 
14:      end if
15:    end for
16:    if  $|\bigcup_{i=1}^{m-h} L_i| = l$  then
17:      return  $H_{G,\Psi} \notin \mathcal{P}_{h,l}$ 
18:    break
19:    end if
20:  end for
21:  remove  $(f_1, f_2, \dots, f_{m-h})$  from  $Q$ 
22: end while
23: return  $H_{G,\Psi} \in \mathcal{P}_{h,l}$ 

```

3 Anti-Ramsey number $Ar(P_{n,k}, C_5)$

In this section, we will determine the anti-Ramsey number $Ar(P_{n,k}, C_5)$ for $1 \leq i \leq \lfloor \frac{n-1}{2} \rfloor$. If G contains no C_5 , then $Ar(P_{n,k}, C_5) = |E(P_{n,k})| = 3n$. Thus by Proposition 1.1(i), we can assume that $n \in \{3, 5\}$ or $k \in \{2, \frac{n}{5}, \frac{2n}{5}, \frac{n-1}{2}\}$.

Lemma 3.1 $Ar(P_{3,1}, C_5) = 7$.

Proof. Let $G_i^1 := u_i u_{i+1} v_{i+1} v_{i+2} u_{i+2} u_i$, $G_i^2 := u_i u_{i+1} v_{i+1} v_{i+2} v_i u_i$ for $0 \leq i \leq 2$. Then $\Psi = \{G_i^1, G_i^2 \mid 0 \leq i \leq 2\}$ is the set of all C_5 in $P_{3,1}$, and then $Ar(P_{3,1}, C_5) = Ar_{P_{3,1}}(\Psi)$.

First we show that $Ar(P_{3,1}, C_5) \geq 7$. Let $\phi : E(P_{3,1}) \rightarrow \{1, 2, \dots, 7\}$ be an edge-coloring of $P_{3,1}$ such that $\phi(u_i v_i) = 1$ for $0 \leq i \leq 2$ and every other edges obtains distinct colorings from $\{2, \dots, 7\}$ (see Fig. 1(a)). Note that each $C_5 \in \Psi$ contains two spokes $u_i v_i$ and $u_{i+1} v_{i+1}$ for some $0 \leq i \leq 2$, and hence $P_{3,1}$ has no rainbow C_5 under the coloring ϕ . So, $Ar(P_{3,1}, C_5) \geq 7$.

Next we show that $Ar(P_{3,1}, C_5) \leq 7$. Let $V(H_{P_{3,1}, \Psi}) = \{x_i, y_i \mid 0 \leq i \leq 2\}$ and $E(H_{P_{3,1}, \Psi}) = \{F_0, \dots, F_8\}$, where x_i and y_i correspond to G_i^1 and G_i^2 in Ψ , respectively, each F_i corresponds to $u_i u_{i+1}$, F_{i+3} corresponds to $u_i v_i$, and F_{i+6} corresponds to $v_i v_{i+1}$ for $0 \leq i \leq 2$. Note that $u_i u_{i+1}$ is contained in G_i^1 , G_{i+1}^1 and G_i^2 , $u_i v_i$ is contained in G_{i+1}^1 , G_{i+2}^1 , G_i^2 and G_{i+2}^2 , $v_i v_{i+1}$ is contained in G_{i+2}^1 , G_{i+1}^2 and G_{i+2}^2 for $0 \leq i \leq 2$. Thus $F_i = \{x_i, x_{i+1}, y_i\}$, $F_{i+3} = \{x_{i+1}, x_{i+2}, y_i, y_{i+2}\}$, $F_{i+6} = \{x_{i+2}, y_{i+1}, y_{i+2}\}$ for $0 \leq i \leq 2$, and so $r(H_{P_{3,1}, \Psi}) = 4$ and $s(H_{P_{3,1}, \Psi}) = 2$. By Lemma 2.2, we have $Ar(P_{3,1}, C_5) = Ar_{P_{3,1}}(\Psi) \leq |E(P_{3,1})| - \lceil \frac{2|\Psi|}{4+2} \rceil = 9 - 2 = 7$. ■




Fig. 1 $P_{3,1}$ and $P_{5,2}$ have no rainbow C_5 under the coloring ϕ

Lemma 3.2 $Ar(P_{5,1}, C_5) = 13$.

Proof. Let $P'_{5,1} = P_{5,1} - E'$, where $E' = \{u_i v_i \mid 0 \leq i \leq 4\}$, and let $G_1 := u_0 u_1 u_2 u_3 u_4 u_0$, $G_2 := v_0 v_1 v_2 v_3 v_4 v_0$. Then $\Psi' = \{G_1, G_2\}$ is the set of all copies of C_5 in $P'_{5,1}$, and then $Ar(P'_{5,1}, C_5) = Ar_{P'_{5,1}}(\Psi')$. Note that each edge in $E(G_i)$ is only contained in G_i . Thus $H_{P'_{5,1}, \Psi'}$ is a graph obtained from two isolated vertices by adding five loops on each isolated vertex, and so $M(H_{P'_{5,1}, \Psi'}) = 0$. By Lemma 2.1, we have $Ar(P'_{5,1}, C_5) = Ar_{P'_{5,1}}(\Psi') = |E(P'_{5,1})| - |\Psi'| + M(H_{P'_{5,1}, \Psi'}) = 8$. Since there is no copy of C_5 in $P_{5,1}$ containing any edge of E' , $Ar(P_{5,1}, C_5) = |E'| + Ar(P'_{5,1}, C_5) = 13$. ■

Lemma 3.3 $Ar(P_{5,2}, C_5) = 10$.

Proof. Let $G_i^1 := u_i u_{i+1} v_{i+1} v_{i+3} v_i u_i$, $G_i^2 := u_i u_{i+1} u_{i+2} v_{i+2} v_i u_i$ for $0 \leq i \leq 4$, and let $G^3 := u_0 u_1 u_2 u_3 u_4$, $G^4 := v_0 v_2 v_4 v_1 v_3 v_0$. Then $\Psi = \{G_i^1, G_i^2, G^3, G^4 \mid 0 \leq i \leq 4\}$ is the set of all copies of C_5 in $P_{5,2}$. Hence $Ar(P_{5,2}, C_5) = Ar_{P_{5,2}}(\Psi)$.

First we show that $Ar(P_{5,2}, C_5) \geq 10$. Let $\phi : E(P_{5,2}) \rightarrow \{1, 2, \dots, 10\}$ be an edge-coloring of $P_{5,2}$ defined as follows:

$$\phi(e) = \begin{cases} 1, & \text{if } e \in E(u_0) \setminus \{u_0 v_0\}; \\ 2, & \text{if } e \in E(v_2); \\ 3, & \text{if } e \in E(v_3), \end{cases}$$

and every other edges obtains distinct colorings from $\{4, \dots, 10\}$ (see Fig. 1(b)). Note that each $C_5 \in \Psi$ contains the edges $\{u_0 u_1, u_0 u_4\}$, or two edges of $E(v_i)$ ($i \in \{2, 3\}$), and so $P_{5,2}$ has no rainbow C_5 . Hence, $Ar(P_{5,2}, C_5) \geq 10$.

Next we show that $Ar(P_{5,2}, C_5) \leq 10$. Let $V(H_{P_{5,2}, \Psi}) = \{x_i, y_i, z, w \mid 0 \leq i \leq 4\}$ and $E(H_{P_{5,2}, \Psi}) = \{F_0, \dots, F_{14}\}$, where x_i, y_i, z and w correspond to G_i^1, G_i^2, G^3 and G^4 in Ψ , respectively, each F_i corresponds to $u_i u_{i+1}$, F_{i+5} corresponds to $u_i v_i$, and F_{i+10} corresponds to $v_i v_{i+2}$ for $0 \leq i \leq 4$. Note that each $u_i u_{i+1}$ is contained in G_i^1, G_i^2, G_{i+4}^2 and G^3 , each spoke $u_i v_i$ is contained in G_i^1, G_{i+4}^1, G_i^2 and G_{i+3}^2 , and each $v_i v_{i+2}$ is contained in $G_{i+2}^1, G_{i+4}^1, G_i^2$ and G^4 . Thus $F_i = \{x_i, y_i, y_{i+4}, z\}$, $F_{i+5} = \{x_i, x_{i+4}, y_i, y_{i+3}\}$ and $F_{i+10} = \{x_{i+2}, x_{i+4}, y_i, w\}$ for $0 \leq i \leq 4$. It follows that $r(H_{P_{5,2}, \Psi}) = 4$ and $s(H_{P_{5,2}, \Psi}) = 2$.

Now we will show that $H_{P_{5,2}, \Psi} \in \mathcal{P}_{4,12}$. Suppose that $H_{P_{5,2}, \Psi} \notin \mathcal{P}_{4,12}$. Then there exists a 11-partition $\{\mathcal{F}_1, \mathcal{F}_2, \dots, \mathcal{F}_{11}\}$ of $E(H_{P_{5,2}, \Psi})$ with $|\bigcup_{i=1}^{11} L_i| = 12$. Assume that $|\mathcal{F}_1| \geq \dots \geq |\mathcal{F}_{11}| \geq 1$. Then $2 \leq |\mathcal{F}_1| \leq 5$ and $\sum_{i=1}^{11} (|\mathcal{F}_i| - 1) = 4$. If $|\mathcal{F}_2| \leq 2$, then by (1), $|\bigcup_{i=1}^{11} L_i| \leq \sum_{i=1}^{11} |L_i| \leq 2|\mathcal{F}_1| + \sum_{i=2}^{11} 2(|\mathcal{F}_i| - 1) = 2 \sum_{i=1}^{11} (|\mathcal{F}_i| - 1) + 2 = 10 < 12$, a contradiction. So $|\mathcal{F}_2| \geq 3$, and then $|\mathcal{F}_1| = |\mathcal{F}_2| = 3$. By (1),

$$12 = \left| \bigcup_{i=1}^{11} L_i \right| \leq \sum_{i=1}^{11} |L_i| \leq 2|\mathcal{F}_1| + 2|\mathcal{F}_2| + \sum_{i=3}^{11} 2(|\mathcal{F}_i| - 1) = 2 \sum_{i=1}^{11} (|\mathcal{F}_i| - 1) + 4 = 12,$$

which implies $|L_i| = 2|\mathcal{F}_i| = 6$ for $i = 1, 2$, $L_1 \cup L_2 = V(H_{P_{5,2},\Psi})$ and $L_1 \cap L_2 = \emptyset$. Hence, each \mathcal{F}_i ($i \in \{1, 2\}$) is a barrier. Without loss of generality, we assume $F_0 \in \mathcal{F}_1$. Then $x_0, y_0, y_4, z \in L_1$, and then $\{F_{10}, F_{11}, F_{13}, F_{14}\} \cap \mathcal{F}_2 = \emptyset$ as $y_0 \in F_{10}, x_0 \in F_{11} \cap F_{13}, y_4 \in F_{14}$. Note that $F(w) = \{F_i \mid 10 \leq i \leq 14\}$, and so $w \notin L_2$ as $|\mathcal{F}_2| = 3$. Then $w \in L_1$, and then $|\mathcal{F}_1| \geq 4$, a contradiction. Therefore, $H_{P_{5,2},\Psi} \in \mathcal{P}_{4,12}$. By Lemma 2.4, $Ar(P_{5,2}, C_5) = Ar_{P_{5,2}}(\Psi) \leq |E(P_{5,2})| - (4 + 1) = 10$. ■

Lemma 3.4 $Ar(P_{10,2}, C_5) = 22$.

Proof. Let $G_i^1 := u_i u_{i+1} u_{i+2} v_{i+2} v_i u_i$ for $0 \leq i \leq 9$, and let $G^2 := v_0 v_2 v_4 v_6 v_8 v_0$, $G^3 := v_1 v_3 v_5 v_7 v_9 v_1$. Then $\Psi = \{G_i^1, G^2, G^3 \mid 0 \leq i \leq 9\}$ is the set of all copies of C_5 in $P_{10,2}$, and then $Ar(P_{10,2}, C_5) = Ar_{P_{10,2}}(\Psi)$.

Note that each edge $u_i u_{i+1}$ is contained in G_i^1 and G_{i+9}^1 ; each spoke $u_i v_i$ is contained in G_i^1 and G_{i+8}^1 ; each $v_i v_{i+2}$ is contained in G_i^1 and G^2 for $i \in \{0, 2, 4, 6, 8\}$; each edge $v_i v_{i+2}$ is contained in G_i^1 and G^3 for $i \in \{1, 3, 5, 7, 9\}$. Thus $H_{P_{10,2},\Psi}$ is a simple graph of order 12 (see Fig. 2). Since there exist four vertex-disjoint cycles $w_0 w_1 w_2 w_0$, $w_3 w_4 w_5 w_3$, $w_6 w_8 w_{11} w_6$ and $w_7 w_9 w_{10} w_7$ in $H_{P_{10,2},\Psi}$, $M(H_{P_{10,2},\Psi}) = 4$. By Lemma 2.1, $Ar(P_{10,2}, C_5) = Ar_{P_{10,2}}(\Psi) = |E(P_{10,2})| - |\Psi| + M(H_{P_{10,2},\Psi}) = 22$. ■




Fig. 2 $P_{10,2}$ and $H_{P_{10,2},\Psi}$

Lemma 3.5 $Ar(P_{n,2}, C_5) = \lfloor \frac{7n}{3} \rfloor$ for $n \geq 6$ and $n \neq 10$.

Proof. Let $G_i := u_i u_{i+1} u_{i+2} v_{i+2} v_i u_i$ for $0 \leq i \leq n-1$. Then $\Psi = \{G_i \mid 0 \leq i \leq n-1\}$ is the set of all copies of C_5 in $P_{n,2}$, and then $Ar(P_{n,2}, C_5) = Ar_{P_{n,2}}(\Psi)$. Note that each edge $u_i u_{i+1}$ is contained in G_i and G_{i+n-1} ; $u_i v_i$ is contained in G_i and G_{i+n-2} ; $v_i v_{i+2}$ is contained in G_i for $0 \leq i \leq n-1$. Thus $H_{P_{n,2},\Psi}$ is a graph obtained from

$C_n = w_0w_1 \cdots w_{n-1}w_0$ by adding a loop on w_i and edge w_iw_{i+2} for all $0 \leq i \leq n-1$. Since there exist $\lfloor \frac{n}{3} \rfloor$ vertex-disjoint cycles $w_{3i}w_{3i+1}w_{3i+2}w_{3i}$ in $H_{P_{n,2},\Psi}$ for $0 \leq i \leq \lfloor \frac{n}{3} \rfloor - 1$, we have $M(H_{P_{n,2},\Psi}) = \lfloor \frac{n}{3} \rfloor$. By Lemma 2.1, $Ar(P_{n,2}, C_5) = Ar_{P_{n,2}}(\Psi) = |E(P_{n,2})| - |\Psi| + M(H_{P_{n,2},\Psi}) = \lfloor \frac{7n}{3} \rfloor$. ■

Lemma 3.6 $Ar(P_{n,k}, C_5) = \frac{14n}{5}$ for $k \geq 3$ and $k \in \{\frac{n}{5}, \frac{2n}{5}\}$.

Proof. Denote $P'_{n,k} := P_{n,k} - E'$, where $E' = \{u_iu_{i+1}, u_iv_i \mid 0 \leq i \leq n-1\}$. Let $G_i := v_iv_{i+k}v_{i+2k}v_{i+3k}v_{i+4k}v_i$ for $0 \leq i \leq \frac{n}{5}-1$. Then $\Psi' = \{G_i \mid 0 \leq i \leq \frac{n}{5}-1\}$ is the set of all copies of C_5 in $P'_{n,k}$ for $k \in \{\frac{n}{5}, \frac{2n}{5}\}$, and then $Ar(P'_{n,k}, C_5) = Ar_{P'_{n,k}}(\Psi')$. Note that each edge in $P'_{n,k}$ is only contained in some G_i . Thus $H_{P'_{n,k},\Psi'}$ is a graph obtained from k isolated vertices by adding five loops on each isolated vertex. Then $M(H_{P'_{n,k},\Psi'}) = 0$. By Lemma 2.1, $Ar(P'_{n,k}, C_5) = Ar_{P'_{n,k}}(\Psi') = |E(P'_{n,k})| - |\Psi'| + M(H_{P'_{n,k},\Psi'}) = \frac{4n}{5}$ for $k \in \{\frac{n}{5}, \frac{2n}{5}\}$. Since there is no copy of C_5 in $P_{n,k}$ containing any edge of E' , we have $Ar(P_{n,k}, C_5) = |E'| + Ar(P'_{n,k}, C_5) = \frac{14n}{5}$. ■

Lemma 3.7 $Ar(P_{n,k}, C_5) = \lfloor \frac{7n}{3} \rfloor$ for $k = \frac{n-1}{2} \geq 3$.

Proof. Note that $k = \frac{n-1}{2} \geq 3$. Hence $n \geq 7$, moreover, $n \geq 11$ if $n \equiv 2 \pmod{3}$.

Let $G_i := u_iu_{i+1}v_{i+1}v_{i+\frac{n+1}{2}}v_iu_i$ for $0 \leq i \leq n-1$. Then $\Psi = \{G_i \mid 0 \leq i \leq n-1\}$ is the set of all copies of C_5 in $P_{n,k}$, and then $Ar(P_{n,k}, C_5) = Ar_{P_{n,k}}(\Psi)$. Note that each u_iu_{i+1} is contained in G_i , u_iv_i is contained in G_i and G_{i+n-1} , $v_iv_{i+\frac{n-1}{2}}$ is contained in $G_{i+\frac{n-1}{2}}$ and G_{i+n-1} . Thus $H_{P_{n,k},\Psi}$ is a graph obtained from the cycle $w_0w_1 \cdots w_{n-1}w_0$ by adding a loop on w_i and an edge $w_{i+\frac{n-1}{2}}w_{i+n-1}$ for all $0 \leq i \leq n-1$. Then $C_j = w_jw_{j+\frac{n-1}{2}}w_{j+\frac{n+1}{2}}w_j$ is a 3-cycle in $H_{P_{n,k},\Psi}$ for $0 \leq j \leq n-1$. Let \mathcal{C} be the set of vertex-disjoint 3-cycles in $H_{P_{n,k},\Psi}$. Then

$$\mathcal{C} = \begin{cases} \{C_{3i} \mid 0 \leq i \leq \frac{n}{3} - 1\}, & \text{if } n \equiv 0 \pmod{3}; \\ \{C_{2+3i}, C_{n-3i} \mid 0 \leq i \leq \frac{n-7}{6}\}, & \text{if } n \equiv 1 \pmod{3}; \\ \{C_0, C_{2+3i}, C_{n-2-3i} \mid 0 \leq i \leq \frac{n-11}{6}\}, & \text{otherwise.} \end{cases}$$

This implies that $M(H_{P_{n,k},\Psi}) = \lfloor \frac{n}{3} \rfloor$. By Lemma 2.1, we have $Ar(P_{n,k}, C_5) = Ar_{P_{n,k}}(\Psi) = |E(P_{n,k})| - |\Psi| + M(H_{P_{n,k},\Psi}) = \lfloor \frac{7n}{3} \rfloor$. ■

By Lemmas 3.1-3.7, we complete the proof of Theorem 1.2.

4 Anti-Ramsey number $Ar(P_{n,k}, C_6)$

In this section, we will determine the anti-Ramsey number $Ar(P_{n,k}, C_6)$. If G contains no C_6 , then $Ar(P_{n,k}, C_6) = |E(P_{n,k})| = 3n$. Thus by Proposition 1.1(ii), it is only necessary to consider the anti-Ramsey number $Ar(P_{n,k}, C_6)$ when $k \in \{1, 3, \frac{n}{6}, \frac{n-1}{3}, \frac{n+1}{3}, \frac{n-2}{2}\}$. Moreover, if $k = 2$, then $n \in \{5, 6, 7, 12\}$.

4.1 $k = 1$

In this subsection, we focus on the case that $k = 1$. Then $n \geq 3$.

Lemma 4.1 $Ar(P_{3,1}, C_6) = 7$.

Proof. Let $G_i := u_i u_{i+1} u_{i+2} v_{i+2} v_{i+1} v_i u_i$ for $0 \leq i \leq 2$. Then $\Psi = \{G_0, G_1, G_2\}$ is the set of all copies of C_6 in $P_{3,1}$, and then $Ar(P_{3,1}, C_6) = Ar_{P_{3,1}}(\Psi)$. Note that each $u_i u_{i+1}$ is contained in G_i and G_{i+2} , $u_i v_i$ is contained in G_i and G_{i+1} , and $v_i v_{i+1}$ is contained in G_i and G_{i+2} for $0 \leq i \leq 2$. Thus $H_{P_{3,1}, \Psi}$ is a graph obtained from the 3-cycle $w_0 w_1 w_2 w_0$ by adding two parallel edges on each edge of C_3 . Then $M(H_{P_{3,1}, \Psi}) = 1$. By Lemma 2.1, $Ar(P_{3,1}, C_6) = Ar_{P_{3,1}}(\Psi) = |E(P_{3,1})| - |\Psi| + M(H_{P_{3,1}, \Psi}) = 7$. ■




Fig. 3 $P_{4,1}$ and $P_{6,1}$ have no rainbow C_6 under the coloring ϕ .

Lemma 4.2 $Ar(P_{4,1}, C_6) = 9$.

Proof. Let $G_i^1 := u_i u_{i+1} v_{i+1} v_{i+2} v_{i+3} v_i u_i$, $G_i^2 := u_i u_{i+1} u_{i+2} v_{i+2} v_{i+1} v_i u_i$, $G_i^3 := u_i u_{i+1} u_{i+2} v_{i+2} v_{i+3} v_i u_i$ and $G_i^4 := u_i u_{i+1} u_{i+2} u_{i+3} v_{i+3} v_i u_i$ for $0 \leq i \leq 3$. Then $\Psi = \{G_i^j \mid 0 \leq i \leq 3, 1 \leq j \leq 4\}$ is the set of all C_6 in $P_{4,1}$, and then $Ar(P_{4,1}, C_6) = Ar_{P_{4,1}}(\Psi)$.

Let $\phi : E(P_{4,1}) \rightarrow \{1, 2, \dots, 9\}$ be an edge-coloring of $P_{4,1}$ such that $\phi(u_i v_i) = 1$ for $0 \leq i \leq 3$ and every other edges having distinct colorings from $\{2, \dots, 9\}$ (see Fig. 3(a)). Note that each $C_6 \in \Psi$ contains two spokes, and hence $P_{4,1}$ has no rainbow C_6 under the coloring ϕ . So, $Ar(P_{4,1}, C_6) \geq 9$.

Now we will show that $Ar(P_{4,1}, C_6) \geq 9$. Let $V(H_{P_{4,1}, \Psi}) = \{x_i, y_i, z_i, w_i \mid 0 \leq i \leq 3\}$ and $E(H_{P_{4,1}, \Psi}) = \{F_0, \dots, F_{11}\}$, where x_i, y_i, z_i, w_i correspond to $G_i^1, G_i^2, G_i^3, G_i^4$ in Ψ , respectively, each F_i corresponds to $u_i u_{i+1}$, F_{i+4} corresponds to $u_i v_i$ and F_{i+8} corresponds to $v_i v_{i+1}$ for $0 \leq i \leq 3$. Note that each $u_i u_{i+1}$ is contained in $G_i^1, G_i^2, G_{i+3}^2, G_i^3, G_{i+3}^3, G_i^4, G_{i+2}^4$ and G_{i+3}^4 , $u_i v_i$ is contained in $G_i^1, G_{i+3}^1, G_i^2, G_{i+2}^2, G_i^3, G_{i+2}^3, G_i^4$ and G_{i+1}^4 , and $v_i v_{i+1}$ is contained in $G_{i+1}^1, G_{i+2}^1, G_{i+3}^1, G_i^2, G_{i+3}^2, G_{i+1}^3, G_{i+2}^3$ and G_{i+1}^4 for $0 \leq i \leq 3$. It follows that $F_i = \{x_i, y_i, y_{i+3}, z_i, z_{i+3}, w_i, w_{i+2}, w_{i+3}\}$, $F_{i+4} = \{x_i, x_{i+3}, y_i, y_{i+2}, z_i, z_{i+2}, w_i, w_{i+1}\}$ and $F_{i+8} = \{x_{i+1}, x_{i+2}, x_{i+3}, y_i, y_{i+3}, z_{i+1}, z_{i+2}, w_{i+1}\}$ for $0 \leq i \leq 3$, and thus $r(H_{P_{4,1}, \Psi}) = 8$. Obviously, $s(H_{P_{4,1}, \Psi}) = 4$. By Lemma 2.2, we have $Ar(P_{4,1}, C_6) = Ar_{P_{4,1}}(\Psi) \leq |E(P_{4,1})| - \lceil \frac{2|\Psi|}{8+4} \rceil = 9$. ■

Lemma 4.3 $Ar(P_{6,1}, C_6) = 14$.

Proof. Let $G_i^1 := u_i u_{i+1} u_{i+2} v_{i+2} v_{i+1} v_i u_i$ for $0 \leq i \leq 5$, $G^2 := u_0 u_1 u_2 u_3 u_4 u_5 u_6 u_0$ and $G^3 := v_0 v_1 v_2 v_3 v_4 v_5 v_6 v_0$. Then $\Psi = \{G_i^1, G^2, G^3 \mid 0 \leq i \leq 5\}$ is the set of all copies of C_6 in $P_{6,1}$, and $Ar(P_{6,1}, C_6) = Ar_{P_{6,1}}(\Psi)$.

Let $\phi : E(P_{6,1}) \rightarrow \{1, 2, \dots, 14\}$ be an edge-coloring of $P_{6,1}$ defined as follows:

$$\phi(e) = \begin{cases} 1, & \text{if } e \in E(u_0); \\ 2, & \text{if } e \in E(v_3), \end{cases}$$

and every other edges obtains distinct colorings from $\{3, \dots, 14\}$ (see Fig. 3(b)). Note that each $C_6 \in \Psi$ contains two edges of $E(u_0)$ (or $E(v_3)$), and so $P_{6,1}$ has no rainbow C_6 . Hence, $Ar(P_{6,1}, C_6) \geq 14$.

On the other hand, let $V(H_{P_{6,1}, \Psi}) = \{x_i, y, z \mid 0 \leq i \leq 5\}$ and $E(H_{P_{6,1}, \Psi}) = \{F_0, \dots, F_{17}\}$, where x_i, y and z correspond to G_i^1, G^2 and G^3 in Ψ , respectively, each F_i corresponds to $u_i u_{i+1}$, F_{i+6} corresponds to $u_i v_i$ and each F_{i+12} corresponds to $v_i v_{i+1}$ for $0 \leq i \leq 5$. Note that each $u_i u_{i+1}$ is contained in G_i^1, G_{i+5}^1 and G^2 , $u_i v_i$ is contained in G_i^1 and G_{i+4}^1 , and $v_i v_{i+1}$ is contained in G_i^1, G_{i+5}^1 and G^3 for $0 \leq i \leq 5$. It follows that $F_i = \{x_i, x_{i+5}, y\}$, $F_{i+6} = \{x_i, x_{i+4}\}$ and $F_{i+12} = \{x_i, x_{i+5}, z\}$ for $0 \leq i \leq 5$, and then $r(H_{P_{6,1}, \Psi}) = 3$ and $s(H_{P_{6,1}, \Psi}) = 2$. By Lemma 2.2, we have $Ar(P_{6,1}, C_6) = Ar_{P_{6,1}}(\Psi) \leq |E(P_{6,1})| - \lceil \frac{2|\Psi|}{3+2} \rceil = 14$. ■

Lemma 4.4 $Ar(P_{n,1}, C_6) = \lfloor \frac{5n}{2} \rfloor$ for $n \geq 5$ and $n \neq 6$.

Proof. Since $n \geq 5$ and $n \neq 6$, $\Psi = \{G_i \mid 0 \leq i \leq n-1\}$ is the set of all copies of C_6 in $P_{n,1}$, where $G_i := u_i u_{i+1} u_{i+2} v_{i+2} v_{i+1} v_i u_i$. Note that each $u_i u_{i+1}$ is contained in G_i and G_{i+n-1} , $u_i v_i$ is contained in G_i and G_{i+n-2} , $v_i v_{i+1}$ is contained in G_i and G_{i+n-1} for $0 \leq i \leq n-1$. Thus $H_{P_{n,1}, \Psi}$ is a graph obtained from $C_n = w_0 w_1 \cdots w_{n-1} w_0$ by adding one parallel edge on each edge of C_n and edge $w_i w_{i+2}$ for $0 \leq i \leq n-1$. Then $M(H_{P_{n,1}, \Psi}) = \lfloor \frac{n}{2} \rfloor$. By Lemma 2.1, $Ar(P_{n,1}, C_6) = |E(P_{n,1})| - |\Psi| + M(H_{P_{n,1}, \Psi}) = \lfloor \frac{5n}{2} \rfloor$. ■

By Lemmas 4.1-4.4, the proof of Theorem 1.3 is completed.

4.2 $k = 2$

In this subsection, we focus on the case that $k = 2$. Then $n \geq 5$. By Proposition 1.1(ii), we have $n \in \{5, 6, 7, 12\}$.




Fig. 4 $P_{5,2}$ and $P_{6,2}$ have no rainbow C_6 under the coloring ϕ .

Lemma 4.5 $Ar(P_{5,2}, C_6) = 11$.

Proof. Let $G_i^1 = u_i u_{i+1} v_{i+1} v_{i+4} v_{i+2} v_i u_i$, $G_i^2 = u_i u_{i+1} u_{i+2} u_{i+3} v_{i+3} v_i u_i$ for $0 \leq i \leq 4$. Then $\Psi = \{G_i^1, G_i^2 \mid 0 \leq i \leq 4\}$ is the set of all copies of C_6 in $P_{5,2}$.

Let $\phi : E(P_{5,2}) \rightarrow \{1, 2, \dots, 11\}$ be an edge-coloring of $P_{5,2}$ such that $\phi(u_i v_i) = 1$ for $0 \leq i \leq 4$ and every other edges obtains distinct colorings from $\{2, \dots, 11\}$ (see Fig. 4(a)). Note that each $C_6 \in \Psi$ contains two spokes, and so $P_{5,2}$ has no rainbow C_6 . Hence, $Ar(P_{5,2}, C_6) \geq 11$.

Next we will show that $Ar(P_{5,2}, C_6) \leq 11$. Let $V(H_{P_{5,2}, \Psi}) = \{x_i, y_i \mid 0 \leq i \leq 4\}$ and $E(H_{P_{5,2}, \Psi}) = \{F_0, \dots, F_{14}\}$, where x_i and y_i correspond to G_i^1 and G_i^2 in Ψ , respectively,

each F_i corresponds to $u_i u_{i+1}$, F_{i+5} corresponds to $u_i v_i$, and F_{i+10} corresponds to $v_i v_{i+2}$ for $0 \leq i \leq 4$. Note that each edge $u_i u_{i+1}$ is contained in G_i^1 , G_i^2 , G_{i+3}^2 and G_{i+4}^2 , $u_i v_i$ is contained in G_i^1 , G_{i+4}^1 , G_i^2 and G_{i+2}^2 , $v_i v_{i+2}$ is contained in G_i^1 , G_{i+1}^1 , G_{i+3}^1 and G_{i+2}^2 for $0 \leq i \leq 4$. It follows that $F_i = \{x_i, y_i, y_{i+3}, y_{i+4}\}$, $F_{i+5} = \{x_i, x_{i+4}, y_i, y_{i+2}\}$ and $F_{i+10} = \{x_i, x_{i+1}, x_{i+3}, y_{i+2}\}$ for $0 \leq i \leq 4$, and thus $r(H_{P_{5,2}, \Psi}) = 4$ and $s(H_{P_{5,2}, \Psi}) = 2$. By Lemma 2.2, we have $Ar(P_{5,2}, C_6) \leq |E(P_{5,2})| - \lceil \frac{2|\Psi|}{4+2} \rceil = 15 - \lceil \frac{10}{3} \rceil = 11$. ■

Lemma 4.6 $Ar(P_{6,2}, C_6) = 14$.

Proof. Let $G_i^1 := u_i u_{i+1} v_{i+1} v_{i+3} v_{i+5} u_{i+5} u_i$ for $0 \leq i \leq 5$, and $G^2 := u_0 u_1 u_2 u_3 u_4 u_5 u_0$. Then $\Psi = \{G_i^1, G^2 \mid 0 \leq i \leq 5\}$ is the set of all C_6 in $P_{6,2}$, and $Ar(P_{6,2}, C_6) = Ar_{P_{6,2}}(\Psi)$.

Let $\phi : E(P_{6,2}) \rightarrow \{1, 2, \dots, 14\}$ be an edge-coloring of $P_{6,2}$ defined as follows:

$$\phi(e) = \begin{cases} 1, & \text{if } e \in E(u_0); \\ 2, & \text{if } e \in E(u_3), \end{cases}$$

and every other edges obtains distinct colorings from $\{3, \dots, 14\}$ (see Fig. 4(b)). Note that each $C_6 \in \Psi$ contains two edges of $E(u_i)$ ($i \in \{0, 3\}$), and so $P_{6,2}$ has no rainbow C_6 . Hence, $Ar(P_{6,2}, C_6) \geq 14$.

Next we will show that $Ar(P_{6,2}, C_6) \leq 14$. Let $V(H_{P_{6,2}, \Psi}) = \{x_i, y \mid 0 \leq i \leq 5\}$ and $E(H_{P_{6,2}, \Psi}) = \{F_0, \dots, F_{17}\}$, where x_i and y correspond to G_i^1 and G^2 in Ψ , respectively, each F_i corresponds to $u_i u_{i+1}$, F_{i+6} corresponds to $u_i v_i$, and F_{i+12} corresponds to $v_i v_{i+2}$ for $0 \leq i \leq 5$. Note that each edge $u_i u_{i+1}$ is contained in G_i , G_{i+1} and G^2 , $u_i v_i$ is contained in G_{i+1} and G_{i+5} , $v_i v_{i+2}$ is contained in G_{i+3} and G_{i+5} for $0 \leq i \leq 5$. Hence $F_i = \{x_i, x_{i+1}, y\}$, $F_{i+6} = \{x_{i+1}, x_{i+5}\}$ and $F_{i+12} = \{x_{i+3}, x_{i+5}\}$ for $0 \leq i \leq 5$, and thus $r(H_{P_{6,2}, \Psi}) = 3$ and $s(H_{P_{6,2}, \Psi}) = 2$. By Lemma 2.3, $H_{P_{6,2}, \Psi} \in \mathcal{P}_{3,7}$, and then by Lemma 2.4, we have $Ar(P_{6,2}, C_6) = Ar_{P_{6,2}}(\Psi) \leq |E(P_{6,2})| - (3 + 1) = 14$. ■

Lemma 4.7 $Ar(P_{7,2}, C_6) = 17$.

Proof. Let $G_i := u_i u_{i+1} v_{i+1} v_{i+3} v_{i+5} v_i u_i$ for $0 \leq i \leq 6$. Then $\Psi = \{G_i \mid 0 \leq i \leq 6\}$ is the set of all copies of C_6 in $P_{7,2}$, and then $Ar(P_{7,2}, C_6) = Ar_{P_{7,2}}(\Psi)$.

Let $\phi : E(P_{7,2}) \rightarrow \{1, 2, \dots, 17\}$ be an edge-coloring of $P_{7,2}$ defined as follows:

$$\phi(e) = \begin{cases} 1, & \text{if } e \in E(v_3); \\ 2, & \text{if } e \in E(v_4), \end{cases}$$

and every other edges obtains distinct colorings from $\{3, \dots, 17\}$ (see Fig. 5(a)). Note that each $C_6 \in \Psi$ contains two edges of $E(v_i)$ ($i \in \{3, 4\}$), and so $P_{7,2}$ has no rainbow C_6 . Hence, $Ar(P_{7,2}, C_6) \geq 17$.

Next we will show that $Ar(P_{7,2}, C_6) \leq 17$. Let $V(H_{P_{7,2}, \Psi}) = \{x_i \mid 0 \leq i \leq 6\}$ and $E(H_{P_{7,2}, \Psi}) = \{F_0, \dots, F_{20}\}$, where x_i corresponds to G_i in Ψ , each F_i corresponds to $u_i u_{i+1}$, F_{i+7} corresponds to $u_i v_i$ and F_{i+14} corresponds to $v_i v_{i+2}$ for $0 \leq i \leq 6$. Note that each $u_i u_{i+1}$ is contained in G_i , $u_i v_i$ is contained in G_i and G_{i+6} , $v_i v_{i+2}$ is contained in G_{i+2} , G_{i+4} and G_{i+6} for $0 \leq i \leq 6$. It follows that $F_i = \{x_i\}$, $F_{i+6} = \{x_i, x_{i+6}\}$ and $F_{i+12} = \{x_{i+2}, x_{i+4}, x_{i+6}\}$ for $0 \leq i \leq 6$, and thus $r(H_{P_{7,2}, \Psi}) = 3$ and $s(H_{P_{7,2}, \Psi}) = 2$. By Lemma 2.3, $H_{P_{7,2}, \Psi} \in \mathcal{P}_{3,7}$, and then by Lemma 2.4, $Ar(P_{7,2}, C_6) = Ar_{P_{7,2}}(\Psi) \leq |E(P_{7,2})| - (3 + 1) = 17$. ■




Fig. 5 $P_{7,2}$ and $P_{8,3}$ have no rainbow C_6 under the coloring ϕ .

Lemma 4.8 $Ar(P_{12,2}, C_6) = 34$.

Proof. Let $P'_{12,2} = P_{12,2} - E'$, where $E' = \{u_i u_{i+1}, u_i v_i \mid 0 \leq i \leq 11\}$, and let $G_i := v_i v_{i+2} v_{i+4} v_{i+6} v_{i+8} v_{i+10} v_i$ for $0 \leq i \leq 1$. Then $\Psi' = \{G_0, G_1\}$ is the set of all C_6 in $P'_{n,k}$, and then $Ar(P'_{12,2}, C_6) = Ar_{P'_{12,2}}(\Psi')$. Note that each edge in $P'_{12,2}$ is only contained in some G_i . Thus $H_{P'_{12,2}, \Psi'}$ is a graph obtained from two isolated vertices by adding six loops at each vertex. Then $M(H_{P'_{12,2}, \Psi'}) = 0$. By Lemma 2.1, $Ar(P'_{12,2}, C_6) = Ar_{P'_{12,2}}(\Psi') = |E(P'_{12,2})| - |\Psi'| + M(H_{P'_{12,2}, \Psi'}) = 10$. Since there is no C_6 in $P_{12,2}$ containing any edge of E' , $Ar(P_{n,k}, C_6) = |E'| + Ar(P'_{n,k}, C_6) = 34$. ■

By Lemmas 4.5-4.8, the proof of Theorem 1.4 is completed.

4.3 $k = 3$

In this subsection, we focus on the case that $k = 3$. Then $n \geq 7$.

Lemma 4.9 $Ar(P_{8,3}, C_6) = 17$.

Proof. Let $G_i^1 := u_i u_{i+1} u_{i+2} u_{i+3} v_{i+3} v_i u_i$, $G_i^2 := u_i u_{i+1} v_{i+1} v_{i+4} v_{i+7} u_{i+7} u_i$ and $G_i^3 := u_i u_{i+1} v_{i+1} v_{i+6} v_{i+3} v_i u_i$ for $0 \leq i \leq 7$. Then $\Psi = \{G_i^1, G_i^2, G_i^3 \mid 0 \leq i \leq 7\}$ is the set of all copies of C_6 in $P_{8,3}$, and then $Ar(P_{8,3}, C_6) = Ar_{P_{8,3}}(\Psi)$.

Let $\phi : E(P_{8,3}) \rightarrow \{1, 2, \dots, 17\}$ be an edge-coloring of $P_{8,3}$ such that $\phi(u_i v_i) = 1$ for $0 \leq i \leq 7$ and every other edges having distinct colorings from $\{2, \dots, 17\}$ (see Fig. 5(b)). Note that each $C_6 \in \Psi$ contains two spokes, and hence $P_{8,3}$ has no rainbow C_6 under the coloring ϕ . So, $Ar(P_{8,3}, C_6) \geq 17$.

Next we will show that $Ar(P_{8,3}, C_6) \leq 17$. Let $V(H_{P_{8,3}, \Psi}) = \{x_i, y_i, z_i \mid 0 \leq i \leq 7\}$ and $E(H_{P_{8,3}, \Psi}) = \{F_i^0, F_i^1, F_i^2 \mid 0 \leq i \leq 7\}$, where x_i, y_i and z_i correspond to G_i^1, G_i^2 and G_i^3 in Ψ , respectively, each F_i^1 corresponds to $u_i u_{i+1}$, F_i^2 corresponds to $u_i v_i$, and F_i^3 corresponds to $v_i v_{i+3}$ for $0 \leq i \leq 7$. Note that each edge $u_i u_{i+1}$ is contained in $G_i^1, G_{i+6}^1, G_{i+7}^1, G_i^2, G_{i+1}^2$ and G_i^3 ; each spoke $u_i v_i$ is contained in $G_i^1, G_{i+5}^1, G_{i+1}^2, G_{i+7}^2, G_i^3$ and G_{i+7}^3 ; each edge $v_i v_{i+3}$ is contained in $G_i^1, G_{i+4}^2, G_{i+7}^2, G_i^3, G_{i+2}^3$ and G_{i+5}^3 for $0 \leq i \leq 7$. It follows that $F_i^1 = \{x_i, x_{i+6}, x_{i+7}, y_i, y_{i+1}, z_i\}$, $F_i^2 = \{x_i, x_{i+5}, y_{i+1}, y_{i+7}, z_i, z_{i+7}\}$ and $F_i^3 = \{x_i, y_{i+4}, y_{i+7}, z_i, z_{i+2}, z_{i+5}\}$ for $0 \leq i \leq 7$, and thus $r(H_{P_{8,3}, \Psi}) = 6$, $s(H_{P_{8,3}, \Psi}) = 3$.

Now we will show that $H_{P_{8,3}, \Psi} \in \mathcal{P}_{6,24}$. Suppose that $H_{P_{8,3}, \Psi} \notin \mathcal{P}_{6,24}$. Then there exists a 18-partition $\{\mathcal{F}_1, \mathcal{F}_2, \dots, \mathcal{F}_{18}\}$ of $E(H_{P_{8,3}, \Psi})$ with $|\bigcup_{i=1}^{18} L_i| = 24$. Assume that $|\mathcal{F}_1| \geq \dots \geq |\mathcal{F}_{18}| \geq 1$. Then $2 \leq |\mathcal{F}_1| \leq 7$ and $\sum_{i=1}^{18} (|\mathcal{F}_i| - 1) = 6$.

Define a graph $Q = (V(Q), E(Q))$ (see Fig. 6) as: $V(Q) = \{F_i^1, F_i^2, F_i^3 \mid 0 \leq i \leq 7\}$, and for $x, y \in V(Q)$, $xy \in E(Q)$ if and only if $|x \cap y| = 3$. The subgraph of Q induced by the elements of \mathcal{F}_i is denoted by Q_i . Note that if \mathcal{F}_i is a barrier, then Q_i is a 2-factor of order $|\mathcal{F}_i|$.




Fig. 6 Q

If $|\mathcal{F}_2| \leq 2$, then by (1), $|\bigcup_{i=1}^{18} L_i| \leq \sum_{i=1}^{18} |L_i| \leq 3|\mathcal{F}_1| + \sum_{i=2}^{18} 3(|\mathcal{F}_i| - 1) \leq 3 \sum_{i=1}^{18} (|\mathcal{F}_i| - 1) + 3 = 21 < 24$, a contradiction. So $|\mathcal{F}_2| \geq 3$, and then $3 \leq |\mathcal{F}_1| \leq 5$.

If $|\mathcal{F}_3| \leq 2$, then $6 \leq |\mathcal{F}_1| + |\mathcal{F}_2| \leq 8$, and by (1),

$$24 = \left| \bigcup_{i=1}^{18} L_i \right| \leq \sum_{i=1}^{18} |L_i| \leq 3|\mathcal{F}_1| + 3|\mathcal{F}_2| + \sum_{i=3}^{18} 3(|\mathcal{F}_i| - 1) = 3 \sum_{i=1}^{18} (|\mathcal{F}_i| - 1) + 6 = 24, \quad (2)$$

which implies $|\mathcal{F}_1| = 5$, $|\mathcal{F}_2| = 3$ and $|\mathcal{F}_3| = 1$, or $|\mathcal{F}_1| = |\mathcal{F}_2| = 4$ and $|\mathcal{F}_3| = 1$, or $|\mathcal{F}_1| = 4$, $|\mathcal{F}_2| = 3$, $|\mathcal{F}_3| = 2$ and $|\mathcal{F}_4| = 1$, or $|\mathcal{F}_1| = |\mathcal{F}_2| = 3$, $|\mathcal{F}_3| = |\mathcal{F}_4| = 2$ and $|\mathcal{F}_5| = 1$. Moreover, $\{L_1, L_2, \dots, L_{18}\}$ is a partition of $V(H_{P_{8,3}, \Psi})$, $\mathcal{F}_1, \mathcal{F}_2$ are barriers. Note that Q contains no K_4 and K_5 , and hence $|\mathcal{F}_1| = |\mathcal{F}_2| = 3$, $|\mathcal{F}_3| = |\mathcal{F}_4| = 2$ and $|\mathcal{F}_5| = 1$. By symmetry, we can assume $\mathcal{F}_1 = \{F_0^1, F_7^1, F_0^2\}$. Then $L_1 = \{x_0, x_5, x_6, x_7, y_0, y_1, y_7, z_0, z_7\}$, and then $\mathcal{F}_2 = \{F_3^1, F_4^1, F_4^2\}$ as $L_1 \cap L_2 = \emptyset$. So $L_2 = \{x_1, x_2, x_3, x_4, y_3, y_4, y_5, z_3, z_4\}$ and $L_3 \cup L_4 = \{y_2, y_6, z_1, z_2, z_5, z_6\}$. On the other hand, by (2), $|L_3| = |L_4| = 3$. Since $x_i \notin L_3 \cup L_4$ for $0 \leq i \leq 7$, $F_i^1 F_{i+1}^1$, $F_i^1 F_i^2$, $F_i^1 F_{i+1}^2$ and $F_i^2 F_i^3$ would be excluded from $E(Q)$. Similarly, each edge on two paths $F_0^3 F_3^3 F_6^3 F_1^3$, $F_4^3 F_7^3 F_2^3 F_5^3$ would be excluded from $E(Q)$ as $z_0, z_3, z_4, z_7 \notin L_3 \cup L_4$. So $\mathcal{F}_3, \mathcal{F}_4 \in \{\{F_0^3, F_5^3\}, \{F_1^3, F_4^3\}\}$, and thus $y_2, y_6 \notin L_3 \cup L_4$, a contradiction.

Thus $|\mathcal{F}_3| \geq 3$, and so $|\mathcal{F}_1| = |\mathcal{F}_2| = |\mathcal{F}_3| = 3$ and $|\mathcal{F}_4| = 1$. Then by (1), $|L_i| \leq 9$ for $1 \leq i \leq 3$. If Q_i is not a K_3 for some $1 \leq i \leq 3$, then we can check that $|L_i| \leq 6$. So we can assume $|L_1| = |L_2| = 9$, and then Q_1, Q_2 are K_3 . If Q_3 is not a K_3 , then $|L_3| = 6$ and F_i are barriers as $|L_1 \cup L_2 \cup L_3| = 24$ for $i = 1, 2, 3$, and thus by an argument similar to the above, we can have a contradiction. Therefore $Q_3 = K_3$ and $|L_3| = 9$. If $L_i \cap L_j = \emptyset$ for some $1 \leq i < j \leq 3$, then by an argument similar to the above, we can have a contradiction. So $L_i \cap L_j \neq \emptyset$ for any $1 \leq i < j \leq 3$. Assume that $\mathcal{F}_1 = \{F_0^1, F_7^1, F_0^2\}$. Then $L_1 = \{x_0, x_5, x_6, x_7, y_0, y_1, y_7, z_0, z_7\}$, and then $\mathcal{F}_i \neq \{F_3^1, F_4^1, F_4^2\}$ as $L_1 \cap L_i = \emptyset$ for $i = 2, 3$. Furthermore, we can check that $|L_i \cap L_1| = 3$ for $i = 2, 3$. Note that Q is 4-regular, and hence there exists some $x \in (L_1 \cap L_2) \setminus (L_1 \cap L_3)$. Thus $|\bigcup_{i=1}^{18} L_i| \leq 9 + 9 + 9 - 3 - 1 = 23 < 24$, a contradiction. Therefore, $H_{P_{8,3}, \Psi} \in \mathcal{P}_{6,24}$. By Lemma 2.4, $Ar(P_{8,3}, C_6) = Ar_{P_{8,3}}(\Psi) \leq |E(P_{8,3})| - (6 + 1) = 17$. ■

Lemma 4.10 $Ar(P_{10,3}, C_6) = 22$.

Proof. Note that $\Psi = \{G_i^1 := u_i u_{i+1} u_{i+2} u_{i+3} v_{i+3} v_i u_i, G_i^2 := u_i u_{i+1} v_{i+1} v_{i+4} v_{i+7} v_i u_i \mid 0 \leq i \leq 9\}$ is the set of all copies of C_6 in $P_{10,3}$. Then $Ar(P_{10,3}, C_6) = Ar_{P_{10,3}}(\Psi)$. Let $\phi : E(P_{10,3}) \rightarrow \{1, 2, \dots, 22\}$ be an edge-coloring of $P_{10,3}$ defined as follows:

$$\phi(e) = \begin{cases} i + 1, & \text{if } e \in E(v_i) \text{ with } 0 \leq i \leq 2; \\ 4, & \text{if } e \in E(u_5) \setminus \{u_5 v_5\}; \\ 5, & \text{if } e \in E(u_7) \setminus \{u_7 v_7\}; \end{cases}$$




Fig. 7 $P_{10,3}$ has no rainbow C_6 under the coloring ϕ .

and every other edges obtains distinct colorings from $\{6, \dots, 22\}$ (see Fig. 7). Note that each $C_6 \in \Psi$ contains two edges of $E(v_i)$ ($0 \leq i \leq 2$) or two edges of $E(u_j) \setminus \{u_jv_j\}$ ($j \in \{5, 7\}$), and hence $P_{10,3}$ has no rainbow C_6 under the coloring ϕ . So, $Ar(P_{10,3}, C_6) \geq 22$.

Next, we will show that $Ar(P_{10,3}, C_6) \leq 22$. Let $V(H_{P_{10,3}, \Psi}) = \{x_i, y_i \mid 0 \leq i \leq 9\}$ and $E(H_{P_{10,3}, \Psi}) = \{F_i^1, F_i^2, F_i^3 \mid 0 \leq i \leq 9\}$, where x_i and y_i correspond to G_i^1 and G_i^2 in Ψ , respectively, each F_i^1 corresponds to u_iu_{i+1} , F_i^2 to u_iv_i , and F_i^3 to v_iv_{i+3} for $0 \leq i \leq 9$. Note that each edge u_iu_{i+1} is contained in $G_i^1, G_{i+8}^1, G_{i+9}^1$ and G_i^2 , u_iv_i is contained in G_i^1, G_{i+7}^1, G_i^2 and G_{i+9}^2 , v_iv_{i+3} is contained in $G_i^1, G_{i+3}^2, G_{i+6}^2$ and G_{i+9}^2 for $0 \leq i \leq 9$. It follows that $F_i^1 = \{x_i, x_{i+8}, x_{i+9}, y_i\}$, $F_i^2 = \{x_i, x_{i+7}, y_i, y_{i+9}\}$ and $F_i^3 = \{x_i, y_{i+3}, y_{i+6}, y_{i+9}\}$ for $0 \leq i \leq 9$, and thus $r(H_{P_{10,3}, \Psi}) = 4$, $s(H_{P_{10,3}, \Psi}) = 2$

Now we will show that $H_{P_{10,3}, \Psi} \in \mathcal{P}_{7,20}$. Suppose that $H_{P_{10,3}, \Psi} \notin \mathcal{P}_{7,20}$. Then there exists a 23-partition $\{\mathcal{F}_1, \mathcal{F}_2, \dots, \mathcal{F}_{23}\}$ of $E(H_{P_{10,3}, \Psi})$ with $|\bigcup_{i=1}^{23} L_i| = 20$. Assume that $|\mathcal{F}_1| \geq \dots \geq |\mathcal{F}_{23}| \geq 1$. Then $2 \leq |\mathcal{F}_1| \leq 8$ and $\sum_{i=1}^{23} (|\mathcal{F}_i| - 1) = 7$.

Define a graph $Q' = (V(Q'), E(Q'))$ as: $V(Q') = \{F_i^1, F_i^2, F_i^3 \mid 0 \leq i \leq 9\}$, and for any $x, y \in V(Q')$, $xy \in E(Q')$ if and only if $|x \cap y| = 2$. Then Q' is 4-regular and contains no K_4 (see Fig. 8). The subgraph of Q' induced by the elements of \mathcal{F}_i is denoted by Q'_i .

If $|\mathcal{F}_3| \leq 2$, then by (1), $\sum_{i=1}^{23} |L_i| \leq 2|\mathcal{F}_1| + 2|\mathcal{F}_2| + \sum_{i=3}^{23} 2(|\mathcal{F}_i| - 1) \leq 2 \sum_{i=1}^{23} (|\mathcal{F}_i| - 1) + 4 = 18 < 20$, a contradiction. So $|\mathcal{F}_3| \geq 3$. Then $3 \leq |\mathcal{F}_1| \leq 4$, $|\mathcal{F}_2| = |\mathcal{F}_3| = 3$ and $|\mathcal{F}_4| \leq 2$. By (1),

$$20 = \left| \bigcup_{i=1}^{23} L_i \right| \leq \sum_{i=1}^{23} |L_i| \leq 2|\mathcal{F}_1| + 2|\mathcal{F}_2| + 2|\mathcal{F}_3| + \sum_{i=4}^{23} 2(|\mathcal{F}_i| - 1) \leq 2 \sum_{i=1}^{23} (|\mathcal{F}_i| - 1) + 6 = 20,$$

which implies $\mathcal{F}_1, \mathcal{F}_2, \mathcal{F}_3$ are barriers and $\{L_1, L_2, \dots, L_{23}\}$ is a partition of $V(H_{P_{10,3}, \Psi})$.

Since Q' contains no K_4 , $|\mathcal{F}_1| \neq 4$. So $|\mathcal{F}_1| = |\mathcal{F}_2| = |\mathcal{F}_3| = 3$ and $|\mathcal{F}_4| = 2$. Then each Q'_i is a K_3 for $1 \leq i \leq 3$. Without loss of generality, assume $\mathcal{F}_1 =$

$\{F_0^1, F_9^1, F_0^2\}$. Then $L_1 = \{x_0, x_7, x_8, x_9, y_0, y_9\}$. Since $L_i \cap L_j = \emptyset$ for $1 \leq i < j \leq 3$, we have $\{F_3^1, F_4^1, F_5^1, F_6^1, F_4^2, F_5^2, F_6^2, F_2^3, F_5^3\} \subseteq \mathcal{F}_2 \cup \mathcal{F}_3$. But in any case, we can check $L_2 \cap L_3 \neq \emptyset$, a contradiction. Therefore, $H_{P_{10,3}, \Psi} \in \mathcal{P}_{7,20}$. By Lemma 2.4, $Ar(P_{10,3}, C_6) = Ar_{P_{10,3}}(\Psi) \leq |E(P_{10,3})| - (7 + 1) = 22$. ■




Fig. 8 Q'

Lemma 4.11 $Ar(P_{18,3}, C_6) = 42$.

Proof. Note that $\Psi = \{G_i^1, G_j^2 \mid 0 \leq i \leq 17, 0 \leq j \leq 2\}$ is the set of all copies of C_6 in $P_{18,3}$, where $G_i^1 := u_i u_{i+1} u_{i+2} u_{i+3} v_{i+3} v_i u_i$, $G_j^2 := v_j v_{j+3} v_{j+6} v_{j+9} v_{j+12} v_{j+15} v_j$. Then $Ar(P_{18,3}, C_6) = Ar_{P_{18,3}}(\Psi)$.

Let $\phi : E(P_{18,3}) \rightarrow \{1, 2, \dots, 42\}$ be an edge-coloring of $P_{18,3}$ defined as follows:

$$\phi(e) = \begin{cases} i + 1, & \text{if } e \in \{u_{2i} u_{2i+1}, u_{2i+1} u_{2i+2}\}, 0 \leq i \leq 8; \\ j + 10, & \text{if } e \in \{v_{j+3} v_j, v_j v_{j-3}\}, 0 \leq j \leq 2; \end{cases}$$

and every other edges obtains distinct colorings from $\{13, \dots, 42\}$ (see Fig. 9). Note that each $C_6 \in \Psi$ contains two edges $u_{2i} u_{2i+1}, u_{2i+1} u_{2i+2}$ for some $0 \leq i \leq 8$, or two edges $v_{j+3} v_j, v_j v_{j-3}$ for some $0 \leq j \leq 2$, and hence $P_{18,3}$ has no rainbow C_6 under the coloring ϕ . So, $Ar(P_{18,3}, C_6) \geq 42$.

Next, we will show that $Ar(P_{18,3}, C_6) \leq 42$. Let $V(H_{P_{18,3}, \Psi}) = \{x_i, y_j \mid 0 \leq i \leq 17, 0 \leq j \leq 2\}$ and $E(H_{P_{18,3}, \Psi}) = \{F_i^1, F_i^2, F_i^3 \mid 0 \leq i \leq 17\}$, where x_i and y_j correspond to G_i^1 and G_j^2 in Ψ , respectively, each F_i^1 corresponds to $u_i u_{i+1}$, F_i^2 to $u_i v_i$, and F_i^3 to $v_i v_{i+3}$ for $0 \leq i \leq 17$. Note that each edge $u_i u_{i+1}$ is contained in G_i^1 , G_{i+16}^1 and G_{i+17}^1 , $u_i v_i$ is contained in G_i^1 and G_{i+15}^1 , $v_{3i+t} v_{3i+t+3}$ is contained in G_{3i+t}^1 and

G_t^2 for $0 \leq i \leq 17$ and $t = \lfloor \frac{i}{6} \rfloor$. Hence $F_i^1 = \{x_i, x_{i+16}, x_{i+17}\}$, $F_i^2 = \{x_i, x_{i+15}\}$ and $F_i^3 = \{x_{3i+t}, y_t\}$ for $0 \leq i \leq 17$ and $t = \lfloor \frac{i}{6} \rfloor$, and thus $r(H_{P_{18,3},\Psi}) = 3$, $s(H_{P_{18,3},\Psi}) = 2$.

Now we will show that $H_{P_{18,3},\Psi} \in \mathcal{P}_{11,21}$. Suppose that $H_{P_{18,3},\Psi} \notin \mathcal{P}_{11,21}$. Then there exists a 43-partition $\{\mathcal{F}_1, \mathcal{F}_2, \dots, \mathcal{F}_{43}\}$ of $E(H_{P_{18,3},\Psi})$ with $|\bigcup_{i=1}^{43} L_i| = 21$, and then $\bigcup_{i=1}^{43} L_i = V(H_{P_{18,3},\Psi})$. Note that $\sum_{i=1}^{43} (|\mathcal{F}_i| - 1) = 11$. By (1), $|L_i| \leq 2(|\mathcal{F}_i| - 1)$ for $1 \leq i \leq 43$. Assume $y_j \in L_{i_j}$ for $j = 0, 1, 2$. Note that $s(H_{P_{18,3},\Psi}) = 2$ and $|F_i^3 \cap F_j^3| \leq 1$ for $0 \leq i < j \leq 17$, and hence, by (1)

$$|L_{i_j}| \leq \begin{cases} \lfloor \frac{3|\mathcal{F}_{i_j}|}{2} \rfloor - 1 \leq 2(|\mathcal{F}_{i_j}| - 1) - 1, & \text{if } |\mathcal{F}_{i_j}| \geq 3; \\ 1, & \text{if } |\mathcal{F}_{i_j}| = 2. \end{cases}$$

Moreover, if $i_t = i_j$ for some $0 \leq t < j \leq 2$, then $|\mathcal{F}_{i_j}| \geq 4$ and $|L_{i_j}| \leq 2(|\mathcal{F}_{i_j}| - 1) - 2$. So $\sum_{i=1}^{43} |L_i| \leq 2 \sum_{i=1}^{43} (|\mathcal{F}_i| - 1) - 2 = 20 < 21$, a contradiction. Therefore, $H_{P_{18,3},\Psi} \in \mathcal{P}_{11,21}$. By Lemma 2.4, $Ar(P_{18,3}, C_6) = Ar_{P_{18,3}}(\Psi) \leq |E(P_{18,3})| - (11 + 1) = 42$. ■




Fig. 9 $P_{18,3}$ has no rainbow C_6 under the coloring ϕ .

Lemma 4.12 $Ar(P_{n,3}, C_6) = \lfloor \frac{5n}{2} \rfloor$ for $n \geq 7$ and $n \notin \{8, 10, 18\}$.

Proof. Let $G_i := u_i u_{i+1} u_{i+2} u_{i+3} v_{i+3} v_i u_i$ for $0 \leq i \leq n - 1$. Then $\Psi = \{G_i \mid 0 \leq i \leq n - 1\}$ is the set of all copies of C_6 in $P_{n,3}$, and then $Ar(P_{n,3}, C_6) = Ar_{P_{n,3}}(\Psi)$.

Let $\phi : E(P_{n,3}) \rightarrow \{1, 2, \dots, \lfloor \frac{5n}{2} \rfloor\}$ be an edge-coloring of $P_{n,3}$ defined as follows:

$$\phi(e) = \begin{cases} i + 1, & \text{if } e \in \{u_{2i} u_{2i+1}, u_{2i+1} u_{2i+2}\}, 0 \leq i \leq \lfloor \frac{n}{2} \rfloor - 1; \\ \lfloor \frac{n}{2} \rfloor, & \text{if } e = u_0 u_{n-1} \text{ and } n \equiv 1 \pmod{2}, \end{cases}$$

and every other edges obtains distinct colorings from $\{\lfloor \frac{n}{2} \rfloor + 1, \dots, \lfloor \frac{5n}{2} \rfloor\}$. Note that if $n \equiv 0 \pmod{2}$, then each $C_6 \in \Psi$ contains two edges $u_{2i}u_{2i+1}, u_{2i+1}u_{2i+2}$ for some $0 \leq i \leq \lfloor \frac{n}{2} \rfloor - 1$; and if $n \equiv 1 \pmod{2}$, then each $C_6 \in \Psi$ contains two edges $u_{2i}u_{2i+1}, u_{2i+1}u_{2i+2}$ for some $0 \leq i \leq \lfloor \frac{n}{2} \rfloor - 2$ or two of $\{u_{n-3}u_{n-2}, u_{n-2}u_{n-1}, u_0u_{n-1}\}$, and hence $P_{n,3}$ has no rainbow C_6 under the coloring ϕ . So, $Ar(P_{n,3}, C_6) \geq \lfloor \frac{5n}{2} \rfloor$.

Next, we will show that $Ar(P_{n,3}, C_6) \leq \lfloor \frac{5n}{2} \rfloor$. Note that each edge u_iu_{i+1} is contained in G_i, G_{i+n-2} and G_{i+n-1} , u_iv_i is contained in G_i and G_{i+n-3} , v_iv_{i+3} is contained in G_i for $0 \leq i \leq n-1$. Then $V(H_{P_{n,3}, \Psi}) = \{x_i \mid 0 \leq i \leq n-1\}$ and $E(H_{P_{n,3}, \Psi}) = \{F_0, \dots, F_{3n-1}\}$ with $F_i = \{x_i, x_{i+n-2}, x_{i+n-1}\}$, $F_{i+n} = \{x_i, x_{i+n-3}\}$ and $F_{i+2n} = \{x_i\}$ for $0 \leq i \leq n-1$. Thus $r(H_{P_{n,3}, \Psi}) = 3$ and $s(H_{P_{n,3}, \Psi}) = 1$. By Lemma 2.2, we have $Ar(P_{n,3}, C_6) = Ar_{P_{n,3}}(\Psi) \leq |E(P_{n,3})| - \lceil \frac{2|\Psi|}{3+1} \rceil = 3n - \lceil \frac{n}{2} \rceil = \lfloor \frac{5n}{2} \rfloor$. ■

By Lemmas 4.9-4.12, the proof of Theorem 1.5 is completed.

4.4 $k \geq 4$

In this subsection, $k \in \{\frac{n}{6}, \frac{n-1}{3}, \frac{n+1}{3}\}$ by Proposition 1.1(ii).

Lemma 4.13 $Ar(P_{n,k}, C_6) = \frac{17n}{6}$ for $k = \frac{n}{6} \geq 4$.

Proof. Denote $P'_{n,k} = P_{n,k} - E'$, where $E' = \{u_iv_{i+1}, u_iv_i \mid 0 \leq i \leq \frac{n}{6} - 1\}$. Let $G_i := v_iv_{i+k}v_{i+2k}v_{i+3k}v_{i+4k}v_{i+5k}v_i$ for $0 \leq i \leq \frac{n}{6} - 1$. Then $\Psi' = \{G_i \mid 0 \leq i \leq \frac{n}{6} - 1\}$ is the set of all copies of C_6 in $P'_{n,k}$, and $Ar(P'_{n,k}, C_6) = Ar_{P'_{n,k}}(\Psi')$. Note that each edge in $P'_{n,k}$ is only contained in some G_i . Thus $H_{P'_{n,k}, \Psi'}$ is a graph obtained from $\frac{n}{6}$ isolated vertices by adding six loops on each isolated vertex. Then $M(H_{P'_{n,k}, \Psi'}) = 0$. By Lemma 2.1, we have $Ar(P'_{n,k}, C_6) = Ar_{P'_{n,k}}(\Psi') = |E(P'_{n,k})| - |\Psi'| + M(H_{P'_{n,k}, \Psi'}) = \frac{5n}{6}$ for $k = \frac{n}{6} \geq 4$. Since there is no copy of C_6 in $P_{n,k}$ containing any edge of F , $Ar(P_{n,k}, C_6) = |E'| + Ar(P'_{n,k}, C_6) = \frac{17n}{6}$. ■

Lemma 4.14 $Ar(P_{n,k}, C_6) = \lfloor \frac{5n}{2} \rfloor$ for $k \in \{\frac{n-1}{3}, \frac{n+1}{3}\}$.

Proof. Let

$$G_i := \begin{cases} u_iv_{i+1}v_{i+1}v_{i+1+k}v_{i+1+2k}v_iu_i, & \text{if } k = \frac{n-1}{3}; \\ u_iv_iv_{i+k}v_{i+2k}v_{i+1}u_{i+1}u_i, & \text{if } k = \frac{n+1}{3}, \end{cases}$$

Then $\Psi = \{G_i \mid 0 \leq i \leq n-1\}$ is the set of all copies of C_6 in $P_{n,k}$ for $k \in \{\frac{n-1}{3}, \frac{n+1}{3}\}$. Then $Ar(P_{n,k}, C_6) = Ar_{P_{n,k}}(\Psi)$. Recall that $|E(P_{n,k})| = 3n$.

If $n \equiv 1 \pmod{2}$, then let $\phi : E(P_{n,k}) \rightarrow \{1, 2, \dots, \lfloor \frac{5n}{2} \rfloor\}$ be an edge-coloring of $P_{n,k}$ such that $\phi(v_0v_k) = \phi(v_kv_{2k}) = \phi(v_{2k}v_{3k}) = 1$, $\phi(v_{ik}v_{(i+1)k}) = \lceil \frac{i}{2} \rceil$ for all




Fig. 10 Illustrations of $P_{13,4}$ and $P_{16,5}$ having no rainbow C_6 under ϕ .

$3 \leq i \leq n-1$ and every other edges obtains distinct colorings from $\{\frac{n+1}{2}, \dots, \lfloor \frac{5n}{2} \rfloor\}$ (Fig. 10(a) illustrates the coloring ϕ of $P_{13,4}$). Then each $C_6 \in \Psi$ contains two edges of $\{v_0v_k, v_kv_{2k}, v_{2k}v_{3k}\}$ or two edges $v_{(2i+1)k}v_{(2i+2)k}, v_{(2i+2)k}v_{(2i+3)k}$ for some $1 \leq i \leq \frac{n-1}{2}-1$, and hence $P_{n,k}$ has no rainbow C_6 under the coloring ϕ .

If $n \equiv 0 \pmod{2}$, then let $\phi : E(P_{n,k}) \rightarrow \{1, 2, \dots, \lfloor \frac{5n}{2} \rfloor\}$ be an edge-coloring of $P_{n,k}$ such that $\phi(v_iv_{(i+1)k}) = \lfloor \frac{i}{2} \rfloor + 1$ for all $0 \leq i \leq n-1$ and every other edges obtains distinct colorings from $\{\frac{n}{2} + 1, \dots, \lfloor \frac{5n}{2} \rfloor\}$ (Fig. 10(b) illustrates the coloring ϕ of $P_{16,5}$). Note that each $C_6 \in \Psi$ contains two edges $v_{2ik}v_{(2i+1)k}, v_{(2i+1)k}v_{(2i+2)k}$ for some $0 \leq i \leq \frac{n}{2}-1$, and hence $P_{n,k}$ has no rainbow C_6 under the coloring ϕ . So, $Ar(P_{n,k}, C_6) \geq \lfloor \frac{5n}{2} \rfloor$ for $n \equiv 0 \pmod{2}$. So, in either case, $Ar(P_{n,k}, C_6) \geq \lfloor \frac{5n}{2} \rfloor$.

Now, we show that $Ar(P_{n,k}, C_6) \leq \lfloor \frac{5n}{2} \rfloor$. Let $V(H_{P_{n,k}, \Psi}) = \{x_i \mid 0 \leq i \leq n-1\}$ and $E(H_{P_{n,k}, \Psi}) = \{F_0, \dots, F_{3n-1}\}$, where each x_i corresponds to G_i in Ψ , each F_i corresponds to u_iu_{i+1} , F_{i+n} corresponds to u_iv_i , F_{i+2n} corresponds to v_iv_{i+k} for $k \in \{\frac{n-1}{3}, \frac{n+1}{3}\}$. Note that for $0 \leq i \leq n-1$, each u_iu_{i+1} is contained in G_i , u_iv_i is contained in G_i, G_{i+n-1} , $v_iv_{i+\frac{n-1}{3}}$ is contained in $G_{i+\frac{n-1}{3}}, G_{i+\frac{2n-2}{3}}, G_{i+n-1}$ when $k = \frac{n-1}{3}$, and $v_iv_{i+\frac{n+1}{3}}$ is contained in $G_i, G_{i+\frac{n-2}{3}}, G_{i+\frac{2n-1}{3}}$ when $k = \frac{n+1}{3}$. It follows that $F_i = \{x_i\}$, $F_{i+n} = \{x_i, x_{i+n-1}\}$ and

$$F_{i+2n} = \begin{cases} \{x_{i+\frac{n-1}{3}}, x_{i+\frac{2n-2}{3}}, x_{i+n-1}\}, & \text{if } k = \frac{n-1}{3}; \\ \{x_i, x_{i+\frac{n-2}{3}}, x_{i+\frac{2n-1}{3}}\}, & \text{if } k = \frac{n+1}{3}. \end{cases}$$

Obviously, $r(H_{P_{n,k}, \Psi}) = 3$ and $s(H_{P_{n,k}, \Psi}) = 1$, then by Lemma 2.2, $Ar(P_{n,k}, C_6) = Ar_{P_{n,k}}(\Psi) \leq |E(P_{n,k})| - \lceil \frac{2|\Psi|}{3+1} \rceil = 3n - \lceil \frac{n}{2} \rceil = \lfloor \frac{5n}{2} \rfloor$. ■

Lemma 4.15 $Ar(P_{n,k}, C_6) = \lfloor \frac{7n}{3} \rfloor$ for $k = \frac{n-2}{2} \geq 4$.

Proof. Let $G_i := u_i u_{i+1} v_{i+1} v_{i+\frac{n}{2}} v_{i+n-1} u_{i+n-1} u_i$ for $0 \leq i \leq n-1$. Then $\Psi = \{G_i \mid 0 \leq i \leq n-1\}$ is the set of all copies of C_6 in $P_{n,k}$, and $Ar(P_{n,k}, C_6) = Ar_{P_{n,k}}(\Psi)$. Note that each edge $u_i u_{i+1}$ is contained in G_i and G_{i+1} ; each spoke $u_i v_i$ is contained in G_{i+1} and G_{i+n-1} ; each edge $v_i v_{i+\frac{n-2}{2}}$ is contained in $G_{i+\frac{n}{2}}$ and G_{i+n-1} for $0 \leq i \leq n-1$. Thus $H_{P_{n,k}, \Psi}$ is a graph obtained from a cycle $w_0 w_1 \cdots w_{n-1} w_0$ by adding edges $w_{i+1} w_{i+n-1}$ and $w_{i+\frac{n}{2}} w_{i+n-1}$ for $0 \leq i \leq n-1$. Since there exist $\lfloor \frac{n}{3} \rfloor$ vertex-disjoint cycles $w_{3i} w_{3i+1} w_{3i+2} w_{3i}$ in $H_{P_{n,k}, \Psi}$ for $0 \leq i \leq \lfloor \frac{n}{3} \rfloor - 1$, we have $M(H_{P_{n,k}, \Psi}) = \lfloor \frac{n}{3} \rfloor$. By Lemma 2.1, $Ar(P_{n,k}, C_6) = Ar_{P_{n,k}}(\Psi) = |E(P_{n,k})| - |\Psi| + M(H_{P_{n,k}, \Psi}) = \lfloor \frac{7n}{3} \rfloor$. ■

By Lemmas 4.13-4.15, we complete the proof of Theorem 1.6.

References

- [1] N. Alon, On a conjecture of Erdős, Simonovits, and Sós concerning anti-Ramsey theorems, *J. Graph Theory*, 7(1)(1983) 91-94.
- [2] M. Axenovich, T. Jiang and A. Kündgen, Bipartite anti-Ramsey numbers of cycles, *J. Graph Theory*, 47(1)(2004) 9-28.
- [3] M. Axenovich, H. Harborth, A. Kemnitz, M. Möller and I. Schiermeyer, Rainbows in the hypercube, *Graphs Combin.*, 23(2007) 123-133.
- [4] P. Erdős, M. Simonovits and V. Sós, Anti-Ramsey theorems. In: Infinite and finite sets (Colloq. Keszthely 1973), *Colloq. Math. Soc. János Bolyai*, 10(1975) 633-643.
- [5] I. Gorgol, Anti-Ramsey numbers in complete split graphs, *Discrete Math.*, 339(2016) 1944-1949.
- [6] R. Gu, J. Li, and Y. Shi, Anti-ramsey numbers of paths and cycles in hypergraphs, *SIAM J. Discrete Math.*, 34(2020) 271-307.
- [7] R. Haas and M. Young, The anti-Ramsey number of perfect matching, *Discrete Math.*, 312(2012) 933-937.
- [8] M. Hornák, S. Jendrol, I. Schiermeyer and R. Soták, Rainbow numbers for cycles in plane triangulations, *J. Graph Theory*, 78(2015) 248-257.
- [9] S. Jahanbekam and D. B. West, Anti-Ramsey problems for t edge-disjoint rainbow spanning subgraphs: cycles, matchings, or trees, *J. Graph Theory*, 82(1)(2016) 75-89.

- [10] S. Jendrol, I. Schiermeyer and J. Tu, Rainbow numbers for matchings in plane triangulations, *Discrete Math.*, 331(2014) 158-164.
- [11] Y. Jia, M. Lu and Y. Zhang, Anti-Ramsey problems in complete bipartite graphs for t edge-disjoint rainbow spanning subgraphs: Cycles and Matchings, *Graphs Combin.*, 35(5)(2019) 1011-1021.
- [12] T. Jiang and D. B. West, On the Erdős-Simonovits-Sós conjecture about the anti-Ramsey number of a cycle, *Combin. Probab. Comput.*, 12(5-6)(2003) 585-598.
- [13] Z. Jin and X. Li, Anti-Ramsey numbers for graphs with independent cycles, *Electron. J. Combin.*, 16(2009) 85.
- [14] Z. Jin, K. Ye, Y. Sun and H. Chen, Rainbow matchings in edge-colored complete split graphs, *European J. Combin.*, 70(2018) 297-316.
- [15] Y. Lan, Y. Shi and Z. Song, Planar anti-ramsey numbers for paths and cycles, *Discrete Math.*, 342(11)(2019) 3216-3224.
- [16] X. Li, J. Tu and Z. Jin, Bipartite rainbow numbers of matchings, *Discrete Math.*, 309(2009) 2575-2578.
- [17] J.J. Montellano-Ballesteros, V. Neumann-Lara, An anti-Ramsey theorem, *Combinatorica*, 22(3)(2002) 445-449.
- [18] J.J. Montellano-Ballesteros, V. Neumann-Lara, An anti-Ramsey theorem on cycles, *Graphs Combin.*, 21(3)(2005) 343-354.
- [19] J. Xu, M. Lu and K. Liu, Anti-Ramsey problems for cycles, *Applied Mathematics and Computation*, 408 (2021) 126345.