

# PFH spectral invariants and $C^\infty$ closing lemmas

Oliver Edtmair and Michael Hutchings\*

## Abstract

We develop the theory of spectral invariants in periodic Floer homology (PFH) of area-preserving surface diffeomorphisms. We use this theory to prove  $C^\infty$  closing lemmas for certain Hamiltonian isotopy classes of area-preserving surface diffeomorphisms. In particular, we show that any area-preserving diffeomorphism of the torus is  $C^\infty$  close to an area-preserving diffeomorphism with dense periodic orbits. Our closing lemmas are quantitative, asserting roughly speaking that for a given Hamiltonian isotopy, within time  $\delta$  a periodic orbit must appear of period  $O(\delta^{-1})$ . We also prove a “Weyl law” describing the asymptotic behavior of PFH spectral invariants.

## 1 Introduction

Throughout this paper, let  $\Sigma$  be a closed connected surface of genus  $g$ , and let  $\omega$  be a symplectic (area) form on  $\Sigma$ . We are interested in area-preserving diffeomorphisms  $\phi : (\Sigma, \omega) \rightarrow (\Sigma, \omega)$ . We are also interested in Hamiltonian isotopy classes in the set of all area-preserving diffeomorphisms of  $(\Sigma, \omega)$ ; we denote the Hamiltonian isotopy class of  $\phi$  by  $[\phi]$ .

### 1.1 Closing lemmas

Our convention is that a *periodic orbit* of  $\phi$  with period  $k$  is a set of  $k$  distinct points in  $\Sigma$  that are cyclically permuted by  $\phi$ .

**Definition 1.1.** Let  $\Phi$  be a Hamiltonian isotopy class of area-preserving diffeomorphisms of  $(\Sigma, \omega)$ . We say that  $\Phi$  has the  $C^\infty$  *closing property* if for every map  $\phi \in \Phi$  and for every nonempty open set  $\mathcal{U} \subset \Sigma$ , there exists a  $C^\infty$  small Hamiltonian isotopy supported in  $\mathcal{U}$  from  $\phi$  to  $\phi'$  such that  $\phi'$  has a periodic orbit intersecting  $\mathcal{U}$ .

Standard arguments, see e.g. [29, §3], show:

**Lemma 1.2.** *If the Hamiltonian isotopy class  $\Phi$  has the  $C^\infty$  closing property, then for a  $C^\infty$ -generic area-preserving diffeomorphism  $\phi \in \Phi$ , the set of periodic orbits of  $\phi$  is dense in  $\Sigma$ .*

---

\*Partially supported by NSF grant DMS-2005437.

It was proved by Asaoka-Irie [1] that for a  $C^\infty$ -generic Hamiltonian diffeomorphism  $\phi$  of  $(\Sigma, \omega)$ , i.e. for a generic  $\phi$  in the Hamiltonian isotopy class  $[\text{id}_\Sigma]$ , the set of periodic orbits of  $\phi$  is dense in  $\Sigma$ . One of our goals is to generalize this result to some other Hamiltonian isotopy classes.

**Theorem 1.3.** *(proved in §7) Let  $\Phi$  be a Hamiltonian isotopy class of area-preserving diffeomorphisms of  $(\Sigma, \omega)$ . Suppose that  $\Phi$  is rational (Definition 1.4) and has the  $U$ -cycle property (Definition 2.20). Then  $\Phi$  satisfies the  $C^\infty$  closing property.*

To explain the rationality hypothesis, we need to introduce a key actor in the story, the *mapping torus* of  $\phi$ . This is a three-manifold defined by

$$Y_\phi = [0, 1] \times \Sigma / \sim, \quad (1, x) \sim (0, \phi(x)). \quad (1.1)$$

The mapping torus is a fiber bundle over  $S^1 = \mathbb{R}/\mathbb{Z}$  with fiber  $\Sigma$ . If  $t$  denotes the  $[0, 1]$  coordinate on  $[0, 1] \times \Sigma$ , then the vector field  $\partial_t$  on  $[0, 1] \times \Sigma$  descends to a vector field on  $Y_\phi$ , which we also denote by  $\partial_t$ . Periodic orbits of the map  $\phi$  of period  $d$  correspond to simple periodic orbits of the vector field  $\partial_t$  whose projection to  $S^1$  has degree  $d$ . Since the map  $\phi$  preserves the symplectic form  $\omega$  on  $\Sigma$ , this form induces a fiberwise symplectic form  $\omega_\phi$  on  $Y_\phi$ . The latter extends to a closed 2-form  $\omega_\phi$  on  $Y_\phi$ , characterized by  $\omega_\phi(\partial_t, \cdot) = 0$ .

We need to consider how the cohomology class  $[\omega_\phi] \in H^2(Y_\phi; \mathbb{R})$  depends on  $\phi$ . Let  $\{\phi_s\}_{s \in [0, 1]}$  be a smooth isotopy of area-preserving diffeomorphisms of  $(\Sigma, \omega)$ , and suppose for the moment that  $\phi_s$  is constant for  $s$  close to 0 or 1. We then obtain a diffeomorphism of mapping tori

$$f : Y_{\phi_0} \xrightarrow{\cong} Y_{\phi_1}.$$

This is induced by the diffeomorphism of  $[0, 1] \times \Sigma$  sending

$$(t, x) \mapsto (t, \phi_t^{-1}(\phi_0(x))). \quad (1.2)$$

If the isotopy  $\{\phi_s\}$  is Hamiltonian, then  $f^*[\omega_{\phi_1}] = [\omega_{\phi_0}] \in H^2(Y_{\phi_0}; \mathbb{R})$ .

**Definition 1.4.** The Hamiltonian isotopy class  $\Phi$  is *rational* if for  $\phi \in \Phi$ , the cohomology class  $[\omega_\phi] \in H^2(Y_\phi; \mathbb{R})$  is a real multiple of a class in the image of  $H^2(Y_\phi; \mathbb{Z})$ .

**Example 1.5.** Suppose that  $\Sigma = T^2 = \mathbb{R}^2/\mathbb{Z}^2$ . Then any area-preserving diffeomorphism  $\phi$  is Hamiltonian isotopic to a map of the form  $\phi(x) = Ax + b$  where  $A \in \text{SL}_2 \mathbb{Z}$  and  $b \in \mathbb{R}^2/\mathbb{Z}^2$ . We have  $H_2(Y_\phi) \simeq \mathbb{Z} \oplus \text{Ker}(A - I)$ . If  $A = I$ , that is if  $\phi$  is smoothly isotopic to the identity, then  $[\phi]$  is rational if and only if  $b \in \mathbb{Q}^2/\mathbb{Z}^2$ , namely  $\phi$  is a rational rotation. If  $\text{Ker}(A - I)$  is one-dimensional, then  $\phi$  is isotopic to a power of a Dehn twist, and  $[\phi]$  is rational when the projection of  $b$  to  $\mathbb{R}^2/\text{Ker}(A - I)$  is rational. In all other cases,  $b_2(Y_\phi) = 1$  so  $[\phi]$  is automatically rational.

We will use Theorem 1.3 and some additional arguments to prove:

**Theorem 1.6.** *(proved in §7) Let  $\Phi$  be a Hamiltonian isotopy class of area-preserving diffeomorphisms of  $T^2$ . If  $\Phi$  is rational, then  $\Phi$  has the  $C^\infty$  closing property.*

**Remark 1.7.** The  $C^\infty$  closing property fails to hold for some non-rational Hamiltonian isotopy classes of area-preserving diffeomorphisms of  $T^2$ . In fact, it follows from a result of Herman [15, Annexe, Thm. 2.2] that if  $\phi$  is a Diophantine rotation of  $T^2$ , then there is a neighborhood of  $\phi$  in the  $C^\infty$  topology in the space of maps Hamiltonian isotopic to  $\phi$  such that any map  $\phi'$  in this neighborhood is smoothly conjugate to  $\phi$ , and hence has no periodic orbits. (Thanks to V. Humilière for this reference.)

Any non-rational area-preserving diffeomorphism of  $T^2$  can be perturbed to a rational one by a  $C^\infty$ -small (non-Hamiltonian) isotopy. Thus we obtain:

**Corollary 1.8.** *Every area-preserving diffeomorphism of  $T^2$  is  $C^\infty$ -close to an area-preserving diffeomorphism whose periodic orbits are dense in  $T^2$ .*

## 1.2 Motivation from previous work

The proof of Theorem 1.3 is inspired by an argument of Irie [29], who proved the following  $C^\infty$  closing lemma for contact forms on closed three-manifolds. See also the survey [16].

**Theorem 1.9** (Irie). *Let  $Y$  be a closed three-manifold, let  $\lambda$  be a contact form on  $Y$ , and let  $\mathcal{U} \subset Y$  be a nonempty open set. Then there exists a contact form  $\lambda'$  which is  $C^\infty$ -close to  $\lambda$  and agrees with  $\lambda$  outside of  $\mathcal{U}$ , such that the Reeb vector field associated to  $\lambda'$  has a periodic orbit intersecting  $\mathcal{U}$ .*

The proof uses the *embedded contact homology* of  $(Y, \lambda)$ ; see [20] for detailed definitions. If  $\lambda$  is nondegenerate, meaning that the periodic orbits of the Reeb vector field are nondegenerate, then the embedded contact homology  $ECH(Y, \lambda)$  is the homology of a chain complex generated (over  $\mathbb{Z}/2$ ) by certain finite sets of Reeb orbits with multiplicities, and whose differential counts certain  $J$ -holomorphic curves in  $\mathbb{R} \times Y$  for a suitable almost complex structure  $J$ . Taubes [36] proved that  $ECH(Y, \lambda)$  is canonically isomorphic to a version of Seiberg-Witten Floer cohomology of  $Y$  as defined by Kronheimer-Mrowka [30], and in particular depends only on the contact structure  $\xi = \text{Ker}(\lambda)$ . For any contact form  $\lambda$  on  $Y$ , possibly degenerate, and for any nonzero class  $\sigma \in ECH(Y, \xi)$ , there is a “spectral invariant”  $c_\sigma(Y, \lambda) \in \mathbb{R}$ , which is the total period of a finite set of Reeb orbits with multiplicities homologically selected by ECH. The spectral invariants, unlike ECH, are highly sensitive to the contact form  $\lambda$ .

For the proof of Theorem 1.9, we can assume without loss of generality that  $Y$  is connected. There is then a well-defined map  $U : ECH(Y, \xi) \rightarrow ECH(Y, \xi)$ ,

which is induced by a chain map counting certain  $J$ -holomorphic curves that are constrained to pass through a base point in  $\mathbb{R} \times Y$ . Define a  $U$ -sequence to be a sequence of nonzero classes  $\{\sigma_k\}_{k \geq 1}$  in  $ECH(Y, \xi)$  such that  $U\sigma_{k+1} = \sigma_k$  for all  $k \geq 1$ . As explained for example in [8, Lem. A.1], results of Kronheimer-Mrowka [30] on Seiberg-Witten Floer homology imply that  $U$ -sequences always exist.

The key ingredient now is the following “Weyl law” for ECH spectral invariants proved<sup>1</sup> in [9].

**Theorem 1.10.** [9] *Let  $Y$  be a closed connected three-manifold, let  $\lambda$  be a contact form on  $Y$ , and let  $\{\sigma_k\}_{k \geq 1}$  be a  $U$ -sequence in  $ECH(Y, \xi)$ . Then*

$$\lim_{k \rightarrow \infty} \frac{c_{\sigma_k}(Y, \lambda)^2}{k} = 2 \operatorname{vol}(Y, \lambda). \quad (1.3)$$

Here the *contact volume* is defined by

$$\operatorname{vol}(Y, \lambda) = \int_Y \lambda \wedge d\lambda.$$

**Remark 1.11.** There is also a more general version of Theorem 1.10 in which the classes  $\sigma_k$  are nonzero and homogeneous and their ECH gradings converge to  $+\infty$ , but  $\{\sigma_k\}$  is not required to be a  $U$ -sequence.

To prove Theorem 1.9, one can define a smooth one-parameter family of contact forms  $\{\lambda_t\}$  such that  $\lambda_0 = \lambda$ , outside of  $\mathcal{U}$  we have  $\lambda_t = \lambda$ , and  $\frac{d}{dt} \operatorname{vol}(Y, \lambda_t) > 0$ . There must then exist Reeb orbits of  $\lambda_t$  passing through  $\mathcal{U}$  for arbitrarily small  $t$ . Otherwise, there exists  $\delta > 0$  such that  $\lambda_t$  has no Reeb orbit passing through  $\mathcal{U}$  for  $t \in [0, \delta]$ . One can deduce that each spectral invariant  $c_\sigma(Y, \lambda_t)$  is independent of  $t \in [0, \delta]$ . It then follows from the Weyl law (1.3) that  $\operatorname{vol}(Y, \lambda_t)$  is independent of  $t \in [0, \delta]$ , which is a contradiction.

Returning to area-preserving surface diffeomorphisms: Asaoka-Irie proved that a  $C^\infty$ -generic Hamiltonian diffeomorphism of  $(\Sigma, \omega)$  has dense periodic orbits by starting with a Hamiltonian diffeomorphism  $\phi$  and constructing a contact three-manifold with an open book decomposition whose page is  $\Sigma$  with a disk removed, and whose monodromy is a slight modification of  $\phi$ . One can then apply Theorem 1.9 to find a  $C^\infty$ -small perturbation of the contact form with dense Reeb orbits, and translate this back to a  $C^\infty$ -small perturbation of  $\phi$  with dense periodic orbits.

It is not obvious how to extend the above argument to other Hamiltonian isotopy classes, because there are cohomological obstructions to defining the desired contact form. We will instead work more directly with *periodic Floer homology* (PFH). This is a theory which is defined analogously to ECH, but using periodic orbits of an area-preserving surface diffeomorphism instead of Reeb orbits of a contact form on a three-manifold. PFH is isomorphic to a version of Seiberg-Witten Floer cohomology

---

<sup>1</sup>This was earlier proved in a special case in [19], and later given a different proof by Sun [35].

of the mapping torus  $Y_\phi$ , as shown by Lee-Taubes [31]. Originally, PFH was defined before ECH; see [17, 23]. Since then there have been many applications of ECH to dynamics of Reeb vector fields in three dimensions and symplectic embeddings in four dimensions. Applications of PFH have only recently begun to appear, including the spectacular proof by Cristofaro-Gardiner, Humilière, and Seyfaddini [5] that the group of compactly supported area-preserving homeomorphisms of the disk is not simple, and additional applications of PFH to area-preserving homeomorphisms of the two-sphere [6].

We will prove a PFH analogue of the “Weyl law” (1.3) in Theorem 8.1 below, replacing the notion of “ $U$ -sequence” by a notion of “ $U$ -cycle”. This Weyl law implies closing lemmas as in Theorem 1.3 (under slightly stronger hypotheses on the  $U$ -cycles), following Irie’s proof of Theorem 1.9.

However, a Weyl law is really much stronger than necessary to detect the creation of periodic orbits. In §6 we introduce a refinement of Irie’s argument which, instead of a Weyl law, uses bounds on “spectral gaps” coming from ball packings in symplectic cobordisms, to obtain stronger results, namely *quantitative* closing lemmas.

### 1.3 Quantitative closing lemmas

The kind of quantitative closing lemma that we can obtain asserts roughly that during a given Hamiltonian isotopy, within time  $\delta$  a periodic orbit must appear with period  $O(\delta^{-1})$ . We now give some examples of precise statements that we can prove.

**Definition 1.12.** Let  $\mathcal{U} \subset \Sigma$  be a nonempty open set, let  $l \in (0, 1)$ , and let  $a \in (0, \text{area}(\mathcal{U}))$ . A  $(\mathcal{U}, a, l)$ -admissible Hamiltonian is a smooth function  $H : [0, 1] \times \Sigma \rightarrow \mathbb{R}$  such that:

- $H(t, x) = 0$  for  $t$  close to 0 or 1.
- $H(t, x) = 0$  for  $x \notin \mathcal{U}$ .
- $H \geq 0$ .
- There is an interval  $I \subset (0, 1)$  of length  $l$  and a disk  $D \subset \mathcal{U}$  of area  $a$  such that  $H \geq 1$  on  $I \times D$ .

Given any Hamiltonian  $H : [0, 1] \times \Sigma \rightarrow \mathbb{R}$  satisfying the first bullet point above, let  $\{\varphi_t\}_{t \in [0, 1]}$  denote the associated Hamiltonian isotopy (see §3 for conventions), and given an area-preserving diffeomorphism  $\phi$  of  $\Sigma$ , write  $\phi_H = \phi \circ \varphi_1$ .

**Theorem 1.13.** (proved in §7) Let  $\phi$  be an area-preserving diffeomorphism of  $(S^2, \omega)$ , write  $A = \int_{S^2} \omega$ , let  $\mathcal{U} \subset S^2$  be a nonempty open set, and let  $H$  be a

$(\mathcal{U}, a, l)$ -admissible Hamiltonian. If  $0 < \delta \leq al^{-1}$ , then for some  $\tau \in [0, \delta]$ , the map  $\phi_{\tau H}$  has a periodic orbit intersecting  $\mathcal{U}$  with period  $d$  satisfying

$$d \leq \lfloor Al^{-1}\delta^{-1} \rfloor. \quad (1.4)$$

**Remark 1.14.** When  $\delta = al^{-1}$  and  $Aa^{-1} \notin \mathbb{Z}$ , the bound (1.4) is sharp. That is, under the hypotheses of Theorem 1.13, one cannot prove the existence of a period orbit intersecting  $\mathcal{U}$  with period less than  $\lfloor Aa^{-1} \rfloor$ . The reason is that if  $a = \text{area}(\mathcal{U}) - \varepsilon$  for  $\varepsilon > 0$  sufficiently small, then the open sets  $\phi^i(\mathcal{U})$  for  $0 \leq i < \lfloor Aa^{-1} \rfloor$  have total area less than  $A$  and thus could be disjoint.

We also obtain a slightly weaker inequality for rational area-preserving diffeomorphisms of the torus:

**Theorem 1.15.** (proved in §7) Let  $\phi$  be an area-preserving diffeomorphism of  $(T^2, \omega)$ , and write  $A = \int_{T^2} \omega$ . Suppose that the Hamiltonian isotopy class  $[\phi]$  is rational, let  $\Omega \in H^2(Y_\phi; \mathbb{Z})$  be an integral cohomology class such that  $[\omega_\phi]$  is a positive multiple of the image of  $\Omega$ , and let  $d_0 = \langle \Omega, [T^2] \rangle$ . Let  $\mathcal{U} \subset \Sigma$  be a nonempty open set, and let  $H$  be a  $(\mathcal{U}, a, l)$ -admissible Hamiltonian. If  $0 < \delta \leq al^{-1}$ , then for some  $\tau \in [0, \delta]$ , the map  $\phi_{\tau H}$  has a periodic orbit intersecting  $\mathcal{U}$  with period  $d$  satisfying

$$d \leq d_0 (\lfloor Ad_0^{-1}l^{-1}\delta^{-1} \rfloor + 1).$$

Theorems 1.13 and 1.15 are special cases of a more general statement, Theorem 7.4 below, which is also applicable to higher genus surfaces when there are enough  $U$ -cycles.

The rest of the paper is organized as follows. In §2 we review the definition of periodic Floer homology. In §3 we discuss invariance of PFH under Hamiltonian isotopy; here we find it useful to relate a Hamiltonian isotopy to the graph of a function  $Y_\phi \rightarrow \mathbb{R}$ . In §4 we explain how to define the PFH spectral numbers we will use. In §5 we prove a key lemma which gives relations between PFH spectral invariants of different maps in the same Hamiltonian isotopy class arising from ball packings in symplectic cobordisms between the graphs of different Hamiltonians. In §6 we use this lemma to show how “spectral gaps” in PFH allow one to detect the creation of periodic orbits. In §7 we use this machinery to prove all of our theorems stated above. Finally, in §8 we state and prove a Weyl for PFH spectral invariants.

**Remark 1.16.** Cristofaro-Gardiner, Prasad, and Zhang [11] have independently proved a related Weyl law in PFH by different methods. This Weyl law (together with a nonvanishing result for Seiberg-Witten Floer cohomology) is used in [11] to prove a  $C^\infty$  closing result for arbitrary diffeomorphisms on surfaces of arbitrary genus, where non-local and non-Hamiltonian perturbations are allowed. The Weyl law in [11], unlike the one here, does not need any hypothesis about “ $U$ -cycles”; cf. Remark 1.11.

We expect that one can use the Seiberg-Witten nonvanishing results in [11] to show that rational Hamiltonian isotopy classes satisfy the  $U$ -cycle property, similarly to Lemma 7.3 below. If so, then combining this with Theorem 1.3 would imply that Theorem 1.6 extends to surfaces of arbitrary genus.

The proof of the Weyl law in [11] uses the Lee-Taubes isomorphism and various careful estimates on solutions to the Seiberg-Witten equations. The proofs of our Weyl law and quantitative closing lemmas also use Seiberg-Witten theory and the Lee-Taubes isomorphism, but more indirectly, as a “black box” to provide cobordism maps on PFH whose nontriviality implies the existence of certain holomorphic curves.

## 2 Periodic Floer homology

We now set up the version of PFH that we will be using, which one might call “twisted PFH”, in more detail. Most of this material is also explained in [17, 23, 18, 31, 5], although we will use a particular bookkeeping formalism of Novikov rings and reference cycles to keep track of areas of holomorphic curves. Apart from this bookkeeping, PFH is extremely similar to ECH, and we will refer to the lecture notes [20] on ECH for some definitions and basic results that do not differ significantly from the PFH case.

### 2.1 PFH generators and holomorphic curves

Let  $(\Sigma, \omega)$  be a closed connected surface of genus  $g$  with a symplectic (area) form, and let  $\phi : (\Sigma, \omega) \rightarrow (\Sigma, \omega)$  be an area-preserving diffeomorphism.

**Definition 2.1.** An *orbit set* is a finite set of pairs  $\alpha = \{(\alpha_i, m_i)\}$  such that:

- The  $\alpha_i$  are distinct periodic orbits of  $\phi$ .
- The  $m_i$  are positive integers.

Let  $Y_\phi$  denote the mapping torus of  $\phi$  as in (1.1). For an orbit set as above, regarding the periodic orbits  $\alpha_i$  as embedded loops in  $Y_\phi$ , we define the homology class

$$[\alpha] = \sum_i m_i [\alpha_i] \in H_1(Y_\phi).$$

A periodic orbit of  $\phi$  of period  $k$  is *nondegenerate* if for  $x \in \Sigma$  in the periodic orbit, the derivative  $d\phi_x^k : T_x \Sigma \rightarrow T_x \Sigma$  does not have 1 as an eigenvalue. A nondegenerate orbit as above is *hyperbolic* if  $d\phi_x^k$  has real eigenvalues. We say that  $\phi$  is nondegenerate if all of its periodic orbits (including multiply covered periodic orbits where the points in  $\Sigma$  are not distinct) are nondegenerate; this holds for  $C^\infty$  generic  $\phi$  in any Hamiltonian isotopy class  $\Phi$ .

**Definition 2.2.** Assume that  $\phi$  is nondegenerate. A *PFH generator* is an orbit set  $\alpha = \{(\alpha_i, m_i)\}$  such that  $m_i = 1$  whenever  $\alpha_i$  is hyperbolic.

**Notation 2.3.** If  $\gamma$  and  $\gamma'$  are 1-cycles<sup>2</sup> in  $Y_\phi$  with  $[\gamma] = [\gamma'] \in H_1(Y_\phi)$ , let  $H_2(Y_\phi, \gamma, \gamma')$  denote the set of relative homology classes of 2-chains  $Z$  in  $Y_\phi$  with  $\partial Z = \gamma - \gamma'$ . This is an affine space over  $H_2(Y_\phi)$ .

To define the differential on the chain complex below, we will need to choose a generic almost complex structure  $J$  on  $\mathbb{R} \times Y_\phi$  satisfying the following conditions. To state them, let  $E \rightarrow Y_\phi$  denote the vertical tangent bundle of  $Y_\phi \rightarrow S^1$ ; this subbundle of  $TY_\phi$  plays an analogous role in PFH to the contact structure  $\xi$  in ECH.

**Definition 2.4.** An almost complex structure  $J$  on  $\mathbb{R} \times Y_\phi$  is *admissible* if:

- $J(\partial_s) = \partial_t$ , where  $s$  denotes the  $\mathbb{R}$  coordinate on  $\mathbb{R} \times Y_\phi$ .
- $J$  is independent of  $s$ , i.e. invariant under translation of the  $\mathbb{R}$  factor in  $\mathbb{R} \times Y_\phi$ .
- $J(E) = E$ , rotating positively with respect to  $\omega$ .

Fix a generic admissible  $J$  as above. If  $\alpha$  and  $\beta$  are PFH generators with  $[\alpha] = [\beta]$ , let  $\mathcal{M}^J(\alpha, \beta)$  denote the moduli space of  $J$ -holomorphic currents (finite positive integer linear combinations of  $J$ -holomorphic curves) in  $\mathbb{R} \times Y_\phi$  which as currents are asymptotic to  $\alpha$  as  $s \rightarrow +\infty$  and to  $\beta$  as  $s \rightarrow -\infty$ . For  $Z \in H_2(Y, \alpha, \beta)$ , let  $\mathcal{M}^J(\alpha, \beta, Z)$  denote the set of such currents that represent the relative homology class  $Z$ . See [20, §3] for more precise definitions in the analogous case of ECH. Note that  $\mathbb{R}$  acts on  $\mathcal{M}^J(\alpha, \beta, Z)$  by translation of the  $\mathbb{R}$  coordinate on  $\mathbb{R} \times Y_\phi$ .

We note for later use that the admissibility conditions on the almost complex structure  $J$  imply that the restriction of  $\omega_\phi$  to any  $J$ -holomorphic curve is pointwise nonnegative. Consequently,

$$\mathcal{M}^J(\alpha, \beta, Z) \neq \emptyset \implies \int_Z \omega_\phi \geq 0. \quad (2.1)$$

Given  $\alpha = \{(\alpha_i, m_i)\}$ ,  $\beta = \{(\beta_j, n_j)\}$ , and  $Z$  as above, the *ECH index*<sup>3</sup> is defined to be

$$I(\alpha, \beta, Z) = c_\tau(Z) + Q_\tau(Z) + \sum_i \sum_{k=1}^{m_i} CZ_\tau(\alpha_i^k) - \sum_j \sum_{k=1}^{n_j} CZ_\tau(\beta_j^k) \in \mathbb{Z}. \quad (2.2)$$

Here  $\tau$  is a homotopy class of trivialization of the bundle  $E$  over the orbits  $\alpha_i$  and  $\beta_j$ , while  $c_\tau$  denotes the relative first Chern class,  $Q_\tau$  denotes the relative

---

<sup>2</sup>In this paper, a “1-cycle” will always be a finite integer linear combination of closed oriented 1-dimensional submanifolds.

<sup>3</sup>Perhaps here it should be called the “PFH index”.



self-intersection number, and  $\text{CZ}_\tau(\gamma^k)$  denotes the Conley-Zehnder index of the  $k^{\text{th}}$  iterate of the periodic orbit  $\gamma$  with respect to  $\tau$ . The detailed definitions of these notions are not important for the present discussion but may be found in [20, §3] for the analogous case of ECH, or in [18, §2] for the more general case of stable Hamiltonian structures which includes both PFH and ECH. If  $C \in \mathcal{M}^J(\alpha, \beta, Z)$ , we write  $I(C) = I(\alpha, \beta, Z)$ .

A key property of the ECH index is that if  $J$  is generic (so that relevant moduli spaces of somewhere injective  $J$ -holomorphic curves are cut out transversely), if  $[\alpha] \cdot [\Sigma] > g$  (see Remark 2.10 below), and if  $I(C) = 1$ , then  $C$  is a (possibly empty) finite linear combination of “trivial cylinders”  $\mathbb{R} \times \eta$  where  $\eta$  is a periodic orbit, plus an embedded holomorphic curve of Fredholm index 1, so that it lives in a 1-dimensional moduli space, which becomes 0 dimensional after modding out by the  $\mathbb{R}$  action above. See e.g. [20, Prop. 3.7].

## 2.2 The chain complex

To define the PFH of  $\phi$  in general, we need to keep track of some information about relative homology classes of holomorphic curves in  $\mathbb{R} \times Y_\phi$ . There are different options for how to do this, resulting in different versions of PFH. We will use a version with Novikov ring coefficients, which depends on the following choice:

**Choice 2.5.** Let  $\text{Ker}([\omega_\phi])$  denote the kernel of  $\langle [\omega_\phi], \cdot \rangle : H_2(Y_\phi) \rightarrow \mathbb{R}$ . Below, fix a subgroup  $G \subset \text{Ker}([\omega_\phi])$ . On a first reading it may be simplest to just consider the case  $G = \{0\}$ , although later we will find it convenient to choose  $G = \text{Ker}([\omega_\phi])$ .

**Definition 2.6.** Let  $q$  be a formal variable and<sup>4</sup> write  $\mathbb{F} = \mathbb{Z}/2$ . Let  $\Lambda_G$  denote the Novikov ring consisting of formal sums

$$\sum_{A \in H_2(Y_\phi)/G} p_A q^A$$

where  $p_A \in \mathbb{F}$ , such that for each  $R \in \mathbb{R}$ , there are only finitely many  $A$  such that  $p_A \neq 0$  and  $\langle [\omega_\phi], A \rangle > R$ .

**Definition 2.7.** A *reference cycle* for  $\phi$  is a 1-cycle  $\gamma$  in  $Y_\phi$ . We define the *degree*  $d(\gamma) = [\gamma] \cdot [\Sigma] \in \mathbb{Z}$ , where  $[\Sigma] \in H_2(Y_\phi)$  denotes the homology class of a fiber of  $Y_\phi \rightarrow S^1$ . Note that if  $\alpha = \{(\alpha_i, m_i)\}$  is an orbit set with  $[\alpha] = [\gamma] \in H_1(Y_\phi)$ , then the total period of the orbits  $\alpha_i$ , counted with their multiplicities  $m_i$ , must equal the degree  $d(\gamma)$ .

**Definition 2.8.** Fix a subgroup  $G$  as above and a reference cycle  $\gamma$  for  $\phi$ . A  $(G, \gamma)$ -*anchored orbit set* is a pair  $(\alpha, Z)$ , where  $\alpha$  is an orbit set with  $[\alpha] = [\gamma] \in H_1(Y_\phi)$ , and  $Z \in H_2(Y_\phi, \alpha, \gamma)/G$ .

---

<sup>4</sup>One can also use  $\mathbb{Z}$  coefficients instead of  $\mathbb{Z}/2$ ; it is explained in [26, §9] how to set up orientations for the differential in ECH, and this carries over to PFH. However the applications in this paper, and all other applications of ECH and PFH so far, only need  $\mathbb{Z}/2$  coefficients.

We define the *symplectic action* of  $(\alpha, Z)$  by

$$\mathcal{A}(\alpha, Z) = \int_Z \omega_\phi.$$

When  $\phi$  is nondegenerate, a  $(G, \gamma)$ -anchored PFH generator is a  $(G, \gamma)$ -anchored orbit set  $(\alpha, Z)$  for which  $\alpha$  is a PFH generator.

We can now define the periodic Floer homology  $HP(\phi, \gamma, G)$ , which is a  $\Lambda_G$ -module.

**Definition 2.9.** If  $\phi$  is nondegenerate and  $\gamma$  is a reference cycle, define  $CP(\phi, \gamma, G)$  to be the set of (possibly infinite) formal sums

$$\sum_{\alpha, Z} n_{\alpha, Z}(\alpha, Z) \tag{2.3}$$

where:

- The sum is over  $(G, \gamma)$ -anchored PFH generators  $(\alpha, Z)$ .
- Each coefficient  $n_{\alpha, Z} \in \mathbb{F}$ .
- For each  $R \in \mathbb{R}$ , there are only finitely many  $(\alpha, Z)$  such that  $n_{\alpha, Z} \neq 0$  and  $\mathcal{A}(\alpha, Z) > R$ .

Then  $CP(\phi, \gamma, G)$  is a  $\Lambda_G$ -module, with the  $\Lambda_G$ -action given by

$$\sum_{A \in H_2(Y_\phi)/G} p_A q^A \cdot \sum_{\alpha, Z} n_{\alpha, Z}(\alpha, Z) = \sum_{\alpha, W} \left( \sum_{A \in H_2(Y_\phi)/G} p_A n_{\alpha, W-A} \right) (\alpha, W).$$

The finiteness conditions imply that the right hand side<sup>5</sup> is a well defined element of  $CP(\phi, \gamma, G)$ .

**Remark 2.10.** In general, to define the differential when  $d(\gamma) \leq g$ , one needs to choose a slight perturbation of an admissible almost complex structure, relaxing the last condition in Definition 2.4. This is because if  $J(E) = E$ , then each fiber of  $\mathbb{R} \times Y_\phi \rightarrow \mathbb{R} \times S^1$  is a  $J$ -holomorphic curve, which is not cut out transversely when  $g > 0$ , and this interferes with compactness arguments to define the PFH differential when  $d(\gamma) \leq g$ . See e.g. [17, §9.5]. For simplicity, **we assume below that**  $d(\gamma) > g$  so that we can stick with admissible almost complex structures. The theory below can be extended to the case  $d(\gamma) \leq g$  with some additional work, but this will not be necessary for the applications here.

---

<sup>5</sup>One can also write the right hand side in more informal notation as  $\sum_{A, \alpha, Z} p_A n_{\alpha, Z}(\alpha, Z + A)$ .

**Definition 2.11.** For generic admissible  $J$ , we define a differential

$$\partial_J : CP(\phi, \gamma, G) \longrightarrow CP(\phi, \gamma, G)$$

by

$$\partial_J \sum_{\alpha, Z} n_{\alpha, Z}(\alpha, Z) = \sum_{\beta, W} \left( \sum_{\alpha} \sum_{\substack{Z \in H_2(Y, \alpha, \beta) \\ I(\alpha, \beta, Z) = 1}} n_{\alpha, W+Z} \# \frac{\mathcal{M}^J(\alpha, \beta, Z)}{\mathbb{R}} \right) (\beta, W). \quad (2.4)$$

Here the first sum on the right hand side is over  $(G, \gamma)$ -anchored PFH generators  $(\beta, W)$ , the second sum on the right hand side is over PFH generators  $\alpha$  homologous to  $\gamma$ , and  $\#$  denotes the mod 2 count.

**Lemma 2.12.**  $\partial_J$  is well defined.

*Proof.* Let  $\sum_{\alpha, Z} n_{\alpha, Z}(\alpha, Z) \in CP(\phi, \gamma, G)$ . We need to show that for each real number  $R$ , there are only finitely many  $(G, \gamma)$ -anchored PFH generators  $(\beta, W)$  with  $\int_W \omega_\phi > R$  such that the sum in parentheses on the right hand side of (2.4) has any nonzero terms; and we need to show that for each such  $(\beta, W)$ , there are only finitely many nonzero terms, and that the mod 2 counts that arise are well-defined.

By (2.1), if  $\int_W \omega_\phi > R$  and  $\mathcal{M}^J(\alpha, \beta, Z)$  is nonempty, then  $\int_Z \omega_\phi \geq 0$ , so  $\int_{W+Z} \omega_\phi > R$ . Write  $V = W + Z$ . By the definition of  $CP(\phi, \gamma, G)$ , there are only finitely many  $(G, \gamma)$ -anchored PFH generators  $(\alpha, V)$  with  $\int_V \omega_\phi > R$  and  $n_{\alpha, V} \neq 0$ . For each such pair, and for each of the finitely many PFH generators  $\beta$  with  $[\beta] = [\gamma]$ , it follows from the compactness result<sup>6</sup> in [17, Thm. 1.8], that the union over  $W$  with  $\int_W \omega_\phi > R$  and  $I(\alpha, \beta, V - W) = 1$  of the moduli spaces  $\mathcal{M}^J(\alpha, \beta, V - W)/\mathbb{R}$  is finite.  $\square$

It follows from minor modifications of [25, Thm. 7.20] (which applies to ECH) that  $\partial_J^2 = 0$ .

**Definition 2.13.** We define the *periodic Floer homology*  $HP(\phi, \gamma, G)$  to be the homology of the chain complex  $(CP(\phi, \gamma, G), \partial_J)$ .

A special case of the isomorphism of Lee-Taubes [31] implies that  $HP(\phi, \gamma, G)$  is isomorphic to a version of Seiberg-Witten Floer cohomology. Together with the proof of [28, Thm. 1.3(a)] (which applies to the similar case of ECH), this shows that up to canonical isomorphism,  $HP(\phi, \gamma, G)$  does not depend on the choice of  $J$ .

---

<sup>6</sup>The compactness result and other results in [17] made an additional hypothesis of “ $d$ -admissibility”, asserting that  $(\phi, J)$  has a nice form near the periodic orbits of period at most  $d(\gamma)$ . This hypothesis is no longer needed thanks to the asymptotic analysis of Siefring [34].

**Remark 2.14.** If  $\gamma'$  is another reference cycle with  $[\gamma] = [\gamma']$ , and if  $Z \in H_2(Y_\phi, \gamma, \gamma')/G$ , then it follows from the definition (2.4) that there is an isomorphism of chain complexes

$$\psi_Z : (CP(\phi, \gamma, G), \partial_J) \xrightarrow{\cong} (CP(\phi, \gamma', G), \partial_J)$$

sending

$$\sum_{\alpha, W} n_{\alpha, W}(\alpha, W) \mapsto \sum_{\alpha, W} n_{\alpha, W+Z}(\alpha, W). \quad (2.5)$$

This induces an isomorphism of  $\Lambda_G$ -modules

$$\Psi_Z : HP(\phi, \gamma, G) \xrightarrow{\cong} HP(\phi, \gamma', G) \quad (2.6)$$

depending only on the relative homology class  $Z$ . Thus, up to noncanonical isomorphism,  $HP(\phi, \gamma, G)$  depends only on the diffeomorphism  $\phi$  and the homology class  $[\gamma]$ , and not on the choice of reference cycle  $\gamma$ .

We will see below that  $HP(\phi, \gamma, G)$  is also invariant (up to noncanonical isomorphism) under Hamiltonian isotopy of  $\phi$ . Thus for a Hamiltonian isotopy class  $\Phi$ , a homology class  $\Gamma \in H_1(Y_\phi)$ , and a subgroup  $G$  of  $\text{Ker}([\omega_\phi])$ , we have a well-defined isomorphism class of  $\Lambda_G$ -modules  $HP(\Phi, \Gamma, G)$ .

### 2.3 Examples of PFH

**Example 2.15.** Let  $\phi$  be the identity map on  $\Sigma$ . Although  $\phi$  is degenerate, one can define its PFH to be the PFH of a nondegenerate Hamiltonian perturbation; see Remark 3.3 below.

The mapping torus  $Y_\phi = S^1 \times \Sigma$ . The Novikov ring  $\Lambda_G$  consists of formal sums

$$\sum_{k \leq k_0} a_k q^{k[\Sigma]} \quad (2.7)$$

where

$$a_k \in \mathbb{F}[(H_1(S^1) \otimes H_1(\Sigma))/G].$$

Write  $[S^1] = [S^1] \times \{\text{pt}\} \in H_1(Y_\phi)$ . If  $d$  is a nonnegative integer and if  $\Gamma = d[S^1]$ , then we can choose the reference cycle  $\gamma$  to be  $d$  circles of the form  $S^1 \times \{x\}$ , and there is a canonical isomorphism

$$HP(\text{id}_\Sigma, d[S^1] \times \{x\}, G) = \text{Sym}^d H_*(\Sigma; \mathbb{F}) \otimes_{\mathbb{F}} \Lambda_G. \quad (2.8)$$

Here  $\text{Sym}^d$  denotes the degree  $d$  part of the graded symmetric product; given a homogeneous basis of  $H_*(\Sigma; \mathbb{F})$ , this is a vector space over  $\mathbb{F}$  with a basis consisting of symmetric degree  $d$  monomials in basis elements of  $H_*(\Sigma; \mathbb{F})$ , where basis elements in  $H_1(\Sigma; \mathbb{F})$  cannot be repeated.

To prove the isomorphism (2.8), one fixes  $d$  and replaces the identity map with the time 1 flow  $\phi$  of a  $C^2$ -small autonomous Hamiltonian  $H : \Sigma \rightarrow \mathbb{R}$  which is

a Morse function. It follows from Definition 2.2 that PFH generators in the class  $\Gamma = d[S^1]$  correspond to degree  $d$  symmetric monomials in critical points of  $H$ , where index 1 critical points cannot be repeated. One can choose a metric  $g_\Sigma$  on  $\Sigma$  making the pair  $(H, g_\Sigma)$  Morse-Smale, along with a corresponding almost complex structure  $J$  on  $\mathbb{R} \times Y_\phi$  for which Morse flow lines give rise to  $J$ -holomorphic cylinders. The  $S^1$  symmetry of the mapping torus can be used to show that no other  $J$ -holomorphic curves contribute to the PFH differential; this argument is worked out in [12, 32] for the very similar problem of computing the ECH of prequantization bundles. In particular, if we choose  $H$  to be a perfect Morse function, then the differential vanishes, and the chain complex agrees with the right hand side of (2.8).

For a homology class  $\Gamma \in H_1(S^1 \times \Sigma)$  which is not of the form  $d[S^1]$  for a nonnegative integer  $d$ , the PFH is zero, because after a small perturbation of the identity as above, there are no PFH generators in the class  $\Gamma$ .

Some more examples of PFH (more precisely untwisted PFH in the monotone case, see §2.6 below) are computed in [23] and [31, Cor. 1.5]. For classes  $\Gamma$  with  $d = \Gamma \cdot [\Sigma] = 1$ , the PFH reduces to Floer homology for symplectic fixed points, which has been computed by Cotton-Clay [3].

## 2.4 The $U$ map

There is also a well-defined map

$$U : HP(\phi, \gamma, G) \longrightarrow HP(\phi, \gamma, G).$$

This is induced by a chain map which is defined analogously to the differential (2.4); but instead of counting  $I = 1$  holomorphic currents modulo  $\mathbb{R}$  translation, it counts  $I = 2$  holomorphic currents that are constrained to pass through a base point in  $\mathbb{R} \times Y_\phi$ . See [27, §2.5] for details in the completely analogous case of ECH. We will see in Proposition 3.1 below that the  $U$  map is invariant (in a certain sense) under Hamiltonian isotopy of  $\phi$ .

**Example 2.16.** Suppose that  $\Sigma = S^2$  and  $\phi$  is Hamiltonian isotopic to the identity. Let  $d$  be a positive integer, and set  $\gamma = d[S^1] \times \{x\}$  as in Example 2.15. Here we must take  $G = \{0\}$ . Under the identification (2.8), denote the generators of  $\text{Sym}^d H_*(S^2; \mathbb{F})$ , in increasing homological degree, as  $e_{d,0}, e_{d,1}, \dots, e_{d,d}$ . Then a calculation as in [20, §4.1] shows that

$$\begin{aligned} Ue_{d,i} &= e_{d,i-1}, & i &= 1, \dots, d, \\ Ue_{d,0} &= q^{-[S^2]} e_{d,d}. \end{aligned}$$

The above example has an important property which we now formalize.

**Definition 2.17.** Let  $\phi$  be a (possibly degenerate) area-preserving diffeomorphism of  $(\Sigma, \omega)$ , let  $\gamma$  be a reference cycle for  $\phi$ , and let  $G$  be as in Choice 2.5. We say

that a nonzero element  $\sigma \in HP(\phi, \gamma, G)$  is *U-cyclic* if there is a positive integer  $m$  such that

$$U^{m(d(\gamma)-g+1)}\sigma = q^{-m[\Sigma]}\sigma. \quad (2.9)$$

We say that  $\sigma$  is *U-cyclic of order  $m$*  if  $m$  is the smallest positive integer with this property.

**Remark 2.18.** In general, if  $U^k\sigma = q^{-m[\Sigma]}\sigma$  for some  $k$ , then we must have  $k = m(d(\gamma)-g+1)$ . The reason is that in the nondegenerate case,  $U$  counts holomorphic currents with ECH index  $I = 2$ ; while if  $\alpha$  is any PFH generator with  $\alpha \cdot [\Sigma] = d$ , then it follows from equation (2.2) that the ECH index  $I(\alpha, \alpha, [\Sigma]) = 2(d - g + 1)$ .

**Example 2.19.** Suppose that  $\Sigma = S^2$  and  $\gamma = d[S^1] \times \{x\}$  where  $d$  is a positive integer. Then it follows from Example 2.16 that  $e_{d,i}$ , and indeed every nonzero element of  $HP(\phi, \gamma, \{0\})$ , is *U-cyclic* (of order 1).

**Definition 2.20.** We say that the Hamiltonian isotopy class  $[\phi]$  has the *U-cycle property* if there exist *U-cyclic* elements with arbitrarily large degree. That is, we require that for all positive integers  $d_0$ , there exist a subgroup  $G \subset \text{Ker}([\omega_\phi])$ , a reference cycle  $\gamma$  for  $\phi$  with  $d(\gamma) \geq d_0$ , and a *U-cyclic* element  $\sigma \in HP(\phi, \gamma, G)$ . (This condition is invariant under Hamiltonian isotopy.)

## 2.5 Filtered PFH

Fix a nondegenerate  $\phi$ , a reference cycle  $\gamma$ , a group  $G$  as in Choice 2.5, and a real number  $L$ . Define  $CP^L(\phi, \gamma, G)$  to be the set of formal sums (2.3) in  $CP(\phi, \gamma, G)$  such that  $\mathcal{A}(\alpha, Z) < L$  whenever  $n_{\alpha, Z} \neq 0$ . For a generic admissible  $J$ , it follows from (2.1) that  $CP^L(\phi, \gamma, G)$  is a subcomplex of  $(CP(\phi, \gamma, G), \partial_J)$ .

**Definition 2.21.** We define the *filtered PFH*, denoted by  $HP^L(\phi, \gamma, G)$ , to be the homology of the subcomplex  $(CP^L(\phi, \gamma, G), \partial_J)$ .

Inclusion of chain complexes induces a map

$$\iota^L : HP^L(\phi, \gamma, G) \longrightarrow HP(\phi, \gamma, G). \quad (2.10)$$

Similarly we have inclusion-induced maps

$$\iota^{L_1, L_2} : HP^{L_1}(\phi, \gamma, G) \longrightarrow HP^{L_2}(\phi, \gamma, G) \quad (2.11)$$

for  $L_1 \leq L_2$ , and with respect to these maps,  $HP(\phi, \gamma, G)$  is the direct limit of  $HP^L(\phi, \gamma, G)$  as  $L \rightarrow \infty$ .

As in [28, Thm. 1.3], the filtered homology  $HP^L(\phi, \gamma, G)$ , as well as the inclusion-induced maps (2.10) and (2.11), do not depend on the choice of  $J$ .

## 2.6 The monotone case

We now recall two alternate versions of PFH which are defined in the following special situation, which is possible when  $[\phi]$  is rational:

**Definition 2.22.** For  $\Gamma \in H_1(Y_\phi)$ , we say that the pair  $(\phi, \Gamma)$  is *monotone* if the cohomology class  $[\omega] \in H^2(Y_\phi; \mathbb{R})$  is a real multiple of the image of the class

$$c_1(E) + 2\text{PD}(\Gamma) \in H^2(Y_\phi; \mathbb{Z}).$$

In this case, one can define a simpler, “untwisted” version<sup>7</sup> of PFH, which we denote here by  $\overline{HP}(\phi, \Gamma)$ . When  $\phi$  is nondegenerate, this is the homology of a chain complex  $\overline{CP}(\phi, \Gamma)$  which is freely generated over  $\mathbb{F}$  by the PFH generators in the homology class  $\Gamma$ . For a generic admissible almost complex structure  $J$ , the differential is defined by

$$\partial_J \alpha = \sum_{\beta} \sum_{\substack{Z \in H_2(Y_\phi, \alpha, \beta) \\ I(\alpha, \beta, Z)=1}} \# \frac{\mathcal{M}^J(\alpha, \beta, Z)}{\mathbb{R}} \beta. \quad (2.12)$$

In the monotone case one can also define a twisted version of PFH without using a Novikov ring<sup>8</sup>, which we denote here by  $\widetilde{HP}(\phi, \gamma, G)$ , where  $\gamma$  is a reference cycle and  $G$  is as in Choice 2.5. This version of PFH is a module over the group ring  $\mathbb{F}[H_2(Y_\phi)/G]$ . Again assuming that  $\phi$  is nondegenerate, it is the homology of a chain complex  $\widetilde{CP}(\phi, \gamma, G)$  which is freely generated over  $\mathbb{F}$  by  $(G, \gamma)$ -anchored PFH generators, and whose differential is defined by

$$\partial_J(\alpha, W) = \sum_{\beta} \sum_{\substack{Z \in H_2(Y_\phi, \alpha, \beta) \\ I(\alpha, \beta, Z)=1}} \# \frac{\mathcal{M}^J(\alpha, \beta, Z)}{\mathbb{R}} (\beta, W - Z). \quad (2.13)$$

The differentials (2.12) and (2.13) are well defined because when computing the differential of a generator, the monotonicity hypothesis implies that there is an upper bound on the integral of  $\omega_\phi$  over all holomorphic currents that one needs to count, so that one obtains a finite count; compare Lemma 2.12.

Suppose now that the reference cycle  $\gamma$  is positively transverse to the fibers of  $Y_\phi \rightarrow S^1$ . A framing  $\tau$  of  $\gamma$  then induces a  $\mathbb{Z}$ -grading on  $\widetilde{HP}(\phi, \gamma, G)$ . The grading of a generator  $(\alpha, Z)$  is defined by

$$|(\alpha, Z)| = I(\alpha, \gamma, Z) \quad (2.14)$$

where the right hand side is defined as in (2.2), but with no Conley-Zehnder terms for  $\gamma$ . The grading (2.14) descends to a  $\mathbb{Z}/N$  grading on  $\overline{HP}(\phi, \Gamma)$ , where  $N$  denotes the divisibility of  $c_1(E) + 2\text{PD}(\Gamma)$  in  $\text{Hom}(H_2(Y_\phi), \mathbb{Z})$ ; note that  $N$  is an even integer.

<sup>7</sup>This is the original version of PFH from [17, 23].

<sup>8</sup>This is analogous to the twisted ECH introduced in [24, §11.2].

The definition of the  $U$  map carries over to  $\overline{HP}$  and  $\widetilde{HP}$ , and with respect to the above gradings, it has degree  $-2$ .

Even in the monotone case, we will need to use a twisted version of PFH with reference cycles in order to define spectral invariants. We will later need the following relation between twisted and untwisted versions:

**Lemma 2.23.** *Suppose that  $\phi$  is nondegenerate and  $(\phi, \Gamma)$  is monotone, let  $\gamma$  be a reference cycle with  $[\gamma] = \Gamma$ , and choose  $G = \text{Ker}([\omega_\phi])$ . Then there is a noncanonical isomorphism of  $\Lambda_G$ -modules*

$$HP(\phi, \gamma, G) \simeq \overline{HP}(\phi, \Gamma) \otimes_F \Lambda_G. \quad (2.15)$$

Under the above isomorphism, if  $d = d(\Gamma)$ , then

$$U^{d-g+1} \longleftrightarrow U^{d-g+1} \otimes q^{-[\Sigma]}. \quad (2.16)$$

*Proof.* Let  $A \in H_2(Y_\phi)/G$  be the unique class such that  $\langle [\omega_\phi], A \rangle$  is positive and minimal. Then the Novikov ring  $\Lambda_G$  is canonically identified with  $\mathbb{F}((q^{-A}))$ , namely the ring of Laurent series in  $q^{-A}$  with coefficients in  $\mathbb{F}$ .

As in Remark 2.14, we can assume without loss of generality that  $\gamma$  is positively transverse to  $E$ . Choose a framing  $\tau$  of  $\gamma$  as needed to define a  $\mathbb{Z}$ -grading on  $\widetilde{HP}(\phi, \gamma, G)$  and a  $\mathbb{Z}/N$  grading on  $\overline{HP}(\phi, \Gamma)$ . It follows from the definitions that there is a canonical isomorphism

$$\widetilde{HP}_i(\phi, \gamma, G) = \overline{HP}_{i \bmod N}(\phi, \Gamma). \quad (2.17)$$

On the left hand side, multiplication by  $q^{-A}$  shifts the grading down by  $N$ . It follows that  $HP(\phi, \gamma, G)$  is canonically identified with the set of sequences  $(\sigma_i)_{i \in \mathbb{Z}}$  where  $\sigma_i \in \widetilde{HP}_i(\phi, \gamma, G)$  and  $\sigma_i = 0$  if  $i$  is sufficiently large. If we choose a right inverse of the projection  $\mathbb{Z} \rightarrow \mathbb{Z}/N$ , then together with (2.17) this defines an identification of the above set of sequences with  $\overline{HP}(\phi, \Gamma) \otimes_F \Lambda_G$ . This gives an isomorphism (2.15).

To prove (2.16), we observe that under the isomorphism (2.15) constructed above,

$$U^{N/2} \longleftrightarrow U^{N/2} \otimes q^{-A}.$$

The positive<sup>9</sup> integer  $2(d - g + 1)$  must be divisible by  $N$ , since  $c_1(E) + 2\text{PD}(\Gamma)$  evaluates to  $2(d - g + 1)$  on  $[\Sigma]$ . It follows that

$$U^{d-g+1} \longleftrightarrow U^{d-g+1} \otimes q^{-(2(d-g+1)/N)A}. \quad (2.18)$$

By monotonicity, we have  $\langle c_1(E) + 2\text{PD}(\Gamma), A \rangle = N$ , and it follows that  $(2(d - g + 1)/N)A = [\Sigma]$  in  $H_2(Y_\phi)/G$ . Putting this into (2.18) proves (2.16).  $\square$

---

<sup>9</sup>Recall from Remark 2.10 that we are making the standing assumption that  $d > g$ .



### 3 Invariance of PFH under Hamiltonian isotopy

We now work out how PFH and the additional structure on it defined above behave under Hamiltonian isotopy of  $\phi$ .

It is useful for our purposes to define a Hamiltonian isotopy of  $\phi$  via a smooth map

$$H : Y_\phi \longrightarrow \mathbb{R}.$$

Under the projection  $[0, 1] \times \Sigma \rightarrow Y_\phi$ , the map  $H$  pulls back to a map  $\tilde{H} : [0, 1] \times \Sigma \rightarrow \mathbb{R}$  satisfying  $\tilde{H}(1, x) = \tilde{H}(0, \phi(x))$ . For  $t \in [0, 1]$ , let  $H_t = \tilde{H}(t, \cdot) : \Sigma \rightarrow \mathbb{R}$ , and let  $X_{H_t}$  denote the associated Hamiltonian vector field on  $\Sigma$ ; we use the sign convention  $\omega(X_{H_t}, \cdot) = dH_t$ . Let  $\{\varphi_t\}_{t \in [0, 1]}$  denote the Hamiltonian isotopy defined by  $\varphi_0 = \text{id}_\Sigma$  and  $\partial_t \varphi_t = X_{H_t} \circ \varphi_t$ . We define  $\phi_H = \phi \circ \varphi_1$ .

As in (1.2), we have a diffeomorphism

$$f_H : Y_\phi \xrightarrow{\cong} Y_{\phi_H}$$

defined by the diffeomorphism of  $[0, 1] \times \Sigma$  sending

$$(t, x) \longmapsto (t, \varphi_t^{-1}(x)).$$

If  $\gamma$  is a reference cycle in  $Y_\phi$ , let  $\gamma_H$  denote its pushforward  $(f_H)_\# \gamma$  in  $Y_{\phi_H}$ .

**Proposition 3.1.** *Let  $\phi$  be a (possibly degenerate) area-preserving diffeomorphism of  $(\Sigma, \omega)$ , let  $\gamma \subset Y_\phi$  be a reference cycle, and fix  $G$  as in Choice 2.5. For  $H_1, H_2 : Y_\phi \rightarrow \mathbb{R}$  with  $H_1 < H_2$  and  $\phi_{H_1}, \phi_{H_2}$  nondegenerate, there is a canonical isomorphism*

$$\Psi_{H_1, H_2} : HP(\phi_{H_2}, \gamma_{H_2}, G) \longrightarrow HP(\phi_{H_1}, \gamma_{H_1}, G) \quad (3.1)$$

with the following properties:

(a) If  $H_2 < H_3$  and if  $\phi_{H_3}$  is also nondegenerate, then

$$\Psi_{H_1, H_3} = \Psi_{H_1, H_2} \circ \Psi_{H_2, H_3} : HP(\phi_{H_3}, \gamma_{H_3}, G) \longrightarrow HP(\phi_{H_1}, \gamma_{H_1}, G).$$

(b)  $U \circ \Psi_{H_1, H_2} = \Psi_{H_1, H_2} \circ U$ .

(c) The isomorphism (3.1) is the direct limit as  $L \rightarrow \infty$  of canonical maps

$$\Psi_{H_1, H_2}^L : HP^L(\phi_{H_2}, \gamma_{H_2}, G) \longrightarrow HP^{L+\Delta}(\phi_{H_1}, \gamma_{H_1}, G) \quad (3.2)$$

where  $\Delta = \int_\gamma (H_2 - H_1) dt$ .

(d) If  $H_2 - H_1$  is a constant  $C > 0$ , so that  $\phi_{H_1} = \phi_{H_2}$  and  $\gamma_{H_1} = \gamma_{H_2}$ , then  $\Psi_{H_1, H_2}^L = \iota^{L, L+dC}$  where  $d = [\gamma] \cdot [\Sigma]$ . In particular,  $\Psi_{H_1, H_2}$  is the identity map.

*Proof.* We proceed in 6 steps.

*Step 1.* To prepare to define the map  $\Psi_{H_1, H_2}$ , we construct a “strong symplectic cobordism of stable Hamiltonian structures” between  $(Y_{\phi_{H_1}}, \omega_{\phi_{H_1}})$  and  $(Y_{\phi_{H_2}}, \omega_{\phi_{H_2}})$  as follows.

Consider the “symplectization” of the mapping torus defined by

$$X = \mathbb{R} \times Y_\phi$$

with the symplectic form

$$\omega_X = ds \wedge dt + \omega_\phi.$$

Here  $s$  denotes the  $\mathbb{R}$  coordinate on  $\mathbb{R} \times Y_\phi$ .

Given  $H : Y_\phi \rightarrow \mathbb{R}$ , define an inclusion

$$\begin{aligned} \iota_H : Y_\phi &\longrightarrow \mathbb{R} \times Y_\phi, \\ z &\longmapsto (H(z), z). \end{aligned}$$

We can then identify the mapping torus  $Y_{\phi_H}$  with a hypersurface in  $\mathbb{R} \times Y_\phi$  via the inclusion

$$\iota_H \circ f_H^{-1} : Y_{\phi_H} \longrightarrow \mathbb{R} \times Y_\phi. \quad (3.3)$$

Note that there is a symplectomorphism

$$\mathbb{R} \times Y_{\phi_H} \longrightarrow \mathbb{R} \times Y_\phi \quad (3.4)$$

induced by the symplectomorphism of  $(\mathbb{R} \times [0, 1] \times \Sigma, ds \wedge dt + \omega)$  sending

$$(s, t, x) \longmapsto (s + H(t, x), t, \varphi_t(x)).$$

The inclusion (3.3) is the restriction of the symplectomorphism (3.4) to  $\{0\} \times Y_{\phi_H}$ . It follows that

$$(\iota_H \circ f_H^{-1})^* \omega_X = \omega_{\phi_H}. \quad (3.5)$$

We note also that under the inclusion (3.3), the reference cycle  $\gamma_H$  corresponds to the graph of  $H$  on  $\gamma$  in  $\mathbb{R} \times Y_\phi$ .

Now if  $H_1 < H_2$ , define

$$M = \{(s, z) \in \mathbb{R} \times Y_\phi \mid H_1(z) \leq s \leq H_2(z)\} \quad (3.6)$$

with the symplectic form  $\omega_M = (\omega_X)|_M$ . It follows from the above calculations that the boundary components of  $M$  have neighborhoods in  $M$  symplectomorphic to  $[0, \varepsilon) \times Y_{\phi_{H_1}}$  and  $(-\varepsilon, 0] \times Y_{\phi_{H_2}}$ , where the latter manifolds are equipped with the restrictions of the symplectic forms on the symplectizations of  $Y_{\phi_{H_1}}$  and  $Y_{\phi_{H_2}}$ . Using these neighborhood identifications, we can glue to form the “symplectization completion” of  $M$ , which is a symplectic four-manifold

$$\overline{M} = \left( (-\infty, 0] \times Y_{\phi_{H_1}} \right) \bigcup_{Y_{\phi_{H_1}}} M \bigcup_{Y_{\phi_{H_2}}} \left( [0, \infty) \times Y_{\phi_{H_2}} \right). \quad (3.7)$$

We note that there is a canonical symplectomorphism

$$\overline{M} \simeq \mathbb{R} \times Y_\phi \quad (3.8)$$

which is the inclusion on  $M$ , and which on the rest of (3.7) is defined using the restrictions of the symplectomorphisms (3.4) for  $H_1$  and  $H_2$ .

*Step 2.* Suppose now that  $\phi_{H_1}$  and  $\phi_{H_2}$  are nondegenerate. Observe that  $V = (\mathbb{R} \times \gamma) \cap M$  defines a 2-chain in the cobordism  $M$  with  $\partial V = \gamma_{H_2} - \gamma_{H_1}$ . The cobordism  $M$ , together with the 2-chain  $V$ , induces the desired map  $\Psi_{H_1, H_2}$  in (3.1), as a special case of a general construction of cobordism maps<sup>10</sup> on PFH by Chen [2, Thm. 1]. The PFH cobordism maps in [2, Thm. 1] satisfy a composition property which implies assertion (a), and they commute with the  $U$  maps, giving assertion (b).

*Step 3.* To prove assertion (c), we will use the fact from [2, Thm. 1] that the map  $\Psi_{H_1, H_2}$  satisfies a crucial “holomorphic curve axiom”. We now state this property.

Let  $J_1$  and  $J_2$  be almost complex structures on  $\mathbb{R} \times Y_{\phi_{H_1}}$  and  $\mathbb{R} \times Y_{\phi_{H_2}}$  as needed to define differentials  $\partial_{J_1}$  on  $CP(\phi_{H_1}, \gamma_{H_1}, G)$  and  $\partial_{J_2}$  on  $CP(\phi_{H_2}, \gamma_{H_2}, G)$ . We can extend  $J_1$  and  $J_2$  to an almost complex structure  $J$  on  $\overline{M}$  compatible with the symplectic form  $\omega_M$ .

Let  $\alpha$  be an orbit set for  $\phi_{H_2}$  and let  $\beta$  be an orbit set for  $\phi_{H_1}$ . Define a *broken  $J$ -holomorphic current* in  $\overline{M}$  from  $\alpha$  to  $\beta$  to be a tuple  $(C_{k_+}, C_{k_+-1}, \dots, C_{k_-})$  where  $k_+ \geq 0 \geq k_-$ , and there are orbit sets  $\alpha = \alpha(k_+), \alpha(k_+ - 1), \dots, \alpha(0)$  for  $\phi_{H_2}$  and orbit sets  $\beta(0), \beta(-1), \dots, \beta(k_-) = \beta$  for  $\phi_{H_1}$ , such that:

- $C_i \in \mathcal{M}^{J_2}(\alpha(i), \alpha(i-1))/\mathbb{R}$  for  $i > 0$ .
- $C_0 \in \mathcal{M}^J(\alpha(0), \beta(0))$ . That is,  $C_0$  is a  $J$ -holomorphic current in  $\overline{M}$  which as a current is asymptotic to  $\alpha(0)$  as  $s \rightarrow \infty$  on  $[0, \infty) \times Y_{\phi_{H_2}}$ , and asymptotic to  $\beta(0)$  as  $s \rightarrow -\infty$  on  $(-\infty, 0] \times Y_{\phi_{H_1}}$ .
- $C_i \in \mathcal{M}^{J_1}(\beta(i+1), \beta(i))/\mathbb{R}$  for  $i < 0$ .

The holomorphic curves axiom now states that the map  $\Psi$  is induced by a (noncanonical) chain map

$$\psi : (CP(\phi_{H_2}, \gamma_{H_2}, G), \partial_{J_2}) \longrightarrow (CP(\phi_{H_1}, \gamma_{H_1}, G), \partial_{J_1})$$

with the following property. Similarly to (2.4), we can write  $\psi$  in the form

$$\psi \sum_{\alpha, Z} n_{\alpha, Z}(\alpha, Z) = \sum_{\beta, W} \left( \sum_{\alpha} \sum_{\substack{Z \in H_2(Y_\phi, \alpha, \beta) \\ I(\alpha, \beta, Z) = 0}} n_{\alpha, W+Z} m_{\alpha, \beta, Z} \right) (\beta, W). \quad (3.9)$$

---

<sup>10</sup>This is related to the construction of cobordism maps on ECH in [28, Thm. 1.9].

Here the first sum on the right hand side is over  $(G, \gamma_{H_1})$ -anchored PFH generators  $(\beta, W)$  for  $\phi_{H_1}$ , the second sum on the right hand side is over PFH generators  $\alpha$  for  $\phi_{H_2}$  in the homology class  $[\gamma_{H_2}]$ , and  $m_{\alpha, \beta, Z} \in \mathbb{F}$ . The key property is now:

- (\*) If the coefficient  $m_{\alpha, \beta, Z} \neq 0$ , then there is a broken  $J$ -holomorphic current<sup>11</sup> in  $\overline{M}$  from  $\alpha$  to  $\beta$  which, under the identification (3.8), represents the relative homology class  $Z$ .

*Step 4.* We claim now that if  $(\beta, W)$  is a  $(G, \gamma_{H_1})$ -anchored PFH generator for  $\phi_{H_1}$ , if  $(\alpha, W + Z)$  is a  $(G, \gamma_{H_2})$ -anchored PFH generator for  $\phi_{H_2}$ , and if there exists a broken  $J$ -holomorphic current in  $\overline{M}$  from  $\alpha$  to  $\beta$  in the relative homology class  $Z$ , then

$$\int_W \omega_{\phi_{H_1}} \leq \int_{W+Z} \omega_{\phi_{H_2}} + \Delta. \quad (3.10)$$

To prove this, by (2.1) we can assume without loss of generality that the  $J$ -holomorphic current has  $k_+ = k_- = 0$ , and thus consists of a single  $J$ -holomorphic current  $C \in \mathcal{M}^J(\alpha, \beta, Z)$ . We can also assume without loss of generality (by a slight modification of the cobordism) that  $C$  is transverse to  $\partial M$ . Under the decomposition (3.7), we can divide  $C$  into three pieces: let

$$\begin{aligned} C_1 &= C \cap \left( (-\infty, 0] \times Y_{\phi_{H_1}} \right), \\ C_0 &= C \cap M, \\ C_2 &= C \cap \left( [0, \infty) \times Y_{\phi_{H_2}} \right). \end{aligned}$$

Since the almost complex structures  $J_1$  and  $J_2$  are admissible, as in (2.1) we have

$$\int_{C_1} \omega_{\phi_{H_1}} \geq 0, \quad (3.11)$$

$$\int_{C_2} \omega_{\phi_{H_2}} \geq 0. \quad (3.12)$$

Also, since  $J$  is  $\omega_M$ -compatible, we have

$$\int_{C_0} \omega_M \geq 0. \quad (3.13)$$

We now deduce (3.10) by applying Stokes's theorem. To start, write  $\eta_1 = C_1 \cap Y_{\phi_{H_1}}$  and  $\eta_2 = C_2 \cap Y_{\phi_{H_2}}$ . Then  $C_1$  projects, via the projection  $(-\infty, 0] \times Y_{\phi_{H_1}} \rightarrow Y_{\phi_{H_1}}$ , to a relative homology class of 2-chain  $[C_1] \in H_2(Y_{\phi_{H_1}}, \eta_1, \beta)$ . Likewise,  $C_2$  projects to a relative homology class of 2-chain  $[C_2] \in H_2(Y_{\phi_{H_2}}, \alpha, \eta_2)$ .

---

<sup>11</sup>The “holomorphic curves axiom” as stated in [2, Thm. 1] implies a slightly weaker statement than (\*), namely that for fixed  $\alpha$  and  $\beta$ , if any of the coefficients  $m_{\alpha, \beta, Z}$  is nonzero, then there exists a broken  $J$ -holomorphic current from  $\alpha$  to  $\beta$ . The property (\*) follows from the same argument, keeping track of the relative homology classes of the holomorphic currents.

It follows from the homological assumption on  $C$  that in  $M$ , the 2-cycle

$$(\iota_{H_1} \circ f_{H_1}^{-1})_{\#}([C_1] + W) + (\iota_{H_2} \circ f_{H_2}^{-1})_{\#}([C_2] - W - Z) + C_0 - V$$

is nullhomologous. Consequently the integral of the closed 2-form  $\omega_M$  over this 2-cycle vanishes. By (3.5), this means that

$$\int_{C_1+W} \omega_{\phi_{H_1}} + \int_{C_2-W-Z} \omega_{\phi_{H_2}} + \int_{C_0} \omega_M - \Delta = 0. \quad (3.14)$$

Combining (3.11), (3.12), (3.13), and (3.14) proves (3.10).

*Step 5.* We now prove (c). It follows from Step 4 that the chain map (3.9) restricts to a chain map

$$\psi^L : (CP^L(\phi_{H_2}, \gamma_{H_2}, G), \partial_{J_2}) \longrightarrow (CP^{L+\Delta}(\phi_{H_1}, \gamma_{H_1}, G), \partial_{J_1})$$

We define the map  $\Psi_{H_1, H_2}^L$  in (3.2) to be the map on filtered PFH induced by  $\psi^L$ . Although the chain map  $\psi$  is defined only up to chain homotopy, the chain homotopies satisfy a version of the “holomorphic curves axiom” which implies that  $\Psi_{H_1, H_2}^L$  depends only on  $J_1$  and  $J_2$ . See e.g. [21, Prop. 6.2] for an analogous argument in the case of ECH. The map  $\Psi_{H_1, H_2}^L$  does not depend on  $J_1$  and  $J_2$  either. An analogous statement for ECH was proved in [28, Thm. 1.9], and this carries over to the case of PFH using the analysis of Chen [2, Thm. 1].

*Step 6.* Finally, the proof of property (d) follows the proof of the “scaling” property for ECH cobordism maps in [28, Thm. 1.9].  $\square$

**Remark 3.2.** Naively one would like to define the chain map (3.9) by taking  $m_{\alpha, \beta, Z}$  to be a count of  $I = 0$  holomorphic currents in  $\mathcal{M}^J(\alpha, \beta, Z)$ . Unfortunately it is not currently known in general<sup>12</sup> how to directly count  $J$ -holomorphic currents with  $I = 0$  in a completed cobordism, due to transversality difficulties with multiply covered holomorphic curves; see [20, §5.5] for explanation in the case of ECH, where there are similar issues. The actual chain map (3.9) is defined instead by counting solutions to the Seiberg-Witten equations on  $\overline{M}$ , using the metric determined by  $J$  and the symplectic form, and perturbed using a large multiple of the symplectic form as in [31].

**Remark 3.3.** It follows from Proposition 3.1 that if  $\phi$  is degenerate, then we can define  $HP(\phi, \gamma, G)$  by first perturbing  $\phi$  to be nondegenerate via a Hamiltonian isotopy; by parts (b) and (d) of the proposition, the PFH modules for such perturbations of  $\phi$  are all canonically isomorphic to each other.

<sup>12</sup>Such a count is possible in some special cases; see e.g. [14], [2, Thm. 2], and [33].

## 4 Spectral invariants in PFH

We now define spectral invariants in PFH, analogously to the spectral invariants in ECH defined in [19, Def. 4.1].

**Definition 4.1.** Suppose that  $\phi$  is nondegenerate, let  $\gamma$  be a reference cycle, fix  $G$  as in Choice 2.5, and let  $\sigma$  be a nonzero class in  $HP(\phi, \gamma, G)$ . Define the *spectral invariant*

$$c_\sigma(\phi, \gamma) \in \mathbb{R}$$

to be the infimum over  $L \in \mathbb{R}$  such that  $\sigma$  is in the image of the inclusion-induced map (2.10).

We now establish some properties of the spectral invariants  $c_\sigma$ . We first consider the dependence of  $c_\sigma$  on basic choices. Given a nonzero Novikov ring element  $\lambda = \sum_{A \in H_2(Y_\phi)/G} p_A q^A \in \Lambda_G$ , define

$$|\lambda| = \max\{\langle [\omega], A \rangle \mid p_A \neq 0\}.$$

Note that this maximum is well-defined by the definition of the Novikov ring  $\Lambda_G$ .

**Proposition 4.2.** *Suppose  $\sigma \in HP(\phi, \gamma, G)$  is nonzero.*

(a) *If  $\lambda \in \Lambda_G$  is invertible, then*

$$c_{\lambda\sigma}(\phi, \gamma) = c_\sigma(\phi, \gamma) + |\lambda|.$$

(b) *In the notation of Remark 2.14, we have*

$$c_{\Psi_Z(\sigma)}(\phi, \gamma') = c_\sigma(\phi, \gamma) - \int_Z \omega.$$

*Proof.* (a) It follows from the definitions that multiplication by  $\lambda$  induces an isomorphism of chain complexes

$$(CP^L(\phi, \gamma, G), \partial_J) \xrightarrow{\simeq} (CP^{L+|\lambda|}(\phi, \gamma, G), \partial_J).$$

The induced isomorphism on homology

$$HP^L(\phi, \gamma, G) \xrightarrow{\simeq} HP^{L+|\lambda|}(\phi, \gamma, G) \tag{4.1}$$

respects the maps (2.10), and it follows that

$$c_{\lambda\sigma}(\phi, \gamma) \leq c_\sigma(\phi, \gamma) + |\lambda|.$$

The inverse of the isomorphism (4.1) is induced by multiplication of chains by  $\lambda^{-1}$ , and this implies the reverse inequality.

(b) This follows by a similar argument.  $\square$

We now begin to explore how spectral invariants behave under Hamiltonian isotopy, using the notation of Proposition 3.1.

**Proposition 4.3.** *Let  $\phi$  be a (possibly degenerate) area-preserving diffeomorphism of  $(\Sigma, \omega)$ , let  $\gamma$  be a reference cycle for  $\phi$ , let  $H_1, H_2 : Y_\phi \rightarrow \mathbb{R}$  with  $H_1 < H_2$ , and suppose that  $\phi_{H_1}$  and  $\phi_{H_2}$  are nondegenerate. Let  $\sigma_2$  be a nonzero class in  $HP(\phi_{H_2}, \gamma_{H_2}, G)$ . Write  $\sigma_1 = \Phi_{H_1, H_2}(\sigma_2) \in HP(\phi_{H_1}, \gamma_{H_1}, G)$ . Then*

$$c_{\sigma_1}(\phi_{H_1}, \gamma_{H_1}) \leq c_{\sigma_2}(\phi_{H_2}, \gamma_{H_2}) + \int_{\gamma} (H_2 - H_1) dt. \quad (4.2)$$

*Proof.* Write  $\Delta = \int_{\gamma} (H_2 - H_1) dt$ . By Proposition 3.1(c), for each real number  $L$  we have a commutative diagram

$$\begin{array}{ccc} HP^L(\phi_{H_2}, \gamma_{H_2}, G) & \xrightarrow{i^L} & HP(\phi_{H_2}, \gamma_{H_2}, G) \\ \Psi_{H_1, H_2}^L \downarrow & & \downarrow \Psi_{H_1, H_2} \\ HP^{L+\Delta}(\phi_{H_1}, \gamma_{H_1}, G) & \xrightarrow{i^{L+\Delta}} & HP(\phi_{H_1}, \gamma_{H_1}, G). \end{array}$$

It follows that if  $\sigma_2$  is in the image of the top arrow, then  $\sigma_1$  is in the image of the bottom arrow.  $\square$

**Corollary 4.4.** (a) *The definition of  $c_\sigma(\phi, \gamma)$  has a unique extension to the case where  $\phi$  is degenerate<sup>13</sup> such that the following continuity property holds: Let  $\sigma \in HP(\phi, \gamma, G)$ , let  $H_1, H_2 : Y_\phi \rightarrow \mathbb{R}$ , and for  $i = 1, 2$  let  $\sigma_i$  denote the corresponding class in  $HP(\phi_{H_i}, \gamma_{H_i}, G)$ . Then*

$$|c_{\sigma_1}(\phi_{H_1}, \gamma_{H_1}) - c_{\sigma_2}(\phi_{H_2}, \gamma_{H_2})| \leq d(\gamma) \max_{Y_\phi} |H_2 - H_1|. \quad (4.3)$$

(b) *The extended spectral invariants satisfy the conclusions of Propositions 4.2 and 4.3.*

*Proof.* This follows from Proposition 4.3 using the formal procedure in [19, §3.1] and [7, §2.5].  $\square$

**Proposition 4.5.** *Let  $\phi$  be an area-preserving diffeomorphism of  $(\Sigma, \omega)$ , possibly degenerate, and suppose that  $[\phi]$  is rational. Then for any reference cycle  $\gamma$ , any  $G$  as in Choice 2.5, and any nonzero class  $\sigma \in HP(\phi, \gamma, G)$ , there exists a  $(G, \gamma)$ -anchored orbit set  $(\alpha, Z)$  such that*

$$c_\sigma(\phi, \gamma) = \mathcal{A}(\alpha, Z).$$

---

<sup>13</sup>Here the PFH of a degenerate map  $\phi$  is defined by Remark 3.3.

*Proof.* To start, note that since  $[\phi]$  is rational, the set of values that  $[\omega] \in H^2(Y_\phi; \mathbb{R})$  takes on  $H_2(Y_\phi)$  is discrete.

Suppose first that  $\phi$  is nondegenerate. Then there are only finitely many PFH generators in the homology class  $[\gamma]$ . Let  $S$  denote the set of actions of  $(G, \gamma)$ -anchored PFH generators; then the set  $S$  is discrete. If  $L < L'$  and the interval  $[L, L')$  does not intersect  $S$ , then the inclusion-induced map

$$HP^L(\phi, \gamma, G) \longrightarrow HP^{L'}(\phi, \gamma, G)$$

is an isomorphism, since it is induced by an isomorphism of chain complexes. It then follows from Definition 4.1 that  $c_\sigma(\phi, \gamma) \in S$ .

Suppose now that  $\phi$  is degenerate. Let  $\{H_i\}_{i \geq 1}$  be a sequence of Hamiltonians converging to 0 in  $C^\infty$  such that each  $\phi_{H_i}$  is nondegenerate. Let  $\sigma_i$  denote the class in  $HP(\phi_{H_i}, \gamma_{H_i}, G)$  corresponding to  $\sigma$ . By the continuity in (4.3), we have

$$c_\sigma(\phi, \gamma) = \lim_{i \rightarrow \infty} c_{\sigma_i}(\phi_{H_i}, \gamma_{H_i}).$$

By the nondegenerate case, for each  $i$  there exists a  $(G, \gamma_{H_i})$ -anchored PFH generator  $(\alpha(i), Z(i))$  for  $\phi_{H_i}$  such that  $c_{\sigma_i}(\phi_{H_i}, \gamma_{H_i}) = \mathcal{A}(\alpha(i), Z(i))$ . Since  $\Sigma$  is compact and each periodic orbit in each  $\alpha(i)$  has period at most  $d(\gamma)$ , we can pass to a subsequence so that  $\alpha(i)$  converges to an orbit set  $\alpha$  for  $\phi$ . Then the distance from the sequence  $\mathcal{A}(\alpha(i), Z(i))$  to the set  $\{\mathcal{A}(\alpha, Z) \mid Z \in H_2(Y, \alpha, \gamma)/G\}$  limits to zero. Since the latter set is discrete by our rationality hypothesis, the sequence  $(\mathcal{A}(\alpha(i), Z(i)))$  converges to an element of it.  $\square$

**Remark 4.6.** Without the hypothesis that  $[\phi]$  is rational, Proposition 4.5 still holds if  $\phi$  is nondegenerate, by [38, Thm. 1.4]. However we do not know whether Proposition 4.5 extends to the case where  $[\phi]$  is not rational and  $\phi$  is degenerate.

## 5 The ball packing lemma

We now prove a key fact, Lemma 5.2 below, which will be needed for the proofs of the main theorems. This lemma gives a relation between the PFH spectral invariants of two different Hamiltonian perturbations of  $\phi$ .

To state the lemma, recall that for any symplectic four-manifold  $X$ , its ECH capacities are a sequence of real numbers

$$0 = c_0^{\text{ECH}}(X) < c_1^{\text{ECH}}(X) \leq c_2^{\text{ECH}}(X) \leq \dots \leq \infty$$

defined in [19]. We will need the following examples. For  $r > 0$ , define the ball

$$B(r) = \{z \in \mathbb{C}^2 \mid \pi|z|^2 \leq r\}$$



with the restriction of the standard symplectic form on  $\mathbb{C}^2 = \mathbb{R}^4$ . Note that the volume of the ball is given by

$$\text{vol}(B(r)) = \frac{r^2}{2}.$$

It is shown in [19] that the ECH capacities of a ball are given by

$$c_k^{\text{ECH}}(B(r)) = dr, \quad (5.1)$$

where  $d$  is the unique nonnegative integer such that

$$d^2 + d \leq 2k \leq d^2 + 3d.$$

Furthermore, the ECH capacities of a disjoint union are given by

$$c_k^{\text{ECH}}\left(\coprod_{i=1}^n X_i\right) = \max_{k_1 + \dots + k_n = k} \sum_{i=1}^n c_{k_i}^{\text{ECH}}(X_i). \quad (5.2)$$

The calculation in [19, Prop. 8.4] deduces from (5.1) and (5.2) that if  $X$  is a finite disjoint union of balls, then<sup>14</sup>

$$\lim_{k \rightarrow \infty} \frac{c_k^{\text{ECH}}(X)^2}{k} = 4 \text{vol}(X). \quad (5.3)$$

To simplify notation we use the following convention:

**Notation 5.1.** If  $\sigma \in \text{HP}(\phi, \gamma, G)$ , and if  $H : Y_\phi \rightarrow \mathbb{R}$ , let  $\sigma_H \in \text{HP}(\phi_H, \gamma_H, G)$  denote the class corresponding to  $\sigma$  under the canonical isomorphism given by Proposition 3.1 and Remark 3.3. Write

$$c_\sigma(\phi, \gamma, H) = c_{\sigma_H}(\phi_H, \gamma_H) \in \mathbb{R}.$$

**Lemma 5.2.** *Let  $\phi$  be an area-preserving diffeomorphism of  $(\Sigma, \omega)$ , and let  $H_1, H_2 : Y_\phi \rightarrow \mathbb{R}$  with  $H_1 \leq H_2$ . Let  $\gamma$  be a reference cycle for  $\phi$ , let  $\sigma \in \text{HP}(\phi, \gamma, G)$ , let  $k$  be a nonnegative integer, and suppose that  $U^k \sigma \neq 0$ . Let  $X$  be a finite disjoint union of balls symplectically embedded in the cobordism  $M$  from (3.6). Then*

$$c_{U^k \sigma}(\phi, \gamma, H_1) \leq c_\sigma(\phi, \gamma, H_2) + \int_\gamma (H_2 - H_1) dt - c_k^{\text{ECH}}(X).$$

**Remark 5.3.** When  $k = 0$ , Lemma 5.2 reduces to the inequality (4.2).

The  $k = 1$  case of Lemma 5.2 is all that we will need for most of our applications. It asserts that under the hypotheses of the lemma, if the ball  $B(r)$  can be symplectically embedded into  $M$ , then

$$c_{U\sigma}(\phi, \gamma, H_1) \leq c_\sigma(\phi, \gamma, H_2) + \int_\gamma (H_2 - H_1) dt - r. \quad (5.4)$$

---

<sup>14</sup>This fact is related to the ECH Weyl law (1.3), and as explained in [19, §8] can be used to prove it for the standard contact structure on  $S^3$ .

*Proof of Lemma 5.2.* By the continuity in Corollary 4.4, we can assume without loss of generality, by slightly decreasing  $H_1$  and increasing  $H_2$  if necessary, that  $H_1 < H_2$  and that  $\phi_{H_1}$  and  $\phi_{H_2}$  are nondegenerate. We now proceed in three steps.

*Step 1.* Choose  $J_1$ ,  $J_2$ , and  $J$  as in Step 3 of the proof of Proposition 3.1. Also, fix  $k$  distinct points  $z_1, \dots, z_k \in M$ .

We claim that the composition

$$U^k \circ \Psi_{H_1, H_2} = \Psi_{H_1, H_2} \circ U^k : HP(\phi_{H_2}, \gamma_{H_2}, G) \longrightarrow HP(\phi_{H_1}, \gamma_{H_1}, G) \quad (5.5)$$

is induced by a (noncanonical) chain map

$$\psi : (CP(\phi_{H_2}, \gamma_{H_2}, G), \partial_{J_2}) \longrightarrow (CP(\phi_{H_1}, \gamma_{H_1}, G), \partial_{J_1})$$

with the following property: Similarly to (3.9), we can write  $\psi$  in the form

$$\psi \sum_{\alpha, Z} n_{\alpha, Z}(\alpha, Z) = \sum_{\beta, W} \left( \sum_{\alpha} \sum_{\substack{Z \in H_2(Y_{\phi, \alpha, \beta}) \\ I(\alpha, \beta, Z) = 2k}} n_{\alpha, W+Z} m_{\alpha, \beta, Z} \right) (\beta, W), \quad (5.6)$$

such that:

- (\*) If the coefficient  $m_{\alpha, \beta, Z} \neq 0$ , then there is a broken  $J$ -holomorphic current in  $\overline{M}$  from  $\alpha$  to  $\beta$  in the relative homology class  $Z$  passing through the points  $z_1, \dots, z_k$ .

The reason is that in the proof of the “holomorphic curves axiom” in [2], the chain map counts solutions to the Seiberg-Witten equations on  $\overline{W}$  perturbed using a large multiple of the symplectic form; and as the perturbation goes to infinity, the zero sets of the spinor components of a sequence of Seiberg-Witten solutions converge to a holomorphic current. Similarly, as in [37], the map (5.5) can be induced by a chain map counting Seiberg-Witten solutions where the spinor is constrained to vanish at the points  $z_1, \dots, z_k$ . As the perturbation goes to infinity, the zero sets of the spinor components of Seiberg-Witten solutions converge to a holomorphic current passing through the points  $z_1, \dots, z_k$ .

*Step 2.* Write  $X = \coprod_{i=1}^n B_i$  where  $B_i$  is symplectomorphic to  $B(r_i)$ . Let  $k_1, \dots, k_n$  be nonnegative integers with  $\sum_i k_i = k$  which maximize  $\sum_{i=1}^n c_{k_i}^{\text{ECH}}(B_i)$ , so that the latter sum agrees with  $c_k^{\text{ECH}}(X)$ . We claim that by making suitable choices of  $J$  and  $z_1, \dots, z_k$ , we can arrange for the chain map (5.6) to have the following property:

- (\*\*) If the coefficient  $m_{\alpha, \beta, Z} \neq 0$ , then the inequality (3.10) can be refined to

$$\int_W \omega_{\phi_{H_1}} \leq \int_{W+Z} \omega_{\phi_{H_2}} + \Delta - c_k^{\text{ECH}}(X).$$

To prove this, observe that for each  $i$ , the boundary of  $B_i$  is a contact type hypersurface in  $M$ ; this means that there exists a contact form  $\lambda_i$  on  $\partial B_i$  with  $d\lambda_i = (\omega_M)|_{\partial B_i}$ . We can perturb the balls  $B_i$  in  $M$  (to slightly larger domains, thus not decreasing their ECH capacities), to arrange that the contact forms  $\lambda_i$  are nondegenerate, and we will still denote the perturbed balls by  $B_i$ . We can then remove the  $B_i$  from  $\overline{M}$  and attach symplectization ends to form a new completed cobordism

$$\overline{M}' = \left( M \setminus \bigsqcup_i B_i \right) \bigcup \bigsqcup_i (-\infty, 0] \times \partial B_i. \quad (5.7)$$

Here  $(-\infty, 0] \times \partial B_i$  has the symplectization form  $d(e^s \lambda_i)$  where  $s$  denotes the  $(-\infty, 0]$  coordinate. For each  $i$ , choose a generic  $\lambda_i$ -compatible<sup>15</sup> almost complex structure  $J'_i$  on  $\mathbb{R} \times \partial B_i$  as needed to define the ECH of  $(\partial B_i, \lambda_i)$ . We can then choose the almost complex structure  $J$  on  $M$  so that it glues to the  $J'_i$  in (5.7), to give a well-defined almost complex structure  $J'$  on  $\overline{M}'$ . We can further choose a sequence of  $\omega_M$ -compatible almost structures  $J(n)$  on  $M$  such that  $J(n)$  agrees with  $J$  outside of the  $B_i$ , and moreover inside each  $B_i$ , the boundary has a neighborhood that can be identified with  $(-n, 0] \times \partial B_i$  so that  $J(n)$  agrees with  $J'_i$ .

By placing  $k_i$  of the points  $z_1, \dots, z_k$  inside  $B_i$  for each  $i$ , and using a compactness argument as in [20, Lem. 5.11], we can arrange the following: For any sequence of broken  $J(n)$ -holomorphic currents as in (\*), with an upper bound on  $\int_{W+Z} \omega_{\phi_{H_1}} - \int_W \omega_{\phi_{H_2}}$ , a subsequence of the holomorphic currents in  $\overline{M}$  converge on compact sets to a  $J'$ -holomorphic current in  $\overline{M}'$ , which on  $(-\infty, 0] \times \partial B_i$  is asymptotic to an ECH generator  $\alpha(i)$  for  $\lambda_i$  with ECH index at least  $2k_i$ . By the definition of the ECH capacities,  $\int_{\alpha(i)} \lambda_i \geq c_{k_i}^{\text{ECH}}(B_i)$ , so that  $\sum_{i=1}^n \int_{\alpha(i)} \lambda_i \geq c_k^{\text{ECH}}(X)$ . Repeating Step 4 of the proof of Proposition 3.1 then shows that we obtain (\*\*) if we take  $J = J(n)$  for  $n$  sufficiently large.

*Step 3.* If we make the choices as in Step 2, then for each  $L \in \mathbb{R}$ , the chain map (5.6) restricts to a chain map

$$\psi : (CP^L(\phi_{H_2}, \gamma_{H_2}, G), \partial_{J_2}) \longrightarrow (CP^{L+\Delta-c_k^{\text{ECH}}(X)}(\phi_{H_1}, \gamma_{H_1}, G), \partial_{J_1})$$

The induced map on homology fits into a commutative diagram

$$\begin{array}{ccc} HP^L(\phi_{H_2}, \gamma_{H_2}, G) & \xrightarrow{\iota^L} & HP(\phi_{H_2}, \gamma_{H_2}, G) \\ \downarrow & & \downarrow U^k \circ \Psi_{H_1, H_2} \\ HP^{L+\Delta-c_k^{\text{ECH}}(X)}(\phi_{H_1}, \gamma_{H_1}, G) & \xrightarrow{\iota^{L+\Delta-c_k^{\text{ECH}}(X)}} & HP(\phi_{H_1}, \gamma_{H_1}, G). \end{array}$$

<sup>15</sup>Here “ $\lambda_i$ -compatible” means that  $J'_i \partial_s = R_i$ , where  $s$  denotes the  $\mathbb{R}$  coordinate and  $R_i$  denotes the Reeb vector field associated to  $\lambda_i$ ; the almost complex structure  $J'_i$  sends the contact structure  $\text{Ker}(\lambda_i)$  to itself, rotating positively with respect to  $d\lambda_i$ ; and  $J'_i$  is invariant under the  $\mathbb{R}$  action on  $\mathbb{R} \times \partial B_i$ .

We are now done as in the proof of Proposition 4.3.  $\square$

**Remark 5.4.** With more work one should be able to generalize Lemma 5.2 to allow  $X$  to be a (possibly disconnected) compact Liouville domain. This would require setting up cobordism maps relating the PFH of  $\phi_{H_1}$  and  $\phi_{H_2}$  and the ECH of the boundary of the Liouville domain. The  $k^{\text{th}}$  ECH capacity of  $M$  is defined in [19] to be the supremum over the  $k^{\text{th}}$  ECH capacity of such  $X$ , and so it would follow that

$$c_{U^k\sigma}(\phi, \gamma, H_1) \leq c_\sigma(\phi, \gamma, H_2) + \int_\gamma (H_2 - H_1) dt - c_k^{\text{ECH}}(M).$$

## 6 From spectral gaps to periodic orbits

We now explain a mechanism for detecting the creation of periodic orbits. The following concept will be useful:

**Definition 6.1.** Let  $\phi$  be an area-preserving diffeomorphism of  $(\Sigma, \omega)$  and let  $d$  be an integer with  $d > g$ . Define the *minimal spectral gap*

$$\text{gap}_d(\phi) \in [0, \infty]$$

to be the infimum, over reference cycles  $\gamma$  for  $\phi$  with  $d(\gamma) = d$ , subgroups  $G \subset \text{Ker}([\omega_\phi])$ , and classes  $\sigma \in HP(\phi, \gamma, G)$  with  $U\sigma \neq 0$ , of  $c_\sigma(\phi, \gamma) - c_{U\sigma}(\phi, \gamma)$ .

We now have the following relation between spectral gaps and creation of periodic orbits.

**Proposition 6.2.** *Let  $\phi$  be an area-preserving diffeomorphism of  $(\Sigma, \omega)$ , and suppose that the Hamiltonian isotopy class  $[\phi]$  is rational. Let  $\mathcal{U} \subset \Sigma$  be a nonempty open set and let  $H$  be a  $(\mathcal{U}, a, l)$ -admissible Hamiltonian as in Definition 1.12. Let  $d$  be an integer with  $d > g$ , and suppose that*

$$\text{gap}_d(\phi) < a. \tag{6.1}$$

*Then for some  $\tau \in [0, l^{-1} \text{gap}_d(\phi)]$ , the map  $\phi_{\tau H}$  has a periodic orbit intersecting  $\mathcal{U}$  with period  $\leq d$ .*

*Proof.* We proceed in four steps.

*Step 1.* We first claim that for any class  $\sigma \in HP(\phi, \gamma, G)$  with  $U\sigma \neq 0$ , and for any  $\delta \geq 0$ , we have

$$c_{U\sigma}(\phi, \gamma) \leq c_\sigma(\phi, \gamma, \delta H) + \delta \int_\gamma H dt - \min(\delta l, a). \tag{6.2}$$

Here we are using the convention of Notation 5.1 on the right hand side.

To prove (6.2), recall from Definition 1.12 that there is a disk  $D \subset \mathcal{U}$  of area  $a$  and an interval  $I \subset (0, 1)$  of length  $l$  such that  $H \geq 1$  on  $I \times D$ . We can regard  $H$

as defined on  $Y_\phi$  as in §3. Let  $M_\delta$  denote the cobordism (3.6) between the mapping torus  $Y_\phi$  and the graph of  $\delta H$ . Then we can symplectically embed the polydisk  $P(a, \delta l)$ , namely the symplectic product of two-disks of areas  $a$  and  $\delta l$ , into  $M_\delta$ . Consequently, we can symplectically embed the ball  $B(\min(\delta l, a))$  into  $M_\delta$ . The inequality (6.2) now follows from the  $k = 1$  case of Lemma 5.2; see Remark 5.3.

*Step 2.* Suppose now that  $\delta \geq 0$  and

(\*) for all  $\tau \in [0, \delta]$ , the map  $\phi_{\tau H}$  has no periodic orbit intersecting  $\mathcal{U}$  with period  $\leq d$ .

We claim that if  $\gamma$  is a reference cycle for  $\phi$  with  $d(\gamma) = d$ , and if  $\sigma \in HP(\phi, \gamma, G)$  is a nonzero class, then

$$c_\sigma(\phi, \gamma, \delta H) = c_\sigma(\phi, \gamma) - \delta \int_\gamma H dt. \quad (6.3)$$

To prove (6.3), let  $S$  denote the set of actions of  $(G, \gamma)$ -anchored orbit sets for  $\phi$ . By the hypothesis (\*), if  $\tau \in [0, \delta]$ , then the set of actions of  $(G, \gamma)$ -anchored orbit sets for  $\phi_{\tau H}$  is  $S - \tau \int_\gamma H dt$ . Since we are assuming that  $[\phi]$  is rational, it follows from Proposition 4.5 that the function

$$\begin{aligned} [0, \delta] &\longrightarrow \mathbb{R}, \\ \tau &\longmapsto c_\sigma(\phi, \gamma, \tau H) + \tau \int_\gamma H dt \end{aligned} \quad (6.4)$$

takes values in the set  $S$ . This function is also continuous by Corollary 4.4. However the set  $S$  has measure zero as in [29, Lem. 2.2]. It follows that the function (6.4) is constant, and this proves (6.3).

*Step 3.* We now show that if  $\delta > l^{-1} \text{gap}_d(\phi)$ , then for some  $\tau \in [0, \delta]$ , the map  $\phi_{\tau H}$  has a periodic orbit intersecting  $\mathcal{U}$  with period  $\leq d$ . Suppose to get a contradiction that (\*) holds. Suppose that  $d(\gamma) = d$  and that  $\sigma \in HP(\phi, \gamma, G)$  satisfies  $U\sigma \neq 0$ . Then combining (6.2) and (6.3) gives

$$c_\sigma(\phi, \gamma) - c_{U\sigma}(\phi, \gamma) \geq \min(\delta l, a).$$

It then follows from Definition 6.1 that

$$\text{gap}_d(\phi) \geq \min(\delta l, a).$$

Since we assumed that  $\delta l > \text{gap}_d(\phi)$ , this contradicts the hypothesis (6.1).

*Step 4.* The proposition follows from Step 3 by replacing  $\mathcal{U}$  by an open set  $\mathcal{V}$  such that  $\overline{\mathcal{V}} \subset \mathcal{U}$  and  $H$  is supported in  $[0, 1] \times \mathcal{V}$ , and using a compactness argument.  $\square$

For example, we have the following corollary, which is a PFH analogue of [10, Lem. 3.1] for ECH:

**Corollary 6.3.** *Let  $\phi$  be an area-preserving diffeomorphism of  $(\Sigma, \omega)$ , and suppose that the Hamiltonian isotopy class  $[\phi]$  is rational. Let  $d$  be an integer with  $d > g$  and suppose that  $\text{gap}_d(\phi) = 0$ . Then every point in  $\Sigma$  is contained in a periodic orbit of  $\phi$  with period  $\leq d$ .*

*Proof.* It follows immediately from Proposition 6.2 that every nonempty open set  $\mathcal{U} \subset \Sigma$  contains a periodic point of period  $\leq d$ . It then follows from a compactness argument that every point in  $\Sigma$  is periodic with period  $\leq d$ .  $\square$

## 7 Proofs of theorems

We now prove all of our theorems stated in §1. We begin with the following simple observation:

**Lemma 7.1.** *Suppose that  $HP(\phi, \gamma, G)$  contains  $U$ -cyclic elements. Write  $d = d(\gamma)$  and  $A = \int_{\Sigma} \omega$ . Then*

$$\text{gap}_d(\phi) \leq \frac{A}{d - g + 1}.$$

*Proof.* We are given that equation (2.9) holds for some positive integer  $m$ . It follows using Proposition 4.2(a) that

$$\begin{aligned} mA &= c_{\sigma}(\phi, \gamma) - c_{U^{m(d-g+1)}\sigma}(\phi, \gamma) \\ &= \sum_{i=1}^{m(d-g+1)} (c_{U^{i-1}\sigma}(\phi, \gamma) - c_{U^i\sigma}(\phi, \gamma)). \end{aligned}$$

Since each of the summands on the right hand side is nonnegative, at least one of them must be less than or equal to  $A/(d - g + 1)$ .  $\square$

*Proof of Theorem 1.3.* Suppose that  $[\phi]$  is rational and satisfies the  $U$ -cycle property. Let  $\mathcal{U} \subset \Sigma$  be a nonempty open set. We need to show that there is a  $C^{\infty}$  small Hamiltonian perturbation, supported in  $\mathcal{U}$ , of  $\phi$  to a map having a periodic orbit intersecting  $\mathcal{U}$ .

Let  $H$  be a  $(\mathcal{U}, a, l)$ -admissible Hamiltonian. It is enough to show that for all  $\delta > 0$ , there exists  $\tau \in [0, \delta]$  such that  $\phi_{\tau H}$  has a periodic orbit intersecting  $\mathcal{U}$ .

Since  $[\phi]$  has the  $U$ -cycle property, it follows from Lemma 7.1 that

$$\liminf_{d \rightarrow \infty} \text{gap}_d(\phi) = 0.$$

Thus we can find  $d > g$  such that  $\text{gap}_d(\phi) < \min(a, l\delta)$ . For such  $d$ , since  $[\phi]$  is rational, Proposition 6.2 implies that for some  $\tau \in [0, \delta]$ , the map  $\phi_{\tau H}$  has a periodic orbit intersecting  $\mathcal{U}$  of period at most  $d$ .  $\square$

**Remark 7.2.** Although this might not be any more general, a similar argument shows that if  $[\phi]$  is rational and  $[\phi^n]$  has the  $U$ -cycle property for some positive integer  $n$ , then  $[\phi]$  satisfies the  $C^\infty$  closing property.

*Proof of Theorem 1.6.* This follows from Theorem 1.3 and Lemma 7.3 below.  $\square$

**Lemma 7.3.** *Let  $\phi$  be an area-preserving diffeomorphism of  $T^2$ , and suppose that the Hamiltonian isotopy class  $[\phi]$  is rational. Then  $[\phi]$  has the  $U$ -cycle property. In fact, if  $[\omega_\phi]$  is a positive multiple of the image of  $\Omega \in H^2(Y_\phi; \mathbb{Z})$ , and if  $\Gamma$  is a positive integer multiple of  $\text{PD}(\Omega)$ , then  $HP(\phi, \Gamma, \text{Ker}([\omega_\phi]))$  contains  $U$ -cyclic elements.*

*Proof.* Let  $\phi$  be an area-preserving diffeomorphism of  $(\Sigma, \omega)$ , of arbitrary genus for now, and assume that  $[\phi]$  is rational. Since the cohomology class  $[\omega_\phi] \in H^2(Y_\phi; \mathbb{R})$  is a real multiple of the image of an integral cohomology class  $\Omega \in H^2(Y_\phi; \mathbb{Z})$ , we can find classes  $\Gamma \in H_1(Y_\phi)$  with  $d(\Gamma)$  arbitrarily large such that the pair  $(\phi, \Gamma)$  is monotone as in Definition 2.22. (Simply take  $\Gamma$  to the Poincaré dual of  $n\Omega - c_1(E)/2$  where  $n$  is a large integer.)

For such a  $\Gamma$ , it is shown in Lee-Taubes [31, Cor. 1.5] that if  $g > 0$  and  $d(\Gamma) > 2g - 2$ , then the untwisted theory  $\overline{HP}(\phi, \Gamma)$  from §2.6 is isomorphic to an instance of the “bar” version of Seiberg-Witten Floer cohomology. The latter is an  $\mathbb{F}[U, U^{-1}]$  module, and the Lee-Taubes isomorphism intertwines the  $U$  maps (as in [37]; this is also implicit in [2]). We deduce that  $U$  is an isomorphism on  $\overline{HP}(\phi, \Gamma)$ . In particular,  $U^{d(\Gamma)-g+1}$  is a permutation of each graded piece  $\overline{HP}_i(\phi, \Gamma)$ .

If  $\Sigma = T^2$ , then for  $\Gamma$  as above, by computations in [30, §35] (see the remark after [31, Cor. 1.5]), we always have  $\overline{HP}(\phi, \Gamma) \neq 0$ , with the graded pieces  $\overline{HP}_i(\phi, \gamma)$  having rank  $\leq 2$ . Then there is a positive integer  $m$  such that  $U^{md(\Gamma)}$  is the identity on all such groups  $\overline{HP}(\phi, \Gamma)$ . It now follows from Lemma 2.23 that  $HP(\phi, \Gamma, \text{Ker}([\omega_\phi]))$  contains  $U$ -cyclic elements.  $\square$

To prove Theorems 1.13 and 1.13, we first prove a more general statement:

**Theorem 7.4.** *Let  $\phi$  be an area-preserving diffeomorphism of  $(\Sigma, \omega)$  such that the Hamiltonian isotopy class  $[\phi]$  is rational. Suppose that there is a positive integer  $d_0$  such that  $\phi$  has  $U$ -cyclic elements of degree  $d$  whenever  $d$  is a positive multiple of  $d_0$  with  $d > g$ . Let  $\mathcal{U} \subset \Sigma$  be a nonempty open set and write  $A = \int_\Sigma \omega$ . Let  $H$  be a  $(\mathcal{U}, a, l)$ -admissible Hamiltonian. If  $\delta l \leq a$ , then for some  $\tau \in [0, \delta]$ , the map  $\phi_{\tau H}$  has a periodic orbit intersecting  $\mathcal{U}$  with period at most  $d_0 k$ , where*

$$k = \left\lfloor \frac{A\delta^{-1}l^{-1} + g - 1}{d_0} \right\rfloor + 1. \quad (7.1)$$

*Proof.* Write  $d = kd_0$ . It follows from (7.1) that

$$d > A\delta^{-1}l^{-1} + g - 1. \quad (7.2)$$

In particular, it follows from (7.2) that  $d > g$ , since  $\delta l \leq a < A$ . Then by Lemma 7.1, we have

$$\text{gap}_d(\phi) \leq \frac{A}{d - g + 1}.$$

Also, it follows from the above inequalities that  $\text{gap}_d(\phi) < \delta l \leq a$ . It then follows from Proposition 6.2 that for some  $\tau \in [0, l^{-1} \text{gap}_d(\phi)] \subset [0, \delta]$ , the map  $\phi_{\tau H}$  has a periodic orbit intersecting  $\mathcal{U}$  with period at most  $d$ .  $\square$

*Proof of Theorem 1.13.* If  $\Sigma = S^2$ , then by Example 2.19,  $\phi$  has  $U$ -cyclic elements of degree  $d$  for all positive integers  $d$ . Thus Theorem 7.4 applies with  $d_0 = 1$  to give the result.  $\square$

*Proof of Theorem 1.15.* This follows from Theorem 7.4 and Lemma 7.3.  $\square$

## 8 Asymptotics of PFH spectral invariants

To conclude, we now prove the following ‘‘Weyl law’’ for PFH spectral invariants.

**Theorem 8.1.** *Let  $\phi$  be a (possibly degenerate) area-preserving diffeomorphism of  $(\Sigma, \omega)$ . Let  $\{G_i\}_{i \geq 1}$  be a sequence of subgroups of  $\text{Ker}([\omega_\phi])$ , let  $\{\gamma_i\}_{i \geq 1}$  be a sequence of reference cycles for  $\phi$ , and for each  $i \geq 1$ , let  $\sigma_i \in HP(\phi, \gamma_i, G_i)$  be a nonzero class. Assume that:*

- $\lim_{i \rightarrow \infty} d(\gamma_i) = \infty$ .
- *There is a positive integer  $m$  such that each class  $\sigma_i$  is  $U$ -cyclic of order  $\leq m$ .*

Let  $H_1, H_2 : Y_\phi \rightarrow \mathbb{R}$ . Write  $A = \int_\Sigma \omega$ . Then

$$\lim_{i \rightarrow \infty} \frac{c_{\sigma_i}(\phi, \gamma_i, H_2) - c_{\sigma_i}(\phi, \gamma_i, H_1) + \int_{\gamma_i} (H_2 - H_1) dt}{d(\gamma_i)} = A^{-1} \int_{Y_\phi} (H_2 - H_1) \omega_\phi \wedge dt.$$

**Example 8.2.** Let  $D$  be a disk with a symplectic form  $\omega$  of area 1, and let  $\phi$  be the time 1 map of a Hamiltonian  $H : [0, 1] \times D \rightarrow \mathbb{R}$  which vanishes on  $[0, 1] \times \{x\}$  when  $x$  is near  $\partial D$ . Then  $\phi$  defines an area-preserving diffeomorphism of  $S^2$ , with a symplectic form of area 1, which is the identity on an open set. Recall from Example 2.16 that if  $\gamma = d[S^1] \times \{x\}$ , where  $x$  corresponds to a point on  $\partial D$ , then  $HP(\phi, \gamma, \{0\})$  is the free  $\Lambda$ -module generated by classes  $e_{d,0}, \dots, e_{d,d}$ ; and each of these classes is  $U$ -cyclic of order 1. Note that if  $\phi$  is the identity, then each spectral invariant  $c_{e_{d,i}}(\phi, \gamma) = 0$ . It then follows from Theorem 8.1 that in general, we have

$$\lim_{d \rightarrow \infty} \frac{c_{e_{d,0}}(\phi, d[S^1] \times \{x\})}{d} = \int_{[0,1] \times D} H \omega \wedge dt. \quad (8.1)$$



Here the right hand side (up to constant factors depending on conventions) is the *Calabi invariant* of  $\phi$ ; see e.g. [13].

The special case of (8.1) where  $\phi$  is a “monotone twist” was proved<sup>16</sup> by direct calculation in [5, Thm. 5.1], and this result plays a key role in the proof of the simplicity conjecture. It is also noted in [5, §7.4] that (8.1) implies that the Calabi invariant extends to a homeomorphism defined on the group of compactly supported “homeomorphisms” of the disk. The latter statement was subsequently proved using different methods in [4, Thm. 1.4].

*Proof of Theorem 8.1.* We use a “ball packing” argument similar to [9, §3.2].

Suppose first that  $H_1 < H_2$ . Let  $X$  be a finite disjoint union of balls symplectically embedded in  $M$  and write  $V = \text{vol}(X)$ .

We know that each  $\sigma_i$  is  $U$ -cyclic of order  $m_i$  where  $m_i \leq m$ . By Proposition 4.2(a), if  $k_i = m_i n_i (d(\gamma_i) - g + 1)$  where  $n_i$  is an integer, then we have

$$c_{U^{k_i} \sigma_i}(\phi, \gamma_i, H_1) = c_{\sigma_i}(\phi, \gamma_i, H_1) - A m_i n_i.$$

Then Lemma 5.2 gives

$$c_{\sigma_i}(\phi, \gamma_i, H_2) - c_{\sigma_i}(\phi, \gamma_i, H_1) + \int_{\gamma_i} (H_2 - H_1) dt \geq c_{k_i}^{\text{ECH}}(X) - A m_i n_i. \quad (8.2)$$

Now choose

$$n_i = \left\lfloor \frac{d(\gamma_i)^2 V}{m_i A^2 (d(\gamma_i) - g + 1)} \right\rfloor.$$

Then it follows from (8.2) that

$$\begin{aligned} \liminf_{i \rightarrow \infty} \frac{c_{\sigma_i}(\phi, \gamma_i, H_2) - c_{\sigma_i}(\phi, \gamma_i, H_1) + \int_{\gamma_i} (H_2 - H_1) dt}{d(\gamma_i)} &\geq \liminf_{i \rightarrow \infty} \frac{c_{k_i}^{\text{ECH}}(X) - A m_i n_i}{d(\gamma_i)} \\ &= A^{-1} V \end{aligned}$$

Here in the second line we have used (5.3) and the hypothesis that  $d(\gamma_i) \rightarrow \infty$ .

Now we can choose  $X$  to make  $V$  arbitrarily close to

$$\begin{aligned} \text{vol}(M, \omega_M) &= \frac{1}{2} \int_M \omega_M \wedge \omega_M \\ &= \int_{Y_\phi} (H_2 - H_1) \omega_\phi \wedge dt. \end{aligned}$$

Thus we obtain

$$\liminf_{i \rightarrow \infty} \frac{c_{\sigma_i}(\phi, \gamma_i, H_2) - c_{\sigma_i}(\phi, \gamma_i, H_1) + \int_{\gamma_i} (H_2 - H_1) dt}{d(\gamma_i)} \geq A^{-1} \int_{Y_\phi} (H_2 - H_1) \omega_\phi \wedge dt.$$

---

<sup>16</sup>The paper [5] writes a slightly different, but equivalent, version of (8.1). That paper defines spectral invariants using the variant  $\widetilde{HP}(\phi, \gamma, \{0\})$  from §2.6, which is possible here since monotonicity holds. Our spectral invariant  $c_{e_{d,i}}(\phi, d[S^1] \times \{x\})$  agrees with the spectral invariant denoted in [5] by  $c_{d,2i-d}(\phi)$ .

Now observe that both sides of the above inequality change by the same amount if one adds a constant to  $H_1$  or  $H_2$ . Thus the above inequality is true for any  $H_1$  and  $H_2$ , without the hypothesis that  $H_1 < H_2$ . In particular, the above inequality is true with  $H_1$  and  $H_2$  switched, which gives

$$\limsup_{i \rightarrow \infty} \frac{c_{\sigma_i}(\phi, \gamma_i, H_2) - c_{\sigma_i}(\phi, \gamma_i, H_1) + \int_{\gamma_i} (H_2 - H_1) dt}{d(\gamma_i)} \leq A^{-1} \int_{Y_\phi} (H_2 - H_1) \omega_\phi \wedge dt.$$

The above two inequalities imply the theorem.  $\square$

**Remark 8.3.** By choosing the ball packings carefully, as in the proof of [22, Thm. 1.1], one can show that the rate of convergence in Theorem 8.1 is  $O(d(\gamma_i)^{-1/2})$ .

## References

- [1] M. Asaoka and K. Irie, *A  $C^\infty$  closing lemma for Hamiltonian diffeomorphisms of closed surfaces*, GAFA **26** (2016), 1245–1254.
- [2] G. Chen, *On cobordism maps on periodic Floer homology*, Alg. Geom. Topol. **21** (2021), 1–103.
- [3] A. Cotton-Clay, *Symplectic Floer homology of area-preserving surface diffeomorphisms*, Geom. Topol. **13** (2009), 2619–2674.
- [4] D. Cristofaro-Gardiner, V. Humilière, C. Y. Mak, S. Seyfaddini, and I. Smith, *Quantitative Heegaard Floer homology and the Calabi invariant*, arXiv:2105.11026.
- [5] D. Cristofaro-Gardiner, V. Humilière, and S. Seyfaddini, *Proof of the simplicity conjecture*, arXiv:2001.01792.
- [6] D. Cristofaro-Gardiner, V. Humilière, and S. Seyfaddini, *PFH spectral invariants on the two-sphere and the large scale geometry of Hofer’s metric*, arXiv:2102.04404.
- [7] D. Cristofaro-Gardiner and M. Hutchings, *From one Reeb orbit to two*, J. Diff. Geom. **102** (2016), 25–36.
- [8] D. Cristofaro-Gardiner, M. Hutchings, and D. Pomerleano, *Torsion contact forms in three dimensions have two or infinitely many Reeb orbits*, Geom. Topol. **23** (2019), 3601–3645.
- [9] D. Cristofaro-Gardiner, M. Hutchings, and V. Ramos, *The asymptotics of ECH capacities*, Invent. Math. **199** (2015), 187–214.

- [10] D. Cristofaro-Gardiner and M. Mazzucchelli, *The action spectrum characterizes closed contact manifolds all of whose Reeb orbits are closed*, arXiv:1901.10617.
- [11] D. Cristofaro-Gardiner, R. Prasad, and B. Zhang, *The smooth closing lemma for area-preserving surface diffeomorphisms*, arXiv:2110.02925.
- [12] D. Farris, *The embedded contact homology of nontrivial circle bundles over Riemann surfaces*, UC Berkeley PhD thesis, 2011.
- [13] J-M. Gambaudo and E. Ghys, *Enlacements asymptotiques*, Topology **36** (1997), 1355–1379.
- [14] C. Gerig, *Taming the pseudoholomorphic beasts in  $\mathbb{R} \times (S^1 \times S^2)$* , Geom. Topol. **24** (2020), 1791–1839.
- [15] M. Herman, *Sur la conjugaison différentiable des difféomorphismes du cercle à des rotations*. Pub. Math. de l’IHÉS, Tome **49** (1970), 5–233.
- [16] V. Humilière, *Un lemme de fermeture  $C^\infty$  [d’après Asaoka et Irie]*, Séminaire Bourbaki, nov 2019, Astérisque **430** (2021).
- [17] M. Hutchings, *An index inequality for embedded pseudoholomorphic curves in symplectizations*, J. Eur. Math. Soc. **4** (2002), 313–361.
- [18] M. Hutchings, *The embedded contact homology index revisited*, in New Perspectives and Challenges in Symplectic Field Theory, CRM Proc. Lecture Notes, vol. 49 (AMS, 2009), pp. 263–297.
- [19] M. Hutchings, *Quantitative embedded contact homology*, J. Diff. Geom. **88** (2011), 231–266.
- [20] M. Hutchings, *Lecture notes on embedded contact homology*, Contact and symplectic topology, 389–484, Bolyai Soc. Math. Stud. bf 26, Springer, 2014.
- [21] M. Hutchings, *Mean action and the Calabi invariant*, J. Mod. Dyn. **10** (2016), 511–539.
- [22] M. Hutchings, *ECH capacities and the Ruelle invariant*, arXiv:1910.08260, to appear in Journal of Fixed Point Theory and Applications.
- [23] M. Hutchings and M. Sullivan, *The periodic Floer homology of a Dehn twist*, Alg. Geom. Topol. **5** (2005), 301–354.
- [24] M. Hutchings and M. Sullivan, *Rounding corners of polygons and the embedded contact homology of  $T^3$* , Geom. Topol. **10** (2006), 169–266.
- [25] M. Hutchings and C. H. Taubes, *Gluing pseudoholomorphic curves along branched covered cylinders I*, J. Sympl. Geom. **5** (2007), 43–137.

- [26] M. Hutchings and C. H. Taubes, *Gluing pseudoholomorphic curves along branched covered cylinders II*, J. Sympl. Geom. **7** (2009), 29–133.
- [27] M. Hutchings and C. H. Taubes, *The Weinstein conjecture for stable Hamiltonian structures*, Geom. Topol. **13** (2009), 901–941.
- [28] M. Hutchings and C. H. Taubes, *Proof of the Arnold chord conjecture in three dimensions II*, Geom. Topol. **17** (2013), 2601–2688.
- [29] K. Irie, *Dense existence of periodic orbits and ECH spectral invariants*, J. Mod. Dyn. **9** (2015), 357–363.
- [30] P. B. Kronheimer and T. S. Mrowka, *Monopoles and three-manifolds*, Cambridge University Press, 2008.
- [31] Y.-J. Lee and C. H. Taubes, *Periodic Floer homology and Seiberg-Witten Floer cohomology*, J. Sympl. Geom. **10** (2012), 81–164.
- [32] J. Nelson and M. Weiler, *Embedded contact homology of prequantization bundles*, arXiv:2007.13883.
- [33] J. Rooney, *Cobordism maps in embedded contact homology*, arXiv:1912.01048.
- [34] R. Siefring, *Relative asymptotic behavior of pseudoholomorphic half-cylinders*, Comm. Pure Appl. Math. **61** (2008), 1631–1684.
- [35] W. Sun, *An estimate on energy of min-max Seiberg-Witten Floer generators*, arXiv:1801.02301.
- [36] C. H. Taubes, *Embedded contact homology and Seiberg-Witten Floer cohomology I*, Geom. Topol. **14** (2010), 2497–2581.
- [37] C. H. Taubes, *Embedded contact homology and Seiberg-Witten Floer cohomology V*, Geom. Topol. **14** (2010), 2961–3000.
- [38] M. Usher, *Spectral numbers in Floer theories*, Compos. Math. **144** (2008), 1581–1592.