

# $\imath$ HALL ALGEBRAS OF WEIGHTED PROJECTIVE LINES AND QUANTUM SYMMETRIC PAIRS

MING LU AND SHIQUAN RUAN

ABSTRACT. The  $\imath$ Hall algebra of a weighted projective line is defined to be the semi-derived Ringel-Hall algebra of the category of 1-periodic complexes of coherent sheaves on the weighted projective line over a finite field. We show that this Hall algebra provides a realization of the  $\imath$ quantum loop algebra, which is a generalization of the  $\imath$ quantum group arising from the quantum symmetric pair of split affine type ADE in its Drinfeld type presentation. The  $\imath$ Hall algebra of the  $\imath$ quiver algebra of split affine type A was known earlier to realize the same algebra in its Serre presentation. We then establish a derived equivalence which induces an isomorphism of these two  $\imath$ Hall algebras, explaining the isomorphism of the  $\imath$ quantum group of split affine type A under the two presentations.

## CONTENTS

1. Introduction	1
2. A Drinfeld type presentation of $\imath$ quantum groups	7
3. $\imath$ Hall algebras	11
4. Weighted projective lines and coherent sheaves	16
5. $\imath$ Hall algebras of cyclic quivers	21
6. $\imath$ Hall algebras and $\imath$ quantum loop algebras	31
7. Relations in tubes and commutative relations	34
8. Relations between star point and $[i, 1]$ , I	38
9. Relations between star point and $[i, 1]$ , II	42
10. Derived equivalence and two presentations of affine type A	45
Appendix A. Proofs of results in Section 8	48
Appendix B. Proofs of results in Section 9	56
References	69

## 1. INTRODUCTION

### 1.1. Backgrounds.

1.1.1. *Quantum groups and Hall algebras.* Ringel [Rin90] in 1990s used the Hall algebra associated to a Dynkin quiver  $Q$  over a finite field  $\mathbb{F}_q$  to realize half a quantum group  $\mathbf{U}^+ = \mathbf{U}_v^+(\mathfrak{g})$ ; see Green [Gr95] for an extension to arbitrary acyclic quivers. Later, inspired by Ringel's construction, Lusztig [Lus90] gave a geometric realization of  $\mathbf{U}^+$ , and additionally

---

2020 *Mathematics Subject Classification.* Primary 17B37, 14A22, 16E60, 18G80.

*Key words and phrases.* Quantum symmetric pairs, Affine  $\imath$ quantum groups, Drinfeld type presentations, Hall algebras, Weighted projective lines.

constructed its canonical basis [Lus90]. These constructions can be regarded as the earliest examples of categorifications of half parts of quantum groups.

The Hall algebra realization of the whole quantum group was achieved by Bridgeland [Br13] who in fact provided a Hall algebra construction of the Drinfeld double  $\tilde{\mathbf{U}}$  of a quantum group  $\mathbf{U}$ . As a further generalization of Bridgeland's construction, the first author and Peng [LP21] formulated the *semi-derived Ringel-Hall algebras* for arbitrary hereditary abelian categories via 2-periodic complexes (also called  $\mathbb{Z}_2$ -graded complexes); compare with the semi-derived Hall algebras defined by Gorsky for *Frobenius categories* [Gor18]. The construction of semi-derived Ringel-Hall algebras was further extended to 1-Gorenstein algebras in [LW22a, Appendix A].

There has been a current realization of the affine quantum groups formulated by Drinfeld [Dr88, Be94, Da15], which plays a crucial role on (algebraic and geometric) representation theory. The Hall algebra of the projective line was studied in a visionary paper by Kapranov [Ka97] and then extended by Baumann-Kassel [BKa01] to realize the current half of quantum affine  $\mathfrak{sl}_2$ . The Hall algebra of a weighted projective line was developed in [Sch04] to realize half an affine quantum group of ADE type, which was then upgraded to the whole quantum group via Drinfeld double techniques [DJX12, BS13] (in fact, they considered more general simply-laced quantum loop algebras of star-shaped graphs). The semi-derived Ringel-Hall algebras formulated in [LP21] are isomorphic to the Drinfeld double Ringel-Hall algebras, and so a geometric realization of quantum loop algebras in Drinfeld presentation was achieved in some sense based on [DJX12] by considering the categories of 2-periodic complexes over weighted projective lines; see [LP21, Remark 4.16].

*1.1.2.  $\imath$ Quantum groups and  $\imath$ Hall algebras.* The  $\imath$ quantum groups  $\mathbf{U}^\imath$  arising from quantum symmetric pairs  $(\mathbf{U}, \mathbf{U}^\imath)$  associated to Satake diagrams [Let99] (see [Ko14]) can be viewed as a vast generalization of Drinfeld-Jimbo quantum groups associated to Dynkin diagrams; see the survey [W22] and references therein.

The universal  $\imath$ quantum group  $\tilde{\mathbf{U}}^\imath = \langle B_i, \mathbb{K}_i^{\pm 1} \mid i \in \mathbb{I} \rangle$  is by definition a subalgebra of  $\tilde{\mathbf{U}}$ , and  $(\tilde{\mathbf{U}}, \tilde{\mathbf{U}}^\imath)$  forms a quantum symmetric pair; cf. [LW22a]. A central reduction of  $\tilde{\mathbf{U}}^\imath$  reproduces the  $\imath$ quantum group  $\mathbf{U}^\imath = \mathbf{U}_\zeta^\imath$  (with parameters  $\zeta$ ) which arises from the construction of quantum symmetric pairs  $(\mathbf{U}, \mathbf{U}^\imath)$  by Letzter [Let99]. Recently, Lu and Wang [LW21b] established a Drinfeld type presentation for the  $\imath$ quantum groups arising from quantum symmetric pairs of split affine ADE type.

In recent years, the first author and Wang [LW22a, LW23] have developed  $\imath$ Hall algebras of  $\imath$ quivers to realize the (universal) quasi-split  $\imath$ quantum groups of Kac-Moody type. The  $\imath$ Hall algebras are constructed in the framework of semi-derived Ringel-Hall algebras of 1-Gorenstein algebras ([LW22a, Appendix A], [LW23]). In particular, Bridgeland's Hall algebra realization of quantum groups is reformulated as  $\imath$ Hall algebras of  $\imath$ quivers of diagonal type; see [LW22a].

Together with Wang, we [LRW23] used the  $\imath$ Hall algebra of the projective line to give a geometric realization of the  $q$ -Onsager algebra (i.e.,  $\imath$ quantum group of split affine  $\mathfrak{sl}_2$ ) in its Drinfeld type presentation, a generalization of the geometric realization of the current half of quantum affine  $\mathfrak{sl}_2$  via the Hall algebra of the projective line [Ka97, BKa01].

We view Letzter's  $\imath$ quantum groups and our universal  $\imath$ quantum groups as a vast generalization of the Drinfeld-Jimbo quantum groups. The  $\imath$ -program as outlined by Bao and Wang

[BW18a] aims at generalizing various fundamental constructions from quantum groups to  $\imath$ quantum groups; in addition to the works mentioned above, see [BW18a, BK19, BW18b] for generalizations of (quasi) R-matrix and canonical bases.

**1.2. Goal.** This is a sequel of [LRW23] in our program devoted to developing a geometric realization of the  $\imath$ quantum groups in Drinfeld type presentations, a vast generalization of [Ka97, BKa01, Sch04, DJX12, BS13]. Explicitly, we use the  $\imath$ Hall algebras of weighted projective lines to give a geometric realization of the so-called simply-laced  $\imath$ quantum loop algebra of a star-shaped graph in Drinfeld type presentation (a generalization of  $\imath$ quantum groups arising from quantum symmetric pairs of split affine ADE type considered in [LW21b]). We further show that the isomorphism of  $\imath$ quantum groups of split affine type A in two (Serre vs Drinfeld) presentations is induced from a derived equivalence of the categories underlying the two (quivers vs weighted projective lines)  $\imath$ Hall algebra realizations.

The construction in this paper should be viewed as a geometric counterpart of the  $\imath$ Hall algebra construction in [LW22a, LW23]. It is our hope that this work will stimulate further connection between the theory of weighted projective lines and the theory of quantum symmetric pairs.

### 1.3. Main results.

**1.3.1.** Let  $\Gamma := T_{p_1, \dots, p_t}$  be a star-shaped graph with its vertex set  $\mathbb{I}_0$  as in (2.27),  $\mathfrak{g}$  be its associated Lie algebra,  ${}^{\text{Dr}}\widetilde{\mathbf{U}}^\imath$  be its associated  $\imath$ quantum loop algebra [LW21b] defined as in Definition 2.4. Note that if  $\Gamma$  is of ADE type, then  ${}^{\text{Dr}}\widetilde{\mathbf{U}}^\imath$  is the Drinfeld type presentation of the  $\imath$ quantum group arising from quantum symmetric pairs of split affine ADE type; see Theorem 2.6. Recall that  ${}^{\text{Dr}}\widetilde{\mathbf{U}}^\imath$  is generated by  $B_{\star, l}, B_{[i, j], l}, H_{\star, m}, H_{[i, j], m}, \Theta_{\star, m}, \Theta_{[i, j], m}$  and central elements  $\mathbb{K}_{\star}^{\pm 1}, \mathbb{K}_{[i, j]}^{\pm 1}, C^{\pm 1}$  for  $l \in \mathbb{Z}, m > 0$ , and  $1 \leq i \leq t, 1 \leq j \leq p_i - 1$ , subject to (2.12)–(2.20).

Let  $\mathbb{X}_{\mathbf{k}}$  be the weighted projective line over a finite field  $\mathbf{k} = \mathbb{F}_q$  associated to  $\Gamma$ . Let  $\mathbf{v} = \sqrt{q}$ . The  $\imath$ Hall algebra  ${}^\imath\widetilde{\mathcal{H}}(\mathbb{X}_{\mathbf{k}})$  considered in this paper is the twisted semi-derived Ringel-Hall algebra of the category of 1-periodic complexes over  $\text{coh}(\mathbb{X}_{\mathbf{k}})$ ; cf. [LP21, LW22a].

**Theorem A** (Theorem 6.2). There exists a  $\mathbb{Q}(\mathbf{v})$ -algebra homomorphism

$$\Omega : {}^{\text{Dr}}\widetilde{\mathbf{U}}^\imath_{|\mathbf{v}=\mathbf{v}} \longrightarrow {}^\imath\widetilde{\mathcal{H}}(\mathbb{X}_{\mathbf{k}}), \quad (1.1)$$

which sends

$$\mathbb{K}_{\star} \mapsto [K_{\mathcal{O}}], \quad \mathbb{K}_{[i, j]} \mapsto [K_{S_{ij}}], \quad C \mapsto [K_{\delta}]; \quad (1.2)$$

$$B_{\star, l} \mapsto \frac{-1}{q-1}[\mathcal{O}(l\bar{c})], \quad \Theta_{\star, r} \mapsto \widehat{\Theta}_{\star, r}, \quad H_{\star, r} \mapsto \widehat{H}_{\star, r}; \quad (1.3)$$

$$B_{[i, j], l} \mapsto \frac{-1}{q-1}\widehat{B}_{[i, j], l}, \quad \Theta_{[i, j], r} \mapsto \widehat{\Theta}_{[i, j], r}, \quad H_{[i, j], r} \mapsto \widehat{H}_{[i, j], r} \quad (1.4)$$

for any  $[i, j] \in \mathbb{I}_0 - \{\star\}, l \in \mathbb{Z}, r > 0$ .

Let us explain the counterparts of the Drinfeld type generators in  ${}^\imath\widetilde{\mathcal{H}}(\mathbb{X}_{\mathbf{k}})$ . First,  $\widehat{\Theta}_{\star, r}$  and  $\widehat{H}_{\star, r}$  are defined by using the corresponding ones  $\widehat{\Theta}_r, \widehat{H}_r$  defined in  ${}^\imath\widetilde{\mathcal{H}}(\mathbb{P}_{\mathbf{k}}^1)$  [LRW23] via the

natural embedding (see (6.5))

$$F_{\mathbb{X}, \mathbb{P}^1} : {}^i\tilde{\mathcal{H}}(\mathbb{P}_{\mathbf{k}}^1) \longrightarrow {}^i\tilde{\mathcal{H}}(\mathbb{X}_{\mathbf{k}}).$$

Second, for  $1 \leq i \leq \mathbf{t}$ , there is an algebra embedding

$$\tilde{\psi}_{C_{p_i}} : \tilde{\mathbf{U}}_{\mathbf{v}}^i(\widehat{\mathfrak{sl}}_{p_i}) \longrightarrow {}^i\tilde{\mathcal{H}}(\mathbf{k}C_{p_i})$$

by [LW23, Theorem 9.6], where  $C_{p_i}$  is the cyclic quiver with  $p_i$  vertices. As a consequence, each branch of  $\Gamma$  corresponds to a subalgebra of  ${}^i\tilde{\mathcal{H}}(\mathbb{X}_{\mathbf{k}})$  isomorphic to  ${}^{\text{Dr}}\tilde{\mathbf{U}}_{\mathbf{v}}^i(\widehat{\mathfrak{sl}}_{p_i})$  by the composition  $\Omega_{C_{p_i}}$  of  $\tilde{\psi}_{C_{p_i}}$  and the isomorphism of two presentations [LW21b] (see (5.3))

$$\Phi : {}^{\text{Dr}}\tilde{\mathbf{U}}_{\mathbf{v}}^i(\widehat{\mathfrak{sl}}_{p_i}) \xrightarrow{\cong} \tilde{\mathbf{U}}_{\mathbf{v}}^i(\widehat{\mathfrak{sl}}_{p_i}).$$

On the other hand, the subcategory  $\mathcal{T}_{\lambda_i}$  of  $\text{coh}(\mathbb{X}_{\mathbf{k}})$ , consisting of torsion sheaves supported at the distinguished point  $\lambda_i$ , is isomorphic to  $\text{rep}_{\mathbf{k}}^{\text{nil}}(C_{p_i})$ , the category of finite-dimensional nilpotent representations of  $C_{p_i}$  over  $\mathbf{k}$ . This isomorphism induces an algebra embedding (see (6.1))

$$\iota_i : {}^i\tilde{\mathcal{H}}(\mathbf{k}C_{p_i}) \longrightarrow {}^i\tilde{\mathcal{H}}(\mathbb{X}_{\mathbf{k}}).$$

So we define  $\widehat{B}_{[i,j],\ell}$ ,  $\widehat{\Theta}_{[i,j],r}$ ,  $\widehat{H}_{[i,j],r}$  to be the images of the Drinfeld generators of  ${}^{\text{Dr}}\tilde{\mathbf{U}}_{\mathbf{v}}^i(\widehat{\mathfrak{sl}}_{p_i})$  under the composition  $\iota_i \circ \Omega_{C_{p_i}}$ ; see (6.2). To show that  $\Omega : {}^{\text{Dr}}\tilde{\mathbf{U}}_{\mathbf{v}}^i \rightarrow {}^i\tilde{\mathcal{H}}(\mathbb{X}_{\mathbf{k}})$  is a homomorphism, we must verify the Drinfeld type relations (2.12)–(2.20) for the  $i$ quantum group  ${}^{\text{Dr}}\tilde{\mathbf{U}}^i$ .

The counterparts in  ${}^i\tilde{\mathcal{H}}(\mathbb{X}_{\mathbf{k}})$  of the relations at the star point  $\star$  follow from [LRW23, Theorem A], and the counterparts in  ${}^i\tilde{\mathcal{H}}(\mathbb{X}_{\mathbf{k}})$  of the relations involving all the vertices  $[i, j]$  follow from the above definitions and the fact any two torsion sheaves supported at distinct points have zero Hom and  $\text{Ext}^1$ -spaces.

In order to check the remaining Drinfeld type relations in  ${}^i\tilde{\mathcal{H}}(\mathbb{X}_{\mathbf{k}})$ , the key part is to verify the relations between  $\star$  and  $[i, j]$  for  $1 \leq i \leq \mathbf{t}$ , especially the ones between  $\star$  and  $[i, 1]$ , which consists of plenty of highly non-trivial computations in §7.3–§9. For these relations we have to determine some of the Drinfeld generators in  ${}^i\tilde{\mathcal{H}}(\mathbb{X}_{\mathbf{k}})$  clearly, especially the ones at  $[i, 1]$ ; see §9.1.

For each  $r > 0$  and  $1 \leq i \leq \mathbf{t}$ ,  $\mathcal{M}_{r\delta \pm \alpha_{i1}}$  is defined in (9.4), and  $\mathcal{M}_{i,r\delta}$  is defined in (9.6).

**Theorem B** (Proposition 9.1, Proposition 9.2). For any  $r > 0$  and  $1 \leq i \leq \mathbf{t}$ , we have

$$\widehat{B}_{[i,1],r} = (q-1) \sum_{[M] \in \mathcal{M}_{r\delta + \alpha_{i1}}} \mathbf{n}(\ell(M) - 1)[[M]]; \quad (1.5)$$

$$\widehat{B}_{[i,1],-r} = (1-q) \sum_{[M] \in \mathcal{M}_{r\delta - \alpha_{i1}}} \mathbf{n}(\ell(M) - 1)[[M]] * [K_{-r\delta + \alpha_{i1}}]; \quad (1.6)$$

$$\widehat{\Theta}_{[i,1],r} = \frac{\mathbf{v}}{q-1} \sum_{|\lambda|=r} \mathbf{n}(\ell(\lambda)) [[S_{i,0}^{(\lambda)}]] + \mathbf{v}^{-1} \sum_{[M] \in \mathcal{M}_{i,r\delta}} \mathbf{n}(\ell(M) - 1)[[M]]. \quad (1.7)$$

It is very helpful and interesting to describe all the root vectors at any  $[i, j]$  in the  $i$ Hall algebra  ${}^i\tilde{\mathcal{H}}(\mathbb{X}_{\mathbf{k}})$ . In fact, all the root vectors can be described in  $i$ Hall algebra with the quiver being suitably oriented, and then in the  $i$ Hall algebra of the cyclic quiver  $C_n$  by using Fourier transformations; see Proposition 5.5.

1.3.2. Let  $\mathbb{X}$  be a weighted projective line of domestic type, and  $Q$  be the affine quiver of the associated type. Then there exists a tilting object in  $\text{coh}(\mathbb{X}_{\mathbf{k}})$  which gives a derived equivalence (see [GL87])

$$\mathcal{D}^b(\text{coh}(\mathbb{X}_{\mathbf{k}})) \xrightarrow{\simeq} \mathcal{D}^b(\text{mod}(\mathbf{k}Q^{op})).$$

In the case  $\mathbb{X}_{\mathbf{k}}$  being of domestic type A, we have  $\mathbf{k}Q \cong \mathbf{k}C_{p_1, p_2}$ ; see (10.3). As a special case of the main result in [LW23], there is a realization of the universal  $\imath$ quantum group  $\tilde{\mathbf{U}}^{\imath}$  (in its Serre type presentation) via the  $\imath$ Hall algebra  ${}^{\imath}\tilde{\mathcal{H}}(\mathbf{k}C_{p_1, p_2}^{op})$  of the quiver  $C_{p_1, p_2}$ , that is, we have an injective homomorphism (see Theorem 3.8)

$$\tilde{\psi} : \tilde{\mathbf{U}}^{\imath}_{|v=\mathbf{v}} \longrightarrow {}^{\imath}\tilde{\mathcal{H}}(\mathbf{k}C_{p_1, p_2}^{op}).$$

The aforementioned derived equivalence induces an isomorphism of  $\imath$ Hall algebras

$$F : {}^{\imath}\tilde{\mathcal{H}}(\mathbb{X}_{\mathbf{k}}) \xrightarrow{\cong} {}^{\imath}\tilde{\mathcal{H}}(\mathbf{k}C_{p_1, p_2}^{op}),$$

providing a realization of the algebra isomorphism [LW21b]

$$\Phi : \text{Dr} \tilde{\mathbf{U}}^{\imath} \xrightarrow{\cong} \tilde{\mathbf{U}}^{\imath}$$

of the  $\imath$ quantum group of split affine type A in two (Serre and Drinfeld type) presentations. We summarize this result as follows.

**Theorem C** (Proposition 10.3, Theorem 10.4). We have the following commutative diagram

$$\begin{array}{ccc} \text{Dr} \tilde{\mathbf{U}}^{\imath}_{|v=\mathbf{v}} & \xrightarrow{\Phi} & \tilde{\mathbf{U}}^{\imath}_{|v=\mathbf{v}} \\ \downarrow \Omega & & \downarrow \tilde{\psi} \\ {}^{\imath}\tilde{\mathcal{H}}(\mathbb{X}_{\mathbf{k}}) & \xrightarrow{F} & {}^{\imath}\tilde{\mathcal{H}}(\mathbf{k}C_{(p_1, p_2)}^{op}). \end{array}$$

In particular, the homomorphism  $\Omega$  is injective.

This result generalizes the corresponding result in [LRW23] for affine  $\mathfrak{sl}_2$ . For  $\mathfrak{g}$  of type DE and  $Q$  the associated affine quiver, we also have a derived equivalence  $\mathcal{D}^b(\text{coh}(\mathbb{X}_{\mathbf{k}})) \simeq \mathcal{D}^b(\mathbf{k}Q^{op})$ , which induces an algebra isomorphism  $F : {}^{\imath}\tilde{\mathcal{H}}(\mathbb{X}_{\mathbf{k}}) \xrightarrow{\cong} {}^{\imath}\tilde{\mathcal{H}}(\mathbf{k}Q^{op})$ . However, the morphism  $\Phi$  can not be the one induced by  $F$ , in fact it can not be induced by any derived equivalent functors by the same argument as in [DJX12, §9.2].

**1.4. Comparison with previous works.** The  $\imath$ Hall algebras used in this paper to give a realization of the whole  $\imath$ quantum loop algebras are categorical and intrinsic [LP21, LW22a], while the Drinfeld double Ringel-Hall algebras to give a realization of the whole quantum loop algebras are constructed from two copies of Ringel-Hall algebras by algebraic technique, and have much unnatural flexibility, such as Grothendieck groups, Hopf pairings, etc. For weighted projective lines, their Ringel-Hall algebras are only *topological bialgebras*, and we need to compute coproducts in completed spaces to obtain products of elements, even though their Drinfeld double Ringel-Hall algebras could be defined without considering completed spaces (it is also not easy to prove this fact); see [BS12a, Cr10].

Compared with the geometric construction of quantum loop algebras in [Ka97, BKa01, Sch04, DJX12, BS13], for  $\imath$ quantum groups, our approach of proving the Drinfeld type relations in  ${}^{\imath}\tilde{\mathcal{H}}(\mathbb{X}_{\mathbf{k}})$  is completely different. In fact, we can not use the same method as in

[DJX12] to define the corresponding root vectors in Hall algebras. In addition, the relations (2.14) (equivalently, (2.22)), (2.17) and (2.18) contain terms involving  $\mathbb{K}_\delta$  which do not arise in the computations *loc. cit.* of similar identities in the (Drinfeld double) Hall algebra of the (weighted) projective line; some new homological computations are needed to determine these  $\mathbb{K}_\delta$  terms. In particular, the complexity of (2.14) makes our computations much more challenging than [Sch04, DJX12].

The root vectors of  $\imath$ quantum loop algebras at  $[i, 1]$  are also described more explicitly. In fact, the counterpart of  $\widehat{B}_{[i,1],l}$  (see (9.5)) for quantum loop algebras is only known when  $l \geq 0$  (see [Sch04, §6.5]), while the counterpart of  $\widehat{\Theta}_{[i,1],r}$  is not known. To our knowledge, it is the first attempt to describe the root vectors of quantum and  $\imath$ quantum loop algebras by using the technique of Fourier transformations as in Proposition 5.5.

The corresponding result of Theorem C for the whole quantum group is not known except for the special case affine  $\mathfrak{sl}_2$ ; see [BS12b]. In fact, for affine  $\mathfrak{sl}_2$ , the Drinfeld-Beck isomorphism  $\Phi$  can be interpreted by Beilinson's derived equivalence [BS12b].

**1.5. Further works.** This work opens up further research directions. In a sequel of this paper, we shall describe the composition algebra of the  $\imath$ Hall algebra of a weighted projective line, prove that the morphism  $\Omega : {}^{\text{Dr}}\widetilde{\mathcal{U}}^\imath \rightarrow {}^\imath\widetilde{\mathcal{H}}(\mathbb{X}_{\mathbf{k}})$  is injective if  $\mathfrak{g}$  is of ADE type, and give a PBW basis for the  $\imath$ quantum group via coherent sheaves. For a cyclic quiver  $C_n$ , its  $\imath$ Hall algebra can be used to realize the  $\imath$ quantum groups of split affine  $\mathfrak{sl}_n$  and  $\mathfrak{gl}_n$ .

The  $\imath$ quantum loop algebra defined in Definition 2.4 is a generalization of the Drinfeld type presentation of  $\imath$ quantum group of split affine ADE type. It will be interesting to study its algebraic structures. In particular, for  $\mathfrak{g}$  of affine type  $D_4, E_6, E_7, E_8$ , this  $\imath$ quantum loop algebra is the  $\imath$ quantum group of elliptic (or 2-toroidal) algebras of types  $D_4^{(1,1)}, E_6^{(1,1)}, E_7^{(1,1)}$  and  $E_8^{(1,1)}$  (in this case,  $\mathbb{X}_{\mathbf{k}}$  is of tubular type), whose algebraic structure and geometric realizations should be studied more deeply. We expect the morphism  $\Omega : {}^{\text{Dr}}\widetilde{\mathcal{U}}^\imath_{|v=\mathbf{v}} \rightarrow {}^\imath\widetilde{\mathcal{H}}(\mathbb{X}_{\mathbf{k}})$  to be injective for arbitrary Kac-Moody algebra  $\mathfrak{g}$ . It will also be interesting to study the  $\imath$ Hall algebras of higher genus curves, in particular, of elliptic curves, which shall be studied deeply in the future.

A Drinfeld type presentation of  $\imath$ quantum groups arising from quasi-split quantum symmetric pairs of affine ADE type is in preparation. We shall develop connections between  $\imath$ Hall algebras of weighted projective lines and the  $\imath$ quantum groups of quasi-split affine ADE type in Drinfeld type presentation, where a new kind of categories (different from the categories of 1-periodic complexes) over weighted projective lines shall be introduced.

As quantum groups are  $\imath$ quantum groups arising from quantum symmetric pairs of diagonal type (which can be viewed as  $\imath$ quantum groups of quasi-split type), we shall reformulate the geometric realization [DJX12] (see also [Sch04, BS13]) of quantum groups in Drinfeld presentation via the twisted semi-derived Hall algebras of weighted projective lines, which consolidates [LP21, Remark 4.16] and strengthens the results of [Ka97, BKa01, Sch04, DJX12, BS12b, BS13].

**1.6. Organization.** The paper is organized as follows. In Section 2, we review the materials on affine  $\imath$ quantum groups of split ADE type,  $\imath$ quantum loop algebras and their Drinfeld type presentations. The  $\imath$ Hall algebra on the categories of 1-periodic complexes and the  $\imath$ quiver

algebras of split type are summarized in Section 3. In Section 4, we recalled weighted projective lines and their categories of coherent sheaves. The canonical algebras, Grothendieck groups and Euler forms are also reviewed.

In Section 5, we studied the  $\iota$ Hall algebra of a cyclic quiver. The map  $\Omega : \mathrm{Dr}\widetilde{\mathbf{U}}_{|v=v}^i \rightarrow {}^i\widetilde{\mathcal{H}}(\mathbb{X}_{\mathbf{k}})$  in Theorem A is defined in Section 6, and the proof of  $\Omega$  being an algebra homomorphism occupies Sections 7–9. In particular, in Subsection 9.1, we describe the root vectors at  $[i, 1]$  in  ${}^i\widetilde{\mathcal{H}}(\mathbb{X}_{\mathbf{k}})$  as stated in Theorem B.

A derived equivalence leading to the isomorphism of  $\iota$ Hall algebras is established and then Theorem C is proved in Section 10.

In order to make the paper readable and the idea of proofs clear, we put a dozen proofs in Appendix.

**1.7. Acknowledgments.** ML deeply thanks Weiqiang Wang for guiding him to study the  $\iota$ quantum groups, and also his continuing encouragement. ML thanks University of Virginia for hospitality and support. We thank Liangang Peng and Jie Xiao for helpful discussions on Ringel-Hall algebras of weighted projective lines. We thank the anonymous referee for very helpful suggestions and comments.

ML is partially supported by the National Natural Science Foundation of China (No. 12171333). SR is partially supported by the National Natural Science Foundation of China (No. 11801473) and the Fundamental Research Funds for Central Universities of China (No. 20720210006).

## 2. A DRINFELD TYPE PRESENTATION OF $\iota$ QUANTUM GROUPS

In this section, we collect materials on (affine) Lie algebras,  $\iota$ quantum groups and their Drinfeld presentations.

**2.1. The loop algebras and affine Lie algebras.** Let  $\mathbb{I}_0$  be an index set. Let  $A = (a_{ij})_{i,j \in \mathbb{I}_0}$  be an irreducible generalized Cartan matrix (GCM) of a Kac-Moody algebra  $\mathfrak{g}$ . Let  $\mathcal{R}_0$  be the set of roots for  $\mathfrak{g}$ , and fix a set  $\mathcal{R}_0^+$  of positive roots with simple roots  $\alpha_i$  ( $i \in \mathbb{I}_0$ ). Let  $L\mathfrak{g}$  be the loop algebra of  $\mathfrak{g}$ , and  $\widehat{\mathfrak{g}} := \mathfrak{g}[t, t^{-1}] \oplus \mathbb{C}c$ . If  $\mathfrak{g}$  is a complex simple Lie algebra, then  $L\mathfrak{g} \cong \widehat{\mathfrak{g}}$ , which is an affine Kac-Moody algebra [Ga80]. In general  $L\mathfrak{g}$  and  $\widehat{\mathfrak{g}}$  are not Kac-Moody algebras, and there is a surjective (not injective) homomorphism  $L\mathfrak{g} \rightarrow \widehat{\mathfrak{g}}$ .

Let  $\widehat{\mathfrak{g}}$  be the (untwisted) affine Kac-Moody algebra with affine Cartan matrix denoted by  $(a_{ij})_{i,j \in \mathbb{I}}$ , where  $\mathbb{I} = \{0\} \cup \mathbb{I}_0$  with the affine node 0. Let  $\alpha_i$  ( $i \in \mathbb{I}$ ) be the simple roots of  $\widehat{\mathfrak{g}}$ , and  $\alpha_0 = \delta - \theta$ , where  $\delta$  denotes the basic imaginary root, and  $\theta$  is the highest root of  $\mathfrak{g}$ . The root system  $\mathcal{R}$  for  $\widehat{\mathfrak{g}}$  is defined to be

$$\mathcal{R} = \{\pm(\beta + k\delta) \mid \beta \in \mathcal{R}_0^+, k \in \mathbb{Z}\} \cup \{m\delta \mid m \in \mathbb{Z} \setminus \{0\}\}. \quad (2.1)$$

For  $\gamma = \sum_{i \in \mathbb{I}} n_i \alpha_i \in \mathbb{N}\mathbb{I}$ , the height  $\mathrm{ht}(\gamma)$  is defined as  $\mathrm{ht}(\gamma) = \sum_{i \in \mathbb{I}} n_i$ .

Let  $P$  and  $Q$  denote the weight and root lattices of  $\mathfrak{g}$ , respectively. Let  $\omega_i \in P$  ( $i \in \mathbb{I}_0$ ) be the fundamental weights of  $\mathfrak{g}$ . Note  $\alpha_i = \sum_{j \in \mathbb{I}_0} a_{ij} \omega_j$ . We define a bilinear pairing  $\langle \cdot, \cdot \rangle : P \times Q \rightarrow \mathbb{Z}$  such that  $\langle \omega_i, \alpha_j \rangle = \delta_{i,j}$ , for  $i, j \in \mathbb{I}_0$ , and thus  $\langle \alpha_i, \alpha_j \rangle = a_{ij}$ .

The Weyl group  $W_0$  of  $\mathfrak{g}$  is generated by the simple reflection  $s_i$ , for  $i \in \mathbb{I}_0$ . It acts on  $P$  by  $s_i(x) = x - \langle x, \alpha_i \rangle \alpha_i$  for  $x \in P$ . The extended affine Weyl group  $\widetilde{W}$  is the semi-direct

product  $W_0 \times P$ , which contains the affine Weyl group  $W := W_0 \times Q = \langle s_i \mid i \in \mathbb{I} \rangle$  as a subgroup; we denote

$$t_\omega = (1, \omega) \in \widetilde{W}, \quad \text{for } \omega \in P.$$

We identify  $P/Q$  with a finite group  $\Omega$  of Dynkin diagram automorphisms, and so  $\widetilde{W} = \Omega \cdot W$ . For  $i \in \mathbb{I}_0$ , as in [Be94], we define

$$\omega'_i := t_{\omega_i} s_i \in \widetilde{W}. \quad (2.2)$$

**2.2. Affine  $\imath$ quantum groups of split type ADE.** For  $n \in \mathbb{Z}, r \in \mathbb{N}$ , denote by

$$[n] = \frac{v^n - v^{-n}}{v - v^{-1}}, \quad \begin{bmatrix} n \\ r \end{bmatrix} = \frac{[n][n-1] \cdots [n-r+1]}{[r]}.$$

For  $A, B$  in a  $\mathbb{Q}(v)$ -algebra, we shall denote  $[A, B]_{v^a} = AB - v^a BA$ , and  $[A, B] = AB - BA$ . We also use the convenient notation throughout the paper

$$[A, B, C]_{v^a} = [[A, B]_{v^a}, C]_{v^a}. \quad (2.3)$$

Recall the Cartan matrix  $A = (a_{ij})_{i,j \in \mathbb{I}}$  of affine type ADE, for  $\mathbb{I} = \mathbb{I}_0 \cup \{0\}$  with the affine node 0. The notion of (quasi-split) universal  $\imath$ quantum groups  $\widetilde{\mathbf{U}}^\imath$  was formulated in [LW22a].

The *universal affine  $\imath$ quantum group of split type ADE* is the  $\mathbb{Q}(v)$ -algebra  $\widetilde{\mathbf{U}}^\imath = \widetilde{\mathbf{U}}^\imath(\widehat{\mathfrak{g}})$  with generators  $B_i, \mathbb{K}_i^{\pm 1}$  ( $i \in \mathbb{I}$ ), subject to the following relations, for  $i, j \in \mathbb{I}$ :

$$\mathbb{K}_i \mathbb{K}_i^{-1} = \mathbb{K}_i^{-1} \mathbb{K}_i = 1, \quad \mathbb{K}_i \text{ is central}, \quad (2.4)$$

$$B_i B_j - B_j B_i = 0, \quad \text{if } a_{ij} = 0, \quad (2.5)$$

$$B_i^2 B_j - [2] B_i B_j B_i + B_j B_i^2 = -v^{-1} B_j \mathbb{K}_i, \quad \text{if } a_{ij} = -1, \quad (2.6)$$

$$\sum_{r=0}^3 (-1)^r \begin{bmatrix} 3 \\ r \end{bmatrix} B_i^{3-r} B_j B_i^r = -v^{-1} [2]^2 (B_i B_j - B_j B_i) \mathbb{K}_i, \quad \text{if } a_{ij} = -2. \quad (2.7)$$

For  $\mu = \mu' + \mu'' \in \mathbb{Z}\mathbb{I} := \bigoplus_{i \in \mathbb{I}} \mathbb{Z}\alpha_i$ , define  $\mathbb{K}_\mu$  such that

$$\mathbb{K}_{\alpha_i} = \mathbb{K}_i, \quad \mathbb{K}_{-\alpha_i} = \mathbb{K}_i^{-1}, \quad \mathbb{K}_\mu = \mathbb{K}_{\mu'} \mathbb{K}_{\mu''}, \quad \mathbb{K}_\delta = \mathbb{K}_0 \mathbb{K}_\theta. \quad (2.8)$$

*Remark 2.1.* The generator  $\mathbb{K}_i$  here, which corresponds to the acyclic complex  $K_{S_i}$  in the  $\imath$ Hall algebra, is related to the generator  $\tilde{k}_i$  used in [LW22a, LW23] (which is natural from the viewpoint of Drinfeld doubles) by a rescaling:  $\mathbb{K}_i = -v^2 \tilde{k}_i$ . The precise relation between the algebra  $\widetilde{\mathbf{U}}^\imath$  and the  $\imath$ quantum group  $\mathbf{U}^\imath$  arising from quantum symmetric pairs [Ko14] is explained *loc. cit.*

*Remark 2.2.* The  $\mathbb{Q}(v)$ -algebra  $\widetilde{\mathbf{U}}^\imath$  is  $\mathbb{Z}\mathbb{I}$ -graded by letting

$$\text{wt}(B_i) = \alpha_i, \quad \text{wt}(\mathbb{K}_i) = 2\alpha_i, \quad \text{for } i \in \mathbb{I}. \quad (2.9)$$

A variant of  $\widetilde{\mathbf{U}}^\imath$ , in which  $\mathbb{K}_i$  is not assumed to be invertible, is  $\mathbb{N}\mathbb{I}$ -graded by (2.9).



**Lemma 2.3** ([LW22b]; also cf. [KP11, BK20]). *For  $i \in \mathbb{I}$ , there exists an automorphism  $\mathbf{T}_i$  of the  $\mathbb{Q}(v)$ -algebra  $\widetilde{\mathbf{U}}^v$  such that  $\mathbf{T}_i(\mathbb{K}_\mu) = \mathbb{K}_{s_i\mu}$  for  $\mu \in \mathbb{Z}\mathbb{I}$ , and*

$$\mathbf{T}_i(B_j) = \begin{cases} \mathbb{K}_i^{-1}B_i, & \text{if } j = i, \\ B_j, & \text{if } a_{ij} = 0, \\ B_jB_i - vB_iB_j, & \text{if } a_{ij} = -1, \\ [2]^{-1}(B_jB_i^2 - v[2]B_iB_jB_i + v^2B_i^2B_j) + B_j\mathbb{K}_i, & \text{if } a_{ij} = -2, \end{cases}$$

for  $j \in \mathbb{I}$ . Moreover,  $\mathbf{T}_i$  ( $i \in \mathbb{I}$ ) satisfy the braid group relations, i.e.,  $\mathbf{T}_i\mathbf{T}_j = \mathbf{T}_j\mathbf{T}_i$  if  $a_{ij} = 0$ , and  $\mathbf{T}_i\mathbf{T}_j\mathbf{T}_i = \mathbf{T}_j\mathbf{T}_i\mathbf{T}_j$  if  $a_{ij} = -1$ .

**2.3. A Drinfeld type presentation.** Let  $A = (a_{ij})_{i,j \in \mathbb{I}_0}$  be a generalized Cartan matrix (GCM) of a simply-laced Kac-Moody algebra  $\mathfrak{g}$ . The  $i$ quantum loop algebra  ${}^{\text{Dr}}\widetilde{\mathbf{U}}^v = {}^{\text{Dr}}\widetilde{\mathbf{U}}^v(L\mathfrak{g})$  of split type is the  $\mathbb{Q}(v)$ -algebra generated by  $\mathbb{K}_i^{\pm 1}$ ,  $C^{\pm 1}$ ,  $H_{i,m}$ ,  $\Theta_{i,l}$  and  $B_{i,l}$ , where  $i \in \mathbb{I}_0$ ,  $m \in \mathbb{Z}_+$ ,  $l \in \mathbb{Z}$ , subject to some relations. In order to give the explicit definition of  ${}^{\text{Dr}}\widetilde{\mathbf{U}}^v$ , we introduce some shorthand notations below.

Let  $k_1, k_2, l \in \mathbb{Z}$  and  $i, j \in \mathbb{I}_0$ . Set

$$\begin{aligned} S(k_1, k_2 \mid l; i, j) &= B_{i,k_1}B_{i,k_2}B_{j,l} - [2]B_{i,k_1}B_{j,l}B_{i,k_2} + B_{j,l}B_{i,k_1}B_{i,k_2}, \\ \mathbb{S}(k_1, k_2 \mid l; i, j) &= S(k_1, k_2 \mid l; i, j) + \{k_1 \leftrightarrow k_2\}. \end{aligned} \quad (2.10)$$

Here and below,  $\{k_1 \leftrightarrow k_2\}$  stands for repeating the previous summand with  $k_1, k_2$  switched if  $k_1 \neq k_2$ , so the sums over  $k_1, k_2$  are symmetric. We also denote

$$\begin{aligned} R(k_1, k_2 \mid l; i, j) &= \mathbb{K}_i C^{k_1} \left( - \sum_{p \geq 0} v^{2p} [2][\Theta_{i,k_2-k_1-2p-1}, B_{j,l-1}]_{v^{-2}} C^{p+1} \right. \\ &\quad \left. - \sum_{p \geq 1} v^{2p-1} [2][B_{j,l}, \Theta_{i,k_2-k_1-2p}]_{v^{-2}} C^p - [B_{j,l}, \Theta_{i,k_2-k_1}]_{v^{-2}} \right), \end{aligned} \quad (2.11)$$

$$\mathbb{R}(k_1, k_2 \mid l; i, j) = R(k_1, k_2 \mid l; i, j) + \{k_1 \leftrightarrow k_2\}.$$

Sometimes, it is convenient to rewrite part of the summands in (2.11) as

$$\begin{aligned} & - \sum_{p \geq 1} v^{2p-1} [2][B_{j,l}, \Theta_{i,k_2-k_1-2p}]_{v^{-2}} C^p - [B_{j,l}, \Theta_{i,k_2-k_1}]_{v^{-2}} \\ & = - \sum_{p \geq 0} v^{2p-1} [2][B_{j,l}, \Theta_{i,k_2-k_1-2p}]_{v^{-2}} C^p + v^{-2} [B_{j,l}, \Theta_{i,k_2-k_1}]_{v^{-2}}. \end{aligned}$$

**Definition 2.4** ( $i$ quantum loop algebras, [LW21b]). *Let  $A = (a_{ij})_{i,j \in \mathbb{I}_0}$  be a generalized Cartan matrix (GCM) of a simply-laced Kac-Moody algebra  $\mathfrak{g}$ . The  $i$ quantum loop algebra  ${}^{\text{Dr}}\widetilde{\mathbf{U}}^v$  of split type is the  $\mathbb{Q}(v)$ -algebra generated by  $\mathbb{K}_i^{\pm 1}$ ,  $C^{\pm 1}$ ,  $H_{i,m}$  and  $B_{i,l}$ , where  $i \in \mathbb{I}_0$ ,  $m > 0$ ,  $l \in \mathbb{Z}$ , subject to the following relations (for  $m, n > 0$  and  $k, l \in \mathbb{Z}$ ):*

$$\mathbb{K}_i, C \text{ are central, } \quad \mathbb{K}_i\mathbb{K}_i^{-1} = 1, \quad CC^{-1} = 1, \quad (2.12)$$

$$[H_{i,m}, H_{j,n}] = 0, \quad (2.13)$$

$$[H_{i,m}, B_{j,l}] = \frac{[ma_{ij}]}{m} B_{j,l+m} - \frac{[ma_{ij}]}{m} B_{j,l-m} C^m, \quad (2.14)$$

$$[B_{i,k}, B_{j,l}] = 0, \quad \text{if } a_{ij} = 0, \quad (2.15)$$

$$[B_{i,k}, B_{j,l+1}]_{v^{-a_{ij}}} - v^{-a_{ij}} [B_{i,k+1}, B_{j,l}]_{v^{a_{ij}}} = 0, \text{ if } i \neq j, \quad (2.16)$$

$$[B_{i,k}, B_{i,l+1}]_{v^{-2}} - v^{-2} [B_{i,k+1}, B_{i,l}]_{v^2} \quad (2.17)$$

$$= v^{-2} \Theta_{i,l-k+1} C^k \mathbb{K}_i - v^{-4} \Theta_{i,l-k-1} C^{k+1} \mathbb{K}_i + v^{-2} \Theta_{i,k-l+1} C^l \mathbb{K}_i - v^{-4} \Theta_{i,k-l-1} C^{l+1} \mathbb{K}_i, \quad (2.18)$$

$$\mathbb{S}(k_1, k_2 \mid l; i, j) = \mathbb{R}(k_1, k_2 \mid l; i, j), \text{ if } a_{ij} = -1.$$

Here we set

$$\Theta_{i,0} = (v - v^{-1})^{-1}, \quad \Theta_{i,m} = 0, \text{ for } m < 0; \quad (2.19)$$

and  $\Theta_{i,m}$  ( $m \geq 1$ ) are related to  $H_{i,m}$  by the following equation:

$$1 + \sum_{m \geq 1} (v - v^{-1}) \Theta_{i,m} u^m = \exp \left( (v - v^{-1}) \sum_{m \geq 1} H_{i,m} u^m \right). \quad (2.20)$$

Let us mention that in spite of its appearance, the RHS of (2.17) typically has two non-zero terms, thanks to the convention (2.19).

**Lemma 2.5** ([LW21b]).

(1) The relation (2.13) is equivalent to

$$[\Theta_{i,m}, \Theta_{j,n}] = 0 \quad (m, n \geq 1). \quad (2.21)$$

(2) The relation (2.14) is equivalent to

$$\begin{aligned} & [\Theta_{i,m}, B_{j,l}] + [\Theta_{i,m-2}, B_{j,l}] C \\ &= v^{a_{ij}} [\Theta_{i,m-1}, B_{j,l+1}]_{v^{-2a_{ij}}} + v^{-a_{ij}} [\Theta_{i,m-1}, B_{j,l-1}]_{v^{2a_{ij}}} C \quad (m \geq 1, l \in \mathbb{Z}). \end{aligned} \quad (2.22)$$

Let  $A$  be a GCM of an affine Lie algebra  $\mathfrak{g}$  of type ADE. Define a sign function

$$o(\cdot) : \mathbb{I}_0 \longrightarrow \{\pm 1\} \quad (2.23)$$

such that  $o(i)o(j) = -1$  whenever  $a_{ij} < 0$  (there are clearly and exactly 2 such functions). We define the following elements in  $\tilde{\mathbf{U}}^v$ , for  $i \in \mathbb{I}_0$ ,  $k \in \mathbb{Z}$  and  $m \geq 1$ :

$$B_{i,k} = o(i)^k \mathbf{T}_{\omega_i}^{-k}(B_i), \quad (2.24)$$

$$\acute{\Theta}_{i,m} = o(i)^m \left( -B_{i,m-1} \mathbf{T}_{\omega_i}'(B_i) + v^2 \mathbf{T}_{\omega_i}'(B_i) B_{i,m-1} + (v^2 - 1) \sum_{p=0}^{m-2} B_{i,p} B_{i,m-p-2} \mathbb{K}_i^{-1} \mathbb{K}_\delta \right),$$

$$\Theta_{i,m} = \acute{\Theta}_{i,m} - \sum_{a=1}^{\lfloor \frac{m-1}{2} \rfloor} (v^2 - 1) v^{-2a} \acute{\Theta}_{i,m-2a} \mathbb{K}_{a\delta} - \delta_{m,ev} v^{1-m} \mathbb{K}_{\frac{m}{2}\delta}. \quad (2.25)$$


and  $H_{i,m}$  is defined via (2.20). Here,  $\delta_{m,ev} = 1$  if  $m$  is even and  $\delta_{m,ev} = 0$  otherwise. In particular,  $B_{i,0} = B_i$ .

**Theorem 2.6** ([LW21b]). *If  $A = (a_{ij})_{i,j \in \mathbb{I}_0}$  is a GCM of a Lie algebra  $\mathfrak{g}$  of type ADE, then there is a  $\mathbb{Q}(v)$ -algebra isomorphism  $\Phi : {}^{Dr} \tilde{\mathbf{U}}^v(L\mathfrak{g}) \rightarrow \tilde{\mathbf{U}}^v(\hat{\mathfrak{g}})$ , which sends*

$$B_{i,k} \mapsto B_{i,k}, \quad H_{i,m} \mapsto H_{i,m}, \quad \Theta_{i,m} \mapsto \Theta_{i,m}, \quad \mathbb{K}_i \mapsto \mathbb{K}_i, \quad C \mapsto \mathbb{K}_\delta, \quad (2.26)$$

for  $i \in \mathbb{I}_0$ ,  $k \in \mathbb{Z}$ , and  $m \geq 1$ .

**2.4. Star-shaped graph.** For  $\mathbf{p} = (p_1, \dots, p_t)$  with  $p_i \geq 2$ ,  $t \geq 1$ , let us consider the following star-shaped graph  $\Gamma = \mathbb{T}_{p_1, \dots, p_t}$ :



The set of vertices is denoted by  $\mathbb{I}_0$ . As marked in the graph, the central vertex is denoted by  $\star$ . Let  $J_1, \dots, J_t$  be the branches, which are subdiagrams of type  $A_{p_1-1}, \dots, A_{p_t-1}$  respectively. Denote by  $[i, j]$  the  $j$ -th vertex in the  $i$ -th branch. These examples include all finite-type Dynkin diagrams as well as the affine Dynkin diagrams of types  $D_4^{(1)}, E_6^{(1)}, E_7^{(1)}$ , and  $E_8^{(1)}$ .

Let  $\mathfrak{g}$  be the Kac-Moody algebra corresponding to  $\Gamma$ , and  $L\mathfrak{g}$  be its loop algebra. Then we have the quantum loop algebra  ${}^{\text{Dr}}\widetilde{\mathbf{U}}^i$ . In view of the graph  $\Gamma$ , for the root system  $\mathcal{R}$  of  $L\mathfrak{g}$ , its simple roots are denoted by  $\alpha_\star$  and  $\alpha_{ij}$ , for  $1 \leq i \leq t$  and  $1 \leq j \leq p_i - 1$ .

**Lemma 2.7.** *For any  $a \in \mathbb{Z}$ ,  ${}^{\text{Dr}}\widetilde{\mathbf{U}}^i$  is generated by  $B_{\star,a}, B_{\star,a+1}, B_{i,0}, \mathbb{K}_i^{\pm 1}$  and  $C^{\pm 1}$  for all  $i \in \mathbb{I}_0 - \{\star\}$ .*

*Proof.* Let  $U$  be the subalgebra of  ${}^{\text{Dr}}\widetilde{\mathbf{U}}^i$  generated by  $B_{\star,a}, B_{\star,a+1}, B_{i,0}, \mathbb{K}_i^{\pm 1}$  and  $C^{\pm 1}$  for  $i \in \mathbb{I}_0 - \{\star\}$ . First, we have  $\Theta_{\star,1} \in U$  by taking  $i = \star, k = a = l$  in (2.17) since  $\Theta_{\star,m} = 0$  for  $m < 0$ . From (2.14) by taking  $i = j = \star$ , since  $H_{\star,1} = \Theta_{\star,1}$ , we have  $B_{\star,a-1} \in U$  by taking  $m = 1$  and  $l = a$ ; and  $B_{\star,a+2} \in U$  by taking  $m = 1$  and  $l = a + 1$ . Inductively, we have  $B_{\star,l} \in U$  for any  $l \in \mathbb{Z}$ .

From (2.17), we have  $\Theta_{\star,2} \in U$  by taking  $k = 0, l = 1$  since  $\Theta_{\star,0} = \frac{1}{v-v^{-1}}$ . Inductively, one can prove that  $\Theta_{\star,m} \in U$  for any  $m \geq 1$ . It follows from (2.20) that  $H_{\star,m} \in U$  for any  $m \geq 1$ .

In order to prove that  $U = {}^{\text{Dr}}\widetilde{\mathbf{U}}^i$ , it is enough to prove that  $\Theta_{j,m}, B_{j,l} \in U$  for any  $m \geq 1, l \in \mathbb{Z}$ , provided  $\Theta_{i,m}, B_{i,l} \in U$  and  $a_{ij} = -1$ .

Assume  $\Theta_{i,m}, B_{i,l} \in U$  for any  $m \geq 1, l \in \mathbb{Z}$  and  $a_{ij} = -1$ . Considering  $\mathbb{S}(k_1, k_1 + 1 | 0; i, j)$ , it follows from (2.18) that  $B_{j,-1} \in U$ . Together with (2.14), we have  $B_{j,1} \in U$ ; using (2.17) by taking  $k = 0 = l$ , we have  $\Theta_{j,1} \in U$ . We prove  $B_{j,\pm m}$  and  $\Theta_{j,m}$  are in  $U$  by induction on  $m \geq 0$ . By considering  $\mathbb{S}(k_1, k_1 + 1 | m; i, j)$ , it follows from (2.18) that  $B_{j,-m-1} \in U$ ; by (2.14), we have  $B_{j,m+1} \in U$ ; by (2.17), we have  $\Theta_{j,m+1} \in U$ .  $\square$

### 3. iHALL ALGEBRAS

For an additive category  $\mathcal{E}$  and  $M \in \mathcal{E}$ , we denote

- $\triangleright$   $\text{add } M$  – subcategory of  $\mathcal{E}$  whose objects are the direct summands of finite direct sums of copies of  $M$ ,
- $\triangleright$   $\text{Iso}(\mathcal{E})$  – set of the isoclasses of objects in  $\mathcal{E}$ ,
- $\triangleright$   $K_0(\mathcal{E})$  – Grothendieck group of  $\mathcal{E}$  if  $\mathcal{E}$  is an exact category,

$\triangleright \widehat{M}$  – the class of  $M \in \mathcal{E}$  in  $K_0(\mathcal{E})$ .

In this paper, we take the field  $\mathbf{k} = \mathbb{F}_q$ , a finite field of  $q$  elements. Let  $\mathcal{A}$  be a hereditary abelian category over  $\mathbf{k}$ . We define the  $\iota$ Hall algebra  ${}^{\iota}\widetilde{\mathcal{H}}(\mathcal{A})$  as a twisted semi-derived Ringel-Hall algebra for the category  $\mathcal{C}_1(\mathcal{A})$  of 1-periodic complexes over  $\mathcal{A}$ .

**3.1. Hall algebras.** Let  $\mathcal{E}$  be an essentially small exact category in the sense of Quillen, linear over  $\mathbf{k}$ . Assume that  $\mathcal{E}$  has finite morphism and extension spaces, i.e.,

$$|\mathrm{Hom}(M, N)| < \infty, \quad |\mathrm{Ext}^1(M, N)| < \infty, \quad \forall M, N \in \mathcal{E}.$$

Given objects  $M, N, L \in \mathcal{E}$ , define  $\mathrm{Ext}^1(M, N)_L \subseteq \mathrm{Ext}^1(M, N)$  as the subset parameterizing extensions whose middle term is isomorphic to  $L$ . We define the *Ringel-Hall algebra*  $\mathcal{H}(\mathcal{E})$  (or *Hall algebra* for short) to be the  $\mathbb{Q}$ -vector space whose basis is formed by the isoclasses  $[M]$  of objects  $M$  in  $\mathcal{E}$ , with the multiplication defined by (see [Br13])

$$[M] \diamond [N] = \sum_{[L] \in \mathrm{Iso}(\mathcal{E})} \frac{|\mathrm{Ext}^1(M, N)_L|}{|\mathrm{Hom}(M, N)|} [L]. \quad (3.1)$$

For any three objects  $L, M, N$ , let

$$F_{MN}^L := |\{X \subseteq L \mid X \cong N, L/X \cong M\}|. \quad (3.2)$$

The Riedtmann-Peng formula states that

$$F_{MN}^L = \frac{|\mathrm{Ext}^1(M, N)_L|}{|\mathrm{Hom}(M, N)|} \cdot \frac{|\mathrm{Aut}(L)|}{|\mathrm{Aut}(M)||\mathrm{Aut}(N)|}.$$

For any object  $M$ , let

$$[[M]] := \frac{[M]}{|\mathrm{Aut}(M)|}. \quad (3.3)$$

Then the Hall multiplication (3.1) can be reformulated to be

$$[[M]] \diamond [N] = \sum_{[L]} F_{M,N}^L [[L]], \quad (3.4)$$

which is the original version of Hall multiplication used in [Rin90].

**3.2. The category of 1-periodic complexes.** Let  $\mathcal{A}$  be a hereditary abelian category which is essentially small with finite-dimensional homomorphism and extension spaces.

A 1-periodic complex  $X^\bullet$  in  $\mathcal{A}$  is a pair  $(X, d)$  with  $X \in \mathcal{A}$  and a differential  $d : X \rightarrow X$ . A morphism  $(X, d) \rightarrow (Y, e)$  is given by a morphism  $f : X \rightarrow Y$  in  $\mathcal{A}$  satisfying  $f \circ d = e \circ f$ . Let  $\mathcal{C}_1(\mathcal{A})$  be the category of all 1-periodic complexes in  $\mathcal{A}$ . Then  $\mathcal{C}_1(\mathcal{A})$  is an abelian category. A 1-periodic complex  $X^\bullet = (X, d)$  is called acyclic if  $\mathrm{Ker}(d) = \mathrm{Im}(d)$ . We denote by  $\mathcal{C}_{1,ac}(\mathcal{A})$  the full subcategory of  $\mathcal{C}_1(\mathcal{A})$  consisting of acyclic complexes. Denote by  $H(X^\bullet) \in \mathcal{A}$  the cohomology group of  $X^\bullet$ , i.e.,  $H(X^\bullet) = \mathrm{Ker}(d)/\mathrm{Im}(d)$ , where  $d$  is the differential of  $X^\bullet$ .

The category  $\mathcal{C}_1(\mathcal{A})$  is Frobenius with respect to the degree-wise split exact structure. The 1-periodic homotopy category  $\mathcal{K}_1(\mathcal{A})$  is obtained as the stabilization of  $\mathcal{C}_1(\mathcal{A})$ , and the 1-periodic derived category  $\mathcal{D}_1(\mathcal{A})$  is the localization of the homotopy category  $\mathcal{K}_1(\mathcal{A})$  with respect to quasi-isomorphisms. Both  $\mathcal{K}_1(\mathcal{A})$  and  $\mathcal{D}_1(\mathcal{A})$  are triangulated categories.

Let  $\mathcal{C}^b(\mathcal{A})$  be the category of bounded complexes over  $\mathcal{A}$  and  $\mathcal{D}^b(\mathcal{A})$  be the corresponding derived category with the shift functor  $\Sigma$ . Then there is a covering functor  $\pi : \mathcal{C}^b(\mathcal{A}) \rightarrow$

$\mathcal{C}_1(\mathcal{A})$ , induces a covering functor  $\pi : \mathcal{D}^b(\mathcal{A}) \rightarrow \mathcal{D}_1(\mathcal{A})$  which is dense (see, e.g., [St17, Lemma 5.1]). The orbit category  $\mathcal{D}^b(\mathcal{A})/\Sigma$  is a triangulated category [Ke05], and we have

$$\mathcal{D}_1(\mathcal{A}) \simeq \mathcal{D}^b(\mathcal{A})/\Sigma. \quad (3.5)$$

For any  $X \in \mathcal{A}$ , denote the stalk complex by

$$C_X = (X, 0)$$

(or just by  $X$  when there is no confusion), and denote by  $K_X$  the following acyclic complex:

$$K_X := (X \oplus X, d), \quad \text{where } d = \begin{pmatrix} 0 & \text{Id} \\ 0 & 0 \end{pmatrix}.$$

**Lemma 3.1** ([LRW23, Lemma 2.2]). *For any acyclic complex  $K^\bullet$  and  $p \geq 2$ , we have*

$$\text{Ext}_{\mathcal{C}_1(\mathcal{A})}^p(K^\bullet, -) = 0 = \text{Ext}_{\mathcal{C}_1(\mathcal{A})}^p(-, K^\bullet). \quad (3.6)$$

For any  $K^\bullet \in \mathcal{C}_{1,ac}(\mathcal{A})$  and  $M^\bullet \in \mathcal{C}_1(\mathcal{A})$ , by [LRW23, Corollary 2.4], define

$$\begin{aligned} \langle K^\bullet, M^\bullet \rangle &= \dim_{\mathbf{k}} \text{Hom}_{\mathcal{C}_1(\mathcal{A})}(K^\bullet, M^\bullet) - \dim_{\mathbf{k}} \text{Ext}_{\mathcal{C}_1(\mathcal{A})}^1(K^\bullet, M^\bullet), \\ \langle M^\bullet, K^\bullet \rangle &= \dim_{\mathbf{k}} \text{Hom}_{\mathcal{C}_1(\mathcal{A})}(M^\bullet, K^\bullet) - \dim_{\mathbf{k}} \text{Ext}_{\mathcal{C}_1(\mathcal{A})}^1(M^\bullet, K^\bullet). \end{aligned}$$

These formulas give rise to well-defined bilinear forms (called *Euler forms*), again denoted by  $\langle \cdot, \cdot \rangle$ , on the Grothendieck groups  $K_0(\mathcal{C}_{1,ac}(\mathcal{A}))$  and  $K_0(\mathcal{C}_1(\mathcal{A}))$ .

Denote by  $\langle \cdot, \cdot \rangle_{\mathcal{A}}$  (also denoted by  $\langle \cdot, \cdot \rangle$  if no confusion arises) the Euler form of  $\mathcal{A}$ , i.e.,

$$\langle M, N \rangle_{\mathcal{A}} = \dim_{\mathbf{k}} \text{Hom}_{\mathcal{A}}(M, N) - \dim_{\mathbf{k}} \text{Ext}_{\mathcal{A}}^1(M, N).$$

Let  $\text{res} : \mathcal{C}_1(\mathcal{A}) \rightarrow \mathcal{A}$  be the restriction functor by forgetting differentials.

**Lemma 3.2** ([LRW23, Lemma 2.7]). *We have*

- (1)  $\langle K_X, M^\bullet \rangle = \langle X, \text{res}(M^\bullet) \rangle_{\mathcal{A}}$ ,  $\langle M^\bullet, K_X \rangle = \langle \text{res}(M^\bullet), X \rangle_{\mathcal{A}}$ , for  $X \in \mathcal{A}$ ,  $M^\bullet \in \mathcal{C}_1(\mathcal{A})$ ;
- (2)  $\langle M^\bullet, N^\bullet \rangle = \frac{1}{2} \langle \text{res}(M^\bullet), \text{res}(N^\bullet) \rangle_{\mathcal{A}}$ , for  $M^\bullet, N^\bullet \in \mathcal{C}_{1,ac}(\mathcal{A})$ .

**Lemma 3.3** ([LRW23, Lemma 2.3]). *For any  $X^\bullet, Y^\bullet \in \mathcal{C}_1(\mathcal{A})$ , we have  $H(X^\bullet) \cong H(Y^\bullet)$  if and only if there exist two short exact sequences*

$$0 \longrightarrow U_1^\bullet \longrightarrow Z^\bullet \longrightarrow X^\bullet \longrightarrow 0, \quad 0 \longrightarrow U_2^\bullet \longrightarrow Z^\bullet \longrightarrow Y^\bullet \longrightarrow 0$$

with  $U_1^\bullet, U_2^\bullet \in \mathcal{C}_{1,ac}(\mathcal{A})$ .

**3.3. Semi-derived Hall algebras and iHall algebras.** Define

$$\mathbf{v} := \sqrt{q}.$$

We continue to work with a hereditary abelian category  $\mathcal{A}$  as in §3.2. Let  $\mathcal{H}(\mathcal{C}_1(\mathcal{A}))$  be the Ringel-Hall algebra of  $\mathcal{C}_1(\mathcal{A})$  over  $\mathbb{Q}(\mathbf{v})$ , i.e.,  $\mathcal{H}(\mathcal{C}_1(\mathcal{A})) = \bigoplus_{[M^\bullet] \in \text{Iso}(\mathcal{C}_1(\mathcal{A}))} \mathbb{Q}(\mathbf{v})[M^\bullet]$ , with multiplication defined by

$$[M^\bullet] \diamond [N^\bullet] = \sum_{[L^\bullet] \in \text{Iso}(\mathcal{C}_1(\mathcal{A}))} \frac{|\text{Ext}^1(M^\bullet, N^\bullet)_{L^\bullet}|}{|\text{Hom}(M^\bullet, N^\bullet)|} [L^\bullet].$$

For any  $M \in \mathcal{A}$ , we denote by  $\widehat{M}$  its class in the Grothendieck group  $K_0(\mathcal{A})$  of  $\mathcal{A}$ . Following [LP21, LW22a, LW23], we consider the ideal  $\mathcal{I}$  of  $\mathcal{H}(\mathcal{C}_1(\mathcal{A}))$  generated by

$$\left\{ [M^\bullet] - [N^\bullet] \mid H(M^\bullet) \cong H(N^\bullet), \quad \widehat{\text{Im}(d_{M^\bullet})} = \widehat{\text{Im}(d_{N^\bullet})} \right\}. \quad (3.7)$$

We denote

$$\mathcal{S} := \{a[K^\bullet] \in \mathcal{H}(\mathcal{C}_1(\mathcal{A}))/\mathcal{I} \mid a \in \mathbb{Q}(\mathbf{v})^\times, K^\bullet \in \mathcal{C}_1(\mathcal{A}) \text{ acyclic}\},$$

a multiplicatively closed subset of  $\mathcal{H}(\mathcal{C}_1(\mathcal{A}))/\mathcal{I}$  with the identity  $[0]$ .

**Lemma 3.4** ([LW22a, Proposition A.5]). *The multiplicatively closed subset  $\mathcal{S}$  is a right Ore, right reversible subset of  $\mathcal{H}(\mathcal{C}_1(\mathcal{A}))/\mathcal{I}$ . Equivalently, there exists the right localization of  $\mathcal{H}(\mathcal{C}_1(\mathcal{A}))/\mathcal{I}$  with respect to  $\mathcal{S}$ , denoted by  $(\mathcal{H}(\mathcal{C}_1(\mathcal{A}))/\mathcal{I})[\mathcal{S}^{-1}]$ .*

*Proof.* It follows by noting that  $[K^\bullet] \diamond [M^\bullet] = q^{\langle M^\bullet, K^\bullet \rangle - \langle K^\bullet, M^\bullet \rangle} [M^\bullet] \diamond [K^\bullet]$  in  $\mathcal{H}(\mathcal{C}_1(\mathcal{A}))/\mathcal{I}$  for any acyclic  $K^\bullet$ .  $\square$

The algebra  $(\mathcal{H}(\mathcal{C}_1(\mathcal{A}))/\mathcal{I})[\mathcal{S}^{-1}]$  is the *semi-derived Ringel-Hall algebra* of  $\mathcal{C}_1(\mathcal{A})$  in the sense of [LP21, LW22a] (also cf. [Gor18]), and will be denoted by  $\mathcal{SDH}(\mathcal{C}_1(\mathcal{A}))$ .

**Definition 3.5.** *The  $\imath$ Hall algebra of a hereditary abelian category  $\mathcal{A}$ , denoted by  ${}^{\imath}\tilde{\mathcal{H}}(\mathcal{A})$ , is defined to be the twisted semi-derived Ringel-Hall algebra of  $\mathcal{C}_1(\mathcal{A})$ , that is, the  $\mathbb{Q}(\mathbf{v})$ -algebra on the same vector space as  $\mathcal{SDH}(\mathcal{C}_1(\mathcal{A})) = (\mathcal{H}(\mathcal{C}_1(\mathcal{A}))/\mathcal{I})[\mathcal{S}^{-1}]$  equipped with the following modified multiplication (twisted via the restriction functor  $\text{res} : \mathcal{C}_1(\mathcal{A}) \rightarrow \mathcal{A}$ )*

$$[M^\bullet] * [N^\bullet] = \mathbf{v}^{\langle \text{res}(M^\bullet), \text{res}(N^\bullet) \rangle_{\mathcal{A}}} [M^\bullet] \diamond [N^\bullet]. \quad (3.8)$$

For any complex  $M^\bullet$  and acyclic complex  $K^\bullet$ , we have

$$[K^\bullet] * [M^\bullet] = [K^\bullet \oplus M^\bullet] = [M^\bullet] * [K^\bullet].$$

For any  $\alpha \in K_0(\mathcal{A})$ , there exist  $X, Y \in \mathcal{A}$  such that  $\alpha = \widehat{X} - \widehat{Y}$ . Define  $[K_\alpha] := [K_X] * [K_Y]^{-1}$ . This is well defined, see e.g., [LP21, §3.2]. It follows that  $[K_\alpha]$  ( $\alpha \in K_0(\mathcal{A})$ ) are central in the algebra  ${}^{\imath}\tilde{\mathcal{H}}(\mathcal{A})$ .

The *quantum torus*  $\tilde{\mathcal{T}}(\mathcal{A})$  is defined to be the subalgebra of  ${}^{\imath}\tilde{\mathcal{H}}(\mathcal{A})$  generated by  $[K_\alpha]$ , for  $\alpha \in K_0(\mathcal{A})$ .

**Proposition 3.6** ([LRW23, Proposition 2.9]). *The following hold in  ${}^{\imath}\tilde{\mathcal{H}}(\mathcal{A})$ :*

- (1) *The quantum torus  $\tilde{\mathcal{T}}(\mathcal{A})$  is a central subalgebra of  ${}^{\imath}\tilde{\mathcal{H}}(\mathcal{A})$ .*
- (2) *The algebra  $\tilde{\mathcal{T}}(\mathcal{A})$  is isomorphic to the group algebra of the abelian group  $K_0(\mathcal{A})$ .*
- (3)  *${}^{\imath}\tilde{\mathcal{H}}(\mathcal{A})$  has an ( $\imath$ Hall) basis given by*

$$\{[M] * [K_\alpha] \mid [M] \in \text{Iso}(\mathcal{A}), \alpha \in K_0(\mathcal{A})\}.$$

For any  $f : X \rightarrow Y$  in  $\mathcal{A}$ , we denote by

$$C_f := \left( Y \oplus X, \begin{pmatrix} 0 & f \\ 0 & 0 \end{pmatrix} \right) \in \mathcal{C}_1(\mathcal{A}).$$

**Lemma 3.7** ([LRW23, Lemma 2.10]). *For any  $M^\bullet = (M, d)$ , we have  $[M^\bullet] = [H(M^\bullet)] * [K_{\text{Im}(d)}]$  in  ${}^{\imath}\tilde{\mathcal{H}}(\mathcal{A})$ . In particular, for any  $f : X \rightarrow Y$ , we have*

$$[C_f] = [\text{Ker}(f) \oplus \text{Coker}(f)] * [K_{\text{Im}(f)}].$$

**3.4. Realization of  $i$ quantum groups via quivers.** Let  $Q = (Q_0, Q_1)$  be a quiver (not necessarily acyclic), and we sometimes write  $\mathbb{I} = Q_0$ . For a quiver  $Q$ , we denote

- ▷  $\text{rep}_{\mathbf{k}}(Q)$  – category of finite-dimensional representations of  $Q$  over  $\mathbf{k}$ ,
- ▷  $\text{rep}_{\mathbf{k}}^{\text{nil}}(Q)$  – subcategory of  $\text{rep}_{\mathbf{k}}(Q)$  consisting of nilpotent representations of  $Q$ .

Note that  $\text{rep}_{\mathbf{k}}(Q) = \text{rep}_{\mathbf{k}}^{\text{nil}}(Q)$  if  $Q$  is acyclic.

Let  $S_i$  be the simple  $\mathbf{k}Q$ -module associated to  $i \in \mathbb{I}$ . We denote by  ${}^i\tilde{\mathcal{H}}(\mathbf{k}Q)$  the twisted semi-derived Ringel-Hall algebra of  $\mathcal{C}_1(\text{rep}_{\mathbf{k}}^{\text{nil}}(Q))$ . Then we have the following result.

**Theorem 3.8** ([LW23]). *Let  $Q$  be an arbitrary quiver. Then there exists a  $\mathbb{Q}(\mathbf{v})$ -algebra embedding*

$$\tilde{\psi}_Q : \tilde{\mathbf{U}}_{|v=\mathbf{v}}^i \longrightarrow {}^i\tilde{\mathcal{H}}(\mathbf{k}Q),$$

which sends

$$B_i \mapsto \frac{-1}{q-1}[S_i], \quad \mathbb{K}_i \mapsto [K_{S_i}], \quad \text{for } i \in \mathbb{I}. \quad (3.9)$$

Let  ${}^i\tilde{\mathcal{C}}(\mathbf{k}Q)$  be the composition subalgebra of  ${}^i\tilde{\mathcal{H}}(\mathbf{k}Q)$  generated by  $[S_i]$  and  $[K_{S_i}]^{\pm 1}$  for  $i \in \mathbb{I}$ . Then (3.9) gives a  $\mathbb{Q}(\mathbf{v})$ -algebra isomorphism

$$\tilde{\psi}_Q : \tilde{\mathbf{U}}_{|v=\mathbf{v}}^i \xrightarrow{\cong} {}^i\tilde{\mathcal{C}}(\mathbf{k}Q). \quad (3.10)$$

**3.5. An  $i$ Hall multiplication formula.** The following multiplication formulas in the  $i$ Hall algebra are useful and applicable.

**Proposition 3.9** ([LW23, Proposition 3.10]). *Let  $\mathcal{A}$  be a hereditary abelian category over  $\mathbf{k}$ . For any  $A, B \in \mathcal{A} \subset \mathcal{C}_1(\mathcal{A})$ , the following formulas hold in  ${}^i\tilde{\mathcal{H}}(\mathcal{A})$ .*

$$\begin{aligned} [A] * [B] &= \sum_{[M] \in \text{Iso}(\mathcal{A})} \sum_{[L], [N] \in \text{Iso}(\mathcal{A})} \mathbf{v}^{-\langle A, B \rangle} q^{\langle N, L \rangle} \frac{|\text{Ext}^1(N, L)_M|}{|\text{Hom}(N, L)|} \\ &\quad \cdot |\{f \in \text{Hom}(A, B) \mid \text{Ker}(f) \cong N, \text{Coker}(f) \cong L\}| [M] * [K_{\hat{A}-\hat{N}}], \end{aligned} \quad (3.11)$$

$$\begin{aligned} [A] * [B] &= \sum_{[M] \in \text{Iso}(\mathcal{A})} \sum_{[L], [N] \in \text{Iso}(\mathcal{A})} \mathbf{v}^{-\langle A, B \rangle} \frac{|\text{Ext}^1(N, L)_M|}{|\text{Ext}^1(N, L)|} \\ &\quad \cdot |\{f \in \text{Hom}(A, B) \mid \text{Ker}(f) \cong N, \text{Coker}(f) \cong L\}| [M] * [K_{\hat{A}-\hat{N}}] \end{aligned} \quad (3.12)$$

*Proof.* The proof of (3.11) is completely same to [LW23, Proposition 3.10], hence omitted here. (3.12) follows from (3.11) by a direct computation.  $\square$

**Corollary 3.10.** *For any  $A, B \in \mathcal{A}$ , the following formulas hold in  ${}^i\tilde{\mathcal{H}}(\mathcal{A})$ .*

(1) *If any non-zero map in  $\text{Hom}(A, B)$  is injective, then*

$$\begin{aligned} [A] * [B] &= \mathbf{v}^{\langle A, B \rangle} \sum_{[M] \in \text{Iso}(\mathcal{A})} F_{A, B}^M [M] \\ &\quad + \mathbf{v}^{-\langle A, B \rangle} \sum_{[M] \in \text{Iso}(\mathcal{A})} \frac{|\text{Ext}^1(M, A)_B|}{|\text{Hom}(M, A)| \cdot |\text{Aut}(A)|} [M] * [K_{\hat{A}}]. \end{aligned}$$

(2) If any non-zero map in  $\text{Hom}(A, B)$  is surjective, then

$$\begin{aligned} [A] * [B] &= \mathbf{v}^{\langle A, B \rangle} \sum_{[M] \in \text{Iso}(\mathcal{A})} F_{A, B}^M [M] \\ &\quad + \mathbf{v}^{-\langle A, B \rangle} \sum_{[M] \in \text{Iso}(\mathcal{A})} \frac{|\text{Ext}^1(B, M)_A|}{|\text{Hom}(B, M)| \cdot |\text{Aut}(B)|} [M] * [K_{\hat{B}}]. \end{aligned}$$

*Proof.* We only prove (1) since (2) is dual. For any non-zero map  $f : A \rightarrow B$ , by assumption  $f$  is injective, i.e.,  $\text{Ker}(f) = 0$ . So it follows from (3.11) that

$$\begin{aligned} [A] * [B] &= \sum_{[M] \in \text{Iso}(\mathcal{A})} \mathbf{v}^{\langle A, B \rangle} \frac{|\text{Ext}^1(A, B)_M|}{|\text{Hom}(A, B)|} [M] \\ &\quad + \sum_{[M] \in \text{Iso}(\mathcal{A})} \mathbf{v}^{-\langle A, B \rangle} |\{0 \neq f \in \text{Hom}(A, B) \mid \text{Coker}(f) \cong M\}| \cdot [M] * [K_{\hat{A}}] \\ &= \sum_{[M] \in \text{Iso}(\mathcal{A})} \left( \mathbf{v}^{\langle A, B \rangle} \frac{|\text{Ext}^1(A, B)_M|}{|\text{Hom}(A, B)|} [M] + \mathbf{v}^{-\langle A, B \rangle} F_{M, A}^B |\text{Aut}(A)| \cdot [M] * [K_{\hat{A}}] \right). \end{aligned}$$

Then the desired formula follows from the Riedtmann-Peng formula and (3.3).  $\square$

#### 4. WEIGHTED PROJECTIVE LINES AND COHERENT SHEAVES

In this section, we recall some basic facts on the category of coherent sheaves on a weighted projective line given by Geigle and Lenzing [GL87].

**4.1. Weighted projective lines.** Recall  $\mathbf{k} = \mathbb{F}_q$ . Fix a positive integer  $\mathbf{s}$  such that  $2 \leq \mathbf{s} \leq q$  in the following. Let  $\mathbf{p} = (p_1, p_2, \dots, p_{\mathbf{s}})$  with  $p_i \geq 1$ . Without loss of generality, we assume  $p_i \geq 2$  for  $1 \leq i \leq \mathbf{t}$ , and  $p_i = 1$  for  $\mathbf{t} + 1 \leq i \leq \mathbf{s}$ ; compare with the star-shaped graph  $\Gamma = \mathbb{T}_{p_1, \dots, p_{\mathbf{t}}}$ .

Let  $L(\mathbf{p})$  be the rank one abelian group on generators  $\vec{x}_1, \vec{x}_2, \dots, \vec{x}_{\mathbf{s}}$  with relations  $p_1 \vec{x}_1 = p_2 \vec{x}_2 = \dots = p_{\mathbf{s}} \vec{x}_{\mathbf{s}}$ . We call  $\vec{c} := p_i \vec{x}_i$  the *canonical element* of  $L(\mathbf{p})$ . Obviously, the polynomial ring  $\mathbf{k}[X_1, \dots, X_{\mathbf{s}}]$  is a  $L(\mathbf{p})$ -graded algebra by setting  $\deg X_i = \vec{x}_i$ , which is denoted by  $\mathbf{S}(\mathbf{p})$ .

Let  $\underline{\lambda} = \{\lambda_1, \dots, \lambda_{\mathbf{s}}\}$  be a collection of distinguished closed points (of degree one) on the projective line  $\mathbb{P}_{\mathbf{k}}^1$ , normalized such that  $\lambda_1 = \infty$ ,  $\lambda_2 = 0$ ,  $\lambda_3 = 1$ . Let  $I(\mathbf{p}, \underline{\lambda})$  be the  $L(\mathbf{p})$ -graded ideal of  $\mathbf{S}(\mathbf{p})$  generated by

$$\{X_i^{p_i} - (X_2^{p_2} - \lambda_i X_1^{p_1}) \mid 3 \leq i \leq \mathbf{s}\}.$$

Then  $\mathbf{S}(\mathbf{p}, \underline{\lambda}) := \mathbf{S}(\mathbf{p})/I(\mathbf{p}, \underline{\lambda})$  is an  $L(\mathbf{p})$ -graded algebra. For any  $1 \leq i \leq \mathbf{s}$ , denote by  $x_i$  the image of  $X_i$  in  $\mathbf{S}(\mathbf{p}, \underline{\lambda})$ .

The *weighted projective line*  $\mathbb{X} := \mathbb{X}_{\mathbf{p}, \underline{\lambda}}$  is the set of all non-maximal prime homogeneous ideals of  $\mathbf{S} := \mathbf{S}(\mathbf{p}, \underline{\lambda})$ , which is also denoted by  $\mathbb{X}_{\mathbf{k}}$  to emphasis the base field  $\mathbf{k}$ . For any homogeneous element  $f \in \mathbf{S}$ , let  $V_f := \{\mathfrak{p} \in \mathbb{X} \mid f \in \mathfrak{p}\}$  and  $D_f := \mathbb{X} \setminus V_f$ . The set  $\{D_f \mid f \in \mathbf{S}\}$  forms a basis of the Zariski topology on  $\mathbb{X}$ . The pair  $(\mathbf{p}, \underline{\lambda})$  (or just  $\mathbf{p}$  if  $\underline{\lambda}$  is obvious) is called the *weight type* of  $\mathbb{X}_{\mathbf{p}, \underline{\lambda}}$ , and  $p_i$  is the weight of the point  $\lambda_i$ . Roughly speaking,  $\mathbb{X}$  is the projective line  $\mathbb{P}_{\mathbf{k}}^1$  with a finite number of marked points  $\lambda_1, \dots, \lambda_{\mathbf{s}}$  with





Let  $C_1$  be the Jordan quiver, i.e., the quiver with only one vertex and one loop arrow.

Then the structure of  $\mathcal{T}$  is described in the following.

**Lemma 4.2** ([GL87]; also see [Sch04]). (1) The category  $\mathcal{T}$  decomposes as a coproduct  $\mathcal{T} = \coprod_{x \in \mathbb{X}} \mathcal{T}_x$ , where  $\mathcal{T}_x$  is the subcategory of torsion sheaves with support at  $x$ .

(2) For any ordinary point  $x$  of degree  $d$ , let  $\mathbf{k}_x$  denote the residue field at  $x$ , i.e.,  $[\mathbf{k}_x : \mathbf{k}] = d$ . Then  $\mathcal{T}_x$  is equivalent to the category  $\text{rep}_{\mathbf{k}_x}^{\text{nil}}(C_1)$ .

(3) For any exceptional point  $\lambda_i$  ( $1 \leq i \leq \mathbf{s}$ ), the category  $\mathcal{T}_{\lambda_i}$  is equivalent to  $\text{rep}_{\mathbf{k}}^{\text{nil}}(C_{p_i})$ .

For any ordinary point  $x$  of degree  $d$ , let  $\pi_x$  be the prime homogeneous polynomial corresponding to  $x$ . The multiplication by  $\pi_x$  gives the exact sequence

$$0 \longrightarrow \mathcal{O} \xrightarrow{\pi_x} \mathcal{O}(d\vec{c}) \longrightarrow S_x \rightarrow 0,$$

where  $S_x$  is the unique (up to isomorphism) simple sheaf in the category  $\mathcal{T}_x$ . Then  $S_x(\vec{l}) = S_x$  for any  $\vec{l} \in L(\mathbf{p})$ .

For any exceptional point  $\lambda_i$ , multiplication by  $x_i$  yields the short exact sequence

$$0 \longrightarrow \mathcal{O}((j-1)\vec{x}_i) \xrightarrow{x_i} \mathcal{O}(j\vec{x}_i) \longrightarrow S_{ij} \rightarrow 0, \text{ for } 1 \leq j \leq p_i;$$

where  $\{S_{ij} \mid j \in \mathbb{Z}_{p_i}\}$  is a complete set of pairwise non-isomorphic simple sheaves in the category  $\mathcal{T}_{\lambda_i}$  for any  $1 \leq i \leq \mathbf{s}$ . Moreover,  $S_{ij}(\vec{l}) = S_{i,j+l_i}$  for any  $\vec{l} = \sum_{1 \leq i \leq \mathbf{s}} l_i \vec{x}_i$ .

In order to describe the indecomposable objects in  $\mathcal{T}$ , we need the following well-known results. For  $\text{rep}_{\mathbf{k}}^{\text{nil}}(C_n)$ , we have  $n$  simple representations  $S_j = S_j^{(1)}$  ( $1 \leq j \leq n$ ). Denote by  $S_j^{(a)}$  the unique indecomposable representation with top  $S_j$  and length  $a$ . Then  $\{S_j^{(a)} \mid 1 \leq j \leq n, a \in \mathbb{N}\}$  is the set of isoclasses of indecomposable objects of  $\text{rep}_{\mathbf{k}}^{\text{nil}}(C_n)$ . For any  $1 \leq j \leq n$ , define  $S_j^{(\mu)} = \bigoplus_i S_j^{(\mu_i)}$  for any partition  $\mu = (\mu_1 \geq \dots \geq \mu_l)$ . Combining with Lemma 4.2, we can give a classification of indecomposable objects in  $\mathcal{T}$ .

We denote by  $S_x^{(a)}$  the unique indecomposable object of length  $a$  in  $\mathcal{T}_x$  for any ordinary point  $x$ ; and denote by  $S_{ij}^{(a)}$  the unique indecomposable object with top  $S_{ij}$  and length  $a$  in  $\mathcal{T}_{\lambda_i}$ .

**4.4. Grothendieck groups and Euler forms.** The Grothendieck group  $K_0(\text{coh}(\mathbb{X}))$  of  $\text{coh}(\mathbb{X})$  satisfies

$$K_0(\text{coh}(\mathbb{X})) \cong \left( \mathbb{Z}\widehat{\mathcal{O}} \oplus \mathbb{Z}\widehat{\mathcal{O}(\vec{c})} \oplus \bigoplus_{i,j} \mathbb{Z}\widehat{S}_{ij} \right) / I, \quad (4.4)$$

where  $I$  is the subgroup generated by  $\{\sum_{j=1}^{p_i} \widehat{S}_{ij} + \widehat{\mathcal{O}} - \widehat{\mathcal{O}(\vec{c})} \mid i = 1, \dots, \mathbf{s}\}$ ; see [GL87]. Let  $\delta := \widehat{\mathcal{O}(\vec{c})} - \widehat{\mathcal{O}} = \sum_{j=1}^{p_i} \widehat{S}_{ij}$ . Note that  $\widehat{S}_{i1} = \delta$  if  $p_i = 1$ . The Euler form  $\langle -, - \rangle$  on  $K_0(\text{coh}(\mathbb{X}))$  is described in [Sch04], which is given by

$$\langle \widehat{\mathcal{O}}, \widehat{\mathcal{O}} \rangle = 1, \quad \langle \widehat{\mathcal{O}}, \delta \rangle = 1, \quad \langle \delta, \widehat{\mathcal{O}} \rangle = -1, \quad (4.5)$$

$$\langle \delta, \delta \rangle = 0, \quad \langle \widehat{S}_{ij}, \delta \rangle = 0, \quad \langle \delta, \widehat{S}_{ij} \rangle = 0, \quad (4.6)$$

$$\langle \widehat{\mathcal{O}}, \widehat{S}_{ij} \rangle = \begin{cases} 1 & \text{if } j = p_i, \\ 0 & \text{if } j \neq p_i, \end{cases} \quad \langle \widehat{S}_{ij}, \widehat{\mathcal{O}} \rangle = \begin{cases} -1 & \text{if } j = 1, \\ 0 & \text{if } j \neq 1, \end{cases} \quad (4.7)$$

$$\langle \widehat{S}_{ij}, \widehat{S}_{i',j'} \rangle = \begin{cases} 1 & \text{if } i = i', j = j', \\ -1 & \text{if } i = i', j \equiv j' + 1 \pmod{p_i}, \\ 0 & \text{otherwise.} \end{cases} \quad (4.8)$$

Recall the root system  $\mathcal{R}$  of  $L\mathfrak{g}$  of the star-shaped graph  $\Gamma = T_{p_1, \dots, p_t}$ . Then there is a natural isomorphism of  $\mathbb{Z}$ -modules  $K_0(\text{coh}(\mathbb{X})) \cong \mathcal{R}$  given as below (for  $1 \leq i \leq t$ ,  $1 \leq j \leq p_i - 1$ ,  $r \in \mathbb{N}$  and  $l \in \mathbb{Z}$ ):

$$\widehat{S}_{ij} \mapsto \alpha_{ij}, \quad \widehat{S}_{i,0} \mapsto \delta - \sum_{j=1}^{p_i-1} \alpha_{ij}, \quad \widehat{S}_{i,0}^{(p_i-1)} \mapsto \delta - \alpha_{i1}, \quad \widehat{S}_{i,0}^{(rp_i)} \mapsto r\delta, \quad \widehat{\mathcal{O}(l\vec{c})} \mapsto \alpha_\star + l\delta.$$

Under this isomorphism, the symmetric Euler form on  $K_0(\text{coh}(\mathbb{X}))$  coincides with the Cartan form on  $\mathcal{R}$ . So we always identify  $K_0(\text{coh}(\mathbb{X}))$  with  $\mathcal{R}$  via this isomorphism in the following.

Let  $p = \text{l.c.m.}(p_1, \dots, p_t)$ . We can define the degree and rank for each coherent sheaf, which induces two functions on  $K_0(\text{coh}(\mathbb{X}))$  denoted by  $\text{deg}$  and  $\text{rank}$  respectively. In fact, they are uniquely determined by

$$\text{deg}(\mathcal{O}) = 0, \quad \text{deg}(S_{ij}) = \frac{p}{p_i}; \quad \text{rank}(\mathcal{O}) = 1, \quad \text{rank}(S_{ij}) = 0; \quad (4.9)$$

for any  $1 \leq i \leq t, 0 \leq j \leq p_i - 1$ .

**4.5. vHall algebras of weighted projective lines.** By [GL87], we know that  $\text{coh}(\mathbb{X})$  is a hereditary abelian category, so we can define the vHall algebra (a.k.a. the twisted semi-derived Ringel-Hall algebra) of  $\mathbb{X}$ , which is denoted by  ${}^v\widetilde{\mathcal{H}}(\mathbb{X}_{\mathbf{k}})$ .

A direct computation in  ${}^v\widetilde{\mathcal{H}}(\mathbb{X}_{\mathbf{k}})$  shows that for any  $l \in \mathbb{Z}$  and  $1 \leq i \leq t$ :

$$\begin{aligned} [\mathcal{O}(l\vec{c})] * [S_{i,1}] &= [S_{i,1} \oplus \mathcal{O}(l\vec{c})], \\ [S_{i,1}] * [\mathcal{O}(l\vec{c})] &= \mathbf{v}^{-1}[S_{i,1} \oplus \mathcal{O}(l\vec{c})] + (\mathbf{v} - \mathbf{v}^{-1})[\mathcal{O}(l\vec{c} + \vec{x}_i)]; \end{aligned} \quad (4.10)$$

and

$$\begin{aligned} [S_{i,0}^{(p_i-1)}] * [\mathcal{O}(l\vec{c})] &= [S_{i,0}^{(p_i-1)} \oplus \mathcal{O}(l\vec{c})], \\ [\mathcal{O}(l\vec{c})] * [S_{i,0}^{(p_i-1)}] &= \mathbf{v}^{-1}[S_{i,0}^{(p_i-1)} \oplus \mathcal{O}(l\vec{c})] + (\mathbf{v} - \mathbf{v}^{-1})[\mathcal{O}((l-1)\vec{c} + \vec{x}_i)] * [K_{\delta - \alpha_i}]. \end{aligned} \quad (4.11)$$

The following results will be used in the proof of Lemma 7.3 and Proposition 8.1.

**Lemma 4.3.** *For any  $1 \leq i \leq t$  and  $1 \leq j < k \leq p_i$ , the following equations hold in  ${}^v\widetilde{\mathcal{H}}(\mathbb{X}_{\mathbf{k}})$ :*

$$[[\mathcal{O}(l\vec{c})], [S_{ij}^{(k)}]] = [[\mathcal{O}(l\vec{c})], [S_{ij}^{(j)} \oplus S_{i,0}^{(k-j)}]], \quad \text{for any } l \in \mathbb{Z}; \quad (4.12)$$

$$[[S_{i,0}^{(rp_i)}], [S_{ij}^{(k)}]] = [[S_{i,0}^{(rp_i)}], [S_{ij}^{(j)} \oplus S_{i,0}^{(k-j)}]], \quad \text{for any } r \geq 1. \quad (4.13)$$

*Proof.* First, we prove (4.12). Observe that we have the following exact sequence

$$0 \rightarrow S_{i,0}^{(k-j)} \rightarrow S_{ij}^{(k)} \rightarrow S_{ij}^{(j)} \rightarrow 0. \quad (4.14)$$

The following computations are in  $\mathcal{C}_1(\text{coh}(\mathbb{X}))$ .

By applying  $\text{Hom}_{\mathcal{C}_1(\text{coh}(\mathbb{X}))}(\mathcal{O}(l\vec{c}), -)$  and  $\text{Hom}_{\mathcal{C}_1(\text{coh}(\mathbb{X}))}(-, \mathcal{O}(l\vec{c}))$  to (4.14), we know that

$$\text{Hom}(\mathcal{O}(l\vec{c}), S_{ij}^{(k)}) \cong \text{Hom}(\mathcal{O}(l\vec{c}), S_{i,0}^{(k-j)}) \cong \mathbf{k}, \quad \text{Ext}^1(S_{ij}^{(k)}, \mathcal{O}(l\vec{c})) \cong \text{Ext}^1(S_{ij}^{(j)}, \mathcal{O}(l\vec{c})) \cong \mathbf{k}.$$

Let  $X \in \{S_{ij}^{(k)}, S_{ij}^{(j)} \oplus S_{i,0}^{(k-j)}\}$ . For any non-zero morphism  $f : \mathcal{O}(l\vec{c}) \rightarrow X$ , there are

$$\mathrm{Im}(f) = S_{i,0}^{(k-j)}, \quad \mathrm{Ker}(f) = \mathcal{O}(l\vec{c} - (k-j)\vec{x}_i), \quad \mathrm{Coker}(f) = S_{ij}^{(j)},$$

while the non-trivial extension  $\mathrm{Ext}^1(X, \mathcal{O}(l\vec{c}))$  is given by

$$0 \rightarrow \mathcal{O}(l\vec{c}) \longrightarrow \mathcal{O}(l\vec{c} + j\vec{x}_i) \oplus S_{i,0}^{(k-j)} \longrightarrow X \rightarrow 0.$$

Therefore, by (3.11) we have

$$\begin{aligned} & [[\mathcal{O}(l\vec{c})], [S_{ij}^{(k)}]] \\ &= [\mathcal{O}(l\vec{c})] * [S_{ij}^{(k)}] - [S_{ij}^{(k)}] * [\mathcal{O}(l\vec{c})] \\ &= \mathbf{v}^{\widehat{(\mathcal{O}(l\vec{c}), S_{ij}^{(k)})}} \cdot \left( \frac{1}{q} [\mathcal{O}(l\vec{c}) \oplus S_{ij}^{(k)}] + \frac{q-1}{q} [\mathcal{O}(l\vec{c} - (k-j)\vec{x}_i) \oplus S_{ij}^{(j)}] * [K_{S_{i,0}^{(k-j)}}] \right) \\ &\quad - \mathbf{v}^{\widehat{(S_{ij}^{(k)}, \mathcal{O}(l\vec{c})}}) \cdot \left( [\mathcal{O}(l\vec{c}) \oplus S_{ij}^{(k)}] + (q-1) [\mathcal{O}(l\vec{c} + j\vec{x}_i) \oplus S_{i,0}^{(k-j)}] \right) \\ &= (\mathbf{v} - \mathbf{v}^{-1}) \left( [\mathcal{O}(l\vec{c} - (k-j)\vec{x}_i) \oplus S_{ij}^{(j)}] * [K_{S_{i,0}^{(k-j)}}] - [\mathcal{O}(l\vec{c} + j\vec{x}_i) \oplus S_{i,0}^{(k-j)}] \right). \end{aligned}$$

Similarly,

$$\begin{aligned} & [[\mathcal{O}(l\vec{c})], [S_{ij}^{(j)} \oplus S_{i,0}^{(k-j)}]] \\ &= (\mathbf{v} - \mathbf{v}^{-1}) \left( [\mathcal{O}(l\vec{c} - (k-j)\vec{x}_i) \oplus S_{ij}^{(j)}] * [K_{S_{i,0}^{(k-j)}}] - [\mathcal{O}(l\vec{c} + j\vec{x}_i) \oplus S_{i,0}^{(k-j)}] \right), \end{aligned}$$

and then (4.12) follows.

For (4.13), observe that the composition factors of  $S_{ij}^{(j)}$  are  $S_{i,1}, S_{i,2}, \dots, S_{ij}$  (from socle to top), while the composition factors of  $S_{i,0}^{(k-j)}$  are  $S_{i,p_i+1-k+j}, \dots, S_{i,p_i-1}, S_{i,0}$ . Hence

$$\mathrm{Hom}_{\mathbb{X}}(S_{i,0}^{(rp_i)}, S_{ij}^{(j)}) = 0 = \mathrm{Ext}_{\mathbb{X}}^1(S_{i,0}^{(rp_i)}, S_{ij}^{(j)}), \quad \mathrm{Hom}_{\mathbb{X}}(S_{ij}^{(k-j)}, S_{i,0}^{(rp_i)}) = 0 = \mathrm{Ext}_{\mathbb{X}}^1(S_{ij}^{(k-j)}, S_{i,0}^{(rp_i)}).$$

By applying  $\mathrm{Hom}_{\mathbb{X}}(S_{i,0}^{(rp_i)}, -)$  and  $\mathrm{Hom}_{\mathbb{X}}(-, S_{i,0}^{(rp_i)})$  to (4.14), we obtain

$$\begin{aligned} \mathrm{Hom}_{\mathbb{X}}(S_{i,0}^{(rp_i)}, S_{ij}^{(k)}) &\cong \mathrm{Hom}(S_{i,0}^{(rp_i)}, S_{i,0}^{(k-j)}), & \mathrm{Ext}_{\mathbb{X}}^1(S_{i,0}^{(rp_i)}, S_{ij}^{(k)}) &\cong \mathrm{Ext}^1(S_{i,0}^{(rp_i)}, S_{i,0}^{(k-j)}), \\ \mathrm{Hom}_{\mathbb{X}}(S_{ij}^{(k)}, S_{i,0}^{(rp_i)}) &\cong \mathrm{Hom}_{\mathbb{X}}(S_{i,0}^{(k-j)}, S_{i,0}^{(rp_i)}), & \mathrm{Ext}_{\mathbb{X}}^1(S_{ij}^{(k)}, S_{i,0}^{(rp_i)}) &\cong \mathrm{Ext}_{\mathbb{X}}^1(S_{i,0}^{(k-j)}, S_{i,0}^{(rp_i)}), \end{aligned}$$

which are of dimension 1. Then (4.13) holds by a similar argument as (4.12).  $\square$

**Lemma 4.4.** *For any  $1 \leq i \leq \mathbf{t}$  and  $l \in \mathbb{Z}$ ,*

$$[[S_{i,1}^{(p_i+1)}] - [S_{i,1} \oplus S_{i,0}^{(p_i)}], [\mathcal{O}(l\vec{c})]]_{\mathbf{v}^{-1}} = \mathbf{v} [[S_{i,1}], [\mathcal{O}((l+1)\vec{c})]]_{\mathbf{v}}.$$

*Proof.* By (4.10), we have

$$[[S_{i,1}], [\mathcal{O}((l+1)\vec{c})]]_{\mathbf{v}} = (\mathbf{v} - \mathbf{v}^{-1}) \left( [\mathcal{O}((l+1)\vec{c} + \vec{x}_i)] - [\mathcal{O}((l+1)\vec{c}) \oplus S_{i,1}] \right).$$

On the other hand, by definition we have

$$\begin{aligned} & [[S_{i,1}^{(p_i+1)}], [\mathcal{O}(l\vec{c})]]_{\mathbf{v}^{-1}} \\ &= [S_{i,1}^{(p_i+1)}] * [\mathcal{O}(l\vec{c})] - \mathbf{v}^{-1} [\mathcal{O}(l\vec{c})] * [S_{i,1}^{(p_i+1)}] \\ &= \mathbf{v}^{-2} \left( [S_{i,1}^{(p_i+1)} \oplus \mathcal{O}(l\vec{c})] + (q-1) [\mathcal{O}(l\vec{c} + \vec{x}_i) \oplus S_{i,0}^{(p_i)}] + (q^2 - q) [\mathcal{O}((l+1)\vec{c} + \vec{x}_i)] \right) \end{aligned}$$

$$\begin{aligned}
 & -\mathbf{v}^{-1} \cdot \frac{\mathbf{v}}{q} \left( [S_{i,1}^{(p_i+1)}] \oplus \mathcal{O}(l\vec{c}) + (q-1)[S_{i,1} \oplus \mathcal{O}((l-1)\vec{c})] * [K_\delta] \right) \\
 & = (q-1)[\mathcal{O}((l+1)\vec{c} + \vec{x}_i)] + \frac{q-1}{q} [\mathcal{O}(l\vec{c} + \vec{x}_i) \oplus S_{i,0}^{(p_i)}] - \frac{q-1}{q} [S_{i,1} \oplus \mathcal{O}((l-1)\vec{c})] * [K_\delta],
 \end{aligned}$$

and

$$\begin{aligned}
 & [[S_{i,1} \oplus S_{i,0}^{(p_i)}], [\mathcal{O}(l\vec{c})]]_{\mathbf{v}^{-1}} \\
 & = [S_{i,1} \oplus S_{i,0}^{(p_i)}] * [\mathcal{O}(l\vec{c})] - \mathbf{v}^{-1} [\mathcal{O}(l\vec{c})] * [S_{i,1} \oplus S_{i,0}^{(p_i)}] \\
 & = \mathbf{v}^{-2} \left( [S_{i,1} \oplus S_{i,0}^{(p_i)}] \oplus \mathcal{O}(l\vec{c}) + (q-1)[\mathcal{O}(l\vec{c} + \vec{x}_i) \oplus S_{i,0}^{(p_i)}] + (q^2 - q)[\mathcal{O}((l+1)\vec{c}) \oplus S_{i,1}] \right) \\
 & \quad - \mathbf{v}^{-1} \cdot \frac{\mathbf{v}}{q} \left( [S_{i,1} \oplus S_{i,0}^{(p_i)}] \oplus \mathcal{O}(l\vec{c}) + (q-1)[S_{i,1} \oplus \mathcal{O}((l-1)\vec{c})] * [K_\delta] \right) \\
 & = (q-1)[\mathcal{O}((l+1)\vec{c}) \oplus S_{i,1}] + \frac{q-1}{q} [\mathcal{O}(l\vec{c} + \vec{x}_i) \oplus S_{i,0}^{(p_i)}] - \frac{q-1}{q} [S_{i,1} \oplus \mathcal{O}((l-1)\vec{c})] * [K_\delta].
 \end{aligned}$$

Hence, by combining the above two formulas we obtain

$$\begin{aligned}
 & [[S_{i,1}^{(p_i+1)}] - [S_{i,1} \oplus S_{i,0}^{(p_i)}], [\mathcal{O}(l\vec{c})]]_{\mathbf{v}^{-1}} \\
 & = (q-1) \left( [\mathcal{O}((l+1)\vec{c} + \vec{x}_i)] - [\mathcal{O}((l+1)\vec{c}) \oplus S_{i,1}] \right) \\
 & = \mathbf{v} [[S_{i,1}], [\mathcal{O}((l+1)\vec{c})]]_{\mathbf{v}}.
 \end{aligned}$$

□

## 5. iHALL ALGEBRAS OF CYCLIC QUIVERS

Let  $C_n$  be the cyclic quiver with  $n$  vertices; see (4.3). Recall that  $\text{rep}_{\mathbf{k}}^{\text{nil}}(C_n)$  is the category of finite-dimensional nilpotent representations of  $C_n$ . For the Grothendieck group  $K_0(\text{rep}_{\mathbf{k}}^{\text{nil}}(C_n))$ , we denote  $\alpha_j = \widehat{S}_j$  and  $\delta = \sum_{j=1}^n \widehat{S}_j$  by abusing the notations. For any positive real root  $\beta$  of  $\widehat{\mathfrak{sl}}_n$ , by Gabriel-Kac Theorem, we denote by  $M(\beta)$  the unique (up to isomorphism)  $\mathbf{k}C_n$ -module with its class  $\beta$  in  $K_0(\text{rep}_{\mathbf{k}}^{\text{nil}}(C_n))$ .

For a finite-dimensional representation  $M$  of  $C_n$ , we denote by  $\text{soc}(M)$  and  $\text{top}(M)$  the socle and top of  $M$  respectively.

**5.1. Root vectors.** Let  $\widetilde{\mathbf{U}}^v := \widetilde{\mathbf{U}}_v^v(\widehat{\mathfrak{sl}}_n)$  be the universal affine  $i$ quantum group of type  $A_{n-1}$ , and  ${}^{\text{Dr}}\widetilde{\mathbf{U}}^v$  be its Drinfeld presentation throughout this section. By Theorem 2.6, there exists an algebra isomorphism  $\Phi : {}^{\text{Dr}}\widetilde{\mathbf{U}}^v \rightarrow \widetilde{\mathbf{U}}^v$ . We choose the sign function  $o(\cdot)$  such that  $o(j) = (-1)^j$  for any  $1 \leq j \leq n-1$  throughout this paper. On the other hand, for  $n \geq 2$ , [LW23, Theorem 9.6] gives an algebra monomorphism:

$$\widetilde{\psi}_{C_n} : \widetilde{\mathbf{U}}_{|v=\mathbf{v}}^v \longrightarrow {}^i\widetilde{\mathcal{H}}(\mathbf{k}C_n) \tag{5.1}$$

$$B_j \mapsto \frac{-1}{q-1} [S_j], \quad \mathbb{K}_j \mapsto [K_{S_j}], \quad \text{for } 0 \leq j \leq n-1. \tag{5.2}$$

Define

$$\Omega_{C_n} := \widetilde{\psi}_{C_n} \circ \Phi : {}^{\text{Dr}}\widetilde{\mathbf{U}}_{|v=\mathbf{v}}^v \longrightarrow {}^i\widetilde{\mathcal{H}}(\mathbf{k}C_n). \tag{5.3}$$

Then  $\Omega_{C_n}$  sends:

$$\mathbb{K}_j \mapsto [K_{S_j}], \quad C \mapsto [K_\delta], \quad B_{j,0} \mapsto \frac{-1}{q-1}[S_j], \quad \forall 1 \leq j \leq n-1. \quad (5.4)$$

For any  $1 \leq j \leq n-1$ ,  $l \in \mathbb{Z}$  and  $r \geq 1$ , we define

$$\widehat{B}_{j,l} := (1-q)\Omega_{C_n}(B_{j,l}), \quad \widehat{\Theta}_{j,r} := \Omega_{C_n}(\Theta_{j,r}), \quad \widehat{H}_{j,r} := \Omega_{C_n}(H_{j,r}). \quad (5.5)$$

In particular,  $\widehat{B}_{j,0} = [S_j]$  for any  $1 \leq j \leq n-1$ .

For  $1 \leq j \leq n$  and any  $\alpha \in K_0(\text{rep}_{\mathbf{k}}^{\text{nil}}(C_n))$ , set

$$\mathcal{M}_{j,\alpha} := \{[M] \mid \widehat{M} = \alpha, \text{soc}(M) \subseteq S_1 \oplus \cdots \oplus S_j\}. \quad (5.6)$$

Define

$$\pi_{j,1} = \frac{-\mathbf{v}^{-j}}{\mathbf{v} - \mathbf{v}^{-1}} \sum_{[M] \in \mathcal{M}_{j,\delta}} (-1)^{\dim \text{End}(M)} [M]. \quad (5.7)$$

Set  $\pi_{0,1} = 0$ .

**Proposition 5.1.** *For any  $1 \leq j \leq n-1$ , we have*

$$\widehat{B}_{j,-1} = \mathbf{v}^{-j+1} \sum_{[M] \in \mathcal{M}_{j+1,\delta-e_j}} (-1)^{\dim \text{End}(M)} [M] * [K_{\delta-\alpha_j}]^{-1}.$$

In order to prove Proposition 5.1, we need to recall the isomorphisms of  $\iota$ Hall algebras induced by the BGP type reflection functors introduced in [LW21a, LW22b].

Let  $Q = (Q_0, Q_1)$  be a general quiver. For any sink  $\ell \in Q_0$ , define the quiver  $\mathbf{s}_\ell^+ Q$  by reversing all the arrows of  $Q$  ending at  $\ell$ . Associated to a sink  $\ell \in Q_0$ , the BGP reflection functor induces a reflection functor (see [LW21a, §3.2]):

$$F_\ell^+ : \mathcal{C}_1(\text{rep}_{\mathbf{k}}(Q)) \longrightarrow \mathcal{C}_1(\text{rep}_{\mathbf{k}}(\mathbf{s}_\ell^+ Q)). \quad (5.8)$$

The functor  $F_\ell^+$  induces an isomorphism  $\Gamma_\ell : {}^\iota\widetilde{\mathcal{H}}(\mathbf{k}Q) \xrightarrow{\sim} {}^\iota\widetilde{\mathcal{H}}(\mathbf{k}\mathbf{s}_\ell^+ Q)$  by [LW21a, Theorem 4.3], and then a commutative diagram

$$\begin{array}{ccc} \widetilde{\mathbf{U}}_{|v=\mathbf{v}}^{\iota} & \xrightarrow{\mathbf{T}_\ell} & \widetilde{\mathbf{U}}_{|v=\mathbf{v}}^{\iota} \\ \downarrow \widetilde{\psi}_Q & & \downarrow \widetilde{\psi}_{\mathbf{s}_\ell^+ Q} \\ {}^\iota\widetilde{\mathcal{H}}(\mathbf{k}Q) & \xrightarrow{\Gamma_\ell} & {}^\iota\widetilde{\mathcal{H}}(\mathbf{k}\mathbf{s}_\ell^+ Q), \end{array} \quad (5.9)$$

where  $\mathbf{T}_\ell$  is defined in Lemma 2.3.

Dually, we can define  $\mathbf{s}_\ell^- Q$  for any source  $\ell$  by reversing all the arrows of  $Q$  starting from  $\ell$ , and a reflection functor  $F_\ell^-$  associated to a source  $\ell \in Q_0$  is also defined. This functor  $F_\ell^-$  also induces an isomorphism  $\Gamma_\ell^- : {}^\iota\widetilde{\mathcal{H}}(\mathbf{k}Q) \xrightarrow{\sim} {}^\iota\widetilde{\mathcal{H}}(\mathbf{k}\mathbf{s}_\ell^- Q)$ . In particular, the action of  $F_\ell^+$  (and  $F_\ell^-$ ) on  $\text{rep}_{\mathbf{k}}(Q) \subseteq \mathcal{C}_1(\text{rep}_{\mathbf{k}}(Q))$  is the same as that of the classic BGP reflection functor.

A sequence  $i_1, \dots, i_t$  in  $Q_0$  is called  $\iota$ -admissible if  $i_r$  is a sink in  $\mathbf{s}_{i_{r-1}} \cdots \mathbf{s}_{i_1}(Q)$ , for each  $1 \leq r \leq t$ .

*Proof of Proposition 5.1.* Fix  $1 \leq j \leq n-1$ . We assume  $Q(j)$  is the following quiver:

$$(5.10)$$

Recall that

$$\omega_j = \sigma^j(s_{n-j} \cdots s_1)(s_{n-j+1} \cdots s_2) \cdots (s_{n-1} \cdots s_j), \quad (5.11)$$

where  $\sigma$  is the automorphism of the underlying graph of  $C_n$  such that  $\sigma(i) = i + 1$ ; see e.g. [DJX12, §6.2]. Note that  $\mathbf{s}_{\omega_j}(Q(j)) = Q(j)$ , and  $j, \dots, n-1, \dots, 2, \dots, n-j+1, 1, \dots, n-j$  is an  $\iota$ -admissible sequence. Then [LW21a, Corollary 3.10, Proposition 4.4] show that

$$\Gamma_{\omega_j}([S_j]) = [M(\delta - \alpha_j)] * [K_{\delta - \alpha_j}]^{-1}, \quad (5.12)$$

where  $M(\delta - \alpha_j)$  is the unique (up to isomorphism)  $\mathbf{k}Q(j)$ -module with  $\delta - \alpha_j$  as its class in  $K_0(\mathbf{k}Q(j))$ .

Let  $N$  be the unique (up to isomorphism)  $\mathbf{k}Q(j)$ -module with  $\widehat{S}_0 + \widehat{S}_{n-1} + \cdots + \widehat{S}_{j+1}$  as its class in  $K_0(\mathbf{k}Q(j))$ . A direct calculation in  ${}^v\mathcal{H}(\mathbf{k}Q(j))$  shows that

$$[M(\delta - \alpha_j)] = \frac{(-1)^{j-1}}{(q-1)^{j-1}} [[S_{j-1}], [S_{j-2}], \dots, [S_1], [N]]_{\mathbf{v}}.$$

Then the commutative diagram (5.9) yields that

$$\begin{aligned} \widetilde{\psi}_{Q(j)}((-1)^j \mathbf{T}_{\omega_j}(B_j)) &= (-1)^j \cdot \frac{-1}{q-1} \Gamma_{\omega_j}([S_j]) \\ &= \frac{1}{(q-1)^j} [[S_{j-1}], [S_{j-2}], \dots, [S_1], [N]]_{\mathbf{v}} * [K_{\delta - \alpha_j}]^{-1}. \end{aligned} \quad (5.13)$$

It is important to note that  $N$  can be viewed as a  $\mathbf{k}C_n$ -module. Together with [LW21a, Lemma 6.3],  $\widetilde{\psi}_{C_n}((-1)^j \mathbf{T}_{\omega_j}(B_j))$  has the same expression as in (5.13). Therefore, by (2.24), a direct computation in  ${}^v\mathcal{H}(\mathbf{k}C_n)$  shows that

$$\begin{aligned} \widehat{B}_{j,-1} &= (1-q) \widetilde{\psi}_{C_n}((-1)^j \mathbf{T}_{\omega_j}(B_j)) \\ &= \frac{-1}{(q-1)^{j-1}} [[S_{j-1}], [S_{j-2}], \dots, [S_1], [N]]_{\mathbf{v}} * [K_{\delta - \alpha_j}]^{-1} \\ &= \mathbf{v}^{-j+1} \sum_{\mathcal{M}_{j+1, \delta - e_j}} (-1)^{\dim \text{End}(M)} [M] * [K_{\delta - \alpha_j}]^{-1}. \end{aligned}$$

□

**Corollary 5.2.** *For any  $1 \leq j \leq n-1$ , we have*

$$\widehat{H}_{j,1} = \widehat{\Theta}_{j,1} = \pi_{j+1,1} - (\mathbf{v} + \mathbf{v}^{-1})\pi_{j,1} + \pi_{j-1,1}.$$

*Proof.* Let

$$\begin{aligned} \mathcal{M}_{j,\delta}^j &:= \{[M] \in \mathcal{M}_{j,\delta} \mid S_j \text{ is a direct summand of } \text{soc}(M)\}, \\ \mathcal{N}_{j+1,\delta}^j &:= \{[M] \in \mathcal{M}_{j+1,\delta} \mid S_j \text{ is not a direct summand of } \text{soc}(M)\}. \end{aligned}$$

Then we have the following disjoint unions

$$\mathcal{M}_{j,\delta} = \mathcal{M}_{j,\delta}^j \sqcup \mathcal{M}_{j-1,\delta}, \quad (5.14)$$

$$\mathcal{M}_{j+1,\delta}^{j+1} = \mathcal{N}_{j+1,\delta}^j \sqcup \{[M \oplus S_j] \mid [M] \in \mathcal{M}_{j+1,\delta-e_j}\}. \quad (5.15)$$

Recall that  $\Phi(\Theta_{j,1}) = (-B_j B_{j,-1} + v^2 B_{j,-1} B_j) \mathbb{K}_{\delta-\alpha_j}$  in  $\tilde{\mathbf{U}}^i$ . By applying  $\Omega_{C_n}$  and using Proposition 5.1, we have

$$\begin{aligned} \widehat{\Theta}_{j,1} &= \frac{1}{(q-1)^2} \left( -\widehat{B}_{j,0} * \widehat{B}_{j,-1} + \mathbf{v}^2 \widehat{B}_{j,-1} * \widehat{B}_{j,0} \right) * [K_{\delta-\alpha_j}] \\ &= \frac{-\mathbf{v}^{-j+1}}{(q-1)^2} \sum_{[M] \in \mathcal{M}_{j+1,\delta-e_j}} (-1)^{\dim \text{End}(M)} \left[ [S_j], [M] \right]_{\mathbf{v}^2} \\ &= \frac{\mathbf{v}^{-j}}{(q-1)^2} \sum_{[M] \in \mathcal{M}_{j+1,\delta-e_j}} (-1)^{\dim \text{End}(M \oplus S_j)} [M \oplus S_j] - \frac{\mathbf{v}^{-j}}{q-1} \sum_{[N] \in \mathcal{N}_{j+1,\delta}^j} (-1)^{\dim \text{End}(N)} [N] \\ &\quad - \frac{\mathbf{v}^{-j+2}}{(q-1)^2} \sum_{[M] \in \mathcal{M}_{j+1,\delta-e_j}} (-1)^{\dim \text{End}(M \oplus S_j)} [M \oplus S_j] + \frac{\mathbf{v}^{-j+2}}{q-1} \sum_{[L] \in \mathcal{M}_{j,\delta}^j} (-1)^{\dim \text{End}(L)} [L] \\ &= -\frac{\mathbf{v}^{-j-1}}{\mathbf{v} - \mathbf{v}^{-1}} \sum_{[M] \in \mathcal{M}_{j+1,\delta-e_j}} (-1)^{\dim \text{End}(M \oplus S_j)} [M \oplus S_j] \\ &\quad - \frac{\mathbf{v}^{-j-1}}{\mathbf{v} - \mathbf{v}^{-1}} \sum_{[N] \in \mathcal{N}_{j+1,\delta}^j} (-1)^{\dim \text{End}(N)} [N] + \frac{\mathbf{v}^{-j+1}}{\mathbf{v} - \mathbf{v}^{-1}} \sum_{[L] \in \mathcal{M}_{j,\delta}^j} (-1)^{\dim \text{End}(L)} [L] \\ &\stackrel{(5.15)}{=} -\frac{\mathbf{v}^{-j-1}}{\mathbf{v} - \mathbf{v}^{-1}} \sum_{[M] \in \mathcal{M}_{j+1,\delta}^{j+1}} (-1)^{\dim \text{End}(M)} [M] + \frac{\mathbf{v}^{-j+1}}{\mathbf{v} - \mathbf{v}^{-1}} \sum_{[L] \in \mathcal{M}_{j,\delta}^j} (-1)^{\dim \text{End}(L)} [L] \\ &\stackrel{(5.14)}{=} -\frac{\mathbf{v}^{-j-1}}{\mathbf{v} - \mathbf{v}^{-1}} \left( \sum_{[M] \in \mathcal{M}_{j+1,\delta}} (-1)^{\dim \text{End}(M)} [M] - \sum_{[M] \in \mathcal{M}_{j,\delta}} (-1)^{\dim \text{End}(M)} [M] \right) \\ &\quad + \frac{\mathbf{v}^{-j+1}}{\mathbf{v} - \mathbf{v}^{-1}} \left( \sum_{[L] \in \mathcal{M}_{j,\delta}} (-1)^{\dim \text{End}(L)} [L] - \sum_{[L] \in \mathcal{M}_{j-1,\delta}} (-1)^{\dim \text{End}(L)} [L] \right) \\ &= \pi_{j+1,1} - (\mathbf{v} + \mathbf{v}^{-1}) \pi_{j,1} + \pi_{j-1,1}. \end{aligned}$$

□

**Corollary 5.3.** *We have*

$$\widehat{\Theta}_{1,1} = \frac{1}{q-1} \left( \mathbf{v}^{-1} [S_1^{(n)}] - \mathbf{v}^{-1} [S_0^{(n-1)} \oplus S_1] - \mathbf{v} [S_0^{(n)}] \right), \quad (5.16)$$

$$\widehat{B}_{1,-1} = -[S_0^{(n-1)}] * [K_{\alpha_1-\delta}], \quad (5.17)$$

$$\widehat{B}_{1,1} = \frac{1}{q} [S_1^{(n+1)}] - \frac{1}{q} [S_1 \oplus S_0^{(n)}], \quad (5.18)$$

$$\widehat{B}_{1,-2} = \left( -\frac{1}{q} [S_0^{(2n-1)}] + \frac{1}{q} [S_0^{(n)} \oplus S_0^{(n-1)}] \right) * [K_{\alpha_1-2\delta}]. \quad (5.19)$$



*Proof.* (5.16)–(5.17) follow from Corollary 5.2 and Proposition 5.1.

For (5.18), a direct computation shows that

$$\begin{aligned} [[S_0^{(n-1)} \oplus S_1], [S_1]] &= \frac{q-1}{q}[S_0^{(n)} \oplus S_1] - \frac{q-1}{q}[S_1^{(n)} \oplus S_1]; \\ [[S_1^{(n)}], [S_1]] &= \frac{1-q}{q}[S_1^{(n)} \oplus S_1] + \frac{q-1}{q}[S_1^{(n+1)}] + \frac{q(q-1)}{q}[S_0^{(n-1)}] * [K_{S_1}]; \\ [[S_0^{(n)}], [S_1]] &= \frac{q-1}{q}[S_0^{(n)} \oplus S_1] - \frac{q-1}{q}[S_1^{(n+1)}] - \frac{q(q-1)}{q}[S_0^{(n-1)}] * [K_{S_1}]. \end{aligned}$$

It follows that

$$\begin{aligned} [\widehat{\Theta}_{1,1}, \widehat{B}_{1,0}] &= \frac{1}{q-1}[\mathbf{v}^{-1}[S_1^{(n)}] - \mathbf{v}^{-1}[S_0^{(n-1)} \oplus S_1] - \mathbf{v}[S_0^{(n)}], [S_1]] \\ &= \frac{[2]_{\mathbf{v}}}{q}[S_1^{(n+1)}] - \frac{[2]_{\mathbf{v}}}{q}[S_1 \oplus S_0^{(n)}] + [2]_{\mathbf{v}}[S_0^{(n-1)}] * [K_{S_1}]. \end{aligned}$$

By (2.14) we have

$$\begin{aligned} \widehat{B}_{1,1} &= \frac{1}{[2]_{\mathbf{v}}}[\widehat{\Theta}_{1,1}, \widehat{B}_{1,0}] + \widehat{B}_{1,-1} * [K_{\delta}] \\ &= \frac{1}{[2]_{\mathbf{v}}} \cdot \left( \frac{[2]_{\mathbf{v}}}{q}[S_1^{(n+1)}] - \frac{[2]_{\mathbf{v}}}{q}[S_1 \oplus S_0^{(n)}] + [2]_{\mathbf{v}}[S_0^{(n-1)}] * [K_{S_1}] \right) - [S_0^{(n-1)}] * [K_{S_1}] \\ &= \frac{1}{q}[S_1^{(n+1)}] - \frac{1}{q}[S_1 \oplus S_0^{(n)}]. \end{aligned}$$

For (5.19), a direct computation shows that

$$\begin{aligned} [[S_0^{(n-1)} \oplus S_1], [S_0^{(n-1)}]] &= \frac{q-1}{q}[S_1^{(n)} \oplus S_0^{(n-1)}] - \frac{q-1}{q}[S_0^{(n)} \oplus S_0^{(n-1)}]; \\ [[S_1^{(n)}], [S_0^{(n-1)}]] &= \frac{q-1}{q}[S_1^{(n)} \oplus S_0^{(n-1)}] - \frac{q-1}{q}[S_0^{(2n-1)}] - \frac{q(q-1)}{q}[S_1] * [K_{S_0^{(n-1)}}]; \\ [[S_0^{(n)}], [S_0^{(n-1)}]] &= \frac{1-q}{q}[S_0^{(n)} \oplus S_0^{(n-1)}] + \frac{q-1}{q}[S_0^{(2n-1)}] + \frac{q(q-1)}{q}[S_1] * [K_{S_0^{(n-1)}}]. \end{aligned}$$

It follows that

$$[\widehat{\Theta}_{1,1}, \widehat{B}_{1,-1}] = \frac{[2]_{\mathbf{v}}}{q}[S_0^{(2n-1)}] * [K_{\alpha_1 - \delta}] - \frac{[2]_{\mathbf{v}}}{q}[S_0^{(n)} \oplus S_0^{(n-1)}] * [K_{\alpha_1 - \delta}] + [2]_{\mathbf{v}}[S_1].$$

Then by (2.14) we have

$$\begin{aligned} \widehat{B}_{1,-2} &= -\frac{1}{[2]_{\mathbf{v}}}[\widehat{\Theta}_{1,1}, \widehat{B}_{1,-1}] * [K_{-\delta}] + \widehat{B}_{1,0} * [K_{-\delta}] \\ &= \left( -\frac{1}{q}[S_0^{(2n-1)}] + \frac{1}{q}[S_0^{(n)} \oplus S_0^{(n-1)}] \right) * [K_{\alpha_1 - 2\delta}]. \end{aligned}$$

□

The following lemma will be used in the proof of Lemma A.5. It is understood that the terms involving  $S^{(r)}$  vanish if  $r < 0$ .

**Lemma 5.4.** *For any  $r \geq 0$ , we have*

$$\begin{aligned} & [[S_0^{(rn)}], \widehat{B}_{1,-1}] + q[[S_0^{((r-2)n)}], \widehat{B}_{1,-1}] * [K_\delta] \\ &= [[S_0^{((r-1)n)}], \widehat{B}_{1,0}]_{\mathbf{v}^2} + q[[S_0^{((r-1)n)}], \widehat{B}_{1,-2}]_{\mathbf{v}^{-2}} * [K_\delta]. \end{aligned} \quad (5.20)$$

*Proof.* By Corollary 5.3, we only need to check that

$$\begin{aligned} & [[S_0^{(rn)}], [S_0^{(n-1)}]] + q[[S_0^{((r-2)n)}], [S_0^{(n-1)}]] * [K_\delta] \\ &= q[[S_1], [S_0^{((r-1)n)}]]_{\mathbf{v}^{-2}} * [K_{\delta-\alpha_{11}}] + [[S_0^{((r-1)n)}], [S_0^{(2n-1)}] - [S_0^{(n)} \oplus S_0^{(n-1)}]]_{\mathbf{v}^{-2}}. \end{aligned} \quad (5.21)$$

For any  $r \geq 0$ , an easy computation shows that

$$\begin{aligned} [S_0^{(n-1)}] * [S_0^{(rn)}] &= [S_0^{(rn)} \oplus S_0^{(n-1)}], \\ [S_0^{(rn)}] * [S_0^{(n-1)}] &= \frac{1}{q}[S_0^{(rn)} \oplus S_0^{(n-1)}] + \frac{q-1}{q}[S_0^{((r+1)n-1)}] + (q-1)[S_1^{((r-1)n+1)}] * [K_{\delta-\alpha_1}]; \\ [S_0^{(rn)}] * [S_1] &= [S_0^{(rn)} \oplus S_1], \\ [S_1] * [S_0^{(rn)}] &= \frac{1}{q}[S_0^{(rn)} \oplus S_1] + \frac{q-1}{q}[S_1^{(rn+1)}] + (q-1)[S_0^{(rn-1)}] * [K_{\alpha_1}]. \end{aligned}$$

Then we obtain

$$[[S_1], [S_0^{(rn)}]]_{\mathbf{v}^{-2}} = \frac{q-1}{q}[S_1^{(rn+1)}] + (q-1)[S_0^{(rn-1)}] * [K_{\alpha_1}],$$

and

$$[[S_0^{(rn)}], [S_0^{(n-1)}]] = \frac{q-1}{q} \left( [S_0^{((r+1)n-1)}] - [S_0^{(rn)} \oplus S_0^{(n-1)}] + q[S_1^{((r-1)n+1)}] * [K_{\delta-\alpha_1}] \right).$$

On the other hand, for  $r \geq 1$ , by Proposition 3.9, we have

$$\begin{aligned} & [S_0^{(2n-1)}] * [S_0^{(rn)}] \\ &= \frac{1}{q}[S_0^{(rn)} \oplus S_0^{(2n-1)}] + \frac{q-1}{q}[S_0^{((r+1)n)} \oplus S_0^{(n-1)}] + \frac{q(q-1)}{q}[S_0^{((r-1)n)} \oplus S_0^{(n-1)}] * [K_\delta], \\ & [S_0^{(rn)}] * [S_0^{(2n-1)}] \\ &= \frac{1}{q^2}[S_0^{(rn)} \oplus S_0^{(2n-1)}] + \frac{q-1}{q^2}[S_0^{((r+1)n-1)} \oplus S_0^{(n)}] + \frac{q(q-1)}{q^2}[S_0^{((r+2)n-1)}] \\ & \quad + \frac{q-1}{q}[S_1^{((r-1)n+1)} \oplus S_0^{(n)}] * [K_{S_0^{(n-1)}}] + \frac{(q-1)^2}{q}[S_1^{(rn+1)}] * [K_{S_0^{(n-1)}}] \\ & \quad + q(q-1)[S_1^{((r-2)n+1)}] * [K_{S_0^{(2n-1)}}]; \end{aligned}$$

and

$$\begin{aligned} & [S_0^{(n)} \oplus S_0^{(n-1)}] * [S_0^{(rn)}] \\ &= \frac{1}{q}[S_0^{(n)} \oplus S_0^{(n-1)} \oplus S_0^{(rn)}] + \frac{q-1}{q}[S_0^{((r+1)n)} \oplus S_0^{(n-1)}] + \frac{q(q-1)}{q}[S_0^{((r-1)n)} \oplus S_0^{(n-1)}] * [K_\delta], \\ & [S_0^{(rn)}] * [S_0^{(n)} \oplus S_0^{(n-1)}] \end{aligned}$$

$$\begin{aligned}
 &= \frac{1}{q^2}[S_0^{(rn)} \oplus S_0^{(n)} \oplus S_0^{(n-1)}] + \frac{q-1}{q^2}[S_0^{((r+1)n-1)} \oplus S_0^{(n)}] + \frac{q(q-1)}{q^2}[S_0^{((r+1)n)} \oplus S_0^{(n-1)}] \\
 &\quad + \frac{q-1}{q}[S_1^{((r-1)n+1)} \oplus S_0^{(n)}] * [K_{S_0^{(n-1)}}] + \frac{(q-1)^2}{q}[S_1^{(rn+1)}] * [K_{S_0^{(n-1)}}] \\
 &\quad + \frac{q(q-1)}{q}[S_0^{((r-1)n)} \oplus S_0^{(n-1)}] * [K_\delta] + \frac{q(q-1)^2}{q}[S_0^{(rn-1)}] * [K_\delta].
 \end{aligned}$$

By combining the above formulas, we obtain

$$\begin{aligned}
 &[[S_0^{(rn)}], [S_0^{(2n-1)}]]_{\mathbf{v}^{-2}} - [[S_0^{(rn)}], [S_0^{(n)} \oplus S_0^{(n-1)}]]_{\mathbf{v}^{-2}} \\
 &= \frac{q-1}{q}[S_0^{((r+2)n-1)}] - \frac{q-1}{q}[S_0^{((r+1)n)} \oplus S_0^{(n-1)}] + q(q-1)[S_1^{((r-2)n+1)}] * [K_{S_0^{(2n-1)}}] \\
 &\quad - (q-1)[S_0^{((r-1)n)} \oplus S_0^{(n-1)}] * [K_\delta] - \frac{q(q-1)^2}{q}[S_0^{(rn-1)}] * [K_\delta].
 \end{aligned}$$

In particular, for  $r = 0$ , we have

$$\left[ [S_0^{(rn)}], [S_0^{(2n-1)}] - [S_0^{(n)} \oplus S_0^{(n-1)}] \right]_{\mathbf{v}^{-2}} = (1 - \mathbf{v}^{-2}) \left( [S_0^{(2n-1)}] - [S_0^{(n)} \oplus S_0^{(n-1)}] \right)$$

since  $S_0^{(0)} = 0$ . Then (5.21) follows since both sides are equal to

$$\begin{aligned}
 &\frac{q-1}{q} \left( [S_0^{((r+1)n-1)}] - [S_0^{(rn)} \oplus S_0^{(n-1)}] + q[S_1^{((r-1)n+1)}] * [K_{S_0^{(n-1)}}] \right) \\
 &\quad + (q-1) \left( [S_0^{((r-1)n-1)}] - [S_0^{((r-2)n)} \oplus S_0^{(n-1)}] + q[S_1^{((r-3)n+1)}] * [K_{S_0^{(n-1)}}] \right) * [K_\delta].
 \end{aligned}$$

□

**5.2. Another description of root vectors.** It is helpful and interesting to compute all root vectors clearly in  ${}^i\mathcal{H}(\mathbf{k}C_n)$ . In particular, the root vectors at the vertex 1 are described in Subsection 9.1. There is another way to interpret the root vectors in  ${}^i$ Hall algebras, which we give in the following.

Let  $Q$  be a quiver with its underlying graph the same as  $C_n$ . Then there exists an isomorphism  $\mathrm{FT}_{C_n, Q} : {}^i\tilde{\mathcal{C}}(\mathbf{k}Q) \xrightarrow{\cong} {}^i\tilde{\mathcal{C}}(\mathbf{k}C_n)$  such that  $\mathrm{FT}_{C_n, Q}([S_i]) = [S_i]$ ,  $\mathrm{FT}_{C_n, Q}([K_{S_i}]) = [K_{S_i}]$  for any  $i \in \mathbb{I}$ ; cf. (3.10). For any positive real root  $\beta$  of  $\widehat{\mathfrak{sl}}_n$ , we denote by  $M(\beta)$  the unique (up to isomorphism)  $\mathbf{k}Q$ -module with its class  $\beta$  in  $K_0(\mathrm{mod}(\mathbf{k}Q))$ .

For any hereditary abelian category  $\mathcal{A}$  and a full subcategory  $\mathcal{S}$ , recall from [GL87] that the right (resp. left) perpendicular category  $\mathcal{S}^\perp$  (resp.  ${}^\perp\mathcal{S}$ ) is defined as follows:

$$\mathcal{S}^\perp := \{X \in \mathcal{A} \mid \mathrm{Hom}_{\mathcal{A}}(C, X) = 0 = \mathrm{Ext}_{\mathcal{A}}^1(C, X) \text{ for any } C \in \mathcal{S}\}; \quad (5.22)$$

$${}^\perp\mathcal{S} := \{X \in \mathcal{A} \mid \mathrm{Hom}_{\mathcal{A}}(X, C) = 0 = \mathrm{Ext}_{\mathcal{A}}^1(X, C) \text{ for any } C \in \mathcal{S}\}. \quad (5.23)$$

Then the category  $\mathcal{S}^\perp$  (resp.  ${}^\perp\mathcal{S}$ ) is a hereditary abelian category, which is extension-closed in  $\mathcal{A}$ ; see [BS13, Proposition 3.2]. If  $\mathcal{S} = \{M\}$  for an object  $M \in \mathcal{A}$ , then we also denote  $M^\perp := \mathcal{S}^\perp$  and  ${}^\perp M := {}^\perp\mathcal{S}$ .

Recall  $Q(j)$  in (5.10).

**Proposition 5.5.** *For any  $l > 0$ , we have*

$$\widehat{B}_{j,-l} = (-1)^{jl} \mathbf{FT}_{C_n, Q(j)}([M(l\delta - \alpha_j)]) * [K_{l\delta - \alpha_j}]^{-1}, \quad (5.24)$$

$$\widehat{B}_{j,l} = (-1)^{jl} \mathbf{FT}_{C_n, Q(j)}([M(l\delta + \alpha_j)]), \quad (5.25)$$

$$\widehat{\Theta}_{j,l} = \frac{(-1)^{jl}}{(q-1)^{2\mathbf{v}^{l-1}}} \mathbf{FT}_{C_n, Q(j)} \left( \sum_{0 \neq f: S_j \rightarrow M(l\delta + \alpha_j)} [\text{Coker}(f)] \right). \quad (5.26)$$

*Proof.* From the proof of Proposition 5.1, in  ${}^i\widetilde{\mathcal{H}}(\mathbf{k}Q(j))$  we have

$$\Gamma_{\omega_j}^l([S_j]) = [M(l\delta - \alpha_j)] * [K_{l\delta - \alpha_j}]^{-1}.$$

Then  $B_{j,-l}$  satisfies that  $\Phi(B_{j,-l}) = (-1)^{jl} \mathbf{T}_{\omega_j}^l(B_j)$  and so

$$\widetilde{\psi}_{Q(j)} \circ \Phi(B_{j,-l}) = (-1)^{jl} \cdot \frac{-1}{q-1} [M(l\delta - \alpha_j)] * [K_{l\delta - \alpha_j}]^{-1}$$

in  ${}^i\widetilde{\mathcal{H}}(\mathbf{k}Q(j))$ . Similarly, one can see that

$$\widetilde{\psi}_{Q(j)} \circ \Phi(B_{j,l}) = (-1)^{jl} \cdot \frac{-1}{q-1} [M(l\delta + \alpha_j)];$$

cf. [LRW23, Lemma 5.5]. So we have (5.24)–(5.25).

For (5.26), let  $\mathcal{S} := \{S_i \in \text{mod}(\mathbf{k}Q(j)) \mid i \neq 0, j\}$ . Then the subcategory  ${}^\perp\mathcal{S}$  is equivalent to  $\text{mod}(\mathbf{k}Q_{\text{kr}})$ , where  $Q_{\text{kr}}$  is the Kronecker quiver  $0 \rightrightarrows 1$ . So there exists an embedding functor  $\mathbb{F} : \text{mod}(\mathbf{k}Q_{\text{kr}}) \simeq {}^\perp\mathcal{S} \rightarrow \text{mod}(\mathbf{k}Q(j))$ . In particular,  $\mathbb{F}(S_1) = S_j$ , and  $\mathbb{F}(M(l\alpha_0 + (l+1)\alpha_1)) = M(l\delta + \alpha_j)$  for any  $l \geq 0$ . So  $\mathbb{F}$  induces an embedding of algebras  $\widetilde{\mathcal{H}}(\mathcal{C}_1(\text{mod}(\mathbf{k}Q_{\text{kr}}))) \rightarrow \widetilde{\mathcal{H}}(\mathcal{C}_1(\text{mod}(\mathbf{k}Q(j))))$ , and then an embedding of algebras  $F : {}^i\widetilde{\mathcal{H}}(\mathbf{k}Q_{\text{kr}}) \rightarrow {}^i\widetilde{\mathcal{H}}(\mathbf{k}Q(j))$ . Using [LRW23, Theorem 5.11], in order to compute  $\widehat{\Theta}_{j,l}$  in  ${}^i\widetilde{\mathcal{H}}(\mathbf{k}Q_{\text{kr}})$ , it is equivalent to compute them in  ${}^i\widetilde{\mathcal{H}}(\mathbb{P}_{\mathbf{k}}^1)$ . By [LRW23, (4.2)], we have

$$\widehat{\Theta}_{1,l} = \frac{1}{(q-1)^{2\mathbf{v}^{l-1}}} \sum_{0 \neq f: S_1 \rightarrow M(l\alpha_0 + (l+1)\alpha_1)} [\text{Coker}(f)]$$

in  ${}^i\widetilde{\mathcal{H}}(\mathbf{k}Q_{\text{kr}})$ . By applying  $F$ , we have

$$\widehat{\Theta}_{j,l} = \frac{(-1)^{jl}}{(q-1)^{2\mathbf{v}^{l-1}}} \sum_{0 \neq f: S_j \rightarrow M(l\delta + \alpha_j)} [\text{Coker}(f)] \quad (5.27)$$

in  ${}^i\widetilde{\mathcal{H}}(\mathbf{k}Q(j))$ . Here the scalar  $(-1)^{jl}$  comes from (5.24)–(5.25). Then (5.26) follows.  $\square$

**5.3. Two useful formulas.** Return to the cyclic quiver  $C_n$ . Let  $\mathcal{S} = \{S_j \mid 2 \leq j \leq n-1\}$ . Observe that any indecomposable object in  ${}^\perp\mathcal{S}$  has the form  $S_0^{(an)}$ ,  $S_1^{(an)}$ ,  $S_0^{(an-1)}$  or  $S_1^{(bp+1)}$ , for some  $a \geq 1$  and  $b \geq 0$ . In particular,  ${}^\perp\mathcal{S}$  is a rank-two tube with two simples  $S_1$  and  $S_0^{(n-1)}$ .

For any nilpotent  $\mathbf{k}C_n$ -module  $N$ , let  $\ell(N)$  be the number of its indecomposable direct summands. In particular,  $\ell(N) = 1$  if and only if  $N$  is indecomposable.

The following two lemmas will be used to prove Lemma 9.4; see Appendix B.4.

**Lemma 5.6.** *Assume  $a \geq 1$  and  $M \in {}^\perp \mathcal{S}$ . We have*

$$\sum_{[N]; \ell(N) = \ell(M) + 1} |\mathrm{Ext}^1(M, S_0^{(an-1)})_N| = \frac{|\mathrm{Hom}(M, S_0^{(an-1)})|}{|\mathrm{Hom}(M, S_0^{(n-1)})|}. \quad (5.28)$$

*Proof.* We prove (5.28) by induction on  $a$ . If  $a = 1$ , then by  $M \in {}^\perp \mathcal{S}$  and  $S_0^{(n-1)}$  is simple in  ${}^\perp \mathcal{S}$ , we see that any extension  $0 \rightarrow S_0^{(n-1)} \rightarrow N \rightarrow M \rightarrow 0$  with  $\ell(N) = \ell(M) + 1$  must be split. Then the result holds by noting

$$\sum_{[N]; \ell(N) = \ell(M) + 1} |\mathrm{Ext}^1(M, S_0^{(n-1)})_N| = 1.$$

For  $a > 1$ , we fix a short exact sequence  $0 \rightarrow S_1^{(n)} \xrightarrow{f_1} S_0^{(an-1)} \xrightarrow{f_2} S_0^{((a-1)n-1)} \rightarrow 0$  in  $\mathrm{rep}_k^{\mathrm{hfil}}(C_n)$ . By applying  $\mathrm{Hom}(M, -)$ , we have the following long exact sequence

$$\begin{aligned} 0 \longrightarrow \mathrm{Hom}(M, S_1^{(n)}) \longrightarrow \mathrm{Hom}(M, S_0^{(an-1)}) \longrightarrow \mathrm{Hom}(M, S_0^{((a-1)n-1)}) \longrightarrow \\ \mathrm{Ext}^1(M, S_1^{(n)}) \longrightarrow \mathrm{Ext}^1(M, S_0^{(an-1)}) \xrightarrow{\phi} \mathrm{Ext}^1(M, S_0^{((a-1)n-1)}) \longrightarrow 0, \end{aligned} \quad (5.29)$$

where the action of  $\phi$  is induced by  $f_2$  via the following pushout diagram

$$\begin{array}{ccccc} S_1^{(n)} & \xrightarrow{f_1} & S_0^{(an-1)} & \xrightarrow{f_2} & S_0^{((a-1)n-1)} \\ \parallel & & \downarrow h_1 & & \downarrow g_1 \\ S_1^{(n)} & \xrightarrow{l_1} & N & \xrightarrow{l_2} & L \\ & & \downarrow h_2 & & \downarrow g_2 \\ & & M & \xlongequal{\quad} & M. \end{array}$$

We claim that  $\ell(N) = \ell(L)$ , or equivalently,  $\ell(N) \neq \ell(L) + 1$ . Suppose for a contradiction that  $\ell(N) = \ell(L) + 1$ . In this case, the short exact sequence in the second row can not be split by observing the one in the first row is not split. Together with the assumption  $\ell(N) = \ell(L) + 1$ ,  $l_1$  should have a component  $S_1^{(n)} \rightarrow X$ , for some indecomposable direct summand  $X$  of  $N$ . Recall that  ${}^\perp \mathcal{S}$  is a rank-two tube with two simples  $S_1$  and  $S_0^{(n-1)}$ . It follows from  $N \in {}^\perp \mathcal{S}$  that  $X = S_1$ . By observing  $\mathrm{Hom}(S_0^{(an-1)}, S_1) = 0$ , the above commutative diagram does not exist, giving a contradiction. So the claim holds.

Therefore,

$$\begin{aligned} & \sum_{[N]; \ell(N) = \ell(M) + 1} |\mathrm{Ext}^1(M, S_0^{(an-1)})_N| \\ &= \sum_{[L]; \ell(L) = \ell(M) + 1} |\phi^{-1}(\mathrm{Ext}^1(M, S_0^{((a-1)n-1)})_L)| \\ &= \sum_{[L]; \ell(L) = \ell(M) + 1} |\mathrm{Ext}^1(M, S_0^{((a-1)n-1)})_L| \cdot |\mathrm{Ker}(\phi)|. \end{aligned}$$

By (5.29) we obtain that

$$\begin{aligned} |\mathrm{Ker}(\phi)| &= \frac{|\mathrm{Ext}^1(M, S_1^{(n)})| \cdot |\mathrm{Hom}(M, S_0^{(an-1)})|}{|\mathrm{Hom}(M, S_1^{(n)})| \cdot |\mathrm{Hom}(M, S_0^{((a-1)n-1)})|} \\ &= \frac{|\mathrm{Hom}(M, S_0^{(an-1)})|}{|\mathrm{Hom}(M, S_0^{((a-1)n-1)})|}. \end{aligned}$$

Now by induction, we have

$$\begin{aligned} &\sum_{[N]; \ell(N)=\ell(M)+1} |\mathrm{Ext}^1(M, S_0^{(an-1)})_N| \\ &= \frac{|\mathrm{Hom}(M, S_0^{((a-1)n-1)})|}{|\mathrm{Hom}(M, S_0^{(n-1)})|} \cdot \frac{|\mathrm{Hom}(M, S_0^{(an-1)})|}{|\mathrm{Hom}(M, S_0^{((a-1)n-1)})|} = \frac{|\mathrm{Hom}(M, S_0^{(an-1)})|}{|\mathrm{Hom}(M, S_0^{(n-1)})|}. \end{aligned}$$

□

**Lemma 5.7.** *Assume  $a \geq 1$  and  $M \in {}^\perp \mathcal{S}$ . We have*

$$\sum_{[N]; \ell(N)=\ell(M)+1} |\mathrm{Ext}^1(M, S_0^{(an)})_N| = \frac{|\mathrm{Hom}(M, S_0^{(an)})|}{|\mathrm{Ext}^1(M, S_0^{(n-1)})|}. \quad (5.30)$$

*Proof.* We fix a short exact sequence  $0 \rightarrow S_1 \xrightarrow{f_1} S_0^{(an)} \xrightarrow{f_2} S_0^{(an-1)} \rightarrow 0$  in  $\mathrm{rep}_{\mathbf{k}}^{\mathrm{nil}}(C_n)$ . By applying  $\mathrm{Hom}(M, -)$ , we have the following long exact sequence

$$\begin{aligned} 0 \longrightarrow \mathrm{Hom}(M, S_1) \longrightarrow \mathrm{Hom}(M, S_0^{(an)}) \longrightarrow \mathrm{Hom}(M, S_0^{(an-1)}) \longrightarrow \\ \mathrm{Ext}^1(M, S_1) \longrightarrow \mathrm{Ext}^1(M, S_0^{(an)}) \xrightarrow{\phi} \mathrm{Ext}^1(M, S_0^{(an-1)}) \longrightarrow 0, \end{aligned} \quad (5.31)$$

where the action of  $\phi$  is induced by  $f_2$  via the following pushout diagram

$$\begin{array}{ccccc} S_1 & \xrightarrow{f_1} & S_0^{(an)} & \xrightarrow{f_2} & S_0^{(an-1)} \\ \parallel & & \downarrow h_1 & & \downarrow g_1 \\ S_1 & \xrightarrow{l_1} & N & \xrightarrow{l_2} & L \\ & & \downarrow h_2 & & \downarrow g_2 \\ & & M & \xlongequal{\quad} & M. \end{array}$$

Since  $S_1$  is simple, we obtain that  $\ell(N) = \ell(L) + 1$  if and only if the short exact sequence in the second row is split, contradicting to the non-splitness of the one in the first row. Hence  $\ell(N) \neq \ell(L) + 1$ , or equivalently,  $\ell(N) = \ell(L)$ . Therefore,

$$\begin{aligned} \sum_{[N]; \ell(N)=\ell(M)+1} |\mathrm{Ext}^1(M, S^{(an)})_N| &= \sum_{[L]; \ell(L)=\ell(M)+1} |\phi^{-1}(\mathrm{Ext}^1(M, S^{(an-1)})_L)| \\ &= \sum_{[L]; \ell(L)=\ell(M)+1} |\mathrm{Ext}^1(M, S^{(an-1)})_L| \cdot |\mathrm{Ker}(\phi)|. \end{aligned}$$

By (5.31) we obtain that

$$\begin{aligned} |\mathrm{Ker}(\phi)| &= \frac{|\mathrm{Ext}^1(M, S_1)| \cdot |\mathrm{Hom}(M, S_0^{(an)})|}{|\mathrm{Hom}(M, S_1)| \cdot |\mathrm{Hom}(M, S_0^{(an-1)})|} \\ &= q^{-\langle M, S_1 \rangle} \cdot \frac{|\mathrm{Hom}(M, S_0^{(an)})|}{|\mathrm{Hom}(M, S_0^{(an-1)})|}. \end{aligned}$$

Now by Lemma 5.6, we have

$$\begin{aligned} &\sum_{[N]; \ell(N) = \ell(M) + 1} |\mathrm{Ext}^1(M, S_0^{(an)})_N| \\ &= \frac{|\mathrm{Hom}(M, S_0^{(an-1)})|}{|\mathrm{Hom}(M, S_0^{(n-1)})|} \cdot q^{-\langle M, S_1 \rangle} \cdot \frac{|\mathrm{Hom}(M, S_0^{(an)})|}{|\mathrm{Hom}(M, S_0^{(an-1)})|} \\ &= q^{-\langle M, S_1 \rangle} \cdot \frac{|\mathrm{Hom}(M, S_0^{(an)})|}{|\mathrm{Hom}(M, S_0^{(n-1)})|} = \frac{|\mathrm{Hom}(M, S_0^{(an)})|}{|\mathrm{Ext}^1(M, S_0^{(n-1)})|}. \end{aligned}$$

Here the last equality uses (5.31) with  $a = 1$ . □

## 6. $\imath$ HALL ALGEBRAS AND $\imath$ QUANTUM LOOP ALGEBRAS

In this section, we shall formulate the main result of this paper.

Let us fix some notations. For a partition  $\lambda = (1^{l_1}, \dots, n^{l_n})$ , we denote  $|\lambda| := \sum_{i=1}^n il_i$ , and  $\ell(\lambda) := \sum_{i=1}^n l_i$ .

**6.1. Embedding of  $\imath$ Hall algebras I.** Let  $\mathbb{X}$  be a weighted projective line of weight type  $(\mathbf{p}, \boldsymbol{\lambda})$ . Recall that  $\boldsymbol{\lambda}_i$  is the exceptional closed point of  $\mathbb{X}$  of weight  $p_i$  for any  $1 \leq i \leq \mathbf{t}$ , and  $\mathcal{T}_{\boldsymbol{\lambda}_i}$  is the Serre subcategory of  $\mathrm{coh}(\mathbb{X})$  consisting of torsion sheaves supported at  $\boldsymbol{\lambda}_i$ . There is an equivalence  $\mathcal{T}_{\boldsymbol{\lambda}_i} \cong \mathrm{rep}_{\mathbf{k}}^{\mathrm{nil}}(C_{p_i})$ , which induces a canonical embedding of Ringel-Hall algebras  $\mathcal{H}(\mathcal{C}_1(\mathrm{rep}_{\mathbf{k}}^{\mathrm{nil}}(C_{p_i}))) \rightarrow \mathcal{H}(\mathcal{C}_1(\mathrm{coh}(\mathbb{X}_{\mathbf{k}})))$ , and then an embedding of  $\imath$ Hall algebras:

$$\iota_i : \imath\tilde{\mathcal{H}}(\mathbf{k}C_{p_i}) \longrightarrow \imath\tilde{\mathcal{H}}(\mathbb{X}_{\mathbf{k}}). \quad (6.1)$$

Inspired by (5.5), we define

$$\widehat{B}_{[i,j],l} := \iota_i(\widehat{B}_{j,l}), \quad \widehat{\Theta}_{[i,j],r} := \iota_i(\widehat{\Theta}_{j,r}), \quad \widehat{H}_{[i,j],r} := \iota_i(\widehat{H}_{j,r}), \quad (6.2)$$

for any  $1 \leq j \leq p_i - 1$ ,  $l \in \mathbb{Z}$  and  $r > 0$ . By (2.20) we have

$$1 + \sum_{m \geq 1} (\mathbf{v} - \mathbf{v}^{-1}) \widehat{\Theta}_{[i,j],m} u^m = \exp \left( (\mathbf{v} - \mathbf{v}^{-1}) \sum_{m \geq 1} \widehat{H}_{[i,j],m} u^m \right). \quad (6.3)$$

For convenience, we set  $\widehat{\Theta}_{[i,j],0} = \frac{1}{\mathbf{v} - \mathbf{v}^{-1}}$  for any  $1 \leq i \leq \mathbf{t}$ ,  $1 \leq j \leq p_i - 1$ .

**6.2. Embedding of  $\iota$ Hall algebras II.** Let  $\mathcal{C}$  be the Serre subcategory of  $\text{coh}(\mathbb{X})$  generated by those simple sheaves  $S$  satisfying  $\text{Hom}(\mathcal{O}, S) = 0$ . Recall from [BS13] that the Serre quotient  $\text{coh}(\mathbb{X})/\mathcal{C}$  is equivalent to the category  $\text{coh}(\mathbb{P}^1)$  and the canonical functor  $\text{coh}(\mathbb{X}) \rightarrow \text{coh}(\mathbb{X})/\mathcal{C}$  has an exact fully faithful right adjoint functor

$$\mathbb{F}_{\mathbb{X}, \mathbb{P}^1} : \text{coh}(\mathbb{P}^1) \rightarrow \text{coh}(\mathbb{X}), \quad (6.4)$$

which sends

$$\mathcal{O}_{\mathbb{P}^1}(l) \mapsto \mathcal{O}(l\vec{c}), \quad S_{\lambda_i}^{(r)} \mapsto S_{i,0}^{(rp_i)}, \quad S_x^{(r)} \mapsto S_x^{(r)}$$

for any  $l \in \mathbb{Z}$ ,  $r \geq 1$ ,  $1 \leq i \leq \mathbf{t}$  and  $x \in \mathbb{P}_{\mathbf{k}}^1 \setminus \{\lambda_1, \dots, \lambda_{\mathbf{t}}\}$ . Then  $\mathbb{F}_{\mathbb{X}, \mathbb{P}^1}$  induces an exact fully faithful functor  $\mathcal{C}_1(\text{coh}(\mathbb{P}^1)) \rightarrow \mathcal{C}_1(\text{coh}(\mathbb{X}))$ , which is also denoted by  $\mathbb{F}_{\mathbb{X}, \mathbb{P}^1}$ . This functor  $\mathbb{F}_{\mathbb{X}, \mathbb{P}^1}$  induces a canonical embedding of Hall algebras  $\mathcal{H}(\mathcal{C}_1(\text{coh}(\mathbb{P}^1))) \rightarrow \mathcal{H}(\mathcal{C}_1(\text{coh}(\mathbb{X})))$ , and then an embedding of  $\iota$ Hall algebras:

$$F_{\mathbb{X}, \mathbb{P}^1} : {}^{\iota}\tilde{\mathcal{H}}(\mathbb{P}_{\mathbf{k}}^1) \longrightarrow {}^{\iota}\tilde{\mathcal{H}}(\mathbb{X}_{\mathbf{k}}). \quad (6.5)$$

Recall from [LRW23, (4.2), Proposition 6.3] that for any  $m \geq 0$ , in  ${}^{\iota}\tilde{\mathcal{H}}(\mathbb{P}_{\mathbf{k}}^1)$

$$\hat{\Theta}_m = \frac{1}{(q-1)^{2\mathbf{v}m-1}} \sum_{0 \neq f: \mathcal{O}(s) \rightarrow \mathcal{O}(m+s)} [\text{Coker}(f)],$$

which is independent of  $s \in \mathbb{Z}$ , and

$$\hat{H}_m = \sum_{x, d_x | m} \frac{[m]_{\mathbf{v}}}{m} d_x \sum_{|\lambda| = \frac{m}{d_x}} \mathbf{n}_x(\ell(\lambda) - 1) \frac{[S_x^{(\lambda)}]}{|\text{Aut}(S_x^{(\lambda)})|} - \delta_{m, ev} \frac{[m]_{\mathbf{v}}}{m} [K_{\frac{m}{2}} \delta],$$

where  $d_x$  is defined in (4.1), and

$$\mathbf{n}_x(l) = \prod_{i=1}^l (1 - \mathbf{v}^{2id_x}), \quad \forall l \geq 1.$$

Hence, we define

$$\hat{\Theta}_{\star, m} := F_{\mathbb{X}, \mathbb{P}^1}(\hat{\Theta}_m) = \frac{1}{(q-1)^{2\mathbf{v}m-1}} \sum_{0 \neq f: \mathcal{O}(s\vec{c}) \rightarrow \mathcal{O}((m+s)\vec{c})} [\text{Coker}(f)], \quad (6.6)$$

and

$$\hat{H}_{\star, m} := F_{\mathbb{X}, \mathbb{P}^1}(\hat{H}_m) = \sum_{x, d_x | m} \frac{[m]_{\mathbf{v}}}{m} d_x \sum_{|\lambda| = \frac{m}{d_x}} \mathbf{n}_x(\ell(\lambda) - 1) \frac{F_{\mathbb{X}, \mathbb{P}^1}([S_x^{(\lambda)}])}{|\text{Aut}(S_x^{(\lambda)})|} - \delta_{m, ev} \frac{[m]_{\mathbf{v}}}{m} [K_{\frac{m}{2}} \delta]. \quad (6.7)$$

Here, for any partition  $\lambda = (\lambda_1, \lambda_2, \dots)$ ,  $F_{\mathbb{X}, \mathbb{P}^1}([S_x^{(\lambda)}]) = [S_x^{(\lambda)}]$  for  $x \in \mathbb{P}_{\mathbf{k}}^1 \setminus \{\lambda_1, \dots, \lambda_{\mathbf{t}}\}$ , while  $F_{\mathbb{X}, \mathbb{P}^1}([S_{\lambda_i}^{(\lambda)}]) = S_{i,0}^{(\lambda)} := \bigoplus_i S_{i,0}^{(\lambda_i p_i)}$  for  $1 \leq i \leq \mathbf{t}$ . Note that  $\hat{\Theta}_{\star, m} = 0$  if  $m < 0$  and  $\hat{\Theta}_{\star, 0} = \frac{1}{\mathbf{v} - \mathbf{v}^{-1}}$ .

**Lemma 6.1.** *For any  $m \geq 1$  and  $x \in \mathbb{P}_{\mathbf{k}}^1$  such that  $d_x | m$ , we define*

$$\hat{H}_{x, m} := \frac{[m]_{\mathbf{v}}}{m} d_x \sum_{|\lambda| = \frac{m}{d_x}} \mathbf{n}_x(\ell(\lambda) - 1) \frac{F_{\mathbb{X}, \mathbb{P}^1}([S_x^{(\lambda)}])}{|\text{Aut}(S_x^{(\lambda)})|} - \delta_{\frac{m}{d_x}, ev} \cdot d_x \mathbf{v}^{-\frac{m}{2}} \frac{[\frac{m}{2}]_{\mathbf{v}}}{m} [K_{\frac{m}{2}} \delta]. \quad (6.8)$$



Then

$$\widehat{H}_{\star,m} = \sum_{x,d_x|m} \widehat{H}_{x,m}. \quad (6.9)$$

*Proof.* Note that for  $m \geq 1$ ,

$$\sum_{x \in \mathbb{P}_k^1; d_x|m} d_x = q^m + 1.$$

A direct computation shows that

$$\begin{aligned} & \sum_{x,d_x|m} (\mathbf{v}_x - \mathbf{v}_x^{-1}) \delta_{\frac{m}{d_x},ev} \mathbf{v}^{-\frac{m}{2}} \frac{[m/(2d_x)]_{\mathbf{v}_x}}{m/d_x} [K_{\frac{m}{2}\delta}] \\ &= \sum_{x,d_x|m} \delta_{\frac{m}{d_x},ev} \mathbf{v}^{-\frac{m}{2}} \frac{\mathbf{v}^{\frac{m}{2}} - \mathbf{v}^{-\frac{m}{2}}}{m} d_x [K_{\frac{m}{2}\delta}] = \delta_{m,ev} \frac{1 - \mathbf{v}^{-m}}{m} \sum_{x,d_x|\frac{m}{2}} d_x [K_{\frac{m}{2}\delta}] \\ &= \delta_{m,ev} \frac{1 - \mathbf{v}^{-m}}{m} (q^{\frac{m}{2}} + 1) [K_{\frac{m}{2}\delta}] = (\mathbf{v} - \mathbf{v}^{-1}) \delta_{m,ev} \frac{[m]_{\mathbf{v}}}{m} [K_{\frac{m}{2}\delta}]. \end{aligned}$$

Then the desired result follows from

$$\frac{\mathbf{v}_x - \mathbf{v}_x^{-1}}{\mathbf{v} - \mathbf{v}^{-1}} \left[ \frac{m}{2d_x} \right]_{\mathbf{v}_x} = \left[ \frac{m}{2} \right]_{\mathbf{v}}.$$

□

**6.3. The homomorphism  $\Omega$ .** Recall the star-shaped graph  $\Gamma = T_{p_1, \dots, p_t}$  in (2.27). The following is one of the main results of this paper.

**Theorem 6.2.** *For any star-shaped graph  $\Gamma$ , let  $\mathfrak{g}$  be the Kac-Moody algebra and  $\mathbb{X}$  be the weighted projective line associated to  $\Gamma$ . Then there exists a  $\mathbb{Q}(\mathbf{v})$ -algebra homomorphism*

$$\Omega : {}^{Dr} \widetilde{\mathbf{U}}_{|v=\mathbf{v}} \longrightarrow {}^i \widetilde{\mathcal{H}}(\mathbb{X}_{\mathbf{k}}), \quad (6.10)$$

which sends

$$\mathbb{K}_{\star} \mapsto [K_{\mathcal{O}}], \quad \mathbb{K}_{[i,j]} \mapsto [K_{S_{ij}}], \quad C \mapsto [K_{\delta}]; \quad (6.11)$$

$$B_{\star,l} \mapsto \frac{-1}{q-1} [\mathcal{O}(l\vec{c})], \quad \Theta_{\star,r} \mapsto \widehat{\Theta}_{\star,r}, \quad H_{\star,r} \mapsto \widehat{H}_{\star,r}; \quad (6.12)$$

$$B_{[i,j],l} \mapsto \frac{-1}{q-1} \widehat{B}_{[i,j],l}, \quad \Theta_{[i,j],r} \mapsto \widehat{\Theta}_{[i,j],r}, \quad H_{[i,j],r} \mapsto \widehat{H}_{[i,j],r}; \quad (6.13)$$

for any  $[i,j] \in \mathbb{I}_0 - \{\star\}$ ,  $l \in \mathbb{Z}$ ,  $r > 0$ .

We shall verify that the relations (2.12)–(2.18) are preserved by  $\Omega$ , and the proof of this theorem consists of the next 3 sections.

The composition subalgebra  ${}^i \widetilde{\mathcal{C}}(\mathbb{X}_{\mathbf{k}})$  is the subalgebra of  ${}^i \widetilde{\mathcal{H}}(\mathbb{X}_{\mathbf{k}})$  generated by the elements  $[\mathcal{O}(l\vec{c})]$  for  $l \in \mathbb{Z}$ ,  $\widehat{\Theta}_{\star,r}$  for  $r \geq 1$ ,  $[S_{ij}]$  for  $1 \leq i \leq \mathbf{t}$  and  $1 \leq j \leq p_i - 1$ , together with  $[K_{\mathcal{O}}]^{\pm 1}$ ,  $[K_{\delta}]^{\pm 1}$  and  $[K_{S_{ij}}]^{\pm 1}$ .

**Corollary 6.3.** *For any star-shaped graph  $\Gamma$ , let  $\mathfrak{g}$  be the Kac-Moody algebra and  $\mathbb{X}$  be the weighted projective line associated to  $\Gamma$ . Then there exists a  $\mathbb{Q}(\mathbf{v})$ -algebra epimorphism*

$$\Omega : {}^{Dr}\widetilde{\mathcal{U}}_{|v=\mathbf{v}}^i \longrightarrow {}^i\widetilde{\mathcal{C}}(\mathbb{X}_{\mathbf{k}}), \quad (6.14)$$

defined as in (6.11)–(6.13).

*Proof.* Using Lemma 2.7, it follows from Theorem 6.2.  $\square$

## 7. RELATIONS IN TUBES AND COMMUTATIVE RELATIONS

In this section, we shall verify the relations (2.12)–(2.18) for the indexes  $\mu, \nu \in \mathbb{I}_0$ , where  $\{\mu, \nu\} \neq \{\star, [i, 1]\}$  for any  $1 \leq i \leq \mathbf{t}$ . More precisely, we will consider the following three cases:  $\mu, \nu \in \mathbb{I}_0 - \{\star\}$ , or  $\mu = \star = \nu$ , or  $\{\mu, \nu\} = \{\star, [i, j]\}$  for  $2 \leq j \leq p_i - 1$  in §7.1, §7.2 and §7.3 respectively.

Let  $k_1, k_2, l \in \mathbb{Z}$  and  $\mu, \nu \in \mathbb{I}_0$ . Inspiring by (2.10)–(2.11), we introduce shorthand notations:

$$\begin{aligned} \widehat{S}(k_1, k_2 | l; \mu, \nu) &:= \widehat{B}_{\mu, k_1} * \widehat{B}_{\mu, k_2} * \widehat{B}_{\nu, l} - [2]_{\mathbf{v}} \widehat{B}_{\mu, k_1} * \widehat{B}_{\nu, l} * \widehat{B}_{\mu, k_2} + \widehat{B}_{\nu, l} * \widehat{B}_{\mu, k_1} * \widehat{B}_{\mu, k_2}, \\ \widehat{\mathbb{S}}(k_1, k_2 | l; \mu, \nu) &:= \widehat{S}(k_1, k_2 | l; \mu, \nu) + \{k_1 \leftrightarrow k_2\}. \end{aligned} \quad (7.1)$$

We also denote

$$\begin{aligned} \widehat{R}(k_1, k_2 | l; \mu, \nu) &:= [K_{k_1 \delta + \alpha_\mu}] * \left( - \sum_{p \geq 0} \mathbf{v}^{2p} [2]_{\mathbf{v}} \cdot [\widehat{\Theta}_{\mu, k_2 - k_1 - 2p - 1}, \widehat{B}_{\nu, l - 1}]_{\mathbf{v}^{-2}} * [K_\delta]^{p+1} \right. \\ &\quad \left. - \sum_{p \geq 1} \mathbf{v}^{2p-1} [2]_{\mathbf{v}} \cdot [\widehat{B}_{\nu, l}, \widehat{\Theta}_{\mu, k_2 - k_1 - 2p}]_{\mathbf{v}^{-2}} * [K_\delta]^p - [\widehat{B}_{\nu, l}, \widehat{\Theta}_{\mu, k_2 - k_1}]_{\mathbf{v}^{-2}} \right), \\ \widehat{\mathbb{R}}(k_1, k_2 | l; \mu, \nu) &:= \widehat{R}(k_1, k_2 | l; \mu, \nu) + \{k_1 \leftrightarrow k_2\}. \end{aligned} \quad (7.2)$$

By direct computations, we have

$$\widehat{\mathbb{R}}(k, k | l; \mu, \nu) = -[\widehat{B}_{\nu, l}, \widehat{\Theta}_{\mu, 0}]_{\mathbf{v}^{-2}} * [K_{k \delta + \alpha_\mu}] = -\mathbf{v}^{-1} \widehat{B}_{\nu, l} * [K_{k \delta + \alpha_\mu}], \quad (7.3)$$

$$\widehat{\mathbb{R}}(k, k + 1 | l; \mu, \nu) = \left( - [2]_{\mathbf{v}} \cdot [\widehat{\Theta}_{\mu, 0}, \widehat{B}_{\nu, l - 1}]_{\mathbf{v}^{-2}} * [K_\delta] - [\widehat{B}_{\nu, l}, \widehat{\Theta}_{\mu, 1}]_{\mathbf{v}^{-2}} \right) * [K_{k \delta + \alpha_\mu}], \quad (7.4)$$

and for  $k_2 \geq k_1 + 2$ ,

$$\begin{aligned} &\widehat{\mathbb{R}}(k_1, k_2 | l; \mu, \nu) - q \widehat{\mathbb{R}}(k_1, k_2 - 2 | l; \mu, \nu) * [K_\delta] \\ &= \left( - [2]_{\mathbf{v}} \cdot [\widehat{\Theta}_{\mu, k_2 - k_1 - 1}, \widehat{B}_{\nu, l - 1}]_{\mathbf{v}^{-2}} * [K_\delta] - [\widehat{B}_{\nu, l}, \widehat{\Theta}_{\mu, k_2 - k_1 - 2}]_{\mathbf{v}^{-2}} * [K_\delta] - [\widehat{B}_{\nu, l}, \widehat{\Theta}_{\mu, k_2 - k_1}]_{\mathbf{v}^{-2}} \right) \\ &\quad * [K_{k_1 \delta + \alpha_\mu}]. \end{aligned} \quad (7.5)$$

### 7.1. Relations for vertices in $\mathbb{I}_0 - \{\star\}$ .

**Proposition 7.1.** *For any  $\mu = [i, j], \nu = [i', j'] \in \mathbb{I}_0 - \{\star\}$ , the following relations hold in  ${}^i\widetilde{\mathcal{H}}(\mathbb{X}_{\mathbf{k}})$ , where  $k, l \in \mathbb{Z}$ ,  $m, n \geq 1$ :*

$$[\widehat{H}_{\mu, m}, \widehat{H}_{\nu, n}] = 0, \quad (7.6)$$

$$[\widehat{H}_{\mu, m}, \widehat{B}_{\nu, l}] = \frac{[ma_{\mu\nu}]_{\mathbf{v}}}{m} \widehat{B}_{\nu, l+m} - \frac{[ma_{\mu\nu}]_{\mathbf{v}}}{m} \widehat{B}_{\nu, l-m} * [K_{m\delta}], \quad (7.7)$$

$$[\widehat{B}_{\mu, k}, \widehat{B}_{\nu, l}] = 0, \text{ if } a_{\mu\nu} = 0, \quad (7.8)$$

$$[\widehat{B}_{\mu,k}, \widehat{B}_{\nu,l+1}]_{\mathbf{v}^{-a_{\mu\nu}}} - \mathbf{v}^{-a_{\mu\nu}} [\widehat{B}_{\mu,k+1}, \widehat{B}_{\nu,l}]_{\mathbf{v}^{a_{\mu\nu}}} = 0, \text{ if } a_{\mu\nu} = -1, \quad (7.9)$$

$$[\widehat{B}_{\mu,k}, \widehat{B}_{\mu,l+1}]_{\mathbf{v}^{-2}} - \mathbf{v}^{-2} [\widehat{B}_{\mu,k+1}, \widehat{B}_{\mu,l}]_{\mathbf{v}^2} = (1-q)^2 \left( \mathbf{v}^{-2} \widehat{\Theta}_{\mu,l-k+1} * [K_{k\delta+\alpha_{ij}}] \right. \quad (7.10)$$

$$\left. - \mathbf{v}^{-4} \widehat{\Theta}_{\mu,l-k-1} * [K_{(k+1)\delta+\alpha_{ij}}] + \mathbf{v}^{-2} \widehat{\Theta}_{\mu,k-l+1} * [K_{l\delta+\alpha_{ij}}] - \mathbf{v}^{-4} \widehat{\Theta}_{\mu,k-l-1} * [K_{(l+1)\delta+\alpha_{ij}}] \right),$$

$$\widehat{\mathbb{S}}(k_1, k_2 \mid l; \mu, \nu) = (1-q)^2 \widehat{\mathbb{R}}(k_1, k_2 \mid l; \mu, \nu), \text{ if } a_{\mu\nu} = -1. \quad (7.11)$$

*Proof.* Recall from (6.1) and (5.3) that we have the following algebra embeddings

$$\mathrm{Dr} \widetilde{\mathbf{U}}_{\mathbf{v}}^i(\widehat{\mathfrak{sl}}_{p_i}) \xrightarrow{\Omega_{C_{p_i}}} {}^i\widetilde{\mathcal{H}}(\mathbf{k}C_{p_i}) \xrightarrow{\iota_i} {}^i\widetilde{\mathcal{H}}(\mathbb{X}_{\mathbf{k}}).$$

Now it follows from (5.5) and (6.2) that all the relations (7.6)–(7.11) hold for  $i' = i$ .

Moreover, for any torsion sheaves  $X, Y$  supported at distinct points, we have  $\mathrm{Ext}_{\mathbb{X}}^1(X, Y) = 0 = \mathrm{Hom}_{\mathbb{X}}(X, Y)$ . It is well known that  $\mathrm{Ext}_{\mathcal{C}_1(\mathrm{coh}(\mathbb{X}))}^1(X, Y) \cong \mathrm{Ext}_{\mathbb{X}}^1(X, Y) \oplus \mathrm{Hom}_{\mathbb{X}}(X, Y)$ , and  $\mathrm{Hom}_{\mathcal{C}_1(\mathrm{coh}(\mathbb{X}))}(X, Y) = \mathrm{Hom}_{\mathbb{X}}(X, Y)$ ; see e.g. [LRW23, Lemma 2.6]. Hence  $[X] * [Y] = [X \oplus Y] = [Y] * [X]$  in  $\widetilde{\mathcal{H}}(\mathcal{C}_1(\mathrm{coh}(\mathbb{X})))$  and in  ${}^i\widetilde{\mathcal{H}}(\mathbb{X}_{\mathbf{k}})$ . Therefore, all the relations (7.6)–(7.11) hold for  $i' \neq i$ .  $\square$

**7.2. Relations for  $\star$ .** The following proposition shows that (2.13) and (2.17) hold in  ${}^i\widetilde{\mathcal{H}}(\mathbb{X}_{\mathbf{k}})$  for  $\mu = \nu = \star$ .

**Proposition 7.2.** *For any  $m, n \geq 1, k, l \in \mathbb{Z}$ , the following relations hold in  ${}^i\widetilde{\mathcal{H}}(\mathbb{X}_{\mathbf{k}})$ :*

$$[\widehat{H}_{\star,m}, \widehat{H}_{\star,n}] = 0, \quad (7.12)$$

$$[\widehat{H}_{\star,m}, [\mathcal{O}(l\vec{c})]] = \frac{[2m]_{\mathbf{v}}}{m} [\mathcal{O}((l+m)\vec{c})] - \frac{[2m]_{\mathbf{v}}}{m} [\mathcal{O}((l-m)\vec{c})] * [K_{m\delta}], \quad (7.13)$$

$$\begin{aligned} & [[\mathcal{O}(k\vec{c}), [\mathcal{O}((l+1)\vec{c})]]_{\mathbf{v}^{-2}} - \mathbf{v}^{-2} [[\mathcal{O}((k+1)\vec{c}), [\mathcal{O}(l\vec{c})]]_{\mathbf{v}^2} \quad (7.14) \\ & = (1-q)^2 \left( \mathbf{v}^{-2} \widehat{\Theta}_{\star,l-k+1} * [K_{\delta}]^k * [K_{\mathcal{O}}] - \mathbf{v}^{-4} \widehat{\Theta}_{\star,l-k-1} * [K_{\delta}]^{k+1} * [K_{\mathcal{O}}] \right. \\ & \left. + \mathbf{v}^{-2} \widehat{\Theta}_{\star,k-l+1} * [K_{\delta}]^l * [K_{\mathcal{O}}] - \mathbf{v}^{-4} \widehat{\Theta}_{\star,k-l-1} * [K_{\delta}]^{l+1} * [K_{\mathcal{O}}] \right). \end{aligned}$$

*Proof.* Recall from §6.2 that there is an embedding

$$F_{\mathbb{X}, \mathbb{P}^1} : {}^i\widetilde{\mathcal{H}}(\mathbb{P}_{\mathbf{k}}^1) \longrightarrow {}^i\widetilde{\mathcal{H}}(\mathbb{X}_{\mathbf{k}}),$$

which sends  $[\mathcal{O}_{\mathbb{P}^1}(l)] \mapsto [\mathcal{O}(l\vec{c})]$ ,  $[S_{\lambda_i}^{(r)}] \mapsto [S_{i,0}^{(rp_i)}]$ ,  $\widehat{\Theta}_r \mapsto \widehat{\Theta}_{\star,r}$  and  $\widehat{H}_r \mapsto \widehat{H}_{\star,r}$  for any  $l \in \mathbb{Z}, r \geq 1$ . Then the formulas follow from [LRW23, Theorem 4.2] and the Drinfeld type presentation of  $\widetilde{\mathbf{U}}_{\mathbf{v}}^i(\widehat{\mathfrak{sl}}_2)$  therein.  $\square$

**7.3. Relations between  $\star$  and  $[i, j]$  with  $j \neq 1$ .** In this subsection, we verify the relations (2.13)–(2.15) between  $\star$  and  $[i, j]$  with  $j \neq 1$ . First we give some preliminary results.

The set  $\mathcal{M}_{j,\alpha}$  defined in (5.6) can be viewed as a set of isoclasses of  $\mathrm{coh}(\mathbb{X})$  via  $\iota_i$ ; see §6.1. Recall from Corollary 5.2 that

$$\widehat{\Theta}_{[i,j],1} = \widehat{H}_{[i,j],1} = \pi_{[i,j+1],1} - (\mathbf{v} + \mathbf{v}^{-1})\pi_{[i,j],1} + \pi_{[i,j-1],1},$$

where

$$\pi_{[i,j],1} = \frac{-\mathbf{v}^{-j}}{\mathbf{v} - \mathbf{v}^{-1}} \sum_{\mathcal{M}_{j,\delta}} (-1)^{\dim \text{End}(M)} [M].$$

**Lemma 7.3.** *For any  $1 \leq i \leq \mathbf{t}$ ,  $1 \leq j \leq p_i - 1$ , and any  $l \in \mathbb{Z}$ ,  $r \geq 1$ , we have*

$$[\pi_{[i,j],1}, [\mathcal{O}(l\vec{c})]] = \frac{\mathbf{v}^{-j}}{\mathbf{v} - \mathbf{v}^{-1}} [[S_{i,0}^{(p_i)}], [\mathcal{O}(l\vec{c})]]; \quad (7.15)$$

$$[\pi_{[i,j],1}, [S_{i,0}^{(rp_i)}]] = 0. \quad (7.16)$$

*Proof.* For any  $1 \leq j', k \leq p_i$ , the torsion sheaf  $S_{i,j'}^{(k)}$  is uniserial, and  $S_{i,1}$  is a composition factor of  $S_{i,j'}^{(k)}$  if and only if  $k \geq j'$ . Hence, for any  $[M] \in \mathcal{M}_{j,\delta}$ , there exists a unique direct summand  $S_{i,j'}^{(k)}$  of  $M$  with  $k \geq j'$ , namely,  $M \cong S_{i,j'}^{(k)} \oplus M'$  for some  $M'$ . We emphasize that

$$[S_{i,j'}^{(k)} \oplus M'] \in \mathcal{M}_{j,\delta} \quad \text{if and only if} \quad [S_{i,j'}^{(j')} \oplus S_{i,0}^{(k-j')} \oplus M'] \in \mathcal{M}_{j,\delta}.$$

For any  $k > j'$ , we have

$$\dim \text{End}(S_{i,j'}^{(j')} \oplus S_{i,0}^{(k-j')} \oplus M') = \dim \text{End}(S_{i,j'}^{(k)} \oplus M') + 1;$$

moreover, there are no homomorphisms and extensions between  $\mathcal{O}(l\vec{c})$  and  $M'$  since both  $S_{i,0}$  and  $S_{i,1}$  are not composition factors of  $M'$ , hence by (4.12) we get

$$[[S_{i,j'}^{(k)} \oplus M'], [\mathcal{O}(l\vec{c})]] = [[S_{i,j'}^{(j')} \oplus S_{i,0}^{(k-j')} \oplus M'], [\mathcal{O}(l\vec{c})]].$$

Therefore, only for the case  $k = j' = p_i$ , i.e., the term  $[S_{i,0}^{(p_i)}]$  in  $\pi_{[i,j],1}$ , has non-trivial contribution to  $[\pi_{[i,j],1}, [\mathcal{O}(l\vec{c})]]$ . Then (7.15) follows immediately.

For (7.16), by similar arguments as above and using (4.13), we obtain that

$$[\pi_{[i,j],1}, [S_{i,0}^{(rp_i)}]] = \frac{\mathbf{v}^{-j}}{\mathbf{v} - \mathbf{v}^{-1}} [[S_{i,0}^{(p_i)}], [S_{i,0}^{(rp_i)}]] = 0,$$

where the last equality follows from [LRW23, Lemma 4.5] and the embedding (6.5).  $\square$

Consequently, we have the following result.

**Lemma 7.4.** *For any  $1 \leq i \leq \mathbf{t}$  and  $2 \leq j \leq p_i - 1$ , we have*

- (1)  $[\widehat{\Theta}_{[i,j],1}, [\mathcal{O}(l\vec{c})]] = 0$  for any  $l \in \mathbb{Z}$ ;
- (2)  $[\widehat{\Theta}_{[i,j],1}, \widehat{\Theta}_{*,m}] = 0$  for any  $m \geq 1$ .

*Proof.* For the statement (1), by (7.15) we have

$$\begin{aligned} & [\widehat{\Theta}_{[i,j],1}, [\mathcal{O}(l\vec{c})]] \\ &= [\pi_{[i,j+1],1} - (\mathbf{v} + \mathbf{v}^{-1})\pi_{[i,j],1} + \pi_{[i,j-1],1}, [\mathcal{O}(l\vec{c})]] \\ &= \left( \frac{\mathbf{v}^{-(j+1)}}{\mathbf{v} - \mathbf{v}^{-1}} - (\mathbf{v} + \mathbf{v}^{-1}) \cdot \frac{\mathbf{v}^{-j}}{\mathbf{v} - \mathbf{v}^{-1}} + \frac{\mathbf{v}^{-(j-1)}}{\mathbf{v} - \mathbf{v}^{-1}} \right) [[S_{i,0}^{(p_i)}], [\mathcal{O}(l\vec{c})]] \\ &= 0. \end{aligned}$$

For the statement (2), observe that there are no non-zero homomorphisms and extensions between distinct tubes, then by the definition of  $\widehat{\Theta}_{\star, m}$ , it suffices to show that

$$[\widehat{\Theta}_{[i, j], 1}, [S_{i, 0}^{(rp_i)}]] = 0, \quad \forall r \geq 1,$$

which then follows from (7.16) immediately.  $\square$

**Lemma 7.5.** *For any  $1 \leq i \leq \mathbf{t}$  and  $2 \leq j \leq p_i - 1$ , we have*

- (1)  $[\widehat{B}_{[i, j], -1}, [\mathcal{O}(l\vec{c})]] = 0$  for any  $l \in \mathbb{Z}$ ;
- (2)  $[\widehat{B}_{[i, j], -1}, \widehat{\Theta}_{\star, m}] = 0$  for any  $m \geq 1$ .

*Proof.* Recall from Proposition 5.1 and (6.1) that

$$\widehat{B}_{[i, j], -1} = \mathbf{v}^{-j+1} \sum_{\mathcal{M}_{j+1, \delta - e_j}} (-1)^{\dim \text{End}(M)} [M] * [K_{\delta - \alpha_j}]^{-1}.$$

For any term  $[M]$  in  $\mathcal{M}_{j+1, \delta - e_j}$ , we know that  $S_{i, 1}$  is a composition factor of  $M$  with multiplicity one. Hence there exists a unique direct summand  $S_{i, j'}^{(k)}$  of  $M$  with  $k \geq j'$ . Then by similar arguments as in the proof of Lemma 7.3 (observing  $[S_{i, 0}^{(rp_i)}] \notin \mathcal{M}_{j+1, \delta - e_j}$ ), we obtain  $[\widehat{B}_{[i, j], -1}, [\mathcal{O}(l\vec{c})]] = 0$  for any  $l \in \mathbb{Z}$ , and  $[\widehat{B}_{[i, j], -1}, [S_{i, 0}^{(rp_i)}]] = 0$  for any  $r \geq 1$ . Then the result follows immediately.  $\square$

7.3.1. *Relation (2.15) between  $\star$  and  $[i, j]$  with  $j \neq 1$ .*

**Proposition 7.6.** *For any  $1 \leq i \leq \mathbf{t}$ ,  $2 \leq j \leq p_i - 1$ , and any  $k, l \in \mathbb{Z}$ , we have*

$$[[\mathcal{O}(k\vec{c})], \widehat{B}_{[i, j], l}] = 0. \quad (7.17)$$

*Proof.* For  $2 \leq j \leq p_i - 1$ , by (7.7) we have

$$[\widehat{H}_{[i, j], 1}, \widehat{B}_{[i, j], l}] = [2]_{\mathbf{v}} \widehat{B}_{[i, j], l+1} - [2]_{\mathbf{v}} \widehat{B}_{[i, j], l-1} * [K_{\delta}].$$

Hence

$$[[\mathcal{O}(k\vec{c})], [\widehat{H}_{[i, j], 1}, \widehat{B}_{[i, j], l}]] = [2]_{\mathbf{v}} [[\mathcal{O}(k\vec{c})], \widehat{B}_{[i, j], l+1}] - [2]_{\mathbf{v}} [[\mathcal{O}(k\vec{c})], \widehat{B}_{[i, j], l-1}] * [K_{\delta}]. \quad (7.18)$$

By Lemma 7.4 (1), we have  $[[\mathcal{O}(k\vec{c})], \widehat{H}_{[i, j], 1}] = 0$ . Hence

$$[[\mathcal{O}(k\vec{c})], [\widehat{H}_{[i, j], 1}, \widehat{B}_{[i, j], l}]] = [\widehat{H}_{[i, j], 1}, [[\mathcal{O}(k\vec{c})], \widehat{B}_{[i, j], l}]].$$

If  $[[\mathcal{O}(k\vec{c})], \widehat{B}_{[i, j], l}] = 0$ , then LHS of (7.18) = 0, hence  $[[\mathcal{O}(k\vec{c})], \widehat{B}_{[i, j], l+1}] = 0$  if and only if  $[[\mathcal{O}(k\vec{c})], \widehat{B}_{[i, j], l-1}] = 0$ . Therefore, by induction, we only need to prove (7.17) for  $l = 0$  and  $-1$ .

Recall that  $\widehat{B}_{[i, j], 0} = [S_{ij}]$ , and there are no non-zero homomorphisms and extensions between  $\mathcal{O}(k\vec{c})$  and  $S_{ij}$  for  $2 \leq j \leq p_i - 1$ . Hence  $[[\mathcal{O}(k\vec{c})], [S_{ij}]] = 0$ . Moreover, by Lemma 7.5 (1), we have  $[[\mathcal{O}(k\vec{c})], \widehat{B}_{[i, j], -1}] = 0$ . The proof is completed.  $\square$

7.3.2. Relation (2.14) between  $\star$  and  $[i, j]$  with  $j \neq 1$ .

**Proposition 7.7.** For any  $1 \leq i \leq \mathbf{t}$ ,  $2 \leq j \leq p_i - 1$ , and any  $m \geq 1, l \in \mathbb{Z}$ , we have

- (1)  $[\widehat{H}_{\star, m}, \widehat{B}_{[i, j], l}] = 0$ ;
- (2)  $[\widehat{H}_{[i, j], m}, [\mathcal{O}(l\vec{c})]] = 0$ .

*Proof.* (1) It is equivalent to prove that  $[\widehat{\Theta}_{\star, m}, \widehat{B}_{[i, j], l}] = 0$ . By using similar arguments as in Proposition 7.6, thanks to (7.7) and Lemma 7.4 (2), we only need to prove the formula for  $l = 0$  and  $-1$ . For  $l = 0$ ,  $[\widehat{\Theta}_{\star, m}, [S_{ij}]] = 0$  is obvious since  $[[S_{i, 0}^{(p_i)}], [S_{ij}]] = 0$  for  $2 \leq j \leq p_i - 1$ ; while for  $l = -1$ , it follows from Lemma 7.5 (2).

(2) By (7.10) we have

$$\begin{aligned} & [\widehat{B}_{[i, j], l+1}, \widehat{B}_{[i, j], l+1}]_{\mathbf{v}^{-2}} - \mathbf{v}^{-2} [\widehat{B}_{[i, j], l+2}, \widehat{B}_{[i, j], l}]_{\mathbf{v}^2} \\ &= (1-q)^2 (\mathbf{v}^{-3} [K_{(l+1)\delta + \alpha_{ij}}] + \mathbf{v}^{-2} \widehat{\Theta}_{[i, j], 2} * [K_{l\delta + \alpha_{ij}}]). \end{aligned} \quad (7.19)$$

Apply the operator  $[-, [\mathcal{O}(l\vec{c})]]$  to (7.19). Then it follows from Proposition 7.6 and Lemma 7.4 (1) that

$$[\widehat{\Theta}_{[i, j], 1}, [\mathcal{O}(l\vec{c})]] = 0 = [\widehat{\Theta}_{[i, j], 2}, [\mathcal{O}(l\vec{c})]].$$

For  $n \geq 2$ , by (7.10) again we obtain

$$\begin{aligned} & [\widehat{B}_{[i, j], l+n}, \widehat{B}_{[i, j], l+1}]_{\mathbf{v}^{-2}} - \mathbf{v}^{-2} [\widehat{B}_{[i, j], l+n+1}, \widehat{B}_{[i, j], l}]_{\mathbf{v}^2} \\ &= (1-q)^2 (\mathbf{v}^{-2} \widehat{\Theta}_{[i, j], n+1} * [K_{l\delta + \alpha_{ij}}] - \mathbf{v}^{-4} \widehat{\Theta}_{[i, j], n-1} * [K_{(l+1)\delta + \alpha_{ij}}]). \end{aligned} \quad (7.20)$$

By applying  $[-, [\mathcal{O}(l\vec{c})]]$  to (7.20), we obtain  $[\widehat{\Theta}_{[i, j], m}, [\mathcal{O}(l\vec{c})]] = 0$  for any  $m \geq 1$  inductively. Then (2) follows from (2.20).  $\square$

7.3.3. Relation (2.13) between  $\star$  and  $[i, j]$  with  $j \neq 1$ .

**Proposition 7.8.** For any  $1 \leq i \leq \mathbf{t}$ ,  $2 \leq j \leq p_i - 1$ , and any  $m, n \geq 1$ ,

$$[\widehat{H}_{\star, m}, \widehat{H}_{[i, j], n}] = 0. \quad (7.21)$$

*Proof.* The proof is similar to that of Proposition 7.7 (2). Apply the operator  $[\widehat{\Theta}_{\star, m}, -]$  to (7.19). By Proposition 7.6 and Lemma 7.4 (2), we have

$$[\widehat{\Theta}_{\star, m}, \widehat{\Theta}_{[i, j], 1}] = 0 = [\widehat{\Theta}_{\star, m}, \widehat{\Theta}_{[i, j], 2}].$$

Now by applying  $[\widehat{\Theta}_{\star, m}, -]$  to (7.20), we obtain  $[\widehat{\Theta}_{\star, m}, \widehat{\Theta}_{[i, j], n}] = 0$  for any  $n \geq 1$ . Then (7.21) follows from (2.20).  $\square$

## 8. RELATIONS BETWEEN STAR POINT AND $[i, 1]$ , I

In this section and Section 9, we shall verify the relations (2.12)–(2.18) between  $\star$  and  $[i, 1]$ . In this section, we shall prove the following formulas in  ${}^v\widetilde{\mathcal{H}}(\mathbb{X}_{\mathbf{k}})$ :

$$[[\mathcal{O}(k\vec{c})], \widehat{B}_{[i, 1], l+1}]_{\mathbf{v}} - \mathbf{v} [[\mathcal{O}((k+1)\vec{c})], \widehat{B}_{[i, 1], l}]_{\mathbf{v}^{-1}} = 0, \quad (8.1)$$

$$\widehat{\mathbb{S}}(k_1, k_2 \mid l; \star, [i, 1]) = (1-q)^2 \widehat{\mathbb{R}}(k_1, k_2 \mid l; \star, [i, 1]), \quad (8.2)$$

$$[\widehat{H}_{\star, m}, \widehat{B}_{[i, 1], l}] = -\frac{[m]_{\mathbf{v}}}{m} \widehat{B}_{[i, 1], l+m} + \frac{[m]_{\mathbf{v}}}{m} \widehat{B}_{[i, 1], l-m} * [K_{m\delta}], \quad (8.3)$$

$$[\widehat{H}_{\star, m}, \widehat{H}_{[i, 1], r}] = 0, \quad (8.4)$$

for any  $k, k_1, k_2, l \in \mathbb{Z}$ ,  $m, r > 0$  and  $1 \leq i \leq \mathbf{t}$ .

Without loss of generality, we assume  $i = 1$  throughout this section.

### 8.1. The relation (8.1).

**Proposition 8.1.** *We have (8.1) holds for any  $k, l \in \mathbb{Z}$ .*

*Proof.* Obviously, (8.1) is equivalent to

$$[[\mathcal{O}(k\vec{c})], \widehat{B}_{[1, 1], l+1}]_{\mathbf{v}} + [\widehat{B}_{[1, 1], l}, [\mathcal{O}((k+1)\vec{c})]]_{\mathbf{v}} = 0, \quad \forall k, l \in \mathbb{Z}. \quad (8.5)$$

First, we prove (8.5) for the case when  $l = 0$  and  $-1$ . Recall from Corollary 5.3 that

$$\widehat{B}_{[1, 1], 0} = [S_{1, 1}], \quad \widehat{B}_{[1, 1], -1} = -[S_{1, 0}^{(p_1-1)}] * [K_{\alpha_{11}-\delta}], \quad \widehat{B}_{[1, 1], 1} = \frac{1}{q}[S_{1, 1}^{(p_1+1)}] - \frac{1}{q}[S_{1, 1} \oplus S_{1, 0}^{(p_1)}].$$

By (4.10) and (4.11) we obtain

$$[[\mathcal{O}(k\vec{c})], [S_{1, 1}]]_{\mathbf{v}} = -(q-1)[\mathcal{O}(k\vec{c} + \vec{x}_1)] = [[S_{1, 0}^{(p_1-1)}] * [K_{\alpha_{11}-\delta}], [\mathcal{O}((k+1)\vec{c})]]_{\mathbf{v}}.$$

Hence (8.5) holds for  $l = -1$ . For  $l = 0$ , (8.5) follows from Lemma 4.4.

For any  $l \in \mathbb{Z}$ , from (7.7), we have

$$[\widehat{\Theta}_{[1, 1], 1}, \widehat{B}_{[1, 1], l}] = [2]_{\mathbf{v}} \widehat{B}_{[1, 1], l+1} - [2]_{\mathbf{v}} \widehat{B}_{[1, 1], l-1} * [K_{\delta}].$$

Moreover, by Corollary 5.2 and Lemma 7.3,

$$\begin{aligned} & [\widehat{\Theta}_{[1, 1], 1}, [\mathcal{O}(k\vec{c})]] \\ &= [\pi_{[1, 2], 1} - (\mathbf{v} + \mathbf{v}^{-1})\pi_{[1, 1], 1}, [\mathcal{O}(k\vec{c})]] \\ &= \left( \frac{\mathbf{v}^{-2}}{\mathbf{v} - \mathbf{v}^{-1}} - \frac{\mathbf{v}^{-1}(\mathbf{v} + \mathbf{v}^{-1})}{\mathbf{v} - \mathbf{v}^{-1}} \right) [[S_{1, 0}^{(p_1)}], [\mathcal{O}(k\vec{c})]] \\ &= -\frac{1}{\mathbf{v} - \mathbf{v}^{-1}} \cdot \mathbf{v}^{-1}(q-1) ([\mathcal{O}((k+1)\vec{c})] - [\mathcal{O}((k-1)\vec{c})] * [K_{\delta}]) \\ &= -[\mathcal{O}((k+1)\vec{c})] + [\mathcal{O}((k-1)\vec{c})] * [K_{\delta}]. \end{aligned} \quad (8.6)$$

Therefore, we have

$$\begin{aligned} & \left[ \widehat{\Theta}_{[1, 1], 1}, [[\mathcal{O}(k\vec{c})], \widehat{B}_{[1, 1], l}]_{\mathbf{v}} + [\widehat{B}_{[1, 1], l-1}, [\mathcal{O}((k+1)\vec{c})]]_{\mathbf{v}} \right] \\ &= \left[ \widehat{\Theta}_{[1, 1], 1}, [[\mathcal{O}(k\vec{c})], \widehat{B}_{[1, 1], l}]_{\mathbf{v}} \right] + \left[ \widehat{\Theta}_{[1, 1], 1}, [\widehat{B}_{[1, 1], l-1}, [\mathcal{O}((k+1)\vec{c})]]_{\mathbf{v}} \right] \\ &= \left[ [\widehat{\Theta}_{[1, 1], 1}, [\mathcal{O}(k\vec{c})]], \widehat{B}_{[1, 1], l} \right]_{\mathbf{v}} + \left[ [\mathcal{O}(k\vec{c})], [\widehat{\Theta}_{[1, 1], 1}, \widehat{B}_{[1, 1], l}]_{\mathbf{v}} \right] \\ &\quad + \left[ [\widehat{\Theta}_{[1, 1], 1}, \widehat{B}_{[1, 1], l-1}], [\mathcal{O}((k+1)\vec{c})] \right]_{\mathbf{v}} + \left[ \widehat{B}_{[1, 1], l-1}, [\widehat{\Theta}_{[1, 1], 1}, [\mathcal{O}((k+1)\vec{c})]]_{\mathbf{v}} \right] \\ &= \left[ -[\mathcal{O}((k+1)\vec{c})] + [\mathcal{O}((k-1)\vec{c})] * [K_{\delta}], \widehat{B}_{[1, 1], l} \right]_{\mathbf{v}} \\ &\quad + [2]_{\mathbf{v}} \left[ [\mathcal{O}(k\vec{c})], \widehat{B}_{[1, 1], l+1} - \widehat{B}_{[1, 1], l-1} * [K_{\delta}] \right]_{\mathbf{v}} \\ &\quad + [2]_{\mathbf{v}} \left[ \widehat{B}_{[1, 1], l} - \widehat{B}_{[1, 1], l-2} * [K_{\delta}], [\mathcal{O}((k+1)\vec{c})] \right]_{\mathbf{v}} \end{aligned}$$

$$\begin{aligned}
& + \left[ \widehat{B}_{[1,1],l-1}, -[\mathcal{O}((k+2)\vec{c})] + [\mathcal{O}(k\vec{c})] * [K_\delta] \right]_{\mathbf{v}} \\
= & - \left( [[\mathcal{O}((k+1)\vec{c})], \widehat{B}_{[1,1],l}]_{\mathbf{v}} + [\widehat{B}_{[1,1],l-1}, [\mathcal{O}((k+2)\vec{c})]]_{\mathbf{v}} \right) \\
& + \left( [[\mathcal{O}((k-1)\vec{c})], \widehat{B}_{[1,1],l}]_{\mathbf{v}} + [\widehat{B}_{[1,1],l-1}, [\mathcal{O}(k\vec{c})]]_{\mathbf{v}} \right) * [K_\delta] \\
& + [2]_{\mathbf{v}} \left( [[\mathcal{O}(k\vec{c})], \widehat{B}_{[1,1],l+1}]_{\mathbf{v}} + [\widehat{B}_{[1,1],l}, [\mathcal{O}((k+1)\vec{c})]]_{\mathbf{v}} \right) \\
& - [2]_{\mathbf{v}} \left( [[\mathcal{O}(k\vec{c})], \widehat{B}_{[1,1],l-1}]_{\mathbf{v}} + [\widehat{B}_{[1,1],l-2}, [\mathcal{O}((k+1)\vec{c})]]_{\mathbf{v}} \right) * [K_\delta].
\end{aligned}$$

Observe that twisting with  $\vec{c}$  induces an automorphism on  ${}^v\mathcal{H}(\mathbb{X}_{\mathbf{k}})$ . Then (8.5) holds for any  $l \in \mathbb{Z}$  by induction.  $\square$

**8.2. The relation (8.2).** We shall prove (8.2) by induction on  $l$ . First we check (8.2) for  $l = 0, -1$ .

**Lemma 8.2.** *For any  $k_1, k_2 \in \mathbb{Z}$ , we have*

$$\widehat{\mathbb{S}}(k_1, k_2 \mid 0; \star, [1, 1]) = (1-q)^2 \widehat{\mathbb{R}}(k_1, k_2 \mid 0; \star, [1, 1]). \quad (8.7)$$

The proof of Lemma 8.2 is provided in Appendix A.1

**Lemma 8.3.** *For any  $k_1, k_2 \in \mathbb{Z}$ , we have*

$$\widehat{\mathbb{S}}(k_1, k_2 \mid -1; \star, [1, 1]) = (1-q)^2 \widehat{\mathbb{R}}(k_1, k_2 \mid -1; \star, [1, 1]). \quad (8.8)$$

The proof of Lemma 8.3 is provided in Appendix A.2.

**Proposition 8.4.** *For any  $k_1, k_2, l \in \mathbb{Z}$ , we have*

$$\widehat{\mathbb{S}}(k_1, k_2 \mid l; \star, [1, 1]) = (1-q)^2 \widehat{\mathbb{R}}(k_1, k_2 \mid l; \star, [1, 1]). \quad (8.9)$$

*Proof.* We only need to consider the case  $l \geq 0$ , since the other case  $l < 0$  is similar. By Lemma 8.2, we do induction on  $l$ , and assume that (8.9) holds for  $0 \leq l' \leq l$ .

Using (7.7) and (8.6), a direct computation shows

$$\begin{aligned}
& [\widehat{\Theta}_{[1,1],1}, \widehat{\mathbb{S}}(k_1, k_2 \mid l; \star, [1, 1])] \quad (8.10) \\
= & -\widehat{\mathbb{S}}(k_1 + 1, k_2 \mid l; \star, [1, 1]) + \widehat{\mathbb{S}}(k_1 - 1, k_2 \mid l; \star, [1, 1]) * [K_\delta] \\
& - \widehat{\mathbb{S}}(k_1, k_2 + 1 \mid l; \star, [1, 1]) + \widehat{\mathbb{S}}(k_1, k_2 - 1 \mid l; \star, [1, 1]) * [K_\delta] \\
& + [2]_{\mathbf{v}} \widehat{\mathbb{S}}(k_1, k_2 \mid l + 1; \star, [1, 1]) - [2]_{\mathbf{v}} \widehat{\mathbb{S}}(k_1, k_2 \mid l - 1; \star, [1, 1]) * [K_\delta] \\
= & (1-q)^2 \left( -\widehat{\mathbb{R}}(k_1 + 1, k_2 \mid l; \star, [1, 1]) + \widehat{\mathbb{R}}(k_1 - 1, k_2 \mid l; \star, [1, 1]) * [K_\delta] \right. \\
& \left. - \widehat{\mathbb{R}}(k_1, k_2 + 1 \mid l; \star, [1, 1]) + \widehat{\mathbb{R}}(k_1, k_2 - 1 \mid l; \star, [1, 1]) * [K_\delta] \right) \\
& + [2]_{\mathbf{v}} \widehat{\mathbb{S}}(k_1, k_2 \mid l + 1; \star, [1, 1]) - [2]_{\mathbf{v}} \widehat{\mathbb{S}}(k_1, k_2 \mid l - 1; \star, [1, 1]) * [K_\delta] \\
= & [2]_{\mathbf{v}} \widehat{\mathbb{S}}(k_1, k_2 \mid l + 1; \star, [1, 1]) - [2]_{\mathbf{v}} \widehat{\mathbb{S}}(k_1, k_2 \mid l - 1; \star, [1, 1]) * [K_\delta].
\end{aligned}$$

Here the second equality uses the inductive assumptions, and the third equality uses the following equality obtained from (7.2):

$$\widehat{\mathbb{R}}(k_1, k_2 \pm 1 \mid l; \star, [1, 1]) = \widehat{\mathbb{R}}(k_1 \mp 1, k_2 \mid l; \star, [1, 1]) * [K_\delta]^{\pm 1}.$$



On the other hand, using the same argument as in Lemma 7.4 (2), one can obtain that  $[\widehat{\Theta}_{[1,1],1}, \widehat{\Theta}_{\star,m}] = 0$  for any  $m > 0$ . Together with (7.7), we have

$$\begin{aligned} & [\widehat{\Theta}_{[1,1],1}, \widehat{\mathbb{R}}(k_1, k_2 \mid l; \star, [1, 1])] \\ &= [2]_{\mathbf{v}} \widehat{\mathbb{R}}(k_1, k_2 \mid l+1; \star, [1, 1]) - [2]_{\mathbf{v}} \widehat{\mathbb{R}}(k_1, k_2 \mid l-1; \star, [1, 1]) * [K_{\delta}]. \end{aligned} \quad (8.11)$$

Comparing (8.10) and (8.11), using the inductive assumption, one can obtain

$$\widehat{\mathbb{S}}(k_1, k_2 \mid l+1; \star, [1, 1]) = (1-q)^2 \widehat{\mathbb{R}}(k_1, k_2 \mid l+1; \star, [1, 1]).$$

The proof is completed.  $\square$

### 8.3. The relations (8.3) and (8.4).

**Proposition 8.5.** *Relation (8.3) holds in  ${}^{\mathbf{v}}\widetilde{\mathcal{H}}(\mathbb{X}_{\mathbf{k}})$  for any  $m > 0$  and  $l \in \mathbb{Z}$ .*

*Proof.* Denote

$$\begin{aligned} \widehat{Y}_{m|l} := & [\widehat{B}_{[1,1],l+1}, \widehat{\Theta}_{\star,m-1}]_{\mathbf{v}^{-2}} - [\widehat{\Theta}_{\star,m-1}, \widehat{B}_{[1,1],l-1}]_{\mathbf{v}^{-2}} * [K_{\delta}] \\ & + \mathbf{v}^{-1} [\widehat{\Theta}_{\star,m}, \widehat{B}_{[1,1],l}] + \mathbf{v}^{-1} [\widehat{\Theta}_{\star,m-2}, \widehat{B}_{[1,1],l}] * [K_{\delta}]. \end{aligned}$$

Using Lemma 2.5 (2), it is equivalent to prove that  $\widehat{Y}_{m|l} = 0$  for any  $m \geq 0$ . Clearly,  $\widehat{Y}_{m|l} = 0$  if  $m \leq 0$ . By induction, we assume that  $\widehat{Y}_{m|l} = 0$  for any  $m \leq k$ .

We also denote for  $n \in \mathbb{Z}$ ,

$$\begin{aligned} \widehat{X}_{n|l} = & \sum_{p \geq 0} \mathbf{v}^{2p+1} [\widehat{B}_{[1,1],l+1}, \widehat{\Theta}_{\star,n-2p-1}]_{\mathbf{v}^{-2}} * [K_{\delta}]^{p+1} + \sum_{p \geq 1} \mathbf{v}^{2p-1} [2]_{\mathbf{v}} [\widehat{\Theta}_{\star,n-2p}, \widehat{B}_{[1,1],l}] * [K_{\delta}]^{p+1} \\ & - \sum_{p \geq 0} \mathbf{v}^{2p+1} [\widehat{\Theta}_{\star,n-2p-1}, \widehat{B}_{[1,1],l-1}]_{\mathbf{v}^{-2}} * [K_{\delta}]^{p+2} + [\widehat{\Theta}_{\star,n}, \widehat{B}_{[1,1],l}] * [K_{\delta}]. \end{aligned}$$

Clearly,  $\widehat{X}_{n|l} = 0$ , for  $n \leq 0$ . By using the same argument as in [LW21b, Lemma 4.10], we have  $\widehat{X}_{n|l} - q\widehat{X}_{n-2|l} = \mathbf{v}\widehat{Y}_{n|l}$ . Then  $\widehat{X}_{n|l} = 0$  for any  $n \leq k$  by using the inductive assumption.

Using the same argument as in [LW21b, Lemma 4.14], we have

$$\begin{aligned} & \widehat{\mathbb{R}}(k_1, k_2 + 1 \mid l; \star, [1, 1]) + \widehat{\mathbb{R}}(k_1 + 1, k_2 \mid l; \star, [1, 1]) - [2]_{\mathbf{v}} \widehat{\mathbb{R}}(k_1, k_2 \mid l+1; \star, [1, 1]) \\ &= [2]_{\mathbf{v}}^2 \widehat{X}_{k-1|l} * [K_{\delta}]^{k_1} * [K_{\mathcal{O}}] + [2]_{\mathbf{v}} \widehat{Y}_{k+1|l} * [K_{\delta}]^{k_1} * [K_{\mathcal{O}}] \\ &+ ( - [\widehat{\Theta}_{\star,k_2-k_1+1}, \widehat{B}_{[1,1],l}]_{\mathbf{v}^{-2}} * [K_{\delta}]^{k_1} + \mathbf{v}^{-2} [\widehat{\Theta}_{\star,k_2-k_1-1}, \widehat{B}_{[1,1],l}]_{\mathbf{v}^{-2}} * [K_{\delta}]^{k_1+1} ) * [K_{\mathcal{O}}] \\ &+ \{k_1 \leftrightarrow k_2\}. \end{aligned} \quad (8.12)$$

With the help of Proposition 8.1 and (7.14), by using the same argument as in [LW21b, Lemma 4.13], we have

$$\begin{aligned} & \widehat{\mathbb{S}}(k_1, k_2 + 1 \mid l; \star, [1, 1]) + \widehat{\mathbb{S}}(k_1 + 1, k_2 \mid l; \star, [1, 1]) - [2]_{\mathbf{v}} \widehat{\mathbb{S}}(k_1, k_2 \mid l+1; \star, [1, 1]) \\ &= (1-q)^2 \left( - [\widehat{\Theta}_{\star,k_2-k_1+1}, \widehat{B}_{[1,1],l}]_{\mathbf{v}^{-2}} * [K_{\delta}]^{k_1} + \mathbf{v}^{-2} [\widehat{\Theta}_{\star,k_2-k_1-1}, \widehat{B}_{[1,1],l}]_{\mathbf{v}^{-2}} * [K_{\delta}]^{k_1+1} \right) \\ & * [K_{\mathcal{O}}] + \{k_1 \leftrightarrow k_2\}. \end{aligned} \quad (8.13)$$

From Proposition 8.4, we have

$$\widehat{\mathbb{S}}(k_1, k_2 + 1 \mid l; \star, [1, 1]) + \widehat{\mathbb{S}}(k_1 + 1, k_2 \mid l; \star, [1, 1]) - [2]_{\mathbf{v}} \widehat{\mathbb{S}}(k_1, k_2 \mid l+1; \star, [1, 1]) \quad (8.14)$$

$$= (1 - q)^2 \left( \widehat{\mathbb{R}}(k_1, k_2 + 1 \mid l; \star, [1, 1]) + \widehat{\mathbb{R}}(k_1 + 1, k_2 \mid l; \star, [1, 1]) - [2]_{\mathbf{v}} \widehat{\mathbb{R}}(k_1, k_2 \mid l + 1; \star, [1, 1]) \right).$$

By combining (8.12), (8.13) and (8.14), it follows that  $\widehat{Y}_{k+1|l} = 0$  since  $\widehat{X}_{k-1|l} = 0$  by the inductive assumption. The proof is completed.  $\square$

**Proposition 8.6.** *Relation (8.4) holds in  ${}^i\widetilde{\mathcal{H}}(\mathbb{X}_{\mathbf{k}})$  for any  $m, r > 0$ .*

*Proof.* With the help of (8.3) proved in Proposition 8.5 and (7.10) for  $\mu = [1, 1]$ , we can apply the same argument as in [LW21b, Proposition 4.8] to prove that (8.4) holds in  ${}^i\widetilde{\mathcal{H}}(\mathbb{X}_{\mathbf{k}})$  for any  $m, r > 0$ .  $\square$

## 9. RELATIONS BETWEEN STAR POINT AND $[i, 1]$ , II

In this section, we shall prove the following formulas in  ${}^i\widetilde{\mathcal{H}}(\mathbb{X}_{\mathbf{k}})$ :

$$\left[ \widehat{H}_{[i,1],m}, [\mathcal{O}(l\vec{c})] \right] = -\frac{[m]_{\mathbf{v}}}{m} [\mathcal{O}((l+m)\vec{c})] + \frac{[m]_{\mathbf{v}}}{m} [\mathcal{O}((l-m)\vec{c})] * [K_{m\delta}], \quad (9.1)$$

$$\widehat{\mathbb{S}}(k_1, k_2 \mid l; [i, 1], \star) = (1 - q)^2 \widehat{\mathbb{R}}(k_1, k_2 \mid l; [i, 1], \star), \quad (9.2)$$

for any  $k_1, k_2, l \in \mathbb{Z}$ ,  $m > 0$ ,  $1 \leq i \leq \mathbf{t}$ .

**9.1. Real and imaginary roots at  $[i, 1]$ .** In this subsection we give explicit descriptions of the real and imaginary roots at  $[i, 1]$  in  ${}^i\widetilde{\mathcal{H}}(\mathbb{X}_{\mathbf{k}})$ .

Without loss of generality, we assume  $i = 1$  in this whole section.

**9.1.1. Real roots at  $[1, 1]$ .** For any  $r > 0$ , denote by

$$\begin{aligned} \mathcal{M}_{r\delta+\alpha_{11}} &= \{[S_{1,1}^{(bp_1+1)} \oplus S_{1,0}^{(\nu)} \mid b \geq 0, |\nu| + b = r\}; \\ \mathcal{M}_{r\delta-\alpha_{11}} &= \{[S_{1,0}^{(ap_1-1)} \oplus S_{1,0}^{(\nu)} \mid a \geq 1, |\nu| + a = r\}; \end{aligned} \quad (9.3)$$

and set

$$[M_{r\delta \pm \alpha_{11}}] := \sum_{[M] \in \mathcal{M}_{r\delta \pm \alpha_{11}}} \mathbf{n}(\ell(M) - 1) \llbracket M \rrbracket, \quad (9.4)$$

where

$$\llbracket M \rrbracket = \frac{[M]}{|\text{Aut}(M)|}, \quad \mathbf{n}(l) = \prod_{i=1}^l (1 - \mathbf{v}^{2i}), \quad \forall l \geq 1.$$

**Proposition 9.1.** *For any  $r > 0$ , we have*

$$\widehat{B}_{[1,1],r} = (q - 1)[M_{r\delta+\alpha_{11}}], \quad \widehat{B}_{[1,1],-r} = (1 - q)[M_{r\delta-\alpha_{11}}] * [K_{-r\delta+\alpha_{11}}]. \quad (9.5)$$

The proof of Proposition 9.1 is provided in Appendix B.1.

9.1.2. *Imaginary roots at [1, 1].* For any  $r > 0$ , set

$$\begin{aligned} \mathcal{M}_{i,r\delta} := & \{ [S_{i,1}^{(ap_1)} \oplus S_{i,0}^{(\nu)}] \mid a \geq 1, a + |\nu| = r \} \\ & \cup \{ [S_{i,0}^{(ap_i-1)} \oplus S_{i,1}^{(bp_i+1)} \oplus S_{i,0}^{(\nu)}] \mid a + b + |\nu| = r \}. \end{aligned} \quad (9.6)$$

Since we focus on  $i = 1$  in the following, we also denote

$$\mathcal{M}_{r\delta} := \mathcal{M}_{1,r\delta}.$$

**Proposition 9.2.** *For any  $r > 0$ , we have*

$$\widehat{\Theta}_{[1,1],r} = \frac{\mathbf{v}}{q-1} \sum_{|\lambda|=r} \mathbf{n}(\ell(\lambda)) [S_{1,0}^{(\lambda)}] + \mathbf{v}^{-1} \sum_{[M] \in \mathcal{M}_{r\delta}} \mathbf{n}(\ell(M) - 1) [M]. \quad (9.7)$$

The proof of Proposition 9.2 is provided in Appendix B.2.

9.2. **The relation (9.1).** In this subsection, we shall prove (9.1). Note that twisting with  $\vec{c}$  induces an automorphism on  ${}^i\widetilde{\mathcal{H}}(\mathbb{X}_{\mathbf{k}})$ . By using Lemma 2.5 (2), it is equivalent to prove that for any  $r > 0$ ,

$$\begin{aligned} & [\widehat{\Theta}_{[1,1],r}, [\mathcal{O}]] + [\widehat{\Theta}_{[1,1],r-2}, [\mathcal{O}]] * [K_{\delta}] \\ &= \mathbf{v}^{-1} [\widehat{\Theta}_{[1,1],r-1}, [\mathcal{O}(\vec{c})]]_{\mathbf{v}^2} + \mathbf{v} [\widehat{\Theta}_{[1,1],r-1}, [\mathcal{O}(-\vec{c})]]_{\mathbf{v}^{-2}} * [K_{\delta}]. \end{aligned} \quad (9.8)$$

The proof of (9.8) needs the following Lemmas 9.3 and 9.4, whose proofs are technical, and we put them in Appendix B.3–B.4.

**Lemma 9.3.** *For any  $r > 0$ , we have*

$$\begin{aligned} \widehat{\Theta}_{[1,1],r} * [\mathcal{O}] &= \frac{\mathbf{v}^{r+1}}{q-1} \sum_{|\mu|=r} \mathbf{n}(\ell(\mu)) [\mathcal{O} \oplus S_{1,0}^{(\mu)}] + \mathbf{v}^{r-1} \sum_{[N] \in \mathcal{M}_{r\delta}} \mathbf{n}(\ell(N) - 1) [\mathcal{O} \oplus N] \\ &\quad - \sum_{k>0} \sum_{|\mu|=r-k} \mathbf{v}^{r-2k+1} \cdot \mathbf{n}(\ell(\mu)) [\mathcal{O}(k\vec{c}) \oplus S_{1,0}^{(\mu)}] \\ &\quad + \sum_{k>0} \sum_{[N] \in \mathcal{M}_{(r-k)\delta}} \mathbf{v}^{r-2k-1} (1-q) \cdot \mathbf{n}(\ell(N) - 1) [\mathcal{O}(k\vec{c}) \oplus N]. \end{aligned}$$

**Lemma 9.4.** *For any  $r > 0$ , we have*

$$\begin{aligned} [\mathcal{O}] * \widehat{\Theta}_{[1,1],r} &= \frac{\mathbf{v}^{r+1}}{q-1} \sum_{|\mu|=r} \mathbf{n}(\ell(\mu)) [\mathcal{O} \oplus S_{1,0}^{(\mu)}] + \mathbf{v}^{r-1} \sum_{[N] \in \mathcal{M}_{r\delta}} \mathbf{n}(\ell(N) - 1) [\mathcal{O} \oplus N] \\ &\quad - \sum_{k>0} \sum_{|\mu|=r-k} \mathbf{v}^{r-2k+1} \cdot \mathbf{n}(\ell(\mu)) [\mathcal{O}(-k\vec{c}) \oplus S_{1,0}^{(\mu)}] * [K_{k\delta}] \\ &\quad + \sum_{k>0} \sum_{[N] \in \mathcal{M}_{(r-k)\delta}} \mathbf{v}^{r-2k-1} (1-q) \cdot \mathbf{n}(\ell(N) - 1) [\mathcal{O}(-k\vec{c}) \oplus N] * [K_{k\delta}]. \end{aligned}$$

**Proposition 9.5.** *Relation (9.8) holds in  ${}^i\widetilde{\mathcal{H}}(\mathbb{X}_{\mathbf{k}})$  for any  $r > 0$ .*

*Proof.* By using Lemmas 9.3 and 9.4, (9.8) follows by easy (but tedious) cancellations, which we omit here.  $\square$

9.3. **The relation (9.2).** In this subsection, we shall prove the relation (9.2). First, let us give a lemma.

**Lemma 9.6.** *The following formula holds in  ${}^v\mathcal{H}(\mathbb{X}_{\mathbf{k}})$  for any  $l \in \mathbb{Z}$ :*

$$\widehat{\mathbb{S}}(0, 0 \mid l; [1, 1], \star) = (1 - q)^2 \widehat{\mathbb{R}}(0, 0 \mid l; [1, 1], \star).$$

*Proof.* Using (4.10), we obtain

$$\begin{aligned} & \widehat{\mathbb{S}}(0, 0 \mid l; [1, 1], \star) \\ &= [S_{1,1}] * [S_{1,1}] * [\mathcal{O}(l\vec{c})] - [2]_{\mathbf{v}}[S_{1,1}] * [\mathcal{O}(l\vec{c})] * [S_{1,1}] + [\mathcal{O}(l\vec{c})] * [S_{1,1}] * [S_{1,1}] \\ &= [S_{1,1}] * (\mathbf{v}^{-1}[\mathcal{O}(l\vec{c})] * [S_{1,1}] + (\mathbf{v} - \mathbf{v}^{-1})[\mathcal{O}(l\vec{c} + \vec{x}_1)]) - [2]_{\mathbf{v}}[S_{1,1}] * [\mathcal{O}(l\vec{c})] * [S_{1,1}] \\ &\quad + (\mathbf{v}[S_{1,1}] * [\mathcal{O}(l\vec{c})] - (q - 1)[\mathcal{O}(l\vec{c} + \vec{x}_1)]) * [S_{1,1}] \\ &= (\mathbf{v} - \mathbf{v}^{-1})[[S_{1,1}], [\mathcal{O}(l\vec{c} + \vec{x}_1)]]_{\mathbf{v}} \\ &= -\mathbf{v}^{-1}(q - 1)^2[\mathcal{O}(l\vec{c})] * [K_{\alpha_{11}}] \\ &= (1 - q)^2 \widehat{\mathbb{R}}(0, 0 \mid l; [1, 1], \star). \end{aligned}$$

□

**Proposition 9.7.** *Relation (9.2) holds in  ${}^v\mathcal{H}(\mathbb{X}_{\mathbf{k}})$  for any  $k_1, k_2, l \in \mathbb{Z}$ .*

*Proof.* We can assume that  $k_1 \leq k_2$ . With the help of Proposition 8.1 and (7.10) for  $\mu = [1, 1]$ , by using the same argument as in the proof of [LW21b, Lemmas 4.9 and 4.13], we have

$$\begin{aligned} & \widehat{\mathbb{S}}(k_1, k_2 + 1 \mid l; [1, 1], \star) + \widehat{\mathbb{S}}(k_1 + 1, k_2 \mid l; [1, 1], \star) - [2]_{\mathbf{v}}\widehat{\mathbb{S}}(k_1 + 1, k_2 + 1 \mid l - 1; [1, 1], \star) \\ &= (1 - q)^2 \left( - [[\mathcal{O}(l\vec{c})], \widehat{\Theta}_{[1,1], k_2 - k_1 + 1}]_{\mathbf{v}^{-2}} * [K_{\delta}]^{k_1} \right. \\ &\quad \left. + \mathbf{v}^{-2} [[\mathcal{O}(l\vec{c})], \widehat{\Theta}_{[1,1], k_2 - k_1 - 1}]_{\mathbf{v}^{-2}} * [K_{\delta}]^{k_1 + 1} \right) * [K_{\alpha_{11}}] + \{k_1 \leftrightarrow k_2\}, \end{aligned}$$

and

$$\begin{aligned} & \widehat{\mathbb{S}}(k_1, k_2 + 1 \mid l; [1, 1], \star) + \widehat{\mathbb{S}}(k_1 + 1, k_2 \mid l; [1, 1], \star) - [2]_{\mathbf{v}}\widehat{\mathbb{S}}(k_1, k_2 \mid l + 1; [1, 1], \star) \\ &= (1 - q)^2 \left( - [\widehat{\Theta}_{[1,1], k_2 - k_1 + 1}, [\mathcal{O}(l\vec{c})]]_{\mathbf{v}^{-2}} * [K_{\delta}]^{k_1} \right. \\ &\quad \left. + \mathbf{v}^{-2} [\widehat{\Theta}_{[1,1], k_2 - k_1 - 1}, [\mathcal{O}(l\vec{c})]]_{\mathbf{v}^{-2}} * [K_{\delta}]^{k_1 + 1} \right) * [K_{\alpha_{11}}] + \{k_1 \leftrightarrow k_2\}. \end{aligned}$$

Denote for  $m \in \mathbb{Z}$ ,

$$\begin{aligned} \widehat{Y}_{m|l} &= [[\mathcal{O}((l + 1)\vec{c})], \widehat{\Theta}_{[1,1], m-1}]_{\mathbf{v}^{-2}} - [\widehat{\Theta}_{[1,1], m-1}, [\mathcal{O}((l - 1)\vec{c})]]_{\mathbf{v}^{-2}} * [K_{\delta}] \\ &\quad + \mathbf{v}^{-1} [\widehat{\Theta}_{[1,1], m}, [\mathcal{O}(l\vec{c})]] + \mathbf{v}^{-1} [\widehat{\Theta}_{[1,1], m-2}, [\mathcal{O}(l\vec{c})]] * [K_{\delta}]. \end{aligned}$$

Clearly,  $\widehat{Y}_{m|l} = 0$  if  $m \leq 0$ . Using Proposition 9.5, we have  $\widehat{Y}_{m|l} = 0$  for any  $m$ .

We also denote for  $n \in \mathbb{Z}$ ,

$$\begin{aligned} \widehat{X}_{n|l} &= \left( \sum_{p \geq 0} \mathbf{v}^{2p+1} [[\mathcal{O}((l + 1)\vec{c})], \widehat{\Theta}_{[1,1], n-2p-1}]_{\mathbf{v}^{-2}} + \sum_{p \geq 1} \mathbf{v}^{2p-1} [2]_{\mathbf{v}} [\widehat{\Theta}_{[1,1], n-2p}, [\mathcal{O}(l\vec{c})]] \right. \\ &\quad \left. - \sum_{p \geq 0} \mathbf{v}^{2p+1} [\widehat{\Theta}_{[1,1], n-2p-1}, [\mathcal{O}((l - 1)\vec{c})]]_{\mathbf{v}^{-2}} * [K_{\delta}] \right) * [K_{\delta}]^{p+1} + [\widehat{\Theta}_{[1,1], n}, [\mathcal{O}(l\vec{c})]] * [K_{\delta}]. \end{aligned}$$

Clearly,  $\widehat{X}_{n|l} = 0$  for  $n \leq 0$ . By using the same argument as in [LW21b, Lemma 4.10] and Proposition 9.5, we have  $\widehat{X}_{n|l} = 0$  for any  $n$ .

Using the same argument as in [LW21b, Lemmas 4.11 and 4.14], we have

$$\begin{aligned}
 & \widehat{\mathbb{R}}(k_1, k_2 + 1 | l; [1, 1], \star) + \widehat{\mathbb{R}}(k_1 + 1, k_2 | l; [1, 1], \star) - [2]_{\mathbf{v}} \widehat{\mathbb{R}}(k_1 + 1, k_2 + 1 | l - 1; [1, 1], \star) \\
 &= -[2]_{\mathbf{v}}^2 \widehat{X}_{k_2 - k_1 | l - 1} * [K_{\delta}]^{k_1} * [K_{\alpha_{11}}] \\
 &\quad - \left( [[\mathcal{O}(l\vec{c})], \widehat{\Theta}_{[1,1], k_2 - k_1 + 1}]_{\mathbf{v}^{-2}} * -\mathbf{v}^{-2} [[\mathcal{O}(l\vec{c})], \widehat{\Theta}_{[1,1], k_2 - k_1 - 1}]_{\mathbf{v}^{-2}} * [K_{\delta}] \right) * [K_{k_1 \delta + \alpha_{11}}] \\
 &\quad + \{k_1 \leftrightarrow k_2\}, \\
 & \widehat{\mathbb{R}}(k_1, k_2 + 1 | l; [1, 1], \star) + \widehat{\mathbb{R}}(k_1 + 1, k_2 | l; [1, 1], \star) - [2]_{\mathbf{v}} \widehat{\mathbb{R}}(k_1, k_2 | l + 1; [1, 1], \star) \\
 &= [2]_{\mathbf{v}}^2 \widehat{X}_{k_2 - k_1 - 1 | l} * [K_{\delta}]^{k_1} * [K_{\alpha_{11}}] + [2]_{\mathbf{v}} \widehat{Y}_{k_2 - k_1 + 1 | l} [K_{\delta}]^{k_1} * [K_{\alpha_{11}}] \\
 &\quad - \left( [\widehat{\Theta}_{[1,1], k_2 - k_1 + 1}, [\mathcal{O}(l\vec{c})]]_{\mathbf{v}^{-2}} - \mathbf{v}^{-2} [\widehat{\Theta}_{[1,1], k_2 - k_1 - 1}, [\mathcal{O}(l\vec{c})]]_{\mathbf{v}^{-2}} * [K_{\delta}] \right) * [K_{k_1 \delta + \alpha_{11}}] \\
 &\quad + \{k_1 \leftrightarrow k_2\}.
 \end{aligned}$$

Combining the above computations, for any  $k_1, k_2, l \in \mathbb{Z}$  we have

$$\widehat{\mathbb{S}}(k_1, k_2 + 1 | l; [1, 1], \star) + \widehat{\mathbb{S}}(k_1 + 1, k_2 | l; [1, 1], \star) - [2]_{\mathbf{v}} \widehat{\mathbb{S}}(k_1 + 1, k_2 + 1 | l - 1; [1, 1], \star) \tag{9.9}$$

$$\begin{aligned}
 &= (1 - q)^2 \left( \widehat{\mathbb{R}}(k_1, k_2 + 1 | l; [1, 1], \star) + \widehat{\mathbb{R}}(k_1 + 1, k_2 | l; [1, 1], \star) \right. \\
 &\quad \left. - [2]_{\mathbf{v}} \widehat{\mathbb{R}}(k_1 + 1, k_2 + 1 | l - 1; [1, 1], \star) \right),
 \end{aligned}$$

$$\widehat{\mathbb{S}}(k_1, k_2 + 1 | l; [1, 1], \star) + \widehat{\mathbb{S}}(k_1 + 1, k_2 | l; [1, 1], \star) - [2]_{\mathbf{v}} \widehat{\mathbb{S}}(k_1, k_2 | l + 1; [1, 1], \star) \tag{9.10}$$

$$\begin{aligned}
 &= (1 - q)^2 \left( \widehat{\mathbb{R}}(k_1, k_2 + 1 | l; [1, 1], \star) + \widehat{\mathbb{R}}(k_1 + 1, k_2 | l; [1, 1], \star) \right. \\
 &\quad \left. - [2]_{\mathbf{v}} \widehat{\mathbb{R}}(k_1, k_2 | l + 1; [1, 1], \star) \right).
 \end{aligned}$$

By Lemma 9.6, we have

$$\widehat{\mathbb{S}}(0, 0 | l; [1, 1], \star) = (1 - q)^2 \widehat{\mathbb{R}}(0, 0 | l; [1, 1], \star).$$

By induction and using (9.9)–(9.10), one can obtain that

$$\widehat{\mathbb{S}}(k_1, k_2 | l; [1, 1], \star) = (1 - q)^2 \widehat{\mathbb{R}}(k_1, k_2 | l; [1, 1], \star)$$

for any  $k_1, k_2, l \in \mathbb{Z}$ . □

## 10. DERIVED EQUIVALENCE AND TWO PRESENTATIONS OF AFFINE TYPE A

Let  $\mathbb{X}$  be a weighted projective line of domestic type. Let  $Q$  be an acyclic quiver of the associated type. In this section, we establish a derived equivalence between  $\mathcal{D}^b(\mathcal{C}_1(\text{coh}(\mathbb{X})))$  and  $\mathcal{D}^b(\mathcal{C}_1(\text{rep}_{\mathbf{k}}^{\text{nil}}(Q^{\text{op}})))$ , which induces an isomorphism of  ${}^i\widetilde{\mathcal{H}}(\mathbb{X}_{\mathbf{k}})$  and  ${}^i\widetilde{\mathcal{H}}(\mathbf{k}Q^{\text{op}})$ . Moreover, if  $\mathbb{X}$  is of weight type  $(p_1, p_2)$ , then it induces the isomorphism of the  $i$ quantum group in the Serre and Drinfeld type presentations.

**10.1. A derived equivalence.** Let  $\mathbb{X}$  be a weighted projective line of domestic type. Then there exists a tilting sheaf  $T$  such that  $\text{End}_{\mathbb{X}}(T)$  is hereditary of affine type ADE. Let  $Q$  be the quiver such that  $\text{End}_{\mathbb{X}}(T) \cong \mathbf{k}Q$ . Then there is a derived equivalence

$$\text{RHom}_{\mathbb{X}}(T, -) : \mathcal{D}^b(\text{coh}(\mathbb{X})) \xrightarrow{\cong} \mathcal{D}^b(\text{rep}_{\mathbf{k}}^{\text{nil}}(Q^{\text{op}})). \quad (10.1)$$

Using the proof of [LRW23, Proposition 5.9], one can obtain the following remark.

*Remark 10.1.* Let  $T$  be the tilting sheaf as above. Then  $K_T = (T \oplus T, d)$  is a tilting object in  $\mathcal{D}^b(\mathcal{C}_1(\text{coh}(\mathbb{X}_{\mathbf{k}})))$ , which gives rise to an equivalence:

$$\text{RHom}(K_T, -) : \mathcal{D}^b(\mathcal{C}_1(\text{coh}(\mathbb{X}_{\mathbf{k}}))) \xrightarrow{\cong} \mathcal{D}^b(\mathcal{C}_1(\text{rep}_{\mathbf{k}}^{\text{nil}}(Q^{\text{op}}))).$$

Let  $\mathcal{U} := \{M \in \text{coh}(\mathbb{X}) \mid \text{Ext}_{\mathbb{X}}^1(T, M) = 0\}$  and  $\mathcal{V} := \{M \in \text{coh}(\mathbb{X}) \mid \text{Hom}_{\mathbb{X}}(T, M) = 0\}$ . Then  $(\mathcal{U}, \mathcal{V})$  is a *torsion pair* of  $\text{coh}(\mathbb{X})$ . Let  $\mathcal{C}_1(\mathcal{U})$  and  $\mathcal{C}_1(\mathcal{V})$  be the categories of all 1-periodic complexes in  $\mathcal{U}$  and  $\mathcal{V}$  respectively.

**Lemma 10.2** ([LRW23, Lemma 5.8]).  *$(\mathcal{C}_1(\mathcal{U}), \mathcal{C}_1(\mathcal{V}))$  is a torsion pair of  $\mathcal{C}_1(\text{coh}(\mathbb{X}))$ . In particular, any  $M \in \mathcal{C}_1(\text{coh}(\mathbb{X}))$  admits a short exact sequence of the form*

$$0 \longrightarrow M \longrightarrow X_M \longrightarrow T_M \longrightarrow 0, \quad (10.2)$$

where  $X_M \in \mathcal{C}_1(\mathcal{U})$  and  $T_M \in \text{add } K_T$ .

**Proposition 10.3.** *Let  $F = \text{Hom}(K_T, -)$ . Then there exists an isomorphism*

$$\begin{aligned} F : {}^v\tilde{\mathcal{H}}(\mathbb{X}_{\mathbf{k}}) &\xrightarrow{\cong} {}^v\tilde{\mathcal{H}}(\mathbf{k}Q^{\text{op}}) \\ [M] &\mapsto [F(T_M)]^{-1} * [F(X_M)], \end{aligned}$$

where  $X_M \in \mathcal{C}_1(\mathcal{U})$  and  $T_M \in \text{add } K_T$ , are defined in the short exact sequence (10.2).

*Proof.* The proof is the same as in [LRW23, Proposition 5.10], hence omitted here.  $\square$

**10.2. A commutative diagram.** In this subsection, we always assume that  $\mathbb{X}$  is of weight type  $(p_1, p_2)$ . Let  $T = \bigoplus_{0 \leq \vec{x} \leq \vec{c}} \mathcal{O}_{\mathbb{X}}(\vec{x})$  be the canonical tilting sheaf in  $\text{coh}(\mathbb{X})$ . Then  $\text{End}_{\mathbb{X}}(T) = \mathbf{k}Q$  is the canonical algebra [Rin84] with  $Q = C_{p_1, p_2}$  as shown below:

$$\begin{array}{ccccccc} & & \begin{array}{c} \circ \\ \uparrow \\ \circ \end{array} & \xrightarrow{[1, 1]} & \cdots & \xrightarrow{[1, p_1 - 2]} & \xrightarrow{[1, p_1 - 1]} & \circ \\ & \searrow & & & & & & \uparrow \\ \star & & \circ & \xrightarrow{[2, 1]} & \cdots & \xrightarrow{[2, p_2 - 2]} & \xrightarrow{[2, p_2 - 1]} & \circ \\ & & & & & & & \downarrow \\ & & & & & & & \circ \end{array} \quad (10.3)$$

Let  $\tilde{\mathbf{U}}^i = \tilde{\mathbf{U}}^i(\widehat{\mathfrak{sl}}_n)$  for  $n = p_1 + p_2$ . Let  $\tilde{\psi} : \tilde{\mathbf{U}}^i_{|v=\mathbf{v}} \rightarrow {}^v\tilde{\mathcal{H}}(\mathbf{k}C_{(p_1, p_2)}^{\text{op}})$  be the algebra embedding obtained in Theorem 3.8. The following result is a generalization of [LRW23, Theorem 5.11] for  $\tilde{\mathbf{U}}^i(\widehat{\mathfrak{sl}}_2)$ .

**Theorem 10.4.** *We have the following commutative diagram of algebra homomorphisms*

$$\begin{array}{ccc} Dr\tilde{\mathbf{U}}|_{v=\mathbf{v}} & \xrightarrow{\Phi} & \tilde{\mathbf{U}}|_{v=\mathbf{v}} \\ \downarrow \Omega & & \downarrow \tilde{\psi} \\ {}^i\tilde{\mathcal{H}}(\mathbb{X}_{\mathbf{k}}) & \xrightarrow{F} & {}^i\tilde{\mathcal{H}}(\mathbf{k}C_{(p_1, p_2)}^{op}). \end{array}$$

*In particular, the homomorphism  $\Omega$  is injective.*

*Proof.* A direct computation shows that

$$\begin{aligned} \tilde{\psi} \circ \Phi(B_{\star, 0}) &= \tilde{\psi}(B_{\star}) = \frac{1}{1-q}[S_{\star}], \\ F \circ \Omega(B_{\star, 0}) &= \frac{1}{1-q}F([\mathcal{O}]) = \frac{1}{1-q}[S_{\star}], \end{aligned}$$

and then  $\tilde{\psi} \circ \Phi(B_{\star, 0}) = F \circ \Omega(B_{\star, 0})$ .

Denote by  $M(l\delta - \alpha_{\star})$  the indecomposable  $\mathbf{k}C_{(p_1, p_2)}^{op}$ -module with dimension vector  $l\delta - \alpha_{\star}$ . Similar to (5.12), we have

$$\Gamma_{\omega_{\star}}([S_{\star}]) = [M(\delta - \alpha_{\star})] * [K_{\delta - \alpha_{\star}}]^{-1},$$

and then

$$\tilde{\psi} \circ \Phi(B_{\star, -1}) = \tilde{\psi}(\mathbf{T}_{\omega_{\star}}(B_{\star})) = \frac{-1}{q-1}\Gamma_{\omega_{\star}}([S_{\star}]) = \frac{-1}{q-1}[M(\delta - \alpha_{\star})] * [K_{\delta - \alpha_{\star}}]^{-1}.$$

On the other hand, there exists a short exact sequence in  $\text{coh}(\mathbb{X})$

$$0 \longrightarrow \mathcal{O}(-\vec{c}) \longrightarrow \mathcal{O} \oplus \mathcal{O} \xrightarrow{(g_1, g_2)} \mathcal{O}(\vec{c}) \longrightarrow 0,$$

and then a short exact sequence in  $\mathcal{C}_1(\text{coh}(\mathbb{X}))$

$$0 \longrightarrow \mathcal{O}(-\vec{c}) \longrightarrow N^{\bullet} := (\mathcal{O} \oplus \mathcal{O} \oplus \mathcal{O}(\vec{c}), d) \longrightarrow K_{\mathcal{O}(\vec{c})} \longrightarrow 0,$$

where  $d = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ g_1 & g_2 & 0 \end{bmatrix}$ . Note that

$$F([N^{\bullet}]) = [(S_{\star} \oplus S_{\star} \oplus P_c, F(d))] = [M(\delta - \alpha_{\star})] * [K_{2\alpha_{\star}}].$$

Here  $P_c$  is the projective  $\mathbf{k}C_{(p_1, p_2)}^{op}$ -module corresponding to the vertex  $c$ . Then

$$\begin{aligned} F([\mathcal{O}(-\vec{c})]) &= F([N^{\bullet}]) * [F(K_{\mathcal{O}(\vec{c})})]^{-1} \\ &= [M(\delta - \alpha_{\star})] * [K_{2\alpha_{\star}}] * [K_{P_c}]^{-1} \\ &= [M(\delta - \alpha_{\star})] * [K_{\delta - \alpha_{\star}}]^{-1} \end{aligned}$$

by Proposition 10.3. Therefore,

$$F \circ \Omega(B_{\star, -1}) = \frac{-1}{q-1}F([\mathcal{O}(-\vec{c})]) = \frac{-1}{q-1}[M(\delta - \alpha_{\star})] * [K_{\delta - \alpha_{\star}}]^{-1} = \tilde{\psi} \circ \Phi(B_{\star, -1}).$$

We consider  $S_{ij} \in \text{coh}(X)$  with  $j \neq 0$  in the following. Then we have a short exact sequence

$$0 \longrightarrow \mathcal{O}((j-1)\vec{x}_i) \xrightarrow{x_i} \mathcal{O}(j\vec{x}_i) \longrightarrow S_{ij} \rightarrow 0.$$

Similarly to the above, we have

$$F([S_{ij}]) = [S_{[i,j]}],$$

where  $S_{[i,j]}$  is the simple  $\mathbf{k}C_{p_1, p_2}^{op}$ -module corresponding to  $[i, j]$ . Then

$$F \circ \Omega(B_{[i,j],0}) = \frac{-1}{q-1} F([S_{ij}]) = \frac{-1}{q-1} [S_{[i,j]}] = \tilde{\psi}(B_{[i,j]}) = \tilde{\psi} \circ \Phi(B_{[i,j],0}).$$

Finally, we verify that

$$\begin{aligned} F \circ \Omega(\mathbb{K}_{[i,j]}) &= F([K_{S_{ij}}]) = [K_{S_{[i,j]}}] = \tilde{\psi}(\mathbb{K}_{[i,j]}) = \tilde{\psi} \circ \Phi(\mathbb{K}_{[i,j]}), \\ F \circ \Omega(\mathbb{K}_\star) &= F([K_{\mathcal{O}}]) = [K_{S_\star}] = \tilde{\psi}(\mathbb{K}_\star) = \tilde{\psi} \circ \Phi(\mathbb{K}_\star), \\ F \circ \Omega(C) &= F([K_{\mathcal{O}(\tilde{c})}] * [K_{\mathcal{O}}]^{-1}) = [K_{P_c}] * [K_{S_\star}]^{-1} = [K_\delta] = \tilde{\psi} \circ \Phi(C). \end{aligned}$$

Summarizing, by using Lemma 2.7, we have proved  $F \circ \Omega = \tilde{\psi} \circ \Phi$ .

The injectivity of  $\Omega$  follows by the injectivity of  $\tilde{\psi}$  and the commutative diagram.  $\square$

*Remark 10.5.* For  $\mathfrak{g}$  of type DE and  $Q$  the associated affine quiver, we also have a derived equivalence  $\mathcal{D}^b(\text{coh}(\mathbb{X}_{\mathbf{k}})) \simeq \mathcal{D}^b(\text{rep}_{\mathbf{k}}(Q^{op}))$ , which induces a derived equivalence on the categories of 1-periodic complexes. Similar to Proposition 10.3, there is an algebra isomorphism  $F : {}^v\tilde{\mathcal{H}}(\mathbb{X}_{\mathbf{k}}) \xrightarrow{\cong} {}^v\tilde{\mathcal{H}}(\mathbf{k}Q^{op})$ . However, the morphism  $\Phi$  can not be the one induced by  $F$ , in fact it can not be induced by any derived equivalent functors by the same argument as in [DJX12, §9.2].

*Remark 10.6.* For the commutative diagram obtained in Theorem 10.4, the corresponding result for quantum groups is only known for quantum affine  $\mathfrak{sl}_2$  [BS12b] by using the approach of Drinfeld double Ringel-Hall algebras. In fact, Burban-Schiffmann [BS12b] interpreted the Drinfeld-Beck isomorphism of quantum affine  $\mathfrak{sl}_2$  via Beilinson's derived equivalence between the Kronecker quiver and the projective line, which gives an isomorphism of their Drinfeld double Ringel-Hall algebras [Cr10].

We also expect the corresponding result of Theorem 10.4 holds for quantum groups of affine type  $A$ , and it will be considered elsewhere by using semi-derived Ringel-Hall algebras.

## APPENDIX A. PROOFS OF RESULTS IN SECTION 8

**A.1. Proof of Lemma 8.2.** We first introduce the following notations. Denote by  $\mathfrak{S}(a, b)$  the set of pairs of coprime homogeneous polynomials  $(J, L) \in \mathbf{k}[X, Y]$  of degree  $a$  and  $b$  respectively, and denote by  $\mathfrak{T}(a, b)$  its subset consisting of those  $(J, L)$ 's such that  $X$  does not divide  $J$ . The cardinalities of  $\mathfrak{S}(a, b)$  and  $\mathfrak{T}(a, b)$  are denoted by  $\mathfrak{s}(a, b)$  and  $\mathfrak{t}(a, b)$  respectively.

**Lemma A.1.** *For any  $a, b \geq 0$ ,*

$$\mathfrak{t}(a, b) = \begin{cases} (q-1)(q^{b+1}-1), & \text{if } a = 0, \\ (q-1)^2 q^{a+b}, & \text{if } a \neq 0. \end{cases}$$

*Consequently,  $\mathfrak{t}(a, b) = \mathfrak{t}(a-k, b+k)$  for  $1 \leq k \leq a-1$ .*



*Proof.* Recall from [BKa01, Lemma 9] that

$$\mathfrak{s}(a, b) = \begin{cases} (q-1)(q^{a+b+1} - 1), & \text{if } ab = 0, \\ (q-1)(q^2 - 1)q^{a+b-1}, & \text{if } ab \neq 0. \end{cases}$$

By definition of  $\mathfrak{S}(a, b)$  and  $\mathfrak{T}(a, b)$ , we know that  $\mathfrak{t}(0, b) = \mathfrak{s}(0, b) = (q-1)(q^{b+1} - 1)$  for  $b \geq 1$ , and  $\mathfrak{t}(a, 0) = \mathfrak{s}(a, 0) - \mathfrak{s}(a-1, 0) = (q-1)(q^{a+1} - q^a) = (q-1)^2 q^a$  for  $a \geq 1$ . Moreover, for any  $a, b \geq 1$ , according to [Sch04, (6.10)], we have

$$\mathfrak{t}(a, b) = \mathfrak{s}(a, b) - \mathfrak{t}(b, a-1).$$

Then by induction on  $a+b$ , we get

$$\mathfrak{t}(a, b) = (q-1)(q^2 - 1)q^{a+b-1} - (q-1)^2 q^{a+b-1} = (q-1)^2 q^{a+b}.$$

The proof is completed.  $\square$

**Corollary A.2.** *For any  $l \geq 1$ , we have*

$$[\mathcal{O}(l\vec{c})] * [\mathcal{O}(\vec{x}_1)] - \mathbf{v}^{-1}[\mathcal{O}(l\vec{c} + \vec{x}_1)] * [\mathcal{O}] \quad (\text{A.1})$$

$$= \mathbf{v}^{-l-1} \left( [\mathcal{O}(l\vec{c}) \oplus \mathcal{O}(\vec{x}_1)] - [\mathcal{O}(l\vec{c} + \vec{x}_1) \oplus \mathcal{O}] \right),$$

$$[\mathcal{O}(l\vec{c} + \vec{x}_1)] * [\mathcal{O}(\vec{c})] - \mathbf{v}^{-1}[\mathcal{O}((l+1)\vec{c})] * [\mathcal{O}(\vec{x}_1)] \quad (\text{A.2})$$

$$= \mathbf{v}^{-l-1} \left( [\mathcal{O}(l\vec{c} + \vec{x}_1) \oplus \mathcal{O}(\vec{c})] - [\mathcal{O}((l+1)\vec{c}) \oplus \mathcal{O}(\vec{x}_1)] \right).$$

*Proof.* For any  $1 \leq k \leq l-1$ , set  $M_k := \mathcal{O}((l-k)\vec{c}) \oplus \mathcal{O}(k\vec{c} + \vec{x}_1)$ . According to [Sch04, Lemma 6.5], we have

$$F_{\mathcal{O}(l\vec{c}), \mathcal{O}(\vec{x}_1)}^{M_k} = \frac{\mathfrak{t}(k, l-k-1)}{q-1}, \quad F_{\mathcal{O}(l\vec{c} + \vec{x}_1), \mathcal{O}}^{M_k} = \frac{\mathfrak{t}(l-k, k)}{q-1}.$$

Then it follows from Riedtmann-Peng formula that

$$|\text{Ext}^1(\mathcal{O}(l\vec{c}), \mathcal{O}(\vec{x}_1))_{M_k}| = \frac{(q-1) \cdot \mathfrak{t}(k, l-k-1)}{|\text{Aut}(M_k)|},$$

$$|\text{Ext}^1(\mathcal{O}(l\vec{c} + \vec{x}_1), \mathcal{O})_{M_k}| = \frac{(q-1) \cdot \mathfrak{t}(l-k, k)}{|\text{Aut}(M_k)|}.$$

So

$$[\mathcal{O}(l\vec{c})] * [\mathcal{O}(\vec{x}_1)] = \mathbf{v}^{-l-1} \left( [\mathcal{O}(l\vec{c}) \oplus \mathcal{O}(\vec{x}_1)] + \sum_{k=1}^{l-1} \frac{(q-1) \cdot \mathfrak{t}(k, l-k-1)}{|\text{Aut}(M_k)|} [M_k] \right), \quad (\text{A.3})$$

$$\begin{aligned} [\mathcal{O}(l\vec{c} + \vec{x}_1)] * [\mathcal{O}] &= \mathbf{v}^{-l} \left( [\mathcal{O}(l\vec{c} + \vec{x}_1) \oplus \mathcal{O}] + (q-1)[\mathcal{O}(l\vec{c}) \oplus \mathcal{O}(\vec{x}_1)] \right. \\ &\quad \left. + \sum_{k=1}^{l-1} \frac{(q-1) \cdot \mathfrak{t}(k, l-k)}{|\text{Aut}(M_k)|} [M_k] \right). \end{aligned} \quad (\text{A.4})$$

By Lemma A.1, we have  $\mathfrak{t}(k, l-k) = q \cdot \mathfrak{t}(k, l-k-1)$  for  $1 \leq k \leq l-1$ , and then (A.1) follows by combining (A.3)–(A.4).

For (A.2), we define

$$\mathcal{S} := \{S_{1,j} \mid 2 \leq j \leq p_1 - 1\} \cup \{S_{ij} \mid 2 \leq i \leq \mathfrak{t}, 1 \leq j \leq p_i - 1\}. \quad (\text{A.5})$$

Then the left perpendicular category  ${}^{\perp}\mathcal{S}$  is equivalent to  $\text{coh}(\mathbb{Y}_{\mathbf{k}})$ , where  $\mathbb{Y}_{\mathbf{k}}$  is a weighted projective line of weight type  $\mathbf{p} = (2, 1)$ . By similar arguments as in §6.2, we have an embedding

$$F_{\mathbb{X}, \mathbb{Y}} : {}^i\tilde{\mathcal{H}}(\mathbb{Y}_{\mathbf{k}}) \longrightarrow {}^i\tilde{\mathcal{H}}(\mathbb{X}_{\mathbf{k}}). \quad (\text{A.6})$$

Observe that each term in (A.1) and (A.2) belongs to the image of  $F_{\mathbb{X}, \mathbb{Y}}$ , hence we can restrict to  $\mathbb{Y}_{\mathbf{k}}$ . In this case,  $2\vec{x}_1 = \vec{c}$ . Then (A.2) follows from (A.1) by twisting with  $\vec{x}_1$ .  $\square$

For any  $m \geq 0$ , define

$$\hat{\Theta}_{*,m}^+ := \frac{1}{(q-1)^2 \mathbf{v}^{m-1}} \sum_{0 \neq f: \mathcal{O}(s\vec{c}) \rightarrow \mathcal{O}((m+s)\vec{c} + \vec{x}_1)} [\text{Coker}(f)], \quad (\text{A.7})$$

$$\hat{\Theta}_{*,m}^- := \frac{1}{(q-1)^2 \mathbf{v}^{m-1}} \sum_{0 \neq f: \mathcal{O}(s\vec{c} + \vec{x}_1) \rightarrow \mathcal{O}((m+s)\vec{c})} [\text{Coker}(f)], \quad (\text{A.8})$$

which are independent of  $s \in \mathbb{Z}$ . For convenience we set  $\hat{\Theta}_{*,m}^{\pm} = 0$  for  $m < 0$ .

**Lemma A.3.** *For any  $l \geq 1$ ,*

$$[[\mathcal{O}(l\vec{c}), [\mathcal{O}(\vec{x}_1)]]_{\mathbf{v}^{-1}} + [[\mathcal{O}], [\mathcal{O}(l\vec{c} + \vec{x}_1)]]_{\mathbf{v}^{-1}}] \quad (\text{A.9})$$

$$\begin{aligned} &= (\mathbf{v} - \mathbf{v}^{-1})^2 (\hat{\Theta}_{*,l}^+ * [K_{\mathcal{O}}] - \hat{\Theta}_{*,l}^- * [K_{\mathcal{O}(\vec{x}_1)}]), \\ &[[\mathcal{O}(l\vec{c} + \vec{x}_1), [\mathcal{O}(\vec{c})]]_{\mathbf{v}^{-1}} + [[\mathcal{O}(\vec{x}_1)], [\mathcal{O}((l+1)\vec{c})]]_{\mathbf{v}^{-1}}] \quad (\text{A.10}) \\ &= (\mathbf{v} - \mathbf{v}^{-1})^2 (\mathbf{v} \hat{\Theta}_{*,l+1}^- * [K_{\mathcal{O}(\vec{x}_1)}] - \mathbf{v}^{-1} \hat{\Theta}_{*,l-1}^+ * [K_{\mathcal{O}(\vec{c})}]), \end{aligned}$$

*Proof.* Observe that

$$\begin{aligned} &[\mathcal{O}(\vec{x}_1)] * [\mathcal{O}(l\vec{c})] \\ &= \mathbf{v}^{-l} [\mathcal{O}(\vec{x}_1) \oplus \mathcal{O}(l\vec{c})] + \mathbf{v}^{-l} \sum_{0 \neq f: \mathcal{O}(\vec{x}_1) \rightarrow \mathcal{O}(l\vec{c})} [\text{Coker}(f)] * [K_{\mathcal{O}(\vec{x}_1)}] \\ &= \mathbf{v}^{-l} [\mathcal{O}(\vec{x}_1) \oplus \mathcal{O}(l\vec{c})] + \mathbf{v}^{-1} (q-1)^2 \hat{\Theta}_{*,l}^- * [K_{\mathcal{O}(\vec{x}_1)}], \end{aligned}$$

and

$$\begin{aligned} [\mathcal{O}] * [\mathcal{O}(l\vec{c} + \vec{x}_1)] &= \mathbf{v}^{-l-1} [\mathcal{O} \oplus \mathcal{O}(l\vec{c} + \vec{x}_1)] + \mathbf{v}^{-l-1} \sum_{0 \neq g: \mathcal{O} \rightarrow \mathcal{O}(l\vec{c} + \vec{x}_1)} [\text{Coker}(g)] * [K_{\mathcal{O}}] \\ &= \mathbf{v}^{-l-1} [\mathcal{O} \oplus \mathcal{O}(l\vec{c} + \vec{x}_1)] + \mathbf{v}^{-2} (q-1)^2 \hat{\Theta}_{*,l}^+ * [K_{\mathcal{O}}]. \end{aligned}$$

Moreover, by (A.1) we have

$$\begin{aligned} &\mathbf{v}^{-1} [\mathcal{O}(l\vec{c} + \vec{x}_1)] * [\mathcal{O}] - [\mathcal{O}(l\vec{c})] * [\mathcal{O}(\vec{x}_1)] \\ &= \mathbf{v}^{-l-1} ([\mathcal{O}(l\vec{c} + \vec{x}_1) \oplus \mathcal{O}] - [\mathcal{O}(l\vec{c}) \oplus \mathcal{O}(\vec{x}_1)]). \end{aligned}$$

Combining the above formulas, (A.9) follows.

For (A.10), similar to the proof of (A.2), we can restrict to consider  $\mathbb{X}_{\mathbf{k}}$  of weight type  $\mathbf{p} = (2, 1)$ . In this case, by definition we have

$$\hat{\Theta}_{*,l}^+(\vec{x}_1) = \mathbf{v} \hat{\Theta}_{*,l+1}^-, \quad \hat{\Theta}_{*,l}^-(\vec{x}_1) = \mathbf{v}^{-1} \hat{\Theta}_{*,l-1}^+.$$

Then (A.10) follows from (A.9) by twisting with  $\vec{x}_1$ .  $\square$

**Lemma A.4.** For any  $m \in \mathbb{Z}$ ,

$$\widehat{\Theta}_{\star, m}^+ = q\widehat{\Theta}_{\star, m-2}^+ * [K_\delta] + \frac{q}{q-1} [[S_{1,1}], \widehat{\Theta}_{\star, m}]_{\mathbf{v}^{-2}} - \frac{q\mathbf{v}}{q-1} [\widehat{\Theta}_{\star, m-1}, [S_{1,0}^{(p_1-1)}]]_{\mathbf{v}^{-2}} * [K_{\alpha_{11}}], \quad (\text{A.11})$$

$$\widehat{\Theta}_{\star, m}^- = q\widehat{\Theta}_{\star, m-2}^- * [K_\delta] + \frac{\mathbf{v}}{q-1} [\widehat{\Theta}_{\star, m-1}, [S_{1,0}^{(p_1-1)}]]_{\mathbf{v}^{-2}} - \frac{q}{q-1} [[S_{1,1}], \widehat{\Theta}_{\star, m-2}]_{\mathbf{v}^{-2}} * [K_{\delta-\alpha_{11}}]. \quad (\text{A.12})$$

*Proof.* For  $m \leq 0$ , the above equalities hold obviously by definition. We assume  $m > 0$  in the following.

For any  $r \geq 0$ , an easy computation shows that

$$[S_{1,0}^{(p_1-1)}] * [S_{1,0}^{(rp_1)}] = [S_{1,0}^{(rp_1)} \oplus S_{1,0}^{(p_1-1)}], \quad (\text{A.13})$$

$$\begin{aligned} [S_{1,0}^{(rp_1)}] * [S_{1,0}^{(p_1-1)}] &= \frac{1}{q} \left( [S_{1,0}^{(rp_1)} \oplus S_{1,0}^{(p_1-1)}] + (q-1)[S_{1,0}^{((r+1)p_1-1)}] \right) \\ &\quad + q(q-1)[S_{1,1}^{((r-1)p_1+1)}] * [K_{\delta-\alpha_{11}}]. \end{aligned} \quad (\text{A.14})$$

Here and in the following, it is understood that the terms involving  $S_{1,1}^{((r-1)p_1+1)}$  do not exist if  $r = 0$ . Hence it follows by combining (A.13) and (A.14) that

$$[[S_{1,0}^{(rp_1)}], [S_{1,0}^{(p_1-1)}]]_{\mathbf{v}^{-2}} = \frac{q-1}{q} [S_{1,0}^{((r+1)p_1-1)}] + (q-1)[S_{1,1}^{((r-1)p_1+1)}] * [K_{\delta-\alpha_{11}}]. \quad (\text{A.15})$$

For any non-zero map  $f : \mathcal{O} \rightarrow \mathcal{O}(k\vec{c})$ , assume  $f = x_1^{rp_1} \cdot g$  for some  $r \geq 0$  and  $x_1 \nmid g$ . So  $\text{Coker}(f) \cong S_{1,0}^{(rp_1)} \oplus M$ , where  $M$  has no direct summand supported at  $\lambda_1 = \infty$ . Note that any two torsion sheaves supported at distinct points have zero Hom and Ext<sup>1</sup>-spaces. Hence  $[M] * [X] = [M \oplus X] = [X] * [M]$  for any  $X \in \mathcal{T}_{\lambda_1}$ . Therefore,

$$\begin{aligned} [[\text{Coker}(f)], [S_{1,0}^{(p_1-1)}]]_{\mathbf{v}^{-2}} &= [M] * [[S_{1,0}^{(rp_1)}], [S_{1,0}^{(p_1-1)}]]_{\mathbf{v}^{-2}} \\ &\stackrel{(\text{A.15})}{=} [M] * \left( \frac{q-1}{q} [S_{1,0}^{((r+1)p_1-1)}] + (q-1)[S_{1,1}^{((r-1)p_1+1)}] * [K_{\delta-\alpha_{11}}] \right) \\ &= \frac{q-1}{q} [M \oplus S_{1,0}^{((r+1)p_1-1)}] + (q-1)[M \oplus S_{1,1}^{((r-1)p_1+1)}] * [K_{\delta-\alpha_{11}}] \\ &= \frac{q-1}{q} [\text{Coker}(f_1)] + (q-1)[\text{Coker}(f_2)] * [K_{\delta-\alpha_{11}}], \end{aligned}$$

where  $f_1 = x_1^{(r+1)p_1-1} \cdot g : \mathcal{O}(\vec{x}_1) \rightarrow \mathcal{O}((k+1)\vec{c})$  and  $f_2 = x_1^{(r-1)p_1+1} \cdot g : \mathcal{O}(\vec{c}) \rightarrow \mathcal{O}(k\vec{c} + \vec{x}_1)$ . Here it is understood that  $f_2$  and then the term involving  $[\text{Coker}(f_2)]$  does not exist if  $r = 0$ . Hence,

$$\begin{aligned} &\sum_{0 \neq f: \mathcal{O} \rightarrow \mathcal{O}(m\vec{c})} [[\text{Coker}(f)], [S_{1,0}^{(p_1-1)}]]_{\mathbf{v}^{-2}} \\ &= \frac{q-1}{q} \sum_{0 \neq f_1: \mathcal{O}(\vec{x}_1) \rightarrow \mathcal{O}((m+1)\vec{c})} [\text{Coker}(f_1)] + (q-1) \sum_{0 \neq f_2: \mathcal{O}(\vec{c}) \rightarrow \mathcal{O}(m\vec{c} + \vec{x}_1)} [\text{Coker}(f_2)] * [K_{\delta-\alpha_{11}}]. \end{aligned}$$

Then it follows from (6.6), (A.7) and (A.8) that

$$[\widehat{\Theta}_{\star, m}, [S_{1,0}^{(p_1-1)}]]_{\mathbf{v}^{-2}} = (\mathbf{v} - \mathbf{v}^{-1})\widehat{\Theta}_{\star, m+1}^- + (\mathbf{v} - \mathbf{v}^{-1})\widehat{\Theta}_{\star, m-1}^+ * [K_{\delta-\alpha_{11}}]. \quad (\text{A.16})$$

Similarly, for any  $r \geq 0$  we have

$$[[S_{1,1}], [S_{1,0}^{(rp_1)}]]_{\mathbf{v}^{-2}} = \frac{q-1}{q} [S_{1,1}^{(rp_1+1)}] + (q-1) [S_{1,0}^{(rp_1-1)}] * [K_{\alpha_{11}}],$$

which implies

$$\begin{aligned} & [[S_{1,1}], \sum_{0 \neq f: \mathcal{O} \rightarrow \mathcal{O}(m\vec{c})} [\text{Coker}(f)]]_{\mathbf{v}^{-2}} \\ &= \frac{q-1}{q} \sum_{0 \neq f_1: \mathcal{O} \rightarrow \mathcal{O}(m\vec{c} + \vec{x}_1)} [\text{Coker}(f_1)] + (q-1) \sum_{0 \neq f_2: \mathcal{O}(\vec{x}_1) \rightarrow \mathcal{O}(m\vec{c})} [\text{Coker}(f_2)] * [K_{\alpha_{11}}]. \end{aligned}$$

Then by definition we have

$$[[S_{1,1}], \widehat{\Theta}_{\star, m}]_{\mathbf{v}^{-2}} = \frac{q-1}{q} \widehat{\Theta}_{\star, m}^+ + (q-1) \widehat{\Theta}_{\star, m}^- * [K_{\alpha_{11}}]. \quad (\text{A.17})$$

Combining (A.16) and (A.17), we obtain (A.11) and (A.12).  $\square$

Now we can give the proof of Lemma 8.2.

*Proof of Lemma 8.2.* For any  $k \in \mathbb{Z}$ , by (4.10), we have

$$[\mathcal{O}(k\vec{c})] * [S_{1,1}] = \mathbf{v} [S_{1,1}] * [\mathcal{O}(k\vec{c})] - (q-1) [\mathcal{O}(k\vec{c} + \vec{x}_1)]. \quad (\text{A.18})$$

Hence,

$$\begin{aligned} & \widehat{S}(k_1, k_2 \mid 0; \star, [1, 1]) \\ &= [S_{1,1}] * [\mathcal{O}(k_1\vec{c})] * [\mathcal{O}(k_2\vec{c})] - [2]_{\mathbf{v}} [\mathcal{O}(k_1\vec{c})] * [S_{1,1}] * [\mathcal{O}(k_2\vec{c})] + [\mathcal{O}(k_1\vec{c})] * [\mathcal{O}(k_2\vec{c})] * [S_{1,1}] \\ &= (\mathbf{v}^{-1} [\mathcal{O}(k_1\vec{c})] * [S_{1,1}] + (\mathbf{v} - \mathbf{v}^{-1}) [\mathcal{O}(k_1\vec{c} + \vec{x}_1)]) * [\mathcal{O}(k_2\vec{c})] \\ &\quad - [2]_{\mathbf{v}} [\mathcal{O}(k_1\vec{c})] * [S_{1,1}] * [\mathcal{O}(k_2\vec{c})] + [\mathcal{O}(k_1\vec{c})] * (\mathbf{v} [S_{1,1}] * [\mathcal{O}(k_2\vec{c})] - (q-1) [\mathcal{O}(k_2\vec{c} + \vec{x}_1)]) \\ &= (\mathbf{v} - \mathbf{v}^{-1}) [\mathcal{O}(k_1\vec{c} + \vec{x}_1)] * [\mathcal{O}(k_2\vec{c})] - (q-1) [\mathcal{O}(k_1\vec{c})] * [\mathcal{O}(k_2\vec{c} + \vec{x}_1)]. \end{aligned}$$

For  $k_1 = k_2$ , the desired formula (8.7) holds by noting that

$$\begin{aligned} \widehat{\mathbb{S}}(k_1, k_1 \mid 0; \star, [1, 1]) &= \widehat{S}(k_1, k_1 \mid 0; \star, [1, 1]) \\ &= - (q-1) [[\mathcal{O}(k_1\vec{c})], [\mathcal{O}(k_1\vec{c} + \vec{x}_1)]]_{\mathbf{v}^{-1}} \\ &= - \mathbf{v}^{-1} (q-1)^2 [S_{1,1}] * [K_{\mathcal{O}(k_1\vec{c})}] \\ &\stackrel{(7.3)}{=} (1-q)^2 \widehat{\mathbb{R}}(k_1, k_1 \mid 0; \star, [1, 1]). \end{aligned}$$

Without loss of generality, we assume  $k_1 < k_2$  in the following. Then we have

$$\begin{aligned} & \widehat{\mathbb{S}}(k_1, k_2 \mid 0; \star, [1, 1]) \\ &= (\mathbf{v} - \mathbf{v}^{-1}) [\mathcal{O}(k_1\vec{c} + \vec{x}_1)] * [\mathcal{O}(k_2\vec{c})] - (q-1) [\mathcal{O}(k_1\vec{c})] * [\mathcal{O}(k_2\vec{c} + \vec{x}_1)] \\ &\quad + (\mathbf{v} - \mathbf{v}^{-1}) [\mathcal{O}(k_2\vec{c} + \vec{x}_1)] * [\mathcal{O}(k_1\vec{c})] - (q-1) [\mathcal{O}(k_2\vec{c})] * [\mathcal{O}(k_1\vec{c} + \vec{x}_1)] \\ &= -(q-1) \left( [[\mathcal{O}(k_2\vec{c})], [\mathcal{O}(k_1\vec{c} + \vec{x}_1)]]_{\mathbf{v}^{-1}} + [[\mathcal{O}(k_1\vec{c})], [\mathcal{O}(k_2\vec{c} + \vec{x}_1)]]_{\mathbf{v}^{-1}} \right) \\ &\stackrel{(A.9)}{=} \mathbf{v}^{-2} (q-1)^3 \left( \widehat{\Theta}_{\star, k_2-k_1}^- * [K_{\mathcal{O}(k_1\vec{c} + \vec{x}_1)}] - \widehat{\Theta}_{\star, k_2-k_1}^+ * [K_{\mathcal{O}(k_1\vec{c})}] \right). \end{aligned}$$

If  $k_2 = k_1 + 1$ , then (8.7) holds since

$$\begin{aligned}
 & \widehat{\mathbb{S}}(k_1, k_1 + 1 \mid 0; \star, [1, 1]) \\
 &= \mathbf{v}^{-2}(q-1)^3 \widehat{\Theta}_{\star,1}^- * [K_{\mathcal{O}(k_1\bar{c}+\bar{x}_1)}] - \mathbf{v}^{-2}(q-1)^3 \widehat{\Theta}_{\star,1}^+ * [K_{\mathcal{O}(k_1\bar{c})}] \\
 &= \mathbf{v}^{-1}[2]_{\mathbf{v}}(q-1)^2 [S_{1,0}^{(p_1-1)}] * [K_{\mathcal{O}(k_1\bar{c}+\bar{x}_1)}] - (q-1)^2 [[S_{1,1}], \widehat{\Theta}_{\star,1}]_{\mathbf{v}^{-2}} * [K_{\mathcal{O}(k_1\bar{c})}] \\
 &= (q-1)^2 \left( -\mathbf{v}^{-1}[2]_{\mathbf{v}} \widehat{B}_{[1,1],-1} * [K_{\delta}] - [\widehat{B}_{[1,1],0}, \widehat{\Theta}_{\star,1}]_{\mathbf{v}^{-2}} \right) * [K_{\mathcal{O}(k_1\bar{c})}] \\
 &\stackrel{(7.4)}{=} (1-q)^2 \widehat{\mathbb{R}}(k_1, k_1 + 1 \mid 0; \star, [1, 1]),
 \end{aligned}$$

where the second equality follows by Lemma A.4.

If  $k_2 > k_1 + 1$ , then by (A.11) and (A.12), we have

$$\begin{aligned}
 & \widehat{\mathbb{S}}(k_1, k_2 \mid 0; \star, [1, 1]) - q\widehat{\mathbb{S}}(k_1, k_2 - 2 \mid 0; \star, [1, 1]) * [K_{\delta}] \\
 &= \mathbf{v}^{-2}(q-1)^3 [K_{\mathcal{O}(k_1\bar{c}+\bar{x}_1)}] * (\widehat{\Theta}_{\star, k_2-k_1}^- - q\widehat{\Theta}_{\star, k_2-k_1-2}^- * [K_{\delta}]) \\
 &\quad - \mathbf{v}^{-2}(q-1)^3 [K_{\mathcal{O}(k_1\bar{c})}] * (\widehat{\Theta}_{\star, k_2-k_1}^+ - q\widehat{\Theta}_{\star, k_2-k_1-2}^+ * [K_{\delta}]) \\
 &= (q-1)^2 \left( \mathbf{v}^{-1} [\widehat{\Theta}_{\star, k_2-k_1-1}, [S_{1,0}^{(p_1-1)}]]_{\mathbf{v}^{-2}} - [[S_{1,1}], \widehat{\Theta}_{\star, k_2-k_1-2}]_{\mathbf{v}^{-2}} * [K_{\delta-\alpha_{11}}] \right) * [K_{\mathcal{O}(k_1\bar{c}+\bar{x}_1)}] \\
 &\quad - (q-1)^2 \left( [[S_{1,1}], \widehat{\Theta}_{\star, k_2-k_1}]_{\mathbf{v}^{-2}} - \mathbf{v} [\widehat{\Theta}_{\star, k_2-k_1-1}, [S_{1,0}^{(p_1-1)}]]_{\mathbf{v}^{-2}} * [K_{S_{1,1}}] \right) * [K_{\mathcal{O}(k_1\bar{c})}] \\
 &= (q-1)^2 \left( -[2]_{\mathbf{v}} [\widehat{\Theta}_{\star, k_2-k_1-1}, \widehat{B}_{[1,1],-1}]_{\mathbf{v}^{-2}} * [K_{\mathcal{O}((k_1+1)\bar{c})}] - [\widehat{B}_{[1,1],0}, \widehat{\Theta}_{\star, k_2-k_1}]_{\mathbf{v}^{-2}} * [K_{\mathcal{O}(k_1\bar{c})}] \right. \\
 &\quad \left. - [\widehat{B}_{[1,1],0}, \widehat{\Theta}_{\star, k_2-k_1-2}]_{\mathbf{v}^{-2}} * [K_{\mathcal{O}((k_1+1)\bar{c})}] \right) \\
 &\stackrel{(7.5)}{=} (q-1)^2 \left( \widehat{\mathbb{R}}(k_1, k_2 \mid 0; \star, [1, 1]) - q\widehat{\mathbb{R}}(k_1, k_2 - 2 \mid 0; \star, [1, 1]) * [K_{\delta}] \right).
 \end{aligned}$$

Therefore, the desired formula (8.7) follows by induction.  $\square$

**A.2. Proof of Lemma 8.3.** In order to prove Lemma 8.3, we need the following lemma.

**Lemma A.5.** *For any  $m \geq 1$ ,*

$$\begin{aligned}
 & [\widehat{\Theta}_{\star, m}, \widehat{B}_{[1,1],-1}] + [\widehat{\Theta}_{\star, m-2}, \widehat{B}_{[1,1],-1}] * [K_{\delta}] \\
 &= \mathbf{v}^{-1} [\widehat{\Theta}_{\star, m-1}, \widehat{B}_{[1,1],0}]_{\mathbf{v}^2} + \mathbf{v} [\widehat{\Theta}_{\star, m-1}, \widehat{B}_{[1,1],-2}]_{\mathbf{v}^{-2}} * [K_{\delta}].
 \end{aligned} \tag{A.19}$$

*Proof.* Recall  $\widehat{\Theta}_{\star, m}$  defined in (6.6). Denote by  $\mathcal{L}_k = \{\text{Coker}(g) \mid g \in \text{Hom}(\mathcal{O}, \mathcal{O}(k\bar{c})), x_1 \nmid g\}$  for any  $k \geq 0$ . Then any  $M \in \mathcal{L}_k$  has no direct summands from  $\mathcal{T}_{\lambda_1}$ , and hence  $[M] * [X] = [M \oplus X] = [X] * [M]$  for any  $X \in \mathcal{T}_{\lambda_1}$ . For any non-zero map  $f : \mathcal{O} \rightarrow \mathcal{O}(k\bar{c})$ , assume  $f = x_1^{rp_1} \cdot g$  for some  $r \geq 0$  and  $x_1 \nmid g$ . Then  $\text{Coker}(f) \cong S_{1,0}^{(rp_1)} \oplus M_{k-r}$ , where  $M_{k-r} \in \mathcal{L}_{k-r}$ . Similar to the proof of Lemma A.4, it suffices to show for any  $r \geq 0$  that

$$\begin{aligned}
 & [[S_{1,0}^{(rp_1)}], \widehat{B}_{[1,1],-1}] + q[[S_{1,0}^{((r-2)p_1)}], \widehat{B}_{[1,1],-1}] * [K_{\delta}] \\
 &= [[S_{1,0}^{((r-1)p_1)}], \widehat{B}_{[1,1],0}]_{\mathbf{v}^2} + q[[S_{1,0}^{((r-1)p_1)}], \widehat{B}_{[1,1],-2}]_{\mathbf{v}^{-2}} * [K_{\delta}],
 \end{aligned}$$

which follows from Lemma 5.4.  $\square$

Using the equalities  $\mathbf{v}^{-1}[2]_{\mathbf{v}}[a, b] = [a, b]_{\mathbf{v}^{-2}} - [b, a]_{\mathbf{v}^{-2}}$  and  $[a, b]_{\mathbf{v}^2} = -\mathbf{v}^2[b, a]_{\mathbf{v}^{-2}}$ , (A.19) can be reformulated as follows

$$\begin{aligned} & [\widehat{\Theta}_{\star, m}, \widehat{B}_{[1,1], -1}]_{\mathbf{v}^{-2}} - [2]_{\mathbf{v}}[\widehat{\Theta}_{\star, m-1}, \widehat{B}_{[1,1], -2}]_{\mathbf{v}^{-2}} * [K_{\delta}] + [\widehat{\Theta}_{\star, m-2}, \widehat{B}_{[1,1], -1}]_{\mathbf{v}^{-2}} * [K_{\delta}] \quad (\text{A.20}) \\ &= [\widehat{B}_{[1,1], -1}, \widehat{\Theta}_{\star, m}]_{\mathbf{v}^{-2}} - [2]_{\mathbf{v}}[\widehat{B}_{[1,1], 0}, \widehat{\Theta}_{\star, m-1}]_{\mathbf{v}^{-2}} + [\widehat{B}_{[1,1], -1}, \widehat{\Theta}_{\star, m-2}]_{\mathbf{v}^{-2}} * [K_{\delta}]. \end{aligned}$$

*Proof of Lemma 8.3.* For any  $k \in \mathbb{Z}$ , by (4.11) we have

$$[S_{1,0}^{(p_1-1)}] * [\mathcal{O}(k\vec{c})] = \mathbf{v}[\mathcal{O}(k\vec{c})] * [S_{1,0}^{(p_1-1)}] - (q-1)[\mathcal{O}((k-1)\vec{c} + \vec{x}_1)] * [K_{\delta-\alpha_{11}}]. \quad (\text{A.21})$$

Hence,

$$\begin{aligned} & [S_{1,0}^{(p_1-1)}] * [\mathcal{O}(k_1\vec{c})] * [\mathcal{O}(k_2\vec{c})] - [2]_{\mathbf{v}}[\mathcal{O}(k_1\vec{c})] * [S_{1,0}^{(p_1-1)}] * [\mathcal{O}(k_2\vec{c})] \\ &+ [\mathcal{O}(k_1\vec{c})] * [\mathcal{O}(k_2\vec{c})] * [S_{1,0}^{(p_1-1)}] \\ &= \left( \mathbf{v}[\mathcal{O}(k_1\vec{c})] * [S_{1,0}^{(p_1-1)}] - (q-1)[\mathcal{O}((k_1-1)\vec{c} + \vec{x}_1)] * [K_{\delta-\alpha_{11}}] \right) * [\mathcal{O}(k_2\vec{c})] \\ &- [2]_{\mathbf{v}}[\mathcal{O}(k_1\vec{c})] * [S_{1,0}^{(p_1-1)}] * [\mathcal{O}(k_2\vec{c})] \\ &+ [\mathcal{O}(k_1\vec{c})] * \left( \mathbf{v}^{-1}[S_{1,0}^{(p_1-1)}] * [\mathcal{O}(k_2\vec{c})] + (\mathbf{v} - \mathbf{v}^{-1})[\mathcal{O}((k_2-1)\vec{c} + \vec{x}_1)] * [K_{\delta-\alpha_{11}}] \right) \\ &= (\mathbf{v} - \mathbf{v}^{-1}) \left( [\mathcal{O}(k_1\vec{c})] * [\mathcal{O}((k_2-1)\vec{c} + \vec{x}_1)] - \mathbf{v}[\mathcal{O}((k_1-1)\vec{c} + \vec{x}_1)] * [\mathcal{O}(k_2\vec{c})] \right) * [K_{\delta-\alpha_{11}}]. \end{aligned}$$

Then

$$\begin{aligned} & \widehat{S}(k_1, k_2 \mid -1; \star, [1, 1]) \\ &= -(\mathbf{v} - \mathbf{v}^{-1}) \left( [\mathcal{O}(k_1\vec{c})] * [\mathcal{O}((k_2-1)\vec{c} + \vec{x}_1)] - \mathbf{v}[\mathcal{O}((k_1-1)\vec{c} + \vec{x}_1)] * [\mathcal{O}(k_2\vec{c})] \right). \end{aligned}$$

For  $k_1 = k_2$ , the desired formula (8.8) holds since

$$\begin{aligned} \widehat{S}(k_1, k_1 \mid -1; \star, [1, 1]) &= \widehat{S}(k_1, k_1 \mid -1; \star, [1, 1]) \\ &= (q-1) [[\mathcal{O}((k_1-1)\vec{c} + \vec{x}_1)], [\mathcal{O}(k_1\vec{c})]]_{\mathbf{v}^{-1}} \\ &= \mathbf{v}^{-1}(q-1)^2 [S_{1,0}^{(p_1-1)}] * [K_{\mathcal{O}((k_1-1)\vec{c} + \vec{x}_1)}] \\ &= -\mathbf{v}^{-1}(q-1)^2 \widehat{B}_{[1,1], -1} * [K_{\mathcal{O}(k_1\vec{c})}] \\ &\stackrel{(7.3)}{=} (1-q)^2 \widehat{\mathbb{R}}(k_1, k_1 \mid -1; \star, [1, 1]). \end{aligned}$$

Without loss of generality, we assume  $k_1 < k_2$  in the following. Then we have

$$\begin{aligned} & \widehat{S}(k_1, k_2 \mid -1; \star, [1, 1]) \\ &= -(\mathbf{v} - \mathbf{v}^{-1})[\mathcal{O}(k_1\vec{c})] * [\mathcal{O}((k_2-1)\vec{c} + \vec{x}_1)] + (q-1)[\mathcal{O}((k_1-1)\vec{c} + \vec{x}_1)] * [\mathcal{O}(k_2\vec{c})] \\ &\quad - (\mathbf{v} - \mathbf{v}^{-1})[\mathcal{O}(k_2\vec{c})] * [\mathcal{O}((k_1-1)\vec{c} + \vec{x}_1)] + (q-1)[\mathcal{O}((k_2-1)\vec{c} + \vec{x}_1)] * [\mathcal{O}(k_1\vec{c})] \\ &= (q-1) [[\mathcal{O}((k_2-1)\vec{c} + \vec{x}_1)], [\mathcal{O}(k_1\vec{c})]]_{\mathbf{v}^{-1}} + (q-1) [[\mathcal{O}((k_1-1)\vec{c} + \vec{x}_1)], [\mathcal{O}(k_2\vec{c})]]_{\mathbf{v}^{-1}} \\ &\stackrel{(A.10)}{=} -\mathbf{v}^{-3}(q-1)^3 \widehat{\Theta}_{\star, k_2-k_1-1}^+ * [K_{\mathcal{O}(k_1\vec{c})}] + \mathbf{v}^{-1}(q-1)^3 \widehat{\Theta}_{\star, k_2-k_1+1}^- * [K_{\mathcal{O}((k_1-1)\vec{c} + \vec{x}_1)}]. \end{aligned}$$

If  $k_2 = k_1 + 1$ , then the desired formula (8.8) holds since

$$\widehat{S}(k_1, k_1 + 1 \mid -1; \star, [1, 1])$$

$$\begin{aligned}
 &= -\mathbf{v}^{-3}(q-1)^3 \widehat{\Theta}_{\star,0}^+ * [K_{\mathcal{O}(k_1\bar{c})}] + \mathbf{v}^{-1}(q-1)^3 \widehat{\Theta}_{\star,2}^- * [K_{\mathcal{O}((k_1-1)\bar{c}+\bar{x}_1)}] \\
 &= -\mathbf{v}^{-3}(q-1)^3 \cdot \frac{1}{\mathbf{v}-\mathbf{v}^{-1}} [S_{1,1}] * [K_{\mathcal{O}(k_1\bar{c})}] \\
 &\quad + \mathbf{v}^{-1}(q-1)^2 \cdot \left( \mathbf{v} [\widehat{\Theta}_{\star,1}, [S_{1,0}^{(p_1-1)}]]_{\mathbf{v}^{-2}} - q [[S_{1,1}], \widehat{\Theta}_{\star,0}]_{\mathbf{v}^{-2}} * [K_{\delta-\alpha_{11}}] \right) * [K_{\mathcal{O}((k_1-1)\bar{c}+\bar{x}_1)}] \\
 &= -\mathbf{v}^{-1} [2]_{\mathbf{v}} (q-1)^2 [S_{1,1}] * [K_{\mathcal{O}(k_1\bar{c})}] + (q-1)^2 [\widehat{\Theta}_{\star,1}, [S_{1,0}^{(p_1-1)}]]_{\mathbf{v}^{-2}} * [K_{\delta-\alpha_{11}}]^{-1} * [K_{\mathcal{O}(k_1\bar{c})}] \\
 &= (1-q)^2 \left( -[2]_{\mathbf{v}} [\widehat{\Theta}_{\star,0}, \widehat{B}_{[1,1],0}]_{\mathbf{v}^{-2}} - [\widehat{\Theta}_{\star,1}, \widehat{B}_{[1,1],-1}]_{\mathbf{v}^{-2}} \right) * [K_{\mathcal{O}(k_1\bar{c})}] \\
 &\stackrel{(A.20)}{=} (1-q)^2 \left( -[2]_{\mathbf{v}} [\widehat{\Theta}_{\star,0}, \widehat{B}_{[1,1],-2}]_{\mathbf{v}^{-2}} * [K_{\delta}] - [\widehat{B}_{[1,1],-1}, \widehat{\Theta}_{\star,1}]_{\mathbf{v}^{-2}} \right) * [K_{\mathcal{O}(k_1\bar{c})}] \\
 &\stackrel{(7.4)}{=} (1-q)^2 \widehat{\mathbb{R}}(k_1, k_1+1 \mid -1; \star, [1, 1]).
 \end{aligned}$$

If  $k_2 > k_1 + 1$ , then by (A.11) and (A.12), we have

$$\begin{aligned}
 &\frac{1}{(q-1)^2} \left( \widehat{\mathbb{S}}(k_1, k_2 \mid -1; \star, [1, 1]) - q \widehat{\mathbb{S}}(k_1, k_2 - 2 \mid -1; \star, [1, 1]) * [K_{\delta}] \right) \\
 &= -\mathbf{v}^{-3}(q-1) \left( \widehat{\Theta}_{\star, k_2-k_1-1}^+ - q \widehat{\Theta}_{\star, k_2-k_1-3}^+ * [K_{\delta}] \right) * [K_{\mathcal{O}(k_1\bar{c})}] \\
 &\quad + \mathbf{v}^{-1}(q-1) \left( \widehat{\Theta}_{\star, k_2-k_1+1}^- - q \widehat{\Theta}_{\star, k_2-k_1-1}^- * [K_{\delta}] \right) * [K_{\mathcal{O}((k_1-1)\bar{c}+\bar{x}_1)}] \\
 &= - \left( \mathbf{v}^{-1} [[S_{1,1}], \widehat{\Theta}_{\star, k_2-k_1-1}]_{\mathbf{v}^{-2}} - [\widehat{\Theta}_{\star, k_2-k_1-2}, [S_{1,0}^{(p_1-1)}]]_{\mathbf{v}^{-2}} * [K_{S_{1,1}}] \right) * [K_{\mathcal{O}(k_1\bar{c})}] \\
 &\quad + \left( [\widehat{\Theta}_{\star, k_2-k_1}, [S_{1,0}^{(p_1-1)}]]_{\mathbf{v}^{-2}} - \mathbf{v} [[S_{1,1}], \widehat{\Theta}_{\star, k_2-k_1-1}]_{\mathbf{v}^{-2}} * [K_{\delta-\alpha_{11}}] \right) * [K_{\mathcal{O}((k_1-1)\bar{c}+\bar{x}_1)}] \\
 &= -[2]_{\mathbf{v}} [[S_{1,1}], \widehat{\Theta}_{\star, k_2-k_1-1}]_{\mathbf{v}^{-2}} * [K_{\mathcal{O}(k_1\bar{c})}] + [\widehat{\Theta}_{\star, k_2-k_1-2}, [S_{1,0}^{(p_1-1)}]]_{\mathbf{v}^{-2}} * [K_{\mathcal{O}(k_1\bar{c}+\bar{x}_1)}] \\
 &\quad + [\widehat{\Theta}_{\star, k_2-k_1}, [S_{1,0}^{(p_1-1)}]]_{\mathbf{v}^{-2}} * [K_{\mathcal{O}((k_1-1)\bar{c}+\bar{x}_1)}] \\
 &= -[2]_{\mathbf{v}} [\widehat{B}_{[1,1],0}, \widehat{\Theta}_{\star, k_2-k_1-1}]_{\mathbf{v}^{-2}} * [K_{\mathcal{O}(k_1\bar{c})}] - [\widehat{\Theta}_{\star, k_2-k_1-2}, \widehat{B}_{[1,1],-1}]_{\mathbf{v}^{-2}} * [K_{\mathcal{O}((k_1+1)\bar{c})}] \\
 &\quad - [\widehat{\Theta}_{\star, k_2-k_1}, \widehat{B}_{[1,1],-1}]_{\mathbf{v}^{-2}} * [K_{\mathcal{O}(k_1\bar{c})}].
 \end{aligned}$$

Therefore,

$$\begin{aligned}
 &\frac{1}{(q-1)^2} \left( \widehat{\mathbb{S}}(k_1, k_2 \mid -1; \star, [1, 1]) - q \widehat{\mathbb{S}}(k_1, k_2 - 2 \mid -1; \star, [1, 1]) * [K_{\delta}] \right) \\
 &= \left( -[2]_{\mathbf{v}} [\widehat{B}_{[1,1],0}, \widehat{\Theta}_{\star, k_2-k_1-1}]_{\mathbf{v}^{-2}} - [\widehat{\Theta}_{\star, k_2-k_1-2}, \widehat{B}_{[1,1],-1}]_{\mathbf{v}^{-2}} * [K_{\delta}] \right. \\
 &\quad \left. - [\widehat{\Theta}_{\star, k_2-k_1}, \widehat{B}_{[1,1],-1}]_{\mathbf{v}^{-2}} \right) * [K_{\mathcal{O}(k_1\bar{c})}] \\
 &\stackrel{(A.20)}{=} \left( -[2]_{\mathbf{v}} [\widehat{\Theta}_{\star, k_2-k_1-1}, \widehat{B}_{[1,1],-2}]_{\mathbf{v}^{-2}} * [K_{\delta}] - [\widehat{B}_{[1,1],-1}, \widehat{\Theta}_{\star, k_2-k_1-2}]_{\mathbf{v}^{-2}} * [K_{\delta}] \right. \\
 &\quad \left. - [\widehat{B}_{[1,1],-1}, \widehat{\Theta}_{\star, k_2-k_1}]_{\mathbf{v}^{-2}} \right) * [K_{\mathcal{O}(k_1\bar{c})}] \\
 &\stackrel{(7.5)}{=} \widehat{\mathbb{R}}(k_1, k_2 \mid -1; \star, [1, 1]) - q \widehat{\mathbb{R}}(k_1, k_2 - 2 \mid -1; \star, [1, 1]),
 \end{aligned}$$

Then (8.8) follows by induction and the proof is completed.  $\square$

## APPENDIX B. PROOFS OF RESULTS IN SECTION 9

For a partition  $\lambda = (1^{l_1}, \dots, n^{l_n})$  and  $u > 0$ , we denote

- $\triangleright u \triangleleft \lambda - u$  is a part of  $\lambda$  (i.e.,  $l_u > 0$ ),
- $\triangleright \lambda \cup \{u\}$  – the partition  $(1^{l_1}, \dots, u^{l_u+1}, \dots, n^{l_n})$ ,
- $\triangleright \lambda \setminus \{u\}$  – the partition  $(1^{l_1}, \dots, u^{l_u-1}, \dots, n^{l_n})$  if  $u \triangleleft \lambda$ .

**B.1. Proof of Proposition 9.1.** Recall definition of  $\widehat{H}_{\star, r}$  in (6.7). Then we have the following results.

**Lemma B.1.** *For any  $r > 0$ , we have*

$$[[S_{1,1}], \widehat{H}_{\star, r}] = \frac{[r]_{\mathbf{v}}}{r} [M_{r\delta+\alpha_1}] + \frac{[r]_{\mathbf{v}}}{r} [M_{r\delta-\alpha_1}] * [K_{S_{1,1}}], \quad (\text{B.1})$$

$$[[\widehat{H}_{\star, r}, [S_{1,0}^{(p_1-1)}]]] = \frac{[r]_{\mathbf{v}}}{r} [M_{(r+1)\delta-\alpha_{11}}] + \frac{[r]_{\mathbf{v}}}{r} [M_{(r-1)\delta+\alpha_{11}}] * [K_{\delta-\alpha_{11}}]. \quad (\text{B.2})$$

*Proof.* We only prove (B.1) since (B.2) can be proved dually.

Let  $\lambda = (1^{l_1}, 2^{l_2}, \dots, n^{l_n})$  be a partition of  $r$ . Since  $\langle S_{1,0}^{(\lambda)}, S_{1,1} \rangle = 0$ , we know that  $\text{Hom}_{\mathbb{X}}(S_{1,0}^{(\lambda)}, S_{1,1}) = 0 = \text{Ext}_{\mathbb{X}}^1(S_{1,0}^{(\lambda)}, S_{1,1})$ . Hence,  $[S_{1,0}^{(\lambda)}] * [S_{1,1}] = [S_{1,0}^{(\lambda)} \oplus S_{1,1}]$ , and then

$$[S_{1,0}^{(\lambda)}] * [S_{1,1}] = q^{\ell(\lambda)} [S_{1,1} \oplus S_{1,0}^{(\lambda)}] \quad (\text{B.3})$$

by noting that  $|\text{Hom}_{\mathbb{X}}(S_{1,1}, S_{1,0}^{(\lambda)})| = q^{\ell(\lambda)}$ . Observe that the middle term of any non-split sequence in  $\text{Ext}_{\mathbb{X}}^1(S_{1,1}, S_{1,0}^{(\lambda)})$  has the form  $S_{1,1}^{(up_1+1)} \oplus S_{1,0}^{(\lambda \setminus \{u\})}$  for some  $u \triangleleft \lambda$ , and

$$F_{S_{1,1}, S_{1,0}^{(\lambda)}}^{S_{1,1}^{(up_1+1)} \oplus S_{1,0}^{(\lambda \setminus \{u\})}} = F_{S_{1,1}, S_{1,0}^{(u)}}^{S_{1,1}^{(up_1+1)}} = 1.$$

On the other hand, for any non-zero map  $f : S_{1,1} \rightarrow S_{1,0}^{(\lambda)}$ , we have  $f$  is injective and  $\text{Coker}(f) = S_{1,0}^{(up_1-1)} \oplus S_{1,0}^{(\lambda \setminus \{u\})}$  for some  $u \triangleleft \lambda$ . Obviously,

$$\text{Ext}_{\mathbb{X}}^1(S_{1,0}^{(up_1-1)} \oplus S_{1,0}^{(\lambda \setminus \{u\})}, S_{1,1}) \cong \text{Ext}_{\mathbb{X}}^1(S_{1,0}^{(up_1-1)}, S_{1,1}) \cong \mathbf{k}.$$

Then by Corollary 3.10 (1), we get

$$[S_{1,1}] * [S_{1,0}^{(\lambda)}] = [S_{1,1} \oplus S_{1,0}^{(\lambda)}] + \sum_{u \triangleleft \lambda} [[S_{1,1}^{(up_1+1)} \oplus S_{1,0}^{(\lambda \setminus \{u\})}]] + \sum_{u \triangleleft \lambda} [[S_{1,0}^{(up_1-1)} \oplus S_{1,0}^{(\lambda \setminus \{u\})}] * [K_{\alpha_{11}}]]. \quad (\text{B.4})$$

Observe that  $[[S_{1,1}], [M]] = 0$  for any indecomposable torsion sheaf  $M \notin \mathcal{T}_{\lambda_1}$ . Therefore, it follows from (B.3)–(B.4) that

$$\begin{aligned} \frac{r}{[r]} [[S_{1,1}], \widehat{H}_{\star, r}] &= \sum_{|\lambda|=r} \mathbf{n}(\ell(\lambda) - 1) [[S_{1,1}], [S_{1,0}^{(\lambda)}]] \\ &= \sum_{|\lambda|=r} \mathbf{n}(\ell(\lambda) - 1) \cdot (1 - q^{\ell(\lambda)}) [S_{1,1} \oplus S_{1,0}^{(\lambda)}] + \sum_{|\lambda|=r} \mathbf{n}(\ell(\lambda) - 1) \sum_{u \triangleleft \lambda} [[S_{1,1}^{(up_1+1)} \oplus S_{1,0}^{(\lambda \setminus \{u\})}]] \\ &\quad + \sum_{|\lambda|=r} \mathbf{n}(\ell(\lambda) - 1) \sum_{u \triangleleft \lambda} [[S_{1,0}^{(up_1-1)} \oplus S_{1,0}^{(\lambda \setminus \{u\})}] * [K_{\alpha_{11}}]] \\ &= [M_{r\delta+\alpha_1}] + [M_{r\delta-\alpha_1}] * [K_{\alpha_{11}}], \end{aligned}$$



by using  $\mathbf{n}(\ell(\lambda) - 1) \cdot (1 - q^{\ell(\lambda)}) = \mathbf{n}(\ell(\lambda))$ . The proof is completed.  $\square$

With the help of Lemma B.1, we can prove Proposition 9.1 now.

*Proof of Proposition 9.1.* We prove the formulas by induction on  $r$ . For  $r = 1$ , they follow from (5.18) and Proposition 5.1.

We assume  $\widehat{B}_{[1,1],l} = (q-1)[M_{l\delta+\alpha_{11}}]$  and  $\widehat{B}_{[1,1],-l} = (1-q)[M_{l\delta-\alpha_{11}}] * [K_{-l\delta+\alpha_{11}}]$  for any  $1 \leq l \leq r$ . Using (8.3) proved in Proposition 8.5, we have

$$[\widehat{H}_{\star,m}, \widehat{B}_{[1,1],l}] = -\frac{[m]_{\mathbf{v}}}{m} \widehat{B}_{[1,1],l+m} + \frac{[m]_{\mathbf{v}}}{m} \widehat{B}_{[1,1],l-m} * [K_{m\delta}]$$

for any  $l \in \mathbb{Z}$ ,  $m > 0$ . Recall that  $\widehat{B}_{[1,1],0} = [S_{1,1}]$ ,  $\widehat{B}_{[1,1],-1} = -[S_{1,0}^{(p_1-1)}] * [K_{-\delta+\alpha_{11}}]$ . Using (B.2), one can see that  $\widehat{B}_{[1,1],-(r+1)} = (1-q)[M_{(r+1)\delta-\alpha_{11}}] * [K_{-(r+1)\delta+\alpha_{11}}]$ . Together with (B.1), we have  $\widehat{B}_{[1,1],r+1} = (q-1)[M_{(r+1)\delta+\alpha_{11}}]$ . The proof is completed.  $\square$

**B.2. Proof of Proposition 9.2.** First, let us give some formulas on the cardinalities of Hom-spaces and automorphism groups.

**Lemma B.2.** *Let  $\lambda = (1^{l_1}, 2^{l_2}, \dots, n^{l_n})$  be a partition and  $u > 0$ . Then*

$$|\mathrm{Aut}(S_{1,0}^{(up_1-1)})| = q^{u-1}(q-1), \quad |\mathrm{Aut}(S_{1,1}^{(up_1+1)})| = q^u(q-1), \quad (\text{B.5})$$

$$|\mathrm{Aut}(S_{1,0}^{(\lambda)})| = q^{|\lambda| + \sum_i il_i(l_i-1) + 2 \sum_{i < j} il_i l_j} \prod_i (1 - q^{-1})(1 - q^{-2}) \cdots (1 - q^{-l_i}), \quad (\text{B.6})$$

$$|\mathrm{Hom}(S_{1,1}^{(up_1+1)}, S_{1,0}^{(\lambda)})| = |\mathrm{Hom}(S_{1,1}^{((u+1)p_1)}, S_{1,0}^{(\lambda)})| = q^{\sum_i \min\{i, u+1\}l_i}, \quad (\text{B.7})$$

$$|\mathrm{Hom}(S_{1,0}^{(\lambda)}, S_{1,1}^{(up_1+1)})| = |\mathrm{Hom}(S_{1,0}^{(\lambda)}, S_{1,0}^{(up_1)})| = q^{\sum_i \min\{i, u\}l_i}, \quad (\text{B.8})$$

$$|\mathrm{Hom}(S_{1,0}^{(up_1-1)}, S_{1,0}^{(\lambda)})| = |\mathrm{Hom}(S_{1,0}^{((u-1)p_1)}, S_{1,0}^{(\lambda)})| = q^{\sum_i \min\{i, u-1\}l_i}, \quad (\text{B.9})$$

$$|\mathrm{Hom}(S_{1,0}^{(\lambda)}, S_{1,0}^{(up_1-1)})| = |\mathrm{Hom}(S_{1,0}^{(\lambda)}, S_{1,0}^{(up_1)})| = q^{\sum_i \min\{i, u\}l_i}, \quad (\text{B.10})$$

$$|\mathrm{Aut}(S_{1,0}^{(\lambda \setminus \{u\})})| = \frac{|\mathrm{Aut}(S_{1,0}^{(\lambda)})| \cdot q^u}{(1 - q^{-lu}) \cdot q^{2 \sum_i \min\{i, u\}l_i}} \quad (\text{B.11})$$

$$|\mathrm{Aut}(S_{1,0}^{(\lambda \cup \{u\})})| = |\mathrm{Aut}(S_{1,0}^{(\lambda)})| \cdot q^u \cdot (1 - q^{-lu-1}) \cdot q^{2 \sum_i \min\{i, u\}l_i}. \quad (\text{B.12})$$

*Proof.* See [Sch12, Lemma 2.8] for (B.6), and the others follow by direct computations.  $\square$

For any  $a, b \geq 1$ , we denote by

$$\phi_{a,b} = \frac{1}{q-1} \cdot |\mathrm{Ext}^1(S_{1,0}^{(ap_1-1)} \oplus S_{1,0}^{(bp_1-1)}, S_{1,1})_{S_{1,0}^{(ap_1-1)} \oplus S_{1,0}^{(bp_1-1)}}|,$$

and

$$\psi_{a,b} = F_{S_{1,0}^{(ap_1-1)} \oplus S_{1,0}^{(bp_1-1)}, S_{1,0}^{((b-1)p_1)}, S_{1,0}^{(p_1-1)}}.$$

A direct calculation shows that

$$\phi_{a,b} = \psi_{b,a} = \begin{cases} q, & \text{if } a < b, \\ 1, & \text{if } a > b, \\ q+1, & \text{if } a = b. \end{cases} \quad (\text{B.13})$$

**Lemma B.3.** *For any  $r \geq 1$ , we have*

$$\begin{aligned} & - \frac{1}{(q-1)^2} [\widehat{B}_{[1,1],0}, \widehat{B}_{[1,1],-r}]_{\mathbf{v}^2} * [K_{r\delta-\alpha_{11}}] \\ & = \mathbf{v}^{-1} \sum_{[M] \in \mathcal{M}_{r\delta}} \mathbf{n}(\ell(M) - 1) \llbracket M \rrbracket + \frac{\mathbf{v}}{q-1} \sum_{|\lambda|=r} \mathbf{n}(\ell(\lambda)) \llbracket S_{1,0}^{(\lambda)} \rrbracket \\ & \quad + \mathbf{v} \sum_{a+|\nu|=r} \mathbf{n}(\ell(\nu)) \sum_{u \triangleleft \nu} \phi_{a,u} \llbracket S_{1,0}^{(ap_1-1)} \oplus S_{1,0}^{(up_1-1)} \oplus S_{1,0}^{(\nu \setminus \{u\})} \rrbracket * [K_{\alpha_{11}}]. \end{aligned}$$

*Proof.* Let  $[M] = [S_{1,0}^{(ap_1-1)} \oplus S_{1,0}^{(\nu)}] \in \mathcal{M}_{r\delta-\alpha_{11}}$ . Then we have  $|\nu| + a = r$  and

$$\langle \widehat{M}, \alpha_{11} \rangle = \langle \alpha_{11}, \widehat{M} \rangle = -\langle \alpha_{11}, \alpha_{11} \rangle = -1.$$

Since  $\text{Hom}(M, S_{1,1}) = 0$  and  $\text{Ext}^1(M, S_{1,1}) \cong \mathbf{k}$ , we get

$$[M] * [S_{1,1}] = \mathbf{v}^{-1} [M \oplus S_{1,1}] + \mathbf{v}^{-1} (q-1) [S_{1,0}^{(\nu \cup \{a\})}].$$

Let  $\nu = (1^{l_1}, 2^{l_2}, \dots, n^{l_n})$ . Then we obtain

$$[M] * [S_{1,1}] = \mathbf{v}^{-1} q^{\ell(\nu)} \llbracket M \oplus S_{1,1} \rrbracket + \mathbf{v}^{-1} \frac{q^{a+1} - 1}{q-1} \cdot q^{\sum_{i>a} l_i} \llbracket S_{1,0}^{(\nu \cup \{a\})} \rrbracket. \quad (\text{B.14})$$

Here we use  $|\text{Hom}(S_{1,1}, M)| = q^{\ell(\nu)}$  and

$$\frac{|\text{Aut}(S_{1,0}^{(\nu \cup \{a\})})|}{|\text{Aut}(M)|} = \frac{q^{a+1} - 1}{q-1} \cdot q^{\sum_{i>a} l_i},$$

which follows from Lemma B.2 by direct computations.

Observe that the middle term of any non-split extension in  $\text{Ext}^1(S_{1,1}, M)$  has the form  $S_{1,1}^{(ap_1)} \oplus S_{1,0}^{(\nu)}$  or  $S_{1,0}^{(ap_1-1)} \oplus S_{1,1}^{(up_1+1)} \oplus S_{1,0}^{(\nu \setminus \{u\})}$  for some  $u \triangleleft \nu$ , and

$$F_{S_{1,1}, M}^{S_{1,1}^{(ap_1)} \oplus S_{1,0}^{(\nu)}} = F_{S_{1,1}, S_{1,0}^{(ap_1-1)}}^{S_{1,1}^{(ap_1)}} = 1, \quad F_{S_{1,1}, M}^{S_{1,0}^{(ap_1-1)} \oplus S_{1,1}^{(up_1+1)} \oplus S_{1,0}^{(\nu \setminus \{u\})}} = F_{S_{1,1}, S_{1,0}^{(up_1+1)}}^{S_{1,1}^{(up_1+1)}} = 1.$$

Moreover, for the split extension we also have  $F_{S_{1,1}, M}^{S_{1,1} \oplus M} = 1$ . On the other hand, for any non-zero map  $f : S_{1,1} \rightarrow M$ , we have that  $f$  is injective and  $\text{Coker}(f) = S_{1,0}^{(ap_1-1)} \oplus S_{1,0}^{(up_1-1)} \oplus S_{1,0}^{(\nu \setminus \{u\})}$  for some  $u \triangleleft \nu$ , hence  $\text{Hom}(\text{Coker}(f), S_{1,1}) = 0$ . Then by Corollary 3.10 (1), we get

$$\begin{aligned} & \llbracket S_{1,1} \rrbracket * \llbracket M \rrbracket \\ & = \mathbf{v}^{-1} \left( \llbracket S_{1,1} \oplus M \rrbracket + \llbracket S_{1,1}^{(ap_1)} \oplus S_{1,0}^{(\nu)} \rrbracket + \sum_{u \triangleleft \nu} \llbracket S_{1,0}^{(ap_1-1)} \oplus S_{1,1}^{(up_1+1)} \oplus S_{1,0}^{(\nu \setminus \{u\})} \rrbracket \right) \\ & \quad + \mathbf{v} \sum_{u \triangleleft \nu} \phi_{a,u} \llbracket S_{1,0}^{(ap_1-1)} \oplus S_{1,0}^{(up_1-1)} \oplus S_{1,0}^{(\nu \setminus \{u\})} \rrbracket * [K_{\alpha_{11}}]. \end{aligned} \quad (\text{B.15})$$

Combining (B.14)–(B.15), we have

$$\begin{aligned} & \llbracket [S_{1,1}], [M] \rrbracket_{\mathbf{v}^2} \\ & = \mathbf{v}^{-1} \left( (1 - q^{\ell(\nu)+1}) \llbracket S_{1,1} \oplus M \rrbracket + \llbracket S_{1,1}^{(ap_1)} \oplus S_{1,0}^{(\nu)} \rrbracket + \sum_{u \triangleleft \nu} \llbracket S_{1,0}^{(ap_1-1)} \oplus S_{1,1}^{(up_1+1)} \oplus S_{1,0}^{(\nu \setminus \{u\})} \rrbracket \right) \end{aligned}$$

$$- \mathbf{v} \frac{q^{l_a+1} - 1}{q - 1} \cdot q^{\sum_{i>a} l_i} \llbracket S_{1,0}^{\nu \cup \{a\}} \rrbracket + \mathbf{v} \sum_{u \triangleleft \nu} \phi_{a,u} \llbracket S_{1,0}^{(ap_1-1)} \oplus S_{1,0}^{(up_1-1)} \oplus S_{1,0}^{\nu \setminus \{u\}} \rrbracket * [K_{\alpha_{11}}].$$

Therefore, by definition,

$$\begin{aligned} & - \frac{1}{(q-1)^2} [\widehat{B}_{[1,1],0}, \widehat{B}_{[1,1],-r}]_{\mathbf{v}^2} * [K_{r\delta - \alpha_{11}}] \\ &= \sum_{[M] \in \mathcal{M}_{r\delta - \alpha_{11}}} \mathbf{n}(\ell(M) - 1) \llbracket [S_{1,1}], [M] \rrbracket_{\mathbf{v}^2} \\ &= \sum_{a+|\nu|=r} \mathbf{n}(\ell(\nu)) \llbracket [S_{1,1}], [S_{1,0}^{(ap_1-1)} \oplus S_{1,0}^{(\nu)}] \rrbracket_{\mathbf{v}^2} \\ &= \mathbf{v}^{-1} \sum_{[M] \in \mathcal{M}_{r\delta}} \mathbf{n}(\ell(M) - 1) \llbracket [M] \rrbracket - \frac{\mathbf{v}}{q-1} \sum_{a+|\nu|=r} \mathbf{n}(\ell(\nu)) (q^{l_a+1} - 1) \cdot q^{\sum_{i>a} l_i} \llbracket S_{1,0}^{\nu \cup \{a\}} \rrbracket \\ & \quad + \mathbf{v} \sum_{a+|\nu|=r} \mathbf{n}(\ell(\nu)) \sum_{u \triangleleft \nu} \phi_{a,u} \llbracket S_{1,0}^{(ap_1-1)} \oplus S_{1,0}^{(up_1-1)} \oplus S_{1,0}^{\nu \setminus \{u\}} \rrbracket * [K_{\alpha_{11}}]. \end{aligned} \tag{B.16}$$

Let  $\lambda = \nu \cup \{a\} = (1^{l'_1}, 2^{l'_2}, \dots, n^{l'_n})$ . Then  $l'_i = l_i$  for  $i \neq a$  and  $l'_a = l_a + 1$ . Hence

$$\begin{aligned} & \sum_{a+|\nu|=r} \mathbf{n}(\ell(\nu)) (q^{l_a+1} - 1) \cdot q^{\sum_{i>a} l_i} \llbracket S_{1,0}^{\nu \cup \{a\}} \rrbracket \\ &= \sum_{|\lambda|=r} \mathbf{n}(\ell(\lambda) - 1) \sum_{a \triangleleft \lambda} (q^{l'_a} - 1) \cdot q^{\sum_{i>a} l'_i} \llbracket S_{1,0}^{(\lambda)} \rrbracket \\ &= \sum_{|\lambda|=r} \mathbf{n}(\ell(\lambda) - 1) \sum_{a \triangleleft \lambda} \left( q^{\sum_{i \geq a} l'_i} - q^{\sum_{i>a} l'_i} \right) \llbracket S_{1,0}^{(\lambda)} \rrbracket \\ &= \sum_{|\lambda|=r} \mathbf{n}(\ell(\lambda) - 1) (q^{\ell(\lambda)} - 1) \llbracket S_{1,0}^{(\lambda)} \rrbracket = - \sum_{|\lambda|=r} \mathbf{n}(\ell(\lambda)) \llbracket S_{1,0}^{(\lambda)} \rrbracket \end{aligned} \tag{B.17}$$

by using  $\mathbf{n}(\ell(\lambda) - 1)(1 - q^{\ell(\lambda)}) = \mathbf{n}(\ell(\lambda))$ .

Then the desired formula follows by plugging (B.17) into (B.16).  $\square$

**Lemma B.4.** *For any  $r \geq 3$ , we have*

$$\begin{aligned} & \frac{1}{(q-1)^2} [\widehat{B}_{[1,1],-r+1}, \widehat{B}_{[1,1],-1}]_{\mathbf{v}^2} * [K_{r\delta - 2\alpha_{11}}] \\ &= \mathbf{v} \sum_{a+|\nu|=r} \mathbf{n}(\ell(\nu)) \sum_{u \triangleleft \nu} \phi_{a,u} \llbracket S_{1,0}^{(ap_1-1)} \oplus S_{1,0}^{(up_1-1)} \oplus S_{1,0}^{\nu \setminus \{u\}} \rrbracket \\ & \quad + \mathbf{v}^{-3} \sum_{[M] \in \mathcal{M}_{(r-2)\delta}} \mathbf{n}(\ell(M) - 1) \llbracket [M] \rrbracket * [K_{\delta - \alpha_{11}}] + \frac{\mathbf{v}}{q(q-1)} \sum_{|\lambda|=r-2} \mathbf{n}(\ell(\lambda)) \llbracket S_{1,0}^{(\lambda)} \rrbracket * [K_{\delta - \alpha_{11}}]. \end{aligned} \tag{B.18}$$

*Proof.* Let  $[M] := [S_{1,0}^{(ap_1-1)} \oplus S_{1,0}^{(\mu)}] \in \mathcal{M}_{(r-1)\delta - \alpha_{11}}$ . Then we have  $|\mu| + a = r - 1$  and

$$\langle \delta - \alpha_{11}, \widehat{M} \rangle = \langle \widehat{M}, \delta - \alpha_{11} \rangle = \langle -\alpha_{11}, -\alpha_{11} \rangle = 1.$$

Observe that  $\text{Ext}^1(S_{1,0}^{(p_1-1)}, M) = 0$  and  $\text{Hom}(S_{1,0}^{(p_1-1)}, M) \cong \mathbf{k}$ . For any non-zero map  $f : S_{1,0}^{(p_1-1)} \rightarrow M$ , we have  $f$  is injective and  $\text{Coker}(f) = S_{1,0}^{(\mu \cup \{a-1\})}$ . Then

$$F_{S_{1,0}^{(\mu \cup \{a-1\}), S_{1,0}^{(p_1-1)}}}^M = 1, \quad F_{S_{1,0}^{(p_1-1)}, M}^{S_{1,0}^{(\mu \cup \{a-1\})} \oplus M} = \begin{cases} q^{\ell(\mu)+1}, & \text{if } a \geq 2, \\ (q+1)q^{\ell(\mu)}, & \text{if } a = 1. \end{cases}$$

Let  $\mu = (1^{l_1}, 2^{l_2}, \dots, n^{l_n})$ . By Lemma B.2, a direct computation shows that

$$\frac{|\text{Aut}(S_{1,0}^{(\mu \cup \{a-1\})})|}{|\text{Aut}(M)|} = \begin{cases} \frac{q^{l_{a-1}+1}-1}{q(q-1)} \cdot q^{-\sum_{i \geq a-1} l_i}, & \text{if } a \geq 2, \\ \frac{1}{q-1} \cdot q^{-\ell(\mu)}, & \text{if } a = 1. \end{cases}$$

Then by Lemma 3.10, we have for  $a \geq 2$ ,

$$\llbracket S_{1,0}^{(p_1-1)} \rrbracket * \llbracket M \rrbracket = \mathbf{v} \cdot q^{\ell(\mu)+1} \llbracket S_{1,0}^{(p_1-1)} \oplus M \rrbracket + \mathbf{v}^{-1} \frac{q^{l_{a-1}+1}-1}{q(q-1)} \cdot q^{-\sum_{i \geq a-1} l_i} \llbracket S_{1,0}^{(\mu \cup \{a-1\})} \rrbracket * \llbracket K_{\delta-\alpha_{11}} \rrbracket, \quad (\text{B.19})$$

while for  $a = 1$ ,

$$\llbracket S_{1,0}^{(p_1-1)} \rrbracket * \llbracket M \rrbracket = \mathbf{v}(q+1) \cdot q^{\ell(\mu)} \llbracket S_{1,0}^{(p_1-1)} \oplus M \rrbracket + \mathbf{v}^{-1} \frac{1}{q-1} \cdot q^{-\ell(\mu)} \llbracket S_{1,0}^{(\mu)} \rrbracket * \llbracket K_{\delta-\alpha_{11}} \rrbracket. \quad (\text{B.20})$$

On the other hand, the middle term of any non-split extension in  $\text{Ext}^1(M, S_{1,0}^{(p_1-1)})$  has the form  $S_{1,0}^{(ap_1-1)} \oplus S_{1,0}^{((u+1)p_1-1)} \oplus S_{1,0}^{(\mu \setminus \{u\})}$  for some  $u \triangleleft \mu$ , and

$$F_{M, S_{1,0}^{(p_1-1)}}^{S_{1,0}^{(ap_1-1)} \oplus S_{1,0}^{((u+1)p_1-1)} \oplus S_{1,0}^{(\mu \setminus \{u\})}} = F_{S_{1,0}^{(ap_1-1)} \oplus S_{1,0}^{((u+1)p_1-1)}, S_{1,0}^{(ap_1-1)} \oplus S_{1,0}^{(u p_1)}, S_{1,0}^{(p_1-1)}} = \psi_{a,u+1};$$

while for the split extension we have

$$F_{M, S_{1,0}^{(p_1-1)}}^{S_{1,0}^{(p_1-1)} \oplus M} = \psi_{a,1} = \begin{cases} q, & \text{if } a \neq 1, \\ q+1, & \text{if } a = 1. \end{cases}$$

For any non-zero map  $f : M \rightarrow S_{1,0}^{(p_1-1)}$ , we have  $f$  is surjective and  $\text{Ker}(f) = S_{1,1}^{((a-1)p_1)} \oplus S_{1,0}^{(\mu)}$  or  $S_{1,0}^{(ap_1-1)} \oplus S_{1,1}^{((u-1)p_1+1)} \oplus S_{1,0}^{(\mu \setminus \{u\})}$  for some  $u \triangleleft \mu$ . It is easy to see that

$$\dim_{\mathbf{k}} \text{Ext}^1(S_{1,0}^{(p_1-1)}, \text{Ker}(f)) = \dim_{\mathbf{k}} \text{Hom}(S_{1,0}^{(p_1-1)}, \text{Ker}(f)) \leq 1,$$

and equals to 0 if and only if  $\text{Ker}(f) = S_{1,0}^{(\mu)}$ . Then by Corollary 3.10 (2), we get for  $a \geq 2$ ,

$$\begin{aligned} & \llbracket M \rrbracket * \llbracket S_{1,0}^{(p_1-1)} \rrbracket \quad (\text{B.21}) \\ &= \mathbf{v} \cdot \psi_{a,1} \llbracket S_{1,0}^{(p_1-1)} \oplus M \rrbracket + \mathbf{v} \sum_{u \triangleleft \mu} \psi_{a,u+1} \llbracket S_{1,0}^{(ap_1-1)} \oplus S_{1,0}^{((u+1)p_1-1)} \oplus S_{1,0}^{(\mu \setminus \{u\})} \rrbracket \\ & \quad + \mathbf{v}^{-3} \left( \llbracket S_{1,1}^{((a-1)p_1)} \oplus S_{1,0}^{(\mu)} \rrbracket + \sum_{u \triangleleft \mu} \llbracket S_{1,0}^{(ap_1-1)} \oplus S_{1,1}^{((u-1)p_1+1)} \oplus S_{1,0}^{(\mu \setminus \{u\})} \rrbracket \right) * \llbracket K_{\delta-\alpha_{11}} \rrbracket; \end{aligned}$$

while for  $a = 1$ ,

$$\llbracket M \rrbracket * \llbracket S_{1,0}^{(p_1-1)} \rrbracket \quad (\text{B.22})$$

$$\begin{aligned}
 &= \mathbf{v} \cdot \psi_{1,1} \llbracket S_{1,0}^{(p_1-1)} \oplus M \rrbracket + \mathbf{v} \sum_{u \triangleleft \mu} \psi_{1,u+1} \llbracket S_{1,0}^{(p_1-1)} \oplus S_{1,0}^{((u+1)p_1-1)} \oplus S_{1,0}^{(\mu \setminus \{u\})} \rrbracket \\
 &\quad + \mathbf{v}^{-1} \frac{1}{q-1} \llbracket S_{1,0}^{(\mu)} \rrbracket * [K_{\delta-\alpha_{11}}] + \mathbf{v}^{-3} \sum_{u \triangleleft \mu} \llbracket S_{1,0}^{(p_1-1)} \oplus S_{1,1}^{((u-1)p_1+1)} \oplus S_{1,0}^{(\mu \setminus \{u\})} \rrbracket * [K_{\delta-\alpha_{11}}].
 \end{aligned}$$

Therefore, by combining with (B.19) and (B.21), for  $a \geq 2$ ,

$$\begin{aligned}
 &\llbracket [M], [S_{1,0}^{(p_1-1)}] \rrbracket_{\mathbf{v}^2} \tag{B.23} \\
 &= \mathbf{v} \cdot \psi_{a,1} (1 - q^{\ell(\mu)+1}) \llbracket S_{1,0}^{(p_1-1)} \oplus M \rrbracket + \mathbf{v} \sum_{u \triangleleft \mu} \psi_{a,u+1} \llbracket S_{1,0}^{(ap_1-1)} \oplus S_{1,0}^{((u+1)p_1-1)} \oplus S_{1,0}^{(\mu \setminus \{u\})} \rrbracket \\
 &\quad + \mathbf{v}^{-3} \left( \llbracket S_{1,1}^{((a-1)p_1)} \oplus S_{1,0}^{(\mu)} \rrbracket + \sum_{u \triangleleft \mu} \llbracket S_{1,0}^{(ap_1-1)} \oplus S_{1,1}^{((u-1)p_1+1)} \oplus S_{1,0}^{(\mu \setminus \{u\})} \rrbracket \right) * [K_{\delta-\alpha_{11}}] \\
 &\quad - \frac{\mathbf{v}}{q(q-1)} \cdot (q^{l_{a-1}+1} - 1) q^{-\sum_{i \geq a-1} l_i} \llbracket S_{1,0}^{(\mu \cup \{a-1\})} \rrbracket * [K_{\delta-\alpha_{11}}];
 \end{aligned}$$

while by combining with (B.20) and (B.22), for  $a = 1$ ,

$$\begin{aligned}
 &\llbracket [M], [S_{1,0}^{(p_1-1)}] \rrbracket_{\mathbf{v}^2} \tag{B.24} \\
 &= \mathbf{v} \cdot \psi_{1,1} (1 - q^{\ell(\mu)+1}) \llbracket S_{1,0}^{(p_1-1)} \oplus M \rrbracket + \mathbf{v} \sum_{u \triangleleft \mu} \psi_{1,u+1} \llbracket S_{1,0}^{(p_1-1)} \oplus S_{1,0}^{((u+1)p_1-1)} \oplus S_{1,0}^{(\mu \setminus \{u\})} \rrbracket \\
 &\quad + \mathbf{v}^{-3} \sum_{u \triangleleft \mu} \llbracket S_{1,0}^{(p_1-1)} \oplus S_{1,1}^{((u-1)p_1+1)} \oplus S_{1,0}^{(\mu \setminus \{u\})} \rrbracket * [K_{\delta-\alpha_{11}}] \\
 &\quad + \frac{\mathbf{v}}{q(q-1)} \cdot (1 - q^{-\ell(\mu)+1}) \llbracket S_{1,0}^{(\mu)} \rrbracket * [K_{\delta-\alpha_{11}}].
 \end{aligned}$$

To sum up with (B.23) and (B.24), we have

$$\begin{aligned}
 &\frac{1}{(q-1)^2} [\widehat{B}_{[1,1],-r+1}, \widehat{B}_{[1,1],-1}]_{\mathbf{v}^2} * [K_{r\delta-2\alpha_{11}}] \\
 &= \sum_{[M] \in \mathcal{M}_{(r-1)\delta-\alpha_{11}}} \mathbf{n}(\ell(M) - 1) \llbracket [M], [S_{1,0}^{(p_1-1)}] \rrbracket_{\mathbf{v}^2} \\
 &= \sum_{a+|\mu|=r-1} \mathbf{n}(\ell(\mu)) \llbracket [S_{1,0}^{(ap_1-1)} \oplus S_{1,0}^{(\mu)}], [S_{1,0}^{(p_1-1)}] \rrbracket_{\mathbf{v}^2} \\
 &= \mathbf{v} \sum_{a+|\mu|=r-1} \mathbf{n}(\ell(\mu) + 1) \psi_{a,1} \llbracket S_{1,0}^{(ap_1-1)} \oplus S_{1,0}^{(p_1-1)} \oplus S_{1,0}^{(\mu)} \rrbracket \tag{B.25}
 \end{aligned}$$

$$+ \mathbf{v} \sum_{a+|\mu|=r-1} \mathbf{n}(\ell(\mu)) \sum_{u \triangleleft \mu} \psi_{a,u+1} \llbracket S_{1,0}^{(ap_1-1)} \oplus S_{1,0}^{((u+1)p_1-1)} \oplus S_{1,0}^{(\mu \setminus \{u\})} \rrbracket \tag{B.26}$$

$$+ \mathbf{v}^{-3} \sum_{[M] \in \mathcal{M}_{(r-2)\delta}} \mathbf{n}(\ell(M) - 1) \llbracket [M] \rrbracket * [K_{\delta-\alpha_{11}}] \tag{B.27}$$

$$+ \sum_{|\mu|=r-2} \mathbf{n}(\ell(\mu)) \cdot \frac{\mathbf{v}}{q(q-1)} \cdot (1 - q^{-\ell(\mu)+1}) \llbracket S_{1,0}^{(\mu)} \rrbracket * [K_{\delta-\alpha_{11}}] \tag{B.28}$$

$$+ \sum_{a \geq 2} \sum_{a+|\mu|=r-1} \mathbf{n}(\ell(\mu)) \cdot \left( -\frac{\mathbf{v}}{q(q-1)} \cdot (q^{l_{a-1}+1} - 1) \cdot q^{-\sum_{i \geq a-1} l_i} \llbracket S_{1,0}^{(\mu \cup \{a-1\})} \rrbracket \right) * [K_{\delta-\alpha_{11}}] \quad (\text{B.29})$$

= RHS of (B.18),

where the last equality follows from the following Lemmas B.5 and B.6.  $\square$

**Lemma B.5.** *Keep the notations as above. For any  $r \geq 2$ , we have*

$$(\text{B.25}) + (\text{B.26}) = \mathbf{v} \sum_{a+|\nu|=r} \mathbf{n}(\ell(\nu)) \sum_{u \triangleleft \nu} \phi_{a,u} \llbracket S_{1,0}^{(ap_1-1)} \oplus S_{1,0}^{(up_1-1)} \oplus S_{1,0}^{(\nu \setminus \{u\})} \rrbracket. \quad (\text{B.30})$$

*Proof.* The right-hand side of (B.30) can be divided into two parts according to  $a = 1$  or not. For  $a = 1$ , we have

$$\begin{aligned} & \sum_{|\nu|=r-1} \mathbf{n}(\ell(\nu)) \sum_{u \triangleleft \nu} \phi_{1,u} \llbracket S_{1,0}^{(p_1-1)} \oplus S_{1,0}^{(up_1-1)} \oplus S_{1,0}^{(\nu \setminus \{u\})} \rrbracket \\ & \stackrel{\mu = \nu \setminus \{u\}}{=} \sum_{u+|\mu|=r-1} \mathbf{n}(\ell(\mu) + 1) \psi_{u,1} \llbracket S_{1,0}^{(up_1-1)} \oplus S_{1,0}^{(p_1-1)} \oplus S_{1,0}^{(\mu)} \rrbracket \\ & \stackrel{u \rightarrow a}{=} \sum_{a+|\mu|=r-1} \mathbf{n}(\ell(\mu) + 1) \psi_{a,1} \llbracket S_{1,0}^{(ap_1-1)} \oplus S_{1,0}^{(p_1-1)} \oplus S_{1,0}^{(\mu)} \rrbracket, \end{aligned}$$

where  $u \rightarrow a$  means substituting  $u$  by  $a$ ; while for  $a \geq 2$ ,

$$\begin{aligned} & \sum_{a \geq 2, \nu; a+|\nu|=r} \mathbf{n}(\ell(\nu)) \sum_{u \triangleleft \nu} \phi_{a,u} \llbracket S_{1,0}^{(ap_1-1)} \oplus S_{1,0}^{(up_1-1)} \oplus S_{1,0}^{(\nu \setminus \{u\})} \rrbracket \\ & \stackrel{\nu \setminus \{u\} = \nu'}{=} \sum_{a \geq 2, u} \sum_{|\nu'|=r-u-a} \mathbf{n}(\ell(\nu') + 1) \phi_{a,u} \llbracket S_{1,0}^{(ap_1-1)} \oplus S_{1,0}^{(up_1-1)} \oplus S_{1,0}^{(\nu')} \rrbracket \\ & \stackrel{a=b+1}{=} \sum_{b,u} \sum_{|\nu'|=r-u-b-1} \mathbf{n}(\ell(\nu') + 1) \phi_{b+1,u} \llbracket S_{1,0}^{((b+1)p_1-1)} \oplus S_{1,0}^{(up_1-1)} \oplus S_{1,0}^{(\nu')} \rrbracket \\ & \stackrel{\mu = \nu' \cup \{b\}}{=} \sum_{b,u} \sum_{|\mu|=r-u-1} \mathbf{n}(\ell(\mu)) \psi_{u,b+1} \llbracket S_{1,0}^{(up_1-1)} \oplus S_{1,0}^{((b+1)p_1-1)} \oplus S_{1,0}^{(\mu \setminus \{b\})} \rrbracket \\ & \stackrel{u \rightarrow a; b \rightarrow u}{=} \sum_{a+|\mu|=r-1} \mathbf{n}(\ell(\mu)) \sum_{u \triangleleft \mu} \psi_{a,u+1} \llbracket S_{1,0}^{(ap_1-1)} \oplus S_{1,0}^{((u+1)p_1-1)} \oplus S_{1,0}^{(\mu \setminus \{u\})} \rrbracket. \end{aligned}$$

Then the result follows.  $\square$

**Lemma B.6.** *Keep the notations as above. For any  $r \geq 2$ , we have*

$$(\text{B.28}) + (\text{B.29}) = \frac{\mathbf{v}}{q(q-1)} \sum_{|\lambda|=r-2} \mathbf{n}(\ell(\lambda)) \llbracket S_{1,0}^{(\lambda)} \rrbracket * [K_{\delta-\alpha_{11}}]. \quad (\text{B.31})$$

*Proof.* Recall  $\mu = (1^{l_1}, 2^{l_2}, \dots, n^{l_n})$ . For  $a \geq 2$ , let  $\lambda = \mu \cup \{a-1\} = (1^{l_1}, 2^{l_2}, \dots, n^{l_n})$ , then  $l'_i = l_i$  for  $i \neq a-1$  and  $l'_{a-1} = l_{a-1} + 1$ . Hence

$$\sum_{a \geq 2} \sum_{|\mu|=r-a-1} \mathbf{n}(\ell(\mu)) (q^{l_{a-1}+1} - 1) \cdot q^{-\sum_{i \geq a-1} l_i} \llbracket S_{1,0}^{(\mu \cup \{a-1\})} \rrbracket$$

$$\begin{aligned}
 &= \sum_{|\lambda|=r-2} \mathbf{n}(\ell(\lambda) - 1) \sum_{a-1 \triangleleft \lambda} (q^{l'_{a-1}} - 1) \cdot q^{-\sum_{i \geq a-1} l'_i + 1} \llbracket S_{1,0}^{(\lambda)} \rrbracket \\
 &= \sum_{|\lambda|=r-2} \mathbf{n}(\ell(\lambda) - 1) \sum_{a-1 \triangleleft \lambda} q \cdot \left( q^{-\sum_{i \geq a} l'_i} - q^{-\sum_{i \geq a-1} l'_i} \right) \llbracket S_{1,0}^{(\lambda)} \rrbracket \\
 &= \sum_{|\lambda|=r-2} \mathbf{n}(\ell(\lambda) - 1) \cdot q \cdot (1 - q^{-\ell(\lambda)}) \llbracket S_{1,0}^{(\lambda)} \rrbracket \\
 &= - \sum_{|\lambda|=r-2} \mathbf{n}(\ell(\lambda)) \cdot q^{-\ell(\lambda)+1} \llbracket S_{1,0}^{(\lambda)} \rrbracket.
 \end{aligned}$$

Then (B.31) follows. □

With the help of Lemmas B.3 and B.4, we can prove Proposition 9.2 now.

*Proof of Proposition 9.2.* First, for  $r = 1$ , it follows from (5.16).

For  $r = 2$ , by Corollary 3.10, we have

$$\llbracket S_{1,0}^{(p_1-1)} \rrbracket * \llbracket S_{1,0}^{(p_1-1)} \rrbracket = \mathbf{v}(q+1) \llbracket S_{1,0}^{(p_1-1)} \oplus S_{1,0}^{(p_1-1)} \rrbracket + \frac{\mathbf{v}^{-1}}{q-1} [K_{\delta-\alpha_{11}}].$$

Hence,

$$\begin{aligned}
 &\frac{1}{(q-1)^2} [\widehat{B}_{[1,1],-1}, \widehat{B}_{[1,1],-1}]_{\mathbf{v}^2} * [K_{2\delta-\alpha_{11}}] \\
 &= (1 - \mathbf{v}^2) \llbracket S_{1,0}^{(p_1-1)} \rrbracket * \llbracket S_{1,0}^{(p_1-1)} \rrbracket * [K_{\alpha_{11}}] \\
 &= \mathbf{v}(1 - q^2) \llbracket S_{1,0}^{(p_1-1)} \oplus S_{1,0}^{(p_1-1)} \rrbracket * [K_{\alpha_{11}}] - \mathbf{v}^{-1} [K_{\delta}].
 \end{aligned}$$

On the other hand, by Lemma B.3, we have

$$\begin{aligned}
 &-\frac{1}{(q-1)^2} [\widehat{B}_{[1,1],0}, \widehat{B}_{[1,1],-2}]_{\mathbf{v}^2} * [K_{2\delta-\alpha_{11}}] = \mathbf{v}^{-1} \sum_{[M] \in \mathcal{M}_2} \mathbf{n}(\ell(M) - 1) \llbracket M \rrbracket \\
 &+ \frac{\mathbf{v}}{q-1} \sum_{|\lambda|=2} \mathbf{n}(\ell(\lambda)) \llbracket S_{1,0}^{(\lambda)} \rrbracket + \mathbf{v}(1 - q^2) \llbracket S_{1,0}^{(p_1-1)} \oplus S_{1,0}^{(p_1-1)} \rrbracket * [K_{\alpha_{11}}].
 \end{aligned}$$

It follows that

$$\begin{aligned}
 &-\frac{1}{(q-1)^2} \left( [\widehat{B}_{[1,1],-1}, \widehat{B}_{[1,1],-1}]_{\mathbf{v}^2} + [\widehat{B}_{[1,1],0}, \widehat{B}_{[1,1],-2}]_{\mathbf{v}^2} \right) * [K_{2\delta-\alpha_{11}}] \\
 &= \mathbf{v}^{-1} \sum_{[M] \in \mathcal{M}_{2\delta}} \mathbf{n}(\ell(M) - 1) \llbracket M \rrbracket + \frac{\mathbf{v}}{q-1} \sum_{|\lambda|=2} \mathbf{n}(\ell(\lambda)) \llbracket S_{1,0}^{(\lambda)} \rrbracket + \mathbf{v}^{-1} [K_{\delta}],
 \end{aligned}$$

which also equals to  $\mathbf{v}^{-1} [K_{\delta}] + \widehat{\Theta}_{[1,1],2}$  by (7.10). This proves the result for  $r = 2$ .

For  $r \geq 3$ , by Lemmas B.3 and B.4, we have

$$\begin{aligned}
 &\frac{1}{(q-1)^2} \left( [\widehat{B}_{[1,1],0}, \widehat{B}_{[1,1],-r}]_{\mathbf{v}^2} + [\widehat{B}_{[1,1],-r+1}, \widehat{B}_{[1,1],-1}]_{\mathbf{v}^2} \right) * [K_{r\delta-\alpha_{11}}] \\
 &= - \left( \mathbf{v}^{-1} \sum_{[M] \in \mathcal{M}_{r\delta}} \mathbf{n}(\ell(M) - 1) \llbracket M \rrbracket + \frac{\mathbf{v}}{q-1} \sum_{|\lambda|=r} \mathbf{n}(\ell(\lambda)) \llbracket S_{1,0}^{(\lambda)} \rrbracket \right)
 \end{aligned}$$

$$+ \mathbf{v}^{-2} \left( \mathbf{v}^{-1} \sum_{[M] \in \mathcal{M}_{(r-2)\delta}} \mathbf{n}(\ell(M) - 1) \llbracket [M] \rrbracket + \frac{\mathbf{v}}{q-1} \sum_{|\lambda|=r-2} q^{-\ell(\lambda)} \mathbf{n}(\ell(\lambda)) \llbracket S_{1,0}^{(\lambda)} \rrbracket \right) * [K_\delta].$$

Furthermore, it follows from (7.10) that

$$\begin{aligned} & [\widehat{B}_{[1,1],0}, \widehat{B}_{[1,1],-r}]_{\mathbf{v}^2} + [\widehat{B}_{[1,1],-r+1}, \widehat{B}_{[1,1],-1}]_{\mathbf{v}^2} \\ &= (q-1)^2 \left( -\widehat{\Theta}_{[1,1],r} + \mathbf{v}^{-2} \widehat{\Theta}_{[1,1],r-2} * [K_\delta] \right) * [K_{-r\delta+\alpha_{11}}]. \end{aligned}$$

Then the result follows by induction.  $\square$

### B.3. Proof of Lemma 9.3.

**Lemma B.7.** *Assume  $0 < k < r$  and  $M, N \in \mathcal{T}_{\lambda_1}$ .*

- (1) *If  $\text{Ext}^1(N, S_{1,0}^{(kp_1)})_M \neq 0$ , then  $[M] \in \mathcal{M}_{r\delta}$  if and only if  $[N] \in \mathcal{M}_{(r-k)\delta}$ .*
- (2) *If  $\text{Ext}^1(N, S_{1,0}^{(kp_1-1)})_M \neq 0$ , then  $[M] \in \mathcal{M}_{r\delta}$  if and only if  $[N] \in \mathcal{M}_{(r-k)\delta+\alpha_{11}}$ .*

*Proof.* Similar to the proof of Corollary A.2, we can restrict to consider the weighted projective line of weight type  $\mathbf{p} = (2, 1)$ . Then the result follows by noting that  $S_{1,1}$  is a composition factor of  $\text{top}(M)$  with multiplicity one for  $[M] \in \mathcal{M}_{r\delta}$ .  $\square$

**Lemma B.8.** *For any  $k > 0$  and any  $N \in \mathcal{T}_{\lambda_1}$ , we have*

$$\sum_{M \in \mathcal{T}_{\lambda_1}; \ell(M) = \ell(N) + 1} F_{M, \mathcal{O}}^{\mathcal{O}(k\vec{x}_1) \oplus N} = \frac{|\text{Hom}(\mathcal{O}, N)|}{|\text{Hom}(\mathcal{O}, \text{top}(N))|}.$$

*Proof.* Consider the exact sequence in  $\text{coh } \mathbb{X}$  of the following forms:

$$0 \longrightarrow \mathcal{O} \xrightarrow{(\phi, \psi)^t} \mathcal{O}(k\vec{x}_1) \oplus N \longrightarrow M \longrightarrow 0. \quad (\text{B.32})$$

Then  $M \in \mathcal{T}_{\lambda_1}$  if and only if  $\phi = \kappa x_1^k$  for some non-zero scalar  $\kappa$ . Note that the map  $(\phi, \psi)^t$  in (B.32) is injective if and only if  $\phi$  is injective; there are  $q-1$  many such  $\phi$ 's. Assume  $N$  has a decomposition  $N = \bigoplus_i N_i$  with each  $N_i$  indecomposable, and write  $\psi = (\psi_i)_i$  with  $\psi_i : \mathcal{O} \rightarrow N_i$ . Then one can see that  $\ell(M) = \ell(N)$  or  $\ell(N) + 1$ . Moreover,  $\ell(M) = \ell(N) + 1$  if and only if each  $\psi_i$  is not surjective; there are  $\frac{|\text{Hom}(\mathcal{O}, N_i)|}{|\text{Hom}(\mathcal{O}, \text{top}(N_i))|}$  many such  $\psi_i$ 's. Therefore,

$$\sum_{M \in \mathcal{T}_{\lambda_1}; \ell(M) = \ell(N) + 1} F_{M, \mathcal{O}}^{\mathcal{O}(k\vec{x}_1) \oplus N} = \frac{q-1}{|\text{Aut}(\mathcal{O})|} \cdot \prod_i \frac{|\text{Hom}(\mathcal{O}, N_i)|}{|\text{Hom}(\mathcal{O}, \text{top}(N_i))|} = \frac{|\text{Hom}(\mathcal{O}, N)|}{|\text{Hom}(\mathcal{O}, \text{top}(N))|}.$$

$\square$

**Corollary B.9.** *Assume  $0 < k < r$ . Then for any  $[N_1] \in \mathcal{M}_{r\delta}$ ,  $[N_2] \in \mathcal{M}_{r\delta-\alpha_{11}}$  and any partition  $\mu$ ,*

$$\sum_{[M] \in \mathcal{M}_{(r+k)\delta}; \ell(M) = \ell(N_1) + 1} F_{M, \mathcal{O}}^{\mathcal{O}(k\vec{c}) \oplus N_1} = q^{r-\ell(N_1)+1}; \quad (\text{B.33})$$

$$\sum_{[M] \in \mathcal{M}_{(r+k)\delta}; \ell(M) = \ell(N_2) + 1} F_{M, \mathcal{O}}^{\mathcal{O}(k\vec{c} + \vec{x}_1) \oplus N_2} = q^{r-\ell(N_2)}; \quad (\text{B.34})$$

$$\sum_{\lambda; \ell(\lambda) = \ell(\mu) + 1} F_{S_{1,0}^{(\lambda)}, \mathcal{O}}^{\mathcal{O}(k\vec{c}) \oplus S_{1,0}^{(\mu)}} = q^{|\mu| - \ell(\mu)}. \quad (\text{B.35})$$



*Proof.* We only prove (B.33), since the proofs for the other two formulas are similar. For any  $M \in \mathcal{T}_{\lambda_1}$ ,  $F_{M,\mathcal{O}}^{\mathcal{O}(k\vec{c}) \oplus N_1} \neq 0$  if and only if  $[M] \in \mathcal{M}_{(r+k)\delta}$  by Lemma B.7. Moreover,  $\dim_{\mathbf{k}} \text{Hom}(\mathcal{O}, N_1) = \langle \widehat{\mathcal{O}}, r\delta \rangle = r$  and  $\dim_{\mathbf{k}} \text{Hom}(\mathcal{O}, \text{top}(N_1)) = \ell(N_1) - 1$ . Then the result follows from Lemma B.8.  $\square$

Now we can prove Lemma 9.3.

*Proof of Lemma 9.3.* Recall  $\mathcal{S}$  defined in (A.5). Recall from (9.6) and (9.7) that each term of  $\widehat{\Theta}_{[1,1],r}$  comes from  ${}^\perp\mathcal{S}$ . Hence each extension term in the product  $\widehat{\Theta}_{[1,1],r} * [\mathcal{O}]$  also comes from  ${}^\perp\mathcal{S}$ . Similar to the proof of Corollary A.2, by considering the degrees and the ranks, we see that only the following terms can appear in  $\widehat{\Theta}_{[1,1],r} * [\mathcal{O}]$ :

$$[\mathcal{O}(k\vec{c}) \oplus S_{1,0}^{(\lambda)}], \quad [\mathcal{O}(k\vec{c}) \oplus M], \quad [\mathcal{O}(k\vec{c} + \vec{x}_1) \oplus N],$$

where  $k > 0$ ,  $\lambda$  is a partition of  $r - k$ ,  $[M] \in \mathcal{M}_{(r-k)\delta}$  and  $[N] \in \mathcal{M}_{(r-k)\delta - \alpha_{11}}$ .

Recall from (9.7) that

$$\widehat{\Theta}_{[1,1],r} = \frac{\mathbf{v}}{q-1} \sum_{|\lambda|=r} \mathbf{n}(\ell(\lambda)) [[S_{1,0}^{(\lambda)}]] + \mathbf{v}^{-1} \sum_{[M] \in \mathcal{M}_{r\delta}} \mathbf{n}(\ell(M) - 1) [[M]].$$

For any partition  $\lambda$  of  $r$ , we have  $\langle \widehat{S}_{1,0}^{(\lambda)}, \widehat{\mathcal{O}} \rangle = -r$ . Hence,

$$\begin{aligned} & \frac{\mathbf{v}}{q-1} \sum_{|\lambda|=r} \mathbf{n}(\ell(\lambda)) [[S_{1,0}^{(\lambda)}]] * [[\mathcal{O}]] & (B.36) \\ &= \frac{\mathbf{v}}{q-1} \sum_{|\lambda|=r} \sum_{k \geq 0} \sum_{|\mu|=r-k} \mathbf{n}(\ell(\lambda)) \cdot \mathbf{v}^{-r} F_{S_{1,0}^{(\lambda)}, \mathcal{O}}^{\mathcal{O}(k\vec{c}) \oplus S_{1,0}^{(\mu)}} [[\mathcal{O}(k\vec{c}) \oplus S_{1,0}^{(\mu)}]] \\ &= \frac{\mathbf{v}^{-r+1}}{q-1} \sum_{k \geq 0} \sum_{|\mu|=r-k} \sum_{|\lambda|=r} \mathbf{n}(\ell(\lambda)) \cdot F_{S_{1,0}^{(\lambda)}, \mathcal{O}}^{\mathcal{O}(k\vec{c}) \oplus S_{1,0}^{(\mu)}} [[\mathcal{O}(k\vec{c}) \oplus S_{1,0}^{(\mu)}]] \\ &= \frac{\mathbf{v}^{-r+1}}{q-1} \sum_{|\mu|=r} \mathbf{n}(\ell(\mu)) \cdot F_{S_{1,0}^{(\mu)}, \mathcal{O}}^{\mathcal{O} \oplus S_{1,0}^{(\mu)}} [[\mathcal{O} \oplus S_{1,0}^{(\mu)}]] + \frac{\mathbf{v}^{-r+1}}{q-1} \sum_{k > 0} \sum_{|\mu|=r-k} \\ & \quad \left( \mathbf{n}(\ell(\mu) + 1) \sum_{\lambda; \ell(\lambda) = \ell(\mu) + 1} F_{S_{1,0}^{(\lambda)}, \mathcal{O}}^{\mathcal{O}(k\vec{c}) \oplus S_{1,0}^{(\mu)}} + \mathbf{n}(\ell(\mu)) \sum_{\lambda; \ell(\lambda) = \ell(\mu)} F_{S_{1,0}^{(\lambda)}, \mathcal{O}}^{\mathcal{O}(k\vec{c}) \oplus S_{1,0}^{(\mu)}} \right) [[\mathcal{O}(k\vec{c}) \oplus S_{1,0}^{(\mu)}]] \\ & \stackrel{(B.35)}{=} \frac{\mathbf{v}^{-r+1}}{q-1} \sum_{|\mu|=r} \mathbf{n}(\ell(\mu)) \cdot q^r [[\mathcal{O} \oplus S_{1,0}^{(\mu)}]] + \frac{\mathbf{v}^{-r+1}}{q-1} \sum_{k > 0} \sum_{|\mu|=r-k} \\ & \quad \left( \mathbf{n}(\ell(\mu) + 1) \cdot q^{|\mu| - \ell(\mu)} + \mathbf{n}(\ell(\mu)) \cdot (q^{|\mu|} - q^{|\mu| - \ell(\mu)}) \right) [[\mathcal{O}(k\vec{c}) \oplus S_{1,0}^{(\mu)}]] \\ &= \frac{\mathbf{v}^{r+1}}{q-1} \sum_{|\mu|=r} \mathbf{n}(\ell(\mu)) [[\mathcal{O} \oplus S_{1,0}^{(\mu)}]] - \sum_{k > 0} \sum_{|\mu|=r-k} \mathbf{n}(\ell(\mu)) \cdot \mathbf{v}^{r-2k+1} [[\mathcal{O}(k\vec{c}) \oplus S_{1,0}^{(\mu)}]]. \end{aligned}$$

For any  $[M] \in \mathcal{M}_{r\delta}$ , we have  $\langle \widehat{M}, \widehat{\mathcal{O}} \rangle = -r$  and then by Lemma B.7,

$$[[M]] * [[\mathcal{O}]] = \mathbf{v}^{-r} \sum_{k \geq 0} \sum_{[N_1] \in \mathcal{M}_{(r-k)\delta}} F_{M, \mathcal{O}}^{\mathcal{O}(k\vec{c}) \oplus N_1} [[\mathcal{O}(k\vec{c}) \oplus N_1]] \quad (B.37)$$

$$+ \mathbf{v}^{-r} \sum_{k \geq 0} \sum_{[N_2] \in \mathcal{M}_{(r-k)\delta - \alpha_{11}}} F_{M, \mathcal{O}}^{\mathcal{O}(k\vec{c} + \vec{x}_1) \oplus N_2} [\mathcal{O}(k\vec{c} + \vec{x}_1) \oplus N_2].$$

For any  $k \geq 0$  and  $[N_2] \in \mathcal{M}_{(r-k)\delta - \alpha_{11}}$ , the coefficient of  $[\mathcal{O}(k\vec{c} + \vec{x}_1) \oplus N_2]$  in  $\widehat{\Theta}_{[1,1],r} * [\mathcal{O}]$  equals to (up to a power of  $\mathbf{v}$ )

$$\begin{aligned} & \sum_{[M] \in \mathcal{M}_{r\delta}} \mathbf{n}(\ell(M) - 1) \cdot F_{M, \mathcal{O}}^{\mathcal{O}(k\vec{c} + \vec{x}_1) \oplus N_2} \\ &= \mathbf{n}(\ell(N_2)) \sum_{[M] \in \mathcal{M}_{r\delta}, \ell(M) = \ell(N_2) + 1} F_{M, \mathcal{O}}^{\mathcal{O}(k\vec{c} + \vec{x}_1) \oplus N_2} + \mathbf{n}(\ell(N_2) - 1) \sum_{[M] \in \mathcal{M}_{r\delta}, \ell(M) = \ell(N_2)} F_{M, \mathcal{O}}^{\mathcal{O}(k\vec{c} + \vec{x}_1) \oplus N_2} \\ &\stackrel{\text{(B.34)}}{=} \mathbf{n}(\ell(N_2)) \cdot q^{r-k-\ell(N_2)} + \mathbf{n}(\ell(N_2) - 1) \cdot (q^{r-k} - q^{r-k-\ell(N_2)}) = 0. \end{aligned} \quad (\text{B.38})$$

Similarly, for  $k > 0$  and  $[N_1] \in \mathcal{M}_{(r-k)\delta}$  we have

$$\begin{aligned} & \sum_{[M] \in \mathcal{M}_{r\delta}} \mathbf{n}(\ell(M) - 1) \cdot F_{M, \mathcal{O}}^{\mathcal{O}(k\vec{c}) \oplus N_1} \\ &= \mathbf{n}(\ell(N_1)) \sum_{[M] \in \mathcal{M}_{r\delta}, \ell(M) = \ell(N_1) + 1} F_{M, \mathcal{O}}^{\mathcal{O}(k\vec{c}) \oplus N_1} + \mathbf{n}(\ell(N_1) - 1) \sum_{[M] \in \mathcal{M}_{r\delta}, \ell(M) = \ell(N_1)} F_{M, \mathcal{O}}^{\mathcal{O}(k\vec{c}) \oplus N_1} \\ &\stackrel{\text{(B.33)}}{=} \mathbf{n}(\ell(N_1)) \cdot q^{r-k-\ell(N_1)+1} + \mathbf{n}(\ell(N_1) - 1) \cdot (q^{r-k} - q^{r-k-\ell(N_1)+1}) \\ &= \mathbf{n}(\ell(N_1) - 1) \cdot q^{r-k-\ell(N_1)+1} (1 - q^{\ell(N_1)} + q^{\ell(N_1)-1} - 1) \\ &= \mathbf{n}(\ell(N_1) - 1) \cdot q^{r-k} (1 - q); \end{aligned} \quad (\text{B.39})$$

while for  $k = 0$  and  $[N_1] \in \mathcal{M}_{r\delta}$ ,

$$\sum_{[M] \in \mathcal{M}_{r\delta}} \mathbf{n}(\ell(M) - 1) \cdot F_{M, \mathcal{O}}^{\mathcal{O} \oplus N_1} = \mathbf{n}(\ell(N_1) - 1) \cdot F_{N_1, \mathcal{O}}^{\mathcal{O} \oplus N_1} = \mathbf{n}(\ell(N_1) - 1) \cdot q^r. \quad (\text{B.40})$$

Combining with (B.37)–(B.40), we obtain

$$\mathbf{v}^{-1} \sum_{[M] \in \mathcal{M}_{r\delta}} \mathbf{n}(\ell(M) - 1) [M] * [\mathcal{O}] \quad (\text{B.41})$$

$$\begin{aligned} &= \mathbf{v}^{-r-1} \sum_{k \geq 0} \sum_{[N] \in \mathcal{M}_{(r-k)\delta}} \left( \sum_{[M] \in \mathcal{M}_{r\delta}} \mathbf{n}(\ell(M) - 1) F_{M, \mathcal{O}}^{\mathcal{O}(k\vec{c}) \oplus N} \right) [\mathcal{O}(k\vec{c}) \oplus N] \\ &= \mathbf{v}^{-r-1} \sum_{[N] \in \mathcal{M}_{r\delta}} \mathbf{n}(\ell(N) - 1) [\mathcal{O} \oplus N] \\ &\quad + \sum_{k > 0} \sum_{[N] \in \mathcal{M}_{(r-k)\delta}} \mathbf{n}(\ell(N) - 1) \cdot (1 - q) \mathbf{v}^{r-2k-1} [\mathcal{O}(k\vec{c}) \oplus N]. \end{aligned} \quad (\text{B.42})$$

Combining (B.41) and (B.36), we finish the proof of Lemma 9.3.  $\square$

**B.4. Proof of Lemma 9.4.** For any  $[M] \in \mathcal{M}_{r\delta}$ , we have  $\langle \widehat{\mathcal{O}}, \widehat{M} \rangle = r$ ,  $\text{Ext}^1(\mathcal{O}, M) = 0$  and  $\text{Hom}(\mathcal{O}, M) \cong \mathbf{k}^r$ . For any non-zero morphism  $f : \mathcal{O} \rightarrow M$ , we have  $\text{Im}(f) \cong S_{1,0}^{(kp_1)}$  or  $S_{1,0}^{(kp_1-1)}$  for some  $k > 0$ .

If  $\text{Im}(f) = S_{1,0}^{(kp_1)}$ , then  $\text{Ker}(f) = \mathcal{O}(-k\vec{c})$  and  $[\text{Coker}(f)] \in \mathcal{M}_{(r-k)\delta}$  by Lemma B.7. For any  $[N] \in \mathcal{M}_{(r-k)\delta}$ , we have

$$\begin{aligned} & |\{f : \mathcal{O} \rightarrow M \mid \text{Ker}(f) \cong \mathcal{O}(-k\vec{c}), \text{Coker}(f) \cong N\}| \\ &= |\{g : S_{1,0}^{(kp_1)} \twoheadrightarrow M \mid \text{Coker}(g) \cong N\}| \\ &= F_{N, S_{1,0}^{(kp_1)}}^M \cdot |\text{Aut}(S_{1,0}^{(kp_1)})|. \end{aligned}$$

Similarly, if  $\text{Im}(f) = S_{1,0}^{(kp_1-1)}$ , then  $\text{Ker}(f) = \mathcal{O}(-k\vec{c} + \vec{x}_1)$  and  $[\text{Coker}(f)] \in \mathcal{M}_{(r-k)\delta + \alpha_{11}}$ . For any  $[N'] \in \mathcal{M}_{(r-k)\delta + \alpha_{11}}$ ,

$$\begin{aligned} & |\{f : \mathcal{O} \rightarrow M \mid \text{Ker}(f) \cong \mathcal{O}(-k\vec{c} + \vec{x}_1), \text{Coker}(f) \cong N'\}| \\ &= |\{g : S_{1,0}^{(kp_1-1)} \twoheadrightarrow M \mid \text{Coker}(g) \cong N'\}| \\ &= F_{N', S_{1,0}^{(kp_1-1)}}^M \cdot |\text{Aut}(S_{1,0}^{(kp_1-1)})|. \end{aligned}$$

Then by (3.11), we have

$$\begin{aligned} [\mathcal{O}] * [M] &= \mathbf{v}^{-r} \sum_{k \geq 0} \sum_{[N] \in \mathcal{M}_{(r-k)\delta}} F_{N, S_{1,0}^{(kp_1)}}^M |\text{Aut}(S_{1,0}^{(kp_1)})| [\mathcal{O}(-k\vec{c}) \oplus N] * [K_{k\delta}] \\ &\quad + \mathbf{v}^{-r} \sum_{k \geq 1} \sum_{[N'] \in \mathcal{M}_{(r-k)\delta + \alpha_{11}}} F_{N', S_{1,0}^{(kp_1-1)}}^M |\text{Aut}(S_{1,0}^{(kp_1-1)})| [\mathcal{O}(-k\vec{c} + \vec{x}_1) \oplus N'] * [K_{k\delta - \alpha_{11}}]; \end{aligned}$$

or equivalently, by using the Riedtmann-Peng formula, we have

$$\begin{aligned} [[\mathcal{O}] * [M]] &= \mathbf{v}^{-r} \sum_{k \geq 0} \sum_{[N] \in \mathcal{M}_{(r-k)\delta}} q^{r-k} \frac{|\text{Ext}^1(N, S_{1,0}^{(kp_1)})_M|}{|\text{Hom}(N, S_{1,0}^{(kp_1)})|} [[\mathcal{O}(-k\vec{c}) \oplus N] * [K_{k\delta}]] \quad (\text{B.43}) \\ &\quad + \mathbf{v}^{-r} \sum_{k \geq 1} \sum_{[N'] \in \mathcal{M}_{(r-k)\delta + \alpha_{11}}} q^{r-k+1} \frac{|\text{Ext}^1(N', S_{1,0}^{(kp_1-1)})_M|}{|\text{Hom}(N', S_{1,0}^{(kp_1-1)})|} [[\mathcal{O}(-k\vec{c} + \vec{x}_1) \oplus N'] * [K_{k\delta - \alpha_{11}}]]. \end{aligned}$$

Similar arguments show that

$$[[\mathcal{O}] * [S_{1,0}^{(\lambda)}]] = \mathbf{v}^{-r} \sum_{k \geq 0} \sum_{|\mu| = r-k} q^{r-k} \frac{|\text{Ext}^1(S_{1,0}^{(\mu)}, S_{1,0}^{(kp_1)})_{S_{1,0}^{(\lambda)}}|}{|\text{Hom}(S_{1,0}^{(\mu)}, S_{1,0}^{(kp_1)})|} \cdot [[\mathcal{O}(-k\vec{c}) \oplus S_{1,0}^{(\mu)}] * [K_{k\delta}]]. \quad (\text{B.44})$$

Now we can prove Lemma 9.4.

*Proof of Lemma 9.4.* For any  $k > 0$  and  $[L] \in \mathcal{M}_{(r-k)\delta + \alpha_{11}}$ , we have  $\langle L, S_{1,0}^{(kp_1-1)} \rangle = -1$  and  $\dim_{\mathbf{k}} \text{Hom}(L, S_{1,0}^{(kp_1-1)}) = \ell(L) - 1$ . Then the coefficient of  $[[\mathcal{O}(-k\vec{c} + \vec{x}_1) \oplus L] * [K_{k\delta - \alpha_{11}}]]$  in the product  $[[\mathcal{O}] * \widehat{\Theta}_{[1,1],r}]$  equals to (up to a power of  $\mathbf{v}$ )

$$\begin{aligned} & \sum_{[M] \in \mathcal{M}_{r\delta}} \mathbf{n}(\ell(M) - 1) \cdot |\text{Ext}^1(L, S_{1,0}^{(kp_1-1)})_M| \quad (\text{B.45}) \\ &= \mathbf{n}(\ell(L)) \sum_{[M]; \ell(M) = \ell(L) + 1} |\text{Ext}^1(L, S_{1,0}^{(kp_1-1)})_M| + \mathbf{n}(\ell(L) - 1) \sum_{[M]; \ell(M) = \ell(L)} |\text{Ext}^1(L, S_{1,0}^{(kp_1-1)})_M| \end{aligned}$$

$$\begin{aligned}
&= \mathbf{n}(\ell(L)) \frac{|\mathrm{Hom}(L, S_{1,0}^{(kp_1-1)})|}{|\mathrm{Hom}(L, S_{1,0}^{(p_1-1)})|} + \mathbf{n}(\ell(L) - 1) \left( |\mathrm{Ext}^1(L, S_{1,0}^{(kp_1-1)})| - \frac{|\mathrm{Hom}(L, S_{1,0}^{(kp_1-1)})|}{|\mathrm{Hom}(L, S_{1,0}^{(p_1-1)})|} \right) \\
&= \mathbf{n}(\ell(L) - 1) \left( \frac{|\mathrm{Hom}(L, S_{1,0}^{(kp_1-1)})|}{|\mathrm{Hom}(L, S_{1,0}^{(p_1-1)})|} (1 - q^{\ell(L)}) + \frac{|\mathrm{Hom}(L, S_{1,0}^{(kp_1-1)})|}{q^{\langle L, S_{1,0}^{(kp_1-1)} \rangle}} - \frac{|\mathrm{Hom}(L, S_{1,0}^{(kp_1-1)})|}{|\mathrm{Hom}(L, S_{1,0}^{(p_1-1)})|} \right) \\
&= \mathbf{n}(\ell(L) - 1) \cdot |\mathrm{Hom}(L, S_{1,0}^{(kp_1-1)})| \cdot \left( \frac{1 - q^{\ell(L)}}{q^{\ell(L)-1}} + q - \frac{1}{q^{\ell(L)-1}} \right) = 0.
\end{aligned}$$

Here the second equality follows from Lemma 5.6.

Similarly, for any  $k > 0$  and  $[N] \in \mathcal{M}_{(r-k)\delta}$ , we have  $\dim_{\mathbf{k}} \mathrm{Ext}^1(N, S_{1,0}^{(p_1-1)}) = \ell(N) - 1$ . By using Lemma 5.7,

$$\begin{aligned}
&\sum_{[M] \in \mathcal{M}_{r\delta}} \mathbf{n}(\ell(M) - 1) \cdot |\mathrm{Ext}^1(N, S_{1,0}^{(kp_1)})_M| \tag{B.46} \\
&= \mathbf{n}(\ell(N)) \sum_{\ell(M)=\ell(N)+1} |\mathrm{Ext}^1(N, S_{1,0}^{(kp_1)})_M| + \mathbf{n}(\ell(N) - 1) \sum_{\ell(M)=\ell(N)} |\mathrm{Ext}^1(N, S_{1,0}^{(kp_1)})_M| \\
&= \mathbf{n}(\ell(N)) \frac{|\mathrm{Hom}(N, S_{1,0}^{(kp_1)})|}{q^{\ell(N)-1}} + \mathbf{n}(\ell(N) - 1) \left( |\mathrm{Ext}^1(N, S_{1,0}^{(kp_1)})| - \frac{|\mathrm{Hom}(N, S_{1,0}^{(kp_1)})|}{q^{\ell(N)-1}} \right) \\
&= \mathbf{n}(\ell(N) - 1) \cdot (1 - q) \cdot |\mathrm{Hom}(N, S_{1,0}^{(kp_1)})|;
\end{aligned}$$

while for  $k = 0$ ,

$$\sum_{[M] \in \mathcal{M}_{r\delta}} \mathbf{n}(\ell(M) - 1) \cdot |\mathrm{Ext}^1(N, S_{1,0}^{(kp_1)})_M| = \mathbf{n}(\ell(N) - 1). \tag{B.47}$$

Therefore, combining (B.43)–(B.47), we obtain

$$\begin{aligned}
&\mathbf{v}^{-1}[\mathcal{O}] * \sum_{[M] \in \mathcal{M}_{r\delta}} \mathbf{n}(\ell(M) - 1)[M] \tag{B.48} \\
&= \mathbf{v}^{-(r+1)} \sum_{k \geq 0} \sum_{[N] \in \mathcal{M}_{(r-k)\delta}} \sum_{[M] \in \mathcal{M}_{r\delta}} \mathbf{n}(\ell(M) - 1) q^{r-k} \frac{|\mathrm{Ext}^1(N, S_{1,0}^{(kp_1)})_M|}{|\mathrm{Hom}(N, S_{1,0}^{(kp_1)})|} [[\mathcal{O}(-k\vec{c}) \oplus N] * [K_{k\delta}]] \\
&= \sum_{[N] \in \mathcal{M}_{r\delta}} \mathbf{n}(\ell(N) - 1) \mathbf{v}^{r-1} [[\mathcal{O} \oplus N] \\
&\quad + \sum_{k > 0} \sum_{[N] \in \mathcal{M}_{(r-k)\delta}} (1 - q) \mathbf{v}^{r-2k-1} \mathbf{n}(\ell(N) - 1) [[\mathcal{O}(-k\vec{c}) \oplus N] * [K_{k\delta}]].
\end{aligned}$$

On the other hand, for any  $k > 0$  and  $|\mu| = r - k$ , by using Lemma 5.7 we obtain

$$\begin{aligned}
&\sum_{|\lambda|=r} \mathbf{n}(\ell(\lambda)) \cdot |\mathrm{Ext}^1(S_{1,0}^{(\mu)}, S_{1,0}^{(kp_1)})_{S_{1,0}^{(\lambda)}}| \tag{B.49} \\
&= \mathbf{n}(\ell(\mu) + 1) \sum_{\ell(\lambda)=\ell(\mu)+1} |\mathrm{Ext}^1(S_{1,0}^{(\mu)}, S_{1,0}^{(kp_1)})_{S_{1,0}^{(\lambda)}}| + \mathbf{n}(\ell(\mu)) \sum_{\ell(\lambda)=\ell(\mu)} |\mathrm{Ext}^1(S_{1,0}^{(\mu)}, S_{1,0}^{(kp_1)})_{S_{1,0}^{(\lambda)}}|
\end{aligned}$$

$$\begin{aligned}
 &= \mathbf{n}(\ell(\mu) + 1) \cdot \frac{|\mathrm{Hom}(S_{1,0}^{(\mu)}, S_{1,0}^{(kp_1)})|}{|\mathrm{Hom}(S_{1,0}^{(\mu)}, S_{1,0}^{(p_1)})|} + \mathbf{n}(\ell(\mu)) \left( |\mathrm{Ext}^1(S_{1,0}^{(\mu)}, S_{1,0}^{(kp_1)})| - \frac{|\mathrm{Hom}(S_{1,0}^{(\mu)}, S_{1,0}^{(kp_1)})|}{|\mathrm{Hom}(S_{1,0}^{(\mu)}, S_{1,0}^{(p_1)})|} \right) \\
 &= \mathbf{n}(\ell(\mu)) \cdot |\mathrm{Hom}(S_{1,0}^{(\mu)}, S_{1,0}^{(kp_1)})| \cdot \left( \frac{1 - q^{\ell(\mu)+1}}{q^{\ell(\mu)}} + 1 - \frac{1}{q^{\ell(\mu)}} \right) \\
 &= (1 - q) \cdot \mathbf{n}(\ell(\mu)) \cdot |\mathrm{Hom}(S_{1,0}^{(\mu)}, S_{1,0}^{(kp_1)})|,
 \end{aligned}$$

while for  $k = 0$  and  $|\mu| = r$ ,

$$\sum_{|\lambda|=r} \mathbf{n}(\ell(\lambda)) \cdot |\mathrm{Ext}^1(S_{1,0}^{(\mu)}, S_{1,0}^{(kp_1)})_{S_{1,0}^{(\lambda)}}| = \mathbf{n}(\ell(\mu)). \quad (\text{B.50})$$

Combining (B.44), (B.49) and (B.50), we obtain

$$\begin{aligned}
 &\frac{\mathbf{v}}{q-1} \llbracket \mathcal{O} \rrbracket * \sum_{|\lambda|=r} \mathbf{n}(\ell(\lambda)) \llbracket S_{1,0}^{(\lambda)} \rrbracket \quad (\text{B.51}) \\
 &= \frac{\mathbf{v}^{-r+1}}{q-1} \sum_{k \geq 0} \sum_{|\mu|=r-k} \sum_{|\lambda|=r} \mathbf{n}(\ell(\lambda)) \cdot q^{r-k} \frac{|\mathrm{Ext}^1(S_{1,0}^{(\mu)}, S_{1,0}^{(kp_1)})_{S_{1,0}^{(\lambda)}}|}{|\mathrm{Hom}(S_{1,0}^{(\mu)}, S_{1,0}^{(kp_1)})|} \cdot \llbracket \mathcal{O}(-k\vec{c}) \oplus S_{1,0}^{(\mu)} \rrbracket * \llbracket K_{k\delta} \rrbracket \\
 &= \frac{\mathbf{v}^{r+1}}{q-1} \sum_{|\mu|=r} \mathbf{n}(\ell(\mu)) \cdot \llbracket \mathcal{O} \oplus S_{1,0}^{(\mu)} \rrbracket - \sum_{k > 0} \sum_{|\mu|=r-k} \mathbf{v}^{r-2k+1} \cdot \mathbf{n}(\ell(\mu)) \cdot \llbracket \mathcal{O}(-k\vec{c}) \oplus S_{1,0}^{(\mu)} \rrbracket * \llbracket K_{k\delta} \rrbracket.
 \end{aligned}$$

Combining (B.48) and (B.51), we finish the proof of Lemma 9.4.  $\square$

## REFERENCES

- [BK19] M. Balagovic and S. Kolb, *Universal K-matrix for quantum symmetric pairs*, J. Reine Angew. Math. **747** (2019), 299–353.
- [BW18a] H. Bao and W. Wang, *A new approach to Kazhdan-Lusztig theory of type B via quantum symmetric pairs*, Astérisque **402**, 2018, vii+134pp.
- [BW18b] H. Bao and W. Wang, *Canonical bases arising from quantum symmetric pairs*, Invent. Math. **213** (2018), 1099–1177.
- [BK20] P. Baseilhac and S. Kolb, *Braid group action and root vectors for the q-Onsager algebra*, Transform. Groups **25** (2020), 363–389.
- [BKa01] P. Baumann and C. Kassel, *The Hall algebra of the category of coherent sheaves on the projective line*, J. Reine Angew. Math. **533** (2001), 207–233.
- [Be94] J. Beck, *Braid group actions and quantum affine algebras*, Commun. Math. Phys. **165** (1994), 555–568.
- [Br13] T. Bridgeland, *Quantum groups via Hall algebras of complexes*, Ann. Math. **177** (2013), 739–759.
- [BS12a] I. Burban and O. Schiffmann, *On the Hall algebra of an elliptic curve, I*, Duke Math. J. **161** (2012), 1171–1231.
- [BS12b] I. Burban and O. Schiffmann, *Two descriptions of the quantum affine algebra  $U_v(\widehat{\mathfrak{sl}}_2)$  via Hall algebra approach*, Glasgow Math. J. **54** (2012), 283–307.
- [BS13] I. Burban and O. Schiffmann, *The composition Hall algebra of a weighted projective line*, J. Reine Angew. Math. **679** (2013), 75–124.
- [Cr10] T. Cramer, *Double Hall algebras and derived equivalences*, Adv. Math. **224** (2010), 1097–1120.
- [Da15] I. Damiani, *From the Drinfeld realization to the Drinfeld-Jimbo presentation of affine quantum algebras: injectivity*, Publ. Res. Inst. Math. Sci. **51** (2015), 131–171.
- [DJX12] R. Dou, Y. Jiang and J. Xiao, *Hall algebra approach to Drinfeld’s presentation of quantum loop algebras*, Adv. Math. **231** (2012), 2593–2625.

- [Dr88] V. Drinfeld, *A new realization of Yangians and quantized affine algebras*, Soviet Math. Dokl. **36** (1988), 212–216.
- [Ga80] H. Garland, *The arithmetic theory of loop groups*, Publ. Math. Inst. Hautes Études Sci. **52** (1980), 5–136.
- [GL87] W. Geigle and H. Lenzing, *A class of weighted projective curves arising in representation theory of finite dimensional algebras*, Singularities, representation of algebras, and vector bundles, Lect. Notes Math. **1273** (1987), 265–297.
- [Gor18] M. Gorsky, *Semi-derived and derived Hall algebras for stable categories*, IMRN, Vol. **2018**, No. 1, 138–159.
- [Gr95] J.A. Green, *Hall algebras, hereditary algebras and quantum groups*, Invent. Math. **120** (1995), 361–377.
- [Hu05] A. Hubery, *Symmetric functions and the centre of the Ringel-Hall algebra of a cyclic quiver*, Math. Z. **251** (2005), 705–719.
- [Ka97] M. Kapranov, *Eisenstein series and quantum affine algebras*, J. Math. Sci. **84** (1997), 1311–1360.
- [Ko14] S. Kolb, *Quantum symmetric Kac-Moody pairs*, Adv. Math. **267** (2014), 395–469.
- [KP11] S. Kolb and J. Pellegrini, *Braid group actions on coideal subalgebras of quantized enveloping algebras*, J. Algebra **336** (2011), 395–416.
- [Ke05] B. Keller, *On triangulated orbit categories*, Doc. Math. **10** (2005), 551–581.
- [Let99] G. Letzter, *Symmetric pairs for quantized enveloping algebras*, J. Algebra **220** (1999), 729–767.
- [LP21] M. Lu and L. Peng, *Semi-derived Ringel-Hall algebras and Drinfeld double*, Adv. Math. **383** (2021), 107668.
- [LRW23] M. Lu, S. Ruan and W. Wang,  *$\iota$ Hall algebra of the projective line and  $q$ -Onsager algebra*, Trans. Amer. Math. Soc. **376** (2) (2023), 1475–1505.
- [LW21a] M. Lu and W. Wang, *Hall algebras and quantum symmetric pairs II: reflection functors*, Commun. Math. Phys. **381** (2021), 799–855.
- [LW21b] M. Lu and W. Wang, *A Drinfeld type presentation of affine  $\iota$ quantum groups I: split ADE type*, Adv. Math. **393** (2021), 108111.
- [LW22a] M. Lu and W. Wang, *Hall algebras and quantum symmetric pairs I: foundations*, Proc. London Math. Soc. **124** (2022), 1–82.
- [LW22b] M. Lu and W. Wang, *Braid group symmetries on quasi-split  $\iota$ quantum groups via  $\iota$ Hall algebras*, Selecta Math. **28** (2022), no. 5, Paper No. 84.
- [LW23] M. Lu and W. Wang, *Hall algebras and quantum symmetric pairs of Kac-Moody type*, Adv. Math. **430** (2023), 109215.
- [Lus90] G. Lusztig, *Canonical bases arising from quantized enveloping algebras*, J. Amer. Math. Soc. **3** (1990), 447–498.
- [MRY90] R.V. Moody, S.E. Rao and T. Yokonuma, *Toroidal Lie algebras and vertex representations*, Geom. Dedicata **35** (1990), 283–307.
- [Rin84] C.M. Ringel, *Tame algebras and quadratic forms*, Lect. Notes in Math., **1099**, Springer, 1984.
- [Rin90] C.M. Ringel, *Hall algebras and quantum groups*, Invent. Math. **101** (1990), 583–591.
- [Sch04] O. Schiffmann, *Noncommutative projective curves and quantum loop algebras*, Duke Math. J. **121** (2004), 113–168.
- [Sch12] O. Schiffmann, *Lectures on Hall algebras*, In: Geometric methods in representation theory. II, 1–141, Sémin. Congr., 24-II, Soc. Math. France, Paris (2012).
- [St17] T. Stai, *The triangulated hull of periodic complexes*, Math. Res. Lett. **25** (2017), 199–236.
- [W22] W. Wang, *Quantum symmetric pairs*, Proceedings of ICM 2022, [arxiv:2112.10911](https://arxiv.org/abs/2112.10911)

DEPARTMENT OF MATHEMATICS, SICHUAN UNIVERSITY, CHENGDU 610064, P.R.CHINA  
*Email address:* luming@scu.edu.cn

SCHOOL OF MATHEMATICAL SCIENCES, XIAMEN UNIVERSITY, XIAMEN 361005, P.R.CHINA  
*Email address:* sqruan@xmu.edu.cn