ON THE DISCRETISED ABC SUM-PRODUCT PROBLEM

TUOMAS ORPONEN

ABSTRACT. Let $\alpha, \beta, \gamma \in (0, 1)$, and let $A, B, C \subset [0, 1]$ be finite sets with $|A| = \delta^{-\alpha}$, $|B| = \delta^{-\beta}$, and $|C| = \delta^{-\gamma}$ for some small $\delta \in 2^{-\mathbb{N}}$. It follows from the Szemerédi-Trotter theorem that if $\gamma > \alpha - \beta$, then there exists $c \in C$ with $|A + cB| \gg |A|$.

This paper studies the δ -discretised version of the problem above, where every occurrence of $|\cdot|$ (cardinality) is replaced by $|\cdot|_{\delta}$ (δ -covering number). Bourgain already solved the problem in the case $\alpha = \beta$. Roughly speaking, he showed that if $\gamma > 0$, and B, C satisfy suitable non-concentration conditions, then $|A + cB|_{\delta} \gg |A|_{\delta}$ for some $c \in C$. I extend Bourgain's theorem to the case $\beta \leq \alpha$, under the assumption $\gamma > (\alpha - \beta)/(1 - \beta)$. I conjecture that the sharp assumption would be $\gamma > \alpha - \beta$.

CONTENTS

1. Introduction	2
1.1. Related work	3
1.2. Proof sketch	5
1.3. Acknowledgements	6
2. Notation and preliminaries	6
2.1. Dyadic cubes and covering numbers	6
2.2. Entropy	7
3. Three initial reductions	8
3.1. Reduction to the case where <i>B</i> has small doubling	8
3.2. Reducing the Frostman constant of ν	11
3.3. Removing reference to subsets	14
3.4. Bonus reduction	17
4. Proof of Theorem 3.25	18
4.1. Preliminaries	18
4.2. Shmerkin's inverse theorem	18
4.3. Applying the inverse theorem	20
4.4. Pruning B' to improve separation I	22
4.5. Intervals with small but non-zero <i>B</i> '-branching	23
4.6. Branching of A' on typical intervals in \mathcal{N}_+	26
4.7. Pruning B' to improve separation II	27
4.8. Elementary projection estimates	29
4.9. Projecting pieces of $A' \times B''$	31

Date: December 6, 2021.

²⁰¹⁰ Mathematics Subject Classification. 11B30 (primary) 28A80 (secondary).

Key words and phrases. Discretised sum-product problem, Projections.

T.O. is supported by the Academy of Finland via the projects *Quantitative rectifiability in Euclidean and non-Euclidean spaces* and *Incidences on Fractals*, grant Nos. 309365, 314172, 321896.

4.10. Final multiscale argument References

1. INTRODUCTION

Let $A, B, C \subset \mathbb{R}$ be large but finite sets. Is it true that there exists some $c \in C$ such that $|A + cB| \gg |A|$? Here $|\cdot|$ refers to cardinality. Not necessarily: consider for example

$$A_n = \left\{ \frac{1}{n^{1/2}}, \frac{2}{n^{1/2}}, \dots, 1 \right\} \quad \text{and} \quad B_n = \left\{ \frac{1}{n^{1/4}}, \frac{2}{n^{1/4}}, \dots, 1 \right\} = C_n.$$
(1.1)

It is not hard to check that for every $\epsilon > 0$, there exists $n \in \mathbb{N}$ such that $|A_n + B_n C_n| \leq n^{\epsilon}|A|$, so in particular $|A_n + cB_n| \leq n^{\epsilon}|A|$ for all $c \in C_n$. The problem can be fixed by adding one assumption: $|B||C| \gg |A|$. Then, a positive answer to the question follows easily from the Szemerédi-Trotter theorem [36] applied to the planar set $A \times B$. The requirement $|B||C| \gg |A|$ is also necessary, as (1.1) shows.

The *ABC* sum-product problem, stated above, also makes sense in contexts where the Szemerédi-Trotter bound is not available, for example if $A, B, C \subset \mathbb{Z}_p$, and $p \in \mathbb{N}$ is prime. Again, it turns out that the lower bound $|B||C| \gg |A|$ yields the existence of $c \in C$ with $|A + cB| \gg |A|$. One way to show this is to adapt elementary techniques of Garaev [10], Glibichuk and Konyagin [11], and Bourgain [4]. The details can be found in [28]. Another way is to apply directly an incidence bound in finite fields due to Stevens and de Zeeuw [35]. The theorem of Stevens and de Zeeuw gives a stronger lower bound for |A + cB| than the elementary approach (see [28, Proposition 1.3] for the details), but ultimately relies on the polynomial method.

The main purpose of this paper is to consider the δ -discretised ABC sum-product problem in \mathbb{R} . This is otherwise the same problem as the one we started with, but instead of counting the cardinality of |A + cB|, we seek lower bounds for the δ -covering number $|A + cB|_{\delta}$ for some small scale $\delta > 0$. We will also assume that the sets A, B, C are δ separated, and have cardinalities $|A| = \delta^{-\alpha}$, $|B| = \delta^{-\beta}$, and $|C| = \delta^{-\gamma}$. In this variant of the problem, hypotheses on |B||C| need to be coupled with additional non-concentration conditions to hope for positive results. The following theorem of Bourgain [5] from 2010 (extending his own work [2] from 2003) treats the case A = B:

Theorem 1.2 (Bourgain). Given $\alpha \in (0, 1)$ and $\gamma, \kappa > 0$, there exist $\epsilon_0, \epsilon > 0$ such that the following holds for $\delta > 0$ sufficiently small.

Let ν be a probability measure on [0,1] satisfying $\nu(B(x,r)) \leq r^{\gamma}$ for all $x \in \mathbb{R}$ and $0 < r \leq \delta^{\epsilon_0}$. Let additionally $A \subset [0,1]$ be a δ -separated set with $|A| \geq \delta^{-\alpha}$, which also satisfies the non-concentration condition $|A \cap B(x,r)| \leq r^{\kappa}|A|$ for $x \in \mathbb{R}$ and $\delta \leq r \leq \delta^{\epsilon_0}$.

Then, there exists a point $c \in \operatorname{spt}(\nu)$ *such that*

$$|A + cA|_{\delta} \ge \delta^{-\alpha - \epsilon}. \tag{1.3}$$

Here $|\cdot|_{\delta}$ *refers to the* δ *-covering number of* A*.*

To see the connection to the *ABC* problem, let $C \subset [0,1]$ be a δ -separated set satisfying $|C \cap B(x,r)| \leq r^{\gamma}|C|$ for all $x \in \mathbb{R}$ and $\delta \leq r \leq \delta^{\epsilon_0}$. Then the uniformly distributed probability measure ν on the δ -neighbourhood of *C* satisfies $\nu(B(x,r)) \leq r^{\gamma}$, and it follows from (1.3) (after a moment of thought) that there exists $c \in C$ with $|A + cA|_{\delta} \geq \delta^{-\alpha-\epsilon}$.

Theorem 1.2 formally only treats the case A = B, but an inspection of its proof (or, more directly, an application of [5, Theorem 3]), reveals that the result remains valid for two different δ -separated sets $A, B \subset [0, 1]$, provided that |A| = |B|, or at least $|B| \approx |A|$. The precise meaning of " \approx " is defined via the various constants appearing in [5, Theorem 3]. To the best of my knowledge, Theorem 1.2 does not cover the case where $|A| = \delta^{-\alpha}$ and $|B| = \delta^{-\beta}$ with $\beta < \alpha$ (the case $\beta > \alpha$ is not relevant here: then $|A + cB|_{\delta} \gtrsim |B| \gg |A|$ for any $c \in \mathbb{R}$ with $|c| \sim 1$).

The following conjecture corresponds to the assumption $|B||C| \gg |A|$ which suffices in the discrete variants (on \mathbb{R} and \mathbb{Z}_p) of the *ABC* problem:

Conjecture 1.4. Let $\alpha, \beta, \gamma \in (0, 1)$ with $\beta \leq \alpha$ and $\gamma > \alpha - \beta$. Assume that $A, B, C \subset [0, 1]$ are δ -separated sets with cardinalities $|A| \leq \delta^{-\alpha}$, $|B| = \delta^{-\beta}$, and $|C| = \delta^{-\gamma}$. Assume moreover that $|B \cap B(x, r)| \leq r^{\beta}|B|$ and $|C \cap B(x, r)| \leq r^{\gamma}|C|$ for all $x \in \mathbb{R}$ and r > 0. Then, there exists $\epsilon = \epsilon(\alpha, \beta, \gamma) > 0$ and a point $c \in C$ such that $|A + cB|_{\delta} \geq_{\alpha,\beta,\gamma} \delta^{-\epsilon}|A|$.

The lower bound for γ in Conjecture 1.4 is necessary, but the non-concentration assumptions on *B* and *C* are quite likely not sharp. The main result of this paper is the following partial result, where the lower bound $\gamma > \alpha - \beta$ is upgraded to $\gamma > (\alpha - \beta)/(1-\beta)$:

Theorem 1.5. Let $0 < \beta \leq \alpha < 1$ and $\kappa > 0$. Then, for every $\gamma \in ((\alpha - \beta)/(1 - \beta), 1]$, there exist $\epsilon, \epsilon_0, \delta_0 \in (0, \frac{1}{2}]$, depending only on $\alpha, \beta, \gamma, \kappa$, such that the following holds. Let $\delta \in 2^{-\mathbb{N}}$ with $\delta \in (0, \delta_0]$, and let $A, B \subset (\delta \cdot \mathbb{Z}) \cap [0, 1]$ satisfy the following hypotheses:

(A) $|A| \leq \delta^{-\alpha}$.

(B) $|B| \ge \delta^{-\beta}$, and B satisfies the following Frostman condition:

$$|B \cap B(x,r)| \leq r^{\kappa}|B|, \qquad \delta \leq r \leq \delta^{\epsilon_0}.$$

Further, let ν be a Borel probability measure with $\operatorname{spt}(\nu) \subset [\frac{1}{2}, 1]$, satisfying the Frostman condition $\nu(B(x, r)) \leq r^{\gamma}$ for $x \in \mathbb{R}$ and $\delta \leq r \leq \delta^{\epsilon_0}$. Then, there exists a point $c \in \operatorname{spt}(\nu)$ such that

$$|A + cB|_{\delta} \ge \delta^{-\epsilon}|A|.$$

Theorem 1.5 with $\alpha = \beta$ recovers Theorem 1.2. The fact that Theorem 1.5 does not impose any non-concentration conditions on *A* is useful in proving Corollary 1.7 below.

1.1. **Related work.** A relevant piece of recent literature is the paper of Guth, Katz, and Zahl [12], where the authors extend an argument (due to Garaev [10]) from finite fields to give a new, relatively simple, proof of Bourgain's Theorem 1.2. Given that the \mathbb{Z}_p analogue of Conjecture 1.4 is known [28], it seems remotely plausible that Conjecture 1.4 can be solved by extending the \mathbb{Z}_p argument in the fashion of Guth, Katz, and Zahl. I was not able to carry this out, and here is why. The proof in [28] is chiefly based on the following lemma: if $A, B \subset \mathbb{Z}_p$ are sets with $|A| = p^{\alpha}$ and $|B| = p^{\beta}$, then for every $\eta > 0$ there exists an integer $n = n(\alpha, \beta, \eta) \in \mathbb{N}$, and choices $a_1, \ldots, a_n \in \pm A$ such that

$$|a_1B + \ldots + a_nB| \gtrsim p^{-\eta} \min\{|A||B|, p\}.$$
(1.6)

I was not able to extend the finite field techniques in [28] to (directly) prove a δ -discretised analogue of (1.6). However, once Theorem 1.5 is known, it can be applied to make partial progress towards a δ -discretised analogue of (1.6) (a sharper result would follow from Conjecture 1.4 in the same way):

Corollary 1.7. Let $\beta, \gamma \in (0, 1)$ and $0 < \eta < \gamma(1 - \beta)$. Then, there exists $\epsilon_0, \delta_0 > 0$ and $n \in \mathbb{N}$, depending on β, γ, η , such that the following holds for all $\delta \in (0, \delta_0]$. Let $B, C \subset (\delta \cdot \mathbb{Z}) \cap [0, 1]$ be non-empty sets satisfying

$$|B \cap B(x,r)| \leq r^{\beta}|B| \quad and \quad |C \cap B(x,r)| \leq r^{\gamma}|C| \tag{1.8}$$

for $x \in \mathbb{R}$ and $\delta \leq r \leq \delta^{\epsilon_0}$. Then, there exist points $c_1, \ldots, c_n \in C$ such that

$$|c_1B + \ldots + c_nB|_{\delta} \ge \delta^{-\gamma - \beta(1-\gamma) + \eta} = \delta^{-\beta - \gamma(1-\beta) + \eta}$$

Remark 1.9. A classical projection theorem of Kaufman [19] would imply the existence of $c \in C$ such that $|B + cB|_{\delta} \gtrsim \max\{\delta^{-\beta}, \delta^{-\gamma}\}$. For $\gamma > \beta$, a recent sharpening of Kaufman's theorem by the author and Shmerkin [27] would even yield $|B + cB|_{\delta} \ge \delta^{-\gamma-\eta}$ for some $\eta = \eta(\beta, \gamma) > 0$, and for $\delta > 0$ small enough. In comparison, Corollary 1.7 gives a more quantitative improvement, but at the cost of adding the number of summands. I give the proof straight away: it is simple, and good preparation for Section 3.3, which contains a slightly more complicated version of the same argument.

Proof of Corollary **1**.7. Start by applying Theorem **1**.5 with

$$\alpha := \beta + \gamma(1 - \beta) - \eta \in (\beta, 1), \quad \beta, \quad \kappa := \beta, \quad \text{and} \quad \gamma$$

Note that $\gamma > (\alpha - \beta)/(1 - \beta)$, so the parameters are admissible. Let $\epsilon, \epsilon_0, \delta_0 \in (0, \frac{1}{2}]$ be the constants given by Theorem 1.5 with $\alpha, \beta, \gamma, \kappa$. Let $\nu := |C|^{-1} \cdot \mathcal{H}^0|_C$ be the normalised counting measure on C, which satisfies the Frostman condition $\nu(B(x, r)) \leq r^{\gamma}$ for all $x \in \mathbb{R}$ and $\delta \leq r \leq \delta^{\epsilon_0}$ by (1.8). We also note that $|B| \geq \delta^{-\beta}$ by (1.8) applied with $r = \delta$, and B satisfies the $\kappa = \beta$ -dimensional Frostman condition required in Theorem 1.5.

We construct a sequence of sets $H_n \subset \delta \cdot \mathbb{Z}$, $n \in \mathbb{N}$, with the following greedy algorithm. We first define $H_1 := (c_1 B)_{\delta}$ arbitrarily, where $A_{\delta} := (\delta \cdot \mathbb{Z}) \cap A(\delta)$. Then, we assume that H_n has already been defined for some $n \ge 1$, and we let

$$H_{n+1} := (H_n + c_{n+1}B)_{\delta} \subset \delta \cdot \mathbb{Z},$$

where $c_{n+1} \in C = \operatorname{spt}(\nu)$ maximises $|H_n + cB|_{\delta}$ among all choices $c \in C$. We observe (by induction) that $H_n \subset (\delta \cdot \mathbb{Z}) \cap [0, n]$, so $|H_n| \leq n\delta^{-1}$. For arbitrary $N \in \mathbb{N}$ with $N \geq 2$, it follows from the pigeonhole principle that there exists $n \in \{1, \ldots, N-1\}$ such that

$$H_{n+1} \leq (N\delta^{-1})^{1/(N-1)} |H_n| \leq 2\delta^{-1/(N-1)} |H_n|$$

By definition of H_{n+1} , this means that

$$|H_n + cB|_{\delta} \lesssim |H_{n+1}| \leqslant 2\delta^{-1/(N-1)}|H_n|, \qquad c \in C.$$
(1.10)

We now choose $N \in \mathbb{N}$ so large that $2\delta^{-1/(N-1)} \leq \delta^{-\epsilon/2}$, where $\epsilon = \epsilon(\alpha, \beta, \gamma) > 0$ was one of the constants produced by Theorem 1.5. Since *B* and ν satisfy the hypotheses of Theorem 1.5, we see from (1.10) that $A := H_n$ must fail the hypotheses. However, the only hypotheses on *A* in Theorem 1.5 are

$$A \subset (\delta \cdot \mathbb{Z}) \cap [0,1]$$
 and $|A| \leq \delta^{-\alpha}$.

Of course $H_n \notin [0,1]$, but this is not really relevant: we may find $k \in \{0, \ldots, n-1\}$ such that $|H_n \cap [k, k+1]| \ge \frac{1}{N} |H_n|$. Now, defining instead $A := (H_n \cap [k, k+1]) - \{k\}$, we have $A \subset (\delta \cdot \mathbb{Z}) \cap [0,1]$, and $|A + cB|_{\delta} \le \delta^{-\epsilon/2} |H_n| \le \delta^{-\epsilon} |A|$ by (1.10), for $\delta > 0$ so small that $\delta^{-\epsilon/2} \ge N$. This violates violate Theorem 1.5, unless

$$|H_n| \ge |A| > \delta^{-\alpha} = \delta^{-\beta - \gamma(1-\beta) + \eta}$$

and this is what the corollary claimed.

The *ABC* problem is closely related to the highly active area of sum-product theory. The main open question is the Erdős-Szemerédi sum-product conjecture [8]: if $A \subset \mathbb{R}$ or $A \subset \mathbb{Z}_p$ is a finite set ($p \in \mathbb{N}$ is prime), the E-S conjecture asks to prove that

$$\max\{|A+A|, |A\cdot A|\} \gtrsim_{\epsilon} |A|^{2-\epsilon}, \qquad \epsilon > 0.$$

The research around this problem is too active to survey here: I only mention the papers [30] of Rudnev-Stevens and [23] of Mohammadi-Stevens for some current world records, and further references. For results on the the δ -discretised variant of the Erdős-Szemerédi problem, see [12] by Guth-Katz-Zahl, and [7] by Dąbrowski, the author, and Villa.

Bourgain's δ -discretised sum-product estimate, Theorem 1.2, has been extended in various ways beyond the real line. For example, He [16] found a version of the theorem in \mathbb{R}^n . Closely related are also the works [3, 6] by Bourgain-Gamburd, [15] by He, [17] by He-de Saxcé, [1] by Benoist-de Saxcé, and [20] by Li. These papers contain δ -discretised sum-product or product theorems in various Lie groups. Viewing the δ -discretised *ABC* sum-product problem as a special case of a δ -discretised incidence problem between points and δ -tubes in \mathbb{R}^2 , the papers [7, 9, 13] are also relevant.

Theorems 1.2 and 1.5 can be viewed as statements concerning linear projections of planar sets (see the next subsection). Starting with this interpretation, one may ask if analogous statements hold for non-linear projections. Examples of particular interest are the *pinned distance projections* $\Delta_x(y) = |x - y|$ and the *radial projections* $\pi_x(y) = (x - y)/|x - y|$. Again, the literature is too broad for a survey, but see the recent papers [34] by Shmerkin and [29] by Raz-Zahl for recent exciting developments and more references.

Finally, Conjecture 1.4 was recently solved by the author [25] for Ahlfors-regular sets $A, B \subset [0, 1]$. In fact, a much stronger result can be obtained for such sets. Let $\alpha, \beta \in (0, 1)$. Assume that $A, B \subset \mathbb{R}$ are closed sets, A is α -Ahlfors-regular and B is β -Ahlfors-regular. Then

$$\dim_{\mathrm{H}} \{ c \in \mathbb{R} : \dim(A + cB) < \alpha + \eta \} = 0$$

for $\eta := \beta(1-\alpha)/(2-\alpha) > 0$. (The paper [25] also contains a δ -discretised version.)

1.2. **Proof sketch.** For this exposition, I make the following additional assumptions on A and B. Both sets have a "tree" (or "Cantor set") structure: for a suitable parameter $m \in \mathbb{N}$, each dyadic interval $I \in \mathcal{D}_{ms}$ intersecting A contains exactly $R_A(s)$ sub-intervals in $\mathcal{D}_{m(s+1)}$ which intersect A. The same is assumed of B. The numbers $R_A(s)$ and $R_B(s)$ are known as the *branching numbers* of A and B, respectively. Assume that the scale parameter $\delta > 0$ has the special form $\delta = 2^{-mN}$ for some $N \in \mathbb{N}$ (thus $R_A(s) = 1 = R_B(s)$ for $s \ge N$, since A, B were assumed to be δ -separated). We make even more assumptions:

(P1) For every $s \in \mathbb{N}$, either $R_B(s) = 1$ or $R_A(s) = 2^m$.

(P2) $|B| = \delta^{-\beta}$, and for every $s \in \mathbb{N}$, either $R_B(s) = 1$ or $R_B(s) = 2^m$.

The proof of Theorem 1.5 will begin by several reductions, which allow us to make (approximate versions of) all the assumptions listed above. Now, the key question turns out to be: given a scale $s \in \mathbb{N}$ with $R_B(s) = 1$, what **upper** bound can we guarantee for $R_A(s)$? It turns out that we can easily use (P1)-(P2) to deduce an answer.

Assume that $R_A(s) \ge 2^{\Gamma m}$ for all $s \in \{0, ..., N-1\} =: [N]$ with $R_B(s) = 1$. Write $\mathcal{N} := \{s \in [N] : R_B(s) = 1\}$, and note that $R_A(s) = 2^m$ for all $s \in [N] \setminus \mathcal{N}$ by assumption

(P1). Now, we may calculate a lower bound on the cardinality of *A* as follows:

$$2^{\alpha mN} \ge |A| = \prod_{s \in [N]} R_A(s) = \prod_{s \in [N] \setminus \mathcal{N}} R_A(s) \cdot \prod_{s \in \mathcal{N}} R_A(s) \ge 2^{m(N-|\mathcal{N}|)} \cdot 2^{\Gamma m|\mathcal{N}|}.$$
 (1.11)

On the other hand, by assumption (P2), we have

$$2^{\beta mN} = |B| = \prod_{s \in [N] \setminus \mathcal{N}} 2^m = 2^{m(N - |\mathcal{N}|)},$$
(1.12)

so may solve $N - |\mathcal{N}| = \beta N$ and $|\mathcal{N}| = (1 - \beta)N$. Plugging this information into (1.11) yields $\Gamma \leq (\alpha - \beta)/(1 - \beta)$. This is where the numerology in Theorem 1.5 comes from. Namely, the argument above shows that if $\Gamma > (\alpha - \beta)/(1 - \beta)$, then there exists at least one scale $s \in [N]$ such that $R_A(s) \leq 2^{\Gamma m}$. In fact, the same must be true for a positive fraction of the scales, say $\mathcal{G} \subset [N]$, where $|\mathcal{G}|/N$ only depends on $\Gamma - (\alpha - \beta)/(1 - \beta)$.

After this observation, we focus attention separately on pieces of $A \times B$ of the form $(A \cap I) \times (B \cap J)$, where $I, J \in \mathcal{D}_{ms}$ are intervals intersecting A, B, respectively, and $s \in \mathcal{G}$. By definition, $|A \cap I|_{m(s+1)} = R_A(s) \leq 2^{\Gamma m}$ for some Γ slightly larger than $(\alpha - \beta)/(1 - \beta)$. To be precise, we choose $(\alpha - \beta)/(1 - \beta) < \Gamma < \gamma$, where γ is the Frostman exponent of the measure ν in Theorem 1.5. If we **additionally** knew that $|B \cap J|_{2^{-m(s+1)}} = R_B(s) \geq 2^{\epsilon m}$ for some $\epsilon > 0$, and the points in $B \cap J$ are well enough separated, we could at this point use an elementary argument (essentially the "potential theoretic method" due to Kaufman [19]) to deduce that

$$|(A \cap I) + c(B \cap J)|_{2^{-m(s+1)}} \ge 2^{\epsilon m} |A \cap I|_{2^{-m(s+1)}}$$
(1.13)

for a generic choice $c \in C = \operatorname{spt}(\nu)$. This argument is crucially based on $\gamma > \Gamma$, see Lemma 4.43 for the details. After this, summing up the increments (1.13) for all $s \in \mathcal{G}$ would complete the proof of Theorem 1.5.

An major problem is that, as a matter of fact, $|B \cap J|_{2^{-m(s+1)}} = R_B(s) = 1$ for all $s \in \mathcal{G}$. This follows from our assumption (P1), since $R_A(s) \leq 2^{\Gamma m} < 2^m$ for all $s \in \mathcal{G}$. To solve the problem, we follow Bourgain's proof of Theorem 1.2 faithfully: instead of considering individual scales $s \in [N]$, we recombine consecutive elements of [N] into longer intervals $\mathcal{I} \subset [N]$ where the branching of B is small but non-trivial, say $R_B(\mathcal{I}) = 2^{\epsilon m |\mathcal{I}|}$. Then, we carry out calculations similar to the ones we saw at (1.11)-(1.12) to make sure that also $R_A(\mathcal{I}) \leq 2^{\Gamma m |\mathcal{I}|}$ for a subset of these intervals \mathcal{I} with substantial total length. At the end of the day, the intervals \mathcal{I} with $R_B(\mathcal{I}) = 2^{\epsilon m |\mathcal{I}|}$ and $R_A(\mathcal{I}) \leq 2^{\Gamma m |\mathcal{I}|}$, will actually play the role we had written for the scales \mathcal{G} in the discussion above.

There are numerous places in the arguments below where I either follow Bourgain's argument for the case A = B, or at least draw heavy inspiration from such an argument. Bourgain's influence on this paper will be treated as an absolute constant, and not spelled out every time separately.

1.3. Acknowledgements. I would like to thank Pablo Shmerkin for clarifying a point about applying his inverse theorem [32, Theorem 2.1], see Remark 4.7.

2. NOTATION AND PRELIMINARIES

2.1. Dyadic cubes and covering numbers. Let \mathcal{D}_n be the family of dyadic cubes in \mathbb{R}^d with side-length 2^{-n} . We will only use this notation for $n \ge 0$. For $Q = x + [0, 2^{-n})^d \in \mathcal{D}_n$, we associate the affine map $T_Q(y) := 2^n(y - x)$, which rescales Q to $[0, 1)^d$.

If μ is a Borel measure on \mathbb{R}^d , and $E \subset \mathbb{R}^d$ is a Borel set with $\mu(E) > 0$, we write $\mu_E := \mu(E)^{-1}\mu|_E$ for the *renormalised restriction of* μ to E. This notation is most commonly used in the case $E = Q \in \mathcal{D}_n$. In this special case, we additionally define the notation

$$\mu^Q := T_Q \mu_Q.$$

Here $f\nu(H) := \nu(f^{-1}H)$ refers, in general, to the push-forward of a measure ν under a map f. For a dyadic rational $r = 2^{-n}$, and a bounded set $A \subset \mathbb{R}^d$, we write $|A|_r$ for the least number of cubes in \mathcal{D}_n required to cover A (in the introduction, we used the same notation for the *r*-covering number, which is comparable up to a multiplicative constant). We will also write A(r) for the open *r*-neighbourhood of A, and (only with $r = \delta$),

$$A_r = (r \cdot \mathbb{Z}) \cap A(r)$$

Finally, for $n \in \mathbb{N}$, we abbreviate $[n] := \{0, \dots, n-1\}$.

2.2. **Entropy.** If (Ω, μ) is a probability space, and \mathcal{F} is a countable μ -measurable partition of Ω , we denote the \mathcal{F} -entropy of μ by

$$H(\mu, \mathcal{F}) := \sum_{F \in \mathcal{F}} \mu(F) \log \frac{1}{\mu(F)},$$

with the convention $0 \cdot \log 0 = 0$. If \mathcal{E}, \mathcal{F} are two countable partitions, we denote the *conditional* \mathcal{F} -entropy of μ relative to \mathcal{E} by

$$H(\mu, \mathcal{F} \mid \mathcal{E}) := \sum_{E \in \mathcal{E}} \mu(E) H(\mu_E, \mathcal{F}).$$
(2.1)

If \mathcal{F} refines \mathcal{E} (each element of \mathcal{E} can be written as a disjoint union of elements of \mathcal{F}), the conditional entropy can be alternatively written as

$$H(\mu, \mathcal{F} \mid \mathcal{E}) = H(\mu, \mathcal{F}) - H(\mu, \mathcal{E}).$$
(2.2)

For a proof, see [26, Proposition 3.3]. In practice, we will only be concerned with \mathcal{D}_n entropies of compactly supported Borel probability measures on \mathbb{R}^d , where \mathcal{D}_n is the partition of \mathbb{R}^d into dyadic cubes of side-length 2^{-n} . In this special case \mathcal{D}_{n+1} always refines \mathcal{D}_n , so the formula (2.2) is available. We record the following simple lemma, whose proof is a combination of [26, Lemma 3.5] and [26, Remark 3.6]:

Lemma 2.3. Let μ be a Borel probability measure on \mathbb{R}^d , and let $\pi \colon \mathbb{R}^d \to \mathbb{R}^D$ be linear. Let $n \in \mathbb{N}$, and let $0 = n_0 < n_1 < \ldots < n_h = n$ be an arbitrary partition of $\{0, \ldots, n\}$. Then,

$$H(\pi\mu, \mathcal{D}_n) \ge \sum_{j=0}^{h-1} \sum_{Q \in \mathcal{D}_{n_j}} \mu(Q) \cdot H(\pi\mu^Q, \mathcal{D}_{n_{j+1}-n_j} \mid \mathcal{D}_0).$$

The inner summation only runs over those $Q \in \mathcal{D}_{n_i}$ with $\mu(Q) > 0$.

A basic fact about entropy (which follows from Jensen's inequality) is that

$$|\{F \in \mathcal{F} : \mu(F) > 0\}| \leq N \quad \Longrightarrow \quad H(\mu, \mathcal{F}) \leq \log N.$$

In particular, if $\pi : \mathbb{R}^d \to \mathbb{R}^D$ is *L*-Lipschitz in Lemma 2.3, then

$$H(\pi\mu^{Q}, \mathcal{D}_{n_{j+1}-n_{j}} \mid \mathcal{D}_{0}) = H(\pi\mu^{Q}, \mathcal{D}_{n_{j+1}-n_{j}}) - H(\pi\mu^{Q}, \mathcal{D}_{0}) \ge H(\pi\mu^{Q}, \mathcal{D}_{n_{j+1}-n_{j}}) - CL,$$

where $C \ge 1$ only depends on d, D. Therefore, the lower bound of Lemma 2.3 can be upgraded to

$$H(\pi\mu, \mathcal{D}_n) \ge \left(\sum_{j=0}^{h-1} \sum_{Q \in \mathcal{D}_{n_j}} \mu(Q) \cdot H(\pi\mu^Q, \mathcal{D}_{n_{j+1}-n_j})\right) - h \cdot CL.$$
(2.4)

We mention two further useful fact about entropy: first, if \mathcal{E}, \mathcal{F} are two countable μ -measurable partitions such that

$$|\{F \in \mathcal{F} : F \cap E_0\}| \leqslant N \quad \text{and} \quad |\{E \in \mathcal{E} : E \cap F_0\}| \leqslant N$$

for all $E_0 \in \mathcal{E}$ and $F_0 \in \mathcal{F}$, then $|H(\mu, \mathcal{E}) - H(\mu, \mathcal{F})| \leq \log N$. Second, entropy (and also conditional entropy) is concave. We will use the convexity of entropy in the following form: if μ, ν are two Borel probability measures on \mathbb{R}^d , then

$$H(\mu * \nu, \mathcal{D}_n) \ge \int H(\mu_x, \mathcal{D}_n) \, d\nu(x), \tag{2.5}$$

where μ_x is the probability measure defined by $\mu_x(H) = \mu(H - x)$. Since $(\mu * \nu)(H) = \int \mu_x(H) d\nu(x)$ for all Borel sets $H \subset \mathbb{R}^d$, one may view $\mu * \nu$ as a convex combination of the measures μ_x . Formally, (2.5) is deduced by applying Jensen's inequality to the concave function $f(r) = r \log(1/r)$ on [0, 1], and the random variable $X : x \mapsto \mu_x(Q)$ in the probability space (\mathbb{R}^d, ν) (for fixed $Q \in \mathcal{D}_n$).

3. THREE INITIAL REDUCTIONS

This section contains a reduction of Theorem 1.5 to a special case, where we additionally assume that $|B + B| \leq \delta^{-\epsilon_B}|B|$, and $\nu(B(x,r)) \leq 40 \cdot r^{\gamma}$ for all $x \in \mathbb{R}$ and $r \geq \delta$ (see Theorem 3.25). It seems difficult to do achieve this reduction in a "single pass": instead, we add the extra assumptions in two separate steps (Sections 3.1 and 3.2). After these steps, we arrive at Theorem 3.12, where the assumptions are present, but unfortunately the conclusion is also a little stronger. Then, the final reduction to Theorem 3.25 "restores" the weaker conclusion, but maintains the additional assumptions. This is the version of Theorem 1.5 we will eventually be able to prove directly.

3.1. Reduction to the case where *B* has small doubling. The purpose of this section is to reduce the proof of Theorem 1.5 to the following version, where the hypothesis $|B + B| \leq \delta^{-\epsilon_B}$ has been added. This does not come for free: the price to pay is that the conclusion of Theorem 3.1 is also a little stronger (that is, more difficult to prove).

Theorem 3.1. Let $0 < \beta \leq \alpha < 1$ and $\kappa > 0$. Then, for every $\gamma \in ((\alpha - \beta)/(1 - \beta), 1]$, there exist $\epsilon_0, \epsilon, \epsilon_B, \delta_0, \rho \in (0, \frac{1}{2}]$, depending only on $\alpha, \beta, \gamma, \kappa$, such that the following holds. Let $\delta \in 2^{-\mathbb{N}}$ with $\delta \in (0, \delta_0]$, and let $A, B \subset (\delta \cdot \mathbb{Z}) \cap [0, 1]$ satisfy the following hypotheses:

(A)
$$|A| \leq \delta^{-\alpha}$$
.

(B) $|B| \ge \delta^{-\beta}$, and B satisfies the following Frostman condition:

$$|B \cap B(x,r)| \leq r^{\kappa}|B|, \qquad \delta \leq r \leq \delta^{\epsilon_0}.$$

Assume moreover that $|B + B| \leq \delta^{-\epsilon_B} |B|$.

Further, let ν be a Borel probability measure with $\operatorname{spt}(\nu) \subset [0, 1]$, and satisfying the Frostman condition $\nu(B(x, r)) \leq r^{\gamma}$ for $x \in \mathbb{R}$ and $\delta \leq r \leq \delta^{\epsilon_0}$. Then, there exists a point $c \in \operatorname{spt}(\nu)$ such that $|A' + cB|_{\delta} \geq \delta^{-\epsilon}|A|$ for all $A' \subset A$ with $|A'| \geq (1 - \rho)|A|$.

The proof will use the following version of the Plünnecke-Ruzsa inequality:

Lemma 3.2 (Plünnecke-Ruzsa inequality). Let $\delta \in 2^{-\mathbb{N}}$, let $A, B_1, \ldots, B_n \subset \mathbb{R}$ be arbitrary sets, and assume that $|A + B_i|_{\delta} \leq K_i |A|_{\delta}$ for all $1 \leq i \leq n$, and for some constants $K_i \geq 1$. Then, for every $\epsilon > 0$, there exists a subset $A' \subset A$ with $|A'|_{\delta} \geq (1 - \epsilon)|A|_{\delta}$ such that

$$|A' + B_1 + \ldots + B_n|_{\delta} \lesssim_{\epsilon,n} K_1 \cdots K_n |A'|_{\delta}.$$

This form of the inequality is due to Ruzsa [31]. For a more general result, see [14, Theorem 1.5], by Gyarmati-Matolcsi-Ruzsa. To be accurate, these statements are not formulated in terms of δ -covering numbers, but one may consult [12, Corollary 3.4] by Guth-Katz-Zahl to see how to handle the reduction.

Proof of Theorem **1.5** *assuming Theorem* **3.1**. Let $\alpha, \beta, \gamma, \kappa$ be the constants given in Theorem **1.5**. Our task is to find $\epsilon, \epsilon_0, \delta_0$ depending on $\alpha, \beta, \gamma, \kappa$, such that the claims of Theorem **3.1** are satisfied. To do this, we fix some $\overline{\beta} < \beta$ slightly smaller than β so that still

$$\gamma > (\alpha - \bar{\beta})/(1 - \bar{\beta})$$

Then, we apply Theorem 3.1 with the parameters $\alpha, \overline{\beta}, \gamma, \kappa/4$, and first extract the constants $\overline{\epsilon}, \overline{\epsilon}_0, \overline{\epsilon}_B, \overline{\delta}_0, \overline{\rho} > 0$, depending only on $\alpha, \overline{\beta}, \gamma, \kappa/4$. Now, we claim that Theorem 1.5 holds with constants

$$\epsilon_0 := \frac{2}{\kappa n} \quad \text{and} \quad \epsilon := 2^{-n-1} \bar{\epsilon},$$
(3.3)

where

$$n := \max\{[2/(\kappa\bar{\epsilon}_0)], [2/\bar{\epsilon}_B]\}$$
(3.4)

and any $\delta_0 \in (0, \overline{\delta}_0]$ with the additional requirements

$$2^{n+1} \leqslant \delta_0^{-\bar{\epsilon}_B/2} \quad \text{and} \quad \delta_0^{\beta-\beta} \geqslant 2^n.$$
(3.5)

At the very end of the proof, there will be an additional requirement for $\delta_0 > 0$, which only depends on $n, \bar{\rho}$ (hence eventually) $\alpha, \beta, \gamma, \kappa$; I will not spell this out explicitly, since the bounds depend on the implicit constants in Lemma 3.2.

To prove Theorem 1.5, fix $\delta \in (0, \delta_0]$, and assume that $A, B \subset (\delta \cdot \mathbb{Z}) \cap [0, 1]$ and ν satisfy the assumptions of Theorem 1.5 with parameters $\alpha, \beta, \gamma, \kappa, \epsilon_0$, as specified in and below (3.3). Thus $|A| \leq \delta^{-\alpha}$, and $|B| \geq \delta^{-\beta}$, and $|B \cap B(x, r)| \leq r^{\gamma}$ for all $x \in \mathbb{R}$ and $\delta \leq r \leq \delta^{\epsilon_0}$. We claim that there exists $c \in \operatorname{spt}(\nu)$ such that $|A + cB|_{\delta} \geq \delta^{-\epsilon}|A|$.

Write *kB* for the *k*-fold sum $B + \ldots + B$. Clearly $kB \subset (\delta \cdot \mathbb{Z}) \cap [0, k]$, so

$$|kB| \leqslant k \cdot \delta^{-1}, \qquad k \geqslant 1. \tag{3.6}$$

We claim that for any $n \ge 1$, there exists $1 \le k \le n$ such that

$$|2^{k}B + 2^{k}B| = |2^{k+1}B| \leq 2\delta^{-1/n} |2^{k}B|.$$
(3.7)

Indeed, if this were not the case, then

$$2^{n}\delta^{-1} \stackrel{(3,6)}{\geqslant} |2^{n}B| > 2\delta^{-1/n}|2^{n-1}B| > 4\delta^{-2/n}|2^{n-2}B| > \ldots > 2^{n}\delta^{-n/n}|B| \ge 2^{n}\delta^{-1},$$

a contradiction. We then apply this observation with $n \in \mathbb{N}$ as in (3.4), and we pick $1 \leq k \leq n$ such that (3.7) holds. Write $\overline{B} := 2^k B$ for this choice of "k", so

$$|\bar{B} + \bar{B}| \le 2\delta^{-1/n} |\bar{B}|. \tag{3.8}$$

Recalling that $1/n \leq \bar{\epsilon}_B/2$, this looks promising for the purpose of applying Theorem 3.1 to the pair of sets *A* and \bar{B} . But does \bar{B} satisfy a Frostman condition? It turns out that it

does, with constants " $\kappa/4$ " and " $\overline{\epsilon}_0$ ". The following argument is copied from [5, Section 8.2]. Assume, to reach a contradiction, that there exists some dyadic scale

$$r \in [\delta, \delta^{\overline{\epsilon}_0}] \stackrel{(3.4)}{\subset} [\delta, \delta^{2/(\kappa n)}] \stackrel{(3.3)}{=} [0, \delta^{\epsilon_0}], \tag{3.9}$$

and a point $a \in \mathbb{R}$, such that $|\bar{B} \cap B(a,r)|_{\delta} \ge C_0 r^{\kappa/2} |\bar{B}|$ for a suitable absolute constant $C_0 \ge 1$. Now, since $r \le \delta^{\epsilon_0}$, we know by hypothesis that B satisfies $|B \cap B(x,r)| \le r^{\kappa}$ for all $x \in \mathbb{R}$. Consequently $|\bar{B}|_r \ge |B|_r \ge r^{-\kappa}$, and

$$\begin{split} |\bar{B} + \bar{B}|_{\delta} \gtrsim |\bar{B}|_{r} \cdot |\bar{B} \cap B(a, r)|_{\delta} \\ \geqslant C_{0}|B|_{r} \cdot r^{\kappa/2}|\bar{B}| \\ \geqslant C_{0}r^{-\kappa} \cdot r^{\kappa/2}|\bar{B}| \\ \geqslant C_{0}\delta^{-1/n}|\bar{B}|, \end{split}$$

using $r \leq \delta^{\tilde{\epsilon}_0} \leq \delta^{2/(\kappa n)}$ in the final inequality. If $C_0 \geq 1$ was chosen large enough, this lower bound contradicts the small doubling property (3.8). We conclude that

$$|\bar{B} \cap B(x,r)| \le C_0 r^{\kappa/2} |\bar{B}|, \qquad \delta \le r \le \delta^{\bar{\epsilon}_0}.$$
(3.10)

There are also a few smaller issues before we can apply Theorem 3.1 to B: evidently $|\bar{B}| \ge |B| \ge \delta^{-\beta}$, but unfortunately $\bar{B} \subset [0, 2^n]$ instead of $\bar{B} \subset [0, 1]$. Regardless, there exists some an interval $I_0 = [m, m+1] \subset [0, 2^n]$ such that $|\bar{B} \cap I_0| \ge |\bar{B}|/2^n$. We define

$$\bar{B}_m := (\bar{B} \cap I_0) - \{m\} \subset (\delta \cdot \mathbb{Z}) \cap [0, 1],$$

so $|\bar{B}_m| \ge |\bar{B}|/2^n$. Then,

$$|\bar{B}_m + \bar{B}_m| \stackrel{(\mathbf{3.8})}{\leqslant} 2\delta^{1/n} |\bar{B}| \stackrel{(\mathbf{3.4})-(\mathbf{3.5})}{\leqslant} \delta^{-\bar{\epsilon}_B} |\bar{B}_m| \quad \text{and} \quad |\bar{B}_m| \ge 2^{-n} \delta^{-\beta} \stackrel{(\mathbf{3.5})}{\geqslant} \delta^{-\bar{\beta}}.$$

Finally,

$$|\bar{B}_m \cap B(x,r)| \stackrel{(3.10)}{\leqslant} C_0 2^n r^{\kappa/2} |\bar{B}_m| \stackrel{(3.5)}{\leqslant} r^{\kappa/4} |\bar{B}_m|, \quad r \in [\delta, \delta^{\bar{\epsilon}_0}].$$

Now we have shown that the triple A, \overline{B}_m, ν satisfies all the hypotheses of Theorem 3.1 with parameters $\alpha, \overline{\beta}, \gamma, \kappa/4$, and $\overline{\epsilon}_0, \overline{\epsilon}_B, \overline{\delta}_0$. It follows that there exists $c \in \operatorname{spt}(\nu)$ with the property that if $A' \subset A$ is any subset with $|A'| \ge (1 - \overline{\rho})|A|$, then

$$|A' + c\bar{B}_m|_{\delta} \ge \delta^{-\bar{\epsilon}}|A|. \tag{3.11}$$

We now claim that $|A + cB| \ge \delta^{-\epsilon} |A|$ for this specific $c \in \operatorname{spt}(\nu)$, which will complete the proof of Theorem 1.5. If this fails, then by the Plünnecke-Ruzsa inequality, Lemma 3.2, we find a subset $A' \subset A$ of cardinality $|A'| \ge (1 - \bar{\rho})|A|$ such that

$$|A' + c\bar{B}_m|_{\delta} \lesssim |A' + c(2^k B)|_{\delta} \lesssim_{n,\bar{\rho}} \delta^{-2^n \epsilon} |A| \stackrel{(3.3)}{=} \delta^{-\bar{\epsilon}/2} |A|.$$

This contradicts (3.11) for $\delta > 0$ small enough, depending on $n, \bar{\rho}$. This contradiction completes the proof of Theorem 1.5.

3.2. Reducing the Frostman constant of ν . Let ν be the measure appearing in the statement of Theorem 1.5 or 3.1. We assumed that $\nu(B(x,r)) \leq r^{\gamma}$ for all scales $r \leq \delta^{\epsilon_0}$. We will need, in fact, is that ν satisfies the Frostman condition $\nu(B(x,r)) \leq Cr^{\gamma}$ for all $r \geq \delta$, and with an absolute constant $C \geq 1$. It turns out that this can be achieved, eventually with C = 40. In this section, we reduce the proof of Theorem 3.1 to the following:

Theorem 3.12. Let $0 < \beta \leq \alpha < 1$ and $\kappa > 0$. Then, for every $\gamma \in ((\alpha - \beta)/(1 - \beta), 1]$, there exist $\epsilon, \epsilon_0, \epsilon_B, \delta_0, \rho \in (0, \frac{1}{2}]$, depending only on $\alpha, \beta, \gamma, \kappa$, such that the following holds. Let $\delta \in 2^{-\mathbb{N}}$ with $\delta \in (0, \delta_0]$, and let $A, B \subset (\delta \cdot \mathbb{Z}) \cap [0, 1]$ satisfy the following hypotheses:

- (A) $|A| \leq \delta^{-\alpha}$.
- (B) $|B| \ge \delta^{-\beta}$, and B satisfies the following Frostman condition:

$$|B \cap B(x,r)| \leqslant r^{\kappa}|B|, \qquad \delta \leqslant r \leqslant \delta^{\epsilon_0}.$$

Assume moreover that $|B + B| \leq \delta^{-\epsilon_B} |B|$.

Further, let ν be a Borel probability measure with $\operatorname{spt}(\nu) \subset [-1, 1]$, and ν satisfies the Frostman condition $\nu(B(x, r)) \leq 20r^{\gamma}$ for $x \in \mathbb{R}$ and $r \geq \delta$. Then, there exists a point $c \in \operatorname{spt}(\nu)$ such that

$$|A' + cB|_{\delta} \ge \delta^{-\epsilon}|A'|$$

for all subsets $A' \subset A$ with $|A'| \ge (1 - \rho)|A|$.

The proof will require the Plünnecke-Ruzsa inequality, that is Lemma 3.2 in the previous section, and also the following [37, Exercise 6.5.12] in the book of Tao and Vu:

Lemma 3.13. Let $A, B \subset \delta \cdot \mathbb{Z}$, and assume that $|A + B| \leq K|A|$ for some $K \geq 1$. Then, for every $N \geq 1$ and $\rho > 0$, there exists a subset $A' \subset A$ with $|A'| \geq (1 - \rho)|A|$ with the property $|A' - B| \leq_{\rho,n} K^{2^N/N} |A|^{1+1/N}$.

The exercise is only stated with constant $\rho = \frac{1}{2}$, but if one reads the subsequent hint about how to solve the exercise, it is clear (based on [37, Exercise 6.5.1]) that any $\rho > 0$ will work, at the cost of making the implicit constant larger. We are then prepared to reduce Theorem 3.1 to Theorem 3.12.

Proof of Theorem 3.1 *assuming Theorem* 3.12. The argument roughly follows [5, Section 5] in Bourgain's paper. Recall the meaning of the constants $\alpha, \beta, \gamma, \kappa$ from the statement of Theorem 3.1. Our task is to find the constants $\epsilon_0, \epsilon, \epsilon_B, \delta_0, \rho \in (0, \frac{1}{2}]$, depending only on $\alpha, \beta, \gamma, \kappa$. This will eventually be accomplished by applying Theorem 3.12 with constants $\alpha, \beta, \gamma, \kappa/2$. Recall from the statement of Theorem 3.12 that there exist constants $\bar{\epsilon}_0, \bar{\epsilon}, \bar{\epsilon}_B, \bar{\delta}_0, \bar{\rho} > 0$, which only depend on the constants $\alpha, \beta, \gamma, \kappa/2$. We begin by choosing $\epsilon > 0$ so small that

$$\frac{C}{\log_2(1/\epsilon)} \leqslant \bar{\epsilon} \quad \Longleftrightarrow \quad \epsilon \leqslant 2^{-C/\bar{\epsilon}} \tag{3.14}$$

for a suitable absolute constant C > 0 to be determined later. We now pick the other constants $\epsilon_0, \epsilon_B \in (0, \frac{1}{2}]$ so that

$$\epsilon_0 + \epsilon_B := \min\{\kappa \bar{\epsilon}_0/8, \bar{\epsilon}_B/2\}. \tag{3.15}$$

We choose $\delta_0 \leq \delta_0$, and additionally δ_0 needs to satisfy a few other restrictions, which we explain on the spot. We finally choose $\rho > 0$ so small that

$$(1-\rho)(1-\rho-\sqrt{\rho}) \ge (1-\bar{\rho}).$$
 (3.16)

With these choices of constants, fix $\delta \in (0, \delta_0]$, and let $A, B \subset (\delta \cdot \mathbb{Z}) \cap [0, 1]$ be sets, and let ν be a Borel probability measure on [0, 1], satisfying the hypotheses in Theorem 3.1. To land in a situation where Theorem 3.12 becomes applicable, we consider, instead, the measure $\bar{\nu} := \nu * (-\nu)$, where $-\nu(A) := \nu(-A)$. Evidently spt $\bar{\nu} \subset [-1, 1]$. As Bourgain shows in [5, (5.5)], the measure $\bar{\nu}$ has the property

$$\bar{\nu}(B(x,r)) \leqslant 4 \cdot \bar{\nu}(B(0,r)) \leqslant 4 \cdot \sup_{y \in \mathbb{R}} \nu(B(y,r)), \qquad x \in \mathbb{R}, \, r > 0.$$
(3.17)

Now, let $c_0 \ge 0$ be the infimum of the numbers such that

$$\bar{\nu}(B(0,c_0)) > 5 \cdot c_0^{\gamma},$$
(3.18)

if any such numbers exist. Evidently $c_0 \in [0, 1]$, since $\bar{\nu}$ is a probability measure on B(0, 1). If no c_0 as in (3.18) exists, then let $c_0 := \max\{|r| : r \in \operatorname{spt} \bar{\nu}\}$, and note that

$$5 \cdot c_0^{\gamma} \ge \bar{\nu}(B(0, c_0)) \ge \bar{\nu}(B(0, 1)) = 1 \implies c_0 \ge 5^{-1/\gamma} \ge \delta^{\epsilon_0}, \tag{3.19}$$

assuming here that $\delta_0 \ge \delta$ is sufficiently small in terms of γ , ϵ_0 . Assume then that c_0 , as in (3.18), exists. Since $\sup_{y \in \mathbb{R}} \nu(B(y, r)) \le r^{\gamma}$ for all $0 < r \le \delta^{\epsilon_0}$ by assumption, (3.17) implies that $c_0 \ge \delta^{\epsilon_0}$. In both cases, $c_0 \ge \delta^{\epsilon_0}$. Moreover, we note that $\operatorname{spt}(\nu) \cap \{c_0, -c_0\} \ne \emptyset$ in both cases (in the non-trivial case, otherwise some smaller value of c_0 would also satisfy (3.18)).

We then consider the re-normalised measure $\bar{\nu}^{c_0}$ defined by

$$\bar{\nu}^{c_0}(H) := \frac{1}{\bar{\nu}(B(0,c_0))} \cdot \bar{\nu}|_{B(0,c_0)}(c_0 \cdot H), \qquad H \subset \mathbb{R},$$

which satisfies $\operatorname{spt}(\bar{\nu}^{c_0}) = c_0^{-1} \cdot (\operatorname{spt} \bar{\nu} \cap \bar{B}(0, c_0)) \subset [-1, 1]$. Clearly $\bar{\nu}^{c_0}$ is a Borel probability measure. Moreover, if $x \in \mathbb{R}$ and $r \in [\delta, 1]$, then, assuming that $c_0 \in [0, 1]$ was defined via (3.18), we have

$$\bar{\nu}^{c_0}(B(x,r)) \leqslant \frac{\bar{\nu}(B(c_0x,c_0r))}{\bar{\nu}(B(0,c_0))} \overset{(3.17)}{\leqslant} 4 \cdot \frac{\bar{\nu}(B(0,c_0r))}{5 \cdot c_0^{\gamma}} \leqslant 4 \cdot \frac{5 \cdot (c_0r)^{\gamma}}{5 \cdot c_0^{\gamma}} = 4 \cdot r^{\gamma}.$$

If c_0 was, instead, defined as $c_0 = \max\{|r| : r \in \operatorname{spt} \bar{\nu}\}$, then $\bar{\nu}(B(0, c_0)) = 1$, so

$$\bar{\nu}^{c_0}(B(x,r)) \leqslant \frac{\bar{\nu}(B(c_0x,c_0r))}{\bar{\nu}(B(0,c_0))} \stackrel{(3.17)}{\leqslant} 4 \cdot \bar{\nu}(B(0,c_0r)) \leqslant 20 \cdot (c_0r)^{\gamma} \leqslant 20 \cdot r^{\gamma}.$$

The same estimates are also true for r > 1, since $\|\bar{\nu}^{c_0}\| = 1$. Therefore, in any case $\bar{\nu}^{c_0}$ satisfies the hypotheses of Theorem 3.12 with Frostman constant 20.

We will not apply Theorem 3.12 directly to the sets A, B, but rather to A, $(c_0B)_{\delta}$, where

$$(c_0 B)_{\delta} = (\delta \cdot \mathbb{Z}) \cap (c_0 B)_{\delta} \subset (\delta \cdot \mathbb{Z}) \cap [0, 1].$$

Evidently $|(c_0B)_{\delta}| \gtrsim c_0|B| \ge \delta^{\epsilon_0}|B|$ by (3.19). It follows that

$$|(c_0B)_{\delta} + (c_0B)_{\delta}| \lesssim |B + B| \leq \delta^{-\epsilon_B}|B| \lesssim \delta^{-\epsilon_0 - \epsilon_B}|(c_0B)_{\delta}|.$$

Since $\epsilon_0 + \epsilon_B \leq \overline{\epsilon}_B/2$ by (3.15), and if $\delta > 0$ is sufficiently small, we conclude that $(c_0B)_{\delta}$ satisfies the small doubling assumption in Theorem 3.12 with constant $\overline{\epsilon}_B$. We moreover claim that $(c_0B)_{\delta}$ satisfies the Frostman condition $|(c_0B)_{\delta} \cap B(x,r)| \leq r^{\kappa/2} |(c_0B)_{\delta}|$ for all

 $\delta \leq r \leq \delta^{\overline{\epsilon}_0}$. To see this, fix $\delta \leq r \leq \delta^{\overline{\epsilon}_0} \leq \delta^{2\epsilon_0} \leq c_0 \delta^{\epsilon_0}$ (by (3.15) and (3.19)), and note that

$$\begin{aligned} |(c_0B)_{\delta} \cap B(x,r)| &\lesssim |B \cap B(x,c_0^{-1}r)| \\ &\leqslant (c_0^{-1}r)^{\kappa} \cdot |B| \\ &\lesssim c_0^{-2} \cdot r^{\kappa} \cdot |(c_0B)_{\delta}| \\ &\leqslant \delta^{-2\epsilon_0} \cdot r^{\kappa/2} \cdot r^{\kappa/2} \cdot |(c_0B)_{\delta}| \\ &\stackrel{(3.15)}{\leqslant} \delta^{2\epsilon_0} \cdot r^{\kappa/2} \cdot |(c_0B)_{\delta}|. \end{aligned}$$

This implies $|(c_0B)_{\delta} \cap B(x,r)| \leq r^{\kappa/2} |(c_0B)_{\delta}|$, provided that $\delta_0 \geq \delta$ is sufficiently small. We have now shown that Theorem 3.12 is applicable with the parameters $\alpha, \beta, \gamma, \kappa/2$, and $\mathbf{C} = 20$ to the the sets $A, (c_0B)_{\delta}$, and the measure $\bar{\nu}^{c_0}$.

Since $\delta \leq \delta_0 \leq \overline{\delta}_0$, Theorem 3.12 implies the existence of a point $c \in \operatorname{spt}(\overline{\nu}^{c_0}) \subset [-1,1] \cap c_0^{-1} \cdot (\operatorname{spt} \nu - \operatorname{spt} \nu)$ such that

$$A' + c(c_0 B)_{\delta} \ge \delta^{-\bar{\epsilon}} |A| \tag{3.20}$$

for all subsets $A' \subset A$ with $|A'| \ge (1 - \bar{\rho})|A|$. Note that the point $c \in \operatorname{spt}(\bar{\nu}^{c_0})$ in (3.20) can be written as $c = c_0^{-1} \cdot (c_1 - c_2)$ for certain points $c_1, c_2 \in \operatorname{spt}(\nu)$. Therefore

$$|A' + (c_1 - c_2)B|_{\delta} = |A' + \frac{c_1 - c_2}{c_0} \cdot c_0 B|_{\delta} \gtrsim |A' + c(c_0 B)_{\delta}|_{\delta} \ge \delta^{-\bar{\epsilon}}|A|$$
(3.21)

for all $A' \subset A$ with $|A'| \ge (1 - \bar{\rho})|A|$. We now claim that there exists $\bar{c} \in \{c_1, c_2\}$ such that

$$|A' + \bar{c}B|_{\delta} \ge \delta^{-\epsilon}|A|, \qquad A' \subset A, \ |A'| \ge (1-\rho)|A|, \tag{3.22}$$

assuming that $\delta_0 \ge \delta$ is small enough, depending on ϵ, ρ . This will prove Theorem 3.1.

Assume that (3.22) fails for both $\bar{c} \in \{c_1, c_2\}$, and let $A'_1, A'_2 \subset A$ be subsets of cardinalities $|A'_j| \ge (1 - \rho), j \in \{1, 2\}$, such that

$$|A'_{1} + c_{1}B| < \delta^{-\epsilon}|A|$$
 and $|A'_{2} + c_{2}B| < \delta^{-\epsilon}|A|$. (3.23)

We first observe from the second inequality in (3.23) that

$$A_2' + c_2 B|_{\delta} \leq \delta^{-\epsilon} |A| \leq 2\delta^{-\epsilon} |A_2'|.$$

By Lemma 3.13, for $N \ge 1$ there exists a subset $A_2'' \subset A_2'$ of cardinality $|A_2''| \ge (1 - \rho)|A_2'|$ such that

$$|A_2'' - c_2 B|_{\delta} \lesssim_{\rho,N} (\delta^{-\epsilon})^{2^N} |A_2'|^{1+1/N} \lesssim \delta^{-\epsilon \cdot 2^{N+2} - 1/N} |A_2''|.$$
(3.24)

We apply this with $N \sim \log_2(1/\epsilon)$ satisfying $\epsilon \cdot 2^{N+2} \sim \sqrt{\epsilon}$. Since with this choice $\epsilon \cdot 2^{N+2} \sim \sqrt{\epsilon} \ll 1/\log_2(1/\epsilon) \sim 1/N$, we have $\epsilon \cdot 2^{N+1} + 1/N \leq 2/N \sim 1/\log_2(1/\epsilon)$, and we deduce from (3.24) that

$$|A_2'' - c_2 B|_{\delta} \lesssim_{\epsilon,\rho} \delta^{-C_0/\log_2(1/\epsilon)} |A_2''|$$

for some absolute constant $C_0 > 0$.

Now, recall that $|A'_1| \ge (1-\rho)|A|$ and $|A''_2| \ge (1-\rho)|A'_2| \ge (1-\sqrt{\rho})|A|$. Consequently, the intersection $A' := A'_1 \cap A''_2$ satisfies $|A'| \ge (1-\rho-\sqrt{\rho})|A|$. Evidently,

$$|A' + c_1 B| \leqslant \delta^{-\epsilon} |A| \lesssim \delta^{-C_0/\log_2(1/\epsilon)} |A'| \quad \text{and} \quad |A' - c_2 B| \lesssim_{\rho,\epsilon} \delta^{-C_0/\log_2(1/\epsilon)} |A'|.$$

By Lemma 3.2, there exists a further subset $A'' \subset A'$ with

$$|A''| \ge (1-\rho)|A'| \ge (1-\rho)(1-\rho-\sqrt{\rho})|A| \stackrel{(3.16)}{\ge} (1-\bar{\rho})|A|$$

such that

$$|A'' + c_1 B - c_2 B|_{\delta} \lesssim_{\epsilon,\rho} \delta^{-2C_0/\log_2(1/\epsilon)} |A''| \stackrel{(3.14)}{\leqslant} \delta^{-\overline{\epsilon}/2} |A|$$

This contradicts (3.21) for $\delta > 0$ small enough, depending on ϵ , ρ , and proves (3.22). The proof of Theorem 3.1 is complete.

3.3. **Removing reference to subsets.** In the previous reductions, we have upgraded the assumptions of Theorem 1.5 in two ways: we have arranged the set *B* to have small doubling, and we have arranged the Frostman constant of ν to be absolute. However, there has been a price: whereas Theorem 1.5 only claims that $|A + cB|_{\delta} \ge \delta^{-\epsilon}|A|$ for some $c \in \operatorname{spt}(\nu)$, Theorem 3.12 claims the existence of $c \in \operatorname{spt}(\nu)$ such that $|A + cB|_{\delta} \ge \delta^{-\epsilon}|A|$ for all $A' \subset A$ with $|A'| \ge (1 - \rho)|A|$. It turns out that this innocent-looking difference makes Theorem 3.12 difficult (impossible for the author) to prove with a direct assault. Therefore, we need a final reduction to the following statement:

Theorem 3.25. Let $0 < \beta \leq \alpha < 1$ and $\kappa > 0$. Then, for every $\gamma \in ((\alpha - \beta)/(1 - \beta), 1]$, there exist $\epsilon, \epsilon_0, \epsilon_B, \delta_0 \in (0, \frac{1}{2}]$, depending only on $\alpha, \beta, \gamma, \kappa$, such that the following holds. Let $\delta \in 2^{-\mathbb{N}}$ with $\delta \in (0, \delta_0]$, and let $A, B \subset (\delta \cdot \mathbb{Z}) \cap [0, 1]$ satisfy the following hypotheses:

(A)
$$|A| \leq \delta^{-\alpha}$$
.

(B) $|B| \ge \delta^{-\beta}$, and B satisfies the following Frostman condition:

 $|B \cap B(x,r)| \leqslant r^{\kappa}|B|, \qquad \delta \leqslant r \leqslant \delta^{\epsilon_0}.$

Assume moreover that $|B + B| \leq \delta^{-\epsilon_B} |B|$.

Further, let ν be a Borel probability measure with $\operatorname{spt}(\nu) \subset [-1,1]$ satisfying the Frostman condition $\nu(B(x,r)) \leq 40 \cdot r^{\gamma}$ for $x \in \mathbb{R}$ and $r \geq \delta$. Then, there exists a point $c \in \operatorname{spt}(\nu)$ such that $|A + cB|_{\delta} \geq \delta^{-\epsilon}|A|$.

Theorem 3.25 only differs from Theorem 3.12 in its (superficially) weaker conclusion, and in that the Frostman constant of ν has increased from 20 to 40.

Proof of Theorem 3.12 *assuming Theorem* 3.25. Fix the parameters $0 < \beta \leq \alpha < 1$, κ , and $\gamma > (\alpha - \beta)/(1 - \beta)$ from Theorem 3.12. As usual, our task is to find the parameters $\epsilon, \epsilon_0, \epsilon_B, \delta_0, \rho$ such that Theorem 3.12 is satisfied. In doing so, we apply Theorem 3.25 to the parameters $0 < \beta \leq \overline{\alpha} < 1$ and κ , where $\overline{\alpha} \in (\alpha, 1)$ is arbitrary such that the key inequality $\gamma > (\overline{\alpha} - \beta)/(1 - \beta)$ still holds. Then, we let

$$\bar{\epsilon}, \bar{\epsilon}_0, \bar{\epsilon}_B, \delta_0 > 0 \tag{3.26}$$

be the constants given by Theorem 3.25 with parameters $\bar{\alpha}$, β , κ , γ . We now begin defining the parameters ϵ , ϵ_0 , ϵ_B , δ_0 , ρ . We set

$$\epsilon_0 := \bar{\epsilon}_0 \quad \text{and} \quad \epsilon_B = \bar{\epsilon}_B.$$
 (3.27)

We will need that $\delta_0 \leq \overline{\delta}_0$, and there will be an additional (simple) dependences on the allowed parameters, which will be explained when they arise. To define the parameters ϵ, ρ , fix a natural number $N \sim 1/\overline{\epsilon}$, so that the following holds:

$$(N-1)^{-1} < \bar{\epsilon}/2. \tag{3.28}$$

Then, let

$$\epsilon := \frac{\bar{\alpha} - \alpha}{2^{N+1}}.\tag{3.29}$$

Finally, define $\rho > 0$, depending only on $\bar{\epsilon}$, so small that

$$\left(2(1-(1-\rho)^N)\right)^{1/2^N} \leqslant \frac{1}{2}.$$
(3.30)

This is possible, since the inequality is clearly true for $\rho = 0$.

We will now make the counter assumption that Theorem 3.12 fails for certain sets $A, B \subset (\delta \cdot \mathbb{Z}) \cap [0, 1]$, and a probability measure ν on [-1, 1], satisfying the hypotheses of Theorem 3.12 with parameters $\alpha, \beta, \kappa, \gamma$, and the constants $\epsilon_0, \epsilon, \delta_0$ described above. This means that for every $c \in C = \operatorname{spt}(\nu)$, there exists a subset $A_c \subset A$ with the properties

$$|A_c| \ge (1-\rho)|A| \quad \text{and} \quad |A_c + cB|_{\delta} \le \delta^{-\epsilon}|A|.$$
(3.31)

The plan is to use this information to construct a new set $A \subset (\delta \cdot \mathbb{Z}) \cap [0, 1]$, and a new probability measure $\bar{\nu}$ on [-1, 1], such that the triple $\bar{A}, B, \bar{\nu}$ satisfies the hypotheses of Theorem 3.25 with parameters $\bar{\alpha}, \beta, \kappa, \gamma$ and constants $\bar{\epsilon}_0, \bar{\epsilon}_B$, but nevertheless $|\bar{A} + cB| < \delta^{-\bar{\epsilon}} |\bar{A}|$ for all $c \in \operatorname{spt}(\bar{\nu})$. This contradiction will complete the proof of Theorem 3.12.

Given such a set $A_c \subset A$ for every $c \in C$, we observe that

$$\int \dots \int |A_{c_1} \cap \dots \cap A_{c_N}| \, d\nu(c_1) \cdots d\nu(c_N) \ge (1-\rho)^N |A| \tag{3.32}$$

by Hölder's inequality. Consider the set

$$\Omega := \{ (c_1, \ldots, c_N) \in C^N : |A_{c_1} \cap \ldots \cap A_{c_N}| \ge \frac{1}{2} |A| \},\$$

If "I" temporarily stands for the integral in (3.32), we have

$$|1-\rho)^N|A| \leq I \leq \nu^N(\Omega^c) \cdot \frac{1}{2}|A| + (1-\nu^N(\Omega^c)) \cdot |A|$$

which can be rearranged to $\nu^N(\Omega^c) \leq 2(1 - (1 - \rho)^N)$. Consequently

$$\nu^{N}(\Omega) \ge 1 - 2(1 - (1 - \rho)^{N}) =: 1 - \theta_{0}.$$
(3.33)

For $c_1, \ldots, c_n \in C$ fixed, we define

$$\Omega_{c_1\cdots c_n} := \{ (c_{n+1}, \ldots, c_N) \in C^{N-n} : (c_1, \ldots, c_N) \in \Omega \}.$$

It follows from Fubini's theorem that

$$\nu^{N-n}(\Omega_{c_1\cdots c_n}) = \int \nu^{N-n-1}(\Omega_{c_1\cdots c_n c}) \, d\nu(c)$$
(3.34)

for all $c_1, \ldots, c_n \in C$, and $0 \le n \le N-2$. The same remains true for n = 0, if the left hand side is interpreted as $\nu^N(\Omega)$. Equation (3.34) also remains valid for n = N-1 if we define the notation $\nu^{N-n-1} = \nu^0$ as follows:

$$\nu^{0}(\Omega_{c_{1}\cdots c_{N-1}c}) := \mathbf{1}_{\Omega}(c_{1},\dots,c_{N-1},c).$$
(3.35)

We will use this notation in the sequel.

For $(c_1, \ldots, c_N) \in \Omega$ fixed, we write

$$A_{c_1 \cdots c_N} := A_{c_1} \cap \ldots \cap A_{c_N} \implies |A_{c_1 \cdots c_N}| \ge \frac{1}{2} |A|.$$
(3.36)

We now construct a sequence of sets $H_n \subset \delta \cdot \mathbb{Z}$, $1 \leq n \leq N$. At the same time, we will construct subsets $C_1, \ldots, C_N \subset C$, and points $c_n \in C_n$, $1 \leq n \leq N$, with the properties

$$\nu^{N-n}(\Omega_{c_1\cdots c_n}) \ge 1 - \theta_n \quad \text{and} \quad \nu(C_n) \ge 1 - \theta_n, \quad 1 \le n \le N,$$
(3.37)

where we define inductively

$$\theta_n := \sqrt{\theta_{n-1}} \ge \theta_{n-1}.$$

In particular, the first part of (3.37) with n = N shows that $(c_1, \ldots, c_N) \in \Omega$, recall the notation (3.35). As a second remark, recalling the definition of $\theta_0 = 2(1 - (1 - \rho)^N)$, and combining this with the definition of ρ in (3.30), one sees that $\theta_n \leq \frac{1}{2}$ for all $1 \leq n \leq N$. To begin with, we define

$$C_1 := \{ c \in C : \nu^{N-1}(\Omega_c) \ge 1 - \theta_1 \},\$$

and we choose an arbitrary element $c_1 \in C_1$. Since

$$1 - \theta_0 \leqslant \nu^N(\Omega) = \int \nu^{N-1}(\Omega_c) \, d\nu(c) \leqslant \nu(C_1^c) \cdot (1 - \theta_1) + (1 - \nu(C_1^c)) = -\theta_1 \cdot \nu(C_1^c) + 1$$

by (3.33), we observe that $\nu(C_1^c) \leq \theta_0/\theta_1 = \theta_1$, and consequently $\nu(C_1) \geq 1 - \theta_1$. In particular $C_1 \neq \emptyset$. We then define

$$H_1 := c_1 B.$$

Assume inductively that H_1, \ldots, H_n and $C_1, \ldots, C_n \subset C$, and $c_j \in C_j$, $1 \leq j \leq n$, have already been constructed, and satisfy (3.37). We pick an element $c_{n+1} \in C_{n+1}$, where

$$C_{n+1} := \{ c \in C : \nu^{N-n-1}(\Omega_{c_1 \cdots c_n c}) \ge 1 - \theta_{n+1} \}, \quad 1 \le n \le N-1.$$

For n = N - 1, the notation $\nu^{N-n-1}(\Omega_{c_1\cdots c_n c})$ should be interpreted as in (3.35), so

$$C_N = \{ c \in C : \mathbf{1}_{\Omega}(c_1, \dots, c_{N-1}, c) \ge 1 - \theta_N \} = \{ c \in C : (c_1, \dots, c_{N-1}, c) \in \Omega \}.$$

For an arbitrary choice $c_{n+1} \in C_{n+1}$, we note that the first part of (3.37) is satisfied with index "n + 1", by the definition of C_{n+1} .

The set C_{n+1} also satisfies the second part of (3.37) with index "n + 1", since

$$1 - \theta_n \stackrel{(3.37)}{\leqslant} \nu^{N-n} (\Omega_{c_1 \cdots c_n}) \stackrel{(3.34)}{=} \int \nu^{N-n-1} (\Omega_{c_1 \cdots c_n c}) \, d\nu(c) \leqslant -\theta_{n+1} \cdot \nu(C_{n+1}^c) + 1,$$

and consequently $\nu(C_{n+1}^c) \leq \theta_n/\theta_{n+1} = \theta_{n+1}$, and $\nu(C_{n+1}) \geq 1 - \theta_{n+1}$.

Whereas $c_1 \in C_1$ was chosen arbitrarily, the element $c_{n+1} \in C_{n+1}$ is chosen in such a way that the quantity $|H_n + c_{n+1}B|_{\delta}$ is maximised, among all possible choices $c_{n+1} \in C_{n+1}$. We then define

$$H_{n+1} := (H_n + c_{n+1}B)_{\delta}.$$

Continuing in this manner produces a distinguished sequence $(c_1, \ldots, c_N) \in \Omega$, which we fix for the remainder of the argument, and the sequence of sets H_1, \ldots, H_N .

Note that $H_n \subset (\delta \cdot \mathbb{Z}) \cap [0, N]$ for all $1 \leq n \leq N$ by a straightforward induction, so $|H_n| \leq N\delta^{-1}$. Therefore, by the pigeonhole principle, there exists an index $n \in \{1, \ldots, N-1\}$ such that

$$|H_{n+1}| \leq (N\delta^{-1})^{1/(N-1)} |H_n| \leq 2\delta^{-1/(N-1)} |H_n|.$$
(3.38)

For this particular index $n \in \{1, ..., N-1\}$, we then have $|H_n + cB|_{\delta} \leq |H_{n+1}| \leq 2\delta^{-1/(N-1)}|H_n|$ for all $c \in C_{n+1}$ by the definition of H_{n+1} , and therefore

$$|H_n + cB|_{\delta} \leqslant \delta^{-\bar{\epsilon}/2} |H_n|, \qquad c \in C_{n+1}, \tag{3.39}$$

recalling (3.28), and assuming that $\delta > 0$ is small enough.

We now claim that (3.39) violates Theorem 3.25 with parameters $\bar{\alpha}$, β , κ , γ , and with the objects

$$A := H_n, \quad B, \quad \text{and} \quad \bar{\nu} := \nu (C_{n+1})^{-1} \cdot \nu|_{C_{n+1}}.$$
 (3.40)

We need to check the following items to contradict Theorem 1.5:

- (a) $|\bar{A}| \leq \delta^{-\bar{\alpha}}$,
- (b) $|B| \ge \delta^{-\beta}$ and $|B + B| \le \delta^{-\overline{\epsilon}_B} |B|$, and *B* satisfies a Frostman condition with exponents κ and $\overline{\epsilon}_0$,
- (c) $\bar{\nu}$ satisfies a Frostman condition with exponent γ and constant 40.

Point (b) is true by assumption (and since we chose $\epsilon_B = \bar{\epsilon}_B$ and $\epsilon_0 = \bar{\epsilon}_0$), so only (a) and (c) need to be verified. We first use the Plünnecke-Ruzsa inequality to establish (a), assuming that $\delta > 0$ is sufficiently small in terms of $N, \bar{\alpha}$. Clearly \bar{A} can be written as a sum of $n \leq N$ sets of the form $c_m B$, for some $1 \leq m \leq n$, where c_m is an index in the (fixed) sequence $(c_1, \ldots, c_N) \in \Omega$. Noting that $A_{c_1 \cdots c_N} \subset A_{c_m} \subset A$, each of these sets individually satisfies

$$|A_{c_1\cdots c_N} + c_m B|_{\delta} \leq |A_{c_m} + c_m B|_{\delta} \stackrel{(3.31)}{\leq} \delta^{-\epsilon} |A| \stackrel{(3.36)}{\leq} 2\delta^{-\epsilon} |A_{c_1\cdots c_N}|.$$

We may therefore infer that

$$|\bar{A}| \lesssim_{N,\rho} \delta^{-2^N \epsilon} |A| \leqslant \delta^{-2^N \epsilon - \alpha}$$

from the Plünnecke-Ruzsa inequality, Lemma 3.2. This inequality implies $|\bar{A}| \leq \delta^{-\bar{\alpha}}$ for small enough $\delta > 0$, recalling our choice of ϵ at (3.29).

We move to (c). Recalling (3.40), and from (3.37) that $\nu(C_{n+1}) \ge 1 - \theta_N \ge \frac{1}{2}$, we have

$$\bar{\nu}(B(x,r)) \leq 2 \cdot \nu(B(x,r)) \leq 40 \cdot r^{\gamma}, \qquad x \in \mathbb{R}, \, r \geq \delta.$$

We have now reached a situation which violates Theorem 3.25 for the choice of parameters $\bar{\alpha}, \beta, \kappa, \gamma$: the objects $\bar{A}, B, \bar{\nu}$ satisfy all the hypotheses (by (a)-(c)), but nevertheless we have $|\bar{A} + cB|_{\delta} \leq \delta^{-\bar{\epsilon}}|\bar{A}|$ for all $c \in \operatorname{spt}(\bar{\nu}) \subset \overline{C}_{n+1}$ (we know this for $c \in C_{n+1}$ by (3.39), but this implies the same for $c \in \overline{C}_{n+1}$, up to a constant factor). This violates Theorem 3.25, since $\bar{\epsilon} > 0$ was the constant associated to $\bar{\alpha}, \beta, \kappa, \gamma$. Therefore the counter assumption (3.31) is false, and the proof of Theorem 3.12 is complete.

To be precise, we have ignored that $\overline{A} \subset [0, N]$ instead of $\overline{A} \subset [0, 1]$. This can be dealt with as in the proof of Corollary 1.7, or below (3.10). We leave this to the reader.

3.4. **Bonus reduction.** We have now reduced the proof of Theorem 1.5 to the proof of Theorem 3.25. For notational convenience in the future, we mention one final reduction: we may assume that $1 \in \operatorname{spt}(\nu)$. Indeed, assume that Theorem 3.25 is known under this extra assumption. Then, let A, B, ν be a general triple as in Theorem 3.25. Since ν is a probability measure, $\operatorname{spt}(\nu) \subset [-1, 1]$ and $\nu(B(x, r)) \leq 40 \cdot r^{\gamma}$, the point $c_0 \in \operatorname{spt}(\nu) \cap [-1, 1]$ with maximal absolute value satisfies

$$|c_0| \ge 40^{-1/\gamma}$$
.

Consider the measure $\bar{\nu}(A) := \nu(c_0 A)$. Observe that $\bar{\nu}(B(x,r)) \leq 40 \cdot r^{\gamma}$ and $\operatorname{spt}(\bar{\nu}) = c_0^{-1} \operatorname{spt}(\nu)$. Therefore $1 \in \operatorname{spt}(\bar{\nu}) \subset [-1,1]$, so $\bar{\nu}$ satisfies the extra assumption. We then apply the (assumedly known) version of Theorem 3.25 to A, $(c_0 B)_{\delta}$, $\bar{\nu}$. The set $(c_0 B)_{\delta}$ will have slightly worse constants than B, in a manner depending on γ only, so the theorem needs to be applied with appropriately modified parameters. Once this has been done, we find a point $c = c_0^{-1}c_1 \in \operatorname{spt}(\bar{\nu})$, where $c_1 \in \operatorname{spt}(\nu)$, such that

$$|A + cB|_{\delta} \gtrsim |A + (c_1/c_0) \cdot (c_0B)_{\delta}|_{\delta} = |A + c(c_0B)_{\delta}|_{\delta} \ge \delta^{-\epsilon}|A|,$$

and the proof of Theorem 3.25 (without the extra assumption) is complete.

4. PROOF OF THEOREM 3.25

4.1. **Preliminaries.** We have now reduced the proof of Theorem 1.5 to the proof of Theorem 3.25. We fix the parameters $\alpha, \beta, \gamma, \kappa$, with $0 < \beta \leq \alpha < 1$ and $(\alpha - \beta)/(1 - \beta) < \gamma \leq 1$. We also fix sets $A, B \subset (\delta \cdot \mathbb{Z}) \cap [0, 1]$ and a Borel probability measure ν with $\operatorname{spt}(\nu) \subset [-1, 1]$, satisfying all the hypotheses of Theorem 3.25 with sufficiently small constants $\epsilon_0, \epsilon_B > 0$ to be determined later. For future reference, we write

$$|A| =: \delta^{-\alpha}, \qquad 0 \leqslant \bar{\alpha} \leqslant \alpha. \tag{4.1}$$

We make a counter assumption: $|A + cB|_{\delta} < \delta^{-\epsilon}|A|$ for all $c \in \operatorname{spt}(\nu)$. Since we may assume that $1 \in \operatorname{spt}(\nu)$ by Section 3.4, we have the assumptions

$$|A+B| \leq \delta^{-\epsilon} |A| \quad \text{and} \quad |B+B| \leq \delta^{-\epsilon_B} |B|. \tag{4.2}$$

If $\epsilon, \epsilon_B > 0$ in (4.2) are small enough, depending only on $\alpha, \beta, \kappa, \gamma$, we will be able to find a point $c \in \operatorname{spt}(\nu)$ such that $|A + cB|_{\delta} \ge \delta^{-\epsilon}|A|$. This will violate the counter assumption, and prove Theorem 3.25. The necessary values of $\epsilon = \epsilon(\alpha, \beta, \gamma, \kappa) > 0$ and $\epsilon_B = \epsilon_B(\alpha, \beta, \gamma, \kappa) > 0$ in (4.2) will be fixed during the proof of Proposition 4.12.

4.2. Shmerkin's inverse theorem. In the case A = B, Bourgain [5] used an assumption of the form (4.2) to obtain, up to passing to a subset, a special multi-scale structure inside A: informally speaking, when passing from one scale to the next, either A has *full branching*, or then *no branching*. Similar statements have, after Bourgain's work, been proved by Hochman [18] and Shmerkin [32] in the case where $A \neq B$, and indeed A and B may have very different sizes. This is our situation, and we will apply Shmerkin's theorem, which we state in Theorem 4.6.

Definition 4.3 (δ -sets and measures, L^2 -norms). Let $\delta \in 2^{-\mathbb{N}}$ be a dyadic rational. A subset of $(\delta \cdot \mathbb{Z}) \cap [0, 1)$ is called a δ -set. A probability measure supported on a δ -set is called a δ -measure. The L^2 -norm of a δ -measure μ is defined by

$$\|\mu\|_{L^2} := \left(\sum_{z \in \delta \cdot \mathbb{Z}} \mu(\{z\})^2\right)^{1/2}$$

We will only be concerned with δ -measures of the form $\mu = |A|^{-1} \mathcal{H}^0|_A$, where $A \subset [0,1)$ is a δ -set. Then $\|\mu\|_{L^2} = |A|^{-1/2}$.

Definition 4.4 (Uniform sets). Let $m, N \in \mathbb{N}$, and set $\delta := 2^{-mN} \in 2^{-\mathbb{N}}$. For $A \subset [0, 1)$ and $s \in \{0, \ldots, N-1\}$, write $\mathcal{I}_{ms}(A) := \{I \in \mathcal{D}_{ms} : A \cap I \neq \emptyset\}$ for the collection of dyadic intervals of side-length 2^{-ms} (these are denoted \mathcal{D}_{ms}) with non-empty intersection with A. We say that A is (m, N)-uniform if

$$R_A(s) := |I \cap A|_{2^{-m(s+1)}}, \qquad I \in \mathcal{I}_{ms}(A),$$

is independent of the choice of $I \in \mathcal{I}_{ms}(A)$. We may also write that A is (m, N, R_A) uniform if the branching numbers R_A need emphasising.

As in [32], we will only consider uniform sets which are also δ -sets. It was observed by Bourgain [5] that every δ -set contains a uniform subsets of "comparable" cardinality. Thus, the possibility of finding uniform subsets has nothing to do with an assumption like (4.2). To explain what (4.2) implies, we introduce the following terminology: **Definition 4.5** (η -polarised pair). Let $m, N \in \mathbb{N}$, $\delta = 2^{-mN}$, and $\eta > 0$. A pair of (m, N)-uniform sets (A, B) is (η, m, N) -polarised, if

$$R_B(s) > 1 \implies R_A(s) \ge 2^{(1-\eta)m}, \qquad s \in \{0, \dots, N-1\}.$$

If A = B, we say that A (instead of (A, A)) is (η, m, N) -polarised.

Note that $R_A(s) \leq 2^m$ for all $s \in \{0, \ldots, N-1\}$, so $R_A(s) \geq 2^{(1-\eta)m}$ means that $R_A(s)$ is nearly maximal. Bourgain [5] proved that if A is a δ -set with $|A + A| \leq \delta^{-\epsilon}|A|$, then A contains a uniform subset A' such that $|A'| \geq \delta^{\eta}|A|$, and A' is η -polarised, where $\eta = o_{\epsilon}(1)$. This means that either $R_{A'}(s) = 1$ or $R_{A'}(s) \geq 2^{(1-\eta)m}$ for all scales "s".

Versions of Bourgain's "polarisation theorem", explained above, for two different sets were found by Hochman [18] and Shmerkin [32]. Hochman first showed that if μ, ν are probability measures on [0, 1), then the entropy inequality $H(\mu * \nu) \leq H(\mu) + \epsilon$ implies a measure-theoretic version of the polarisation phenomenon for μ, ν . The set version, below, was established by Shmerkin [32] (with a proof very different from [18]):

Theorem 4.6 (Shmerkin). Let $\eta > 0$, and let $m(\eta) \in \mathbb{N}$ be sufficiently large, depending on η . Then, for all $m \ge m(\eta)$ there exists $\epsilon = \epsilon(\eta, m) > 0$ such that the following holds for all large enough $N \in \mathbb{N}$. Let $\delta = (2^{-m})^N$, and let $A, B \subset [0, 1]$ be δ -sets such that

$$|A + B| \leq \delta^{-\epsilon} |A|.$$

Then, there exist (m, N)-uniform sets $A' \subset A$ and $B' \subset B$ such that $|A'| \ge \delta^{\eta} |A|, |B'| \ge \delta^{\eta} |B|$, and (A', B') is (η, m, N) -polarised.

Remark 4.7. To be accurate, Theorem 4.6 is a slight refinement of Shmerkin's theorem: [32, Theorem 2.1] literally contains the following statement: *if* $\eta > 0$ *and* $m_0 \in \mathbb{N}$, *then there exists* $m = m(\eta, m_0) \ge m_0$ and $\epsilon = \epsilon(\eta, m_0) > 0$ as in Theorem 4.6. However, if one inspects the proof of [32, Theorem 2.1], one observes that the only dependence of $m = m(\eta, m_0)$ on m_0 is " $m \ge m_0$ ", and any choice of $m \ge m_0$ works, provided that m is also sufficiently large in terms of η . This is precisely what Theorem 4.6 says.

As another remark, Shmerkin's theorem actually concerns a pair of δ -measures μ_1, μ_2 instead of δ -sets: the measures of interest for our application are simply $\mu_1 = |A|^{-1} \mathcal{H}^0|_A$ and $\mu_2 = |B|^{-1} \mathcal{H}^0|_B$, and with such choices [32, Theorem 2.1] implies Theorem 4.6.

Remark 4.8. We will be applying Theorem 4.6 to dyadic scales of the form $\delta = 2^{-\ell mN}$, where $\ell, m, N \in \mathbb{N}$. Since $\delta = (2^{-m})^{\ell N} = (2^{-\ell m})^N$, a δ -set $A \subset [0, 1)$ may be $(m, \ell N)$ -uniform, or both. The former condition means that the branching numbers $R_A^m(s) = |A \cap I|_{2^{-m(s+1)}}$ are well-defined for $m \in [\ell N]$, whereas the latter means that the branching numbers $R_A^{\ell m}(\sigma) = |A \cap I|_{2^{-\ell m(\sigma+1)}}$ are well-defined for $\sigma \in [N]$. It is clear that every $(m, \ell N)$ -uniform $2^{-\ell mN}$ -set is $(\ell m, N)$ -uniform, and indeed

$$R_A^{\ell m}(\sigma) = \prod_{s=\ell\sigma}^{\ell(\sigma+1)-1} R_A^m(s), \qquad \sigma \in [N].$$

The converse is generally not true, so $(m, \ell N)$ -uniformity is a strictly stronger property than $(\ell m, N)$ -uniformity. We will also be interested in pairs (A, B) which are sometimes $(\eta, m, \ell N)$ -polarised, and sometimes $(\eta, \ell m, N)$ -polarised. In contrast to uniformity, there is no simple implication between these two properties.

In addition to Shmerkin's theorem, we will also need a lemma from its proof:

Lemma 4.9. Let $m, \ell, N \in \mathbb{N}$, $\delta = 2^{-\ell m N}$, and let $A \subset [0, 1)$ be an $(m, \ell N)$ -uniform δ set. Then A is also $(\ell m, N, R_A^{\ell m})$ -uniform for some $R_A \colon [N] \to \{1, \ldots, 2^{\ell m}\}$. If $S \subset [N]$ is arbitrary, there exists $A' \subset A$ which is $(m, \ell N)$ -uniform, and also $(\ell m, N, R_A^{\ell m})$ -uniform with

$$|A'| \ge |A| \cdot \prod_{\sigma \in \mathcal{S}} R_A^{\ell m}(\sigma)^{-1}, \quad and \quad R_{A'}^{\ell m}(\sigma) = \begin{cases} 1, & \sigma \in \mathcal{S}, \\ R_A^{\ell m}(\sigma), & \sigma \notin \mathcal{S}. \end{cases}$$

A similar statement holds true if $S \subset [\ell N]$, with the only difference that " $R_A^{\ell m}(\sigma)$ " and " $R_{A'}^{\ell m}(\sigma)$ " should be replaced by " $R_A^m(s)$ " and " $R_{A'}^m(s)$ " for $s \in [\ell N]$.

The lemma above is [32, Lemma 3.7]. To be accurate, the statement about A' remaining $(m, \ell N)$ -uniform is not part of the statement of [32, Lemma 3.7], but the 3.8-line proof quickly reveals that $(m, \ell N)$ -uniformity is not violated when passing between A and A'; the only point is to "collapse" all the branching of A for levels corresponding to $\sigma \in S$, or equivalently for $s \in \{\ell \sigma, \ell(\sigma + 1) - 1\}$ for all $\sigma \in S$.

4.3. Applying the inverse theorem. We start by fixing the following parameters:

$$\begin{cases} \ell = \ell(\alpha, \beta, \gamma, \kappa) \in \mathbb{N}, \\ \eta = \eta(\alpha, \beta, \gamma, \kappa) \in (0, 1), \\ m_0 \in \mathbb{N} \text{ with } m_0 \ge (40 + C_0)/\eta. \end{cases}$$

$$(4.10)$$

Here $C_0 > 0$ is an absolute constant to be specified later. The values of all these constants will be specified later, but as indicated above, all of them only depend on $\alpha, \beta, \gamma, \kappa$. For the reader interested in seeing specific choices, we refer to (4.23) and the discussion afterwards. Recall that the set *B* satisfies the Frostman condition $|B \cap B(x, r)| \leq r^{\kappa}|B|$ for all $r \leq \delta^{\epsilon_0}$, where we may freely choose $\epsilon_0 = \epsilon_0(\alpha, \beta, \kappa, \gamma)$. We choose

$$\epsilon_0 := \eta. \tag{4.11}$$

We will assume that η , ϵ , $\epsilon_B < 1/1000$ in the sequel (but these upper bounds will generally not suffice). This section is devoted to the proof of the following proposition, whose proof will also finalise the choice of the parameters ϵ , $\epsilon_B > 0$, relative to η :

Proposition 4.12. There exist $\epsilon, \epsilon_B > 0$ and $m \ge m_0$, depending on $\alpha, \beta, \gamma, \kappa$, such that the following holds for all $\delta \in 2^{-\mathbb{N}}$ of the form $\delta = 2^{-\ell m N}$, $N \in \mathbb{N}$. Assume that $A, B \subset [0, 1]$ are δ -sets satisfying the small doubling assumptions (4.2). Then there exist subsets $A' \subset A$ and $B' \subset B$ with the following properties:

- (1) A' and B' are $(m, \ell N)$ -uniform with $|A'| \ge \delta^{\eta} |A|$ and $|B'| \ge \delta^{\eta/2} |B|$.
- (2) The pair (A', B') is $(\eta, m, \ell N)$ -polarised.
- (3) The set B' is $(\eta/2, \ell m, N)$ -polarised.

Remark 4.13. In the sequel, we will always work with scales of the form $\delta = 2^{-\ell mN}$ with the fixed parameters ℓ, m , which depend on $\alpha, \beta, \gamma, \kappa$. In other words, we initially prove Theorem 3.25 (and find the constants $\epsilon, \epsilon_0, \epsilon_B$) for only scales of this special form. After this has been accomplished, it is easy to check that the case of general scales $\delta \in 2^{-\mathbb{N}}$ is a corollary, assuming that the upper bound $\delta_0 = \delta_0(\alpha, \beta, \gamma, \kappa) > 0$ for δ is sufficiently small. The reason is that if $\delta \in 2^{-\mathbb{N}}$ is arbitrary, then there exists a scale of the form $\overline{\delta} = 2^{-\ell mN}$ with $\delta \leq \overline{\delta} \leq_{\alpha,\beta,\gamma,\kappa} \delta$. We leave the rest of this reduction to the reader. As another remark, we will later in the paper need to assume that $\epsilon = \epsilon(\alpha, \gamma) > 0$ is sufficiently small that

$$\frac{\epsilon}{1-\alpha-\epsilon} \leqslant \frac{\gamma}{2}.\tag{4.14}$$

This requirement should be combined with the one coming from Proposition 4.12.

Proof of Proposition 4.12. We begin by applying Theorem 4.6 with constant $\eta^3 > 0$ to the pair (B, B), for which we assumed in (4.2) that $|B + B| \leq \delta^{-\epsilon_B}|B|$. Assume that $m \geq m(\eta^3) \in \mathbb{N}$ is sufficiently large that Theorem 4.6 applies. Assume additionally that $m \geq m_0$, where m_0 is the constant from (4.10). Then, if $\epsilon_B = \epsilon_B(\eta^3, \ell m) = \epsilon_B(\alpha, \beta, \gamma, \kappa) > 0$ and $\delta = (2^{-\ell m})^N$ are sufficiently small, we find an $(\ell m, N)$ -uniform subset $B' \subset B$ such that $|B'| \geq \delta^{\eta^3}|B|$, and B' is $(\eta^3, \ell m, N)$ -polarised. We have now fixed the value of the parameter $\epsilon_B > 0$ in (4.10) (and hence in Theorem 3.25)!

Next, note that $|A + B'| \leq |A + B| \leq \delta^{-\epsilon}|A|$. We therefore may apply Theorem 4.6 again to the pair (A, B'), again with parameter $\eta^3 > 0$. If $\epsilon = \epsilon(\eta^3, m) > 0$ is sufficiently small, we find an $(m, \ell N)$ -uniform subset $A' \subset A$ with $|A'| \geq \delta^{\eta^3}|A| \geq \delta^{\eta}|A|$, and an $(m, \ell N)$ -uniform subset $B'' \subset B'$ such that

$$|B''| \ge \delta^{\eta^3} |B'|, \tag{4.15}$$

and (A', B'') is $(\eta^3, m, \ell N)$ -polarised. In particular (A', B'') is $(\eta, m, \ell N)$ -polarised. We have now fixed the value of the parameter $\epsilon > 0$ in (4.2)!

Are we done with properties (1)-(3) above? Not quite: while passing from B' to B'', we might have lost the $(\eta^3, \ell m, N)$ -polarisation of B'. The plan will be to pass to a final $(m, \ell N)$ -uniform subset $B''' \subset B''$ which is $(\eta/2, \ell m, N)$ -polarised, and such that $|B'''| \ge \delta^{\eta/4}|B''|$. Then finally

$$|B'''| \ge \delta^{\eta/4} |B''| \ge \delta^{\eta/4 + \eta^3} |B'| \ge \delta^{\eta/4 + 2\eta^3} |B| \ge \delta^{\eta/2} |B|.$$

Also (A', B''') remains $(\eta, m, \ell N)$ -polarised, since this property is not violated by replacing B'' by an $(m, \ell N)$ -uniform subset, for example B'''.

Write

$$S_0 := \{ \sigma \in [N] : R_{B'}^{\ell m}(\sigma) = 1 \}$$
 and $S_1 := \{ \sigma \in [N] : R_{B'}^{\ell m}(\sigma) \ge 2^{(1-\eta^3)\ell m} \}.$

Since B' was constructed to be $(\eta^3, \ell m, N)$ -polarised, we have $[N] = S_0 \cup S_1$, and

$$|B'| = \prod_{\sigma \in \mathcal{S}_1} R_{B'}^{\ell m}(\sigma) \ge 2^{(1-\eta^3)\ell m|\mathcal{S}_1|}.$$

Now, let $S_{\text{bad}} := \{ \sigma \in S_1 : R_{B''}^{\ell m}(\sigma) < 2^{(1-\eta/2)\ell m} \}$, and $S_{\text{good}} := S_1 \setminus S_{\text{bad}}$. (Note that the numbers $R_{B''}^{\ell m}(\sigma)$ are well-defined, since B'' is $(m, \ell N)$ -uniform, hence $(\ell m, N)$ -uniform.) Then, since evidently $R_{B''}^{\ell m}(\sigma) \leq R_{B'}^{\ell m}(\sigma) = 1$ for all $\sigma \in S_0$, we have

$$|B''| = \prod_{\sigma \in \mathcal{S}_{\text{bad}}} R_{B''}^{\ell m}(\sigma) \cdot \prod_{\sigma \in \mathcal{S}_{\text{good}}} R_{B''}^{\ell m}(\sigma) \leqslant 2^{(1-\eta/2)\ell m|\mathcal{S}_{\text{bad}}|} \cdot 2^{\ell m(|\mathcal{S}_1| - |\mathcal{S}_{\text{bad}}|)}$$
$$= 2^{\ell m|\mathcal{S}_1| - (\eta/2)\ell m|\mathcal{S}_{\text{bad}}|}.$$

On the other hand, recalling that $\delta = 2^{\ell m N}$, we have

$$|B''| \stackrel{(4.15)}{\geqslant} \delta^{\eta^3} |B'| = \delta^{\eta^3} \cdot \prod_{\sigma \in \mathcal{S}_1} R^{\ell m}_{B'}(\sigma) \ge 2^{-\eta^3 \ell m N + (1-\eta^3) \ell m |\mathcal{S}_1|}.$$

Combining these inequalities and dividing both sides by $2^{\ell m |S_1|}$ yields

$$2^{-(\eta/2)\ell m|\mathcal{S}_{\text{bad}}|} \ge 2^{-\eta^3\ell mN - \eta^3\ell m|\mathcal{S}_1|} \implies |\mathcal{S}_{\text{bad}}| \le 2\eta^2 N + 2\eta^2|\mathcal{S}_1| \le (\eta/4)N$$

noting that $16\eta^2 \leq \eta$ since $\eta < 1/1000$. At this point, we simply apply the "collapsing" Lemma 4.9 to the the $(m, \ell N)$ -uniform set B'', and the set of scales $S := S_{\text{bad}}$. The result is an $(m, \ell N)$ -uniform subset $B''' \subset B''$, which is also $(\ell m, N, R_{B'''}^{\ell m})$ -uniform, with

$$R_{B''}^{\ell m}(\sigma) = \begin{cases} R_{B''}^{\ell m}(\sigma), & \sigma \notin \mathcal{S}_{\text{bad}}, \\ 1, & \sigma \in \mathcal{S}_{\text{bad}}, \end{cases}$$

and

$$|B'''| \ge |B''| \cdot \prod_{\sigma \in \mathcal{S}_{\text{bad}}} R_{B''}^{\ell m}(\sigma)^{-1} \ge |B''| \cdot 2^{-\ell m |\mathcal{S}_{\text{bad}}|} \ge \delta^{\eta/4} \cdot |B''|.$$

Then B''' is $(\eta/2, \ell m, N)$ -polarised, since if $R_{B'''}^{\ell m}(\sigma) > 1$, then necessarily $\sigma \in S_{\text{good}}$, hence $R_{B''}^{\ell m}(\sigma) = R_{B''}^{\ell m}(\sigma) \ge 2^{(1-\eta/2)\ell m}$ by definition. Now the pair of sets A', B''' (in place of A', B') satisfies all the requirements (1)-(3) in Proposition 4.12.

4.4. **Pruning** B' to improve separation I. It will be useful to reduce B' to a further subset, in order to gain a little extra separation. We prove the following proposition:

Proposition 4.16. Let $B' \subset B$ be the $(m, \ell N)$ -uniform $(\eta/2, \ell m, N)$ -polarised set found in Proposition 4.12. Then, there exists an $(m, \ell N)$ -uniform $(\eta, \ell m, N)$ -polarised subset $B'' \subset B'$ with $|B''| \ge \delta^{\eta/2}|B'|$, and which satisfies the following separation property:

$$s \in [\ell N], I_1, I_2 \in \mathcal{I}_{ms}(B''), I_1 \neq I_2 \implies \text{dist}(I_1, I_2) \ge 2^{-ms}.$$

$$(4.17)$$

Proof. We perform a straightforward "top down" reduction of B'. At scale s = 0, there is only one interval $[0,1) \in \mathcal{I}_0(B')$, so (4.17) is clear. At scale s = 1, remove $I \cap B'$ from B' for at most every second interval $I \in \mathcal{I}_m(B')$. This results in a new set $B'_1 \subset B'$ with $|B'_1| \ge \frac{1}{2}|B'|$ points, and

$$R^m_{B'_1}(0) := |B'_1|_{2^{-m}} \ge \frac{1}{2} R^m_{B'}(0).$$

Next, for s = 2, consider every (remaining) interval in $\mathcal{I}_m(B'_1)$. For each $I \in \mathcal{I}_m(B'_1)$, at most $\frac{1}{2}$ of the intervals from $\mathcal{I}_{2m}(B'_1)$ contained in I need removal to ensure (4.17) at level s = 2. However, the (minimal) number may vary depending on the choice of $I \in \mathcal{I}_m(B'_1)$. Fortunately, by removing some extra $\mathcal{I}_{2m}(B'_1)$ -intervals if necessary, we may make the number independent of $I \in \mathcal{I}_m(B'_1)$. This way, the number of remaining points in B'_1 again gets reduced by at most a factor of $\frac{1}{2}$. For the remaining points, say B'_2 , the new branching numbers $R^m_{B'_2}(2) := |B'_2 \cap I|_{2^{-2m}}$ are independent of the choice of $I \in \mathcal{I}_m(B'_2)$, and of course $R^m_{B'_2}(0) = R^m_{B'_1}(0)$ (in general, the removal process at level s never alters the branching numbers at levels < s - 1). Hence B'_2 is again $(m, \ell N)$ -uniform.

Once the deletion process has been executed successively at all levels $s \in \{1, ..., \ell N\}$, the remaining set $B'' := B'_N$ satisfies

$$|B''| \ge 2^{-\ell N} |B'| = \delta^{1/m} \cdot |B'| \ge \delta^{\eta/2} \cdot |B'|,$$

recalling that $m \ge m_0 \ge 2/\eta \ge 1/\eta$ by Proposition 4.12 and (4.10). Also, B'' remains $(m, \ell N)$ -uniform, with $R^m_{B''}(s) \ge \frac{1}{2}R^m_{B'}(s)$ for $s \in [\ell N]$. This implies that if $R^{\ell m}_{B''}(\sigma) > 1$, then

$$R_{B''}^{\ell m}(\sigma) \ge 2^{-\ell} R_{B'}^{\ell m}(\sigma) \ge 2^{-\ell + (1 - \eta/2)\ell m} = 2^{(1 - \eta/2 - 1/m)\ell m} \ge 2^{(1 - \eta)\ell m}$$

by the $(\eta/2, \ell m, N)$ -polarisation of B', and using that $m \ge m_0 \ge 2/\eta$, recall (4.10). In other words, B'' is $(\eta, \ell m, N)$ -polarised, as claimed.

Since $|B'| \ge \delta^{\eta/2}|B|$ by Proposition 4.16, we have $|B''| \ge \delta^{\eta/2}|B'| \ge \delta^{\eta}|B|$. Also, since (A', B') is $(\eta, m, \ell N)$ -polarised, the same is true of (A', B''). To simplify notation, we will remove one prime, that is, assume that B' already satisfies the separation property (4.17) constructed for B'' in the proposition above. To summarise the progress so far:

- $|A'| \ge \delta^{\eta} |A|$ and $|B'| \ge \delta^{\eta} |B|$,
- (A', B') is $(\eta, m, \ell N)$ -polarised and B' is $(\eta, \ell m, N)$ -polarised,
- *B*′ satisfies the separation property (4.17).

4.5. Intervals with small but non-zero *B*'-branching. For intervals $\mathcal{I} \subset [\ell N]$ and $\mathcal{J} \subset [N]$, and for $D' \in \{A', B'\}$, we define

$$R_{D'}^m(\mathcal{I}) := \prod_{s \in \mathcal{I}} R_{D'}^m(s) \quad \text{and} \quad R_{D'}^{\ell m}(\mathcal{J}) := \prod_{\sigma \in \mathcal{J}} R_{D'}^{\ell m}(\sigma).$$

We also define $\mathcal{N}_{B'}$ to consist of the maximal intervals $\mathcal{I} \subset [N]$ such that $R_{B'}^{\ell m}(\mathcal{I}) = 1$. Thus the intervals in $\mathcal{N}_{B'}$ partition the set $\{\sigma \in [N] : R_{B'}^{\ell m}(\sigma) = 1\}$.

Lemma 4.18. Write $|B| := \delta^{-\beta_1}$, where $\beta_1 \ge \beta$ by the hypothesis of Theorem 3.25. Then the following lower bound holds for the total length of the intervals in $\mathcal{N}_{B'}$:

$$\sum_{\mathcal{I}\in\mathcal{N}_{B'}} |\mathcal{I}| \ge (1 - \frac{\beta_1}{1 - \eta}) \cdot N$$

Remark 4.19. We remark that $\beta_1 \leq \alpha + \epsilon < 1$ (recall (4.14) for the second inequality). Otherwise $|A + B| \geq |B| = \delta^{-\beta_1} > \delta^{-\epsilon} |A|$, violating our counter assumption (4.2). Therefore $1 - \beta_1 > 0$, which will be (tacitly) needed several times below.

Proof. Let $N_{B'} := \bigcup N_{B'} \subset [N]$. Note that if $\sigma \in [N] \setminus N_{B'}$, then $R_{B'}^{\ell m}(\sigma) > 1$, and consequently $R_{B'}^{\ell m}(\sigma) \ge 2^{(1-\eta)\ell m}$, by the $(\eta, \ell m, N)$ -polarisation of B'. It follows that

$$2^{\beta_1 \ell m N} = \delta^{-\beta_1} \ge |B'| = \prod_{\sigma \in [N] \setminus N_{B'}} R^{\ell m}_{B'}(\sigma) \ge 2^{(1-\eta)\ell m \cdot (N-|N_{B'}|)}.$$

Consequently $N - |N_{B'}| \leq (\beta_1/(1 - \eta)) \cdot N$, and finally

$$|N_{B'}| \ge N - \frac{\beta_1}{1-\eta} \cdot N = (1 - \frac{\beta_1}{1-\eta}) \cdot N,$$

as claimed.

We would next like to extend the intervals $\mathcal{I} \in \mathcal{N}_{B'}$ to the left in such a manner that the B'-branching numbers of the extended intervals are relatively small, but not vanishingly small; say, we keep extending left until the extension $\overline{\mathcal{I}} \supset \mathcal{I}$ satisfies $R_{B'}^{\ell m}(\overline{\mathcal{I}}) \sim 2^{\zeta \ell m |\overline{\mathcal{I}}|}$ for some small parameter $\zeta > 0$. Unfortunately, this is not always possible: consider for example a scenario where the elements in $\mathcal{N}_{B'}$ are singletons. Then, as soon as an interval (a singleton) $\mathcal{I} \in \mathcal{N}_{B'}$ is extended to the left by a single element $\sigma \in [N]$, we have $R_{B'}^{\ell m}(\sigma) \approx 2^{\ell m}$ by the polarisation of B', and hence

$$R_{B'}^{\ell m}(\mathcal{I} \cup \{\sigma\}) = R_{B'}^{\ell m}(\sigma) \approx 2^{\ell m} = 2^{\ell m |\mathcal{I} \cup \{\sigma\}|/2}.$$

The right hand side is far too large for our purposes. This issue is not possible to overcome as long as we are fixated with the single scale partition [N], and here arises the need to play with the two different scale partitions [N] and $[\ell N]$.

The solution is to identify every interval $\mathcal{I} \in \mathcal{N}_{B'}$ with another, ℓ -times longer, subinterval of $[\ell N]$. More precisely, for every $\mathcal{I} = \{\sigma, \ldots, \tau\} \subset [N]$, we define the interval

$$\mathcal{I} := \{\ell\sigma, \ell\sigma + 1, \dots, \ell(\tau+1) - 1\} \subset [\ell N].$$

Thus, for example ℓ {0} = $[\ell]$, and $\ell[N] = [\ell N]$. It is easy to check that the branching numbers interact with this operation as follows:

$$R^m_{D'}(\ell \mathcal{I}) = R^{\ell m}_{D'}(\mathcal{I}), \qquad D' \in \{A', B'\}, \, \mathcal{I} \subset [N]$$

In particular:

$$\mathcal{I} \in \mathcal{N}_{B'} \implies R_{B'}^{\ell m}(\mathcal{I}) = 1 \implies R_{B'}^{m}(\ell \mathcal{I}) = 1.$$
(4.20)

Motivated by this observation, we define

$$\ell \mathcal{N}_{B'} := \{\ell \mathcal{I} : \mathcal{I} \in \mathcal{N}_{B'}\}.$$

The intervals in $\mathcal{N}_{B'}$ partition { $\sigma \in [N]$: $R_{B'}^{\ell m}(\sigma) = 1$ } by definition. The intervals in $\ell \mathcal{N}_{B'}$ are contained in the set { $s \in [\ell N]$: $R_{B'}^m(s) = 1$ } by (4.20), but may not cover it. However, we may infer the following lower bound for their total length form Lemma 4.18:

Corollary 4.21. The following lower bound holds for the total length of the intervals in $\mathcal{N}_{B'}^m$:

$$\sum_{\mathcal{J} \in \ell \mathcal{N}_{B'}} |\mathcal{J}| \ge \left(1 - \frac{\beta_1}{1 - \eta}\right) \cdot \ell N.$$

Proof. This follows immediately from Lemma 4.18 and the relation $|\ell \mathcal{I}| = \ell |\mathcal{I}|$ (noting also that if $\mathcal{I}_1, \mathcal{I}_2 \subset [N]$ with $\mathcal{I}_1 \cap \mathcal{I}_2 = \emptyset$, then $\ell \mathcal{I}_1 \cap \ell \mathcal{I}_2 = \emptyset$).

Next, instead of extending the intervals in $\mathcal{N}_{B'}$ to the left, as we first proposed, we do this to the intervals in $\ell \mathcal{N}_{B'}$. More precisely, fix a small parameter

$$\zeta = \zeta(\alpha, \beta, \gamma, \kappa) > 0 \quad \text{with} \quad \eta \ll_{\alpha, \beta, \gamma, \kappa} \zeta < \kappa/2.$$
(4.22)

It will eventually turn out that $\eta, \zeta > 0$ will need to be chosen so small that

$$\left(1 - \frac{\beta_1}{1 - \eta} - \frac{2\eta}{\kappa}\right) - \left(\frac{\alpha - (1 - \eta)(\beta_1 - \eta - 2\zeta)}{\Gamma}\right) \ge \frac{1}{2} \cdot \left[(1 - \beta_1) - (\alpha - \beta_1)/\Gamma\right],\tag{4.23}$$

where $(\alpha - \beta)/(1 - \beta) < \Gamma < \gamma$ is a constant to be fixed in (4.29), which satisfies $\Gamma \ge \gamma/2$. It might look suspicious that the requirement in (4.23) depends on β_1 , and not just $\alpha, \beta, \kappa, \gamma$. To see that this is not a problem, denote the left hand side $L(\beta_1)$ and the right hand side $\frac{1}{2} \cdot R(\beta_1)$. With this notation, one can easily check that

$$|L(\beta_1) - R(\beta_1)| \lesssim \frac{\eta}{\kappa} + \frac{\eta + \zeta}{\gamma} \text{ and } R(\beta_1) \ge R(\beta) \gtrsim_{\alpha,\beta,\gamma} 1.$$

The lower bound on $R(\beta)$ follows from $\Gamma > (\alpha - \beta)/(1 - \beta)$. From these estimates, one sees that if $\eta, \zeta > 0$ are chosen small enough depending only on $\alpha, \beta, \gamma, \kappa$, then $|L(\beta_1) - R(\beta_1)| \leq \frac{1}{2}R(\beta) \leq \frac{1}{2}R(\beta_1)$, and hence $L(\beta_1) \geq \frac{1}{2} \cdot R(\beta_1)$, as in (4.23).

In addition to the constraint in (4.23), we will finally (after (4.59)) need to take η sufficiently small in terms of $\zeta(\gamma - \Gamma)/4$. Of course (4.23) is compatible with such a constraint.

Enumerate $\ell N_{B'} = {\mathcal{I}_1, \mathcal{I}_2, ..., \mathcal{I}_k}$, where $\max \mathcal{I}_j < \min \mathcal{I}_{j+1}$. Start with \mathcal{I}_k , and recall that $R_{B'}^m(\mathcal{I}_k) = 1$ by (4.20). Begin extending \mathcal{I}_k to the left, adding elements of $[\ell N]$ one by one, until (the newly defined interval) \mathcal{I}_k satisfies

- (a) $R_{B'}^m(\mathcal{I}_k) \ge 2^{\zeta m |\mathcal{I}_k|}$, or
- (b) $0 \in \mathcal{I}_k$, and $R^m_{B'}(\mathcal{I}_k) < 2^{\zeta m |\mathcal{I}_k|}$.

In both cases (a)-(b) we have the upper bound

$$R_{B'}^m(\mathcal{I}_k) < 2^{\zeta m(|\mathcal{I}_k|-1)} \cdot 2^m \leqslant 2^{m(\zeta|\mathcal{I}_k|+1)} \leqslant 2^{m(2\zeta)|\mathcal{I}_k|},\tag{4.24}$$

choosing here $\ell = \ell(\alpha, \beta, \gamma) \ge 1$ sufficient large that

$$\ell \ge \zeta^{-1} \implies \zeta |\mathcal{I}_k| \ge \ell \zeta \ge 1.$$

This will be legitimate, since ζ only depends on α , β , γ , κ . Later, in (4.37), we will need to couple this requirement with $\ell \ge \epsilon^{-1}$, where $\epsilon = \epsilon(\alpha, \beta, \gamma) > 0$.

The extended interval \mathcal{I}_k may have "swallowed" a number of the previous intervals $\mathcal{I}_j \in \ell \mathcal{N}_{B'}$: note however that if $\mathcal{I}_j \cap \mathcal{I}_k \neq \emptyset$ for some $1 \leq j < k$, then actually $\mathcal{I}_j \subset \mathcal{I}_k$, since $R_{B'}^m(\mathcal{I}_j) = 1$ (in other words, there is no reason why the extension algorithm would terminate in the middle of \mathcal{I}_j). Let $\mathcal{I}_{k_1} \in \ell \mathcal{N}_{B'}$ be the right-most interval which does not intersect (equivalently: is not contained in) the extension \mathcal{I}_k . If no such interval remains, the algorithm terminates. Otherwise, repeat the extension procedure with \mathcal{I}_{k_1} . Continue in this manner until all the intervals in $\ell \mathcal{N}_{B'}$ are contained in (precisely) one of the extensions. Then rename the extensions as $\{\mathcal{J}_1, \ldots, \mathcal{J}_l\}$.

Since all the intervals in $\ell N_{B'}$ are contained in one of the extensions \mathcal{J}_i , we have $\sum |\mathcal{J}_i| \ge (1 - \beta_1/(1 - \eta)) \cdot \ell N$ by Corollary 4.21. Unfortunately, the leftmost interval \mathcal{J}_1 will be "useless" to us in case it was generated by case (b) (of course the intervals $\mathcal{J}_2, \ldots, \mathcal{J}_l$, if any exist, were generated by case (a)). We will next argue that \mathcal{J}_1 is so short in this case that its removal makes virtually no difference for the sum $\sum |\mathcal{J}_i|$.

If the interval \mathcal{J}_1 was indeed generated by case (b), then $\mathcal{J}_1 = \{0, \ldots, h\}$ for some $h \in \{0, \ldots, \ell N - 1\}$. Writing $r := 2^{-m(h+1)} \ge \delta$, then

$$|B'|_r \leqslant R^m_{B'}(\mathcal{J}_1) < 2^{\zeta m|\mathcal{J}_1|} = 2^{\zeta m(h+1)} = r^{-\zeta}.$$
(4.25)

If $r \ge \delta^{\epsilon_0} = 2^{-\epsilon_0 \ell m N}$, then

$$|\mathcal{J}_1| = h + 1 \leqslant \epsilon_0 \cdot \ell N \stackrel{(4.11)}{\leqslant} \eta \cdot \ell N.$$
(4.26)

On the other hand, if $\delta \leq r < \delta^{\epsilon_0}$, then the Frostman condition for *B* yields

$$|B' \cap B(x,r)| \leqslant |B \cap B(x,r)| \leqslant r^{\kappa}|B| \leqslant r^{\kappa} \cdot \delta^{-\eta}|B'|, \quad x \in \mathbb{R}$$

and consequently

$$r^{-\zeta} \stackrel{(4.25)}{>} |B'|_r \ge r^{-\kappa} \cdot \delta^{\eta}$$

This yields $r^{\kappa-\zeta} \ge \delta^{\eta}$, and since $\zeta < \kappa/2$ by (4.22), we have $r \ge \delta^{2\eta/\kappa}$. Recalling that $r = 2^{-m(h+1)}$ and $\delta = 2^{-\ell mN}$, this rearranges to

$$|\mathcal{J}_1| = h + 1 \leq \frac{2\eta}{\kappa} \cdot \ell N.$$

Combining this estimate with (4.26), we reach the following conclusion: if \mathcal{N}_+ stands for the intervals among $\{\mathcal{J}_1, \ldots, \mathcal{J}_l\}$ which were generated by case (a), then

$$\sum_{\mathcal{J}\in\mathcal{N}_{+}} |\mathcal{J}| \ge \sum_{i=1}^{l} |\mathcal{J}_{i}| - |\mathcal{J}_{1}| \ge \left(1 - \frac{\beta_{1}}{1 - \eta} - \frac{2\eta}{\kappa}\right) \cdot \ell N.$$
(4.27)

To recap, the intervals in \mathcal{N}_+ are subsets of $[\ell N]$, they are roughly compositions of a few intervals in $\ell \mathcal{N}_{B'}$, plus a little extra, and

$$2^{\zeta m|\mathcal{J}|} \leqslant R_{B'}^m(\mathcal{J}) \stackrel{(4.24)}{\leqslant} 2^{m(2\zeta)|\mathcal{J}|}, \qquad \mathcal{J} \in \mathcal{N}_+.$$
(4.28)

4.6. Branching of A' on typical intervals in \mathcal{N}_+ . In this section we are concerned with upper bounding the numbers $R^m_{A'}(\mathcal{J})$ for $\mathcal{J} \in \mathcal{N}_+$. We already sketched these computations in Section 1.2. Recall that $\gamma > (\alpha - \beta)/(1 - \beta)$ is one of the parameters specified in the statement of Theorem 3.25. Write

$$\Gamma := \frac{1}{2} \cdot \frac{\alpha - \beta}{1 - \beta} + \frac{1}{2} \cdot \gamma \in ((\alpha - \beta)/(1 - \beta), \gamma),$$
(4.29)

and decompose $\mathcal{N}_+ = \mathcal{N}_+^{low} \cup \mathcal{N}_+^{high}$, where

$$\mathcal{N}^{\text{low}}_{+} := \{ \mathcal{J} \in \mathcal{N}_{+} : R^{m}_{A'}(\mathcal{J}) \leq 2^{\Gamma m |\mathcal{J}|} \} \text{ and } \mathcal{N}^{\text{high}}_{+} := \mathcal{N}_{+} \setminus \mathcal{N}^{\text{low}}_{+}.$$
(4.30)

We remark that $\Gamma \ge \gamma/2 > 0$, since $\beta \le \alpha$.

We claim that the total length of intervals in $\mathcal{N}^{\text{low}}_+$ must be reasonably large; in the sequel, these will be the only "useful" intervals for us. More precisely, if $\eta, \zeta > 0$ are sufficiently small (as chosen in (4.23)), then

$$\sum_{\mathcal{J}\in\mathcal{N}_{+}^{\text{low}}} |\mathcal{J}| \ge \frac{1}{2} [(1-\beta) - (\alpha-\beta)/\Gamma] \cdot \ell N.$$
(4.31)

Note that $(1 - \beta) - (\alpha - \beta)/\Gamma > 0$, since $\Gamma > (\alpha - \beta)/(1 - \beta)$. To prove (4.31), set

$$\mathfrak{l} := \bigcup \mathcal{N}_+ \cup \{ s \in [\ell N] : R^m_{B'}(s) = 1 \} \subset [\ell N].$$

We start by claiming that

$$|[\ell N] \setminus \mathfrak{N}| \ge (\beta_1 - \eta - 2\zeta) \cdot \ell N.$$
(4.32)

To see this, start with the estimate

$$\delta^{\eta-\beta_1} \leqslant |B'| \leqslant \prod_{s \in \mathfrak{N}} R^m_{B'}(s) \cdot \prod_{s \in [\ell N] \setminus \mathfrak{N}} 2^m = 2^{m|[\ell N] \setminus \mathfrak{N}|} \cdot \prod_{s \in \mathfrak{N}} R^m_{B'}(s).$$
(4.33)

The last factor can further be decomposed to those indices "*s*" with $R_{B'}^m(s) = 1$ (which in total contribute "1" to the product), and then a product over the intervals $\mathcal{J} \in \mathcal{N}_+$:

$$\prod_{s \in \mathfrak{N}} R_{B'}^m(s) = \prod_{\mathcal{J} \in \mathcal{N}_+} R_{B'}^m(\mathcal{J}) \stackrel{(4.28)}{\leqslant} \prod_{\mathcal{J} \in \mathcal{N}_+} 2^{m(2\zeta)|\mathcal{J}|} \leqslant 2^{m(2\zeta)\ell N}.$$
(4.34)

Recalling that $\delta = 2^{-\ell m N}$, and combining (4.33)-(4.34), leads to

$$2^{m|[\ell N] \setminus \mathfrak{N}|} \ge 2^{\ell m N(\beta_1 - \eta - 2\zeta)}.$$

which is equivalent to (4.32).

We continue with the proof of (4.31). Note that, by the $(\eta, m, \ell N)$ -polarisation of (A', B'), we have $R^m_{A'}(s) \ge 2^{(1-\eta)m}$ for all $m \in [\ell N] \setminus \mathfrak{N} \subset \{s \in [\ell N] : R^m_{B'}(s) > 1\}$. Consequently,

$$\delta^{-\alpha} \ge |A'| = \prod_{s \in \mathfrak{N}} R^m_{A'}(s) \cdot \prod_{s \in [\ell N] \setminus \mathfrak{N}} 2^{(1-\eta)m} \ge 2^{(1-\eta)m|[\ell N] \setminus \mathfrak{N}|} \cdot \prod_{\mathcal{J} \in \mathcal{N}_+} R^m_{A'}(\mathcal{J}).$$
(4.35)

For the first factor, we will derive a lower bound from (4.32). Regarding the second factor, recall the high and low branching families from (4.30), and write $H := \bigcup \mathcal{N}^{\text{high}}_{+}$. Then,

$$\prod_{\mathcal{J}\in\mathcal{N}_+} R^m_{A'}(\mathcal{J}) \ge \prod_{\mathcal{J}\in\mathcal{N}^{\text{high}}_+} 2^{\Gamma m|\mathcal{J}|} = 2^{\Gamma m|H|}$$

Consequently, combining (4.35) with (4.32) and the estimate above, we find that

$$2^{\alpha \ell m N} \ge 2^{(1-\eta)(\beta_1 - \eta - 2\zeta)\ell m N} \cdot 2^{\Gamma m |H|}.$$

or equivalently

$$|H| \leq \frac{\alpha - (1 - \eta)(\beta_1 - \eta - 2\zeta)}{\Gamma} \cdot \ell N$$

If $\eta, \zeta = 0$, then we would have just shown that $|H| \leq \Gamma^{-1}(\alpha - \beta_1)\ell N$. Since, on the other hand, the intervals in \mathcal{N}_+ have total length at least $(1 - \beta_1)\ell N$ by (4.27) (still assuming $\eta, \zeta = 0$), we may conclude that the intervals in $\mathcal{N}_+^{\text{low}}$ have total length at least $[(1 - \beta_1) - (\alpha - \beta_1)/\Gamma] \cdot \ell N$. Finally, if the parameters η, ζ are chosen appropriately, more precisely as in (4.23), then the slightly weaker estimate (4.31) holds, namely

$$\sum_{\mathcal{J}\in\mathcal{N}_+^{\text{low}}} |\mathcal{J}| \ge \frac{1}{2} \cdot \left[(1-\beta_1) - (\alpha-\beta_1)/\Gamma \right] \cdot \ell N \ge \frac{1}{2} \cdot \left[(1-\beta) - (\alpha-\beta)/\Gamma \right] \cdot \ell N.$$

The final inequality only uses $\beta \leq \beta_1$ and $\Gamma \leq \gamma \leq 1$.

4.7. **Pruning** B' to improve separation II. Fix an interval $\mathcal{J} = \{t-r, t-r+1, \ldots, t\} \in \mathcal{N}_+$, as defined above (4.28). Then, (4.28) means that if $I \in \mathcal{I}_{m(t-r)}(B')$ is a fixed interval of length $2^{-m(t-r)}$ intersecting B', we have $2^{\zeta m r} \leq |\{J \in \mathcal{I}_{m(t+1)}(B') : J \subset I\}| \leq 2^{(2\zeta)mr}$. How well are these intervals $J \subset I$ separated? By (4.17), we already know that any two distinct intervals in $\mathcal{I}_{m(t+1)}(B')$ are separated by at least $2^{-m(t+1)}$, but this is far too weak for our purposes: for purposes to become apparent later, we would like the intervals J to be closer to $2^{-m(t-r)}$ -separated, and the only control for "r" we have is the lower bound $r \geq \ell$ (recalling that each interval in \mathcal{N}_+ contains an interval in $\ell \mathcal{N}_{B'}$).

The better separation is "morally true" for the following reason: the interval \mathcal{J} was created by combining levels with *almost* trivial branching, until *roughly* the first moment we saw some non-trivial branching. If the words "almost" and "roughly" could be omitted, we would be done: then each interval $I \in \mathcal{I}_{m(t+1)}(B')$ would be a "single child" of its parent in $\mathcal{I}_{m(t-r)}(B')$, and since the intervals in $\mathcal{I}_{m(t-r)}(B')$ are $2^{-m(t-r)}$ -separated by (4.17), the same would be true of the intervals in $\mathcal{I}_{m(t+1)}(B')$.

The words "almost" and "roughly" cannot be omitted, so we need to force the separation by trimming B' to a further $(\eta, m, \ell N)$ -uniform subset $B'' \subset B'$. Write

$$\xi := (\gamma - \Gamma)/4, \tag{4.36}$$

where Γ was defined in (4.29). In particular, $\xi \gtrsim_{\alpha,\beta,\gamma} 1$. We also impose the following additional condition on the constant $\ell \in \mathbb{N}$ selected at (4.10):

$$\ell \geqslant \xi^{-1} = \frac{4}{\gamma - \Gamma}.\tag{4.37}$$

Recall that $\mathcal{J} = \{t - r, \dots, t\} \in \mathcal{N}_+$ was the shortest extension (to the left) of a certain interval $\mathcal{J}_0 \in \ell \mathcal{N}_{B'}$ with the property $R_{B'}^m(\mathcal{J}) \ge 2^{\zeta m |\mathcal{J}|} = 2^{\zeta m (r+1)}$. Consequently, the

subinterval $\mathcal{J}_{\xi} = \{t - \lfloor (1 - 2\xi)r \rfloor, \dots, t\}$ does not yet have this property, that is,

$$R_{B'}^m(\mathcal{J}_{\xi}) < 2^{\zeta m |\mathcal{J}_{\xi}|} = 2^{\zeta m(\lfloor (1-2\xi)r \rfloor + 1)}.$$
(4.38)

Here we used that $\lfloor (1 - 2\xi)r \rfloor < r$, which is true because $\xi r \ge \xi \ell \ge 1$ by (4.37). Now, it follows from a combination of (4.38), and $R_{B'}^m(\mathcal{J}) \ge 2^{\zeta m|\mathcal{J}|} = 2^{\zeta m(r+1)}$, that

$$R_{B'}^m(\mathcal{J} \setminus \mathcal{J}_{\xi}) \ge 2^{\zeta m(r-\lfloor (1-2\xi)r \rfloor)} \ge 2^{2\xi\zeta mr} \ge 2^{\xi\zeta m(r+1)} = 2^{\xi\zeta m|\mathcal{J}|}.$$
(4.39)

We are then prepared to define the desired subset $B'' \subset B'$. Let $S_{\xi} := \bigcup \{\mathcal{J}_{\xi} : \mathcal{J} \in \mathcal{N}_+\}$, and apply the "collapsing" Lemma 4.9 to the $(m, \ell N)$ -uniform set B', and the set of scales $S_{\xi} \subset [\ell N]$. The product is an $(m, \ell N)$ -uniform subset $B'' \subset B'$ such that

$$R^m_{B''}(s) = \begin{cases} R^m_{B'}(s), & s \notin \mathcal{S}_{\xi}, \\ 1, & s \in \mathcal{S}_{\xi}. \end{cases}$$

In particular, $R_{B''}^m(s) = R_{B'}^m(s)$ for all $s \in \mathcal{J} \setminus \mathcal{J}_{\xi}$, for $\mathcal{J} \in \mathcal{N}_+$, so (4.39) remains valid for the set B'':

$$R^m_{B''}(\mathcal{J}) \ge R^m_{B'}(\mathcal{J} \setminus \mathcal{J}_{\xi}) \ge 2^{\xi \zeta m |\mathcal{J}|}.$$
(4.40)

In fact, the first inequality is an equation, since $R_{B''}^m(s) = 1$ for all $s \in \mathcal{J}_{\xi} \subset \mathcal{S}_{\xi}$. Curiously, we will have no use for a "global" lower bound for |B''|, although it would be easy to deduce from (4.28) that $|B''| \ge \delta^{2\zeta} \cdot |B'|$. From now on, only the "local" branching estimate (4.40) will be needed, and "global" lower bound $|B'| \ge \delta^{-\beta}$ has already been fully exploited in previous sections (where the relation between γ , α and β appeared).

The point of reducing B' to B'' was to improve the $2^{-m(t+1)}$ -separation of distinct intervals $I \in \mathcal{I}_{m(t+1)}(B')$ to something resembling $2^{-m(t-r)}$ -separation. This has now been accomplished. More precisely, assume that $\mathcal{J} = \{t - r, \ldots, t\} \in \mathcal{N}_+$, let $I \in \mathcal{I}_{m(t-r)}(B'')$, and and let $I_1, I_2 \in \mathcal{I}_{m(t+1)}(B'')$ be distinct. Then, since

$$R^m_{B''}(s) = 1, \qquad s \in \mathcal{J}_{\xi} = \{t - \lfloor (1 - \xi)r \rfloor, \dots, t\},\$$

the intervals I_1, I_2 are contained inside distinct intervals

$$I_1, I_2 \in \mathcal{I}_{m(t-\lfloor (1-\xi)r \rfloor)}(B'') \subset \mathcal{I}_{m(t-\lfloor (1-\xi)r \rfloor)}(B').$$

Consequently, using also that $\lfloor (1 - \xi)r \rfloor \ge (1 - \xi)r - 1$, and $\xi r \ge \xi \ell \ge 1$,

$$dist(I_1, I_2) \ge dist(\hat{I}_1, \hat{I}_2) \stackrel{(4.17)}{\ge} 2^{-m(t - \lfloor (1 - \xi)r \rfloor)}$$
$$\ge 2^{-\xi mr - m} \cdot 2^{-m(t - r)}$$
$$\ge 2^{-2\xi m(r+1)} \cdot 2^{-m(t - r)}. \tag{4.41}$$

Inequality (4.41) is more clearly phrased in the following way:

Lemma 4.42. Let $\mathcal{J} = \{t - r, \ldots, t\} \in \mathcal{N}_+, \Delta_{\mathcal{J}} := 2^{-m(t-r)}$, and $\delta_{\mathcal{J}} := 2^{-m(t+1)}$. Let $I \in \mathcal{I}_{m(t-r)}(B'')$ be a dyadic interval of length $\Delta_{\mathcal{J}}$ intersecting B'', and let $I_1, I_2 \in \mathcal{I}_{m(t+1)}(B'')$ be distinct with $I_1, I_2 \subset I$. Then,

dist
$$(I_1, I_2) \ge \left(\frac{\delta_{\mathcal{J}}}{\Delta_{\mathcal{J}}}\right)^{2\xi} \cdot |\Delta_{\mathcal{J}}|$$

Proof. Observing that $\delta_{\mathcal{J}}/\Delta_{\mathcal{J}} = 2^{-m(r+1)}$, this inequality is just a rewording of (4.41).

4.8. Elementary projection estimates. The plan is to prove lower bounds for $|A' + cB''|_{\delta}$ by, roughly speaking, establishing separately lower bounds for $|(A' \cap I) + c(B'' \cap J)|_{\delta}$, where $I, J \subset [0, 1)$ are suitable dyadic intervals intersecting A', B'', and then combining the results. In this section, we will prove an auxiliary result which will imply the required lower bounds for $|(A' \cap I) + c(B'' \cap J)|_{\delta}$. To be more accurate, instead of proving lower bounds for $|(A' \cap I) + c(B'' \cap J)|_{\delta}$ directly, we prove (stronger) lower bounds for the entropies of suitable measures supported on $(A' \cap I) + c(B'' \cap J)$ (see (4.58)). This is (only!) done for the reason that such "multi-scale" information about entropy is cleaner to combine than "multi-scale" information about cardinalities.

We introduce the following notation. Dyadic cubes in \mathbb{R}^d of side-length 2^{-n} are denoted \mathcal{D}_n . If μ is a Borel probability measure on \mathbb{R}^d , and $n \in \mathbb{N}$, we write

$$\mu^{(n)} := \sum_{Q \in \mathcal{D}_n} \frac{\mu(Q)}{\mathcal{L}^d(Q)} \cdot \mathcal{L}^d|_Q.$$

Thus $\mu^{(n)}$ is a "2⁻ⁿ-discretisation of μ ". Note that $\mu^{(n)} \in L^2(\mathbb{R}^d) \cap L^\infty(\mathbb{R}^d)$. We also define the projections $\pi_c(x, y) := x + cy$ for $(x, y) \in \mathbb{R}^2$ and $c \in \mathbb{R}$.

Lemma 4.43. Let $\Delta = 2^{-n} \in 2^{-\mathbb{N}}$, and let $\gamma, \gamma_A, \gamma_B \in (0, 1]$, and $\mathbb{C} \ge 1$. Let $\mathcal{A}, \mathcal{B} \subset \mathcal{D}_n$ be collections of dyadic Δ -intervals with $|\mathcal{A}| = \Delta^{-\gamma_A}$ and $|\mathcal{B}| = \Delta^{-\gamma_B}$. We assume the following separation from \mathcal{B} , for some $\xi \in (0, 1]$:

dist
$$(I_1, I_2) \ge \Delta^{\xi}$$
 for distinct $I_1, I_2 \in \mathcal{B}$. (4.44)

Let

- Let μ be a probability measure with spt $\mu \subset (\cup \mathcal{A}) \times (\cup \mathcal{B})$ with the property that $\mu(Q) \leq C\Delta^{\gamma_A + \gamma_B}$ for $Q \in \mathcal{D}_n$.
- Let ν be a probability measure on [-1,1] such that $\nu(I) \leq \mathbb{C}\Delta^{\gamma}$ for all $I \in \mathcal{D}_n$.

Then,

$$\int_{-1}^{1} \|(\pi_c \mu)^{(n)}\|_2^2 \, d\nu(c) \lesssim \mathbf{C} \cdot \max\{\Delta^{\gamma_A + \gamma_B - 1}, \Delta^{\gamma - 1 - \xi}\}.$$
(4.45)

Remark 4.46. The help interpreting the upper bound (4.45), let us mention the "trivial" estimate $\|(\pi_c \mu)^{(n)}\|_2^2 \leq \Delta^{\gamma_A - 1}$ for **every** $c \in [0, 1)$. This could be deduced rather easily from (4.47) below. Therefore, (4.45) beats the trivial bound whenever $\gamma > \gamma_A + \xi$.

Proof of Lemma 4.43. By definition,

$$(\pi_c \mu)^{(n)} = \sum_{I \in \mathcal{D}_n} \frac{\pi_c \mu(I)}{\Delta} \cdot \mathcal{L}^1|_I = \sum_{I \in \mathcal{D}_n} \frac{\mu(\pi_c^{-1}(I))}{\Delta} \cdot \mathcal{L}^1|_I. \qquad c \in [-1, 1].$$

Consequently,

$$\|(\pi_c \mu)^{(n)}\|_2^2 = \sum_{I \in \mathcal{D}_n} \left(\frac{\mu(\pi_c^{-1}(I))}{\Delta}\right)^2 \cdot \Delta = \frac{1}{\Delta} \cdot \sum_{I \in \mathcal{D}_n} (\mu \times \mu)(\{(p,q) : p, q \in \pi_c^{-1}(I)\}).$$
(4.47)

Therefore,

$$\begin{split} \Delta \cdot \int_{-1}^{1} \|(\pi_c \mu)^{(n)}\|_2^2 \, d\nu(c) &\sim \int_{-1}^{1} \sum_{I \in \mathcal{D}_n} (\mu \times \mu) (\{(p,q) : p, q \in \pi_c^{-1}(I)\}) \, d\nu(c) \\ &= \iint \int \sum_{I \in \mathcal{D}_n} \mathbf{1}_{\{p,q \in \pi_c^{-1}(I)\}}(c) \, d\nu(c) \, d\mu(p) \, d\mu(q). \end{split}$$

We split the outer integration into

$$\Omega_{\text{near}} := \{(p,q) : |p-q| < 10\Delta\} \text{ and } \Omega_{\text{far}} := \{(p,q) : |p-q| \ge 10\Delta\}.$$

Regarding Ω_{near} , we only use the observation that if $p, q \in A \times B$ and $c \in [0, 1]$ are fixed, then there is at most one interval $I \in \mathcal{D}_n$ such that $p, q \in \pi_c^{-1}(I)$. Since μ, ν are probability measures, and $\mu(B(x, 10\Delta)) \lesssim \mathbf{C}\Delta^{\gamma_A + \gamma_B}$ for every $x \in \mathbb{R}^2$, this leads to

$$\iint_{\Omega_{\text{near}}} \sum_{I \in \mathcal{D}_n} \mathbf{1}_{\{p, q \in \pi_c^{-1}(I)\}}(c) \, d\nu(c) \, d\mu(p) \, d\mu(q) \lesssim (\mu \times \mu)(\Omega_{\text{near}}) \lesssim \mathbf{C} \Delta^{\gamma_A + \gamma_B}.$$
(4.48)

We then consider integral over the domain Ω_{far} . A basic, easy to verify, observation is this: if $p, q \in \mathbb{R}^2$ are fixed and distinct, then the set

$$I(p,q) := \{ c \in [0,1] : p,q \in \pi_c^{-1}(I) \text{ for some } I \in \mathcal{D}_n \}$$

is contained in an interval of length $\leq \Delta/|p-q|$, and in particular can be covered by $\leq |p-q|^{-1}$ dyadic intervals of length Δ .

We combine this with the following additional observation. Note that all the tubes $\pi_c^{-1}(I)$ make an angle $\leq \pi/4$ with the *y*-axis (this is attained for c = 1, and for c = 0, the tubes $\pi_c^{-1}(I)$ are vertical). Therefore,

$$p, q \in A \times B, |p-q| \ge 10\Delta \text{ and } \exists c \in [-1, 1] \text{ s.t. } p, q \in \pi_c^{-1}(I) \implies |p_y - q_y| > \Delta.$$

Here $p_y, q_y \in B$ refer to the second coordinates of p, q. Namely, if $|p - q| \ge 10\Delta$ and $|p_y - q_y| < \Delta$, then $|p_x - q_x| \ge 9\Delta$, which makes the pair p, q too "horizontal" to be contained in any common tube $\pi_c^{-1}(I)$, with $c \in [-1, 1]$ and $I \in \mathcal{D}_n$. Now, recalling our assumption (4.44) that $\operatorname{dist}(I_1, I_2) \ge \Delta^{\xi}$ for distinct $I_1, I_2 \in \mathcal{B}$, the conclusion $|p_y - q_y| > \Delta$ can be amplified substantially: $|p_y - q_y| > \Delta$ implies that p_y, q_y lie in distinct intervals in \mathcal{B} , hence $|p - q| \ge |p_y - q_y| \ge \Delta^{\xi}$. Therefore:

$$\iint_{\Omega_{\mathrm{far}}} \int \sum_{I \in \mathcal{D}_n} \mathbf{1}_{\{p,q \in \pi_c^{-1}(I)\}}(c) \, d\nu(c) \, d\mu(p) \, d\mu(q) = \iint_{\Omega_{\mathrm{FAR}}} \int_{I(p,q)} \dots \, d\nu(c) \, d\mu(p) \, d\mu(q),$$

with $\Omega_{\text{FAR}} = \{(p,q) : |p-q| \ge \Delta^{\xi}\}$. Now, for every pair $(p,q) \in \Omega_{\text{FAR}}$, we note that the set $I(p,q) \subset [0,1]$ can be covered by $\leq |p-q|^{-1} \leq \Delta^{-\xi}$ dyadic intervals of length Δ , and for each $c \in I(p,q)$, there is exactly one $I \in \mathcal{D}_n$ such that $p, q \in \pi_c^{-1}(I)$. Therefore,

$$\int_{I(p,q)} \sum_{I \in \mathcal{D}_n} \mathbf{1}_{\{p,q \in \pi_c^{-1}(I)\}}(c) \, d\nu(v) = \nu(I(p,q)) \lesssim \mathbf{C} \Delta^{\gamma-\xi}, \qquad (p,q) \in \Omega_{\mathrm{FAR}}$$

and consequently

$$\iint_{\Omega_{\text{far}}} \int \sum_{I \in \mathcal{D}_n} \mathbf{1}_{\{p, q \in \pi_c^{-1}(I)\}}(c) \, d\nu(c) \, d\mu(p) \, d\mu(q) \lesssim \mathbf{C} \Delta^{\gamma - \xi}.$$

Combining this estimate with (4.48), we arrive at (4.45).

30

We will next deduce, as a corollary, an entropy version of Lemma 4.43. For this purpose, we record the following [33, Lemma 3.6] by Shmerkin:

Lemma 4.49. Let μ be a Borel probability measure on \mathbb{R}^d . The following relation holds between the \mathcal{D}_n -entropy $H(\mu, \mathcal{D}_n)$ of μ , and the L^2 -norm of $\mu^{(n)}$:

$$H(\mu, \mathcal{D}_n) \ge dn - \log \|\mu^{(n)}\|_2^2.$$
 (4.50)

Here, and below, "log" refers to logarithm in base 2.

Corollary 4.51. Let $\Delta = 2^{-n} \in 2^{-\mathbb{N}}$, and assume that $\mathcal{A}, \mathcal{B}, \mu, \nu, \gamma_A, \gamma_B, \gamma, \mathbb{C}$, and ξ have the same meaning as in Lemma 4.43. Then,

$$\int_{-1}^{1} H(\pi_c \mu, \mathcal{D}_n) \, d\nu(c) \ge n \cdot \min\{\gamma_A + \gamma_B, \gamma - \xi\} - \log \mathbf{C} - \log C_0, \tag{4.52}$$

where $C_0 > 0$ is an absolute constant.

Proof. First combine (4.50) (with d = 1) and Jensen's inequality to deduce that

$$\int_{-1}^{1} H(\pi_{c}\mu, \mathcal{D}_{n}) \, d\nu(c) \ge n - \int_{-1}^{1} \log \|(\pi_{c}\mu)^{(n)}\|_{2}^{2} \, d\nu(c) \ge n - \log \left(\int_{0}^{1} \|(\pi_{c}\mu)^{(n)}\|_{2}^{2} \, d\nu(c)\right).$$

Here,

Н

$$\int_{-1}^{1} \|(\pi_c \mu)^{(n)}\|_2^2 d\nu(c) \leq C_0 \mathbf{C} \max\{2^{n(1-\gamma_A - \gamma_B)}, 2^{n(1+\xi-\gamma)}\}$$

for some absolute constant $C_0 > 0$, by Lemma 4.43. These inequalities give (4.52).

4.9. **Projecting pieces of** $A' \times B''$. We next put Corollary 4.51 to work in our "real-world" situation. We introduce the following notation. Assume that μ is a Borel probability measure on \mathbb{R}^d (we will use this for both d = 1 and d = 2), and let $Q \in \mathcal{D}_n$ be a dyadic cube of side-length 2^{-n} such that $\mu(Q) > 0$. Let $T_Q \colon Q \to [0,1)^d$ be the rescaling map with $T_Q(Q) = [0, 1)^d$. We define the measures

$$\mu_Q := \frac{1}{\mu(Q)} \cdot \mu|_Q \quad \text{and} \quad \mu^Q := T_Q \mu_Q. \tag{4.53}$$

In this section, $\mu = \mu_{A'} \times \mu_{B''}$, where $\mu_{A'}$ is the normalised counting measure on A', and $\mu_{B''}$ is the normalised counting measure on B'' (defined in Section 4.7). For $s \in [\ell N]$, we will write

$$\mathcal{D}_{ms}(\mu) := \{ I \times J : I \in \mathcal{I}_{ms}(A') \text{ and } J \in \mathcal{I}_{ms}(B'') \} = \{ Q \in \mathcal{D}_{ms} : \mu(Q) > 0 \}.$$

Fix

$$\mathcal{J} = \{t - r, \dots, t\} \in \mathcal{N}_+^{\text{low}}.$$

As defined in (4.30), this means that $R^m_{A'}(\mathcal{J}) \leq 2^{\Gamma m |\mathcal{J}|}$, where $\Gamma \in ((\alpha - \beta)/(1 - \beta), \gamma) \subset$ $(\gamma/2, \gamma)$ was the parameter specified in (4.29). For now, it is only important to remember that $\gamma - \Gamma \gtrsim_{\alpha,\beta,\gamma} 1$. Fix intervals $I_0 \in \mathcal{I}_{m(t-r)}(A')$ and $J_0 \in \mathcal{I}_{m(t-r)}(B'')$. Write

$$\mathcal{A}_{I_0} := \{ I' \in \mathcal{I}_{m(t+1)}(A') : I' \subset I_0 \} \text{ and } \mathcal{B}_{J_0} := \{ J' \in \mathcal{I}_{m(t+1)}(B'') : J' \subset J_0 \}.$$

Then

$$|\mathcal{A}_{I_0}| = R^m_{A'}(\mathcal{J}) \leqslant 2^{\Gamma m |\mathcal{J}|} \quad \text{and} \quad |\mathcal{B}_{J_0}| = R^m_{B''}(\mathcal{J}) \stackrel{(4.40)}{\geqslant} 2^{\xi \zeta m |\mathcal{J}|}.$$
(4.54)

In particular, we may write

$$R_{A'}^m(\mathcal{J}) = |\mathcal{A}_{I_0}| = 2^{\gamma_A m |\mathcal{J}|} \quad \text{and} \quad |\mathcal{B}_{J_0}| = 2^{\gamma_B m |\mathcal{J}|}$$
(4.55)

for some $0 \leq \gamma_A \leq \Gamma$ and $\gamma_B \geq \xi \zeta$. Then, write

$$Q_0 := I \times J \in \mathcal{D}_{m(t-r)}(\mu), \, n := m|\mathcal{J}| = m(r+1) \quad \text{and} \quad \Delta := 2^{-n} = \frac{2^{-m(t+1)}}{2^{-m(t-r)}} \ge \delta$$

Consider the normalised measure μ^{Q_0} , as in (4.53). The measure μ^{Q_0} is supported on a product of the form $(\cup A) \times (\cup B) = T_{Q_0}((\cup A_{I_0}) \times (\cup B_{J_0}))$, where A, B are the families of Δ -intervals obtained by normalising the intervals in A_{I_0} and B_{J_0} by a factor of $2^{m(t-r)}$. It follows from the $(m, \ell N)$ -uniformity of A' and B'' that

$$\mu^{Q_0}(Q) \leq (|\mathcal{A}||\mathcal{B}|)^{-1} = (|\mathcal{A}_{I_0}||\mathcal{B}_{J_0}|)^{-1} \stackrel{(4.55)}{=} \Delta^{\gamma_A + \gamma_B}, \qquad Q \in \mathcal{D}_n.$$

Moreover, the intervals in \mathcal{B} satisfy the following separation property by Lemma 4.42:

$$I_1, I_2 \in \mathcal{B}, I_1 \neq I_2 \implies \operatorname{dist}(I_1, I_2) \ge \Delta^{2\xi}.$$

These facts place us in a position to apply Corollary 4.51 to the measure μ^{Q_0} :

$$\int_{-1}^{1} H(\pi_c \mu^{Q_0}, \mathcal{D}_n) \, d\nu(c) \ge n \cdot \min\{\gamma_A + \gamma_B, \gamma - 2\xi\} - \log 40 - \log C_0.$$
(4.56)

The parameter " ξ " was initially chosen (see (4.36)) so that $2\xi \leq (\gamma - \Gamma)/2$. Since $\gamma_A \leq \Gamma$, this leads to

 $\gamma - 2\xi \ge \gamma_A + (\gamma - \Gamma) - 2\xi \ge \gamma_A + (\gamma - \Gamma)/2.$

Recalling also that $\gamma_B \ge \xi \zeta$ by (4.54), and $\zeta \in (0, 1)$ (see (4.22) for a reminder), we find that

 $\min\{\gamma_A + \gamma_B, \gamma - 2\xi\} \ge \min\{\gamma_A + \xi\zeta, \gamma_A + (\gamma - \Gamma)/2\} = \gamma_A + \xi\zeta.$ (4.57) Before the final conclusion, let us recall that $n = m(r+1) = m|\mathcal{J}|$, and observe that

$$\gamma_A \cdot n = \gamma_A \cdot m |\mathcal{J}| \stackrel{(4.55)}{=} \log R^m_{A'}(\mathcal{J}).$$

Therefore, (4.56)-(4.57) yield

$$\int_{-1}^{1} H(\pi_c \mu^{Q_0}, \mathcal{D}_{m|\mathcal{J}|}) d\nu(c) \ge n \cdot (\gamma_A + \xi\zeta) - \log 40 - \log C_0$$

$$= \log R_{A'}^m(\mathcal{J}) + \xi\zeta \cdot m|\mathcal{J}| - \log 40 - \log C_0$$
(4.58)

for all $\mathcal{J} = \{t - r, \dots, t\} \in \mathcal{N}^{\text{low}}_+$ and for all $Q_0 = I \times J \in \mathcal{D}_{m(t-r)}(\mu)$.

4.10. Final multiscale argument. As in the previous section, let $\mu_{A'}$ be the normalised counting measure on the set A', let $\mu_{B''}$ be the normalised counting measure on the set B'', and let $\mu = \mu_{A'} \times \mu_{B''}$. Recall also that $\mathcal{D}_{ms}(\mu) = \{Q \in \mathcal{D}_{ms} : \mu(Q) > 0\}$ for $s \in [\ell N]$. We warn the reader that the notation " \mathcal{D}_{ms} " will in this section refer to both dyadic squares in \mathbb{R}^2 , and dyadic intervals in \mathbb{R} . The meaning should always be clear from context.

The purpose fo this section is to show that there exists $c \in \operatorname{spt}(\nu)$ such that

$$\frac{1}{\ell m N} \cdot H(\pi_c \mu, \mathcal{D}_{\ell m N}) \geqslant \bar{\alpha} + \xi \zeta \cdot \frac{1}{2} [(1 - \beta) - (\alpha - \beta)/\Gamma] - 2\eta.$$
(4.59)

Here $\bar{\alpha}$ was the constant (defined in (4.1)) such that $|A| = \delta^{-\bar{\alpha}}$. The lower bound in (4.59) yields a lower bound for $|A' + cB''|_{\delta}$, and consequently $|A + cB|_{\delta}$: since $H(\pi_c \mu, \mathcal{D}_{\ell m N}) \leq \log |A' + cB''|_{\delta} \leq |A + cB|_{\delta}$, and $\ell m N = -\log \delta$, we deduce from (4.59) that

$$\frac{\log |A + cB|_{\delta}}{-\log \delta} \ge \bar{\alpha} + \xi \zeta \cdot \frac{1}{2} [(1 - \beta) - (\alpha - \beta)/\Gamma] - 2\eta$$

If $\eta > 0$ is sufficiently small, depending only on $\alpha, \beta, \gamma, \kappa$, this implies $|A + cB|_{\delta} \ge \delta^{-\bar{\alpha}-\eta} = \delta^{-\eta}|A|$. Of course it is important here that the values of $\xi = (\gamma - \Gamma)/4$ (see (4.36)) and $\zeta > 0$ (see (4.22)) are independent of $\bar{\alpha}$, although they may depend on α . This proves Theorem 3.25: either (4.2) fails, and $|A + cB|_{\delta} \ge \delta^{-\epsilon}|A|$ with $c = 1 \in \operatorname{spt}(\nu)$, or (4.2) holds, and in this case $|A + cB|_{\delta} \ge \delta^{-\eta}|A|$ for the point $c \in \operatorname{spt}(\nu)$ provided by (4.59).

It remains to prove (4.59). This will be accomplished by combining (4.58) with the following uniform lower bound:

Lemma 4.60. Let
$$\mathcal{J} = \{s, \dots, t\} \subset [\ell N]$$
, and let $Q_0 \in \mathcal{D}_{ms}(\mu)$. Then,

$$H(\pi_c \mu^{Q_0}, \mathcal{D}_{m|\mathcal{J}|}) \ge \log R^m_{A'}(\mathcal{J}) - 1, \qquad c \in [0, 1].$$
(4.61)

Proof. Let $I \in \mathcal{I}_{ms}(A')$ and $J \in \mathcal{I}_{ms}(B'')$ such that $Q_0 = I \times J$. Then $\mu^{Q_0} = \mu^I_{A'} \times \mu^J_{B''}$, hence $\pi_c \mu^Q = \mu^I_{A'} * \mu^J_{B''}$, and finally

$$H(\pi_{c}\mu^{Q_{0}}, \mathcal{D}_{m|\mathcal{J}|}) = H(\mu_{A'}^{I} * \mu_{B''}^{J}, \mathcal{D}_{m|\mathcal{J}|}) \ge \int H((\mu_{A'}^{I})_{x}, \mathcal{D}_{m|\mathcal{J}|}) d\mu_{B''}^{J}(x),$$
(4.62)

where the inequality follows from the concavity of entropy (we discussed this at (2.5)), and where $(\mu_{A'})_x^I(H) := \mu_{A'}^I(H-x)$ for $H \subset \mathbb{R}$. From the definition of entropy, one has

$$H((\mu_{A'}^{I})_{x}, \mathcal{D}_{m|\mathcal{J}|}) = H(\mu_{A'}^{I}, \mathcal{D}_{m|\mathcal{J}|} - x),$$

where $\mathcal{D}_{m|\mathcal{J}|} - x$ refers to the family of (-x)-translated dyadic intervals. Now, for $x \in \mathbb{R}$ fixed, every intervals in $\mathcal{D}_{m|\mathcal{J}|}$ can be covered by 2 intervals in $\mathcal{D}_{m|\mathcal{J}|} - x$ and vice versa. This implies that

$$|H(\mu_{A'}^{I}, \mathcal{D}_{m|\mathcal{J}|} - x) - H(\mu_{A'}^{I}, \mathcal{D}_{m|\mathcal{J}|})| \leq \log 2 = 1, \qquad x \in \mathbb{R}.$$
(4.63)

Furthermore, by definition,

$$H(\mu_{A'}^{I}, \mathcal{D}_{m|\mathcal{J}|}) = -\sum_{L \in \mathcal{D}_{m|\mathcal{J}|}} \mu_{A'}^{I}(L) \log \mu_{A'}^{I}(L).$$

Since A' is $(m, \ell N, R_{A'}^m)$ -uniform, either $\mu_{A'}^I(L) = 0$, or then $\mu_{A'}^I(L) = R_{A'}^m(\mathcal{J})^{-1}$ for every $L \in \mathcal{D}_{m|\mathcal{J}|}$. Therefore

$$H(\mu_{A'}^I, \mathcal{D}_{m|\mathcal{J}}) = R_{A'}^m(\mathcal{J})^{-1}.$$

In combination with (4.62)-(4.63), this yields (4.61).

Recall the intervals $\mathcal{N}_{+}^{\text{low}} \subset \mathcal{N}_{+}$, defined in (4.30). In this section, the properties of these intervals will be used via the formula (4.58), and we additionally need to recall that

$$\sum_{\mathcal{J}\in\mathcal{N}_{+}^{\text{low}}} |\mathcal{J}| \ge \frac{1}{2} [(1-\beta) - (\alpha-\beta)/\Gamma] \cdot \ell N$$
(4.64)

by (4.31). Let \mathcal{P} be the partition of $[\ell N]$ which is induced by the intervals in $\mathcal{N}^{\text{low}}_+$. In other words, \mathcal{P} consists of the intervals in $\mathcal{N}^{\text{low}}_+$, and the maximal complementary intervals. We write

$$\mathcal{P}_{\text{useless}} := \mathcal{P} \setminus \mathcal{N}_{+}^{\text{low}},$$

and we enumerate $\mathcal{P} = \{\mathcal{J}_1, \mathcal{J}_2, \dots, \mathcal{J}_h\}$, where $1 \leq h \leq \ell N$. We write $\mathcal{J}_j = \{s_j, \dots, t_j\}$ for $1 \leq j \leq h$, so $s_1 = 0$, $t_h + 1 = \ell N$, and $s_{j+1} = t_j + 1$ for all $1 \leq j < h$. We artificially define $s_{h+1} := \ell N$, so the relation $s_{j+1} = t_j + 1$ also remains valid for j = h.

We abbreviate

$$\mathcal{D}_j := \mathcal{D}_{ms_j}(\mu) := \{ I \times J : I \in \mathcal{I}_{ms_j}(A') \text{ and } J \in \mathcal{I}_{ms_j}(B'') \}.$$

We then apply the entropy lower bound in Lemma 2.3, and its corollary (2.4), to the partition $0 = ms_1 < \ldots < ms_h < ms_{h+1} = \ell mN$ of $\{0, \ldots, \ell mN\}$, and the 2-Lipschitz maps $\pi_c \colon \mathbb{R}^2 \to \mathbb{R}$ with $c \in [-1, 1]$:

$$\int_{-1}^{1} H(\pi_{c}\mu, \mathcal{D}_{\ell m N}) \, d\nu(c) = \sum_{j=1}^{h} \sum_{Q \in \mathcal{D}_{j}} \mu(Q) \int_{-1}^{1} H(\pi_{c}\mu^{Q}, \mathcal{D}_{ms_{j+1}-ms_{j}} \mid \mathcal{D}_{0}) \, d\nu(c)$$

$$\geq -C_{0}h + \sum_{j=1}^{h} \sum_{Q \in \mathcal{D}_{j}} \mu(Q) \int_{-1}^{1} H(\pi_{c}\mu^{Q}, \mathcal{D}_{m(t_{j}-s_{j}+1)}) \, d\nu(c).$$
(4.65)

Above, $t_j - s_j + 1 = |\mathcal{J}_j|$. For $\mathcal{J}_j \in \mathcal{N}_+^{\text{low}}$, and $Q \in \mathcal{D}_j$, we recall from (4.58) that

$$\int_{-1}^{1} H(\pi_c \mu^Q, \mathcal{D}_m|\mathcal{J}_j|) \, d\nu(c) \ge \log R^m_{A'}(\mathcal{J}_j) + \xi \zeta \cdot m|\mathcal{J}_j| - \log 40 - \log C_0$$

For $\mathcal{J} \in \mathcal{P}_{useless}$ we have to settle with the estimate

$$\int_0^1 H(\pi_c \mu^Q, \mathcal{D}_{m|\mathcal{J}_j|}) \, d\nu(c) \ge \log R^m_{A'}(\mathcal{J}_j) - 1$$

from Lemma 4.60. Plugging these bounds into (4.65) (and redefining C_0 as $C_0 + 1$) yields

$$\int_{-1}^{1} H(\pi_{c}\mu, \mathcal{D}_{\ell m N}) d\nu(c) \geq -(40 + C_{0})h + \sum_{\mathcal{J} \in \mathcal{P}} \log R_{A'}^{m}(\mathcal{J}) + \xi\zeta \sum_{\mathcal{J} \in \mathcal{N}_{+}^{\mathrm{low}}} |\mathcal{J}|$$

$$\stackrel{(4.64)}{\geq} \log |A'| + \xi\zeta \cdot \frac{1}{2} [(1 - \alpha) - (\alpha - \beta)/\Gamma] \cdot \ell m N - h(40 + C_{0})$$

Recalling that $|A'| \ge \delta^{\eta} |A| \ge 2^{(\bar{\alpha} - \eta)\ell mN}$, there exists $c \in \operatorname{spt}(\nu)$ with

$$H_{\ell m N}(\pi_c \mu) \ge \left(\bar{\alpha} + \xi \zeta \cdot \frac{1}{2} \left[(1 - \alpha) - (\alpha - \beta) / \Gamma \right] - \eta \right) - \frac{h(40 + C_0)}{\ell m N}$$

Here $h(40 + C_0)/(\ell m N) \le (40 + C_0)/m_0 \le \eta$ by the choice of m_0 at (4.10), and since we chose $m \ge m_0$ in Proposition 4.12. Therefore we have established (4.59), and completed the proof of Theorem 3.25.

References

- [1] Yves Benoist and Nicolas de Saxcé. A spectral gap theorem in simple Lie groups. *Invent. Math.*, 205(2):337–361, 2016.
- [2] J. Bourgain. On the Erdös-Volkmann and Katz-Tao ring conjectures. *Geom. Funct. Anal.*, 13(2):334–365, 2003.
- [3] J. Bourgain and A. Gamburd. A spectral gap theorem in SU(d). J. Eur. Math. Soc. (JEMS), 14(5):1455– 1511, 2012.
- [4] Jean Bourgain. Multilinear exponential sums in prime fields under optimal entropy condition on the sources. *Geom. Funct. Anal.*, 18(5):1477–1502, 2009.
- [5] Jean Bourgain. The discretized sum-product and projection theorems. J. Anal. Math., 112:193–236, 2010.
- [6] Jean Bourgain and Alex Gamburd. On the spectral gap for finitely-generated subgroups of SU(2). *Invent. Math.*, 171(1):83–121, 2008.
- [7] Damian Dabrowski, Tuomas Orponen, and Michele Villa. Integrability of orthogonal projections, and applications to Furstenberg sets. *arXiv e-prints*, page arXiv:2107.04471, July 2021.

- [8] P. Erdős and E. Szemerédi. On sums and products of integers. In *Studies in pure mathematics*, pages 213–218. Birkhäuser, Basel, 1983.
- [9] Yuqiu Fu, Shengwen Gan, and Kevin Ren. An incidence estimate and a Furstenberg type estimate for tubes in ℝ². arXiv e-prints, page arXiv:2107.11937, July 2021.
- [10] M. Z. Garaev. An explicit sum-product estimate in \mathbb{F}_p . Int. Math. Res. Not. IMRN, (11):Art. ID rnm035, 11, 2007.
- [11] A. A. Glibichuk and S. V. Konyagin. Additive properties of product sets in fields of prime order. In Additive combinatorics, volume 43 of CRM Proc. Lecture Notes, pages 279–286. Amer. Math. Soc., Providence, RI, 2007.
- [12] Larry Guth, Nets Hawk Katz, and Joshua Zahl. On the discretized sum-product problem. *Int. Math. Res. Not. IMRN*, (13):9769–9785, 2021.
- [13] Larry Guth, Noam Solomon, and Hong Wang. Incidence estimates for well spaced tubes. Geom. Funct. Anal., 29(6):1844–1863, 2019.
- [14] Katalin Gyarmati, Máté Matolcsi, and Imre Z. Ruzsa. Plünnecke's inequality for different summands. In *Building bridges*, volume 19 of *Bolyai Soc. Math. Stud.*, pages 309–320. Springer, Berlin, 2008.
- [15] Weikun He. Discretized sum-product estimates in matrix algebras. J. Anal. Math., 139(2):637–676, 2019.
- [16] Weikun He. Orthogonal projections of discretized sets. J. Fractal Geom., 7(3):271–317, 2020.
- [17] Weikun He and Nicolas de Saxcé. Sum-product for real Lie groups. J. Eur. Math. Soc. (JEMS), 23(6):2127– 2151, 2021.
- [18] Michael Hochman. On self-similar sets with overlaps and inverse theorems for entropy. *Ann. of Math.* (2), 180(2):773–822, 2014.
- [19] Robert Kaufman. On Hausdorff dimension of projections. Mathematika, 15:153–155, 1968.
- [20] Jialun Li. Discretized Sum-product and Fourier decay in \mathbb{R}^n . J. Anal. Math., 143(2):763–800, 2021.
- [21] J. M. Marstrand. Some fundamental geometrical properties of plane sets of fractional dimensions. *Proc. London Math. Soc.* (3), 4:257–302, 1954.
- [22] P. Mattila. *Geometry of sets and measures in Euclidean spaces. Fractals and rectifiability. 1st paperback ed.* Cambridge: Cambridge University Press, 1st paperback ed. edition, 1999.
- [23] Ali Mohammadi and Sophie Stevens. Attaining the exponent 5/4 for the sum-product problem in finite fields. *arXiv e-prints*, page arXiv:2103.08252, March 2021.
- [24] Daniel M. Oberlin. Some toy Furstenberg sets and projections of the four-corner Cantor set. Proc. Amer. Math. Soc., 142(4):1209–1215, 2014.
- [25] Tuomas Orponen. On arithmetic sums of Ahlfors-regular sets. Geom. Funct. Anal. (to appear).
- [26] Tuomas Orponen. On the distance sets of Ahlfors-David regular sets. Adv. Math., 307:1029–1045, 2017.
- [27] Tuomas Orponen and Pablo Shmerkin. On the Hausdorff dimension of Furstenberg sets and orthogonal projections in the plane. *arXiv e-prints*, page arXiv:2106.03338, June 2021.
- [28] Tuomas Orponen and Laura Venieri. A note on expansion in prime fields. *arXiv e-prints*, page arXiv:1801.09591, January 2018.
- [29] Orit E. Raz and Joshua Zahl. On the dimension of exceptional parameters for nonlinear projections, and the discretized Elekes-Rónyai theorem. *arXiv e-prints*, page arXiv:2108.07311, August 2021.
- [30] Misha Rudnev and Sophie Stevens. An update on the sum-product problem. *arXiv e-prints*, page arXiv:2005.11145, May 2020.
- [31] Imre Z. Ruzsa. An application of graph theory to additive number theory. *Sci. Ser. A Math. Sci. (N.S.)*, 3:97–109, 1989.
- [32] Pablo Shmerkin. On Furstenberg's intersection conjecture, self-similar measures, and the L^q norms of convolutions. *Ann. of Math.* (2), 189(2):319–391, 2019.
- [33] Pablo Shmerkin. On the Hausdorff dimension of pinned distance sets. *Israel J. Math.*, 230(2):949–972, 2019.
- [34] Pablo Shmerkin. A nonlinear version of bourgain's projection theorem. Preprint, arXiv:2003.01636, 2020.
- [35] Sophie Stevens and Frank de Zeeuw. An improved point-line incidence bound over arbitrary fields. *Bull. Lond. Math. Soc.*, 49(5):842–858, 2017.
- [36] Endre Szemerédi and William T. Trotter, Jr. Extremal problems in discrete geometry. Combinatorica, 3(3-4):381–392, 1983.
- [37] Terence Tao and Van Vu. Additive combinatorics, volume 105 of Cambridge Studies in Advanced Mathematics. Cambridge University Press, Cambridge, 2006.

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF JYVÄSKYLÄ, P.O. BOX 35 (MAD), FI-40014 UNIVERSITY OF JYVÄSKYLÄ, FINLAND *Email address*: tuomas.t.orponen@jyu.fi