Bounds for the Twin-width of Graphs

Jungho Ahn^{1,2}, Kevin Hendrey², Donggyu Kim^{1,2}, and Sang-il Oum^{2,1}

¹Department of Mathematical Sciences, KAIST,

Daejeon, South Korea

²Discrete Mathematics Group, Institute for Basic Science (IBS), Daejeon, South Korea

Email addresses: junghoahn@kaist.ac.kr, kevinhendrey@ibs.re.kr, donggyu@kaist.ac.kr, sangil@ibs.re.kr

October 8, 2021

Abstract

Bonnet, Kim, Thomassé, and Watrigant [8] introduced the twinwidth of a graph. We show that the twin-width of an *n*-vertex graph is less than $(n + \sqrt{n \ln n} + \sqrt{n} + 2 \ln n)/2$, and the twin-width of an *m*-edge graph is less than $\sqrt{3m} + m^{1/4}\sqrt{\ln m}/(4 \cdot 3^{1/4}) + 3m^{1/4}/2$. Conference graphs of order *n* (when such graphs exist) have twinwidth at least (n - 1)/2, and we show that Paley graphs achieve this lower bound. We also show that the twin-width of the Erdős-Rényi random graph G(n,p) with $1/n \leq p = p(n) \leq 1/2$ is larger than $2p(1-p)n - (2\sqrt{2}+\varepsilon)\sqrt{p(1-p)n \ln n}$ asymptotically almost surely for any positive ε . Lastly, we calculate the twin-width of random graphs G(n,p) with $p \leq c/n$ for a constant c < 1, determining the thresholds at which the twin-width jumps from 0 to 1 and from 1 to 2.

1 Introduction

Bonnet, Kim, Thomassé, and Watrigant [8] introduced a new width parameter, called the *twin-width*, of a graph G, denoted by tww(G). The twin-width of a graph G captures a way of reducing G into a one-vertex graph by iteratively contracting pairs of vertices with similar neighborhoods (see Section 2.1 for the formal definiton). It has been shown that every

²All authors are supported by the Institute for Basic Science (IBS-R029-C1).

graph class of bounded twin-width is *small*³ [4], χ -bounded [5], and admits a linear-time algorithm for first-order model checking [8] (if we are given a certificate for an input graph to have small twin-width). Despite being a relatively new concept, twin-width has already generated a large amount of interests [8, 4, 5, 6, 13, 7, 1, 10, 17, 9].

Twin-width can be seen as a generalization of both tree-width and rankwidth in the sense that graph classes of bounded tree-width or rank-width also have bounded twin-width [8]. However, there are graph classes that have bounded twin-width and unbounded tree-width or rank-width. For instance, the $n \times n$ grid has tree-width n and rank-width n - 1 [15], but twin-width at most 4 [8, 3]. Other interesting classes of bounded twinwidth include proper minor-closed classes, map graphs, and K_t -free unit d-dimensional ball graphs [8, 4].

Given a graph parameter, it is natural to ask for the highest possible value of the parameter. The maximum tree-width of an *n*-vertex graph is equal to n-1. By definition, the rank-width of an *n*-vertex graph is bounded above by $\lfloor n/3 \rfloor$, and Lee, Lee, and Oum $\lfloor 16 \rfloor$ showed that random graphs G(n,p) with a constant $p \in (0,1)$ is of rank-width $\lfloor n/3 \rfloor - O(1)$ asymptotically almost surely. One can observe from definition that the twin-width of an *n*-vertex graph is at most n-2, and currently no nontrivial upper bound for twin-width has been studied.

Here are our main results. Firstly, we present an upper bound for the twin-width of an n-vertex graph.

Theorem 1.1. For a positive integer n and an n-vertex graph G,

$$\operatorname{tww}(G) < \frac{1}{2}(n + \sqrt{n \ln n} + \sqrt{n} + 2 \ln n).$$

Secondly, we present an upper bound for the twin-width of a graph G in terms of its number of edges. This refines the bound in Theorem 1.1 when G has at most $|V(G)|^2/12$ edges.

Theorem 1.2. For a positive integer m and an m-edge graph G,

$$\operatorname{tww}(G) < \sqrt{3m} + \frac{m^{1/4}\sqrt{\ln m}}{4 \cdot 3^{1/4}} + \frac{3m^{1/4}}{2}.$$

Thirdly, we present an asymptotic lower bound for the twin-width of random graphs.

³A graph class C is *small* if C has at most $n!c^n$ graphs labeled by [n] for a constant c.

Theorem 1.3. Let G := G(n,p) be a random graph where p := p(n) is a function such that $1/n \leq p \leq 1/2$ and ε be a positive real. Then

$$\operatorname{tww}(G) > 2p(1-p)n - (2\sqrt{2} + \varepsilon)\sqrt{p(1-p)n\ln n}$$

asymptotically almost surely.

By Theorems 1.1 and 1.3 with p = 1/2,

$$\lim_{n \to \infty} \max\left\{ \frac{\operatorname{tww}(G)}{|V(G)|} : |V(G)| = n \right\} = \frac{1}{2}.$$

In terms of the number of edges, we can state the following by Theorems 1.2 and 1.3 with p = 1/3:

$$1.088 < \frac{4\sqrt{6}}{9} \leq \limsup_{m \to \infty} \max\left\{\frac{\operatorname{tww}(G)}{\sqrt{|E(G)|}} : |E(G)| = m\right\} \leq \sqrt{3} < 1.733.$$

It would be interesting to determine this value exactly.

The best lower bound we have for the maximum twin-width of an *n*-vertex graph comes from conference graphs and is better than the lower bound for random graphs from Theorem 1.3. As we will discuss in Section 3, every *n*-vertex conference graph has twin-width at least (n-1)/2. In particular, we prove that Paley graphs achieve this lower bound. It is an open problem to determine whether there is an *n*-vertex graph having twin-width at least n/2.

Theorem 1.4. For each prime power q with $q \equiv 1 \pmod{4}$, the Paley graph P(q) has twin-width exactly (q-1)/2.

We calculate the twin-width of random graphs G(n, p) with $p \leq c/n$ for a constant c < 1. This determines the thresholds at which the twin-width jumps from 0 to 1 and from 1 to 2.

Theorem 1.5. For a random graph G := G(n, p) with a function p := p(n), the following hold asymptotically almost surely.

- 1. tww(G) = 0 if $p = o(n^{-4/3})$.
- 2. tww(G) = 1 if $p = \omega(n^{-4/3})$ and $p = o(n^{-7/6})$.
- 3. tww(G) = 2 if $p = \omega(n^{-7/6})$ and $p \leq c/n$ for a constant $c \in (0, 1)$.

We organize this paper as follows. In Section 2, we present some terminology from graph theory and probability theory, and formally define twin-width. In Section 3, we determine the twin-width of Paley graphs. We prove Theorem 1.1 in Section 4 and Theorem 1.2 in Section 5. In Section 6, we prove Theorems 1.3 and 1.5.

2 Preliminaries

In this paper, all (tri)graphs are simple and finite. For a set X and a positive integer t, $\binom{X}{t}$ be the set of t-element subsets of X, $[t] := \{1, \ldots, t\}$, and $[0] := \emptyset$. For sets X and Y, the symmetric difference of X and Y, denoted by $X \triangle Y$, is the union of $X \backslash Y$ and $Y \backslash X$. A partition of X is a set $\{X_1, \ldots, X_t\}$ of pairwise disjoint nonempty subsets of X such that $X = \bigcup_{i=1}^t X_i$. For a graph G and disjoint vertex sets X and Y of G, X is complete to Y if every vertex in X is adjacent to all vertices of Y.

A trigraph is a triple G = (V, B, R) where B and R are disjoint subsets of $\binom{V}{2}$. Given a trigraph G = (V, B, R), elements in B are black edges of G and elements in R are red edges of G. We denote the set of black edges of G by B(G), the set of red edges of G by R(G), and the set of edges of G by E(G), that is, $E(G) = B(G) \cup R(G)$. For each vertex v of G, the degree of v, denoted by $\deg_G(v)$, is the number of edges of G incident with v, and the red-degree of v, denoted by $\operatorname{rdeg}_G(v)$, is the number of red edges of G incident with v. We denote by $\Delta^R(G)$ the maximum red-degree of a vertex of G. The average degree of G is 2|E(G)|/|V(G)|. For a subset A of V(G), $G \setminus A$ is a trigraph obtained from G by removing all vertices in A and all edges incident with vertices in A. Let $\delta_G(A)$ be the set of edges of G having one end in A and the other end in $V(G) \setminus A$. If $A = \{v\}$ for a vertex v of G, then we may write $G \setminus v$ and $\delta_G(v)$ instead of $G \setminus \{v\}$ and $\delta_G(\{v\})$, respectively. We identify a graph H = (V, E) with a trigraph $H = (V, E, \emptyset)$.

2.1 Twin-width

Let d be a nonnegative real. A trigraph G = (V, B, R) is a d-trigraph if $\Delta^R(G) \leq d$. For distinct vertices u and v of G, not necessarily adjacent, $G/\{u, v\} = (V', B', R')$ is a trigraph obtained from G such that $V' = (V \setminus \{u, v\}) \cup \{w\}$ for a new vertex $w, G \setminus \{u, v\} = (G/\{u, v\}) \setminus w$, and for every vertex x in $V' \setminus \{w\}$, the following hold:

- (i) $wx \in B'$ if and only if $\{ux, vx\} \subseteq B$,
- (ii) $wx \notin B' \cup R'$ if and only if $\{ux, vx\} \cap (B \cup R) = \emptyset$, and
- (iii) $wx \in R'$ otherwise.

We say that $G/\{u, v\}$ is obtained from G by contracting u and v into a new vertex w. We often identify w with either u or v. A partial contraction sequence from G_1 to G_t is a sequence G_1, \ldots, G_t of trigraphs such that for

each $i \in [t-1]$, G_{i+1} is obtained from G_i by contracting two distinct vertices. A contraction sequence of G is a partial contraction sequence from G to a one-vertex graph. A *(partial) d-contraction sequence* is a (partial) contraction sequence such that every trigraph in the sequence is a *d*-trigraph. The *twin-width* of G, denoted by tww(G), is the minimum d such that G admits a *d*-contraction sequence.

A partial contraction sequence σ from G_1 to G_t defines a function from $V(G_1)$ to $V(G_t)$, which we denote also by σ , as follows: for a vertex v of G_1 , $\sigma(v)$ is the vertex of G_t to which v is contracted. If v was never contracted, then $\sigma(v) = v$. For a subset X of $V(G_t)$, we define

$$\sigma^{-1}(X) = \{ v \in V(G_1) : \sigma(v) \in X \}.$$

For a vertex w of G_t , we may write $\sigma^{-1}(w)$ for $\sigma^{-1}(\{w\})$. In other words, $\sigma^{-1}(w)$ is the set of vertices of G_1 which are eventually contracted into w in G_t . We have the following two simple lemmas.

Lemma 2.1. Let σ be a partial contraction sequence from G to G'. Then for every vertex v of G',

$$\deg_{G'}(v) \leq |\delta_G(\sigma^{-1}(v))| \leq \sum_{w \in \sigma^{-1}(v)} \deg_G(w).$$

Proof. For each $e := vw \in \delta_{G'}(v)$, there is at least one edge f(e) of G having one end in $\sigma^{-1}(v)$ and the other end in $\sigma^{-1}(w)$. Since $\sigma^{-1}(v)$ and $\sigma^{-1}(w)$ are disjoint, f(e) is in $\delta_G(\sigma^{-1}(v))$. For $e' := vw' \in \delta_{G'}(v)$ distinct from $e, \sigma^{-1}(w)$ and $\sigma^{-1}(w')$ are disjoint, and therefore f(e) and f(e') are distinct. Thus, $f : \delta_{G'}(v) \to \delta_G(\sigma^{-1}(v))$ is an injection, and this completes the proof. \Box

The following lemma is immediate from the definition.

Lemma 2.2. Let σ be a partial contraction sequence from G to G'. If v is a vertex of G' with $|\sigma^{-1}(v)| = 1$, then $\operatorname{rdeg}_{G'}(v) \leq \operatorname{rdeg}_{G}(v) + t$, where t is the number of vertices w of G' with $|\sigma^{-1}(w)| > 1$.

2.2 Random graphs

For a positive integer n and $0 \leq p \leq 1$, the *Erdős-Rényi random graph* G := G(n, p) is a probability space over the set of graphs with vertex set [n] such that for all $i, j \in [n]$ with i < j, the event that i is adjacent to j occurs with probability p, and these events are mutually independent. For a function $p : \mathbb{N} \to [0, 1]$, we say that the random graph G(n, p(n)) has a

graph property A asymptotically almost surely if $\Pr[G(n, p(n)) \models A] \to 1$ as $n \to \infty$.

We will use the following inequality to investigate the asymptotic behavior of the twin-width of random graphs.

Lemma 2.3 (Chernoff bound; see [14, Theorem 2.1]). Let X_1, \ldots, X_n be mutually independent random variables such that for each $i \in [n]$, $\Pr[X_i = 1] = p$ and $\Pr[X_i = 0] = 1 - p$. Let $X := \sum_{i=1}^n X_i$. Then

$$\Pr[X - \mathbb{E}[X] \leq -t] \leq \exp\left(-\frac{t^2}{2np}\right)$$

for any $t \ge 0$.

2.3 Lattice paths

For points **x** and **y** in \mathbb{Z}^2 , a *lattice path* from **x** to **y** is a sequence $\mathbf{x}_1, \ldots, \mathbf{x}_k$ of points in \mathbb{Z}^2 such that $\mathbf{x}_1 = \mathbf{x}$, $\mathbf{x}_k = \mathbf{y}$, and $\mathbf{x}_{i+1} - \mathbf{x}_i \in \{(0, \pm 1), (\pm 1, 0)\}$ for each $i \in [k-1]$. A lattice path $\mathbf{x}_1, \ldots, \mathbf{x}_k$ is *North-East* if $\mathbf{x}_{i+1} - \mathbf{x}_i \in \{(0, 1), (1, 0)\}$ for each $i \in [k-1]$. We say that a lattice path $\mathbf{x}_1, \ldots, \mathbf{x}_k$ *intersects* a line y = ax + b if it intersects both $\{(x, y) : y \ge ax + b\}$ and $\{(x, y) : y \le ax + b\}$.

Proposition 2.4 (See [2, Theorem 10.3.1]). Let a, b, and t be nonnegative integers with a + t > b. Then $\binom{a+b}{a+t}$ is the number of North-East lattice paths from (0,0) to (a,b) intersecting the line y = x + t.⁴

The following corollary will help us to prove Theorem 1.1.

Corollary 2.5. Let a and b be nonnegative integers and t be a positive integer. Let A be the event that a North-East lattice path from (0,0) to (a,b) chosen uniformly at random intersects the line y = x + t. Then $\Pr[A] \leq (b/(a+t))^t$.

Proof. We may assume that $0 < t \le b$, because otherwise $\Pr[A] = 0$. We may assume that a + t > b, because otherwise $(b/(a + t))^t \ge 1$. Since $\binom{a+b}{a}$ is the number of North-East lattice paths from (0,0) to (a,b), by Proposition 2.4,

$$\Pr[A] = \frac{\binom{a+b}{a+t}}{\binom{a+b}{a}} = \frac{a!}{(a+t)!} \cdot \frac{b!}{(b-t)!} = \prod_{i=0}^{t-1} \frac{b-i}{a+t-i} \leqslant \left(\frac{b}{a+t}\right)^t,$$

where the last inequality holds because a + t > b.

⁴We remind the readers that for integers n and m with $m > n \ge 0$, $\binom{n}{m}$ is defined as 0.

3 Twin-width of Paley graphs

From the definition of the twin-width, we deduce the following lemma.

Lemma 3.1. Every n-vertex graph G with $n \ge 2$ has twin-width at least $\min\{|(N_G(v) \triangle N_G(w)) \setminus \{v, w\}| : v, w \in V(G), v \ne w\}.$

The following proposition allows us to choose a pair of vertices whose contraction yields a small red-degree.

Proposition 3.2. For an *n*-vertex graph G, every subset L of V(G) with $|L| \ge 2$ contains distinct vertices v and w such that

$$|(N_G(v) \triangle N_G(w)) \setminus \{v, w\}| \leq \frac{n}{2} + \frac{n-1}{2|L|-2} - 1.$$

Proof. Let $\ell := |L|$. The number of elements $\{u, v\}$ in $\binom{L}{2}$ with $w \in (N_G(u) \triangle N_G(v)) \setminus \{u, v\}$ is $|L \cap N_G(w)| \cdot (|L \setminus N_G(w)| - 1)$ if $w \in L$, and is $|L \cap N_G(w)| \cdot |L \setminus N_G(w)|$ otherwise. We choose a pair $\{u, v\}$ from $\binom{L}{2}$ uniformly at random. Then

$$\begin{split} \mathbb{E}[|(N_G(u) \triangle N_G(v)) \setminus \{u, v\}|] \\ &= \sum_{w \in L} \frac{|L \cap N_G(w)| \cdot (|L \setminus N_G(w)| - 1)}{\binom{\ell}{2}} + \sum_{w \in V(G) \setminus L} \frac{|L \cap N_G(w)| \cdot |L \setminus N_G(w)|}{\binom{\ell}{2}} \\ &\leq \ell \cdot \frac{\left(\frac{\ell-1}{2}\right)^2}{\binom{\ell}{2}} + (n-\ell) \cdot \frac{\left(\frac{\ell}{2}\right)^2}{\binom{\ell}{2}} = \frac{n}{2} + \frac{n-1}{2\ell-2} - 1. \end{split}$$

Thus, $\binom{L}{2}$ contains an element $\{u', v'\}$ with $|(N_G(u') \triangle N_G(v')) \setminus \{u', v'\}| \leq n/2 + (n-1)/(2\ell-2) - 1.$

By applying Proposition 3.2 with L = V(G) we obtain the following corollary which indicates that the lower bound in Lemma 3.1 is at most (n-1)/2 for an *n*-vertex graph with $n \ge 2$.

Corollary 3.3. Every n-vertex graph G with $n \ge 2$ has distinct vertices v and w such that $|(N_G(v) \triangle N_G(w)) \setminus \{v, w\}| \le (n-1)/2$.

We provide an infinite class of graphs such that every *n*-vertex graph in the class has twin-width at least (n-1)/2. A conference graph is a graph G on n vertices for a positive integer n with $n \equiv 1 \pmod{4}$ such that G is ((n-1)/2)-regular and for each pair $\{v, w\}$ of distinct vertices of G, $|N_G(v) \cap N_G(w)|$ is (n-5)/4 if v and w are adjacent, and otherwise (n-1)/4. Observe that in an *n*-vertex conference graph G, for distinct vertices v and w of G, $|(N_G(v) \triangle N_G(w)) \setminus \{v, w\}| = (n-1)/2$. Thus, by Lemma 3.1, every *n*-vertex conference graph has twin-width at least (n-1)/2.

For each prime power q with $q \equiv 1 \pmod{4}$, the Paley graph P(q) is a graph on the field \mathbb{F}_q of order q such that vertices a and b are adjacent in P(q) if and only if $a - b = c^2$ for some $c \in \mathbb{F}_q \setminus \{0\}$. We remark that -1 is a square in \mathbb{F}_q for a prime power q with $q \equiv 1 \pmod{4}$, so that the Paley graph P(q) is well defined as an undirected graph. Erdős and Rényi [12] showed that every Paley graph is a conference graph.

We now prove Theorem 1.4.

Theorem 1.4. For each prime power q with $q \equiv 1 \pmod{4}$, the Paley graph P(q) has twin-width exactly (q-1)/2.

Proof. Since every Paley graph is a conference graph, by Lemma 3.1, it suffices to show that tww(P(q)) $\leq (q-1)/2$. Let $u_1, \ldots, u_{(q-1)/2}$ be elements in $\mathbb{F}_q \setminus \{0\}$ such that for all distinct $i, j \in [(q-1)/2], u_i \neq u_j$ and $u_i \neq -u_j$. Let $G_0 := P(q)$ and for each $k \in [(q-1)/2]$, let $G_k := G_{k-1}/\{u_k, -u_k\}$. We will identify the vertex created by contracting u_k and $-u_k$ with u_k . Since $|V(G_{(q-1)/2})| = (q+1)/2$, it is enough to show that $\Delta^R(G_k) \leq (q-1)/2$ for each $k \in [(q-1)/2]$.

By Lemma 2.2, for each $u \in V(G_k) \setminus \{u_1, \ldots, u_k\}$, we have $\operatorname{rdeg}_{G_k}(u) \leq k \leq (q-1)/2$. Let $i \in [k]$ and $Q_i := G_0/\{u_i, -u_i\}$. We will again identify the new vertex created by contracting u_i and $-u_i$ with u_i . Since G_0 is a conference graph, we have $\operatorname{rdeg}_{Q_i}(u_i) = (q-1)/2$. We show that for each $j \in [k] \setminus \{i\}$, one of the following holds.

- (i) Q_i has black edges from u_i to both u_j and $-u_j$.
- (ii) u_i is nonadjacent to both u_j and $-u_j$ in Q_i .
- (iii) Q_i has at least one red edge from u_i to u_j or $-u_j$.

Suppose that Q_i has no red edges from u_i to u_j or $-u_j$. Then each of u_j and $-u_j$ is either complete or anti-complete to $\{u_i, -u_i\}$ in G_0 . Since $u_i - u_j = (-u_j) - (-u_i)$, by the definition of the Paley graph, u_i and u_j are adjacent in G_0 if and only if $-u_i$ and $-u_j$ are adjacent in G_0 . Therefore, $\{u_j, -u_j\}$ is either complete or anti-complete to $\{u_i, -u_i\}$ in G_0 . This implies that (i) or (ii) holds. There is a partial contraction sequence from Q_i to G_k by contracting pairs of the form $\{u_j, -u_j\}$ for all $j \in [k] \setminus \{i\}$. We observe from (i), (ii), and (iii) that in this sequence, the red-degree of u_i does not increase. Therefore, $\operatorname{rdeg}_{G_k}(u_i) \leq \operatorname{rdeg}_{Q_i}(u_i)$, so that $\Delta^R(G_k) \leq (q-1)/2$, and this completes the proof.

4 Twin-width versus the number of vertices

For an *n*-vertex graph, our strategy to prove Theorem 1.1 will involve applying Proposition 3.2 to find more than $(n - \sqrt{n})/2$ disjoint pairs of vertices such that contracting any one of these pairs creates a trigraph whose maximum red-degree is at most $(n + \sqrt{n} - 1)/2$. Given an ordering of these pairs, the following simple lemma presents an upper bound for the twin-width.

Lemma 4.1. Let G be an n-vertex graph for $n \ge 2$ and p_1, \ldots, p_s be pairwise disjoint elements in $\binom{V(G)}{2}$. For each $i \in [s]$, let v_i be the new vertex of G/p_i and $\mathbf{x}_i : [s] \to \{-1, 0, 1\}$ be the function such that

$$\mathbf{x}_{i}(j) = \begin{cases} \operatorname{rdeg}_{G/p_{i}/p_{j}}(v_{i}) - \operatorname{rdeg}_{G/p_{i}}(v_{i}) & \text{if } i \neq j, \\ 0 & \text{otherwise.} \end{cases}$$

Then the twin-width of G is at most

$$\max\left(\{n-s-1\}\cup\left\{\mathrm{rdeg}_{G/p_i}(v_i)+\sum_{k=1}^{j}\mathbf{x}_i(k):1\leqslant i\leqslant j\leqslant s-1\right\}\right).$$

Proof. Let $G_0 := G$ and $G_i := G_{i-1}/p_i$ for each $i \in [s]$. We also denote the new vertex of G_i by v_i . Since G_0, \ldots, G_s is a partial contraction sequence from G to G_s , tww $(G) \leq \max\{\Delta^R(G_0), \ldots, \Delta^R(G_{s-1}), \operatorname{tww}(G_s)\}$. Since G_s has n-s vertices, tww $(G_s) < |V(G_s)| = n-s$. Let $j \in [s-1]$. Since $\Delta^R(G) = 0$, by Lemma 2.2, for each vertex v in $V(G_j) \setminus \{v_1, \ldots, v_j\}$, $\operatorname{rdeg}_{G_j}(v) \leq j$. For each $i \in [j]$, by the definition of \mathbf{x}_i , $\operatorname{rdeg}_{G_j}(v_i) = \operatorname{rdeg}_{G/p_i}(v_i) + \sum_{k=1}^j \mathbf{x}_i(k)$. Therefore,

$$\Delta^{R}(G_{j}) \leq \max\left(\{j\} \cup \left\{ \operatorname{rdeg}_{G/p_{i}}(v_{i}) + \sum_{k=1}^{j} \mathbf{x}_{i}(k) : i \in [j] \right\} \right),$$

and this completes the proof, because $j \leq s - 1 \leq n - s - 1$.

We now prove Theorem 1.1.

Theorem 1.1. For a positive integer n and an n-vertex graph G,

$$\operatorname{tww}(G) < \frac{1}{2}(n + \sqrt{n \ln n} + \sqrt{n} + 2 \ln n).$$

Proof. We may assume that $n \ge 3$, since every graph on at most two vertices has twin-width 0. Let M be a maximal set of pairwise disjoint elements in $\binom{V(G)}{2}$ such that

$$|(N_G(u) \triangle N_G(v)) \setminus \{u, v\}| \leq \frac{n}{2} + \frac{n-1}{2(\sqrt{n}-1)} - 1 = \frac{n+\sqrt{n}-1}{2}$$

and each $\{u, v\} \in M$. By Proposition 3.2 with $L = V(G) \setminus \bigcup_{Q \in M} Q$ and the maximality of M, we have $|L| < \sqrt{n}$, and therefore $|M| = (n - |L|)/2 > (n - \sqrt{n})/2$.

We show that there is an ordering of the elements in M such that if we contract the elements in M in this order, then the maximum red-degree is not too large. Let s := |M| and p_1, \ldots, p_s be the elements in M and for each $i \in [s]$, v_i be the new vertex of G/p_i and $\mathbf{x}_i : [s] \to \{-1, 0, 1\}$ be the function such that

$$\mathbf{x}_{i}(j) = \begin{cases} \operatorname{rdeg}_{G/p_{i}/p_{j}}(v_{i}) - \operatorname{rdeg}_{G/p_{i}}(v_{i}) & \text{if } i \neq j, \\ 0 & \text{otherwise.} \end{cases}$$

Let $d := (n + \sqrt{n \ln n} + \sqrt{n} + 2 \ln n)/2$. Let σ be a permutation of [s] chosen uniformly at random. For each $i \in [s]$, let A_i be the event that $i \neq \sigma(s)$ and

$$\operatorname{rdeg}_{G/p_i}(v_i) + \max\left\{\sum_{k=1}^{j} \mathbf{x}_i(\sigma(k)) : \sigma^{-1}(i) \leq j \leq s-1\right\} \ge d.$$

Note that if $i \neq \sigma(s)$, then there is some j with $\sigma^{-1}(i) \leq j \leq s-1$, so A_i is well defined.

We first show that $\Pr[A_i] < 1/s$ for each $i \in [s]$. We fix an element i in [s]. Let $\alpha := |\{j \in [s] : \mathbf{x}_i(j) = -1\}|, \beta := |\{j \in [s] : \mathbf{x}_i(j) = 1\}|$, and $\gamma := s - \alpha - \beta$. Since

$$\max\left\{\sum_{k=1}^{j} \mathbf{x}_{i}(\sigma(k)) : \sigma^{-1}(i) \leq j \leq s-1\right\} \leq \beta \leq s-1,$$

we may assume that $\operatorname{rdeg}_{G/p_i}(v_i) \ge d - s + 1$. Let B_i be the event that

$$\operatorname{rdeg}_{G/p_i}(v_i) + \max\left\{\sum_{k=1}^{j} \mathbf{x}_i(\sigma(k)) : 1 \leq j \leq s\right\} \ge d.$$

We observe that $A_i \subseteq B_i$ and that $\Pr[B_i]$ is equal to the probability that a North-East lattice path from (0,0) to (α,β) chosen uniformly at random intersects the line y = x + t where

$$t := -\mathrm{rdeg}_{G/p_i}(v_i) + \lceil d \rceil.$$

Since $\operatorname{rdeg}_{G/p_i}(v_i) \leq (n + \sqrt{n} - 1)/2$, we have $t \geq d - (n + \sqrt{n} - 1)/2 = (\sqrt{n \ln n} + 1)/2 + \ln n > 0$. By Corollary 2.5, $\Pr[B_i] \leq (\beta/(\alpha + t))^t$. Thus, it suffices to show that $(\beta/(\alpha + t))^t < 1/s$. For $i, j \in [s]$, we observe that $\mathbf{x}_i(j) = -1$ if and only if G/p_i has two red edges from v_i to p_j . In addition, if i = j or G/p_i has exactly one red edge from v_i to p_j , then $\mathbf{x}_i(j) = 0$. Therefore, the number of vertices in $\bigcup_{Q \in M \setminus \{p_i\}} Q$ which are adjacent to v_i by red edges in G/p_i is at most $2\alpha + \gamma - 1$. Thus,

$$d \leq t + \operatorname{rdeg}_{G/p_i}(v_i) \leq t + 2\alpha + \gamma - 1 + \left| V(G) \setminus \bigcup_{Q \in M} Q \right|$$
$$= t + \alpha - \beta - 1 + n - s,$$

so that $\beta \leq t + \alpha - d - 1 + n - s$. Then since $1 - x \leq e^{-x}$ for all real x,

$$\left(\frac{\beta}{\alpha+t}\right)^t \leqslant \left(1 - \frac{d+s+1-n}{\alpha+t}\right)^t \leqslant \exp\left(-\frac{t(d+s+1-n)}{\alpha+t}\right).$$

Since 1/n < 1/s, it suffices to show that $t(d+s+1-n) \ge (\alpha+t) \ln n$, that is, $t(d+s+1-n-\ln n) \ge \alpha \ln n$. Since $2\alpha \le \operatorname{rdeg}_{G/p_i}(v_i) \le (n+\sqrt{n}-1)/2$, we have $\alpha \ln n \le (n \ln n + \sqrt{n} \ln n - \ln n)/4$. Note that $t \ge (\sqrt{n \ln n} + 1)/2 + \ln n$. Since $s > (n - \sqrt{n})/2$, we have $d+s > \sqrt{n \ln n}/2 + n + \ln n$. Thus,

$$\begin{split} t(d+s+1-n-\ln n) &-\alpha \ln n \\ &> \left(\frac{\sqrt{n\ln n}+1}{2} + \ln n\right) \frac{\sqrt{n\ln n}+2}{2} - \frac{n\ln n + \sqrt{n}\ln n - \ln n}{4} \\ &> \left(\frac{\sqrt{n\ln n}}{2} + \ln n\right) \frac{\sqrt{n\ln n}}{2} - \frac{n\ln n + \sqrt{n}\ln n}{4} \\ &= \frac{\sqrt{n}(\ln n)^{3/2}}{2} - \frac{\sqrt{n}\ln n}{4} > 0, \end{split}$$

and therefore $\Pr[A_i] \leq \Pr[B_i] < 1/s$.

By the union bound, $\Pr[\bigcap_{i=1}^{s} \overline{A_i}] > 0$, that is, there exists a permutation τ of [s] such that $\operatorname{rdeg}_{G/p_i}(v_i) + \max\{\sum_{k=1}^{j} \mathbf{x}_i(\tau(k)) : \tau^{-1}(i) \leq j \leq s-1\} < d$ for each $i \in [s] \setminus \{\tau(s)\}$. Note that $n - s - 1 < (n + \sqrt{n} - 2)/2 < d$. Therefore, by Lemma 4.1, the twin-width of G is less than d.

5 Twin-width versus the number of edges

We now present an upper bound for the twin-width of a graph in terms of the number of edges. This bound improves the bound in Theorem 1.1 for any *n*-vertex graph having at most $n^2/12$ edges. We will need the following lemma. While it is equivalent to restrict k to be an integer in the following lemma, we use the current form for convenience.

Lemma 5.1. Let a_1, \ldots, a_n be a sequence of nonnegative reals and $d := \sum_{i=1}^n a_i/n$. For a positive real k, there exist a subset R of [n] with |R| < k and a partition $\{B_1, \ldots, B_{k'}\}$ of $[n]\setminus R$ such that $k' \leq [k]$ and $\sum_{j \in B_i} a_j \leq dn/k$ for each $i \in [k']$.

Proof. We proceed by induction on [k]. We may assume that k > 1, because otherwise we can take $B_1 = [n]$. We may assume that $k \leq n$, because otherwise we can take R = [n]. Let B be a maximal subset of [n] such that $\sum_{j \in B} a_j \leq dn/k$. Since $k \leq n$, there is an integer j in [n] with $a_j \leq dn/k$, because otherwise $\sum_{j=1}^n a_j > dn^2/k \geq dn$. Thus, B exists and is nonempty. In addition, since k > 1, B is a proper subset of [n]. Let r be an element in $[n] \setminus B$. Without loss of generality, we may assume that $[n] \setminus (B \cup \{r\})$ is equal to [n'] for some positive integer n'. Let $d' := \sum_{j=1}^{n'} a_j/n'$. By the maximality of B, $dn - d'n' = \sum_{j \in B \cup \{r\}} a_j > dn/k$, and therefore d'n'/(k - 1) < dn/k. By the inductive hypothesis, there exist a subset R of [n'] with |R| < k - 1 and a partition $\{B_1, \ldots, B_t\}$ of $[n'] \setminus R$ such that $t \leq [k - 1]$ and $\sum_{j \in B_i} a_j \leq d'n'/(k - 1) < dn/k$ for each $i \in [t]$. Then the subset $R \cup \{r\}$ and the partition $\{B_1, \ldots, B_t, B\}$ of $[n] \setminus (R \cup \{r\})$ satisfy the required conditions. \Box

We now prove Theorem 1.2.

Theorem 1.2. For a positive integer m and an m-edge graph G,

$$\operatorname{tww}(G) < \sqrt{3m} + \frac{m^{1/4}\sqrt{\ln m}}{4 \cdot 3^{1/4}} + \frac{3m^{1/4}}{2}.$$

Proof. It is easy to check that if $m \leq 3$, then tww(G) ≤ 1 . Thus, we may assume that $m \geq 4$. Let n := |V(G)| and $q := \sqrt{4m/3}$. Note that 2 < q < n, because $4 \leq m < n^2/2$. Let

$$k := q - \frac{\sqrt{q \ln q} + \sqrt{q}}{6} - \frac{\ln q}{9},$$

$$\alpha := \frac{3}{2} \left(q + \frac{\sqrt{q \ln q} + \sqrt{q}}{6} + \frac{5 \ln q}{9} \right),$$

and d be the average degree of G. Since $q > \ln q$, we have $q > \sqrt{q \ln q} > \ln q$, and therefore 1 < q/2 < k < q. By Lemma 5.1, there exist a subset R of V(G) with r := |R| < k and a partition $\{B_1, \ldots, B_{k'}\}$ of $V(G) \setminus R$ such that $k' \leq [k]$ and $\sum_{v \in B_i} \deg_G(v) \leq dn/k$ for each $i \in [k']$. We may assume that R is nonempty by moving one vertex in B_i into R for some $i \in [k']$ if necessary. Since k < q < n, R is a proper subset of V(G) and $k' \geq 1$. For each $i \in [k']$, let $\ell_i := |B_i|$ and $v_{i,1}, \ldots, v_{i,\ell_i}$ be the vertices in B_i .

Now, we construct an $\lceil \alpha \rceil$ -contraction sequence of G. Let $G_1 := G$ and for $i \in [k']$ and $j \in [\ell_i] \setminus \{1\}$,

$$G_{j+\sum_{h=1}^{i-1}(\ell_h-1)} := G_{j-1+\sum_{h=1}^{i-1}(\ell_h-1)}/\{v_{i,1}, v_{i,j}\}.$$

Here, the new vertex created by contracting $v_{i,1}$ and $v_{i,j}$ will be called $v_{i,1}$ for convenience. Let $t := \sum_{h=1}^{k'} (\ell_h - 1) + 1 = n - k' - r + 1$. We observe that $G[R] = G_t[R]$. By Theorem 1.1, G[R] admits an $((r + \sqrt{r \ln r} + \sqrt{r} + 2 \ln r)/2)$ -contraction sequence H_1, \ldots, H_r . For each $i \in [r-1]$, let x_i and y_i be the vertices in H_i such that $H_{i+1} = H_i/\{x_i, y_i\}$ and define

$$G_{t+i} := G_{t+i-1} / \{x_i, y_i\}.$$

Finally, we take an arbitrary contraction sequence G_{t+r-1}, \ldots, G_n of G_{t+r-1} . We show that $\Delta^R(G_s) \leq [\alpha]$ for each $s \in [n]$. For each $i \in [n]$, let σ_i be the partial contraction sequence G_1, \ldots, G_i . Let $s \in [n]$ and v be a vertex of G_s .

Firstly, suppose that $s \leq t$. If $|\sigma_s^{-1}(v)| = 1$, then by Lemma 2.2, $\operatorname{rdeg}_G(v) \leq k' \leq [k] < \alpha$. If $|\sigma_s^{-1}(v)| > 1$, then by the construction of G_s , $v = v_{i,1}$ for some $i \in [k']$, and therefore by Lemma 2.1,

$$\mathrm{rdeg}_{G_s}(v) \leqslant \mathrm{deg}_{G_s}(v) \leqslant \sum_{w \in \sigma_s^{-1}(v)} \mathrm{deg}_G(w) \leqslant \sum_{w \in B_i} \mathrm{deg}_G(w) \leqslant \frac{dn}{k}$$

Since

$$\begin{split} k\alpha &= \frac{3}{2} \left(q - \frac{\sqrt{q \ln q} + \sqrt{q}}{6} - \frac{\ln q}{9} \right) \left(q + \frac{\sqrt{q \ln q} + \sqrt{q}}{6} + \frac{5 \ln q}{9} \right) \\ &= \frac{3}{2} \left(\left(q + \frac{2 \ln q}{9} \right)^2 - \left(\frac{\sqrt{q \ln q} + \sqrt{q}}{6} + \frac{\ln q}{3} \right)^2 \right) \\ &> \frac{3}{2} \left(\left(q + \frac{2 \ln q}{9} \right)^2 - \frac{4q \ln q}{9} \right) \\ &= \frac{3}{2} \left(q^2 + \frac{4 \ln^2 q}{81} \right) = dn + \frac{2 \ln^2 q}{27} > dn, \end{split}$$

we have $dn/k < \alpha$, so that $\operatorname{rdeg}_{G_s}(v) < \alpha$. Therefore, $\Delta^R(G_s) < \alpha$ for every $s \leq t$.

Secondly, suppose that t < s < t + r. By the construction of G_s , if $v = v_{i,1}$ for some $i \in [k']$, then $\sigma_s^{-1}(v) = \sigma_t^{-1}(v) = B_i$. Therefore, by Lemma 2.1,

$$\operatorname{rdeg}_{G_s}(v) \leqslant \operatorname{deg}_{G_s}(v) \leqslant \sum_{w \in \sigma_s^{-1}(v)} \operatorname{deg}_G(w) = \sum_{w \in B_i} \operatorname{deg}_G(w) \leqslant \frac{dn}{k} < \alpha.$$

If $v \in V(G_s) \setminus \{v_{i,1} : i \in [k']\}$, then since $G_s \setminus \{v_{i,1} : i \in [k']\} = H_{s-t+1}$,

$$\begin{aligned} \operatorname{rdeg}_{G_s}(v) &\leqslant |\{v_{i,1} : i \in [k']\}| + \operatorname{rdeg}_{H_{s-t+1}}(v) \\ &< k+1 + \frac{r + \sqrt{r \ln r} + \sqrt{r} + 2\ln r}{2} \\ &< \frac{3k}{2} + \frac{\sqrt{q \ln q} + \sqrt{q} + 2\ln q}{2} + 1 \\ &= \frac{3}{2} \left(q - \frac{\sqrt{q \ln q} + \sqrt{q}}{6} - \frac{\ln q}{9}\right) + \frac{\sqrt{q \ln q} + \sqrt{q} + 2\ln q}{2} + 1 \\ &= \alpha + 1. \end{aligned}$$

Since $\operatorname{rdeg}_{G_s}(v)$ is an integer, we have $\operatorname{rdeg}_{G_s}(v) \leq \lceil \alpha \rceil$. Therefore, $\Delta^R(G_s) \leq \lceil \alpha \rceil$ for every s with t < s < t + r.

Thirdly, if $s \ge t + r$, then

$$\Delta^{\mathbb{R}}(G_s) < |V(G_s)| = n - s + 1 \leq n - t - r + 1 = k' \leq \lceil k \rceil < \alpha,$$

and therefore tww(G) $\leq [\alpha]$. By the mean value theorem,

$$\frac{\sqrt{\ln(4m/3)} - \sqrt{\ln m}}{m/3} = \frac{1}{2m'\sqrt{\ln m'}}$$

for some $m' \in (m, 4m/3)$. Hence, $\sqrt{\ln(4m/3)} - \sqrt{\ln m} < 1/(6\sqrt{\ln m})$. Then

$$\begin{split} &\alpha - \sqrt{3m} + 1 \\ &= \frac{\sqrt{q \ln q}}{4} + \frac{\sqrt{q}}{4} + \frac{5 \ln q}{6} + 1 \\ &= \frac{\sqrt{\sqrt{m/3} \cdot \ln(4m/3)}}{4} + \frac{\sqrt{2}m^{1/4}}{4 \cdot 3^{1/4}} + \frac{5(\ln m + \ln(4/3))}{12} + 1 \\ &< \frac{1}{4 \cdot 3^{1/4}} \left(m^{1/4}\sqrt{\ln m} + \frac{m^{1/4}}{6\sqrt{\ln m}} \right) + \frac{\sqrt{2}m^{1/4}}{4 \cdot 3^{1/4}} + \frac{5(\ln m + \ln(4/3))}{12} + 1 \\ &< \frac{m^{1/4}\sqrt{\ln m}}{4 \cdot 3^{1/4}} + \frac{m^{1/4}}{24 \cdot 3^{1/4}\sqrt{\ln m}} + \frac{\sqrt{2}m^{1/4}}{4 \cdot 3^{1/4}} + \frac{5\ln m}{12} + \frac{9}{8} \\ &= \frac{m^{1/4}\sqrt{\ln m}}{4 \cdot 3^{1/4}} + m^{1/4} \left(\frac{1}{24 \cdot 3^{1/4}\sqrt{\ln m}} + \frac{\sqrt{2}}{4 \cdot 3^{1/4}} + \frac{5\ln m}{12m^{1/4}} + \frac{9}{8m^{1/4}} \right) \\ &< \frac{m^{1/4}\sqrt{\ln m}}{4 \cdot 3^{1/4}} + \frac{3m^{1/4}}{2}, \end{split}$$

where the last inequality holds because

$$\frac{1}{24 \cdot 3^{1/4}\sqrt{\ln x}} + \frac{\sqrt{2}}{4 \cdot 3^{1/4}} + \frac{5\ln x}{12x^{1/4}} + \frac{9}{8x^{1/4}}$$

is decreasing if $x \ge 4$. Thus,

$$\mathrm{tww}(G) < \alpha + 1 < \sqrt{3m} + \frac{m^{1/4}\sqrt{\ln m}}{4\cdot 3^{1/4}} + \frac{3m^{1/4}}{2}.$$

6 Twin-width and random graphs

We now prove Theorem 1.3. Since a graph and its complement have the same twin-width [8], we focus on random graphs G(n, p) with 0 .

Theorem 1.3. Let G := G(n,p) be a random graph where p := p(n) is a function such that $1/n \leq p \leq 1/2$ and ε be a positive real. Then

$$\operatorname{tww}(G) > 2p(1-p)n - (2\sqrt{2} + \varepsilon)\sqrt{p(1-p)n\ln n}$$

asymptotically almost surely.

Proof. By Lemma 3.1, it suffices to show that for all distinct $i, j \in [n]$,

$$|(N_G(i) \triangle N_G(j)) \setminus \{i, j\}| > 2p(1-p)n - (2\sqrt{2} + \varepsilon)(p(1-p)n\ln n)^{1/2}$$

asymptotically almost surely. Let $s := 1 - p^2 - (1 - p)^2 = 2p(1 - p)$. For all distinct $i, j \in [n]$ and each $k \in [n] \setminus \{i, j\}$, let $X_k^{\{i, j\}}$ be the random variable such that

$$X_k^{\{i,j\}} = \begin{cases} 1 & \text{if exactly one of } i \text{ or } j \text{ is adjacent to } k \text{ in } G, \\ 0 & \text{otherwise,} \end{cases}$$

which indicates whether k is incident with a red edge in $G/\{i, j\}$. For all distinct $i, j \in [n]$, let $X^{\{i, j\}} := \sum_{\ell \in [n] \setminus \{i, j\}} X_{\ell}^{\{i, j\}}$. It is readily seen that $\mathbb{E}[X^{\{i, j\}}] = s(n-2)$ for all distinct $i, j \in [n]$. Let $C := 2 + \varepsilon/\sqrt{2}$ and $D := 2 + \varepsilon/2$. For all distinct $i, j \in [n]$, let $A^{\{i, j\}}$ be the event that $X^{\{i, j\}} > sn - C(sn \ln n)^{1/2}$. Then it suffices to show that $\Pr[\bigcap_{1 \leq i < j \leq n} A^{\{i, j\}}] \to 1$ as $n \to \infty$. Since $1/n \leq p \leq 1/2$, we have $sn \geq 1$. Thus, for all sufficiently large n,

$$C(sn\ln n)^{1/2} - D(s(n-2)\ln n)^{1/2} > (C-D)(sn\ln n)^{1/2} \ge 2 \ge 2s.$$

That is,

$$-C(sn\ln n)^{1/2} < -2s - D(s(n-2)\ln n)^{1/2}$$

Then by the union bound and Chernoff bound,

$$\begin{split} \Pr\left[\bigcup_{1\leqslant i < j\leqslant n} \overline{A^{\{i,j\}}}\right] &\leqslant \binom{n}{2} \Pr\left[\overline{A^{\{1,2\}}}\right] \\ &= \binom{n}{2} \Pr[X^{\{1,2\}} \leqslant sn - C(sn\ln n)^{1/2}] \\ &\leqslant \binom{n}{2} \Pr[X^{\{1,2\}} \leqslant s(n-2) - D(s(n-2)\ln n)^{1/2}] \\ &\leqslant \binom{n}{2} \exp\left(-\frac{D^2 s(n-2)\ln n}{2s(n-2)}\right) \\ &= \binom{n}{2} n^{-D^2/2} < n^{2-D^2/2} \to 0 \end{split}$$

as $n \to \infty$, and this completes the proof.

As a corollary, we remark that if $(2 + \delta)n^{-1} \ln n \leq p \leq 1/2$ for some positive real δ , then tww $(G(n, p)) = \Omega(\ln n)$ asymptotically almost surely.

We now prove Theorem 1.5. We will use the following three theorems of Erdős and Rényi [11].

Theorem 6.1 (Erdős and Rényi [11]). Let c be a constant in (0,1). The random graph G := G(n, c/n) asymptotically almost surely has the property that every component of G has at most one cycle.

A graph G is *balanced* if for every non-null subgraph H of G, the average degree of H is at most the average degree of G. We remark that all trees and cycles are balanced.

Theorem 6.2 (Erdős and Rényi [11]). Let $p : \mathbb{N} \to [0,1]$ be a function and H be a balanced graph with v vertices and e edges. If $p = o(n^{-v/e})$, then G(n,p) has no subgraph isomorphic to H asymptotically almost surely. If $p = \omega(n^{-v/e})$, then G(n,p) has a subgraph isomorphic to H asymptotically almost surely.

Theorem 6.3 (Erdős and Rényi [11]). For a function $p : \mathbb{N} \to [0,1]$ with $p = o(n^{-1}), G(n,p)$ is a forest asymptotically almost surely.

We also show that if a connected graph has at most one cycle, then its twin-width is at most 2. We will use the following 2-contraction sequence of a tree [8].

Proposition 6.4 (Bonnet et al. [8]). Every rooted tree admits a 2-contraction sequence such that the root is contracted only in the last contraction.

Proposition 6.5. If every component of a graph G has at most one cycle, then $tww(G) \leq 2$.

Proof. Since the twin-width of a graph is the maximum twin-width of its components [8], we may assume that G is connected. By Proposition 6.4, we may assume that G has exactly one cycle C. We are going to construct a 2-contraction sequence of G. Let v_1, \ldots, v_ℓ be the vertices of C such that $v_i v_j \in E(C)$ if either |i - j| = 1 or $\{i, j\} = \{1, \ell\}$. For each $i \in [\ell]$, let T_i be the component of $G \setminus \{v_{i-1}, v_{i+1}\}$ containing v_i where $v_0 := v_n$ and $v_{\ell+1} := v_1$ which is a tree rooted at v_i . For each $i \in [\ell]$, by Proposition 6.4, there exists a partial 2-contraction sequence σ_i from T_i to some T'_i such that $\sigma_i^{-1}(v_i) = \{v_i\}$ and T'_i has at most two vertices. By applying $\sigma_1, \ldots, \sigma_\ell$ in sequence, we obtain a partial contraction sequence σ from G to some trigraph G'. Note that for each $i \in [\ell]$, v_i is not contracted during the process of constructing G', so no edge of C is a red edge of G' and σ is a partial 2-contraction sequence.

Now, we present a 2-contraction sequence of G'. Since taking induced subgraph does not increase twin-width, we may assume that for each $i \in [\ell]$, T'_i has two vertices, say v_i and w_i . For convenience, if we contract



Figure 1: A 1-subdivision of $K_{1,3}$.

vertices v_i and w_j for some $i, j \in [\ell]$, then we call the new vertex v_i . Let $G_1 := G'/\{v_1, w_1\}$. For each $i \in [\ell] \setminus \{1\}$, let $G_i := G_{i-1}/\{v_{i-1}, w_i\}$. One may observe that for each $i \in [\ell]$, $\Delta^R(G_i) = 2$, and G_ℓ is a cycle. Thus, by contracting adjacent vertices of G_ℓ iteratively, we find a 2-contraction sequence of G'.

A caterpillar is a tree that contains a path P such that every vertex outside P is a leaf whose neighbor is in V(P). It is easy to observe that a tree is a caterpillar if and only if it has no 1-subdivision of $K_{1,3}$ (see Figure 1) as a subgraph.

Lemma 6.6. For a tree T, $tww(T) \leq 1$ if and only if T is a caterpillar.

Proof. We first show that $tww(T) \leq 1$ if T is a caterpillar. We may assume that no pair of leaves are adjacent to the same vertex. Hence, for some integer n, T is isomorphic to an induced subgraph of the tree obtained from the path $v_2v_4\cdots v_{2n}$ by appending a leaf v_{2i-1} to the vertex v_{2i} for each $i \in [n]$. We find a 1-contraction sequence for this graph by iteratively contracting v_i and v_{i+1} into v_{i+1} for $i \in [2n-1]$.

For the forward direction, it suffices to show that the 1-subdivision of $K_{1,3}$ has twin-width 2. Let H be the 1-subdivision of $K_{1,3}$. It is readily seen that for distinct vertices v and w of H, $|(N_H(v) \triangle N_H(w)) \setminus \{v, w\}| \leq 1$ if and only if v and w are adjacent and one of them is a leaf. Thus, by Lemma 3.1, we may assume that we first contract such vertices, say v and w. Then $H/\{v, w\}$ has no two vertices whose contraction yields a 1-trigraph.

We now prove Theorem 1.5.

Theorem 1.5. For a random graph G := G(n, p) with a function p := p(n), the following hold asymptotically almost surely.

- 1. tww(G) = 0 if $p = o(n^{-4/3})$.
- 2. tww(G) = 1 if $p = \omega(n^{-4/3})$ and $p = o(n^{-7/6})$.
- 3. tww(G) = 2 if $p = \omega(n^{-7/6})$ and $p \leq c/n$ for a constant $c \in (0, 1)$.

Proof. If $p = o(n^{-4/3})$, then by Theorems 6.2 and 6.3, G has no induced path of length 3 asymptotically almost surely. A graph has twin-width 0 if and only if it has no induced path of length 3 [8], and therefore tww(G) = 0 asymptotically almost surely.

If $p = \omega(n^{-4/3})$ and $p = o(n^{-7/6})$, then by Theorems 6.2 and 6.3, G is a forest and contains a path of length 3 and no 1-subdivision of $K_{1,3}$ as a subgraph asymptotically almost surely. Then by Lemma 6.6, tww(G) = 1 asymptotically almost surely.

If $p = \omega(n^{-7/6})$ and $p = o(n^{-1})$, then by Theorems 6.2 and 6.3, G is a forest and contains the 1-subdivision of $K_{1,3}$ as a subgraph asymptotically almost surely. Then by Proposition 6.4 and Lemma 6.6, tww(G) = 2 asymptotically almost surely.

Now, suppose that $p = \omega(n^{-9/8})$ and $p \leq c/n$ for a constant $c \in (0, 1)$. By Theorem 6.1, every component of G has at most one cycle asymptotically almost surely, and therefore tww $(G) \leq 2$ asymptotically almost surely by Proposition 6.5. By Theorem 6.2, G has the 1-subdivision of $K_{1,4}$ as a subgraph asymptotically almost surely, and therefore has the 1-subdivision of $K_{1,3}$ as an induced subgraph. Since the 1-subdivision of $K_{1,3}$ has twinwidth 2, we have tww $(G) \geq 2$ asymptotically almost surely.

References

- Jakub Balabán and Petr Hliněný, Twin-width is linear in the poset width, 16th International Symposium on Parameterized and Exact Computation (IPEC 2021), arXiv:2106.15337, 2021, accepted.
- [2] Miklós Bóna (ed.), Handbook of enumerative combinatorics, Discrete Mathematics and its Applications (Boca Raton), CRC Press, Boca Raton, FL, 2015. MR 3408702
- [3] Édouard Bonnet, Twin-width I: tractable FO model checking, YouTube, Nov. 3, 2020, https://www.youtube.com/watch?v=DFMEiVz_A3Q.
- [4] Édouard Bonnet, Colin Geniet, Eun Jung Kim, Stéphan Thomassé, and Rémi Watrigant, *Twin-width II: small classes*, 2021 ACM-SIAM Symposium on Discrete Algorithms (SODA 2021), SIAM, 2021, pp. 1977– 1996.
- [5] Édouard Bonnet, Colin Geniet, Eun Jung Kim, Stéphan Thomassé, and Rémi Watrigant, Twin-width III: Max Independent Set, Min Dominating Set, and Coloring, 48th International Colloquium on Automata,

Languages, and Programming (ICALP 2021) (Dagstuhl, Germany) (Nikhil Bansal, Emanuela Merelli, and James Worrell, eds.), Leibniz International Proceedings in Informatics (LIPIcs), vol. 198, Schloss Dagstuhl – Leibniz-Zentrum für Informatik, 2021, pp. 35:1–35:20.

- [6] Édouard Bonnet, Ugo Giocanti, Patrice Ossona de Mendez, Pierre Simon, Stéphan Thomasé, and Szymon Toruńczyk, *Twin-width IV: ordered graphs and matrices*, 2022 ACM-SIAM Symposium on Discrete Algorithms (SODA 2022), SIAM, arXiv:2102.03117, 2022, accepted.
- [7] Edouard Bonnet, Eun Jung Kim, Amadeus Reinald, Stéphan Thomassé, and Rémi Watrigant, *Twin-width and polynomial kernels*, 16th International Symposium on Parameterized and Exact Computation (IPEC 2021), arXiv:2107.02882, 2021, accepted.
- [8] Édouard Bonnet, Eun Jung Kim, Stéphan Thomassé, and Rémi Watrigant, Twin-width I: tractable FO model checking, 2020 IEEE 61st Annual Symposium on Foundations of Computer Science (FOCS 2020), IEEE Computer Soc., Los Alamitos, CA, 2020, pp. 601–612. MR 4232070
- [9] Édouard Bonnet, Jaroslav Nešetřil, Patrice Ossona de Mendez, Sebastian Siebertz, and Stéphan Thomassé, Twin-width and permutations, arXiv:2102.06880, 2021.
- [10] Jan Dreier, Jakub Gajarsky, Yiting Jiang, Patrice Ossona de Mendez, and Jean-Florent Raymond, *Twin-width and generalized coloring num*bers, arXiv:2104.09360, 2021.
- [11] Paul Erdős and Alfréd Rényi, On the evolution of random graphs, A Magyar Tudományos Akadémia. Matematikai Kutató Intézetének Közleményei 5 (1960), 17–61. MR 125031
- [12] _____, Asymmetric graphs, Acta Math. Acad. Sci. Hungar. 14 (1963), 295–315. MR 156334
- [13] Jakub Gajarský, Michał Pilipczuk, and Szymon Toruńczyk, Stable graphs of bounded twin-width, arXiv:2107.03711, 2021.
- [14] Svante Janson, Tomasz Łuczak, and Andrzej Rucinski, Random graphs, Wiley-Interscience Series in Discrete Mathematics and Optimization, Wiley-Interscience, New York, 2000. MR 1782847
- [15] Vít Jelínek, The rank-width of the square grid, Discrete Appl. Math.
 158 (2010), no. 7, 841–850. MR 2602833

- [16] Choongbum Lee, Joonkyung Lee, and Sang-il Oum, Rank-width of random graphs, J. Graph Theory 70 (2012), no. 3, 339–347. MR 2946080
- [17] Pierre Simon and Szymon Toruńczyk, Ordered graphs of bounded twinwidth, arXiv:2102.06881, 2021.