



# Transverse spherocity dependence of azimuthal anisotropy in heavy-ion collisions at the LHC using a multi-phase transport model

Neelkamal Mallick,<sup>*a*,\*</sup> Raghunth Sahoo,<sup>*a*,*b*</sup> Sushanta Tripathy<sup>*c*</sup> and Antonio Ortiz<sup>*d*</sup>

<sup>a</sup>Department of Physics, Indian Institute of Technology Indore, Simrol, Indore 453552, India

<sup>b</sup>CERN, CH 1211, Geneva 23, Switzerland

<sup>c</sup> INFN - sezione di Bologna, via Irnerio 46, 40126 Bologna BO, Italy

<sup>d</sup> Instituto de Ciencias Nucleares, Universidad Nacional Autónoma de México, México Distrito Federal 04510, México

*E-mail:* Neelkamal.Mallick@cern.ch, Raghunath.Sahoo@cern.ch, Sushanta.Tripathy@cern.ch, Antonio.Ortiz.Velasquez@cern.ch

One of the event shape observables, the transverse spherocity ( $S_0$ ), has been studied successfully in small collision systems such as proton-proton collisions at the LHC as a tool to separate jetty and isotropic events. It has a unique capability to distinguish events based on their geometrical shapes. In our work, we report the first implementation of transverse spherocity in heavyion collisions using a multi-phase transport model (AMPT). We have performed an extensive study of azimuthal anisotropy of charged particles produced in heavy-ion collisions as a function of transverse spherocity ( $S_0$ ). We have followed the two-particle correlation (2PC) method to estimate the elliptic flow ( $v_2$ ) in different centrality classes in Pb-Pb collisions at  $\sqrt{s_{NN}} = 5.02$  TeV for high- $S_0$ ,  $S_0$ -integrated and low- $S_0$  events. We found that transverse spherocity successfully differentiates heavy-ion collisions' event topology based on their geometrical shapes, i.e., high and low values of spherocity. The high- $S_0$  events have nearly zero elliptic flow, while the low- $S_0$ events contribute significantly to the elliptic flow of spherocity-integrated events.

The Ninth Annual Conference on Large Hadron Collider Physics - LHCP2021 7-12 June 2021 Online


#### \*Speaker

<sup>©</sup> Copyright owned by the author(s) under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License (CC BY-NC-ND 4.0).

# 1. Introduction

The production of a hot and dense, deconfined state of matter known as the quark-gluon plasma (QGP) has already been established in heavy-ion collisions at the Large Hadron Collider (LHC) at CERN, Switzerland, and Relativistic Heavy Ion Collider (RHIC) at BNL, USA. Recent studies at the LHC show heavy-ion-like features such as ridge-like structures [1] and strangeness enhancements [2] in *pp* collisions. To understand the system dynamics and production of jets in *pp* collisions, an event shape observable, known as the transverse spherocity ( $S_0$ ), has been introduced recently at the LHC [3–7]. These studies show that transverse spherocity ( $S_0$ ) has unique capabilities to distinguish events based on its geometrical shapes *i.e.* jetty and isotropic events. The study of transverse spherocity in heavy-ion collisions may reveal new and unique physics results where the formation of QGP is already known. This study in heavy-ion collisions shall also complement the current event shape approach based on flow vector analysis at the LHC [8, 9].

In this work [10], we report the first implementation of transverse spherocity in heavy-ion collisions using a multi-phase transport (AMPT) model [11]. We have performed an extensive study of azimuthal anisotropy of charged particles produced in Pb-Pb collisions at  $\sqrt{s_{NN}} = 5.02$  TeV as a function of transverse spherocity ( $S_0$ ). We have followed the two-particle correlation (2PC) method to estimate the elliptic flow ( $v_2$ ) and subtract non-flow from our calculations by following standard experimental procedures.



**Figure 1:** (Color Online) Transverse momentum space correlation ( $p_y$  vs.  $p_x$ ) for different spherocity classes in (40-50)% central Pb-Pb collisions at  $\sqrt{s_{NN}} = 5.02$  TeV using AMPT model. (Fig. 2 [10])

Transverse spherocity  $(S_0)$  is an event property which is also a collinear and infrared safe quantity [3–5] and defined as follows,

$$S_0 = \frac{\pi^2}{4} \left( \frac{\sum_i |\vec{p}_{T_i} \times \hat{n}|}{\sum_i p_{T_i}} \right)^2.$$
(1)


Here,  $\hat{n}(n_T, 0)$  is a unit vector known as the spherocity axis, which minimizes the ratio in Eq. 1. For jetty events  $S_0 \approx 0$  and for isotropic events  $S_0 \approx 1$  [7]. We have used the mid-rapidity ( $|\eta| < 0.8$ ) spherocity distribution with  $p_T > 0.15$  GeV/c with events having at least five such tracks to meet similar conditions as in the ALICE experiment at the LHC. The low- $S_0$  and high- $S_0$  events represent the events lying in the bottom 20% and top 20% in the  $S_0$  distributions whereas  $S_0$ -integrated events take all events into account.

Figure 1 represents the transverse momentum space correlation  $(p_y \text{ vs. } p_x)$  for high- $S_0$ ,  $S_0$ integrated and low- $S_0$  events in (40-50)% central Pb-Pb collisions at  $\sqrt{s_{NN}} = 5.02$  TeV using AMPT model. The elliptic flow in  $S_0$ -integrated events can be clearly seen from the elliptic shape of the correlation plot. This is indeed credited to the initial pressure gradient caused due to the initial almond-shaped nuclear overlap region in semi-central collisions, which is then translated to the momentum space  $(p_x > p_y)$  correlations. The interesting thing is that the high- $S_0$  events show almost spherical momentum correlation indicating the presence of nearly zero elliptic flow in such events. Whereas the events with low- $S_0$  show a greater elliptical shape correlation. That indicates low- $S_0$  events should be more elliptic and therefore contribute more towards  $v_2$ .

## 2. Results and Discussions

To estimate the elliptic flow we have used the two-particle correlation method (2PC) [8, 12]. The two particle correlation function  $C(\Delta \eta, \Delta \phi)$  is constructed by taking the ratios of same-event pairs  $S(\Delta \eta, \Delta \phi)$  to mixed-event pairs  $B(\Delta \eta, \Delta \phi)$  given by,

$$C(\Delta\eta, \Delta\phi) = \frac{S(\Delta\eta, \Delta\phi)}{B(\Delta\eta, \Delta\phi)}.$$
(2)



**Figure 2:** (Color Online) First column: One dimensional azimuthal correlation of charged particles, second column: two particle elliptic flow coefficient  $(v_{2,2}(p_T^a, p_T^b))$  of charged particles as a function of  $p_T^a$ , and third column: single particle elliptic flow coefficient  $(v_2(p_T^a))$  of charged particles as a function of  $p_T^a$  for low- $S_0$ , high- $S_0$  and  $S_0$ -integrated events in Pb–Pb collisions at  $\sqrt{s_{NN}} = 5.02$  TeV for 0%–10% (top), 40%–50% (bottom) centrality classes using AMPT model. (Figs. 4-6 [10])

The one dimensional correlation function  $(C(\Delta \phi))$  is then calculated by integrating all the pairs with pseudorapidity gap 2.0 <  $|\Delta \eta|$  < 4.8. This step helps in reducing the contributions from

non-flow effects [8]. From the  $C(\Delta \phi)$  distribution, the two particle flow coefficient  $(v_{2,2}(p_T^a, p_T^b))$  could be easily obtained as [13–15],

$$v_{n,n}(p_{\rm T}^{a}, p_{\rm T}^{b}) = \langle \cos(n\Delta\phi) \rangle = \frac{\sum_{m=1}^{N} \cos(n\Delta\phi_m) \times C(\Delta\phi_m)}{\sum_{m=1}^{N} C(\Delta\phi_m)}.$$
(3)

where, N = 200 is the number of  $\Delta \phi$  bins in the range  $-\pi/2 < \Delta \phi < 3\pi/2$ . Here,  $v_{n,n}$  are symmetric functions with respect to  $p_T^a$  and  $p_T^b$ . The single particle flow coefficient could be obtained as,

$$v_{n,n}(p_{\rm T}^a, p_{\rm T}^b) = v_n(p_{\rm T}^a)v_n(p_{\rm T}^b).$$
 (4)

Figure 2 first column shows that the shape and nature of the 1D azimuthal correlation vary with spherocity, and one can observe that the amplitude of the correlation is greater in low- $S_0$ , intermediate in  $S_0$ -integrated and lower in high- $S_0$  events. Again, the amplitude of the correlation is more in semi-central collisions compared to most central collisions. Figure 2 second column represents the two particle elliptic flow co-efficient  $v_{2,2}(p_T^a, p_T^b)$  as a function of spherocity. Here,  $v_{2,2}(p_T^a, p_T^b)$  has strong centrality dependence with showing greater values towards semi-central collisions. As far as spherocity is concerned, the high- $S_0$  events are found to have nearly zero elliptic flow while the low- $S_0$  events contribute significantly to elliptic flow of spherocity-integrated events. This is evident from the third column of Fig. 2.

### 3. Summary

In summary, we found that transverse spherocity successfully differentiates heavy-ion collisions' event topology based on their geometrical shapes, *i.e.* high and low values of spherocity  $(S_0)$ . The high- $S_0$  events are found to have nearly zero elliptic flow, while the low- $S_0$  events contribute significantly to the elliptic flow of spherocity-integrated events. Transverse spherocity is anti-correlated to elliptic flow.

#### Acknowledgements

R.S. acknowledges the financial supports under the CERN Scientific Associateship and the financial grants from DAE-BRNS Project No. 58/14/29/2019-BRNS. The authors would like to acknowledge the usage of resources of the LHC grid computing facility at VECC, Kolkata, and the computing farm at ICN-UNAM. S.T. acknowledges the support from INFN postdoctoral fellowship in experimental physics. A.O. acknowledges the financial support from CONACyT under Grant No. A1-S-22917.

#### References

- [1] V. Khachatryan et al. [CMS], Phys. Lett. B 765 (2017), 193
- [2] J. Adam et al. [ALICE], Nature Phys. 13 (2017), 535

- [3] E. Cuautle, R. Jimenez, I. Maldonado, A. Ortiz, G. Paic and E. Perez, [arXiv:1404.2372 [hep-ph]].
- [4] A. Ortiz, G. Paić and E. Cuautle, Nucl. Phys. A 941 (2015), 78
- [5] G. P. Salam, Eur. Phys. J. C 67 (2010), 637
- [6] G. Bencédi [ALICE], Nucl. Phys. A 982 (2019), 507
- [7] S. Acharya et al. [ALICE], Eur. Phys. J. C 79 (2019), 857
- [8] G. Aad et al. [ATLAS], Phys. Rev. C 92 (2015), 034903
- [9] M. Masera, G. Ortona, M. G. Poghosyan and F. Prino, Phys. Rev. C 79 (2009), 064909
- [10] N. Mallick, R. Sahoo, S. Tripathy and A. Ortiz, J. Phys. G 48 (2021), 045104
- [11] Z. W. Lin, C. M. Ko, B. A. Li, B. Zhang and S. Pal, Phys. Rev. C 72 (2005), 064901
- [12] S. A. Voloshin, A. M. Poskanzer and R. Snellings, Landolt-Bornstein 23 (2010), 293
- [13] G. Aad et al. [ATLAS], Phys. Rev. C 86 (2012), 014907
- [14] S. Chatrchyan et al. [CMS], Phys. Rev. C 89 (2014), 044906
- [15] K. Aamodt et al. [ALICE], Phys. Lett. B 708 (2012), 249