Odd-dimensional solvmanifolds are contact

CHRISTOPH BOCK

MSC 2020: Primary: 57R17; Secondary: 53D15.

Abstract

Bourgeois proved in [5] that odd-dimensional tori admit a contact structure. We shall prove a more general result: Any odd-dimensional parallelisable closed manifold admits a contact structure. This implies that a solvmanifold $\Gamma \setminus G$ is contact, where Γ is a lattice in a connected and simply-connected solvable Lie group G of odd dimension.

Throughout this short note a manifold is assumed to be smooth, connected, oriented and to have no boundary.

A contact manifold is a pair (M, ξ) , where M is a (2n+1)-manifold, $n \in \mathbb{N}$, and ξ can be written as $\xi = \ker \alpha$, where $\alpha \in \Omega^1(M)$ is a differential 1-form with $\alpha_p \wedge (\mathrm{d}\alpha)^n_{\ p} \neq 0$ for all $p \in M$. This implies that the structure group of the tangent bundle of M reduces to $\mathrm{U}(n) \times \{1\}$, see [6, p. 68]. Therefore, it makes sense to call a manifold almost contact, if the structure group of the tangent bundle reduces to $\mathrm{U}(n) \times \{1\}$. There is the following important consequence of the deep result [4, Corollary 1.3] of Borman, Eliashberg and Murphy:

Proposition 1. Let M be an almost contact closed manifold. Then M admits the structure of a contact manifold.

Therefore, in order to obtain the next theorem, it is enough to prove the next proposition.

Theorem 2. Any odd-dimensional parallelisable closed manifold admits a contact structure.

Proposition 3. Any odd-dimensional parallelisable closed manifold is almost contact.

Proof. Let $n \in \mathbb{N}$, M be a (2n+1)-dimensional closed manifold equipped with a parallelisation (X_1, \ldots, X_{2n+1}) . Then,

$$JX_1 := X_2, JX_2 := -X_1, \dots, JX_{2n-1} := X_{2n}, JX_{2n} := -X_{2n-1}$$

gives rise to an almost contact structure $(\langle X_1, \dots, X_{2n} \rangle_{\mathbb{R}}, J, \langle X_{2n+1} \rangle_{\mathbb{R}})$ on M, see again [6, p. 68], and the structure group of the tangent bundle of M reduces to $U(n) \times \{1\}$.

In [3], we asked whether every five-dimensional solvmanifold admits a contact structure.¹ Using Theorem 2, we can give the answer for any odd-dimensional solvmanifold.

A solvmanifold is a homogeneous space $\Gamma \backslash G$, where G is a connected and simply-connected solvable Lie group and Γ a lattice in G. Recall that a lattice in a Lie group is a discrete co-compact subgroup. Therefore, tori are solvmanifolds.

Remark. It is important to note that there is a more general notion of solvmanifold, namely a compact quotient of a connected and simply-connected solvable Lie group by a (possibly non-discrete) closed Lie subgroup (see [1]), but we are only considering solvmanifolds as in the last definition. Sometimes, such are called *special solvmanifolds* in the literature.

By [7, Theorem 2.3.11], a solvmanifold in our sense is necessary parallelisable. E.g. the Klein bottle (which can be written as the compact homogeneous space

$$\{(\frac{k}{2},l+iy)\,|\,k,l\in\mathbb{Z}\,\wedge\,y\in\mathbb{R}\}\backslash(\mathbb{R}\ltimes_{\mu}\mathbb{C}),\text{ where }\forall_{t\in\mathbb{R}}\forall_{z\in\mathbb{C}}\,\mu(t)(z):=\mathrm{e}^{2\pi\mathrm{i}t}z,$$

of the three-dimensional connected and simply-connected solvable Lie group $\mathbb{R} \ltimes_{\mu} \mathbb{C}$) is not a solvmanifold covered by our definition. (The manifold M in [2, Section 3] is not a solvmanifod covered by our definition, too.)

[7, Theorem 2.3.11] and Theorem 2 imply:

Theorem 4. Odd-dimensional solvmanifolds admit a contact structure. \Box

References

- [1] L. Auslander: An exposition of the structure of solvmanifolds part I, Bull. Amer. Math. Soc. **79** (1973), no. 2, 227–261.
- [2] L. Auslander, R. H. Szczarba: Characteristic classes of compact solvmanifolds, Ann. of Math. (2) **76** (1962), 1–8.
- [3] Ch. Bock: The solution on the geography problem of non-formal (almost) contact manifolds, Topology appl. **317** (2022), Article 108186.
- [4] M. S. Borman, Y. Eliashberg, E. Murphy, Existence and classification of overtwisted contact structures in all dimensions, Acta Math. 215 (2015), 281–361.
- [5] F. Bourgeois: Odd dimensional tori are contact manifolds, Int. Math. Res. Notices 30 (2002), 1571–1574.
- [6] H. Geiges: An Introduction to Contact Geometry, Cambridge University Press (2008).

¹There, by another approach, we saw that most five-dimensinal solvmanifolds are contact.

[7] J. Oprea, A. Tralle: Symplectic Manifolds with no Kähler Structure, Lecture Notes in Math. 1661, Springer (1997).

CHRISTOPH BOCK
DEPARTMENT MATHEMATIK
UNIVERSITÄT ERLANGEN-NÜRNBERG
CAUERSTRASSE 11
91058 ERLANGEN
GERMANY
e-mail: bock@mi.uni-erlangen.de