

Eigenvalues and cycles of consecutive lengths

Binlong Li* Bo Ning†

Abstract: As the counterpart of classical theorems on cycles of consecutive lengths due to Bondy and Bollobás in spectral graph theory, Nikiforov proposed the following open problem in 2008: What is the maximum C such that for all positive $\varepsilon < C$ and sufficiently large n , every graph G of order n with spectral radius $\rho(G) > \sqrt{\lfloor \frac{n^2}{4} \rfloor}$ contains a cycle of length ℓ for each integer $\ell \in [3, (C - \varepsilon)n]$. We prove that $C \geq \frac{1}{4}$, improving the existing bounds. Besides several novel ideas, our proof technique is partly inspired by the recent research on Ramsey numbers of star versus large even cycles due to Allen, Łuczak, Polcyn and Zhang, and with aid of a powerful spectral inequality. We also derive an Erdős-Gallai-type edge number condition for even cycles, which may be of independent interest.

Keywords: Ramsey Theory; spectral radius; cycles

AMS classification: 05C50; 05C35

This note is devoted to an open problem on cycles with consecutive lengths due to Nikiforov [8]. Throughout this note, for a graph G , $V(G)$ denotes the vertex set of G , $|G|$ and $e(G)$ denotes the number of vertices and edges in G , respectively. We denote by $\rho(G)$ the spectral radius of G . Let G_1 and G_2 be two graphs. We use $G_1 \vee G_2$ to denote the new graph obtained from the disjoint union of G_1 and G_2 by adding all edges xy where $x \in V(G_1)$ and $y \in V(G_2)$.

A classical theorem due to Bondy [3] says that every hamiltonian graph G on n vertices contains cycles of all lengths $\ell \in [3, n]$ if $e(G) \geq \frac{n^2}{4}$, unless n is even and G is isomorphic to $K_{\frac{n}{2}, \frac{n}{2}}$. If one drops the condition that “ G is hamiltonian” in Bondy’s theorem, a result in Bollobás’ textbook [2, Corrolary 5.4] states that such a graph contains all cycles C_ℓ for each $\ell \in [3, \lfloor \frac{n+3}{2} \rfloor]$. Nikiforov [8] considered cycles of consecutive lengths from a spectral perspective and posed the following open problem.

Problem 1 (Nikiforov [8]). What is the maximum C such that for all positive $\varepsilon < C$ and sufficiently large n , every graph G of order n with $\rho(G) > \sqrt{\lfloor \frac{n^2}{4} \rfloor}$ contains a cycle of length ℓ for every integer $3 \leq \ell \leq (C - \varepsilon)n$.

One may guess $C = \frac{1}{2}$. However, the class of graphs $G = K_s \vee (n - s)K_1$ where $s = \lfloor \frac{(3-\sqrt{5})n}{4} \rfloor$ (see [8]) shows $C \leq \frac{(3-\sqrt{5})}{2}$. Nikiforov [8] proved that $C \geq \frac{1}{320}$. Ning and Peng [9] slightly refined this as $C \geq \frac{1}{160}$. Only very recently, Zhai and Lin [14] improved these results to $C \geq \frac{1}{7}$.

The purpose of this note is to show $C \geq \frac{1}{4}$ by completely different methods.

*School of Mathematics and Statistics, Northwestern Polytechnical University, Xi’an, Shaanxi 710072, P.R. China. Email: binlongli@nwpu.edu.cn. Partially supported by the NSFC grant (No. 12171393) and Natural Science Foundation of Shaanxi Provincial Department of Education (2021JM-040).

†Corresponding author. College of Computer Science, Nankai University, Tianjin 300350, P.R. China. Email: bo.ning@nankai.edu.cn. Partially supported by the NSFC grant (No. 11971346).

One key ingredient is from Ramsey Theory. Very recently, Allen, Łuczak, Polcyn and Zhang [1] studied the Ramsey numbers of large even cycles versus stars. Their key technique of finding large even cycles is to combine a theorem on cycle lengths with given parity under minimum degree condition due to Voss and Zuluaga [13] and a theorem on the distribution of cycles with consecutive lengths due to Gould, Haxell, and Scott [4]. The proof of our main theorem is somewhat inspired by this idea.

Theorem 1. ¹ Let ε be real with $0 < \varepsilon < \frac{1}{4}$. Then there exists an integer $N := N(\varepsilon)$, such that if G is a graph on n vertices with $n \geq N$ and $\rho(G) > \sqrt{\lfloor \frac{n^2}{4} \rfloor}$, then G contains all cycles C_ℓ with $\ell \in [3, (\frac{1}{4} - \varepsilon)n]$.

Before the proof, we first collect various known results on cycles that will be used in our arguments. We use $ec(G)$ and $oc(G)$ to denote the length of a longest even cycle and the length of a longest odd cycle, of G , respectively.

Theorem A (Voss and Zuluaga [13]). (1) Every 2-connected graph G with $\delta(G) \geq k \geq 3$ having at least $2k+1$ vertices satisfies that $ec(G) \geq 2k$. (2) Every 2-connected non-bipartite graph G with $\delta(G) \geq k \geq 3$ having at least $2k+1$ vertices satisfies that $oc(G) \geq 2k-1$.

Theorem B (Ore [11]). Let G be a graph on n vertices. If G contains no Hamilton cycle, then $e(G) \leq \binom{n-1}{2} + 1$.

Theorem C (Gould, Haxell and Scott [4]). ² For every real number $c > 0$, there exists a constant $K := K(c)$ such that the following holds. Let G be a graph with $n \geq \frac{45K}{c^4}$ vertices and minimum degree at least cn . Then G contains a cycle of length t for every even $t \in [4, ec(G) - K]$ and every odd $t \in [K, oc(G) - K]$.

Another tools are the following two spectral inequalities, of which the second one was originally conjectured by Guo et al. [5].

Theorem D (Hong [6]). Let G be a graph on n vertices and m edges. If $\delta(G) \geq 1$ then $\rho(G) \leq \sqrt{2m - n + 1}$.

Theorem E (Sun and Das [12]). Let G be a graph with minimum degree $\delta(G) \geq 1$. For any $v \in V(G)$, we have $\rho^2(G - v) \geq \rho^2(G) - 2d(v) + 1$.

Theorem F (Nosal [10]). Let G be a graph on m edges. If $\rho(G) > \sqrt{m}$, then G contains a triangle.

Our proof of Theorem 1 uses the following three lemmas. A graph is called a *theta-graph* if it consists of three paths starting and ending with two same vertices and internally vertex-disjoint. The first lemma is very basic.

Lemma 1. Let G be a graph containing no theta-graphs. Then each block of G is an isolated vertex, or an edge (a K_2), or a cycle.

Lemma 2. Let G be a graph on n vertices. If $ec(G) \leq 2k$ where $k \geq 1$ is an integer, then $e(G) \leq \frac{(2k+1)(n-1)}{2}$.

¹If $0 < \varepsilon < 10^{-6}$, then we can choose $N = 2.5 \times 10^{10} \varepsilon^{-1}$.

²In [4], the authors set $K = \frac{7.5 \times 10^5}{c^5}$.

Proof. If $n \leq 2k + 1$, then $e(G) \leq \binom{n}{2} \leq \frac{(2k+1)(n-1)}{2}$. If $n = 2k + 2$, then by Theorem B, we have $e(G) \leq \binom{2k+1}{2} + 1 \leq \frac{(2k+1)(n-1)}{2}$. Next, we assume $n \geq 2k + 3$.

Let $k = 1$. We shall prove that if a graph on n vertices contains no even cycles then $e(G) \leq \frac{3(n-1)}{2}$. By Lemma 1, every block of G is an isolated vertex, an edge or an odd cycle. Let c be the number of blocks which are odd cycles. By induction on c , we can prove that $e(G) \leq n + c - 1 \leq n - 1 + \frac{n-1}{2} = \frac{3(n-1)}{2}$. In the following, we suppose $k \geq 2$.

Let $v \in V(G)$ with $d_G(v) = \delta(G)$, and $G' := G - v$. Note that G' satisfies that $|G'| \geq 2k + 2$ and G' contains no even cycle of length more than $2k$. By induction hypothesis, if $d(v) \leq k$, then we have $e(G) = e(G') + \delta \leq \frac{(2k+1)(n-2)}{2} + k < \frac{(2k+1)(n-1)}{2}$, as required. Thus, $\delta(G) \geq k + 1 \geq 3$. If G is 2-connected, then by Theorem A, G contains an even cycle of length at least $2k + 2$, a contradiction. Thus, G contains a cut-vertex or is disconnected. In such a case, let $G = G_1 \cup G_2$, where $|V(G_1) \cap V(G_2)| \leq 1$, and we use induction to G_1 and G_2 to get the required result. The proof is complete. \blacksquare

Lemma 3. *Let G be a graph. For any $v \in V(G)$, we have $\rho^2(G) \leq \rho^2(G - v) + 2d_G(v)$.*

Proof. If $\delta(G) \geq 1$, then by Theorem E, the result follows. Suppose that $\delta(G) = 0$. If $d_G(v) = 0$, then $\rho^2(G) = \rho^2(G - v) \leq \rho^2(G - v) + 2d_G(v)$. Now assume that $d_G(v) \geq 1$. Let I be the set of isolated vertices in G and $G' := G - I$. Then $\delta(G') \geq 1$. By Theorem E, $\rho^2(G) = \rho^2(G') \leq \rho^2(G' - v) + 2d_{G'}(v) = \rho^2(G - v) + 2d_G(v)$. This proves Lemma 3. \blacksquare

Now we give the proof of Theorem 1.

Proof of Theorem 1. If G is disconnected, then we use $\omega(G)$ to denote the number of components in G . Then we can add $\omega(G) - 1$ edges to make it connected. Note that $\rho(G') > \rho(G)$. For any integer $k \geq 3$, G contains a C_ℓ if and only if G' contains a C_ℓ . Thus, we can assume G is connected.

By Theorem D, we have

$$\frac{n^2 - 1}{4} \leq \rho^2(G) \leq 2m - n + 1.$$

One can derive that $2m \geq \frac{n^2 + 4n - 5}{4}$. Thus, the average degree $d(G) := \frac{2m}{n} > \frac{n}{4}$.

Let H be a subgraph of G defined by a sequence of graphs G_0, G_1, \dots, G_k such that:

- (1) $G = G_0$ and $H = G_k$;
- (2) for every $i \in [0, k - 1]$, there is $v_i \in V(G_i)$ such that $d_{G_i}(v_i) \leq \frac{n}{8}$ and $G_{i+1} = G_i - v_i$;
- (3) for every $v \in V(G_k)$, $d_{G_k}(v) > \frac{n}{8}$.

We claim that $d(H) > \frac{n}{4}$. Suppose not the case. Then there is a least integer $i \in [1, k]$ with $d(G_i) \leq \frac{n}{4}$. This implies that

$$d(G_{i-1}) = \frac{2d_{G_{i-1}}(v_{i-1}) + |G_i|d(G_i)}{|G_i| + 1} \leq \frac{n}{4},$$

a contradiction. Thus, we conclude that $d(H) > \frac{n}{4}$ and $\delta(H) > \frac{n}{8}$.

Case A: Even cycle. Note that $e(H) = \frac{d(H)|H|}{2} > \frac{\frac{n}{4}(|H|-1)}{2}$. By Lemma 2, $ec(H) > \lfloor \frac{n}{4} \rfloor \geq \frac{n}{4} - 1$. Recall that $\delta(H) > \frac{n}{8}$. By Theorem C, H contains all even cycles C_ℓ with $\ell \in [4, ec(H) - K]$ if $|H| \geq 45 \cdot 8^4 \cdot K$, where $K = K(\frac{1}{8}) = \frac{7.5 \times 10^5}{(\frac{1}{8})^5}$ corresponds to the constant $K(c)$ in Theorem C. Clearly $|H| > \frac{n}{4}$. Let n_1 be an integer satisfying

$$(i) \frac{n_1}{4} \geq 45 \cdot 8^4 \cdot K; \quad (ii) \varepsilon n_1 \geq K + 1.$$

Now if $n \geq n_1$ then G contains C_ℓ with every even $\ell \in [4, (\frac{1}{4} - \varepsilon)n]$.

Case B: Odd cycle. Set $h = |H|$. By Lemma 3, we have

$$\rho^2(G) \leq \rho^2(H) + 2 \sum_{i=0}^{k-1} d_{G_i}(v_i) \leq \rho^2(H) + 2k \cdot \frac{n}{8} = \rho^2(H) + \frac{kn}{4},$$

where G_i, v_i are those in the definition of H , and $k = n - h$. This implies that $\rho(H) \geq \frac{\sqrt{nh-1}}{2}$. Since $h \leq n$ and $\rho(H) > \frac{\sqrt{nh-1}}{2} \geq \sqrt{\lfloor \frac{h^2}{4} \rfloor}$, by Theorem F and Mantel's theorem, H contains a triangle, and so is non-bipartite.

Let F be a subgraph of H defined by a sequence of graphs H_0, H_1, \dots, H_s such that:

- (1) $H_0 = H$ and $H_s = F$;
- (2) for every $i \in [0, s-1]$, there is a cut-vertex v_i of H_i and $H_{i+1} := H_i - v_i$;
- (3) H_s has no cut-vertex.

Note that $w(H_{i+1}) \geq w(H_i) + 1$. Clearly $w(H) \leq 8$, for otherwise H will have a vertex of degree less than $\frac{n}{8}$. We claim that $w(F) \leq 8$. Suppose to the contrary that there is a least integer $i \in [0, s]$ such that $w(H_i) \geq 9$. Notice that for $i \leq 8$, $\delta(H_i) > \frac{n}{8} - 8$. As $w(H_i) \geq 9$, H_i has a vertex with degree less than $\frac{|H_i|}{9} < \frac{n}{9}$, a contradiction when $n \geq 577$. Thus we conclude that $w(F) \leq 8$, and specially, $|V(F)| \geq h - 7$.

By Lemma 3, we have

$$\rho^2(H) \leq \rho^2(F) + 2 \sum_{i=0}^{s-1} d_{H_i}(v_i) \leq \rho^2(F) + 2s(h-1) \leq \rho^2(F) + 14(h-1).$$

Since $d(H) > \frac{n}{4}$, we obtain $h > \frac{n}{4} + 1$. Thus,

$$\begin{aligned} \rho(F) &\geq \sqrt{\rho^2(H) - 14(h-1)} \geq \sqrt{\frac{nh-1}{4} - 14(h-1)} \\ &= \sqrt{\left(\frac{n}{4} - 14\right)h + \frac{55}{4}} > \sqrt{\left(\frac{n}{4} - 14\right)\left(\frac{n}{4} + 1\right) + \frac{55}{4}} \geq \frac{n}{4} - 7 \end{aligned}$$

when $n \geq 197$.

Recall that F has no cut-vertex and $\delta(F) \geq \delta(H) - 7 \geq \frac{n}{8} - 7 \geq 2$, i.e., every component of F is 2-connected. Let F_1 be a component of F with $\rho(F_1) = \rho(F)$. Thus we have $\delta(F_1) \geq \frac{n}{8} - 7$ and $\rho(F_1) > \frac{n}{4} - 7$. Specially $|F_1| > \frac{n}{4} - 6$.

We claim that $\delta(F_1) \geq \frac{|F_1|}{8}$. Recall that $\delta(H) \geq \frac{n}{8} \geq \frac{|H|}{8}$, we assume that $F_1 \neq H$. This implies that F has a second component, say F_2 . Since $\delta(F) \geq \frac{n}{8} - s$, we have $|F_2| \geq \frac{n}{8} - s + 1$ (here s is the subscript of $F = H_s$). This implies that $|F_1| \leq h - s - (\frac{n}{8} - s + 1) < \frac{7h}{8}$. Thus $\delta(F_1) \geq \frac{n}{8} - 7 \geq \frac{7n/8}{8} \geq \frac{|F_1|}{8}$ when $n \geq 448$.

Now we show that F_1 is non-bipartite. Recall that H is non-bipartite. So we assume that $F_1 \neq H$. By the analysis above we have $|F_1| < \frac{7h}{8}$. Thus

$$\rho^2(F_1) = \rho^2(F) \geq \left(\frac{n}{4} - 14\right)h + \frac{55}{4} \geq \left(\frac{h}{4} - 14\right)h + \frac{55}{4} > \frac{(7h/8)^2}{4} > \frac{|F_1|^2}{4}$$

when $h \geq 238$. Recall that $h > \frac{n}{4} + 1$, we have that F is non-bipartite when $n \geq 944$.

By Theorem A, $oc(F_1) \geq \min\{2\delta(F_1) - 1, |F_1|\} \geq \min\{\frac{n}{4} - 15, \rho(F_1)\} = \frac{n}{4} - 15$. By Theorem C, F_1 contains all odd cycles C_ℓ for $\ell \in [K, \frac{n}{4} - 15 - K]$ if $|F_1| \geq 45 \cdot 8^4 \cdot K$, where $K = K(\frac{1}{8})$ is the constant as in Theorem C. A theorem of Nikiforov [8, Theorem 1]³

³By refining the proof, one can let $N = 8400$.

states that there exists a sufficiently large N such that any graph of order $n \geq N$ such that $\rho(G) > \sqrt{\lfloor \frac{n^2}{4} \rfloor}$ has a cycle of length ℓ for every $\ell \in [3, \frac{n}{320}]$. Let n_2 be an integer such that

$$(i) \frac{n_2}{4} - 6 \geq 45 \cdot 8^4 \cdot K; (ii) n_2 \geq \max\{944, N\}; (iii) \frac{n_2}{320} \geq K; (iv) \varepsilon n_2 \geq K + 15.$$

Now if $n \geq \max\{n_1, n_2\}$, then G contains all cycles C_ℓ with $\ell \in [3, (\frac{1}{4} - \varepsilon)n]$. The proof is complete. ■

Note added

The result presented in this note was a part of work in [7] due to the current authors. Till now, the problem of determining the value of C is still open.

After the first version of this paper was written and submitted for publication, we have learned our paper was cited in [15], and the method developed here (together with some powerful technique developed by Nikiforov in [8]) has been used by Zhang and Zhao [15] to obtain a spectral condition for the existence of cycles with consecutive odd lengths in non-bipartite graphs.

Acknowledgment

The authors express their gratitude to the anonymous reviewers for their detailed and constructive comments which are very helpful to the improvement of the presentation of this note, and to Prof. Agelos Georgakopoulos, the Managing Editor, for his excellent editorial job. The second author is also grateful to Xing Peng who gave an online talk on Ramsey Theory for Tsinghua University. After Xing's talk, the second author drew his attention to the topic of Ramsey Theory of books versus even cycles and realized the technique in [1].

References

- [1] P. Allen, T. Łuczak, J. Polcyn, Y.-B. Zhang, The Ramsey number of a long even cycle versus a star, arXiv:2003.03310.
- [2] B. Bollobás, Extremal graph theory. London Mathematical Society Monographs, 11. Academic Press, Inc., London-New York, 1978. xx+488 pp. ISBN: 0-12-111750-2
- [3] J.A. Bondy, Pancyclic graphs. I, *J. Combinatorial Theory Ser. B* **11** (1971), 80–84.
- [4] R.J. Gould, P.E. Haxell, A.D. Scott, A note on cycle lengths in graphs, *Graphs Combin.* **18** (2002), no. 3, 491–498.
- [5] J.M. Guo, Z.W. Wang, X. Li, Sharp upper bounds of the spectral radius of a graph, *Discrete Math.* **342** (2019), no. 9, 2559–2563.
- [6] Y. Hong, Bounds of eigenvalues of graphs, *Discrete Math.* **123** (1993), no. 1-3, 65–74.
- [7] B. Li, B. Ning, The stability method, eigenvalues and cycles of consecutive lengths, arXiv:2102.03855v1.

- [8] V. Nikiforov, A spectral condition for odd cycles in graphs, *Linear Algebra Appl.* **428** (2008), no. 7, 1492–1498.
- [9] B. Ning, X. Peng, Extensions of the Erdős-Gallai theorem and Luo’s theorem, *Combin. Probab. Comput.* **29** (2020), no. 1, 128–136.
- [10] E. Nosal, Eigenvalues of Graphs, Master Thesis, University of Calgary, 1970.
- [11] O. Ore, Arc coverings of graphs, *Ann. Mat. Pura Appl.*, **55** (1961), 315–321.
- [12] S.W. Sun, K. Das, A conjecture on the spectral radius of graphs, *Linear Algebra Appl.* **588** (2020), 74–80.
- [13] H. Voss, C. Zuluaga, Maximale gerade und ungerade Kreise in Graphen I (German), *Wiss. Z. - Tech. Hochsch. Ilmenau* **23** (1977) 57–70.
- [14] M. Zhai, H. Lin, A strengthening of the spectral chromatic critical edge theorem: books and theta graphs, see <https://doi.org/10.1002/jgt.22883>, to appear in JGT.
- [15] Z. Zhang, Y. Zhao, A spectral condition for the existence of cycles with consecutive odd lengths in non-bipartite graphs, arXiv:2110.11345.