THE SUBSPACE STRUCTURE OF MAXIMUM CLIQUES IN PSEUDO-PALEY GRAPHS FROM UNIONS OF CYCLOTOMIC CLASSES

SHAMIL ASGARLI AND CHI HOI YIP

ABSTRACT. Blokhuis showed that all maximum cliques in Paley graphs of square order have a subfield structure. Recently it is shown that in Peisert-type graphs, some maximum cliques do not have a subfield structure, and yet all maximum cliques have a subspace structure. In this paper, we investigate the existence of a clique of size \sqrt{q} with a subspace structure in pseudo-Paley graphs of order q from unions of semi-primitive cyclotomic classes. We show that such a clique must have an equal contribution from each cyclotomic class and that most pseudo-Paley graphs do not admit such cliques, suggesting that the Delsarte bound \sqrt{q} on the clique number can be improved in general. We also prove that generalized Peisert graphs are not isomorphic to Paley graphs or Peisert graphs, confirming a conjecture of Mullin.

1. INTRODUCTION

Throughout the paper, p denotes an odd prime, q denotes a positive power of p, and \mathbb{F}_q denotes the finite field with q elements. For a finite field \mathbb{F}_q , we write \mathbb{F}_q^+ for its additive group, and \mathbb{F}_q^* for its multiplicative group. We always assume that g is a fixed primitive root of \mathbb{F}_q .

Let $q \equiv 1 \pmod{4}$. The *Paley graph* of order q, denoted P_q , is the Cayley graph $\operatorname{Cay}(\mathbb{F}_q^+, (\mathbb{F}_q^*)^2)$, where $(\mathbb{F}_q^*)^2$ is the set of squares in \mathbb{F}_q^* . Paley graphs are well-studied, and they connect many branches of mathematics [Jon20], notably combinatorics and number theory.

In this paper, we will consider *pseudo-Paley graphs*, that is, graphs that share the same spectrum with some Paley graphs. More precisely, we will consider pseudo-Paley graphs arising from unions of semi-primitive cyclotomic classes, following a general construction of strongly regular Cayley graphs by Brouwer, Wilson, and Xiang [BWX99]. We shall first recall the definition of cyclotomic classes and revisit a special case of their construction.

Let $N \mid (q-1)$. Let C_0 be the subgroup of \mathbb{F}_q^* with index N, and let C_1, \ldots, C_{N-1} be all the cosets of C_0 , where $C_j = g^j C_0$. The sets $C_0, C_1, \ldots, C_{N-1}$ are called the *N*-th cyclotomic classes of \mathbb{F}_q . Furthermore, we say that *N*-th cyclotomic classes are *semi-primitive* if -1 is a power of p modulo N.

Definition 1.1. Suppose q is a prime power, d a positive integer such that $2d \mid (q-1)$, and $I = \{m_1, \ldots, m_d\} \subset \{0, 1, \ldots, 2d-1\}$ with |I| = d. Let $C_0, C_1, \ldots, C_{2d-1}$ be the 2d-th cyclotomic classes of \mathbb{F}_q . The graph PP(q, 2d, I) is defined to be the Cayley graph $Cay(\mathbb{F}_q^+, D)$ where

$$D = \bigcup_{j=1}^{d} C_{m_j}.$$
 (1)

²⁰²⁰ *Mathematics Subject Classification*. Primary 05C25, 11T22; Secondary 11T24, 11B30, 05E30, 05C60. *Key words and phrases*. Paley graph, Peisert graph, maximum clique, clique number, cyclotomy.

Furthermore, we say that X = PP(q, 2d, I) is a semi-primitive pseudo-Paley graph if the cyclotomic classes building X are semi-primitive (or equivalently, -1 is a power of p modulo 2d). In this case, we always write $q = p^{2rt}$ where t is the smallest integer satisfying $p^t \equiv -1 \pmod{2d}$.

Note that in the above definition, C_j clearly depends on the choice of g; however, the isomorphism class of the resulting graph will only depend on q, d, and the set I. Indeed, any two primitive roots g and g' can be interchanged by a unique field automorphism which gives rise to a graph isomorphism. For a similar reason, we can assume $0 \in I$ without loss of generality.

The notation **PP** stands for **P**seudo-**P**aley. Indeed, X = PP(q, 2d, I) is a pseudo-Paley graph when -1 is a power of p modulo 2d; this immediately follows from [**BWX99**, Theorem 2]. Note that Paley graphs can also be obtained from the above construction, since $P_q = PP(q, 2, \{0\})$. We refer to Example 2.2 for other pseudo-Paley graphs that are of particular interest, including Peisert graphs and generalized Peisert graphs.

One important open problem in additive combinatorics is to obtain good estimates on the clique number of Paley graphs and other similar graphs [CL07, Section 2.7]. Since P_q is strongly regular, the Delsarte bound [Del73] implies that $\omega(P_q) \leq \sqrt{q}$; in fact, the same upper bound applies to any pseudo-Paley graph of order q. This square root upper bound is known as the *trivial upper bound* on the clique number of P_q and it is notoriously difficult to improve this trivial upper bound when q is a non-square. Recently, a minor improvement on the trivial upper bound (from \sqrt{q} to $\sqrt{q} - 1$) has been made by Greaves and Soicher [GS18] for infinitely many pseudo-Paley graphs with non-square order. More recently, Hanson and Petridis [HP21] and Yip [Yip22c] improved the \sqrt{q} bound to $\sqrt{q/2}(1 + o(1))$ for $\omega(P_q)$ when q is a non-square using polynomial method. This is still far from the conjectural bound; see the related discussion in [Yip22c, Section 1] and [Yip22a, Section 1.3]. Nevertheless, it is known that $\omega(P_q) = \sqrt{q}$ when q is a square since the subfield $\mathbb{F}_{\sqrt{q}}$ forms a clique; in fact, Blokhuis [Blo84] showed all maximum cliques have a subfield structure.

Theorem 1.2 ([Blo84]). Let $q = p^{2r}$, where p is an odd prime. Then the only maximum clique in the Paley graph P_q containing $\{0, 1\}$ is the subfield $\mathbb{F}_{\sqrt{q}}$. Consequently, each maximum clique is the image of an affine transformation on the subfield $\mathbb{F}_{\sqrt{q}}$.

Let us consider the same question for a semi-primitive pseudo-Paley graph X = PP(q, 2d, I)constructed above. Without loss of generality, we assume that $0 \in I$. As always, we write $q = p^{2rt}$, where t is the smallest integer satisfying $p^t \equiv -1 \pmod{2d}$. Again, the Delsarte bound implies that the trivial upper bound $\omega(X) \leq \sqrt{q}$ holds. We continue the discussion on $\omega(X)$ according to the parity of r.

We first consider the case when r is odd. Note that in this case, $\mathbb{F}_{\sqrt{q}}^*$ is contained in the cyclotomic class $C_0 = \langle g^{2d} \rangle$ because $\sqrt{q} = p^{rt} \equiv -1 \pmod{2d}$; therefore, $\mathbb{F}_{\sqrt{q}}$ forms a clique in X and $\omega(X) = \sqrt{q}$. Indeed, in this case, the connection set D of X can be written as a union of $\mathbb{F}_{\sqrt{q}}^*$ -cosets in \mathbb{F}_q^* , and thus, by definition X is a *Peisert-type graph*; see [AY22, Definition 1.1]. The structure of maximum cliques in a Peisert-type graph has been studied recently in [AY22, AGLY22]. In particular, each maximum clique in X has a subspace structure [AY22, Theorem 1.2].

Theorem 1.3 ([AY22, Theorem 1.2]). Let X = PP(q, 2d, I) be a semi-primitive pseudo-Paley graph with $q = p^{2rt}$ and odd r. Each maximum clique in X containing 0 is an \mathbb{F}_p -subspace of \mathbb{F}_q .

There are counterexamples for which X admits a maximum clique with no subfield structure; see for example [AY22, Example 2.21] and [AGLY22, Section 5]. However, under extra assumptions, the *subspace* structure of maximum cliques can be upgraded to the *subfield* structure; in this direction, Asgarli and Yip managed to obtain an analogue of Theorem 1.2 in [AY22, Theorem 1.3].

In this paper we shall focus on the case when r is even. This change in parity introduces a huge difference: there is no obvious choice of a clique of size \sqrt{q} in general since $\mathbb{F}_{\sqrt{q}}$ fails to be a clique in X in general. Additionally, Lemma 2.7 shows how a naive analogue of $\mathbb{F}_{\sqrt{q}}$ fails to be a valid clique. In particular, when $X = PP(p^{2r}, 4, \{0, 1\})$ (a Peisert graph, see Section 2.1) with $p \equiv 3 \pmod{4}$ and r even, it is conjectured that $\omega(X) \leq \sqrt{q} - 1$ by Yip [Yip22b]; see the related discussions by Kisielewicz and Peisert [KP04, Theorem 5.1], and Mullin [Mul09, Section 3.4]. Note that a simple (yet crucial) fact in all recent improvements of the clique number of Paley graphs and related graphs (of non-square order q) is that \sqrt{q} is not an integer when q is a non-square [GS18, HP21, Yip21, Yip22c], so that the small gap between \sqrt{q} and its integer part can be "blown up" to gain an improvement on the trivial upper bound. In our case, q is a square, which prevents all such techniques from being applied. To our best knowledge, there is no single infinite family of semi-primitive pseudo-Paley graphs (of square order) for which the trivial upper bound has been improved.

The above discussion, together with Theorem 1.3, leads to the following conjecture:

Conjecture 1.4. Let X = PP(q, 2d, I) be a semi-primitive pseudo-Paley graph with $q = p^{2rt}$ where r is even. If $\omega(X) = \sqrt{q}$, then every maximum clique in X is an \mathbb{F}_{p^t} -affine subspace.

We refer to Section 2.3 for the further motivation behind this conjecture. Conjecture 1.4 is computationally useful in understanding the clique number: it provides a polynomial-time algorithm to check if the clique number is given by \sqrt{q} . According to Conjecture 1.4, the only candidates for maximum cliques of size \sqrt{q} are \mathbb{F}_{p^t} -subspaces of a fixed dimension, whose number is bounded by a polynomial function of q. We are thus led to investigate the existence of cliques with a subspace structure and with size \sqrt{q} in X.

Problem 1.5. Suppose that X = PP(q, 2d, I) is a semi-primitive pseudo-Paley graph with $q = p^{2rt}$ where r is even. Determine if X has a clique of size \sqrt{q} with a subspace structure.

We will show that the answer to the above question is "no" in a probabilistic sense (see Proposition 5.7), thus lending some evidence to the inequality $\omega(X) \le \sqrt{q} - 1$ in general. However, numerical experiments suggest that there is an infinite family of semi-primitive pseudo-Paley graphs (other than Paley graphs) where the trivial upper bound is tight; see Conjecture 5.8.

1.1. **Main results.** Our first main result guarantees that a maximum clique of size \sqrt{q} has an equal contribution from each cyclotomic class building the graph.

Theorem 1.6. Let X = PP(q, 2d, I) be a semi-primitive pseudo-Paley graph with $q = p^{2rt}$ where r is even. Then $\omega(X) \leq \sqrt{q}$. Moreover, if $\omega(X) = \sqrt{q}$, and A is a maximum clique in X such that $0 \in A$, then

$$|A \cap C_{m_1}| = |A \cap C_{m_2}| = \dots = |A \cap C_{m_d}| = \frac{\sqrt{q-1}}{d}.$$
 (2)

When X is a Paley graph, Theorem 1.6 follows from Theorem 1.2; see Lemma 2.6. Furthermore, Theorem 1.6 is a genuine extension: most graphs PP(q, 2d, I) are not isomorphic to the Paley graph P_q by Proposition 4.7. This latter fact also helps us to prove a conjecture by Mullin on generalized Peisert graphs (see Corollary 4.8). It is crucial to assume that r is even in the hypothesis of Theorem 1.6. Indeed, when r is odd, we have mentioned that $\mathbb{F}_{\sqrt{q}} \subset C_0 \cup \{0\}$ forms a clique in X and yet the contribution solely comes from a single cyclotomic class. On the other hand, when r is even, any clique of size \sqrt{q} has an equal contribution from each cyclotomic class according to our main theorem. We see again that cliques behave very differently when r is odd and when r is even.

It is worth pointing out that neither Conjecture 1.4 nor Theorem 1.6 imply each other. However, let us discuss how Theorem 1.6 shows that a maximum clique containing 0 behaves like a subspace, thus lending some evidence towards Conjecture 1.4. Indeed, if $V \subset \mathbb{F}_q$ is a vector space over \mathbb{F}_{p^t} , we expect that the contribution to V from each cyclotomic class forming the connection set D would be *approximately* equal. This is due to the general principle that the intersection of a subspace and a multiplicative subgroup can be modelled as the intersection of two random sets [Kop19, Lemma 11]. On the other hand, Theorem 1.6 asserts that any maximum clique A in X has *exactly* equal contribution from each cyclotomic class, so Conjecture 1.4 alone does not imply Theorem 1.6.

Assuming Conjecture 1.4, we apply Theorem 1.6 to improve the upper bound on the clique number of most semi-primitive pseudo-Paley graphs (unconditionally, we prove that there is no clique of size \sqrt{q} with a subspace structure for most graphs). First, we show that the density of such graphs with clique number \sqrt{q} is zero in Proposition 5.7. Next, we prove an effective version of this statement. The condition that p is sufficiently large in the result below is necessary; see Table 1 for counterexamples.

Theorem 1.7. Let X = PP(q, 2d, I) be a semi-primitive pseudo-Paley graph with $q = p^{2rt}$ where r is even, and $I \neq \{0, 2, ..., 2d - 2\}$ and $I \neq \{1, 3, ..., 2d - 1\}$. Assuming Conjecture 1.4, $\omega(X) \leq \sqrt{q} - 1$ for $p^t > 10.2r^2d$.

As a concrete application, we can conditionally improve the Delsarte bound for the clique number of certain generalized Peisert graphs $GP^*(q, 2d) := PP(q, 2d, \{0, 1, \dots, d-1\})$.

Theorem 1.8. Suppose that Conjecture 1.4 is true. If $d \ge 2$, and t is the smallest integer such that $p^t \equiv -1 \pmod{2d}$ and $p^t > 3$, then $\omega(GP^*(p^{4t}, 2d)) \le p^{2t} - 1$.

We also prove the following two theorems on the number of maximum cliques. Comparing these with Theorem 1.2, we see that pseudo-Paley graphs in general behave much differently compared to Paley graphs. Note that Theorem 1.9 generalizes Mullin's theorem [Mul09, Lemma 3.3.6] for the Peisert graph.

Theorem 1.9. Let X = PP(q, 2d, I) be a semi-primitive pseudo-Paley graph with $q = p^{2rt}$ where r is even. Suppose that $0 \in I$ and $I \neq \{0, 2, ..., 2d - 2\}$. If $\omega(X) = \sqrt{q}$, then there are at least 2 maximum cliques in X that contain $\{0, 1\}$.

Theorem 1.10. Let X = PP(q, 2d, I) be a semi-primitive pseudo-Paley graph with $q = p^{4t}$. Suppose that $0 \in I$ and $I \neq \{0, 2, ..., 2d - 2\}$. Then X has an even number of cliques (possibly zero) of size \sqrt{q} that are subspaces defined over \mathbb{F}_{p^t} containing 1. Furthermore, these maximum cliques come in pairs that are interchanged by the map $x \mapsto x^{\sqrt{q}}$.

1.2. **Structure of the paper.** We describe the flowchart of the paper. In Section 2, we introduce the relevant notations, provide further background, and present preliminary properties. We present the proof of Theorem 1.6 and Theorem 1.9 in Section 3. In Section 4, we prove that the families of graphs considered in this paper are almost always not isomorphic to Peisert graphs or Paley

graphs. In Section 5, we prove Theorem 1.7, Theorem 1.8, and Theorem 1.10. In Appendix A, we present all the necessary details on Gauss sums, and give a detailed proof of Proposition 3.2 in Appendix B. Finally, in Appendix C, we report results from our numerical experiments, including two algorithms.

2. BACKGROUND AND OVERVIEW OF THE PAPER

2.1. Peisert graphs and generalized Peisert graphs. We first recall the definitions of Peisert graphs [Pei01]. The *Peisert graph* of order $q = p^r$, where p is a prime such that $p \equiv 3 \pmod{4}$ and r is even, denoted by P_q^* , is the Cayley graph $\operatorname{Cay}(\mathbb{F}_q^+, M_q)$ with $M_q := \{g^j : j \equiv 0, 1 \pmod{4}\}$, where g is a primitive root of the field \mathbb{F}_q . We will discuss more properties of Peisert graphs in Section 4.

Motivated by the similarity shared among Paley graphs, generalized Paley graphs, and Peisert graphs, Mullin introduced generalized Peisert graphs; see [Mul09, Section 5.3] and [AY22, Definition 2.8]. Observe that $P_q = GP^*(q, 2)$ and $P_q^* = GP^*(q, 4)$.

Definition 2.1 (generalized Peisert graph). Let d be a positive integer, and q a prime power such that $q \equiv 1 \pmod{4d}$. The 2d-th power Peisert graph of order q, denoted $GP^*(q, 2d)$, is the Cayley graph $\operatorname{Cay}(\mathbb{F}_q^+, M_{q,2d})$, where $M_{q,2d} = \{g^{2dk+j} : 0 \leq j \leq d-1, k \in \mathbb{Z}\}$, and g is a primitive root of \mathbb{F}_q .

Next we will see that our Definition 1.1 is general enough to encompass both Peisert graphs and generalized Peisert graphs. In the following discussion, when we are talking about a cyclotomic class C_i in PP(q, 2d, I), we always assume C_i is a 2*d*-th cyclotomic class of \mathbb{F}_q .

Example 2.2. By specializing the choice of the set *I*, we recover several important families of Cayley graphs we mentioned previously:

- (1) $PP(q, 2, \{0\})$ is the Paley graph $P_q = Cay(\mathbb{F}_q^+, C_0)$.
- (2) $PP(q, 4, \{0, 1\})$ is the Peisert graph $P_q^* = \text{Cay}(\mathbb{F}_q^+, C_0 \cup C_1)$.
- (3) $PP(q, 2d, \{0, 2, 4, ..., 2d 2\})$ is also the Paley graph $P_q = Cay(\mathbb{F}_q^+, C_0 \cup C_2 \cup ... \cup C_{2d-2}).$
- (4) $PP(q, 2d, \{1, 3, 5, ..., 2d 1\})$ is the complement of the Paley graph P_q , and it is isomorphic to P_q .
- (5) $PP(q, 2d, \{0, 1, 2, ..., d-1\})$ is the generalized Peisert graph $GP^*(q, 2d) = Cay(\mathbb{F}_q^+, C_0 \cup C_1 \cup ... \cup C_{d-1}).$
- (6) Suppose that d = 2m is even. If $I = \{i, i+4, \dots, i+4(m-1)\} \cup \{j, j+4, \dots, j+4(m-1)\}$ for some $i < j \in \{0, 1, 2, 3\}$, then PP(q, 2d, I) is isomorphic to the Paley graph P_q when j - i = 2, and PP(q, 2d, I) is isomorphic to the Peisert graph P_q^* when $j - i \in \{1, 3\}$.

From Example 2.2, we see that two different choices of (d, I) may represent the same graph. In general, if $d' \mid d$ and $I = \bigcup_{j \in I'} \{j + 2d'k : 0 \le k \le \frac{d}{d'} - 1\}$, then PP(q, 2d, I) = PP(q, 2d', I'). It is natural to work with the representation that has the smallest value of d. This motivates the following definition.

Definition 2.3 (Minimal representation). We say X = PP(q, 2d, I) is in its minimal representation if there is no $d' \mid d$ and $I' \subset \{0, 1, \ldots, 2d' - 1\}$ such that d' < d and X = PP(q, 2d', I'). Note that the minimal representation of X is unique.

Define $I + k := \{j + k \pmod{2d} : j \in I\}$. The next lemma gives a necessary and sufficient condition for the equality I = I + k to hold.

Lemma 2.4. Let X = PP(q, 2d, I) and let PP(q, 2d', I') be its minimal representation. Then I = I + k if and only k is a multiple of 2d'.

Proof. Since $d' \mid d$ and $I = \bigcup_{j \in I'} \{j + 2d'r : 0 \le r \le \frac{d}{d'} - 1\}$, it follows that I = I + k for any k that is a multiple of 2d'.

If I = I + k, then I is a union of arithmetic progressions each with common difference gcd(2d, k). The length of each arithmetic progression appearing in the union is $\frac{2d}{gcd(2d,k)}$. Therefore, d = |I| is a multiple of $\frac{2d}{gcd(2d,k)}$. It follows that gcd(2d, k) is even, that is, gcd(2d, k) = 2e for some $e \in \mathbb{N}$. In particular, X = PP(q, 2d, I) = PP(q, 2e, J), where $J = \{j \pmod{2e} : j \in I\}$. Since PP(q, 2d', I') is the minimal representation of X, it follows that $2d' \mid 2e$, and so $2d' \mid k$. \Box

Remark 2.5. The condition I = I + k is connected to the symmetries of the graph X = PP(q, 2d, I). More precisely, the map $x \mapsto g^k x$ induces a graph automorphism of X if and only if I + k = I.

2.2. Semi-primitive pseudo-Paley graph. One of our aims in the present paper is to draw parallels between Paley graphs and the graphs of the form PP(q, 2d, I). One special property Paley graphs enjoy is the following "equal contribution" statement. Our main result, Theorem 1.6, extends this property to a larger family of pseudo-Paley graphs.

Lemma 2.6. Let X = PP(q, 2d, I) be a semi-primitive pseudo-Paley graph with $q = p^{2rt}$ where r is even and $I = \{0, 2, ..., 2d - 2\}$, so that X is the Paley graph P_q . If A is a maximum clique in X such that $0 \in A$, then

$$|A \cap C_0| = |A \cap C_2| = \dots = |A \cap C_{2d-2}| = \frac{\sqrt{q-1}}{d}.$$

Proof. Note that $g^{p^{rt}+1}$ is a primitive root of the subfield $\mathbb{F}_{\sqrt{q}}$. Since $p^t \equiv -1 \pmod{2d}$ and r is even, $p^{rt} + 1 \equiv 2 \pmod{2d}$. Let A be a maximum clique in X such that $0 \in A$. Since $0 \in A$, by Theorem 1.2, $A = a\mathbb{F}_{\sqrt{q}}$ for some square $a \in \mathbb{F}_q^*$. Observe that for each $0 \leq j \leq d-1$,

$$\mathbb{F}_{\sqrt{q}} \cap C_{2j} = \{ (g^{p^{rt}+1})^k : k \equiv j \pmod{d} \} \implies |\mathbb{F}_{\sqrt{q}} \cap C_{2j}| = \frac{\sqrt{q-1}}{d}.$$
(3)

The same conclusion holds for $A = a \mathbb{F}_{\sqrt{q}}$.

From Lemma 2.6, we see that the subfield $\mathbb{F}_{\sqrt{q}}$ admits the following presentation

$$\mathbb{F}_{\sqrt{q}} = \{0\} \cup \bigcup_{j=0}^{d-1} \{ (g^{p^{rt}+1})^k : k \equiv j \pmod{d} \} = \{0\} \cup \bigcup_{j=0}^{d-1} g^{2j} \langle g^{d(\sqrt{q}+1)} \rangle.$$

This motivates the following construction of a set in PP(q, 2d, I):

$$A(q, 2d, I) = \{0\} \cup \bigcup_{j=1}^{d} g^{m_j} \langle g^{d(\sqrt{q}+1)} \rangle.$$

$$\tag{4}$$

The following lemma shows this "naive construction" of A(q, 2d, I) does not give a clique in PP(q, 2d, I) except for the Paley graph.

Lemma 2.7. Let X = PP(q, 2d, I) be a semi-primitive pseudo-Paley graph with $q = p^{2rt}$ where r is even. Then A(q, 2d, I) defined in equation (4) is a clique in X if and only if $I = \{0, 2, ..., 2d-2\}$ or $I = \{1, 3, ..., 2d-1\}$.

Proof. If $I = \{0, 2, ..., 2d - 2\}$, then the conclusion follows from Lemma 2.6. The case $I = \{1, 3, ..., 2d - 1\}$ is similar.

Conversely, assume that I is different from $\{0, 2, ..., 2d-2\}$ and $\{1, 3, ..., 2d-1\}$. Given X = PP(q, 2d, I), without loss of generality, we may assume that $m_1 = 0$, and that X is already in its minimal representation: indeed, if PP(q, 2d, I) = PP(q, 2d', I'), then A(q, 2d', I') = A(q, 2d, I). Thus, by hypothesis, $d \ge 2$.

Assume, to the contrary, that A(q, 2d, I) is a clique in X. Let $B_j = A(q, 2d, I) \cap (C_j \cup \{0\})$ for each $0 \le j \le 2d - 1$. If $x, y \in B_0$ such that $x - y \in C_k$ with $0 \le k \le 2d - 1$, then for each $1 \le j \le d$, we have $g^{m_j}x, g^{m_j}y \in B_{m_j}$ and $g^{m_j}x - g^{m_j}y \in C_{k+m_j}$. As a result, $I + k = \{m_j + k \pmod{2d} : 1 \le j \le d\} = I$. By Lemma 2.4, 2d|k as PP(q, 2d, I) is the minimal representation of X. Thus, k = 0 and $B_0 - B_0 \subset B_0$.

From $B_0 - B_0 \subset B_0$ we infer that B_0 is a subgroup of \mathbb{F}_q^+ , and must have cardinality $|B_0| = p^n$ for some $n \ge 1$. We show this is impossible by analyzing the following Diophantine equation:

$$\frac{\sqrt{q} - 1}{d} + 1 = |B_0| = p^n.$$

Writing $\sqrt{q} = p^{rt}$, and rearranging the equation, we obtain $p^{rt} - 1 = d(p^n - 1)$. In particular, $d \equiv 1 \pmod{p^n}$, implying that $d \ge p^n + 1$. Combining $d \ge p^n + 1$ and $p^{rt} - 1 = d(p^n - 1)$, we obtain $rt \ge 2n$. Thus, $d \ge p^{rt/2} + 1$. Since $p^t \equiv -1 \pmod{2d}$, we have $p^t \ge 2d - 1 > 2p^{rt/2} \ge 2p^t$, a contradiction. This proves that A(q, 2d, I) is not a clique in X.

We will later see in the proof of Theorem 1.9 how Lemma 2.7 can help us distinguish X from the Paley graph. Indeed, we will show that most semi-primitive pseudo-Paley graphs are not isomorphic to the Paley graph; see Corollary 3.4 for a precise criterion.

2.3. **Further motivation.** Variants of Erdős-Ko-Rado (EKR) theorem have been studied intensively; the book by Godsil and Meagher [GM16] provides an excellent survey. Typically EKR-type results state that the extremal configurations must be canonical in the sense that they have a remarkably simple structure compared to the complicated ground space. Theorem 1.2 can be regarded as the EKR theorem of Paley graphs [GM16, Section 5.9]: if we view cliques with a subfield structure as canonical cliques, then Theorem 1.2 states that all maximum cliques are canonical.

In our setting, X = PP(q, 2d, I) is a Cayley graph $\operatorname{Cay}(\mathbb{F}_q^+, D)$, so $A \subset \mathbb{F}_q$ forms a clique in X if and only if $A - A \subset D \cup \{0\}$, where $A - A = \{a - a' : a, a' \in A\}$. We see that additive combinatorics plays an important role here because A - A has a rich additive structure, whereas D, being a union of cyclotomic classes, has a rich multiplicative structure. This partially explains why estimating the clique number of Paley graphs and other related graphs is a central open problem in additive combinatorics [CL07, Section 2.7]: the study of a clique is essentially the study of the interaction between additive structure and multiplicative structure of a set [AB14, Shk14]. All results in this paper conditional on Conjecture 1.4 can be reformulated unconditionally based on such interaction; for instance, compare Theorem 1.7 and Corollary 5.13.

Let K be a subfield of \mathbb{F}_q . Note that if V is a K-subspace, then V - V = V; thus, the set V forms a clique in X if and only if $V \subset D \cup \{0\}$. Therefore, it is much easier for a subspace to form a clique in X, compared to a generic subset of \mathbb{F}_q . We would like to choose the subfield K so that it serves the role of the prime subfield \mathbb{F}_p . In our case $q = p^{2rt} = (p^t)^{2r}$ and t is the smallest integer with $p^t \equiv -1 \pmod{2d}$. In view of Stickelberger's Theorem (Theorem A.3), the choice $K = \mathbb{F}_{p^t}$ is a natural replacement of the prime subfield \mathbb{F}_p . Therefore, we call V a *canonical clique* in X if V is a \mathbb{F}_{p^t} -affine subspace with $|V| = \sqrt{q}$ such that V forms a clique in X. In contrast to

a typical EKR-type result, in our situation canonical objects may not exist, because X may have clique number less than \sqrt{q} . Using this terminology, Conjecture 1.4 states that each clique of size \sqrt{q} in X is canonical.

We remark that our conjecture is an analogue of the celebrated Chvátal's conjecture [Chv74] on families of set systems, which is a variant of the EKR theorem and is still widely open (see Chvátal's webpage [Chv11] for related discussion and references on the conjecture). Chvátal's conjecture states that for any family \mathcal{F} of subsets of a finite set that is closed under taking subsets (known as an *ideal*), there is a largest intersecting subfamily of \mathcal{F} that is a star. In other words, in any nice set system, there is a canonical maximum intersecting family; our conjecture is actually stronger because it implies that in any nice semi-primitive pseudo-Paley graph (that is, with clique number \sqrt{q}), every maximum clique is canonical. In summary, Conjecture 1.4 can be viewed as a combination of both the EKR theorem and Chvátal's conjecture in the context of pseudo-Paley graphs. The following table is a short dictionary between concepts in Chvátal's conjecture (first column) and in the present work (second column).

uniform set system	Paley graph
set system	semi-primitive pseudo-Paley graph
nice set system	nice semi-primitive pseudo-Paley graph
(ideal, i.e., closed under taking subsets)	(with clique number \sqrt{q})
largest intersecting family	maximum clique
canonical intersecting family (star)	canonical clique (affine subspace)
The original EKR theorem	Blokhuis theorem (Theorem 1.2)
Chvátal's conjecture + EKR	Conjecture 1.4

3. PROOF OF THEOREM 1.6

The novel idea in the proof of Theorem 1.6 is to establish a system of linear equations in the expressions $|A \cap C_{m_1}|, |A \cap C_{m_1}|, \dots, |A \cap C_{m_d}|$ and then find a way to solve the system. A key ingredient is the following Fourier analytic characterization of maximum cliques in a semiprimitive pseudo-Paley graph, which is of independent interest. We need the following definition of exponential sums (scaled Fourier coefficients of the indicator function on A). Recall that $e_p(x) = \exp(2\pi i x/p)$ for all $x \in \mathbb{F}_p$, and $\operatorname{Tr}_{\mathbb{F}_q}(\cdot) \colon \mathbb{F}_q \to \mathbb{F}_p$ is the absolute trace map.

Definition 3.1. Let A be a subset of \mathbb{F}_q . For each $c \in \mathbb{F}_q^*$, we define

$$S(q, A; c) = \sum_{a \in A} e_p \big(\operatorname{Tr}_{\mathbb{F}_q}(ac) \big).$$
(5)

Proposition 3.2. Let X = PP(q, 2d, I) be a semi-primitive pseudo-Paley graph with $q = p^{2rt}$. If A is a clique in X, then $|A| \le \sqrt{q}$; moreover, $|A| = \sqrt{q}$ if and only if S(q, A; c) = 0 for all $c \in D'$, where S(q, A; c) is defined in equation (5) and $D' = \bigcup_{\ell=1}^{d} C_{-m_{\ell}}$.

The proof of Proposition 3.2 is technical, and is deferred to Appendix B. We remark that a similar characterization for maximum cliques in generalized Paley graphs has appeared in [Yip22a, Theorem 1.4].

Now we present the proof of Theorem 1.6 using Proposition 3.2 and the novel idea mentioned in the beginning of this section.

Proof of Theorem 1.6. Let A be a clique of size \sqrt{q} such that $0 \in A$. Let $A_j = A \cap C_j$ for each $0 \le j \le 2d - 1$. Since A is a clique in X and $0 \in A$, we have $A \setminus \{0\} = \bigcup_{j=1}^d A_{m_j}$, and thus

$$\sum_{j=1}^{d} |A_{m_j}| = \sqrt{q} - 1.$$
(6)

By Proposition 3.2, S(q, A; c) = 0 for all $c \in D'$ where $D' = C_{-m_1} \cup C_{-m_2} \cup \cdots \cup C_{-m_d}$. Without loss of generality, we may assume that $m_1 = 0$. In particular, we have S(q, A; c) = 0 for each $c \in C_0$, implying the following:

$$0 = \sum_{c \in C_0} S(q, A; c) = \sum_{c \in C_0} \sum_{a \in A} e_p \big(\operatorname{Tr}_{\mathbb{F}_q}(ac) \big) = \sum_{a \in A} \sum_{c \in C_0} e_p \big(\operatorname{Tr}_{\mathbb{F}_q}(ac) \big)$$
$$= |C_0| + \sum_{j=1}^d \sum_{a \in A_{m_j}} \sum_{c \in C_0} e_p \big(\operatorname{Tr}_{\mathbb{F}_q}(ac) \big).$$

Note that if $a \in A_{m_i}$, then as c runs over C_0 , ac runs over C_{m_i} . Therefore,

$$0 = |C_0| + \sum_{j=1}^d |A_{m_j}| \sum_{c \in C_{m_j}} e_p \big(\operatorname{Tr}_{\mathbb{F}_q}(c) \big).$$

We can apply the similar argument to C_{-m_k} for each $k \ge 2$ to obtain:

$$0 = \sum_{c \in C_{-m_k}} S(q, A; c) = \sum_{a \in A} \sum_{c \in C_{-m_k}} e_p \big(\operatorname{Tr}_{\mathbb{F}_q}(ac) \big)$$
$$= |C_{-m_k}| + \sum_{j=1}^d |A_{m_j}| \sum_{c \in C_{m_j-m_k}} e_p \big(\operatorname{Tr}_{\mathbb{F}_q}(c) \big).$$

Recall that the index j in C_j is defined up to modulo 2d, which allows us to consider $C_{m_j-m_k}$ even when $m_j - m_k < 0$. For convenience, we define

$$\lambda := \sum_{c \in C_0} e_p \big(\operatorname{Tr}_{\mathbb{F}_q}(c) \big), \quad \text{and} \quad \mu := \sum_{c \in C_k} e_p \big(\operatorname{Tr}_{\mathbb{F}_q}(c) \big)$$

for each $1 \le k \le 2d - 1$. The value of μ is well-defined in view of Corollary A.6.

Given a fixed integer k satisfying $1 \le k \le d$, there is a unique value of j such that $m_j - m_k \equiv 0 \pmod{2d}$, namely, j = k. Therefore, the equation

$$0 = |C_{-m_k}| + \sum_{j=1}^d |A_{m_j}| \sum_{c \in C_{m_j-m_k}} e_p(\operatorname{Tr}_{\mathbb{F}_q}(c))$$

becomes

$$0 = |C_0| + \lambda |A_{m_k}| + \sum_{\substack{1 \le j \le d \\ j \ne k}} \mu |A_{m_j}|$$
(7)

where we also used the fact that $|C_{-m_k}| = |C_0|$. Note that equation (7) is valid for each $1 \le k \le d$, and so these d linear equations correspond to the matrix equation Mx = b where

$$\mathbf{M} = \begin{pmatrix} \lambda & \mu & \mu & \dots & \mu \\ \mu & \lambda & \mu & \dots & \mu \\ \mu & \mu & \lambda & \dots & \mu \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ \mu & \mu & \mu & \dots & \lambda \end{pmatrix}, \quad \mathbf{x} = \begin{pmatrix} |A_{m_1}| \\ |A_{m_2}| \\ |A_{m_3}| \\ \vdots \\ |A_{m_d}| \end{pmatrix} \text{ and } \mathbf{b} = - \begin{pmatrix} |C_0| \\ |C_0| \\ |C_0| \\ \vdots \\ |C_0| \end{pmatrix}$$

Note that $\mathbf{M} = (\lambda - \mu)I_d + \mu J_d$, where I_d is the $d \times d$ identity matrix and J_d is the $d \times d$ all 1's matrix. It is straightforward to see that \mathbf{M} has eigenvalue $\lambda - \mu$ with multiplicity d - 1, and eigenvalue $\lambda + (d - 1)\mu$ with multiplicity 1. By Corollary A.6, we know that $\lambda \neq \mu$ and

$$\lambda + (d-1)\mu = -\frac{(2d-1)\sqrt{q}+1}{2d} + (d-1)\cdot\frac{\sqrt{q}-1}{2d} = -\frac{\sqrt{q}+1}{2}.$$

Therefore, M is invertible. Thus, Mx = b has a unique solution given by

$$\mathbf{x} = \begin{pmatrix} \gamma \\ \gamma \\ \vdots \\ \gamma \end{pmatrix}$$

where $\gamma = -|C_0|/(\lambda + (d-1)\mu)$. We deduce that

$$|A_{m_j}| = \gamma = -\frac{|C_0|}{\lambda + (d-1)\mu} = \frac{2|C_0|}{\sqrt{q}+1} = \frac{2(q-1)}{2d(\sqrt{q}+1)} = \frac{\sqrt{q}-1}{d}$$

for each $1 \le j \le d$. It is easy to verify that $|A_{m_1}|, |A_{m_2}|, \ldots, |A_{m_d}|$ satisfy the equation (6). \Box

As a quick application, Theorem 1.6 immediately implies the following non-trivial upper bound on the clique number of a particular type of Cayley graphs.

Corollary 3.3. Let X = PP(q, 2d, I) be a semi-primitive pseudo-Paley graph with $q = p^{2rt}$ where r is even. Pick subsets $D_{m_j} \subset C_{m_j}$ for each $1 \leq j \leq d$ such that $|D_{m_k}| < (\sqrt{q} - 1)/d$ holds for some k. Consider the Cayley graph $X' = \operatorname{Cay}\left(\mathbb{F}_q^+, \bigcup_{j=1}^d D_{m_j}\right)$, which is a subgraph of X. Then $\omega(X') \leq \sqrt{q} - 1$.

Proof. By Theorem 1.6, $\omega(X') \leq \omega(X) \leq \sqrt{q}$. So, if $\omega(X') = \sqrt{q}$, then $\omega(X) = \sqrt{q}$ as well. Now, take a maximum clique A in X' containing 0, so that A is also a maximum clique in X. By Theorem 1.6, $|D_{m_j}| \geq |A \cap C_{m_j}| = \frac{\sqrt{q}-1}{d}$ for each $1 \leq j \leq d$, contradicting the hypothesis. \Box

We continue with another application of Theorem 1.6 on the number of maximum cliques. Recall that in the Paley graph P_q , there is a unique maximum clique that contains $\{0, 1\}$, namely the subfield $\mathbb{F}_{\sqrt{q}}$. In contrast, Mullin [Mul09, Lemma 3.3.6] shows that if r is even and $q = p^{2r}$ for some prime $p \equiv 3 \pmod{4}$ such that $\omega(P_q^*) = \sqrt{q}$, then there are at least 2 maximum cliques in P_q^* that contain $\{0, 1\}$. Next we present the proof of Theorem 1.9, which generalizes Mullin's result.

Proof of Theorem 1.9. Suppose $\omega(X) = \sqrt{q}$, and that there is a unique maximum clique A in X that contains $\{0, 1\}$. We will show that A = A(q, 2d, I) defined in equation (4), which contradicts Lemma 2.7 and thus establishes the theorem.

Let $A_j = A \cap C_j$ for each $0 \le j \le 2d - 1$. By Theorem 1.6, $|A_{m_j}| = \frac{\sqrt{q}-1}{d}$ for each $1 \le j \le d$. For each $y \in A_0$, the map $x \mapsto y^{-1}x$ induces a graph automorphism of X, so $y^{-1}A$ is a maximum clique; moreover, $1 = y^{-1}y \in y^{-1}A$, and so $y^{-1}A = A$ by the uniqueness of the maximum clique containing $\{0, 1\}$. In particular, for each $x, y \in A_0$, we have $xy^{-1} \in y^{-1}A_0 = A_0$. Hence A_0 is a subgroup of \mathbb{F}_q^* . Since $|A_0| = \frac{\sqrt{q}-1}{d}$, we conclude that $A_0 = \langle g^{d(\sqrt{q}+1)} \rangle$. For each $1 \le j \le d$, we can apply the same argument to $g^{-m_j}A_{m_j}$ to conclude that $A_{m_j} = g^{m_j} \langle g^{d(\sqrt{q}+1)} \rangle$. Therefore,

$$A = \{0\} \cup \bigcup_{j=1}^{d} A_{m_j} = \{0\} \cup \bigcup_{j=1}^{d} g^{m_j} \langle g^{d(\sqrt{q}+1)} \rangle = A(q, 2d, I)$$

as desired.

Corollary 3.4. Let X = PP(q, 2d, I) be a semi-primitive pseudo-Paley graph with $q = p^{2rt}$ where r is even. Then X is isomorphic to the Paley graph P_q if and only if $I = \{0, 2, ..., 2d - 2\}$ or $I = \{1, 3, ..., 2d - 1\}$.

Proof. If $I = \{0, 2, \dots, 2d - 2\}$ or $I = \{1, 3, \dots, 2d - 1\}$, then $X \cong P_q$ by Example 2.2.

Conversely, suppose that I is different $\{0, 2, ..., 2d-2\}$ and $\{1, 3, ..., 2d-1\}$. We may assume that $0 \in I$. Assume, to the contrary, that X is isomorphic to the Paley graph P_q . Then $\omega(X) = \sqrt{q}$. Let $f: X \to P_q$ be a graph isomorphism. Since $0 \in I$ and $1 \in C_0$, it follows that 0 and 1 are adjacent in X, and f(0) and f(1) are adjacent in P_q . After composing f with the automorphism $x \mapsto (x - f(0))/(f(1) - f(0))$ of the Paley graph P_q , we obtain an isomorphism $f': X \to P_q$ fixing 0 and 1. This implies that the number of maximum cliques containing $\{0, 1\}$ must be the same in both graphs; however, there is a unique such maximum clique in P_q by Theorem 1.2, while there are at least two such cliques in X by Theorem 1.9. This contradiction shows that X is not isomorphic to P_q .

The corollary above shows that almost all the graphs of the form PP(q, 2d, I) in the semiprimitive case are not isomorphic to the Paley graph. In Section 4 below, we will prove a stronger result without the semi-primitive assumption using different tools. We included the weaker statement above because its proof is self-contained.

4. Non-isomorphism

The goal of this section is to show that most graphs of the form PP(q, 2d, I) are not isomorphic to the Paley graph P_q or the Peisert graph P_q^* . In order to achieve this result, we will borrow the deep results obtained by Peisert, Foulser, Kallaher, and others.

Peisert [Pei01] proved that P_q and P_q^* are non-isomorphic except when $q = 3^2$ by studying their respective automorphism groups. Another approach to distinguish these graphs is to study their *p*-ranks [BvE92]; this was carried out successfully in [WQWX07, Proposition 3.6]. We are going to use ideas in Peisert's paper [Pei01] to establish our key criterion in Proposition 4.7.

Throughout this section, we assume that $q = p^n$ for some positive integer n. For any field F and a positive integer m, we can consider the general semilinear group $\Gamma L_m(F)$ [dLF11, Section 3.6]. We are primarily interested in the case when $F = \mathbb{F}_q$ and m = 1. In this case, we often write $\Gamma L_1(q)$ instead of $\Gamma L_1(\mathbb{F}_q)$.

Definition 4.1 (general semilinear group). Let $\Gamma L_1(q) := \mathbb{F}_q^* \rtimes \operatorname{Gal}(\mathbb{F}_q/\mathbb{F}_p)$ be the semi-direct product obtained via the natural action of $\operatorname{Gal}(\mathbb{F}_q/\mathbb{F}_p)$ on \mathbb{F}_q^* . This group is called the general semilinear group of degree 1 over the finite field \mathbb{F}_q .

Let g be a fixed primitive root of \mathbb{F}_q , and let $\zeta : \mathbb{F}_q \to \mathbb{F}_q$ be the map $x \mapsto gx$. Since $\zeta^k : \mathbb{F}_q \to \mathbb{F}_q$ is given by $x \mapsto g^k x$, we see that $\langle \zeta \rangle \cong \mathbb{F}_q^*$. Thus, the elements of $\Gamma L_1(q)$ can be viewed as functions $f : \mathbb{F}_q \to \mathbb{F}_q$ satisfying f(x + y) = f(x) + f(y) for all $x, y \in \mathbb{F}_q$ and $f(cx) = c^{p^j} x$ for some j and for all $x, c \in \mathbb{F}_q$. Since the Galois group $\operatorname{Gal}(\mathbb{F}_q/\mathbb{F}_p) = \langle \alpha \rangle$ is generated by the Frobenius map $\alpha : x \mapsto x^p$, we can also write $\Gamma L_1(q) = \mathbb{F}_q^* \rtimes \langle \alpha \rangle = \langle \zeta \rangle \rtimes \langle \alpha \rangle$. Thus, $\Gamma L_1(q) = \langle \zeta, \alpha \rangle$ and in particular, $\Gamma L_1(q)$ is a permutation group. We remark that other authors (for example Peisert [Pei01], and Foulser and Kallaher [FK78]) use ω to denote the map $x \mapsto gx$. We use ζ for this purpose as ω is reserved for the clique number throughout the paper.

Peisert [Pei01] classified all self-complementary symmetric graphs. Recall that a graph is selfcomplementary if it is isomorphic to its complement, and a graph is symmetric if its automorphism group acts transitively on the ordered pairs of adjacent vertices.

Theorem 4.2 ([Pei01]). A graph G is self-complementary and symmetric if and only if $|G| = p^n$ for some prime p with $p^n \equiv 1 \pmod{4}$, and G is isomorphic to a Paley graph, a Peisert graph, or the exceptional graph $G(23^2)$.

The key to proving Theorem 4.2 is the classification of the automorphism groups of all possible self-complementary symmetric graphs [Pei01, Theorem 4.2]. For our purposes, we state a slight variation of this result.

Proposition 4.3. Let X be a self-complementary symmetric Cayley graph defined on \mathbb{F}_{p^n} with $\mathscr{A} = \operatorname{Aut}(X)$, and \mathscr{A}_0 be the subgroup of \mathscr{A} which fixes 0. If $p^n \notin \{9, 49, 81, 529\}$, then \mathscr{A}_0 is a subgroup of $\Gamma L_1(p^n)$ and $\mathscr{A} \cong \mathbb{F}_{p^n} \times \mathscr{A}_0$.

Proof. Each automorphism $f \in \mathscr{A}$ can be decomposed as $f_2 \circ f_1$ where $f_1 \in \mathscr{A}_0$ and f_2 is a translation by an element of \mathbb{F}_{p^n} . This explains the identity $\mathscr{A} \cong \mathbb{F}_{p^n} \times \mathscr{A}_0$.

The proof given in [Pei01, Theorem 4.2] classifies the automorphism group of any such graph X. There are four different cases in Peisert's proof. Case 1 corresponds exactly to $\mathscr{A}_0 \leq \Gamma L_1(p^n)$. Next, Case 2 corresponds to $p^n = 9$, while Cases 3 and 4 correspond to $p^n \in \{49, 81, 529\}$. By the hypothesis, we must be in Case 1.

The following lemma shows that a subgroup of $\Gamma L_1(p^n)$ has a simple structure. The sentence starting with "Moreover" is not literally part of [FK78, Lemma 2.1], but appears on page 117 of [FK78] and also on page 220 of [Pei01].

Lemma 4.4 ([FK78, Lemma 2.1]). If \mathscr{A}_0 is a subgroup of $\Gamma L_1(p^n)$, then $\mathscr{A}_0 = \langle \zeta^d, \zeta^e \alpha^s \rangle$ for some integers d, e, s satisfying the following conditions:

$$s \mid n, d \mid (p^n - 1), e\left(\frac{p^n - 1}{p^s - 1}\right) \equiv 0 \pmod{d}$$

Such a representation $\mathscr{A}_0 = \langle \zeta^d, \zeta^e \alpha^s \rangle$ is unique when d, s > 0 and $0 \le e < d$ and it is called the standard form. Moreover, if $\mathscr{A}_0 = \langle \zeta^d, \zeta^e \alpha^s \rangle$ is in standard form, then $|\mathscr{A}_0| = n (p^n - 1) / sd$.

The automorphism group of Paley graphs and Peisert graphs were first determined by Carlitz [Car60] and Peisert [Pei01], respectively.

Theorem 4.5 ([Car60]). Let $q = p^n$. Then $\operatorname{Aut}(P_q) \cong \mathbb{F}_{p^n} \times \langle \zeta^2, \alpha \rangle$ and $|\operatorname{Aut}(P_q)| = np^n (p^n - 1)/2$.

Theorem 4.6 ([Pei01, Corollary 6.3]). Let $q = p^n$, where $p \equiv 3 \pmod{4}$ and n is even. If $q \neq 3^2, 7^2, 3^4, 23^2$, then $\operatorname{Aut}(P_q^*) \cong \mathbb{F}_{p^n} \times \langle \zeta^4, \zeta \alpha \rangle$ and $|\operatorname{Aut}(P_q^*)| = np^n (p^n - 1)/4$.

For a generalized Paley graph, it is much more difficult to determine its automorphism group. In [LP09], Lim and Praeger used association scheme to study the automorphism groups of certain generalized Paley graphs. We expect in general it is even harder to determine the automorphism group of X = PP(q, 2d, I) since it is a union of copies of generalized Paley graphs. While determining the automorphism group of X remains open, we show that X is not isomorphic to P_q or P_q^* in general. The following proposition is a stronger version of Corollary 3.4.

Proposition 4.7. Let X = PP(q, 2d, I) be in its minimal representation such that $q = p^n \notin \{3^2, 7^2, 3^4, 23^2\}$.

- (1) If d = 1, then X is isomorphic to the Paley graph P_q .
- (2) If d = 2, then X is isomorphic to the Peisert graph P_a^*
- (3) If $d \ge 3$, then X is not isomorphic to P_q or P_q^* .

Proof. It is straightforward to prove the cases d = 1 and d = 2. See Example 2.2.

Next we assume $d \ge 3$. Assume, to the contrary, that X is isomorphic to P_q or to P_q^* . In particular, X is symmetric and self-complementary by Theorem 4.2. Let $\mathscr{A} = \operatorname{Aut}(X)$ and $\mathscr{A}_0 \le \mathscr{A}$ be the subgroup fixing 0. By Proposition 4.3, \mathscr{A}_0 is a subgroup of $\Gamma L_1(\mathbb{F}_q)$.

By Lemma 2.4 and Remark 2.5, $\zeta^k : x \mapsto g^k x$ induces a graph automorphism of X if and only $2d \mid k$. Therefore, each $\zeta^k \in \mathscr{A}_0$ with k > 0 must satisfy $k \ge 2d \ge 6$. Using Lemma 4.4, we have $\mathscr{A}_0 = \langle \zeta^{d'}, \zeta^e \alpha^s \rangle$ in its standard form for some $d' \ge 6$, $e \ge 0$ and $s \ge 1$. Consequently,

$$|\mathscr{A}| = |\mathbb{F}_{p^n}||\mathscr{A}_0| = \frac{np^n(p^n - 1)}{sd'} \le \frac{np^n(p^n - 1)}{6}.$$

The automorphism group of X has a smaller size than the automorphism group of both the Paley graph P_q and the Peisert graph P_q^* (if n is even) shown in Theorem 4.5 and Theorem 4.6, which is a contradiction.

Mullin [Mul09, Chapter 8] conjectured that generalized Peisert graphs are distinct from Peisert and Paley graphs for infinitely many prime powers. She verified the conjecture computationally for small prime powers [Mul09, Section 5.4]. We confirm this conjecture below.

Corollary 4.8. If $d \ge 3$ and $q \equiv 1 \pmod{4d}$ and $q \notin \{3^2, 7^2, 3^4, 23^2\}$, then $GP^*(q, 2d)$ is not isomorphic to P_q or P_q^* . In fact, $GP^*(q, 2d)$ is self-complementary but not symmetric.

Proof. By Example 2.2, we identify $GP^*(q, 2d)$ with $PP(q, 2d, \{0, 1, ..., d - 1\})$ which is in its minimal representation, and the complement of $GP^*(q, 2d)$ corresponds to $PP(q, 2d, \{d, d + 1, ..., 2d - 1\})$. The first assertion follows directly from Proposition 4.7. Consider the graph homomorphism:

$$f: PP(q, 2d, \{0, 1, \dots, d-1\}) \to PP(q, 2d, \{d, d+1, \dots, 2d-1\})$$

induced by $x \mapsto g^d x$ on the set of vertices. Then f is an isomorphism from $GP^*(q, 2d)$ to its complement. Thus, $GP^*(q, 2d)$ is self-complementary. Since $GP^*(q, 2d)$ is not isomorphic to the Paley graph and the Peisert graph, $GP^*(q, 2d)$ cannot be symmetric by Theorem 4.2.

Mathon [Mat88] asked whether there is an infinite family of self-complementary strongly regular graphs of non-Paley type. This was already settled affirmatively by Peisert in Theorem 4.2 as Peisert graphs are of non-Paley type. Klin, Kriger, and Woldar [KKW16] also answered Mathon's question for graphs of order p^2 . Corollary 4.8 gives another answer to this question. Indeed, when -1 is a power of p modulo 2d, the generalized Peisert graph $GP^*(q, 2d)$ is a self-complementary strongly regular graph, but not isomorphic to P_q or P_q^* .

5. THE SUBSPACE STRUCTURE OF MAXIMUM CLIQUES

In previous sections, we saw that a semi-primitive pseudo-Paley graph PP(q, 2d, I) is not isomorphic to P_q unless there is an obvious reason. In this section, we continue to study these graphs and further investigate how they compare to Paley graphs from different perspectives.

5.1. Maximum cliques come in pairs. In this subsection, we investigate the correspondence between maximum cliques in a semi-primitive pseudo-Paley graph on $q = p^{4t}$ vertices. The goal is to prove Theorem 1.10 on the parity of the number of canonical cliques: we will see that they come in pairs, and the two cliques in each pair are interchanged by the Galois automorphism $x \mapsto x^{\sqrt{q}}$.

Lemma 5.1. If $p^t \equiv -1 \pmod{2d}$, and $q = p^{2rt}$, then $\mathbb{F}_{p^t}^* \subset C_0 = \langle g^{2d} \rangle$.

Proof. Note that g^k is a primitive root of \mathbb{F}_{p^t} , where

$$k = \frac{p^{2rt} - 1}{p^t - 1} = \sum_{j=0}^{2r-1} p^{jt} \equiv \sum_{j=0}^{2r-1} (-1)^j \equiv 0 \pmod{2d}$$

since $p^t \equiv -1 \pmod{2d}$. Thus, $g^k \in \langle g^{2d} \rangle$, and it follows that $\mathbb{F}_{p^t}^* \subset C_0$.

The following lemma is a simple consequence of Theorem 1.6 and Lemma 2.6.

Lemma 5.2. Let X = PP(q, 2d, I) be a semi-primitive pseudo-Paley graph with $q = p^{2rt}$ where r is even, and $I \neq \{0, 2, ..., 2d - 2\}$. Then the subfield $\mathbb{F}_{\sqrt{q}}$ is not a clique in X.

Proof. If $\mathbb{F}_{\sqrt{q}}$ were a clique in X, then it would be a maximum clique. Comparing equations (2) and (3), we get $I = \{0, 2, \dots, 2d - 2\}$, a contradiction.

Lemma 5.3. Let g be a primitive root of \mathbb{F}_q where $q = p^{4t}$ and $p^t \equiv -1 \pmod{2d}$. Set $C_0 = \langle g^{2d} \rangle$, and pick $x \in C_0$. If $\{x, x^{p^{2t}}\}$ is linearly dependent over \mathbb{F}_{p^t} , then $x \in \langle g^{(p^{2t}+1)d} \rangle$.

Proof. If $\{x, x^{p^{2t}}\}$ is linearly dependent over \mathbb{F}_{p^t} , then $x^{p^{2t}-1} \in \mathbb{F}_{p^t}^*$. After expressing $x = g^k$ for some integer k, we obtain $g^{(p^{2t}-1)k} \in \mathbb{F}_{p^t}^*$. As $\mathbb{F}_{p^t}^* = \langle g^{(q-1)/(p^t-1)} \rangle$, and $q = p^{4t}$ we must have

$$\frac{q-1}{p^t-1} \mid (p^{2t}-1)k \implies (p^{2t}+1)(p^t+1) \mid (p^{2t}-1)k \implies p^{2t}+1 \mid (p^t-1)k$$

Using $gcd(p^{2t} + 1, p^t - 1) = 2$, we obtain $\frac{p^{2t}+1}{2} \mid k$. Next, $x = g^k \in C_0 = \langle g^{2d} \rangle$ implies that $2d \mid k$. We will now combine these two divisibility relations $\frac{p^{2t}+1}{2} \mid k$ and $2d \mid k$.

Observe that $p^t \equiv -1 \pmod{2d}$ implies $p^{2t} \equiv 1 \pmod{4d}$, so that $\frac{p^{2t}+1}{2} \equiv 1 \pmod{2d}$. Therefore, $gcd(\frac{p^{2t}+1}{2}, 2d) = 1$. Combining $\frac{p^{2t}+1}{2} \mid k$ and $2d \mid k$, we conclude $(p^{2t}+1)d \mid k$ and hence $x = g^k \in \langle g^{(p^{2t}+1)d} \rangle$.

We are now ready to present the proof of Theorem 1.10.

Proof of Theorem 1.10. Without loss of generality, we may assume $m_1 = 0$. By assumption, X is a Cayley graph on $q = p^{4t}$ many vertices where $p^t \equiv -1 \pmod{2d}$. Note that the map $f \colon \mathbb{F}_q \to \mathbb{F}_q$ given by $f(x) = x^{\sqrt{q}}$ preserves the edges of the graph: if $x, y \in \mathbb{F}_q$ and $x - y \in D$, then

$$f(x) - f(y) = x^{\sqrt{q}} - y^{\sqrt{q}} = (x - y)^{\sqrt{q}} \in D$$

because $\sqrt{q} = p^{2t} \equiv 1 \pmod{2d}$. Note that f fixes each element of $\mathbb{F}_{\sqrt{q}}$ pointwise; in particular, f fixes \mathbb{F}_{p^t} . It is also evident that f is a bijection on the vertex set \mathbb{F}_q . We conclude that f is a

graph automorphism of X which fixes \mathbb{F}_{p^t} , and hence should send maximum cliques containing \mathbb{F}_{p^t} to maximum cliques containing \mathbb{F}_{p^t} . Let A be a clique of size \sqrt{q} , such that A is a \mathbb{F}_{p^t} -subspace containing \mathbb{F}_{p^t} . We claim that $f(A) \neq A$. The theorem follows immediately from this claim since f is an involution.

We assume, to the contrary, that f(A) = A. Since I is different from $\{0, 2, \ldots, 2d - 2\}$, the subfield $\mathbb{F}_{\sqrt{q}}$ is not a clique in X by Lemma 5.2. Since $A \subset \mathbb{F}_q = \mathbb{F}_{p^{4t}}$ is a 2-dimensional subspace over \mathbb{F}_{p^t} containing \mathbb{F}_{p^t} and $A \neq \mathbb{F}_{\sqrt{q}} = \mathbb{F}_{p^{2t}}$, we must have $A \cap \mathbb{F}_{\sqrt{q}} = \mathbb{F}_{p^t}$. We will estimate the number of $x \in C_0 \cap A$ such that x and $f(x) = x^{p^{2t}}$ are linearly independent over \mathbb{F}_{p^t} , and use Theorem 1.6 to show that there are too many such elements when f(A) = A.

Let V be the set of elements x in $C_0 \cap A$ such that $\{x, x^{p^{2t}}\}$ are dependent over \mathbb{F}_{p^t} . By Lemma 5.3, we have $V \subset \langle g^{(p^{2t}+1)d} \rangle$. As $g^{(p^{2t}+1)d} \in \langle g^{\sqrt{q}+1} \rangle = \mathbb{F}_{\sqrt{q}}$, we have $V \subset \mathbb{F}_{\sqrt{q}}$. In particular, $V \subset A \cap \mathbb{F}_{\sqrt{q}} = \mathbb{F}_{p^t}$. On the other hand, it is clear that $\mathbb{F}_{p^t}^* \subset V$ and $0 \notin V$. Therefore, $V = \mathbb{F}_{p^t}^*$. Theorem 1.6 states that $|C_0 \cap A| = \frac{p^{2t}-1}{d}$, which implies that there are $\frac{p^{2t}-1}{d} - (p^t - 1)$ elements x in $C_0 \cap A$ such that $\{x, x^{p^{2t}}\}$ are independent over \mathbb{F}_{p^t} , that is, the complement $V' := C_0 \cap (A \setminus \mathbb{F}_{p^t}^*)$ has $\frac{p^{2t}-1}{d} - (p^t - 1)$ elements.

Since f fixes A, and A is a \mathbb{F}_{p^t} -subspace, we can apply Lemma 5.1 to obtain the following inclusion of sets:

$$\bigcup_{x \in V'} \left(x \mathbb{F}_{p^t}^* \cup x^{p^{2t}} \mathbb{F}_{p^t}^* \right) \subset C_0 \cap (A \setminus \mathbb{F}_{p^t}^*) = V'.$$

Note that $|x\mathbb{F}_{p^t}^* \cup x^{p^{2t}}\mathbb{F}_{p^t}^*| = 2(p^t - 1)$ for each $x \in V'$. If $x, y \in V'$, such that $x\mathbb{F}_{p^t}^* \cup x^{p^{2t}}\mathbb{F}_{p^t}^* = y\mathbb{F}_{p^t}^* \cup y^{p^{2t}}\mathbb{F}_{p^t}^*$, then $y \in x\mathbb{F}_{p^t}^* \cup x^{p^{2t}}\mathbb{F}_{p^t}^*$. Therefore,

$$|V'| \ge \left| \bigcup_{x \in V'} \left(x \mathbb{F}_{p^t}^* \cup x^{p^{2t}} \mathbb{F}_{p^t}^* \right) \right| \ge 2(p^t - 1) \left\lceil \frac{|V'|}{2(p^t - 1)} \right\rceil$$

This implies that $\frac{|V'|}{2(p^t-1)}$ is an integer. However, $p^t \equiv -1 \pmod{2d}$ implies that $\frac{|V'|}{2(p^t-1)} = \frac{p^t+1}{2d} - \frac{1}{2} \notin \mathbb{Z}$, a contradiction. We conclude that $f(A) \neq A$, which completes the proof.

Example 5.4. If X is the Peisert graph with order $q = p^4$ where p > 3, then there are 0 maximum cliques of size \sqrt{q} with the subspace structure because \mathbb{F}_p forms a maximal clique in P_q^* [AY22, Theorem 1.5].

Example 5.5 (Graphs of order 625 and 2401). If $X = PP(5^4, 6, I)$ with $I \neq \{0, 2, 4\}, \{1, 3, 5\}$ then X contains either 0 or 2 maximum cliques of the kind described in Theorem 1.10. A similar conclusion holds for $X = PP(7^4, 8, I)$.

With the help of SageMath, we noticed that in many examples of semi-primitive pseudo-Paley graphs there are exactly 2 cliques of size \sqrt{q} containing \mathbb{F}_{p^t} . These examples are presented in Appendix C. This leads us to the following conjecture which simultaneously strengthens and generalizes Theorem 1.10.

Conjecture 5.6. Let X = PP(q, 2d, I) be a semi-primitive pseudo-Paley graph with $q = p^{2rt}$ where r is even. Assume that $0 \in I$ and $I \neq \{0, 2, ..., 2d - 2\}$. If $\omega(X) = \sqrt{q}$, then the number of maximum cliques containing \mathbb{F}_{p^t} is 2.

5.2. **The density result.** In this subsection, we discuss the density of the graphs achieving the trivial upper bound on the clique number.

Fix a positive integer N. Define $\mathcal{PP}(N)$ to be collection of all the semi-primitive pseudo-Paley graphs of the form X = PP(q, 2d, I) with $q = p^{2rt} \leq N$ and r even. Let $\mathcal{F}(N)$ be the subset of $\mathcal{PP}(N)$ consisting of those graphs X = PP(q, 2d, I) with $\omega(X) = \sqrt{q}$. Since $\omega(X) \leq \sqrt{q}$ always holds, the graphs in $\mathcal{F}(N)$ are precisely those that attain the trivial upper bound on the clique number.

Proposition 5.7. Assume that Conjecture 1.4 is true. The density of graphs attaining the trivial upper bound on the clique number inside the family of semi-primitive pseudo-Paley graphs is zero, that is,

$$\lim_{N \to \infty} \frac{\#\mathcal{F}(N)}{\#\mathcal{P}\mathcal{P}(N)} = 0.$$

Proof. Given a prime power $q = p^{2rt} \leq N$, with r even, let $\mathcal{PP}(N, p, t) \subset \mathcal{PP}(N)$ be the subset consisting of graphs PP(q, 2d, I) with $q = p^{2rt}$ vertices such that t is the smallest positive integer such that $p^t \equiv -1 \pmod{2d}$. Similarly, define $\mathcal{F}(N, p, t) = \mathcal{F}(N) \cap \mathcal{PP}(N, p, t)$. We have the following decomposition depending on p and t:

$$\mathcal{PP}(N) = \bigsqcup_{p^t \le N^{1/4}} \mathcal{PP}(N, p, t), \quad \text{and} \quad \mathcal{F}(N) = \bigsqcup_{p^t \le N^{1/4}} \mathcal{F}(N, p, t).$$

This is a disjoint union because for any $PP(q, 2d, I) \in \mathcal{PP}(N)$, there is a unique (p, t) such that $PP(q, 2d, I) \in \mathcal{PP}(N, p, t)$. Indeed, q determines the value of p, and the congruence $p^t \equiv -1 \pmod{2d}$ uniquely determines the value of smallest such t.

Recall that the Grassmannian $\operatorname{Gr}(r, 2r)(\mathbb{F}_{p^t})$ is the set of all r-dimensional \mathbb{F}_{p^t} -subspaces inside $\mathbb{F}_{(p^t)^{2r}} = \mathbb{F}_q$. Given a graph $X = PP(q, 2d, I) \in \mathcal{F}(N, p, t)$ with $q = p^{2rt}$, Conjecture 1.4 implies that the set of maximum cliques in X containing 0 gives a subset $S_X \subset \operatorname{Gr}(r, 2r)(\mathbb{F}_{p^t})$ with $S_X \neq \emptyset$. If $Y \in \mathcal{F}(N, p, t)$ with Y = PP(q, 2d, J), and $I \neq J$, then we claim that $S_X \cap S_Y = \emptyset$. Indeed, by Theorem 1.6, a maximum clique $V \in S_X$ uniquely determines I because $I = \{j : 0 \leq j \leq 2d - 1 \text{ and } |C_j \cap V| = \frac{\sqrt{q}-1}{d}\}$, and hence the graph X. If $V \in S_X \cap S_Y$, then X = Y. The claim is proved, and we obtain:

$$\#\mathcal{F}(N,p,t) \le \# \bigsqcup_{p^{2rt} \le N} \operatorname{Gr}(r,2r)(\mathbb{F}_{p^t}).$$

The cardinality of the Grassmannian can be bounded as follows:

$$\#\operatorname{Gr}(r,2r)(\mathbb{F}_{p^t}) = \frac{(p^{2rt}-1)(p^{2rt}-p^t)\cdots(p^{2rt}-p^{(r-1)t})}{(p^{rt}-1)(p^{rt}-p^t)\cdots(p^{rt}-p^{(r-1)t})} \le p^{2r^2t}.$$

Thus,

$$\#\mathcal{F}(N, p, t) \le \sum_{p^{2rt} \le N} (p^t)^{2r^2} \le \ln(N) \cdot N^r \le \ln(N) \cdot N^{\ln(N)} = \ln(N) \cdot e^{(\ln(N))^2},$$

which implies,

$$\#\mathcal{F}(N) = \sum_{p^{4t} \le N} \#\mathcal{F}(N, p, t) \le N \cdot \ln(N) \cdot e^{(\ln(N))^2}.$$
(8)

On the other hand, the number of graphs of the form $PP(q, 2d, I) \in \mathcal{PP}(N, p, t)$ with $q = p^{2rt}$ and $d = (p^t + 1)/2$ is given by the binomial coefficient $\binom{2d}{d} = \binom{p^t+1}{(p^t+1)/2}$ since $I \subset \{0, 1, \dots, 2d-1\}$ with #I = d. Using Stirling's approximation,

$$#\mathcal{PP}(N, p, t) \ge \binom{p^t + 1}{(p^t + 1)/2} \ge \frac{4^{p^t}}{\sqrt{4p^t}}.$$

By Chebyshev's Theorem, there exists a prime p_0 such that $\frac{1}{2}N^{1/4} < p_0 < N^{1/4}$ for N large. As a result,

$$#\mathcal{PP}(N) \ge #\mathcal{PP}(N, p_0, 1) \ge \frac{4^{p_0}}{\sqrt{4p_0}} \ge \frac{2^{N^{1/4}}}{\sqrt{2N^{1/4}}}.$$
(9)

Combining inequalities (8) and (9),

$$\limsup_{N \to \infty} \frac{\#\mathcal{F}(N)}{\#\mathcal{P}\mathcal{P}(N)} \le \limsup_{N \to \infty} \frac{N \cdot \ln(N) \cdot e^{(\ln(N))^2}}{\frac{2^{N^{1/4}}}{\sqrt{2N^{1/4}}}} = 0$$

as desired.

We remark that Proposition 5.7 can be phrased unconditionally. The argument shows that a union of at most d distinct 2d-th semi-primitive cyclotomic classes of \mathbb{F}_q together with $\{0\}$, almost surely does not contain a subspace of size \sqrt{q} , as $q \to \infty$.

5.3. An infinite family of graphs (of non-Paley type) achieving the square root upper bound. Despite the density of $\mathcal{F}(N)$ being zero, we still expect that $\mathcal{F}(N)$ will contain infinitely many elements that are non-isomorphic to the Paley graph. Specializing to the case $q = p^4$ and 2d = p + 1, the proof of Proposition 5.7 can be modified to show that $\omega(PP(p^4, p + 1, I)) < p^2$ for a generic choice of I. Nonetheless, we expect that there are infinitely many exceptions of non-Paley type. This expectation is made precise in the following conjecture.

Conjecture 5.8. For each prime $p \ge 3$, there are exactly $\frac{p^2+3}{4}$ subsets $I \subset \{0, 1, \dots, p\}$ with $0 \in I$ and $|I| = \frac{p+1}{2}$ such that $\omega(PP(p^4, p+1, I)) = p^2$. Furthermore, in these graphs, the maximum cliques containing $\{0, 1\}$ are of the form $\mathbb{F}_p \oplus a\mathbb{F}_p$, where $a = g^{(p+1)k}$ and k is an odd integer.

Conjecture 5.8 is impractical to verify using a computer. Instead, we consider the following weaker conjecture, which can be verified in polynomial time via Algorithm 2 in Appendix C.

Conjecture 5.9. Let V be a 2-dimensional subspace in \mathbb{F}_{p^4} , such that $1 \in V$. Then V is a clique in $PP(p^4, p+1, I)$ for some I if and only if $V = \mathbb{F}_p \oplus a\mathbb{F}_p$, where $a = g^{(p+1)k}$ and k is an odd integer.

The next proposition states that Conjecture 5.9 is essentially equivalent to Conjecture 5.8 if we assume the main Conjecture 1.4.

Proposition 5.10. Conjecture 1.4, Conjecture 5.6, and Conjecture 5.9 together imply Conjecture 5.8.

Proof. Suppose $I \subset \{0, 1, ..., p\}$ with $0 \in I$ and $|I| = \frac{p+1}{2}$ such that $\omega(PP(p^4, p+1, I)) = p^2$. Let V be a maximum clique of size p^2 with $0, 1 \in V$; see the beginning of the proof of Theorem 1.7 for the existence of such a clique. Conjecture 1.4 implies that V is a 2-dimensional \mathbb{F}_p -subspace. We apply Conjecture 5.9 to express $V = \mathbb{F}_p \oplus a\mathbb{F}_p$ for some $a = g^{(p+1)k} \in \mathbb{F}_{p^4}$, where k is an odd integer. Recall that $C_0 = \langle g^{p+1} \rangle$ is a (p+1)-th cyclotomic class. Lemma 5.1 states that $\mathbb{F}_p^* \subset C_0$. Observe that $V \cap C_0 \supset \mathbb{F}_p^* \cup a\mathbb{F}_p^*$. By Theorem 1.6, $|V \cap C_0| = \frac{p^2-1}{(p+1)/2} = 2(p-1)$. Hence, $V \cap C_0 = \mathbb{F}_p^* \cup a\mathbb{F}_p^*$. Let $V' = \mathbb{F}_p \oplus b\mathbb{F}_p$ be another clique of the same type. The same argument shows that $V' \cap C_0 = \mathbb{F}_p^* \cup b\mathbb{F}_p^*$. Thus, we have V = V' if and only if $b/a \in \mathbb{F}_p^*$.

Given $V = \mathbb{F}_p^* \oplus a\mathbb{F}_p^*$ with $a = g^{(p+1)k}$ with k odd, the number of possible k's is $\frac{p^4-1}{2(p+1)} = (p^2+1)(p-1)/2$. The argument above shows that different choices of a in the same \mathbb{F}_p^* -coset give the same space V. Therefore, the number of choices for V is:

$$\frac{1}{2}\frac{(p^2+1)(p-1)}{p-1} = \frac{p^2+1}{2}$$

If $I = \{0, 2, 4, \dots, p-1\}$ then $PP(p^4, p+1, I)$ is the Paley graph and the only possible V in this case is $V = \mathbb{F}_{p^2}$. If $I \neq \{0, 2, 4, \dots, p-1\}$, then Conjecture 5.6 yields 2 such subspaces V. By Theorem 1.6, different choices of I give rise to different subspaces V. In summary, there are exactly

$$\frac{1}{2}\left(\frac{p^2+1}{2}-1\right)+1 = \frac{p^2+3}{4}$$

many I with $0 \in I$ satisfying $\omega(PP(p^4, p+1, I)) = p^2$.

5.4. Conditional improvement on the trivial upper bound. In this subsection, we prove Theorem 1.7 and Theorem 1.8, which improve the previously known upper bound \sqrt{q} on the clique number conditional on Conjecture 1.4.

The following lemma is useful for obtaining a lower bound on the character sum as we will see in the proof of Theorem 1.7 below.

Lemma 5.11. Let $d \ge 2$ be a positive integer and θ be a primitive 2d-th root of unity. Given a subset $I = \{m_1, m_2, \ldots, m_d\} \subset \{0, 1, \ldots, 2d - 1\}$, there exists an integer $1 \le k \le 2d - 1$ such that $|\sum_{j=1}^d \theta^{km_j}| > \sqrt{d/2}$.

Proof. We have

$$\sum_{k=0}^{2d-1} \left| \sum_{j=1}^{d} \theta^{km_j} \right|^2 = \sum_{k=0}^{2d-1} \sum_{1 \le j, \ell \le d} \theta^{k(m_j - m_\ell)} = \sum_{1 \le j, \ell \le d} \sum_{k=0}^{2d-1} \theta^{k(m_j - m_\ell)} = 2d^2.$$

It follows that

$$\max_{1 \le k \le 2d-1} \left| \sum_{j=1}^{d} \theta^{km_j} \right| \ge \sqrt{\frac{2d^2 - d^2}{2d - 1}} = \sqrt{\frac{d^2}{2d - 1}} > \sqrt{\frac{d}{2}}.$$

The following theorem is a consequence of the main result in a recent paper by Reis [Rei22]. It states that there will be a lot of cancellations in a character sum over a generic subspace V of \mathbb{F}_q provided that V is not the subfield $\mathbb{F}_{\sqrt{q}}$ and q is sufficiently large.

Theorem 5.12 ([AY22, Corollary 3.6]). Let n be an integer such that $n \ge 2$, and q an odd prime power. Let $V \subseteq \mathbb{F}_{q^{2n}}$ be an \mathbb{F}_q -subspace of dimension n, with $1 \in V$, and $V \neq \mathbb{F}_{q^n}$. Then for any non-trivial multiplicative character χ of $\mathbb{F}_{q^{2n}}$,

$$\left|\sum_{x\in V}\chi(x)\right| < \frac{2n}{\sqrt{q}} \cdot |V|.$$
(10)

In other words, if the absolute value of the character sum over a subset $S \subset \mathbb{F}_q$ with size \sqrt{q} is too large, then we can conclude that either S is not a subspace or S is the subfield $\mathbb{F}_{\sqrt{q}}$. We remark that the second alternative (the subfield case) was one of the main tools used in [AY22] to establish a new proof of Theorem 1.2 as well as its generalization. Next, we use the first alternative to prove Theorem 1.7.

Proof of Theorem 1.7. Assume, to the contrary, that $\omega(X) = \sqrt{q}$ and $p^t > 10.2r^2d$. Without loss of generality, we may assume $m_1 = 0$. Since X is a Cayley graph, we can restrict attention to maximum cliques containing 0. If A is a maximum clique in X containing 0, Theorem 1.6 implies that $C_0 \cap A \neq \emptyset$ and thus we can further assume that $1 \in A$. Indeed, any $y \in C_0 \cap A$ yields another maximum clique $B = y^{-1}A$ containing both 0 and 1. By Lemma 5.2, $A \neq \mathbb{F}_{\sqrt{q}}$.

By Lemma 5.11, we can find $1 \le k \le 2d - 1$, such that $|\sum_{j=1}^{d} \theta^{km_j}| > \sqrt{d/2}$. Let χ be the multiplicative character of \mathbb{F}_q such that $\chi(g) = \theta^k$. Then χ is a nontrivial character with order dividing 2d. Suppose that A is a subspace over \mathbb{F}_{p^t} ; then by Theorem 5.12 and Theorem 1.6, we have

$$\frac{\sqrt{q}-1}{d} \left| \sum_{j=1}^{d} \theta^{km_j} \right| = \left| \sum_{j=1}^{d} \sum_{x \in C_{m_j}} \chi(x) \right| = \left| \sum_{x \in A} \chi(x) \right| < \frac{2r}{\sqrt{p^t}} \sqrt{q}, \tag{11}$$

Thus,

$$\frac{\sqrt{q}-1}{\sqrt{2d}} \le \frac{\sqrt{q}-1}{d} \left| \sum_{j=1}^{d} \theta^{km_j} \right| < \frac{2r}{\sqrt{p^t}} \sqrt{q}.$$

However, the above inequality implies $p^t \leq 10.2r^2d$, a contradiction. This finishes the proof. \Box

While Theorem 1.7 is stated conditionally in terms of the clique number of certain Cayley graphs, it can also be phrased unconditionally with additive combinatorics flavor as follows.

Corollary 5.13. Let $q = p^{2rt}$, where r is even, and t is the smallest positive integer such that $p^t \equiv -1 \pmod{2d}$. Let D be the union of at most d distinct 2d-th cyclotomic classes of \mathbb{F}_q . Assume that D contains at least one square and one non-square element. If $p^t > 10.2r^2d$, then there is no \mathbb{F}_{p^t} -subspace V of \mathbb{F}_q , with dimension at least r, such that $V \subset D \cup \{0\}$.

When the set I has a simple structure, the lower bound on p in Theorem 1.7 can be further improved. We illustrate this by analyzing the case of generalized Peisert graphs, where we can compute the sum of roots of unity explicitly.

Corollary 5.14. Let $q = p^{2rt}$, where r is even and t is the smallest positive integer such that $p^t \equiv -1 \pmod{2d}$. Assuming Conjecture 1.4, the generalized Peisert graph $GP^*(q, 2d)$ has clique number less than \sqrt{q} when $p^t \ge 12.5r^2$.

Proof. We will assume that $\omega(GP^*(q, 2d)) = \sqrt{q}$ and show that $p^t < 12.5r^2$ must hold. Recall that $GP^*(q, 2d) = PP(q, 2d, I)$, where $I = \{0, 1, \dots, d-1\}$; see Example 2.2.

Let $\theta = e^{2\pi i/2d} = e^{i\pi/d}$. Let χ be the multiplicative character of \mathbb{F}_q such that $\chi(g) = \theta$. We have

$$\sum_{j=1}^{d} \theta^{m_j} = \sum_{j=1}^{d} \theta^{j-1} = \frac{1-\theta^d}{1-\theta} = \frac{2}{1-\theta}.$$

Using the inequality $\cos x \ge 1 - \frac{x^2}{2}$ for all real x, we have

$$|1 - \theta| = \sqrt{2 - 2\cos\frac{\pi}{d}} \le \sqrt{2 - 2\left(1 - \frac{\pi^2}{2d^2}\right)} = \frac{\pi}{d}$$

and so,

 $\left|\frac{2}{1-\theta}\right| \ge \frac{2d}{\pi}.\tag{12}$

Combining the inequalities (11) and (12) yields

$$\frac{\sqrt{q}-1}{d} \cdot \frac{2d}{\pi} \le \frac{\sqrt{q}-1}{d} \left| \sum_{j=1}^{d} \theta^{m_j} \right| < \frac{2r}{\sqrt{p^t}} \sqrt{q}.$$

This last inequality rearranges into:

$$\sqrt{p^t} < \frac{r\pi\sqrt{q}}{\sqrt{q}-1} \quad \Rightarrow \quad p^t < \frac{r^2\pi^2 q}{(\sqrt{q}-1)^2} \le \left(\frac{81}{64}\right)r^2\pi^2 < 12.5r^2 \tag{13}$$

since $q \ge 81$.

To prove Theorem 1.8, we apply the above corollary, and then use SageMath [Sag21] to verify the cases when q is small.

Proof of Theorem 1.8. We assume the clique number is \sqrt{q} and apply the contrapositive of Corollary 5.14 in the special case r = 2 to conclude that $p^t < 50$. Note that when $p^t \ge 41$, $q = p^{4t} \ge 41^4$, and inequality (13) implies that

$$p^t < 4\pi^2 \left(\frac{1681}{1680}\right)^2 < 40$$

a contradiction. Therefore,

 $p^t \in \{5, 7, 9, 11, 13, 17, 19, 23, 25, 27, 29, 31, 37\}.$

Recall that $p^t \equiv -1 \pmod{2d}$ and $d \geq 2$. However, using Algorithm 1 (described in Appendix C), we verified that the corresponding graph for each choice of (p^t, d) in the following list has clique number less than $\sqrt{q} = p^{2t}$ leading to a contradiction:

$$\{(5,3), (7,2), (7,4), (9,5), (11,2), (11,3), (11,6), (13,7), (17,3), (17,9), (19,5), (19,10), (23,2), (23,3), (23,4), (23,6), (23,12), (25,13), (27,7), (27,14), (29,3), (29,5), (29,15), (31,2), (31,4), (31,8), (31,16), (37,19) \}.$$

The largest case $(p^t, d) = (37, 19)$ took about 16 hours in SageMath.

ACKNOWLEDGEMENTS

The authors thank Gabriel Currier, Greg Martin, Zinovy Reichstein, József Solymosi, Zixiang Xu, and Joshua Zahl for helpful discussions. During the preparation of this manuscript, the first author was supported by a postdoctoral research fellowship from the University of British Columbia and the NSERC PDF award. The research of the second author was supported in part by a Four Year Doctoral Fellowship from the University of British Columbia.

REFERENCES

- [AB14] N. Alon and J. Bourgain, *Additive patterns in multiplicative subgroups*, Geom. Funct. Anal. **24** (2014), no. 3, 721–739.
- [AGLY22] S. Asgarli, S. Goryainov, H. Lin, and C. H. Yip, *The EKR-module property of pseudo-Paley graphs of square order*, arXiv e-prints (2022), available at https://arxiv.org/abs/2201.03100.
 - [AY22] S. Asgarli and C. H. Yip, Van Lint-MacWilliams' conjecture and maximum cliques in Cayley graphs over finite fields, J. Combin. Theory Ser. A 192 (2022), Paper No. 105667, 23pp.
- [BEW98] B. C. Berndt, R. J. Evans, and K. S. Williams, *Gauss and Jacobi sums*, Canadian Mathematical Society Series of Monographs and Advanced Texts, John Wiley & Sons, Inc., New York, 1998.
 - [Blo84] A. Blokhuis, On subsets of $GF(q^2)$ with square differences, Nederl. Akad. Wetensch. Indag. Math. **46** (1984), no. 4, 369–372.
- [BM72] L. D. Baumert and R. J. McEliece, Weights of irreducible cyclic codes, Information and Control 20 (1972), 158–175.
- [BvE92] A. E. Brouwer and C. A. van Eijl, *On the p-rank of the adjacency matrices of strongly regular graphs*, J. Algebraic Combin. **1** (1992), no. 4, 329–346.
- [BWX99] A. E. Brouwer, R. M. Wilson, and Q. Xiang, Cyclotomy and strongly regular graphs, J. Algebraic Combin. 10 (1999), no. 1, 25–28.
- [Car60] L. Carlitz, A theorem on permutations in a finite field, Proc. Amer. Math. Soc. 11 (1960), 456–459.
- [CGW89] F. R. K. Chung, R. L. Graham, and R. M. Wilson, *Quasi-random graphs*, Combinatorica **9** (1989), no. 4, 345–362.
- [Chv11] V. Chvátal, A conjecture in extremal combinatorics, 2011. https://users.encs.concordia.ca/~chvatal/cc
- [Chv74] _____, Intersecting families of edges in hypergraphs having the hereditary property, Hypergraph Seminar (Proc. First Working Sem., Ohio State Univ., Columbus, Ohio, 1972; dedicated to Arnold Ross), 1974, pp. 61–66. Lecture Notes in Math., Vol. 411.
- [CL07] E. S. Croot III and V. F. Lev, *Open problems in additive combinatorics*, Additive combinatorics, 2007, pp. 207–233.
- [Del73] P. Delsarte, An algebraic approach to the association schemes of coding theory, Philips Res. Rep. Suppl. 10 (1973), vi+97.
- [dLF11] W. de Launey and D. Flannery, *Algebraic design theory*, Mathematical Surveys and Monographs, vol. 175, American Mathematical Society, Providence, RI, 2011.
- [FK78] D. A. Foulser and Michael J. Kallaher, *Solvable, flag-transitive, rank 3 collineation groups*, Geometriae Dedicata **7** (1978), no. 1, 111–130.
- [GM16] C. Godsil and K. Meagher, Erdős-Ko-Rado theorems: algebraic approaches, Cambridge Studies in Advanced Mathematics, vol. 149, Cambridge University Press, Cambridge, 2016.
- [GR17] C. Godsil and B. Rooney, *Hardness of computing clique number and chromatic number for Cayley graphs*, European J. Combin. **62** (2017), 147–166.
- [GS18] G. R. W. Greaves and L. H. Soicher, On the clique number of a strongly regular graph, Electron. J. Combin. 25 (2018), no. 4, Paper No. 4.15, 15.
- [GS71] R. L. Graham and J. H. Spencer, *A constructive solution to a tournament problem*, Canad. Math. Bull. **14** (1971), 45–48.
- [HP21] B. Hanson and G. Petridis, *Refined estimates concerning sumsets contained in the roots of unity*, Proc. Lond. Math. Soc. (3) **122** (2021), no. 3, 353–358.
- [Jon20] G. A. Jones, *Paley and the Paley graphs*, Isomorphisms, symmetry and computations in algebraic graph theory, 2020, pp. 155–183.
- [KKW16] M. Klin, N. Kriger, and A. Woldar, On the existence of self-complementary and non-self-complementary strongly regular graphs with Paley parameters, J. Geom. 107 (2016), no. 2, 329–356.
- [Kop19] S. Kopparty, Fourier analysis, 2019. https://sites.math.rutgers.edu/~sk1233/courses/finitefiel
- [KP04] A. Kisielewicz and W. Peisert, *Pseudo-random properties of self-complementary symmetric graphs*, J. Graph Theory **47** (2004), no. 4, 310–316.
- [LN97] R. Lidl and H. Niederreiter, *Finite fields*, Second, Encyclopedia of Mathematics and its Applications, vol. 20, Cambridge University Press, Cambridge, 1997.

- [LP09] T. K. Lim and C. E. Praeger, On generalized Paley graphs and their automorphism groups, Michigan Math. J. 58 (2009), no. 1, 293–308.
- [Mat88] R. Mathon, On self-complementary strongly regular graphs, Discrete Math. 69 (1988), no. 3, 263–281.
- [Mul09] N. Mullin, *Self-complementary arc-transitive graphs and their imposters* (2009). Master's thesis, University of Waterloo.
- [Pei01] W. Peisert, All self-complementary symmetric graphs, J. Algebra 240 (2001), no. 1, 209–229.
- [Rei22] L. Reis, Character sums over affine spaces and applications, Finite Fields Appl. 83 (2022), Paper No. 102067.
- [Sag21] Sage Developers, SageMath, the Sage Mathematics Software System (Version 9.4), 2021.
- [Shk14] I. D. Shkredov, Sumsets in quadratic residues, Acta Arith. 164 (2014), no. 3, 221–243.
- [Sti90] L. Stickelberger, Ueber eine Verallgemeinerung der Kreistheilung, Math. Ann. 37 (1890), no. 3, 321– 367.
- [WQWX07] G. Weng, W. Qiu, Z. Wang, and Q. Xiang, *Pseudo-Paley graphs and skew Hadamard difference sets from presemifields*, Des. Codes Cryptogr. **44** (2007), no. 1-3, 49–62.
 - [Yip21] C. H. Yip, On the directions determined by Cartesian products and the clique number of generalized *Paley graphs*, Integers **21** (2021), Paper No. A51, 31pp.
 - [Yip22a] _____, Gauss sums and the maximum cliques in generalized Paley graphs of square order, Funct. Approx. Comment. Math. 66 (2022), no. 1, 119–128.
 - [Yip22b] _____, On maximal cliques of Cayley graphs over fields, J. Algebraic Combin. 56 (2022), no. 2, 323–333.
 - [Yip22c] _____, On the clique number of Paley graphs of prime power order, Finite Fields Appl. 77 (2022), Paper No. 101930.

APPENDIX A. GAUSS SUMS AND CHARACTER SUMS

In this section, we work with Gauss sums and character sums over finite fields. We refer to [BEW98] and [LN97, Chapter 5] for a detailed discussion. This appendix contains preparatory material for Appendix B on the proof of Proposition 3.2.

Let χ be a multiplicative character of \mathbb{F}_q , and ψ an additive character of \mathbb{F}_q . As usual, we extend χ to be defined on \mathbb{F}_q by setting $\chi(0) = 0$. The *Gauss sum* associated to χ and ψ is defined to be $G(\chi, \psi) = \sum_{c \in \mathbb{F}_q} \chi(c)\psi(c)$. When ψ is the canonical additive character $e_p(\operatorname{Tr}_{\mathbb{F}_q}(\cdot))$, we use $G(\chi)$ to denote $G(\chi, \psi)$. The first non-trivial example is when χ is a quadratic character; in this case, the Gauss sum is well-understood according to the following classical theorem.

Theorem A.1 ([BEW98, Theorem 11.5.4]). Let \mathbb{F}_q be a finite field with $q = p^s$, where p is an odd prime and s is a positive integer. Let χ be the quadratic character of \mathbb{F}_q . Then

$$G(\chi) = \begin{cases} (-1)^{s-1}\sqrt{q}, & \text{if } p \equiv 1 \pmod{4}, \\ (-1)^{s-1}i^s\sqrt{q}, & \text{if } p \equiv 3 \pmod{4}. \end{cases}$$

Corollary A.2. Suppose p is a prime such that $p^t \equiv -1 \pmod{2d}$ with t minimal. Let $q = p^{2rt}$, where r is even. If χ be the quadratic character of \mathbb{F}_q , then $G(\chi) = -\sqrt{q}$.

Proof. We can write $q = p^s$ where s = 2rt. Note that $i^{2rt} = 1$ since r is even. The corollary follows from Theorem A.1.

Stickelberger's theorem (see for example [BEW98, Theorem 11.6.3]) provides an explicit formula to compute semi-primitive Gauss sums. The original proof by Stickelberger [Sti90] was based on algebraic geometry.

Theorem A.3 (Stickelberger's theorem). Let p be an odd prime and let d > 2 be an integer. Suppose that there exists a positive integer t such that $p^t \equiv -1 \pmod{d}$, with t chosen to be minimal. Let χ be a multiplicative character of \mathbb{F}_{p^v} with order d. Then v = 2ts for some positive integer s, and $p^{-v/2}G(\chi) = (-1)^{s-1+(p^t+1)s/d}$.

According to [BWX99, Proposition 1], the following corollary is a consequence of Stickelberger's theorem and Davenport-Hasse theorem: an equivalent statement in terms of the generating function of Gauss periods first appeared (implicitly) in [BM72, Section 5]. For the sake of completeness, we present a self-contained proof below.

Corollary A.4. Let $q = p^{2rt}$, where $p^t \equiv -1 \pmod{2d}$ is a prime with t minimal. Let χ be a multiplicative character of \mathbb{F}_q with order 2d. Then for each $1 \leq k \leq 2d - 1$, $G(\chi^k) = -\sqrt{q}$.

Proof. When k = d, then χ^d is the quadratic character and $G(\chi^d) = -\sqrt{q}$ by Corollary A.2.

Next we assume that $d \nmid k$. Then χ^k is a character with order $e = 2d/\gcd(2d, k) > 2$, where $e \mid 2d$. Since -1 is a power of $p \pmod{2d}$, -1 is also a power of $p \pmod{e}$. Let t' be the smallest positive integer such that $p^{t'} \equiv -1 \pmod{e}$; then it is easy to check that the order of $p \pmod{e}$ is 2t'. Note that $p^t \equiv -1 \pmod{2d}$; in particular, $p^t \equiv -1 \pmod{e}$ and $p^{t-t'} \equiv 1 \pmod{e}$. Thus, $2t' \mid (t - t')$ so that t/t' is an odd integer. Let r' be defined by the equation rt = r't' so that $q = p^{2rt} = p^{2r't'}$. It is clear that r and r' have the same parity.

We claim that $p^t \equiv p^{t'} \pmod{2e}$. We consider two cases.

- (i) Suppose that $p^{t'} \equiv -1 \pmod{2e}$. Since t/t' is odd, we also have $p^t \equiv -1 \pmod{2e}$.
- (ii) Suppose that $p^{t'} \equiv e 1 \pmod{2e}$.
 - (a) When e is even, observe that $(e-1)^2 \equiv 1 \pmod{2e}$, and so $(e-1)^{t/t'} \equiv e-1 \pmod{2e}$.
 - (b) When e is odd, $(e-1)^2 \equiv e^2 + 1 \equiv e+1 \pmod{2e}$; now, $(e+1)^2 \equiv e^2 + 1 \equiv e+1 \pmod{2e}$ and so $(e-1)^{t/t'} \equiv (e+1)(e-1) \equiv e^2 1 \equiv e-1 \pmod{2e}$.

In both subcases, $(e-1)^{t/t'} \equiv e-1 \pmod{2e}$. As a result, $p^t \equiv e-1 \pmod{2e}$.

This proves the claim. Consequently,

- (i) If $p^t \equiv -1 \pmod{4d}$, then $p^{t'} \equiv p^t \equiv -1 \pmod{2e}$.
- (ii) If $p^t \equiv 2d 1 \pmod{4d}$, then we consider two subcases.
 - (a) if $e \mid d$ (i.e., k is even), then $p^{t'} \equiv p^t \equiv -1 \pmod{2e}$.
 - (b) if $e \nmid d$ (i.e., k is odd), then $p^{t'} \equiv p^t \equiv e 1 \pmod{2e}$.

Finally, we apply Theorem A.3 to obtain:

$$G(\chi^k)/\sqrt{q} = (-1)^{r'-1+(p^{t'}+1)r'/e} = (-1)^{r'-1} = (-1)^{r-1} = -1.$$

as desired.

The following theorem, which can be proved directly using orthogonality relations, expresses a complete character sum with a monomial argument as a linear combination of Gauss sums.

Theorem A.5 ([LN97, Theorem 5.30]). Let ψ be a nontrivial additive character of \mathbb{F}_q , $n \in \mathbb{N}$, and χ a multiplicative character of \mathbb{F}_q of order $e = \gcd(n, q - 1)$. Then for any $a \in \mathbb{F}_q^*$,

$$\sum_{c \in \mathbb{F}_q} \psi(ac^n) = \sum_{j=1}^{e-1} \bar{\chi}^j(a) G(\chi^j, \psi).$$

Theorem A.5 allows us to explicitly compute the following *Gauss periods*, that is, the character values of cyclotomic classes.

Corollary A.6. Let $q = p^{2rt}$, where $p^t \equiv -1 \pmod{2d}$ is a prime with t minimal and r is even. Let $C_0, C_1, \ldots, C_{2d-1}$ be the list of 2*d*-th cyclotomic classes of \mathbb{F}_q . Then

$$\sum_{c \in C_0} e_p \left(\operatorname{Tr}_{\mathbb{F}_q}(c) \right) = -\frac{(2d-1)\sqrt{q}+1}{2d}, \quad and \quad \sum_{c \in C_k} e_p \left(\operatorname{Tr}_{\mathbb{F}_q}(c) \right) = \frac{\sqrt{q}-1}{2d}$$

for each $1 \leq k \leq 2d - 1$.

Proof. Let ψ denote the canonical additive character of \mathbb{F}_q and let χ be a multiplicative character such that $\chi(g) = \theta$, where θ is a primitive 2*d*-th root of unity. Then χ has order 2*d*, and Corollary A.4 implies that $G(\chi^j) = -\sqrt{q}$ for each $1 \le j \le 2d - 1$.

Using Theorem A.5 with n = e = 2d, for any $a \in \mathbb{F}_q^*$, we have

$$\sum_{c \in \mathbb{F}_q} e_p(\operatorname{Tr}_{\mathbb{F}_q}(ac^{2d})) = \sum_{j=1}^{2d-1} \bar{\chi}^j(a) G(\chi^j) = -\sqrt{q} \sum_{j=1}^{2d-1} \bar{\chi}^j(a) = \sqrt{q} - \sqrt{q} \sum_{j=0}^{2d-1} \bar{\chi}^j(a).$$
(14)

By the orthogonality relations,

$$\sum_{j=0}^{2d-1} \bar{\chi}^j(a) = \begin{cases} 1 & \text{if } a = 1\\ 0 & \text{if } a \neq 1 \end{cases}$$
(15)

Combining (14) and (15), we obtain

$$\sum_{c \in \mathbb{F}_q} e_p(\operatorname{Tr}_{\mathbb{F}_q}(c^{2d})) = -(2d-1)\sqrt{q}, \quad \text{and} \quad \sum_{c \in \mathbb{F}_q} e_p(\operatorname{Tr}_{\mathbb{F}_q}(g^k c^{2d})) = \sqrt{q}$$

for each $1 \le k \le 2d - 1$. The desired conclusion follows immediately from $e_p(\operatorname{Tr}_{\mathbb{F}_q}(0)) = 1$. \Box

The following lemma allows us to compute character values using the associated Gauss sum.

Lemma A.7 ([LN97, Theorem 5.12]). Let χ be a multiplicative character of \mathbb{F}_q . Then for any $a \in \mathbb{F}_q$,

$$\overline{\chi(a)} = \frac{1}{G(\chi)} \sum_{c \in \mathbb{F}_q} \chi(c) e_p \big(\operatorname{Tr}_{\mathbb{F}_q}(ac) \big).$$

The lemma below, which is an application of Lemma A.7 to double character sums, appeared implicitly in [Yip22a, Proposition 4.2].

Lemma A.8. Let q be an odd prime power. Let A be a subset of \mathbb{F}_{q} , and let χ be a nontrivial multiplicative character of \mathbb{F}_q . Then

$$\sum_{a,b\in A} \overline{\chi(a-b)} = \frac{1}{G(\chi)} \sum_{c\in \mathbb{F}_q^*} \chi(c) |S(q,A;c)|^2.$$

Proof. Recall that the trace map is linear. By Lemma A.7,

$$\sum_{a,b\in A} \overline{\chi(a-b)} = \sum_{a,b\in A} \frac{1}{G(\chi)} \sum_{c\in\mathbb{F}_q^*} \chi(c) e_p \big(\operatorname{Tr}_{\mathbb{F}_q}((a-b)c) \big)$$
$$= \frac{1}{G(\chi)} \sum_{c\in\mathbb{F}_q^*} \chi(c) \sum_{a,b\in A} e_p \big(\operatorname{Tr}_{\mathbb{F}_q}((a-b)c) \big)$$

$$= \frac{1}{G(\chi)} \sum_{c \in \mathbb{F}_q^*} \chi(c) \left(\sum_{a \in A} e_p \left(\operatorname{Tr}_{\mathbb{F}_q}(ac) \right) \right) \left(\sum_{b \in A} e_p \left(\operatorname{Tr}_{\mathbb{F}_q}(-bc) \right) \right)$$
$$= \frac{1}{G(\chi)} \sum_{c \in \mathbb{F}_q^*} \chi(c) \left(\sum_{a \in A} e_p \left(\operatorname{Tr}_{\mathbb{F}_q}(ac) \right) \right) \left(\sum_{b \in A} e_p \left(- \operatorname{Tr}_{\mathbb{F}_q}(bc) \right) \right)$$
$$= \frac{1}{G(\chi)} \sum_{c \in \mathbb{F}_q^*} \chi(c) |S(q, A; c)|^2.$$

Recall that S(q, A; c), defined in equation (5), is a scaled Fourier coefficient of the indicator function on A. Thus, Plancherel's identity implies the following lemma.

Lemma A.9 ([Yip22a, Lemma 4.1]). *For any* $A \subset \mathbb{F}_q$ *, we have*

$$\sum_{c \in \mathbb{F}_q^*} |S(q, A; c)|^2 = q|A| - |A|^2$$

APPENDIX B. PROOF OF PROPOSITION 3.2

Let A be a maximum clique in X. Our aim is to show the following two statements:

(1) $|A| \leq \sqrt{q}$; (2) $|A| = \sqrt{q}$ if and only if S(q, A; c) = 0 for all $c \in D'$.

We have mentioned earlier that (1) can be proved independently using the Delsarte bound. The proof given here is genuinely different, and we include it as (1) and (2) naturally fit together. In the following, we use S(c) to denote S(q, A; c) for each $c \in \mathbb{F}_q^*$, and use $T_k = \sum_{c \in C_k} |S(c)|^2$ as a shorthand for each $0 \le k \le 2d - 1$.

Let θ be a primitive 2*d*-th root of unity. Let χ be the multiplicative character of \mathbb{F}_q such that $\chi(g) = \theta$. In order to study the structure of *A*, we associate to the graph *X* the following interpolation function $f: \mathbb{F}_q \to \mathbb{R}$ defined by

$$f(x) = \sum_{j=0}^{2d-1} c_j \overline{\chi^j(x)} \qquad \text{where } c_j = \sum_{\ell=1}^d \theta^{jm_\ell}$$

We claim that f/2d is the indicator function on the connection set D of the Cayley graph X:

$$f(x) = \begin{cases} 2d, & \text{if } x \in D, \\ 0, & \text{if } x \notin D, \end{cases}$$
(16)

where D is defined in equation (1). To prove this claim, recall the orthogonality relations for the group of order |G| = 2d generated by the character χ :

$$\frac{1}{2d}\sum_{j=0}^{2d-1}\chi^j(g^k)\overline{\chi^j(g^\ell)} = \begin{cases} 1 & \text{if } k = \ell, \\ 0 & \text{if } k \neq \ell. \end{cases}$$

Next,

$$f(x) = \sum_{j=0}^{2d-1} \left(\sum_{\ell=1}^{d} \theta^{jm_{\ell}} \right) \overline{\chi^{j}(x)} = \sum_{j=0}^{2d-1} \left(\sum_{\ell=1}^{d} \chi^{j}(g^{m_{\ell}}) \right) \overline{\chi^{j}(x)} = \sum_{\ell=1}^{d} \left(\sum_{j=0}^{2d-1} \chi^{j}(g^{m_{\ell}}) \overline{\chi^{j}(x)} \right)$$

Given $x \in D$, we have $x = g^{2dk+m_{\ell}}$ for some $k \ge 0$ and for some $1 \le \ell \le d$; in this case, there will be exactly one non-zero inner sum, and so $f(x) = f(g^{m_{\ell}}) = 2d$. If x = 0, then f(x) = 0. Given $x \notin D$ such that $x \neq 0$, we have $x = g^{2dk+m}$ for some $k \ge 0$ and $m \neq m_{\ell}$ for any ℓ ; in this case, all the inner sums will vanish and we get $f(x) = f(g^m) = 0$, establishing the claim above.

Since A is a clique, for any $a, b \in A$, if $a \neq b$, we have $a - b \in D$; otherwise f(a - b) = 0. Thus, equation (16) implies that

$$\sum_{a,b\in A} f(a-b) = 2d(|A|^2 - |A|).$$
(17)

On the other hand, using Lemma A.8, we have

$$\begin{split} \sum_{a,b\in A} f(a-b) &= \sum_{a,b\in A} \sum_{j=0}^{2d-1} c_j \overline{\chi^j(a-b)} \\ &= \sum_{a,b\in A} c_0 \overline{\chi^0(a-b)} + \sum_{j=1}^{2d-1} \sum_{a,b\in A} c_j \overline{\chi^j(a-b)} \\ &= d(|A|^2 - |A|) + \sum_{j=1}^{2d-1} \frac{c_j}{G(\chi^j)} \left(\sum_{c\in \mathbb{F}_q^*} \chi^j(c) |S(c)|^2\right) \\ &= d(|A|^2 - |A|) + \sum_{j=1}^{2d-1} \frac{c_j}{G(\chi^j)} \left(\sum_{k=0}^{2d-1} \sum_{c\in C_k} \chi^j(c) |S(c)|^2\right) \\ &= d(|A|^2 - |A|) + \sum_{j=1}^{2d-1} \frac{c_j}{G(\chi^j)} \left(\sum_{k=0}^{2d-1} \frac{c_j}{c\in C_k} \chi^j(c) |S(c)|^2\right) \end{split}$$

By Corollary A.4, $G(\chi^j) = -\sqrt{q}$ for each $1 \le j \le 2d - 1$. Thus,

$$\sum_{a,b\in A} f(a-b) = d(|A|^2 - |A|) - \frac{1}{\sqrt{q}} \sum_{k=0}^{2d-1} \left(\sum_{j=1}^{2d-1} c_j \theta^{jk}\right) T_k.$$
(18)

Observe that

$$\sum_{j=1}^{2d-1} c_j \theta^{jk} = \sum_{j=1}^{2d-1} \sum_{\ell=1}^d \theta^{jm_\ell} \theta^{jk} = \sum_{\ell=1}^d \sum_{j=1}^{2d-1} \theta^{j(m_\ell+k)}$$
$$= -d + \sum_{\ell=1}^d \sum_{j=0}^{2d-1} \theta^{j(m_\ell+k)}$$
$$= \begin{cases} d, & \text{if } k \equiv -m_1, -m_2, \dots, -m_d \pmod{2d}, \\ -d, & \text{otherwise.} \end{cases}$$
(19)

For the last step, we used the following elementary fact based on the geometric series:

$$\sum_{j=0}^{2d-1} \theta^{j\ell} = \begin{cases} 0 & \text{if } \ell \not\equiv 0 \pmod{2d}, \\ 2d & \text{if } \ell \equiv 0 \pmod{2d}. \end{cases}$$

Combining equations (18) and (19), we obtain that

$$\sum_{a,b\in A} f(a-b) = d(|A|^2 - |A|) + \frac{d}{\sqrt{q}} \sum_{k\neq -m_\ell} T_k - \frac{d}{\sqrt{q}} \sum_{\ell=1}^d T_{-m_\ell}$$
$$\leq d(|A|^2 - |A|) + \frac{d}{\sqrt{q}} \sum_{k=0}^{2d-1} T_k$$
$$= d(|A|^2 - |A|) + \frac{d}{\sqrt{q}} (q|A| - |A|^2).$$

In the last step, we used Lemma A.9. Comparing the inequality above with equation (17), it follows that for a clique A, we have $|A|^2 - |A| \le \frac{1}{\sqrt{q}}(q|A| - |A|^2)$, which implies that $|A| \le \sqrt{q}$. This completes the proof of (1). Moreover, the equality $|A| = \sqrt{q}$ holds if and only if

$$T_{-m_1} = T_{-m_2} = \ldots = T_{-m_d} = 0.$$
 (20)

This implies that S(q, A; c) = 0 for all $c \in D' = \bigcup_{\ell=1}^{d} C_{-m_{\ell}}$, proving (2).

APPENDIX C. ALGORITHMS AND NUMERICAL COMPUTATIONS

In this section, we present algorithms to find and classify maximum cliques in semi-primitive pseudo-Paley graphs. We also report results from our experiments using SageMath [Sag21] to support the conjectures in this paper.

Using SageMath, we can find several examples of pseudo-Paley graphs of order q coming from unions of semi-primitive cyclotomic classes with clique number \sqrt{q} . The following table summarizes some of these examples. The set I indicates the choice of C_i 's in the connection set D. For example, the second row corresponds to the Cayley graph $Cay(\mathbb{F}_{5^4}, D)$ where $D = C_0 \cup C_1 \cup C_3$ and $C_j = \{g^{6i+j} \mid 0 \le i \le \frac{5^4-1}{6}\}$ and g is a primitive root of \mathbb{F}_{5^4} .

q	d	Ι	$\omega(X)$
5^{4}	3	$\{0, 1, 3\}$	25
7^{4}	4	$\{0, 1, 2, 4\}$	49
7^{4}	4	$\{0, 1, 3, 6\}$	49
3^{8}	5	$\{0, 1, 2, 3, 5\}$	81
3^{8}	5	$\{0, 1, 3, 6, 7\}$	81
11^{4}	6	$\{0, 1, 2, 3, 6, 7\}$	121
11^{4}	6	$\{0, 1, 2, 4, 5, 8\}$	121
11^{4}	6	$\{0, 1, 2, 3, 5, 10\}$	121
13^{4}	7	$\{0, 1, 2, 3, 5, 6, 9\}$	169
13^{4}	7	$\{0, 1, 2, 4, 7, 8, 10\}$	169
13^{4}	7	$\{0, 1, 2, 5, 7, 9, 10\}$	169

TABLE 1. Graphs X = PP(q, 2d, I) with $\omega(X) = \sqrt{q}$

In each of the graphs listed above, there are exactly 2 cliques of size \sqrt{q} containing \mathbb{F}_{p^t} , which supports Conjecture 5.6. Both cliques are subspaces over \mathbb{F}_{p^t} which supports Conjecture 1.4.

In general, finding the clique number of a Cayley graph is NP-hard [GR17]. It is not possible to exactly compute the clique number of X = PP(q, 2d, I) for q > 2000 using the current computational power. In contrast, finding the common neighbors of \mathbb{F}_p inside X takes only polynomial-time, and SageMath can return the clique number of a graph with < 700 vertices in a few minutes. Therefore, we employ the following algorithm.

Algorithm 1: Find the size of a maximum clique containing \mathbb{F}_p inside $PP(q, 2d, I)$				
$k = \min\{n \in \mathbb{N} : q/2^n < 700\}$				
$V = \{0, 1, \dots, k-1\} \cup \bigcap_{i=0}^{k-1} \operatorname{Neighbors}(j)$	// Finding the set of common neighbors			
Y = PP(q, 2d, I)[V]	// Taking the induced subgraph			
return $\omega(Y)$				

Since the graph is $\frac{q-1}{2}$ -regular, we expect that the subgraph Y has roughly $q/2^k$ many vertices because the graphs we consider are pseudo-random [CGW89, KP04]. This heuristic can be proved rigorously using character sums and Weil's bound; see [GS71] for a precise version. Since $q/2^k < 700$, SageMath can return the clique number of Y quickly. We used this algorithm to handle the exceptional cases in Theorem 1.8.

In general, it suffices to consider maximum cliques containing $\{0, 1\}$. The reasoning is given at the beginning of the proof of Theorem 1.7. When $q \le 7^4$, we have $q/2^2 < 700$, and we can perform the algorithm outlined above with k = 2 to find the clique number of all possible PP(q, 2d, I). This verifies the main Conjecture 1.4 for values of q up to 7^4 .

In the final part of the paper, we present an efficient algorithm to verify Conjecture 5.9 for a given prime p. The strategy is to fix an \mathbb{F}_p -subspace $V \subset \mathbb{F}_{p^4}$ with basis $\{1, a\}$ with $a \in C_0$, and search for an index set $I \subset \{0, 1, \ldots, p\}$ with |I| = (p+1)/2 such that $V \setminus \{0\} \subset \bigcup_{j \in I} C_j$. In order for V to form a clique in $PP(p^4, p+1, I)$, by Theorem 1.6, we must have

$$I = \left\{ 0 \le j \le p : |V \cap C_j| = \frac{p^2 - 1}{(p+1)/2} = 2(p-1) \right\}.$$
 (21)

Since $1 \in V$, it follows that $0 \in I$. Note that $V \setminus \{0\}$ is a disjoint union of (p + 1) many \mathbb{F}_p^* cosets with representatives $R = \{1, a, a + 1, a + 2, \dots, a + (p - 1)\}$. By Lemma 5.1, \mathbb{F}_p^* is a subgroup of C_0 , and so each cyclotomic class C_j is a disjoint union of \mathbb{F}_p^* -cosets. As a result, we can decompose $(V \setminus \{0\}) \cap C_j$ into \mathbb{F}_p^* -cosets. Therefore, equation (21) is equivalent to the statement that C_j contains exactly two \mathbb{F}_p^* -coset representatives in R for each $j \in I$. Applying this idea, we can design a polynomial-time algorithm as follows:

Algorithm 2: Count the number of I such that $0 \in I$ and $\omega(PP(p^4, p+1, I)) = p^2$.

```
ValidSets \leftarrow \emptyset
for k \leftarrow 0 to p^2 do
    a \leftarrow q^{(p+1)k}
     L \leftarrow [] // initiating an empty list
    R \leftarrow \{1, a, a+1, \dots, a+(p-1)\}
    for r \in R do
        t \leftarrow \log_{q}(r) // finding exponent t with g^{t} = r using the discrete logarithm
      L \leftarrow L \text{ append } \{t \pmod{p+1}\}
    I \leftarrow \operatorname{set}(L)
                             // remove duplicates from the list L to get the set I
                   // assume I is a valid set
    flag \leftarrow 1
    for j \in I do
         if L. count(j) \neq 2 then
              flag \leftarrow 0 // I is not valid
           _ break
    if flag = 1 then
      ValidSets \leftarrow ValidSets \cup {I}
```

return #ValidSets // return cardinality

Using SageMath, Conjecture 5.9 has been checked to hold for all odd primes p < 100. We also found that each index set I (except for $\{0, 2, ..., p - 1\}$) appeared exactly twice, which is consistent with Conjecture 5.6.

As an example, SageMath outputs the following index set when p = 97 and $a = g^{p+1} = g^{98}$:

46, 50, 51, 53, 56, 59, 62, 70, 72, 75, 76, 78, 80, 82, 84, 87, 88, 90, 92, 93, 94, 95, 96, 97

In other words, $\omega(PP(97^4, 98, I_0)) = 97^2$ with a maximum clique given by $V = \mathbb{F}_{97} \oplus g^{98}\mathbb{F}_{97}$.

DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE, SANTA CLARA UNIVERSITY, 500 EL CAMINO REAL, USA 95053

Email address: sasgarli@scu.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF BRITISH COLUMBIA, 1984 MATHEMATICS ROAD, CANADA V6T 1Z2

Email address: kyleyip@math.ubc.ca