

# Lorentz-boost-driven magneto-optics in Dirac matter

J. Wyzula,<sup>1</sup> X. Lu,<sup>2</sup> D. Santos-Cottin,<sup>3</sup> D. K. Mukherjee,<sup>2,4</sup> I. Mohelský,<sup>1</sup> F. Le Maréchal,<sup>3</sup> J. Novák,<sup>5</sup> M. Novák,<sup>6</sup> R. Sankar,<sup>7</sup> Y. Krupko,<sup>1,8</sup> B. A. Piot,<sup>1</sup> W.-L. Lee,<sup>7</sup> A. Akrap,<sup>3</sup> M. Potemski,<sup>1,9</sup> M. O. Goerbig,<sup>2</sup> and M. Orlita<sup>1,10,\*</sup>

<sup>1</sup>LNCMI-EMFL, CNRS UPR3228, Univ. Grenoble Alpes, Univ. Toulouse,  
Univ. Toulouse 3, INSA-T, Grenoble and Toulouse, France

<sup>2</sup>Laboratoire de Physique des Solides, Université Paris Saclay,  
CNRS UMR 8502, F-91405 Orsay Cedex, France

<sup>3</sup>Department of Physics, University of Fribourg, Chemin du Musée 3, 1700 Fribourg, Switzerland

<sup>4</sup>Department of Physics, Indiana University, Bloomington, Indiana 47405, USA

<sup>5</sup>Department of Condensed Matter Physics, Masaryk University, Kotlářská 2, 611 37 Brno, Czech Republic

<sup>6</sup>Department of Physics, Faculty of Science, University of Zagreb, 10000 Zagreb, Croatia

<sup>7</sup>Institute of Physics, Academia Sinica, Nankang, 11529 Taipei, Taiwan

<sup>8</sup>Institut d'Electronique et des Systemes, UMR CNRS 5214,  
Université de Montpellier, 34000, Montpellier, France

<sup>9</sup>Institute of Experimental Physics, Faculty of Physics,  
University of Warsaw, ul. Pasteura 5, 02-093 Warszawa, Poland

<sup>10</sup>Charles University, Faculty of Mathematics and Physics,  
Institute of Physics, Ke Karlovu 5, 121 16 Prague 2, Czech Republic

Optical response of crystalline solids is to a large extent driven by excitations that promote electrons among individual bands. This allows one to apply optical and magneto-optical methods to determine experimentally the energy band gap – a fundamental property crucial to our understanding of any solid – with a great precision. Here we show that such conventional methods, applied with great success to many materials in the past, do not work in topological Dirac semimetals with a dispersive nodal line. There, the optically deduced band gap depends on how the magnetic field is oriented with respect to the crystal axes. Such highly unusual behaviour is explained in terms of band-gap renormalization driven by Lorentz boosts.

Ever since the advent of graphene and topological materials, relativistic physics has become an integral part of condensed-matter sciences [1–3]. While emergent, it is important to stress that this type of relativity is pertinent beyond the dispersion of the low-energy excitations in many different solids. Klein tunneling [4–6] and the chiral anomaly [7–14] represent well-known examples. One of the most salient aspects of relativity is probably the particular dependence of energy on the frame of reference: for a particle of mass  $m$  moving at a speed  $u$  lower than the speed of light  $c$ , a Lorentz boost to the comoving frame of reference changes the particle's energy from  $E$  to  $E/\gamma = mc^2$ , in terms of the Lorentz factor  $\gamma = 1/\sqrt{1-\beta^2}$  and the rapidity  $\beta = u/c$ . Can one observe this relativistic renormalization equally in topological materials governed by the Dirac Hamiltonian or a variant of it, upon the replacement of  $c$  by a characteristic velocity  $v$ ?

While such effects of Lorentz covariance have been theoretically studied namely in the framework of materials with tilted conical bands both in 2D [15–17] and 3D materials [18–20], an experimental verification is yet lacking. The key finding of this paper is that Lorentz boosts are experimentally relevant in Dirac materials. Most notably and unexpectedly, we find that the optical band gap extracted from magneto-optical measurements in the dispersive nodal-line Dirac semimetal niobium diarsenide (NbAs<sub>2</sub>) depends on the orientation of the explored crystal. As we show below, this orientation defines a particu-

lar Lorentz boost by fixing the angle between the applied magnetic field and the direction of the relevant part of the nodal line.

In order to appreciate the link between magnetic-field orientation and Lorentz boosts, let us first consider a 2D cone with a gap of  $2\Delta$  and an asymptotic velocity of  $v$  which is tilted by an additional velocity parameter  $u$  (Fig. 1a). In the presence of an out-of-plane magnetic field, with  $\mathbf{B} \perp \mathbf{u}$ , the tilt can be formally viewed as a drift velocity of electrons in the crossed magnetic and effective electric field,  $\mathcal{E} = uB$ . In this particular case, the problem of the electron motion in a tilted cone becomes mathematically equivalent to the dynamics of a relativistic charge carrier in the crossed electric and magnetic fields, and therefore, governed by fully Lorentz-covariant Dirac and Maxwell equations. This covariant formulation, and thus the use of Lorentz transformations, then allows us to calculate the energy spectrum in a reference frame where the (effective) electric field vanishes, meaning  $u = 0$  [16, 17, 21]. Notably, a similar relativistic-like approach has been invoked in the past, in order to understand behaviour of narrow-gap semiconductors [22–24] in real crossed electric and magnetic fields.

The impact of the tilt  $\mathbf{u}$  – or in the sense of reasoning above, the impact of the Lorentz boost – on the spectrum of Landau-quantized 2D Dirac electrons is profound:

$$E_n = \pm \sqrt{(\Delta/\gamma)^2 + 2eB\hbar v^2/\gamma^3}, \quad n = 0, 1, 2, \dots \quad (1)$$

The energy band gap and velocity parameter are renor-




FIG. 1. (a) Schematic sketch of a hyperbolic dispersion for (massive) Dirac electrons with an additional tilt:  $E(\mathbf{k}) = -\hbar v k_x \pm \sqrt{\Delta^2 + \hbar^2 v^2 k^2}$ . The dashed lines correspond to asymptotic dispersions. (b) Brillouin zone of NbAs<sub>2</sub> with the crossings of the nodal lines with the Fermi level and selected high-symmetry points marked. The blue lines show approximate position of two nodal lines, reflecting the DFT calculations and overall conclusions in Ref. [25]. (c) Schematic view of the nodal-line dispersion over a half of the Brillouin zone along the direction of propagation. The band gaps at the crossing points with the Fermi energy, in the dispersive and the (approximately) flat parts of the nodal line, are  $2\Delta_D$  and  $2\Delta_F$ , respectively. (d) Infrared reflectivity, and the extracted optical conductivity, measured on (001)-oriented facet of NbAs<sub>2</sub>, using light polarized along the  $a$  and  $b$  crystallographic axes.

malized by the Lorentz factor  $\gamma$ . For large tilts,  $u \geq v$ , the spectrum collapses and marks transitions between regimes referred to as magnetic and electric [19, 26]. In the semi-classical picture, this crossover corresponds to transitions from closed towards open cyclotron orbits in type-I and type-II conical bands (Figs. 1a).

Let us now discuss how a Lorentz boost allows us to understand the magneto-optical response of NbAs<sub>2</sub> – a nearly compensated 3D nodal-line semimetal [27] with a monoclinic crystal lattice (space group C2/m [28]). Theory, in agreement with experimental studies performed so far [25, 27], indicate the existence of two nodal lines. Mathematically, these are 1D objects (curves) in momentum space,  $\mathbf{k}_{\text{line}}$ , along which the band gap vanishes, or at least, reaches its local minimum. In NbAs<sub>2</sub>, the nodal lines are open, propagating through the Brillouin zone and located symmetrically with respect to the  $\Gamma$ -Y-Z mirror plane (Fig. 1b).

Consider first a plane perpendicular to the local direction  $\boldsymbol{\tau}$  of the nodal lines, in which electrons can be well described using a model for 2D massive Dirac electrons (left panel of Fig. 1a). The corresponding velocity parameter  $v$  and the spin-orbit gap  $2\Delta$ , typical of all currently studied nodal-line or nodal-loop semimetals [29–

31], may vary along the line. Essentially for this work, the nodal lines in NbAs<sub>2</sub> disperse with momentum and each approaches the Fermi energy four times within the Brillouin zone (Fig. 1b,c). There are two types of such crossings, associated respectively with a dispersive and a flat part [25]. They are located at  $\mathbf{k}_{\text{line}}^D$  and  $\mathbf{k}_{\text{line}}^F$  and characterized by the local directions  $\boldsymbol{\tau}_D$  and  $\boldsymbol{\tau}_F$  (Fig. 1c).

The optical response of NbAs<sub>2</sub> at low photon energies comprises direct signatures of nodal lines [25, 32, 33]. Two steps are clearly visible at the onset of interband absorption in the optical conductivity (Fig. 1d down). They correspond to the local band gaps:  $2\Delta_D = (88 \pm 2)$  meV and  $2\Delta_F = (113 \pm 2)$  meV, at the dispersive and flat parts, respectively. Above this onset, the optical conductivity increases linearly in  $\omega$ . While such behavior resembles systems with 3D conical bands (see, e.g., Refs. [34–36]), in NbAs<sub>2</sub>, it is due to the occupation effect (Pauli-blocking) along the dispersive part of the nodal line [25, 32, 37, 38]. The anisotropy of the optical response (Fig. 1d) reflects the orientation of the nodal lines, implying significantly higher Drude-type and interband absorption strength for the radiation polarized along the  $b$  axis (approximately perpendicular to  $\boldsymbol{\tau}$ ) [25]. Strong anisotropy, implying elongated Fermi surfaces, was also

observed in magneto-transport experiments[33, 39–41].

To explore the magneto-optical response, we have identified a number of crystallographic facets on several NbAs<sub>2</sub> monocrystals using the standard  $x$ -ray technique [33]. Then, a series of infrared reflectivity measurements has been carried out, using non-polarized radiation and the Faraday configuration. To keep the geometry as simple as possible, we selected facets with a zero middle index ( $n0m$ ). Thus the vector  $\mathbf{B}$  always lied in the mirror plane of the NbAs<sub>2</sub> crystals and made the identical angles  $\theta_F$  and  $\theta_D$  with the local directions,  $\boldsymbol{\tau}_F$  and  $\boldsymbol{\tau}_D$ , of the two mirror-symmetric nodal lines at the flat and dispersive parts, respectively, where the Fermi level is crossed.

The relative magneto-reflectivity of NbAs<sub>2</sub>,  $R_B/R_0$ , measured with  $\mathbf{B}$  applied perpendicular to the (101), (201), (20 $\bar{1}$ ) and (001) crystallographic planes are plotted in Fig. 2. Data measured on other facets are shown in Supplementary Materials [33]. The observed response comprises a series of sharp resonances with a weakly sub-linear dependence on  $B$  that can be directly associated with interband inter-Landau-level (inter-LL) excitations. The observed response – in position, spacing and relative/absolute intensity of inter-LL transitions – strongly varies with the explored facet. Across the board, this response includes two characteristic sets of interband inter-LL excitations: (i) the upper set, with transitions that always extrapolate to the energy of  $2\Delta_F$  in the zero-field limit and are thus clearly associated with the flat crossing of the nodal line and (ii) the lower set of transitions that extrapolate, depending on the particular facet, to the energy equal to or lower than  $2\Delta_D$  (see  $B = 0$  extrapolations using yellow dashed lines in Fig. 2).

This observation – an orientation-dependent gap – is the main finding of this work and it ventures beyond the common knowledge in LL spectroscopy of solids. In conventional materials, the slope  $d(\hbar\omega)/dB$ , and the spacing of inter-LL excitations may depend on the orientation of the crystal with respect to the magnetic field, as well as on the experimental configuration (e.g., Faraday versus Voigt). Nevertheless, the apparent band gap obtained from the zero-field extrapolation of interband inter-LL excitations is widely used as an unambiguous estimate of the separation between electronic bands. Quite unexpectedly, such an apparently evident approach fails in our case here.

To show that this orientation-dependent gap is a signature of the relativistic properties in the present material, let us consider the following minimal Hamiltonian for a dispersive nodal line:

$$\hat{H} = (\varepsilon_0 + \hbar w q_{\text{line}})\mathbb{1} + \begin{bmatrix} \Delta & \hbar v(q_x - iq_y) \\ \hbar v(q_x + iq_y) & -\Delta \end{bmatrix}, \quad (2)$$

where the wave vector  $\mathbf{q} = (q_x, q_y, q_{\text{line}}) = \mathbf{k} - \mathbf{k}_{\text{line}}$  is defined with respect to any point,  $\mathbf{k}_{\text{line}}$ , on the nodal line, where the midgap energy is  $\varepsilon_0$  and the gap  $2\Delta$ . In

the plane  $(q_x, q_y)$  perpendicular to the local direction  $\boldsymbol{\tau}$ , the Hamiltonian (2) becomes that of a 2D massive Dirac electron.

When a magnetic field  $\mathbf{B}$  is applied, making an angle  $\theta$  with the local direction  $\boldsymbol{\tau}$ , Hamiltonian (2) becomes equivalent [33] to the one describing 2D massive Dirac electrons in a comoving frame subjected to a magnetic field,  $B_{\perp} = B \cos \theta$ . This is due to the diagonal dispersive term which can be viewed as an effective in-plane electric field,  $\mathcal{E}_{\parallel} = wB \sin \theta$ , or alternatively, as an effective in-plane drift (tilt) velocity,  $w \tan \theta$ , perpendicular to the magnetic field. Apart from the addition of a dispersive term  $\hbar w q_B / \cos \theta$ , where  $q_B$  is the wave vector in the  $B$ -field direction, the LL spectrum has the Lorentz-boost-renormalized form of Eq. 1. One only has to replace  $B$  by its perpendicular component  $B_{\perp}$  and set the rapidity as  $\beta = \mathcal{E}_{\parallel} / (B_{\perp} v) = (w/v) \tan \theta$  which then defines the Lorentz factor  $\gamma$ . Depending on the orientation of the magnetic field with respect to the nodal line, we then expect a pseudo-relativistic decrease of the band gap and the velocity parameter

$$2\Delta \rightarrow \frac{2\Delta}{\gamma} \quad \text{and} \quad v \rightarrow \frac{v}{\gamma^{3/2}}. \quad (3)$$

For angles exceeding the critical value of  $\tan^{-1}(v/w)$ , the quantization into LLs collapses in a way analogous to overtilted 3D conical bands. [19] As compared to 2D Dirac systems,  $\gamma$  is now tunable by the angle  $\theta$ , which thus enables us to continuously monitor the gap renormalization until the extinction of discrete LLs (see Fig. 2).

To analyze our experimental data quantitatively in view of the above theoretical picture, we have focused on the lowest observed line in both sets and assign it to the inter-LL excitation  $0 \leftrightarrow 1$ . Even though the pseudo-relativistic renormalization may profoundly alter selection rules [17, 33], this transition is theoretically [33] expected to remain strong for any  $\beta < 1$  and its energy reads:

$$\hbar\omega_{0 \leftrightarrow 1} = \Delta^{\text{eff}} + \sqrt{(\Delta^{\text{eff}})^2 + 2e\hbar B (v^{\text{eff}})^2}. \quad (4)$$

Here,  $\Delta^{\text{eff}}$  and  $v^{\text{eff}}$  are effective parameters which include the impact of the Lorentz boost and of the geometrical factor  $\cos \theta$ .

In our data analysis, we associated the maxima in  $R_B/R_0$  spectra with positions of inter-LL excitations, a solid assumption in the vicinity of the plasma edge (cf. Fig. 1d and see Methods). The effective values of the band gap and velocity parameter derived for all explored facets are presented in Figs. 3a-d. The response of the flat part matches perfectly the expectations for a Landau-quantized 2D massive Dirac system – the effective band gap  $2\Delta_F^{\text{eff}} = (113 \pm 2)$  meV stays constant within the experimental error (Fig. 3a). The variation of the effective velocity (Fig. 3b) with the facet reflects the geometrical




FIG. 2. Relative magneto-reflectivity of NbAs<sub>2</sub>,  $R_B/R_0$ , in the magnetic field applied along four different reciprocal space directions: (101), (201), (20 $\bar{1}$ ) and (001), which make angles 62°, 51°, -7° and 90° with the  $a$  crystallographic axis, respectively. Horizontal gray bars correspond to positions of the two steps in the onset of interband excitations at  $2\Delta_D$  and  $2\Delta_F$  at  $B = 0$ , cf. Fig. 1d. The yellow values indicate the apparent band gap,  $2\Delta_D^{\text{eff}}$ , deduced using a simple linear zero-field extrapolation of inter-LL resonances belonging to the lower set (yellow dashed lines).

factor,  $v_F^{\text{eff}} = v_F \sqrt{\cos \theta}$ , where  $v_F = (6.3 \pm 0.3) \times 10^5$  m/s, due to mutual orientation of  $\mathbf{B}$  and  $\boldsymbol{\tau}_F$ . This allows us to deduce the local direction of the flat part. The best agreement has been found for  $\boldsymbol{\tau}_F$  in the mirror  $a$ - $c$  plane,  $(0 \pm 2)^\circ$ , but inclined by  $(-23 \pm 2)^\circ$  from the  $(a$ - $b$ ) plane, in perfect agreement with conclusions reached by Shao et al. [25].

In contrast, the effective gap  $2\Delta_D^{\text{eff}}$  deduced for the dispersive part spreads over an interval greater than 10 meV (Fig. 3c). To compare this behaviour with our Lorentz-boost induced gap renormalization (3), we have minimized the difference between the theoretical expectations and experimental values of the effective gap and velocity, deduced for all facets, by varying parameters  $\Delta_D$ ,  $v_D$ ,  $w$  as well as the local direction  $\boldsymbol{\tau}_D$ . A very good agreement was obtained (dashed lines in Figs. 3c,d) for the following parameters:  $v_D = (5.3 \pm 0.5) \times 10^5$  m/s,  $w = (1.5 \pm 0.5) \times 10^5$  m/s,  $2\Delta_D = (89 \pm 2)$  meV, and the local direction  $\boldsymbol{\tau}_D$  that deviates by  $(25 \pm 10)^\circ$  from the mirror  $(a$ - $c$ ) plane and by  $(-5 \pm 2)^\circ$  from the  $a$ - $b$  plane. These parameters agree well with the previous estimates given by Shao et al. [25]. Slightly larger values were found for  $w$  and the angle of  $\boldsymbol{\tau}_D$  with respect to the  $a$ - $c$  plane. Importantly, the angle  $\theta_D$  reaches large values for certain explored facets, thus implying a rapidity exceeding unity. In such cases, the lower set of inter-LL excitations disappears entirely from the magneto-optical response, suggesting the complete collapse of the LL spectrum. This happens for the data collected on the (001)-oriented facet ( $\theta_D \approx 85^\circ$ ) presented in Fig. 2d, and also

the (20 $\bar{3}$ )-oriented facet ( $\theta_D \approx 64^\circ$ ), see Supplementary Materials [33].

In addition to the Lorentz-boost renormalization of the spectrum, our model for the dispersive nodal line implies a departure from the conventional electric-dipole selection rules,  $n \rightarrow n \pm 1$ , which are generally valid for all isotropic systems [42]. To illustrate this, we have numerically evaluated the matrix elements for electric-dipole interband excitations between different pairs of LLs ( $n = 0 \dots 6$ ) and visualized them graphically in Fig. 3e. We have chosen two particular angles,  $\theta_D = 61$  and  $28^\circ$ , which correspond to the magnetic field oriented perpendicular to the (101) and (20 $\bar{1}$ ) planes, respectively. For small angles  $\theta_D$ , the magneto-optical response is dominated by  $n \rightarrow n \pm 1$  transitions, although, other excitations emerge as well (e.g.,  $n \rightarrow n \pm 2$ ). In contrast, for larger angles  $\theta_D$ , one finds a plethora of optical transitions. The dominant ones follow the rule-of-thumb selection rules  $n \rightarrow \alpha n$  and  $n \rightarrow n/\alpha$ , where  $\alpha$  is an integer ( $\alpha = 4 - 6$  in the left panel of Fig. 3e), in agreement with preceding works on tilted 3D cones [17, 19].

As seen in Fig. 3e, the lowest in energy transitions  $1 \leftrightarrow 0$  stay strong as long as  $\beta < 1$  and were used to deduce the effective parameters  $2\Delta_D^{\text{eff}}$  and  $v_D^{\text{eff}}$ . These parameters may now, in turn, be used to identify excitations between LLs with higher indices and thus get experimental insights into the selection rules. To this end, we compare in Fig. 3f the experimental data collected with  $B$  perpendicular to the (101) and (20 $\bar{1}$ ) crystallographic planes, with the expected positions of selected






FIG. 3. Effective band gap and velocity parameter obtained by a fit of the massive-Dirac model to the lowest inter-LL transition in sets belonging to the flat and dispersive crossings with the Fermi energy: (a),(b) and (c),(d), respectively. (e) Absolute values of matrix elements (the darker color, the stronger the transition) for electric-dipole inter-LL interband excitations (for LLs  $n = 0 \dots 6$  in both conduction and valence bands) calculated for  $2\Delta_D = 89$  meV,  $v_D = 5.3 \times 10^5$  m/s,  $w = 1.5 \times 10^5$  m/s and for two different angles  $\theta_D = 61^\circ$  and  $28^\circ$  between  $\tau_D$  and  $\mathbf{B}$ . These two angles correspond to the configuration with  $\mathbf{B}$  perpendicular to the crystallographic planes (101) and  $(20\bar{1})$ , respectively, for which the experimental  $R_B/R_0$  traces are plotted in (f) in a form of false-color plots. The dotted lines show selected inter-LL excitations from the dispersive part of the nodal line, assumed to be electric-dipole active based on the matrix elements presented in (e). We use the same color-framing/coding in (e) and (f) to facilitate the identification of individual transitions. For instance, the lowest (red-dotted) line in (f) correspond to the  $0 \leftrightarrow 1$  transitions. The gray dashed lines show the expected transitions in the flat part (selection rules  $n \rightarrow n \pm 1$ , for  $v_F = 6.3 \times 10^5$  m/s and  $\Delta_F = 113$  meV).

interband inter-LL excitations (dotted lines) calculated using the corresponding effective gap and velocity parameters. To facilitate the comparison, we use the color-framing/coding introduced in Fig. 3e. In line with our expectations, we identify  $n \rightarrow n \pm 1$  and  $n \rightarrow n \pm 2$  excitations in the response on the  $(20\bar{1})$ -oriented facet which implies a relatively small angle  $\theta_D$  (Fig. 3f right). In contrast, when the magnetic field is applied perpendicularly to the (101) crystallographic plane (Fig. 3f left), we identify transitions with a greater change of the LL index, such as  $1 \rightarrow 4$  or  $1 \rightarrow 3$ , and no line following the standard  $n \rightarrow n \pm 1$  selection rule is found, except for the

lowest one,  $0 \leftrightarrow 1$ . In both cases, the gray dashed lines show the expected response of the flat part of the nodal line that follow the standard  $n \rightarrow n \pm 1$  selection rules.

In conclusion, we have found that the optical band gap of the nodal-line semimetal NbAs<sub>2</sub> measured via magneto-optical spectroscopy depends on the facet explored in the experiment. This observation is understood as a consequence of the pseudo-relativistic renormalization of the band gap within a Lorentz boost determined by the slope of the dispersive nodal line. The slope defines, together with the direction of the applied magnetic field, the tilt of the conical dispersion of a massive 2D

Dirac electron in the plane perpendicular to the applied magnetic field. Our findings show that the emergent relativistic description of topological quantum materials in terms of Dirac Hamiltonians or its variants as well as the use of Lorentz transformations can be pushed surprisingly far. The observed Lorentz-boost-driven renormalization can be also viewed as an analogue of the well-known Franz-Keldysh effect in the magnetic field [22, 43], nevertheless, in our case, with no real electric field applied.

### Acknowledgments

We acknowledge discussions with T. Brauner and S. Tchoumakov. The work has been supported by the ANR DIRAC3D project (ANR-17-CE30-0023) and exchange programme PHC ORCHID (47044XE). A. A. acknowledges funding from the Swiss National Science Foundation through Project No. PP00P2-170544. M.N. acknowledges the support of CeNIKS project co-financed by the Croatian Government and the EU through the European Regional Development Fund - Competitiveness and Cohesion Operational Program (Grant No. KK.01.1.1.02.0013). R.S. acknowledges financial support provided by the Ministry of Science and Technology in Taiwan under project numbers MOST-110-2112-M-001-065-MY3 as well as Academia Sinica for the budget of AS-iMate-109-13. D.K.M. acknowledges partial support from NSF Grant No. DMR-1914451, and the Research Corporation for Science Advancement through a Cottrell SEED award. This work was also supported by CNRS through IRP "TeraMIR" and by ANR Colector (ANR-19-CE30-0032). We also acknowledge the support of the LNCMI-CNRS in Grenoble, a member of the European Magnetic Field Laboratory (EMFL).

---

\* milan.orlita@lncmi.cnrs.fr

- [1] A. Bansil, H. Lin, and T. Das, Colloquium: Topological band theory, *Rev. Mod. Phys.* **88**, 021004 (2016).
- [2] C.-K. Chiu, J. C. Y. Teo, A. P. Schnyder, and S. Ryu, Classification of topological quantum matter with symmetries, *Rev. Mod. Phys.* **88**, 035005 (2016).
- [3] N. P. Armitage, E. J. Mele, and A. Vishwanath, Weyl and Dirac semimetals in three-dimensional solids, *Rev. Mod. Phys.* **90**, 015001 (2018).
- [4] O. Klein, Die reflexion von elektronen an einem potenzsprung nach der relativistischen dynamik von Dirac, *Zeitschrift für Physik* **53**, 157 (1929).
- [5] M. I. Katsnelson, K. S. Novoselov, and A. K. Geim, Chiral tunnelling and the Klein paradox in graphene, *Nature Phys.* **2**, 620 (2006).
- [6] A. F. Young and P. Kim, Quantum interference and Klein tunnelling in graphene heterojunctions, *Nature Phys.* **5**, 222 (2009).
- [7] H. B. Nielsen and M. Ninomiya, The Adler-Bell-Jackiw anomaly and Weyl fermions in a crystal, *Physics Letters B* **130**, 389 (1983).
- [8] D. T. Son and B. Z. Spivak, Chiral anomaly and classical negative magnetoresistance of Weyl metals, *Phys. Rev. B* **88**, 104412 (2013).
- [9] J. Xiong, S. K. Kushwaha, T. Liang, J. W. Krizan, M. Hirschberger, W. Wang, R. J. Cava, and N. P. Ong, Evidence for the chiral anomaly in the Dirac semimetal Na<sub>3</sub>Bi, *Science* **350**, 413 (2015).
- [10] X. Huang, L. Zhao, Y. Long, P. Wang, D. Chen, Z. Yang, H. Liang, M. Xue, H. Weng, Z. Fang, X. Dai, and G. Chen, Observation of the chiral-anomaly-induced negative magnetoresistance in 3D Weyl semimetal TaAs, *Phys. Rev. X* **5**, 031023 (2015).
- [11] A. Burkov, Chiral anomaly without relativity, *Science* **350**, 378 (2015).
- [12] R. D. dos Reis, M. O. Ajeesh, N. Kumar, F. Arnold, C. Shekhar, M. Naumann, M. Schmidt, M. Nicklas, and E. Hassinger, On the search for the chiral anomaly in Weyl semimetals: the negative longitudinal magnetoresistance, *New J. Phys.* **18**, 085006 (2016).
- [13] S. Jia, S.-Y. Xu, and M. Z. Hasan, Weyl semimetals, Fermi arcs and chiral anomalies, *Nature Mater.* **15**, 1140 (2016).
- [14] N. P. Ong and S. Liang, Experimental signatures of the chiral anomaly in Dirac-Weyl semimetals, *Nature Rev. Phys.* **3**, 394 (2021).
- [15] S. Katayama, A. Kobayashi, and Y. Suzumura, Pressure-induced zero-gap semiconducting state in organic conductor  $\alpha$ -(BEDT-TTF)<sub>2</sub>I<sub>3</sub> salt, *Journal of the Physical Society of Japan* **75**, 054705 (2006).
- [16] M. O. Goerbig, J.-N. Fuchs, G. Montambaux, and F. Piéchon, Electric-field-induced lifting of the valley degeneracy in  $\alpha$ -(BEDT-TTF)<sub>2</sub>I<sub>3</sub> Dirac-like Landau levels, *EPL (Europhysics Letters)* **85**, 57005 (2009).
- [17] J. Sári, M. O. Goerbig, and C. Tóke, Magneto-optics of quasirelativistic electrons in graphene with an in-plane electric field and in tilted Dirac cones in  $\alpha$ -(BEDT TTF)<sub>2</sub>I<sub>3</sub>, *Phys. Rev. B* **92**, 035306 (2015).
- [18] A. A. Soluyanov, D. Gresch, Z. Wang, Q. Wu, M. Troyer, X. Dai, and B. A. Bernevig, Type-II Weyl semimetals, *Nature* **527**, 495 (2015).
- [19] S. Tchoumakov, M. Civelli, and M. O. Goerbig, Magnetic-field-induced relativistic properties in type-I and type-II Weyl semimetals, *Phys. Rev. Lett.* **117**, 086402 (2016).
- [20] L. Zhang, Y. Jiang, D. Smirnov, and Z. Jiang, Landau quantization in tilted Weyl semimetals with broken symmetry, *J. Appl. Phys.* **129**, 105107 (2021).
- [21] V. Lukose, R. Shankar, and G. Baskaran, Novel electric field effects on Landau levels in graphene, *Phys. Rev. Lett.* **98**, 116802 (2007).
- [22] A. G. Aronov and G. E. Pikus, Light absorption in semiconductors in crossed electric and magnetic fields, *Soviet Physics JETP* **24**, 339 (1967).
- [23] W. Zawadzki, S. Klahn, and U. Merkt, Semirelativistic behavior of electrons in InSb in crossed magnetic and electric fields, *Phys. Rev. Lett.* **55**, 983 (1985).
- [24] W. Zawadzki, S. Klahn, and U. Merkt, Inversion electrons on narrow-band-gap semiconductors in crossed electric and magnetic fields, *Phys. Rev. B* **33**, 6916 (1986).
- [25] Y. Shao, Z. Sun, Y. Wang, C. Xu, R. Sankar, A. J. Breindel, C. Cao, M. M. Fogler, A. J. Millis, F. Chou, Z. Li, T. Timusk, M. B. Maple, and D. N. Basov, Optical signatures of Dirac nodal lines in NbAs<sub>2</sub>, *PNAS* **116**, 1168

- (2019).
- [26] S. Takeuchi, Relativistic  $E \times B$  acceleration, *Phys. Rev. E* **66**, 037402 (2002).
  - [27] C. Xu, J. Chen, G.-X. Zhi, Y. Li, J. Dai, and C. Cao, Electronic structures of transition metal dipnictides  $XPn_2$  ( $X = \text{Ta, Nb}$ ;  $Pn = \text{P, As, Sb}$ ), *Phys. Rev. B* **93**, 195106 (2016).
  - [28] W. Bensch and W. Heid,  $\text{NbAs}_2$ , *Acta Crystallographica Section C* **51**, 2205 (1995).
  - [29] Y. Shao, A. N. Rudenko, J. Hu, Z. Sun, Y. Zhu, S. Moon, A. J. Millis, S. Yuan, A. I. Lichtenstein, D. Smirnov, *et al.*, Electronic correlations in nodal-line semimetals, *Nature Phys.* **16**, 636 (2020).
  - [30] L. M. Schoop, M. N. Ali, C. Straßer, A. Topp, A. Varykhalov, D. Marchenko, V. Duppel, S. S. P. Parkin, B. V. Lotsch, and C. R. Ast, Dirac cone protected by non-symmorphic symmetry and three-dimensional Dirac line node in  $\text{ZrSiS}$ , *Nature Comm.* **7**, 11696 (2016).
  - [31] D. Takane, Z. Wang, S. Souma, K. Nakayama, C. X. Trang, T. Sato, T. Takahashi, and Y. Ando, Dirac-node arc in the topological line-node semimetal  $\text{HfSiS}$ , *Phys. Rev. B* **94**, 121108 (2016).
  - [32] E. J. Mele, Dowsing for nodal lines in a topological semimetal, *PNAS* **116**, 1084 (2019).
  - [33] See Supplementary Materials.
  - [34] T. Timusk, J. P. Carbotte, C. C. Homes, D. N. Basov, and S. G. Sharapov, Three-dimensional Dirac fermions in quasicrystals as seen via optical conductivity, *Phys. Rev. B* **87**, 235121 (2013).
  - [35] M. Orlita, D. M. Basko, M. S. Zholudev, F. Teppe, W. Knap, V. I. Gavrilenko, N. N. Mikhailov, S. A. Dvoretiskii, P. Neugebauer, C. Faugeras, A.-L. Barra, G. Martinez, and M. Potemski, Observation of three-dimensional massless Kane fermions in a zinc-blende crystal, *Nature Phys.* **10**, 233 (2014).
  - [36] D. Neubauer, J. P. Carbotte, A. A. Nateprov, A. Löhle, M. Dressel, and A. V. Pronin, Interband optical conductivity of the [001]-oriented Dirac semimetal  $\text{Cd}_3\text{As}_2$ , *Phys. Rev. B* **93**, 121202 (2016).
  - [37] A. V. Pronin and M. Dressel, Nodal semimetals: A survey on optical conductivity, *physica status solidi (b)* **258**, 2000027 (2021).
  - [38] D. Santos-Cottin, M. Casula, L. d. Medici, F. L. Mardelé, J. Wyzula, M. Orlita, Y. Klein, A. Gauzzi, A. Akrap, and R. P. S. Lobo, Universal optical features of Dirac cone networks: The case of  $\text{BaNiS}_2$ , arXiv:2104.05521 (2021).
  - [39] B. Shen, X. Deng, G. Kotliar, and N. Ni, Fermi surface topology and negative longitudinal magnetoresistance observed in the semimetal  $\text{NbAs}_2$ , *Phys. Rev. B* **93**, 195119 (2016).
  - [40] Y.-Y. Wang, Q.-H. Yu, P.-J. Guo, K. Liu, and T.-L. Xia, Resistivity plateau and extremely large magnetoresistance in  $\text{NbAs}_2$  and  $\text{TaAs}_2$ , *Phys. Rev. B* **94**, 041103 (2016).
  - [41] G. Peramaiyan, R. Sankar, I. P. Muthuselvam, and W.-L. Lee, Anisotropic magnetotransport and extremely large magnetoresistance in  $\text{NbAs}_2$  single crystals, *Sci. Rep.* **8**, 1 (2018).
  - [42] G. Landwehr and E. I. Rashba, *Landau level spectroscopy* (Elsevier, 2012).
  - [43] W. Franz, Einfluß eines elektrischen Feldes auf eine optische Absorptionskante, *Zeitschrift für Naturforschung A* **13**, 484 (1958).

## Supplementary Information for "Lorentz-boost-driven magneto-optics in Dirac matter"

J. Wyzula,<sup>1</sup> X. Lu,<sup>2</sup> D. Santos-Cottin,<sup>3</sup> D. K. Mukherjee,<sup>2,4</sup> I. Mohelský,<sup>1</sup> F. Le Mardelé,<sup>3</sup> J. Novák,<sup>5</sup> M. Novak,<sup>6</sup> R. Sankar,<sup>7</sup> Y. Krupko,<sup>1,8</sup> B. A. Piot,<sup>1</sup> W.-L. Lee,<sup>7</sup> A. Akrap,<sup>3</sup> M. Potemski,<sup>1,9</sup> M. O. Goerbig,<sup>2</sup> and M. Orlita<sup>1,10,\*</sup>

<sup>1</sup>*LNCMI-EMFL, CNRS UPR3228, Univ. Grenoble Alpes, Univ. Toulouse,  
Univ. Toulouse 3, INSA-T, Grenoble and Toulouse, France*

<sup>2</sup>*Laboratoire de Physique des Solides, Université Paris Saclay,  
CNRS UMR 8502, F-91405 Orsay Cedex, France*

<sup>3</sup>*Department of Physics, University of Fribourg, Chemin du Musée 3, 1700 Fribourg, Switzerland*

<sup>4</sup>*Department of Physics, Indiana University, Bloomington, Indiana 47405, USA*

<sup>5</sup>*Department of Condensed Matter Physics, Masaryk University, Kotlářská 2, 611 37 Brno, Czech Republic*

<sup>6</sup>*Department of Physics, Faculty of Science, University of Zagreb, 10000 Zagreb, Croatia*

<sup>7</sup>*Institute of Physics, Academia Sinica, Nankang, 11529 Taipei, Taiwan*

<sup>8</sup>*Institut d'Electronique et des Systemes, UMR CNRS 5214,  
Université de Montpellier, 34000, Montpellier, France*

<sup>9</sup>*Institute of Experimental Physics, Faculty of Physics,  
University of Warsaw, ul. Pasteura 5, 02-093 Warszawa, Poland*

<sup>10</sup>*Charles University, Faculty of Mathematics and Physics,  
Institute of Physics, Ke Karlovu 5, 121 16 Prague 2, Czech Republic*

### CONTENTS

I. Sample growth and x-ray characterization	1
II. Optical response of NbAs <sub>2</sub> at $B = 0$	2
III. Nodal lines in magnetic field	5
A. Landau levels	5
B. Velocity operators	8
C. Selection rules	8
IV. Infrared magneto-reflectivity technique and complementary magneto-optical data	9
V. Magneto-transport experiments	9
References	10

### I. SAMPLE GROWTH AND X-RAY CHARACTERIZATION

NbAs<sub>2</sub> single crystals explored in this work were grown using a chemical vapor transport method. The as-grown crystals usually have a several facets with different crystallographic orientations with shiny surfaces suitable for infrared reflectivity experiments. In total, four NbAs<sub>2</sub> single crystal samples (No. 1...4) were used to collect the magneto-optical data presented in this work and characterized using standard  $x$ -ray technique. To this end, Bragg-Brentano diffractometer equipped with Cu  $x$ -ray tube, channel-cut germanium monochromator and scintillation detector were employed. The  $x$ -ray diffraction patterns of each explored sample and facet are presented in Fig. S1. For the purpose of optical experiments on the (001) facet at  $B = 0$ , the orientation  $a$  and  $b$  crystallographic axes was determined using pole diagrams. The magneto-reflectivity data with  $B$  oriented perpendicular to the (101) and (607) crystallographic planes were collected using the nearby (403) facet on Sample 1 rotated respectively by 5 and 7° in the ( $a$ - $c$ ) plane.

---

\* milan.ortita@lncmi.cnrs.fr






FIG. S1. Normalized diffraction patterns of all facets studied in our magneto-optical experiments. The (001)-oriented facet of Sample 1 was also used for reflectance measurements at  $B = 0$ .

## II. OPTICAL RESPONSE OF NBAS<sub>2</sub> AT $B = 0$

To characterize the optical response of NbAs<sub>2</sub> at  $B = 0$ , infrared reflectivity was measured on the (001)-oriented facet using radiation polarized linearly along the  $a$  and  $b$  crystallographic axes. To this end, the Vertex 70v FTIR spectrometer was used, equipped with custom-built in situ gold evaporation. At high photon energies, the phase was fixed by ellipsometry. Then, the standard Kramers-Kronig analysis was employed to obtain the frequency-dependent complex optical conductivity. The measured reflectivity and the deduced optical conductivity are presented in Fig. 1d in the main part of the manuscript and in Figs. S3a,c,e.

To describe the optical response of NbAs<sub>2</sub> at  $B = 0$  theoretically, we use the simple model for electronic states in a nodal-line semimetal proposed in the main text. The proposed Hamiltonian, now with both line index  $\xi = \pm 1$  and spin included, reads:

$$\hat{H} = (\varepsilon_0 + \hbar w q_{\text{line}}) \mathbb{1} + \hbar v (\xi q_x \sigma_x + q_y \sigma_y) + \xi \Delta \sigma_z \tau_z, \quad (\text{S1})$$

where  $\sigma_i$  ( $i = x, y, z$ ) and  $\tau_z$  are standard Pauli matrices for orbital and spin degrees of freedom, respectively. This Hamiltonian describes electronic states in the vicinity of any point at the nodal line,  $\mathbf{k} = \mathbf{k}_{\text{line}} + \mathbf{q}$ . The used orthogonal coordinate system,  $\mathbf{q} = (q_x, q_y, q_{\text{line}} \equiv q_z)$ , has the third component always aligned with the local direction of the nodal line  $\boldsymbol{\tau}$  (see Fig. S2) which is roughly parallel with the  $a$  crystallographic axis [1]. The other nodal line is located symmetrically in momentum space with respect to the  $\Gamma$ -Y-Z mirror plane. Similar to the local direction  $\boldsymbol{\tau}$ , the Hamiltonian parameters  $\varepsilon_0$ ,  $v$ ,  $\Delta$  and  $w$  also vary, smoothly and weakly along the nodal line, and therefore, can be viewed as a function of  $q_{\text{line}}$ .

To simplify the problem at hand, we have divided the nodal line zone into four segments I, II, III and IV, which approximately describe its propagation through a half of the Brillouin zone. The locations of these segments, as well as their conduction/valence band energies (at  $\mathbf{k}_{\text{line}}$ ), are shown in Figs. S3b,d,f. The suggested profile of the nodal lines approach the results of DFT simulations and overall conclusions presented by Shao et al. [1]. In each segment, the nodal line is reduced down to a straight line in momentum space – having the length of  $k^i$  – and characterized by a set of corresponding parameters:  $\varepsilon_0^i$ ,  $v^i$ ,  $\Delta^i$  and  $w^i$ , where  $i = \text{I, II, III and IV}$ , and two angles  $\theta_a^i$  and  $\theta_b^i$  which




FIG. S2. The position of two nodal lines in NbAs<sub>2</sub> with respect to the real space crystal axes  $a, b, c$  and the corresponding reciprocal space axes  $k_a, k_b, k_c$ . The local axes along the nodal line:  $\mathbf{q}_{\text{line}}$  is parallel to the nodal line and  $\mathbf{q}_{x,y}$  is always orthogonal to it.

encodes the local direction of the given segment  $\tau^i$  with respect to  $a$  and  $b$  crystallographic axes, cf. Figs. S3d,f. Within each segment, these parameters are supposed to be constant except  $\Delta^{\text{III}}$  which we consider to vary linearly with  $q_{\text{line}}$  to ensure the continuity of the nodal line in energy.

With such simplifications, the diagonal component of optical conductivity is obtained by integration along the chosen segment  $i$ , using the local basis of the Hamiltonian:

$$\Re[\sigma_{xx}^i(\omega)] = \frac{Ne^2}{8h} \int dq_{\text{line}} \left( 1 + \frac{4(\Delta^i)^2}{(\hbar\omega)^2} \right) [f(\epsilon_0^i + \hbar w^i q_{\text{line}} - \hbar\omega/2) - f(\epsilon_0^i + \hbar w^i q_{\text{line}} + \hbar\omega/2)] \Theta(\hbar\omega - 2\Delta^i) \quad (\text{S2})$$

and, by the isotropy of the model,  $\sigma_{xx}^i(\omega) = \sigma_{yy}^i(\omega)$ . The other diagonal component reads:

$$\begin{aligned} \Re[\sigma_{zz}^i(\omega)] = \\ \frac{Ne^2}{4h(v^i)^2} \int dq_{\text{line}} \left( \frac{\partial \Delta^i}{\partial q_{\text{line}}} \right)^2 \left[ 1 - \left( \frac{\Delta^i}{\hbar\omega} \right)^2 \right] [f(\epsilon_0^i + \hbar w^i q_{\text{line}} - \hbar\omega/2) - f(\epsilon_0^i + \hbar w^i q_{\text{line}} + \hbar\omega/2)] \Theta(\hbar\omega - 2\Delta^i). \end{aligned} \quad (\text{S3})$$

In the equations above,  $N = 2$  is the number of the nodal lines in the Brillouin zone (double degeneracy due to spin already included),  $f$  is the Fermi-Dirac distribution and  $\Theta$  is the Heaviside function. The main contribution to the optical conductivity comes from Eq. (S2). Eq. (S3) leads to a non-zero contribution to optical conductivity only when the gap parameter changes with  $q_{\text{line}}$ . Such a possibility is not explicitly included in the Hamiltonian (S1), nevertheless, as mentioned above, it is relevant in our approximation for the segment III (see Fig. S3b). Anyway, in practice, the contribution from (S3) is negligible due to the small ratio between  $\hbar^{-1} \partial_{q_{\text{line}}} \Delta^i$  and  $v^i$ .

In our reflectivity measurements at  $B = 0$  (Fig. 1d top in the main text), the incident radiation was polarized linearly along the  $a$  or  $b$  axes. So, to calculate the experimentally probed optical conductivities  $\Re[\sigma_{aa}(\omega)]$  and  $\Re[\sigma_{bb}(\omega)]$ , one has to make a corresponding projection for each segment ( $i = \text{I, II, III and IV}$ ) and sum their contributions:

$$\Re[\sigma_{aa}(\omega)] = \sum_i (\cos^2 \theta_a^i \Re[\sigma_{zz}^i(\omega)] + \sin^2 \theta_a^i \Re[\sigma_{xx}^i(\omega)]) \quad (\text{S4})$$

$$\Re[\sigma_{bb}(\omega)] = \sum_i (\cos^2 \theta_b^i \Re[\sigma_{zz}^i(\omega)] + \sin^2 \theta_b^i \Re[\sigma_{xx}^i(\omega)]) \quad (\text{S5})$$

Even though we use a relatively crude approximation for the profile of the nodal lines, there still remains a number of free parameters to be tuned/set. For each segment, we have its length  $k_i$  in momentum space, four Hamiltonian parameters  $v^i$ ,  $\Delta^i$ ,  $w^i$  and  $\epsilon_0^i$ , two angles  $\theta_a$  and  $\theta_b$ . The Fermi energy is an additional parameter common to all segments. To reduce this number, let us consider the following points: (i) When the anisotropy of the dispersion perpendicular to the nodal line is neglected (i.e.,  $v_x^i = v_y^i$ ), the main contribution to optical conductivity (S2) becomes independent of the velocity parameter  $v^i$ . (ii) For the parameters  $\Delta^i$  and  $w^i$  as well as the corresponding angles,  $\theta_a^i$  and  $\theta_b^i$ , for  $i = \text{II and IV}$ , we have solid estimates coming from the analysis of the magneto-optical response, due to




FIG. S3. Parts (a),(c) and (e): Real part of optical conductivity (solid lines) deduced via Kramers-Kronig analysis of the NbAs<sub>2</sub> reflectivity response measured on the (001)-oriented facet at  $T = 10$  K using radiation polarized linearly along the  $a$  and  $b$  crystallographic axes. The dashed and dotted lines show results of theoretical modelling described in the text. Parts (b), (d) and (f): Energy (at  $k_{\text{line}}$ ) and momentum profiles of the nodal lines considered in the simplified (segment-based) model for optical conductivity of NbAs<sub>2</sub> at  $B = 0$ .

the dispersive and flat parts of the nodal line, respectively: e.g.,  $\Delta^{\text{II}} = \Delta_D$ ,  $\Delta^{\text{IV}} = \Delta_F$ ,  $w^{\text{II}} = w$ ,  $w^{\text{IV}} = 0$ . (iii) Additional constraints on  $w^i$ ,  $\Delta^i$ ,  $\varepsilon_0^i$  and  $k^i$ , appear because the nodal lines are continuous in momentum as well as in energy. For instance, the size of the Brillouin zone approximately fixes the sum of segments lengths and the gap parameter  $\Delta^{\text{III}}$  has to evolve smoothly from  $\Delta^{\text{II}} = \Delta_F$  to  $\Delta^{\text{IV}} = \Delta_D$ . In the latter case, we choose the simplest linear in  $q_{\text{line}}$  approximation. (iv) The expected profile of the nodal line [1] allows us to neglect – due to Pauli occupation effect – the contribution of interband excitations coming from the segment I. This is because we are only interested in the optical response at relatively low photon energies (below  $\hbar\omega \approx 400$  meV, see Fig. S3a).

Using the constraints (i) to (iv), the number of freely tunable parameters is strongly reduced, basically down to the segments' lengths. The theoretically calculated conductivity approaches fairly well the experimentally measured curves, see Fig. S3a. The contributions of individual segments are shown using dotted lines and the corresponding color coding in Figs. S3c,e. Notably, the agreement could be further improved by considering velocity anisotropy ( $v_x^i \neq v_y^i$ ) which alters the relative strength of individual segments. Such an approach has been successfully used by Shao et al. [1], nevertheless, we prefer not introducing (three) additional tuning parameters which cannot be directly determined from our magneto-optical experiments.

The obtained values of the used tuning parameters – the lengths of the segments  $k^i$  and the angles  $\theta_a^{\text{III}}$  and  $\theta_b^{\text{III}}$  – are visualized graphically in Figs. S3d,f in which the nodal line location was projected to the  $(a-c)$  and  $(b-c)$

crystallographic planes, respectively. Let us remind that the vectors  $\mathbf{k}_b$  and  $\mathbf{k}_b \times \mathbf{k}_c$  lies in the  $a$ - $b$  plane while the vectors  $\mathbf{k}_c$ ,  $\mathbf{k}_a$  as well as  $\mathbf{k}_b \times \mathbf{k}_c$  lie in the  $a$ - $c$  crystallographic plane, see Fig. S2.

### III. NODAL LINES IN MAGNETIC FIELD

To model a nodal line subjected to an externally applied magnetic field, we use the (zero-field) Hamiltonian introduced, together with all its parameters, in the main text:

$$\hat{H} = (\varepsilon_0 + \hbar w q_{\text{line}}) \mathbb{1} + \begin{bmatrix} \Delta & \hbar v(q_x - iq_y) \\ \hbar v(q_x + iq_y) & -\Delta \end{bmatrix}, \quad (\text{S6})$$

where we consider only one spin sector (spin up) of one of the two nodal lines ( $\xi = +1$ ) [cf. Eq. (S1)]. The calculations for spin down or the other nodal line follow exactly the same procedure as we are about to show below.

#### A. Landau levels

The Hamiltonian (S6) is isotropic in the  $x$ - $y$  plane, hence, we may set, with no loss of generality, that  $\mathbf{B} = B(\cos \theta \mathbf{e}_{\text{line}} + \sin \theta \mathbf{e}_x) = B\mathbf{e}_B$  for  $0 < \theta < \pi/2$  and define a new orthogonal basis with the third component aligned with the magnetic field:

$$\{\mathbf{e}_{x'}, \mathbf{e}_{y'}, \mathbf{e}_{z'} \equiv \mathbf{e}_B\} = \{\cos \theta \mathbf{e}_x - \sin \theta \mathbf{e}_{\text{line}}, -\mathbf{e}_y, \cos \theta \mathbf{e}_{\text{line}} + \sin \theta \mathbf{e}_x\}. \quad (\text{S7})$$

When this new basis is used, the Hamiltonian reads:

$$H = \hbar w (q_B \cos \theta - q_{x'} \sin \theta) + \hbar v (q_{x'} \cos \theta + q_B \sin \theta) \sigma_x - \hbar v q_{y'} \sigma_y + \Delta \sigma_z, \quad (\text{S8})$$

and our goal is to find the spectrum in the presence of a magnetic field.

Let us first illustrate the relativistic properties embedded in the above Hamiltonian, which will inspire the solution of the problem. When the dispersion along the direction of the magnetic field is neglected ( $q_B = 0$ ), the Hamiltonian becomes that of a tilted 2D massive Dirac electron

$$H = \hbar w q_{x'} \sin \theta + \hbar v q_{x'} \cos \theta \sigma_x - \hbar v q_{y'} \sigma_y + \Delta \sigma_z, \quad (\text{S9})$$

where the tilt velocity along the  $x'$  direction is  $w \sin \theta$  and the Fermi velocity along the  $x'$  and  $y'$  directions are  $v \cos \theta$  and  $-v$ , respectively. However, this Hamiltonian is not that for the Lorentz-invariant Dirac equation but a variant of it due to the tilt term. The relativistic properties are revealed once the magnetic field is introduced using the standard Landau gauge,  $\mathbf{A} = -By' \mathbf{e}_{x'}$ , so that

$$H - \hbar w q_{x'} \sin \theta = -\hbar w \sin \theta e B y' + \hbar v \cos \theta (q_{x'} - e B y') \sigma_x - \hbar v q_{y'} \sigma_y + \Delta \sigma_z, \quad (\text{S10})$$

where one can identify an effective electric field  $Bw \sin \theta$  in the  $y'$  direction. Most importantly, the right hand side is now that of the standard Lorentz invariant Dirac equation except the anisotropy of the speed of light in the  $x'$  and  $y'$  directions. Therefore, one encounters now the problem of a relativistic 2D massive Dirac electron in the presence of an electric field of  $Bw \sin \theta$  in the  $y'$  direction and a magnetic field of  $B$  in the  $z'$  direction, thus moving with a drift velocity of  $w \sin \theta$  in the  $x'$  direction assuming a small drift velocity. Remark that the speed of light in the  $x'$  direction is replaced by  $v \cos \theta$ , so the rapidity is defined as  $\beta = w \tan \theta / v$ . It is well-known in the theory of special relativity that one can always shut down the electric field in the co-moving frame of an electron subjected to a crossed electric and magnetic field if the drift velocity is smaller than the speed of light, i.e.,  $\beta < 1$  [2, 3]. Working in the co-moving frame, with the drift velocity, and using a Lorentz boost, the problem is thus simplified to that of an electron subjected only to magnetic field, i.e., the usual Landau quantization problem. Once one finds the Landau levels (LL) in the co-moving frame, the spectrum in the lab frame, i.e., the original frame of reference, follows immediately by doing the inverse Lorentz boost from the co-moving frame back to the lab frame. Technically, this can be conveniently accomplished using hyperbolic transformation, which is the 2D representation of Lorentz boost in the Lorentz group. In our case, the hyperbolic transformation of the aforementioned Lorentz boost reads  $M = \exp(\phi \sigma_x / 2)$ , with the rapidity  $\beta = \tanh \phi = w \tan \theta / v$ .

Now we are ready to solve the problem described the Hamiltonian below

$$H = \hbar w [q_B \cos \theta - (q_{x'} - e B y') \sin \theta] + \hbar v [(q_{x'} - e B y') \cos \theta + q_B \sin \theta] \sigma_x - \hbar v q_{y'} \sigma_y + \Delta \sigma_z, \quad (\text{S11})$$






FIG. S4. Relative strength of inter-band transitions between different pairs of Landau levels (for  $n = 0 \dots 6$ ) plotted for angles  $\theta_D = 0, 20, 40, 60^\circ$ , and parameters ( $w$ ,  $v_D$  and  $\Delta_D$ ) deduced experimentally for the dispersive part of the nodal line in NbAs<sub>2</sub>. For  $\theta_D = 0^\circ$ , we obtain standard selection rules  $n \rightarrow n \pm 1$  for electric-dipole transitions, typical of all isotropic systems. With increasing  $\theta_D$ , gradually, additional transitions become allowed ( $n \rightarrow n \pm 2, \pm 3 \dots$ ). For large angles  $\theta_D$ , but still for  $\beta < 1$ , one finds a plethora of optically active transitions. The dominant ones follow the rule-of-thumb selection rules  $n \rightarrow \alpha n$  and  $n \rightarrow n/\alpha$ , where  $\alpha$  is an integer ( $\alpha = 2 - 4$  for  $\theta_D = 60^\circ$ ).

where the magnetic field is already incorporated by the above Landau gauge. Using the same hyperbolic transformation,  $M = \exp(\phi \sigma_x / 2)$ , with the rapidity  $\beta = \tanh \phi = w \tan \theta / v$ , we obtain the Hamiltonian

$$\begin{aligned} H_T &= M H M \\ &= \hbar \frac{wv}{v^*} q_B + \hbar \frac{w^2 + v^2}{v^*} q_B \sin \theta \cos \theta \sigma_x + \hbar v^* (q_{x'} - eB y') \sigma_x - \hbar v q_{y'} \sigma_y + \Delta \sigma_z \end{aligned} \quad (\text{S12})$$

where the velocity  $v^*$  is defined as

$$v^{*2} = v^2 \cos^2 \theta - w^2 \sin^2 \theta \quad \text{with} \quad \gamma = \frac{1}{\sqrt{1 - \beta^2}} = \cosh \phi = \frac{v \cos \theta}{v^*} \quad (\text{S13})$$

where  $\gamma > 1$  is the relativistic Lorentz factor.

Note that  $H_T$  does not have the same spectrum as  $H$  for the reason that we are now working in the co-moving frame. In the basis of  $|\psi_T\rangle = \gamma^{-1/2} M^{-1} |\psi\rangle$  given the eigenstate  $|\psi\rangle$  of  $H$ , one can construct from  $H_T$  another Hamiltonian  $H_E = (H_T - E \sinh \phi \sigma_x) / \cosh \phi$  with the same spectrum as  $H$

$$H_E = \frac{\hbar w q_B}{\cos \theta} + \frac{1}{\gamma} \begin{bmatrix} \Delta & \hbar \frac{\sqrt{2vv^*}}{\ell_B} a_E^\dagger \\ \hbar \frac{\sqrt{2vv^*}}{\ell_B} a_E & -\Delta \end{bmatrix} \quad (\text{S14})$$

where  $\ell_B = \sqrt{\hbar / (eB)}$  is the magnetic length and a pair of energy-dependent ladder operators is defined as:

$$\begin{aligned} a_E &= -\frac{1}{\sqrt{2vv^*}} \left( \frac{v^*}{\ell_B} (y' - \langle y' \rangle_E) + i \ell_B v q_{y'} \right) \\ a_E^\dagger &= -\frac{1}{\sqrt{2vv^*}} \left( \frac{v^*}{\ell_B} (y' - \langle y' \rangle_E) - i \ell_B v q_{y'} \right) \\ \langle y' \rangle_E &= \frac{\ell_B^2}{v^*} \left( v^* q_{x'} + q_B \frac{w^2 + v^2}{v^*} \sin \theta \cos \theta - E \frac{w \sin \theta}{\hbar v^*} \right) \end{aligned} \quad (\text{S15})$$

with  $[a_E, a_E^\dagger] = 1$ . The subscript  $E$  indicates the dependence on energy  $E$ . In particular, the center of cyclotron  $\langle y' \rangle_E$  shifts with energy.

Since the energy-dependent term in  $H_E$  is absorbed in the definition of the ladder operators, the energy spectrum is self-consistently found in the eigenstates of energy  $E_n^\lambda$  in the form

$$|\psi_{T,n,\lambda}\rangle = \begin{pmatrix} \cos \zeta_{n,\lambda} |n, E_n^\lambda\rangle \\ \sin \zeta_{n,\lambda} |n-1, E_n^\lambda\rangle \end{pmatrix} \quad (\text{S16})$$




FIG. S5. Parts (a-f): Relative magneto-reflectivity of NbAs<sub>2</sub>,  $R_B/R_0$ , in the magnetic field applied perpendicular to six different crystallographic planes:  $(40\bar{1})$ ,  $(100)$ ,  $(10\bar{1})$ ,  $(403)$ ,  $(607)$  and  $(20\bar{3})$ , which make angles 12, 29, 39, 58, 65 and 56° with the  $a$  crystallographic axis, respectively. Horizontal gray bars correspond to positions of two steps in the onset of interband excitations at  $2\Delta_D$  and  $2\Delta_F$  at  $B = 0$ , cf. Fig. 1d in the main text or Fig. S3. The yellow values indicate the apparent band gap,  $2\Delta_D^{\text{eff}}$ , deduced using a linear zero-field extrapolation of inter-LL resonance belonging to the lower set (yellow dashed lines).

where  $\zeta_{n,\lambda}$  is an angle depending on  $n$  and the sign of energy  $\lambda = \pm$ ,  $|n', E_n^\lambda\rangle$  is the wave function of the one-dimensional quantum harmonic oscillator defined by the previous ladder operators. Given the Landau level index  $n$ ,  $n' = n$  or  $n - 1$ .

Finally, we obtain the LL spectrum:

$$\begin{aligned} E_n^\pm &= \hbar w q_B / \cos \theta \pm \sqrt{(\Delta/\gamma)^2 + 2n\hbar e B v^2 \cos \theta / \gamma^3} & \text{for } n > 0 \\ E_0 &= \hbar w q_B / \cos \theta + \frac{\Delta}{\gamma} & \text{for } n = 0, \end{aligned} \quad (\text{S17})$$

which has the form typical of massive Dirac electrons, nevertheless, with the gap and velocity renormalized by the pseudo-relativistic Lorentz factor:  $2\Delta \rightarrow 2\Delta/\gamma$  and  $v^2 \rightarrow v^2/\gamma^3$ . Importantly, for the other line index  $\xi = -1$  or the spin down we obtain an analogous LL spectrum replacing  $\Delta$  by  $-\Delta$ .

It is worth noting that the discussed Lorentz-boost-driven renormalization of the apparent band gap may, to certain extent, resemble the well-known Franz-Keldysh effect [4, 5] which finds its practical use in electro-optical modulation, see e.g. Ref. [6]. Even a closer analogy appears when the Franz-Keldysh effect is explored in crossed electric and magnetic fields [7]. Nevertheless, in our case, there is no real electric field applied to the explored system. It only appears in an effective way, in the pseudo-relativistic Dirac-type Hamiltonian, being proportional to the magnetic-field component perpendicular to  $\tau_D$ .

## B. Velocity operators

To study the magneto-optical properties of NbAs<sub>2</sub> theoretically, one has to evaluate the matrix element of the corresponding velocity operators:  $\langle \psi_n | \hat{v}_{\mathbf{k}} | \psi_m \rangle$ . In this case, it is more practical to work with the basis  $|\psi_{T,n}\rangle$  already in hands by remarking that

$$\langle \psi_n | \nabla_{\mathbf{k}} H | \psi_{n'} \rangle = \gamma \langle \psi_{T,n} | \nabla_{\mathbf{k}} H_T | \psi_{T,n'} \rangle \quad (\text{S18})$$

thanks to the fact that  $M$  is  $k$ -independent. So, in the basis  $|\psi_T\rangle$ , the velocity operators for  $H_T$  are

$$\hat{v}_{T,x} = v^* \sigma_x \quad (\text{S19})$$

$$\hat{v}_{T,y} = -v \sigma_y \quad (\text{S20})$$

$$\hat{v}_{T,z} = \frac{wv}{v^*} + \frac{w^2 + v^2}{v^*} \sin \theta \cos \theta \quad (\text{S21})$$

where one notices an emergent anisotropy of the velocity parameter induced by the applied magnetic field.

## C. Selection rules

With the velocity operators, one can *a priori* derive the selection rules for electric-dipole inter-LL transitions which are active in the Faraday configuration (the configuration with the wave vector of light parallel to  $B$ ). We proceed in a way analogous Landau-quantized tilted 3D cones [3]. When calculating the matrix elements such as  $\langle \psi_{T,n} | \hat{v}_{T,x} | \psi_{T,m} \rangle$ , one has to deal with terms such as  $\langle n', E_n | m', E_m \rangle$  which are no longer 0 or 1. This is due to the mismatch of their energy-dependent orbital center [see Eq. (S15)]. Therefore, we do not obtain, in general, the selection rules,  $n \rightarrow n \pm 1$ , typical of isotropic solids [8]. Instead, all direct transitions become in principle possible, as long as the occupation of states (Pauli principle) allows. The transitions stemming from the selection rule other than  $n \rightarrow n \pm 1$  proliferates in the optical conductivity. By the sum rule, this would dilute the prominent Landau fan which reflects the  $n \rightarrow n \pm 1$  rule. Therefore, the usual clear-cut Landau fan is blurred.

To illustrate the evolution of electric-dipole selection rules with the angle  $\theta_D$  between the magnetic field and the local nodal-line direction of the dispersive part  $\tau$ , we proceed in the way analogous to [3] and [9]. In Fig. S4, we plot normalized squares of matrix elements, such as  $|\langle \psi_{T,n} | \hat{v}_{T,x} / v^* | \psi_{T,m} \rangle|^2$ , for different interband inter-LL excitations (for indices  $n = 0 \dots 6$ ) and parameters deduced for the dispersive part of the nodal line in NbAs<sub>2</sub>. As expected, one obtains the standard selection rules in isotropic systems for  $\theta_D = 0^\circ$ :  $n \rightarrow n \pm 1$ . This is because the energy-dependence in the cyclotron center is canceled by  $\sin \theta$  [see in Eq. (S15)]. With increasing  $\theta_D$ , additional inter-LL excitations emerge and their relative strength gradually increases. At relatively large angles  $\theta_D$ , the selection rules differ significantly and transitions,  $n \rightarrow n \pm 1$ , originally dominating the response, effectively disappear, except the transition  $0 \rightarrow 1$  that stays well-defined and relatively strong (as long as  $\beta < 1$ ).

#### IV. INFRARED MAGNETO-REFLECTIVITY TECHNIQUE AND COMPLEMENTARY MAGNETO-OPTICAL DATA

The magneto-reflectivity of NbAs<sub>2</sub> was explored in the Faraday configuration, with  $\mathbf{B}$  applied perpendicular to the chosen crystallographic plane. During experiments, a macroscopic area of the sample (typically a few mm<sup>2</sup>), placed in a superconducting coil and kept at  $T = 4.2$  K in the helium exchange gas, was exposed to radiation of a globar, which was analyzed by the Vertex 80v Fourier-transform spectrometer and delivered to the sample via light-pipe optics. The reflected light was detected by a liquid-helium-cooled bolometer placed outside the magnet. The reflectivity  $R_B$  recorded at a given magnetic field  $B$  was normalized by  $R_{B=0}$ . To facilitate the data analysis, we assumed that the maxima in relative magneto-reflectivity,  $R_B/R_0$ , directly correspond to the positions of inter-LL resonances. This is justified when the imaginary part of the dielectric function exceeds the absolute value of the real part – a condition fulfilled at photon energies around and slightly above the plasma edge (Fig. 1d). A more detailed analysis indicates that, in this way, we slightly overestimate/underestimate the positions of resonances at lower/higher part of the explored range.

In total, we have explored the magneto-optical response of NbAs<sub>2</sub> with  $B$  applied perpendicular to ten different crystallographic planes. The results for (101), (201), (20 $\bar{1}$ ) and (001) planes are presented in the main part of the manuscript (Fig. 2), the rest of our data has been plotted, again in a form of false-color plots of relative magneto-reflectivity,  $R_B/R_0$ , in Fig. S5. Notably, the explored NbAs<sub>2</sub> monocrystals did not have sufficiently large (101) and (607) facets that would allow us to collect magneto-optical data with a sufficiently high quality. Therefore, we used a larger-in-size (403) facet on Sample 1 rotated respectively by 5° and 7°.

To analyze the magneto-reflectivity data quantitatively, we used the procedure described in the main text. We have focused on the lowest observed line in both sets and assign it to the inter-LL excitation  $0 \leftrightarrow 1$  and extracted the effective value of the gap  $2\Delta^{\text{eff}}$  and velocity  $v^{\text{eff}}$  using the formula:

$$\hbar\omega_{0\leftrightarrow 1} = \Delta^{\text{eff}} + \sqrt{(\Delta^{\text{eff}})^2 + 2e\hbar B(v^{\text{eff}})^2}. \quad (\text{S22})$$

The extracted values of  $2\Delta^{\text{eff}}$  and velocity  $v^{\text{eff}}$  were then plotted for flat and dispersive parts/crossings in Figs. 3a,b and c,d, respectively, in the main part of the manuscript. Notably, for certain facets, inter-LL transitions belonging only to one of two sets are observed. For the upper set, originating in the flat part of the nodal line, this happens for angles  $\theta_F$  approaching  $\pi/2$  and nearly vanishing perpendicular field component  $B \cos \theta_F$ . This is the case, for instance of the (607) crystallographic plane (Fig. S5e) with  $\theta_F \approx 90^\circ$ . In contrast, the lower set of inter-LL excitations – stemming from the dispersive part of the nodal line – vanishes at significantly lower angles  $\theta_D$ , due to Lorentz-boost-driven collapse of the LL spectrum. For instance, the lower set of excitations of completely missing in the response measured with  $B$  applied perpendicular to the (203 $\bar{1}$ ) plane, see Fig. S5f, for which  $\theta_D \approx 64^\circ$ .

#### V. MAGNETO-TRANSPORT EXPERIMENTS

Magneto-transport experiments provided us with another, complementary characterization of the explored NbAs<sub>2</sub> crystals. To this end, we have chosen a monocrystal from the same batch as Sample 1. The electrical contacts were deposited using silver paint in the corners of the (001)-oriented facet of a rectangular shape. The current was always applied along  $b$  axis of the sample and the longitudinal magneto-resistance  $R_{xx}$  measured at selected (low) temperatures. Two sets of experiments were performed, with the magnetic field applied perpendicular to the  $a$ - $b$  plane and along the  $a$  axis, see Figs. S6a and b, respectively. In both cases, a pronounced magneto-resistance  $R_{xx}(B)$  was observed [10]. Shubnikov-de Haas oscillations became visible in the background removed data  $\Delta R_{xx}$ . The corresponding fast Fourier-transform applied to  $\Delta R_{xx}(1/B)$  implies several characteristic oscillation frequencies and harmonics, see Figs. S6c and d. While the frequency  $F_a \approx 300$  T seems to be rather independent of the magnetic field direction, other frequencies,  $F_b$  and  $F_c$ , indicate a large degree of anisotropy and correspondingly non-spherical Fermi surfaces. The damping of oscillations with temperature has been used to get estimates of the effective (cyclotron) masses  $m^*$ . The damping of the most prominent frequencies, i.e.,  $F_a = 276$  T and  $F_c = 120$  T for the magnetic field applied perpendicular to the (001) plane and  $F_a = 280$  T with  $B$  parallel to the  $a$  axis, was analyzed using Lifshitz-Kosevich formula:

$$\Delta_0/\Delta(T_0) = \frac{\alpha T m^* / \bar{B}}{\sinh(\alpha T m^* / \bar{B})}, \quad (\text{S23})$$

where  $\bar{B}$  stands for the mean magnetic field,  $T_0$  is the lowest measured temperature and  $\alpha = 2\pi^2 k_B m_0 / (e\hbar) = 14.69$  T/K. The fitted effective masses are similar for all three frequencies  $m^* = (0.25 \pm 0.05)m_0$  where  $m_0$  is the bare electron mass. This result is in agreement with values reported in the literature [10–13].






FIG. S6. Parts (a) and (b): Longitudinal magneto-resistance,  $R_{xx}$  and  $\Delta R_{xx}(B)$ , measured at indicated temperatures. The corresponding Fourier-transform applied  $\Delta R_{xx}(1/B)$  are plotted in parts (c) and (d): together with insets showing the directions of the magnetic field and current. Parts (e) and (f): Sh-dH oscillation frequencies measured at  $T = 1.5$  K as a function of the angle  $\theta$  between the magnetic field and the  $b$  and  $a$  axes, respectively, see insets.

To explore this (an)isotropy in greater detail, we followed the longitudinal magneto-resistance of NbAs<sub>2</sub> as a function of the magnetic-field direction with respect to the crystal, while keeping the current flowing along the  $b$  axis. In Figs. S6e,f, the magnetic field was applied perpendicular to the  $a$  and  $b$  axes and the sample rotated around the  $a$  and  $b$  crystallographic axes, respectively. In the latter case, the frequencies  $F_b$  and  $F_c$  exhibit a fairly pronounced angle dependence, with the maximum appearing roughly at  $\theta_{B,a} \approx 90^\circ$  (i.e.,  $B \perp a$ ) and with the maximum-to-minimum ratio reaching nearly 10. This indicates strongly elongated, cigar-like, Fermi surfaces oriented approximately along the  $a$  axis. In contrast, the frequencies  $F_b$  and  $F_c$  remain nearly constant when rotating the sample around the  $a$  axis. Such behavior agrees with conclusions of the preceding magneto-transport studies [10–13], and importantly, also with the presence of the nodal lines propagating roughly parallel to the  $a$  crystallographic axis.

- 
- [1] Y. Shao, Z. Sun, Y. Wang, C. Xu, R. Sankar, A. J. Breindel, C. Cao, M. M. Fogler, A. J. Millis, F. Chou, Z. Li, T. Timusk, M. B. Maple, and D. N. Basov, Optical signatures of Dirac nodal lines in NbAs<sub>2</sub>, *PNAS* **116**, 1168 (2019).
  - [2] S. Tchoumakov, M. Civelli, and M. O. Goerbig, Magnetic-field-induced relativistic properties in type-I and type-II Weyl semimetals, *Phys. Rev. Lett.* **117**, 086402 (2016).
  - [3] J. Sári, M. O. Goerbig, and C. Tóke, Magneto-optics of quasirelativistic electrons in graphene with an inplane electric field and in tilted Dirac cones in  $\alpha - (\text{BEDT TTF})_2\text{I}_3$ , *Phys. Rev. B* **92**, 035306 (2015).
  - [4] W. Franz, Einfluß eines elektrischen Feldes auf eine optische Absorptionskante, *Zeitschrift für Naturforschung A* **13**, 484 (1958).
  - [5] L. V. Keldysh, Behavior of non-metallic crystals in strong electric fields, *Soviet Journal of Experimental and Theoretical Physics* **6**, 763 (1958).

- [6] K. Liu, C. R. Ye, S. Khan, and V. J. Sorger, Review and perspective on ultrafast wavelength-size electro-optic modulators, *Laser & Photonics Reviews* **9**, 172 (2015).
- [7] A. G. Aronov and G. E. Pikus, Light absorption in semiconductors in crossed electric and magnetic fields, *Soviet Physics JETP* **24**, 339 (1967).
- [8] G. Landwehr and E. I. Rashba, *Landau level spectroscopy* (Elsevier, 2012).
- [9] The technical trick to analytically evaluate the matrix element is described in the Supplementary information of [2] or in [14].
- [10] Z. Yuan, H. Lu, Y. Liu, J. Wang, and S. Jia, Large magnetoresistance in compensated semimetals TaAs<sub>2</sub> and NbAs<sub>2</sub>, *Phys. Rev. B* **93**, 184405 (2016).
- [11] B. Shen, X. Deng, G. Kotliar, and N. Ni, Fermi surface topology and negative longitudinal magnetoresistance observed in the semimetal NbAs<sub>2</sub>, *Phys. Rev. B* **93**, 195119 (2016).
- [12] Y.-Y. Wang, Q.-H. Yu, P.-J. Guo, K. Liu, and T.-L. Xia, Resistivity plateau and extremely large magnetoresistance in NbAs<sub>2</sub> and TaAs<sub>2</sub>, *Phys. Rev. B* **94**, 041103 (2016).
- [13] G. Peramaiyan, R. Sankar, I. P. Muthuselvam, and W.-L. Lee, Anisotropic magnetotransport and extremely large magnetoresistance in NbAs<sub>2</sub> single crystals, *Sci. Rep.* **8**, 1 (2018).
- [14] M. O. Scully, M. S. Zubairy, *et al.*, Quantum optics cambridge university press, Cambridge, CB2 2RU, UK (1997).