

# **$M|G|\infty$ QUEUE BUSY PERIOD WITH PME DISTRIBUTION<sup>1</sup>**

Prof. Dr. **MANUEL ALBERTO M. FERREIRA**  
Instituto Universitário de Lisboa (ISCTE – IUL), BRU - IUL, Lisboa, Portugal  
[manuel.ferreira@iscte.pt](mailto:manuel.ferreira@iscte.pt)

## **ABSTRACT**

In this article it is shown that if the busy period of a  $M|G|\infty$  queue system is PME distributed, the service time is a random variable with a long-tail distribution. The result is obtained through Laplace transforms analysis.

**Keywords:**  $M|G|\infty$ , busy period, PME distribution, long-tail distribution, Laplace transform.

## **1. INTRODUCTION**

In a  $M|G|\infty$  queue system,  $\lambda$  is the Poisson process arrivals rate,  $\alpha$  is the mean service time,  $G(\cdot)$  is the service time distribution function and so  $\alpha = \int_0^\infty [1 - G(t)] dt$ . The traffic intensity is  $\rho = \lambda\alpha$  and  $B$  is the busy period length.

Note the importance of the busy period study, for this queuing system, because in it any customer, when it arrives, finds immediately an available server. So the problem is “for how long the servers – and how many servers? – must be available? That is: how long is the busy period length?”

It will be supposed that the  $B$  probability density function is

$$g_r(t) = \int_{\frac{r-1}{r}}^\infty f_r(y) y^{-1} e^{-\frac{t}{y}} dy, r > 1 \quad (1.1)$$

where  $f_r(y) = r \left(\frac{r-1}{r}\right)^r x^{-(r+1)}$ ,  $x \geq \frac{r-1}{r}$ , is a Pareto distribution probability density function. It is a PME<sup>2</sup>, see (1), and it is long-tail type. The  $g_r$  moments are

$$m_n = n! \frac{r}{r-n} \left[\frac{r-1}{r}\right]^n, n = 1, 2, \dots \quad (1.2).$$

---

<sup>1</sup> This work was financially supported by FCT through the Strategic Project PEst-OE/EGE/UI0315/2011.

<sup>2</sup> Pareto Mixture of Exponentials.

Through Laplace transform analysis it will be emphasized that in these circumstances the service time is a random variable with a long-tail distribution.

## 2. THE PME LAPLACE TRANSFORM

Calling  $\hat{g}_r(s)$  the Laplace transform of a PME with parameter  $r$ , see again (1),

$$\hat{g}_r(s) = r \left( \frac{r-1}{r} \right)^r \int_0^{\frac{r}{r-1}} \frac{x^r}{s+x} dx, r > 1 \quad (2.1).$$

$$\begin{aligned} \text{But } \hat{g}_r(s) &= r \left( \frac{r-1}{r} \right)^r \left( \int_0^{\frac{r}{r-1}} \left( x^{r-1} - \frac{sx^{r-1}}{s+x} \right) dx \right) = r \left( \frac{r-1}{r} \right)^r \left( \int_0^{\frac{r}{r-1}} x^{r-1} dx - \right. \\ & s \int_0^{\frac{r}{r-1}} \frac{x^{r-1}}{s+x} dx \left. \right) = r \left( \frac{r-1}{r} \right)^r \left( \left[ \frac{x^r}{r} \right]_0^{\frac{r}{r-1}} - s \int_0^{\frac{r}{r-1}} \frac{x^{r-1}}{s+x} dx \right) = r \left( \frac{r-1}{r} \right)^r \left( \frac{1}{r} \left( \frac{r}{r-1} \right)^r - \right. \\ & \left. s \int_0^{\frac{r}{r-1}} \frac{x^{r-1}}{s+x} dx \right) = 1 - sr \left( \frac{r-1}{r} \right)^r \int_0^{\frac{r}{r-1}} \frac{x^{r-1}}{s+x} dx. \end{aligned}$$

If  $\hat{h}_r(s)$  is the PME with parameter  $r$  tail Laplace transform, as  $\hat{h}_r(s) = \frac{1}{s} - \frac{1}{s} \hat{g}_r(s)$ ,

$$\hat{h}_r(s) = r \left( \frac{r-1}{r} \right)^r \int_0^{\frac{r}{r-1}} \frac{x^{r-1}}{s+x} dx, r > 1 \quad (2.2).$$

So, after (2.2),

$$\hat{h}_r^{(n)}(s) = (-1)^n n! r \left( \frac{r-1}{r} \right)^r \int_0^{\frac{r}{r-1}} \frac{x^{r-1}}{(s+x)^{n+1}} dx, n = 0, 1, 2, \dots \quad (2.3)$$

where  $\hat{h}_r^{(n)}$  is the  $n^{\text{th}}$  order derivative of  $\hat{h}_r$ .

$$\begin{aligned} \text{Then } \hat{h}_r^{(n)}(0) &= (-1)^n n! r \left( \frac{r-1}{r} \right)^r \int_0^{\frac{r}{r-1}} x^{r-n-2} dx, n = \\ 0, 1, 2, \dots \quad \text{But } \int_0^{\frac{r}{r-1}} x^{r-n-2} dx &= \begin{cases} \left[ \frac{x^{r-n-1}}{r-n-1} \right]_0^{\frac{r}{r-1}}, n \neq r-1 \\ \left[ \log|x| \right]_0^{\frac{r}{r-1}}, n = r-1 \end{cases} = \begin{cases} \left( \frac{r}{r-1} \right)^{r-(n+1)}, r > n+1 \\ -\infty, r \leq n+1 \end{cases} \end{aligned}$$

$$\text{So } \hat{h}_r^{(n)}(0) = \begin{cases} (-1)^n n! r \left( \frac{r-1}{r} \right)^r \frac{\left( \frac{r}{r-1} \right)^{r-(n+1)}}{r-(n+1)}, r > n+1 \text{ or, equivalently,} \\ (-1)^n (-\infty), 1 < r \leq n+1 \end{cases}$$

$$\hat{h}_r^{(n)}(0) = \begin{cases} (-1)^n n! r \left(\frac{r-1}{r}\right)^r \frac{\left(\frac{r}{r-1}\right)^{r-(n+1)}}{r-(n+1)}, & n < r+1, r > 1 \\ (-1)^n (-\infty), & n \geq r-1 \end{cases} \quad (2.4).$$

### 3. $M|G|\infty$ BUSY PERIOD TAIL LAPLACE TRANSFORM

Call  $U(t)$  the  $M|G|\infty$  busy period tail and  $u(s)$  the respective Laplace transform so, see (2),

$$\frac{d}{dt} \left( \frac{1 - e^{-\lambda \int_0^t [1-G(v)] dv}}{1 - e^{-\rho}} \right) = TL^{-1} \left( \frac{1}{1 - e^{-\rho}} \frac{\lambda u(s)}{\lambda u(s) + 1} \right) \quad (3.1)$$

and

$$\frac{e^{-\lambda \int_0^t [1-G(v)] dv} \lambda (1 - G(t))}{1 - e^{-\rho}} = TL^{-1} \left( \frac{1}{1 - e^{-\rho}} \frac{\lambda u(s)}{\lambda u(s) + 1} \right) \quad (3.2)$$

being  $TL^{-1}(\cdot)$  the inverse Laplace transform.

$$\begin{aligned} \text{So,} \quad & \int_0^\infty t^n \frac{e^{-\lambda \int_0^t [1-G(v)] dv} \lambda (1-G(t))}{1 - e^{-\rho}} dt = \\ (-1)^n \frac{1}{1 - e^{-\rho}} \left( \frac{\lambda u(s)}{\lambda u(s) + 1} \right)_{s=0}^{(n)} & \Leftrightarrow \int_0^\infty t^n e^{-\lambda \int_0^t [1-G(v)] dv} \lambda (1-G(t)) dt = \\ (-1)^n \left( \frac{\lambda u(s)}{\lambda u(s) + 1} \right)_{s=0}^{(n)}. & \end{aligned}$$

As  $e^{-\lambda \int_0^t [1-G(v)] dv} \leq 1$  the consequence is that  $\int_0^\infty t^n \lambda (1-G(t)) dt \geq (-1)^n \left( \frac{\lambda u(s)}{\lambda u(s) + 1} \right)_{s=0}^{(n)} \Leftrightarrow \int_0^\infty \frac{1-G(t)}{\alpha} dt \geq \frac{(-1)^n}{\rho} \left( \frac{\lambda u(s)}{\lambda u(s) + 1} \right)_{s=0}^{(n)}$  this entire happening if  $\frac{1}{1 - e^{-\rho}} \frac{\lambda u(s)}{\lambda u(s) + 1}$  is in fact a probability density function Laplace transform, being enough so that this happens that  $(-1)^n \left( \frac{\lambda u(s)}{\lambda u(s) + 1} \right)_{s=0}^{(n)} \geq 0, n = 0, 1, 2, \dots$ .

### 4. $M|G|\infty$ BUSY PERIOD WITH PME DISTRIBUTION

Note that in the  $\frac{\lambda u(s)}{\lambda u(s)+1}$   $n^{\text{th}}$  order derivative,  $u^{(n)}(s)$  always appears with a positive sign in the numerator and, for  $s = 0$ , if  $u(s)$  is given by (2.1), in that order  $n$  derivative the denominator is  $(\lambda + 1)^{n+1}$ . It is enough to take in account the quotient derivative expression and that  $\hat{h}_r(0) = 1$ .

So, after (2.4), it is concluded that

-If there is a service distribution such that the  $M|G|\infty$  queue busy period is distributed as a PME distribution with parameter  $r$ , the service equilibrium distribution moments of order greater than  $r - 1$ , centered in the origin, are infinite.

Note that

-The service equilibrium distribution, with these moments, is a long-tail distribution, see again (1),

-As  $\frac{u(s)}{\frac{\rho-1}{\lambda}}$  is the  $M|G|\infty$  queue busy period equilibrium distribution

Laplace transform it is also concluded that if this has moments of order greater than  $n, n \in \mathbb{N}$ , infinite, the same happens with the service equilibrium distribution, that is: they are both long –tail distributions.

## 5. CONCLUDING REMARKS

As it is stated in (1) the PME are long-tail distributions. So it is checked in this work that there is an uncontested association between long –tail service distributions and long-tail busy period distributions for the  $M|G|\infty$  queue, as it was shown in (3).

The PME were introduced in (1). There they were a tool to investigate properties of waiting times tail probabilities in queues with long-tail service-time distributions. For this investigation the authors developed algorithms for computing the waiting time distribution by Laplace transform inversion when the Laplace transforms of the inter-arrival time and service time distributions are known. The procedure here followed is similar.

## REFERENCES

1. J. Abate, G. L. Choudhury, W. Whitt. Waiting-time tail probabilities in queues with long-tail service-time distributions. *Queueing systems*, 16, 311-338, 1994.
2. M. A. M. Ferreira, M. Andrade.  $M|G|\infty$  queue busy period tail. *Journal of Mathematics and Technology*, 1(3), 11-16, 2010.
3. M. A. M. Ferreira, M. Andrade. Infinite servers queue systems busy period length moments. *Journal of Mathematics and Technology*, 3(2), 5-9, 2012.
4. B. Murteira. *Probabilidades e estatística*. Vol. I. Editora McGraw-Hill de Portugal, Lda. Lisboa, 1979.
5. F. P. Kelly. *Reversibility and stochastic networks*. New York, John Wiley and Sons, 1979.
6. J. Figueira, M. A. M. Ferreira. Representation of a pensions fund by a stochastic network with two nodes: an exercise. *Portuguese Revue of Financial Markets*, 1(3), 1999.
7. J. C. Hershey, E. N. Weiss, A. C. Morris. A stochastic service network model with application to hospital facilities. *Operations Research*, 29(1), 1-22, 1981.
8. L. Kleinrock. *Queueing systems*. Vol. I and Vol. II. Wiley- New York, 1985.
9. L. Tackács. *An introduction to queueing theory*. Oxford University Press, New York, 1962.
10. M. Andrade. A note on foundations of probability. *Journal of Mathematics and Technology*, 1(1), 96-98, 2010.
11. M. J. Carrillo. Extensions of Palm's theorem: a review. *Management Science*, 37(6), 739-744, 1991.
12. M. A. M. Ferreira. Simulação de variáveis aleatórias-método de monte carlo. *Revista Portuguesa de Gestão*, I (III/IV), 119-127. ISCTE, 1994.
13. M. A. M. Ferreira. Application of Ricatti equation to the busy period study of the  $M|G|\infty$  system. *Statistical Review*, 1<sup>st</sup> Quadrimester, INE, 23-28, 1998.
14. M. A. M. Ferreira. Computational simulation of infinite servers systems. *Statistical Review*, 3<sup>rd</sup> Quadrimester, INE, 23-28, 1998.
15. M. A. M. Ferreira.  $M|G|\infty$  queue heavy-traffic situation busy period length distribution (power and Pareto service distributions). *Statistical Review*, 1<sup>st</sup> Quadrimester, INE, 27-36, 2001.
16. M. A. M. Ferreira. The exponentiality of the  $M|G|\infty$  queue busy period. *Actas das XII Jornadas Luso-Espanholas de Gestão Científica*, Volume VIII-Economia da Empresa e Matemática Aplicada. UBI, Covilhã, Portugal, 267-272, 2002.
17. M. A. M. Ferreira. Differential equations important in the  $M|G|\infty$  queue system transient behavior and busy period study. *Proceedings of 4th International Conference APLIMAT 2005*, Bratislava, Slovakia, 119-132, 2005.
18. M. A. M. Ferreira, M. Andrade. The ties between the  $M|G|\infty$  queue system transient behavior and the busy period. *International Journal of Academic Research*, 1(1), 84-92, 2009.

19. M. A. M. Ferreira, M. Andrade. Looking to a  $M|G|\infty$  system occupation through a Ricatti equation. *Journal of Mathematics and Technology*, 1(2), 58-62, 2010.
20. M. A. M. Ferreira, M. Andrade.  $M|G|\infty$  system transient behavior with time origin at the beginning of a busy period mean and variance. *APLIMAT- Journal of Applied Mathematics*, 3(3), 213-221, 2010.
21. M. A. M. Ferreira, M. Andrade. Fundamentals of theory of queues. *International Journal of Academic Research*, 3(1), part II, 427-429, 2011.
22. M. A. M. Ferreira, M. Andrade. Busy period and busy cycle distributions and parameters for a particular  $M|G|\infty$  queue system. *American Journal of Mathematics and Statistics*, 2(2), 10-15, 2012.
23. M. A. M. Ferreira, M. Andrade. Queue networks models with more general arrival rates. *International Journal of Academic Research*, 4(1), part A, 5-11, 2012.
24. M. A. M. Ferreira, M. Andrade. Transient behavior of the  $M|G|m$  and  $M|G|\infty$  system. *International Journal of Academic Research*, 4(3), part A, 24-33, 2012.
25. M. A. M. Ferreira, M. F. Ramalhoto. Estudo dos parâmetros básicos do período de ocupação da fila de espera  $M|G|\infty$ . *A Estatística e o Futuro e o Futuro da Estatística. Actas do I Congresso Anual da S.P.E.. Edições Salamandra, Lisboa, 1994.*
26. M. A. M. Ferreira, M. Andrade, J. A. Filipe. The Ricatti equation in the  $M|G|\infty$  busy cycle study. *Journal of Mathematics, Statistics and Allied Fields* 2(1), 2008.
27. M. A. M. Ferreira, M. Andrade, J. A. Filipe. Networks of queues with infinite servers in each node applied to the management of a two echelons repair system. *China-USA Business Review*, 8(8), 39-45 and 62, 2009.
28. M. A. M. Ferreira, M. Andrade, J. A. Filipe. The age or excess of the  $M|G|\infty$  queue busy cycle mean value. *Computer and Information Science*, 5(5), 93-97, 2012.
29. M. F. Ramalhoto. A note on the variance of the busy period of the  $M|G|\infty$  systems. *Centro de Estatística e Aplicações, CEAUL, do INIC and IST, 1983.*
30. M. F. Ramalhoto, M. A. M. Ferreira. Some further properties of the busy period of an  $M|G|\infty$  queue. *Central European Journal of Operations Research and Economics*, 4(4), 251-278, 1996.
31. M. G. Kendall and A. Stuart: *The advanced theory of statistics. Distributions theory.* London, Charles Griffin and Co., Ltd. 4<sup>th</sup> Edition, 1979.
32. N. Metropolis, S. Ulan. The monte carlo method. *Journal of American Statistical Association*, 44(247), 335-341, 1949.
33. R. Syski. *Introduction to congestion theory in telephone systems.* Oliver and Boyd- London, 1960.
34. R. Syski. *Introduction to congestion theory in telephone systems.* North Holland, Amsterdam, 1986.
35. W. Stadge. The busy period of the queueing system  $M|G|\infty$ . *Journal of Applied Probability*, 22, 697-704, 1985.

