

Isolation of the diamond graph

Jingru Yan *

Department of Mathematics, East China Normal University, Shanghai 200241, China

Abstract

A graph is H -free if it does not contain H as a subgraph. The diamond graph is the graph obtained from K_4 by deleting one edge. We prove that if G is a connected graph with order $n \geq 10$, then there exists a subset $S \subseteq V(G)$ with $|S| \leq n/5$ such that the graph induced by $V(G) \setminus N[S]$ is diamond-free, where $N[S]$ is the closed neighborhood of S . Furthermore, the bound is sharp.

Keywords. Diamond graph, isolating set, isolation number

Mathematics Subject Classification. 05C35, 05C69

1 Introduction

Let G be a finite simple graph with vertex set $V(G)$ and edge set $E(G)$. The order and the size of a graph G , denoted $|V(G)|$ and $|E(G)|$, are its number of vertices and edges, respectively. For a subset $S \subseteq V(G)$, the neighborhood of S is the set $N_G(S) = \{u \in V(G) \setminus S \mid uv \in E(G), v \in S\}$ and closed neighborhood of S is the set $N_G[S] = N_G(S) \cup S$. Thus $N_G(v)$ and $N_G[v]$ denote the neighborhood and closed neighborhood of $v \in V(G)$, respectively. The degree of v is $d_G(v) = |N_G(v)|$. If the graph G is clear from the context, we will omit it as the subscript. $\delta(G)$ and $\Delta(G)$ denote the minimum and maximum degree of a graph G , respectively. Denote by $G[S]$ the subgraph of G induced by $S \subseteq V(G)$. For terminology and notations not explicitly described in this paper, readers can refer to related books [1, 9].

Given graphs G and H , the notation $G + H$ means the *disjoint union* of G and H . Then tG denotes the disjoint union of t copies of G . For graphs we will use equality up to isomorphism, so $G = H$ means that G and H are isomorphic. A graph is H -free if it does not contain H as a subgraph. $\kappa(G)$ and $\gamma(G)$ denote the connectivity and domination number of a graph G , respectively. P_n, C_n, K_n and $K_{p,q}$ stand for the *path*, *cycle*, *complete graph* of order n and *complete bipartite graph* with partition sets of p and q vertices, respectively.

*E-mail address: mathyjr@163.com




Fig. 1. Diamond graph




Fig. 2. Y

Let G be a graph and \mathcal{F} a family of graphs. A subset $S \subseteq V(G)$ is called an \mathcal{F} -isolating set of G if $G - N[S]$ contains no subgraph isomorphic to any $F \in \mathcal{F}$. The minimum cardinality of an \mathcal{F} -isolating set of a graph G will be denoted $\iota(G, \mathcal{F})$ and called the \mathcal{F} -isolation number of G . When $\mathcal{F} = \{H\}$, we simply write $\iota(G, H)$ for $\iota(G, \{H\})$.

The definition of isolation set is a natural extension of the commonly defined dominating set, which was introduced by Caro and Hansberg [5]. Indeed, if $\mathcal{F} = \{K_1\}$, then an \mathcal{F} -isolating set coincides with a dominating set and $\iota(G, \mathcal{F}) = \gamma(G)$. A classical result of Ore [7] is that the domination number of a graph G with order n and $\delta(G) \geq 1$ is at most $n/2$. In other words, if G is a connected graph of order $n \geq 2$, then $\iota(G, K_1) \leq n/2$. Caro and Hansberg [5] focused mainly on $\iota(G, K_2)$ and $\iota(G, K_{1,k+1})$ and gave some basic properties, examples concerning $\iota(G, \mathcal{F})$ and the relation between \mathcal{F} -isolating sets and dominating sets. They [5] proved that if G is a connected graph of order $n \geq 3$ that is not a C_5 , then $\iota(G, K_2) \leq \frac{n}{3}$. Since then, Borg [2] showed that if G is a connected graph of order n , then $\iota(G, \{C_k : k \geq 3\}) \leq \frac{n}{4}$ unless G is K_3 . After that, Borg, Fenech and Kaemawichanurat [3] proved that if G is a connected graph of order n , then $\iota(G, K_k) \leq \frac{n}{k+1}$ unless G is K_k , or $k = 2$ and G is C_5 . Both the bounds are sharp. Then Zhang and Wu [10] gave the result that if G is a connected graph of order n , then $\iota(G, P_3) \leq \frac{2n}{7}$ unless $G \in \{P_3, C_3, C_6\}$, and this bound can be improved to $\frac{n}{4}$ if $G \notin \{P_3, C_7, C_{11}\}$ and the girth of G at least 7. For more research on isolation set, refer to [4, 6, 8].

The *diamond graph* is the graph obtained from K_4 by deleting one edge (see Figure 1). The *book graph* with p pages, denoted B_p , is the graph that consists of p triangles sharing a common edge. Obviously, B_2 is the diamond graph. For the convenience of expression, we use B_2 to represent the diamond graph in the sequel.

In this paper, we consider the isolation number of the diamond graph in a connected graph of a given order.

Theorem 1. *If G is a connected graph of order n , then, unless G is the diamond graph, K_4 , or Y ,*

$$\iota(G, B_2) \leq \frac{n}{5},$$

where Y is shown in the Figure 2.

2 Main results

From the proof of Theorem 3.8 in this paper [5], we obtain Lemma 2 and give an example that satisfies the Lemma, see Figure 3. The minimum cardinality of a B_2 -isolating set of the graph H of order 15 is 3.

Lemma 2. *There exists a connected graph G of order n such that $\iota(G, B_2) = \frac{n}{5}$.*




Fig. 3. H

We start with two lemmas that will be used repeatedly.

Lemma 3. [2] *If G is a graph, \mathcal{F} is a set of graphs, $A \subseteq V(G)$, and $B \subseteq N[A]$, then*

$$\iota(G, \mathcal{F}) \leq |A| + \iota(G - B, \mathcal{F}).$$

In particular, if $A = \{v\}$ and $B = N[A]$, then $\iota(G, \mathcal{F}) \leq 1 + \iota(G - N[v], \mathcal{F})$.

Lemma 4. [2] *If G_1, \dots, G_k are the distinct components of a graph G , then*

$$\iota(G, \mathcal{F}) = \sum_{i=1}^k \iota(G_i, \mathcal{F}).$$

For any graph G , let $A, B \subseteq V(G)$ and $A \cap B = \emptyset$. Denote by $E(A, B)$ the set of edges of G with one end in A and the other end in B and $e(A, B) = |E(A, B)|$. We abbreviate $E(\{x\}, B)$ to $E(x, B)$ and $e(\{x\}, B)$ to $e(x, B)$.

Now, we first prove Theorem 1 when the order $n \leq 9$.

Lemma 5. *Let G be a connected graph of order $n \leq 9$. Then $\iota(G, B_2) \leq \frac{n}{5}$ except for $G \in \{B_2, K_4, Y\}$.*

Proof. Let G be a connected graph of order n . The result is trivial if $n \leq 4$ or $\iota(G, B_2) = 0$. Suppose $5 \leq n \leq 9$ and $\iota(G, B_2) \geq 1$. Then we need to show that G has a B_2 -isolating set S with $|S| = 1$ except for $G = Y$.

Since $\iota(G, B_2) \geq 1$, it follows G contains B_2 and $\Delta(G) \geq 3$. Let $x \in V(G)$ such that $d(x) = \Delta(G)$. Of course, $S = \{x\}$ is a B_2 -isolating set of G if $G - N[x]$ is B_2 -free. Otherwise, it implies that $n = 8$ with $\Delta(G) = 3$ or $n = 9$ with $3 \leq \Delta(G) \leq 4$. We distinguish two cases.

Case 1. $\Delta(G) = 3$ and $n = 8$ or 9

Let $u \in V(G)$ such that $d(u) = 3$ and $G[N[u]] = B_2$. If $G - N[u]$ is B_2 -free, then $\iota(G, B_2) = 1 \leq \frac{n}{5}$. So, suppose that $G - N[u]$ contains B_2 . For $n = 8$, obviously, $G - N[u] = B_2$. Since G is a connected graph and $\Delta(G) = 3$, there is an edge $e = yz$ with $y \in N(u)$ and $z \in V(G) \setminus N[u]$. It is easy to check that $\{y\}$ or $\{z\}$ is a B_2 -isolating set of G . Hence $\iota(G, B_2) = 1 \leq \frac{n}{5}$. Now we prove the case of $n = 9$. Let us consider a copy H of B_2 in $G - N[u]$ and let w be the remaining vertex of $G - N[u] - V(H)$. If there is an edge $e = yz$ with $y \in N(u)$ and $z \in V(H)$, then $\{y\}$ or $\{z\}$ is a B_2 -isolating set of G . Otherwise, w is a cut-vertex of G and $G - N[w]$ is B_2 -free. Hence $\iota(G, B_2) = 1 \leq \frac{n}{5}$.

Case 2. $\Delta(G) = 4$ and $n = 9$

Let $u \in V(G)$ such that $d(u) = 4$ and let $F = G - N[u]$. We have $\iota(G, B_2) = 1$ if F is B_2 -free. Assume that F contains B_2 . Since $|V(F)| = 4$, then $F = B_2$ or $F = K_4$. The vertices are labeled as shown in the Figure 4. We distinguish two subcases.




Fig. 4.




Fig. 5.

Subcase 2.1. $F = K_4$. Note that $e(N(u), V(F)) \neq 0$. Without loss of generality, suppose u_1 is adjacent to v . If $G - N[v]$ is B_2 -free, $\iota(G, B_2) = 1$. Otherwise, $G - N[v] = K_4$ or B_2

since $|V(G) \setminus N[v]| = 4$. For $G - N[v] = K_4$, we have $G - \{u, u_1, v\}$ is B_2 -free. Thus, $\{u_1\}$ is a B_2 -isolating set of G and $\iota(G, B_2) \leq \frac{n}{5}$. For $G - N[v] = B_2$, $G - \{u, u_1, v\}$ contains B_2 if and only if u_2 or u_4 is adjacent to at least two vertices of $\{v_1, v_2, v_3\}$. Now we have $\{u_2\}$ or $\{u_4\}$ is a B_2 -isolating set of G and hence $\iota(G, B_2) \leq \frac{n}{5}$.

Subcase 2.2. $F = B_2$. The proof for this case is similar to Subcase 2.1. First suppose F has a vertex of degree 3 that is adjacent to one vertex of $N(u)$. Without loss of generality, suppose v , $d_F(v) = 3$, is adjacent to u_1 . Then we have $\iota(G, B_2) = 1$ if $G - N[v]$ is B_2 -free. Otherwise, $G - N[v]$ contains B_2 . For $G - N[v] = K_4$, by the proof of Subcase 2.1, $\iota(G, B_2) \leq \frac{n}{5}$. For $G - N[v] = B_2$, $G - \{u, u_1, v\}$ contains B_2 when one of the following four cases is true. (1) u_2 is adjacent to v_1 and v_2 and u_3 is adjacent to v_1 . (2) u_2 is adjacent to v_2 and v_3 and u_3 is adjacent to v_3 . (3) u_3 is adjacent to v_1 and u_4 is adjacent to v_1 and v_2 . (4) u_3 is adjacent to v_3 and u_4 is adjacent to v_2 and v_3 . We can see from Figure 5 that the proof methods are similar for the four cases. So let us just consider the first case. Note that $G - N[u_2]$ contains B_2 if and only if $G[u_1, u_4, v_3] = K_3$. Observe that $G = Y$.

Next suppose that only the vertices of degree 2 of F are adjacent to the vertices of $N(u)$. Suppose v_1 , $d_F(v_1) = 2$, is adjacent to u_1 . Then $\iota(G, B_2) = 1$ if $G - \{u, u_1, v_1\}$ is B_2 -free. Otherwise, since $e(v, N(u)) = e(v_2, N(u)) = 0$, we have $G[u_2, u_3, u_4, v_3]$ contains B_2 as a subgraph. Recall that $\Delta(G) = 4$, then $G[u_2, u_3, u_4, v_3] = B_2$. Moreover, $G - N[v_3]$ is B_2 -free. Thus, $\iota(G, B_2) \leq \frac{n}{5}$.

Hence, in all cases we obtain $\iota(G, B_2) \leq \frac{n}{5}$ with $n \leq 9$ except for $\iota(B_2, B_2) = 1$, $\iota(K_4, B_2) = 1$ and $\iota(Y, B_2) = 2$. \square

Next, we prove Theorem 1 when $\Delta(G) = 3$.

Lemma 6. *Let G be a connected graph of order n . $\iota(G', B_2) = \iota(G, B_2)$ if*

- (1) G' is obtained from G by attaching one edge to any vertex of G ,
- (2) G' is obtained from G by identifying one vertex of a triangle and a vertex of G ,
- (3) G' is obtained from $G + K_3$ by adding an edge joining a vertex of K_3 and a vertex of G .

Proof. (1) Let S be a minimum B_2 -isolating set of G . Then, clearly, S is a B_2 -isolating set of G' and thus $\iota(G', B_2) \leq \iota(G, B_2)$. Let S' be a minimum B_2 -isolating set of G' and let x be the vertex of $V(G') \setminus V(G)$. Note that $S' \setminus \{x\}$ is a B_2 -isolating set of G . Thus, $\iota(G, B_2) \leq \iota(G', B_2)$. Now the both inequalities imply the result.

(2) and (3) can be proved similarly as (1). \square

Lemma 7. *Let G be a connected graph of order n . If $\Delta(G) = 3$, then*

$$\iota(G, B_2) \leq \frac{n}{5}$$

except for $G \in \{B_2, K_4\}$.

Proof. Let G be a connected graph of order n with $\Delta(G) = 3$. The proof is by induction on n . By Lemma 5, the result is trivial if $n \leq 9$ or $\iota(G, B_2) = 0$. Thus, suppose that $n \geq 10$ and $\iota(G, B_2) \geq 1$. Since G contains B_2 , it follows that there exists at least one vertex $u \in V(G)$ such that $d(u) = 3$ and $G[N[u]] = B_2$. Let $N(u) = \{u_1, u_2, u_3\}$ and let u_2 be the another vertex of the B_2 with degree 3. As G is connected and $\Delta(G) = 3$, then either $d(u_1) = 3$ or $d(u_3) = 3$. We distinguish the following two cases.

Case 1. $d(u_1) = 3$ and $d(u_3) = 2$ or $d(u_1) = 2$ and $d(u_3) = 3$

Without loss of generality, suppose $d(u_1) = 3$ and $d(u_3) = 2$. Let $w \in V(G - N[u])$ and w is adjacent to u_1 . Define $G' = G - N[u] - w$. Note that $|V(G')| = n - 5 \geq 5$. Clearly, $\{u_1\}$ is a B_2 -isolating set of G if G' is B_2 -free. Suppose G' contains B_2 . If G' is connected, by the induction hypothesis, $\iota(G', B_2) \leq \frac{n-5}{5}$. Then by Lemma 3 and 4, we have $\iota(G, B_2) \leq |\{u_1\}| + \iota(G', B_2) \leq 1 + \frac{n-5}{5} = \frac{n}{5}$.

Suppose that G' is disconnected. It is easy to check that $d(w) = 3$ and G' has exactly two components. Let $G' = G_1 + G_2$. If $G_1 \neq B_2$ and $G_2 \neq B_2$, the union of a minimum B_2 -isolating set of G_1 , a minimum B_2 -isolating set of G_2 and $\{u_1\}$ is a B_2 -isolating set of G . By the induction hypothesis and Lemma 4,

$$\iota(G, B_2) \leq 1 + \iota(G_1, B_2) + \iota(G_2, B_2) \leq 1 + \frac{|V(G_1)|}{5} + \frac{|V(G_2)|}{5} = \frac{n}{5}.$$

If $G_1 = B_2$ and $G_2 = B_2$, we have $n = 13$. Observe that $\{w\}$ is a B_2 -isolating set of G . Hence $\iota(G, B_2) \leq \frac{n}{5}$. So, it remains to consider the case of exactly one of $\{G_1, G_2\}$ is isomorphic to B_2 . Suppose that $G_1 = B_2$ and $G_2 \neq B_2$. Let w_1 be the neighbor of w in G_2 and let $G'' = G' - V(G_1) - w_1$. Note that $|V(G'')| = n - 10$.

If G'' is connected and $G'' \neq B_2$, by the induction hypothesis, $\iota(G'', B_2) \leq \frac{n-10}{5}$. Then the union of $\{w\}$ and a minimum B_2 -isolating set of G'' is a B_2 -isolating set of G . Thus, $\iota(G, B_2) \leq 1 + \frac{n-10}{5} \leq \frac{n}{5}$. Observe that $n = 14$ and $\{w, w_1\}$ is a B_2 -isolating set of G when $G'' = B_2$. We also have $\iota(G, B_2) \leq \frac{n}{5}$. Suppose that G'' is disconnected. Recall that $\Delta(G) = 3$, then $d(w_1) = 3$ and G'' has exactly two components. Let $G'' = G'_1 + G'_2$ (see Figure 6). Now let us consider the components G'_1 and G'_2 . If $G'_1 \neq B_2$ and $G'_2 \neq B_2$, then




Fig. 6.

the union of a minimum B_2 -isolating set of G'_1 , a minimum B_2 -isolating set of G'_2 and $\{w\}$ is a B_2 -isolating set of G . Thus, by Lemma 3, 4 and the induction hypothesis,

$$\iota(G, B_2) \leq 1 + \iota(G'_1, B_2) + \iota(G'_2, B_2) \leq 1 + \frac{|V(G'_1)|}{5} + \frac{|V(G'_2)|}{5} \leq \frac{n}{5}.$$

If $G'_1 = B_2$ and $G'_2 = B_2$, we have $n = 18$ and $\{w, w_1\}$ is a B_2 -isolating set of G . Hence $\iota(G, B_2) \leq \frac{n}{5}$. So, it remains to consider the case of exactly one of $\{G'_1, G'_2\}$ is isomorphic to B_2 . Suppose that $G'_1 = B_2$ and $G'_2 \neq B_2$. Note that the union of a minimum B_2 -isolating set of G'_2 , the neighbor of w_1 in G'_1 and $\{w\}$ is a B_2 -isolating set of G . Therefore, $\iota(G, B_2) \leq 1 + 1 + \frac{|V(G'_2)|}{5} \leq \frac{n}{5}$. This completes the proof of Case 1.

Case 2. $d(u_1) = 3$ and $d(u_3) = 3$

If $N(u_1) = N(u_3)$, denote $G^* = G \setminus (N[u_1] \cup \{u_3\})$. Then G^* is connected and $|G^*| = n - 5 \geq 5$ since $\Delta(G) = 3$ and $n \geq 10$. Observe that the union of $\{u_1\}$ and a minimum B_2 -isolating set of G^* is a B_2 -isolating set of G . Hence, by the induction hypothesis, $\iota(G, B_2) \leq 1 + \frac{|V(G^*)|}{5} = \frac{n}{5}$. Otherwise, there exist two vertices $w, z \in V(G)$ such that u_1 is adjacent to w and u_3 is adjacent to z .

We first prove $G - N[u]$ is connected. Let $G' = G - N[u] - w$. Note that this case differs from Case 1 only in that there is an edge between u_3 and G' . By Lemma 6 and the proof of Case 1, we have $\iota(G, B_2) \leq \frac{n}{5}$. Therefore, we omit the proof. Next we treat $G - N[u]$ is disconnected. Since $\Delta(G) = 3$, then $G - N[u]$ contains exactly two components and w and z belong to different components. Define $G - N[u] = G_w + G_z$, where G_w

contains w and G_z contains z . Obviously, if $G_w = B_2$ or $G_z = B_2$, the union of $\{u_1\}$ and a minimum B_2 -isolating set of G_z or the union of $\{u_3\}$ and a minimum B_2 -isolating set of G_w is a B_2 -isolating set of G , respectively. Hence $\iota(G, B_2) \leq \frac{n}{5}$. So, suppose that $G_w \neq B_2$ and $G_z \neq B_2$. Let $G' = G - V(G_z) - N[u] - w$. By Lemma 6 (3), we have $\iota(G_z, B_2) = \iota(G[V(G_z) \cup \{u, u_2, u_3\}], B_2)$. Similarly, using the same method of Case 1, we have $\iota(G, B_2) \leq \frac{n}{5}$. This completes the proof of Lemma 7. \square

So far, it remains to consider Theorem 1 when $\Delta(G) \geq 4$.

Lemma 8. *The connected graph Y of order 9 has the following properties:*

- (1) $\kappa(Y) = 4$,
- (2) $\Delta(Y) = \delta(Y) = 4$,
- (3) for any two vertices $u, v \in V(Y)$, $|N(u) \cap N(v)| \leq 2$,
- (4) for any vertex $u \in V(Y)$, there exists a vertex $v \in V(Y) \setminus \{u\}$ such that the graph induced by $V(Y) \setminus (\{u\} \cup N[v])$ is P_3 .

Proof. It is easy to check these properties of the graph Y (see Figure 2). \square

Lemma 9. [5] *Let G be a graph on n vertices and \mathcal{F} a family of graphs and let $A \cup B$ be a partition of $V(G)$. Then*

$$\iota(G, \mathcal{F}) \leq \iota(G[A], \mathcal{F}) + \gamma(G[B]).$$

Lemma 10. *Let G be a connected graph of order n . If $\Delta(G) \geq 4$, then*

$$\iota(G, B_2) \leq \frac{n}{5}$$

except for $G = Y$.

Proof. Let G be a connected graph of order n with $\Delta(G) \geq 4$. The proof is by induction on n . By Lemma 5, the result is trivial if $n \leq 9$ or $\iota(G, B_2) = 0$. Thus, suppose that $n \geq 10$ and $\iota(G, B_2) \geq 1$. Denote by $d(u) = \Delta(G)$ and $H = G - N[u]$. Obviously, $\iota(G, B_2) = 1$ if H is B_2 -free. If $H = B_2$ or K_4 , $\iota(G, B_2) \leq 1 + 1 = 2 \leq \frac{n}{5}$ for $n \geq 10$. If $H = Y$, then $\Delta(G) \geq 5$. Hence we have $n \geq 15$ and $\iota(G, B_2) \leq 1 + \iota(Y, B_2) = 3 \leq \frac{n}{5}$. Suppose that $H \neq B_2, K_4, Y$. By Lemma 7 and the induction hypothesis, it is easy to check that $\iota(G, B_2) \leq \frac{n}{5}$ when H is connected. Therefore, let $H = G_1 + G_2 + \cdots + G_k$ with $k \geq 2$ and $|V(G_i)| = n_i$ for $i = 1, 2, \dots, k$. If H does not contain B_2, K_4 or Y as a component, by Lemma 3, 4, 7 and the induction hypothesis, we have

$$\iota(G, B_2) \leq |\{u\}| + \sum_{i=1}^k \iota(G_i, B_2) \leq 1 + \frac{n_1}{5} + \frac{n_2}{5} + \cdots + \frac{n_k}{5} = \frac{n - \Delta(G) + 4}{5}.$$

Since $\Delta(G) \geq 4$, then $\iota(G, B_2) \leq \frac{n}{5}$.

Next suppose that at least one component of H is B_2, K_4 or Y . We sort the components of H in the order of K_4, Y, B_2 with one vertex of degree 3 of B_2 is adjacent to one vertex of $N(u)$, B_2 with only vertices of degree 2 of B_2 are adjacent to vertices of $N(u)$, and others. Then G_1 is isomorphic to K_4, Y , or B_2 . Let $N(u) = \{u_1, u_2, \dots, u_{\Delta(G)}\}$. Since G is connected, without loss of generality, suppose $N(u_1) \cap V(G_1) \neq \phi$. Denote by $G^* = G - u_1 - V(G_1)$. Obviously, $|V(G^*)| \geq 5$.

Case 1. G^* is connected.

Subcase 1.1. $G_1 = K_4$. If $G^* = Y$, we have $n = 14$ and $\Delta(G) = 5$. By Lemma 8 (4), there exists a vertex $v \in V(G^*)$ such that the graph induced by $G^* - u - N[v]$ is P_3 . Since $\Delta(G) = 5$, we have $\{u_1, v\}$ is a B_2 -isolating set of G and hence $\iota(G, B_2) \leq 2 \leq \frac{n}{5}$. If $G^* \neq Y$, by the induction hypothesis and Lemma 7, $\iota(G^*, B_2) \leq \frac{n-5}{5}$. Then, by Lemma 9, $\iota(G, B_2) \leq \gamma(G[V(G_1) \cup \{u_1\}]) + \iota(G^*, B_2) \leq 1 + \frac{n-5}{5} = \frac{n}{5}$.

Subcase 1.2. $G_1 = Y$. Let x be a neighbor of u_1 in $V(G_1)$. If $G^* = Y$, we have $n = 19$ and $\Delta(G) = 5$. Then, by Lemma 8 (4), there exist a vertex $v_1 \in V(G^*)$ such that the graph induced by $G^* - u - N[v_1]$ is P_3 and a vertex $v_2 \in V(G_1)$ such that the graph induced by $G_1 - x - N[v_2]$ is P_3 . Then $\{v_1, u_1, v_2\}$ is a B_2 -isolating set of G and $\iota(G, B_2) \leq 3 \leq \frac{n}{5}$. If $G^* \neq Y$, similar to Subcase 1.1, $\iota(G, B_2) \leq \gamma(G[V(G_1) \cup \{u_1\}]) + \iota(G^*, B_2) \leq 2 + \frac{n-10}{5} = \frac{n}{5}$.

Subcase 1.3. $G_1 = B_2$ and there is one vertex of degree 3 of $V(G_1)$ is adjacent to u_1 . Let x be a neighbor of u_1 in $V(G_1)$ and $d_{G[V(G_1)]}(x) = 3$. If $G^* = Y$, we have $n = 14$ and $\Delta(G) = 5$. Then, similarly, there exists $v \in V(G^*)$ such that $G^* - u - N[v] = P_3$. Define $P = P_3$. If $G^* - N[u_1] - N[v]$ has no B_2 , then $\iota(G, B_2) \leq 2 \leq \frac{n}{5}$. Otherwise, let $N(x) = \{x_1, x_2, x_3\}$ and let $d_{G[V(G_1)]}(x_2) = 3$. Observe that $G^* - N[u_1] - N[v]$ contains B_2 if and only if $e(V(P), x_1) = 3$ or $e(V(P), x_3) = 3$. Assume that $e(V(P), x_3) = 3$, then $d(x_3) = 5$. By Lemma 8 (1), $G - (N[x_3] \setminus \{x\})$ is a connected graph of order 9. Since $G - (N[x_3] \setminus \{x\}) \neq Y$, by Lemma 3 and 5,

$$\iota(G, B_2) \leq |\{x_3\}| + \iota(G - (N[x_3] \setminus \{x\}), B_2) \leq 1 + 1 \leq \frac{n}{5}.$$

If $G^* \neq Y$, similar to Subcase 1.1, $\iota(G, B_2) \leq \gamma(G[V(G_1) \cup \{u_1\}]) + \iota(G^*, B_2) \leq 1 + \frac{n-5}{5} = \frac{n}{5}$.

Subcase 1.4. $G_1 = B_2$ and only vertices of degree 2 of $V(G_1)$ are adjacent to u_1 . Let x be a neighbor of u_1 in $V(G_1)$ and $d_{G[V(G_1)]}(x) = 2$ and Let $d_{G[V(G_1)]}(x_2) = 2$. Note that the two remaining vertices of $V(G_1) \setminus \{x, x_2\}$ have degrees of 3 in G . First we prove the case of

$u_1 \in N(x_2)$ and the case $u_1 \notin N(x_2)$ and $|N(x_2) \cap N(u)| \leq 1$. If $G^* = Y$, we have $n = 14$ and $\Delta(G) = 5$. By Lemma 8 (4), there exists $v \in V(G^*)$ such that $G^* - u - N[v] = P_3$. Since $\Delta(G) = 5$, then $\{u_1, v\}$ is a B_2 -isolating set of G and $\iota(G, B_2) \leq 2 \leq \frac{n}{5}$. If $G^* \neq Y$, by the induction hypothesis, Lemma 6 (1) and 7, $\iota(G, B_2) \leq 1 + \frac{n-5}{5} = \frac{n}{5}$. It remains the case of $u_1 \notin N(x_2)$ and $|N(x_2) \cap N(u)| \geq 2$, we will deal with it later.

Case 2. G^* is disconnected.

It implies that $E(V(G_i), N(u)) = E(V(G_i), u_1)$ for some $i \in \{2, 3, \dots, k\}$. Let us denote the components satisfying $E(V(G_i), N(u)) = E(V(G_i), u_1)$ as $G_{11}, G_{12}, \dots, G_{1t}$, $t \geq 1$. Let G_u be the component contains u in G^* . Then $G^* = G_{11} + G_{12} + \dots + G_{1t} + G_u$. Assume that there are $s_1 B_2$, $s_2 K_4$ and $s_3 Y$ in $\{G_{11}, G_{12}, \dots, G_{1t}\}$.

Subcase 2.1. $G_1 = K_4$. Let x be a neighbor of u_1 in $V(G_1)$ and let $N(x) = \{x_1, x_2, x_3\}$. It is easy to check that $\iota(G, B_2) \leq 1 + \frac{|V(G_{11})|}{5} + \dots + \frac{|V(G_{1t})|}{5} + \frac{|V(G_u)|}{5} = \frac{n}{5}$ if $G_{11}, G_{12}, \dots, G_{1t}, G_u \notin \{B_2, K_4, Y\}$. If $G_u = K_4$, then $\Delta(G) = 4$. Hence, by Lemma 3, 4, 7 and the induction hypothesis,

$$\iota(G, B_2) \leq |\{u_1\}| + \sum_{i=1}^t \iota(G_{1i} - N[u_1], B_2) \leq 1 + s_3 + \frac{n - (5 + 4 + 4s_1 + 4s_2 + 9s_3)}{5} \leq \frac{n}{5}.$$

If $G_u = B_2$, then $\Delta(G) = 4$. Note that $G[N[u] \cup V(G_1)] - \{u, u_1, x\}$ contains B_2 if and only if $e(u_2, \{x_1, x_2, x_3\}) = 2$ or $e(u_4, \{x_1, x_2, x_3\}) = 2$. Without loss of generality, suppose $e(u_2, \{x_1, x_2, x_3\}) = 2$. Then $d(u_2) = 4$ and $G - N[u_2]$ is a connected graph of order $n - 5$ or the union of a connected graph of order $n - 6$ and an isolated vertex. By Lemma 8, $G - N[u_2]$ does not contain Y as an induced subgraph. Hence, by the induction hypothesis and Lemma 7,

$$\iota(G, B_2) \leq |\{u_2\}| + \iota(G - N[u_2], B_2) \leq 1 + \frac{n-5}{5} = \frac{n}{5}.$$

If $G_u = Y$, by Lemma 8 (4), there exists $v \in V(G_u)$ such that $G_u - u - N[v] = P_3$. Note that $\Delta(G) = 5$, then

$$\iota(G, B_2) \leq |\{u_1, v\}| + \sum_{i=1}^t \iota(G_{1i} - N[u_1], B_2) \leq 2 + s_3 + \frac{n - (5 + 9 + 4s_1 + 4s_2 + 9s_3)}{5} \leq \frac{n}{5}.$$

Suppose $G_u \neq B_2, K_4, Y$. Then at least one of $\{s_1, s_2, s_3\}$ is not less than one. Obviously,

$$\iota(G, B_2) \leq |\{u_1\}| + \sum_{i=1}^t \iota(G_{1i} - N[u_1], B_2) + \iota(G_u, B_2) \leq 1 + s_3 + \frac{n - (5 + 4s_1 + 4s_2 + 9s_3)}{5} \leq \frac{n}{5}$$

when $e(V(G_u), V(G_1) \setminus \{x\}) = 0$. For $e(V(G_u), V(G_1) \setminus \{x\}) > 0$, $G[V(G_u) \cup \{u_1\} \cup V(G_1)] - \{u_1, x\} \neq B_2, K_4$. If $G[V(G_u) \cup \{u_1\} \cup V(G_1)] - \{u_1, x\} = Y$, by Lemma 8 (4), there exists v such that $G[V(G_u) \cup \{u_1\} \cup V(G_1)] - \{u_1, x, u\} - N[v] = P_3$. Then we have

$$\iota(G, B_2) \leq |\{u_1, v\}| + \sum_{i=1}^t \iota(G_{1i} - N[u_1], B_2) \leq 2 + s_3 + \frac{n - (2 + 9 + 4s_1 + 4s_2 + 9s_3)}{5} \leq \frac{n}{5}.$$

Otherwise,

$$\begin{aligned} \iota(G, B_2) &\leq |\{u_1\}| + \iota(G[V(G_u) \cup V(G_1) \setminus \{x\}], B_2) + \sum_{i=1}^t \iota(G_{1i} - N[u_1], B_2) \\ &\leq 1 + s_3 + \frac{n - (2 + 4s_1 + 4s_2 + 9s_3)}{5} \leq \frac{n}{5}. \end{aligned}$$

Subcase 2.2. $G_1 = Y$. Note that none of the components of H is K_4 . It follows that $s_2 = 0$. Let x be a neighbor of u_1 in $V(G_1)$. It is easy to check that $\iota(G, B_2) \leq 2 + \frac{|V(G_{11})|}{5} + \dots + \frac{|V(G_{1t})|}{5} + \frac{|V(G_u)|}{5} = \frac{n}{5}$ if $G_{11}, G_{12}, \dots, G_{1t}, G_u \notin \{B_2, K_4, Y\}$. Since $\Delta(G) \geq 5$, we have $G_u \neq B_2, K_4$. If $G_u = Y$, then $\Delta(G) = 5$. Similar to the proofs of Subcase 1.2 and Subcase 2.1, we have

$$\iota(G, B_2) \leq 3 + \sum_{i=1}^t \iota(G_{1i} - N[u_1], B_2) \leq 3 + s_3 + \frac{n - (19 + 4s_1 + 9s_3)}{5} \leq \frac{n}{5}.$$

Suppose $G_u \neq Y$. Then at least one of $\{s_1, s_3\}$ is not less than one. If $e(V(G_u), V(G_1) \setminus \{x\}) = 0$, we have

$$\iota(G, B_2) \leq \iota(G_1 + u_1, B_2) + \iota(G_u, B_2) + \sum_{i=1}^t \iota(G_{1i} - N[u_1]) \leq 2 + s_3 + \frac{n - (10 + 4s_1 + 9s_3)}{5} \leq \frac{n}{5}.$$

Otherwise,

$$\begin{aligned} \iota(G, B_2) &\leq |\{u_1\}| + \iota(G[V(G_u) \cup V(G_1) \setminus \{x\}], B_2) + \sum_{i=1}^t \iota(G_{1i} - N[u_1], B_2) \\ &\leq 1 + s_3 + \frac{n - (2 + 4s_1 + 9s_3)}{5} \leq \frac{n}{5} \end{aligned}$$

since the component of $G[V(G_1) \cup \{u_1\} \cup V(G_u)] - \{u_1, x\}$ is not B_2, K_4 or Y .

Subcase 2.3. $G_1 = B_2$ and there is one vertex of degree 3 of $V(G_1)$ is adjacent to u_1 . Note that none of the components of H is K_4 or Y . It follows that $s_2 = s_3 = 0$. Let x be a neighbor of u_1 in $V(G_1)$ and $d_{G[V(G_1)]}(x) = 3$. It is easy to check that $\iota(G, B_2) \leq \frac{n}{5}$

if $G_{11}, G_{12}, \dots, G_{1t}, G_u \notin \{B_2, K_4, Y\}$. If $G_u = K_4$, by the proof of the case of $G_u = B_2$ in Subcase 2.1, we have $\iota(G, B_2) \leq \frac{n}{5}$. If $G_u = B_2$, then $\Delta(G) = 4$. Let $N(x) = \{x_1, x_2, x_3\}$ and let $d_{G[V(G_1)]}(x_2) = 3$. Define $G' = G[V(G_u) \cup \{u_1\} \cup V(G_1)] - \{u, u_1, x\}$. Note that G' contains B_2 when one of the following four cases is true. (1) u_2 is adjacent to x_1 and x_2 and u_3 is adjacent to x_1 . (2) u_2 is adjacent to x_2 and x_3 and u_3 is adjacent to x_3 . (3) u_3 is adjacent to x_1 and u_4 is adjacent to x_1 and x_2 . (4) u_3 is adjacent to x_3 and u_4 is adjacent to x_2 and x_3 . We can see that the proof methods are similar for the four cases. So let us just consider the first case. Then $G - N[u_2]$ is a connected graph with order $n - 5$ or the union of a connected graph with order $n - 6$ and a isolated vertex. By Lemma 8, $G - N[u_2]$ does not contain Y as an induced subgraph. Thus

$$\iota(G, B_2) \leq |\{u_2\}| + \iota(G - N[u_2], B_2) \leq \frac{n}{5}.$$

If $G_u = Y$, we have $\Delta(G) = 5$. By Lemma 8 (4), there exists $v \in V(G_u)$ such that $G_u - u - N[v] = P_3$. Denote $P = P_3$. Then $G[V(G_u) \cup \{u_1\} \cup V(G_1)] - \{u, u_1, x\} - N[v]$ contains B_2 if and only if $e(x_1, V(P)) = 3$ or $e(x_3, V(P)) = 3$. Suppose $e(x_1, V(P)) = 3$. Then $d(x_1) = 5$. By Lemma 8 (1), $G - N[x_1] \setminus \{x\}$ is a connected graph with order $n - 5$ and $G - N[x_1] \setminus \{x\} \neq Y$. Therefore,

$$\iota(G, B_2) \leq |\{x_1\}| + \iota(G - (N[x_1] \setminus \{x\}), B_2) \leq \frac{n}{5}.$$

Next suppose $G_u \notin \{B_2, K_4, Y\}$. Then $s_1 \geq 1$. Obviously,

$$\iota(G, B_2) \leq |\{u_1\}| + \iota(G_u, B_2) + \sum_{i=1}^t \iota(G_{1i} - N[u_1], B_2) \leq 1 + \frac{n - 5 - 4s_1}{5} \leq \frac{n}{5}$$

when $e(V(G_u), V(G_1) \setminus \{x\}) = 0$. If $e(V(G_u), V(G_1) \setminus \{x\}) > 0$, then $G[V(G_1) \cup \{u_1\} \cup V(G_u)] - \{u_1, x\} \neq B_2, K_4$. If $G[V(G_1) \cup \{u_1\} \cup V(G_u)] - \{u_1, x\} = Y$, by Lemma 8 (4), there exists v such that $G[V(G_1) \cup \{u_1\} \cup V(G_u)] - \{u, u_1, x\} - N[v] = P_3$. Then we have

$$\iota(G, B_2) \leq |\{u_1, v\}| + \sum_{i=1}^t \iota(G_{1i} - N[u_1], B_2) \leq 2 + \frac{n - (2 + 9 + 4s_1)}{5} \leq \frac{n}{5}.$$

Otherwise, $\iota(G, B_2) \leq |\{u_1\}| + \iota(G - u_1 - x, B_2) \leq 1 + \frac{n - (2 + 4s_1)}{5} \leq \frac{n}{5}$.

Subcase 2.4. $G_1 = B_2$ and only vertex of degree 2 of $V(G_1)$ is adjacent to u_1 . Let x be a neighbor of u_1 in $V(G_1)$ and $d_{G[V(G_1)]}(x) = 2$ and Let $d_{G[V(G_1)]}(x_2) = 2$. Note that the two remaining vertices of $V(G_1) \setminus \{x, x_2\}$ have degrees of 3 in G . First we prove the case of $u_1 \in N(x_2)$ and the case $u_1 \notin N(x_2)$ and $|N(x_2) \cap N(u)| = 1$. It is easy to check that

$\iota(G, B_2) \leq \frac{n}{5}$ if $G_2, \dots, G_t, G_u \notin \{B_2, K_4, Y\}$. If $G_u = B_2$ or K_4 , then $\Delta(G) \leq 4$ and hence

$$\iota(G, B_2) \leq |\{u_1\}| + \sum_{i=1}^t \iota(G_{1i} - N[u_1], B_2) \leq 1 + \frac{n-9-4s_1}{5} \leq \frac{n}{5}.$$

If $G_u = Y$, by Lemma 6 (3), $\iota(G_u, B_2) = \iota(V(G_u) \cup (V(G_1) \setminus \{x\}), B_2)$. Furthermore, by Lemma 8 (4), there exists $v \in V(G_u)$ such that $G_u - u - N[v] = P_3$. Then

$$\iota(G, B_2) \leq |\{u_1, v\}| + \sum_{i=1}^t \iota(G_{1i} - N[u_1], B_2) \leq 2 + \frac{n-5-9-4s_1}{5} \leq \frac{n}{5}.$$

Suppose $G_u \neq B_2, K_4, Y$. Then $s_1 \geq 1$. We have $\iota(G, B_2) \leq |\{u_1\}| + \frac{n-5-4s_1}{5} \leq \frac{n}{5}$. It remains the case of $u_1 \notin N(x_2)$ and $|N(x_2) \cap N(u)| \geq 2$.

In the end, we deal with the case of $u_1 \notin N(x_2)$ and $|N(x_2) \cap N(u)| \geq 2$, whether G^* is connected or not. Assume that $u_2, u_3 \in N(x_2)$. Denote by $G'' = G - \{u_2, u_3, x_1, x_2, x_3\}$. Obviously, if G'' is connected, then $G'' \notin \{B_2, K_4, Y\}$ and we have $\iota(G, B_2) \leq 1 + \frac{n-5}{5} = \frac{n}{5}$. If G'' is disconnected, it implies that $E(V(G_i), N(u)) = E(V(G_i), \{u_2, u_3\})$ for some $i \in \{2, 3, \dots, k\}$. Let us denote the components satisfying $E(V(G_i), N(u)) = E(V(G_i), \{u_2, u_3\})$ as $G''_{11}, G''_{12}, \dots, G''_{1t}$, $t \geq 1$ and let G''_u be the component contains u in G'' . Then $G'' = G''_{11} + G''_{12} + \dots + G''_{1t} + G''_u$. Clearly, $G''_u \neq K_4$. By Lemma 8 (3), $G''_u \neq Y$. And from the proof of above Subcase 2.4, we have $\iota(G, B_2) \leq \frac{n}{5}$ if any component of $\{G''_{11}, G''_{12}, \dots, G''_{1t}\}$ is B_2 . In other words, $G''_{1i} \notin \{B_2, K_4, Y\}$ for $i \in \{1, \dots, t\}$. Now, we distinguish two cases. If $G''_u \neq B_2$, then

$$\iota(G, B_2) \leq |\{x_2\}| + \iota(G''_u, B_2) + \sum_{i=1}^t \iota(G''_{1i}, B_2) \leq 1 + \frac{n-5}{5} = \frac{n}{5}.$$

If $G''_u = B_2$, then $\Delta(G) = 4$. Note that $|V(G''_u) \cup N[x_2]| = 9$ and $\{u_2\}$ is a B_2 -isolating set of $G[V(G''_u) \cup N[x_2]]$. Hence

$$\iota(G, B_2) \leq |\{u_2\}| + \iota(G - (V(G''_u) \cup N[x_2]) \setminus \{u_3\}, B_2) \leq 1 + \frac{n-8}{5} \leq \frac{n}{5}.$$

This completes the proof of Lemma 10. □

Proof of Theorem 1. From Lemma 2, we can see that the bound is sharp. Combining the results in Lemma 5, 7, 10, we obtain Theorem 1. □

Acknowledgement This research was supported Science and Technology Commission of Shanghai Municipality (STCSM) grant 18dz2271000.

References

- [1] J. A. Bondy, U.S.R. Murty, Graph Theory, in: GTM, vol. 244, Springer, 2008.
- [2] P. Borg, Isolation of cycles, *Graphs Combin.* 36 (2020), no. 3, 631–637.
- [3] P. Borg, K. Fenech, P. Kaemawichanurat, Isolation of k -cliques, *Discrete Math.* 343 (2020), no. 7, 111879, 5 pp.
- [4] P. Borg, P. Kaemawichanurat, Partial domination of maximal outerplanar graphs, *Discrete Appl. Math.* 283 (2020), 306–314.
- [5] Y. Caro, A. Hansberg, Partial domination - the isolation number of a graph, *Filomat* 31 (2017), no. 12, 3925–3944.
- [6] O. Favaron, P. Kaemawichanurat, Inequalities between the K_k -isolation number and the independent K_k -isolation number of a graph, *Discrete Appl. Math.* 289 (2021), 93–97.
- [7] O. Ore, *Theory of Graphs*, in: American Mathematical Society Colloquium Publications, vol. 38, American Mathematical Society, Providence, RI, 1962.
- [8] S. Tokunaga, T. Jiarasuksakun, P. Kaemawichanurat, Isolation number of maximal outerplanar graphs, *Discrete Appl. Math.* 267 (2019), 215–218.
- [9] D. B. West, *Introduction to Graph Theory*, Prentice Hall, Inc., 1996.
- [10] G. Zhang, B. Wu, $K_{1,2}$ -isolation in graphs, *Discrete Appl. Math.* 304 (2021), 365–374.