

A geometric approach to Lieb-Schultz-Mattis theorem without translation symmetry under inversion or rotation symmetry

Yuan Yao^{1,*} and Akira Furusaki^{1,2}

¹Condensed Matter Theory Laboratory, RIKEN CPR, Wako, Saitama 351-0198, Japan

²Quantum Matter Theory Research Team, RIKEN CEMS, Wako, Saitama 351-0198, Japan

(Dated: October 19, 2021)

We propose a geometric framework to study quantum many-body systems with discrete spin-rotation symmetries and lattice inversion or rotation symmetry, but without translation symmetry assumed. Under symmetry-twisting on a $(d-1)$ -dimensional plane, we find that any inversion-symmetric spin system possesses a doubly degenerate spectrum when it hosts a half-integer spin at the inversion-symmetric point in d -dimensional space. We also show that any rotation-symmetric spin model with a half-integer spin at the rotation center has a similar degeneracy under symmetry-twisting. We argue that these degeneracies imply either a gapless low-energy spectrum or degenerate ground states in the original systems before symmetry twisting — generalized inversional/rotational Lieb-Schultz-Mattis theorems without lattice translation symmetry imposed. The traditional Lieb-Schultz-Mattis theorems with translations also fit in the proposed framework.

Introduction.— Strongly interacting many-body systems are central topics in condensed matter and statistical physics. An important concept in the study of these systems is the Lieb-Schultz-Mattis (LSM) theorem [1] and its generalizations [2–5] that state an *ingappability* — either a ground-state degeneracy or a gapless spectrum above ground state(s) in the thermodynamic limit — of the systems respecting U(1) and translation symmetries with a fractional filling. Recent works study the ingappabilities from the interplay between translations and other symmetries, e.g., SU(2) [6, 7] and SU(N) [2, 8] symmetries, or even discrete subgroup symmetries [9–14].

The lattice translation symmetry is essential in these LSM-type theorems. However, several generalizations are recently proposed to systems with other lattice symmetries than translations [11–13, 15, 16]. Such extensions have been made in spin systems with rotation or inversion symmetry in low dimensions by employing lattice homotopy principles [15, 16] or even proven in a rigorous manner in one dimension [12, 13], and the ingappabilities of these systems are also related to quantum anomaly of field theories [7, 16, 17]. Nevertheless, higher dimensional generalizations are not well established and require further studies, e.g., inversions beyond one and two dimensions in a unifying way. Moreover, convincing lattice-based arguments are still lacking for systems with general lattice rotational and inversional symmetries.

In this Letter, we propose a geometric picture to study LSM-type ingappabilities for generic lattice systems without translation symmetries. As concrete cases, we first focus on spin systems with discrete spin-rotation symmetries (rather than the full SO(3) for generality) and site-centered inversions in arbitrary dimensions, and general lattice rotations in two dimensions. We consider closed geometry of finite lattice systems by identifying boundary spins, consistently with the lattice symmetries. We then twist the boundary condition using spin-rotation symmetries in a certain geometric pattern. Assuming

that bulk properties are insensitive to such twisting at the boundary [14, 18], we can extract ingappabilities of the original (untwisted) system from the interplay between the geometric pattern and lattice symmetries. The traditional LSM theorems with translation symmetries are also re-derived in our simple paradigm, which is thus expected to apply to broader classes of systems, e.g., with nonsymmorphic symmetries [11, 19].

Geometric patterns of symmetry-twisted boundary conditions.— In order to study bulk properties such as the ground-state degeneracies relevant to LSM-type arguments, we need to “close” the boundary to avoid edge/surface modes by imposing proper boundary conditions. Quite often, periodic boundary conditions (PBCs) are applied in most Hamiltonians on square lattices with sizes L_i by identifying sites $\vec{r}_i \sim \vec{r}_i + L_i$. However, the situation differs here because, except for certain symmetry requirements, we do not specify any concrete form of Hamiltonian in advance. It implies that the chosen boundary condition should be compatible with all possible Hamiltonians respecting the required symmetries. For instance, when the imposed symmetry is the inversion $\vec{r} \rightarrow -\vec{r}$ about the origin $\vec{r} = \vec{0}$ without translation symmetries required, an inversion-symmetric Hamiltonian that cannot be closed by PBC is shown in FIG. 1 (a); PBC is inapplicable when only the inversion is imposed [20].

When the system also possesses internal symmetries, such as U(1) or discrete \mathbb{Z}_m symmetry, we can “mix” them into the boundary condition as follows. Let us take \mathbb{Z}_m twisting in tight-binding models with charged operator c_i^\dagger for illustration. In one dimension, $(2\pi n/m)$ -angle twisting with an integer n can be done by substituting the boundary hopping term $[c_L c_1^\dagger + \text{h.c.}]$ with $[\exp(i2\pi n/m)c_L c_1^\dagger + \text{h.c.}]$. For general coupling terms involving multiple sites crossing the bond between sites L and 1, we can apply the U(1) transformation $c_j^\dagger \rightarrow$




FIG. 1. (a) An inversion-symmetric Hamiltonian with coupling constants $\pm J$ has a boundary that cannot be closed by PBC. (b) The boundary closing by identifications $\vec{r} \sim -\vec{r}$ is compatible with any inversion-symmetric Hamiltonian. The resultant manifold in the continuum limit is a real projective plane \mathbb{RP}^2 .




FIG. 2. Two equivalent representations of $n \in \mathbb{Z}_m$ twisting in one dimension (a) and higher dimensions (b). We will use either of them whichever is convenient.

$\exp(i2\pi n/m)c_j^\dagger$ only on the sites j on the “right” of the bond [21]. This twisting has two equivalent geometric presentations in FIG. 2 (a). The first one is the obvious type that we can make such a boundary bond in bold with a direction indicating the “right”. The second type is the dual one: an arrowed dot is drawn on this bond to indicate that interaction terms across this dot get twisted by the phase specified with the integer n . These two dual pictures are almost the same in one dimension, but their differences will be clear in higher dimensions, making one of them more convenient than the other depending on situations.

In two dimensions, we can similarly twist all bonds crossing the boundary line by the internal symmetries as shown in FIG. 2 (b). The dual picture is exactly the boundary line which transversally intersects those bond centers, and the line is labelled by $n \in \mathbb{Z}_m$ with oriented arrows. In general d dimensions, we twist the bonds intersected by the $(d-1)$ -dimensional boundary hypersurface. The dual picture has a certain hypersurface with an orientation and its symmetry label, which is a special case of the Poincaré duality.

Since the boundary of the boundary is always empty, the boundary line/face (transversal to all the twisted bonds) must be closed themselves, i.e., closed loops/surfaces. It exactly means that the total net symmetry twisting around any plaquette must be zero. In the dual picture of symmetry twisting, we can show that this mixture of twisting into the boundary condition does




FIG. 3. The $n \in \mathbb{Z}_m$ twisting can be moved/deformed by a gauge transformation: $\mathcal{H}_{\text{tw}} \rightarrow U^{-1}\mathcal{H}_{\text{tw}}U$ with $U = \exp(i2\pi c_{1(A)}^\dagger c_{1(A)} n/m)$.

not introduce any boundary modes at the twisted bonds. It is a direct consequence of the following gauge invariance. We can use a gauge freedom to rename the charged operator by a unitary operator. For example in one dimension, the transformation $c_1^\dagger \rightarrow c_1^\dagger \exp(-i2\pi n/m)$ with a unitary operator $U = \exp(i2\pi c_1^\dagger c_1 n/m)$ undoes the earlier twisting of the bond between sites L and 1 . However, it creates a new twisting between sites 1 and 2 , so effectively the twisting is moved by one site as shown in FIG. 3. Thus we can move the twisting freely by unitary transformations, and its position has no physical significance, which implies the absence of physical boundary modes there.

The above argument using gauge degrees of freedom applies to higher dimensions as shown in FIG. 3: the loop/surface transversal to the twisted bonds can be deformed arbitrarily by local unitary transformation, thereby unchanging any energy-spectrum property. It should be noted that such deformations are performed only locally in that they cannot make a non-contractible loop into a contractible one. These gauge properties of twisting induce the concept of bulk insensitivity. It states that the symmetry twisting does not affect the low-energy bulk spectrum, such as the gap, when the original pre-twisted system has a unique gapped ground state with a large system size. The bulk insensitivity is physically reasonable although not proven in general. Nevertheless, it is proven under certain assumptions on excited spectra in one dimension [18] and justified by a quantum-transfer-matrix formalism [14] in higher dimensions. In the following discussion, we will assume such a spectrum insensitivity to twisted boundary conditions to derive LSM-type theorems.

Inversional LSM theorems.— Let us consider a spin chain \mathcal{H}_0 with length L respecting $\mathbb{Z}_2 \times \mathbb{Z}_2$ discrete spin-rotation symmetry and a site-centered inversion symme-

try:

$$R_x^\pi = \exp\left(i\pi \sum_{j=1}^L S_j^x\right); R_z^\pi = \exp\left(i\pi \sum_{j=1}^L S_j^z\right); \quad (1)$$

$$I : j \rightarrow -j \text{ with } I^{-1} \vec{S}_j I = \vec{S}_{-j}, \quad (2)$$

where the spin operator $\vec{S}_j \equiv [S_j^x, S_j^y, S_j^z]$ is not necessarily of the same spin representation for all site j , except for those related by the inversion symmetry, and we denote the spin of $\vec{S}_{j=0}$ at the inversion center as spin- s . Considering the inversion symmetry, we can close the chain by identifying the boundary sites at the two ends that are related by inversion to each other [22]. We then twist the chain by \mathbb{Z}_2 symmetry: R_x^π . Since the twisting position can be changed freely by gauge invariance, we put the twisting on the bond between the sites -1 and 0 and denote the twisted Hamiltonian as \mathcal{H}_{tw} . A typical example is the XYZ model:

$$\begin{aligned} \mathcal{H}_{\text{tw}}^{\text{XYZ}} = & \sum_{j \neq 0} (J_X S_j^x S_{j+1}^x + J_Y S_j^y S_{j+1}^y + J_Z S_j^z S_{j+1}^z) \\ & - J_X S_{-1}^x S_0^x - J_Y S_{-1}^y S_0^y + J_Z S_{-1}^z S_0^z. \end{aligned} \quad (3)$$

The twisted Hamiltonian \mathcal{H}_{tw} is represented by FIG. 4 (up-left), where the arrows are irrelevant in case of \mathbb{Z}_2 -twisting. It still respects $\mathbb{Z}_2 \times \mathbb{Z}_2$ symmetry but explicitly breaks the inversion symmetry I since the twisting is moved to the bond between sites 0 and 1 by I in FIG. 4 (up-right). Nevertheless, we can do a gauge transformation $r_{z,0}^\pi \equiv \exp(i\pi S_{j=0}^z)$ to map it back. It implies that \mathcal{H}_{tw} respects the following inversion symmetry modified by the gauge transformation:

$$\tilde{I} \equiv I \cdot r_{z,0}^\pi; [\tilde{I}, \mathcal{H}_{\text{tw}}] = 0. \quad (4)$$

Therefore, \mathcal{H}_{tw} possesses \tilde{I} , R_x^π and R_z^π symmetries, but two of them obey the algebra

$$\tilde{I} R_x^\pi = (-1)^{2s} R_x^\pi \tilde{I}, \quad (5)$$

where the factor $(-1)^{2s}$ comes from the commutator $r_{z,0}^\pi r_{x,0}^\pi = (-1)^{2s} r_{x,0}^\pi r_{z,0}^\pi$. It implies that \mathcal{H}_{tw} must have a doubly degenerate energy spectrum if $s = 1/2, 3/2, \dots$, i.e., a half-integer spin at the origin. Thus, the original Hamiltonian \mathcal{H}_0 cannot have a unique gapped ground state when $s = 1/2, 3/2, \dots$ — either a gapless or degenerate ground states should result in the thermodynamic limit, because, otherwise, the bulk insensitivity would mean that \mathcal{H}_{tw} also had a unique gapped ground state that contradicts the above double degeneracy.

This approach can be generalized to arbitrary higher dimensions with $\mathbb{Z}_2 \times \mathbb{Z}_2$ and inversion I about the origin $\vec{r} \rightarrow -\vec{r}$, but the boundary closing needs more consideration. For instance, in two dimensions, PBC is not consistent as shown in FIG. 1 (a), and the only sensible way of closing is identifying boundary sites $\vec{S}_{\vec{r}} \sim \vec{S}_{-\vec{r}}$ in FIG. 1 (b) that is compatible with any inversion-symmetric Hamiltonian \mathcal{H}_0 . Then we twist \mathcal{H}_0 by \mathbb{Z}_2 :




FIG. 4. The twisted Hamiltonians \mathcal{H}_{tw} on the left side are transformed by inversions to $I^{-1} \mathcal{H}_{\text{tw}} I$ on the right side. $I^{-1} \mathcal{H}_{\text{tw}} I$ can be transformed back by the gauge transformation: $(r_{z,0}^\pi)^{-1} (I^{-1} \mathcal{H}_{\text{tw}} I) r_{z,0}^\pi = \mathcal{H}_{\text{tw}}$, which implies \mathcal{H}_{tw} is symmetric under \tilde{I} in Eq. (4). The sites on the boundary are identified by $\vec{r} \sim -\vec{r}$, so the continuum limit of lattice in d dimensions is a real projective hyperplane $\mathbb{R}P^d$.

R_z^π as in FIG. 4 (middle-left) with a series of twisted bonds or, in the dual picture, a closed loop. The twisted Hamiltonian \mathcal{H}_{tw} also softly breaks I and has the modified inversion \tilde{I} in the same form as Eq. (4). Therefore, by employing a similar argument as above, we see that \mathcal{H}_0 cannot have a unique gapped ground state when the spin s at $\vec{r} = 0$ is half integer.

The twisted Hamiltonian \mathcal{H}_{tw} in three dimensions is sketched in FIG. 4 (bottom), where the dual picture is suppressed for clarity. The twisting satisfies the closed-form condition that the total twisting around any plaquette is zero. The same result as in the lower-dimensional cases follows since \mathcal{H}_{tw} also respects \tilde{I} . The geometric situation in arbitrary dimensions is that when we perform the inversion transformation, the spin at $\vec{r} = 0$ is wrapped by the hypersurface (in the dual picture) spanned by the centers of twisted bonds together with the hypersurface inverted by I . The wrapping is exactly cancelled by the gauge transformation $r_{z,0}^\pi$, which gives the commutator as in Eq. (5). It follows that \mathcal{H}_0 respecting $\mathbb{Z}_2 \times \mathbb{Z}_2$ and I is ingappable in arbitrary dimensions when s is not an integer.

Rotational LSM theorems. — In two dimensions, the inversion $\vec{r} \rightarrow -\vec{r}$ is equivalent to lattice rotation by 180 degrees. It is natural to generalize I to be N -fold rotations C_N generated by a lattice rotation by $2\pi/N$ around




FIG. 5. The only boundary closing compatible with any C_3 -symmetric Hamiltonian is to identify three boundary edges as above. The continuum limit is a pseudo-projective plane of order 3 with a three-page intersection (up-right) which is realized by a simple identification of three edges at the lattice scale (bottom-right).

the origin $\vec{r} = \vec{0}$, e.g., C_3 on a honeycomb lattice:

$$C_N : \vec{r} \rightarrow \begin{bmatrix} \cos(2\pi/N) & -\sin(2\pi/N) \\ \sin(2\pi/N) & \cos(2\pi/N) \end{bmatrix} \vec{r}. \quad (6)$$

However, even with translation symmetries, spin-1/2 systems on the honeycomb lattice admits a unique gapped ground state with C_3 and $SO(3)$ spin-rotation symmetries [7, 23]. It motivates us to consider $SU(3)$ “spin” degrees of freedom or generally $SU(N)$ “spins” obeying an $\mathfrak{su}(N)$ algebra [24, 25]. However, the only condition to be used here is that there is a discrete “spin”-rotation symmetry $\mathbb{Z}_N \times \mathbb{Z}_N$ (also known as shift symmetries [26, 27]) by $V_N = \prod_{\vec{r}} v_{N,\vec{r}}$ and $W_N = \prod_{\vec{r}} w_{N,\vec{r}}$ generalizing R_z^π and R_x^π of the $N = 2$ case in Eq. (1) [28]. There is a number b called Young-tableau box number [29], analogous to the spin s above, to characterize the $SU(N)$ “spin” at the origin by the following commutator:

$$v_{N,\vec{r}=0} w_{N,\vec{r}=0} = \exp\left(i \frac{2\pi b}{N}\right) w_{N,\vec{r}=0} v_{N,\vec{r}=0}. \quad (7)$$

Thus $b = 2s$ when $N = 2$ and we do not need detailed forms of $v_{N,\vec{r}}$ and $w_{N,\vec{r}}$.

Let us first consider an $SU(3)$ “spin” system on a honeycomb lattice respecting C_3 and $\mathbb{Z}_3 \times \mathbb{Z}_3$ symmetries with a “spin” of b at the rotation center $\vec{r} = 0$. In closing the boundary, PBC is incompatible for a similar reason as before, and the only consistent way is to identify or paste the boundary sites by C_3 as in FIG. 5 (left). In the continuum limit, the resultant space is called a pseudo-projective plane of order 3 and it has three-fold intersection at the boundary closing, which is realized at the lattice scale by gluing the three edges together, i.e., boundary-site identifications $\vec{r} \sim C_3(\vec{r})$ as in FIG. 5. Then we twist the Hamiltonian by \mathbb{Z}_3 , V_3 symmetry, and deform the twisting configuration as FIG. 6 (up-left). The twisted Hamiltonian \mathcal{H}_{tw} preserves $\mathbb{Z}_3 \times \mathbb{Z}_3$ symmetry but breaks C_3 . Nevertheless, as shown in FIG. 6 (up-




FIG. 6. The twisted Hamiltonians \mathcal{H}_{tw} is presented on the left side, which is transformed to $C_M^{-1} \mathcal{H}_{tw} C_M$ on the right side and can be transformed back by the gauge transformation: $v_{N,0}^{-\tau} C_M^{-1} \mathcal{H}_{tw} C_M (v_{N,0})^\tau = \mathcal{H}_{tw}$ with $\tau = N/\text{g.c.d.}(M, N)$. The twisting numbers should be understood as mod- N , so a p -twisting is equal to a $(N-p)$ -twisting with the opposite arrow direction.

right) it preserves a modified rotation $\tilde{C}_3 \equiv C_3 v_{N=3,\vec{r}=0}$, which implies a nontrivial commutator between symmetries \tilde{C}_3 and W_3 :

$$\tilde{C}_3 W_3 = \exp\left(i \frac{2\pi b}{3}\right) W_3 \tilde{C}_3. \quad (8)$$

Thus, when 3 does not divide b , \mathcal{H}_{tw} must possess a triply degenerate spectrum and the bulk insensitivity implies that the original Hamiltonian \mathcal{H}_0 cannot have a unique gapped ground state when the central “spin” b is not a multiple of 3.

For general N , we consider an $SU(N)$ spin system with $\mathbb{Z}_N \times \mathbb{Z}_N$ and general C_M symmetries, where M and N are unnecessarily equal and there is an $SU(N)$ spin of box b at $\vec{r} = 0$. As shown in FIG. 6 (bottom-left), we close the lattice to form a pseudo-projective plane of order M and introduce M lines of V_N -twisting to the τ -th power radiating from $\vec{r} = 0$ with $\tau \equiv N/\text{g.c.d.}(M, N)$ to have a well-defined twisting, i.e., twisting around any plaquette is 0 mod N . Here “g.c.d.” denotes the greatest common divisor. The modified $\tilde{C}_M = C_M (V_N)^\tau$ is respected by \mathcal{H}_{tw} and

$$\tilde{C}_M W_N = \exp\left(i \frac{2\pi b}{\text{g.c.d.}(M, N)}\right) W_N \tilde{C}_M. \quad (9)$$

Together with the bulk insensitivity, the above phase factor gives the rotational LSM theorem: when a two-dimensional $SU(N)$ system preserves $\mathbb{Z}_N \times \mathbb{Z}_N$ and C_M symmetries, a unique gapped ground state is forbidden when the box number b of the “spin” at the rotational center is not a multiple of $\text{g.c.d.}(M, N)$. It explains why




FIG. 7. The twisted Hamiltonians \mathcal{H}_{tw} is presented on the left side, where we suppress the arrows and numbers of twisting for simplicity. \mathcal{H}_{tw} is transformed by the translation T_1 and can be mapped back by a following gauge transformation $v_{N, \vec{C}}$ with \vec{C} indicated above. This argument is generalizable to arbitrary dimensions.

C_3 and spin-rotation symmetries cannot ensure any ingappability on the spin-1/2 honeycomb lattice [7, 23] where $b = 2s \in \mathbb{Z}$ and $N = 2, M = 3$ coprime.

A general framework and conclusions. — In this work, we have focused on LSM-type theorems for lattice Hamiltonians without any translation symmetry imposed. We have developed a geometric approach using symmetry-twisted boundary conditions to studying ingappability of spin systems or $SU(N)$ generalizations from rotation, inversion and discrete spin-rotation symmetries. In fact, such a geometric approach can also be applied to traditional LSM theorems with translation symmetries and b boxes per unit cell reviewed in FIG. 7 [14], where we close the lattice by a tilted boundary condition which is compatible with translations [30, 31]. The modified translation symmetry $\tilde{T}_1 = T_1 v_{N, \vec{C}}$, where \vec{C} is the spin position imposed by the gauge transformation $v_{N, \vec{C}}$, is preserved by the twisted Hamiltonian and has a commutator with W_N symmetry:

$$\tilde{T}_1 W_N = \exp\left(i2\pi \frac{b}{N}\right) W_N \tilde{T}_1. \quad (10)$$

When b is indivisible by N , the LSM-type ingappability is concluded from the bulk insensitivity.

Furthermore, our geometric paradigm suggests the following general framework in arbitrary dimensions: 1) we close the lattice tentatively in all compatible ways with lattice symmetries; 2) then we do the symmetry twisting on the bonds transversal to a *non-contractible* hypersurface of co-dimension 1; 3) finally, the ingappability is extracted from the algebra of the modified lattice symmetry. Thus, it is not restricted to the symmetries discussed in this work and we expect that it can be applied to more general settings, e.g., the systems with nonsymmorphic symmetries [11, 19] and other rotation-like symmetries such as space dihedral symmetries [7, 15, 16].

Finally, we note that it is the first time that pseudo-projective planes appear in LSM-type theorems. The

related quantum anomaly of field theories on such self-intersecting *non-manifolds* and various fermionic extensions [32, 33] can be of future interest.

The authors thank Masaki Oshikawa, Yasuhiro Tada, and Yunqin Zheng for stimulating discussions. This work was supported in part by JSPS KAKENHI (Grant No. 19K03680) and JST CREST (Grant No. JPMJCR19T2).

* yuan.yao@riken.jp

- [1] E. Lieb, T. Schultz, and D. Mattis, *Ann. Phys.* **16**, 407 (1961).
- [2] I. Affleck and E. H. Lieb, *Lett. Math. Phys.* **12**, 57 (1986).
- [3] M. Oshikawa, M. Yamanaka, and I. Affleck, *Phys. Rev. Lett.* **78**, 1984 (1997).
- [4] M. Oshikawa, *Phys. Rev. Lett.* **84**, 1535 (2000).
- [5] M. B. Hastings, *Phys. Rev. B* **69**, 104431 (2004).
- [6] S. C. Furuya and M. Oshikawa, *Phys. Rev. Lett.* **118**, 021601 (2017).
- [7] M. A. Metlitski and R. Thorngren, *Phys. Rev. B* **98**, 085140 (2018).
- [8] Y. Yao, C.-T. Hsieh, and M. Oshikawa, *Phys. Rev. Lett.* **123**, 180201 (2019).
- [9] X. Chen, Z.-C. Gu, and X.-G. Wen, *Phys. Rev. B* **83**, 035107 (2011).
- [10] Y. Fuji, *Phys. Rev. B* **93**, 104425 (2016).
- [11] H. Watanabe, H. C. Po, A. Vishwanath, and M. Zaletel, *Proc. Natl. Acad. Sci. USA* **112**, 14551 (2015).
- [12] Y. Ogata and H. Tasaki, arXiv preprint arXiv:1808.08740 (2018).
- [13] Y. Ogata, Y. Tachikawa, and H. Tasaki, arXiv preprint arXiv:2004.06458 (2020).
- [14] Y. Yao and M. Oshikawa, *Phys. Rev. Lett.* **126**, 217201 (2021).
- [15] H. C. Po, H. Watanabe, C.-M. Jian, and M. P. Zaletel, *Phys. Rev. Lett.* **119**, 127202 (2017).
- [16] D. V. Else and R. Thorngren, *Phys. Rev. B* **101**, 224437 (2020).
- [17] M. Cheng, M. Zaletel, M. Barkeshli, A. Vishwanath, and P. Bonderson, *Phys. Rev. X* **6**, 041068 (2016).
- [18] H. Watanabe, *Phys. Rev. B* **98**, 155137 (2018).
- [19] S. A. Parameswaran, A. M. Turner, D. P. Arovas, and A. Vishwanath, *Nature Physics* **9**, 299 (2013).
- [20] If we also impose lattice translation symmetries, then PBC is permitted. It is consistent since the Hamiltonian in FIG. 1 breaks translations thereby out of our consideration.
- [21] Because the range l of interaction terms $l \ll L$, there is a well-defined “left/right” only locally around the boundary.
- [22] This boundary closing has an advantage that the resultant chain has an even number of sites. If, otherwise, there are an odd number of sites, the spectrum must be degenerate once the spin at the inversion center has half-integer spins by a Kramers-type argument in quantum mechanics. However, such a degeneracy due to a single-body effect is already well-known and not in our interest, so we need to avoid this special situation.
- [23] I. Kimchi, S. A. Parameswaran, A. M. Turner, F. Wang, and A. Vishwanath, *Proc. Natl. Acad. Sci. USA* **110**,

- 16378 (2013).
- [24] I. Affleck and F. D. M. Haldane, *Phys. Rev. B* **36**, 5291 (1987).
- [25] I. Affleck, *Nucl. Phys. B* **305**, 582 (1988).
- [26] H. Shimizu and K. Yonekura, *Phys. Rev. D* **97**, 105011 (2018).
- [27] Y. Tanizaki and Y. Kikuchi, *JHEP* **2017**, 102 (2017).
- [28] The only basis-independent property of $v_{N,\vec{r}}$ and $w_{N,\vec{r}}$ is the commutator such as Eq. (7). A matrix expression of $v_{N,\vec{r}}$ and $w_{N,\vec{r}}$ can be found, for example, in Eq. (2.2) in Ref. [34].
- [29] H. Georgi, *Lie Algebras In Particle Physics* (Boca Raton: CRC Press, 2000).
- [30] Y. Yao and M. Oshikawa, *Phys. Rev. X* **10**, 031008 (2020).
- [31] S. C. Furuya and Y. Horinouchi, *Phys. Rev. B* **100**, 174435 (2019).
- [32] M. Cheng, *Phys. Rev. B* **99**, 075143 (2019).
- [33] Ö. M. Aksoy, A. Tiwari, and C. Mudry, *Phys. Rev. B* **104**, 075146 (2021).
- [34] Y. Yao and A. Furusaki, *JHEP* **2021**, 285 (2021).