

Gallai's path decomposition in planar graphs

Alexandre Blanché¹, Marthe Bonamy¹, and Nicolas Bonichon¹

¹CNRS, LaBRI, Université de Bordeaux, Bordeaux, France

June 22, 2022

Abstract

In 1968, Gallai conjectured that the edges of any connected graph with n vertices can be partitioned into $\lceil \frac{n}{2} \rceil$ paths. We show that this conjecture is true for every planar graph. More precisely, we show that every connected planar graph except K_3 and K_5^- (K_5 minus one edge) can be decomposed into $\lfloor \frac{n}{2} \rfloor$ paths.

1 Introduction

Given a finite undirected connected graph G , a k -path decomposition of G is a partition of the edges of G into k paths. In 1968, Gallai stated this simple but surprising conjecture [1]: every graph on n vertices admits a $\lceil \frac{n}{2} \rceil$ -path decomposition. Gallai's conjecture is still unsolved as of today, and has only been confirmed on very specific graph classes: graphs in which each vertex has degree 2 or 4 [2], graphs whose vertices of even degree induce a forest [3], graphs for which each block of the subgraph induced by the vertices of even degree is triangle-free with maximum degree at most 3 [4, 5], series-parallel graphs [6], or planar 3-trees [7]. Bonamy and Perrett confirmed the conjecture on graphs with maximum degree at most 5 [8], and Chu, Fan and Liu for graphs of maximum degree 6, under the condition that the vertices of degree 6 form an independent set [9]. Recently, Botler, Jiménez and Sambinelli proved that the conjecture is true in the case of triangle-free planar graphs [10].

We confirm the conjecture for the whole class of planar graphs.

Theorem 1.1. *Every connected planar graph on n vertices can be decomposed into $\lceil \frac{n}{2} \rceil$ paths.*

Our result is actually a little more precise. An *odd semi-clique* is a graph obtained from a clique on $2k + 1$ vertices by deleting at most $k - 1$ edges. In 2016, Bonamy and Perrett asked the following question [8, Question 1.1]: does every connected graph on n vertices that is not an odd semi-clique admit a $\lfloor \frac{n}{2} \rfloor$ -path decomposition? In other words, is it possible to save one path in the decomposition when n is odd?

We answer this question positively for planar graphs. Only two odd semi-cliques are planar: the triangle K_3 and K_5 minus one edge, which we denote by K_5^- (see Figure 1). We can therefore state our result as follows:

Theorem 1.2. *Every connected planar graph on n vertices, except K_3 and K_5^- , can be decomposed into $\lfloor \frac{n}{2} \rfloor$ paths.*




Figure 1: A 2-path decomposition of K_3 (left) and a 3-path decomposition of K_5^- (right)

To prove this result, we proceed with a standard approach for coloring problems, by considering a planar graph that is a counterexample to our theorem and is vertex-minimum with respect to this property. We prove that such a *minimum counterexample* (MCE) does not contain a certain set of configurations, by providing for each of these configurations a reduction rule that takes advantage of the properties of the MCE and yields a contradiction. This technique is widely used in the literature on graph coloring and on Gallai's conjecture ([8, 9, 10, 11]). More precisely, these reducible configurations deal with vertices of small degree (at most 5), and after showing that our MCE cannot contain any of these configurations (Lemma 3.1, p. 3, in Section 3), we know that it has mostly vertices of degree at least 6. We finally use Euler's formula and structural arguments to prove that there is no such graph (Lemma 5.1, p. 3, in Section 5).

2 Preliminaries

The graphs we consider are undirected, finite and without loops or multi-edges. In a graph $G = (V, E)$ and given a subset $X \subseteq V$ of vertices, we denote $N(X) = \{v \in V \mid \exists u \in X, uv \in E\}$ the *neighborhood* of X . The subgraph of G **induced by** X is the graph with vertices X and edges $\{uv \in E \mid u, v \in X\}$, denoted $G[X]$. An *i -vertex* is a vertex v of degree i in G , and we denote $i = d_G(v)$.

A *path* of a graph G is a finite sequence of edges $(v_1v_2, v_2v_3, \dots, v_{k-1}v_k)$ of G , denoted $P = (v_1, v_2, \dots, v_k)$, and such that all v_1, \dots, v_k are pairwise distinct. We denote $V(P) = \{v_1, \dots, v_k\}$. We say that a vertex **touches** P if it belongs to $V(P)$. The *internal vertices* of P are v_2, \dots, v_{k-1} and its *ends* are v_1, v_k . We say that two paths are *internally disjoint* if they have no internal vertex in common. We also call P a (v_1, v_k) -*path*. A *section* $Q = (v_i, \dots, v_j)$ of a path $P = (v_1, \dots, v_k)$, $1 \leq i < j \leq k$ is a subsequence of consecutive edges of P . The *length* $l(P)$ of a path $P = (v_1, \dots, v_k)$ is k . We say that two vertices u, v are *at distance* k if the minimum length of a (u, v) -path is k . For simplicity, we denote some paths by a sequence of subpaths: if $(Q_i)_{i=1..k}$ is a family of edge-disjoint paths, we may denote the path $P = (Q_1, Q_2, \dots, Q_k)$; we may also denote it by an alternation of vertices and subpaths: $P = (v_1, Q_1, v_2, \dots, v_{k-1}, Q_k, v_k)$.

A path $P = (v_1, \dots, v_k)$ of a graph G has a *chord* if there is an edge $v_iv_j \in E(G)$ such that v_i and v_j belong to $V(P)$ but are not consecutive in P . We say that P is *chordless* if it has no chord.

A *cycle* is a finite sequence of edges (v_1v_2, \dots, v_kv_1) , denoted $C = (v_1, v_2, \dots, v_k)$. Its *length* is k .

Given two graphs G, K , a *K -subdivision* in G is a subgraph of G that is isomorphic to the graph obtained by replacing the edges of K by internally disjoint paths of positive length. The *roots* of the subdivision are the images in G of the vertices of K , and the *paths* of the subdivision are the images in G of the edges of K . We say that a K -subdivision is *chordless* if its paths are chordless. Given a set U of vertices of a graph G , we say that a K -subdivision of

G is *rooted on* U if its roots are exactly the vertices of U . Given a 4-family $U = \{u_1, u_2, u_3, u_4\}$, the subdivisions we use in this paper are K_4 -subdivisions and C_{4+} -subdivisions rooted on U , where K_4 is the clique on 4 vertices and C_{4+} is the graph made up of a cycle on 4 vertices $U = \{x_1, x_2, x_3, x_4\}$ and two additional parallel edges x_1x_3, x_2x_4 . We say that two subdivisions S, S' have the same *type* if S, S' are both K_4 -subdivisions or both C_{4+} -subdivision. For $u_i, u_j \in U$, we denote $u_i \sim u_j$ the (u_i, u_j) -path of a K_4 - or C_{4+} -subdivision when there is no ambiguity.

To talk about path decompositions, the framework we adopt deals with paths as colors. We say that a (k) -coloring of a graph $G = (V, E)$ is a function $c : E \rightarrow \llbracket 1, k \rrbracket$, with $k \in \mathbb{N} \setminus \{0\}$, such that the edges with the same color form a path. We denote $|c| = k$ the number of colors used in the coloring. We say that a color x *induces* a path P if $E(P) = c^{-1}(\{x\})$. A *good coloring* of a connected graph G with n vertices is a $\lfloor \frac{n}{2} \rfloor$ -coloring. A color x *ends* on a vertex v if the path induced by the color x ends on v .

To describe the 2-coloring of a K_4 -subdivision or a C_{4+} -subdivision S rooted on $\{u_1, u_2, u_3, u_4\}$, we use the notation $\{red = (u_{i_1} \rightarrow u_{i_2} \rightarrow u_{i_3} \rightarrow u_{i_4}), blue = (u_{j_1} \rightarrow u_{j_2} \rightarrow u_{j_3} \rightarrow u_{j_4})\}$ for i_1, i_2, i_3, i_4 and j_1, j_2, j_3, j_4 two permutations of 1, 2, 3, 4. This notation means that we decompose S into two paths $P_{red} = (u_{i_1} \sim u_{i_2}, u_{i_2} \sim u_{i_3}, u_{i_3} \sim u_{i_4})$ and $P_{blue} = (u_{j_1} \sim u_{j_2}, u_{j_2} \sim u_{j_3}, u_{j_3} \sim u_{j_4})$. The decompositions of this kind that we consider throughout the paper feature each edge of S exactly once. We sometimes insert vertices that are not roots in between the vertices from U to describe the paths we take more precisely: the notation $(\dots \rightarrow u_i \rightarrow v \rightarrow u_j \rightarrow \dots)$ means that the path $u_i \sim u_j$ considered is the (u_i, u_j) -path of the subdivision that contains the vertex v .

We call *minimum counterexample* (MCE) a planar graph that is distinct from K_3 or K_5^- , which does not admit a good coloring and is vertex minimum with respect to this property.

Given a planar graph G , a *2-family* is a set U of two vertices of G of degree at most 4. A *4-family* is a set of four 5-vertices. We say that a graph G with a 4-family U is *almost 4-connected* w.r.t. U if it does not contain a 3-cut $A = \{a_1, a_2, a_3\} \subseteq V(G)$ that separates two vertices $u, u' \in U$ or two neighbors of some vertex $u \in U \cap A$.

3 Minimal counterexamples properties

A *configuration* C is a property satisfied by a graph. The configurations we consider are usually defined locally as specifications on the neighborhoods of some vertices in the graph, and the possible presence (or absence) of some paths between them. By abuse of language, we say that a graph G *contains* a configuration C when G satisfies the properties of C .

The following lemma is the main result of this section and the next, and helps us prove the main theorem in Section 5. We prove the first property as Lemma 3.2 at the end of this section and the second property as Lemma 4.1 (p. 35) in Section 4.

Lemma 3.1. *An MCE does not contain any of the following configurations:*

- (C_I) : a 2-family;
- (C_{II}) : a 4-family with respect to which the MCE is almost 4-connected.

To prove that an MCE G does not contain a configuration C , we proceed by contradiction: we assume that G does contain the configuration, then use its nature of MCE to build a good coloring of G . The only configurations C we consider are specifications of (C_I) and (C_{II}) . To build a good coloring of G , we start by removing the two or four *special vertices* forming the 2- or 4-family of C , along with their incident edges. Depending on the case, we add or remove some edges from the obtained graph, and define the resulting graph as the *reduced graph* G' . Ideally, this graph is connected and not a K_3 or K_5^- : in this case, since it is smaller than the

minimum counterexample G , it admits a good coloring, that we call *pre-coloring*. We then adapt this pre-coloring to the original graph G , and since G has two or four more vertices than G' , we may use 1 additional color (for (C_I)) or 2 (for (C_{II})) to color G . The cases where G' is disconnected are equally easy, unless G' has some K_3 or K_5^- connected components, in which case a complementary method is used to combine a “bad” coloring of these components with the rest of the coloring.

In the present section, we split the configuration (C_I) into simpler, specified cases, and for each of them provide a *reduction rule* describing the adaptation of the pre-coloring to a good coloring of G . The general method consists in fixing a shortest path P between the two special vertices. The edges of P are removed alongside the special vertices and possibly some additional edges, in order to obtain the reduced graph G' . In G , the edges of P are then colored with the extra color. This has the advantage of having an end of this extra color on each of the two special vertices, and the path induced by the extra color can be conveniently extended to help cover all the edges of G that were missing in G' .

This is the method used when the two special vertices are sufficiently distant from each other, and in this case the adapting methods for each are independent and can be defined separately. Figure 2 depicts a (C_I) configuration where the two special vertices u_1, u_2 are at distance at least 3, and their neighborhoods are taken care of with two independent *elementary partial rules* that when combined form a complete reduction rule. The extra color induces the path P in red. These rules are defined in Subsection 3.1 on page 20.




Figure 2: A (C_I) reduction rule made up of two elementary partial rules

When instead the special vertices are too close to each other and share some neighbors, two elementary partial rules would interfere with each other and possibly create some cycles in the decomposition. In these cases, we discard the shortest path and instead use a custom reduction rule to treat the neighborhoods of both special vertices at once. These rules correspond to rules $(a), (b), \dots, (u)$ in Subsection 3.1 on page 8. Figure 3 features an example of such a rule, and the extra color is again represented in red as the path P .




Figure 3: A (C_I) reduction rule treating close special vertices

The next subsection introduces the reduction rules we need to treat the (C_I) configurations. The generalization to this method for (C_{II}) configurations is presented at the beginning of Section 4, on page 35.

3.1 Reduction rules for (C_I) configurations

This notion of reducible subgraph has been used in previous works [5, 7, 8, 10, 11]. We present here a formalism appropriate for our subgraphs.

A reduction rule $\mathcal{R} = (\mathcal{C}, f^r, f^c)$ is composed of a configuration \mathcal{C} , a reduction function f^r and a recoloring function f^c . The configuration distinguishes a 2-family or 4-family U , that we call **special vertices**. Given a planar graph G that contains the configuration \mathcal{C} , we call $f^r(G)$ the *reduced graph* G' , whose vertex set is $V(G') = V(G) \setminus U$ and some of its edges were added or removed from G .

Given G and a coloring pc (called *pre-coloring*) of the reduced graph $G' = f^r(G)$, the recoloring function $f^c(G, pc)$ gives a coloring of G .

For instance, let us consider the rule shown in Figure 4, whose formalism will be fully described in the next subsection. The configuration of the rule is the following: the graph contains at least 5 vertices u_1, u_2, v, v_1, v_2 and potentially a vertex v_3 such that u_1 is a 3-vertex adjacent to v_1, v and u_2 ; the vertex u_2 is adjacent to u_1, v, v_2 and potentially v_3 , but not to other vertices; and v_2 is a vertex of even degree. The reduction function consists in removing every edge incident to u_1 and u_2 . As v_2 has an even degree in G , it has an odd degree in the reduced graph G' . Hence it is the end of a path Q . The recoloring function is the following: color the edges (v_2, u_2) and (u_2, u_1) with the color of Q , use a new color to color the edges $(v_1, u_1), (u_1, v), (v, u_2)$ and (u_2, v_3) if u_2 is a 4-vertex, to form a new path P . For all other edges, use colors of pc .




Figure 4: Example of a reduction rule. The formalism used to describe reduction rules is given in Subsection 3.1.

A reduction rule $\mathcal{R} = (\mathcal{C}, f^r, f^c)$ is *valid* if for any planar graph G that contains the configuration \mathcal{C} , then the reduced graph $G' = f^r(G)$ is planar; for any path coloring pc of G' , $f^c(G, pc)$ is a path coloring; and for any coloring pc of G' , $|f^c(G, pc)| - |pc| \leq \lfloor \frac{|V(G)| - |V(G')|}{2} \rfloor$. One can easily check that the rule of Figure 4 is valid.

The rest of this section and the entirety of Section 4 are dedicated to describing a set of configurations that cover all the cases of Lemma 3.1, and providing a resolution rule for each of these configurations. For each rule we provide, we justify that it is valid. The existence of a valid rule $\mathcal{R} = (\mathcal{C}, f^r, f^c)$ is in itself not enough to guarantee that an MCE cannot contain the configuration \mathcal{C} . Indeed, the reduced graph could contain a K_3 or K_5^- connected component, and therefore not admit a good coloring. However, we argue in Lemma 3.6 (on page 24, for rules associated with (C_I) configurations) and Lemma 4.15 (on page 59, for rules associated with (C_{II}) configurations) that our rules are sufficient to build a good coloring of the MCE regardless of the presence of K_3 or K_5^- components in the reduced graph.

The rest of this section is dedicated to proving the first part of Lemma 3.1 (p. 3), which we reformulate as the following lemma.

Lemma 3.2. *An MCE does not contain a configuration (C_I) .*

To prove this result let us introduce a first set of reduction rules, defined over local conditions. We show that each reduction rule is valid in Lemmas 3.3, p. 18, and 3.4, p. 23. We then prove that the application of each of these rules on an MCE G is sufficient to provide a good coloring of G (Lemma 3.6, p. 24). Finally, we show (Lemma 3.7, p. 26) that the configuration (C_I) is of the form of at least one of the reducible configurations that we list below.

We define the rules with both a graphical and a textual formalism, each of them being self-sufficient.

Graphical formalism Each rule (\mathcal{C}, f^r, f^c) is formally defined by a triplet of drawings (see for instance Figure 4). The first drawing defines the configuration \mathcal{C} of the rule, the reduction function f^r of defined by the difference between the first two drawings, and finally the third drawing defines the recoloring function f^c . Let us first describe the graphical formalism used to define the configuration of a rule.

The vertices involved in a configuration are represented by circles (\circ or \bullet), diamonds (\diamond) or squares (\square). The existence of an edge is materialized by a solid line ($\bullet\text{---}\bullet$) between the vertices. The absence of an edge is materialized by a dashed line ($\bullet\text{--}\bullet$). A waved line ($\bullet\text{~}\bullet$) represents a path between two vertices that avoids other represented vertices (unless specified). When an edge is represented by a dash-dotted edge ($\bullet\text{-}\cdot\text{-}\bullet$), this means that we consider the cases when this edge is present and when it is absent. A solid line with a gray vertex in the middle ($\bullet\text{---}\bullet\text{---}\bullet$) represents a path of length 1 or 2. If it is of length 2, the middle vertex is distinct from the other vertices on the figure.

When all the incident edges of a vertex are explicitly drawn (with solid or dash-dotted edges), the vertex is represented by a black circle (\bullet). Vertices of odd (resp. even) degree are represented by diamonds (\diamond) (resp. squares (\square)). A dashed waved line ($\bullet\text{~}\cdot\text{~}\bullet$) means that the graph does not contain a path between its endpoints that avoids every vertex represented.

For the second drawing of the triplet, we need a few other conventions, because it also encodes information on the pre-coloring. Letters Q, R denote paths from the pre-coloring, and the two letters may represent the same path. If an edge is colored Q and another is colored \overline{Q} , this means that they have different colors. If an edge stays black in the second drawing, it means that the edge keeps its color in the recoloring of G . An incoming arrow colored Q at a vertex means that this vertex is the end of a path colored Q . If this arrow is on a half-edge, this means that the last edge of the path is not determined by the figure, and one of the drawn vertices could be the other end of the edge.

A black solid (resp. dashed) arrow between a vertex and a path means that the vertex is (resp. is not) on the path (see for example the cases (h_5) and (h_6) of rule (h)).

The definition of the reduction function is quite straightforward. Every edge that is in the first (resp. second) drawing but not in the second (resp. first) is deleted (resp. added) by the reduction function and both vertices u_1, u_2 are deleted (together with their incident edges).

For rules (h) and (r) , a case analysis is needed to define the recoloring function: the second drawing is split into several copies of the same graph, depicting the different possible properties of the pre-coloring of the reduced graph.

The third graph encodes directly the recoloring function by giving colors explicitly to the edges of $G \setminus G'$ from the pre-coloring of the reduced graph G' .

Textual formalism For each rule (\mathcal{C}, f^r, f^c) , we first define textually the configuration \mathcal{C} , over local conditions around the special vertices. We then define the reduction function f^r by specifying the edges that are added or removed to form the reduced graph G' . The special vertices u_1, u_2 and their incident edges are removed in each rule and omitted in the descriptions. Finally, we define the recoloring function f^c by describing the operations applied to a coloring of G' to color all the edges of G .


We call *deviation* the recoloring operation that consists in replacing a color inducing a path P' in G' by a color inducing a path P in G , such that P is obtained from P' by replacing an edge vv' with a section (v, u, v') of length 2 or (v, u, u', v') of length 3, using only special vertices u, u' as internal vertices. An example of deviation is the path Q in rule (a).

We call *extension* the recoloring operation that extends a path Q induced by a color of G' on several additional edges in the coloring of G , those edges having only the endpoints of Q and special vertices as ends. In particular, when a non-special vertex has an odd degree in G' , a color must end on it and we may extend this color in G . The rules are frequently defined so as to “force” some vertices to have an odd degree in the reduced graph.


When the rule involves 2 special vertices (which is the case for all of them except (a)), we may use an *extra color* (inducing the red path P on the drawings) to color the graph G , as by definition of valid rule, one $\left(\left\lfloor \frac{|V(G)| - |V(G')|}{2} \right\rfloor\right)$ additional color is allowed when adapting the pre-coloring of G' to a coloring of G .

With all this formalism in mind, we can introduce our first set of rules: (a), (b), \dots , (u). Note that we justify the planarity of the reduced graph and the validity of all the rules in Lemma 3.3 after the definitions.


List of the rules (a), (b), ..., (u):




- (a): The special vertex u_1 has degree 2: it has two non-adjacent neighbors v_1, v_2 .
Reduction: In the reduced graph, we add the edge v_1v_2 .
Recoloring: In G , we deviate the color of v_1v_2 on u_1 .



- (b): The two special vertices u_1, u_2 are adjacent and they have precisely one common neighbor v . The special vertex u_1 has degree 3: it has another neighbor v_1 , and u_2 has degree 3 or 4, with a neighbor v_2 and maybe another v_3 . The vertex v_2 has an even degree in G .
Reduction: In the reduced graph, v_2 has an odd degree: let Q be a path of the coloring of G' that ends on v_2 .
Recoloring: In G , we extend Q to the edges v_2u_2, u_2u_1 . We use the extra color on the path $P = (v_1, u_1, v, u_2)$ and maybe the edge u_2v_3 if it is in G .




- (c): Each of the two special vertices u_1, u_2 has degree 3 or 4. They are adjacent, they have precisely one common neighbor v and each of u_1, u_2 has another neighbor, v_1, v_2 respectively. Each of u_1, u_2 may have a third neighbor v_3, v_4 respectively. The vertices v_1, v_2 are non-adjacent.
Reduction: In the reduced graph, we add the edge v_1v_2 .
Recoloring: In G , we deviate the color of v_1v_2 on u_1, u_2 . We use the extra color on the path $P = (u_1, v, u_2)$, and maybe on the edges v_3u_1 and u_2v_4 if they belong to G .



- (d): The two special vertices u_1, u_2 have degree 3. They are adjacent, they have precisely one common neighbor v and each of u_1, u_2 has another neighbor, v_1, v_2 respectively, which is adjacent to v . The vertices v_1, v_2 are adjacent, and v has an odd degree in G .

Reduction: In the reduced graph, we remove the edge v_1v_2 . The vertex v keeps an odd degree: let Q be a path of the coloring of G' that ends on it.


Recoloring: In G , we extend Q on the edges vu_1 and u_1u_2 . We use the extra color on the path (v, u_2, v_2, v_1, u_1) .



- (e): The two special vertices u_1, u_2 have degree 3. They are adjacent, they have precisely one common neighbor v and each of u_1, u_2 has another neighbor, v_1, v_2 respectively. The vertex v_1 is adjacent to v . The vertex v has an even degree in G .

Reduction: In the reduced graph, we remove the edge vv_1 . The vertex v has an odd degree in G' , so let Q be a path of the coloring of G' that ends on v .


Recoloring: In G , we extend Q on the edges vu_1 and u_1u_2 . We use the extra color on the path $P = (u_1, v_1, v, u_2, v_2)$.



- (f): Each of the two special vertices u_1, u_2 has degree 3 or 4. They are non-adjacent, they have precisely two common neighbors v_1, v_2 and each of u_1, u_2 has another neighbor v_3, v_4 respectively. Both v_3, v_4 are adjacent to both v_1, v_2 . The vertices v_1, v_2 are non-adjacent. Each of u_1, u_2 may have another neighbor, v_5, v_6 respectively.

Reduction: In the reduced graph, we add the edge v_1v_2 and remove the edges v_1v_3 and v_2v_4 .


Recoloring: In G , we deviate the color of v_1v_2 on u_1 , and the color of v_1v_4 on u_2 . We use the extra color on the path $P = (u_1, v_3, v_1, v_4, v_2, u_2)$, and maybe on the edges v_5u_1 and u_2v_6 if they belong to G .



- (g) : Each of the two special vertices u_1, u_2 has degree 3 or 4. They are non-adjacent, they have precisely two common neighbors v_1, v_2 and each of u_1, u_2 has another neighbor v_3, v_4 respectively. Both v_3, v_4 are adjacent to both v_1, v_2 . The vertices v_1, v_2 are adjacent. Each of u_1, u_2 may have another neighbor, v_5, v_6 respectively. There is a path P_{34} between v_3 and v_4 in G that is vertex-disjoint from the other 6 vertices.

Reduction: In the reduced graph, we remove the edges of the path P_{34} .


Recoloring: In G , we deviate the color of v_1v_2 on u_1 , and the color of v_1v_4 on u_2 . We use the extra color on the path $P = (u_1, v_3, P_{34}, v_4, v_1, v_2, u_2)$, and maybe on the edges v_5u_1 and u_2v_6 if they belong to G .




- (h) : Each of the two special vertices u_1, u_2 has degree 3 or 4. They are non-adjacent and have precisely two common neighbors v_1, v_2 . Each of u_1, u_2 has another neighbor v_3, v_4 respectively, both adjacent to both v_1, v_2 . The set $\{v_1, v_2\}$ is a 2-cut that separates u_1, u_2 . Each of u_1, u_2 may have another neighbor v_5, v_6 respectively.

Reduction: In the reduced graph, we add the edge v_3v_4 . Let Q be the path of the coloring of G' induced by the color of v_3v_4 . We denote it $Q = (Q_1, v_3, v_4, Q_2)$.

Recoloring: We distinguish between six cases, depending on the properties of the path Q . Rule (h_1) covers the case where the path Q avoids the vertex v_2 . In rules (h_2) and (h_3) , the vertices v_2, v_3, v_4 and v_1 appear in that order on the path Q . Rule (h_2) covers the case where v_5 is on the subpath between v_2 and v_3 , and v_6 is on the subpath between v_4 and v_1 (the rules are a bit more general and only require v_5 and v_6 to avoid some subpaths of Q). Rule (h_3) covers the case where the vertex v_5 (if it exists) avoids the subpath of Q between v_2 and v_3 . By symmetry, this also covers the case where v_6 (if it exists) avoids the subpath of Q between v_4 and v_1 . In rules $(h_4), (h_5)$ and (h_6) , the vertices v_1, v_2, v_3 and v_4 appear in this order on the path Q . Rule (h_4) covers the case where the vertex v_5 avoids the subpath of Q between v_2 and v_3 , while in rules (h_5) and (h_6) the vertex v_5 avoids the subpath of Q between v_1 and v_2 . In rule (h_5) , the vertex v_6 avoids the subpath of Q after v_1 , and in rule (h_6) it avoids the subpath between v_1 and v_2 . Since each of v_5, v_6 avoids at least one subpath of Q , this covers all possible cases.




1. Q does not touch v_2 in G' .
In G , we replace the path Q with the path $Q' = (Q_1, v_3, u_1, v_2, u_2, v_4, Q_2)$. We use the extra color on the path $P = (u_1, v_1, u_2)$ and maybe on the edges v_5u_1 and u_2v_6 if they belong to G .




2. Q_1 touches v_2 and Q_2 touches v_1 . The vertex v_5 does not touch Q_2 , and v_6 does not touch Q_1 .

In G , we replace Q with a path $Q' = (Q_1, v_3, u_1, v_1, u_2)$ and maybe extend Q' on u_2v_6 . We use the extra color on the path $P = (Q_2, v_4, u_2, v_2, u_1)$ and maybe on the edge u_1v_5 .




3. Q_1 touches v_2 and Q_2 touches v_1 . We denote $Q_1 = (v_3, R_1, v_2, R'_1)$. The vertex v_5 does not touch R_1 . Note that by planarity, v_6 cannot touch R_1 .

In G , we replace Q with a path $Q' = (R'_1, v_2, u_2, v_4, Q_2)$ and we deviate the color of v_3v_1 in G' on u_1 . We use the extra color on the path $P = (v_5, u_1, v_2, R_1, v_3, v_1, u_2)$ and maybe on the edge u_2v_6 .




4. Q_1 touches both v_1, v_2 : we denote it $Q_1 = (R_1, v_1, R'_1, v_2, R''_1, v_3)$. The vertex v_5 does not touch R''_1 . Again by planarity, v_6 cannot touch R''_1 .


In G , we replace Q with a path $Q' = (u_1, v_2, R''_1, v_3, v_1, u_2)$ and maybe extend Q' on the edges v_5u_1 and u_2v_6 . We deviate the color of v_3v_1 on u_1 and we use the extra color on the path $P = (Q_2, v_4, u_2, v_2, R'_1, v_1, R_1)$.



5. Q_1 touches both v_1, v_2 : we denote it $Q_1 = (R_1, v_1, R'_1, v_2, R''_1, v_3)$. The vertex v_5 does not touch R'_1 , and v_6 does not touch R_1 . Again, note that by planarity v_5 cannot touch Q_2 and v_6 cannot touch R''_1 .
 In G , we replace Q with a path $Q' = (u_1, v_2, R'_1, v_1, u_2, v_4, Q_2)$ and maybe extend Q' on the edge v_5u_1 . We deviate the color of v_3v_1 on u_1 and we use the extra color on the path $P = (R_1, v_1, v_3, R''_1, v_2, u_2)$ and maybe on the edge u_2v_6 .




6. Q_1 touches both v_1, v_2 : we denote it $Q_1 = (R_1, v_1, R'_1, v_2, R''_1, v_3)$. None of v_5, v_6 touch R'_1 .
 In G , we replace Q with a path $Q' = (R_1, v_1, u_1, v_3, R''_1, v_2, u_2, v_4, Q_2)$. We use the extra color on the path $P = (u_1, v_2, R'_1, v_1, u_2)$ and maybe on the edges v_5u_1 and u_2v_6 .



- (i): The special vertex u_1 has degree 3 or 4, and u_2 has degree 4. The special vertices u_1, u_2 have (at least) two common neighbors v, v' that are non-adjacent. The special vertex u_2 has another neighbor v_2 , non-adjacent to v' nor u_1 , and u_1 may have another neighbor v_1 , non-adjacent to u_2 . If u_1, u_2 are non-adjacent, they have another common neighbor w ; let us denote P_{12} the path (u_1, u_2) if u_1, u_2 are adjacent, and (u_1, w, u_2) otherwise.

Reduction: In the reduced graph, we add the edges vv' and v_2v' .


Recoloring: In G , we deviate the color of vv' on u_1 and the color of v_2v' on u_2 . We use the extra color on the path $P = (v, u_2, P_{12}, u_1)$ and maybe on the edge u_1v_1 if it belongs to G .



- (j): The special vertex u_1 has degree 3 or 4, and u_2 has degree 4. The special vertices u_1, u_2 have (at least) two common neighbors v, v' that are non-adjacent. The special vertex u_2 has another neighbor v_2 , adjacent to v, v' but not u_1 , and u_1 may have another neighbor v_1 , non-adjacent to u_2 . If u_1, u_2 are non-adjacent, they have another common neighbor w ; let us denote P_{12} the path (u_1, u_2) if u_1, u_2 are adjacent, and (u_1, w, u_2) otherwise.

Reduction: In the reduced graph, we add the edge vv' and remove the edge vv_2 .


Recoloring: In G , we deviate the color of vv' on u_1 and the color of v_2v' on u_2 . We use the extra color on the path $P = (v', v_2, v, u_2, P_{12}, u_1)$ and maybe on the edge u_1v_1 .



- (k): The special vertices u_1, u_2 have degree 3, with at least two common neighbors v, v' that are non-adjacent. If u_1, u_2 are non-adjacent, they have another common neighbor w ; let us denote P_{12} the path (u_1, u_2) if u_1, u_2 are adjacent, and (u_1, w, u_2) otherwise. The vertex v' has an even degree in G .

Reduction: In the reduced graph, we add the edge vv' . The vertex v' now has an odd degree, so let R be a path of the coloring of G' that ends on v' .


Recoloring: In G , we deviate the color of vv' on u_1 , and extend R on the edge $v'u_2$. We use the extra color on the path $P = (u_1, P_{12}, u_2, v)$.



- (l): The special vertices u_1, u_2 have degree 3, with at least two common neighbors v, v' that are non-adjacent and both have an odd degree in G . If u_1, u_2 are non-adjacent, they have another common neighbor w ; let us denote P_{12} the path (u_1, u_2) if u_1, u_2 are adjacent, and (u_1, w, u_2) otherwise.


Reduction: In the reduced graph, v, v' now have an odd degree, so let Q, R be paths of the coloring of G' that end on v, v' respectively.

Recoloring: In G , we extend Q on the edge vu_1 and R on $v'u_2$. We use the extra color on the path $P = (v', u_1, P_{12}, u_2, v)$.



- (m_1) : The two special vertices u_1, u_2 have degree 3, they are non-adjacent and have three common neighbors v, v', v'' , such that v' is adjacent to v and v'' .


Recoloring: In G , we deviate the color of vv' in G' on u_2 and the color of $v'v''$ on u_1 . We use the extra color on the path $P = (u_1, v, v', v'', u_2)$.



- (m_2) : The special vertex u_1 has degree 4, and u_2 has degree 3 or 4: they are non-adjacent and have precisely three common neighbors v, v', v'' , such that v' is adjacent to v and v'' . The special vertex u_1 has another neighbor v_1 , and u_2 may have another neighbor v_2 , such that $v_1 \neq v_2$.


Reduction: In the reduced graph, we add the edge vv_1 if it does not already belong to G .

Recoloring: In G , we deviate the color of vv_1 in G' on u_1 , and the color of vv' on u_2 . We use the extra color on the path $P = (v, v', u_1, v'', u_2)$ and maybe on the edges vv_1 and u_2v_2 if they belong to G .



- (n) : Each of the two special vertices u_1, u_2 has degree 3 or 4. They are adjacent and have precisely two common neighbors v, v' that are adjacent. Each of u_1, u_2 may have another neighbor, v_1, v_2 respectively.


Recoloring: In G , we deviate the color of vv' in G' on u_1, u_2 . We use the extra color on the path $P = (u_1, v', v, u_2)$ and maybe on the edges v_1u_1 and u_2v_2 if they belong to G .



- (o): The special vertices u_1, u_2 have degree 4, with at least three common neighbors v, v', v'' such that v' is non-adjacent to v and v'' . If u_1, u_2 are non-adjacent, they have another common neighbor w ; let us denote P_{12} the path (u_1, u_2) if u_1, u_2 are adjacent, and (u_1, w, u_2) otherwise.

Reduction: In the reduced graph, we add the edges vv' and $v'v''$.


Recoloring: In G , we deviate the color of vv' on u_1 and the color of $v'v''$ on u_2 . We use the extra color on the path $P = (v'', u_1, P_{12}, u_2, v)$.



- (p): The special vertices u_1, u_2 have degree 4, with at least three common neighbors v, v', v'' such that v' is adjacent to v and non-adjacent to v'' . If u_1, u_2 are non-adjacent, they have another common neighbor w ; let us denote P_{12} the path (u_1, u_2) if u_1, u_2 are adjacent, and (u_1, w, u_2) otherwise.


Reduction: In the reduced graph, we add the edge $v'v''$.

Recoloring: In G , we deviate the color of vv' on u_1 and the color of $v'v''$ on u_2 . We use the extra color on the path $P = (v'', u_1, P_{12}, u_2, v, v')$.



- (q): The special vertex u_1 has degree 2, and u_2 has degree 3 or 4. The special vertices u_1, u_2 are adjacent and have precisely one common neighbor v . The special vertex u_2 has another neighbor v_1 adjacent to v , and maybe another neighbor v_2 .

Recoloring: In G , we deviate the color of the edge vv_1 in G' on u_1, u_2 . We use the extra color on the path $P = (u_2, v, v_1)$, and maybe on the edge v_2u_2 if it belongs to G .



- (r): The two special vertices u_1, u_2 have degree 2: they are adjacent and have one common neighbor v . The vertex v has at least one other neighbor v_1 .


In the reduced graph, we examine two cases.

- In the first case, a path Q of the coloring of G' ends on v (through the edge v_1v).

Recoloring: In G , we extend Q on the edge vu_1 . We use the extra color on the path $P = (u_1, u_2, v)$.

- In the second case, no path of the coloring of G' ends on v . Let Q be a path of the coloring such that $Q = (Q_1, v_1, v, v_2, Q_2)$, with v_2 another neighbor of v .


Recoloring: In G , we replace Q with a path $Q' = (Q_1, v_1, v, u_1)$ and we use the extra color on the path $P = (u_1, u_2, v, v_2, Q_2)$.



- (s): The two special vertices u_1, u_2 have degree 2: they are non-adjacent and have two common neighbors v_1, v_2 that are adjacent. The vertex v_2 has an odd degree in G .

Reduction: In the reduced graph, v_2 keeps an odd degree: let Q be a path of the coloring that ends on v_2 .


Recoloring: In G , we deviate the color of v_1v_2 in G' on u_1 , and we extend the path Q on the edge v_2u_2 .



- (t): The two special vertices u_1, u_2 have degree 2: they are non-adjacent and have two common neighbors v_1, v_2 that are adjacent and have an even degree in G .

Reduction: In the reduced graph, we remove the edge v_1v_2 . The vertices v_1, v_2 have odd degrees in G' : let Q, R be two paths of the coloring that end on v_2, v_1 respectively.

Recoloring: In G , we extend the path Q on the edge v_2u_2 and the path R on v_1u_1 . We use the extra color on the path $P = (u_1, v_2, v_1, u_2)$.



- (u): The special vertex u_1 has degree 2, and u_2 has degree 3 or 4. The special vertices u_1, u_2 are non-adjacent and have two common neighbors v_1, v_2 that are adjacent. The vertex u_2 has another neighbor v_3 adjacent to v_2 . The vertex u_2 may have another neighbor v_4 .

Recoloring: In G , we deviate the color of the edge v_1v_2 in G' on u_1 and the color of v_2v_3 in G' on u_2 . We use the extra color on the path $P = (v_3, v_2, v_1, u_2)$ and maybe on the edge u_2v_4 if it belongs to G .

Lemma 3.3. *The reduction rules of configurations (a), (b), ..., (u) are valid.*

Proof. For each rule, we need to check three properties: when the rule applied to a planar graph G , the reduced graph G' produced is planar; the recoloring function yields a path coloring (i.e. does not introduce cycles); and the number of additional colors used in the recoloring function is at most $\left\lfloor \frac{|V(G)| - |V(G')|}{2} \right\rfloor$.

For the first property, observe that in all considered rules except rule (h), an edge ab added by the reduction function replaces a deleted path of length 2 or 3 between a and b that goes through 1 or 2 special vertices. Hence the planarity is preserved in these cases. Now, let us consider rule (h). Let G'' be the graph obtained from G by removing the vertices u_1 and u_2 and their incident edges. When removing u_1 and its incident edges, we obtain a face f_1 incident with v_3, v_1 and v_2 . For the same reason, v_4, v_1 and v_2 have a common face f_2 . If $f_1 \neq f_2$, since v_1, v_2 is a separating pair that separates v_3 from v_4 , there exists a planar embedding of G'' such that v_3 and v_4 are on the same face. Hence in both cases there is an embedding of G'' such that v_3 and v_4 are incident with a common face. Hence we can add the edge v_3v_4 to obtain G' while preserving the planarity.

The third property is easy to check, since the rule (a) does not introduce any new color, and all other rules have 2 special vertices and introduce exactly 1 new color (the color red on the figures, inducing the new path P).

Finally, let us check the second property. We can easily check on each rule that when the recoloring function f^c is applied on a planar graph G and a coloring pc of the reduced graph G' , the coloring $f^c(G, pc)$ provides a color for each edge of G (as a reminder, the edges drawn in black in the second and third drawings of a rule keep their color from the pre-coloring pc in G).

In G , the only edges of the new path P are the ones represented in the third drawing of each rule. One can easily check for each rule that these edges form a path. For colors used in the pre-coloring pc , except in rules (h) and (r), we only perform two types of modification, deviations and extensions. Since each special vertex is only involved in one such modification, each color of pc induces a path in G .

In the first case of rule (r), we do a simple path extension as before. In the second case, we first split an existing path into two paths, using the new color (the path P in red). Then we extend these two subpaths toward special vertices, hence we obtain a path coloring.

For rule (h), heavier modifications are made on the path Q in G' . In rule (h₁), the edge v_3v_4 is replaced by the path $(v_3, u_1, v_2, u_2, v_4)$. Since the vertex v_2 explicitly avoids the path Q , the resulting coloring does not introduce cycles. In rule (h₂), the vertex v_5 avoids the subpath Q_2 of Q , i.e. the section after v_4 , and v_6 avoids Q_1 , i.e. the section before v_2 . Hence in the final coloring, P and Q' are paths. In rule (h₃), we can easily check that Q' is a path. Moreover, v_5 avoids the section R_1 of Q , i.e. the section between v_2 and v_3 . Since by planarity v_6 cannot touch R_1 , then P is a path. The same argument applies to rule (h₄) with both v_5 and v_6 avoiding the section R'_1 , the former by hypothesis and the latter by planarity. In rule (h₅), Q' is a path since v_5 cannot touch the section R'_1 of Q by hypothesis, and the section Q_2 by planarity. Similarly, P is a path since v_6 cannot touch the section R_1 of Q by hypothesis, and the section R''_1 by planarity. In rule (h₆), we can easily check that Q' is a path. Since both v_5 and v_6 avoid the section R'_1 of Q , then P is a path. This completes the examination of all cases. \square

The second group of rules we consider are given by the neighborhood of two vertices u_1 and u_2 of degree at most 4 together with a shortest path S joining them, such that $|N(u_1) \cap N(u_2)| \leq 1$. Note that this includes the case where u_1 and u_2 are adjacent, with no common neighbors. These rules can be described as the product of two so-called *elementary partial rules*, that specifies the behavior of the rule around each endpoint of the path. In Section 4, we will introduce more general rules and partial rules to deal with (C_{II}) configurations.

More formally, an *elementary partial configuration* \mathcal{C}_i is a configuration defined over the neighborhood of one *special vertex* u_i , with one identified incident edge, called *subdivision edge*. We say that a neighbor v of u_i is a *remaining neighbor* of u_i if u_iv is not the subdivision edge.

Given a graph G , two vertices u_1, u_2 of G , a shortest (u_1, u_2) -path S , and two elementary partial configurations \mathcal{C}_1 and \mathcal{C}_2 , the *path composite configuration* $(\{\mathcal{C}_1(u_1), \mathcal{C}_2(u_2)\}, S)$ is defined as the following configuration: u_1 (resp. u_2) satisfies the partial configuration \mathcal{C}_1 (resp. \mathcal{C}_2), the path S contains the subdivision edges of \mathcal{C}_1 and \mathcal{C}_2 and does not touch the other neighbors of u_1 and u_2 . For ease of notation, we may simply write $\mathcal{C}_1 \oplus \mathcal{C}_2$.

An *elementary partial rule* is a rule $\mathcal{R}_i = (\mathcal{C}_i, f_i^r, f_i^c)$ associated with an elementary partial configuration \mathcal{C}_i , a partial reduction function f_i^r and a partial recoloring function f_i^c . The partial reduction function f_i^r of the rule is encoded by a set $\mathcal{O}_i \subseteq \{\text{add, remove}\} \times E(\mathcal{C}_i)$ with straightforward semantics. In particular, we can identify all the vertices of $V(G) \setminus N(u_i)$

between G and $f_i^r(G)$. The partial recoloring function defines the coloring of the edges, based on existing colors plus an extra color (represented as red on the figures) used in part for the edges of S .

If $\mathcal{R}_1 = (\mathcal{C}_1, f_1^r, f_1^c)$ and $\mathcal{R}_2 = (\mathcal{C}_2, f_2^r, f_2^c)$ are two elementary partial rules, u_1, u_2 two vertices and S a shortest (u_1, u_2) -path, the *path composite rule* $(\{\mathcal{R}_1(u_1), \mathcal{R}_2(u_2)\}, S)$ is the reduction rule $(\mathcal{C}_c, f_c^r, f_c^c)$ associated with the path composite configuration $(\{\mathcal{C}_1(u_1), \mathcal{C}_2(u_2)\}, S)$, and is defined as follows. Let $U = \{u_1, u_2\}$. The reduction function f_c^r is defined by $f_c^r(G) = (f_1^r \circ f_2^r(G)) \setminus (U \cup E(S))$, i.e. the successive application of the operations in \mathcal{O}_2 and \mathcal{O}_1 and the removal of the special vertices and the edges of S to form the reduced graph G' .

Let pc be a coloring of $G' = f_c^r(G)$, and c_S a 1-coloring of the path S . The recoloring function f_c^c is defined by $f_c^c(G, pc) = f_2^c(G, f_1^c(f_2^r(G), pc \cup c_S))$; in other words, the path S is added to G' and colored with c_S , then the reduction of \mathcal{C}_1 is undone, the edges in the neighborhood of u_1 are colored according to the partial recoloring function f_1^c , and finally the reduction of \mathcal{C}_2 is undone (to obtain G) and the edges in the neighborhood of u_2 are colored according to the partial recoloring function f_2^c .

Let us present the list of elementary partial rules that we consider in this section. We extend our graphical formalism by representing the subdivision edge as a red edge with a double arrow ($\bullet \leftarrow \leftarrow$).

Note again that we justify the planarity of the reduced graph and the validity of the rules formed by two elementary partial rules in Lemma 3.4 after the definitions.


List of the elementary partial configurations:



(\mathcal{C}_{EXT})

- (\mathcal{C}_{EXT}) : The special vertex u_1 has exactly one remaining neighbor v_1 .

Recoloring: In G , we extend the extra color on the edge u_1v_1 .




(\mathcal{C}_V)


- (\mathcal{C}_V) : The special vertex u_1 has exactly two remaining neighbors v_1, v_2 that are non-adjacent.

Reduction: In the reduced graph, we add the edge v_1v_2 .


Recoloring: In G , we deviate the color of v_1v_2 on u_1 .




- (C_{Ne}): The special vertex u_1 has exactly two remaining neighbors v_1, v_2 that are adjacent. The vertex v_1 has an even degree in G .
Reduction: In the reduced graph, v_1 has an odd degree: let R be a path of the coloring of G' that ends on v_1 .
Recoloring: In G , we extend the path R on the edge v_1u_1 .



- (C_{No}): The special vertex u_1 has exactly two remaining neighbors v_1, v_2 that are adjacent. The vertex v_1 has an odd degree in G .
Reduction: In the reduced graph, we remove the edge v_1v_2 . The vertex v_1 keeps an odd degree in G' : let R be a path of the coloring of G' that ends on v_1 .
Recoloring: In G , we extend the path R on the edge v_1u_1 , and we extend the extra color on the edges u_1v_2 and v_2v_1 .



- (C_{4a}): The special vertex u_1 has exactly three remaining neighbors v_1, v_2, v_3 , such that v_1, v_2 are non-adjacent (remark that v_1, v_2 are not necessarily consecutive in the cyclic order of the neighbors of u_1).
Reduction: In the reduced graph, we add the edge v_1v_2 .
Recoloring: In G , we deviate the color of v_1v_2 on u_1 and extend the extra color on the edge u_1v_3 .




- (C_{4b}): The special vertex u_1 has exactly three remaining neighbors v_1, v_2, v_3 , such that the edges v_1v_2 and v_2v_3 belong to G .


Reduction: In the reduced graph, we remove the edge v_2v_3 .

Recoloring: In G , we deviate the color of v_1v_2 on u_1 and extend the extra color on the edges u_1v_3, v_3v_2, v_2v_1 .


For convenience, we define some aliases which group several elementary partial configurations together. Note that these aliases are not disjoint: the configuration (\mathcal{C}_V) appears in both (\mathcal{C}_V^+) and (\mathcal{C}_N) , and these two aliases are particular cases of (\mathcal{C}_N^+) .



- (\mathcal{C}_V^+) : The special vertex u_1 has either 0, 2 or 3 remaining neighbors v_1, v_2, v_3 , such that at least two of them are non-adjacent (not necessarily consecutive in the cyclic order of the neighbors). If it has 0, this is configuration (\mathcal{C}_{1a}) ; if it has exactly 2 and they are non-adjacent this is configuration (\mathcal{C}_V) ; and if it has three remaining neighbors, and at least two of them are non-adjacent, this is configuration (\mathcal{C}_{4a}) .



- (\mathcal{C}_N) : The special vertex u_1 has 2 remaining neighbors v_1, v_2 . If v_1, v_2 are non-adjacent, this is configuration (\mathcal{C}_V) . Otherwise, if one of v_1, v_2 has an even degree in G , this is configuration (\mathcal{C}_{Ne}) , and if both have an odd degree in G , this is configuration (\mathcal{C}_{No}) .



- (\mathcal{C}_N^+) : The special vertex u_1 has between 0 and 3 remaining neighbors v_1, v_2, v_3 . If it has 0, 1 or 2, this is configuration (\mathcal{C}_{1a}) , (\mathcal{C}_{EXT}) and (\mathcal{C}_N) respectively. If it has 3 remaining

neighbors, then if two of them (not necessarily consecutive in the cyclic order) are non-adjacent this is configuration (\mathcal{C}_{4a}) and otherwise (\mathcal{C}_{4b}) .

Since two partial rules must be applied on the same graph, we need to make sure that they are compatible and do not interfere with each other. For example, if two (\mathcal{C}_V) rules were to be applied on the same non-edge v_1v_2 , the color of v_1v_2 in the reduced graph would have to be deviated to two different special vertices, creating a cycle in the path decomposition. Alternatively, if two (\mathcal{C}_{Ne}) or (\mathcal{C}_{No}) configurations share a remaining neighbor v , then the extra color may be extended to v from both remaining neighbors, again creating a cycle in the decomposition. The following lemma provides sufficient conditions of compatibility between partial rules, which are satisfied in the composite configurations given in the proof of Lemma 3.7 (p. 26).

Lemma 3.4 (Sufficient conditions of compatibility between partial rules). *Let u_1, u_2 be two special vertices, and S a shortest (u_1, u_2) -path, let $\mathcal{C}_a \in (\mathcal{C}_N^+)$ and $\mathcal{C}_b \in (\mathcal{C}_N^+)$, and let $\mathcal{R}_a, \mathcal{R}_b$ be the elementary partial rules associated with $\mathcal{C}_a, \mathcal{C}_b$ respectively. Then the path composite rule $(\{\mathcal{R}_a(u_1), \mathcal{R}_b(u_2)\}, S)$ is valid if the following conditions are satisfied:*

- *If none of $\mathcal{C}_a, \mathcal{C}_b$ are (\mathcal{C}_V) or (\mathcal{C}_{4a}) configurations, then u_1, u_2 share no remaining neighbors;*
- *If \mathcal{C}_a is a configuration (\mathcal{C}_V) or (\mathcal{C}_{4a}) , then u_1 shares at most one remaining neighbor v with u_2 , and v is non-adjacent to at least one other remaining neighbor of u_1 (i.e. v is not v_3 in the definition of (\mathcal{C}_{4a})).*

Proof. We need to check three properties: when the path composite rule is applied on a planar graph G , the reduced graph G' produced is planar; the recoloring function does not introduce cycles; and the number of additional colors used in the recoloring function is at most $\lfloor \frac{|V(G)| - |V(G')|}{2} \rfloor$.

The first property is easy to check, as in all considered elementary partial rules, an edge v_1v_2 added by the reduction function replaces a deleted path of length 2 between v_1 and v_2 that goes through a special vertex, and each special vertex is involved in at most one such operation.

For the the third property, observe that the rule $(\{\mathcal{R}_a(u_1), \mathcal{R}_b(u_2)\}, S)$ involves two special vertices. The recoloring function only uses colors from the pre-coloring, plus 1 new color (c_S in the definition, the path P drawn in red on the figures) which is exactly $\lfloor \frac{|V(G)| - |V(G')|}{2} \rfloor$.

For colors used in the pre-coloring, we only perform two types of modification: deviations and extensions. Since each special vertex is involved in at most one such modification, the recoloring function does not introduce cycles involving these colors.

It remains to check that the set P of edges colored with the new color in c_S is indeed a path. It is made up of a shortest path S between the two special vertices, and maybe some edges incident with only remaining neighbors and special vertices. Let P_a (resp. P_b) be the edges of P in the recoloring of \mathcal{R}_a (resp. \mathcal{R}_b). So $P = S \cup P_a \cup P_b$. We can check that each elementary partial rule does not introduce cycles within the elementary partial configuration, i.e. $S \cup P_a$ and $S \cup P_b$ are paths. If \mathcal{C}_a and \mathcal{C}_b do not share remaining neighbors, then P_a and P_b are vertex-disjoint, and thus P induces a path. Now assume that \mathcal{C}_a is a configuration (\mathcal{C}_V) or (\mathcal{C}_{4a}) , that (w.l.o.g.) the vertex v_1 of \mathcal{C}_a (w.r.t. the notations in the definition of (\mathcal{C}_V) and (\mathcal{C}_{4a})) also belongs to \mathcal{C}_b , and that the other remaining neighbors of \mathcal{C}_a and \mathcal{C}_b are disjoint. Since v_1 does not touch the edges of P_a , then P_a and P_b are vertex-disjoint and P is a path. This concludes the proof. \square

3.2 Sufficiency of the (C_I) rules

We prove with Lemma 3.6 that, because of the reduction rules associated with configurations $(a), (b), \dots, (u), (\mathcal{C}_N^+) \oplus (\mathcal{C}_N^+)$, an MCE G cannot contain any of them. We first show that it is easy to derive a contradiction if the reduced graph G' does not contain K_3 or K_5^- components, by building a good coloring of G . Then, we assume that G' contains some K_3 or K_5^- connected components, and we build once again a good (path-)coloring of G , with the help of an intermediate coloring of G' with paths and cycles. The proof then combines cycles and paths together to save colors. Let us introduce a relevant lemma from [9]. The *exceptional graph* is a graph consisting of a cycle C of length 5 and a path P , such that $V(C) \subseteq V(P)$ and $V(C)$ induces a K_5^- . We restate here Lemma 2.1. from [9].




Figure 5: The exceptional graph

Lemma 3.5 ([9]). *Let C be a cycle and P a path, such that C and P are edge-disjoint. If $1 \leq |V(C) \cap V(P)| \leq 5$, then $E(C) \cup E(P)$ can be decomposed into 2 paths, unless $C \cup P$ is the exceptional graph.*

Remark: Lemma 3.5 cannot be applied to K_5^- , as in any decomposition of K_5^- into a cycle of length 5 and a path of length 4, the path and the cycle form the exceptional graph.

There are several possible decompositions of the exceptional graph into a path and a cycle. The following observation states that in any such decomposition, the path and the cycle satisfy the properties of the definition.

Observation 1. *Let C be a cycle and P a path, such that $C \cup P$ forms the exceptional graph. Then C has length 5, $V(C) \subseteq V(P)$ and $(C \cup P)[V(C)]$ (the subgraph of $C \cup P$ induced by the vertices of C) is a K_5^- .*

Proof. There are exactly 5 vertices of degree at least 3 in $C \cup P$. Since the vertices of $(C \setminus P) \cup (P \setminus C)$ have degree 1 or 2, we deduce that $|C \cap P| = 5$.

$(C \cup P)[V(C \cap P)]$ is a K_5^- , so if there is an edge $e \in C$ that does not belong to $E(C \cap P)$, then there is a 2-path decomposition of K_5^- , a contradiction. So $E(C) \subseteq E(C \cap P)$, hence $V(C) = V(C \cap P)$. We deduce that $V(C) \subseteq V(P)$, C has length 5 and $(C \cup P)[V(C)]$ is a K_5^- . \square

Lemma 3.6. *An MCE does not contain any of the configurations $(a), (b), \dots, (u)$, and does not contain a path composite configuration $(\mathcal{C}_N^+) \oplus (\mathcal{C}_N^+)$ that satisfies the conditions of Lemma 3.4 (p. 23).*

Proof. Let us consider such a configuration X and let $\mathcal{R}_X = (X, f_X^r, f_X^c)$ be its associated reduction rule. By Lemmas 3.3 (p. 18) and 3.4 (p. 23), \mathcal{R}_X is valid. Let G be an MCE containing the configuration X , and $G' = f^r(G)$ its reduced graph.

Let n'_1, \dots, n'_p be the sizes of the connected components G'_1, \dots, G'_p of G' . Observe that $\sum_{j \leq k} n'_j = |V(G')|$. Observe that each G'_i that is not a K_3 nor a K_5^- component is a connected

planar graph that is smaller than the minimum counterexample G , hence these G'_i each admit a good coloring, using $\lfloor \frac{n'_i}{2} \rfloor$ colors. Each K_3 component can be decomposed into 1 cycle and each K_5^- component into 1 path and 1 cycle. Then G' admits a coloring pc into paths and cycles with $\sum_{i=1}^p \lfloor \frac{n'_i}{2} \rfloor \leq \lfloor \frac{|V(G')|}{2} \rfloor$ colors. Since the reduction rule is valid, G admits a coloring c_0 into paths and cycles with at most $\lfloor \frac{|V(G')|}{2} \rfloor + \lfloor \frac{|V(G)| - |V(G')|}{2} \rfloor \leq \lfloor \frac{|V(G)|}{2} \rfloor$ colors, such that no new cycles are created.

Observe that all the cycles in c_0 are vertex-disjoint. Indeed, the cycles in pc are vertex-disjoint because they belong to different connected components of G' ; the cycles may have been deviated into longer cycles in G , but since the internal vertices of the deviated sections are all special vertices, and since each special vertex is involved in at most one deviation, then no vertex of G can belong to the intersection of two cycles of c_0 .

We build iteratively a good coloring c of G , by starting from c_0 and using Lemma 3.5 (p. 24) at each iteration to replace a cycle and a path of c by two new paths that decompose the same set of edges. We first consider the case where $X \neq (a)$ and $X \neq (h)$.

We successively treat the K_5^- and K_3 components in G' . First, let us consider a component K in G' that is a K_5^- , colored with a cycle C' of length 5 and a path P' of length 4 in pc . C' is turned into a cycle C_0 from c_0 after possibly some deviations, and Lemma 3.5 has not been applied to it yet, so C_0 is also induced by a color of c . P' may have been extended (such as with path Q in rule (\mathcal{C}_{Ne})) and deviated to special vertices, into a path P_0 of G induced by a color of c_0 . So $V(P_0) \subseteq V(P') \cup U$, and since each special vertex is involved in at most one deviation or one extension, then P_0 is disjoint from the cycles of c_0 different from C_0 , so Lemma 3.5 has not been applied to it in previous iterations, and thus P_0 is also induced by a color of c . Finally, let \hat{P} be a path induced by a color of c such that $V(\hat{P}) \cap V(C_0) \neq \emptyset$ and $\hat{P} \neq P_0$. Such a path \hat{P} exists because G is connected and the cycles in c are disjoint.

Observe that $V(P_0) \cap V(C_0) = V(K)$, so $|V(P_0) \cap V(C_0)| = 5$. If there is at least one deviation on C' , then by Observation 1 (p. 24), $C \cup P$ does not form the exceptional graph. We may then apply Lemma 3.5 (p. 24) on C and P . Otherwise there is no deviation on C' , so $C_0 = C'$ (of length 5). At least 2 edges of $G[V(C_0)]$ belong to P_0 , and thus by Observation 1 (p. 24), $C_0 \cup \hat{P}$ does not form the exceptional graph. We can apply Lemma 3.5 (p. 24) on C_0 and \hat{P} , as $|V(C_0) \cap V(\hat{P})| \leq |V(C_0)| = 5$.

Now consider the case where K is a K_3 , colored by a cycle C' of length 3 in G' . Again let C be the cycle in G that is induced by the same color in c as C' in pc , after possibly some deviations, and let \hat{P} be a path induced by a color of c such that $V(\hat{P}) \cap V(C) \neq \emptyset$. At most two deviations were performed on C' , so $|V(C)| \leq 5$. We apply Lemma 3.5 (p. 24) on C and \hat{P} , unless they form the exceptional graph. Assume it is the case and C is disjoint from any other path in the coloring of G . By Observation 1 (p. 24), C has length 5 and thus contains both special vertices. Since G is not a K_5^- , \hat{P} has at least one other edge, necessarily from one special vertex to a vertex outside $V(C)$. G fits the description of the configuration (m_2) . In the rule associated with (m_2) , if the reduced graph contains a K_3 component, it contains only the vertex v_2 (in the definition of the rule), and its cycle is not deviated to the special vertices, a contradiction.

In all cases, we were able to find a cycle and a path that do not form the exceptional graph. We apply to them Lemma 3.5 (p. 24) and obtain two new paths that decompose them. We replace the cycle and the path in c by the two new paths, to obtain a coloring of G that uses the same number of colors and contains one less cycle. After this method has been successively applied to all K_3 and K_5^- appearing in G' , c has the right number of colors and contains only paths, a contradiction with G being a counterexample.

We are left with rules (a) and (h). If $X = (a)$, observe that the reduced graph G' is connected. If $G' = K_3$ or $G' = K_5^-$, then G is the same graph with one subdivided edge, so we can easily find a good coloring of G . Finally, let us consider the case $X = (h)$. First note that if a K_3 appears in G' , it contains only v_5 or only v_6 among the vertices represented in the figures. It cannot contain both as there would be a path from v_3 to v_4 in G , a contradiction with the definition of (h). A K_5^- in G' contains only v_5 , only v_6 , or only v_1, v_2, v_3, v_4 among the vertices represented. In the latter case, the K_5^- must be on vertices v_1, v_2, v_3, v_4 and another vertex v_7 . If there is a non-edge between v_1 and v_2 , then v_7 is adjacent to v_1, v_2, v_3, v_4 . Then the path v_3, v_7, v_4 contradicts the definition of (h). Thus, there is an edge v_1v_2 , and we may assume w.l.o.g. that v_7 is adjacent to v_1, v_2, v_3 .

If a K_3 (resp. K_5^-) component appears in G' and contains (only) v_5 or v_6 , then the red path P or the blue path Q in $f_X^c(G, pc)$ can easily be extended to color both the path and the K_3 (resp. K_5^-) with 2 colors (resp. 3 colors) (which is equivalent to applying Lemma 3.5, p. 24). Now assume that v_1, v_2, v_3, v_4, v_7 form a K_5^- in G' . We color it in G' with the path $R = (v_7, v_3, v_1, v_2, v_4)$ and a cycle $Q = (v_7, v_1, v_4, v_3, v_2)$. We thus place ourselves in case 3 of (h), treated with rule (h₃). When applying the recoloring function, we change the color of the edge u_2v_4 from Q (blue) to R (green), to turn Q into a path. We thus built a good coloring of G in all cases, a contradiction. \square

3.3 The rules cover all cases

Lemma 3.7. *If a graph contains a configuration (C_I) and is different from K_3 , then it contains at least one configuration among $(a), (b), \dots, (u), (C_N^+) \oplus (C_N^+), (C_V^+) \oplus (C_N^+)$, in which case the conditions of Lemma 3.4 (p. 23) are satisfied.*

Proof. We make a case analysis, described by the tree of Figure 6. For each inner node, we define the configurations of its children, and we show that if a graph contains the configuration described by the inner node, then it contains (at least) one of its children configurations. The configurations drawn inside a frame are the leaves of the tree, and the others are inner nodes.






Figure 6: Tree of implications between configurations.




- $4^- \oplus 4^-$: In the following configurations, let u_1, u_2 be two special vertices that form a configuration (C_I) and let P be a shortest path between u_1 and u_2 .
 - $(0C)$ (0 common remaining neighbor): the vertices u_1 and u_2 have degree 1, 2, 3 or 4; the vertices u_1 and u_2 have no common remaining neighbor.
 - $(1C)$ (1 common remaining neighbor): the special vertices have degree 2, 3 or 4; the path P has length 1 or 2; the special vertices have exactly one common remaining neighbor v .
 - $(2C)$ (2 common remaining neighbors): the special vertices have degree 3 or 4; the path P has length 1 or 2; the special vertices have exactly two common remaining neighbors v and v' .
 - $(3C)$ (3 common remaining neighbors): the special vertices have degree 4; the path P has length 1 or 2; the special vertices have exactly three common remaining neighbors v, v' and v'' .

Depending on the number of common remaining neighbors between u_1 and u_2 , we have one of the configurations $(0C)$, $(1C)$, $(2C)$ or $(3C)$.

Since P is a shortest path between u_1 and u_2 , when u_1 and u_2 have a common neighbor, P has length at most 2, hence all cases are covered.




- $(0C)$: The neighborhoods of both special vertices match the elementary partial configuration (\mathcal{C}_N^+) . Since u_1 and u_2 have no common remaining neighbors, we have a path composite configuration $(\mathcal{C}_N^+) \oplus (\mathcal{C}_N^+)$ that satisfies the conditions of Lemma 3.4 (p. 23).




- (1C):
 - (1Ca): The special vertex u_1 has degree 3 or 4 and the special vertex u_2 has degree 2, 3 or 4; the special vertices have one common remaining neighbor v and u_1 and u_2 are linked by a path P of length 1 or 2; moreover u_1 has a remaining neighbor v_1 that is not adjacent to v .
 - (2x): The special vertex u_1 has degree 2 and the special vertex u_2 has degree 2, 3 or 4; the special vertices have one common remaining neighbor v and u_1 and u_2 are linked by a path P of length 1 or 2; if the degree of u_2 is greater than 2, let v_1 and possibly v_2 be its other remaining neighbors and these neighbors are adjacent to v .
 - (Rx): The special vertices u_1 and u_2 have degree 3 or 4; u_1 and u_2 have 2 common neighbors v_1 and v_2 ; each special vertex has a remaining neighbor (v_3 and v_4 respectively) that is adjacent to v_1 and v_2 .
 - (33b): The special vertices u_1 and u_2 have degree 3; u_1 is adjacent to u_2 and they have a common neighbor v .
 - (44b): The special vertex u_1 has degree 3 or 4 and the special vertex u_2 has degree 4; u_1 is adjacent to u_2 and they have a common neighbor v ; the remaining neighbors of u_1 are adjacent to v .

If there is a remaining neighbor of a special vertex that is not adjacent with the common remaining neighbor, then we have the configuration (1Ca). Otherwise, every remaining neighbor is adjacent to the common neighbor v . If at least one of the special vertices has degree 2, then we have the configuration (2x). Otherwise, every special vertex has degree 3 or 4. If the distance between the special vertices is 2, then u_1 and u_2 have 2 common neighbors (let us call them v and v'). In this case, all the other neighbors are adjacent to both of them (otherwise we are back in case (1Ca), possibly changing the role of v and v'). In this case, we have the configuration (Rx). Otherwise, the special vertices are adjacent. If both vertices have degree 2, we have the configuration (33b), otherwise we have the configuration (44b).




- (2x):
 - (22n): The two special vertices u_1 and u_2 have degree 2; they have two common neighbors v_1 and v_2 ; moreover v_1 and v_2 are not adjacent.
 - (23n): The special vertex u_1 has degree 2 and u_2 has degree 3 or 4; u_1 and u_2 have two common neighbors v_1 and v_2 ; v_1 and v_2 are not adjacent; u_2 has a neighbor v_3 that is adjacent to v_2 .


In this case, u_1 has degree 2, and u_1, u_2 are either adjacent or share a neighbor w in addition to their common remaining neighbor v . First let us assume that u_2 has degree 2. If u_1, u_2 are adjacent, and since the graph is different from K_3 , then v has degree at least 3 and this is configuration (r). Otherwise, u_1, u_2 are non-adjacent. If v, w are non-adjacent, this is configuration (22n), and otherwise this is configuration (s) if one among v, w has an odd degree, and configuration (t) if both have an even degree. Now assume u_2 has degree at least 3. If u_1, u_2 are adjacent this is configuration (q), and otherwise this is configuration (23n) if v, w are non-adjacent, or (u) otherwise.




- (22n): In this case have the configuration (a) around special vertex u_1 .




- (23n): In this case have the configuration (a) around special vertex u_1 .



- (1Ca): The neighborhood of the first special vertex matches the elementary partial configuration (\mathcal{C}_V^+) and the neighborhood of the second one matches the elementary partial configuration (\mathcal{C}_N^+) . Moreover both special vertices have only one remaining neighbor in common, v , which is non-adjacent to v_1 . Hence the conditions of Lemma 3.4 (p. 23) are satisfied.




- (44b): Since the graph is planar, at least one of the remaining neighbors of u_2 is not adjacent to the remaining neighbor of u_1 . Hence we have the configuration (c).




• (33b):

- (33o): The special vertices u_1 and u_2 have degree 3; they are adjacent and they have a common neighbor v of odd degree; each special vertex has another neighbor of odd degree (v_1 and v_2 respectively) that is adjacent to v .


If there is a special vertex with an even non-common remaining neighbor, we are in the configuration (b). Otherwise, depending of the parity of the common neighbor we are in configuration (33o) (odd) or in configuration (e) (even).



- (33o): We split the case depending on whether there is an edge between v_1 and v_2 , the associated rules are (c) if there is not and (d) otherwise.




- (Rx) : If the two common neighbors are non-adjacent we are in configuration (f) , Otherwise, if the two common neighbors form a separating pair, then we are in configuration (h) , otherwise we are in configuration (g) .




- $(2C)$:
 - (C_r) : the special vertex u_1 has degree 3 or 4 and the special vertex u_2 has degree 4; u_1 and u_2 are linked by a path P of length 1 or 2; they have two common remaining neighbors v and v' ; moreover, v and v' are not adjacent.

If there are two common neighbors that are non-adjacent, then we are in configuration (C_r) if at least one special vertex has degree 4, and otherwise in configuration (k) or (l) depending on whether one of these neighbors has an even degree. Otherwise, all common neighbors are pairwise adjacent. If the special vertices are adjacent, then we are in configuration (n) . Otherwise, the path P has length 2 and a middle-vertex v'' . If both special vertices have degree 3, then we are in configuration (m_1) , and if at least one has degree 4, we are in configuration (m_2) .

Remark: Note that if the graph is a K_5^- , then it is a configuration (m_1) , but in this case the associated rule defines a coloring of the graph with 3 colors. This case does not occur if the graph is an MCE.



- (C_r) : If there is a non-common remaining neighbor that is not adjacent to a common remaining neighbor we are in configuration (i), otherwise at least one remaining neighbor is adjacent to both common remaining neighbors and we are in configuration (j).



- $(3C)$: Since the graph is planar, there are at least two non-adjacent common remaining neighbors. If none of them are adjacent we are in configuration (o), otherwise we are in configuration (p).

□

The proof of the main lemma of this section is now straightforward.

Proof of Lemma 3.2 (p. 6). Let G be an MCE, and assume it contains a configuration (C_I) . By Lemma 3.7, G contains a configuration among (a) , (b) , \dots , (u) , or a path composite configuration $(C_V^+) \oplus (C_N^+)$ or $(C_N^+) \oplus (C_N^+)$. Lemma 3.6 (p. 24) provides a contradiction. □

4 Configuration (C_{II})

In this section, we prove the following lemma, which constitutes the second property of Lemma 3.1.

Lemma 4.1. *An MCE does not contain a configuration (C_{II}).*

As a reminder, a planar graph G has a configuration (C_{II}) if it is almost 4-connected w.r.t. a 4-family U , i.e. a set of four vertices of degree 5; we say that G has a (C_{II}) configuration w.r.t. U . The proof of Lemma 4.1 is given at the end of this section, and uses a method similar to the proof of Lemma 3.2 (p. 6) in the previous section.

In this section, we take care of our four special vertices by generalizing the tools of the previous section. Instead of considering a shortest path like in (C_I) rules, we use a *subdivision*, either a K_4 or a C_{4+} -subdivision (see Figure 9), and again remove it in the reduction, then color it with our extra colors. These structures have the same convenient properties as the shortest path of the (C_I) rules: they can be colored with 2 extra colors, and there is an end of an extra color on each of the four special vertices, which is again helpful to take care of all the missing edges. We generalize the concept of elementary partial rule and consider *patterns* that recolor the neighborhoods of one or two special vertices at once. Similarly to the distant special vertices in the (C_I) rules, when the remaining neighbors of the special vertices are disjoint, we combine four “normal” patterns (called C_N like in Section 3) to form a complete reduction rule. Figure 7 shows four special vertices u_1, u_2, u_3, u_4 linked by a K_4 -subdivision S , with disjoint remaining neighbors and thus treated with the pattern C_N .




Figure 7: A (C_{II}) reducible configuration featuring C_N patterns

Several problems can occur however, which we classify in two types, *distant* and *close* problems. A special vertex forms a distant problem when its remaining neighbors are adjacent and touch the subdivision. If left untreated, a distant problem may cause a C_N pattern to create a cycle in the final coloring. The distant problems are eliminated in two ways: either by modifying the subdivision to assign new remaining neighbors to the problematic special vertices, or by **inactivating** them by finding a 2-coloring of the subdivision that is compatible with the C_N patterns treating the remaining distant problems. Figure 8 features a (C_{II}) configuration with a K_4 -subdivision S , where the special vertex u_3 causes a distant problem on a (u_2, u_4) -path of S . This problem is inactivated by a carefully chosen 2-coloring of S .

The close problems occur when two special vertices share some remaining neighbors, with these remaining neighbors possibly touching the subdivision as well. These problems are




Figure 8: A (C_{II}) configuration where u_1, u_2 form a close problem and u_3 a distant problem. The close problem is eliminated by redirection of the subdivision S , and the distant problem is inactivated by the 2-coloring of S .

treated in a custom manner, by redirecting the subdivision to assign new remaining neighbors to the special vertices and finding a compatible set of patterns to treat all four special vertices. In Figure 8, u_1 and u_2 initially form a close problem, which is eliminated by a redirection of the subdivision. The special vertices u_1, u_2 are then both treated with the C_V pattern.

More precisely, we first treat the cases with at least 3 distant problems (and no close ones), in the *distant lemma* (Lemma 4.21, p. 67), then the cases with at most 2 distant problems and no close ones in the *semi-distant lemma* (Lemma 4.23, p. 76), and finally the cases with at most 2 distant problems and some close problems in the *close lemma* (Lemma 4.25, p. 86).

The following claim is a corollary of the properties of almost 4-connectivity, and is useful in various proofs of this section.

Claim 4.2. *Let G be a planar graph that is almost 4-connected w.r.t. a 4-family U . Then G does not have a special vertex $u \in U$ that forms a K_4 with three of its neighbors.*

Proof. Let $v_1, v_2, v_3 \in N(u)$, such that $\{u, v_1, v_2, v_3\}$ form an induced K_4 in G . Since $d(u) = 5$, u has a neighbor v_4 distinct from v_1, v_2, v_3 . W.l.o.g., v_4 belongs to the face delimited by $\{u, v_1, v_2\}$. Then $\{u, v_1, v_2\}$ is a 3-cut that separates v_3 from v_4 , a contradiction to the definition of almost 4-connectivity. \square

4.1 K_4, C_{4+} -subdivisions

In Section 3, the composite rules that we considered were associated with a path between the two special vertices. In order to apply a similar method to the (C_{II}) configurations made up of 4 special vertices, we consider more complex structures: K_4 -subdivisions and C_{4+} -subdivisions (defined in the preliminaries section).

The general idea is to color the edges of the subdivision with 2 new “extra” colors such that each special vertex is the endpoint of one path. We can then proceed as in the case of the configuration (C_I) to color the neighborhood of the special vertices. However, not only are the remaining neighbors of the special vertices still not necessarily disjoint, but they can




Figure 9: The two subdivisions considered in the proof of Lemma 4.4: K_4 -subdivision and C_{4+} -subdivision

now be touched by paths of the subdivision, which forces us to consider a large number of subcases.

A result by Yu [12] gives us a K_4 -subdivision (see Figure 9a) under the condition of almost 4-connectivity, with two exceptions. We show in Lemma 4.4 below that we are able to extract from these two exceptions a C_{4+} -subdivision (see Figure 9b) with some additional properties, which we call C_{4+}^* -subdivision.

In a subdivision S rooted on a 4-family U , let us say that two special vertices u_i, u_j are k -linked, $k \in \{0, 1, 2\}$, if there are k (u_i, u_j) -paths in S with no special vertex as an internal vertex. In a K_4 -subdivision, all special vertices u_i, u_j are pairwise 1-linked, while in a C_{4+} -subdivision there are two pairs of 0-linked, two pairs of 1-linked, and two pairs of 2-linked special vertices. Note that it is sufficient to specify one pair of 1-linked and one pair of 2-linked special vertices to deduce the link of all pairs. If u_i, u_j are 1-linked, we call the (u_i, u_j) -path a *solo path* of S . If u_i, u_j are 2-linked, we call the two (u_i, u_j) -paths *parallel paths* of S .

Just like in the previous section, if $u_i \in U$ is a special vertex and v one of its neighbors, we say that v is a *remaining neighbor* of u_i if the edge $u_i v$ does not belong to S .

Definition 4.3 (C_{4+}^* -subdivision). *Let G be a planar graph with a C_{4+} -subdivision S rooted on a 4-family U . S is a C_{4+}^* if it satisfies the following three conditions:*

- Property “0-linked”: *Two 0-linked special vertices have no common remaining neighbor;*
- Property “1-linked”: *No internal vertex of a solo (u_i, u_j) -path of S is a remaining neighbor of some $u_k \in U \setminus \{u_i, u_j\}$;*
- Property “2-linked”: *If $u_i, u_j \in U$ are 2-linked, then u_i, u_j have at most one common remaining neighbor, and it belongs to a parallel path of S that is not incident with u_i, u_j .*

We say that S is a \mathcal{K} -subdivision if it is a K_4 -subdivision or a C_{4+}^* -subdivision. In the next lemma, we show how we find such a subdivision in a planar graph G with a (C_{II}) configuration U . In order to reduce the number of cases in the rest of the proof, we want to guarantee the additional property of chordlessness of the subdivision (as defined in the Preliminaries section).

Lemma 4.4. *Let H be a planar graph that is almost 4-connected w.r.t. a 4-family $U = \{u_1, u_2, u_3, u_4\}$. Then H contains a chordless \mathcal{K} -subdivision rooted on U .*

In order to prove this lemma, we use a result by Yu [12], which deals with graphs with two types of structure constraints. Let us introduce them, as N_1 -graphs and N_2 -graphs. The following definitions are taken straight from the beginning of Section 4 of [12], as the two *obstructions*, pictured in Figure 7 of [12] on page 36.

An N_1 -graph (Figure 10) is a planar graph H that has a 4-family $U = \{u_1, u_2, u_3, u_4\}$ and a facial cycle C (that we assume is the outer cycle), such that for each $i \in \{1, 2, 3, 4\}$ either $u_i \in C$ or H has a 4-cut X_i separating u_i from $U \setminus \{u_i\}$ (so $u_j \notin X_j$ for $j \neq i$), and $|X_i \cap C| = 2$. Moreover, if H_i is the component of $H \setminus X_i$ containing u_i , then the components H_i for $i \in \{1, \dots, 4\}$ are disjoint.




Figure 10: An N_1 -graph, with u_2, u_4 on the outer cycle, and u_1, u_3 surrounded by 4-cuts

An N_2 -graph (Figure 11) is a planar graph H with a 4-family $U = \{u_1, u_2, u_3, u_4\}$ and distinct (but not necessarily disjoint) 4-cuts $T_i, i \in \{1, \dots, m\}$. The 4-cuts are such that each T_i separates two vertices of U , say $\{u_1, u_2\}$, from $T_{i+1} - T_i \neq \emptyset$; $T_1 = \{a_1, a_2, a_3, a_4\}$, $T_m = \{b_1, b_2, b_3, b_4\}$ and H contains 4 disjoint paths S_i from a_i to b_i for $i \in \{1, \dots, 4\}$ respectively. Additionally, H has no 4-cut T separating $T_i \setminus T \neq \emptyset$ from $T_{i+1} \setminus T \neq \emptyset$, or separating $\{u_1, u_2\}$ from $T_1 \setminus T \neq \emptyset$, or separating $\{u_3, u_4\}$ from $T_m \setminus T \neq \emptyset$; and either $T_i \cap T_{i+1} \neq \emptyset$ or two vertices of T_i and two vertices of T_{i+1} are cofacial in H . Finally, the ten following paths exist and are internally disjoint: a (u_1, u_2) -path P_{12} , a (u_1, a_1) -path Q_1 , a (u_1, a_2) -path Q_2 , a (u_2, a_3) -path Q_3 , a (u_2, a_4) -path Q_4 , a (u_3, u_4) -path P_{34} , a (u_3, b_1) -path Q'_1 , a (u_3, b_2) -path Q'_2 , a (u_4, b_3) -path Q'_3 , and a (u_4, b_4) -path Q'_4 . This last property is not part of the exact definition from [12], but deduced from the first remark in Section 3 on page 20 of [12].

Let us restate Theorem 4.2 from [12] in the new formalism. The definitions in Yu's paper have stronger constraints and the theorem is an equivalence, but in the present paper we only need one implication. Our definitions of N_1 -graph and N_2 -graph are slightly simplified and the theorem is restated as an implication.

Theorem 4.5 (Theorem 4.2 of [12]). *Let G be a 3-connected planar graph and $U = \{u_1, u_2, u_3, u_4\} \subseteq V(G)$ be such that G has no 3-cut separating two vertices in U . Then G has a K_4 -subdivision rooted on U , or G is an N_1 -graph or an N_2 -graph.*

We argue that Theorem 4.5 does not in fact require the 3-connectivity assumption.

Theorem 4.6. *Let G be a planar graph and $U = \{u_1, u_2, u_3, u_4\} \subseteq V(G)$ be such that G has no 3-cut separating two vertices in U . Then G has a K_4 -subdivision rooted on U , or G is an N_1 -graph or an N_2 -graph.*




Figure 11: An N_2 -graph, with three 4-cuts represented: $\{a_1, a_2, a_3, a_4\}$, $\{t_1, t_2, t_3, t_4\}$, $\{b_1, b_2, b_3, b_4\}$

Proof of Theorem 4.6. We proceed by induction on the size of G . Assume that G is not 3-connected. Then there is a cut X of size at most 2. We may assume that X is a minimal cut. We consider the connected components C_1, \dots, C_p of $G \setminus X$, and observe that all special vertices in U belong to the same one, say C_1 . If $|X| = 1$, we consider $G' = G \setminus \{C_2 \cup \dots \cup C_p\}$. If $|X| = 2$, we consider $G' = G \setminus \{C_2 \cup \dots \cup C_p\} + xy$, where x and y are the two vertices in X . If $xy \in E(G)$ then adding the edge xy does not create a double edge.

In both cases, G' has fewer vertices than G , and is almost 4-connected with respect to U . By Theorem 4.5, G' has a K_4 -subdivision rooted on U , or G is an N_1 -graph or an N_2 -graph. Note that each of those three properties extend to all of G (up to modifying the embedding, in the case of an N_1 -graph with $|X| = 1$ so as to maintain that the cycle is facial). \square

We can now tackle the proof of Lemma 4.4 (p. 37).

Proof of Lemma 4.4. First, let us define the operation that turns a subdivision into a chordless one. Given a K_4 - or C_{4+} -subdivision S rooted on a 4-family U , we say that we *eliminate the chords* of S if we apply the following operation exhaustively: if a path $P = (v_1, \dots, v_k)$ of S has a chord $v_i v_j$, we replace P in S with the path $P' = (v_1, \dots, v_i, v_j, \dots, v_k)$, except if $v_i, v_j \in U$ and the edge $v_i v_j$ already constitutes a path of S . Since each elimination decreases the number of edges in S , the process terminates, and the obtained subdivision S' is well-defined and chordless. Since the vertices of S' are a subset of the vertices of S and the ends of the paths are preserved, S' has the same type (K_4 or C_{4+}) as S .

If G has a K_4 -subdivision, eliminating its chords gives us the result. Using Theorem 4.6 (p. 38), it now suffices to show that any N_1 -graph and any N_2 -graph contain a chordless C_{4+}^* -subdivision to prove our lemma.

Let us first take care of the case where H is an N_2 -graph, defined as in the definition above. Let S be the union of the paths $P_{12}, Q_1, Q_2, Q_3, Q_4, P_{34}, Q'_1, Q'_2, Q'_3, Q'_4, S_1, S_2, S_3, S_4$. S is a C_{4+} -subdivision rooted on U , so let S' be a chordless C_{4+} -subdivision obtained by eliminating the chords of S . Let us prove that S' is a C_{4+}^* -subdivision.

By definition, the 4-cut T_1 is a subset of $V(S)$, and we show that $T_1 \subseteq V(S')$. Let us assume for contradiction that there is a vertex $t \in T_1$ which belongs to a path P of S on which a chord elimination was performed: there are two vertices v, v' on P , such that the edge $vv' \in E(H)$ is not in P and t is on the (v, v') -section P' of P . But then the path $(P \setminus P') \cup \{vv'\}$ is a path between two 1-linked special vertices in $H \setminus T_1$, which is a contradiction; so $T_1 \subseteq V(S')$.

Assume for contradiction that two 0-linked special vertices u_i, u_j have a common remaining neighbor v w.r.t. S' . If v belongs to $V(S')$, then the edge $u_i v$ or $u_j v$ forms a chord of S' since all paths of S' are incident with u_i or u_j . It is a contradiction, so v does not belong to $V(S')$. However, since u_i and u_j are 0-linked, they are separated by the 4-cut T_1 , and so v is a vertex of T_1 , hence in S . This is a contradiction, which gives us the property “0-linked” of S' .

To obtain property “1-linked”, observe that no internal vertex of a solo (u_i, u_j) -path is a remaining neighbor of a $u_k \in U \setminus \{u_i, u_j\}$, since this would create a path between 0-linked special vertices that would be disjoint from the 4-cut T_1 , a contradiction.

Let us now prove the property “2-linked”. By definition of the cut T_1 , the common remaining neighbors of two 2-linked special vertices u_i, u_j belong to parallel paths of S' . If a common remaining neighbor v of u_i, u_j belongs to a (u_i, u_j) -path of S' , the two edges $u_i v, u_j v$ form chords in S' , a contradiction, so the common remaining neighbors of u_i, u_j belong to parallel (u_k, u_l) -paths that are not incident with u_i, u_j . If u_1, u_3 (2-linked) have two common remaining neighbors v, v' , they belong to different parallel (u_2, u_4) -paths by planarity (otherwise $\{u_1, u_4, v\}$ and $\{u_2, u_3, v'\}$ form a $K_{3,3}$ -minor in H if v, v' belong to a path $P = (u_2, \dots, v, \dots, v', \dots, u_4)$). Then we claim that $\{u_1, v, v'\}$ is a 3-cut of H that separates u_2 from u_3 . If not, there is a (u_2, u_3) -path P_{23} in H that is vertex-disjoint from u_1, v, v' , and by definition of T_1 , P_{23} contains a vertex $t \in T_1$ that is on a (u_1, u_3) -path of S' , and the (u_2, t) -section of P_{23} does not contain u_4 . Then $\{u_1, u_2, v, t, u_4 = v'\}$ induce a K_5 -minor in H by contracting the (u_4, v') -path of S' into a vertex. Hence $\{u_1, v, v'\}$ is a 3-cut of H that separates u_2 from u_3 , which is a contradiction with the almost 4-connectivity of H w.r.t. U . Property “2-linked” follows.

Now let us consider the case where H is an N_1 -graph with outer cycle C . We build a C_{4+} -subdivision S rooted on U as follows. First, for each $u_i \notin C$, we find four internally-disjoint paths $p_1^i, p_2^i, p_3^i, p_4^i$ from u_i to the four vertices of its associated 4-cut X_i . Such paths exist since there is no 3-cut separating u_i from $U \setminus \{u_i\}$ in H . We assume p_1^i and p_2^i each have an end on the outer cycle C . We add $(E(C) \setminus \bigcup_{u_i \notin C} E(H_i)) \cup \bigcup_{u_i \notin C} (E(p_1^i) \cup E(p_2^i))$ to S , where H_i is the component of $H \setminus X_i$ containing u_i . To obtain the two remaining paths of S , we consider the graph H' formed by removing $(V(C) \cup \bigcup_{u_i \notin C} V(H_i)) \setminus U$ from H , and add back p_3^i and p_4^i to H' . We look at the outer face of H' . Let P_{13} be the outer (u_1, u_3) -path and P_{24} the outer (u_2, u_4) -path of H' . We claim that these paths are vertex-disjoint. To see it, observe that in H there are at least 4 internally-disjoint paths from u_1 to u_3 . At least two of them are disjoint from C , hence belong to H' . Therefore, H' cannot contain a 1-cut separating $\{u_1, u_2\}$ from $\{u_3, u_4\}$. Therefore, let us add P_{13} and P_{24} to S to form our C_{4+} -subdivision S .

Let S' be a chordless C_{4+} -subdivision obtained by eliminating the chords of S . Let us now prove that S' is a C_{4+}^* -subdivision.

We first check property “1-linked”. Let u_i, u_j be 1-linked special vertices with a path P_{ij} of S' , and $u_k \in U \setminus \{u_i, u_j\}$. If there is an internal vertex v of P_{ij} that is a remaining neighbor of u_k , then $\{u_k, v\}$ is a 2-cut of H if u_k is on C , otherwise there is a vertex x in the 4-cut X_k of u_k such that $\{x, v\}$ is a 2-cut of H , contradicting its 3-connectivity.

To check properties “0-linked” and “2-linked”, we show that there is no remaining neighbor in common between u_1 and $\{u_3, u_4\}$ (respectively 2-linked and 0-linked to u_1), because of the properties of almost 4-connectivity of H . Each special vertex u_i either belongs to C or there is a 4-cut $X_i = \{x_1, x_2, x_3, x_4\}$ separating H into a component H_i containing u_i and a $H \setminus (X_i \cup H_i)$ containing $U \setminus \{u_i\}$. If $u_1, u_k, k \in \{3, 4\}$, belong to C and share a remaining neighbor v , then $\{u_1, u_k, v\}$ is a 3-cut that separates two neighbors of u_1 (if $k = 3$) or separates u_2 from u_3 (if $k = 4$). If u_1 belongs to C and u_k has a 4-cut X_k , then their common remaining neighbor is the

only vertex $x \in X_k$ that does not belong to S' . Then there is a vertex $x' \in X_k \cap C$, such that $\{u_1, x, x'\}$ is a 3-cut that separates u_2 from u_3 (whether $k = 3$ or 4). If both u_1, u_k have 4-cuts X_1, X_k , then their common remaining neighbor is again the only $x \in X_1 \cap X_k$ that does not belong to S' , and there are $x'_1 \in X_1 \cap C$ and $x'_k \in X_k \cap C$ such that $\{x'_1, x'_k, x\}$ is a 3-cut that separates u_2 from u_3 . In all cases, we obtain a contradiction with the almost 4-connectivity of H . Properties “0-linked” and “2-linked” follow, which completes the proof. \square

4.2 Patterns

Although almost all the subdivisions that we consider throughout the paper are regular K_4 -subdivision or C_{4+} -subdivisions, we occasionally consider a more convenient structure, which we call *semi- C_{4+} -subdivision*, that consists in a C_{4+} -subdivision where two parallel paths with disjoint ends intersect on one vertex.

Let W_4 be the wheel graph on 5 vertices u_1, u_2, u_3, u_4, w , i.e. the graph where u_1, u_2, u_3, u_4 form a cycle and w is adjacent to the other four vertices.

Definition 4.7 (Semi- C_{4+} -subdivision). *A semi- C_{4+} -subdivision rooted on a 4-family U in a graph G is a W_4 -subdivision rooted on $U \cup \{w\}$, where w is a vertex of G .*

By abuse of notation and by analogy with the C_{4+} -subdivision, we arbitrarily pick two pairs of special vertices $(u_i, u_j), (u_k, u_l)$ and we view the union of the (u_i, w) -path and the (u_j, w) -path as a (u_i, u_j) -path P_{ij} , and the union of the (u_k, w) -path and the (u_l, w) -path as a (u_k, u_l) -path P_{kl} . We say that there is a *contact* between P_{ij} and P_{kl} .

Observe that a semi- C_{4+} -subdivision is always 2-colorable, since we can simply swap the colors of two parallel paths in a 2-coloring, in order to give different colors to the two paths in contact (see Figure 12).




Figure 12: A 2-colored semi- C_{4+} -subdivision

The following definition regroups the different kinds of structures that we consider for our reduction rules.

Definition 4.8 (Semi-subdivision). *A semi-subdivision S in a graph G is a K_4 -subdivision, a C_{4+} -subdivision or a semi- C_{4+} -subdivision rooted on a 4-family U .*

Our overall goal is to find an individual reduction rule for each of the four special vertices, and combine them into a general rule for the whole configuration. We define these rules by extending the formalism of Section 3.

A *subdivision partial configuration* (or **pattern**) \mathcal{C}_i is a configuration defined over the neighborhood of one special vertex u_j or two special vertices u_j, u_l , with three identified incident edges per special vertex, called *subdivision edges*. We denote it by $\mathcal{C}_i(u_j)$ if it involves one special vertex, and $\mathcal{C}_i(u_j, u_k)$ otherwise.

Observe that an elementary partial configuration can be turned into a subdivision partial configuration by adding two subdivision edges. The subdivision partial configuration (\mathcal{C}_U) defined below is an example of partial configuration that involves two special vertices.

We say that a set of patterns $\mathcal{M} = \{\mathcal{C}_1, \dots, \mathcal{C}_k\}$, $k \in \{2, 3, 4\}$, is a *mapping* of a 4-family U , w.r.t. a semi-subdivision S rooted on U , if in this semi-subdivision there is a bijection between the special vertices of $\mathcal{C}_1, \dots, \mathcal{C}_k$ and the special vertices of U ; i.e. each special vertex u of $\mathcal{C}_1, \dots, \mathcal{C}_k$ can be associated with a special vertex $u' \in U$, and the neighborhoods of u and u' are isomorphic. If $u' \in U$ is associated to the special vertex u of a pattern \mathcal{C}_i , we say that u' *forms* a pattern \mathcal{C}_i w.r.t. S .


Given a mapping $\mathcal{M} = \{\mathcal{C}_1, \dots, \mathcal{C}_k\}$ and $\mathcal{C}_i \in \mathcal{M}$, we denote by $V(\mathcal{C}_i)$ the set containing the special vertices associated with \mathcal{C}_i in \mathcal{M} and their remaining neighbors. We say that a pattern \mathcal{C}_i *touches* another pattern \mathcal{C}_j if $V(\mathcal{C}_i) \cap V(\mathcal{C}_j) \neq \emptyset$. We say that \mathcal{C}_i touches S if at least one non-special vertex of $V(\mathcal{C}_i)$ belongs to a path of S .

A *subdivision composite configuration* (\mathcal{M}, S) is the following configuration: the graph contains a 4-family U and a semi-subdivision S rooted on U , while \mathcal{M} is a mapping of U w.r.t. S .

In the previous section, we defined elementary partial rules over elementary partial configurations. This definition can be directly extended to define *subdivision partial rules* over subdivision partial configurations, i.e. as a rule $\mathcal{R}_i = (\mathcal{C}_i, f_i^r, f_i^c)$ associated with a pattern \mathcal{C}_i , a partial reduction function encoded by a set $\mathcal{O}_i \subseteq \{\text{add}, \text{remove}\} \times E(\mathcal{C}_i)$ and the partial recoloring function f_i^c .

Let $\mathcal{M} = \{\mathcal{C}_1, \dots, \mathcal{C}_k\}$, $k \in \{2, 3, 4\}$, be a mapping of a 4-family $U = \{u_1, u_2, u_3, u_4\}$, w.r.t. a semi-subdivision S rooted on U . For each $i \in \{1, \dots, k\}$, let $\mathcal{R}_i = (\mathcal{C}_i, f_i^r, f_i^c)$ be a subdivision partial rule associated with the pattern \mathcal{C}_i . Let c_S be a 2-coloring of S . The *subdivision composite rule* $\mathcal{R}_c = (\mathcal{C}_c, f_c^r, f_c^c)$, denoted by $(\{\mathcal{R}_1, \dots, \mathcal{R}_k\}, S, c_S)$, is the reduction rule associated with the subdivision composite configuration $\mathcal{C}_c = (\mathcal{M}, S)$ and is defined as follows. The reduction function f_c^r is defined by $f_c^r(G) = (f_1^r \circ \dots \circ f_k^r(G)) \setminus (U \cup E(S))$, i.e. the successive application of the operations in \mathcal{O}_i in reverse order and the removal of the special vertices U and the edges of the semi-subdivision S , to form the reduced graph G' .


In order to provide a semantics of f_c^c , we define the *intermediate graphs* $G_{\text{int}}^i = f_{i+1}^r \circ \dots \circ f_k^r(G)$, for $0 \leq i \leq k$. We use these graphs to define a sequence of colorings c_{int}^i that lead to a coloring c of G . Let pc be a coloring of G' . Let $c_{\text{int}}^0 = pc \cup c_S$ be a coloring of $G_{\text{int}}^0 = G' \cup (U \cup E(S))$. We define $c_{\text{int}}^i = f_i^c(G_{\text{int}}^i, c_{\text{int}}^{i-1})$ for $i \in \{1, \dots, k\}$. We finally define $f_c^c(G, pc, c_S) = c_{\text{int}}^k$ for any planar graph G , coloring pc of $f_c^r(G)$ and good coloring c_S of S . In other words, the semi-subdivision S is added to G' and colored with c_S , then for each pattern \mathcal{C}_i considered in ascending order, the reduction of \mathcal{C}_i is undone and the edges in the neighborhood of \mathcal{C}_i are colored according to the partial recoloring function f_i^c . This definition is motivated by the fact that whenever pc is a good coloring of the reduced graph G' , and the partial rules \mathcal{R}_i are valid and do not interfere with each other, each intermediate coloring c_{int}^i is a good coloring of the intermediate graph G_{int}^i , which allows to build step by step a good coloring of G . The 2-coloring c_S of S is specified only when necessary.

In the figures, the two paths P_1, P_2 induced by the 2-coloring of S are represented in red and blue. The purple color is used to color the whole subdivision when its 2-coloring is not specified. An edge represented in black does not belong to the subdivision. A red vertex (\ominus) (resp. blue \oplus) represents a vertex that may be touched by a red (resp. blue) subdivision path. A purple vertex (\odot) may be touched by either a red or a blue subdivision path. When a path of the subdivision ends on a special vertex, it is represented by , respectively if it is the red path, the blue path, or if the color is not specified.


Let us now introduce the patterns we use in the rest of the proof. For each pattern, we describe the associated partial configuration, as well as the conditions on the colors of a 2-coloring of the associated subdivision S . We then provide a definition of the partial reduction and recoloring functions. The patterns (\mathcal{C}_V) , (\mathcal{C}_{Ne}) , (\mathcal{C}_{No}) are taken from Section 3 and their definitions are omitted.

List of the patterns:

The following patterns involve one special vertex.




- (\mathcal{C}_V) : Identical to the elementary partial configuration (\mathcal{C}_V) from Section 3.




- (\mathcal{C}'_V) : The special vertex u_1 has two adjacent remaining neighbors v_1, v_2 , and the edge v_1v_2 belongs to S .

Reduction: In the reduced graph, we add the edge v_1v_2 .

Recoloring: In G , we deviate the color of v_1v_2 on u_1 . The color of v_1v_2 in the recoloring of G is given by the subdivision.




- (\mathcal{C}_{Ne}) : Identical to the elementary partial configuration (\mathcal{C}_{Ne}) from Section 3.
Color requirements: If the remaining neighbor v_1 is even, then the other remaining neighbor v_2 of u_1 cannot touch the color of S that ends on u_1 .




- (\mathcal{C}_{No}) : Identical to the elementary partial configuration (\mathcal{C}_{No}) from Section 3.
Color requirements: None of the remaining neighbors v_1, v_2 of u_1 can touch the color of S that ends on u_1 .


The following patterns involve two special vertices.



- (\mathcal{C}_U) : The two special vertices u_1, u_2 are adjacent but the edge u_1u_2 does not belong to S . Let v_1 (resp. v_2) be the remaining neighbor of u_1 (resp. u_2) distinct from u_2 (resp. u_1). The vertices v_1, v_2 are distinct, non-adjacent and disjoint from U .
Reduction: In the reduced graph, we add the edge v_1v_2 .
Recoloring: In G , we deviate the color of v_1v_2 on u_1, u_2 .




- (\mathcal{C}_{Da}) : The two special vertices u_1, u_2 are adjacent and the edge u_1, u_2 belongs to S . The special vertices u_1, u_2 have precisely one common remaining neighbor v , and u_1, u_2 have v_1, v_2 respectively as their other remaining neighbor. The vertices v_1, v_2 are adjacent and both are adjacent to v . The vertices v, v_1, v_2 are disjoint from S .
Reduction: In the reduced graph, we remove the edge v_1v_2 .
Recoloring: In G , we deviate the color of vv_1 on the edges vu_2, u_2u_1 and u_1v_1 . We redirect the path $u_1 \sim u_2$ of S through the edges u_1v, vv_1, v_1v_2 and v_2u_2 .



- (\mathcal{C}_{Db}) : The two special vertices u_1, u_2 are adjacent and the edge u_1, u_2 belongs to S . The special vertices u_1, u_2 have precisely one common remaining neighbor v , and u_1, u_2 have v_1, v_2 respectively as their other remaining neighbor. The vertices v_1, v_2 are not adjacent and both are adjacent to v . The vertex v does not belong to S .

Reduction: In the reduced graph, we add the edge v_1v_2 .

Recoloring: In G , we deviate the color of v_1v_2 on the edges v_1u_1, u_1u_2 and u_2v_2 . We redirect the path $u_1 \sim u_2$ of S to make it go through the edges u_1v and vv_2 .




- (\mathcal{C}_{T1a}) : The two special vertices u_1, u_2 have precisely one remaining neighbor v in common. We denote v_1, v_2 the other remaining neighbor of u_1, u_2 respectively. Both v_1 and v_2 are adjacent to v . The vertices v, v_1, v_2 are disjoint from U . The vertex v_1 has an even degree.

Color requirements: The vertices v, v_2 cannot touch the color of S that ends on u_1 .

Reduction: In the reduced graph, v_1 has an odd degree: let Q be a path of the coloring of G' that ends on v_1 .

Recoloring: In G , we deviate the color of vv_2 on u_2 , we extend the path Q on the edge v_1u_1 , and we extend the extra color that ends on u_1 on the edges u_1v and vv_2 .




- (C_{T1b}): The two special vertices u_1, u_2 have precisely one remaining neighbor v in common. We denote v_1, v_2 the other remaining neighbor of u_1, u_2 respectively. Both v_1 and v_2 are adjacent to v . The vertices v, v_1, v_2 are disjoint from U . The vertices v_1, v_2 both have an odd degree in G .

Color requirements: The colors ending on u_1, u_2 in a 2-coloring of S must be different. The vertex v_1 (resp. v_2) cannot touch the color that ends on u_1 (resp. u_2).

Reduction: In the reduced graph, we remove the edges vv_1 and vv_2 . The vertices v_1, v_2 keep an odd degree in G' : let Q, R be paths of the coloring of G' that end on v_1, v_2 respectively.

Recoloring: In G , we extend the paths Q, R on the edges v_1u_1 and v_2, u_2 respectively. We extend the extra color ending on u_1 on the edges u_1v and vv_1 , and we extend the extra color ending on u_2 on the edges u_2v and vv_2 .




- (C_{T2Aa}): The two special vertices u_1, u_2 have both their remaining neighbors v, v' in common, which are adjacent and disjoint from U . The vertex v has an odd degree in G .

Color requirements: There is a color of S ending on u_1 or u_2 (let us say u_1) that does not touch v nor v' .

Reduction: In the reduced graph, v keeps an odd degree: let Q be a path of the coloring of G' that ends on v .

Recoloring: In G , we extend the path Q on the edge vu_1 , we deviate the color of vv' on u_2 , and we extend the extra color ending on u_1 on the edges u_1v' and $v'v$.




- (\mathcal{C}_{T2Ab}): The two special vertices u_1, u_2 have both their remaining neighbors v, v' in common, which are adjacent and disjoint from U . Both v and v' have an even degree in G .

Color requirements: The colors of S that end on u_1 and u_2 must be different. At least one of v, v' (let us say v') does not touch at least one of the two colors of S (let us say the one ending on u_1).

Reduction: In the reduced graph, we remove the edge vv' . The vertices v, v' have an odd degree in G' : let T, R be paths of the coloring of G' that end on v, v' respectively.

Recoloring: In G , we extend the paths T, R on the edges vu_1 and $v'u_2$ respectively. We extend the extra color ending on u_1 on the edges u_1v' and $v'v$, and we extend the extra color ending on u_2 on the edge u_2v .




- (\mathcal{C}_{T2NAa}): The two special vertices u_1, u_2 have both their remaining neighbors v, v' in common, which are not adjacent and are disjoint from U . The vertex v has an even degree in the G .

Color requirements: One of v, v' (let us say v') does not touch a color of S that ends on u_1 or u_2 (let us say u_1).

Reduction: In the reduced graph, we add the edge vv' . The vertex v has an odd degree in G' : let Q be a path of the coloring of G' that ends on v .

Recoloring: In G , we extend the path Q on the edge vu_1 , we deviate the color of vv' on u_2 , and we extend the extra color ending on u_1 on the edge u_1v' .




- $(\mathcal{C}_{T2NA b})$: The two special vertices u_1, u_2 have both their remaining neighbors v, v' in common, which are not adjacent and are disjoint from U . Both v and v' have an odd degree in G .

Color requirements: One of v, v' (let us say v') does not touch the color ending on u_1 , the other (v) does not touch the one ending on u_2 .


Reduction: In the reduced graph, v, v' keep an odd degree: let T, R be paths of the coloring of G' that end on v, v' respectively.

Recoloring: In G , we extend the paths T, R on the edges vu_1 and $v'u_2$ respectively. We extend the extra colors ending on u_1, u_2 on the edges u_1v' and u_2v respectively.


For convenience, we define some aliases which group several patterns together.




- (\mathcal{C}_N) : The special vertex u_1 has 2 remaining neighbors v_1, v_2 . If v_1, v_2 are non-adjacent, this is configuration (\mathcal{C}_V) . Otherwise, if one of v_1, v_2 has an even degree in G , this is configuration (\mathcal{C}_{Ne}) , and if both have an odd degree in G , this is configuration (\mathcal{C}_{No}) .



- (\mathcal{C}_{T1}) : The two special vertices u_1, u_2 have one remaining neighbor v in common. We denote v_1, v_2 the other remaining neighbor of u_1, u_2 respectively. Both v_1 and v_2 are adjacent to v . The vertices v, v_1, v_2 are disjoint from U .



- (\mathcal{C}_{T2A}) : The two special vertices u_1, u_2 have both their remaining neighbors v, v' in common, which are adjacent and disjoint from U .



- (\mathcal{C}_{T2NA}) : The two special vertices u_1, u_2 have both their remaining neighbors v, v' in common, which are not adjacent and are disjoint from U .

From now on, when we talk about patterns, we refer exclusively to patterns from this list.

Obviously the partial rules associated with the patterns of the considered mapping may conflict with each other. We now address the conditions of compatibility between patterns.

Definition 4.9 (Compatible patterns). *Let G be a planar graph with a semi-subdivision S rooted on a 4-family U .*

Let $\mathcal{C}_i, \mathcal{C}_j$ be two patterns on U w.r.t. S . $\mathcal{C}_i, \mathcal{C}_j$ are compatible if:

- \mathcal{C}_i or \mathcal{C}_j is a $\mathcal{C}_V, \mathcal{C}'_V$ or \mathcal{C}_U , and $|V(\mathcal{C}_i) \cap V(\mathcal{C}_j)| \leq 1$; or
- $\mathcal{C}_i, \mathcal{C}_j$ are among $\mathcal{C}_N, \mathcal{C}_{T1}, \mathcal{C}_{T2A}, \mathcal{C}_{T2NA}, \mathcal{C}_{Da}, \mathcal{C}_{Db}$, and $V(\mathcal{C}_i) \cap V(\mathcal{C}_j) = \emptyset$.

Let $\mathcal{M} = \{\mathcal{C}_1, \dots, \mathcal{C}_k\}$ be a mapping of U w.r.t. a semi-subdivision S , and $2 \leq k \leq 4$.

We say that \mathcal{M} is a compatible mapping w.r.t. S if it satisfies the following conditions:

- *The \mathcal{C}_i patterns in \mathcal{M} are pairwise compatible;*
- *There exists a 2-coloring c_S of S that fits the color requirements in the definition of each $\mathcal{C}_i \in \mathcal{M}$.*

We justify this notion of compatible patterns and compatible mapping with the following claim.

Claim 4.10. *Let G be a planar graph with a (\mathcal{C}_{II}) configuration w.r.t. a 4-family U . Let $\mathcal{M} = \{\mathcal{C}_1, \dots, \mathcal{C}_k\}$ be a mapping of U w.r.t. a semi-subdivision S , with $2 \leq k \leq 4$, and let \mathcal{R}_i be a subdivision partial rule associated with \mathcal{C}_i for $i \in \{1, \dots, k\}$.*

If \mathcal{M} is a compatible mapping w.r.t. S , then the 2-coloring c_S of S associated with \mathcal{M} is such that the subdivision composite rule $(\{\mathcal{R}_1, \dots, \mathcal{R}_k\}, S, c_S)$ associated with (\mathcal{M}, S) is valid.

Proof. Each pattern $\mathcal{C}_i \in \{\mathcal{C}_V, \mathcal{C}'_V, \mathcal{C}_U\}$ has a deviated edge, v_1v_2 in the previous definitions. Since \mathcal{C}_i is compatible with all other patterns, it shares at most one vertex with each of them, thus the edge v_1v_2 cannot be used as a deviated edge by another pattern. Hence the resolution rules of the $\mathcal{C}_V, \mathcal{C}'_V, \mathcal{C}_U$ patterns can be applied independently.

A pattern \mathcal{C}_i in $\{\mathcal{C}_N, \mathcal{C}_{T1}, \mathcal{C}_{T2A}, \mathcal{C}_{T2NA}, \mathcal{C}_{Da}, \mathcal{C}_{Db}\}$ can only share at most one vertex with each $\mathcal{C}_V, \mathcal{C}'_V$ or \mathcal{C}_U pattern, as they do not prevent the resolution rule of \mathcal{C}_i from being applied.

We emphasize that the parities involved in the resolution rules of $\mathcal{C}_N, \mathcal{C}_{T1}, \mathcal{C}_{T2A}, \mathcal{C}_{T2NA}$ are preserved no matter how many $\mathcal{C}_V, \mathcal{C}'_V$ or \mathcal{C}_U patterns touch them. Say we have a pattern \mathcal{C}_i in $\{\mathcal{C}_N, \mathcal{C}_{T1}, \mathcal{C}_{T2A}, \mathcal{C}_{T2NA}\}$, and a non-special vertex v of \mathcal{C}_i . In the descriptions of the patterns and their resolution rules, we may specify the parity of v in the graph G , then which edges we add or remove to obtain that v has an odd degree in the reduced graph G' . These definitions do not take into account the $\mathcal{C}_V, \mathcal{C}'_V, \mathcal{C}_U$ patterns or a path from S that may touch v , but we argue that they do not interfere with the parity of v in the reduced graph G' .

If a path from S touches v in G and does not form a \mathcal{C}'_V pattern, then the reduction from G to G' removes two edges incident with v , which preserves the parity of v . If a \mathcal{C}_V or \mathcal{C}_U pattern touches v in G , then one edge vu' ($u' \in U$) is removed and one edge vv' ($v' \notin U$) is added, which preserves the parity of v . Finally if a \mathcal{C}'_V pattern $\{u', v, v'\}$ touches v , then the edge vu' is removed, as well as an edge vw from the path of S that contains the edge vv' . The edge vv' is kept in the reduced graph. Thus, v has lost two incident edges, and so its degree is preserved.

In conclusion, we may apply the resolution rules of compatible patterns in any order.

Since the definitions of patterns do not create cycles, do not use additional colors, and since the 2-coloring c_S of S fits the color requirements of all patterns in \mathcal{M} , the subdivision composite rule $(\{\mathcal{R}_1, \dots, \mathcal{R}_k\}, S, c_S)$ associated with (\mathcal{M}, S) is valid. \square

Let us introduce the notion of *settled* special vertex, to characterize the special vertices that are already compatible with the rest of the configuration and whose remaining neighbors do not need to be further altered.

Definition 4.11 (Settled vertices). *Let G be a planar graph with a (C_{II}) configuration w.r.t. a 4-family U . Let S be a semi-subdivision rooted on U .*

We say that a special vertex u is lone-settled w.r.t. S if:

- *u forms a \mathcal{C}_V or \mathcal{C}'_V pattern and shares at most one remaining neighbor with each of the other special vertices, and the two remaining neighbors of u are not the two remaining neighbors of a \mathcal{C}_U pattern; or*
- *u forms a \mathcal{C}_N pattern and its remaining neighbors are disjoint from S and from the remaining neighbors of other special vertices.*

A special vertex is settled if it is lone-settled or forms a \mathcal{C}_{T2NA} pattern $\{u, u', v, v'\}$ with another special vertex u' , such that v, v' are disjoint from S and from the remaining neighbors of other special vertices.

Note that in this definition, the \mathcal{C}_V pattern formed by a lone-settled special vertex can touch the subdivision S . By Claim 4.10 (p. 49), we deduce immediately that if all four vertices of U are settled w.r.t. S , there exists a mapping \mathcal{M} of U compatible w.r.t. S .

Figure 13 provides a few examples of unsettled special vertices: on the left a \mathcal{C}_N pattern touching the subdivision, in the middle two \mathcal{C}_N patterns sharing a remaining neighbor, and on the right a \mathcal{C}_U and a \mathcal{C}_V pattern sharing both remaining neighbors. None of the depicted special vertices are settled.




Figure 13: Examples of **unsettled** vertices

4.3 Redirection procedure

To further decrease the number of problematic cases to consider in the rest of the proof, we consider a set of local transformations that, when applied exhaustively to a \mathcal{K} -subdivision S rooted on a 4-family U , return another \mathcal{K} -subdivision S' rooted on U , of the same type and which does not contain some inconvenient configurations. The special vertices of U can be more easily mapped to patterns in S' than in S .

This procedure does not preserve the chordlessness of a subdivision it is applied to, so let us consider the following weaker properties that are preserved by the procedure (as is proven in Claim 4.13, p. 54).

Given a \mathcal{K} -subdivision S rooted on U , an *A-chord* of S is a chord on a path of S incident with a special vertex $u \in U$ that is unsettled or forms a \mathcal{C}'_V pattern w.r.t. S (see Figure 14a); and a *B-chord* is a chord between two remaining neighbors of $u \in U$ on a path of S that is **not** incident with u (see Figure 14b). We say respectively that u has an A-chord, a B-chord.

A \mathcal{K} -subdivision satisfies *property A* (resp. *property B*) if it does not have an A-chord (resp. a B-chord). A chordless \mathcal{K} -subdivision obviously satisfies properties A and B.





Figure 14: The configurations that are avoided by properties A and B

Let us now define the procedure that helps take care of problematic cases in a \mathcal{K} -subdivision.

Definition 4.12 (Redirection procedure). *Let G be a planar graph with a (C_{II}) configuration w.r.t. a 4-family U , and S be a \mathcal{K} -subdivision rooted on U , such that S satisfies properties A and B. The redirection procedure consists in applying as many times as possible the redirection operations \mathcal{C}_{X1} , \mathcal{C}_{X2} , \mathcal{C}_{X3} and \mathcal{C}_{X4} to S .*

Note that these configurations are defined on u_1, u_2 but may be in contact with unspecified remaining neighbors of u_3, u_4 , in which case we apply the redirections anyway. The contacts with paths from S that would prevent us from applying the redirections are specified.


Remark: the \mathcal{C}_V patterns on the drawings illustrating the redirections could turn out to be \mathcal{C}_{T2NA} patterns, and are only featured as an indication.



Redirection \mathcal{C}_{X1} , when $w_1 = w_2$

- \mathcal{C}_{X1} : The vertices u_1, u_2 are linked by a path $u_1 \sim u_2$ of S of the form (u_1, w_1, Q, w_2, u_2) , with $w_1 = w_2$ if $l(Q) = 0$. The vertices u_1, u_2 have exactly one common remaining neighbor v and have another remaining neighbor v_1, v_2 respectively, both adjacent to v . The vertices v_1 and w_1 are non-adjacent. No path from S touches v, v_1 or v_2 .


Redirection protocol: We replace the path $u_1 \sim u_2$ in S with the path (u_1, v, u_2) .




Redirection \mathcal{C}_{X2} , when $w_1 = w_2$

- \mathcal{C}_{X2} : The vertices u_1, u_2 are linked by a path $u_1 \sim u_2$ of S of the form (u_1, w_1, Q, w_2, u_2) , with $w_1 = w_2$ if $l(Q) = 0$. The vertices u_1, u_2 have exactly one common remaining neighbor v and have another remaining neighbor v_1, v_2 respectively, both adjacent to v . The vertices v_1 and w_1 (resp. v_2 and w_2) are adjacent. No path from S touches v, v_1 or v_2 .

Redirection protocol: We replace the path $u_1 \sim u_2$ in S with the path (u_1, v_1, v, u_2) . The vertices v, w_1 are the new remaining neighbors of u_1 and are not adjacent, otherwise $\{u_1, v, v_1, w_1\}$ would form a K_4 , contradicting Claim 4.2 (p. 36).



Redirection \mathcal{C}_{X3} , when w_1, v_1 are not adjacent




Redirection \mathcal{C}_{X3} when w_1, v are not adjacent

- \mathcal{C}_{X3} : The vertices u_1, u_2 are linked by a path $u_1 \sim u_2$ of S , they have exactly one remaining neighbor v in common, and u_1, u_2 have another remaining neighbor v_1, v_2 respectively. v_1, v_2 are both adjacent to v . There is another special vertex u_3 such that there is a path $u_1 \sim u_3$ in S of the form $P = (u_1, w_1, Q, v_2, Q', u_3)$, with $l(Q) \geq 0$ (so w_1 may be equal to v_2) and $l(Q') \geq 1$. No path from S touches v nor v_1 .

Redirection protocol: The vertex w_1 cannot be adjacent to both v and v_1 , otherwise $\{u_1, v, v_1, w_1\}$ would induce a K_4 in the graph, a contradiction by Claim 4.2 (p. 36). We distinguish between two cases:

- If w_1, v_1 are **not adjacent**: we replace the path P in S with the path $P' = (u_1, v, v_2, Q', u_3)$;
- If w_1, v_1 are **adjacent**: then necessarily w_1, v are not, and in this case we replace P in S with the path $P' = (u_1, v_1, v, v_2, Q', u_3)$.

In both cases, the two new remaining neighbors of u_1 are not adjacent.



Redirection \mathcal{C}_{X4}

- \mathcal{C}_{X4} : The vertices u_1, u_2 are linked by a path $u_1 \sim u_2$ of the form (u_1, w_1, Q, w_2, u_2) , with $l(Q) \geq 0$ (with $w_1 = w_2$ if $l(Q) = 0$). The vertices u_1, u_2 have two adjacent remaining neighbors v, v' in common. No path from S touches v nor v' .

Redirection protocol: There must be a non-edge e among $vv_1, v'w_1$, otherwise $\{u_1, v, v', w_1\}$ form an induced K_4 , which contradicts the fact that G has a (\mathcal{C}_{II}) configuration by Claim 4.2 (p. 36). Let us say that $e = vv_1$. We replace the path $u_1 \sim u_2$ in S with the path (u_1, v', u_2) .

In all cases, the special vertex u_1 is given a new set of remaining neighbors that are not adjacent. The procedure terminates, as each redirection requires two special vertices that have adjacent remaining neighbors to be applied, and increases the number of special vertices with non-adjacent remaining neighbors.

We define the associated property: a subdivision satisfies **Property C** if no redirection can be applied. We say that a subdivision is **strong** if it satisfies properties A, B and C.

To justify the choice of this procedure and the notion of strong subdivision, we prove that it preserves properties A and B and the structure of \mathcal{K} -subdivision.

Claim 4.13. *The redirection procedure preserves properties A and B.*

Proof. Let G be a planar graph with a 4-family U and a \mathcal{K} -subdivision S rooted on U , such that S satisfies properties A and B. Let S' be the subdivision obtained by applying the redirection procedure to S . Let us prove that S' satisfies properties A and B as well.

- $\mathcal{C}_{X1}, \mathcal{C}_{X4}$: Redirection configurations \mathcal{C}_{X1} and \mathcal{C}_{X4} feature two special vertices u_1, u_2 and only modify one (u_1, u_2) -path in the subdivision, by replacing it with another of length 2. No path of length 2 has a B-chord, and since an edge $u_1u_2 \in E(G)$ already constitutes a path of S , the new path does not have an A-chord either. Since the paths of S are internally disjoint, the vertices w_1, w_2 do not belong to S' and thus cannot be part of an A-chord or a B-chord.
- \mathcal{C}_{X2} : In redirection configuration \mathcal{C}_{X2} , the new (u_1, u_2) -path P' has length 3, hence does not contain B-chords. Since the remaining neighbors of u_1, u_2 w.r.t. S' do not belong to S' except one (v), the other paths of S' do not contain B-chords either. The remaining neighbors of u_2 are disjoint from P' , hence u_2 does not have an A-chord, but u_1 does however. We claim that property A is still satisfied because u_1 is (lone-)settled w.r.t. S' . The remaining neighbors of u_1 are not adjacent, hence u_1 is unsettled only if it forms a \mathcal{C}_{T2NA} pattern with u_3 or u_4 (this pattern would then touch S'), since u_3, u_4 do not form a \mathcal{C}_U pattern as S satisfies property A.

Let us first consider the case where S is a K_4 -subdivision, and assume u_3 has w_1, v as its remaining neighbors w.r.t. S' (it has the same remaining neighbors w.r.t. S , since only the remaining neighbors of u_1, u_2 were modified). Then $\{u_1, w_1, v\}$ and $\{v_1, u_2 = u_4, u_3\}$ induce a $K_{3,3}$ -minor of G (by contracting the path $u_2 \sim u_4$ to a vertex, see Figure 15), contradicting the planarity of G .




Figure 15: The $K_{3,3}$ -minor formed by $\{u_1, w_1, v\}$ and $\{v_1, u_2 = u_4, u_3\}$ if S is a K_4 -subdivision (the path $u_2 \sim u_3$ is not pictured)

Now let us take a look at the case where S is a C_{4+} -subdivision and assume that u_3 or u_4 has w_1, v as its remaining neighbors w.r.t. S' (thus w.r.t. S). If u_1, u_2 are 1-linked, u_3 or u_4 has a remaining neighbor in the solo (u_1, u_2) -path P , contradicting the property “1-linked” of S . If u_1, u_2 are 2-linked, then $\{u_1, w_1, v\}$ and $\{v_1, u_2, u_3\}$ induce a $K_{3,3}$ -minor (see Figure 16), again a contradiction. Thus, u_1 cannot form a \mathcal{C}_{T2NA} pattern, so forms a \mathcal{C}_V pattern and is lone-settled.




Figure 16: The $K_{3,3}$ -minor formed by $\{u_1, w_1, v\}$ and $\{v_1, u_2, u_3\}$ if S is a C_{4+} -subdivision

- \mathcal{C}_{X3} : The remaining neighbors v, v_2 of u_2 w.r.t. S' belong to the new (u_1, u_3) -path P' of S' and the edge vv_2 belongs to P' , so u_2 does not have an A-chord nor a B-chord. The remaining neighbors of u_3, u_4 are not modified by the redirection. The paths of S incident with u_4 are not modified, so u_4 does not have an A-chord, and a B-chord of u_4 could only belong to the new path P' of S' . An A-chord of u_3 could only belong to P' , as its other incident paths of S were not modified, and u_3 does not have a B-chord, as its non-incident paths of S were not modified. We claim that none of u_3 and u_4 have an A-chord or B-chord on P' in S' if they did not in S .

We examine the case of u_1 at the end of this proof. Let us make several observations that will help us prove our claims.

Whether S is a K_4 - or a C_{4+} -subdivision, it contains a (u_1, u_2) -path P_{12} , a (u_2, u_4) -path P_{24} , a (u_3, u_4) -path P_{34} , as well as the path $P = P_{13}$, split into a (u_1, v_2) -section Q_{13} and a (v_2, u_3) -section T_{13} , all these paths having no special vertex as internal vertex.

Observation 2. *The special vertex u_3 does not have v_1 as a remaining neighbor w.r.t. S .*

Proof. If it were the case, then G would contain a $K_{3,3}$ -minor induced by $\{u_1, u_3, v\}$ and $\{u_2, v_1, v_2\}$, obtained by contracting P_{12} , P_{34} , Q_{13} and T_{13} to one edge, and P_{24} to one vertex (see Figure 17). This would contradict the planarity of G . \square




Figure 17: The $K_{3,3}$ -minor formed by $\{u_1, u_3, v\}$ and $\{u_2, v_1, v_2\}$ in Observation 2

Observation 3. *The special vertex u_3 does not have v as a remaining neighbor w.r.t. S .*

Proof. Because the paths P_{24} and P_{34} are disjoint from v, v_1, v_2 , the vertices u_1, u_3 would then belong to two different regions of the plane delimited by the three edges u_2v, u_2v_2, vv_2 (see Figure 18). This is a contradiction with the almost 4-connectivity of G . \square




Figure 18: The planar embedding of the graph of Observation 3

Observation 4. *The special vertex u_4 does not have two remaining neighbors (w.r.t. S) v_4 in T_{13} , and $v'_4 \in \{v, v_1\}$.*

Proof. If $v'_4 = v_1$, contracting the (v_2, v_4) -section of T_{13} gives us a $K_{3,3}$ -minor of G induced by $\{u_1, v, u_4\}$ and $\{u_2, v_1, v_2\}$, a contradiction (see Figure 19a). If $v'_4 = v$, then by planarity u_1, u_4 must belong to two different regions of the plane delimited by the three edges u_2v, vv_2 and u_2v_2 (see Figure 19b), again a contradiction to the almost 4-connectivity of G . \square




Figure 19: The $K_{3,3}$ -minor and the planar embedding of Observation 4

Observation 5. *The special vertex u_4 does not have w_1, v as its remaining neighbors w.r.t. S if w_1, v are non-adjacent.*

Proof. If it were the case, there would be a $K_{3,3}$ -minor in G induced by $\{u_1, u_4, v_2\}$ and $\{u_2, w_1, v\}$ (see Figure 20), a contradiction with the planarity of G (note that by assumption w_1, v are non-adjacent, hence $w_1 \neq v_2$). \square




Figure 20: The $K_{3,3}$ -minor formed by $\{u_1, u_4, v_2\}$ and $\{u_2, w_1, v\}$ in Observation 5

Since only the (u_1, u_3) -path P changes into the path P' , and the remaining neighbors of u_3, u_4 are not changed by the redirection, u_3 (resp. u_4) may only have an A-chord (resp. B-chord) on the new path P' . This (u_1, u_3) -path is incident with u_3 so u_3 does not have

a B-chord on it, and no A-chord either by Observations 2 and 3. Note that if u_3 has an A-chord in S , then u_3 forms a \mathcal{C}_V pattern w.r.t. S , and the same pattern w.r.t. S' .

P' is not incident with u_4 , so u_4 does not have an A-chord on it, and Observation 4 tells us that it does not have a B-chord either.

As for u_1 , note that in the version with w_1, v_1 non-adjacent, the new remaining neighbors of u_1 are disjoint from S , thus u_1 forms a \mathcal{C}_V or \mathcal{C}_{T2NA} pattern disjoint from S (since no pair of special vertices form a \mathcal{C}_U pattern by property A), hence is settled. In the version with w_1, v non-adjacent, the edge u_1v is an A-chord of u_1 , but Observations 3 and 5 tell us that neither u_3 nor u_4 can form a \mathcal{C}_{T2NA} pattern with u_1 . Again, since no pair of special vertices form a \mathcal{C}_U pattern by property A of S , u_1 is left lone-settled by the redirection, and property A is satisfied by S' . □

Claim 4.14. *Let G be a planar graph with a (C_{II}) configuration w.r.t. a 4-family U . Then G contains a strong \mathcal{K} -subdivision rooted on U .*

Proof. By Lemma 4.4 (p. 37), G contains a chordless \mathcal{K} -subdivision S rooted on U . In particular, S satisfies properties A and B. Since by Claim 4.13 (p. 54) the redirection procedure preserves properties A and B, and since the type of subdivision (K_4 or C_{4+}) is preserved by each redirection operation, then the result follows if S is a K_4 -subdivision.

Now assume that S is a C_{4+}^* -subdivision. We prove that after application of any redirection operation to S , the obtained subdivision S' is a C_{4+}^* -subdivision, and the result follows by induction. By Claim 4.13 (p. 54), S' satisfies properties A and B. Observe that the redirection operations preserve the ends of the paths of S ; in particular, vertices that are k -linked stay k -linked after application of an operation. Observe that in each redirection configuration, the two special vertices u_1, u_2 involved have a common remaining neighbor that is disjoint from S . Therefore, by property “2-linked”, the two special vertices u_1, u_2 involved in the configuration cannot be 2-linked: they are necessarily 1-linked.

Let us prove the three properties of C_{4+}^* of S' in order.

- *Property “0-linked”:* The only special vertices whose remaining neighbors are modified by a redirection operation are the special vertices u_1 and u_2 involved in the redirection configuration. Therefore, if u_j, u_k are 0-linked special vertices, then their remaining neighbors w.r.t. S' are the same as w.r.t. S , and property “0-linked” of S' is implied by the same property of S .
- *Property “1-linked”:* Let u_i, u_j be 1-linked special vertices associated with a path P_{ij} of S , and $u_k \in U \setminus \{u_i, u_j\}$. Assume for contradiction that u_k has a remaining neighbor v_k w.r.t. S' that is an internal vertex of the (u_i, u_j) -path P'_{ij} of S' . If v_k is not a remaining neighbor of u_k w.r.t. S , then the operation applied to S involves u_k , and the edge u_kv_k belongs to S . However, we can check in all redirection operations that when an edge uv incident with a special vertex u is removed from the subdivision, then v does not belong to the subdivision after the operation is applied. This is a contradiction with the definition of v_k , therefore v_k is indeed a remaining neighbor of u_k w.r.t. S . Thus, $P'_{ij} \neq P_{ij}$.

Since u_k is 0-linked with one of u_i, u_j , the vertex v_k cannot be a remaining neighbor of both w.r.t. S , by “0-linked” property of S . So the operation applied to S cannot be \mathcal{C}_{X1} or \mathcal{C}_{X4} , as their new path has length 2.

The operation cannot be \mathcal{C}_{X3} either, since in this case u_i, u_j must be the vertices u_1, u_3 in the definition of this configuration (so that P'_{ij} is the new path); u_i, u_j are 1-linked, and the operation is applied on the vertices u_1, u_2 , which are also 1-linked as mentioned above. Thus, u_i or u_j is 1-linked to two different special vertices, a contradiction.

If the operation is \mathcal{C}_{X2} , then v_k cannot be the common remaining neighbor v of u_1, u_2 in the definition, since u_k is 0-linked to one of them. If v_k is the v_1 of \mathcal{C}_{X2} , then $\{u_k, v, w_1\}$ and $\{u_1, u_2, v_1\}$ induce a $K_{3,3}$ -minor in G (by contracting the (w_1, u_2) -section of $u_1 \sim u_2$ to an edge, and contracting a (u_k, u_2) -path of S or a (u_k, u_m) -path and a (u_m, u_2) -path of S to an edge, for $u_m \in U \setminus \{u_1, u_2, u_k\}$), a contradiction to the planarity of G .

- *Property “2-linked”*: Let u_i, u_j be two 2-linked special vertices. Since the operations are applied on 1-linked special vertices, exactly one of these two special vertices, say u_i , is involved in the operation (as u_1 or u_2 in the definitions) applied to S . Let v be a common remaining neighbor of u_i, u_j w.r.t. S' . If v was not a common remaining neighbor of u_i, u_j w.r.t. S , then it was a remaining neighbor of u_j and not u_i , as u_j is not involved in the operation: the edge $u_i v$ belongs to S , and more precisely belongs to a solo (u_i, u_k) -path P_{ik} of S between the two 1-linked special vertices u_i and $u_k \in U \setminus \{u_i, u_j\}$. Thus, u_j has a remaining neighbor (w.r.t. S) that is an internal vertex of a solo path of S , a contradiction with the “1-linked” property of S .

Therefore, the potential common remaining neighbors v, v' of u_i, u_j w.r.t. S' are also their common remaining neighbors w.r.t. S . So by property “2-linked” of S , u_i, u_j have at most one remaining neighbor v , and it belongs to a parallel (u_k, u_l) -path P_{kl} of S that is not incident with u_i, u_j . If the path P_{kl} was not modified by the operation, the result follows. If P_{kl} was modified into a path P'_{kl} of S' , then the operation is necessarily \mathcal{C}_{X3} (as in the others, only a solo path is modified). The special vertex u_i is the u_2 in the definition, as it keeps the same special neighbor v_2 in S and S' . This vertex belongs to both P_{kl} and P'_{kl} , and the result follows.

This proves that S' is a C_{4+}^* -subdivision, and the result follows by induction. \square

4.4 Sufficiency of the (C_{II}) rules

As with the (C_I) configurations, we need to make sure the composite rules we define in this section can indeed be applied, and yield a contradiction with the existence of their associated configuration in an MCE. Let us first prove that the rules we will define throughout this section allow us to find a good coloring of an MCE, thus a contradiction, similarly to Lemma 3.6 (p. 24).

Lemma 4.15. *An MCE with a 4-family U does not contain a subdivision composite configuration made up of a semi-subdivision S rooted on U and a mapping \mathcal{M} of U compatible w.r.t. S .*

Proof. Let G be such an MCE, and assume it contains the composite configuration $X = (\{\mathcal{C}_1, \dots, \mathcal{C}_k\}, S)$ where $\{\mathcal{C}_1, \dots, \mathcal{C}_k\}$ is a compatible mapping of U . The associated composite rule $\mathcal{R}_X = (X, f_X^r, f_X^c)$ is thus valid. We build a good coloring c of G to show a contradiction.

Let us first build a coloring pc of G' using the right number of colors. Similarly to the proof of Lemma 3.6 (p. 24), let us color each K_3 component of G' with a cycle of length 3 and each K_5^- component with a cycle of length 5 and a path of length 4. We color each other components with a good coloring. Thus, pc uses the right number of colors $(\lfloor \frac{|V(G')|}{2} \rfloor)$, with a mix of cycles and paths. Let $c_0 = f_X^c(G, pc)$ be the coloring of G obtained from pc by \mathcal{R}_X . Since \mathcal{R}_X is valid, c_0 has the right number of colors $(\lfloor \frac{|V(G)|}{2} \rfloor)$ again with some cycles instead of paths. We build iteratively a good coloring c of G , by starting from c_0 and using Lemma 3.5 (p. 24) to successively replace a pair of colors, inducing a path and a cycle, by another pair of colors inducing two paths.

Observe that the cycles induced by colors of c_0 are disjoint in G . Indeed, the cycles in pc are disjoint because they belong to different connected components; the cycles of pc may have been deviated into longer cycles in c_0 , but since the internal vertices of the deviated sections

are all special vertices, and since each special vertex is involved in at most one deviation, then no vertex of G can belong to the intersection of two cycles in c_0 .

First let us prove that no K_3 connected component can appear in G' . Let K be such a component, colored with a cycle in pc , and let C be the cycle of G induced by the same color in c , after some deviations. Observe that if one pattern involved in the deviations is \mathcal{C}_{T1} , \mathcal{C}_{T2A} or \mathcal{C}_{T2NA} , one vertex of K is supposed to have odd degree in G' , which is impossible since K is a K_3 . Since K is a connected component of G' , the vertices of C on $V(K)$ are only incident with edges of $G[V(K)]$, edges between special vertices and their remaining neighbors, and edges from the subdivision S . A semi-subdivision has at most one pair of intersecting paths, sharing exactly one vertex. Hence, there are at least two vertices in $V(K)$ which are incident with 2 edges of C and 0 or 2 edges of S , and so these two vertices have a degree of at most 4 in G . This contradicts Lemma 3.2, so no K_3 component can be created in G' .

Now let K be a K_5^- component of G' , and let C', P' be the cycle and path coloring it in G' . Let C, P be the cycle and path induced in c_0 by the same colors as C', P' . C has not been treated yet and is thus induced by the same color in c as in c_0 . If $P = P'$, then it is disjoint from the cycles treated in previous iterations of c , and thus has not been involved in a replacement of a pair of colors. Otherwise, P results from deviations of P' on special vertices, or an extension of P' by at most two edges (one for each of its endpoints) to special vertices. If u is a special vertex touching P , since u belongs to exactly one pattern of the mapping of X , then u does not touch a cycle treated in any previous iteration of c . Thus P is again disjoint from the cycles treated in the previous iterations. In both cases, P is induced by the same color in c as in c_0 .

Observe that $V(C) \cap V(P) \subseteq V(K)$, so $|V(C) \cap V(P)| \leq 5$. We distinguish between three cases:

- **C results from a deviation of C' :** then its length is different from 5. By Observation 1 (p. 24), $C \cup P$ does not form the exceptional graph.
- **C' has not been deviated, but P' has:** then $V(C) = V(K)$, and there is an edge of K that does not belong to $(C \cup P)[V(C)]$. Thus $(C \cup P)[V(C)] \subsetneq (C' \cup P')[V(C')] = K$, and so $(C \cup P)[V(C)]$ does not form a K_5^- , and by Observation 1 (p. 24), $C \cup P$ does not form the exceptional graph.
- **Neither C' nor P' have been deviated:** then $(C \cup P)[V(C)] = K$, which is a K_5^- , and so $E(K) \subseteq E(G)$. The edges of K split G into 6 regions bounded by triangles. Since the graph is almost 4-connected w.r.t. all the special vertices, they belong to the same region of the graph. Hence at most 3 vertices from K are neighbors of special vertices. Since there is at most one vertex of degree at most 4 in G , all vertices from K , except maybe one, have their degree changed between G and G' . Only 3 of them can belong in patterns, thus at least one of them is touched by a path Q induced by a color of c and different from P . Since $C \cup P$ forms the exceptional graph, by Observation 1 (p. 24) $C \cup Q$ does not, and since C' has not been deviated, $|C| = 5$, and so $|V(C) \cap V(Q)| \leq 5$.

In all three cases, Lemma 3.5 (p. 24) gives us a decomposition of $C \cup P$ or $C \cup Q$ into two new paths Q', Q'' . We replace C, P or C, Q in c , depending on the case, with Q', Q'' .

The coloring c contains the same number of colors as c_0 in all iterations, with one less cycle at each iteration. When all K_3 and K_5^- components have been treated, the resulting coloring c is a good path decomposition of G , a contradiction. \square

In order to show a contradiction, we now need to prove that an MCE containing a configuration (C_{II}) indeed contains a subdivision composite configuration made up of a semi-subdivision rooted on its 4-family and a compatible mapping.

Lemma 4.16. *Let G be a planar graph with a (C_{II}) configuration w.r.t. a 4-family U , that admits a strong \mathcal{K} -subdivision rooted on U . Then G contains a subdivision composite configuration made up of a semi-subdivision S rooted on U and a compatible mapping w.r.t. S .*

Note that the subdivision is a semi-subdivision and so it may have missing edges or having two paths crossing, but it is 2-colorable, and therefore sufficient to produce a good coloring of G if G is an MCE, and therefore show a contradiction.

The rest of this section constitutes a proof of this lemma, before the conclusion in which we show that Lemma 4.1 (p. 35) ensues.

4.5 Distant problems

In the following, we prove that the graph admits a composite configuration made up of a subdivision and a set of compatible patterns. We will distinguish two types of “problems” that could occur and prevent us from applying directly a reduction $\{\mathcal{C}_N, \mathcal{C}_N, \mathcal{C}_N, \mathcal{C}_N\}$. First, a \mathcal{C}_N pattern could cause a “distant problem” by touching a path of the subdivision, and the associated reduction rule could create a cycle in the coloring. Then, some special vertices from U could cause a “close problem” by sharing some of their remaining neighbors and the \mathcal{C}_N patterns would not be compatible. We first treat the cases with at least 3 distant problems (Lemma 4.21, p. 67), then the cases with at most 2 distant problems and no close problems (Lemma 4.23, p. 76) and finally the cases with at most 2 distant problems and some close problems (Lemma 4.25, p. 86).

Definition 4.17 (Distant problem). *Let G be a planar graph with a 4-family U and let S be a strong \mathcal{K} -subdivision rooted on U . Let $u \in U$ be a special vertex and P be a path of S that is not incident with u . We say that u causes a distant problem on P if the three conditions are satisfied:*

- u has two adjacent remaining neighbors v, v' that are disjoint from U ;
- exactly one of its remaining neighbors belongs to P ;
- if some other special vertex u' has one of v, v' as a remaining neighbor, then u' is settled.




Figure 21: Distant problem caused by a special vertex u

Figure 21 provides an example of distant problem. Only the path P from the definition is represented.

This definition is motivated by the fact that because of property B, an unsettled special vertex u cannot have both its remaining neighbors belong to a path P not incident with u .

Note that if three special vertices cause distant problems, the fourth one is lone-settled, because of the last condition in the definition of distant problem and by properties A and B.

Once we 2-color the subdivision S such that each vertex of U is at the end of a color, a distant problem is *active* if the color that ends on u is the same as the color of the path P . A

distant problem is otherwise *inactive*. We can treat inactive distant problems as \mathcal{C}_N patterns, as the coloring fits the color requirements of the pattern (i.e. the color requirements of the two patterns (\mathcal{C}_{Ne}) and (\mathcal{C}_{No})). Since the distant problems are caused by special vertices u which have their remaining neighbors v, v' adjacent, we refer to $\{u, v, v'\}$ as the *triangle of u* .

We prove here two results that are convenient for the rest of the proof.

Claim 4.18. *A K_4 -subdivision can be 2-colored so as to inactivate up to 2 distant problems that are on different paths of S .*

Proof. Let S be a K_4 -subdivision rooted on $\{u_1, u_2, u_3, u_4\}$. Let us assume w.l.o.g. that u_1 causes a distant problem on the path $u_2 \sim u_3$. The following 2-coloring of S inactivates the distant problem of u_1 : $\{(u_3 \rightarrow u_2 \rightarrow u_1 \rightarrow u_4), (u_1 \rightarrow u_3 \rightarrow u_4 \rightarrow u_2)\}$.

If S has a second distant problem on a different path, then there are several possible cases. Either u_3 causes a distant problem on $u_2 \sim u_4$ (case A) or on $u_1 \sim u_4$ (case B). These cases are symmetric with the ones where u_2 causes a distant problem on $u_3 \sim u_4$ and $u_1 \sim u_4$ respectively. If instead the second distant problem is caused by u_4 on (w.l.o.g.) $u_1 \sim u_3$, then this is equivalent to case A: just replace $(1, 2, 3, 4)$ with $(3, 4, 2, 1)$ to obtain case A. The coloring of the previous case inactivates the two distant problems of case A, and the coloring $\{(u_4 \rightarrow u_1 \rightarrow u_3 \rightarrow u_2), (u_1 \rightarrow u_2 \rightarrow u_4 \rightarrow u_3)\}$ of S inactivates those of case B. \square

Claim 4.19. *Let G be a planar graph with a (C_{II}) configuration w.r.t. a 4-family U and a strong C_{4+}^* -subdivision rooted on U , such that G does not have a K_4 -subdivision rooted on U . If $u_i, u_j \in U$ are 1-linked, then at most one of them has a remaining neighbor belonging to S .*

Proof. By property “1-linked” of S and property A, the remaining neighbors of u_i, u_j are on parallel paths of S . Let us assume for contradiction that the 1-linked special vertices u_1, u_2 are such that u_1 has a remaining neighbor v_1 on a (u_2, u_4) -path P_2 of S and u_2 has a remaining neighbor v_2 on a (u_1, u_3) -path P_1 of S : let $P_1 = (u_1, Q_1, v_2, Q_3, u_3)$ and $P_2 = (u_2, Q_2, v_1, Q_4, u_4)$.

Let S' be the set of paths of S different from P_1, P_2 , and let $P'_1 = (u_1, v_1, Q_4, u_4)$ and $P'_2 = (u_2, v_2, Q_3, u_3)$, as depicted on Figure 22. These (u_1, u_4) -path and (u_2, u_3) -path are internally disjoint from the paths of S' , hence $S' \cup \{P'_1, P'_2\}$ forms a K_4 -subdivision of G rooted on U , a contradiction. Hence, two 1-linked special vertices cannot both have a remaining neighbor in S .

Finally, S cannot have three distant problems or more, as two of them would be caused by a pair of 1-linked special vertices. \square

We define below the “distant configurations” that correspond to subdivisions with at least 3 distant problems. We first introduce a *routing operation* that helps us take care of these distant problems. For each distant configuration, we perform a redirection of the subdivision S into a subdivision S' , to turn each special vertex causing a distant problem into a settled one. When S has a distant problem caused by u_i and S' has a new path of the form (u_i, v_i, Q, u_j) , the goal is to turn the distant problem on u_i into a \mathcal{C}_V pattern. To do so, we use the following operation.

Definition 4.20 (Routing operation). *Let G be a planar graph with a 4-family U and a strong \mathcal{K} -subdivision S rooted on U . Let $u \in U$ have two adjacent remaining neighbors v_1, v_2 w.r.t. S , and let w be a neighbor of u that belongs to S . Assume that u causes a distant problem, with v_1 touching a path of S . One of v_1, v_2 , say v' , is not adjacent to w , otherwise $\{u, v_1, v_2, w\}$ would form an induced K_4 in the graph, which contradicts the fact that G has a (C_{II}) configuration by Claim 4.2 (p. 36).*




Figure 22: Display of the underlying K_4 -subdivision of G rooted on U

The paths of S are redirected to create a new subdivision S' , containing a path $P' = (u, v_1, Q, u')$, and such that w does not belong to S' . Applying the routing operation on u consists in replacing the edge uv_1 in P' by the edges uv_2, v_2v_1 if $v' = v_1$, and leaving P' as it is otherwise.




Figure 23: On the left: u causes a distant problem on a path $P = u' \sim u''$ of the subdivision. In the middle: a new subdivision is considered for some reduction rule, with a new path $P' = u' \sim u$. On the right: the routing operation modifies the path $P' = u' \sim u$ in order to choose two non-adjacent vertices (v_1, w) as remaining neighbors for u .

This routing operation ensures that the two remaining neighbors of u in S' are w, v' and are thus non-adjacent. The vertex u now forms a \mathcal{C}_V pattern w.r.t. S' and we justify for each application of the routing operation that u is left settled. For all cases, we provide a subdivision S and describe a mapping of compatible patterns that settles all vertices.

List of the distant configurations:

The distant configurations are the configurations D_1, D_2, D_3, D_4 listed below.

Each configuration describes a 4-family U and a strong K -subdivision S , such that at least 3 special vertices of U cause a distant problem on S . For each configuration, we describe a new semi-subdivision S' . The routing operation is applied to S' for all unsettled special vertices.

We provide for each configuration a subdivision composite rule made up of C_V and C_N patterns. We justify for each rule that the mapping is compatible w.r.t. S' .




Figure 24: Semi-subdivision of D_1

Configuration D_1

Properties:

- The graph has a strong K_4 -subdivision S rooted on u_1, u_2, u_3, u_4
- u_1 causes a distant problem on the path $u_2 \sim u_4$: it has a remaining neighbor v_1 such that $u_2 \sim u_4 = (P_2^1, v_1, P_4^1)$
- u_2 causes a distant problem on the path $u_3 \sim u_4$: it has a remaining neighbor v_2 such that $u_3 \sim u_4 = (P_3^2, v_2, P_4^2)$
- u_3 causes a distant problem on the path $u_1 \sim u_4$: it has a remaining neighbor v_3 such that $u_1 \sim u_4 = (P_1^3, v_3, P_4^3)$

We transform the K_4 -subdivision S into another K_4 -subdivision S' , by removing the paths $u_1 \sim u_4, u_2 \sim u_4, u_3 \sim u_4$, and adding the paths $(u_1, v_1, P_4^1), (u_2, v_2, P_4^2), (u_3, v_3, P_4^3)$.

After the routing operation is applied, all unsettled special vertices are turned into C_V patterns.

Remark: u_4 could not form a C'_V pattern in S by property A and planarity, hence it stays lone-settled in S' .




Figure 25: Semi-subdivision of D_2

Configuration D_2

Properties:

- The graph has a strong K_4 -subdivision S rooted on u_1, u_2, u_3, u_4
- u_2 causes a distant problem on the path $u_1 \sim u_3$: it has a remaining neighbor v_2 such that $u_1 \sim u_3 = (P_1^2, v_2, P_3^2)$
- u_3 causes a distant problem on the path $u_2 \sim u_4$: it has a remaining neighbor v_3 such that $u_2 \sim u_4 = (P_2^3, v_3, P_4^3)$
- u_4 causes a distant problem on the path $u_1 \sim u_2$: it has a remaining neighbor v_4 such that $u_1 \sim u_2 = (P_1^4, v_4, P_2^4)$

We transform the K_4 -subdivision S into a C_{4+} -subdivision S' , by removing the paths $u_1 \sim u_2, u_1 \sim u_3$ and $u_2 \sim u_4$, and adding the paths $(u_2, v_2, P_1^2, u_1), (u_3, v_3, P_2^3, u_2), (u_4, v_4, P_1^4, u_1)$.

After the routing operation is applied, all unsettled special vertices are turned into \mathcal{C}_V patterns.

Remark: u_1 could only form a \mathcal{C}'_V pattern in S on the path $u_3 \sim u_4$ by property A and planarity, and this path was not modified in S' . Hence, u_1 remains lone-settled in S' .




Figure 26: Reduction of configuration D_3

Configuration D_3

Properties:

- The graph has a strong K_4 -subdivision S rooted on u_1, u_2, u_3, u_4
- u_1 and u_2 cause distant problems on the path $u_3 \sim u_4$: they have remaining neighbors v_1, v_2 respectively, such that $u_3 \sim u_4 = (P_3^{12}, v_1, P_4^{12}, v_2, P_4^{12})$
- u_3 causes a distant problem on the path $u_1 \sim u_2$: it has a remaining neighbor v_3 such that $u_1 \sim u_2 = (P_1^3, v_3, P_2^3)$
- **Remark:** u_4 may cause a distant problem on $u_1 \sim u_2$; it has a remaining neighbor v_4 that may belong to P_1^3 or P_2^3

We transform the K_4 -subdivision S into a C_{4+} -subdivision S' by removing the paths $u_1 \sim u_2$ and $u_3 \sim u_4$, and adding the paths (u_1, P_1^3, v_3, u_3) and $(u_2, v_2, P_4^{12}, u_4)$. We consider the following 2-coloring of S' : $\{red = (u_1 \rightarrow u_3 \rightarrow u_2 \rightarrow u_4), blue = (u_2 \rightarrow v_2 \rightarrow u_4 \rightarrow u_1 \rightarrow v_3 \rightarrow u_3)\}$. There is no need to apply the routing operation.

The special vertex u_3 is turned into a C_V pattern. The special vertices u_1, u_4 are treated as C_N patterns. The C_N pattern of u_4 may cause a distant problem on the new path (u_1, P_1^3, v_3, u_3) , but this path is colored blue and u_4 uses the color red, hence the distant problem is inactive.

The patterns used are $C_N(u_1), C_V(u_2), C_N(u_3), C_N(u_4)$.




Figure 27: Semi-subdivision of D_4

Configuration D_4

Properties:

- The graph has a strong K_4 -subdivision S rooted on u_1, u_2, u_3, u_4
- u_1 causes a distant problem on the path $u_2 \sim u_4$: it has a remaining neighbor v_1 such that $u_2 \sim u_4 = (P_2^1, v_1, P_4^1)$
- u_2 causes a distant problem on the path $u_1 \sim u_3$: it has a remaining neighbor v_2 such that $u_1 \sim u_3 = (P_1^2, v_2, P_3^2)$
- u_3 causes a distant problem on the path $u_1 \sim u_4$: it has a remaining neighbor v_3 such that $u_1 \sim u_4 = (P_1^3, v_3, P_4^3)$
- u_4 causes a distant problem on the path $u_2 \sim u_3$: it has a remaining neighbor v_4 such that $u_2 \sim u_3 = (P_2^4, v_4, P_3^4)$

We transform the K_4 -subdivision S into another K_4 -subdivision S' by keeping the paths $u_1 \sim u_2$ and $u_3 \sim u_4$ from S and adding the paths (u_1, v_1, P_4^1, u_4) , (u_2, v_2, P_3^2, u_3) , (u_3, v_3, P_1^3, u_1) , (u_4, v_4, P_2^4, u_2) .

By planarity and definition of distant problem, the routing operation does not need to be applied for all the special vertices to be turned into C_V patterns.

The following lemma shows how a planar graph with (C_{II}) configuration can be treated with one of the distant configurations if the associated subdivision has at least three distant problems.

Lemma 4.21 (Distant lemma). *Let G be a planar graph with a (C_{II}) configuration w.r.t. a 4-family U , with a strong K -subdivision S rooted on U . If G has at least 3 distant problems w.r.t. S , then G contains a configuration among $\{D_1, D_2, D_3, D_4\}$.*

Proof. If S is a C_{4+} -subdivision, we may assume that G does not have a K_4 -subdivision rooted on U . Then Claim 4.19 (p. 62) tells us that S cannot have 3 distant problems or more, which contradicts our hypothesis. Hence S is a K_4 -subdivision.

Let us call i -path a path of S that touches exactly i triangles of special vertices. We consider the three quantities, for $i \in \{0, 1, 2\}$, $p_i := |\{i\text{-paths}\}|$. By property A, a special vertex u can

only cause a distant problem on one of the three paths of S that are not incident with it: we call these paths the *potential paths* of u . For the same reasons, a path of S can touch at most 2 triangles of special vertices.

We have $p_0 + p_1 + p_2 = 6$ and $p_1 + 2 \cdot p_2 = \text{number of distant problems} = 3$ or 4 . Hence we need to consider five cases, depending on whether there are 3 or 4 distant problems and on the values the p_i parameters.

- **3 distant problems**, $p_1 = 3, p_2 = 0$. We have $p_0 = 3$. If all three 0-paths are incident with say u_4 , then there cannot be three 1-paths. Indeed, u_3 would have to cause a distant problem on $u_1 \sim u_2$, then by planarity none of u_1, u_2 could cause a distant problem on $u_2 \sim u_3, u_1 \sim u_3$ respectively.

Now let us assume that the three 0-paths form a subdivision of a triangle rooted on $\{u_1, u_2, u_3\}$. Thus u_4 cannot cause any distant problem. The three 1-paths are $u_1 \sim u_4, u_2 \sim u_4, u_3 \sim u_4$. Let us say w.l.o.g. that u_3 causes a distant problem on $u_1 \sim u_4$. Thus by planarity u_2 causes a distant problem on $u_3 \sim u_4$, and u_1 causes a distant problem on $u_2 \sim u_4$. This is the configuration D_1 .

Let us finally assume that the three 0-paths form a subdivision of a path on three edges, rooted on U . Let us say that the 0-paths are $u_1 \sim u_4, u_3 \sim u_4, u_2 \sim u_3$. The path $u_2 \sim u_4$ touches the triangle of either u_1 or u_3 . If it touches the triangle of u_3 , then u_1 cannot cause any distant problem (as its two other potential paths are 0-paths), thus in this case both u_2 and u_4 cause a distant problem. There is only one possibility for u_2 : it causes a distant problem on $u_1 \sim u_3$; then there is only one possibility for u_4 , the path $u_1 \sim u_2$. This is the configuration D_2 . Now assume that instead, $u_2 \sim u_4$ touches the triangle of u_1 . By planarity and property A, only u_3 can touch the 1-path $u_1 \sim u_2$. Again by planarity, only u_4 can touch the 1-path $u_1 \sim u_3$. This case is equivalent to D_2 : (u_1, u_2, u_3, u_4) in D_2 correspond to (u_2, u_1, u_4, u_3) in this case, in this order.

- **3 distant problems**, $p_1 = 1, p_2 = 1$. We have $p_0 = 4$: the four 0-paths can only either form a subdivision of the “paw” graph (a triangle with an additional edge attached to one vertex) or a subdivision of the cycle on four vertices. We can easily see that the first case is impossible: let us say the non-0-paths are $u_1 \sim u_2, u_1 \sim u_3$; there must be a path that touches two triangles of U , say it is $u_1 \sim u_2$, that necessarily touches the triangles of u_3 and u_4 . Then by planarity, the path $u_1 \sim u_3$ cannot touch the triangle of u_2 and thus cannot be a 1-path. Hence the 0-paths cannot form a paw.

Now let us assume that the 0-paths are $u_1 \sim u_3, u_2 \sim u_3, u_1 \sim u_4, u_2 \sim u_4$. Let us assume w.l.o.g. that each of u_1, u_2 causes a distant problem on $u_3 \sim u_4$, and u_3 causes a distant problem on $u_1 \sim u_2$. This case can be treated as configuration D_3 .

- **4 distant problems**, $p_1 = 4, p_2 = 0$. We have $p_0 = 2$, so the two 0-paths can either be incident on one vertex or disjoint. Let us consider the first case. Assume that $u_1 \sim u_4, u_2 \sim u_4$ are the 0-paths; the other four paths are 1-paths and must each touch one triangle of U , so all of u_1, u_2, u_3, u_4 cause a distant problem. The vertex u_3 causes a distant problem on $u_1 \sim u_2$, as it is its only potential path. Then w.l.o.g. u_4 causes a distant problem on $u_2 \sim u_3$, and u_1 can only cause a distant problem on $u_3 \sim u_4$. Finally, by planarity the triangle of u_2 cannot touch the path $u_1 \sim u_3$, its only potential path left. Hence, the 0-paths cannot be incident.

Now let us assume that the two 0-paths are $u_1 \sim u_2, u_3 \sim u_4$. Assume w.l.o.g. that u_1 causes a distant problem on $u_2 \sim u_4$. Then u_3 causes a distant problem on $u_1 \sim u_4$ as it is its last potential path. In the same way, u_2 causes a distant problem on $u_1 \sim u_3$ and u_4 on $u_2 \sim u_3$. This is the configuration D_4 .

- **4 distant problems**, $p_1 = 2, p_2 = 1$. Let us assume that u_1 and u_2 cause distant

problems on the same 2-path $u_3 \sim u_4$. By planarity, the triangle of u_3 can only reach the path $u_1 \sim u_2$, and so does the triangle of u_4 . Therefore, there cannot be two distinct 1-paths. Hence, this case is impossible.

- **4 distant problems**, $p_1 = 0, p_2 = 2$. If we assume that the path $u_3 \sim u_4$ touches the triangles of u_1 and u_2 , then necessarily the path $u_1 \sim u_2$ touches the triangles of u_3 and u_4 . This is again configuration D_3 .

This concludes the proof. □

4.6 Semi-distant configurations

We can now focus on the cases where the subdivision has up to 2 distant problems. Let us define another type of problem that we have to deal with in order to finish the proof of Lemma 4.16 (p. 61).

Definition 4.22 (Close problem). *Let G be a planar graph with a 4-family U and let S be a \mathcal{K} -subdivision rooted on U . A special vertex $u \in U$ causes a close problem if it is unsettled w.r.t. S and shares at least one of its remaining neighbors with at least one other unsettled special vertex.*

Note that by definition there are either zero or at least two special vertices causing a close problem; there cannot be a single special vertex causing a close problem on its own. Also, note that by definition, an unsettled special vertex that does not cause a distant nor a close problem forms a \mathcal{C}_N pattern that is disjoint from S and that touches only \mathcal{C}_V patterns, hence its reduction rule can be applied safely.

Let us first deal with subdivisions that have at most 2 distant problems and no close problem, with the following *semi-distant configurations* and their associated subdivision composite rules. We will then deal with subdivisions with close problems in Subsection 4.7.

List of the semi-distant configurations:

The semi-distant configurations are the configurations $J_1, J_2, J_3, J_4, J_5, J_6$ listed below.

Each configuration describes a 4-family U and a strong \mathcal{K} -subdivision S , such that at most 2 special vertices of U cause a distant problem on S , and none cause close problems. For each configuration, we describe a new semi-subdivision S' . The routing operation is not applied to S' unless stated otherwise.

We provide for each configuration a subdivision composite rule. We justify for each rule that the mapping is compatible w.r.t. S' .




Figure 28: J_1 in a case where u_1 and u_3 cause distant problems

Configuration J_1

Properties:

- The graph has a strong K_4 -subdivision S rooted on u_1, u_2, u_3, u_4
- At most 2 special vertices cause distant problems
- If there are two distant problems, they are not on the same path of S
- The special vertices that do not cause distant problems are settled

We consider a 2-coloring of S given by Claim 4.18 (p. 62) to inactivate the two potential distant problems.

The patterns used are \mathcal{C}_N for all special vertices.




Figure 29: J_2 in a case where the length of $u_1 \sim u_2$ is at least 2

Configuration J_2

Properties:

- The graph has a strong K_4 -subdivision S rooted on u_1, u_2, u_3, u_4
- u_1 and u_2 cause distant problems on the path $u_3 \sim u_4$: the path $u_3 \sim u_4 = (u_3, P_1, v_1, P_2, v_2, P_3, u_4)$, with v_1, v_2 being remaining neighbors of u_1, u_2 respectively, and $l(P_1), l(P_2), l(P_3) \geq 1$; we denote v'_1, v'_2 the other remaining neighbor of u_1, u_2 respectively
- u_3, u_4 are settled
- Either $l(u_1 \sim u_2) \geq 2$ or neither u_3 nor u_4 has v'_1, v'_2 as remaining neighbors
- If $u_1 \sim u_2$ has length 2 (u_1, w, u_2), then w has at most 1 neighbor among u_3, u_4 , or at least one of v'_1, v'_2 does not have a neighbor in $\{u_3, u_4\}$

We transform the K_4 -subdivision S into a C_{4+} -subdivision S' , by removing the paths $u_1 \sim u_2$ and $u_3 \sim u_4$ from S , and adding the paths (u_1, v_1, P_1, u_3) and (u_2, v_2, P_3, u_4) .

The special vertices u_1, u_2 are thus turned into C_V patterns, unless the path $u_1 \sim u_2$ from S has length 1, in which case u_1, u_2 form a C_U pattern. By the fourth condition of this configuration, neither u_3 nor u_4 has v'_1, v'_2 as remaining neighbors, thus they remain settled (the case where one has v'_1, v'_2 as remaining neighbors is treated as J_3).

Instead of C_V patterns, u_1 or u_2 may form C_{T2NA} patterns with u_3 or u_4 . If there are two such patterns, for instance $C_{T2NA}(u_1, u_3)$ and $C_{T2NA}(u_2, u_4)$, they may only intersect if they have a common vertex in the path $u_1 \sim u_2$ of S . By the last condition of the configuration, this is not the case (this case is treated as J_4).

The patterns used are $C_V(u_1), C_V(u_2)$, or $C_{T2NA}(u_i, u_j)$ for $i \in \{1, 2\}$ and $j \in \{3, 4\}$, or $C_U(u_1, u_2), C_N(u_3), C_N(u_4)$.




Figure 30: Reduction of configuration J_3

Configuration J_3

Properties:

- The graph has a strong K_4 -subdivision S rooted on u_1, u_2, u_3, u_4
- u_1 and u_2 cause distant problems on the path $u_3 \sim u_4$: the path $u_3 \sim u_4 = (u_3, P_1, v_1, P_2, v_2, P_3, u_4)$, with v_1, v_2 being remaining neighbors of u_1, u_2 respectively, and $l(P_1), l(P_2), l(P_3) \geq 1$; we denote v'_1, v'_2 the other remaining neighbor of u_1, u_2 respectively
- $l(u_1 \sim u_2) = 1$
- u_4 has v'_1, v'_2 as remaining neighbors
- u_3 is settled

Remark: u_4 is initially settled, but the rule that follows changes its \mathcal{C}_V nature into a \mathcal{C}_N one.

We transform the K_4 -subdivision S into a \mathcal{C}_{4+} -subdivision S' , by removing the paths $u_1 \sim u_2$ and $u_3 \sim u_4$ from S , and adding the paths (u_1, v_1, P_1, u_3) and (u_2, v_2, v'_2, u_4) .

The special vertices u_1, u_2 are thus turned into a \mathcal{C}_U pattern, while u_4 is turned into a \mathcal{C}_N . If v_2 is adjacent to u_4 , it becomes one of its remaining neighbors in S' , and in this case u_4 causes a distant problem. We inactivate this problem by maybe swapping the colors of the paths $u_2 \sim u_4$ and (u_2, v_2, v'_2, u_4) in a 2-coloring of S' .

The patterns used are $\mathcal{C}_U(u_1, u_2), \mathcal{C}_N(u_3), \mathcal{C}_N(u_4)$.




Figure 31: Reduction of configuration J_4 . Example of a 2-coloring of S'

Configuration J_4

Properties:

- The graph has a strong K_4 -subdivision S rooted on u_1, u_2, u_3, u_4
- u_1 has two adjacent remaining neighbors v_1, v_1' , with $v_1 \in u_3 \sim u_4$
- u_2 has two adjacent remaining neighbors v_2, v_2' with $v_2 \in u_3 \sim u_4$
- v_1, v_2 may be equal, or come in any order on $u_3 \sim u_4$
- $u_1 \sim u_2$ has length 2: call the third vertex w
- u_3 has v_2', w as remaining neighbors
- u_4 has v_1', w as remaining neighbors

We transform the K_4 -subdivision S into another K_4 -subdivision S' by replacing the path $u_3 \sim u_4$ by the path $(u_3, v_2', v_2, \dots, u_4)$. The vertices u_3, u_4 are turned into \mathcal{C}_V patterns and u_2 into a \mathcal{C}'_V pattern. Depending on the order of v_1, v_2 on the path $u_3 \sim u_4$ of S , u_1 forms a \mathcal{C}_N that may cause a distant problem on the new path $(u_3, v_2', v_2, \dots, u_4)$. We consider a coloring of S' given by Claim 4.18 (p. 62) to inactivate it.

The patterns used are $\mathcal{C}_N(u_1), \mathcal{C}'_V(u_2), \mathcal{C}_V(u_3), \mathcal{C}_V(u_4)$.




Figure 32: J_5 when u_1 and u_4 cause distant problems

Configuration J_5

Properties:

- The graph has a strong C_{4+}^* -subdivision S rooted on u_1, u_2, u_3, u_4 , such that u_1, u_2 are 1-linked and u_1, u_3 are 2-linked
- There are at most 2 distant problems: if there is at least one, we may assume w.l.o.g. that u_1 causes a distant problem on a (u_2, u_4) -path P_{24}
- u_2 is settled
- u_3 is settled or causes a distant problem on the (u_2, u_4) -path P'_{24} of S different from P_{24}

We consider a 2-coloring of S that inactivates the distant problems: $\{red = (u_3 \rightarrow u_1 \rightarrow u_2 \rightarrow u_4), blue = (u_1 \rightarrow u_3 \rightarrow u_4 \rightarrow u_2)\}$ in such a way that P_{24} receives the color red. The distant problem of u_1 is thus inactivated. Since the colors ending on u_1 and u_3 are different, and since the colors of P_{24} and P'_{24} are different, the potential distant problem of u_3 is inactivated. If u_4 causes a distant problem instead, we inactivate it by maybe swapping the colors of the two paths between u_1 and u_3 .

The patterns used are \mathcal{C}_N for all the special vertices, or possibly $\mathcal{C}_{T2NA}(u_1, u_2)$ and $\mathcal{C}_{T2NA}(u_3, u_4)$.




Figure 33: Reduction of configuration J_6

Configuration J_6

Properties:

- The graph has a strong C_{4+}^* -subdivision S rooted on u_1, u_2, u_3, u_4 , such that u_1, u_2 are 1-linked and u_1, u_3 are 2-linked
- u_1 and u_3 cause distant problems on a path P_{24} between u_2 and u_4
- u_2 and u_4 are settled and their remaining neighbors are disjoint from S

Assume w.l.o.g. that the path $P_{24} = (u_2, Q_1, v_1, Q_2, v_3, Q_3, u_4)$, where v_1, v_3 are remaining neighbors of u_1, u_3 respectively, and $l(Q_1), l(Q_2), l(Q_3) \geq 1$. Each of u_1, u_3 has another remaining neighbor v'_1, v'_3 respectively, adjacent to v_1, v_3 respectively. The vertices v'_1, v'_3 belong to a region of the graph delimited by the four paths $P_{24}, u_1 \sim u_2, u_3 \sim u_4$ and a path P_{13} of S between u_1 and u_3 . Let P'_{13} be the other path of S between u_1 and u_3 . We transform the C_{4+} -subdivision S into another C_{4+} -subdivision S' , by removing the paths P'_{13} and P_{24} , and adding the paths (u_1, v_1, Q_1, u_2) and (u_3, v_3, Q_3, u_4) .

The remaining neighbors of u_1 (resp. u_3) w.r.t. S' are non-adjacent, and since the remaining neighbors of u_2 (resp. u_4) are disjoint from S , (u_1, u_2) (resp. (u_3, u_4)) cannot form a \mathcal{C}_{T2NA} pattern.

The special vertices u_1, u_3 are thus turned into \mathcal{C}_V patterns, unless the path P'_{13} has length 1, in which case they form a \mathcal{C}_U pattern. In the latter case, by property “0-linked”, none of u_2, u_4 can have both v'_1, v'_3 as remaining neighbors, and by property “2-linked”, the remaining neighbors of u_2, u_4 are disjoint, so u_2 and u_4 remain settled w.r.t. S' .

The patterns used are $\mathcal{C}_V(u_1), \mathcal{C}_N(u_2), \mathcal{C}_V(u_3), \mathcal{C}_N(u_4)$, or $\mathcal{C}_U(u_1, u_3)$.

The following lemma shows that we can treat any subdivision that has at most 2 distant problems and no close problem with one of the semi-distant configurations.

Lemma 4.23 (Semi-distant lemma). *Let G be a planar graph with a (C_{II}) configuration w.r.t. a 4-family U , with a strong \mathcal{K} -subdivision S rooted on U . If G has at most 2 distant problems and no close problem w.r.t. S , then G contains a configuration among $\{J_1, J_2, J_3, J_4, J_5, J_6\}$.*

Proof. Let us consider the case where S is a K_4 -subdivision. If it has at most one distant problem, or two distant problems on different paths of S , then this is configuration J_1 . If $u_1, u_2 \in U$ both cause distant problems on the same path $u_3 \sim u_4$ of S , then we distinguish between 3 cases. Let v_1, v_2 be the remaining neighbors of u_1, u_2 respectively that are on the path $u_3 \sim u_4$, and let v'_1, v'_2 be their other remaining neighbors. If $l(u_1 \sim u_2) = 1$ and u_3 or u_4 has both v'_1, v'_2 as remaining neighbors, then this is configuration J_3 . If $l(u_1 \sim u_2) = 2$, with w as the middle vertex, w is adjacent to u_3 and u_4 , and v'_1, v'_2 are remaining neighbors of u_3 or u_4 , then this is configuration J_4 . Otherwise, this is configuration J_2 .

Now let us consider the case where S is a C_{4+} -subdivision. By property “1-linked” and property A of S , the distant problems occur on parallel paths of S . Thus, if there is at most one distant problem, this is configuration J_5 . By Claim 4.19 (p. 62), two distant problems cannot be caused by 1-linked special vertices. Therefore, if there are two distant problems on different paths of S , then this is configuration J_5 . If there are two distant problems caused by (w.l.o.g.) u_1, u_3 on the same parallel (u_2, u_4) -path of S , by Claim 4.19 (p. 62) the remaining neighbors of u_2, u_4 are disjoint from S , and this is configuration J_6 . This concludes the proof. \square




Figure 34: Semi-distant lemma trees of cases

4.7 Close configurations

For simplicity, we define some macros that encapsulate several patterns and configurations from the redirection procedure.

- \mathcal{C}_{D1} : In this configuration, $u_1, u_2 \in U$ are linked by a path $u_1 \sim u_2$ in S . The vertices u_1, u_2 have a common remaining neighbor v and have another remaining neighbor v_1, v_2 respectively, both adjacent to v . The vertices v and v_1 are disjoint from S , and if v_2 is in S , it belongs to a path of S incident with u_1 and not u_2 .

First let us assume that v_2 is not in S . If the path $u_1 \sim u_2$ has length 1, then it is a \mathcal{C}_{Da} pattern if v_1, v_2 are not adjacent, or a \mathcal{C}_{Db} pattern if they are. If $u_1 \sim u_2$ has length at least two, then this is forbidden by the redirection procedure, as this is a \mathcal{C}_{X1} or \mathcal{C}_{X2} configuration depending on whether v_1 is adjacent to the neighbor w_1 of u_1 on $u_1 \sim u_2$. Now if there is a path $u_1 \sim u'$ in S that touches v_2 , with $u' \neq u_2$, then it is a \mathcal{C}_{X3} configuration, forbidden by property C.

- \mathcal{C}_{D2} : The vertices u_1, u_2 are linked by a path $u_1 \sim u_2$ in S . The vertices u_1, u_2 have two remaining neighbors v, v' in common. No path of S touches v, v' .

If v, v' are not adjacent, then this is (\mathcal{C}_{T2NAa}) or (\mathcal{C}_{T2NAb}) depending on the parity of v, v' . Now assume v, v' are adjacent, and let l be the length of $u_1 \sim u_2$. If $l = 1$, then $\{u_1, u_2, v, v'\}$ form an induced K_4 , contradicting the fact that G has a (C_{II}) configuration by Claim 4.2 (p. 36). Then $l \geq 2$ and this is configuration \mathcal{C}_{X4} from the redirection procedure, hence forbidden by property C.

To summarize, apart from forbidden configurations removed by the redirection procedure, a \mathcal{C}_{D1} macro is a \mathcal{C}_{Da} or \mathcal{C}_{Db} pattern. A \mathcal{C}_{D2} is a \mathcal{C}_{T2NA} pattern and in this case u_1, u_2 are therefore settled if no unsettled special vertex shares remaining neighbors with them. See Figure 35.




Figure 35: Possible patterns for each macro

Let us now introduce the remaining configurations, with which we treat all the cases of \mathcal{K} -subdivisions with close problems.

List of the close configurations:

The close configurations are the configurations R_1, \dots, R_9 listed below.

Each configuration describes a strong \mathcal{K} -subdivision S rooted on a 4-family $U = \{u_1, u_2, u_3, u_4\}$, such that at most two special vertices cause distant problems, and some special vertices cause close problems. We describe for each a subdivision composite rule made up of a semi-subdivision S' (if not specified, $S' = S$) and a compatible mapping w.r.t. S' .

Remark: When two remaining neighbors of a special vertex form a \mathcal{C}_N pattern or a \mathcal{C}'_V pattern, we denote it by \mathcal{C}_N for simplicity. This does not change the case analysis.




Figure 36: Reduction of configuration R_1 . u_3, u_4 may cause distant problems

Configuration R_1

Properties:

- The graph has a strong K_4 -subdivision S rooted on u_1, u_2, u_3, u_4 , with 2 special vertices involved in a close problem: u_1, u_2 share a remaining neighbor v
- $v \notin S$
- u_1, u_2 each have another remaining neighbor v_1, v_2 respectively, and $v_1 \neq v_2$
- **Remark:** each of u_3, u_4 is either settled or causes a distant problem

We consider a 2-coloring of S given by Claim 4.18 (p. 62) to inactivate the two potential distant problems on u_3 and u_4 . If one of v_1, v_2 is not adjacent to v , then its associated special vertex forms a \mathcal{C}'_V pattern and is thus settled: a contradiction, as u_1 and u_2 are the ones causing a close problem. Hence v_1, v_2 are both adjacent to v . The vertices u_1 and u_2 form a \mathcal{C}_{D1} configuration, hence a \mathcal{C}_{Da} or \mathcal{C}_{Db} pattern. Note that v_2 cannot belong to the path $u_1 \sim u_3$ and v_1 cannot belong to $u_2 \sim u_3$, as this would form a \mathcal{C}_{X3} configuration, forbidden by the redirection procedure.

The patterns used are \mathcal{C}_{Da} or $\mathcal{C}_{Db}(u_1, u_2), \mathcal{C}_N(u_3), \mathcal{C}_N(u_4)$.




Figure 37: Reduction of configuration R_2

Configuration R_2

Properties:

- The graph has a strong K_4 -subdivision S rooted on u_1, u_2, u_3, u_4 , without distant problems and such that 2 special vertices are involved in a close problem: u_1, u_2 share two remaining neighbors v, v'
- $v \notin S$ and $v' \in S$
- **Remark:** each of u_3, u_4 is either settled or causes a distant problem

By planarity and property A, there is at most one distant problem, caused by u_3 or u_4 on the path $u_1 \sim u_2$. If it is the case, we assume w.l.o.g. that this is u_3 .

Let us color S with this 2-coloring: $\{red = (u_2 \rightarrow u_1 \rightarrow u_3 \rightarrow u_4), blue = (u_1 \rightarrow u_4 \rightarrow u_2 \rightarrow u_3)\}$. The colors ending on u_1, u_2 are different, so u_1, u_2 form a \mathcal{C}_{T2NA} pattern that crosses the red path $u_3 \sim u_4$. This is authorized by the definition of \mathcal{C}_{T2NA} . The potential distant problem of u_3 is inactive in this coloring of S .

The patterns used are $\mathcal{C}_{T2NA}(u_1, u_2), \mathcal{C}_N(u_3)$ (or \mathcal{C}'_V), $\mathcal{C}_N(u_4)$.

Configuration R_3

Properties:

- The graph has a strong K_4 -subdivision S
- u_1, u_2 each have two remaining neighbors v_1, v'_1 and v_2, v'_2 respectively
- v_1, v_2 belong to $u_3 \sim u_4$; by convention $u_3 \sim u_4 = (u_3, P_1, v_1, P_2, v_2, P_3, u_4)$, with $l(P_1), l(P_3) \geq 1$ and $l(P_2) \geq 0$ (so v_1 may equal v_2)
- If $v_1 \neq v_2, v'_1, v'_2$ are disjoint from S
- **Remark:** if $v_1 = v_2$, then v'_1, v'_2 may belong to $u_3 \sim u_4$ and v'_1 may equal v'_2
- If v'_1 (resp. v'_2) does not belong to S , then it is adjacent to v_1 (resp. v_2)
- u_3, u_4 each have two remaining neighbors v_3, v'_3 and v_4, v'_4 respectively
- The path $u_1 \sim u_2$ does not have length 1
- If $u_1 \sim u_2$ has length 2, let w be its middle vertex. Then w has at most 1 neighbor among u_3, u_4 , or at least one of v'_1, v'_2 does not have a neighbor in $\{u_3, u_4\}$




Figure 38: R_3 when $v_1 \neq v_2, v'_1 \neq v'_2$

We transform the K_4 -subdivision S into a C_{4+} -semi-subdivision S' by removing the paths $u_1 \sim u_2$ and $u_3 \sim u_4$, and adding the paths (u_1, v_1, P_1, u_3) and (u_2, v_2, P_3, u_4) if $v_1 \neq v_2$ or if $v_1 = v_2$ and $v'_1 \neq v'_2$. Otherwise, we have $v_1 = v_2$ and $v'_1 = v'_2$, and we assume $u_3 \sim u_4 = (u_3, Q_1, v'_1, Q_2, v_1, Q_3, u_4)$. In this case, we add the paths (u_1, v'_1, Q_1, u_3) and (u_2, v_1, Q_3, u_4) to S' instead. This semi-subdivision has two paths that intersect if $v_1 = v_2$ and $v'_1 \neq v'_2$ ($l(P_2) = 0$), but it is 2-colorable with the coloring $\{red = (u_2 \rightarrow u_4 \rightarrow u_1 \rightarrow v_1 \rightarrow u_3), blue = (u_1 \rightarrow u_3 \rightarrow u_2 \rightarrow v_2 \rightarrow u_4)\}$.

Since the path $u_1 \sim u_2$ of S does not have length 1, the special vertices u_1, u_2 form \mathcal{C}_V patterns in S' .

The special vertices u_3, u_4 may form \mathcal{C}_{T_2NA} patterns in S' with each of u_1, u_2 . These patterns may only touch each other on the path $u_1 \sim u_2$ from S , but the last condition of the configuration ensures that this is not the case (this case is treated as configuration J_4).

Otherwise, the vertices u_3, u_4 form \mathcal{C}_N patterns if they do not have common remaining neighbors. These \mathcal{C}_N patterns are compatible with the \mathcal{C}_V patterns of u_1, u_2 .

Remark: If $v'_1 \neq v'_2$ and u_4 has both as its remaining neighbors, then u_4 forms a \mathcal{C}_V pattern which touches the \mathcal{C}_V patterns of u_1 and u_2 . The precise case where u_4 has v_1, v_2 as remaining neighbors and $u_1 \sim u_2$ has length 1 in S is treated as configuration J_3 .

The special vertices u_3, u_4 may also form a \mathcal{C}_{T_1} , \mathcal{C}_{T_2A} or \mathcal{C}_{T_2NA} pattern, since the colors ending on each vertex are different in any 2-coloring of S' .

The patterns used are $\mathcal{C}_V(u_1), \mathcal{C}_V(u_2)$ and $\mathcal{C}_{T_1}, \mathcal{C}_{T_2A}$ or $\mathcal{C}_{T_2NA}(u_3, u_4)$, or \mathcal{C}_N for u_3, u_4 .




Figure 39: R_4 when $l(u_1 \sim u_2) \geq 2$ in S

Configuration R_4

Properties:

- The graph has a strong K_4 -subdivision S rooted on u_1, u_2, u_3, u_4 , with 3 special vertices involved in a close problem: u_1, u_2, u_3 .
- The vertices u_1, u_3 share a remaining neighbor $v_{13} \notin S$
- u_1, u_2, u_3 share a remaining neighbor v_{123} (different from v_{13})
- u_2 has another remaining neighbor v_2 adjacent to v_{123}
- **Remark:** u_4 is either settled or causes a distant problem

By planarity and property A, v_{13} and v_{123} do not belong to S . The special vertices u_1, u_3 thus form a \mathcal{C}_{D2} configuration, which is therefore a \mathcal{C}_{T2NA} pattern: the vertices v_{13} and v_{123} are non-adjacent. However, this pattern is not compatible with a \mathcal{C}_N pattern applied to u_2 .

The vertex u_4 may cause a distant problem or cause a \mathcal{C}'_V pattern on the path $u_1 \sim u_3$ or (w.l.o.g.) $u_2 \sim u_3$. We replace the K_4 -subdivision S with another K_4 -subdivision S' by replacing the path $u_1 \sim u_2$ with the path (u_1, v_{123}, u_2) . If u_4 causes a distant problem on $u_1 \sim u_3$ or $u_2 \sim u_3$, we use Claim 4.18 (p. 62) and consider a 2-coloring of S' that inactivates it.

By planarity and property A, v_2 does not belong to S . The vertices u_1 and u_2 form either a \mathcal{C}_V and a \mathcal{C}_N in S' , or a \mathcal{C}_U pattern depending on the length of the original $u_1 \sim u_2$ in S .

The pattern used are $\mathcal{C}_V(u_1), \mathcal{C}_N(u_2)$, or $\mathcal{C}_U(u_1, u_2), \mathcal{C}_V(u_3), \mathcal{C}_N(u_4)$ (or \mathcal{C}'_V).




Figure 40: Reduction of configuration R_5 . The special vertex u_4 may cause a distant problem on $u_1 \sim u_2$ or $u_2 \sim u_3$

Configuration R_5

Properties:

- The graph has a strong K_4 -subdivision S rooted on u_1, u_2, u_3, u_4 , with 3 vertices involved in a close problem: u_1, u_2, u_3 .
- u_1, u_3 share a remaining neighbor v_{13} (which is not a neighbor of u_2)
- u_1, u_2 share a remaining neighbor v_{12} (which is not a neighbor of u_3)
- u_2, u_3 each have another remaining neighbor v_2, v_3 respectively, and $v_2 \neq v_3$
- v_3, v_{12} are not adjacent
- None of v_{13}, v_{12}, v_2, v_3 belong to S
- The graph contains the edges $v_{13}v_3, v_{13}v_{12}, v_{12}v_2$
- **Remark:** u_4 is either settled or causes a distant problem

At least one edge among v_2v_{13}, v_3v_{12} does not exist by planarity (otherwise $\{(u_1, v_2, v_3), (u_2 = u_3, v_{12}, v_{13})\}$ form a $K_{3,3}$ minor by contracting the path $u_2 \sim u_3$). Assume w.l.o.g. that the edge v_3v_{12} is absent from the graph.

The vertices u_1, u_3 form a \mathcal{C}_{D1} configuration, and since v_3v_{12} does not exist, they form a \mathcal{C}_{Db} pattern. By planarity and property A, v_2 does not belong to S . Hence, u_2 can be treated as a \mathcal{C}_N pattern that is compatible with the \mathcal{C}_{Db} pattern, since it touches only v_{12} . We use Claim 4.18 (p. 62) and consider a 2-coloring of G that inactivates the potential distant problem caused by u_4 .

The patterns used are $\mathcal{C}_{Db}(u_1, u_3), \mathcal{C}_N(u_2), \mathcal{C}_N(u_4)$.




Figure 41: R_6 when u_1, u_2 and u_3, u_4 form \mathcal{C}_{D1} configurations

Configuration R_6

Properties:

- The graph has a strong K_4 -subdivision S rooted on u_1, u_2, u_3, u_4 , with all 4 special vertices involved in close problems
- u_1, u_2 share a remaining neighbor $v_{12} \notin S$
- u_3, u_4 share a remaining neighbor $v_{34} \notin S$
- u_1, u_2, u_3, u_4 each have another remaining neighbor v_1, v_2, v_3, v_4 respectively
- v_1, v_2 are disjoint from v_3, v_4
- **Remark:** it may be that $v_1 = v_2$ or $v_3 = v_4$
- v_1, v_2, v_3, v_4 are disjoint from S

This case is straightforward: each of (u_1, u_2) and (u_3, u_4) forms a \mathcal{C}_{D1} or \mathcal{C}_{D2} configuration, which can be a \mathcal{C}_{Da} , \mathcal{C}_{Db} or \mathcal{C}_{T2NA} pattern. By definition, these two patterns are disjoint.




Figure 42: R_7 when $l(u_2 \sim u_4) \geq 2$ in S

Configuration R_7

Properties:

- The graph has a strong K_4 -subdivision S rooted on u_1, u_2, u_3, u_4 , with all 4 special vertices involved in one close problem
- u_1, u_2, u_3 share a remaining neighbor $v_{123} \notin S$
- u_1, u_2, u_4 share a remaining neighbor $v_{124} \notin S$
- u_3, u_4 do not share a remaining neighbor
- u_3, u_4 each have another remaining neighbor v_3, v_4 respectively, adjacent to v_{123}, v_{124} respectively

We first claim that v_4 cannot be adjacent to v_{123} . If it is the case, then $\{(u_1, u_2, v_4), (u_4, v_{123}, v_{124})\}$ form a $K_{3,3}$ -minor, a contradiction with the planarity of G .

We replace the K_4 -subdivision S with another K_4 -subdivision S' by removing the path $u_2 \sim u_4$ and adding the path (u_2, v_{124}, u_4) .

The special vertex u_1 forms a \mathcal{C}_V pattern and u_3 a \mathcal{C}_N pattern disjoint from S' by property A and planarity. u_2 forms a \mathcal{C}_V and u_4 a \mathcal{C}_N pattern, unless the length of $u_2 \sim u_4$ in S is 1. In this case u_2, u_4 form a \mathcal{C}_U pattern, since v_4, v_{123} are non-adjacent. The \mathcal{C}_N patterns are disjoint and may only touch \mathcal{C}_V patterns, hence the mapping is compatible.

The patterns used are $\mathcal{C}_V(u_1), \mathcal{C}_N(u_3)$ and either $\mathcal{C}_V(u_2)$ and $\mathcal{C}_N(u_4)$ or $\mathcal{C}_U(u_2, u_4)$.




Figure 43: R_8 when u_3 causes a distant problem and u_1, u_2 form a \mathcal{C}_{D1} configuration. Example of a 2-coloring of S

Configuration R_8

Properties:

- The graph has a strong C_{4+}^* -subdivision S rooted on u_1, u_2, u_3, u_4 , such that u_1, u_2 are 1-linked and u_1, u_3 are 2-linked
- There is at most one distant problem, caused by u_3 on a (u_2, u_4) -path of S if there is one
- u_1, u_2 share a remaining neighbor
- There is no remaining neighbor in common between one of u_1, u_2 and one of u_3, u_4
- **Remark:** if there is no distant problem, u_3 and u_4 may share a remaining neighbor

If u_3 causes a distant problem, it is necessarily on a parallel (u_2, u_4) -path of S by property “1-linked” of S and property A, and then we may swap the colors of the two (u_2, u_4) -paths in a 2-coloring of S to inactivate this distant problem.

By the last property of this configuration, u_1, u_2 form a \mathcal{C}_{D1} or \mathcal{C}_{D2} pattern, and u_3, u_4 as well if u_3 does not cause a distant problem.

The patterns used are thus $\mathcal{C}_{Da}, \mathcal{C}_{Db}$ or \mathcal{C}_{T2NA} for (u_1, u_2) and maybe for (u_3, u_4) , or $\mathcal{C}_N(u_3), \mathcal{C}_N(u_4)$.

Configuration R_9

Properties:

- The graph has a strong C_{4+}^* -subdivision S rooted on u_1, u_2, u_3, u_4 , such that u_1, u_2 are 1-linked and u_1, u_3 are 2-linked
- u_2, u_4 share exactly remaining neighbor v_{24} , and it belongs to a parallel (u_1, u_3) -path P_{13} of S
- The other remaining neighbors v_2, v_4 of u_2, u_4 respectively are both adjacent to v_{24}
- The remaining neighbors of u_1, u_3 are disjoint from S
- **Remark:** there is no distant problem

Let us write $P_{13} = (u_1, Q_1, v_{24}, Q_2, u_3)$. We transform the C_{4+} -subdivision S into another C_{4+} -(semi-)subdivision S' by removing the paths P_{13} and any of the two (u_2, u_4) -paths of S , P_{24} , and adding (u_1, Q_1, v_{24}, u_2) and (u_3, Q_2, v_{24}, u_4) . This semi-subdivision has a contact between the two new paths on v_{24} , so in a 2-coloring of S' we may swap the colors of the two




Figure 44: Reduction of configuration R_9

(u_1, u_2) -paths to have different colors on the new paths.

The special vertices u_2, u_4 form \mathcal{C}_V patterns in S' by planarity, and since the remaining neighbors of u_1, u_3 are disjoint from S , none of (u_1, u_2) or (u_3, u_4) form \mathcal{C}_{T2NA} patterns.

The patterns used are $\mathcal{C}_N(u_1), \mathcal{C}_V(u_2), \mathcal{C}_N(u_3), \mathcal{C}_V(u_4)$, or maybe $\mathcal{C}_{T2NA}(u_1, u_2)$ or $\mathcal{C}_{T2NA}(u_3, u_4)$.

Before entering the proof of the final lemma of this section, let us show a useful claim.

Claim 4.24. *Let G be a planar graph with a (C_{II}) configuration w.r.t. a 4-family U , with a \mathcal{K} -subdivision S rooted on U .*

If $u \in U$ has no remaining neighbors in common with other special vertices, then either u causes a distant problem in S or u is lone-settled.

Proof. Let us assume that u does not cause a distant problem in S . Since it does not share remaining neighbors with other special vertices, it cannot form a \mathcal{C}_{T2NA} pattern w.r.t. S . Hence, since no pair of special vertices forms a \mathcal{C}_U pattern by property A, if the remaining neighbors of u are non-adjacent, u forms a \mathcal{C}_V pattern and is lone-settled.

If its remaining neighbors are adjacent, since u does not share remaining neighbors and does not cause a distant problem, either none or both of its remaining neighbors belong to S . Then either its remaining neighbors are disjoint from S and u forms a \mathcal{C}_N pattern, or by property B and planarity u has both remaining neighbors in S and forms a \mathcal{C}'_V pattern. In both cases, it is lone-settled. \square

We can now show how all the remaining cases of \mathcal{K} -subdivisions with close problems can be taken care of with the previous close configurations.

Lemma 4.25 (Close lemma). *Let G be a planar graph with a (C_{II}) configuration w.r.t. a 4-family U , with a \mathcal{K} -subdivision S rooted on U , with at most 2 distant problems and some close problems w.r.t. S . Then G has a configuration among $\{R_1, R_2, R_3, R_4, R_5, R_6, R_7, R_8, R_9, J_4\}$.*

Proof. We start by solving **the \mathcal{C}_{4+} cases**: see Figure 45 for the tree of cases.

If S is a \mathcal{C}_{4+}^* -subdivision, let $U = \{u_1, u_2, u_3, u_4\}$ be such that u_1, u_2 are 1-linked, and u_1, u_3 are 2-linked. We can assume that G does not have a K_4 -subdivision rooted on U .

If S has distant problems, they are on parallel paths by property “1-linked” and property A. By Claim 4.19 (p. 62), S does not have two distant problems caused by 1-linked special vertices. If S has two distant problems caused by two 0-linked special vertices, then by property “0-linked”, the other two 0-linked special vertices do not share remaining neighbors, and are thus

lone-settled by definition of close problem and Claim 4.24 (p. 86). So S does not have close problems, which is a contradiction. If S has two distant problem caused by 2-linked special vertices, then by Claim 4.19 (p. 62), the other two special vertices have no remaining neighbor on S . By property “2-linked”, this means that they do not share remaining neighbors, which means that they are (lone-)settled and there is no close problem, a contradiction.

So S has at most one distant problem, caused by u_3 if so. Let us first consider the case where remaining neighbors of $\{u_1, u_2\}$ are disjoint from the ones of $\{u_3, u_4\}$. If u_3 causes a distant problem, then by Claim 4.24 (p. 86) u_4 is lone-settled, and thus (u_1, u_2) must form a \mathcal{C}_{D1} or \mathcal{C}_{D2} configuration to be unsettled. If u_3 does not cause a distant problem, at least one pair among (u_1, u_2) and (u_3, u_4) forms a \mathcal{C}_{D1} or \mathcal{C}_{D2} configuration. In all cases, this is configuration R_8 .

Now, let us consider the case where $\{u_1, u_2\}$ share some remaining neighbors with $\{u_3, u_4\}$. By property “0-linked”, two 0-linked special vertices cannot share remaining neighbors. If u_2, u_4 share remaining neighbors, by property “2-linked” they share exactly one and it belongs to a parallel (u_1, u_3) -path of S . Therefore, by Claim 4.19 (p. 62), the remaining neighbors of u_1, u_3 are disjoint from S , thus disjoint, and this is configuration R_9 .




Figure 45: Close lemma: tree of C_{4+} cases

Let us now deal with **the K_4 cases**: see Figure 46 for the tree of cases.

We first examine the cases in which there are only two vertices involved in a close problem: we assume w.l.o.g. that these two vertices are u_1, u_2 , they share a remaining neighbor v , and u_3, u_4 are either settled or cause distant problems. One may form a \mathcal{C}_N pattern disjoint from S and touching only patterns from settled vertices, but we treat it as settled.

We first examine the case where $v \notin S$. If u_1, u_2 share only one remaining neighbor, this is configuration R_1 . So now assume u_1, u_2 share another remaining neighbor v' . If $v' \notin S$, then this is a \mathcal{C}_{D2} configuration where u_1, u_2 form a \mathcal{C}_{T2NA} pattern, as no other special vertex can cause a close problem by hypothesis, and by property A no path of S can touch v or v' . Thus u_1, u_2 are settled, a contradiction. Otherwise, $v' \in S$ and this is configuration R_2 .

Now let us assume that $v \in S$. By property A, v necessarily belongs to the path $u_3 \sim u_4$. We can immediately see that if u_1, u_2 share another remaining neighbor $v' \notin S$, then this is a case that has already been treated, by swapping v and v' . Now u_1, u_2 may have another common remaining neighbor $v' \in S$, necessarily in the path $u_3 \sim u_4$, or each of u_1, u_2 can have another remaining neighbor v'_1, v'_2 respectively, both adjacent to v to be unsettled. All these cases are treated as configuration R_3 , except in one particular case: there is a vertex w on $u_1 \sim u_2$ that is a common neighbor of u_1, u_2, u_3, u_4 , the special vertex u_4 is adjacent to v'_1 , while u_3 is adjacent to v'_2 , and in this case it is configuration J_4 . This concludes the cases with two special vertices involved in a close problem.

Let us now examine the cases with 3 vertices involved in a close problem, say u_1, u_2, u_3 . We can assume w.l.o.g. that u_1, u_3 share a remaining neighbor v_{13} . There is at most one distant problem, caused by u_4 if so; otherwise, u_4 is lone-settled by definition of distant problem. The

special vertices u_1, u_2, u_3 each have another remaining neighbor v_1, v_2, v_3 respectively, and u_2 has another remaining neighbor v'_2 (these vertices are not necessarily distinct).

We first examine the case where $v_{13} \notin S$.

If $v_1 = v_3$, we can denote this vertex by v'_{13} . If u_2 does not have v_{13} or v'_{13} as a remaining neighbor, it is lone-settled by Claim 4.24 (p. 86), a contradiction with the hypothesis on u_2 ; so one of v_{13}, v'_{13} is also a remaining neighbor of u_2 . By planarity, v_{13}, v'_{13} cannot both be remaining neighbors of u_2 . So let us say that v_{13} is not a remaining neighbor of u_2 , that the vertices u_1, u_2, u_3 have a common remaining neighbor v_{123} , and that $v_2 \neq v_{123}$. Since u_2 is unsettled, v_2, v_{123} are adjacent and this is configuration R_4 .

Now let us take a look at the case where $v_1 \neq v_3$. We distinguish between the case where v_{13} is a shared remaining neighbor of u_1, u_2, u_3 or not.

- v_{13} is also a remaining neighbor of u_2 , and we call it v_{123} . If w.l.o.g. $v_1 = v_2$, then this case is equivalent to the previous one, by swapping u_2 and u_3 ; hence we assume that v_1, v_2, v_3 are pairwise distinct. If one of v_1, v_2, v_3 is not adjacent to v_{123} , then its special vertex is lone-settled, a contradiction; so all three of v_1, v_2, v_3 are adjacent to v . There are three \mathcal{C}_{D1} configurations on $(u_1, u_2), (u_1, u_3), (u_2, u_3)$, since by planarity and property A v_1, v_2, v_3 are disjoint from S . Each of these configurations forms a \mathcal{C}_{Da} or \mathcal{C}_{Db} pattern by property C, so the three paths $u_1 \sim u_2, u_1 \sim u_3, u_2 \sim u_3$ have length 1. Then $\{u_1, u_2, u_3\}$ is a 3-cut in G that separates two neighbors of u_1 , a contradiction to the almost 4-connectivity of G w.r.t. the special vertices.
- v_{13} is not a remaining neighbor of u_2 . Since by Claim 4.24 (p. 86) u_2 has a remaining neighbor in common with u_1 or u_3 , we can assume w.l.o.g. that $v_1 = v_2$ and we call this vertex v_{12} .

First, assume $v'_2 = v_3$ and call it v_{23} . Since u_1, u_2, u_3 are unsettled, the graph contains the edges $v_{13}v_{23}, v_{23}v_{12}, v_{12}v_{13}$. Thus, the three \mathcal{C}_{D1} configurations on $(u_1, u_2), (u_1, u_3)$ and (u_2, u_3) are all \mathcal{C}_{Da} patterns by property C. Thus the three paths between u_1, u_2, u_3 all have length 1, which is again a contradiction to the almost 4-connectivity of G w.r.t. the special vertices.

Now assume $v'_2 \neq v_3$. Due to u_1, u_2, u_3 being unsettled, the graph contains the edges $v_{13}v_3, v_{13}v_{12}, v_{12}v'_2$. By planarity, there is at most one edge among $\{v_{13}v'_2, v_{12}v_3\}$, so we can assume w.l.o.g. that $v_{12}v_3$ is a non-edge and in this case we have configuration R_5 .

We can now examine the case where $v_{13} \in S$.

By property A, v_{13} necessarily belongs to $u_2 \sim u_4$, thus u_2 cannot have it as a remaining neighbor by property A. The vertex u_2 must be involved in the close problem, so it is adjacent to v_1, v_3 , or both.

First assume u_2 is adjacent to both v_1, v_3 . Since u_1, u_2, u_3 are unsettled, we need to have the edges $v_1v_3, v_1v_{13}, v_3v_{13}$, otherwise one of them is a settled \mathcal{C}_V pattern. However, in this case $\{v_{13}, v_1, v_3, u_2, (u_1 = u_3)\}$ form a K_5 -minor, by contracting the path $u_1 \sim u_3$ to a vertex, a contradiction with the planarity of G .

So now we assume w.l.o.g. that u_2 is adjacent to v_3 but not to v_1 : we say that $v_3 = v'_2$ and we call it v_{23} . Since u_2, u_3 are unsettled, we need the edges v_2v_{23} and $v_{13}v_{23}$. But then this is a \mathcal{C}_{X3} configuration, which contradicts property C of the subdivision. This concludes the cases of 3 special vertices involved in the close problem.

Let us finally examine the cases where all four special vertices are involved in one or two close problems. By definition, there are no distant problems in S .

For a special vertex to cause a close problem, it needs to share some of its remaining neighbors with another special vertex. Two cases may occur: either there are two independent close problems each involving two special vertices, or there is one close problem involving all four

special vertices. We start with the first case: u_1, u_2 are involved in a close problem, as well as u_3, u_4 , but the remaining neighbors of $\{u_1, u_2\}$ and those of $\{u_3, u_4\}$ are disjoint. We examine all combinations of cases:

- u_1, u_2 share **exactly one** remaining neighbor v_{12} ; u_3, u_4 share **exactly one** remaining neighbor v_{34} . For all special vertices to be unsettled, we assume that the other remaining neighbors v_1, v_2 of u_1, u_2 are adjacent to v_{12} , and the other remaining neighbors v_3, v_4 of u_3, u_4 are adjacent to v_{34} . If $v_{12}, v_{34} \notin S$, then none of v_1, v_2, v_3, v_4 can belong to S , otherwise by property A the graph contains a redirection configuration \mathcal{C}_{X3} , forbidden by property C. So this is configuration R_6 . By planarity, if one of v_{12}, v_{34} belongs to S , then the other one does too. If they both belong to S , this is configuration R_3 .
- u_1, u_2 share **two** remaining neighbors v_{12}, v'_{12} ; u_3, u_4 share **exactly one** remaining neighbor v_{34} . Since u_3, u_4 are unsettled, their other remaining neighbors v_3, v_4 are adjacent to v_{34} . If none of v_{12}, v'_{12} belongs to S , then neither do v_3, v_4, v_{34} by planarity. This is again configuration R_6 . Now if say v_{12} belongs to S , it belongs to $u_3 \sim u_4$ by property A, and then by planarity v_{34} belongs to the path $u_1 \sim u_2$. Thus, by planarity, v'_{12} must also belong to S . This is once more configuration R_3 .
- u_1, u_2 share **two** remaining neighbors v_{12}, v'_{12} ; u_3, u_4 share **two** remaining neighbors v_{34}, v'_{34} . Using the same argument, either none of $v_{12}, v'_{12}, v_{34}, v'_{34}$ belong to S , or they all do. In the former case this is again configuration R_6 , in the latter this is configuration R_3 .

This concludes the case with two independent close problems. Let us now assume that all four special vertices are involved in the same close problem. Let us decompose according to the case (P_1 or P_2) of the remaining neighbors of u_1, u_2 .

- u_1, u_2 share **two** remaining neighbors v_{12}, v'_{12} . We note that if at least one of v_{12}, v'_{12} belongs to S , then it is impossible by planarity and property A to have both u_3, u_4 involved. So $v_{12}, v'_{12} \notin S$. Assume w.l.o.g. that v_{12} is also a remaining neighbor of u_3 , and call it v_{123} . If u_3, u_4 share a remaining neighbor v_{34} (different from v_{123} by planarity), then by property A if $v_{34} \in S$ it can only belong to the path $u_1 \sim u_2$, but it is impossible in this case by planarity. Thus $v_{34} \notin S$, and by planarity v_{34}, v_{123} are not adjacent. But then u_3 is a settled \mathcal{C}_V pattern, a contradiction. So u_3, u_4 do not share a remaining neighbor. For u_4 to be involved, it must have (by planarity) v'_{12} as a remaining neighbor. This is configuration R_7 .
- u_1, u_2 share **exactly one** remaining neighbor v_{12} , and each have another remaining neighbor v_1, v_2 , both adjacent to v_{12} . We first assume that $v_{12} \notin S$. If u_3 has v_1 as a remaining neighbor, then by planarity v_1, v_2, v_{12} must belong to the region of the graph delimited by the paths $u_1 \sim u_2, u_2 \sim u_3, u_1 \sim u_3$, and none of v_1, v_2, v_{12} can belong to S (by property A and planarity). It is then impossible for u_4 to be involved with the close problem. So necessarily u_3 has v_{12} as a remaining neighbor, we call it v_{123} . By the same argument it is impossible for u_4 to be involved without making u_3 a \mathcal{C}_V pattern, thus settled, a contradiction. So finally assume that v_{12} belongs to S : by property A it belongs to $u_3 \sim u_4$. We can assume w.l.o.g. that u_3 has v_1 as a remaining neighbor (it cannot have v_{12} by property A). By an argument used above, u_3 cannot share a remaining neighbor with u_4 without being a \mathcal{C}_V pattern, so u_4 has v_2 as a remaining neighbor. This is again configuration R_3 .

This concludes the proof. □




Figure 46: Close lemma: tree of K_4 cases

Let us conclude this section by proving Lemma 4.16 (p. 61) and using it to prove Lemma 4.1 (p. 35).

Proof of Lemma 4.16 (p. 61). Let G be a planar graph with a (C_{II}) configuration w.r.t. a 4-family U , that admits a strong \mathcal{K} -subdivision rooted on U . We prove that G contains a subdivision composite configuration made up of a semi-subdivision S rooted on U and a compatible mapping w.r.t. S .

By Lemma 4.21 (*Distant lemma*, p. 67), if G has at least 3 distant problems w.r.t. its strong subdivision S , then it contains a configuration among $\{D_1, D_2, D_3, D_4\}$. Each of these configurations is defined along with a semi-subdivision and a compatible mapping.

So let us assume G has at most 2 distant problems and no close problem w.r.t. S . By Lemma 4.23 (*Semi-distant lemma*, p. 76), G admits a semi-distant configurations among $\{J_1, J_2, J_3, J_4, J_5, J_6\}$, and we provided a semi-subdivision and a compatible mapping for each of these configurations. If G has at most 2 distant problems and some close problems w.r.t. S , then by Lemma 4.25 (*Close lemma*, p. 86), it contains a configuration among $\{R_1, R_2, R_3, R_4, R_5, R_6, R_7, R_8, R_9, J_4\}$, again associated with a semi-subdivision and a compatible mapping for each. \square

Proof of Lemma 4.1 (p. 35). Let G be a minimum counterexample. By Lemma 3.2 (p. 6), it does not contain a configuration (C_I) .

Assume that G contains a (C_{II}) configuration w.r.t. a 4-family U . By Claim 4.14 (p. 58), G admits a strong \mathcal{K} -subdivision S rooted on U . By Lemma 4.16 (p. 61), G contains a semi-subdivision S' rooted on U and a compatible mapping w.r.t. S' , which is a contradiction by Lemma 4.15 (p. 59). \square

Lemma 3.1 (p. 3) ensues by combining Lemmas 3.2 (p. 6) and 4.1 (p. 35).

5 There is no minimal counterexample

We complete the proof of Theorem 1.2 with the following lemma:

Lemma 5.1. *Every connected planar graph on at least 3 vertices contains a configuration (C_I) or (C_{II}) .*

Lemmas 3.1 (p. 3) and 5.1 together guarantee that every connected planar graph other than K_3 and K_5^- admits a good path decomposition – it is trivial to verify for connected planar graphs on at most 2 vertices. To prove Lemma 5.1, we first need some definitions and structural observations on planar graphs.

Given a planar graph G , a *1-contraction* of G is an induced subgraph H of G on at least 2 vertices together with a vertex $u_1 \in V(H)$ such that all vertices have the same degree in G and H , except possibly for u_1 . A graph H on at least 3 vertices is a *2-contraction* of G if there exists an edge $u_1u_2 \in E(H)$ such that $H \setminus u_1u_2$ is a subgraph of G and every vertex $v \notin \{u_1, u_2\}$ satisfies $d_H(v) = d_G(v)$. Additionally, there exists a (u_1, u_2) -path in G with all internal vertices in $V(G) \setminus V(H)$.

The *damaged vertices* of a p -contraction ($p \in \{1, 2\}$) are the vertices $\{u_1\}, \{u_1, u_2\}$ above. Note that any induced subgraph of G (on at least 2 vertices) can be turned into a 1-contraction by selecting an arbitrary vertex as u_1 . Note that a 1-contraction with damaged vertex u_1 can be turned into a 2-contraction by selecting an arbitrary neighbour of u_1 in H , unless the vertex u_1 has no neighbour in H . Note also that any p -contraction ($p \in \{1, 2\}$) of G is a minor of G hence is planar.

A 2-contraction H' of G is *smaller* than a 2-contraction H of G with damaged vertices $\{u_1, u_2\}$ if $V(H') \subsetneq V(H)$ and each of u_1 and u_2 either does not belong to $V(H')$ or is a damaged vertex of H' . We may simply refer to a *smaller 2-contraction than H* if the damaged vertices of H are clear from context. A 2-contraction H of G is *minimal* if G admits no smaller 2-contraction.

Claim 5.2. *Let G be a connected planar graph. Any minimal 2-contraction of G either contains a non-damaged vertex of degree at most 4 or is 3-connected.*

Proof. Assume for a contradiction that H with damaged vertices u_1, u_2 is a counterexample to the statement: H is minimal, not 3-connected and every vertex in $V(H) \setminus \{u_1, u_2\}$ has degree at least 5 in H (hence in G). By definition of a 2-contraction, $V(H) \setminus \{u_1, u_2\}$ is non-empty. We note that it suffices to exhibit a 2-contraction H' of G smaller than H .

Since H is minimal, it is connected. Note that H is in fact 2-connected. Otherwise, let x be a cut-vertex in H . There is a connected component C of $H \setminus \{x\}$ containing none of $\{u_1, u_2\}$. The graph $G[C \cup \{x\}]$ is a 1-contraction of G with damaged vertex x . Note that C contains at least 5 vertices, as C is non-empty and every vertex in C has degree at least 5 in G . We select an arbitrary neighbour y of x in C , and note that $G[C \cup \{x\}]$ is a 2-contraction of G with damaged vertices x, y and $u_1, u_2 \notin C$. This yields a smaller 2-contraction than H , a contradiction.

Therefore, H is 2-connected but not 3-connected. Let x_1, x_2 be a vertex cut of H . Since $u_1u_2 \in E(H)$, u_1 and u_2 do not belong to different connected components of $H \setminus \{x_1, x_2\}$. Let C be a connected component $H \setminus \{x_1, x_2\}$ that contains no u_i . Since H is 2-connected, there is a path between x_1 and x_2 whose internal vertices belong to $V(H) \setminus C$. We obtain a 2-contraction of G with damaged vertices $\{x_1, x_2\}$ where each u_i either does not belong to it or is a damaged vertex, hence a contradiction. \square

Claim 5.3. *Let H be a minimal 2-contraction of a planar graph G . Either H contains a non-damaged vertex of degree at most 4, or there are four non-damaged vertices $\{v_1, v_2, v_3, v_4\}$ of degree 5 with respect to which H is almost 4-connected.*

Proof. Assume that H does not contain any non-damaged vertex of degree 4 or less, and let u_1, u_2 be the damaged vertices of H .

We first assume that H is 4-connected. Then it suffices to argue that there are four vertices of degree 5 in $V(H) \setminus \{u_1, u_2\}$. Since H is 4-connected, we have $d_H(u_1), d_H(u_2) \geq 4$. By Euler's formula, we have $\sum_{x \in V(H)} (d(x) - 6) \leq -12$, hence $\sum_{x \in V(H) \setminus \{u_1, u_2\}} (d(x) - 6) \leq -8$. Since every non-damaged vertex of H has degree at least 5, the conclusion follows.

Therefore, we may assume the graph H is not 4-connected. Consider an embedding of H where u_1 and u_2 lie on the outer-face. Since $u_1 u_2 \in E(H)$ and H is planar since it is a 2-contraction of G , such an embedding exists. For any vertex cut X of H that has size 3, let p be the number of connected components in $H \setminus X$. Consider the minor of H obtained by contracting each connected component into a single vertex. Since H is 3-connected (by Claim 5.2), every resulting vertex is adjacent to all three vertices in X , which yields a $K_{3,p}$ -minor where p is the number of connected components. Since H is planar, there is no $K_{3,3}$ -minor, hence $p = 2$. At most one of the two connected components of $H \setminus X$ contains damaged vertices. If one of them contains damaged vertices, we let $I(X)$ be the connected component of $H \setminus X$ that does not contain damaged vertices and $E(X)$ be the connected component of $H \setminus X$ that contains a damaged vertex. If neither component contains damaged vertices, then $u_1, u_2 \in X$. Since H is 3-connected and u_1, u_2 belong to the outer-face, there is exactly one connected component that contains no vertex of the outer-face; We let $I(X)$ be that component. Let $E(X)$ be the other component. Observe that $E(X)$ contains at least one vertex of the outer-face.

Among all vertex cuts of H that have size 3, we select a vertex cut $\{x_1, x_2, x_3\}$ which minimizes $|I(\{x_1, x_2, x_3\})|$. We first argue that $C = I(\{x_1, x_2, x_3\})$ contains at least four vertices of degree 5. Indeed, since $\{x_1, x_2, x_3\}$ is minimal, every x_i has at least 2 neighbors in C . Hence by Euler's formula on the graph C , we have $\sum_{x \in C} (d(x) - 6) + 3 \times 2 \leq -12$. Since every vertex in C has degree at least 5 in H , there are four non-damaged vertices $\{v_1, v_2, v_3, v_4\}$ in C that have degree 5 in H .

It remains to argue that H is almost 4-connected with respect to them. We show that no 3-cut of H separates two vertices of $X \cup C$. Observe that this proves the two properties of almost 4-connectivity. Assume for a contradiction that there is a vertex cut $Y = \{y_1, y_2, y_3\}$ such that two vertices $z_1, z_2 \in X \cup C$ are in different connected components of $H \setminus Y$. Without loss of generality, consider $z_1 \in I(Y)$ and $z_2 \in E(Y)$.

Note that $H[X \cup C]$ is connected and $z_1, z_2 \in X \cup C$. However, z_1 and z_2 are in different connected components of $H \setminus Y$. Since z_1, z_2 cannot be separated by a vertex cut contained in $X \cup E(X)$, at least one of $\{y_1, y_2, y_3\}$, say y_1 , belongs to C . Since $\{x_1, x_2, x_3\}$ minimizes $|I(\{x_1, x_2, x_3\})|$ over all vertex cuts of H of size 3, at least one of $\{y_1, y_2, y_3\}$, say y_3 , belongs to $E(\{x_1, x_2, x_3\})$.

We consider two cases depending on the cardinal of $\{x_1, x_2, x_3\} \cap I(Y)$.

- Assume that $I(Y)$ contains exactly one vertex in $\{x_1, x_2, x_3\}$, say x_1 .

If $y_2 \in E(X)$, then we claim that $\{x_1, y_1\}$ is a 2-cut that separates z_1 from z_2 , a contradiction with the 3-connectivity of H . To prove it, we note that in $Y \cup I(Y)$, there is a path between z_1 and each of $\{y_1, y_2, y_3\}$ whose internal vertices belong to $I(Y)$. However, since $\{x_1, x_2, x_3\}$ is a vertex cut of H and neither x_2 nor x_3 belongs to $Y \cup I(Y)$, every path between z_1 and y_2 whose internal vertices belong to $I(Y)$ involves the vertex x_1 . Therefore, $\{x_1, y_1\}$ separates z_1 and y_2 , hence the conclusion.

So $y_2 \in X \cup C$, and we now claim that $\{x_1, y_1, y_2\}$ is a vertex cut of H , by the same argument: each path between z_1 and y_3 with internal vertices in $I(Y)$ involves x_1 . Observe that it contradicts the choice of $\{x_1, x_2, x_3\}$.

- Assume from now on that $I(Y)$ contains both x_2 and x_3 , while $x_1 \in E(Y)$. As above, we argue that $\{y_1, x_2, x_3\}$ is a vertex cut of H , which contradicts the choice of $\{x_1, x_2, x_3\}$. For completeness, we include the adapted proof. We note that in $\{x_1, x_2, x_3\} \cup I(\{x_1, x_2, x_3\})$, there is a path between z_1 and each of $\{x_1, x_2, x_3\}$ whose internal vertices belong to $I(\{x_1, x_2, x_3\})$. However, since Y is a vertex cut of H and neither y_2 nor y_3 belongs to $\{x_1, x_2, x_3\} \cup I(\{x_1, x_2, x_3\})$, every path between z_1 and x_1 whose internal vertices belong to $I(\{x_1, x_2, x_3\})$ involves the vertex y_1 . Therefore, $\{y_1, x_2, x_3\}$ separates z_1 and x_1 , hence the conclusion. □

We can now use Claim 5.3 to obtain Lemma 5.1 (p. 92).

Proof of Lemma 5.1. Let G be a non-empty connected planar graph which contains no (C_I) configuration. Let u_1 be the only vertex with degree at most 4 in G if any, and an arbitrary vertex otherwise. Note that since G is connected and contains at least 3 vertices, the graph G with damaged vertex u_1 is a 1-contraction of G . Furthermore, the vertex u_1 has at least one neighbour u_2 , and the graph G with damaged vertices u_1 and u_2 is a 2-contraction of G . Let H be a minimal 2-contraction of G that is either precisely G with damaged vertices u_1, u_2 or smaller than it.

Every non-damaged vertex in H has the same degree in H and in G , and u_1 is not a non-damaged vertex in H . Therefore, Claim 5.3 applied to H yields that H is almost 4-connected w.r.t. a 4-family $\{v_1, v_2, v_3, v_4\}$. Hence G is almost 4-connected w.r.t. $\{v_1, v_2, v_3, v_4\}$, as desired. □

References

- [1] László Lovász. “On covering of graphs”. In: *Theory of Graphs (ed. P. Erdős, G. Katona)*. Akadémiai Kiadó, Budapest, 1968, pp. 231–236.
- [2] Odile Favaron and Mekkia Kouider. “Path partitions and cycle partitions of Eulerian graphs of maximum degree 4”. In: *Studia Scientiarum Mathematicarum Hungarica* 23 (1988), pp. 237–244.
- [3] László Pyber. “Covering the Edges of a Connected Graph by Paths”. In: *Journal of Combinatorial Theory, Series B* 66.1 (1996), pp. 152–159. ISSN: 0095-8956. DOI: <https://doi.org/10.1006/jctb>
- [4] Genghua Fan. “Path decompositions and Gallai’s conjecture”. In: *Journal of Combinatorial Theory, Series B* 93.2 (2005), pp. 117–125. ISSN: 0095-8956. DOI: <https://doi.org/10.1016/j.jctb>
- [5] Fábio Botler and Maycon Sambinelli. “Towards Gallai’s path decomposition conjecture”. In: *Journal of Graph Theory* 97.1 (2021), pp. 161–184. DOI: <https://doi.org/10.1002/jgt.22647>.
- [6] Philipp Kindermann, Lena Schlipf, and André Schulz. “On Gallai’s conjecture for series-parallel graphs and planar 3-trees”. In: *CoRR* abs/1706.04130 (2017). arXiv: 1706.04130.
- [7] Fábio Botler, Maycon Sambinelli, Rafael S. Coelho, and Orlando Lee. “Gallai’s path decomposition conjecture for graphs with treewidth at most 3”. In: *Journal of Graph Theory* 93.3 (2020), pp. 328–349. DOI: [10.1002/jgt.22489](https://doi.org/10.1002/jgt.22489).

- [8] Marthe Bonamy and Thomas J. Perrett. “Gallai’s path decomposition conjecture for graphs of small maximum degree”. In: *Discrete Mathematics* 342.5 (2019), pp. 1293 – 1299. ISSN: 0012-365X. DOI: <https://doi.org/10.1016/j.disc.2019.01.005>.
- [9] Yanan Chu, Genghua Fan, and Qinghai Liu. “On Gallai’s conjecture for graphs with maximum degree 6”. In: *Discrete Mathematics* 344.2 (2021), p. 112212. ISSN: 0012-365X. DOI: <https://doi.org/10.1016/j.disc.2020.112212>.
- [10] Fábio Botler, Andrea Jiménez, and Maycon Sambinelli. “Gallai’s path decomposition conjecture for triangle-free planar graphs”. In: *Discrete Mathematics* 342.5 (2019), pp. 1403 – 1414. ISSN: 0012-365X. DOI: <https://doi.org/10.1016/j.disc.2019.01.020>.
- [11] Fábio Botler and Andrea Jiménez. “On path decompositions of $2k$ -regular graphs”. In: *Discrete Mathematics* 340.6 (2017), pp. 1405–1411. DOI: [10.1016/j.disc.2016.09.029](https://doi.org/10.1016/j.disc.2016.09.029).
- [12] Xingxing Yu. “Subdivisions in Planar Graphs”. In: *Journal of Combinatorial Theory, Series B* 72.1 (1998), pp. 10 – 52. ISSN: 0095-8956. DOI: <https://doi.org/10.1006/jctb.1997.1774>.