

# FIXED POINTS OF KOCH'S MAPS

VAN TU LE

ABSTRACT. We study endomorphisms constructed by Sarah Koch in [Koc13] and we focus on the eigenvalues of the differential of such maps at its fixed points. In [Koc13], to each post-critically finite unicritical polynomial, Koch associated a post-critically algebraic endomorphism of  $\mathbb{C}\mathbb{P}^k$ . Koch showed that the eigenvalues of the differentials of such maps along periodic cycles outside the post-critical sets have modulus strictly greater than 1. In this article, we show that the eigenvalues of the differentials at fixed points are either 0 or have modulus strictly greater than 1. This confirms a conjecture proposed by the author in his thesis. We also provide a concrete description of such values in terms of the multiplier of a unicritical polynomial.

## 1. INTRODUCTION

Let  $M$  be either  $\mathbb{C}^n$  or  $\mathbb{C}\mathbb{P}^n$  and  $f: M \rightarrow M$  be a holomorphic endomorphism. Denote by  $f^{\circ m} = f \circ f \circ \dots \circ f$  the  $m$ -th composition of  $f$ . A point  $z \in M$  is called a *preperiodic point* of *preperiod*  $k$  and of *period*  $m$  if  $f^{\circ(k+m)}(z) = f^{\circ k}(z)$  and  $k, m$  are the smallest integers satisfying such a property. A preperiodic point of preperiod 0 is called a *periodic point*. A periodic point of period 1 is called a *fixed point*. Given a periodic point  $z$  of period  $m$ , a value  $\lambda \in \mathbb{C}$  is called an *eigenvalue of  $f$  along the orbit of  $z$*  (or *at the fixed point  $z$* ) if  $\lambda$  is an eigenvalue of the differential  $D_z f^{\circ m}: T_z M \rightarrow T_z M$ .

A point  $z \in M$  is called a *critical point* if the differential  $D_z f: T_z M \rightarrow T_{f(z)} M$  is not invertible. The set  $C(f)$  containing all critical points of  $f$  is called the *critical set* of  $f$ . The set

$$PC(f) := \bigcup_{j \geq 1} f^{\circ j}(C(f))$$

is called the *post-critical set* of  $f$ . The endomorphism  $f$  is called *post-critically algebraic* if  $PC(f)$  is an algebraic set of codimension one in  $M$ . When  $\dim M = 1$ , post-critically algebraic rational maps are called *post-critically finite* rational maps.

The family of post-critically finite rational maps is one of the most important families of maps in the theory of one dimensional complex dynamics. In higher dimension, post-critically algebraic endomorphisms are interesting family of maps since many results, which are well-known for post-critically finite rational maps,

*Date:* October 19, 2021.

2020 Mathematics Subject Classification: 32H50, 37F99.

*Keywords:* holomorphic dynamics, holomorphic endomorphisms, fixed points, eigenvalues.

remain unknown. We refer to [Ron08],[Ast20],[IRS19],[GV19], [Ji20], [Le20b] for some recent studies about post-critically algebraic endomorphisms. In this article, we focus on the following conjecture proposed by the author in his thesis [Le20a].

**Conjecture 1.** Let  $f$  be a post-critically algebraic endomorphism of  $\mathbb{C}\mathbb{P}^n$ ,  $n \geq 2$  of degree  $d \geq 2$  and  $\lambda$  be an eigenvalue of  $f$  along a periodic cycle. Then either  $\lambda = 0$  or  $|\lambda| > 1$ .

The conjecture has been verified by the author in the case  $n = 2$  and in the case in any dimension with the periodic cycles outside the post-critical set. In this article, we shall verify the conjecture for the family of post-critically algebraic endomorphisms associated to unicritical polynomials constructed by Sarah Koch in [Koc13], or *Koch maps* for short.

We shall now describe the family of Koch maps we want to study and we refer to [Koc13] for the original construction. Throughout this article, we fix

$$d \in \mathbb{N}, d \geq 2 \text{ and } \beta^d = 1, \beta \neq 1.$$

The maps  $\{G_{k,m}: \mathbb{C}^{k+m-1} \rightarrow \mathbb{C}^{k+m-1} \mid (k,m) \in \mathbb{N} \times \mathbb{N}^*\}$  constructed by Sarah Koch are of the following forms (see [Koc13, Proposition 6.1 - 6.2]),

- if  $k = 0$ ,

$$G_{0,m}: \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_{m-1} \end{pmatrix} \mapsto \begin{pmatrix} -x_{m-1}^d \\ x_1^d - x_{m-1}^d \\ \vdots \\ x_{m-2}^d - x_{m-1}^d \end{pmatrix},$$

- if  $k \neq 0$ ,

$$G_{k,m}: \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_{k+m-1} \end{pmatrix} \mapsto \begin{pmatrix} \left(-\frac{\beta x_{k+m-1} - x_{k-1}}{\beta - 1}\right)^d \\ \left(x_1 - \frac{\beta x_{k+m-1} - x_{k-1}}{\beta - 1}\right)^d \\ \vdots \\ \left(x_{k+m-2} - \frac{\beta x_{k+m-1} - x_{k-1}}{\beta - 1}\right)^d \end{pmatrix}.$$

The map  $G_{k,m}$  induces a holomorphic endomorphism of  $\mathbb{C}\mathbb{P}^{k+m-2}$  which is closely related to maps on moduli spaces used in Thurston's topological characterization of rational maps. We refer to [Koc13],[Koc08],[DH93] for further discussion. In [Koc13], Koch showed that  $G_{k,m}$  is post-critically algebraic. It is natural to ask whether Conjecture 1 is true for  $G_{k,m}$ . The eigenvalues of Koch maps along a periodic cycle outside the post-critical set are well understood. It is a consequence of its construction that those values has modulus strictly bigger than 1.

**Theorem 1** (Corollary 7.2 [Koc13]). Let  $\mu$  be an eigenvalue of  $G_{k,m}$  along a periodic cycle outside the post-critical set. Then  $|\mu| > 1$ .

We refer also to [BEK20] for a further discussion about the arithmetics of such values. In [Koc13], Koch asked whether we have the same conclusion for eigenvalues along a cycle inside the post-critical set. In this article, we answer this question in the positive (and hence verify Conjecture 1) for the case when the cycle is a fixed point.

**Theorem A.** Let  $\mu$  be an eigenvalue of a map  $G_{k,m}$  at a fixed point. Then, either  $\mu = 0$  or  $|\mu| > 1$ .

In fact, we can have even better understanding about the values of such eigenvalues. Thanks to Theorem 1, we only need to study the eigenvalues at a fixed point inside the post-critical set of  $G_{k,m}$ . However, the original construction does not provide much information about fixed points inside post-critical set. In order to explain our result, let us take a closer look at  $G_{k,m}$ .

Let  $(k, m) \in \mathbb{N} \times \mathbb{N}^*$ . To a point  $z \in \mathbb{C}^{k+m-1}$ , we can associate a polynomial  $P_z$  of the following form

$$P_z(t) = \begin{cases} t^d - z_{m-1}^d & \text{if } k = 0 \\ \left(t - \frac{\beta z_{k+m-1} - z_{k-1}}{\beta-1}\right)^d & \text{if } k \neq 0. \end{cases}$$

If  $z$  is a fixed point of  $G_{k,m}$ , then for all  $1 \leq i \leq k+m-1$ ,  $z_i = P_z^{\circ i}(0)$ . Moreover,  $P_z$  is post-critically finite. Indeed,

- if  $k = 0$  then  $P_z(z_{m-1}) = 0$ ,
- if  $k \neq 0$  then

$$\begin{aligned} P_z(z_{k+m-1}) &= \left(z_{k+m-1} - \frac{\beta z_{k+m-1} - z_{k-1}}{\beta-1}\right)^d = \left(\frac{-z_{k+m-1} + z_{k-1}}{\beta-1}\right)^d \\ &= \left(\frac{\beta z_{k-1} - \beta z_{k+m-1}}{\beta-1}\right)^d = P_z(z_{k-1}) \end{aligned}$$

We shall call  $P_z$  the *polynomial associated to  $z$*  since  $P_z$  plays an important role in the study of the eigenvalues of  $G_{k,m}$  at a fixed point  $z$ . More precisely, our main result, which completes the description of eigenvalues of Koch maps at fixed points, is the following.

**Theorem A'.** Let  $\mu$  be an eigenvalue of a map  $G_{k,m}$  at a fixed point  $z = (z_1, \dots, z_{k+m-1})$ . Let  $P_z$  be the polynomial associated to  $z$  and  $z_1$  is preperiodic of preperiod  $k'$  and of period  $m'$  to a cycle of multiplier  $\lambda$  under  $P_z$ .

Only one of the following cases happens:

- (1)  $\mu = 0$ .
- (2)  $\mu$  is an eigenvalue of a map  $G_{k',m'}$  at a fixed point outside the post-critical set  $PC(G_{k',m'})$ .
- (3) We have

$$\mu^m = \lambda^{\frac{m}{m'}}, \mu^{m'} \neq \lambda.$$

We can see that Theorem A is a direct consequence of Theorem A'. Indeed, if Case 1 or Case 2 happens, Theorem A follows from Theorem 1. If Case 3 happens, since the polynomial  $P_z$  is a post-critically finite polynomial, Theorem A follows from the equation  $\mu^m = \lambda^{\frac{m}{m'}}$  and the fact that a non-vanishing multiplier of a post-critically finite polynomial has modulus strictly bigger than 1 (see [Mil11, Corollary 14.5])

Let us explain briefly our approach. Instead of using the original construction of  $G_{k,m}$ , we introduce

- a partial order  $\preceq$  on  $\mathbb{N} \times \mathbb{N}^*$ ,
- a dynamically equivalent family of maps, that we denote by

$$\{F_{k,m} : \mathcal{M}_{k,m} \rightarrow \mathcal{M}_{k,m}, (k,m) \in \mathbb{N} \times \mathbb{N}^*\},$$

where  $\mathcal{M}_{k,m}$  is a subspace of the vector spaces of complex sequences  $\mathbb{C}^{\mathbb{N}^*}$ .

More precisely, with the convention  $x_0 := 0$ , the space  $\mathcal{M}_{k,m}$  is defined as

$$\mathcal{M}_{k,m} = \{\mathbf{x} = (x_i)_{i \geq 1} \mid \forall i \geq k+1, x_i = x_{i+m} \text{ and } \beta x_{k+m} - x_k = 0\}$$

and the map  $F_{k,m} : \mathcal{M}_{k,m} \rightarrow \mathcal{M}_{k,m}$  is defined as

$$F_{k,m}(\mathbf{x}) = \mathbf{y} \Leftrightarrow \begin{cases} \beta y_{k+m} - y_k = 0 \\ y_i = x_{i-1}^d + y_1 \quad \text{for all } i \geq 2. \end{cases}$$

The construction shall be presented in details in Section 2. For each  $(k,m) \in \mathbb{N} \times \mathbb{N}^*$ , the maps  $F_{k,m}$  and  $G_{k,m}$  are conjugate. Thus, to prove Theorem A', we need to study an eigenvalue  $\mu$  of  $F_{k,m}$  at a fixed point  $\mathbf{z} \in \mathcal{M}_{k,m}$ . The full statement of what we can prove is Theorem B'. Briefly, to each fixed point  $\mathbf{z}$  in  $\mathcal{M}_{k,m}$ , we associate a pair of integers  $(k', m')$  and a post-critically finite polynomial  $P_z$  whose critical value is preperiodic of preperiod  $k'$  to a cycle of period  $m'$ . The associated polynomial  $P_z$  for a  $\mathbf{z} = (z_i)_{i \geq 1} \in \mathcal{M}_{k,m}$  is simply  $P_z(t) = t^d + z_1$ . The partial order characterizes the following property of the family  $\{F_{k,m}\}$ :

$$(k', m') \preceq (k, m) \Leftrightarrow \begin{cases} \mathcal{M}_{k',m'} \subseteq \mathcal{M}_{k,m}, \\ F_{k,m}|_{\mathcal{M}_{k',m'}} = F_{k',m'}. \end{cases}$$

Moreover, we shall show that  $\mathbf{z} \notin PC(F_{k',m'})$ . Thus, if  $(k', m') = (k, m)$ , our fixed point  $\mathbf{z}$  is outside the post-critical set  $PC(F_{k,m})$  and we are in Case 2.

If  $(k', m') \neq (k, m)$ , then  $\mathcal{M}_{k',m'}$  is a proper subset of  $\mathcal{M}_{k,m}$  which is invariant under  $F_{k,m}$  and the restriction of  $F_{k,m}$  to  $\mathcal{M}_{k',m'}$  is exactly  $F_{k',m'}$ . If  $\mu$  has associated eigenvectors tangent to  $\mathcal{M}_{k',m'}$ , since  $\mathbf{z} \notin PC(F_{k',m'})$ , we are again in the Case 2. Otherwise,  $\mu$  is the eigenvalue of the transpose  $D_{\mathbf{z}}F_{k,m}^*$  of the derivative  $D_{\mathbf{z}}F_{k,m}$  acting on the annihilator  $\mathcal{M}_{k',m'}^0 = \{\omega \in \mathcal{M}_{k,m}^* \mid \omega|_{\mathcal{M}_{k',m'}} = 0\}$ . In such a case, either  $\mu = 0$  or we will show that  $\mathcal{M}_{k',m'}^0$  has a set of generators on which  $D_{\mathbf{z}}F_{k,m}^*$  acts cyclically and we obtain Case 3 by solving a linear algebra problem.

**Acknowledgement:** This article is the improvement of Chapter 2 of the author's thesis [Le20a]. The author is grateful to his supervisors Xavier Buff and

Jasmin Raissy for their support, suggestions and encouragement. The author would like to thank also Valentin Huguin for his comments and useful discussions. This work is supported by the fellowship of Centre International de Mathématiques et d'Informatique de Toulouse (CIMI).

## 2. AN ALTERNATIVE CONSTRUCTION OF KOCH MAPS

2.1. **Construction of  $F_{k,m}$  and  $\mathcal{M}_{k,m}$ .** Recall that in this article, we fix

$$d \in \mathbb{N}, d \geq 2 \text{ and } \beta^d = 1, \beta \neq 1$$

Let  $(k, m) \in \mathbb{N} \times \mathbb{N}^*$  and denote by  $\mathcal{E} = \mathbb{C}^{\mathbb{N}^*}$  the vector space of complex sequences  $\mathbf{x} = (x_i)_{i \geq 1}$ . Set  $\mathbf{0} := (0, 0, \dots)$ . Let  $\mathcal{L} \subset \mathcal{E}$  be the one-dimensional subspace consisting of constant sequences,

$$\mathcal{L} = \{\mathbf{x} \in \mathcal{E} \mid \forall i, j \geq 1, \quad x_i = x_j\},$$

and  $\mathcal{H}_{k,m} \subset \mathcal{E}$  be the hyperspace of  $\mathcal{E}$  defined by

$$\mathcal{H}_{k,m} := \{\mathbf{x} \in \mathcal{E} \mid \beta x_{k+m} - x_k = 0\}.$$

with the convention  $x_0 := 0$ . In particular, when  $k = 0$ ,

$$\mathcal{H}_{0,m} = \{\mathbf{x} \in \mathcal{E} \mid x_m = 0\}.$$

**Lemma 2.1.** *Given  $(k, m) \in \mathbb{N} \times \mathbb{N}^*$ , we have  $\mathcal{E} = \mathcal{H}_{k,m} \oplus \mathcal{L}$ .*

*Proof.* On the one hand, given  $\mathbf{x} \in \mathcal{E}$ , define  $\mathbf{y} \in \mathcal{E}$  by

$$y_i := x_i - \kappa \text{ with } \kappa = \begin{cases} x_m & \text{if } k = 0 \\ \frac{\beta x_{k+m} - x_k}{\beta - 1} & \text{if } k \geq 1 \end{cases}.$$

Then  $\beta y_{k+m} - y_k = 0$  hence  $\mathbf{y} \in \mathcal{H}_{k,m}$ . Note that  $\mathbf{x} - \mathbf{y} \in \mathcal{L}$  hence

$$\mathcal{E} = \mathcal{H}_{k,m} + \mathcal{L}.$$

On the other hand, assume  $\mathbf{x} \in \mathcal{H}_{k,m} \cap \mathcal{L}$ . Then  $\beta x_{k+m} - x_k = 0$  and  $x_{k+m} = x_k$ . Since  $\beta \neq 1$ , we have  $x_k = x_{k+m} = 0$ . Moreover,  $\mathbf{x}$  is a constant sequence. Thus,  $\mathbf{x}$  vanishes identically; that is

$$\mathcal{H}_{k,m} \cap \mathcal{L} = \{\mathbf{0}\}. \quad \square$$

Denote by

$$\pi_{k,m}: \mathcal{E} \rightarrow \mathcal{E}$$

the projection to  $\mathcal{H}_{k,m}$  parallel to  $\mathcal{L}$ . In particular,  $\pi_{k,m}(\mathcal{E}) = \mathcal{H}_{k,m}$ . Consider the map  $\mathcal{Q}: \mathcal{E} \rightarrow \mathcal{E}$  defined by

$$\mathcal{Q}(\mathbf{x}) = \mathbf{y} \quad \text{with} \quad y_1 := 0 \text{ and } y_i = x_{i-1}^d, i \geq 1.$$

**Definition 2.2.** Given  $(k, m) \in \mathbb{N} \times \mathbb{N}^*$ , the map  $F_{k,m} := \mathcal{E} \rightarrow \mathcal{E}$  is defined as

$$F_{k,m} := \pi_{k,m} \circ \mathcal{Q}.$$

We shall now study some important properties of  $F_{k,m}$ .

### 2.1.1. Properties of $F_{k,m}$ .

**Lemma 2.3.** Given  $(k, m) \in \mathbb{N} \times \mathbb{N}^*$ , for every  $\mathbf{x}, \mathbf{y} \in \mathcal{E}$ , we have

$$F_{k,m}(\mathbf{x}) = \mathbf{y} \Leftrightarrow \begin{cases} \beta y_{k+m} - y_k = 0 \\ y_i = x_{i-1}^d + y_1 \quad \text{for all } i \geq 2. \end{cases}$$

In particular, with the convention  $x_0 := 0$ , we have

$$y_1 = \begin{cases} -x_{m-1}^d & \text{if } k = 0, \\ -\frac{\beta x_{k+m-1}^d - x_{k-1}^d}{\beta - 1} & \text{if } k \geq 1. \end{cases}$$

*Proof.* Assume  $\mathbf{x} \in \mathcal{E}$  and  $\mathbf{y} = \pi_{k,m}(\mathcal{Q}(\mathbf{x})) = F_{k,m}(\mathbf{x})$ . On the one hand,  $\mathbf{y} \in \pi_{k,m}(\mathcal{E}) = \mathcal{H}_{k,m}$ , i.e.

$$\beta y_{k+m} - y_k = 0.$$

On the other hand, set  $\mathbf{z} = \mathcal{Q}(\mathbf{x})$ , i.e.  $z_1 = 0$  and  $z_i = x_{i-1}^d$  for all  $i \geq 1$ . Then since  $\mathbf{y} = \pi_{k,m}(\mathbf{z})$ , for all  $i \geq 1$ ,

$$y_i = z_i - \kappa \text{ with } \kappa := \begin{cases} z_m & \text{if } k = 0 \\ \frac{\beta z_{k+m} - z_k}{\beta - 1} & \text{if } k \geq 1 \end{cases}.$$

In particular,

$$y_1 = z_1 - \kappa = -\kappa = \begin{cases} -x_{m-1}^d & \text{if } k = 0, \\ -\frac{\beta x_{k+m-1}^d - x_{k-1}^d}{\beta - 1} & \text{if } k \geq 1. \end{cases}$$

whence for all  $i \geq 2$ ,  $y_i = z_i + y_1$ , i.e.

$$y_i = x_{i-1}^d + y_1.$$

Conversely, assume  $\mathbf{x}, \mathbf{y} \in \mathcal{E}$  such that  $\beta y_{k+m} - y_k = 0$  and for all  $i \geq 2$ ,  $y_i = x_{i-1}^d + y_1$ . In particular,  $\mathbf{y} \in \mathcal{H}_{k,m}$ . Set  $\mathbf{z} = \mathbf{y} - \mathcal{Q}(\mathbf{x})$ . Then for all  $i \geq 2$ ,

$$z_i = y_i - x_{i-1}^d = y_1$$

Moreover,  $z_1 = x_0^d + y_1 = y_1$ . Hence  $\mathbf{z} \in \mathcal{L}$ . In other words,

$$\mathbf{y} = \pi_{k,m}(\mathcal{Q}(\mathbf{x})) = F_{k,m}(\mathbf{x}). \quad \square$$

Although  $F_{k,m}$  is defined on a vector space of infinite dimension, we will now see that the dynamics of  $F_{k,m}$  is captured entirely by some finite dimensional vector space. Given a sequence  $\mathbf{x} \in \mathcal{E}$ ,  $\mathbf{x}$  is preperiodic of preperiod  $k$  to a cycle of period  $m$  if for all  $i \geq k + 1$ ,  $x_i = x_{i+m}$  and  $k, m$  are the smallest integers satisfying such

a property. Given  $(k, m) \in \mathbb{N} \times \mathbb{N}^*$ , let  $\mathcal{P}_{k,m} \subset \mathcal{E}$  be the subspace of preperiodic sequences of preperiod at most  $k$  to a cycle of period dividing  $m$ , i.e.

$$\mathcal{P}_{k,m} := \{\mathbf{x} \in \mathcal{E} \mid x_{i+m} = x_i \text{ for } i \geq k+1\}.$$

Since sequences in  $\mathcal{P}_{k,m}$  are uniquely determined by the first  $k+m$  entries, the vector space  $\mathcal{P}_{k,m}$  has finite dimension  $k+m$ .

**Definition 2.4.** Given  $(k, m) \in \mathbb{N} \times \mathbb{N}^*$ , define

$$\mathcal{M}_{k,m} := \mathcal{P}_{k,m} \cap \mathcal{H}_{k,m}.$$

Note that the constant sequence  $(1, 1, \dots)$  is in  $\mathcal{P}_{k,m} \setminus \mathcal{H}_{k,m}$  and that  $\mathcal{H}_{k,m}$  has codimension one in  $\mathcal{E}$ . Hence,  $\mathcal{M}_{k,m}$  is a vector space of dimension  $k+m-1$ . The following two lemmas show the importance of  $\mathcal{M}_{k,m}$  to the dynamics of  $F_{k,m}$ .

**Lemma 2.5.** *We have  $F_{k,m}(\mathcal{M}_{k,m}) = \mathcal{M}_{k,m}$  and  $F_{k,m}: \mathcal{M}_{k,m} \rightarrow \mathcal{M}_{k,m}$  is a nondegenerate homogeneous map of degree  $d$ .*

*Proof.* Let us first prove that  $\mathcal{Q}(\mathcal{M}_{k,m}) \subseteq \mathcal{P}_{k,m}$ . Assume  $\mathbf{x} \in \mathcal{M}_{k,m}$ . Since  $\mathbf{x} \in \mathcal{P}_{k,m} \cap \mathcal{H}_{k,m}$ ,  $\mathbf{x}$  has preperiod  $k$  and period dividing  $m$  and  $\beta x_{k+m} - x_m = 0$ . In particular, since  $\beta^d = 1$ , we have  $x_{k+m}^d = x_m^d$ . Consequently, setting  $\mathbf{y} := \mathcal{Q}(\mathbf{x})$ ,

$$\begin{cases} y_{k+1} = x_k^d + y_1 = x_{k+m}^d + y_1 = y_{k+m+1} \\ y_i = x_{i-1}^d + y_1 = x_{m+i-1}^d + y_1 = y_{i+m} \end{cases} \quad \text{for all } i \geq k+2.$$

Thus  $\mathcal{Q}(\mathbf{x}) \in \mathcal{P}_{k,m}$ .

Since  $\pi_{k,m}(\mathcal{P}_{k,m}) \subset \mathcal{P}_{k,m}$ ,  $F_{k,m}(\mathcal{M}_{k,m}) \subset \mathcal{P}_{k,m}$ . Since  $F_{k,m}(\mathcal{E}) \subset \mathcal{H}_{k,m}$ ,

$$F_{k,m}(\mathcal{M}_{k,m}) \subset \mathcal{P}_{k,m} \cap \mathcal{H}_{k,m} = \mathcal{M}_{k,m}.$$

Clearly, the map  $\mathcal{Q}$  is homogeneous of degree  $d$  and the map  $\pi_{k,m}$  is homogeneous of degree 1, thus  $F_{k,m}$  is homogeneous of degree  $d$ .

Let us now prove that  $F_{k,m}$  is nondegenerate, i.e.  $F_{k,m}^{-1}(\mathbf{0}) = \{\mathbf{0}\}$ . Assume  $\mathbf{x} \in \mathcal{M}_{k,m}$  and

$$\pi_{k,m} \circ \mathcal{Q}(\mathbf{x}) = F_{k,m}(\mathbf{x}) = \mathbf{0}.$$

Then  $\mathbf{y} := \mathcal{Q}(\mathbf{x}) \in \text{Ker}(\pi_{k,m}) = \mathcal{L}$ . By definition of  $\mathcal{Q}$ ,  $y_1 = 0$ . Since  $\mathbf{y} \in \mathcal{L}$ ,  $\mathbf{y}$  is a constant sequence thus  $\mathbf{y} = \mathbf{0}$ . This implies that  $x_i = 0$  for all  $i \geq 1$ , i.e.  $\mathbf{x} = \mathbf{0}$ .

Since  $\mathcal{M}_{k,m}$  has finite dimension and  $F_{k,m}: \mathcal{M}_{k,m} \rightarrow \mathcal{M}_{k,m}$  is homogeneous and nondegenerate,  $F_{k,m}$  is surjective, i.e.  $F_{k,m}(\mathcal{M}_{k,m}) = \mathcal{M}_{k,m}$ .  $\square$

**Lemma 2.6.** *We have that  $\bigcap_{n \geq 1} F_{k,m}^{\circ n}(\mathcal{E}) = \mathcal{M}_{k,m}$ .*

*Proof.* According to the previous lemma,  $F_{k,m}(\mathcal{M}_{k,m}) = \mathcal{M}_{k,m}$ . Thus,

$$\mathcal{M}_{k,m} \subseteq \bigcap_{n \geq 1} F_{k,m}^{\circ n}(\mathcal{E}).$$

Conversely, it is enough to prove that

$$(2.1) \quad \bigcap_{n \geq 2} F_{k,m}^{\circ n}(\mathcal{E}) \subset \mathcal{P}_{k,m}.$$

Indeed, since  $F_{k,m}(\mathcal{E}) \subset \mathcal{H}_{k,m}$  and  $\mathcal{M}_{k,m} = \mathcal{H}_{k,m} \cap \mathcal{P}_{k,m}$ , the inclusion (2.1) implies that

$$F_{k,m}(\mathcal{E}) \cap \bigcap_{n \geq 2} F_{k,m}^{\circ n}(\mathcal{E}) \subseteq \mathcal{H}_{k,m} \cap \mathcal{P}_{k,m},$$

and hence

$$\bigcap_{n \geq 1} F_{k,m}^{\circ n}(\mathcal{E}) \subseteq \mathcal{M}_{k,m} \subseteq \bigcap_{n \geq 1} F_{k,m}^{\circ n}(\mathcal{E}).$$

To prove (2.1), we show the following claim: for all  $n \geq 2$ , if  $\mathbf{y} \in F_{k,m}^{\circ n}(\mathcal{E})$ , then  $y_{i+m} = y_i$  for all  $i \in \{k+1, \dots, k+n-1\}$ .

Let us prove this claim by induction in  $n$ . If  $n = 2$ , assume  $\mathbf{y} \in F_{k,m}^{\circ 2}(\mathcal{E})$ , i.e.  $\mathbf{y} = F_{k,m}^{\circ 2}(\mathbf{x})$  for some  $\mathbf{x} \in \mathcal{E}$ . Setting  $\mathbf{z} = F_{k,m}(\mathbf{x})$ , according to Lemma 2.3, we have  $\beta z_{k+m} - z_k = 0$ . Thus, since  $\mathbf{y} = F_{k,m}(\mathbf{z})$  and since  $\beta^d = 1$ , also according to Lemma 2.3, we have

$$y_{k+1+m} = z_{k+m}^d + y_1 = z_k^d + y_1 = y_{k+1},$$

i.e. the claim is true for  $n = 2$ .

Assume that it holds for some  $n > 2$ . Assume  $\mathbf{y} \in F_{k,m}^{\circ(n+1)}(\mathcal{E})$ , i.e.  $\mathbf{y} = F_{k,m}(\mathbf{x})$  with  $\mathbf{x} \in F_{k,m}^{\circ n}(\mathcal{E})$ . The induction hypothesis implies that

$$x_{i+m-1} = x_{i-1} \text{ for all } i \in \{k+2, \dots, k+n\},$$

so that

$$y_{i+m} = y_i \text{ for all } i \in \{k+2, \dots, k+n\}.$$

In addition, since  $\mathbf{x} \in \mathcal{H}_{k,m}$ ,  $\beta x_{k+m} = x_k$ , so that  $y_{k+m+1} = y_{k+1}$ . Thus the claim is true for  $n+1$ .  $\square$

**2.2. The main result about  $F_{k,m}$ .** Lemma 2.5 and Lemma 2.6 allow us to restrict our study to the dynamics of  $F_{k,m}$  on  $\mathcal{M}_{k,m}$ . With a slight abuse of notations, from now on, we shall denote by  $F_{k,m}$  the restriction of  $F_{k,m} : \mathcal{E} \rightarrow \mathcal{E}$  to  $\mathcal{M}_{k,m}$ . The following result sums up the properties of  $F_{k,m} : \mathcal{M}_{k,m} \rightarrow \mathcal{M}_{k,m}$  which are important for us.

**Theorem B'.** Given  $(k, m) \in \mathbb{N} \times \mathbb{N}^*$  and let  $\mathbf{z} \in \mathcal{M}_{k,m} \setminus \{\mathbf{0}\}$  be a fixed point of the map  $F_{k,m} : \mathcal{M}_{k,m} \rightarrow \mathcal{M}_{k,m}$ . Let  $k'$  be the preperiod of the sequence  $\mathbf{z}$  and  $m'$  be its period. Then,

- (1) the polynomial  $P(t) = t^d + z_1 \in \mathbb{C}[t]$  is post-critically finite and we have  $\mathbf{z} = (P^{\circ j}(0))_{j \geq 1}$ ; in particular, the critical value  $z_1$  of  $P$  is preperiodic of preperiod  $k'$  to a cycle of period  $m'$  of multiplier  $\lambda$ ,

- (2) there exists a partial order  $\preceq$  on  $\mathbb{N} \times \mathbb{N}^*$  such that  $(k', m') \preceq (k, m)$  if and only if  $\mathcal{M}_{k', m'} \subseteq \mathcal{M}_{k, m}$  is a  $F_{k, m}$ -invariant subspace and the restriction of  $F_{k, m}$  to  $\mathcal{M}_{k', m'}$  is  $F_{k', m'}$ ,
- (3) if  $k + m - 1 \geq 2$ ,  $F_{k, m}: \mathcal{M}_{k, m} \rightarrow \mathcal{M}_{k, m}$  and  $G_{k, m}: \mathbb{C}^{k+m-1} \rightarrow \mathbb{C}^{k+m-1}$  are holomorphically conjugate,
- (4)  $\text{Spec } D_z F_{k', m'} \subset \mathbb{C} \setminus \overline{\mathbb{D}}$ ,
- (5) If  $(k, m) \neq (k', m')$ ,

$$\text{Spec } (D_z F_{k, m})^* |_{(\mathcal{M}_{k', m'})^0} = \begin{cases} \{0\} & \text{if } k' = 0 \\ \{\mu \mid \mu^m = \lambda^{\frac{m}{m'}}, \mu^{m'} \neq \lambda\} & \text{if } k' \neq 0 \end{cases} .$$

where  $\mathcal{M}_{k', m'}^0 = \{\omega \in \mathcal{M}_{k, m}^* \mid \omega|_{\mathcal{M}_{k', m'}} \equiv 0\}$  is the annihilator of  $\mathcal{M}_{k', m'}$  in  $\mathcal{M}_{k, m}$

In particular, we shall see that Theorem A' is a direct consequence of Theorem B'. The rest of this article is devoted to the proof of Theorem B'. Item 1 is proved in Proposition 4.4. The partial order  $\preceq$  will be introduced in Definition 3.2 and item 2 will be proved in Proposition 3.3. Item 3 is proved in Lemma 3.1. Item 4 is due to Koch, and we recall its proof in Proposition 4.9 for the sake of completeness. Our main contribution is the proof of item 5 which will be proved in Proposition 4.12. Finally, we prove Theorem A' by using Theorem B'.

### 3. DYNAMICS OF $F_{k, m}$

3.1.  $F_{k, m}$  is conjugate to  $G_{k, m}$ . The following lemma assures that the class of maps  $F_{k, m}$  for  $(k, m) \in \mathbb{N} \times \mathbb{N}^*$  is a good alternative when one wants to study  $G_{k, m}$ .

**Lemma 3.1.** *Let  $(k, m) \in \mathbb{N} \times \mathbb{N}^*$  be such that  $k + m - 1 \geq 2$ , the maps  $F_{k, m}$  and  $G_{k, m}$  are holomorphically conjugate.*

*Proof.* Recall that when  $k = 0, m \geq 3$ , we have

$$G_{0, m} : \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_{m-1} \end{pmatrix} \mapsto \begin{pmatrix} -x_{m-1}^d \\ x_1^d - x_{m-1}^d \\ \vdots \\ x_{m-2}^d - x_{m-1}^d \end{pmatrix} .$$

and when  $k \geq 1, m \geq 2$ , we have

$$G_{k, m} : \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_{k+m-1} \end{pmatrix} \mapsto \begin{pmatrix} \left(-\frac{\beta x_{k+m-1} - x_{k-1}}{\beta - 1}\right)^d \\ \left(x_1 - \frac{\beta x_{k+m-1} - x_{k-1}}{\beta - 1}\right)^d \\ \vdots \\ \left(x_{k+m-2} - \frac{\beta x_{k+m-1} - x_{k-1}}{\beta - 1}\right)^d \end{pmatrix} .$$

Let  $i_{k,m}: \mathcal{M}_{k,m} \rightarrow \mathbb{C}^{k+m-1}$ ,  $i_{k,m}(\mathbf{x}) = (x_1, \dots, x_{k+m-1})$  and set

$$\widetilde{F}_{k,m} = i_{k,m} \circ F_{k,m} \circ i_{k,m}^{-1}.$$

Then we have when  $k = 0$ ,

$$\widetilde{F}_{0,m} : \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_{m-1} \end{pmatrix} \mapsto \begin{pmatrix} -x_{m-1}^d \\ x_1^d - x_{m-1}^d \\ \vdots \\ x_{m-2}^d - x_{m-1}^d \end{pmatrix}.$$

and when  $k \geq 1$ ,

$$\widetilde{F}_{k,m} : \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_{k+m-1} \end{pmatrix} \mapsto \begin{pmatrix} -\frac{\beta x_{k+m-1}^d - x_{k-1}^d}{\beta-1} \\ x_1^d - \frac{\beta x_{k+m-1}^d - x_{k-1}^d}{\beta-1} \\ \vdots \\ x_{k+m-2}^d - \frac{\beta x_{k+m-1}^d - x_{k-1}^d}{\beta-1} \end{pmatrix}.$$

It is enough to show that  $G_{k,m}$  and  $\widetilde{F}_{k,m}$  are conjugate. Indeed, let  $\tau: \mathbb{C}^{k+m-1} \rightarrow \mathbb{C}^{k+m-1}$  be a linear map of the following form

$$\tau \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_{k+m-1} \end{pmatrix} \mapsto \begin{pmatrix} \tau_1 \\ x_1 + \tau_1 \\ \vdots \\ x_{k+m-2} + \tau_1 \end{pmatrix} \text{ with } \tau_1 = \begin{cases} -x_{m-1} & \text{when } k = 0 \\ -\frac{\beta x_{k+m-1} - x_{k-1}}{\beta-1} & \text{when } k \geq 1 \end{cases}$$

and set

$$\mathfrak{d}(x_1, \dots, x_{k+m-1}) = (x_1^d, \dots, x_{k+m-1}^d)$$

then  $G_{k,m} = \mathfrak{d} \circ \tau$ ,  $\widetilde{F}_{k,m} = \tau \circ \mathfrak{d}$ . Thus,

$$\tau \circ G_{k,m} = \widetilde{F}_{k,m} \circ \tau$$

Note that  $\tau$  is an isomorphism, whence  $G_{k,m}$  and  $\widetilde{F}_{k,m}$  are conjugate. □

**3.2. Comparing  $F_{k,m}$  by a partial order  $\preceq$ .** Our initial expectation was that for arbitrary pairs  $(k_1, m_1)$  and  $(k_2, m_2)$  in  $\mathbb{N} \times \mathbb{N}^*$ , the maps  $F_{k_1, m_1}$  and  $F_{k_2, m_2}$  would agree on the intersection  $\mathcal{M}_{k_1, m_1} \cap \mathcal{M}_{k_2, m_2}$ . However this is not true as shown in the following example. Consider the case  $d = 2$  and  $\beta = -1$ , the sequence

$$\mathbf{x} := \{2, 0, 0, \dots, 0, \dots\}$$

Then  $\mathbf{x} \in \mathcal{M}_{2,1} \cap \mathcal{M}_{3,1}$  and

$$\mathcal{Q}(\mathbf{x}) = \{0, 4, 0, 0, 0, 0, \dots\}$$

However,

$$F_{2,1}(\mathbf{x}) = \{-2, 2, -2, -2, -2, -2, \dots\} \text{ and } F_{3,1}(\mathbf{x}) = \{0, 4, 0, 0, 0, 0, \dots\}.$$

We will now see that if some order  $(k', m') \preceq (k, m)$  is satisfied, with the fixed  $d$  and  $\beta$ , we have  $\mathcal{M}_{k', m'} \subseteq \mathcal{M}_{k, m}$  and  $F_{k', m'}$  is the restriction of  $F_{k, m}$  to  $\mathcal{M}_{k', m'}$ .

**Definition 3.2.** Let  $\preceq$  be the partial order on  $\mathbb{N} \times \mathbb{N}^*$  defined by

$$(k', m') \preceq (k, m) \Leftrightarrow \begin{cases} m' \text{ divides } m, \\ \text{either } k' = k \text{ or } (k' = 0 \text{ and } m' \text{ divides } k). \end{cases}$$

The strict order  $\prec$  is defined by

$$(k', m') \prec (k, m) \Leftrightarrow (k', m') \preceq (k, m) \text{ and } (k', m') \neq (k, m).$$

**Proposition 3.3.** For two pairs of integers  $(k, m), (k', m') \in \mathbb{N} \times \mathbb{N}^*$ , we have that

$$(k', m') \preceq (k, m) \Leftrightarrow \mathcal{M}_{k', m'} \subseteq \mathcal{M}_{k, m}$$

Moreover, if  $\mathcal{M}_{k', m'} \subseteq \mathcal{M}_{k, m}$  then  $F_{k, m}|_{\mathcal{M}_{k', m'}} = F_{k', m'}$ .

*Proof.* We first prove that

$$(k, m) \preceq (k', m') \Leftrightarrow \mathcal{M}_{k', m'} \subseteq \mathcal{M}_{k, m}.$$

Assume that  $(k', m') \preceq (k, m)$ . We shall prove that  $\mathcal{M}_{k', m'} \subseteq \mathcal{M}_{k, m}$ . Assume  $\mathbf{x} \in \mathcal{M}_{k', m'}$ . Then  $\mathbf{x}$  is preperiodic of period less than  $k'$  to a cycle of period dividing  $m'$ . Since  $k' \leq k$  and  $m' \mid m$ , we deduce that  $\mathbf{x} \in \mathcal{P}_{k, m}$ . We need to show that  $\beta x_{k+m} - x_k = 0$ . Indeed,

- if  $k' = k$ , since  $m' \mid m$  and  $\mathbf{x} \in \mathcal{M}_{k', m'}$ , we have  $x_{k+m} = x_{k'+m'}$ . Thus  $\beta x_{k+m} - x_k = \beta x_{k'+m'} - x_{k'} = 0$ ,
- if  $k' = 0$ , in that case  $m' \mid k$  and  $\mathbf{x}$  is periodic of period  $m'$ . Thus  $x_{k+m} = x_k = 0$ .

Let us now assume that  $\mathcal{M}_{k', m'} \subseteq \mathcal{M}_{k, m}$ . We claim that  $(k', m') \preceq (k, m)$ . Indeed,

- either  $k' = 0$ ; in this case, consider  $\mathbf{x} \in \mathcal{M}_{0, m'}$  given by  $x_i = 0$  if  $m' \mid i$  and 1 otherwise. Since  $\mathbf{x} \in \mathcal{M}_{k, m}$ ,  $\beta x_{k+m} - x_k = 0$  with  $\beta \neq 1$ . Then necessarily,  $x_{k+m} = x_k = 0$  thus  $m'$  divides  $k$  and  $m$ .
- or  $k' \geq 1$ ; in this case, consider  $\mathbf{x} \in \mathcal{M}_{k', m'}$  given by  $x_{k'} = \beta, x_{k'+jm'} = 1$  for  $j \geq 1$  and  $x_i = 0$  otherwise. If  $\beta x_{k+m} - x_k = 0$  then  $x_k = \beta$  and  $x_{k+m} = 1$ . Hence  $k = k'$  and  $m' \mid m$ .

We assume now  $\mathcal{M}_{k', m'} \subseteq \mathcal{M}_{k, m}$ , or equivalently,  $(k', m') \preceq (k, m)$ . Let us prove that the restriction of  $F_{k, m}$  to  $\mathcal{M}_{k', m'}$  is  $F_{k', m'}$ . Assume  $\mathbf{x} \in \mathcal{M}_{k', m'}$ . Set  $\mathbf{y} = F_{k', m'}(\mathbf{x})$  and  $\mathbf{z} = F_{k, m}(\mathbf{x})$ . According to Lemma 2.3, for all  $i \geq 2$ ,  $y_i = x_{i-1}^d + y_1, z_i = x_{i-1}^d + z_1$  where

$$y_1 = \begin{cases} -x_{m'-1}^d & \text{if } k' = 0 \\ -\frac{\beta x_{k'+m'-1}^d - x_{k'-1}^d}{\beta - 1} & \text{if } k' \geq 1 \end{cases} \quad \text{and} \quad z_1 = \begin{cases} -x_{m-1}^d & \text{if } k = 0 \\ -\frac{\beta x_{k+m-1}^d - x_{k-1}^d}{\beta - 1} & \text{if } k \geq 1. \end{cases}$$

It is enough to prove that  $y_1 = z_1$ .

- Case  $k' = 0$ . In that case,  $k$  and  $m$  are multiples of  $m'$ . If  $k = 0$  then

$$y_1 = -x_{m'-1}^d = -x_{m-1}^d = z_1.$$

If  $k \neq 0$ ,  $x_{k+m-1} = x_{k-1} = x_{m'-1}$  hence

$$y_1 = -x_{m'-1}^d = -\frac{\beta x_{k+m-1}^d - x_{k-1}^d}{\beta - 1} = z_1.$$

• Case  $k' \neq 0$ . In that case,  $k' = k$  and  $m$  is a multiple of  $m'$ . Then

$$x_{k+m-1} = \begin{cases} \frac{1}{\beta} x_{k'+m'-1} & \text{if } m' - 1 = 0 \text{ and } m - 1 \geq 1 \\ x_{k'+m'-1} & \text{otherwise.} \end{cases}$$

<sup>1</sup> Then

$$y_1 = -\frac{\beta x_{k'+m'-1}^d - x_{k'-1}^d}{\beta - 1} = -\frac{\beta x_{k+m-1}^d - x_{k-1}^d}{\beta - 1} = z_1. \quad \square$$

**3.3. The post-critical set of  $F_{k,m}$ .** In this section, we fix a pair of integers  $(k, m) \in \mathbb{N} \times \mathbb{N}^*$ . Recall that

$$C(F_{k,m}) := \text{the critical set of } F_{k,m}: \mathcal{M}_{k,m} \rightarrow \mathcal{M}_{k,m},$$

$$CV(F_{k,m}) := \text{the critical value set of } F_{k,m}: \mathcal{M}_{k,m} \rightarrow \mathcal{M}_{k,m}$$

and

$$PC(F_{k,m}) := \text{the post-critical set of } F_{k,m}: \mathcal{M}_{k,m} \rightarrow \mathcal{M}_{k,m}.$$

**Lemma 3.4.** *We have that  $C(F_{k,m}) = \{\mathbf{x} \in \mathcal{M}_{k,m} \mid x_i = 0 \text{ for some } 1 \leq i \leq k+m-1\}$*

*Proof.* Recall that  $F_{k,m} = \pi_{k,m} \circ \mathcal{Q}$ . Differentiating both sides, we see that for any  $\mathbf{x} \in \mathcal{M}_{k,m}$  and for any  $\mathbf{v} \in T_{\mathbf{x}}\mathcal{E} = \mathcal{E}$ ,

$$D_{\mathbf{x}}F_{k,m}(\mathbf{v}) = \pi_{k,m} \circ D_{\mathbf{x}}\mathcal{Q}(\mathbf{v}) = \pi_{k,m}(0, dx_1^{d-1}v_1, dx_2^{d-1}v_2, \dots)$$

On the one hand, assume  $\mathbf{x} \in C(F_{k,m})$ . Then there exists  $\mathbf{v} \in T_{\mathbf{x}}\mathcal{M}_{k,m} \setminus \{\mathbf{0}\}$  such that  $x_i^{d-1}v_i = 0$  for all  $i \geq 1$ . Observe that there exists  $i \in \{1, \dots, k+m-1\}$  such that  $v_i \neq 0$  whence  $x_i = 0$ . Indeed otherwise,  $v_{k+m} = \frac{1}{\beta}v_k = 0$  and by preperiodicity,  $v_i = 0$  for all  $i \geq 1$ .

On the other hand, given  $\mathbf{x} \in \mathcal{E}$ , define  $\mathbf{v} \in \mathcal{E}$  by

$$v_j = \begin{cases} \beta & \text{if } x_j = 0 \text{ and } j = k \\ 1 & \text{if } x_j = 0 \text{ and } j \neq k \\ 0 & \text{if } x_j \neq 0 \end{cases}$$

Then  $x_j^{d-1}v_j = 0$  for all  $j \geq 1$  so that  $D_{\mathbf{x}}F_{k,m}(\mathbf{v}) = \mathbf{0}$ . Moreover, if  $\mathbf{x} \in \mathcal{M}_{k,m}$  then  $\mathbf{v} \in \mathcal{M}_{k,m}$ . Finally, if there exists  $i \in \{1, \dots, k+m-1\}$  such that  $x_i = 0$  then  $\mathbf{v} \neq \mathbf{0}$  whence  $\mathbf{x} \in C(F_{k,m})$ .  $\square$

**Definition 3.5.** Denote by

$$\Delta_{k,m} = \{\mathbf{x} \in \mathcal{M}_{k,m} \mid \text{there exists } 1 \leq i < j \leq k+m \text{ such that } x_i = x_j\}$$

The set  $\Delta_{k,m}$  consists of  $\binom{k+m}{2}$  hyperplanes.

<sup>1</sup>Note that if  $m' = 1$ ,  $\mathbf{x} \in \mathcal{M}_{k',m'}$  implies that for all  $i \geq 1$ ,  $x_{k'+i} = x_{k'+1}$ . In particular, with  $k = k'$ ,  $x_{k+m-1} = x_{k'+1} = \frac{1}{\beta}x_{k'}$

**Proposition 3.6.** *We have that  $CV(F_{k,m}) \subseteq \Delta_{k,m}$  and  $F_{k,m}(\Delta_{k,m}) \subseteq \Delta_{k,m}$ . Consequently,  $PC(F_{k,m}) \subseteq \Delta_{k,m}$ .*

*Proof.* Let  $\mathbf{x} \in C(F_{k,m})$  and set  $\mathbf{y} = F_{k,m}(\mathbf{x})$ . Then by Lemma 3.4, there exists  $i \in \{1, \dots, k+m-1\}$  such that  $x_i = 0$ . By Lemma 2.3, we have

$$y_{i+1} = x_i^d + y_1 = y_1$$

Thus  $\mathbf{y} \in \Delta_{k,m}$ , whence  $CV(F_{k,m}) \subseteq \Delta_{k,m}$ .

Now we prove that  $\Delta_{k,m}$  is invariant under  $F_{k,m}$ . Assume  $\mathbf{x} \in \Delta_{k,m}$  and set  $\mathbf{y} = F_{k,m}(\mathbf{x})$ . Then there exist  $1 \leq i < j \leq k+m$  such that  $x_i = x_j$ . By Lemma 2.3, for every  $l \geq 2$ ,  $y_l = x_{l-1}^d + y_1$ . Note that since  $\mathbf{x} \in \mathcal{M}_{k,m}$ ,  $\beta x_{k+m} - x_k = 0$  with the convention  $x_0 := 0$ .

- If  $j \leq k+m-1$ , we have  $y_{i+1} = x_i^d + y_1 = x_j^d + y_1 = y_{j+1}$ .
- If  $j = k+m$ , then
  - either  $i = k$  so that  $x_k = x_i = x_j = x_{k+m}$ ; since  $\beta x_{k+m} - x_k = 0$  and  $\beta \neq 1$ ,  $x_k = 0$  whence  $\mathbf{x} \in C(F_{k,m})$  and  $\mathbf{y} \in \Delta_{k,m}$ ;
  - or  $i \neq k$  so that  $i+1 \neq k+1$ ; since  $x_k = \beta x_{k+m} = \beta x_i$ , we have

$$y_{i+1} = x_i^d + y_1 = x_k^d + y_1 = y_{k+1}.$$

Hence, in any case, we have  $\mathbf{y} \in \Delta_{k,m}$ , i.e.  $F_{k,m}(\Delta_{k,m}) \subseteq \Delta_{k,m}$  and the lemma is proved.  $\square$

#### 4. FIXED POINTS OF KOCH MAPS

In this section, we shall study the eigenvalues of the derivative of  $F_{k,m}$  at its fixed points and we will prove Theorem B'. Then, we deduce Theorem A' by using Theorem B'.

**4.1. Relation with post-critically finite polynomials.** There is a close connection between fixed points  $F_{k,m}$  and post-critically finite polynomials. More precisely, we will consider monic centered unicritical polynomials of degree  $d \geq 2$ ,

$$P(t) = t^d + c \in \mathbb{C}[t], c \in \mathbb{C}$$

The critical orbit of such a polynomial is the sequence  $\mathbf{c}_P \in \mathcal{E}$  defined by

$$\mathbf{c}_P = (c_i)_{i \geq 1} \in \mathcal{E} \text{ where } c_i = P^{o_i}(0).$$

Since the preperiod and the period of a preperiodic sequence will be extensively discussed in this chapter, we introduce the following notions.

**Definition 4.1.** Given integers  $k \geq 0, m \geq 1$ , a sequence  $\mathbf{x} \in \mathcal{E}$  is called *preperiodic of type  $(k, m)$*  if for every  $i \geq k+1$ ,  $x_{i+m} = x_i$ , *preperiodic of exact type  $(k, m)$*  if, additionally,  $k$  and  $m$  are the smallest integers satisfying such conditions.

For a sequence of exact type  $(k, m)$ , the pair  $(k, m)$  consists of the preperiod  $k$  and the period  $m$ . The vector space  $\mathcal{P}_{k,m}$  is the space of preperiodic sequences of type  $(k, m)$ .

**Definition 4.2.** A degree  $d$  polynomial of (exact) type  $(k, m)$  is a monic centered unicritical polynomial  $P$  of degree  $d \geq 2$  whose critical orbit  $\mathbf{c}_P$  is of (exact) type  $(k, m)$ .

In other words, a polynomial is of type  $(k, m)$  if and only if its critical orbit belongs to  $\mathcal{P}_{k,m}$ . Note that a polynomial of type  $(k, m)$  is post-critically finite.

*Remark 4.3.* Let  $P$  be a polynomial of type  $(k, m)$  of degree  $d$ . If  $k = 0$ , then the critical value  $c$  of  $P$  is a periodic point of period dividing  $m$ , i.e.  $P^{\circ m}(c) = c$ . In other words,  $P^{\circ(m-1)}(c) \in P^{-1}(c)$ . However, since  $P$  is a unicritical polynomial,  $P^{-1}(c)$  consists of exactly one point which is the critical point of  $P$ . This means that  $P^{\circ(m-1)}(c)$  is in fact the critical point of  $P$ . This is the case if and only if the critical point of  $P$  is also a periodic point of type  $(0, m)$ .

**Proposition 4.4.** Given  $(k, m) \in \mathbb{N} \times \mathbb{N}^*$ . Let  $\mathbf{z} \in \mathcal{M}_{k,m}$  be a fixed point of  $F_{k,m}$ . Set  $P(t) = t^d + z_1$ . Then  $\mathbf{z} = \mathbf{c}_P$  and  $P$  is of exact type  $(k', m')$ . Moreover,  $(k', m') \preceq (k, m)$ .

*Proof.* First, let us prove that  $P$  is of type  $(k, m)$ . According to Lemma 2.3, for every  $i \geq 2$ , we have that  $z_i = z_{i-1}^d + z_1$  hence

$$z_i = P(z_{i-1}).$$

In other words,  $\mathbf{z}$  is the sequence of iterates of  $z_1$  under  $P$ . Recall that by Lemma 2.6,  $\mathcal{M}_{k,m}$  contains every fixed point of  $F_{k,m}$  hence  $\mathbf{z} \in \mathcal{M}_{k,m} \subset \mathcal{P}_{k,m}$ . Therefore,  $\mathbf{z} = \mathbf{c}_P$  and the polynomial  $P$  is a polynomial of type  $(k, m)$ .

Second, let  $(k', m')$  be the exact type of  $P$ . We prove that  $(k', m') \preceq (k, m)$ , i.e.

$$\begin{cases} m' \mid m \\ \text{either } k' = k \text{ or } (k' = 0 \text{ and } m' \mid k) \end{cases}$$

Since  $(k', m')$  is the exact type of the orbit of  $z_1$ ,  $k' \leq k$  and  $m' \mid m$ . If  $k' = k$ , we are done. If  $k' \neq k$ , we need to prove that  $k' = 0$  and  $m' \mid k$ . Since  $(k', m')$  is the exact type of  $\mathbf{z}$ , we have  $\mathbf{z} \in \mathcal{P}_{k',m'}$ . Thus,  $k' + 1 \leq k$  and  $m' \mid m$ ; and since  $\mathbf{z} \in \mathcal{P}_{k',m'}$ , this implies that  $z_{k+m} = z_k$ . Moreover,  $\mathbf{z} \in \mathcal{M}_{k,m}$  implies that  $\beta z_{k+m} - z_k = 0$ . Therefore,  $P^{\circ k}(0) = z_k = 0$ . In other words, 0 is a periodic point of  $P$ , i.e.  $k' = 0$ , and the period of 0 is  $m'$ . Moreover,  $P^{\circ k}(0) = 0$  also implies that  $k$  is a multiple of  $m'$ . Thus, we can conclude that  $(k', m') \preceq (k, m)$ .  $\square$

*Remark 4.5.* The converse statement of Proposition 4.4 is true under some assumptions on the choice of the root of unity  $\beta$ . More precisely, given a post-critically finite unicritical polynomial  $P(t) = t^d + z_1$  of type  $(k, m)$ , there exists a  $d$ -th root of unity  $\beta' \neq 1$  such that the critical orbit  $\mathbf{c}_P$  of  $P$  is a fixed point of  $F_{k,m}$ .

The partial order  $\preceq$  enables us to study the relative positions of the fixed points of  $F_{k,m}$  and  $\Delta_{k,m}$ .

**Lemma 4.6.** Let  $\mathbf{z}$  be a fixed point of  $F_{k,m}$  and let  $(k', m')$  be the exact type of  $\mathbf{z}$ . Then,  $\mathbf{z} \in \Delta_{k,m}$  if and only if  $(k', m') \prec (k, m)$ .

*Proof.* Assume  $\mathbf{z} \in \Delta_{k,m}$ , i.e. there exists  $1 \leq i < j \leq k+m$  such that  $z_i = z_j$ . In particular,  $\mathbf{z}$  is a preperiodic sequence of preperiod at most  $i-1$  and of period dividing  $j-i$ . Whence, since  $(k', m')$  is the exact type of  $\mathbf{z}$ , we have  $k' \leq i-1$  and  $m'$  divides  $j-i$ .

- If  $i \leq k$  then  $k' \leq i-1 < k$ .
- If  $i \geq k+1$  then  $j-i \leq k+m-(k+1) < m$ . Since  $m' \mid j-i$ , we have  $m' < m$ .

In both cases, we have  $(k', m') \neq (k, m)$ . Note that, according to Proposition 4.4,  $(k', m) \preceq (k, m)$ . Hence  $(k', m') \prec (k, m)$ .

Conversely, assume  $(k', m') \prec (k, m)$ . In particular,  $k' \leq k$ ,  $m' \leq m$  and  $(k', m') \neq (k, m)$ . Note that  $\mathbf{z}$  is of exact type  $(k', m')$ . Hence,

$$z_{k'+1} = z_{k'+m'+1}.$$

If  $k' \neq k$  then  $k' < k$ . Whence  $k'+1$  and  $k'+m'+1$  are integers in  $\{1, \dots, k+m\}$ . If  $k' = k$  then  $m' < m$ . In this case,  $k+1, k+m'+1$  are also in  $\{1, \dots, k+m\}$ . Therefore, in both cases, we deduce by that  $\mathbf{z} \in \Delta_{k,m}$ .  $\square$

**4.2. Eigenvalues of moduli maps at fixed points.** In order to study the eigenvalues of the derivative of moduli maps at one of its fixed point, we will in fact study its transpose. Note that when  $k+m=1$ ,  $\mathcal{M}_{k,m} = \{\mathbf{0}\}$  and  $F_{k,m}$  is trivial. Let us fix  $(k, m) \in \mathbb{N} \times \mathbb{N}^*$  such that  $k+m \geq 2$ . Assume that  $\mathbf{z} \in \mathcal{M}_{k,m}$  is a fixed point of  $F_{k,m}$ . We will describe the transpose of the derivative  $D_{\mathbf{z}}F_{k,m}: T_{\mathbf{z}}\mathcal{M}_{k,m} \rightarrow T_{\mathbf{z}}\mathcal{M}_{k,m}$ . Since  $\mathcal{M}_{k,m}$  is a vector space, there is a canonical identification of  $T_{\mathbf{z}}\mathcal{M}_{k,m}$  with  $\mathcal{M}_{k,m}$ , the derivative  $D_{\mathbf{z}}F_{k,m}: T_{\mathbf{z}}\mathcal{M}_{k,m} \rightarrow T_{\mathbf{z}}\mathcal{M}_{k,m}$  identifies with a linear map

$$L: \mathcal{M}_{k,m} \rightarrow \mathcal{M}_{k,m},$$

and the transpose identifies with the pull-back map of  $L$

$$L^*: \mathcal{M}_{k,m}^* \rightarrow \mathcal{M}_{k,m}^*.$$

4.2.1. *The dual space  $\mathcal{M}_{k,m}^*$ .* For  $i \geq 1$ , let  $\omega_i \in \mathcal{M}_{k,m}^*$  be the linear form defined by for all  $\mathbf{v} \in \mathcal{M}_{k,m}$ ,

$$\omega_i(\mathbf{v}) := v_i.$$

**Lemma 4.7.** *The family  $\{\omega_i, 1 \leq i \leq k+m-1\}$  is a basis of  $\mathcal{M}_{k,m}^*$ .*

*Proof.* Note that  $\dim \mathcal{M}_{k,m} = k+m-1$  hence it is enough to prove that  $\{\omega_1, \dots, \omega_{k+m-1}\}$  are linearly independent. Assume that

$$\sum_{1 \leq i \leq k+m-1} \lambda_i \omega_i = 0 \text{ with } \lambda_i \in \mathbb{C}.$$

Let  $i \geq 1$ . To prove that  $\lambda_i = 0$ , consider the vector  $\mathbf{v} \in \mathcal{M}_{k,m}$  defined by

- if  $i < k$ ,  $v_j = \begin{cases} 1 & \text{if } j = i \\ 0 & \text{otherwise,} \end{cases}$
- if  $i = k$ ,  $v_j = \begin{cases} 1 & \text{if } j = i = k \\ \frac{1}{\beta} & \text{if } j > k \text{ and } j \equiv k \pmod{m} \\ 0 & \text{otherwise,} \end{cases}$
- if  $i > k$ ,  $v_j = \begin{cases} 1 & \text{if } j \geq i \text{ and } j \equiv i \pmod{m} \\ 0 & \text{otherwise.} \end{cases}$

In any case, we have

$$0 = \sum_{1 \leq i \leq k+m-1} \lambda_i \omega_i(\mathbf{v}) = \lambda_i v_i = \lambda_i. \quad \square$$

4.2.2. *The transpose of the derivative.* Observe that  $L^*: \mathcal{M}_{k,m}^* \rightarrow \mathcal{M}_{k,m}^*$  is the pull-back of forms, i.e. for all  $\omega \in \mathcal{M}_{k,m}^*$ ,

$$L^* \omega = \omega \circ L.$$

For all  $i \geq 1$ , set

$$\delta_i = dz_i^{d-1}$$

where  $\mathbf{z} = (z_1, z_2, \dots) \in \mathcal{M}_{k,m}$  is the considered fixed point of  $F_{k,m}$ .

**Lemma 4.8.** *We have that*

$$L^* \omega_1 = \begin{cases} -\delta_{m-1} \omega_{m-1} & \text{if } k = 0 \\ -\frac{\beta \delta_{k+m-1} \omega_{k+m-1} - \delta_{k-1} \omega_{k-1}}{\beta-1} & \text{otherwise,} \end{cases}$$

and for all  $i \geq 2$ ,

$$L^* \omega_i = \delta_{i-1} \omega_{i-1} + L^* \omega_1.$$

*Proof.* Recall that for all  $\mathbf{v} \in \mathcal{M}_{k,m}$ ,

$$L(\mathbf{v}) = \pi_{k,m} \circ D_{\mathbf{z}} \mathcal{Q}(\mathbf{v}).$$

Set  $\mathbf{u} = D_{\mathbf{z}} \mathcal{Q}(\mathbf{v})$  then

$$u_1 = 0 \quad \text{and for all } i \geq 2, \quad u_i = dz_{i-1}^{d-1} v_{i-1} = \delta_{i-1} v_{i-1}.$$

In addition, if  $\mathbf{w} := L(\mathbf{v}) = \pi_{k,m}(\mathbf{u})$  then

$$\text{for all } i \geq 2, \quad w_i = u_i + w_1 \text{ with } w_1 = \begin{cases} -u_m & \text{if } k = 0 \\ -\frac{\beta u_{k+m-1} - u_{k-1}}{\beta-1} & \text{otherwise,} \end{cases}$$

Combining those formulas, we obtain that  $\mathbf{w} = L(\mathbf{v})$  and for all  $i \geq 2$ ,

$$(4.1) \quad w_i = \delta_{i-1} v_{i-1} + w_1 \text{ with } w_1 = \begin{cases} -\delta_{m-1} v_{m-1} & \text{if } k = 0 \\ -\frac{\beta \delta_{k+m-1} v_{k+m-1} - \delta_{k-1} v_{k-1}}{\beta-1} & \text{otherwise,} \end{cases}$$

We deduce that for all  $\mathbf{v} \in \mathcal{M}_{k,m}$ ,

$$L^*\omega_1(\mathbf{v}) = \omega_1 \circ L(\mathbf{v}) = w_1 = \begin{cases} -\delta_{m-1}v_{m-1} & \text{if } k = 0 \\ -\frac{\beta\delta_{k+m-1}\omega_{k+m-1} - \delta_{k-1}\omega_{k-1}}{\beta-1} & \text{otherwise,} \end{cases}$$

hence

$$L^*\omega_1 = \begin{cases} -\delta_{m-1}\omega_{m-1} & \text{if } k = 0 \\ -\frac{\beta\delta_{k+m-1}\omega_{k+m-1} - \delta_{k-1}\omega_{k-1}}{\beta-1} & \text{otherwise.} \end{cases}$$

In addition, for all  $i \geq 2$  and for all  $\mathbf{v} \in \mathcal{M}_{k,m}$ , we have

$$L^*\omega_i(\mathbf{v}) = \omega_i \circ L(\mathbf{v}) = \delta_{i-1}v_{i-1} + w_1 = \delta_{i-1}v_{i-1} + \omega_1 \circ L(\mathbf{v}),$$

hence

$$L^*\omega_i = \delta_{i-1}\omega_{i-1} + L^*\omega_1. \quad \square$$

**4.2.3. Fixed points outside the post-critical set.** According to Section 3.1, the map  $F_{k,m}$  is conjugate to the map  $G_{k,m}$  constructed by Koch [Koc13]. By [Koc13, Corollary 7.2], the derivative of  $G_{k,m}$  at its fixed points outside the post-critical set has only eigenvalues of modulus strictly greater than 1, whence so does  $F_{k,m}$ . For the sake of completeness, we give here the proof of this property. For further discussion about the arithmetics of such eigenvalues, we refer to [BEK20]. The main content of this paragraph is the following result.

**Proposition 4.9.** *Let  $(k, m) \in \mathbb{N} \times \mathbb{N}^*$  and  $\mathbf{z} \notin PC(F_{k,m})$  be a fixed point of the moduli map  $F_{k,m}$ . Then every eigenvalue of  $D_{\mathbf{z}}F_{k,m}$  has modulus strictly greater than 1.*

*Proof.* Since  $\mathcal{M}_{k,m}$  has finite dimension, it is suffice to prove that every eigenvalue of the transpose  $L^*$  of  $D_{\mathbf{z}}F_{k,m}$  has modulus strictly bigger than 1.

Recall that, by Lemma 4.7, the family  $\{\omega_i: \mathcal{M}_{k,m} \rightarrow \mathbb{C}\}_{i \in \{1, \dots, k+m-1\}}$  is a basis of  $\mathcal{M}_{k,m}^*$ . According to Lemma 4.8, setting  $\delta_i = dz_i^{d-1}$ , we have

$$L^*\omega_1 = \begin{cases} -\delta_{m-1}\omega_{m-1} & \text{if } k = 0 \\ -\frac{\beta\delta_{k+m-1}\omega_{k+m-1} - \delta_{k-1}\omega_{k-1}}{\beta-1} & \text{if } k \geq 1, \end{cases}$$

and for all  $i \geq 2$ ,

$$L^*\omega_i = \delta_{i-1}\omega_{i-1} + L^*\omega_1.$$

According to Lemma 4.6, the point  $\mathbf{z}$  is a fixed point of  $F_{k,m}$  of exact type  $(k, m)$ . Therefore,  $\delta_i \neq 0$  for all  $i \in \{1, \dots, k+m-1\}$ . Indeed, note that according to Proposition 4.4, the sequence  $\mathbf{z}$  is the critical orbit of  $P(t) = t^d + z_1$ . Assume that  $dz_i^{d-1} = \delta_i = 0$  for some  $i \in \{1, \dots, k+m-1\}$ . Then  $P^{\circ i}(0) = z_i = 0$ . This implies that  $k = 0$  and  $m$  divides  $i$ . However  $i \leq k+m-1 = m-1 < m$ , hence contradiction.

We may therefore define a linear map  $L_*: \mathcal{M}_{k,m}^* \rightarrow \mathcal{M}_{k,m}^*$  by

$$(4.2) \quad \forall i \in \{1, \dots, k+m-1\} \quad L_*(\omega_i) = \frac{\omega_{i+1} - \omega_1}{\delta_i}.$$

**Lemma 4.10.** *The linear map  $L^*$  is invertible and its inverse is  $L_*$ .*

*Proof.* We need to prove that  $L_* \circ L^* = L^* \circ L_* = \text{id}$ . First, observe that for all  $i \in \{1, \dots, k+m-1\}$ ,

$$\begin{aligned} L^* \circ L_*(\omega_i) &= L^* \left( \frac{\omega_{i+1} - \omega_1}{\delta_i} \right) = \frac{1}{\delta_i} (L^*(\omega_{i+1}) - L^*(\omega_1)) \\ &= \frac{1}{\delta_i} (\delta_i \omega_i + L^*(\omega_1) - L^*(\omega_1)) = \omega_i. \end{aligned}$$

Second, we prove  $L_* \circ L^* = \text{Id}_{\mathcal{M}_{k,m}^*}$ . Note that, by the definition of  $\mathcal{M}_{k,m}$ , we have that

$$\omega_{k+m} = \begin{cases} 0 & \text{if } k = 0 \\ \frac{1}{\beta} \omega_k & \text{if } k \geq 1 \end{cases} \quad \text{and} \quad \forall i \geq k+m+1 \quad \omega_{i+m} = \omega_i.$$

To compute  $L_* \circ L^*(\omega_1)$ , observe that if  $k = 0$ , then

$$L_* \circ L^*(\omega_1) = L_*(-\delta_{m-1} \omega_{m-1}) = -\delta_{m-1} L_*(\omega_{m-1}) = -(\omega_m - \omega_1) = \omega_1$$

and if  $k \geq 1$ , then

$$\begin{aligned} L_* \circ L^*(\omega_1) &= L_* \left( -\frac{\beta \delta_{k+m-1} \omega_{k+m-1} - \delta_{k-1} \omega_{k-1}}{\beta - 1} \right) \\ &= - \left( \frac{\beta}{\beta - 1} (\omega_{k+m} - \omega_1) - \frac{1}{\beta - 1} (\omega_k - \omega_1) \right) = \omega_1. \end{aligned}$$

In both cases,  $L_* \circ L^*(\omega_1) = \omega_1$ . For  $L_* \circ L^*(\omega_i)$  with  $i \in \{2, \dots, k+m-1\}$ ,

$$L_* \circ L^*(\omega_i) = L_*(\delta_{i-1} \omega_{i-1} + L^*(\omega_1)) = \delta_{i-1} \frac{\omega_i - \omega_1}{\delta_{i-1}} + L_* L^*(\omega_1) = \omega_i.$$

Thus, the linear map  $L_*: \mathcal{M}_{k,m}^* \rightarrow \mathcal{M}_{k,m}^*$  is indeed the inverse of  $L^*$  □

In order to prove Proposition 4.9, it is therefore enough to prove that every eigenvalue of  $L_*: \mathcal{M}_{k,m}^* \rightarrow \mathcal{M}_{k,m}^*$  is contained in the open unit disc  $\mathbb{D}$ . Inspired by the proof of [Koc13, Corollary 7.2], we will show that  $L_*$  is conjugate to a linear transformation on a space of meromorphic quadratic differentials on  $\mathbb{C}$ , whose eigenvalues are all contained in  $\mathbb{D}$ .

Consider the quadratic polynomial  $P(t) := t^d + z_1$ , so that  $z_i = P^{oi}(0)$  for all  $i \geq 1$ . Following Milnor [Mil14], denote by  $\mathfrak{Q}(\mathbb{C})$  the space of meromorphic quadratic differentials on  $\mathbb{C}$  which have at worst simple poles and let us use the notation  $Q \in \mathfrak{Q}(\mathbb{C})$  with

$$Q = q(t) dt^2.$$

and  $q(t)$  is a meromorphic function. Let  $U \subset \mathbb{C}$  be a sufficiently large disk so that  $P^{-1}(U)$  is compactly contained in  $U$  and for  $Q \in \mathfrak{Q}(\mathbb{C})$ , consider the norm

$$\|Q\|_U := \iint_U |q(t) dt^2|.$$

The pushforward of  $Q$  by  $P$  is the quadratic differential  $P_*Q \in \mathfrak{Q}(\mathbb{C})$  defined by

$$P_*Q := \sum_{P(u)=t} \frac{q(u)}{(P'(u))^2} dt^2.$$

It follows from the triangle inequality that

$$\|P_*Q\|_U \leq \|Q\|_{P^{-1}(U)} < \|Q\|_U.$$

For  $i \geq 1$ , let  $Q_i \in \mathfrak{Q}(\mathbb{C})$  be the quadratic differential defined by

$$Q_i := \frac{dt^2}{t - z_i}.$$

The following lemma generalizes a result due to Milnor, [Mil14, Lemma 1] in the case  $d = 2$ .

**Lemma 4.11.** *For all  $i \in \{1, \dots, k + m - 1\}$ ,*

$$(4.3) \quad P_*Q_i = \frac{Q_{i+1} - Q_1}{\delta_i}.$$

*Proof of Lemma 4.11.* Set  $\xi := e^{\frac{2\pi i}{d}}$ . Observe that for a given  $z \in \mathbb{C}$  and for a given  $w \in \mathbb{C}$  such that  $P(w) = z$ , we have  $\{P(u) = z\} = \{w, \xi w, \dots, \xi^{d-1}w\}$ . Thus, for a given  $i \in \{1, \dots, k + m - 1\}$ ,

$$\begin{aligned} P_*Q_i(z) &= \sum_{P(u)=z} \frac{1}{u - z_i} \frac{1}{(P'(u))^2} dt^2 \\ &= \sum_{j=0}^{d-1} \frac{1}{\xi^j w - z_i} \frac{1}{(d(\xi^j w)^{d-1})^2} dt^2 \\ &= \frac{1}{d^2 w^{2d-2}} \left( \sum_{j=0}^{d-1} \frac{1}{\xi^{-j} w - \xi^{-2j} z_i} \right) dt^2. \end{aligned}$$

Note that  $\sum_{j=0}^{d-1} \frac{1}{\xi^{-j} w - \xi^{-2j} z_i} = \frac{dz_i w^{d-2}}{w^d - z_i^d}$ <sup>2</sup>. Therefore,

$$\begin{aligned} P_*Q_i(z) &= \frac{z_i}{dw^d(w^d - z_i^d)} dt^2 \\ &= \frac{1}{\delta_i} \frac{z_i^d}{w^d(w^d - z_i^d)} dt^2 \end{aligned}$$

Since  $w^d = z - z_1$ ,  $z_i^d = z_{i+1} - z_1$ , we have

$$P_*Q_i(z) = \frac{1}{\delta_i} \frac{z_{i+1} - z_1}{(z - z_1)(z - z_{i+1})} dt^2 = \frac{1}{\delta_i} \left( \frac{1}{z - z_{i+1}} - \frac{1}{z - z_1} \right) dt^2.$$

---

<sup>2</sup>This equality is equivalent to the equality  $\sum_{j=0}^{d-1} \frac{1}{\xi^{-j} \chi - \xi^{-2j}} = \frac{d\chi^{d-2}}{\chi^d - 1}$ . The later follows from an elementary computation by comparing the partial fraction decomposition.

$$\text{Thus, } P_i Q_i = \frac{Q_{i+1} - Q_i}{\delta_i} \quad \square$$

The quadratic differentials  $(Q_i)_{1 \leq i \leq k+m-1}$  span a vector space  $\mathcal{Q}_P \subset \mathfrak{Q}(\mathbb{C})$  of dimension  $k+m-1$ . According to Equation (4.3), this subspace is invariant by  $P_*$ . According to Equations (4.2) and (4.3), the linear map  $\iota : \mathcal{Q}_P \rightarrow \mathcal{M}_{k,m}$  which sends  $Q_i \in \mathfrak{Q}(\mathbb{C})$  to  $\omega_i \in \mathcal{M}_{k,m}$  is an isomorphism which conjugates  $P_* : \mathcal{Q}_P \rightarrow \mathcal{Q}_P$  to  $L_* : \mathcal{M}_{k,m}^* \rightarrow \mathcal{M}_{k,m}^*$ .

Since  $\|P_* Q\|_U < \|Q\|_U$  for all  $Q \in \mathcal{Q}_P$ , the spectrum of  $P_* : \mathcal{Q}_P \rightarrow \mathcal{Q}_P$  is contained in the unit disk. It follows that the spectrum of  $L_* : \mathcal{M}_{k,m}^* \rightarrow \mathcal{M}_{k,m}^*$  is contained in the unit disk as required.  $\square$

4.2.4. *Fixed points inside the post-critical set.* We will now study the derivatives of moduli maps at fixed points which are inside the post-critical set. Let  $\mathbf{z} \in PC(F_{k,m})$  be a fixed point of  $F_{k,m}$  and let  $(k', m')$  be the exact type of  $\mathbf{z}$ .

According to Lemma 4.6,  $(k', m') \prec (k, m)$  and, by Proposition 3.3,  $\mathcal{M}_{k',m'} \subsetneq \mathcal{M}_{k,m}$  is invariant under  $F_{k,m}$ . Since  $\mathcal{M}_{k',m'}$  is invariant under  $D_{\mathbf{z}} F_{k,m}$ , the vector space

$$\mathcal{M}_{k',m'}^0 = \{\omega \in \mathcal{M}_{k,m}^* \mid \omega|_{\mathcal{M}_{k',m'}} \equiv 0\},$$

which is called *the annihilator of  $\mathcal{M}_{k',m'}$*  in  $\mathcal{M}_{k,m}$ , is invariant under the transpose  $L^*$  of  $D_{\mathbf{z}} F_{k,m}$  and we have the following decomposition

$$(4.4) \quad \text{Spec } L = \text{Spec}(L|_{\mathcal{M}_{k',m'}}) \cup \text{Spec}\left(L^*|_{\mathcal{M}_{k',m'}^0}\right).$$

Moreover, according to Proposition 3.3, we have

$$L|_{\mathcal{M}_{k',m'}} = D_{\mathbf{z}} F_{k',m'}.$$

Whence, by Proposition 4.9,  $L|_{\mathcal{M}_{k',m'}}$  has only eigenvalues of modulus strictly greater than 1. In order to describe  $\text{Spec } L$ , we need to study  $\text{Spec}\left(L^*|_{\mathcal{M}_{k',m'}^0}\right)$ . We will prove the following result.

**Proposition 4.12.** *Let  $(k, m) \in \mathbb{N} \times \mathbb{N}^*$  and  $\mathbf{z} \in PC(F_{k,m})$  be a fixed point of  $F_{k,m}$  of exact type  $(k', m') \prec (k, m)$ . Let  $\lambda$  be the multiplier of the polynomial  $P(t) = t^d + z_1 \in \mathbb{C}[t]$  along the cycle of  $P^{\circ k'}(z_1)$ . Then*

$$\text{Spec}\left((D_{\mathbf{z}} F_{k,m})^*|_{\mathcal{M}_{k',m'}^0}\right) = \begin{cases} \{0\} & \text{if } k' = 0 \\ \{\mu \mid \mu^m = \lambda^{\frac{m}{m'}}, \mu^{m'} \neq \lambda\} & \text{if } k' \neq 0. \end{cases}$$

The rest of this section is devoted to the proof of this proposition. To simplify the notation, we denote by  $L^*$  the restriction of  $(D_{\mathbf{z}} F_{k,m})^*$  to  $\mathcal{M}_{k',m'}^0$ . The study of the transpose  $L^* : \mathcal{M}_{k',m'}^0 \rightarrow \mathcal{M}_{k',m'}^0$  is divided into two cases,  $k' = 0$  and  $k' \neq 0$ , and each case will be treated separately.

*Proof of Proposition 4.12 when  $k' = 0$ .* Since  $(0, m') \preceq (k, m)$ ,  $m'$  divides  $k$  and  $m$ . It is enough to prove that  $L^* : \mathcal{M}_{0, m'}^0 \rightarrow \mathcal{M}_{0, m'}^0$  is nilpotent. Recall that for  $i \geq 1$ , the form  $\omega_i : \mathcal{M}_{k, m} \rightarrow \mathbb{C}$  is defined by  $\omega_i(\mathbf{v}) = v_i$ . For  $i \geq 1$ , set  $\alpha_i : \mathcal{M}_{k, m} \rightarrow \mathbb{C}$  defined by

$$\alpha_i = \omega_i - \omega_{i+m'}.$$

Recall that  $\mathcal{M}_{0, m'}^0 = \{\omega : \mathcal{M}_{k, m} \rightarrow \mathbb{C} \text{ such that } \omega|_{\mathcal{M}_{0, m'}} \equiv 0\}$ .

**Lemma 4.13.** *We have  $\mathcal{M}_{0, m'}^0 = \text{Span} \{\alpha_i, 1 \leq i \leq k + m\}$ .*

*Proof.* By duality, it is equivalent to show that

$$\mathcal{M}_{0, m'} = \bigcap_{1 \leq i \leq k+m} \text{Ker } \alpha_i.$$

Assume  $\mathbf{v} \in \mathcal{M}_{0, m'}$ . Then for all  $j \geq 1, v_j = v_{j+m'}$ . Given  $i \in \{1, \dots, k + m\}$ , we have

$$\alpha_i(\mathbf{v}) = \omega_i(\mathbf{v}) - \omega_{i+m'}(\mathbf{v}) = v_i - v_{i+m'} = 0.$$

Hence  $\mathcal{M}_{0, m'} \subseteq \bigcap_{1 \leq i \leq k+m} \text{Ker } \alpha_i$ .

Conversely, assume  $\mathbf{v} \in \bigcap_{1 \leq i \leq k+m} \text{Ker } \alpha_i$ , i.e. for all  $i \in \{1, \dots, k + m\}, v_i = v_{i+m'}$ .

In order to prove that  $\mathbf{v} \in \mathcal{M}_{0, m'}$ , we will prove that for all  $j \geq k + m + 1, v_j = v_{j+m'}$  and that  $v_{m'} = 0$ . Given  $j \geq k + m + 1$ , there exists an integer  $j' \in \{k + 1, \dots, k + m\}$  such that  $j \equiv j' \pmod{m}$ . Since  $\mathbf{v} \in \mathcal{M}_{k, m}$ , we have  $v_j = v_{j'}$  and  $v_{j+m'} = v_{j'+m'}$ . Moreover, the fact that  $\mathbf{v} \in \text{Ker } \alpha_{j'}$  implies that  $v_{j'} = v_{j'+m'}$ . Thus

$$v_j = v_{j'} = v_{j'+m'} = v_{j+m'}.$$

In order to conclude, we need to show that  $v_{m'} = 0$ . Note that the previous argument shows that  $\mathbf{v}$  is a periodic sequence of period dividing  $m'$ . Since  $m'$  divides  $k$  and  $m$ ,

$$v_{m'} = v_k = v_{k+m}.$$

Since  $\mathbf{v} \in \mathcal{M}_{k, m}$ , we have  $\beta v_{k+m} - v_k = 0$  with  $\beta \neq 1$ , whence

$$v_{m'} = v_k = v_{k+m} = 0.$$

□

**Lemma 4.14.** *We have  $L^* \alpha_1 = 0$  and for  $i \geq 2, L^* \alpha_i = \delta_{i-1} \alpha_{i-1}$ .*

*Proof.* According to Lemma 4.8, for all  $i \geq 2$ ,

$$L^* \omega_i = \delta_{i-1} \omega_{i-1} + L^* \omega_1.$$

Hence, if  $i \geq 2$ ,

$$L^* \alpha_i = L^*(\omega_i - \omega_{i+m'}) = \delta_{i-1} \omega_{i-1} - \delta_{i+m'-1} \omega_{i+m'-1}.$$

Since  $\mathbf{z} \in \mathcal{M}_{0, m'}$ , we have  $\delta_{i-1} = dz_{i-1}^{d-1} = dz_{i+m'-1}^{d-1} = \delta_{i+m'-1}$ . Hence

$$L^* \alpha_{i-1} = \delta_{i-1} (\omega_{i-1} - \omega_{i+m'-1}) = \delta_{i-1} \alpha_{i-1}.$$

If  $i = 1$ , since  $m' \geq 1$ , we have  $1 + m' \geq 2$  so that

$$L^* \omega_{1+m'} = \delta_{m'} \omega_{m'} + L^* \omega_1.$$

Hence

$$L^* \alpha_1 = L^*(\omega_1 - \omega_{1+m'}) = -\delta_{m'} \omega_{m'}.$$

Since  $\mathbf{z} \in \mathcal{M}_{0,m'}$ , we have  $z_{m'} = 0$ . Therefore,  $\delta_{m'} = dz_{m'}^{d-1} = 0$  and  $L^* \alpha_1 = 0$ .  $\square$

It follows from Lemma 4.13 and Lemma 4.14 that  $L^*: \mathcal{M}_{0,m'} \rightarrow \mathcal{M}_{0,m'}$  is nilpotent.  $\square$

*Proof of Proposition 4.12 when  $k' \neq 0$ .* In this case, since  $(k', m') \prec (k, m)$ , we have

$$k' = k \text{ and } m = pm' \text{ with } p \geq 2.$$

Let  $\lambda$  be the multiplier of  $P(t) = t^d + z_1$  at  $P^{\circ k}(0)$ . Note that, according to Proposition 4.4,  $\mathbf{z}$  is the critical orbit of  $P$ . Since  $\mathbf{z}$  is preperiodic of preperiod  $k > 0$ , the critical point 0 of  $P$  is preperiodic, i.e.  $\lambda \neq 0$ . We will show that

$$\text{Spec}(L^* : \mathcal{M}_{k,m'}^0 \rightarrow \mathcal{M}_{k,m'}^0) = \{\mu \mid \mu^m = \lambda^p, \mu^{m'} \neq \lambda\}.$$

Given  $j \in \mathbb{Z}/m\mathbb{Z}$ , denote by  $\underline{j}$  the representative of  $j$  in  $\{k+1, \dots, k+m\}$ , define a linear form  $\beta_j: \mathcal{M}_{k,m} \rightarrow \mathbb{C}$  by

$$\beta_j := \omega_{\underline{j}} - \omega_{\underline{j+m'}}.$$

Note that for all  $j \in \mathbb{Z}/m\mathbb{Z}$ ,  $\beta_j: \mathcal{M}_{k,m} \rightarrow \mathbb{C}$  is non trivial. Indeed, for a given  $j \in \mathbb{Z}/m\mathbb{Z}$ , define  $\mathbf{u} \in \mathcal{M}_{k,m}$  by

$$u_i = \begin{cases} 1 & \text{if } i \geq k+1 \text{ and } i \equiv \underline{j} \pmod{m} \\ \frac{1}{\beta} & \text{if } i = k \text{ and } \underline{j} = k+m \\ 0 & \text{otherwise.} \end{cases}$$

Since  $m' < m$ ,  $\beta_j(\mathbf{u}) = u_{\underline{j}} = 1 \neq 0$ .

We will show that these forms span  $\mathcal{M}_{k,m'}^0 \subsetneq \mathcal{M}_{k,m}$  and use them to study the linear map  $L^* : \mathcal{M}_{k,m'}^0 \rightarrow \mathcal{M}_{k,m'}^0$ . The properties we need are provided by the following lemmas.

**Lemma 4.15.** *We have  $\mathcal{M}_{k,m'}^0 = \text{Span}\{\beta_j, j \in \mathbb{Z}/m\mathbb{Z}\}$ .*

*Proof.* By duality, it is equivalent to show that

$$\mathcal{M}_{k,m'} = \bigcap_{j \in \mathbb{Z}/m\mathbb{Z}} \text{Ker } \beta_j.$$

Assume  $\mathbf{v} \in \mathcal{M}_{k,m'}$ . Then for all  $i \geq k+1$ ,  $v_i = v_{i+m'}$ . Since  $\underline{j} \in j$  and  $m'$  divides  $m$ , we have  $\underline{j} \equiv \underline{j+m'} \pmod{m'}$ . Moreover,  $\underline{j} \geq k+1$  and  $\underline{j+m'} \geq k+1$ , whence

$$\beta_j(\mathbf{v}) = v_{\underline{j}} - v_{\underline{j+m'}} = 0.$$

This shows that  $\mathcal{M}_{k,m'} \subseteq \bigcap_{j \in \mathbb{Z}/m\mathbb{Z}} \text{Ker } \beta_j$ .

Conversely, assume  $\mathbf{v} \in \bigcap_{j \in \mathbb{Z}/m\mathbb{Z}} \text{Ker } \beta_j$ . We want to prove that for all integer  $i \geq k+1$ ,  $v_i = v_{i+m'}$  and that  $\beta v_{k+m'} - v_k = 0$ . First, assume  $i \geq k+1$  and let  $\underline{j} \in \mathbb{Z}/m\mathbb{Z}$  be the congruence class of  $i$ . Since  $\underline{j} \in j$ , we have  $i \equiv \underline{j} \pmod{m}$ . Moreover  $\mathbf{v} \in \text{Ker } \beta_j$ , and so

$$v_{\underline{j}} - v_{\underline{j+m'}} = \beta_j(\mathbf{v}) = 0.$$

From the fact that  $\mathbf{v} \in \mathcal{M}_{k,m}$ , we therefore deduce that  $v_i = v_{\underline{j}}$  and  $v_{\underline{j+m'}} = v_{i+m'}$ . Thus,

$$v_i = v_{i+m'}.$$

Second, let us show  $\beta v_{k+m'} - v_k = 0$ . The previous argument shows that  $\mathbf{v}$  is preperiodic of preperiod at most  $k$  and of period dividing  $m'$ . Since  $m'$  divides  $m$ , we deduce that  $v_{k+m'} = v_{k+m}$ . Since  $\mathbf{v} \in \mathcal{M}_{k,m}$ , we have  $\beta v_{k+m} - v_k = 0$ . Thus  $\beta v_{k+m'} - v_k = 0$ .  $\square$

Thus, it is now important to understand the how  $L^*$  acts on  $\{\beta_j, j \in \mathbb{Z}/m\mathbb{Z}\}$ . Given  $\underline{j} \in \mathbb{Z}/m\mathbb{Z}$ , set

$$\sigma_{\underline{j}} = \delta_{\underline{j}} = dz_{\underline{j}}^{d-1}.$$

**Lemma 4.16.** *For  $\underline{j} \in \mathbb{Z}/m\mathbb{Z}$ , we have  $L^*\beta_{\underline{j}} = \sigma_{\underline{j-1}}\beta_{\underline{j-1}}$ .*

*Proof.* For  $\underline{j} \in \mathbb{Z}/m\mathbb{Z}$ , recall that  $\underline{j}$  is the representative of  $j$  in  $\llbracket k+1, k+m \rrbracket$ . Let us first prove that for all  $\underline{j} \in \mathbb{Z}/m\mathbb{Z}$ ,

$$(4.5) \quad L^*\omega_{\underline{j}} = \sigma_{\underline{j-1}}\omega_{\underline{j-1}} + L^*\omega_1.$$

Indeed, if  $\underline{j} = k+1$  then  $\underline{j-1} = k+m$ , whence  $\sigma_{\underline{j-1}}\omega_{\underline{j-1}} = \delta_{k+m}\omega_{k+m}$ . According to Lemma 4.8, we have  $L^*\omega_{k+1} = \delta_k\omega_k + L^*\omega_1$ . Since  $\mathbf{z} \in \mathcal{M}_{k,m}$ , we have

$$\delta_k = dz_k^{d-1} = d\beta^{d-1}z_{k+m}^{d-1} = \beta^{d-1}\delta_{k+m}.$$

Moreover,  $\omega_k = \beta\omega_{k+m}$ , whence  $\delta_k\omega_k = \beta^d\delta_{k+m}\omega_{k+m}$ . Since  $\beta^d = 1$ ,

$$L^*\omega_{k+1} = \delta_k\omega_k + L^*\omega_1 = \delta_{k+m}\omega_{k+m} + L^*\omega_1.$$

If  $\underline{j} \neq k+1$  then  $\underline{j-1} = \underline{j} - 1$ . According to Lemma 4.8, we have

$$L^*\omega_{\underline{j}} = \delta_{\underline{j-1}}\omega_{\underline{j-1}} + L^*\omega_1 = \delta_{\underline{j-1}}\omega_{\underline{j-1}} + L^*\omega_1 = \sigma_{\underline{j-1}}\omega_{\underline{j-1}} + L^*\omega_1.$$

In any case, we have the equality (4.5). Hence

$$L^*\beta_{\underline{j}} = L^*(\omega_{\underline{j}} - \omega_{\underline{j+m'}}) = \sigma_{\underline{j-1}}\omega_{\underline{j-1}} - \sigma_{\underline{j+m'-1}}\omega_{\underline{j+m'-1}}.$$

Note that  $\underline{j-1}$  and  $\underline{j+m'-1}$ , which are congruence modulo  $m'$ , are two integers at least  $k+1$ . Since  $\mathbf{z} \in \mathcal{M}_{k,m'}$ , we have  $\sigma_{\underline{j-1}} = \delta_{\underline{j-1}} = \delta_{\underline{j+m'-1}} = \sigma_{\underline{j+m'-1}}$ . Thus

$$L^*\beta_{\underline{j}} = \sigma_{\underline{j-1}}(\omega_{\underline{j-1}} - \omega_{\underline{j+m'-1}}) = \sigma_{\underline{j-1}}\beta_{\underline{j-1}}. \quad \square$$

Recall that  $\lambda$  is the multiplier of  $P(t) = t^d + z_1$  at the periodic point  $z_{k+1}$  of period  $m' = \frac{m}{p}$ .

**Lemma 4.17.** For all  $j \in \mathbb{Z}/m\mathbb{Z}$ ,

$$(L^*)^{m'}(\beta_j) = \lambda\beta_{j-m'} \quad \text{and} \quad (L^*)^m(\beta_j) = \lambda^p\beta_j.$$

*Proof.* The second equality is the straightforward consequence of the first one. Hence, it is enough to prove the first equality. According to Proposition 4.4,  $\mathbf{z}$  is the critical orbit of the polynomial  $P(t) = t^d + z_1$ , i.e.  $z_i = P^i(0)$ . In particular, for any  $j \in \mathbb{Z}/m\mathbb{Z}$ , we have  $P(z_{j-1}) = z_j$  and  $P'(z_{j-1}) = dz_{j-1}^{d-1} = \delta_{j-1}$ . Since  $\mathbf{z}$  is of type  $(k, m')$ , the multiplier  $\lambda$  of the cycle of  $\underline{z}_j$  is

$$\lambda = \prod_{i \in \llbracket 1, m' \rrbracket} P'(z_{j-i}) = \prod_{i \in \llbracket 1, m' \rrbracket} \delta_{j-i} = \prod_{i \in \llbracket 1, m' \rrbracket} \sigma_{j-i}.$$

Hence, by Lemma 4.16, we have

$$(L^*)^{m'}(\beta_j) = \left( \prod_{i \in \llbracket 1, m' \rrbracket} \sigma_{j-i} \right) \beta_{j-m'} = \lambda\beta_{j-m'},$$

and

$$(L^*)^m(\beta_j) = (L^*)^{m'p}(\beta_j) = \lambda^p\beta_j. \quad \square$$

Let  $\nu$  be a  $m'$ -th root of  $\lambda$ . Set

$$T = \frac{L^*}{\nu} : \mathcal{M}_{k, m'}^0 \rightarrow \mathcal{M}_{k, m'}^0.$$

We shall prove that  $T$  is diagonalizable with simple eigenvalues and the eigenvalues of  $T$  are  $m'$ -th roots of unity except 1. According to Lemma 4.17, for all  $j \in \mathbb{Z}/m\mathbb{Z}$ ,  $T^{m'}(\beta_j) = \beta_{j-m'}$ . In addition,

$$\begin{aligned} \sum_{n \in m'\mathbb{Z}/m\mathbb{Z}} \beta_n &= \sum_{n \in m'\mathbb{Z}/m\mathbb{Z}} (\omega_n - \omega_{n+m'}) = 0 \\ &= \sum_{n \in m'\mathbb{Z}/m\mathbb{Z}} \omega_n - \sum_{n \in m'\mathbb{Z}/m\mathbb{Z}} \omega_{n+m'} = 0. \end{aligned}$$

Recall that  $p = \frac{m}{m'}$ . Hence

$$\beta_0 + T^{m'}(\beta_0) + \dots + T^{m'(p-1)}(\beta_0) = 0$$

Applying  $m' - 1$  times  $T$  and adding the results, we deduce that

$$\beta_0 + T(\beta_0) + T^2(\beta_0) + \dots + T^{m'p-1}(\beta_0) = 0$$

According to Lemma 4.15 and Lemma 4.16, the set  $\{\beta_0, L^*(\beta_0), (L^*)^2(\beta_0), \dots\}$  generates  $\mathcal{M}_{k, m'}^0$ . Hence  $\{\beta_0, T(\beta_0), T^2(\beta_0), \dots\}$  also generates  $\mathcal{M}_{k, m'}^0$ . Therefore,

$$(4.6) \quad \text{Id} + T + T^2 + \dots + T^{m-1} = 0$$

This means that the minimal polynomial of  $T$  divides the polynomial  $1 + X + X^2 + \dots + X^{m-1}$ . Consequently,  $T$  is diagonalizable and the eigenvalues of  $T$  are

roots of unity which are not 1. We now show that  $T$  has only simple eigenvalues. Assume  $\zeta \in \text{Spec } T$ . Let  $v \in \mathcal{M}_{k,m'}$  be an eigenvector associated to  $\zeta$ . Set

$$H_\zeta = \frac{1}{m} \left( \text{Id} + \frac{T}{\zeta} + \dots + \frac{T^{m-1}}{\zeta^{m-1}} \right).$$

The equality (4.6) implies that  $T^m = \text{Id}$ . Additionally,  $\zeta^m = 1$ . Hence

$$H_\zeta \circ \frac{T}{\zeta} = H_\zeta \quad \text{so that} \quad H_\zeta \circ T^j = \zeta^j H_\zeta \quad \forall j \geq 1.$$

In addition,  $\{\beta_0, T(\beta_0), \dots\}$  generates  $\mathcal{M}_{k,m'}^0$ , hence

$$\text{Im } H_\zeta \subseteq \text{Span } H_\zeta(\beta_0).$$

Note that  $H_\zeta(v) = v$ . Hence

$$v \subseteq \text{Im } H_\zeta \subseteq \text{Span } H_\zeta(\beta_0).$$

Thus the eigenspace associated to the eigenvalue  $\zeta$  of  $T$  has dimension 1, i.e.  $T$  has only simple eigenvalues.

Since  $T = \frac{L^*}{\nu}$ ,  $L^*$  is diagonalizable with simple eigenvalues which are  $m$ -th roots of  $\nu^m = \lambda^p$ . In addition, 1 is not an eigenvalue of  $T$  hence  $\nu$  is not an eigenvalue of  $L^*$ . Since  $\nu$  is an arbitrary  $m'$ -th root of  $\lambda$ , we deduce that

$$\text{Spec } L^* \subseteq \{\mu^m = \lambda^p, \mu^{m'} \neq \lambda\}.$$

Since  $L^*$  has only simple eigenvalues,  $\#\text{Spec } L^* = \dim \mathcal{M}_{k,m'}^0 = m - m'$ . Hence

$$\text{Spec}(L^* : \mathcal{M}_{k,m'}^0 \rightarrow \mathcal{M}_{k,m'}^0) = \{\mu^m = \lambda^p, \mu^{m'} \neq \lambda\} \quad \square$$

*Proof of Theorem A'.* According to Lemma 3.1,  $F_{k,m}$  and  $G_{k,m}$  are conjugate. Hence it is enough to consider an eigenvalue  $\mu$  of  $F_{k,m}$  at a fixed point  $z \in \mathcal{M}_{k,m}$ . Denote by  $(k', m')$  the exact type of  $z$ . Then by Proposition 4.4, we have  $(k', m') \preceq (k, m)$ , or by Definition 3.2, we have

$$m' \mid m, \text{ (either } k' = 0 \text{ or } k' = k, m' \mid k).$$

Regarding  $\mu$  as an eigenvalue of  $F_{k,m}$  at  $z$ . We recall argument at the beginning of 4.2.4, according to Lemma 4.6,  $(k', m') \preceq (k, m)$  and, by Proposition 3.3,  $\mathcal{M}_{k',m'} \subseteq \mathcal{M}_{k,m}$  is invariant under  $F_{k,m}$ . Since  $\mathcal{M}_{k',m'}$  is invariant under  $D_z F_{k,m}$ , the annihilator  $\mathcal{M}_{k',m'}^0$  of  $\mathcal{M}_{k',m'}$  in  $\mathcal{M}_{k,m}$  is invariant under the transpose  $L^*$  of  $D_z F_{k,m}$  and we have the following decomposition

$$(4.7) \quad \text{Spec } L = \text{Spec}(L|_{\mathcal{M}_{k',m'}}) \cup \text{Spec} \left( L^*|_{\mathcal{M}_{k',m'}^0} \right).$$

Moreover, according to Proposition 3.3, we have

$$L|_{\mathcal{M}_{k',m'}} = D_z F_{k',m'}.$$

If  $\mu \in \text{Spec}(L|_{\mathcal{M}_{k',m'}})$ , note that  $L|_{\mathcal{M}_{k',m'}} = D_z F_{k',m'}$  and  $z \notin PC_{k',m'}$ , thus  $\mu$  is an eigenvalue of  $G_{k',m'}$  at a fixed point outside its post-critical set.

If  $\mu \in \text{Spec} \left( L^*|_{\mathcal{M}_{k',m'}^0} \right)$ , then we are done by Proposition 4.12.  $\square$

## REFERENCES

- [Ast20] Matthieu Astorg. Dynamics of post-critically finite maps in higher dimension. *Ergodic Theory and Dynamical Systems*, 40(2):289–308, 2020.
- [BEK20] Xavier Buff, Adam L. Epstein, and Sarah Koch. Eigenvalues of the thurston operator. *Journal of Topology*, 13(3):969–1002, 2020.
- [DH93] Adrien Douady and John H Hubbard. A proof of thurston’s topological characterization of rational functions. *Acta Mathematica*, 171(2):263–297, 1993.
- [GV19] Thomas Gauthier and Gabriel Vigny. The geometric dynamical northcott and bogomolov properties. *arXiv preprint arXiv:1912.07907*, 2019.
- [IRS19] Patrick Ingram, Rohini Ramadas, and Joseph H Silverman. Post-critically finite maps on  $\mathbb{P}^n$  for  $n \geq 2$  are sparse. *arXiv preprint arXiv:1910.11290*, 2019.
- [Ji20] Zhuchao Ji. Structure of julia sets for post-critically finite endomorphisms on  $\mathbb{P}^2$ . *arXiv preprint arXiv:2010.11094*, 2020.
- [Koc08] Sarah Colleen Koch. *A new link between Teichmüller theory and complex dynamics*. PhD thesis, Cornell University, 2008.
- [Koc13] Sarah Koch. Teichmüller theory and critically finite endomorphisms. *Advances in Mathematics*, 248:573–617, 2013.
- [Le20a] Van Tu Le. *Dynamique des endomorphismes post-critiquement algébriques*. PhD thesis, Université de Toulouse, Université Toulouse III-Paul Sabatier, 2020.
- [Le20b] Van Tu Le. Periodic points of post-critically algebraic holomorphic endomorphisms. *Ergodic Theory and Dynamical Systems*, pages 1–33, 2020.
- [Mil11] John Milnor. *Dynamics in One Complex Variable. (AM-160):(AM-160)*. Princeton University Press, 2011.
- [Mil14] John Milnor. Tsujii’s monotonicity proof for real quadratic maps. *Collected Papers of John Milnor: VII. Dynamical Systems (1984-2012)*, page 39, 2014.
- [Ron08] Feng Rong. The fatou set for critically finite maps. *Proceedings of the American Mathematical Society*, 136(10):3621–3625, 2008.

VAN TU LE, DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DEGLI STUDI DI ROMA "TOR VERGATA", VIA DELLA RICERCA SCIENTIFICA 1 - 00133 ROMA, ITALY

*Email address:* levantu.hp@gmail.com